

GLOBAL CLASSICAL SOLUTIONS TO THE IONIC VLASOV-POISSON-BOLTZMANN SYSTEM NEAR MAXWELLIANS

FUCAI LI AND YICHUN WANG

ABSTRACT. In a plasma, the ionic Vlasov-Poisson-Boltzmann system models the evolution of ions interacting with themselves through the self-consistent electrostatic potential and collisions. It distinguishes the electric Vlasov-Poisson-Boltzmann system via an extra exponential nonlinearity in the coupled Poisson-Poincaré equation which creates some essential mathematical difficulties. Despite its physical importance, the global well-posedness to this system remains completely open. This gap is filled in this article in the three dimensional period box case. We show that any smooth, small initial perturbation of a global Maxwellian satisfying the conservations of mass, momentum and energy leads to a unique global-in-time classical solution with an exponential decay for the whole range of collision potential $r \in (-3, 1]$. The construction relies on the nonlinear energy method, new nonlinear elliptic regularity estimates and coercivity inequality of the linearized collision operator \mathcal{L} novelly deduced for the ion model.

1. INTRODUCTION

At the kinetic level, the dynamics of dilute ions can be described by the so-called ionic Vlasov-Poisson-Boltzmann system in absence of magnetic field, where the ions interact with themselves through the self-consistent electric field and collisions. Due to the analytical difficulties raising from the nonlinear Poisson equation, until now there has been no rigorous mathematical study on the well-posedness problem of this model. The purpose of this article is to fill the gap in the 3D periodic box case. First, for reader's convenience, we provide some necessary backgrounds on the ion dynamics below.

1.1. Background and models. Plasma exists commonly in the universe as a state of matter, which can be found, for example, in lightning, ionosphere, corona, accretion disk and interstellar medium [33,35,36]. It is a multi-particle system consisting of electrons and ions whose global dynamical behaviors are mainly dominated by electromagnetic fields. Let us take the stellar atmosphere plasmas as an example which constitute electrons, protons, helium ions and few heavy ions [31]. In physical literature, plasma kinetic theory is usually based on the statistical mechanics viewpoints which lies between the single particle orbital theory and the magnetohydrodynamic theory.

The collisions among charged particles are a kind of electromagnetic interactions. It is extensively considered that a charged particle always interacts with the others within a ball centered on itself with a radius of the *Debye length*, which refers to be the well-known Coulomb collision. In this case, the collision frequencies of electron-electron (ee), electron-ion (ei) and ion-ion (ii) can be presented as (for example, see [35]):

$$\begin{cases} f_{ee} = \frac{n_e e^4 \ln \Lambda}{3\sqrt{6}\pi \varepsilon_0^2 m_e^{\frac{1}{2}} (k_B T_e)^{\frac{3}{2}}}, \\ f_{ei} = \frac{n_e Z_i^2 e^4 \ln \Lambda}{6\sqrt{3}\pi \varepsilon_0^2 m_e^{\frac{1}{2}} (k_B T_e)^{\frac{3}{2}}}, \\ f_{ii} = \frac{n_i Z_i^4 e^4 \ln \Lambda}{3\sqrt{6}\pi \varepsilon_0^2 m_i^{\frac{1}{2}} (k_B T_i)^{\frac{3}{2}}}. \end{cases}$$

Here, for the electrons, n_e is the number density, e and m_e are the magnitude of the charge and mass, and T_e is the temperature. Similarly, we have n_i , $Z_i e$, m_i and T_i for the (one-specie) ions. The parameter $\varepsilon_0 > 0$ is the vacuum

2020 *Mathematics Subject Classification.* 82D10, 35Q20, 35Q83.

Key words and phrases. Ionic Vlasov-Poisson-Boltzmann system, Poisson-Poincaré equation, Global classical solution, exponential decay.

permittivity, $k_B > 0$ is the Boltzmann constant and $\ln \Lambda$ is the Coulomb logarithm. From the representations of f_{ee} , f_{ei} and f_{ii} , it holds that

$$\frac{f_{ee}}{f_{ii}} = \left(\frac{m_i}{m_e}\right)^{1/2} \left(\frac{T_i}{T_e}\right)^{3/2}, \quad f_{ee} \approx f_{ei}.$$

Since the fact that m_i is much larger than m_e , the collision frequencies f_{ee} and f_{ei} are drastically faster than f_{ii} . Therefore, under the electrostatic approximation, the timescales of electrons and ions are naturally separated. On one hand, on the electron's timescale, the distribution of ions is regarded as the positive charged constant background. On the other hand, on the ion's timescale, the electrons are at the thermal equilibrium state and the number density n_e obeys the well-known Boltzmann distribution:

$$n_e = n_{e0} \exp\left\{\frac{e\phi}{k_B T_e}\right\},$$

where ϕ is the electrostatic potential and n_{e0} is the number density of electrons as $\phi = 0$. Then ϕ satisfies the Poisson equation

$$\varepsilon_0 \Delta \phi = -(\rho_i + \rho_e) = en_{e0} \exp\left\{\frac{e\phi}{k_B T_e}\right\} - \rho_i,$$

where ρ_e and ρ_i are the charge density of electrons and ions respectively, and $\rho_e = -en_e$, $\rho_i = Z_i en_i$. Normalization of the parameters yields the so-called Poisson-Poincaré equation:

$$\Delta \phi = e^\phi - \rho, \tag{1.1}$$

which is a semilinear second order elliptic equation for ϕ .

By physical references, the *grazing collision* is dominant among the Coulomb collision in the contributions of momentum and energy transfer. When a particle deviates for 90° after several grazing collisions, the energy loss is approximated to be

$$\frac{\delta E}{E_0} \approx \frac{2m_1 m_2}{(m_1 + m_2)^2},$$

where m_1 and m_2 are the masses of two-species particles respectively, δE is the transferred energy, and E_0 is the initial energy. This means that, $\frac{\delta E}{E_0} \approx \frac{1}{2}$ for ee-collision or ii-collision, while $\frac{\delta E}{E_0} \approx 2\frac{m_e}{m_i} \ll 1$ for ei-collision. Therefore, the relaxation of electrons or ions to the thermal equilibriums is easy to reach through the ee-collisions or ii-collisions, but the equilibrium between electrons and ions is difficult to reach via the ei-collisions. Hence we only consider the collisions within ions when we study the relaxation mechanism of the ion dynamics.

In this article, we concentrate on the evolution of ion particles at the kinetic level. For the simplified collision model, it can be described by the so-called *ionic Vlasov-Poisson-Boltzmann system*. More precisely, it consists of a Vlasov-Boltzmann equation whose unknown is the distribution function $F(t, x, v)$ of particles and a Poisson-Poincaré equation whose unknown is the self-consistent potential $\phi(t, x)$, reading as

$$\begin{cases} \partial_t F + v \cdot \nabla_x F + E \cdot \nabla_v F = Q(F, F) & \text{in } (0, \infty) \times \mathbb{T}^3 \times \mathbb{R}^3, \\ E = -\nabla \phi, \quad \Delta \phi = e^\phi - \rho & \text{in } (0, \infty) \times \mathbb{T}^3, \\ F|_{t=0} = F_0 & \text{in } \mathbb{T}^3 \times \mathbb{R}^3, \end{cases} \tag{1.2}$$

where the position $x \in \mathbb{T}^3$ and the velocity $v \in \mathbb{R}^3$ at time $t > 0$, and $\rho = \int_{\mathbb{R}^3} F dv$ is the macroscopic density of ions. The initial distribution F_0 is assumed to be nonnegative. The Boltzmann collision term $Q(F, F)$ on the right-hand side of (1.2)₁ is given as the bilinear form

$$\begin{aligned} Q(F_1, F_2) &= \iint_{\mathbb{R}^3 \times \mathbb{S}^2} B(v - u, \theta) [F_1(u') F_2(v') - F_1(u) F_2(v)] d\omega du \\ &=: Q_{\text{gain}}(F_1, F_2) - Q_{\text{loss}}(F_1, F_2), \end{aligned}$$

where the relationship between the pre-collision velocity pair (u, v) of two particles and the post-collision velocity pair (u', v') obeys

$$u' = u + [(v - u) \cdot \omega]\omega, \quad v' = v - [(v - u) \cdot \omega]\omega,$$

for $\omega \in \mathbb{S}^2$, which can be determined by the conservation laws of momentum and energy, i.e.

$$u + v = u' + v', \quad |u|^2 + |v|^2 = |u'|^2 + |v'|^2.$$

The Boltzmann collision kernel $B(v - u, \theta)$ depends only on the relative velocity $|v - u|$ and the angular deviation θ with $\cos \theta = (v - u) \cdot \omega / |v - u|$. Specifically,

$$B(v - u, \theta) = \underbrace{|v - u|^r}_{\text{kinetic part}} \underbrace{b\left(\frac{(v - u) \cdot \omega}{|v - u|}\right)}_{\text{angular part}}.$$

Throughout this article, we consider the whole range of inverse power law $r \in (-3, 1]$ (which includes hard potentials, soft potentials and Maxwell molecules cases) with an angular cutoff assumption:

$$0 \leq b(\cos \theta) \leq C|\cos \theta|$$

for some positive constant $C > 0$. For notation simplification, we shall denote $b_0(\theta) := b(\cos \theta)$.

Formally, the system (1.2) can be derived from the following non-dimensional two-species (electron and ion) Vlasov-Poisson-Boltzmann system

$$\begin{cases} \partial_t f_- + w \cdot \nabla_x f_- - \frac{q_e}{m_e} E \cdot \nabla_w f_- = Q_-(f_-, f_-), \\ \partial_t f_+ + v \cdot \nabla_x f_+ - \frac{q_i}{m_i} E \cdot \nabla_v f_+ = Q_+(f_+, f_+), \\ \varepsilon_0 \nabla \cdot E = \int_{\mathbb{R}^3} \{q_i f_+(v) - q_e f_-(v)\} dv, \end{cases}$$

by taking the limit $\sqrt{\frac{m_e}{m_i}} \rightarrow 0$. This limiting process is usually referred to as *massless electron limit*. We highlight the work [2] for the discussion of the formal derivations.

For the classical solutions to the system (1.2), the following conservation laws of mass, momentum and energy hold:

$$\begin{aligned} \frac{d}{dt} \iint_{\mathbb{R}^3 \times \mathbb{S}^2} F(t, x, v) dx dv &= 0, \\ \frac{d}{dt} \iint_{\mathbb{R}^3 \times \mathbb{S}^2} v F(t, x, v) dx dv &= 0, \\ \frac{d}{dt} \left\{ \iint_{\mathbb{R}^3 \times \mathbb{S}^2} |v|^2 F(t, x, v) dx dv + \int_{\mathbb{T}^3} 2\phi e^\phi + |\nabla \phi|^2 dx \right\} &= 0. \end{aligned}$$

1.2. Relevant literature. Over the past several decades, there has been a significant amount of mathematical advances on the well-posedness theory of plasma kinetic equations of Vlasov-Poisson type, mostly focusing on the electron dynamics. On the Vlasov-Poisson system for electrons, a series of works [22, 23, 29, 30, 32] proved the global existence and uniqueness of the classical solutions in domains without boundary, [6, 24, 25] studied the same problem considering the boundary effects, and [1, 23, 34] investigated the long time behaviors of the classical solutions.

Taking the collisions between electrons into account, the Vlasov-Poisson-Boltzmann system for electrons has been extensively studied. Basically, in the L^2 framework near equilibrium, Guo [17], Yang-Zhao [40] and Yang-Yu-Zhao [39] established the classical solutions in \mathbb{T}^3 and \mathbb{R}^3 , respectively. Duan-Strain [9] studied the optimal time decay of classical solutions in \mathbb{R}^3 . Li-Yang-Zhong [27, 28] used the spectral analysis and Green's function to investigate the refined properties of the classical solutions in \mathbb{R}^3 . The above works are confined in hard-sphere collision kernel case. Meanwhile, for the hard and soft potentials with angular cutoff, Duan-Yang-Zhao [10, 11] first dealt with the hard potentials and moderately soft potentials cases, and then Xiao-Xiong-Zhao [38] established the well-posedness result in the case of very soft potentials, see also [37] for the non-cutoff hard potentials case.

If the interactions between particles and boundaries are considered simultaneously, the L^2 - L^∞ method developed by Guo [20] is involved. In term of this framework, Cao-Kim-Lee [4] constructed global strong solutions for hard-sphere case with homogenous Neumann condition for the electric potential and the diffuse boundary condition for the particle distribution functions. See also [26] for soft potentials case with homogenous Neumann boundary condition for the electric potential and the incoming boundary condition for the particle distribution functions.

In contrast, for the Vlasov-Poisson type equations of ion dynamics, the mathematical research is less due to the additional exponential nonlinearity in the Poisson-Poincaré equation (1.1), which is the key difference between the ionic and electronic Vlasov-Poisson type systems. Bouchut [3] first constructed the global weak solutions to the ionic Vlasov-Poisson system in \mathbb{R}^3 . In addition, Bardos, et al. [2] proved the existence and uniqueness of global weak solutions with the conservation of energy to the ionic Vlasov-Poisson system in both a bounded domain $\Omega \subset \mathbb{R}^2$ and \mathbb{T}^3 . For classical solutions, global well-posedness results of the ionic Vlasov-Poisson system have been established in \mathbb{T}^2 and in \mathbb{T}^3 (see [15]), and in \mathbb{R}^3 (see [14]) by Griffin-Pickering and Iacobelli, and more recently in a bounded domain $\Omega \subset \mathbb{R}^3$ by Cesbron-Iacobelli [6]. If taking the ion-ion collisions into account, rigorous results are lack so far. Notice however that Flynn [12] proved the local well-posedness of the ionic Vlasov-Poisson-Landau system in \mathbb{T}^3 for large initial data.

To the best of our knowledge, the global well-posedness result on the ionic Vlasov-Poisson-Boltzmann system (1.2)₁–(1.2)₂ is missing from literature. The goal of this article is to establish the first result on global classical solutions to the problem (1.2) near Maxwellians in \mathbb{T}^3 , which will be proposed in what follows.

1.3. Perturbed equations near the equilibrium. In this article, we shall construct a classical solution $F(t, x, v)$ to the ionic Vlasov-Poisson-Boltzmann system (1.2)₁–(1.2)₂ near the global Maxwellian

$$\mu \equiv \mu(v) = \frac{1}{(2\pi)^{\frac{3}{2}}} \exp\left\{-\frac{|v|^2}{2}\right\}.$$

For this purpose, we define the standard perturbation $f(t, x, v)$ to μ as

$$F = \mu + \sqrt{\mu}f,$$

and the initial datum $f_0(x, v)$ as $F_0 = \mu + \sqrt{\mu}f_0 \geq 0$.

We rewrite the problem (1.2) by using the unknown f . Thus, the ionic Vlasov-Poisson-Boltzmann system now becomes

$$\{\partial_t + v \cdot \nabla_x + E \cdot \nabla_v\} f - E \cdot v \sqrt{\mu} - \frac{v}{2} \cdot E f = -\mathcal{L}f + \Gamma(f, f), \quad (1.3)$$

coupled with

$$\begin{cases} E = -\nabla\phi, \\ \Delta\phi = e^\phi - \int_{\mathbb{R}^3} \sqrt{\mu}f dv - 1, \end{cases} \quad (1.4)$$

and supplemented with the initial datum

$$f(0, x, v) = f_0(x, v). \quad (1.5)$$

Here, \mathcal{L} is the linearized Boltzmann collision operator defined as

$$\mathcal{L}f = \nu f - \mathcal{K}f,$$

where $\nu(v) = c \int_{\mathbb{R}^3} |v - u|^r \mu(u) du$, and \mathcal{K} is split into $\mathcal{K} := \mathcal{K}_2 - \mathcal{K}_1$ with

$$\begin{aligned} \mathcal{K}_1 f(v) &:= \iint_{\mathbb{R}^3 \times \mathbb{S}^2} |v - u|^r b_0(\theta) \mu^{1/2}(u) \mu^{1/2}(v) f(u) dud\omega, \\ \mathcal{K}_2 f(v) &:= \iint_{\mathbb{R}^3 \times \mathbb{S}^2} |v - u|^r b_0(\theta) \mu^{1/2}(u) \{\mu^{1/2}(u')g(v') + \mu^{1/2}(v')g(u')\} dud\omega. \end{aligned}$$

We also define the bilinear collision operator $\Gamma(\cdot, \cdot)$ as

$$\Gamma(g_1, g_2)(v) := \mu^{-1/2}(v) Q(\mu^{1/2}g_1, \mu^{1/2}g_2)(v)$$

$$\begin{aligned}
&= \iint_{\mathbb{R}^3 \times \mathbb{S}^2} |u-v|^r b_0(\theta) \mu^{1/2}(u) g_1(u') g_2(v') dud\omega \\
&\quad - \iint_{\mathbb{R}^3 \times \mathbb{S}^2} |u-v|^r b_0(\theta) \mu^{1/2}(u) g_1(u) g_2(v) dud\omega \\
&=: \Gamma_{\text{gain}}(g_1, g_2)(v) - \Gamma_{\text{loss}}(g_1, g_2)(v).
\end{aligned}$$

By assuming that $[F_0, \phi_{F_0}]$ has the same mass, momentum and total energy as the steady state $[\mu, 0]$, we can then rewrite the conservation laws in terms of $[f, \phi_f]$ as

$$\iint_{\mathbb{T}^3 \times \mathbb{R}^3} f(t, x, v) \sqrt{\mu(v)} dx dv \equiv 0, \quad (1.6)$$

$$\iint_{\mathbb{T}^3 \times \mathbb{R}^3} v f(t, x, v) \sqrt{\mu(v)} dx dv \equiv 0, \quad (1.7)$$

$$\iint_{\mathbb{T}^3 \times \mathbb{R}^3} \frac{1}{2} |v|^2 f(t, x, v) \sqrt{\mu(v)} dx dv + \int_{\mathbb{T}^3} \phi_f e^{\phi_f} dx + \int_{\mathbb{T}^3} \frac{1}{2} |\nabla \phi_f|^2 dx \equiv 0. \quad (1.8)$$

Here we have used the notations ϕ_{F_0} and ϕ_f to emphasise that ϕ depends on F_0 and f , respectively.

1.4. Notations and energy functional. We use (\cdot, \cdot) to denote the standard L^2 inner product in $\mathbb{T}_x^3 \times \mathbb{R}_v^3$ for a pair of functions $g, h \in L^2(\mathbb{T}_x^3 \times \mathbb{R}_v^3)$ and $\langle \cdot, \cdot \rangle$ to denote the L^2 product in \mathbb{R}_v^3 . In the sequel, we use $\|\cdot\|$ to denote L^2 norms in $\mathbb{T}_x^3 \times \mathbb{R}_v^3$ or \mathbb{T}_x^3 , and $|\cdot|$ in \mathbb{R}_v^3 unless otherwise specified, as in Section 3. We also denote

$$(h_1, h_2)_\nu := (\nu(v) h_1, h_2)$$

to be the weighted L^2 inner product in $\mathbb{T}_x^3 \times \mathbb{R}_v^3$, and $\|\cdot\|_\nu$ to be the corresponding weighted L^2 norm.

Let the multi-indices γ and β be

$$\gamma = [\gamma_0, \gamma_1, \gamma_2, \gamma_3], \quad \beta = [\beta_1, \beta_2, \beta_3],$$

and

$$\partial_\beta^\gamma := \partial_t^{\gamma_0} \partial_{x_1}^{\gamma_1} \partial_{x_2}^{\gamma_2} \partial_{x_3}^{\gamma_3} \partial_{v_1}^{\beta_1} \partial_{v_2}^{\beta_2} \partial_{v_3}^{\beta_3}.$$

We denote $\tilde{\alpha} \leq \alpha$ if each component of $\tilde{\alpha}$ is not greater than that of α 's, while $\tilde{\alpha} < \alpha$ means that $\tilde{\alpha} \leq \alpha$ and $|\tilde{\alpha}| < |\alpha|$.

For a fixed $N \in \mathbb{N}$, we define

$$\begin{aligned}
\|f\| &:= \sum_{|\gamma|+|\beta| \leq N} \|\partial_\beta^\gamma f(t)\|, \\
\|f\|_\nu &:= \sum_{|\gamma|+|\beta| \leq N} \|\nu^{\frac{1}{2}} \partial_\beta^\gamma f(t)\|, \\
\|f\|_{\langle v \rangle} &:= \sum_{|\gamma|+|\beta| \leq N} \|\langle v \rangle^{\frac{1}{2}} \partial_\beta^\gamma f(t)\|,
\end{aligned}$$

where $\langle v \rangle = (1 + |v|^2)^{1/2}$. We also define the high order energy functional for $f(t, x, v)$ as

$$\mathcal{E}(t) := \frac{1}{2} \|f\|^2(t) + \int_0^t \|f\|_{\langle v \rangle}^2(s) ds. \quad (1.9)$$

Given the initial datum $f_0(x, v)$, we define the initial energy functional:

$$\mathcal{E}(0) := \frac{1}{2} \sum_{|\gamma|+|\beta| \leq N} \|\partial_\beta^\gamma f_0\|^2 \triangleq \frac{1}{2} \|f_0\|^2,$$

where the temporal derivatives of f_0 are defined naturally through (1.3)–(1.4).

Throughout this article, the constant $C > 0$ in the relationship $a \leq Cb$ may be different with each other and depend on a variable collection of parameters in the context. This notation is used mostly inside the proofs of lemmas, propositions and theorems. The symbol $a \sim b$ means that there exist two constants $c, C > 0$ such that $cb \leq a \leq Cb$.

1.5. Main result. We state our main result as follows.

Theorem 1.1. *Let $N \geq 8$ be an integer. Assume that $F_0(x, v) = \mu(v) + \sqrt{\mu(v)}f_0(x, v) \geq 0$ and f_0 satisfies the conservation laws (1.6)–(1.8). Then, there exists a $\varkappa > 0$ such that if*

$$\mathcal{E}(0) \leq \varkappa,$$

the ionic Vlasov-Poisson-Boltzmann system (1.3)–(1.4) with the initial datum (1.5) has a unique global classical solution $f(t, x, v)$ satisfying $F(t, x, v) = \mu(v) + \sqrt{\mu(v)}f(t, x, v) \geq 0$, and

$$\sup_{0 \leq t \leq \infty} \mathcal{E}(t) \leq C_0 \mathcal{E}(0)$$

for some constant $C_0 > 0$. Moreover, there exist two constants $C_1, C_2 > 0$ such that

$$\|f\|^2(t) \leq C_1 e^{-C_2 t} \mathcal{E}(0).$$

Remark 1.2. *The condition $N \geq 8$ in Theorem 1.1 is required in the proof of Lemma 2.4 for the soft potentials collision kernel case $r \in (-3, 0)$. Meanwhile, $N \geq 4$ is sufficient to deal with the hard potentials case $r \in [0, 1]$.*

1.6. Key points of the proof. For reader's conveniences, we now explain our main ideas and key points developed to the proof of Theorem 1.1.

1.6.1. Nonlinear energy method. Our main strategy aimed to the nonlinear stability of global Maxwellians for the ionic Vlasov-Poisson-Boltzmann system (1.3)–(1.4) is based on the nonlinear L^2 energy method first devised by Guo [16]. It has inspired many subsequent works, such as [17] for the electronic Vlasov-Poisson-Boltzmann system, [18] for the Vlasov-Maxwell-Boltzmann system, and [7, 21] for the Vlasov-Poisson-Landau system.

This method can be applied to many kinetic models of the form

$$\partial_t f + v \cdot \nabla_x f + \mathcal{L}f = \Gamma(f, f). \quad (1.10)$$

The main idea is to find an energy norm $\|\cdot\|_{\mathcal{E}}$, a dissipation norm $\|\cdot\|_{\mathcal{D}}$ and a suitable constant $\delta > 0$ such that

$$\frac{d}{dt} \|f\|_{\mathcal{E}}^2 + \delta \|f\|_{\mathcal{D}}^2 \leq 0. \quad (1.11)$$

The choice of the energy and dissipation norms is delicate. For the model (1.3) in hard sphere case, the energy norm is actually $\|\cdot\|$, and the dissipation norm is actually $\|\cdot\|_{\nu}$ which comes from the linear part of the Boltzmann collision operator $\mathcal{L}f = \nu f - \mathcal{K}f$, and satisfies $|\langle f, \Gamma(f, f) \rangle_{\mathcal{E}}| \leq C \|f\|_{\mathcal{E}} \|f\|_{\mathcal{D}}^2$ by Lemma 2.4. Then (1.11) can be deduced from a suitable small assumption of $\|f\|_{\mathcal{E}}$ in Section 6, combining with the positivity of \mathcal{L} in Section 5. Moreover, noting that the norm $\|\cdot\|_{\mathcal{E}}$ is weaker than $\|\cdot\|_{\mathcal{D}}$, then the solution f decays exponentially in $\|\cdot\|_{\mathcal{E}}$ by (1.11) and Gronwall's inequality.

1.6.2. Elliptic estimates of the Poisson-Poincaré equation (1.1). To study the Poisson-Poincaré equation (1.1) in \mathbb{T}^3 , we follow [15] to decompose the electric potential ϕ into the form $\phi = \bar{\phi} + \hat{\phi}$ satisfying

$$\Delta \bar{\phi} = 1 - \rho, \quad \Delta \hat{\phi} = e^{\bar{\phi} + \hat{\phi}} - 1.$$

Similar to [15], we invoke calculus of variations, Newtonian potential, Poincaré's inequality and Sobolev's embedding to obtain the existence and uniqueness of ϕ to (1.1). To establish the L^2 estimates of ϕ in terms of f , a crucial observation is that $\phi \in H^2(\mathbb{T}^3)$ whenever

$$\rho := \int_{\mathbb{R}^3} \sqrt{\mu} f dv + 1 \in L^2(\mathbb{T}^3).$$

It follows that $\phi \in C(\mathbb{T}^3) \cap L^\infty(\mathbb{T}^3)$ and thus e^{ϕ} possesses positive upper and lower bounds. By mean value theorem,

$$\Delta \phi = e^{\phi} - 1 + 1 - \rho = \phi e^{\phi_\xi} - \int_{\mathbb{R}^3} \sqrt{\mu} f dv,$$

hence we use the standard linear elliptic estimates to get

$$\|\phi\|_{L^\infty(\mathbb{T}^3)} \leq C \|\phi\|_{H^2(\mathbb{T}^3)} \leq C e^{2\|\phi\|_{L^\infty(\mathbb{T}^3)}} \|f\|_{L^2(\mathbb{T}^3 \times \mathbb{R}^3)}.$$

Since $\|f\|_{L^2(\mathbb{T}^3 \times \mathbb{R}^3)}$ is controlled by $\|f\|_{\mathcal{E}}$ which is assumed to be sufficiently small, we know that $\|\phi\|_{L^\infty(\mathbb{T}^3)}$ is also small. Therefore, we obtain $\|\phi\|_{H^2(\mathbb{T}^3)} \leq C\|f\|_{L^2(\mathbb{T}^3 \times \mathbb{R}^3)}$, which is quite analogous to the case of electronic linear Poisson equation. The stability of (1.1) also follows, so the well-posedness of (1.1) is proven.

Furthermore, taking ∂^γ derivatives to (1.1) yields

$$\Delta \partial^\gamma \phi - e^\phi \partial^\gamma \phi = S - \int_{\mathbb{R}^3} \sqrt{\mu} \partial^\gamma f dv,$$

where S is constituted by lower order derivatives of ϕ . Inductively we have

$$\sum_{|\gamma| \leq N} \|\partial^\gamma \phi\|_{H^2(\mathbb{T}^3)} \leq C \sum_{|\gamma| \leq N} \|\partial^\gamma f\|_{L^2(\mathbb{T}^3 \times \mathbb{R}^3)} \leq C\|f\|_{\mathcal{E}}.$$

The above L^2 estimates for the Poisson-Poincaré equation (1.1) are sufficient to establish a priori arguments about the local well-posedness of the problem (1.2), as in Section 4. Meanwhile, the elliptic estimates are strong enough to make $\|\phi\|_{\mathcal{E}}$ be absorbed by $\|f\|_{\mathcal{E}}$. This is the reason why we only need to define $\|\cdot\|_{\mathcal{E}}$ by $\|f\|$ rather than $\|f\| + \|\phi\|$. As a contrast, one can refer to [18] for the explanations on the construction of energy functional for the Vlasov-Maxwell-Boltzmann system where the electronic and magnetic fields are included.

1.6.3. *Coercivity of the linearized operator \mathcal{L} .* Heuristically, we denote $\mathcal{B} = -\mathcal{L} - v \cdot \nabla_x$, then the integral form of (1.10) read as (cf. [5] for the Boltzmann equation case)

$$f(t) = e^{t\mathcal{B}} f(0) + \int_0^t e^{(t-s)\mathcal{B}} \Gamma(f(s), f(s)) ds.$$

If we could prove that

$$\|e^{t\mathcal{B}}\| \leq C e^{-Ct} \quad \text{and} \quad \|\Gamma(f, f)\| \leq C\|f\|^2$$

hold in some norm $\|\cdot\|$ for some $C > 0$, then we have

$$\|f(t)\| \leq C e^{-Ct} \|f(0)\| + \int_0^t C e^{-C(t-s)} \|f(s)\|^2 ds.$$

Denote $\check{f} = \sup_t \|f(t) e^{ct}\|$. It holds

$$\check{f} \leq C\|f(0)\| + C\check{f}^2,$$

which implies the boundedness of \check{f} under some smallness conditions. However, the five dimensional null space of \mathcal{L} prevents us utilizing this strategy naturally. In fact, $e^{-t\mathcal{L}} h = h$ as h belongs to the kernel of \mathcal{L} , so the desired property $\|e^{-t\mathcal{L}}\| \leq e^{-Ct}$ does not hold.

Henceforth, following the similar argument in [18], we devote to establishing the coercivity estimates of \mathcal{L} in terms of the solution f to the ionic Vlasov-Poisson-Boltzmann system (1.3)–(1.4) with the conservation laws (1.6)–(1.8). The key point is to control the macroscopic part of $\partial^\gamma f$ by its microscopic part in $\|\cdot\|_\nu$ norm. Denote

$$\mathbf{P} \partial^\gamma f(t, x, v) = \{\partial^\gamma a(t, x) + \partial^\gamma b(t, x) \cdot v + \partial^\gamma c(t, x) |v|^2\} \sqrt{\mu},$$

where the orthogonal projection $\mathbf{P}: L_v^2 \rightarrow \ker(\mathcal{L})(=: \mathcal{N})$. Due to the nonlinear exponential term in the Poisson-Poincaré equation, it raises new difficulties for estimating the pure spatial derivatives of $a(t, x)$ satisfying (5.7):

$$\partial^0 b_i + \partial^i a - E_i = l_{bi} + h_{bi}.$$

A first glimpse leads to $\|\partial^i a\| \leq \|\partial^0 b_i\| + \|l_{bi}\| + \|h_{bi}\| + \|E_i\|$, but $\|E_i\|$ cannot be controlled by

$$C \sum_{|\gamma| \leq N} \|\{\mathbf{I} - \mathbf{P}\} \partial^\gamma f(t)\|_\nu + C \sqrt{M_0} \sum_{|\gamma| \leq N} \|\partial^\gamma f(t)\|_\nu,$$

required by (5.2), where $M_0 \ll 1$. We thus ought to combine the density of ion, $\varrho_0 a + \varrho_2 c$ ($\varrho_0 = \int \mu dv$, $\varrho_2 = \int |v|^2 \mu dv$), with the density of electron, e^ϕ . In fact, from (5.4) and (5.7) we have three identities:

$$\begin{aligned} \text{(i). } & -\Delta a + \nabla \cdot E = \nabla \cdot \partial^0 b - \sum_i \partial^i (l_{bi} + h_{bi}), \\ \text{(ii). } & -\Delta c = -\nabla \cdot (l_c + h_c), \\ \text{(iii). } & -\Delta \phi - \nabla \cdot E = 0. \end{aligned}$$

For some sufficiently large constant $\Theta > 0$, we operate (i) $\times (\rho_0 a + \rho_2 c)$ + (ii) $\times \Theta c$ + (iii) $\times (e^\phi - 1)$ and then integrate the result over \mathbb{T}^3 to get

$$\begin{aligned} & \int_{\mathbb{T}^3} |\nabla a|^2 + \Theta |\nabla c|^2 + |\nabla \phi|^2 e^\phi + (\nabla \cdot E)^2 + (\nabla a \cdot \nabla c) dx \\ & \leq C(\|\partial^0 b\| + \|l_{bi} + h_{bi}\| + \|l_c + h_c\|)(\|\nabla a\| + \|\nabla c\|). \end{aligned}$$

It follows that

$$\|\nabla a\| + \|\nabla c\| \leq C(\|\partial^0 b\| + \|l_{bi} + h_{bi}\| + \|l_c + h_c\|),$$

which can be controlled by $C \sum_{|\gamma| \leq N} \|\{\mathbf{I} - \mathbf{P}\} \partial^\gamma f(t)\|_\nu + C\sqrt{M_0} \sum_{|\gamma| \leq N} \|\partial^\gamma f(t)\|_\nu$. Through the above deductions, we remark that the identity (5.4) of c plays a crucial role in estimating ∇a . The high-order derivative analogy of this estimate is presented in Section 5 with some extra techniques.

1.6.4. *A unified energy functional for different interaction kernels.* Since the collision frequency $\nu(v) \sim \langle v \rangle^r$ holds for the whole range $r \in (-3, 1]$, the inner product

$$\begin{aligned} & \left\langle \sum_{\gamma_1 \neq 0} C_{\beta^1}^{\beta_1} C_{\gamma_1}^{\gamma_1} \partial^{\gamma_1} E \cdot \partial_{\beta_1} \left(\frac{v}{2}\right) \partial_{\beta - \beta_1}^{\gamma - \gamma_1} f - \sum_{\gamma_1 \neq 0} C_{\gamma_1}^{\gamma_1} \partial^{\gamma_1} E \cdot \nabla_v \partial_{\beta}^{\gamma - \gamma_1} f, \partial_{\beta}^{\gamma} f \right\rangle \\ & \leq C \int_{\mathbb{R}^3} |\partial^{\gamma_1} E| dx \int_{\mathbb{R}^3} (|v| |\partial_{\beta - \beta_1}^{\gamma - \gamma_1} f| + |\nabla_v \partial_{\beta}^{\gamma - \gamma_1} f|) |\partial_{\beta}^{\gamma} f| dv \end{aligned}$$

is not naturally bounded by $\|f\| \cdot \|f\|_\nu^2$ except for $r = 1$. Hence it requires to enhance the L^2 dissipation $\|f\|_\nu^2$ for $-3 < r < 1$. We introduce the weight function $W \equiv W(x, v) := e^{-q \frac{x \cdot v}{\langle v \rangle}}$ (see [8]) with large constant $q \gg 1$. For $x \in \mathbb{T}^3$, it holds $W \sim 1$, which is crucial for simplifying the suitable energy norms. Involving with the transport operator, it holds (see (6.4) below)

$$-\frac{\{\partial_t + v \cdot \nabla_x + E \cdot \nabla_v\} W^2}{2W^2} = q \frac{|v|^2}{\langle v \rangle} + qE \cdot \left(\frac{x}{\langle v \rangle} + \frac{(x \cdot v)v}{\langle v \rangle^3} \right) := \Xi,$$

where $\|E\|_\infty \leq C\|E\|_{H^2} \leq C\|f\| \ll 1$. Under the L^2 inner product, this weight function would contribute an extra term $\Xi |W \partial_{\beta}^{\gamma} f|^2$. Combining with $\nu |W \partial_{\beta}^{\gamma} f|^2$ yields a new dissipation term

$$(\nu + \Xi) |W \partial_{\beta}^{\gamma} f|^2 \sim \langle v \rangle^{1/2} W \partial_{\beta}^{\gamma} f|^2 \sim \langle v \rangle^{1/2} \partial_{\beta}^{\gamma} f|^2,$$

which provides a motivation for the choice of energy functional $\mathcal{E}(t)$ in (1.9).

For the hard and soft potentials cases, we revisit in Section 2 some basic estimates of the Boltzmann operators $\mathcal{K} = \nu - \mathcal{L}$ and Γ , without restrictions of extra weight functions. The coercivity of \mathcal{L} in Section 5 also relies on different potentials, we shall present a unified proof covering the whole ranges.

1.7. **Organization of the article.** The rest of this article is organized as follows. In Section 2, we list some basic estimates of the Boltzmann operator with necessary details of proofs. In Section 3, we establish the L^2 regularity estimates for the Poisson-Poincaré equation (1.1). Section 4 is devoted to the local well-posedness of (1.2) with small initial datum in Picard's iteration schemes. In Section 5 we prove Theorem 5.1 to verify the coercivity of the linearized operator \mathcal{L} . Finally, a global-in-time solution with exponential decay is constructed in Section 6.

2. PRELIMINARY

In this section we present some basic facts about the linearized operator $\mathcal{L} = \nu - \mathcal{K}$ and the nonlinear operator Γ .

Lemma 2.1. *It holds $\langle \mathcal{L}g, h \rangle = \langle \mathcal{L}h, g \rangle$, $\langle \mathcal{L}g, g \rangle \geq 0$. And $\mathcal{L}g = 0$ if and only if*

$$g = \mathbf{P}g,$$

where $\mathbf{P} : L_v^2(\mathbb{R}^3) \rightarrow \mathcal{N}$ is the orthogonal projection in $L_v^2(\mathbb{R}^3)$ to the null space \mathcal{N} of \mathcal{L} , which is a five dimensional space spanned by $\{\sqrt{\mu}, v_1\sqrt{\mu}, v_2\sqrt{\mu}, v_3\sqrt{\mu}, |v|^2\sqrt{\mu}\}$.

Moreover, there exists a constant $\delta > 0$ such that

$$\langle \mathcal{L}g, g \rangle \geq \delta \|\{\mathbf{I} - \mathbf{P}\}g\|_v^2.$$

For the proofs, one can refer to [17, 19].

Lemma 2.2. *For the potential index $r \in (-3, 1]$ and the multi-index β , the collision frequency $\nu(v)$ satisfies*

$$\partial_\beta \nu(v) \leq C(1 + |v|)^{r-|\beta|}. \quad (2.1)$$

In addition, for any $\eta > 0$, there exists a $C_\eta > 0$ such that

$$\langle \partial_\beta[\nu g], \partial_\beta g \rangle \geq |\partial_\beta g|_v^2 - \eta \sum_{|\beta_1| \leq |\beta|} |\partial_{\beta_1} g|_v^2 - C_\eta |g|_v^2, \quad \beta \neq 0, \quad (2.2)$$

$$|\langle \partial_\beta[\mathcal{K}g_1], \partial_\beta g_2 \rangle| \leq \left\{ \eta \sum_{|\beta_1| \leq |\beta|} |\partial_{\beta_1} g_1|_v + C_\eta |g_1|_v \right\} |\partial_\beta g_2|_v. \quad (2.3)$$

Proof. The proof of (2.1) can be found in [16]. By using (2.1) and compact interpolation between ∂_β and ∂_0 , we have

$$\begin{aligned} \langle \partial_\beta[\nu g], \partial_\beta g \rangle &= |\partial_\beta g|_v^2 + \sum_{0 < |\beta_1| \leq |\beta|} C_\beta^{\beta_1} \langle \partial_{\beta_1} \nu \partial_{\beta-\beta_1} g, \partial_\beta g \rangle \\ &\geq |\partial_\beta g|_v^2 - \eta \sum_{|\beta_1| \leq |\beta|} |\partial_{\beta_1} g|_v^2 - C_\eta |g|_v^2 \end{aligned}$$

for any $\eta > 0$, thus (2.2) holds. For the proof of (2.3), Lemma 4 in [19] solved the case $r \in (-3, 0)$. For the case $r \in [0, 1]$, it only needs to note that

$$|\partial_\beta[\mathcal{K}g]| \leq \eta \sum_{|\beta_1|=|\beta|} |\partial_{\beta_1} g| + C_\eta |g|,$$

which was presented in Lemma 2.1 in [10]. Therefore, (2.3) is valid. \square

Remark 2.3. *From the above proofs, we actually obtain that, for $r \in [0, 1]$,*

$$\begin{aligned} \langle \partial_\beta[\nu g], \partial_\beta g \rangle &\geq |\partial_\beta g|_v^2 - \eta \sum_{|\beta_1| \leq |\beta|} |\partial_{\beta_1} g|_v^2 - C_\eta |g|_v^2, \quad \beta \neq 0, \\ |\langle \partial_\beta[\mathcal{K}g_1], \partial_\beta g_2 \rangle| &\leq \left\{ \eta \sum_{|\beta_1| \leq |\beta|} |\partial_{\beta_1} g_1|_v + C_\eta |g_1|_v \right\} |\partial_\beta g_2|_v. \end{aligned}$$

Lemma 2.4. *Let the indices $\beta_0 + \beta_1 + \beta_2 = \beta$, $\gamma_1 + \gamma_2 = \gamma$. It holds*

$$\partial_\beta^\gamma \Gamma(g_1, g_2) = \sum C_\beta^{\beta_0 \beta_1 \beta_2} C_\gamma^{\gamma_1 \gamma_2} \Gamma^0(\partial_{\beta_1}^{\gamma_1} g_1, \partial_{\beta_2}^{\gamma_2} g_2),$$

where

$$\begin{aligned} \Gamma^0(\partial_{\beta_1}^{\gamma_1} g_1, \partial_{\beta_2}^{\gamma_2} g_2) &= \iint_{\mathbb{R}^3 \times \mathbb{S}^2} |u - v|^r b_0(\theta) \partial_{\beta_0}[\sqrt{\mu(u)}] \partial_{\beta_1}^{\gamma_1} g_1(u') \partial_{\beta_2}^{\gamma_2} g_2(v') dx d\omega \\ &\quad - \iint_{\mathbb{R}^3 \times \mathbb{S}^2} |u - v|^r b_0(\theta) \partial_{\beta_0}[\sqrt{\mu(u)}] \partial_{\beta_1}^{\gamma_1} g_1(u) \partial_{\beta_2}^{\gamma_2} g_2(v) dx d\omega \end{aligned}$$

$$=: \Gamma_{\text{gain}}^0 - \Gamma_{\text{loss}}^0.$$

Moreover, for $|\beta| + |\gamma| \leq N (N \geq 8)$, we have

$$\begin{aligned} & \left| \left(\Gamma^0(\partial_{\beta_1}^{\gamma_1} g_1, \partial_{\beta_2}^{\gamma_2} g_2), \partial_{\beta}^{\gamma} g_3 \right) \right| \\ & \leq C \left\{ \sum_{|\beta_k| + |\gamma_k| \leq N} \|\partial_{\beta_k}^{\gamma_k} g_1\| \right\} \|\partial_{\beta_2}^{\gamma_2} g_2\|_{\nu} \|\partial_{\beta}^{\gamma} g_3\|_{\nu} + C \left\{ \sum_{|\beta_k| + |\gamma_k| \leq N} \|\partial_{\beta_k}^{\gamma_k} g_1\|_{\nu} \right\} \|\partial_{\beta_2}^{\gamma_2} g_2\| \|\partial_{\beta}^{\gamma} g_3\|_{\nu} \\ & \quad + C \left\{ \sum_{|\beta_k| + |\gamma_k| \leq N} \|\partial_{\beta_k}^{\gamma_k} g_2\| \right\} \|\partial_{\beta_1}^{\gamma_1} g_1\|_{\nu} \|\partial_{\beta}^{\gamma} g_3\|_{\nu} + C \left\{ \sum_{|\beta_k| + |\gamma_k| \leq N} \|\partial_{\beta_k}^{\gamma_k} g_2\|_{\nu} \right\} \|\partial_{\beta_1}^{\gamma_1} g_1\| \|\partial_{\beta}^{\gamma} g_3\|_{\nu}. \end{aligned} \quad (2.4)$$

Remark 2.5. For $|\beta| + |\gamma| = 0$, it holds that $|\beta_1| + |\gamma_1| = 0$ and $|\beta_2| + |\gamma_2| = 0$. Then (2.4) can be further refined as:

$$|\langle \Gamma(g_1, g_2), g_3 \rangle| \leq C \left\{ \sum_{|\beta_k| + |\gamma_k| \leq N} \|\partial_{\beta_k}^{\gamma_k} g_1\| \right\} \|g_2\|_{\nu} \|g_3\|_{\nu} + C \left\{ \sum_{|\beta_k| + |\gamma_k| \leq N} \|\partial_{\beta_k}^{\gamma_k} g_1\|_{\nu} \right\} \|g_2\| \|g_3\|_{\nu},$$

or

$$|\langle \Gamma(g_1, g_2), g_3 \rangle| \leq C \left\{ \sum_{|\beta_k| + |\gamma_k| \leq N} \|\partial_{\beta_k}^{\gamma_k} g_2\| \right\} \|g_1\|_{\nu} \|g_3\|_{\nu} + C \left\{ \sum_{|\beta_k| + |\gamma_k| \leq N} \|\partial_{\beta_k}^{\gamma_k} g_2\|_{\nu} \right\} \|g_1\| \|g_3\|_{\nu}.$$

Proof of Lemma 2.4. To verify (2.4), we divide the potential index $r \in (-3, 1]$ into $(-3, 0) \cup [0, 1]$.

Case 1. For $r \in [0, 1]$, we only need to show that

$$\left| \left\langle \Gamma^0(\partial_{\beta_1}^{\gamma_1} g_1, \partial_{\beta_2}^{\gamma_2} g_2), \partial_{\beta}^{\gamma} g_3 \right\rangle \right| \leq C |\partial_{\beta_1}^{\gamma_1} g_1|_{\nu} |\partial_{\beta_2}^{\gamma_2} g_2| |\partial_{\beta}^{\gamma} g_3|_{\nu} + C |\partial_{\beta_2}^{\gamma_2} g_2|_{\nu} |\partial_{\beta_1}^{\gamma_1} g_1| |\partial_{\beta}^{\gamma} g_3|_{\nu}. \quad (2.5)$$

Recall $\langle \cdot, \cdot \rangle$ is the inner product in $L^2(\mathbb{R}_v^3)$ and (\cdot, \cdot) is the inner product in $L^2(\mathbb{T}_x^3 \times \mathbb{R}_v^3)$. For the proof of (2.5), we notice that $|v - u|^r \leq C(|v|^r + |u|^r)$ for $r \geq 0$, and

$$|v - u|^r e^{-\frac{|u|^2}{8}} \leq C(|v|^r + |u|^r) e^{-\frac{|u|^2}{8}} \leq C \langle v \rangle^r e^{-\frac{|u|^2}{16}}.$$

Then

$$\begin{aligned} \left| \left\langle \Gamma_{\text{loss}}^0(\partial_{\beta_1}^{\gamma_1} g_1, \partial_{\beta_2}^{\gamma_2} g_2), \partial_{\beta}^{\gamma} g_3 \right\rangle \right| & \leq C \iint_{\mathbb{R}^3 \times \mathbb{R}^3} |u - v|^r e^{-\frac{|u|^2}{8}} |\partial_{\beta_1}^{\gamma_1} g_1(u) \partial_{\beta_2}^{\gamma_2} g_2(v) \partial_{\beta}^{\gamma} g_3(v)| du dv \\ & \leq C \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \langle v \rangle^r e^{-\frac{|u|^2}{16}} |\partial_{\beta_1}^{\gamma_1} g_1(u)| |\partial_{\beta_2}^{\gamma_2} g_2(v)| |\partial_{\beta}^{\gamma} g_3(v)| du dv \\ & \leq C \left\{ \int_{\mathbb{R}^3} |\partial_{\beta_1}^{\gamma_1} g_1(u)|^2 du \right\}^{1/2} \left\{ \int_{\mathbb{R}^3} \nu(v) |\partial_{\beta_2}^{\gamma_2} g_2(v)|^2 dv \right\}^{1/2} \left\{ \int_{\mathbb{R}^3} \nu(v) |\partial_{\beta}^{\gamma} g_3(v)|^2 dv \right\}^{1/2}, \end{aligned}$$

and

$$\begin{aligned} & \left| \left\langle \Gamma_{\text{gain}}^0(\partial_{\beta_1}^{\gamma_1} g_1, \partial_{\beta_2}^{\gamma_2} g_2), \partial_{\beta}^{\gamma} g_3 \right\rangle \right| \\ & \leq C \iint_{\mathbb{R}^3 \times \mathbb{R}^3} |u - v|^r e^{-\frac{|u|^2}{8}} |\partial_{\beta_1}^{\gamma_1} g_1(u') \partial_{\beta_2}^{\gamma_2} g_2(v') \partial_{\beta}^{\gamma} g_3(v)| du dv \\ & \leq C \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \langle v \rangle^r e^{-\frac{|u|^2}{16}} |\partial_{\beta_1}^{\gamma_1} g_1(u') \partial_{\beta_2}^{\gamma_2} g_2(v')| |\partial_{\beta}^{\gamma} g_3(v)| du dv \\ & \leq C \left\{ \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \langle v \rangle^r e^{-\frac{|u|^2}{16}} |\partial_{\beta_1}^{\gamma_1} g_1(u') \partial_{\beta_2}^{\gamma_2} g_2(v')|^2 du dv \right\}^{1/2} \left\{ \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \langle v \rangle^r e^{-\frac{|u|^2}{16}} |\partial_{\beta}^{\gamma} g_3(v)|^2 du dv \right\}^{1/2} \\ & \leq C \left\{ \iint_{\mathbb{R}^3 \times \mathbb{R}^3} (\langle u' \rangle^r + \langle v' \rangle^r) |\partial_{\beta_1}^{\gamma_1} g_1(u') \partial_{\beta_2}^{\gamma_2} g_2(v')|^2 du' dv' \right\}^{1/2} \left\{ \int_{\mathbb{R}^3} \langle v \rangle^r |\partial_{\beta}^{\gamma} g_3(v)|^2 dv \right\}^{1/2} \\ & \leq C \left\{ \int_{\mathbb{R}^3} \nu(u') |\partial_{\beta_1}^{\gamma_1} g_1(u')|^2 du' \right\}^{1/2} \left\{ \int_{\mathbb{R}^3} |\partial_{\beta_2}^{\gamma_2} g_2(v')|^2 dv' \right\}^{1/2} \left\{ \int_{\mathbb{R}^3} \langle v \rangle^r |\partial_{\beta}^{\gamma} g_3(v)|^2 dv \right\}^{1/2} \\ & \quad + C \left\{ \int_{\mathbb{R}^3} |\partial_{\beta_1}^{\gamma_1} g_1(u')|^2 du' \right\}^{1/2} \left\{ \int_{\mathbb{R}^3} \nu(v') |\partial_{\beta_2}^{\gamma_2} g_2(v')|^2 dv' \right\}^{1/2} \left\{ \int_{\mathbb{R}^3} \langle v \rangle^r |\partial_{\beta}^{\gamma} g_3(v)|^2 dv \right\}^{1/2}, \end{aligned}$$

where we have used the fact that $\langle v \rangle^r \leq C(\langle u \rangle^r + \langle v \rangle^r)$, $r \geq 0$. Then (2.5) follows. Recalling $H^2(\mathbb{T}^3) \subset L^\infty(\mathbb{T}^3)$, we have

$$\begin{aligned} \sup_x \int_{\mathbb{R}^3} |g(x, u)|^2 du &\leq \int_{\mathbb{R}^3} \sup_x |g(x, u)|^2 du \leq C \sum_{|\gamma| \leq 2} \|\partial^\gamma g\|^2, \\ \sup_x \int_{\mathbb{R}^3} \nu(u) |g(x, u)|^2 du &\leq C \sum_{|\gamma| \leq 2} \|\partial^\gamma g\|_\nu^2. \end{aligned}$$

By using the above embedding, we take the $L^\infty(\mathbb{T}^3)$ norm of g_1 or g_2 terms on the right-hand side of (2.5) whichever has less total derivatives ($|\beta_i| + |\gamma_i| \leq N/2, i = 1, 2$). Then, integrating (2.5) over $x \in \mathbb{T}^3$ yields (2.4).

Case 2. For $r \in (-3, 0)$, we first consider Γ_{loss}^0 . If $|\beta_1| + |\gamma_1| \leq N/2$, then

$$\begin{aligned} &\int_{\mathbb{R}^3} |u - v|^r |\partial_{\beta_0} [\sqrt{\mu(u)}] \partial_{\beta_1}^{\gamma_1} g_1(x, u)| du \\ &\leq C \left\{ \int_{\mathbb{R}^3} |u - v|^r e^{-\frac{|u|^2}{8}} |\partial_{\beta_1}^{\gamma_1} g_1(x, u)|^2 du \right\}^{1/2} \left\{ \int_{\mathbb{R}^3} |u - v|^r e^{-\frac{|u|^2}{8}} du \right\}^{1/2} \\ &\leq C \sup_{x, u} |\partial_{\beta_1}^{\gamma_1} g_1(x, u)| \left\{ \int_{\mathbb{R}^3} |u - v|^r e^{-\frac{|u|^2}{8}} du \right\} \\ &\leq C \left\{ \sum_{|\beta_k| + |\gamma_k| \leq N} \|\partial_{\beta_k}^{\gamma_k} g_1\| \right\} \langle v \rangle^r, \end{aligned}$$

where we have used $H^4(\mathbb{T}^3 \times \mathbb{R}^3) \subset L^\infty(\mathbb{T}^3 \times \mathbb{R}^3)$ in the last inequality. Hence we have

$$\begin{aligned} \left| \left(\Gamma_{\text{loss}}^0(\partial_{\beta_1}^{\gamma_1} g_1, \partial_{\beta_2}^{\gamma_2} g_2), \partial_{\beta_3}^{\gamma_3} g_3 \right) \right| &\leq C \left\{ \sum_{|\beta_k| + |\gamma_k| \leq N} \|\partial_{\beta_k}^{\gamma_k} g_1\| \right\} \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \langle v \rangle^r |\partial_{\beta_2}^{\gamma_2} g_2(x, v) \partial_{\beta_3}^{\gamma_3} g_3(x, v)| dv dx \\ &\leq C \left\{ \sum_{|\beta_k| + |\gamma_k| \leq N} \|\partial_{\beta_k}^{\gamma_k} g_1\| \right\} \|\partial_{\beta_2}^{\gamma_2} g_2\|_\nu \|\partial_{\beta_3}^{\gamma_3} g_3\|_\nu, \end{aligned}$$

which completes the case of $|\beta_1| + |\gamma_1| \leq N/2$ for Γ_{loss}^0 .

On the other hand, if $|\beta_2| + |\gamma_2| \leq N/2$, then we consider $\{|v - u| \leq |v|/2\} \cup \{|v - u| \geq |v|/2\}$ separately.

(i). $|v - u| \leq |v|/2$ implies that $|u| \geq |v| - |v - u| \geq |v|/2$, thus $e^{-\frac{|u|^2}{8}} \leq e^{-\frac{|u|^2}{16}} e^{-\frac{|v|^2}{64}}$. We have

$$\begin{aligned} &\left| \left(\Gamma_{\text{loss}}^0(\partial_{\beta_1}^{\gamma_1} g_1, \partial_{\beta_2}^{\gamma_2} g_2), \partial_{\beta_3}^{\gamma_3} g_3 \right) \right| \\ &\leq C \iiint_{\mathbb{T}^3 \times \mathbb{R}^3 \times \mathbb{R}^3} |u - v|^r e^{-\frac{|u|^2}{16}} e^{-\frac{|v|^2}{64}} |\partial_{\beta_1}^{\gamma_1} g_1(u) \partial_{\beta_2}^{\gamma_2} g_2(v) \partial_{\beta_3}^{\gamma_3} g_3(v)| dudv dx \\ &\leq C \left\{ \iiint_{\mathbb{T}^3 \times \mathbb{R}^3 \times \mathbb{R}^3} |u - v|^r e^{-\frac{|u|^2}{16}} e^{-\frac{|v|^2}{64}} |\partial_{\beta_1}^{\gamma_1} g_1(u)|^2 dudv dx \right\}^{1/2} \\ &\quad \times \left\{ \iiint_{\mathbb{T}^3 \times \mathbb{R}^3 \times \mathbb{R}^3} |u - v|^r e^{-\frac{|u|^2}{16}} e^{-\frac{|v|^2}{64}} |\partial_{\beta_2}^{\gamma_2} g_2(v) \partial_{\beta_3}^{\gamma_3} g_3(v)|^2 dudv dx \right\}^{1/2} \\ &\leq C \|\partial_{\beta_1}^{\gamma_1} g_1\|_\nu \left\{ \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \langle v \rangle^r e^{-\frac{|v|^2}{64}} |\partial_{\beta_2}^{\gamma_2} g_2(v) \partial_{\beta_3}^{\gamma_3} g_3(v)|^2 dv dx \right\}^{1/2} \\ &\leq C \|\partial_{\beta_1}^{\gamma_1} g_1\|_\nu \left\{ \sum_{|\beta_k| + |\gamma_k| \leq N} \|\partial_{\beta_k}^{\gamma_k} g_2\| \right\} \|\partial_{\beta_3}^{\gamma_3} g_3\|_\nu, \end{aligned}$$

where we have used $H^4(\mathbb{T}^3 \times \mathbb{R}^3) \subset L^\infty(\mathbb{T}^3 \times \mathbb{R}^3)$ again.

(ii). $|v - u| \geq |v|/2$ implies that $|v - u|^r \leq C|v|^r$ for $r < 0$. When $|v| \geq 1$, it holds

$$\begin{aligned} &\left| \left(\Gamma_{\text{loss}}^0(\partial_{\beta_1}^{\gamma_1} g_1, \partial_{\beta_2}^{\gamma_2} g_2), \partial_{\beta_3}^{\gamma_3} g_3 \right) \right| \\ &\leq C \iiint_{\mathbb{T}^3 \times \mathbb{R}^3 \times \mathbb{R}^3} |u - v|^r e^{-\frac{|u|^2}{8}} |\partial_{\beta_1}^{\gamma_1} g_1(u) \partial_{\beta_2}^{\gamma_2} g_2(v) \partial_{\beta_3}^{\gamma_3} g_3(v)| dudv dx \end{aligned}$$

$$\begin{aligned}
&\leq C \int_{\mathbb{T}^3} \left\{ \int_{\mathbb{R}^3} e^{-\frac{|u|^2}{8}} |\partial_{\beta_1}^{\gamma_1} g_1(u)| du \right\} \left\{ \int_{\{|v| \geq 1\}} |v|^r |\partial_{\beta_2}^{\gamma_2} g_2(v) \partial_{\beta}^{\gamma} g_3(v)| dv \right\} dx \\
&\leq C \int_{\mathbb{T}^3} |\partial_{\beta_1}^{\gamma_1} g_1|_{\nu} \left\{ \int_{\mathbb{R}^3} |\partial_{\beta_2}^{\gamma_2} g_2|^2 dv \right\}^{1/2} \left\{ \int_{\{|v| \geq 1\}} \langle v \rangle^r |\partial_{\beta}^{\gamma} g_3|^2 dv \right\}^{1/2} dx \\
&\leq C \|\partial_{\beta_1}^{\gamma_1} g_1\|_{\nu} \left\{ \sum_{|\beta_k| + |\gamma_k| \leq N} \|\partial_{\beta_k}^{\gamma_k} g_2\| \right\} \|\partial_{\beta}^{\gamma} g_3\|_{\nu}.
\end{aligned}$$

When $|v| \leq 1$, it holds $\langle v \rangle^r \geq C > 0$. Then,

$$\begin{aligned}
&\left| \left(\Gamma_{\text{loss}}^0(\partial_{\beta_1}^{\gamma_1} g_1, \partial_{\beta_2}^{\gamma_2} g_2), \partial_{\beta}^{\gamma} g_3 \right) \right| \\
&\leq C \int_{\mathbb{T}^3} \left\{ \int_{\mathbb{R}^3} |u - v|^{r/2} e^{-\frac{|u|^2}{8}} |\partial_{\beta_1}^{\gamma_1} g_1(u)| du \right\} \left\{ \int_{\{|v| \leq 1\}} |v|^{r/2} |\partial_{\beta_2}^{\gamma_2} g_2(v) \partial_{\beta}^{\gamma} g_3(v)| dv \right\} dx \\
&\leq C \int_{\mathbb{T}^3} \left\{ \int_{\mathbb{R}^3} |u - v|^r e^{-\frac{|u|^2}{8}} du \right\}^{1/2} \left\{ \int_{\mathbb{R}^3} e^{-\frac{|u|^2}{8}} |\partial_{\beta_1}^{\gamma_1} g_1(u)|^2 du \right\}^{1/2} \\
&\quad \times \left\{ \int_{\{|v| \leq 1\}} |v|^r |\partial_{\beta_2}^{\gamma_2} g_2(v)|^2 dv \right\}^{1/2} \left\{ \int_{\{|v| \leq 1\}} |\partial_{\beta}^{\gamma} g_3(v)|^2 dv \right\}^{1/2} dx \\
&\leq C \|\partial_{\beta_1}^{\gamma_1} g_1\|_{\nu} \left\{ \sum_{|\beta_k| + |\gamma_k| \leq N} \|\partial_{\beta_k}^{\gamma_k} g_2\| \right\} \|\partial_{\beta}^{\gamma} g_3\|_{\nu}.
\end{aligned}$$

In summary, we have

$$\left| \left(\Gamma_{\text{loss}}^0(\partial_{\beta_1}^{\gamma_1} g_1, \partial_{\beta_2}^{\gamma_2} g_2), \partial_{\beta}^{\gamma} g_3 \right) \right| \leq C \|\partial_{\beta_1}^{\gamma_1} g_1\|_{\nu} \left\{ \sum_{|\beta_k| + |\gamma_k| \leq N} \|\partial_{\beta_k}^{\gamma_k} g_2\| \right\} \|\partial_{\beta}^{\gamma} g_3\|_{\nu}.$$

We next consider Γ_{gain}^0 for $r \in (-3, 0)$. Split $\mathbb{R}_u^3 \times \mathbb{R}_v^3$ into $\{|u| \geq |v|/2\} \cup \{|u| \leq |v|/2\}$.

(i). When $|u| \geq |v|/2$, we have $e^{-|u|^2/8} \leq e^{-|u|^2/16} e^{-|v|^2/64}$. Thus,

$$\begin{aligned}
&\left| \left(\Gamma_{\text{gain}}^0(\partial_{\beta_1}^{\gamma_1} g_1, \partial_{\beta_2}^{\gamma_2} g_2), \partial_{\beta}^{\gamma} g_3 \right) \right| \\
&\leq C \iiint_{\mathbb{T}^3 \times \mathbb{R}^3 \times \mathbb{R}^3} |u - v|^r e^{-\frac{|u|^2}{16}} e^{-\frac{|v|^2}{64}} |\partial_{\beta_1}^{\gamma_1} g_1(u') \partial_{\beta_2}^{\gamma_2} g_2(v') \partial_{\beta}^{\gamma} g_3(v)| dudvdx \\
&\leq C \left\{ \iiint_{\mathbb{T}^3 \times \mathbb{R}^3 \times \mathbb{R}^3} |u - v|^r e^{-\frac{|u|^2}{16}} e^{-\frac{|v|^2}{64}} |\partial_{\beta_1}^{\gamma_1} g_1(u') \partial_{\beta_2}^{\gamma_2} g_2(v')|^2 dudvdx \right\}^{1/2} \\
&\quad \times \left\{ \iiint_{\mathbb{T}^3 \times \mathbb{R}^3 \times \mathbb{R}^3} |u - v|^r e^{-\frac{|u|^2}{16}} e^{-\frac{|v|^2}{64}} |\partial_{\beta}^{\gamma} g_3(v)|^2 dudvdx \right\}^{1/2} \\
&\leq C \left\{ \iiint_{\mathbb{T}^3 \times \mathbb{R}^3 \times \mathbb{R}^3} |u - v|^r e^{-\frac{|u|^2}{64}} e^{-\frac{|v|^2}{64}} |\partial_{\beta_1}^{\gamma_1} g_1(u) \partial_{\beta_2}^{\gamma_2} g_2(v)|^2 dudvdx \right\}^{1/2} \|\partial_{\beta}^{\gamma} g_3\|_{\nu} \\
&\leq C \|\partial_{\beta_1}^{\gamma_1} g_1\|_{\nu} \left\{ \sum_{|\beta_k| + |\gamma_k| \leq N} \|\partial_{\beta_k}^{\gamma_k} g_2\| \right\} \|\partial_{\beta}^{\gamma} g_3\|_{\nu} + \|\partial_{\beta_2}^{\gamma_2} g_2\|_{\nu} \left\{ \sum_{|\beta_k| + |\gamma_k| \leq N} \|\partial_{\beta_k}^{\gamma_k} g_1\| \right\} \|\partial_{\beta}^{\gamma} g_3\|_{\nu},
\end{aligned}$$

where $du'dv' = dudv$ and $H^4(\mathbb{T}^3 \times \mathbb{R}^3) \subset L^\infty(\mathbb{T}^3 \times \mathbb{R}^3)$ have been used.

(ii). When $|u| \leq |v|/2$, we have $|v - u| \geq |v|/2$ and $|u'| + |v'| \leq C|v|$. We then consider cases (iia) and (iib) as follows:

(iia). If $|v| \geq 1$, then $|v|^r \sim \langle v \rangle^r$. We have

$$\begin{aligned}
&\left| \left(\Gamma_{\text{gain}}^0(\partial_{\beta_1}^{\gamma_1} g_1, \partial_{\beta_2}^{\gamma_2} g_2), \partial_{\beta}^{\gamma} g_3 \right) \right| \\
&\leq C \iiint_{\mathbb{T}^3 \times \mathbb{R}^3 \times \mathbb{R}^3} |u - v|^r e^{-\frac{|u|^2}{8}} |\partial_{\beta_1}^{\gamma_1} g_1(u') \partial_{\beta_2}^{\gamma_2} g_2(v') \partial_{\beta}^{\gamma} g_3(v)| dudvdx
\end{aligned}$$

$$\begin{aligned}
&\leq C \left\{ \iiint_{\mathbb{T}^3 \times \mathbb{R}^3 \times \mathbb{R}^3} \langle v \rangle^r e^{-\frac{|u|^2}{8}} |\partial_{\beta_1}^{\gamma_1} g_1(u') \partial_{\beta_2}^{\gamma_2} g_2(v')|^2 dudvdx \right\}^{1/2} \\
&\quad \times \left\{ \iiint_{\mathbb{T}^3 \times \mathbb{R}^3 \times \mathbb{R}^3} \langle v \rangle^r e^{-\frac{|u|^2}{8}} |\partial_{\beta}^{\gamma} g_3(v)|^2 dudvdx \right\}^{1/2} \\
&\leq C \left\{ \iiint_{\mathbb{T}^3 \times \mathbb{R}^3 \times \mathbb{R}^3} \min\{\langle u' \rangle^r, \langle v' \rangle^r\} |\partial_{\beta_1}^{\gamma_1} g_1(u')|^2 |\partial_{\beta_2}^{\gamma_2} g_2(v')|^2 dudvdx \right\}^{1/2} \|\partial_{\beta}^{\gamma} g_3\|_{\nu} \\
&\leq C \|\partial_{\beta_1}^{\gamma_1} g_1\|_{\nu} \left\{ \sum_{|\beta_k|+|\gamma_k| \leq N} \|\partial_{\beta_k}^{\gamma_k} g_2\| \right\} \|\partial_{\beta}^{\gamma} g_3\|_{\nu} + \|\partial_{\beta_2}^{\gamma_2} g_2\|_{\nu} \left\{ \sum_{|\beta_k|+|\gamma_k| \leq N} \|\partial_{\beta_k}^{\gamma_k} g_1\| \right\} \|\partial_{\beta}^{\gamma} g_3\|_{\nu} \\
&\quad + C \|\partial_{\beta_1}^{\gamma_1} g_1\| \left\{ \sum_{|\beta_k|+|\gamma_k| \leq N} \|\partial_{\beta_k}^{\gamma_k} g_2\| \right\} \|\partial_{\beta}^{\gamma} g_3\|_{\nu} + \|\partial_{\beta_2}^{\gamma_2} g_2\| \left\{ \sum_{|\beta_k|+|\gamma_k| \leq N} \|\partial_{\beta_k}^{\gamma_k} g_1\| \right\} \|\partial_{\beta}^{\gamma} g_3\|_{\nu}.
\end{aligned}$$

(iib). If $|v| \leq 1$, then $|u| \leq 1/2$, which implies that $|u'|$ and $|v'| \leq C'$, and thus $1 \leq C' \langle u' \rangle^r$ and $1 \leq C' \langle v' \rangle^r$ for some $C' > 0$. Noticing that

$$\int_{\{|u'| \leq C'\}} |u'|^r du' \leq C, \quad \int_{\{|v'| \leq C'\}} |v'|^r dv' \leq C,$$

we have

$$\begin{aligned}
&\left| \left(\Gamma_{\text{gain}}^0(\partial_{\beta_1}^{\gamma_1} g_1, \partial_{\beta_2}^{\gamma_2} g_2), \partial_{\beta}^{\gamma} g_3 \right) \right| \\
&\leq C \iiint_{\mathbb{T}^3 \times \mathbb{R}^3 \times \mathbb{R}^3} |u-v|^r e^{-\frac{|u|^2}{8}} |\partial_{\beta_1}^{\gamma_1} g_1(u') \partial_{\beta_2}^{\gamma_2} g_2(v') \partial_{\beta}^{\gamma} g_3(v)| dudvdx \\
&\leq C \iint_{\mathbb{T}^3 \times \mathbb{R}^3} \left\{ \int_{\mathbb{R}^3} |v|^r |\partial_{\beta_1}^{\gamma_1} g_1(u') \partial_{\beta_2}^{\gamma_2} g_2(v')|^2 du \right\}^{1/2} \left\{ \int_{\mathbb{R}^3} |u-v|^r e^{-\frac{|u|^2}{4}} du \right\}^{1/2} |\partial_{\beta}^{\gamma} g_3(v)| dvdx \\
&\leq C \left\{ \iiint_{\{|u'|+|v'| \leq C'\}} \min\{|u'|^r, |v'|^r\} |\partial_{\beta_1}^{\gamma_1} g_1(u')|^2 |\partial_{\beta_2}^{\gamma_2} g_2(v')|^2 du' dv' dx \right\}^{1/2} \|\partial_{\beta}^{\gamma} g_3\|_{\nu} \\
&\leq C \|\partial_{\beta_1}^{\gamma_1} g_1\|_{\nu} \left\{ \sum_{|\beta_k|+|\gamma_k| \leq N} \|\partial_{\beta_k}^{\gamma_k} g_2\| \right\} \|\partial_{\beta}^{\gamma} g_3\|_{\nu} + \|\partial_{\beta_2}^{\gamma_2} g_2\|_{\nu} \left\{ \sum_{|\beta_k|+|\gamma_k| \leq N} \|\partial_{\beta_k}^{\gamma_k} g_1\| \right\} \|\partial_{\beta}^{\gamma} g_3\|_{\nu},
\end{aligned}$$

where $du'dv' = dudv$ and $H^4(\mathbb{T}^3 \times \mathbb{R}^3) \subset L^\infty(\mathbb{T}^3 \times \mathbb{R}^3)$ have been used.

In summary, we get

$$\begin{aligned}
&\left| \left(\Gamma_{\text{gain}}^0(\partial_{\beta_1}^{\gamma_1} g_1, \partial_{\beta_2}^{\gamma_2} g_2), \partial_{\beta}^{\gamma} g_3 \right) \right| \\
&\leq C \|\partial_{\beta_1}^{\gamma_1} g_1\|_{\nu} \left\{ \sum_{|\beta_k|+|\gamma_k| \leq N} \|\partial_{\beta_k}^{\gamma_k} g_2\| \right\} \|\partial_{\beta}^{\gamma} g_3\|_{\nu} + \|\partial_{\beta_2}^{\gamma_2} g_2\|_{\nu} \left\{ \sum_{|\beta_k|+|\gamma_k| \leq N} \|\partial_{\beta_k}^{\gamma_k} g_1\| \right\} \|\partial_{\beta}^{\gamma} g_3\|_{\nu} \\
&\quad + C \|\partial_{\beta_1}^{\gamma_1} g_1\| \left\{ \sum_{|\beta_k|+|\gamma_k| \leq N} \|\partial_{\beta_k}^{\gamma_k} g_2\| \right\} \|\partial_{\beta}^{\gamma} g_3\|_{\nu} + \|\partial_{\beta_2}^{\gamma_2} g_2\| \left\{ \sum_{|\beta_k|+|\gamma_k| \leq N} \|\partial_{\beta_k}^{\gamma_k} g_1\| \right\} \|\partial_{\beta}^{\gamma} g_3\|_{\nu}.
\end{aligned}$$

Combining the above discussions in Case 1 for $r \in [0, 1]$ and Case 2 for $r \in (-3, 0)$, we conclude that (2.4) holds. \square

Lemma 2.6. *For the whole range $r \in (-3, 1]$, we have*

$$\begin{aligned}
\left| \int_{\mathbb{R}^3} \Gamma(f, g) h dv \right| &\leq C \sup_v |\mu^{1/4} h| \|f\|_{\nu} \|g\|_{\nu}, \\
\|\nu^{-1/2} \Gamma(f, g)\| &\leq C \left\{ \sup_{x,v} |e^{\vartheta|v|^2} f| \|g\|_{\nu} + \sup_{x,v} |e^{\vartheta|v|^2} g| \|f\|_{\nu} \right\}, \quad \vartheta > 0, \\
\|\nu^{-1/2} \mathcal{L}f\| &\leq C \|f\|_{\nu}.
\end{aligned}$$

Proof. It is easy to know that

$$\left| \int_{\mathbb{R}^3} \Gamma_{\text{gain}}(f, g) h dv \right| \leq \iint_{\mathbb{R}^3 \times \mathbb{R}^3} |u-v|^r \mu^{1/2}(u) f(u') g(v') h(v) dudv$$

$$\begin{aligned}
&\leq \sup_v |\mu^{1/4} h| \iint_{\mathbb{R}^3 \times \mathbb{R}^3} |u-v|^r \mu^{1/4}(u) \mu^{1/4}(v) f(u) g(v) dudv \\
&\leq \sup_v |\mu^{1/4} h| \left(\iint_{\mathbb{R}^3 \times \mathbb{R}^3} |u-v|^r \mu^{1/4}(u) \mu^{1/4}(v) f^2(u) dudv \right)^{1/2} \\
&\quad \times \left(\iint_{\mathbb{R}^3 \times \mathbb{R}^3} |u-v|^r \mu^{1/4}(u) \mu^{1/4}(v) g^2(v) dudv \right)^{1/2} \\
&\leq \sup_v |\mu^{1/4} h| \left(\int_{\mathbb{R}^3} \langle u \rangle^r \mu^{1/4}(u) f^2(u) du \right)^{1/2} \left(\int_{\mathbb{R}^3} \langle v \rangle^r \mu^{1/4}(v) g^2(v) dv \right)^{1/2} \\
&\leq C \sup_v |\mu^{1/4} h| \|f\|_\nu \|g\|_\nu,
\end{aligned}$$

and for $w := e^{\vartheta|v|^2}$, it holds

$$\begin{aligned}
&\left| \int_{\mathbb{R}^3} \nu^{-1/2} \Gamma_{\text{gain}}(f, g) h dv \right| \\
&\leq \iint_{\mathbb{R}^3 \times \mathbb{R}^3} |u-v|^r \mu^{1/2}(u) f(u') g(v') \nu^{-1/2}(v) h(v) dudv \\
&\leq \left(\iint_{\mathbb{R}^3 \times \mathbb{R}^3} |u-v|^r \mu^{1/2}(u) f^2(u') g^2(v') dudv \right)^{1/2} \left(\iint_{\mathbb{R}^3 \times \mathbb{R}^3} |u-v|^r \mu^{1/2}(u) \nu^{-1}(v) h^2(v) dudv \right)^{1/2} \\
&\leq \left(\iint_{\mathbb{R}^3 \times \mathbb{R}^3} |u-v|^r \mu^{1/2}(u') w^{-2}(u) w^2(u) f^2(u) g^2(v) dudv \right)^{1/2} \left(\int_{\mathbb{R}^3} h^2(v) dv \right)^{1/2} \\
&\leq C \|wf\|_{L^\infty_\nu} \|\nu^{1/2} g\|_{L^2_\nu} \|h\|_{L^2_\nu},
\end{aligned}$$

which implies that

$$\|\nu^{-1/2} \Gamma_{\text{gain}}(f, g)\| \leq C \sup_{x,v} |wf| \|g\|_\nu.$$

Similarly, $\Gamma_{\text{loss}}(\cdot, \cdot)$ also satisfies the above estimates. Hence, the first and second inequalities hold. Taking $g = \sqrt{\mu}$ in the second inequality yields the third one. Hence the proof is completed. \square

3. L^2 ESTIMATES OF THE POISSON-POINCARÉ EQUATION IN \mathbb{T}^3

In this section, we investigate the well-posedness and regularity theories of the Poisson-Poincaré equation:

$$\Delta U = e^U - \rho \quad \text{in } \mathbb{T}^3. \quad (3.1)$$

The main result of this section reads

Theorem 3.1. *Suppose that $\rho \in L^2(\mathbb{T}^3)$. Then, there exists a unique solution $U \in H^2(\mathbb{T}^3)$ to the Poisson-Poincaré equation (3.1) in \mathbb{T}^3 .*

Proof. We split $U = \bar{U} + \hat{U}$ such that (3.1) is decomposed into two equations:

$$\Delta \bar{U} = 1 - \rho, \quad \Delta \hat{U} = e^{\bar{U} + \hat{U}} - 1.$$

Without loss of generality, we can assume that

$$(\bar{U})_{\mathbb{T}^3} := \frac{1}{|\mathbb{T}^3|} \int_{\mathbb{T}^3} \bar{U} dx = 0.$$

Firstly, let us consider the equation

$$\begin{cases} \Delta \bar{U} = h & \text{in } \mathbb{T}^3, \\ (\bar{U})_{\mathbb{T}^3} = 0, \end{cases} \quad (3.2)$$

where $h \in L^2(\mathbb{T}^3)$. We say that $\bar{U} \in H^1(\mathbb{T}^3)$ is a weak solution to (3.2) provided that

$$-\int_{\mathbb{T}^3} \nabla \bar{U} \cdot \nabla \psi dx = \int_{\mathbb{T}^3} h \psi dx, \quad \forall \psi \in H^1(\mathbb{T}^3). \quad (3.3)$$

To prove the existence and uniqueness of the weak solution to (3.2), we only need to notice that

$$\|\overline{U}\|_{L^2(\mathbb{T}^3)} = \|\overline{U} - (\overline{U})_{\mathbb{T}^3}\|_{L^2(\mathbb{T}^3)} \leq C\|\nabla\overline{U}\|_{L^2(\mathbb{T}^3)}$$

by using Poincaré's inequality. Then by Lax-Milgram's Theorem, there exists a unique $\overline{U} \in H^1(\mathbb{T}^3)$ satisfying (3.3) and hence the weak solution to (3.2). In addition, it holds that $\|\overline{U}\|_{H^1(\mathbb{T}^3)} \leq C\|h\|_{L^2(\mathbb{T}^3)}$. For the regularity of \overline{U} , recall Section 9.4 in [13] for the Newtonian potential $w := \mathbb{N}h$ of h . Whenever $h \in L^2(\mathbb{T}^3)$, we have $w \in H^2(\mathbb{T}^3)$ and $\Delta w = h$ holds a.e. \mathbb{T}^3 . Moreover, there exists a constant C such that $\|D^2 w\|_{L^2(\mathbb{T}^3)} \leq C\|h\|_{L^2(\mathbb{T}^3)}$. By the uniqueness of \overline{U} , we know that $\overline{U} \equiv w - (w)_{\mathbb{T}^3}$. Hence we have $\|D^2 \overline{U}\|_{L^2(\mathbb{T}^3)} \leq C\|h\|_{L^2(\mathbb{T}^3)}$, which implies that $\|\overline{U}\|_{H^2(\mathbb{T}^3)} \leq C\|h\|_{L^2(\mathbb{T}^3)}$.

Therefore, when $h = 1 - \rho$, we have $\|\overline{U}\|_{H^2(\mathbb{T}^3)} \leq C\|1 - \rho\|_{L^2(\mathbb{T}^3)}$.

Secondly, to solve the equation

$$\Delta \hat{U} = e^{\overline{U} + \hat{U}} - 1 \quad (3.4)$$

with $\overline{U} \in H^2(\mathbb{T}^3)$, in light of [15], we use a direct method of calculus of variations. Define the *Lagrangian*

$$\mathcal{L}(p, z, x) = \frac{1}{2}|p|^2 + e^{\overline{U} + z} - z \in L^1(\mathbb{T}^3),$$

and consider the energy functional $\mathfrak{J}[\cdot]$ on $H^1(\mathbb{T}^3)$,

$$h \mapsto \mathfrak{J}[h] := \int_{\mathbb{T}^3} \frac{1}{2}|\nabla h|^2 + (e^{\overline{U} + h} - h)dx.$$

Then (3.4) is the corresponding Euler-Lagrange equation. Since $\mathcal{L}(p, z, x)$ is uniformly convex in the variable p and convex in the joint variables (p, z) , solutions of the Euler-Lagrange equation are the minimisers of $\mathfrak{J}[\cdot]$ and the minimiser is unique. Thus we seek for the *unique* minimiser which solves (3.4). To this end, noting that

$$\mathfrak{J}[h] = \int_{\mathbb{T}^3} \frac{1}{2}|\nabla h|^2 + e^{\overline{U} + h} - (\overline{U} + h) + \overline{U}dx \geq \int_{\mathbb{T}^3} \overline{U}dx \geq -C\|\overline{U}\|_{L^2(\mathbb{T}^3)},$$

which means that $\mathfrak{J}[\cdot]$ possesses a lower bound and hence the infimum

$$\inf_{H^1(\mathbb{T}^3)} \mathfrak{J}[\cdot] := \alpha > -\infty.$$

Considering a minimising sequence $\{h_k\}_{k \in \mathbb{N}}$ such that

$$\lim_{k \rightarrow \infty} \mathfrak{J}[h_k] = \inf_{H^1(\mathbb{T}^3)} \mathfrak{J}[\cdot] = \alpha,$$

we now prove that $\{h_k\}_{k \in \mathbb{N}}$ is uniformly bounded in $H^1(\mathbb{T}^3)$ and its weak limit is exactly the minimiser of $\mathfrak{J}[\cdot]$.

Indeed, on one hand,

$$\alpha \leq \mathfrak{J}[-\overline{U}] = \int_{\mathbb{T}^3} \frac{1}{2}|\nabla \overline{U}|^2 + 1 + \overline{U}dx \leq C$$

yields

$$\mathfrak{J}[h_k] \leq \alpha + 1 \leq C + 1 =: C_1$$

for sufficiently large k . On the other hand, noticing that $\|\overline{U}\|_{L^\infty(\mathbb{T}^3)} \leq C\|\overline{U}\|_{H^2(\mathbb{T}^3)} < \infty$, then it holds

$$e^{\overline{U} + s} - s \geq e^{-\|\overline{U}\|_{L^\infty(\mathbb{T}^3)}} e^s - s \geq |s| - C_2$$

for some large constant $C_2 > 0$. Thus,

$$\mathfrak{J}[h_k] \geq \int_{\mathbb{T}^3} \left(\frac{1}{2}|\nabla h_k|^2 + |h_k| - C_2 \right) dx.$$

In addition, by Poincaré's inequality and Young's inequality, we have

$$\begin{aligned} \|h_k\|_{L^2(\mathbb{T}^3)} &\leq C\|\nabla h_k\|_{L^2(\mathbb{T}^3)} + \|(h_k)_{\mathbb{T}^3}\|_{L^2(\mathbb{T}^3)} \\ &\leq \epsilon\|\nabla h_k\|_{L^2(\mathbb{T}^3)}^2 + C_\epsilon + C \int_{\mathbb{T}^3} |h_k| dx \end{aligned}$$

for some small $\epsilon > 0$. Hence,

$$\mathfrak{J}[h_k] \geq \int_{\mathbb{T}^3} \left(\frac{1}{4} |\nabla h_k|^2 + |h_k|^2 - C_2 - C_\epsilon \right) dx.$$

Combining the upper and lower bounds of $\mathfrak{J}[h_k]$ yields

$$\|h_k\|_{H^1(\mathbb{T}^3)}^2 \leq C(C_1 + C_2 + C_\epsilon) < \infty.$$

Denote the weak limit of $\{h_k\}$ (up to a subsequence) by \hat{U} : $h_k \rightharpoonup \hat{U}$ in $H^1(\mathbb{T}^3)$. It follows from $H^1(\mathbb{T}^3) \subset\subset L^2(\mathbb{T}^3)$ that $h_k \rightarrow \hat{U}$ in $L^2(\mathbb{T}^3)$ and hence $h_k \rightarrow \hat{U}$ a.e. \mathbb{T}^3 .

We claim that \hat{U} is a minimiser. In fact, by the weak lower semi-continuity of $\{\nabla h_k\}$ in L^2 , strong convergence of $\{h_k\}$ in L^1 and Fatou's lemma for $\{e^{\bar{U}+h_k}\}$, it holds

$$\alpha = \lim_{k \rightarrow \infty} \mathfrak{J}[h_k] = \liminf_{k \rightarrow \infty} \mathfrak{J}[h_k] \geq \mathfrak{J}[\hat{U}] \geq \alpha.$$

Therefore, $\hat{U} \in H^1(\mathbb{T}^3)$ solves (3.4) in weak sense. For any $\psi \in C^\infty(\mathbb{T}^3)$, invoking the dominated convergence theorem concludes that

$$0 = \lim_{\tau \rightarrow 0} \frac{\mathfrak{J}[\hat{U} + \tau\psi] - \mathfrak{J}[\hat{U}]}{\tau} = \int_{\mathbb{T}^3} \left\{ \nabla \hat{U} \cdot \nabla \psi + \left(e^{\bar{U} + \hat{U}} - 1 \right) \psi \right\} dx. \quad (3.5)$$

Note that if $e^{\bar{U} + \hat{U}} \in L^2(\mathbb{T}^3)$, then (3.5) is valid for $\psi \in H^1(\mathbb{T}^3)$. Thus, \hat{U} is proved to be the unique weak solution to (3.4). Moreover, by (3.4), the regularity of \hat{U} can be enhanced:

$$\|\hat{U}\|_{H^2(\mathbb{T}^3)} \leq C \|e^{\bar{U} + \hat{U}} - 1\|_{L^2(\mathbb{T}^3)} < \infty.$$

For this purpose, it suffices to show that $e^{\hat{U}} \in L^2(\mathbb{T}^3)$. Actually, for $\hat{U} \in H^1(\mathbb{T}^3)$, consider the truncated function $\hat{U}_k := \min\{\hat{U}, k\}$, $k \in \mathbb{N}$. Thus $e^{\hat{U}_k} \in L^\infty \cap H^1(\mathbb{T}^3)$, which can be tested by the equation (3.4). It follows that

$$0 = \int_{\mathbb{T}^3} \left(\nabla \hat{U} \cdot \nabla e^{\hat{U}_k} + e^{\bar{U} + \hat{U}} e^{\hat{U}_k} - e^{\hat{U}_k} \right) dx \geq C \int_{\mathbb{T}^3} e^{\hat{U} + \hat{U}_k} dx - \int_{\mathbb{T}^3} e^{\hat{U}_k} dx.$$

By using the monotone convergence theorem, we can take $k \rightarrow \infty$ to get

$$\int_{\mathbb{T}^3} e^{\hat{U}} dx \geq C \int_{\mathbb{T}^3} e^{2\hat{U}} dx.$$

Meanwhile,

$$0 = \int_{\mathbb{T}^3} \Delta \hat{U} dx = \int_{\mathbb{T}^3} e^U - 1 dx$$

implies that

$$1 \geq e^{-\|\bar{U}\|_{L^\infty}} \int_{\mathbb{T}^3} e^{\hat{U}} dx.$$

Hence $e^{\hat{U}} \in L^1(\mathbb{T}^3)$ and then $e^{\hat{U}} \in L^2(\mathbb{T}^3)$.

In conclusion, we obtain that both \bar{U} and \hat{U} are in $H^2(\mathbb{T}^3)$, so $U = \bar{U} + \hat{U} \in H^2(\mathbb{T}^3)$ is the unique strong solution to (3.1) for $\rho \in L^2(\mathbb{T}^3)$. \square

Based on the above regularity result of U , i.e. $U \in H^2(\mathbb{T}^3) \leftrightarrow L^\infty(\mathbb{T}^3) \cap C(\mathbb{T}^3)$, we can further deduce the following

Proposition 3.2. *Suppose that $U \in H^2(\mathbb{T}^3)$ solves (3.1) with $\rho \in L^2(\mathbb{T}^3)$. Then it holds*

$$\|U\|_{H^2(\mathbb{T}^3)} \leq C \|\rho - 1\|_{L^2(\mathbb{T}^3)}$$

for some constant $C = C(\|U\|_{L^\infty(\mathbb{T}^3)}) > 0$. Moreover, when $\|\rho - 1\|_{L^2(\mathbb{T}^3)}$ is sufficiently small, the constant C is independent of $\|U\|_{L^\infty(\mathbb{T}^3)}$.

Proof. Since $U \in L^\infty(\mathbb{T}^3) \cap C(\mathbb{T}^3)$, then for each $x \in \mathbb{T}^3$, there exists a uniformly bounded function $U_\xi : x \mapsto U_\xi(x)$ such that $e^{U(x)} - 1 = U(x)e^{U_\xi(x)}$, i.e., $e^U - 1 = Ue^{U_\xi}$ in \mathbb{T}^3 , where $e^{-\|U\|_{L^\infty(\mathbb{T}^3)}} \leq e^{U_\xi} \leq e^{\|U\|_{L^\infty(\mathbb{T}^3)}}$. The equation (3.1) can thus be rewritten as

$$\Delta U = Ue^{U_\xi} + 1 - \rho.$$

Taking L^2 inner product of the above equation with U yields

$$\int_{\mathbb{T}^3} |\nabla U|^2 + |U|^2 e^{U_\xi} dx = \int_{\mathbb{T}^3} U(\rho - 1) dx,$$

which implies that $\|U\|_{H^1(\mathbb{T}^3)} \leq e^{\|U\|_{L^\infty(\mathbb{T}^3)}} \|\rho - 1\|_{L^2(\mathbb{T}^3)}$. With the standard elliptic estimates, we deduce that

$$\begin{aligned} \|D^2 U\|_{L^2(\mathbb{T}^3)} &\leq C \|U\|_{L^2(\mathbb{T}^3)} e^{\|U\|_{L^\infty(\mathbb{T}^3)}} + C \|\rho - 1\|_{L^2(\mathbb{T}^3)} \\ &\leq C(e^{2\|U\|_{L^\infty(\mathbb{T}^3)}} + 1) \|\rho - 1\|_{L^2(\mathbb{T}^3)}. \end{aligned}$$

Hence we get

$$\|U\|_{H^2(\mathbb{T}^3)} \leq C e^{2\|U\|_{L^\infty(\mathbb{T}^3)}} \|\rho - 1\|_{L^2(\mathbb{T}^3)},$$

which implies that

$$\|U\|_{L^\infty(\mathbb{T}^3)} \leq C e^{2\|U\|_{L^\infty(\mathbb{T}^3)}} \|\rho - 1\|_{L^2(\mathbb{T}^3)}.$$

When $\|\rho - 1\|_{L^2(\mathbb{T}^3)}$ is of order $\varpi \ll 1$, we have $\|U\|_{L^\infty(\mathbb{T}^3)} = O(\varpi)$. Then we have

$$\|U\|_{H^2(\mathbb{T}^3)} \leq C e^{2C\varpi} \|\rho - 1\|_{L^2(\mathbb{T}^3)} \leq C \|\rho - 1\|_{L^2(\mathbb{T}^3)}.$$

□

Corollary 3.3. *Suppose that $U_i \in H^2(\mathbb{T}^3)$ solves (3.1) with $\rho_i \in L^2(\mathbb{T}^3)$ for $i = 1, 2$. Then,*

$$\|U_1 - U_2\|_{H^2(\mathbb{T}^3)} \leq C \|\rho_1 - \rho_2\|_{L^2(\mathbb{T}^3)}$$

holds for some constant $C > 0$.

Proof. Take the difference for U_1 and U_2 to get

$$\Delta(U_1 - U_2) = e^{U_1} - e^{U_2} - (\rho_1 - \rho_2) = (U_1 - U_2)e^{\bar{U}_\xi} - (\rho_1 - \rho_2),$$

where the value of $\bar{U}_\xi(x)$ lies between $U_1(x)$ and $U_2(x)$ for every $x \in \mathbb{T}^3$ and hence $|\bar{U}_\xi|$ is uniformly bounded. Through the similar arguments to that in Proposition 3.2, we obtain that

$$\|D^2(U_1 - U_2)\|_{L^2(\mathbb{T}^3)} + \|U_1 - U_2\|_{H^1(\mathbb{T}^3)} \leq C \|\rho_1 - \rho_2\|_{L^2(\mathbb{T}^3)}.$$

□

Our next intention is to iterate the argument above to deduce that the solution U lies in higher order Sobolev spaces provided that ρ has sufficient regularities.

Proposition 3.4. *Let m be a nonnegative integer and the multi-index $\gamma = [\gamma_0, \gamma_1, \gamma_2, \gamma_3]$ with $|\gamma| \leq m$. Assume that for any $t \geq 0$,*

$$\sum_{|\gamma| \leq m} \|\partial^\gamma \rho(t, \cdot)\|_{L^2(\mathbb{T}^3)} < \infty. \quad (3.6)$$

Suppose that $U \in H^1(\mathbb{T}^3)$ is a weak solution to the equation (3.1). Then for any $t \geq 0$, $\partial^\gamma U \in H^2(\mathbb{T}^3)$, and

$$\sum_{|\gamma| \leq m} \|\partial^\gamma U(t, \cdot)\|_{H^2(\mathbb{T}^3)} \leq C \sum_{|\gamma| \leq m} \|\partial^\gamma (\rho(t, \cdot) - 1)\|_{L^2(\mathbb{T}^3)}, \quad (3.7)$$

where the constant C is independent of U whenever $\|\rho - 1\|_{L^2(\mathbb{T}^3)} \ll 1$.

Proof. We adopt the principle of mathematical induction. Since $\Delta U - e^U + \rho \equiv 0$ holds for every $t \geq 0$ and almost every $x \in \mathbb{T}^3$ by Proposition 3.2, we take ∂^γ derivatives ($|\gamma| = 1$) to the identity in the sense of distribution to get

$$\Delta(\partial^\gamma U) - (\partial^\gamma U)e^U + \partial^\gamma \rho = 0.$$

By applying Lax-Milgram's Theorem and elliptic estimates as in Proposition 3.2, we obtain that there exists a unique solution $\psi_\gamma \in H^2(\mathbb{T}^3)$ satisfying

$$\Delta\psi_\gamma - e^U\psi_\gamma + \partial^\gamma \rho \equiv 0,$$

for every $t \geq 0$ and almost every $x \in \mathbb{T}^3$. We thus denote ψ_γ by $\partial^\gamma U$. Then we have

$$\|\partial^\gamma U\|_{H^2(\mathbb{T}^3)} \leq C\|\partial^\gamma \rho\|_{L^2(\mathbb{T}^3)}, \quad \text{for } \forall t \geq 0, \quad |\gamma| = 1.$$

Whenever $\|\rho - 1\|_{L^2(\mathbb{T}^3)}$ is sufficiently small, the above constant C is independent of U , as in Proposition 3.2.

Assume now the assertion (3.7) is valid for some integer $m \geq 1$. Suppose that (3.6) holds for $m + 1$. Then, it follows

$$\Delta\partial^\gamma U - e^U\partial^\gamma U = [\partial^\gamma(e^U) - e^U\partial^\gamma U] - \partial^\gamma \rho =: S - \partial^\gamma \rho,$$

where $|\gamma| = m + 1$, and

$$S := e^U \sum_{\substack{0 < \beta_1 < \gamma \\ 0 < \beta_2 < \gamma - \beta_1 \\ \gamma = \beta_1 + \dots + \beta_k}} C_\gamma^{\beta_1} \dots C_{\gamma - (\beta_1 + \dots + \beta_{k-1})}^{\beta_k} \prod_{i=1}^k \partial^{\beta_i} U, \quad k \geq 2.$$

By using (3.7) for the integer m , we know that $\|S\|_{L^2(\mathbb{T}^3)}$ is also controlled by the right-hand side of (3.7). For every $v \in H^1(\mathbb{T}^3)$, it holds

$$\int_{\mathbb{T}^3} \nabla\partial^\gamma U \cdot \nabla v + e^U\partial^\gamma U v dx = \int_{\mathbb{T}^3} (S - \partial^\gamma \rho)v dx.$$

Applying Lax-Milgram's Theorem and elliptic estimates again, we obtain that there exists a unique $\partial^\gamma U \in H^2(\mathbb{T}^3)$ satisfying

$$\|\partial^\gamma U\|_{H^2(\mathbb{T}^3)} \leq C\|S - \partial^\gamma \rho\|_{L^2(\mathbb{T}^3)} \leq C \left\{ \sum_{1 \leq |\gamma| \leq m+1} \|\partial^\gamma \rho(t, \cdot)\|_{L^2(\mathbb{T}^3)} + \|\rho - 1\|_{L^2(\mathbb{T}^3)} \right\},$$

which indicates (3.7) holds for $|\gamma| = m + 1$, combining with the m case. Inductively, this completes the proof. \square

4. LOCAL-IN-TIME WELL-POSEDNESS TO THE SYSTEM (1.3)–(1.4)

In this section, we construct a unique local solution to the ionic Vlasov-Poisson-Boltzmann system (1.3)–(1.4) with a uniform energy estimate as follows.

Theorem 4.1. *There exist $M_0 > 0$ and $T_0 > 0$ such that if $T_0 \leq M_0/2$ and $\mathcal{E}(0) \leq M_0/2$, then the ionic Vlasov-Poisson-Boltzmann system (1.3)–(1.4) has a unique solution $f(t, x, v)$ in $[0, T_0] \times \mathbb{T}^3 \times \mathbb{R}^3$ satisfying*

$$\sup_{0 \leq t \leq T_0} \mathcal{E}(t) \leq M_0,$$

and $\mathcal{E}(t)$ is continuous over $[0, T_0]$. If $F_0(x, v) = \mu(v) + \sqrt{\mu(v)}f_0(x, v) \geq 0$, then $F(t, x, v) = \mu(v) + \sqrt{\mu(v)}f(t, x, v) \geq 0$. Furthermore, the conservation laws (1.6)–(1.8) are valid for all $0 < t \leq T_0$ if they hold initially.

Proof. Consider the iterating sequence $\{F^n\}_{n \in \mathbb{N}}$ which solves

$$\begin{aligned} \{\partial_t + v \cdot \nabla_x + E^n \cdot \nabla_v + R(F^n)\}F^{n+1} &= Q_{\text{gain}}(F^n, F^n) \\ &= \iint_{\mathbb{R}^3 \times \mathbb{S}^2} |u - v|^r b_0(\theta) F^n(v') F^n(u') dud\omega \end{aligned} \quad (4.1)$$

with the initial datum $F^{n+1}(0, x, v) = F_0(x, v)$, $n \in \mathbb{N}$. Here,

$$R(F^n) := \iint_{\mathbb{R}^3 \times \mathbb{S}^2} |u - v|^r b_0(\theta) F^n(u) dud\omega.$$

We take $F^0(t, x, v) \equiv F_0(x, v)$. Since $F^n = \mu + \sqrt{\mu} f^n$ for $n \in \mathbb{N}$, we have $f^0(t, x, v) \equiv f_0(x, v)$ and $f^{n+1}(0, x, v) = f_0(x, v)$. Equivalently to (4.1), f^{n+1} satisfies

$$\left\{ \partial_t + v \cdot \nabla_x + E^n \cdot \nabla_v + \nu(v) - \frac{v}{2} \cdot E^n \right\} f^{n+1} - E^n \cdot v \sqrt{\mu} = \mathcal{K} f^n + \Gamma_{\text{gain}}(f^n, f^n) - \Gamma_{\text{loss}}(f^n, f^{n+1}). \quad (4.2)$$

We want to show that, there exist $M_0 > 0$ and $T_0 > 0$ such that if $T_0 \leq M_0/2$, $\mathcal{E}(0) \leq M_0/2$ and $\sup_{0 \leq t \leq T_0} \mathcal{E}_n(t) \leq M_0$, then $\sup_{0 \leq t \leq T_0} \mathcal{E}_{n+1}(t) \leq M_0$. Hence we can verify the local-in-time existence in Theorem 4.1 inductively, due to the fact that $\sup_{0 \leq t \leq T_0} \mathcal{E}_0(t) = \mathcal{E}(0) \leq M_0/2$.

Taking ∂_β^γ derivatives of (4.2), we have

$$\begin{aligned} & \left\{ \partial_t + v \cdot \nabla_x + E^n \cdot \nabla_v - \frac{v}{2} \cdot E^n \right\} \partial_\beta^\gamma f^{n+1} + \partial_\beta [\nu \partial^\gamma f^{n+1}] \\ &= \partial^\gamma E^n \cdot \partial_\beta (v \sqrt{\mu}) - \sum_{\beta_1 \neq 0} C_\beta^{\beta_1} \partial_{\beta_1} v \cdot \nabla_x \partial_{\beta - \beta_1}^\gamma f^{n+1} + \partial_\beta \{ \mathcal{K} \partial^\gamma f^n \} \\ &+ \sum_{\gamma_1, \beta_1 \neq 0} C_\beta^{\beta_1} C_\gamma^{\gamma_1} \partial^{\gamma_1} E^n \cdot \partial_{\beta_1} \left(\frac{v}{2} \right) \partial_{\beta - \beta_1}^{\gamma - \gamma_1} f^{n+1} - \sum_{\gamma_1 \neq 0} C_\gamma^{\gamma_1} \partial^{\gamma_1} E^n \cdot \nabla_v \partial_\beta^{\gamma - \gamma_1} f^{n+1} \\ &+ \sum C_\gamma^{\gamma_1} \partial_\beta \{ \Gamma_{\text{gain}}(\partial^{\gamma_1} f^n, \partial^{\gamma - \gamma_1} f^n) - \Gamma_{\text{loss}}(\partial^{\gamma_1} f^n, \partial^{\gamma - \gamma_1} f^{n+1}) \}. \end{aligned} \quad (4.3)$$

To deal with the possible degeneration raised from the collision potentials, we define the weight function $W \equiv W(x, v) := e^{-q \frac{x \cdot v}{\langle v \rangle}}$ for $q \gg 1$. Now we take L^2 inner product of (4.3) with $W^2 \partial_\beta^\gamma f^{n+1}$, $|\gamma| + |\beta| \leq N$, in $\mathbb{T}^3 \times \mathbb{R}^3$ and deal with the result term by term as follows.

(i). For the left-hand side of (4.3), from

$$-\frac{\{\partial_t + v \cdot \nabla_x + E^n \cdot \nabla_v\} W^2}{2W^2} = q \frac{|v|^2}{\langle v \rangle} + q E^n \cdot \left(\frac{x}{\langle v \rangle} + \frac{(x \cdot v)v}{\langle v \rangle^3} \right),$$

we have

$$\begin{aligned} & W^2 \partial_\beta^\gamma f^{n+1} \left\{ \partial_t + v \cdot \nabla_x + E^n \cdot \nabla_v - \frac{v}{2} \cdot E^n \right\} \partial_\beta^\gamma f^{n+1} \\ &= \{\partial_t + v \cdot \nabla_x + E^n \cdot \nabla_v\} \left(\frac{1}{2} |W \partial_\beta^\gamma f^{n+1}|^2 \right) \\ &+ \left\{ 2q \frac{|v|^2}{\langle v \rangle} + 2E^n \cdot \left(q \frac{x}{\langle v \rangle} + q \frac{(x \cdot v)v}{\langle v \rangle^3} - \frac{v}{4} \right) \right\} |W \partial_\beta^\gamma f^{n+1}|^2 \\ &=: \{\partial_t + v \cdot \nabla_x + E^n \cdot \nabla_v\} \left(\frac{1}{2} |W \partial_\beta^\gamma f^{n+1}|^2 \right) + \Xi^n |W \partial_\beta^\gamma f^{n+1}|^2. \end{aligned}$$

Integrating the above equality over $\mathbb{T}^3 \times \mathbb{R}^3$ yields

$$\begin{aligned} & \iint_{\mathbb{T}^3 \times \mathbb{R}^3} W^2 \partial_\beta^\gamma f^{n+1} \left\{ \partial_t + v \cdot \nabla_x + E^n \cdot \nabla_v - \frac{v}{2} \cdot E^n \right\} \partial_\beta^\gamma f^{n+1} dx dv \\ &= \frac{1}{2} \frac{d}{dt} \|W \partial_\beta^\gamma f^{n+1}\|^2(t) + \iint_{\mathbb{T}^3 \times \mathbb{R}^3} \Xi^n |W \partial_\beta^\gamma f^{n+1}|^2 dx dv. \end{aligned}$$

By (2.2) in Lemma 2.2 and $W \sim 1$, we have

$$\begin{aligned} & \iint_{\mathbb{T}^3 \times \mathbb{R}^3} W^2 \partial_\beta^\gamma f^{n+1} \partial_\beta [\nu \partial^\gamma f^{n+1}] dx dv \\ & \geq \|\partial_\beta^\gamma f^{n+1}\|_\nu^2 - \eta \sum_{|\beta_1| \leq |\beta|} \|\partial_{\beta_1}^\gamma f^{n+1}\|^2 - C_\eta \|\partial^\gamma f^{n+1}\|^2 \end{aligned}$$

for some small $\eta > 0$. Then the contribution on the left-hand side of (4.3) is

$$\frac{1}{2} \frac{d}{dt} \|W \partial_\beta^\gamma f^{n+1}(t)\|^2 + \iint_{\mathbb{T}^3 \times \mathbb{R}^3} [\nu + \Xi^n] |W \partial_\beta^\gamma f^{n+1}|^2 dx dv - \eta \sum_{|\beta_1| \leq |\beta|} \|\partial_{\beta_1}^\gamma f^{n+1}\|^2 - C_\eta \|\partial^\gamma f^{n+1}\|^2.$$

Since $\nu(v) \sim \langle v \rangle^r$ and

$$\left| E^n \cdot \left(q \frac{x}{\langle v \rangle} + q \frac{(x \cdot v)v}{\langle v \rangle^3} - \frac{v}{2} \right) \right| \leq C \|E^n\|_\infty \langle v \rangle \leq CM_0 \langle v \rangle,$$

then, for $M_0 \ll 1$ and $q \gg 1$, it holds

$$\nu + \Xi^n := \nu + q \frac{|v|^2}{\langle v \rangle} + E^n \cdot \left(q \frac{x}{\langle v \rangle} + q \frac{(x \cdot v)v}{\langle v \rangle^3} - \frac{v}{2} \right) \sim \langle v \rangle.$$

Thus

$$\iint_{\mathbb{T}^3 \times \mathbb{R}^3} [\nu + \Xi^n] |W \partial_\beta^\gamma f^{n+1}|^2 dx dv \sim \iint_{\mathbb{T}^3 \times \mathbb{R}^3} \langle v \rangle |W \partial_\beta^\gamma f^{n+1}|^2 dx dv.$$

(ii). For the right-hand side of (4.3), we handle them one by one.

(ia). The first term contributes

$$\iint_{\mathbb{T}^3 \times \mathbb{R}^3} \partial^\gamma E^n \cdot \partial_\beta (v \sqrt{\mu}) \partial_\beta^\gamma f^{n+1} dx dv \leq C \| \|E^n\| \| \|f^{n+1}\| \|.$$

(iib). By using Lemma 2.2, the inner product of $\sum_{\beta_1 \neq 0} C_{\beta_1}^{\beta_1} \partial_{\beta_1} v \cdot \nabla_x \partial_{\beta - \beta_1}^\gamma f^{n+1}$ and $\partial_\beta \{K \partial^\gamma f^n\}$ can be controlled by $(\| \|f^n\| \| + \| \|f^{n+1}\| \|) \| \|f^{n+1}\| \|$.

(iic). The second line on the right-hand side is controlled by

$$\begin{aligned} & C \sum_{\gamma_1 \neq 0, |\beta_1|=1} \int_{\mathbb{T}^3} |\partial^{\gamma_1} E^n| \left(\int_{\mathbb{R}^3} |\partial_{\beta - \beta_1}^{\gamma - \gamma_1} f^{n+1}|^2 dv \right)^{\frac{1}{2}} \left(\int_{\mathbb{R}^3} |\partial_\beta^\gamma f^{n+1}|^2 dv \right)^{\frac{1}{2}} dx \\ & + C \sum_{\gamma_1 \neq 0} \int_{\mathbb{T}^3} |\partial^{\gamma_1} E^n| \left(\int_{\mathbb{R}^3} |\nabla_v \partial_\beta^{\gamma - \gamma_1} f^{n+1}|^2 dv \right)^{\frac{1}{2}} \left(\int_{\mathbb{R}^3} |\partial_\beta^\gamma f^{n+1}|^2 dv \right)^{\frac{1}{2}} dx. \end{aligned} \quad (4.4)$$

Note that

$$\sup_x \int_{\mathbb{R}^3} |g(x, u)|^2 du \leq C \sum_{|\gamma| \leq 2} \|\partial^\gamma g\|^2.$$

Thus, when $|\gamma_1| \geq 3$, $|\gamma - \gamma_1| + 2 \leq |\gamma| - 1$, the second term of (4.4) is controlled by

$$C \|\partial^{\gamma_1} E^n\| \sum_{|\gamma - \gamma_1| + 1 + 2 \leq |\gamma|} \|\nabla_v \partial_\beta^{\gamma - \gamma_1} f^{n+1}\| \| \|f^{n+1}\| \| \leq C \|\partial^{\gamma_1} E^n\| \| \|f^{n+1}\| \|^2 \leq C \| \|E^n\| \| \| \|f^{n+1}\| \|^2.$$

When $|\gamma_1| \leq 2$, $|\gamma_1| + 2 \leq 4 \leq N$, the second term of (4.4) is controlled by

$$C \|\partial^{\gamma_1} E^n\|_{H^2(\mathbb{T}^3)} \| \|f^{n+1}\| \|^2 \leq C \| \|E^n\| \| \| \|f^{n+1}\| \|^2.$$

Adopting the similar argument to the first term of (4.4), we deduce that (4.4) can be controlled by

$$C \| \|E^n\| \| \| \|f^{n+1}\| \|^2.$$

(iid). By using Lemma 2.4, the contribution of the last line on (4.3) is

$$C (\| \|f^n\| \| \| \|f^n\|_\nu \| \|f^{n+1}\|_\nu \| + \| \|f^{n+1}\| \| \| \|f^n\|_\nu \| \|f^{n+1}\|_\nu \| + \| \|f^n\| \| \| \|f^{n+1}\|_\nu \|^2).$$

Collecting the above estimates up and summing over $|\beta| + |\gamma| \leq N$, we get

$$\begin{aligned} \mathcal{E}'_{n+1}(t) &= \frac{1}{2} \frac{d}{dt} \| \|f^{n+1}\| \|^2 + \| \|f^{n+1}\|_{\langle v \rangle}^2 \\ &\leq C \| \|E^n\| \| (\| \|f^{n+1}\| \| + \| \|f^{n+1}\| \|^2) + C (\| \|f^n\| \| + \| \|f^{n+1}\| \|) \| \|f^{n+1}\| \| \\ &\quad + C (\| \|f^n\| \| \| \|f^n\|_\nu \| \| \|f^{n+1}\|_\nu \| + \| \|f^{n+1}\| \| \| \|f^n\|_\nu \| \| \|f^{n+1}\|_\nu \| + \| \|f^n\| \| \| \|f^{n+1}\|_\nu \|^2) \\ &\leq C \{ \| \|f^n\| \|^2 + \| \|f^{n+1}\| \|^2 + \| \|f^n\| \| (\| \|f^n\|_\nu \|^2 + \| \|f^{n+1}\|_\nu \|^2 + \| \|f^{n+1}\| \|^2) \} \\ &\quad + C \| \|f^{n+1}\| \| \| \|f^n\|_\nu \| \| \|f^{n+1}\|_\nu \|, \end{aligned} \quad (4.5)$$

where the last inequality relies on the elliptic estimates established in Proposition 3.4 that $\|E^n\| \leq C\|f^n\|$. To investigate the term $\|f^{n+1}\| \|f^n\|_\nu \|f^{n+1}\|_\nu$, we notice that

$$\begin{aligned} \int_0^t \|f^{n+1}\| \|f^n\|_\nu \|f^{n+1}\|_\nu ds &\leq \sup_{0 \leq s \leq t} \|f^{n+1}\|(s) \left\{ \int_0^t \|f^n\|_{\langle v \rangle}^2 ds \right\}^{1/2} \left\{ \int_0^t \|f^{n+1}\|_{\langle v \rangle}^2 ds \right\}^{1/2} \\ &\leq M_0^{1/2} \sup_{0 \leq s \leq t} \mathcal{E}_{n+1}(s). \end{aligned}$$

Hence, integrating (4.5) over $[0, t]$ yields

$$\mathcal{E}_{n+1}(t) \leq \mathcal{E}_{n+1}(0) + C \left\{ tM_0 + t \sup_{0 \leq s \leq t} \mathcal{E}_{n+1}(s) + M_0^{3/2} + M_0^{1/2} \sup_{0 \leq s \leq t} \mathcal{E}_{n+1}(s) \right\}.$$

With the initial datum $f^{n+1}(0, x, v) = f_0(x, v)$, we inductively deduce that $\partial_\beta^\gamma f^{n+1}(0, x, v) = \partial_\beta^\gamma f_0(x, v)$ over the number of temporal derivatives. Thus $\mathcal{E}_{n+1}(0) \equiv \mathcal{E}(f_0) \leq M_0/2$. For $0 < t \leq T_0$, it holds

$$\begin{aligned} (1 - CT_0 - CM_0^{1/2}) \sup_{0 \leq t \leq T_0} \mathcal{E}_{n+1}(t) &\leq \mathcal{E}_{n+1}(0) + C\{M_0T_0 + M_0^{3/2}\} \\ &\leq M_0/2 + C\{M_0T_0 + M_0^{3/2}\}. \end{aligned}$$

If $T_0 \leq M_0/2 \ll 1$, the local-in-time existence in Theorem 4.1 is proved for sufficiently large $n \in \mathbb{N}$. We denote the solution to the ionic Vlasov-Poisson-Boltzmann system (1.3)–(1.4) by f .

Next, we sketch the argument for proving the uniqueness of solution f . If there is another solution g such that $\sup_{0 \leq t \leq T_0} \mathcal{E}(g)(t) \leq M_0$, we investigate the L^2 energy of $f - g$ via the equation

$$\begin{aligned} &\{\partial_t + v \cdot \nabla_x + E_f \cdot \nabla_v + \nu\}(f - g) \\ &= (E_f - E_g) \cdot v \sqrt{\mu} - (E_f - E_g) \cdot \nabla_v g + \frac{v}{2} \cdot E_f(f - g) + \frac{v}{2} \cdot (E_f - E_g)g \\ &\quad + \mathcal{K}(f - g) + \Gamma(f - g, f) + \Gamma(g, f - g) \end{aligned} \tag{4.6}$$

with $(f - g)|_{t=0} = f(0, x, v) - g(0, x, v) = 0$, where (E_f, ϕ_f) and (E_g, ϕ_g) satisfy

$$\begin{cases} E_f = -\nabla_x \phi_f, \\ \Delta \phi_f = e^{\phi_f} - 1 - \int_{\mathbb{R}^3} \sqrt{\mu} f dv, \end{cases}$$

and

$$\begin{cases} E_g = -\nabla_x \phi_g, \\ \Delta \phi_g = e^{\phi_g} - 1 - \int_{\mathbb{R}^3} \sqrt{\mu} g dv, \end{cases}$$

respectively. Recalling Corollary 3.3, we get the following crucial estimate

$$\|\phi_f - \phi_g\|_{H^2(\mathbb{T}^3)} \leq C\|f - g\|_{L^2(\mathbb{T}^3 \times \mathbb{R}^3)}.$$

On the other hand, for the weight function $W = e^{-q\frac{x \cdot v}{\langle v \rangle}} \sim 1$, we write

$$-\frac{\{\partial_t + v \cdot \nabla_x + E_f \cdot \nabla_v\}W^2}{2W^2} = q\frac{|v|^2}{\langle v \rangle} + qE_f \cdot \left(\frac{x}{\langle v \rangle} + \frac{(x \cdot v)v}{\langle v \rangle^3} \right) := \Xi_f,$$

and it holds that $\nu + \Xi_f \sim \langle v \rangle$.

Taking the inner product of (4.6) with $W^2(f - g)$ and noticing Remark 2.5, we deduce that

$$\begin{aligned} &\frac{1}{2} \frac{d}{dt} \|f - g\|^2 + \|f - g\|_{\langle v \rangle}^2 \\ &\leq \|E_f - E_g\| \|f - g\| + \sqrt{M_0} \|f - g\|_{\langle v \rangle}^2 + \|g\|_{\langle v \rangle} \|E_f - E_g\| \|f - g\|_{\langle v \rangle} + C\|f - g\|^2 \\ &\quad + \left\{ \sum_{|\beta|+|\gamma| \leq 4} [\|\partial_\beta^\gamma f\| + \|\partial_\beta^\gamma g\|] \right\} \|f - g\|_\nu^2 + \left\{ \sum_{|\beta|+|\gamma| \leq 4} [\|\partial_\beta^\gamma f\|_\nu + \|\partial_\beta^\gamma g\|_\nu] \right\} \|f - g\| \|f - g\|_\nu \\ &\leq C(\|f\|_\nu^2 + \|g\|_\nu^2 + 1) \|f - g\|^2 + (C\sqrt{M_0} + \frac{3}{4}) \|f - g\|_{\langle v \rangle}^2. \end{aligned}$$

For $\eta \ll 1$ and $M_0 \ll 1$, it follows that

$$\frac{1}{2}\|f - g\|^2(t) \leq \frac{1}{2}\|f - g\|^2(0) + C \int_0^t (\|f\|_\nu^2 + \|g\|_\nu^2 + 1) \|f - g\|^2(s) ds,$$

which indicates that $f(t) \equiv g(t)$ by Gronwall's inequality.

The continuity of $\mathcal{E}(t)$ comes from (4.5), and the positivity of $F = \mu + \sqrt{\mu}f$ comes from a simple induction over $F^n, n \in \mathbb{N}$. This completes the proof of Theorem 4.1. \square

5. COERCIVITY OF \mathcal{L}

To extend the local solution of the ionic Vlasov-Poisson-Boltzmann system (1.3)–(1.4) to be a global one, a crucial step is to establish the coercivity of the linear collision operator \mathcal{L} . We have

Theorem 5.1. *Let $f(t, x, v)$ be a classical solution to the system (1.3)–(1.4) satisfying (1.6)–(1.8). There exist $M_0 \ll 1$ and $\delta_0 = \delta_0(M_0) > 0$ such that if*

$$\sum_{|\gamma| \leq N} \|\partial^\gamma f(t)\|^2 \leq M_0,$$

then

$$\sum_{|\gamma| \leq N} (\mathcal{L}\partial^\gamma f(t), \partial^\gamma f(t)) \geq \delta_0 \sum_{|\gamma| \leq N} \|\partial^\gamma f(t)\|_\nu^2. \quad (5.1)$$

Proof. Recall the decomposition $f = \mathbf{P}f + \{\mathbf{I} - \mathbf{P}\}f$ in Section 2 and expand the macroscopic part of f as

$$\mathbf{P}f(t, x, v) := \{a^f(t, x) + b^f(t, x) \cdot v + c^f(t, x)|v|^2\} \sqrt{\mu}.$$

In the sequel, we denote $[a, b, c] := [a^f, b^f, c^f]$ for notational simplification. For the time-space derivatives $\partial^\gamma = \partial_t^{\gamma_0} \partial_{x_1}^{\gamma_1} \partial_{x_2}^{\gamma_2} \partial_{x_3}^{\gamma_3}$, we immediately know that $\partial^\gamma \mathbf{P}f = \mathbf{P}\partial^\gamma f$, and

$$\|\partial^\gamma \mathbf{P}f\|^2 + \|\partial^\gamma \{\mathbf{I} - \mathbf{P}\}f\|^2 = \|\partial^\gamma f\|^2.$$

Moreover, there exists $C > 1$ such that

$$\frac{1}{C} \|\partial^\gamma \mathbf{P}f\|_\nu^2 \leq \|\partial^\gamma a\|^2 + \|\partial^\gamma b\|^2 + \|\partial^\gamma c\|^2 \leq C \|\partial^\gamma \mathbf{P}f\|^2.$$

To prove (5.1) in Theorem 5.1, since there exists a $\delta > 0$ such that

$$(\mathcal{L}\partial^\gamma f, \partial^\gamma f) \geq \delta \|\{\mathbf{I} - \mathbf{P}\}\partial^\gamma f\|_\nu^2,$$

it suffices to show that

$$\sum_{|\gamma| \leq N} \|\mathbf{P}\partial^\gamma f(t)\|_\nu \leq C \sum_{|\gamma| \leq N} \|\{\mathbf{I} - \mathbf{P}\}\partial^\gamma f(t)\|_\nu.$$

We thus only need to verify that

$$\sum_{|\gamma| \leq N} \{\|\partial^\gamma a\| + \|\partial^\gamma b\| + \|\partial^\gamma c\|\} \leq C \sum_{|\gamma| \leq N} \|\{\mathbf{I} - \mathbf{P}\}\partial^\gamma f(t)\|_\nu + C\sqrt{M_0} \sum_{|\gamma| \leq N} \|\partial^\gamma f(t)\|_\nu \quad (5.2)$$

holds for $M_0 \ll 1$ for hard and soft potentials cases $\nu(v)$. Recall that $\nu(v) \sim \langle v \rangle^r, r \in (-3, 1]$.

To observe (5.2), we present several lemmas below.

Lemma 5.2. *Under the same condition as in Theorem 5.1, we have*

$$\begin{aligned} \left| \int_{\mathbb{T}^3} a(t, x) dx \right| + \left| \int_{\mathbb{T}^3} c(t, x) dx \right| &\leq C \{\|\phi\|^2 + \|\nabla_x \phi\|^2\}, \\ \int_{\mathbb{T}^3} b(t, x) dx &\equiv 0. \end{aligned}$$

Proof. Plugging $\mathbf{P}f = \{a + b \cdot v + c|v|^2\}\sqrt{\mu}$ into the conservation laws (1.6)–(1.8), we deduce that

$$\begin{aligned} \varrho_0 \int_{\mathbb{T}^3} a dx + \varrho_2 \int_{\mathbb{T}^3} c dx &= 0, \\ \varrho_2 \int_{\mathbb{T}^3} b dx &= 0, \\ \varrho_2 \int_{\mathbb{T}^3} a dx + \varrho_4 \int_{\mathbb{T}^3} c dx + 2 \int_{\mathbb{T}^3} \phi e^\phi dx + \int_{\mathbb{T}^3} |\nabla \phi|^2 dx &= 0, \end{aligned}$$

where $\varrho_i = \int_{\mathbb{R}^3} |v|^i \mu(v) dv$, $i = 1, \dots, 4$. Note that $\varrho_0 \varrho_4 \neq \varrho_2^2$ and $\varrho_0 \varrho_4, \varrho_2^2 > 0$. First, we have $\int_{\mathbb{T}^3} b dx = 0$ and

$$\left| \int_{\mathbb{T}^3} a dx \right| + \left| \int_{\mathbb{T}^3} c dx \right| \leq C \int_{\mathbb{T}^3} \phi e^\phi dx + \int_{\mathbb{T}^3} |\nabla \phi|^2 dx.$$

By Proposition 3.2, when $M_0 \ll 1$, it holds

$$\|\phi\|_{L^\infty(\mathbb{T}^3)} \leq C \|\phi\|_{H^2(\mathbb{T}^3)} \leq C \|f\|_{L^2(\mathbb{T}^3 \times \mathbb{R}^3)} \leq C \sqrt{M_0} \ll 1.$$

Thus,

$$\begin{aligned} \int_{\mathbb{T}^3} \phi e^\phi dx &= \int_{\mathbb{T}^3} (\phi e^\phi - e^\phi + 1) dx = \int_{\mathbb{T}^3} \{\phi^2 + (\phi - 1)[e^\phi - \phi - 1]\} dx \\ &\leq \int_{\mathbb{T}^3} \phi^2 + C(\|\phi\|_\infty + 1)|\phi|^2 dx \leq C \int_{\mathbb{T}^3} \phi^2 dx, \end{aligned}$$

which implies the desired result. \square

Rewrite the perturbed Vlasov-Boltzmann equation (1.3) as

$$\{\partial_t + v \cdot \nabla_x\} \mathbf{P}f - E \cdot v \sqrt{\mu} = l(\{\mathbf{I} - \mathbf{P}\}f) + h(f), \quad (5.3)$$

where

$$\begin{aligned} l(\{\mathbf{I} - \mathbf{P}\}f) &:= -\{\partial_t + v \cdot \nabla_x + \mathcal{L}\}\{\mathbf{I} - \mathbf{P}\}f, \\ h(f) &:= -E \cdot \nabla_v f + \frac{v}{2} \cdot E f + \Gamma(f, f). \end{aligned}$$

We plug $\mathbf{P}f = \{a + b \cdot v + c|v|^2\}\sqrt{\mu}$ into the left-hand side of (5.3) (denote $\partial^0 = \partial_t$, $\partial^j = \partial_{x_j}$, $\partial^i = \partial_{x_i}$) to get

$$\left[\{\partial^i c\} v_i |v|^2 + \{\partial^0 c + \partial^i b_i\} v_i^2 + \sum_{i < j} \{\partial^i b_j + \partial^j b_i\} v_i v_j + \{\partial^0 b_i + \partial^i a - E_i\} v_i + \partial^0 a \right] \sqrt{\mu}.$$

Under the basis $[1, v_i, v_i v_j, v_i^2, v_i |v|^2] \sqrt{\mu}$ ($1 \leq i < j \leq 3$), we denote

$$l + h := (l_c + h_c) \cdot v |v|^2 \sqrt{\mu} + (l_i + h_i) v_i^2 \sqrt{\mu} + \sum_{i < j} (l_{ij} + h_{ij}) v_i v_j \sqrt{\mu} + (l_{bi} + h_{bi}) v_i \sqrt{\mu} + (l_a + h_a) \sqrt{\mu}.$$

Comparing with the coefficients on both sides of (5.3), we deduce that

$$\nabla_x c = l_c + h_c, \quad (5.4)$$

$$\partial^0 c + \partial^i b_i = l_i + h_i, \quad (5.5)$$

$$\partial^i b_j + \partial^j b_i = l_{ij} + h_{ij}, \quad i \neq j, \quad (5.6)$$

$$\partial^0 b_i + \partial^i a - E_i = l_{bi} + h_{bi}, \quad (5.7)$$

$$\partial^0 a = l_a + h_a. \quad (5.8)$$

Lemma 5.3. *Let $\gamma = [\gamma_0, \gamma_1, \gamma_2, \gamma_3]$. Then for any $1 \leq i \neq j \leq 3$,*

$$\sum_{|\gamma| \leq N-1} \|\partial^\gamma l_c\| + \|\partial^\gamma l_i\| + \|\partial^\gamma l_{ij}\| + \|\partial^\gamma l_{bi}\| + \|\partial^\gamma l_a\| \leq C \sum_{|\gamma| \leq N} \|\{\mathbf{I} - \mathbf{P}\} \partial^\gamma f\|_\nu.$$

Proof. Denote $\{\epsilon_n(v), n = 1, \dots, 13\}$ to be the basis $[1, v_i, v_i^2, v_i v_j, v_i |v|^2] \sqrt{\mu}$. Then for fixed (t, x) , $\partial^\gamma l_c, \partial^\gamma l_i, \partial^\gamma l_{ij}, \partial^\gamma l_{bi}$ and $\partial^\gamma l_a$ take the forms of the linear combinations of

$$\int_{\mathbb{R}^3} \partial^\gamma l(\{\mathbf{I} - \mathbf{P}\}f)(t, x, v) \cdot \epsilon_n(v) dv.$$

By Hölder's inequality and Lemma 2.6, it holds

$$\begin{aligned} & \left\| \int_{\mathbb{R}^3} \partial^\gamma l(\{\mathbf{I} - \mathbf{P}\}f)(t, x, v) \cdot \epsilon_n(v) dv \right\|^2 \\ &= \left\| \int_{\mathbb{R}^3} \{\partial_t + v \cdot \nabla_x + \mathcal{L}\}[\{\mathbf{I} - \mathbf{P}\} \partial^\gamma f](t, x, v) \cdot \epsilon_n(v) dv \right\|^2 \\ &\leq \int_{\mathbb{T}^3 \times \mathbb{R}^3} |\epsilon_n(v)| \left(|\{\mathbf{I} - \mathbf{P}\} \partial_t \partial^\gamma f|^2 + |v|^2 |\{\mathbf{I} - \mathbf{P}\} \nabla_x \partial^\gamma f|^2 + |\mathcal{L}\{\mathbf{I} - \mathbf{P}\} \partial^\gamma f|^2 \right) \\ &\leq C \|\{\mathbf{I} - \mathbf{P}\} \partial^0 \partial^\gamma f\|_\nu^2 + C \|\{\mathbf{I} - \mathbf{P}\} \nabla_x \partial^\gamma f\|_\nu^2 + \int_{\mathbb{T}^3 \times \mathbb{R}^3} |\nu \epsilon_n(v)| \left(|\nu^{-1/2} \mathcal{L}\{\mathbf{I} - \mathbf{P}\} \partial^\gamma f|^2 \right) \\ &\leq C \left\{ \|\{\mathbf{I} - \mathbf{P}\} \partial^0 \partial^\gamma f\|_\nu^2 + \|\{\mathbf{I} - \mathbf{P}\} \nabla_x \partial^\gamma f\|_\nu^2 + \|\{\mathbf{I} - \mathbf{P}\} \partial^\gamma f\|_\nu^2 \right\}. \end{aligned}$$

Hence the lemma is proved. \square

Lemma 5.4. *Let $\gamma = [\gamma_0, \gamma_1, \gamma_2, \gamma_3]$. Suppose that $\sum_{|\gamma| \leq N} \|\partial^\gamma f(t)\|^2 \leq M_0$ holds for some $M_0 > 0$, then for any $1 \leq i \neq j \leq 3$,*

$$\sum_{|\gamma| \leq N} \|\partial^\gamma h_c\| + \|\partial^\gamma h_i\| + \|\partial^\gamma h_{ij}\| + \|\partial^\gamma h_{bi}\| + \|\partial^\gamma h_a\| \leq C \sqrt{M_0} \sum_{|\gamma| \leq N} \|\partial^\gamma f\|_\nu.$$

Proof. Recall the basis $\{\epsilon_n(v), n = 1, \dots, 13\}$. Then for fixed (t, x) , $\partial^\gamma h_c, \partial^\gamma h_i, \partial^\gamma h_{ij}, \partial^\gamma h_{bi}$ and $\partial^\gamma h_a$ take the forms of the linear combinations of

$$\int_{\mathbb{R}^3} \partial^\gamma h(f)(t, x, v) \cdot \epsilon_n(v) dv = \int_{\mathbb{R}^3} \partial^\gamma \left[-E \cdot \nabla_v f + \frac{v}{2} \cdot E f + \Gamma(f, f) \right] (t, x, v) \cdot \epsilon_n(v) dv.$$

Note that

$$\begin{aligned} - \int_{\mathbb{R}^3} \partial^\gamma (E \cdot \nabla_v f) \cdot \epsilon_n(v) dv &= \int_{\mathbb{R}^3} \partial^\gamma (E f) \cdot \nabla_v \epsilon_n(v) dv \\ &\leq \sum_{\gamma_1} C_{\gamma_1} \int_{\mathbb{R}^3} |\partial^{\gamma_1} E| |\partial^{\gamma - \gamma_1} f| |\nabla_v \epsilon_n(v)| dv \\ &\leq C \sum_{\gamma_1} |\partial^{\gamma_1} E| \left\{ \int_{\mathbb{R}^3} |\mu^{1/8} \partial^{\gamma - \gamma_1} f|^2 dv \right\}^{1/2}. \end{aligned}$$

Thus

$$\int_{\mathbb{T}^3} \left| \int_{\mathbb{R}^3} \partial^\gamma \left[-E \cdot \nabla_v f + \frac{v}{2} \cdot E f \right] \cdot \epsilon_n(v) dv \right|^2 dx \leq C \sum_{\gamma_1} \iint_{\mathbb{T}^3 \times \mathbb{R}^3} |\partial^{\gamma_1} E|^2 |\mu^{1/8} \partial^{\gamma - \gamma_1} f|^2 dv dx.$$

If $|\gamma_1| \leq N/2$ ($N \geq 4$), by $H^2(\mathbb{T}^3) \hookrightarrow L^\infty(\mathbb{T}^3)$ and Proposition 3.4, we have

$$\sup_x |\partial^{\gamma_1} E| \leq C \sum_{|\gamma| \leq N} \|\partial^\gamma E\| \leq C \sum_{|\gamma| \leq N} \|\partial^\gamma f\| \leq C \sqrt{M_0},$$

and

$$\left(\iint_{\mathbb{T}^3 \times \mathbb{R}^3} |\mu^{1/8} \partial^{\gamma - \gamma_1} f|^2 dv dx \right)^{1/2} \leq C \sum_{|\gamma| \leq N} \|\partial^\gamma f\|_\nu.$$

If $|\gamma_1| \geq N/2$, then $|\gamma - \gamma_1| \leq N/2$. Thus, by $H^2(\mathbb{T}^3) \hookrightarrow L^\infty(\mathbb{T}^3)$, it holds

$$\sup_x \int_{\mathbb{R}^3} |\partial^{\gamma - \gamma_1} f|^2 dv \leq \int_{\mathbb{R}^3} \sup_x |\partial^{\gamma - \gamma_1} f|^2 dv \leq C \sum_{|\gamma| \leq N} \|\partial^\gamma f\|^2 \leq C (\sqrt{M_0})^2,$$

and

$$\sup_x |\partial^{\gamma_1} E| \leq C \sum_{|\gamma| \leq N} \|\partial^\gamma E\| \leq C \sum_{|\gamma| \leq N} \|\partial^\gamma f\|_\nu.$$

Hence the contribution of $-E \cdot \nabla_v f + \frac{v}{2} \cdot E f$ is $C\sqrt{M_0} \sum_{|\gamma| \leq N} \|\partial^\gamma f\|_\nu$.

For the term $\Gamma(f, f)$, by Lemma 2.6, we deduce that

$$\begin{aligned} \left\| \int_{\mathbb{R}^3} \partial^\gamma \Gamma(f, f) \cdot \epsilon_n(v) dv \right\| &\leq \sum C_\gamma^{\gamma_1} \left\| \int_{\mathbb{R}^3} \Gamma(\partial^{\gamma_1} f, \partial^{\gamma - \gamma_1} f) \cdot \epsilon_n(v) dv \right\| \\ &\leq C\sqrt{M_0} \sum_{|\gamma| \leq N} \|\partial^\gamma f\|_\nu. \end{aligned}$$

Hence, the proof is completed. \square

Based on the above facts, we come back to prove (5.2), i.e.,

$$\sum_{|\gamma| \leq N} \{\|\partial^\gamma a\| + \|\partial^\gamma b\| + \|\partial^\gamma c\|\} \leq C \sum_{|\gamma| \leq N} \|\{\mathbf{I} - \mathbf{P}\} \partial^\gamma f(t)\|_\nu + C\sqrt{M_0} \sum_{|\gamma| \leq N} \|\partial^\gamma f(t)\|_\nu$$

holds for $M_0 \ll 1$. Towards this goal, we divide the proof into several steps as follows.

Step 1. The estimate of $\nabla \partial^\gamma b$, $|\gamma| \leq N - 1$.

By using the identities (5.5) and (5.6), we deduce that (as in [18])

$$\begin{aligned} \Delta \partial^\gamma b_i &= \sum_{j \neq i} [\partial^{jj} \partial^\gamma b_i] + \partial^{ii} \partial^\gamma b_i \\ &= \sum_{j \neq i} [-\partial^{ij} \partial^\gamma b_j + \partial^j \partial^\gamma l_{ij} + \partial^j \partial^\gamma h_{ij}] + [\partial^i \partial^\gamma l_i + \partial^i \partial^\gamma h_i - \partial^0 \partial^i \partial^\gamma c] \\ &= \sum_{j \neq i} [\partial^0 \partial^i \partial^\gamma c - \partial^i \partial^\gamma l_j - \partial^i \partial^\gamma h_j] - \partial^0 \partial^i \partial^\gamma c \\ &\quad + \sum_{j \neq i} [\partial^j \partial^\gamma l_{ij} + \partial^j \partial^\gamma h_{ij}] + \partial^i \partial^\gamma l_i + \partial^i \partial^\gamma h_i \\ &= \partial^0 \partial^i \partial^\gamma c + \sum_{j \neq i} [-\partial^i \partial^\gamma l_j - \partial^i \partial^\gamma h_j + \partial^i \partial^\gamma l_{ij} + \partial^i \partial^\gamma h_{ij}] + \partial^i \partial^\gamma l_i + \partial^i \partial^\gamma h_i \\ &= -\partial^{ii} \partial^\gamma b_i + \sum_{j \neq i} [-\partial^i \partial^\gamma l_j - \partial^i \partial^\gamma h_j + \partial^i \partial^\gamma l_{ij} + \partial^i \partial^\gamma h_{ij}] + 2\{\partial^i \partial^\gamma l_i + \partial^i \partial^\gamma h_i\}. \end{aligned}$$

Multiplying the above equality by $\partial^\gamma b_i$, we get

$$\begin{aligned} \|\nabla \partial^\gamma b_i\| &\leq C \sum_{i,j} \{\|\partial^\gamma l_j\| + \|\partial^\gamma h_j\| + \|\partial^\gamma l_{ij}\| + \|\partial^\gamma h_{ij}\| + \|\partial^\gamma l_i\| + \|\partial^\gamma h_i\|\} \\ &\leq C \sum_{|\gamma| \leq N} \|\{\mathbf{I} - \mathbf{P}\} \partial^\gamma f(t)\|_\nu + C\sqrt{M_0} \sum_{|\gamma| \leq N} \|\partial^\gamma f(t)\|_\nu, \end{aligned}$$

where we have used Lemmas 5.3 and 5.4.

Step 2. The estimate of $\partial^\gamma c$, $|\gamma| \leq N$.

For $|\gamma'| \leq N - 1$, from the identities (5.4) and (5.5), it holds

$$\|\partial^0 \partial^{\gamma'} c\| + \|\nabla \partial^{\gamma'} c\| \leq \|\partial^i \partial^{\gamma'} b_i\| + \|\partial^{\gamma'} l_i\| + \|\partial^{\gamma'} h_i\| + \|\partial^{\gamma'} l_c\| + \|\partial^{\gamma'} h_c\|.$$

In addition, applying Poincaré's inequality, Lemma 5.2 and Proposition 3.2, we get

$$\|c\| \leq C \left\{ \|\nabla c\| + \left| \int_{\mathbb{T}^3} c dx \right| \right\} \leq C \{ \|\nabla c\| + \|\phi\|^2 + \|\nabla_x \phi\|^2 \} \leq C \{ \|\nabla c\| + \|f\|^2 \}.$$

Combining the above estimates on $c(t, x)$ and its temporal and spatial derivatives yields

$$\sum_{|\gamma| \leq N} \|\partial^\gamma c\| \leq C \sum_{|\gamma| \leq N} \|\{\mathbf{I} - \mathbf{P}\} \partial^\gamma f(t)\|_\nu + C\sqrt{M_0} \sum_{|\gamma| \leq N} \|\partial^\gamma f(t)\|_\nu$$

by using Lemmas 5.3 and 5.4.

Step 3. The estimate of $\partial^\gamma a$, $|\gamma| \leq N$.

For $|\gamma'| \leq N-1$, from the identity (5.8), we have

$$\|\partial_t \partial^{\gamma'} a\| \leq \|\partial^{\gamma'} l_a\| + \|\partial^{\gamma'} h_a\| \leq C \sum_{|\gamma| \leq N} \|\{\mathbf{I} - \mathbf{P}\} \partial^\gamma f(t)\|_\nu + C\sqrt{M_0} \sum_{|\gamma| \leq N} \|\partial^\gamma f(t)\|_\nu.$$

We now focus on purely spatial derivatives of $a(t, x)$. Let $\gamma' = [0, \gamma'_1, \gamma'_2, \gamma'_3] \neq 0$ ($0 < |\gamma'| \leq N-1$). Taking $\partial^i \partial^{\gamma'}$ of the identities (5.8) and (5.4), and then summing over $i = 1, 2, 3$, we get

$$-\Delta \partial^{\gamma'} a + \nabla \cdot \partial^{\gamma'} E = \nabla \cdot \partial^0 \partial^{\gamma'} b - \sum_i \partial^i \partial^{\gamma'} (l_{bi} + h_{bi}), \quad (5.9)$$

$$\Delta \partial^{\gamma'} c = \nabla \cdot \partial^{\gamma'} (l_c + h_c), \quad (5.10)$$

and

$$\Delta \partial^{\gamma'} \phi = -\nabla \cdot \partial^{\gamma'} E. \quad (5.11)$$

We take (5.9) $\times [\varrho_0 \partial^{\gamma'} a + \varrho_2 \partial^{\gamma'} c]$ - (5.10) $\times [K \partial^{\gamma'} c]$ - (5.11) $\times [\partial^{\gamma'} (e^\phi - 1)]$, where the constant $K > 0$ is sufficiently large, and then integrate the result by parts over \mathbb{T}^3 to get

$$\begin{aligned} & \int_{\mathbb{T}^3} |\nabla \partial^{\gamma'} a|^2 + K |\nabla \partial^{\gamma'} c|^2 + (\nabla \partial^{\gamma'} a \cdot \nabla \partial^{\gamma'} c) + (\nabla \cdot \partial^{\gamma'} E)^2 - \Delta \partial^{\gamma'} \phi \partial^{\gamma'} (e^\phi) dx \\ & \leq C \left(\|\partial^0 \partial^{\gamma'} b\| + \sum_i \|\partial^{\gamma'} (l_{bi} + h_{bi})\| + \|\partial^{\gamma'} (l_c + h_c)\| \right) \left(\|\nabla \partial^{\gamma'} a\| + \|\nabla \partial^{\gamma'} c\| \right). \end{aligned} \quad (5.12)$$

To deal with the term $\Delta \partial^{\gamma'} \phi \partial^{\gamma'} (e^\phi)$ on the left-hand side of (5.12), we use integration by parts to arrive at

$$\begin{aligned} - \int_{\mathbb{T}^3} \Delta \partial^{\gamma'} \phi \partial^{\gamma'} (e^\phi) dx &= \int_{\mathbb{T}^3} \nabla \partial^{\gamma'} \phi \cdot \partial^{\gamma'} (\nabla \phi e^\phi) dx \\ &= \int_{\mathbb{T}^3} |\nabla \partial^{\gamma'} \phi|^2 e^\phi dx + \int_{\mathbb{T}^3} \nabla \partial^{\gamma'} \phi \cdot \bar{S} e^\phi dx, \end{aligned}$$

where every component of \bar{S} is a finite sum: each element therein is a product of at least two terms of $\partial^{\gamma''} \phi$, $0 < |\gamma''| \leq |\gamma'|$. For example, if $|\gamma'| = 1$, the corresponding $\bar{S} = \nabla \phi \partial^{\gamma'} \phi$; if $|\gamma'| = 2$, the corresponding $\bar{S} = \sum \partial^i \nabla \phi \partial^k \phi + \partial^{ij} \phi \nabla \phi + \partial^i \phi \partial^j \phi \nabla \phi$. Recalling Proposition 3.4, we have

$$\|\partial^{\gamma''} \phi\|_{L^\infty(\mathbb{T}^3)} \leq C \|\partial^{\gamma''} \phi\|_{H^2(\mathbb{T}^3)} \leq C\sqrt{M_0} \ll 1.$$

Hence it holds

$$\begin{aligned} \int_{\mathbb{T}^3} \nabla \partial^{\gamma'} \phi \cdot \bar{S} e^\phi dx &\leq C\sqrt{M_0} \sum_{0 < |\gamma''| \leq |\gamma'|} \int_{\mathbb{T}^3} |\nabla \partial^{\gamma'} \phi| |\partial^{\gamma''} \phi| e^\phi dx \\ &\leq C\sqrt{M_0} \int_{\mathbb{T}^3} |\nabla \partial^{\gamma'} \phi|^2 e^\phi dx + C\sqrt{M_0} \sum_{0 < |\gamma''| \leq |\gamma'|} \int_{\mathbb{T}^3} |\partial^{\gamma''} \phi|^2 e^\phi dx. \end{aligned}$$

Note that $\int_{\mathbb{T}^3} \partial^{\gamma''} \phi dx = 0$. By Poincaré's inequality and $e^\phi \sim 1$, it holds

$$\begin{aligned} \int_{\mathbb{T}^3} |\partial^{\gamma''} \phi|^2 e^\phi dx &\leq C \|\partial^{\gamma''} \phi\|^2 \leq C \|\nabla \partial^{\gamma''} \phi\|^2 \leq \dots \leq C \|\nabla^{(|\gamma'|+1)} \phi\|^2 \\ &\leq C \sum_{|\varpi|=|\gamma'|} \int_{\mathbb{T}^3} |\nabla \partial^\varpi \phi|^2 e^\phi dx, \end{aligned}$$

which implies that

$$\sum_{0 < |\gamma''| \leq |\gamma'|} \int_{\mathbb{T}^3} |\partial^{\gamma''} \phi|^2 e^\phi dx \leq C \sum_{|\varpi|=|\gamma'|} \int_{\mathbb{T}^3} |\nabla \partial^\varpi \phi|^2 e^\phi dx.$$

Hence we obtain that

$$- \int_{\mathbb{T}^3} \Delta \partial^{\gamma'} \phi \partial^{\gamma'} (e^\phi) dx \geq \int_{\mathbb{T}^3} |\nabla \partial^{\gamma'} \phi|^2 e^\phi dx - \left| \int_{\mathbb{T}^3} \nabla \partial^{\gamma'} \phi \cdot \bar{S} e^\phi dx \right|$$

$$\begin{aligned}
&\geq (1 - C\sqrt{M_0}) \int_{\mathbb{T}^3} |\nabla \partial^{\gamma'} \phi|^2 e^\phi dx - C\sqrt{M_0} \sum_{0 < |\gamma''| \leq |\gamma'|} \int_{\mathbb{T}^3} |\partial^{\gamma''} \phi|^2 e^\phi dx \\
&\geq (1 - C\sqrt{M_0}) \int_{\mathbb{T}^3} |\nabla \partial^{\gamma'} \phi|^2 e^\phi dx - C\sqrt{M_0} \sum_{|\varpi| = |\gamma'|} \int_{\mathbb{T}^3} |\nabla \partial^\varpi \phi|^2 e^\phi dx.
\end{aligned}$$

Summing the above inequality over $0 < |\gamma'| \leq N - 1$ yields

$$\begin{aligned}
&\sum_{0 < |\gamma'| \leq N-1} \int_{\mathbb{T}^3} -\Delta \partial^{\gamma'} \phi \partial^{\gamma'} (e^\phi) dx \\
&\geq (1 - C\sqrt{M_0}) \sum_{0 < |\gamma'| \leq N-1} \int_{\mathbb{T}^3} |\nabla \partial^{\gamma'} \phi|^2 e^\phi dx - C\sqrt{M_0} \sum_{0 < |\gamma'| \leq N-1} \sum_{|\varpi| = |\gamma'|} \int_{\mathbb{T}^3} |\nabla \partial^\varpi \phi|^2 e^\phi dx \\
&\geq \frac{1}{2} \sum_{0 < |\gamma'| \leq N-1} \int_{\mathbb{T}^3} |\nabla \partial^{\gamma'} \phi|^2 e^\phi dx.
\end{aligned}$$

Therefore, if we take the sum of (5.12) over $0 < |\gamma'| \leq N - 1$, we deduce that

$$\begin{aligned}
&\sum_{0 < |\gamma'| \leq N-1} \int_{\mathbb{T}^3} |\nabla \partial^{\gamma'} a|^2 + K |\nabla \partial^{\gamma'} c|^2 + (\nabla \partial^{\gamma'} a \cdot \nabla \partial^{\gamma'} c) + (\nabla \cdot \partial^{\gamma'} E)^2 + \frac{1}{2} |\nabla \partial^{\gamma'} \phi|^2 e^\phi dx \\
&\leq C \sum_{0 < |\gamma'| \leq N-1} \left(\|\partial^0 \partial^{\gamma'} b\| + \sum_i \|\partial^{\gamma'} (l_{bi} + h_{bi})\| + \|\partial^{\gamma'} (l_c + h_c)\| \right) \sum_{0 < |\gamma'| \leq N-1} \left(\|\nabla \partial^{\gamma'} a\| + \|\nabla \partial^{\gamma'} c\| \right). \quad (5.13)
\end{aligned}$$

It immediately follows from (5.13) that (with $K \gg 1$)

$$\begin{aligned}
\sum_{0 < |\gamma| \leq N-1} (\|\nabla \partial^\gamma a\| + \|\nabla \partial^\gamma c\|) &\leq C \sum_{0 < |\gamma'| \leq N-1} \left(\|\partial^0 \partial^{\gamma'} b\| + \sum_i \|\partial^{\gamma'} (l_{bi} + h_{bi})\| + \|\partial^{\gamma'} (l_c + h_c)\| \right) \\
&\leq C \sum_{|\gamma| \leq N} \|\{\mathbf{I} - \mathbf{P}\} \partial^\gamma f(t)\|_\nu + C\sqrt{M_0} \sum_{|\gamma| \leq N} \|\partial^\gamma f(t)\|_\nu.
\end{aligned}$$

If $\gamma' = 0$, the deduction becomes simpler. In this case, (5.12) is exactly as

$$\begin{aligned}
&\int_{\mathbb{T}^3} |\nabla a|^2 + K |\nabla c|^2 + (\nabla a \cdot \nabla c) + (\nabla \cdot E)^2 + |\nabla \phi|^2 e^\phi dx \\
&\leq C \left(\|\partial^0 b\| + \sum_i \|l_{bi} + h_{bi}\| + \|l_c + h_c\| \right) (\|\nabla a\| + \|\nabla c\|), \quad (5.14)
\end{aligned}$$

which implies that

$$\|\nabla a\| + \|\nabla c\| \leq C \sum_{|\gamma| \leq N} \|\{\mathbf{I} - \mathbf{P}\} \partial^\gamma f(t)\|_\nu + C\sqrt{M_0} \sum_{|\gamma| \leq N} \|\partial^\gamma f(t)\|_\nu,$$

and hence

$$\sum_{0 < |\gamma| \leq N} \|\partial^\gamma a\| \leq C \sum_{|\gamma| \leq N} \|\{\mathbf{I} - \mathbf{P}\} \partial^\gamma f(t)\|_\nu + C\sqrt{M_0} \sum_{|\gamma| \leq N} \|\partial^\gamma f(t)\|_\nu.$$

Furthermore, by using Poincaré's inequality, Lemma 5.2 and Proposition 3.2, we get

$$\|a\| \leq C \left\{ \|\nabla a\| + \left| \int_{\mathbb{T}^3} a dx \right| \right\} \leq C \{ \|\nabla a\| + \|\phi\|^2 + \|\nabla_x \phi\|^2 \} \leq C \{ \|\nabla a\| + \|f\|^2 \}.$$

In summary, $\sum_{|\gamma| \leq N} \|\partial^\gamma a\|$ is controlled by the right-hand side of (5.2).

Step 4. The estimate of $\partial^\gamma b(t, x)$, $\gamma = [\gamma_0, 0, 0, 0]$, $|\gamma| \leq N$.

By Lemma 5.2, we have $\int_{\mathbb{T}^3} b dx = 0$. It thus follows that $\int_{\mathbb{T}^3} \partial^\gamma b dx = 0$. If $|\gamma| \leq N - 1$, we use Poincaré's inequality to get

$$\|\partial^\gamma b\| \leq C \|\nabla \partial^\gamma b\| \leq C \sum_{|\gamma| \leq N} \|\{\mathbf{I} - \mathbf{P}\} \partial^\gamma f(t)\|_\nu + C\sqrt{M_0} \sum_{|\gamma| \leq N} \|\partial^\gamma f(t)\|_\nu.$$

If $|\gamma| = N$, we take $\partial^{\gamma'}$ derivative ($|\gamma'| = N - 1$) on the identity (5.7) to get

$$\partial^0 \partial^{\gamma'} b_i + \partial^i \partial^{\gamma'} a - \partial^{\gamma'} E_i = \partial^{\gamma'} (l_{bi} + h_{bi}).$$

Thus,

$$\sum_{|\gamma|=N} \|\partial^{\gamma} b\| \leq \sum_{|\gamma|=N} \|\partial^{\gamma} a\| + \sum_{|\gamma'|=N-1} \|\partial^{\gamma'} E\| + \sum_{|\gamma'|=N-1} \sum_i \|\partial^{\gamma'} (l_{bi} + h_{bi})\|.$$

Recalling Proposition 3.4, we know that

$$\sum_{|\gamma'|=N-1} \|\partial^{\gamma'} E\| \leq \sum_{|\gamma|\leq N} \|\partial^{\gamma} \phi\| \leq C \sum_{|\gamma|\leq N} \left\| \partial^{\gamma} \int_{\mathbb{R}^3} f \sqrt{\mu} dv \right\| \leq C \sum_{|\gamma|\leq N} \{\|\partial^{\gamma} a\| + \|\partial^{\gamma} c\|\}.$$

Therefore, we obtain that

$$\begin{aligned} \sum_{|\gamma|=N} \|\partial^{\gamma} b\| &\leq C \sum_{|\gamma|\leq N} \{\|\partial^{\gamma} a\| + \|\partial^{\gamma} c\|\} + \sum_{|\gamma'|=N-1} \sum_i \|\partial^{\gamma'} (l_{bi} + h_{bi})\| \\ &\leq C \sum_{|\gamma|\leq N} \|\{\mathbf{I} - \mathbf{P}\} \partial^{\gamma} f(t)\|_{\nu} + C \sqrt{M_0} \sum_{|\gamma|\leq N} \|\partial^{\gamma} f(t)\|_{\nu}. \end{aligned}$$

In summary, $\sum_{|\gamma|\leq N} \|\partial^{\gamma} b\|$ is controlled by the right-hand side of (5.2).

Combining the results from the above four steps, the crucial estimate (5.2) is verified, which completes the proof of Theorem 5.1. \square

6. GLOBAL EXISTENCE AND EXPONENTIAL DECAY OF CLASSICAL SOLUTIONS TO THE SYSTEM (1.3)–(1.4)

In this section, we prove the global existence and exponential decay of classical solutions to the ionic Vlasov-Poisson-Boltzmann system (1.3)–(1.4). We first establish a refined energy estimate of classical solutions.

Lemma 6.1. *Suppose that $f(t, x, v)$ is the unique solution to the system (1.3)–(1.4) constructed in Theorem 4.1 satisfying the conservation laws (1.6)–(1.8). Let $\sum_{|\gamma|\leq N} \|\partial^{\gamma} f(t)\|^2 \leq M_0 \ll 1$ be valid. Then for any given $0 \leq m \leq N$, $|\beta| \leq m$, there exist constants $C_{|\beta|} > 0$, $\delta_m > 0$ and $C_m > 0$ such that*

$$\sum_{|\beta|\leq m, |\gamma|+|\beta|\leq N} \left\{ \frac{d}{dt} \left\{ C_{|\beta|} \|W \partial_{\beta}^{\gamma} f\|^2 + \|\partial^{\gamma} \nabla \phi\|^2 + \|\partial^{\gamma} \phi e^{\frac{\phi}{2}}\|^2 \right\} (t) + \delta_m \|\partial_{\beta}^{\gamma} f\|_{\langle v \rangle}^2 (t) \right\} \leq C_m \|f\| \|f\|_{\langle v \rangle}^2 (t), \quad (6.1)$$

where $W = e^{-q \frac{x \cdot v}{\langle v \rangle}}$, $q \gg 1$.

Proof. Firstly, we take ∂^{γ} ($|\beta| = 0$) derivatives to get

$$\begin{aligned} &\{\partial_t + v \cdot \nabla_x + E \cdot \nabla_v\} \partial^{\gamma} f - \partial^{\gamma} E \cdot v \sqrt{\mu} + \mathcal{L}\{\partial^{\gamma} f\} \\ &= - \sum_{\gamma_1 \neq 0} C_{\gamma_1}^{\gamma_1} \partial^{\gamma_1} E \cdot \nabla_v \partial^{\gamma - \gamma_1} f + \sum_{\gamma_1} C_{\gamma_1}^{\gamma_1} \left\{ \partial^{\gamma_1} E \cdot \frac{v}{2} \partial^{\gamma - \gamma_1} f + \Gamma(\partial^{\gamma_1} f, \partial^{\gamma - \gamma_1} f) \right\}. \end{aligned} \quad (6.2)$$

When taking inner product of the above identity with $\partial^{\gamma} f$, we investigate the term $-\iint_{\mathbb{T}^3 \times \mathbb{R}^3} \partial^{\gamma} E \cdot v \sqrt{\mu} \partial^{\gamma} f dv dx$ and rewrite it as

$$\begin{aligned} - \iint_{\mathbb{T}^3 \times \mathbb{R}^3} \partial^{\gamma} E \cdot v \sqrt{\mu} \partial^{\gamma} f dv dx &= \int_{\mathbb{T}^3} \partial^{\gamma} \nabla \phi \cdot \partial^{\gamma} \left(\int_{\mathbb{R}^3} v \sqrt{\mu} f dv \right) dx \\ &= - \int_{\mathbb{T}^3} \partial^{\gamma} \phi \partial^{\gamma} \nabla \cdot (\rho u) dx \\ &= \int_{\mathbb{T}^3} \partial^{\gamma} \phi \partial^{\gamma} \partial_t \rho dx \\ &= \int_{\mathbb{T}^3} \partial^{\gamma} \phi \partial^{\gamma} (\partial_t \phi e^{\phi} - \partial_t \Delta \phi) dx \\ &= \frac{1}{2} \frac{d}{dt} \left(\int_{\mathbb{T}^3} |\nabla \partial^{\gamma} \phi|^2 dx \right) + \int_{\mathbb{T}^3} \partial^{\gamma} \phi \partial^{\gamma} (\partial_t \phi e^{\phi}) dx \end{aligned}$$

$$= \frac{1}{2} \frac{d}{dt} \left(\int_{\mathbb{T}^3} |\nabla \partial^\gamma \phi|^2 dx \right) + \frac{1}{2} \frac{d}{dt} \left(\int_{\mathbb{T}^3} |\partial^\gamma \phi|^2 e^\phi dx \right) + \int_{\mathbb{T}^3} \tilde{S} e^\phi dx,$$

where \tilde{S} is a finite sum: each element herein is a product of at least three terms of $\partial^{\tilde{\gamma}} \phi$, $0 < |\tilde{\gamma}| \leq |\gamma|$. Recalling Proposition 3.4, we thus have

$$\int_{\mathbb{T}^3} \tilde{S} e^\phi dx \leq C \|f\| \|f\|_{\langle v \rangle}^2.$$

Taking the similar procedures to those in Section 4, and combining it with Theorem 5.1, we deduce that (6.2) contributes

$$\frac{d}{dt} \sum_{|\gamma| \leq N} \left(\frac{1}{2} \|W \partial^\gamma f\|^2 + \|\partial^\gamma \nabla \phi\|^2 + \|\partial^\gamma \phi e^{\frac{\phi}{2}}\|^2 \right) (t) + \delta_0 \sum_{|\gamma| \leq N} \|\partial^\gamma f\|_{\langle v \rangle}^2(t) \leq C \sum_{|\gamma| \leq N} \|f\| \|f\|_{\langle v \rangle}^2.$$

To verify (6.1) for $|\beta| = 0$, we multiply (6.2) by $W^2 \partial^\gamma f$ to get

$$\begin{aligned} & \{\partial_t + v \cdot \nabla_x + E \cdot \nabla_v\} \left(\frac{1}{2} W^2 |\partial^\gamma f|^2 \right) - W^2 \partial^\gamma E \cdot v \sqrt{\mu} \partial^\gamma f + W^2 \mathcal{L}\{\partial^\gamma f\} \partial^\gamma f + \Xi (Wf)^2 \\ & = - \sum_{\gamma_1 \neq 0} W^2 C_{\gamma_1}^{\gamma_1} \partial^{\gamma_1} E \cdot \nabla_v \partial^{\gamma - \gamma_1} f + \sum_{\gamma_1} W^2 C_{\gamma_1}^{\gamma_1} \left\{ \partial^{\gamma_1} E \cdot \frac{v}{2} \partial^{\gamma - \gamma_1} f + \Gamma(\partial^{\gamma_1} f, \partial^{\gamma - \gamma_1} f) \right\}, \end{aligned} \quad (6.3)$$

where

$$\Xi := -\frac{1}{2} \frac{\{\partial_t + v \cdot \nabla_x + E \cdot \nabla_v\} W^2}{W^2} = q \frac{|v|^2}{\langle v \rangle} + qE \cdot \left(\frac{x}{\langle v \rangle} + \frac{(x \cdot v)v}{\langle v \rangle^3} \right) \quad (6.4)$$

and $\nu + \Xi \sim \langle v \rangle$. Since $W \sim 1$, there exist two constants $\tilde{c}, C > 0$ such that

$$-\tilde{c} \iint_{\mathbb{T}^3 \times \mathbb{R}^3} \partial^\gamma E \cdot v \sqrt{\mu} \partial^\gamma f dv dx \leq - \iint_{\mathbb{T}^3 \times \mathbb{R}^3} W^2 \partial^\gamma E \cdot v \sqrt{\mu} \partial^\gamma f dv dx \leq -C \iint_{\mathbb{T}^3 \times \mathbb{R}^3} \partial^\gamma E \cdot v \sqrt{\mu} \partial^\gamma f dv dx,$$

which implies that

$$- \iint_{\mathbb{T}^3 \times \mathbb{R}^3} W^2 \partial^\gamma E \cdot v \sqrt{\mu} \partial^\gamma f dv dx \sim \frac{d}{dt} \left(C \int_{\mathbb{T}^3} |\nabla \partial^\gamma \phi|^2 dx \right) + \frac{d}{dt} \left(C \int_{\mathbb{T}^3} |\partial^\gamma \phi|^2 e^\phi dx \right) + C \int_{\mathbb{T}^3} \tilde{S} e^\phi dx.$$

Therefore, by summing up $|\gamma| \leq N$, (6.3) contributes

$$\frac{d}{dt} \sum_{|\gamma| \leq N} \left(\frac{1}{2} \|W \partial^\gamma f\|^2 + \|\partial^\gamma \nabla \phi\|^2 + \|\partial^\gamma \phi e^{\frac{\phi}{2}}\|^2 \right) (t) + \delta_0 \sum_{|\gamma| \leq N} \|\partial^\gamma f\|_{\langle v \rangle}^2(t) \leq C_0 \sum_{|\gamma| \leq N} \|f\| \|f\|_{\langle v \rangle}^2.$$

Secondly, assume that (6.1) is valid for $|\beta| \leq m$. We shall verify (6.1) holds for the case $|\beta| = m + 1$. Take ∂_β^γ derivatives of (1.3), it holds

$$\begin{aligned} \{\partial_t + v \cdot \nabla_x + E \cdot \nabla_v + \nu\} \partial_\beta^\gamma f & = \partial^\gamma E \cdot \partial_\beta (v \sqrt{\mu}) - \sum_{\beta_1 \neq 0} C_{\beta_1}^{\beta_1} \{\partial_{\beta_1} v \cdot \nabla_x + \partial_{\beta_1} \nu\} \partial_{\beta - \beta_1}^\gamma f \\ & + \sum_{\beta_1} C_{\beta_1}^{\beta_1} C_{\gamma_1}^{\gamma_1} \partial^{\gamma_1} E \cdot \partial_{\beta_1} \left(\frac{v}{2} \right) \partial_{\beta - \beta_1}^{\gamma - \gamma_1} f - \sum_{\gamma_1 \neq 0} C_{\gamma_1}^{\gamma_1} \partial^{\gamma_1} E \cdot \nabla_v \partial_{\beta - \beta_1}^{\gamma - \gamma_1} f \\ & + \partial_\beta [\mathcal{K} \partial^\gamma f] + \sum_{\gamma_1} C_{\gamma_1}^{\gamma_1} \partial_\beta \Gamma(\partial^{\gamma_1} f, \partial^{\gamma - \gamma_1} f). \end{aligned} \quad (6.5)$$

We consider the inner product of (6.5) with $W^2 \partial_\beta^\gamma f$. By direct calculations, the contribution of the term $\partial^\gamma E \cdot \partial_\beta (v \sqrt{\mu})$ in (6.5) is

$$(\partial^\gamma E \cdot \partial_\beta (v \sqrt{\mu}), \partial_\beta^\gamma f) = (-1)^{|\beta|} (\partial^\gamma E \cdot \partial_\beta^2 (v \sqrt{\mu}), \partial^\gamma f) \leq C \|\partial^\gamma E\| \|\partial^\gamma f\| \leq C \sum_{|\gamma| \leq N} \|\partial^\gamma f\|^2.$$

By (2.1) in Lemma 2.2 and the compact interpolation for $\partial_v^{\beta - \beta_1}$ between ∂_v^γ ($|\gamma| = |\beta|$) and ∂_v^0 , the contribution of the term $\sum_{\beta_1 \neq 0} C_{\beta_1}^{\beta_1} \{\partial_{\beta_1} v \cdot \nabla_x + \partial_{\beta_1} \nu\} \partial_{\beta - \beta_1}^\gamma f$ in (6.5) is

$$\eta \|\partial_\beta^\gamma f\|^2 + C_\eta \sum_{|\beta_1|=1} \|\nabla_x \partial_{\beta - \beta_1}^\gamma f\|^2.$$

By Remark 2.3, the contribution of the term $\partial_\beta[\mathcal{K}\partial^\gamma f]$ in (6.5) is

$$\eta \sum_{|\beta'|=|\beta|} \|\partial_{\beta'}^\gamma f\|^2 + C_\eta \|\partial^\gamma f\|^2.$$

By simple modifications of the arguments in Section 4, the contribution of other terms on the right-hand side of (6.5) is

$$C \|f\| \|f\|_{\langle v \rangle}^2.$$

In summary, we deduce that

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|W\partial_\beta^\gamma f\|^2(t) + \|\partial_\beta^\gamma f\|_{\langle v \rangle}^2(t) &\leq C \|\partial^\gamma E\| \|\partial^\gamma f\| + \eta \|\partial_\beta^\gamma f\|^2 + C_\eta \sum_{|\beta_1|=1} \|\nabla_x \partial_{\beta-\beta_1}^\gamma f\|^2 \\ &+ \eta \sum_{|\beta'|=|\beta|} \|\partial_{\beta'}^\gamma f\|^2 + C_\eta \|\partial^\gamma f\|^2 + C \|f\| \|f\|_{\langle v \rangle}^2 \end{aligned} \quad (6.6)$$

for sufficiently small constant $\eta > 0$.

Summing (6.6) over $|\beta| = m + 1$ and $|\gamma| + |\beta| \leq N$ yields

$$\sum_{\substack{|\beta|=m+1 \\ |\gamma|+|\beta|\leq N}} \left\{ \frac{d}{dt} \|W\partial_\beta^\gamma f\|^2(t) + \|\partial_\beta^\gamma f\|_{\langle v \rangle}^2(t) \right\} \leq C_\eta \sum_{\substack{|\beta|=m \\ |\gamma|+|\beta|\leq N}} \|\partial_\beta^\gamma f\|^2 + C_\eta \sum_{|\gamma|\leq N} \|\partial^\gamma f\|^2 + C \|f\| \|f\|_{\langle v \rangle}^2. \quad (6.7)$$

Multiplying (6.7) by a sufficiently small number and add it to (6.1) for $|\beta| \leq m$, we obtain

$$\sum_{\substack{|\beta|\leq m+1 \\ |\gamma|+|\beta|\leq N}} \left\{ \frac{d}{dt} \left\{ C_{|\beta|} \|W\partial_\beta^\gamma f\|^2 + \|\partial^\gamma \nabla \phi\|^2 + \|\partial^\gamma \phi e^{\frac{\phi}{2}}\|^2 \right\} (t) + \delta_{m+1} \|\partial_\beta^\gamma f\|_{\langle v \rangle}^2(t) \right\} \leq C_{m+1} \|f\| (t) \|f\|_{\langle v \rangle}^2(t), \quad (6.8)$$

which inductively concludes the proof of the lemma. \square

Proof of Theorem 1.1 (continued). With (6.1) in hand, we now prove the global existence and exponential decay of the solution to (1.3)–(1.4) with a bootstrap argument. Fix $M_0 \ll 1$ such that Theorems 4.1 and 5.1 hold. In the estimate (6.1), we choose $m = N$, and then define

$$\begin{aligned} y(t) &:= \sum_{|\gamma|+|\beta|\leq N} \left\{ C_{|\beta|} \|W\partial_\beta^\gamma f\|^2 + \|\partial^\gamma \nabla \phi\|^2 + \|\partial^\gamma \phi e^{\phi/2}\|^2 \right\} (t) \\ &\sim \sum_{|\gamma|+|\beta|\leq N} \left\{ C_{|\beta|} \|\partial_\beta^\gamma f\|^2 + \|\partial^\gamma \nabla \phi\|^2 + \|\partial^\gamma \phi\|^2 \right\} (t) \\ &\sim \sum_{|\gamma|+|\beta|\leq N} \|\partial_\beta^\gamma f\|^2(t) \\ &= \|f\|^2(t), \end{aligned}$$

where the third line is obtained from the elliptic estimates in Proposition 3.4. Then (6.1) is equivalent to

$$y'(t) + \delta_N \|f\|_{\langle v \rangle}^2(t) \leq C_N \|f\| \|f\|_{\langle v \rangle}^2(t) \leq C\sqrt{M_0} \|f\|_{\langle v \rangle}^2(t),$$

which implies that

$$y'(t) + \frac{\delta_N}{2} \|f\|_{\langle v \rangle}^2(t) \leq 0. \quad (6.9)$$

It follows that

$$y(t) + \frac{\delta_N}{2} \int_0^t \|f\|_{\langle v \rangle}^2(s) ds \leq y(0) \leq C \|f_0\|^2. \quad (6.10)$$

Recall the energy functional defined in (1.9), i.e., $\mathcal{E}(t) = \frac{1}{2} \|f\|^2(t) + \int_0^t \|f\|_{\langle v \rangle}^2(s) ds$. Then (6.10) indicates that

$$\mathcal{E}(t) \leq C \|f_0\|^2 \leq C_* \mathcal{E}(0),$$

for some $C_* > 0$. For a constant $M \leq M_0/(2C_*)$, we choose the initial datum $\mathcal{E}(0) \leq M$ and define

$$T^* := \sup_t \{t : \mathcal{E}(t) \leq M_0\} > 0,$$

then we have

$$\sup_{0 \leq t \leq T^*} \mathcal{E}(t) \leq C_* M < 2C_* M = M_0,$$

which contradicts the definition of T^* . Hence the maximal existence time T^* can be extended to the infinity.

Moreover, by the fact that $y(t) \sim \|f\|^2(t)$, (6.9) indicates that

$$y'(t) + Cy(t) \leq 0,$$

for some $C > 0$, and thus

$$y(t) \leq e^{-Ct}y(0).$$

Therefore, it holds

$$\|f\|^2(t) \leq Ce^{-Ct}\|f_0\|^2.$$

Hence, the whole proof of Theorem 1.1 is completed. \square

Acknowledgements: Li is supported in part by NSFC (Grant Nos. 12331007, 12071212) and the “333 Project” of Jiangsu Province. Wang is supported in part by NSFC (Grant No. 12401288) and the Jiangsu Provincial Research Foundation for Basic Research (Grant No. BK20241259).

REFERENCES

- [1] C. Bardos and P. Degond, Global existence for the Vlasov-Poisson equation in 3 space variables with small initial data, *Ann. Inst. H. Poincaré Anal. Non Linéaire*, **2**(1985), 101–118.
- [2] C. Bardos, F. Golse, T. T. Nguyen, and R. Sentis, The Maxwell-Boltzmann approximation for ion kinetic modeling, *Phys. D*, **376/377**(2018), 94–107.
- [3] F. Bouchut, Global weak solution of the Vlasov-Poisson system for small electrons mass, *Comm. Partial Differential Equations*, **16**(1991), 1337–1365.
- [4] Y.-B. Cao, C. Kim and D. Lee, Global strong solutions of the Vlasov-Poisson-Boltzmann system in bounded domains, *Arch. Ration. Mech. Anal.*, **233**(2019), 1027–1130.
- [5] C. Cercignani, R. Illner, M. Pulvirenti, *The Mathematical Theory of Dilute Gases*, Applied Mathematical Sciences, 106. Springer-Verlag, New York, 1994.
- [6] L. Cesbron and M. Iacobelli, Global well-posedness of Vlasov-Poisson-type systems in bounded domains, *Anal. PDE*, **16**(2023), 2465–2494.
- [7] S. Chaturvedi, J. Luk and T. T. Nguyen, The Vlasov-Poisson-Landau system in the weakly collisional regime, *J. Amer. Math. Soc.*, **36**(2023), 1103–1189.
- [8] D.-Q. Deng and R.-J. Duan, Spectral gap formation to kinetic equations with soft potentials in bounded domain, *Comm. Math. Phys.*, **397**(2023), 1441–1489.
- [9] R.-J. Duan and R. Strain, Optimal time decay of the Vlasov-Poisson-Boltzmann system in \mathbb{R}^3 , *Arch. Ration. Mech. Anal.*, **199**(2011), 291–328.
- [10] R.-J. Duan, T. Yang and H.-J. Zhao, The Vlasov-Poisson-Boltzmann system in the whole space: the hard potential case, *J. Differential Equations*, **252**(2012), 6356–6386.
- [11] R.-J. Duan, T. Yang and H.-J. Zhao, The Vlasov-Poisson-Boltzmann system for soft potentials, *Math. Models Methods Appl. Sci.*, **23**(2013), 979–1028.
- [12] P. Flynn, Local well-posedness of the Vlasov-Poisson-Landau system and related models, *Kinet. Relat. Models*, published online.
- [13] D. Gilbarg and N. S. Trudinger, *Elliptic Partial Differential Equations of Second Order. Second edition*, Grundlehren der mathematischen Wissenschaften, Springer-Verlag, Berlin, 1983.
- [14] M. Griffin-Pickering and M. Iacobelli, Global strong solutions in \mathbb{R}^3 for ionic Vlasov-Poisson systems, *Kinet. Relat. Models*, **14**(2021), 571–597.
- [15] M. Griffin-Pickering and M. Iacobelli, Global well-posedness for the Vlasov-Poisson system with massless electrons in the 3-dimensional torus, *Comm. Partial Differential Equations*, **46**(2021), 1892–1939.
- [16] Y. Guo, The Landau equation in a period box, *Comm. Math. Phys.*, **231**(2002), 391–434.

- [17] Y. Guo, The Vlasov-Poisson-Boltzmann system near Maxwellians, *Comm. Pure Appl. Math.*, **55**(2002), 1104–1135.
- [18] Y. Guo, The Vlasov-Maxwell-Boltzmann system near Maxwellians, *Invent. Math.*, **153**(2003), 593–630.
- [19] Y. Guo, Classical solutions to the Boltzmann equation for molecules with an angular cutoff, *Arch. Ration. Mech. Anal.*, **169**(2003), 305–353.
- [20] Y. Guo, Decay and continuity of the Boltzmann equation in bounded domains, *Arch. Ration. Mech. Anal.*, **197**(2010), 713–809.
- [21] Y. Guo, The Vlasov-Poisson-Landau system in a period box, *J. Amer. Math. Soc.*, **25**(2012), 759–812.
- [22] E. Horst, On the asymptotic growth of the solutions of the Vlasov-Poisson system, *Math. Methods Appl. Sci.*, **16**(1993), 75–86.
- [23] H. Hwang, A. Rendall and J. Velázquez, Optimal gradient estimates and asymptotic behavior for the Vlasov-Poisson system with small initial data, *Arch. Ration. Mech. Anal.*, **200**(2011), 313–360.
- [24] H. J. Hwang and J. J. Velázquez, On global existence for the Vlasov-Poisson system in a half space, *J. Differential Equations*, **247**(2009), 1915–1948.
- [25] H. J. Hwang and J. J. Velázquez, Global existence for the Vlasov-Poisson system in bounded domains, *Arch. Ration. Mech. Anal.*, **195**(2010), 763–796.
- [26] F.-C. Li and Y.-C. Wang, Global strong solutions to the Vlasov-Poisson-Boltzmann system with soft potential in a bounded domain, *J. Differential Equations*, **305**(2021), 143–205.
- [27] H.-L. Li, T. Yang and M.-Y. Zhong, Green’s function and pointwise space-time behaviors of the Vlasov-Poisson-Boltzmann system, *Arch. Ration. Mech. Anal.*, **235**(2020), 1011–1057.
- [28] H.-L. Li, T. Yang and M.-Y. Zhong, Spectrum analysis for the Vlasov-Poisson-Boltzmann system, *Arch. Ration. Mech. Anal.*, **241**(2021), 311–355.
- [29] P. L. Lions and B. Perthame, Propagation of moments and regularity for the 3-dimensional Vlasov-Poisson system, *Invent. Math.*, **105**(1991), 415–430.
- [30] K. Pfaffelmoser, Global classical solutions of the Vlasov-Poisson system in three dimensions for general initial data, *J. Differential Equations*, **95**(1992), 281–303.
- [31] A. C. Phillips, *The Physics of Stars (second edition)*, Wiley, 2013.
- [32] J. Schaeffer, Global existence of smooth solutions to the Vlasov-Poisson system in three dimensions, *Comm. Partial Differential Equations*, **16**(1991), 1313–1335.
- [33] R. Sentis, *Mathematical Models and Methods for Plasma Physics. Vol. 1. Fluid Models*, Birkhäuser/Springer, Cham, 2014.
- [34] J. Smulevici, Small data solutions of the Vlasov-Poisson system and the vector field method, *Ann. PDE*, **2**(2016), 1–55.
- [35] B.-L. Tan, J.-T. Su, Y.-T. Li, J. Huang and C.-M. Tan, *Plasma Astrophysics* (in Chinese), Beijing, Science Press, 2024.
- [36] A. W. Trivelpiece and N.A. Krall, *Principles of Plasma Physics*, McGraw-Hill, Englewood Cliffs, 1973.
- [37] Q.-H. Xiao, L.-J. Xiong and H.-J. Zhao, The Vlasov-Poisson-Boltzmann system for non-cutoff hard potentials, *Sci. China Math.*, **57**(2014), 515–540.
- [38] Q.-H. Xiao, L.-J. Xiong and H.-J. Zhao, The Vlasov-Poisson-Boltzmann system for the whole range of cutoff soft potentials, *J. Funct. Anal.*, **272**(2017), 166–226.
- [39] T. Yang, H.-J. Yu and H.-J. Zhao, Cauchy problem for the Vlasov-Poisson-Boltzmann system. *Arch. Ration. Mech. Anal.*, **182**(2006), 415–470.
- [40] T. Yang and H.-J. Zhao, Global existence of classical solutions to the Vlasov-Poisson-Boltzmann system, *Commun. Math. Phys.*, **268**(2006), 569–605.

DEPARTMENT OF MATHEMATICS, NANJING UNIVERSITY, NANJING 210093, P. R. CHINA

Email address: fli@nju.edu.cn

DEPARTMENT OF MATHEMATICS, SOUTHEAST UNIVERSITY, NANJING, 211189, P. R. CHINA

Email address: yichunwang@seu.edu.cn