L_q-MINKOWSKI PROBLEM OF ANISOTROPIC *p*-TORSIONAL RIGIDITY

CHAO LI AND BIN CHEN

ABSTRACT. In this paper, the L_q -Minkowski problem of anisotropic *p*-torsional rigidity is considered. The existence of the solution of the L_q -Minkowski problem of anisotropic *p*-torsional rigidity with 0 < q < 1 and $1 < q \neq \frac{p}{p-1} + n$ is given.

1. INTRODUCTION AND MAIN RESULTS

The Minkowski problem has long been a pivotal issue in convex geometry, holding a central position in areas such as nonlinear partial differential equations, differential geometry, and optimal transportation theory. The classical Minkowski problem primarily addresses the existence, uniqueness, and regularity of solutions, to which scholars like Aleksandrov, Fenchel and Jessen etc. [38, Notes for Section 8.2] have made seminal contributions. Since Lutwak [35] introduced and studied the L_q -Minkowski problem, it has become evident that the varying value of the parameter q not only encompasses the classical Minkowski problem but also introduces significant challenges in terms of uniqueness, existence, and regularity when q < 1. This has attracted considerable attention and led to a series of distinguished research achievements.

In the process of exploring the Minkowski problem, various variant forms have also attracted widespread academic interest and demonstrated significant research value. These variants include the Orlicz-Minkowski problem proposed by Haberl, Lutwak, Yang and Zhang [14], the dual Minkowski problem studied by Huang, Lutwak, Yang and Zhang [19], the chord Minkowski problem researched by Lutwak, Xi, Yang and Zhang [36], as well as Minkowski problems related to electrostatic capacity [24, 42], gaussian measure [21, 34, 10, 4], hyperbolic space [32], gauss image measure [3], weighted [26, 30, 27], harmonic measure [1, 22, 23, 31], coconvex sets of finite volume [39, 28], among others.

In this paper, we continue to explore a novel type of Minkowski problem, specifically the L_q -Minkowski problem of anisotropic *p*-torsional rigidity. Let F^p be strictly convex, K be bounded open set of \mathbb{R}^n , $1 and <math>n \geq 2$, let u be the solution of the boundary-value problem

$$\begin{cases} \Delta_p^F u = -1 & \text{in } K, \\ u = 0 & \text{on } \partial K, \end{cases}$$

where Δ_p^F is the anisotropic *p*-Laplacian operator which is given by

$$\Delta_p^F u = \operatorname{div} \left(F^{p-1}(\nabla u) \nabla_{\xi} F(\nabla u) \right), \qquad (1.1)$$

then the anisotropic *p*-torsional rigidity $\tau_{F,p}(K)$ of K (the number $\tau_{F,p}(K) > 0$) is defined by (see [9])

$$\tau_{F,p}(K) = \int_{K} F^{p}(\nabla u) \, dx = \int_{K} u dx, \qquad (1.2)$$

²⁰²⁰ Mathematics Subject Classification. 35N25, 52A20, 53C21, 31A15.

Key words and phrases. anisotropic p-Laplacian operator, anisotropic p-torsional rigidity, L_q -Minkowski problem.

from this and Anisotropic Pohožaev identity [2], Li [29] derived the following integral expression

$$\tau_{F,p}(K) = \frac{p-1}{n(p-1)+p} \int_{\mathbb{S}^{n-1}} h_K(\xi) F^p(\nabla u(\mathbf{g}_K^{-1}(\xi))) dS(K,\xi),$$

and define the anisotropic *p*-torsional measure, $S_{F,p}(K, \cdot)$, of K is a Borel measure on the unit sphere \mathbb{S}^{n-1} defined for a Borel set $\eta \subset \mathbb{S}^{n-1}$ by

$$S_{F,p}(K,\eta) = \int_{x \in \mathbf{g}_K^{-1}(\eta)} F^p\left(\nabla u(x)\right) d\mathcal{H}^{n-1}(x)$$

where $u \in W_0^{1,p}(K) \setminus \{0\}$ and $\mathbf{g}_K : \partial' K \to \mathbb{S}^{n-1}$ is the Gauss map of K, defined on $\partial' K$, the set of points of ∂K that have a unique outer unit normal. In recent years, significant advancements have been made in the field of anisotropic *p*-torsional rigidity research. For detailed information, please refer to Reference [2, 9, 12, 13].

Inspired by Lutwak's [35] formulation of the L_q -Minkowski problem, we naturally turn our attention to the investigation of the L_q -Minkowski problem for anisotropic *p*-torsional rigidity. In this paper, we provide a proof for the following variational formulate (see Section 2.2)

$$\lim_{t \to 0} \frac{\tau_{F,p}\left([h_t]\right) - \tau_{F,p}(K)}{t} = \frac{1}{q} \int_{\mathbb{S}^{n-1}} f^q(v) dS_{F,p,q}(K,v)$$

where $h_t(v) = (h_K(v)^q + tf(v)^q)^{\frac{1}{q}}$ and $q \neq 0$. This naturally leads to the definition of L_q anisotropic *p*-torsional measure: $1 and <math>q \neq 0$, then the L_q anisotropic *p*-torsional measure, $S_{F,p,q}(K, \cdot)$, of K is a Borel measure on the unit sphere \mathbb{S}^{n-1} defined for a Borel set $\eta \subset \mathbb{S}^{n-1}$ by

$$S_{F,p,q}(K,\eta) = \frac{p-1}{n(p-1)+p} \int_{x \in \mathbf{g}_K^{-1}(\eta)} \langle x, \mathbf{g}_K(x) \rangle^{1-q} F^p(\nabla u(x)) \, d\mathcal{H}^{n-1}(x).$$

The L_q -Minkowski problem of anisotropic *p*-torsional rigidity is expressed as: The L_q -Minkowski problem of anisotropic *p*-torsional rigidity: Let 1 , $<math>q \neq 0$ and μ be a Borel measure on \mathbb{S}^{n-1} . What properties must μ have that we may guarantee that it is the L_q anisotropic *p*-torsional measure $S_{F,p,q}(K, \cdot)$ of some convex body of \mathbb{R}^n , i.e.,

$$S_{F,p,q}(K,\cdot) = \mu? \tag{1.3}$$

In the context of the anisotropic *p*-Laplacian operator (1.1), this framework not only subsumes the anisotropic Laplacian but also extends to encompass both the *p*-Laplacian and the classical Laplacian operators. Within the scope of L_q -Minkowski problem of anisotropic *p*-torsional rigidity (1.3), when q = 1, the problem reduces to the Minkowski problem of anisotropic *p*-torsional rigidity, which has been studied by Li [29]. When q = 0, the problem transforms into the log-Minkowski problem of anisotropic *p*-torsional rigidity [29]. Moreover, when $F(\xi) = \sum_k |\xi_k|$, is exactly L_q -Minkowski problem of *p*-torsional rigidity (see [5, 40]). Meanehile, when $F(\xi) = \sum_k |\xi_k|$ and p = 2, we encounter the L_q -Minkowski problem of torsional rigidity (see [6, 17]), if q = 1, p = 2 and $F(\xi) = \sum_k |\xi_k|$, the problem simplifies to the most classical Minkowski problem of torsional rigidity [7, 8]. For further exploration of the research on the Minkowski problem of torsional rigidity, refer to references [15, 16, 18, 20, 33, 40, 41]

Firstly, the existence of the L_q -Minkowski problem of anisotropic *p*-torsional rigidity with q > 1 is established by the variational methods.

Theorem 1.1. Let $1 , <math>1 < q \neq \frac{p}{p-1} + n$ and μ be a nonzero finite Borel measure on \mathbb{S}^{n-1} that is not concentrated in any closed hemisphere, then there exists a convex body $K \in \mathscr{K}_o^n$ such that

$$S_{F,p,q}(K,\cdot) = \mu.$$

Remark 1.2. If we set $F(\xi) = \sum_{k} |\xi_{k}|$ and p = 2 in Theorem 1.1, the resulting conclusion aligns with Chen and Dai's [6] findings.

Then we establish the existence of the L_q -Minkowski problem of anisotropic *p*-torsional rigidity with 0 < q < 1 through the application of polyhedral approximation methods.

Theorem 1.3. Let 1 , <math>0 < q < 1 and μ be a finite Borel measure on \mathbb{S}^{n-1} which is not concentrated in any closed hemisphere. Then there exists a convex body K in \mathbb{R}^n , such that

$$d\mu = \lambda dS_{F,p,q}(K, \cdot),$$

where λ is a positive constant.

Remark 1.4. If we take $F(\xi) = \sum_k |\xi_k|$ in Theorem 1.3, then it is exactly Chen, Zhao, Wang and Zhao's [5, Theorem 1.6] result, and if we add q = 2 on this basis again, we get the result of Hu and Liu's [17, Theorem 1].

2. NOTATIONS AND BACKGROUND MATERIALS

2.1. Convex Geometry. Let \mathscr{K}^n denote the set of convex bodies (compact, convex subsets with non-empty interiors) in Euclidean space \mathbb{R}^n . Let \mathscr{K}_o^n denote the set of convex bodies containing the origin in their interiors in \mathbb{R}^n . For the set of star bodies (about the origin) in \mathbb{R}^n , we write \mathscr{S}_o^n . Besides, let \mathbb{S}^{n-1} denote the unit sphere, \mathbb{B}^n denote the unit ball and V(K) denote the *n*-dimensional volume of K.

A convex body $K \in \mathscr{K}^n$, is uniquely determined by its support function, $h(K, \cdot) = h_K : \mathbb{R}^n \to \mathbb{R}$, which is defined by

$$h(K, x) = \max\{\langle x, y \rangle : y \in K\}, \quad x \in \mathbb{R}^n,$$
(2.1)

where $\langle x, y \rangle$ denotes the standard inner product of x and y in \mathbb{R}^n . It is also clear from the definition that $h(K, u) \leq h(L, u)$ for $u \in \mathbb{S}^{n-1}$, if and only if $K \subseteq L$.

For $q \neq 0$ and $\alpha, \beta \geq 0$ (not both zero) and $K, L \in \mathscr{K}_o^n$, the L_q -Minkowski combination [37], $\alpha \cdot K +_q \beta \cdot L$, of K and L is defined by

$$\alpha \cdot K +_q \beta \cdot L = \bigcap_{y \neq 0} \{ x \in \mathbb{R}^n : x \cdot y \le (\alpha h_K(u)^q + \beta h_L(u)^q)^{\frac{1}{q}} \},$$
(2.2)

for $q \ge 1$ is the common L_q -Minkowski combination in (2.2), is defined by (see [38])

$$h(\alpha \cdot K +_q \beta \cdot L, \cdot)^q = \alpha h(K, \cdot)^q + \beta h(L, \cdot)^q, \qquad (2.3)$$

where " $+_q$ " denotes the L_q -sum and $\alpha \cdot K = \alpha^{1/q} K$ is the L_q -Minkowski scalar multiplication. If q = 1, (2.3) is just the classical Minkowski linear combination.

If K is a compact star-shaped set (about the origin) in \mathbb{R}^n , then its radial function, $\rho_K = \rho(K, \cdot) : \mathbb{R}^n \setminus \{0\} \to [0, \infty)$, is defined by

$$\rho(K, x) = \max\{c \ge 0 : cx \in K\}, \quad x \in \mathbb{R}^n \setminus \{0\}.$$

$$(2.4)$$

If ρ_K is positive and continuous, K will be called a star body (with respect to the origin).

The radial map $r_K : \mathbb{S}^{n-1} \to \partial K$ is

$$r_K(\xi) = \rho_K(\xi)\xi,$$

for $\xi \in \mathbb{S}^{n-1}$, i.e. $r_K(\xi)$ is the unique point on ∂K located on the ray parallel to ξ and emanating from the origin.

Using this we see that the outer unit normal vector to ∂K at x, denoted by $\mathbf{g}(x)$, is well defined for \mathscr{H}^{n-1} almost all $x \in \partial K$. The map $\mathbf{g} : \partial K \to \mathbb{S}^{n-1}$ is called the Gauss map of K. For $\omega \subset \mathbb{S}^{n-1}$, let

$$\mathbf{g}^{-1}(\omega) = \{x \in \partial K : \mathbf{g}(x) \text{ is defined and } \mathbf{g}(x) \in \omega\}.$$

If ω is a Borel subset of \mathbb{S}^{n-1} , then $\mathbf{g}^{-1}(\omega)$ is \mathscr{H}^{n-1} -measurable. The Borel measure S_K , on \mathbb{S}^{n-1} , is defined for Borel $\omega \subset \mathbb{S}^{n-1}$ by

$$S_K(\omega) = \mathscr{H}^{n-1}\left(\mathbf{g}^{-1}(\omega)\right)$$

and is called the surface area measure of K. For every $f \in C(\mathbb{S}^{n-1})$,

$$\int_{\mathbb{S}^{n-1}} f(\xi) dS_K(\xi) = \int_{\partial K} f(\mathbf{g}(x)) d\mathcal{H}^{n-1}(x).$$
(2.5)

Let $C^+(\mathbb{S}^{n-1})$ denote a positive continuous function on \mathbb{S}^{n-1} , and E be a closed subset of \mathbb{S}^{n-1} that is not contained in a closed hemisphere, if $f \in C^+(\mathbb{S}^{n-1})$, the Wulff shape [f] associated with f is defined as

$$[f] = \{ x \in \mathbb{R}^n : x \cdot u \le f(u) \text{ for all } u \in E \}.$$

The Aleksandrov body can also be defined as the unique maximal (with respect to set inclusion) element of

$$\{K \in \mathscr{K}_0^n : h_K(u) \le f(u), u \in E\}.$$
(2.6)

In fact,

$$[f] = \bigcap_{u \in E} H^-(u, f(u)),$$

where

$$H^{-}(u,t) = \{x \in \mathbb{R}^{n} : x \cdot u \le t\}.$$

Lemma 2.1 (Porposition 2.5 of [26]). Let $K \in \mathscr{K}_0^n$, the Jacobian of $r_K : \mathbb{S}^{n-1} \to \partial K$ is $\frac{\rho_K^n(v)}{h_K(\mathbf{g}_K(r_K(v)))}$ up to a set of measure zero.

Lemma 2.2 (Lemma 2.1 of [34]). For $q \neq 0$, let $K \in \mathscr{K}_o^n$, and $f : \mathbb{S}^{n-1} \to \mathbb{R}$ be a continuous function. For small enough $\delta > 0$, and each $t \in (-\delta, \delta)$, we define the continuous function $h_t : \mathbb{S}^{n-1} \to (0, \infty)$ as

$$h_t(v) = (h_K(v)^q + tf(v)^q)^{\frac{1}{q}}, \quad v \in \mathbb{S}^{n-1}.$$

Then

$$\lim_{t \to 0} \frac{\rho_{[h_t]}(v) - \rho_K(v)}{t} = \frac{f(\mathbf{g}_K(r_K(v)))^q}{qh_K(\mathbf{g}_K(r_K(v)))^q} \rho_K(v),$$

holds for almost all $u \in \mathbb{S}^{n-1}$. In addition, there exists M > 0 such that

$$\left|\rho_{[h_t]}(u) - \rho_K(u)\right| \le M|t|,$$

for all $u \in \mathbb{S}^{n-1}$ and $t \in (-\delta, \delta)$.

2.2. Anisotropic *p*-torsional rigidity. Let F^p be strictly convex, K be bounded open set of \mathbb{R}^n , $1 and <math>n \ge 2$,

$$\begin{cases} \Delta_p^F u = -1 & \text{in } K, \\ u = 0 & \text{on } \partial K, \end{cases}$$
(2.7)

where Δ_p^F is the Finsler *p*-Laplacian (or anisotropic *p*-Laplacian) operator which is given by

$$\Delta_p^F u = \operatorname{div} \left(F^{p-1}(\nabla u) \nabla_{\xi} F(\nabla u) \right).$$
(2.8)

Let K be a convex body with boundary of class C^2 such that the Gauss curvature is positive at every point of their boundary; let u be the solution of problem (2.7) in K, then the anisotropic p-torsional rigidity $\tau_{F,p}(K)$ of K is defined by

$$\tau_{F,p}(K) = \frac{p-1}{n(p-1)+p} \int_{\mathbb{S}^{n-1}} h_K(\xi) F^p(\nabla u(\mathbf{g}_K^{-1}(\xi))) dS(K,\xi),$$
(2.9)

where \mathscr{H}^{n-1} is (n-1)-dimensional Hausdorff measure.

Clearly, let p = 2 and $F(\xi) = \sum_{k} |\xi_k|$ in (2.7) and (2.9), then $\tau_{F,p}(K)$ becomes the classical torsional rigidity $\tau(K)$ of K (see [7]).

Furthermore, Li [29] demonstrated that the satisfaction of anisotropic *p*-torsional rigidity is analogous to that of classical torsional rigidity. Let *u* be a solution of (2.7), $1 and <math>K, L \in \mathscr{K}^n$. Then the anisotropic *p*-torsional rigidity $\tau_{F,p}(K)$ is positively homogeneous of order $\frac{p}{p-1} + n$, that is,

$$\tau_{F,p}(\lambda K) = \lambda^{\frac{p}{p-1}+n} \tau_{F,p}(K).$$
(2.10)

The anisotropic p-torsional rigidity $\tau_{F,p}(K)$ is translation invariant, namely,

$$\tau_{F,p}(K+x) = \tau_{F,p}(K), \quad x \in \mathbb{R}^n.$$
(2.11)

The anisotropic p-torsional measure $S_{F,p}(K, \cdot)$ is translation invariant.

If $K, L \in \mathscr{K}^n$ and $K \subseteq L$, then

$$\tau_{F,p}(K) \le \tau_{F,p}(L).$$

Let F be a norm in the class

$$\mathscr{I}_p = \left\{ F \in C^{2,\alpha}\left(\mathbb{R}^n \setminus \{0\}\right), \quad \frac{1}{p} F^p \in C^2_+\left(\mathbb{R}^n \setminus \{0\}\right) \right\},$$

with $1 and for some <math>\alpha \in (0, 1)$, that is a regular norm F where F^p is a twice continuously differentiable function in $\mathbb{R}^n \setminus \{0\}$ whose Hessian matrix is strictly positive definite.

Suppose $K, K_i \subset \mathbb{R}^n$ are two a bounded domain with the boundary of class $C^{2,\alpha}$ and F is a norm in \mathscr{I}_p . If K_i converges to K in Hausdorff metric as $i \to \infty$. Then

$$S_{F,p}(K_i, \cdot) \to S_{F,p}(K, \cdot), \quad \text{weakly on} \quad \mathbb{S}^{n-1}.$$
 (2.12)

and $\tau_{F,p}$ is continuous, i.e.,

$$\tau_{F,p}(K_i) \to \tau_{F,p}(K). \tag{2.13}$$

In addition, Pietra and Gavitone [9, Remark 2.1] used the anisotropic Pólya-Szegö inequality characterized the upper bound estimate of the anisotropic p-torsional rigidity.

Lemma 2.3 ([9]). Let $1 and K be a bounded open set of <math>\mathbb{R}^n$. Then

$$\tau_{F,p}(K) \le \frac{p-1}{n(p-1)+p} n^{-\frac{1}{p-1}} \kappa_n^{-\frac{p}{n(p-1)}} V(K)^{\frac{n(p-1)+p}{n(p-1)}},$$
(2.14)

where $\kappa_n = |\mathcal{W}|$ and $|\mathcal{W}|$ denotes the Lebesgue measure of \mathcal{W} . The set

$$\mathcal{W} = \left\{ \xi \in \mathbb{R}^n : H^o(\xi) < 1 \right\},\,$$

is the so-called Wulff shape centered at the origin.

Now, we give the variational formula of the anisotropic p-torsional rigidity with respect to the L_q -Minkowski combination.

Lemma 2.4. Let $1 , <math>q \neq 0$ and K be a convex body containing the origin in its interior, such that ∂K up to set of (n-1)-dimensional Hausdorff measure zero, and $f: \mathbb{S}^{n-1} \to \mathbb{R}$ be continuous function. For sufficiently small $\delta > 0$ and each $t \in (-\delta, \delta)$, the continuous function $h_t: \mathbb{S}^{n-1} \to (0, \infty)$ is defined by

$$h_t(v) = (h_K(v)^q + tf(v)^q)^{\frac{1}{q}}, \quad v \in \mathbb{S}^{n-1},$$

then

$$\lim_{t \to 0} \frac{\tau_{F,p}\left([h_t]\right) - \tau_{F,p}(K)}{t} = \frac{1}{q} \int_{\mathbb{S}^{n-1}} f^q(v) dS_{F,p,q}(K,v).$$

Proof. Let $h_t(v) = (h_K(v)^q + tf(v)^q)^{\frac{1}{q}}$. By invoking equation (1.2) and applying the polar coordinate transformation formula, we derive

$$\tau_{F,p}\left([h_t]\right) = \int_K F^p(\nabla u(x)) dx = \int_{\mathbb{S}^{n-1}} \int_0^{\rho_{[h_t]}(v)} F^p(\nabla u(\xi v)) \xi^{n-1} d\xi du = \int_{\mathbb{S}^{n-1}} \phi_t(v) dv,$$

where $\phi_t(v) = \int_0^{\rho_{[h_t]}(v)} F^p(\nabla u(\xi v))\xi^{n-1}d\xi$. According to the Lebesgue differentiation theorem, it follows that $\frac{d\phi_t}{dt}\Big|_{t=0} = \phi'_0$ exists. Let $\mathcal{I} = [\rho_K(v), \rho_{[h_t]}(v)]$, for all $v \in \mathbb{S}^{n-1}$, note $F^p(\nabla u(\xi v))$ exists from the assumption on F, by Lemma 2.2, we obtain

$$\lim_{t \to 0} \frac{\phi_t(v) - \phi_0(v)}{t} = \lim_{t \to 0} \frac{1}{t} \int_{\mathcal{I}} F^p(\nabla u(\xi v)) \xi^{n-1} d\xi$$
$$= \lim_{t \to 0} \frac{\rho_{[h_t]}(v) - \rho_K(v)}{t} \frac{1}{|\mathcal{I}|} \int_{\mathcal{I}} F^p(\nabla u(\xi v)) \xi^{n-1} d\xi$$
$$= F^p(\nabla u(r_K(v))) \frac{f(\mathbf{g}_K(r_K(v)))^q}{qh_K(\mathbf{g}_K(r_K(v)))^q} \rho_K^n(v).$$

Since for some M > 0,

$$\frac{f(\mathbf{g}_{K}(r_{K}(v)))^{q}}{qh_{K}(\mathbf{g}_{K}(r_{K}(v)))^{q}}\rho_{K}(v) = \lim_{t \to 0} \frac{\rho_{[h_{t}]}(v) - \rho_{K}(v)}{t} \le M,$$

from Lemma 2.2, we can conclude that the derivative is dominated by the integrable function

$$M\left(\max_{u\in\mathbb{S}^{n-1}}\rho_K(v)\right)^{n-1}F^p(\nabla u(r_K(v))).$$

By applying the dominated convergence theorem to differentiate under the integral sign and combining Lemma 2.1 with (2.5), we obtain

$$\int_{\mathbb{S}^{n-1}} \phi_0'(v) du = \int_{\mathbb{S}^{n-1}} F^p(\nabla u(r_K(v))) \frac{f(\mathbf{g}_K(r_K(v)))^q}{qh_K(\mathbf{g}_K(r_K(v)))^q} \rho_K^n(v) dv$$
$$= \frac{1}{q} \int_{\partial K} F^p(\nabla u(x)) f^q(\mathbf{g}_K(x)) h_K^{1-q}(\mathbf{g}_K(r_K(v))) dx$$
$$= \frac{1}{q} \int_{\mathbb{S}^{n-1}} f^q(v) dS_{F,p,q}(K,v).$$

This concludes the proof of Lemma 2.4.

From Lemma 2.4, if $1 , <math>q \neq 0$ and K is a convex body in \mathbb{R}^n that contains the origin in its interior, then the L_q anisotropic p-torsional measure, $S_{F,p,q}(K, \cdot)$, of K is a Borel measure on the unit sphere \mathbb{S}^{n-1} defined for a Borel set $\eta \subset \mathbb{S}^{n-1}$ by

$$S_{F,p,q}(K,\eta) = \int_{x \in \mathbf{g}_K^{-1}(\eta)} \langle x, \mathbf{g}_K(x) \rangle^{1-q} F^p(\nabla u(x)) \, d\mathcal{H}^{n-1}(x).$$
(2.15)

Obviously, the L_q anisotropic *p*-torsional measure $S_{F,p,q}(K, \cdot)$ is absolutely continuous with respect to anisotropic *p*-torsional measure $S_{F,p}(K, \cdot)$, and has Radon-Nikodym derivative

$$\frac{dS_{F,p,q}(K,\cdot)}{dS_{F,p}(K,\cdot)} = h^{1-q}(K,\cdot).$$
(2.16)

Based on the continuity of support function and the continuity of anisotropic *p*-torsional measure, by (2.16), it can be deduced that the L_q anisotropic *p*-torsional measure exhibits continuity as well. Therefore, we conclude that

$$S_{F,p,q}(K_i, \cdot) \to S_{F,p,q}(K, \cdot), \quad \text{weakly on} \quad \mathbb{S}^{n-1}.$$
 (2.17)

If q = 1 in Lemma 2.4, we can obtain the variational formula of the anisotropic *p*-torsional rigidity with respect to the Minkowski combination.

Corollary 2.5. Let $1 and K be a convex body containing the origin in its interior, such that <math>\partial K$ up to set of (n-1)-dimensional Hausdorff measure zero, and $f: \mathbb{S}^{n-1} \to \mathbb{R}$ be continuous function. For sufficiently small $\delta > 0$ and each $t \in (-\delta, \delta)$, the continuous function $h_t: \mathbb{S}^{n-1} \to (0, \infty)$ is defined by

$$h_t(v) = h_K(v) + tf(v), \quad v \in \mathbb{S}^{n-1},$$

then

$$\lim_{t \to 0} \frac{\tau_{F,p}\left([h_t]\right) - \tau_{F,p}(K)}{t} = \int_{\mathbb{S}^{n-1}} f(v) dS_{F,p}(K,v)$$

3. A variational proof of L_q -Minkowski problem of anisotropic *p*-torsional measure

Let 1 and <math>q > 0. For each non-zero finite Borel measure μ on \mathbb{S}^{n-1} , define the functional $\Psi_{F,p,q}: C^+(\mathbb{S}^{n-1}) \to \mathbb{R}^n$ by

$$\Psi_{F,p,q}(f) = \frac{1}{q} \log \int_{\mathbb{S}^{n-1}} f(v)^q d\mu(v) - \frac{p-1}{n(p-1)+p} \log \tau_{F,p}([f]).$$

It is easy to check that $\Psi_{F,p,q}(f)$ is homogeneous of degree 0, i.e.,

$$\Psi_{F,p,q}(tf) = \Psi_{F,p,q}(f), \quad \forall t > 0, f \in C^+ \left(\mathbb{S}^{n-1} \right)$$

Note that for each $f \in C^+(\mathbb{S}^{n-1})$, from the definition of wulff shapes, we have $h_{[f]} \leq f$ and $[f] = [h_{[f]}]$, then

$$\Psi_{F,p,q}(f) \ge \Psi_{F,p,q}(h_{[f]}).$$

Lemma 3.1. Let $1 , <math>1 < q \neq \frac{p}{p-1} + n$ and μ be a nonzero finite Borel measure on \mathbb{S}^{n-1} that is not concentrated in any closed hemisphere. If the minimization problem

$$\inf \left\{ \Psi_{F,p,q}(h) : \quad \tau_{F,p}([h]) = |\mu|, \quad h \in C^+(\mathbb{S}^{n-1}) \right\},$$
(3.1)

has a solution $h_0 \in C^+(\mathbb{S}^{n-1})$, then there exists $K \in \mathscr{K}_o^n$ such that

$$S_{F,p,q}\left(K,\cdot\right) = \mu. \tag{3.2}$$

Proof. Now, we may restrict our attention in the search of a minimizer to the set of all support functions. That is, h_{K_0} is a minimizer to the optimization problem (3.1) if and only if K_0 is a minimizer to the following optimization problem:

$$\inf \left\{ \Psi_{F,p,q}(K_0) : \quad \tau_{F,p}(K_0) = |\mu|, \quad K_0 \in \mathscr{K}_o^n \right\},$$
(3.3)

where $\Psi_{F,p,q}: \mathscr{K}_o^n \to \mathbb{R}$ is defined by

$$\Psi_{F,p,q}(K_0) = \frac{1}{q} \log \int_{\mathbb{S}^{n-1}} h_{K_0}(v)^q d\mu(v) - \frac{p-1}{n(p-1)+p} \log \tau_{F,p}(K_0),$$

for each $K_0 \in \mathscr{K}_o^n$. Clearly, the functional $\Psi_{F,p,q}$ is continuous with respect to the Hausdorff metric.

Suppose K_0 is a minimizer to (3.3), or equivalently h_{K_0} is a minimizer to (3.1), i.e., $\tau_{F,p}(K_0) = |\mu|$ and

$$\Psi_{F,p,q}(h_{K_0}) = \inf \left\{ \Psi_{F,p,q}(h) : \tau_{F,p}([h]) = |\mu|, h \in C^+(\mathbb{S}^{n-1}) \right\},\$$

this yields that

$$\Psi_{F,p,q}(h_{K_0}) \le \Psi_{F,p,q}(h),$$

for each $h \in C^+(\mathbb{S}^{n-1})$. Let $f \in C^+(\mathbb{S}^{n-1})$ be an arbitrary continuous function. For $\delta > 0$ small enough and $t \in (-\delta, \delta)$, define $h_t : \mathbb{S}^{n-1} \to \mathbb{R}$ by $h_t(v) = (h_{K_0}(v)^q + tf(v)^q)^{\frac{1}{q}}$. Using the fact that h_{K_0} is a minimizer of (3.3) and Lemma 2.4, we have

$$0 = \frac{d}{dt}\Big|_{t=0} \Psi_{F,p,q}(h_t)$$

= $\frac{1}{q} \frac{d}{dt}\Big|_{t=0} \left(\log \int_{\mathbb{S}^{n-1}} h_t(v)^q d\mu(v)\right) - \frac{p-1}{n(p-1)+p} \frac{d}{dt}\Big|_{t=0} \left(\log \tau_{F,p}([h_t])\right)$
= $\frac{1}{q} \frac{\int_{\mathbb{S}^{n-1}} f^q(v) d\mu(v)}{\int_{\mathbb{S}^{n-1}} h_{K_0}^q d\mu(v)} - \frac{(p-1)}{q(n(p-1)+p)\tau_{F,p}([h_{K_0}])} \int_{\mathbb{S}^{n-1}} f^q(v) dS_{F,p,q}([h_{K_0}], v).$

Since $f \in C^+(\mathbb{S}^{n-1})$ is arbitrary and using the fact that $[h_{K_0}] = K_0$, we have

$$\frac{S_{F,p,q}(K_0,\cdot)}{|\mu|} = \frac{(n(p-1)+p)}{(p-1)\int_{\mathbb{S}^{n-1}}h_{K_0}^q d\mu}\mu(\cdot).$$

Since measure $S_{F,p,q}(K_0, \cdot)$ is homogeneous of degree $\frac{p}{p-1} + n - q$ in K_0 . Let

$$K = \left(\frac{(p-1)\int_{\mathbb{S}^{n-1}}h_{K_0}^q d\mu(v)}{(n(p-1)+p)|\mu|}K_0\right)^{1/(\frac{p}{p-1}+n-q)}$$

then

$$S_{F,p,q}\left(K,\cdot\right) = \mu.$$

Theorem 3.2. Let $1 , <math>1 < q \neq \frac{p}{p-1} + n$ and μ be a nonzero finite Borel measure on \mathbb{S}^{n-1} that is not concentrated on a great subsphere, then there exists a convex body $K \in \mathscr{K}_o^n$ such that

$$F_{p,q}(K,\cdot) = \mu.$$

Proof. Suppose $\{K_{0_i}\} \subset \mathscr{K}_o^n$ is a minimal sequence, i.e.,

$$\inf \left\{ \Psi_{F,p,q}(K_{0_i}) : \quad \tau_{F,p}(K_{0_i}) = |\mu|, \quad K_{0_i} \in \mathscr{K}_o^n \right\},$$
(3.4)

we claim that K_{0_i} is uniformly bounded. If not, then there exists $u_i \in \mathbb{S}^{n-1}$ such that $\rho_{K_{0_i}}(u_i) \to \infty$ as $i \to \infty$. By the definition of support function, $\rho_{K_{0_i}}(u_i)(u_i \cdot v)_+ \leq h_{K_{0_i}}(v)$, then

$$\Psi_{F,p,q}(K_{0_i}) = \frac{1}{q} \log \int_{\mathbb{S}^{n-1}} h_{K_{0_i}}(v)^q d\mu(v) - \frac{p-1}{n(p-1)+p} \log \tau_{F,p}(K_{0_i})$$

$$\geq \frac{1}{q} \log \int_{\mathbb{S}^{n-1}} \rho_{K_{0_i}} (u_i)^q (u_i \cdot v)_+^q d\mu(v) - \frac{(p-1)\log(|\mu|)}{n(p-1)+p}$$

$$= \frac{1}{q} \log \left(\rho_{K_{0_i}}(u_i)^q \int_{\mathbb{S}^{n-1}} (u_i \cdot v)_+^q d\mu(v) \right) - \frac{(p-1)\log(|\mu|)}{n(p-1)+p},$$

where $(t)_+ = \max\{t, 0\}$ for any $t \in \mathbb{R}$. Since μ is not concentrated on any closed hemisphere, there exists a positive constant c_0 such that

$$\int_{\mathbb{S}^{n-1}} \left(u_i \cdot v \right)_+^q d\mu(v) > c_0.$$

Therefore,

$$\Psi_{F,p,q}(K_{0_i}) > \frac{1}{q} \log \left(\rho_{K_{0_i}}(u_i)^q c_0 \right) - \frac{(p-1)\log(|\mu|)}{n(p-1)+p} \to \infty$$

as $i \to \infty$. But this contradicts (3.4). So we conclude that K_{0_i} is uniformly bounded. By Blaschke selection theorem [38, Theorem 1.8.7], there is a convergent subsequence of K_{0_i} , still denoted by K_{0_i} , converges to a compact convex set K of \mathbb{R}^n . By the continuity of anisotropic *p*-torsional rigidity, we get

$$|\mu| = \lim_{i \to \infty} \tau_{F,p} \left(K_{0_i} \right) = \tau_{F,p} (K_0).$$

From (2.14), we have

$$|\mu| = \tau_{F,p}(K_0) \le \frac{p-1}{n(p-1)+p} n^{-\frac{1}{p-1}} \kappa_n^{-\frac{p}{n(p-1)}} V(K_0)^{\frac{n(p-1)+p}{n(p-1)}}.$$

Consequently, we have,

$$V(K_0) \ge \left(|\mu| \frac{n(p-1) + p}{p-1} n^{\frac{1}{p-1}} \kappa_n^{\frac{p}{n(p-1)}} \right)^{\frac{n(p-1)}{n(p-1) + p}} > 0.$$

Thus, K_0 is a convex body in \mathbb{R}^n .

Therefore, K_0 is a minimizer for the minimum problem (3.3), Because of the minimum problem (3.3) and (3.1) are equivalent. So h_{K_0} is the solution to problem (3.1). Combining with Lemma 3.1, we can get the desired result.

4. The L_q -Minkowski problem of anisotropic p-torsional rigidity for general measure with 0 < q < 1

Suppose that μ is a finite discrete measure on \mathbb{S}^{n-1} which is not concentrated on any closed hemisphere of \mathbb{S}^{n-1} , that is,

$$\mu = \sum_{k=1}^{N} \alpha_k \delta_{u_k},$$

where δ_{u_k} is Kronecker delta, $\alpha_1, \ldots, \alpha_N > 0$, and $u_1, \ldots, u_N \in \mathbb{S}^{n-1}$ are not concentrated on any closed hemisphere. For 0 < q < 1, we consider the following minimizing problem

$$\inf\left\{\sup_{\xi\in[f]_{\mu}}\Phi_{f,\mu}(\xi):f\in C^{+}\left(\mathbb{S}^{n-1}\right) \text{ and } \tau_{F,p}([f]_{\mu})=1\right\},$$
(4.1)

where $[f]_{\mu}$ denotes the Aleksandrov body associated to $(f, \operatorname{supp}(\mu))$, and $\Phi_{f,\mu} : [f]_{\mu} \to \mathbb{R}$ is defined by

$$\Phi_{f,\mu}(\xi) = \int_{\mathbb{S}^{n-1}} (f(u) - \xi \cdot u)^q d\mu(u) = \int_{\mathrm{supp}(\mu)} (f(u) - \xi \cdot u)^q d\mu(u).$$
(4.2)

In order to prove that this minimum problem is exactly the solution of L_q -Minkowski problem, the following lemma is necessary. This lemma was proved by Jian and Lu [25, Lemma 3.3] for the first time.

Lemma 4.1. If 0 < q < 1. Let μ be a finite discrete Borel measure on \mathbb{S}^{n-1} which is not concentrated in any closed hemisphere of \mathbb{S}^{n-1} . Then $\Phi_{f,\mu}(\xi)$ is strictly concave for every non-negative continuous function f on \mathbb{S}^{n-1} with $[f]_{\mu}$ has a nonempty interior, there is an unique point $\xi_f \in [f]_{\mu}$, such that

$$\Phi_{f,\mu}(\xi_f) = \sup_{\xi \in [f]_{\mu}} \Phi_{f,\mu}(\xi),$$
(4.3)

where ξ_f depends continuously on f.

Lemma 4.2. Let 1 , <math>0 < q < 1 and μ be a finite discrete measure on \mathbb{S}^{n-1} which is not concentrated in any closed hemisphere of \mathbb{S}^{n-1} , then there exists a function $h \in C^+(\mathbb{S}^{n-1})$ with $\xi_h = o$ and $\tau_{F,p}([h]_{\mu}) = 1$ such that

$$\Phi_{h,\mu}(o) = \inf \left\{ \sup_{\xi \in [f]_{\mu}} \Phi_{f,\mu}(\xi) : f \in C^+ \left(\mathbb{S}^{n-1} \right) \text{ and } \tau_{F,p}([f]_{\mu}) = 1 \right\}$$

Proof. Let $\{f_k\} \subset C^+(\mathbb{S}^{n-1}), \tau_{F,p}([f_k]_\mu) = 1$, and

$$\lim_{k \to +\infty} \Phi_{f_k,\mu}(\xi_{f_k}) = \inf \left\{ \sup_{\xi \in [f]_{\mu}} \Phi_{f,\mu}(\xi) : g \in C^+(\mathbb{S}^{n-1}) \text{ and } \tau_{F,p}([f]_{\mu}) = 1 \right\}.$$
(4.4)

For simplicity, write $h_k = h_{[f_k]_{\mu}}$. Thus

$$h_k(u) \le f_k(u)$$

for $u \in \operatorname{supp}(\mu)$, then $[h_k]_{\mu} = [h_{[f_k]_{\mu}}]_{\mu} = [f_k]_{\mu}$. Since $o \in \operatorname{int}[f_k]_{\mu}$, we have $h_k \in C^+(\mathbb{S}^{n-1})$. Thus, for any $\xi \in [h_k]_{\mu} = [f_k]_{\mu}$, by (4.2), we get

$$\Phi_{h_k,\mu}(\xi) = \int_{\operatorname{supp}(\mu)} (h_k(u) - \xi \cdot u)^q d\mu(u)$$

$$\leq \int_{\operatorname{supp}(\mu)} (f_k(u) - \xi \cdot u)^q d\mu(u)$$

$$= \Phi_{f_k,\mu}(\xi).$$
(4.5)

Thus by (4.5), we obtain

$$\sup_{\xi\in[h_k]_{\mu}}\Phi_{h_k,\mu}(\xi)\leq \sup_{\xi\in[f_k]_{\mu}}\Phi_{f_k,\mu}(\xi).$$

Consequently,

$$\lim_{k \to +\infty} \sup_{\xi \in [h_k]_{\mu}} \Phi_{h_k,\mu}(\xi) \le \lim_{k \to +\infty} \sup_{\xi \in [f_k]_{\mu}} \Phi_{f_k,\mu}(\xi).$$

$$(4.6)$$

Due to $h_k \in C^+(\mathbb{S}^{n-1})$ and $\tau_{F,p}([h_k]_{\mu}) = \tau_{F,p}([f_k]_{\mu}) = 1$, by (4.4), we get $\lim_{k \to +\infty} \sup_{\xi \in [h_k]_{\mu}} \Phi_{h_k,\mu}(\xi) \ge \lim_{k \to +\infty} \sup_{\xi \in [f_k]_{\mu}} \Phi_{f_k,\mu}(\xi),$ from this and (4.6), we deduce

$$\lim_{k \to +\infty} \sup_{\xi \in [h_k]_{\mu}} \Phi_{h_k,\mu}(\xi) = \lim_{k \to +\infty} \sup_{\xi \in [f_k]_{\mu}} \Phi_{f_k,\mu}(\xi).$$

$$(4.7)$$

From Lemma 4.1, (4.7) and (4.4), it follows that

$$\lim_{k \to +\infty} \Phi_{h_k,\mu}(\xi_{h_k}) = \lim_{k \to +\infty} \sup_{\xi \in [h_k]_{\mu}} \Phi_{h_k,\mu}(\xi)$$
$$= \lim_{k \to +\infty} \sup_{\xi \in [f_k]_{\mu}} \Phi_{f_k,\mu}(\xi)$$
$$= \inf \left\{ \sup_{\xi \in [f]_{\mu}} \Phi_{f,\mu}(\xi) : g \in C^+ \left(\mathbb{S}^{n-1} \right) \text{ and } \tau_{F,p}([f]_{\mu}) = 1 \right\}.$$

In view of $[h_k]_{\mu} = [f_k]_{\mu}$, we obtain $h_k = h_{[f_k]_{\mu}} = h_{[h_k]_{\mu}}$, where h_k is the support function of $[h_k]_{\mu}$. Since $\tau_{F,p}$ is translation invariant, then for $x \in \mathbb{R}^n$, $\tau_{F,p}([h_k]_{\mu} + x) = \tau_{F,p}([h_k]_{\mu}) = 1$. By (4.2), we derive

$$\Phi_{h_{([h_k]_{\mu+x})},\mu}(\xi_{h_k}+x) = \int_{\mathrm{supp}(\mu)} \left(h_{([h_k]_{\mu}+x)}(u) - (\xi_{h_k}+x) \cdot u\right)^q d\mu(u) = \Phi_{h_k,\mu}(\xi_{h_k}),$$

for $x \in \mathbb{R}^n$.

Therefore, we can select a sequence, which will also be represented by $\{h_k\} \subset C^+(\mathbb{S}^{n-1})$, with $\tau_{F,p}([h_k]_{\mu}) = 1$ and $\xi_{h_k} = o$, such that

$$\lim_{k \to +\infty} \Phi_{h_k,\mu}(o) = \inf \left\{ \sup_{\xi \in [f]_{\mu}} \Phi_{f,\mu}(\xi) : g \in C^+ \left(\mathbb{S}^{n-1} \right) \text{ and } \tau_{F,p}([f]_{\mu}) = 1 \right\}.$$

We assert that the sequence $\{h_k\}$ is uniformly bounded on \mathbb{S}^{n-1} . Should this assertion not hold, it would be possible to extract a subsequence of $\{h_k\}$, which we shall also denote as $\{h_k\}$, such that

$$\lim_{k \to +\infty} \sup_{u \in \mathbb{S}^{n-1}} h_k(u) = +\infty.$$

Let $R_k = \sup_{u \in \mathbb{S}^{n-1}} h_k(u) = h_k(u_k)$ for some $u_k \in \mathbb{S}^{n-1}$. Given that $\{u_k\} \subset \mathbb{S}^{n-1}$, it follows from the compactness of \mathbb{S}^{n-1} that there exists a convergent subsequence, denoted by $\{u_k\}$, under the assumption that

$$\lim_{k \to +\infty} u_k = u_0 \in \mathbb{S}^{n-1}.$$

Since $\operatorname{supp}(\mu)$ is not concentrated on any closed hemisphere, there exists some $\zeta \in \operatorname{supp}(\mu)$ such that

$$\zeta \cdot u_0 > 0$$

Define $\gamma = \frac{1}{2} (\zeta \cdot u_0) > 0$. It follows that there exists $k_0 \in \mathbb{N}$ such that for all $k \geq k_0$,

$$\zeta \cdot u_k > \gamma.$$

Note that $R_k u_k \in [h_k]_{\mu}$. As a result, for all $k \ge k_0$, it follows that

$$h_k(\zeta) \ge R_k(\zeta \cdot u_k) > R_k\gamma.$$

Given that μ is a finite discrete measure, it follows that for all $k \ge k_0$ and 0 < q < 1, we obtain

$$\lim_{k \to +\infty} \Phi_{h_k,\mu}(o) = \lim_{k \to +\infty} \int_{\mathbb{S}^{n-1}} h_k^q(u) d\mu(u)$$

$$\geq \lim_{k \to +\infty} h_k^q(\zeta) \mu(\zeta)$$

$$> \lim_{k \to +\infty} (R_k \gamma)^q \mu(\zeta) = +\infty.$$
(4.8)

Let $h' \in C^+(\mathbb{S}^{n-1})$ and $\tau_{F,p}([h']_{\mu}) = 1$. Then

$$\lim_{k \to +\infty} \Phi_{h_k,\mu}(o) \le \Phi_{h',\mu}(\xi_{h'}) = \int_{\mathbb{S}^{n-1}} \left(h'(u) - \xi_{h'} \cdot u \right)^q d\mu(u) < +\infty,$$

which contradicts to (4.8). Thus, $\{h_k\}$ is uniformly bounded.

By the Blaschke Selection Theorem [38, Theorem 1.8.7], $\{h_k\}$ has a convergent subsequence, likewise denoted by $\{h_k\}$, letting $h_k \to h$ on \mathbb{S}^{n-1} as $k \to +\infty$. This yields $h \ge 0$, $[h_k]_{\mu} \to [h]_{\mu}$ and $h = h_{[h]_{\mu}}$. The continuity of $\tau_{F,p}$ assures that $\tau_{F,p}([h]_{\mu}) = 1$. From (2.14), we have

$$1 = \tau_{F,p}([h]_{\mu}) \le \frac{p-1}{n(p-1)+p} n^{-\frac{1}{p-1}} \kappa_n^{-\frac{p}{n(p-1)}} V([h]_{\mu})^{\frac{n(p-1)+p}{n(p-1)}}.$$

Consequently, we have

$$V([h]_{\mu}) \ge \left(\frac{n(p-1)+p}{p-1}n^{\frac{1}{p-1}}\kappa_n^{\frac{p}{n(p-1)}}\right)^{\frac{n(p-1)}{n(p-1)+p}} > 0$$

This indicates that $[h]_{\mu}$ is a convex body in \mathbb{R}^n , and $h = h_{[h]_{\mu}}$. Furthermore, it directly follows from Lemma 4.1 that

$$o = \lim_{k \to +\infty} \xi_{h_k} = \xi_h \in \operatorname{int}[h]_{\mu}.$$

Namely, h > 0. Consequently, based on (4.4), we obtain

$$\Phi_{h,\mu}(o) = \inf \left\{ \sup_{\xi \in [f]_{\mu}} \Phi_{f,\mu}(\xi) : g \in C^+ (\mathbb{S}^{n-1}) \text{ and } \tau_{F,p}([f]_{\mu}) = 1 \right\}.$$

This concludes the proof.

Lemma 4.3. Let 1 , <math>0 < q < 1 and μ be a finite discrete measure on \mathbb{S}^{n-1} which is not concentrated in any closed hemisphere of \mathbb{S}^{n-1} , then there exists a function $h \in C^+(\mathbb{S}^{n-1})$ and a constant c > 0 such that

$$d\mu = \lambda dS_{F,p,q}\left([h]_{\mu}, \cdot\right),$$

where

$$\lambda = \frac{p-1}{n(p-1)+p} \int_{\mathbb{S}^{n-1}} h^q(u) d\mu(u).$$

Proof. By Lemma 4.1, we infer the existence of a function $h \in C^+(\mathbb{S}^{n-1})$ with $\xi_h = o$ and $\tau_{F,p}([h]_{\mu}) = 1$ such that

$$\Phi_{h,\mu}(o) = \inf \left\{ \sup_{\xi \in [f]_{\mu}} \Phi_{f,\mu}(\xi) : f \in C^+ \left(\mathbb{S}^{n-1} \right) \text{ and } \tau_{F,p}([f]_{\mu}) = 1 \right\}.$$

Let $\varepsilon > 0$ be sufficiently small, for any $f \in C^+(\mathbb{S}^{n-1})$ and $t \in (-\varepsilon, \varepsilon)$, define

$$\vartheta_t = h + tf,$$

ensuring that $\vartheta_t \in C^+(\mathbb{S}^{n-1})$. From Corollary 2.5, it follows that

$$\lim_{t \to 0} \frac{\tau_{F,p}([\vartheta_t]_{\mu}) - \tau_{F,p}([h]_{\mu})}{t} = \int_{\mathbb{S}^{n-1}} f(u) dS_{F,p}([h]_{\mu}, u)$$
$$= \int_{\mathrm{supp}(\mu)} f(u) dS_{F,p}([h]_{\mu}, u). \tag{4.9}$$

Let $\psi_t = \varphi(t)\vartheta_t$, where

$$\varphi(t) = \tau_{F,p} ([\vartheta_t]_{\mu})^{-\frac{p-1}{n(p-1)+p}}$$

Obviously, $\psi_t \in C^+(\mathbb{S}^{n-1}), \tau_{F,p}([\psi_t]_{\mu}) = 1$ and $\psi_0 = h$. By (4.9), we deduce

$$\lim_{t \to 0} \frac{\psi_t - \psi_0}{t} = -h(u) \frac{p-1}{n(p-1)+p} \tau_{F,p}([h]_{\mu})^{-\frac{p-1}{n(p-1)+p}-1} \int_{\mathbb{S}^{n-1}} f(u) dS_{F,p}([h]_{\mu}, u) + f$$
$$= -h(u) \frac{p-1}{n(p-1)+p} \int_{\mathbb{S}^{n-1}} f(u) dS_{F,p}([h]_{\mu}, u) + f.$$
(4.10)

Let $\xi(t) = \xi_{\psi_t}$ and

$$\Phi_{\mu}(t) = \sup_{\xi \in [\psi_t]_{\mu}} \int_{\mathbb{S}^{n-1}} (\psi_t(u) - \xi \cdot u)^q \, d\mu(u)$$

=
$$\int_{\mathbb{S}^{n-1}} (\psi_t(u) - \xi(t) \cdot u)^q \, d\mu(u).$$
(4.11)

Since $\xi(t) \in \operatorname{int} [\psi_t]_{\mu}$, by (4.11), we obtain

$$\int_{\mathbb{S}^{n-1}} \left(\psi_t(u) - \xi(t) \cdot u \right)^{q-1} u_k d\mu(u) = 0, \tag{4.12}$$

for $k = 1, \ldots, n$, where $u = (u_1, \ldots, u_n)^T$. Noting $\xi(0) = \xi_h = o$ and taking t = 0 in (4.12), we get

$$\int_{\mathbb{S}^{n-1}} h^{q-1}(u) u_k d\mu(u) = 0, \qquad (4.13)$$

for $k = 1, \ldots, n$. Hence

$$\int_{\mathbb{S}^{n-1}} h^{q-1}(u) u d\mu(u) = 0.$$
(4.14)

Let

$$H_k(t,\xi_1,\ldots,\xi_n) = \int_{\mathbb{S}^{n-1}} \left(\psi_t(u) - (\xi_1 u_1 + \cdots + \xi_n u_n)\right)^{q-1} u_k d\mu(u),$$

for $k = 1, \ldots, n$. Then

$$\frac{\partial H_k}{\partial \xi_l} = (1-q) \int_{\mathbb{S}^{n-1}} \left(\psi_t(u) - (\xi_1 u_1 + \dots + \xi_n u_n) \right)^{q-2} u_k u_l d\mu(u).$$

Let $H = (H_1, \ldots, H_n)$ and $\xi = (\xi_1, \ldots, \xi_n)$. Thus,

$$\left(\left.\frac{\partial H}{\partial \xi}\right|_{(0,\dots,0)}\right)_{n\times n} = (1-q)\int_{\mathbb{S}^{n-1}} h^{q-2}(u)uu^T d\mu(u),$$

where uu^T is an $n \times n$ matrix.

Since μ is not concentrated on any closed hemisphere, $\operatorname{supp}(\mu)$ spans the whole space \mathbb{R}^n . Consequently, for any $x \in \mathbb{R}^n$ with $x \neq 0$, there exists a $u_{i_0} \in \operatorname{supp}(\mu)$ such that $u_{k_0} \cdot x \neq 0$. Therefore, for 0 < q < 1, we derive that

$$x^{T} \left(\frac{\partial H}{\partial \xi} \Big|_{(0,...,0)} \right) x = x^{T} \left((1-q) \int_{\mathbb{S}^{n-1}} h^{q-2}(u) u u^{T} d\mu(u) \right) x$$
$$= (1-q) \int_{\mathbb{S}^{n-1}} h^{q-2}(u) (x \cdot u)^{2} d\mu(u)$$
$$\geq (1-q) h^{p-2} (u_{k_{0}}) (x \cdot u_{k_{0}})^{2} \mu (u_{k_{0}}) > 0.$$

This indicates that $\left(\frac{\partial H}{\partial \xi}\Big|_{(0,...,0)}\right)$ is positive definite, which consequently implies that

$$\det\left(\left.\frac{\partial H}{\partial \xi}\right|_{(0,\dots,0)}\right) \neq 0.$$

From this, the facts that for $k = 1, ..., n, H_k(0, ..., 0) = 0$ follows by equation (4.13) and $\frac{\partial H_k}{\partial \xi_l}$ is continuous on a neighborhood of (0, ..., 0) for all $1 \le k, l \le n$, and the implicit function theorem, we can conclude that

$$\xi'(0) = (\xi'_1(0), \dots, \xi'_n(0))$$

exists.

Note that $\Phi_{\mu}(0) = \Phi_{h,\mu}(o)$ and $\Phi_{\mu}(t) = \Phi_{\psi_t,\mu}(\xi_{\psi_t})$ with $\psi_t \in C^+(\mathbb{S}^{n-1})$ and $\tau_{F,p}([\psi_t]_{\mu}) = 1$. By Lemma 4.2, it follows that

$$\Phi_{\mu}(t) \ge \Phi_{\mu}(0),$$

i.e., $\Phi_{\mu}(0)$ is an extreme value of $\Phi_{\mu}(t)$. Therefore, from (4.10) and (4.14), we obtain

$$\begin{split} 0 &= \frac{1}{q} \frac{d}{dt} \Big|_{t=0} \Phi_{\mu}(t) \\ &= \int_{\mathbb{S}^{n-1}} h^{q-1}(u) \left(-h(u) \frac{p-1}{n(p-1)+p} \int_{\mathbb{S}^{n-1}} f(u) dS_{F,p}([h]_{\mu}, u) + f(u) - \xi'(0) \cdot u \right) d\mu(u) \\ &= -\frac{p-1}{n(p-1)+p} \int_{\mathbb{S}^{n-1}} h^{q}(u) d\mu(u) \int_{\mathbb{S}^{n-1}} f(u) dS_{F,p}([h]_{\mu}, u) + \int_{\mathbb{S}^{n-1}} h^{p-1}(u) f(u) d\mu(u) \\ &- \int_{\mathbb{S}^{n-1}} \xi'(0) \cdot h^{p-1}(u) u d\mu(u) \\ &= \int_{\mathbb{S}^{n-1}} h^{p-1}(u) f(u) d\mu(u) - \frac{p-1}{n(p-1)+p} \int_{\mathbb{S}^{n-1}} h^{q}(u) d\mu(u) \int_{\mathbb{S}^{n-1}} f(u) dS_{F,p}([h]_{\mu}, u). \end{split}$$

Namely, for all $f \in C^+(\mathbb{S}^{n-1})$,

$$\int_{\mathbb{S}^{n-1}} h^{p-1}(u) f(u) d\mu(u) = \frac{p-1}{n(p-1)+p} \int_{\mathbb{S}^{n-1}} h^q(u) d\mu(u) \int_{\mathbb{S}^{n-1}} f(u) dS_{F,p}([h]_{\mu}, u).$$

Since $h = h_{[h]_{\mu}}$, then

$$d\mu(u) = \lambda dS_{F,p,q}\left([h]_{\mu}, u\right),$$

where

$$\lambda = \frac{p-1}{n(p-1)+p} \int_{\mathbb{S}^{n-1}} h^q(u) d\mu(u).$$

Theorem 4.4. Let 1 , <math>0 < q < 1 and μ be a finite Borel measure on \mathbb{S}^{n-1} which is not concentrated in any closed hemisphere. Then there exists a convex body K in \mathbb{R}^n , such that

$$d\mu = \lambda dS_{F,p,q}(K, \cdot),$$

where λ is a positive constant.

Proof. In accordance with the proof detailed in [38, Theorem 8.2.2], given a finite Borel measure μ on \mathbb{S}^{n-1} that is not concentrated within any closed hemisphere, it is feasible to construct a sequence of finite discrete measures $\{\mu_k\}$ on \mathbb{S}^{n-1} such that $\mu_k(\mathbb{S}^{n-1}) = \mu(\mathbb{S}^{n-1})$ and μ_k converges weakly to μ as k approaches infinity. Specifically, for sufficiently large values of k, μ_k will also not be concentrated in any closed hemisphere. According

to Lemma 4.3, for each μ_k , there exists a function $h_k \in C^+(\mathbb{S}^{n-1})$ and a positive constant λ_k , satisfying the conditions that

$$\mu_k = \lambda_k S_{F,p,q}([h_k]_{\mu_k}, \cdot), \tag{4.15}$$

where

$$\lambda_k = \frac{p-1}{n(p-1)+p} \int_{\mathbb{S}^{n-1}} h_k^q(u) d\mu_k(u).$$
(4.16)

Moreover, h_k satisfies that $\xi_{h_k} = o, \tau_{F,p}([h_k]_{\mu_k}) = 1$, and

$$\Phi_{h_k,\mu_k}(o) = \inf \left\{ \sup_{\xi \in [f]_{\mu_k}} \Phi_{f,\mu_k}(\xi) : f \in C^+ \left(\mathbb{S}^{n-1} \right) \text{ and } \tau_{F,p}([f]_{\mu_k}) = 1 \right\},$$

where

$$[f]_{\mu_k} = \bigcap_{u \in \text{supp}(\mu_k)} \left\{ \xi \in \mathbb{R}^n : \xi \cdot u \le f(u) \right\},\$$

and

$$\Phi_{f,\mu_k}(\xi) = \int_{\mathbb{S}^{n-1}} (f(u) - \xi \cdot u)^q d\mu_k(u) = \int_{\mathrm{supp}(\mu_k)} (f(u) - \xi \cdot u)^q d\mu_k(u)$$

The subsequent proof closely mirrors the approach outlined in [25, 11]. Nonetheless, in the interest of ensuring a comprehensive and rigorous demonstration, we provide a detailed proof process.

Define $g_k = \Phi_{h_k,\mu_k}(o)$. We aim to demonstrate that g_k is uniformly bounded. Regarding the Aleksandrov body corresponding to $(1, \operatorname{supp}(\mu_k))$, we denote it by $[1]_{\mu_k}$. Let

$$\bar{f}_k = (\tau_{F,p}([1]_{\mu_k}))^{-\frac{p-1}{n(p-1)+p}}.$$

We observe that $[\bar{f}_k]_{\mu_k} = \bar{f}_k[1]_{\mu_k}$. It follows directly that $\tau_{F,p}([\bar{f}_k]_{\mu_k}) = 1$. Given that $\mu_k(\mathbb{S}^{n-1}) = \mu(\mathbb{S}^{n-1})$, we get

$$g_{k} = \Phi_{h_{k},\mu_{k}}(o)$$

$$\leq \sup_{\xi \in \left[\bar{f}_{k}\right]_{\mu_{k}}} \int_{\mathbb{S}^{n-1}} \left(\bar{f}_{k}(u) - \xi \cdot u\right)^{q} d\mu_{k}(u)$$

$$\leq \int_{\mathbb{S}^{n-1}} \operatorname{diam}(\left[\bar{f}_{k}\right]_{\mu_{k}})^{q} d\mu_{k}(u)$$

$$= \operatorname{diam}\left(\left[\bar{f}_{k}\right]_{\mu_{k}}\right)^{q} \mu\left(\mathbb{S}^{n-1}\right)$$

$$= \bar{f}_{k}^{q} \operatorname{diam}\left(\left[1\right]_{\mu_{k}}\right)^{q} \mu\left(\mathbb{S}^{n-1}\right).$$

$$(4.17)$$

To demonstrate that g_k is uniformly bounded, we will establish that the diameter of $[1]_{\mu_k}$, denoted as diam $([1]_{\mu_k})$, is uniformly bounded. If this were not the case, it would be possible to identify a sequence $\{\xi_k\}$ such that $\xi_k \in [1]_{\mu_k}$ and

$$\lim_{k \to +\infty} |\xi_k| = +\infty$$

Let $\bar{\xi}_k = \frac{\xi_k}{|\xi_k|} \in \mathbb{S}^{n-1}$. By the compactness of \mathbb{S}^{n-1} , we may assume, without loss of generality, that

$$\lim_{k \to +\infty} \bar{\xi}_k = \bar{\xi} \in \mathbb{S}^{n-1}.$$

Given that $\operatorname{supp}(\mu)$ is not concentrated in any closed hemisphere, there exists $\bar{x} \in \operatorname{supp}(\mu)$, such that

$$\bar{\xi} \cdot \bar{x} > 0. \tag{4.18}$$

Let $O(\bar{x}) \subset \mathbb{S}^{n-1}$ denote an arbitrary neighborhood of \bar{x} . Consequently, it follows that $\liminf_{k \to +\infty} \mu_k(O(\bar{x})) \ge \mu(O(\bar{x})) > 0.$

Observe that for sufficiently large values of k,

 $O(\bar{x}) \bigcap \operatorname{supp}(\mu_k) \neq \emptyset$, for infinitely many k,

which allows us to construct a sequence $\{\bar{x}_{k_l}\}$ such that

$$\bar{x}_{k_l} \in \operatorname{supp}\left(\mu_{k_l}\right)$$
 and $\lim_{i \to \infty} \bar{x}_{k_l} = \bar{x}_{k_l}$

It is important to note that $\xi_{k_l} \in [1]_{\mu_{k_l}}$. Consequently,

$$\xi_{k_l} \cdot \bar{x}_{k_l} \le h_{[1]_{\mu_{k_l}}} (\bar{x}_{k_l}) \le 1$$

which implies that,

$$\bar{\xi}_{k_l} \cdot \bar{x}_{k_l} \le \frac{1}{|\xi_{k_l}|}.$$

By taking the limit, we obtain

$$\bar{\xi} \cdot \bar{x} \le 0$$

which contradicts (4.18). Thus, there exists a positive constant C > 0, such that for all $k \in \mathbb{N}$,

$$\operatorname{diam}\left([1]_{\mu_k}\right) \le C. \tag{4.19}$$

Given that $\mathbb{B}^n \subset [1]_{\mu_k}$ for each $k \in \mathbb{N}$, it follows that for 1 , we have

$$\bar{f}_k \le (\tau_{F,p}(\mathbb{B}^n))^{-\frac{p-1}{n(p-1)+p}}.$$
(4.20)

From (4.17), (4.19) and (4.20), it can be deduced that for 0 < q < 1 and all $k \in \mathbb{N}$,

$$g_k \le C^q \tau_{F,p}(\mathbb{B}^n))^{-\frac{q(p-1)}{n(p-1)+p}} \mu\left(\mathbb{S}^{n-1}\right).$$

$$(4.21)$$

Thus, the sequence g_k is uniformly bounded.

Now, we shall prove that the sequence $\{h_k\}$ is uniformly bounded on \mathbb{S}^{n-1} . Otherwise, that there exists a subsequence $\{h_{k_l}\} \subset \{h_k\}$ such that

$$\lim_{l \to +\infty} \sup_{u \in \mathbb{S}^{n-1}} h_{k_l}(u) = +\infty.$$

Let $R_{k_l} = \sup_{u \in \mathbb{S}^{n-1}} h_{k_l}(u) = h_{k_l}(u_{k_l})$. Since $\{u_{k_l}\} \subset \mathbb{S}^{n-1}$, from the compactness of \mathbb{S}^{n-1} , without loss of generality we can assume

$$\lim_{l \to +\infty} u_{k_l} = u_0 \in \mathbb{S}^{n-1}$$

Since $\operatorname{supp}(\mu)$ is not concentrated in any closed hemisphere, there exists $v_0 \in \operatorname{supp}(\mu)$ such that

$$v_0 \cdot u_0 > 0.$$

Let $O(v_0)$ be a small neighborhood of v_0 such that for all $u \in O(v_0)$, we have that

$$u \cdot u_0 > 0$$

Let $\gamma = \frac{1}{2} (u \cdot u_0) > 0$ for $u \in O(v_0)$, noting that $R_{k_l} u_{k_l} \in [h_{k_l}]_{\mu_{k_l}}$. For sufficiently large l, it follows that for all $u \in O(v_0)$,

 $u \cdot u_{k_l} > \gamma,$

and

$$\mu_{k_l}\left(O\left(v_0\right)\right) \ge \mu\left(O\left(v_0\right)\right) > 0.$$

Thus,

$$h_{k_l}(u) \ge R_{k_l} \left(u \cdot u_{k_l} \right) > R_{k_l} \gamma.$$

Therefore, when l is sufficiently large, we obtain

$$g_{k_l} = \int_{\mathbb{S}^{n-1}} h_{k_l}^q(u) d\mu_{k_l}(u)$$

$$\geq \int_{O(v_0)} h_{k_l}^q(u) d\mu_{k_l}(u)$$

$$\geq R_{k_l}^q \int_{O(v_0)} \gamma^q d\mu_{k_l}(u)$$

$$\geq R_{k_l}^q \int_{O(v_0)} \gamma^q d\mu(u),$$

which implies that $g_{k_l} \to +\infty$ as $l \to +\infty$. This contradicts (4.21). That is, $\{h_k\}$ is uniformly bounded on \mathbb{S}^{n-1} .

Since $\{h_k\}$ is support function of $\{[h_k]_{\mu}\}$, thus, $\{[h_k]_{\mu}\}$ is also uniformly bounded. By the Blaschke Selection Theorem [38, Theorem 1.8.7], there is a subsequence of $\{[h_k]_{\mu}\}$, also written as $\{[h_k]_{\mu}\}$, converges to a compact convex set $[h]_{\mu}$ in \mathbb{R}^n .

The continuity of $\tau_{F,p}$ assures that $\tau_{F,p}([h]_{\mu}) = 1$. From (2.14), we have

$$1 = \tau_{F,p}([h]_{\mu}) \le \frac{p-1}{n(p-1)+p} n^{-\frac{1}{p-1}} \kappa_n^{-\frac{p}{n(p-1)}} V([h]_{\mu})^{\frac{n(p-1)+p}{n(p-1)}}.$$

Consequently, we have,

$$V([h]_{\mu}) \ge \left(\frac{n(p-1)+p}{p-1}n^{\frac{1}{p-1}}\kappa_n^{\frac{p}{n(p-1)}}\right)^{\frac{n(p-1)}{n(p-1)+p}} > 0$$

This implies that $[h]_{\mu}$ is a convex body in \mathbb{R}^n .

Thus, $[h_k]_{\mu} \to [h]_{\mu} \in \mathscr{K}_0^n$ as $k \to +\infty$, and $h = h_{[h]_{\mu}} \ge 0$ is support function of $[h]_{\mu}$, and

$$\lim_{k \to +\infty} \lambda_k = \lim_{k \to +\infty} \left(\frac{p-1}{n(p-1)+p} \int_{\mathbb{S}^{n-1}} h_k^q(u) d\mu_k(u) \right)$$
$$= \frac{p-1}{n(p-1)+p} \int_{\mathbb{S}^{n-1}} h^q(u) d\mu(u)$$
$$= \lambda \ge 0.$$

Thus, by (4.15), (4.16) and (2.17), we obtain

$$d\mu = \lambda dS_{F,p,q}\left([h]_{\mu}, \cdot\right).$$

Obviously, $\lambda \neq 0$, then $\lambda > 0$, let $K = [h]_{\mu}$, we have

$$d\mu = \lambda dS_{F,p,q}(K, \cdot),$$

this yields the desired result.

References

- M. Akman and S. Mukherjee, On the Minkowski problem for p-harmonic measures, Calc. Var. Partial Differential Equations, 64 (2025), 1: 36. 1
- [2] C. Bianchin and G. Ciraolo, Wulff shape characterizations in overdetermined anisotropic elliptic problems, Comm. Partial Differential Equations, 43 (2018), 5: 790-820. 2
- [3] K. J. Böröczky, E. Lutwak, D. Yang and G. Zhang, The Gauss image problem, Comm. Pure Appl. Math., 73 (2020), 7: 1406-1452. 1
- [4] B. Chen, C Li and W Wang, The L_p Gauss dual Minkowski problem, arXiv:2412.13557, 2024. 1
- [5] B. Chen, X. Zhao, W. Wang and P. Zhao, The L_p Minkowski problem for q-torsional rigidity, Adv. Calc. Var., 17 (2024), 3: 587-603. 2, 3

CHAO LI AND BIN CHEN

- [6] Z. Chen and Q. Dai, The L_p Minkowski problem for torsion, J. Math. Anal. Appl., 488, (2020), DOI: 10.1016/j.jmaa.2020.124060. 2, 3
- [7] A. Colesanti, Brunn-Minkowski inequalities for variational functionals and related problems, Adv. Math., 194 (2005), 105-140. 2, 5
- [8] A. Colesanti and M.Fimiani, The Minkowski problem for torsional rigidity, Indiana Univ. Math. J., 59 (2010), 3: 1013-1039.
- [9] F. Della Pietra and N. Gavitone, Sharp bounds for the first eigenvalue and the torsional rigidity related to some anisotropic operators, Math. Nachr., **287** (2014), 194-209. 1, 2, 5
- [10] Y. Feng, Y. Li and L. Xu, Existence of solutions to the Gaussian dual Minkowski problem, J. Differential Equations, 416 (2025), 268-298. 1
- [11] Y. Feng, Y. Zhou and B. He, The L_p electrostatic q-capacitary Minkowski problem for general measures, J. Math. Anal. Appl. 487 (2020), 123959. 15
- [12] D. Francesco, G. Nunzia and G. Serena, On functionals involving the torsional rigidity related to some classes of nonlinear operators, J. Differential Equations, 265 (2018), 12: 6424-6442. 2
- [13] A. Greco and B. Mebrate, An overdetermined problem related to the Finsler p-Laplacian, Mathematika, 70 (2024), 4: e12267. 2
- [14] C. Haberl, E. Lutwak, D. Yang and G. Zhang, The even Orlicz Minkowski problem, Adv. Math., 224 (2010), 2485-2510. 1
- [15] J. Hu, A Gauss curvature flow approach to the torsional Minkowski problem, J. Differential Equations, 385 (2024), 254-279. 2
- [16] J. Hu, The torsion log-Minkowski Problem, J. Geom. Anal., 34 (2024). DOI: 10.1007/s12220-024-01670-1. 2
- [17] J. Hu and J. Liu, On the L_p torsional Minkowski problem for 0 , Adv. Appl. Math., 128 (2021), 102188. 2, 3
- [18] J. Hu and J. Liu and D. Ma, A Gauss curvature flow to the Orlicz-Minkowski problem for torsional rigidity, J. Geom. Anal., 32 (2022), 2: 63. 2
- [19] Y. Huang, E. Lutwak, D. Yang and G. Zhang, Geometric measures in the dual Brunn-Minkowski theory and their associated Minkowski problems, Acta Math., 216 (2016), 325-388. 1
- [20] Y. Huang, C. Song and L. Xu, Hadamard variational formulas for p-torsion and p-eigenvalue with applications, Geom. Dedicata., 197 (2018), 61-76. 2
- [21] Y. Huang, D. Xi and Y. Zhao, The Minkowski problem in Gaussian probability space, Adv. Math., 385 (2021), 107769. 1
- [22] D. Jerison, Harmonic measure in convex domains, Bull. Amer. Math. Soc. (N.S.), 21 (1989), 2: 255-260. 1
- [23] D. Jerison, Prescribing harmonic measure on convex domains, Invent. Math., 105 (1991), 1: 375-400. 1
- [24] D. Jerison, A Minkowski problem for electrostatic capacity, Acta Math., 176 (1996), 1: 1-47. 1
- [25] H. Jian and J. Lu, Existence of solutions to the Orlicz-Minkowski problem, Adv. Math., 344 (2019), 262-288. 10, 15
- [26] L. Kryvonos and D. Langharst, Measure theoretic Minkowski's existence theorem and projection bodies, Trans. Amer. Math. Soc., 376 (2023), 8447-8493. 1, 4
- [27] G. V. Livshyts, An extension of Minkowski's theorem and its applications to questions about projections for measures, Adv. Math., 356 (2019), 106803. 1
- [28] N. Li, D. Ye and B. Zhu, The dual Minkowski problem for unbounded closed convex sets, Math. Ann., 388 (2024), 2001-2039. 1
- [29] C. Li, Minkowski problem of anisotropic p-torsional rigidity, arXiv:2501.00687v2, 2025. 2, 5
- [30] C. Li and G. Wei, Existence of solution for L_p -Minkowski problem of $0 with measures in <math>\mathbb{R}^n$, Int. J. Math., **34** (2023), 3: 2350009. 1
- [31] C. Li and X. Zhao, Flow by Gauss curvature to the Minkowski problem of p-harmonic measure, arXiv:2404.18757v2, 2024. 1
- [32] H. Z. Li and B. T. Xu, Hyperbolic p-sum and Horospherical p-Brunn-Minkowski theory in hyperbolic space, (2022), arXiv preprint arXiv:2211.06875. 1
- [33] N. Li and B. Zhu, The Orlicz-Minkowski problem for torsional rigidity, J. Differential Equations, 269 (2020), 10: 8549–8572. 2
- [34] J. Liu, The L_p-Gaussian Minkowski problem, Calc. Var. Partial Differential Equations, 61 (2022), 28. 1, 4
- [35] E. Lutwak, The Brunn-Minkowski-Firey theory I: Mixed volumes and the Minkowski problem, J. Differential Geom., 38 (1993), 1: 131-150. 1, 2

- [36] E. Lutwak, D. Xi, D. Yang and G. Zhang, Chord measures in integral geometry and their Minkowski problems, Comm. Pure Appl. Math., 77 (2024), 7: 3277-3330.1
- [37] E. Lutwak, D. Yang and G. Y. Zhang, L_p dual curvature measure, Adv. Math., **329** (2018), 85-132. 3
- [38] R. Schneider, Convex Bodies: The Brunn-Minkowski theory, 2nd edn, Cambridge Univ. Press, Cambridge, 2014. 1, 3, 9, 12, 14, 17
- [39] R. Schneider, A Brunn-Minkowski theory for coconvex sets of finite volume, Adv. Math., 332 (2018), 199-234. 1
- [40] G. Sun, L. Xu and P. Zhang, The uniqueness of the L_p Minkowski problem for q-torsional rigidity, Acta Math. Sci., 41 (2021), 1405-1416. 2
- [41] X. Zhao and P. Zhao, The dual Minkowski problem for q-torsional rigidity, arXiv:2411.00779, 2024.
- [42] D. Zou and G. Xiong, The L_p Minkowski problem for the electrostatic p-capacity, J. Differential Geom., 116 (2020), 3: 555-596. 1

Chao Li: School of Mathematics and Statistics, Ningxia University, Yinchuan, Ningxia, 750021, China.

NINGXIA BASIC SCIENCE RESEARCH CENTER OF MATHEMATICS, NINGXIA UNIVERSITY, YINCHUAN 750021, CHINA.

Email address: lichao@nxu.edu.cn, lichao166298@163.com

BIN CHEN: DEPARTMENT OF MATHEMATICS, LANZHOU UNIVERSITY OF TECHNOLOGY, LANZHOU, GANSU, 730050, CHINA.

Email address: chenb121223@163.com