ON LOCAL SMOOTHING ESTIMATES FOR WAVE EQUATIONS

SHENGWEN GAN AND SHUKUN WU

ABSTRACT. We prove sharp local smoothing estimates for wave equations on compact Riemannian manifolds in 3+1 dimensions and obtained improved estimates in higher dimensions. This is achieved by deriving local smoothing estimates for certain Fourier integral operators. We also obtain improved local smoothing estimates for wave equations in Euclidean spaces.

1. INTRODUCTION

1.1. **Overview.** This paper proves local smoothing estimates for wave equations on Euclidean spaces and on general compact Riemannian manifold. Let (M,g) be \mathbb{R}^n or a smooth, compact *n*-dimensional Riemannian manifold with the associated Laplace–Beltrami operator Δ_q . Let *u* be the solution of the Cauchy problem

(1.1)
$$\begin{cases} (\partial_t^2 - \Delta_g)u(x,t) = 0, \quad (x,t) \in M \times \mathbb{R}, \\ u(x,0) = f_0(x), \quad \partial_t u(x,0) = f_1(x). \end{cases}$$

When $M = \mathbb{R}^n$ with flat metric, Peral [17] showed that for fixed time t,

$$||u(\cdot,t)||_{L^p(M)} \lesssim_{M,g} ||f_0||_{L^p_{s_p}(M)} + ||f_1||_{L^p_{s_n-1}(M)}$$

for $p \in (1, \infty)$ and $s_p = (n-1)|\frac{1}{2} - \frac{1}{p}|$ (for general (M, g), this was proved by Seeger-Sogge-Stein in [19]). Sogge [20] observed that with an averaging over $t \sim 1$, there is a gain of regularity in L^p for 2 . In the same paper, he made thefollowing local smoothing conjecture:

Conjecture 1.1. Let $M = \mathbb{R}^n$ equipped with the flat metric, let $s_p = \frac{(n-1)}{2} - \frac{(n-1)}{p}$, and let u solve the Cauchy problem (1.1). Then for all $p \ge 2 + \frac{2}{n-1}$ and $\sigma < \frac{1}{p}$,

(1.2)
$$\|u\|_{L^{p}(\mathbb{R}^{n}\times[1,2])} \leq C_{p,\sigma} \left(\|f_{0}\|_{L^{p}_{s_{p}-\sigma}(\mathbb{R}^{n})} + \|f_{1}\|_{L^{p}_{s_{p}-1-\sigma}(\mathbb{R}^{n})}\right)$$

The local smoothing phenomenon also exists on a general compact manifold (M, g). However, the range of exponents is quite different from the flat case in Conjecture 1.1. Miniccozi and Sogge [16] constructed examples to show that (1.2) fails for all $\sigma < \frac{1}{p}$ when $p < p_n^+$, where

$$p_n^+ := \begin{cases} 2 + \frac{8}{3n-3}, & \text{if } n \text{ is odd,} \\ 2 + \frac{8}{3n-2}, & \text{if } n \text{ is even.} \end{cases}$$

This leads naturally to the following conjecture:

Conjecture 1.2. For $n \ge 2$, let (M, g) be an n-dimensional compact Riemannian manifold, and let u solve the Cauchy problem (1.1). Then for all $p \ge p_n^+$ and $\sigma < \frac{1}{p}$,

(1.3)
$$\|u\|_{L^p(M\times[1,2])} \le C_{p,\sigma} \left(\|f_0\|_{L^p_{s_p-\sigma}(M)} + \|f_1\|_{L^p_{s_p-1-\sigma}(M)} \right).$$

Our main result is the following:

Theorem 1.3. Conjectures 1.1 and 1.2 are true when $p \ge p(n)$, where

(1.4)
$$p(n) := \begin{cases} 2 + \frac{4}{3}, & \text{if } n = 3, \\ 2 + \frac{8}{3n-4}, & \text{if } n \ge 4. \end{cases}$$

Since $p(3) = p_3^+ = \frac{10}{3}$, Theorem 1.3 verifies Conjecture 1.2 when n = 3. Note also that for larger n, p(n) and p_n^+ differ by $O(\frac{1}{n^2})$.

Remark 1.4. One may form a stronger conjecture by including the endpoint $\sigma = \frac{1}{p}$. For example, it was proved by Heo, Nazarov, and Seeger [13] that Conjecture 1.1 is true when $p \ge 2 + \frac{4}{n-3}$ and $\sigma = \frac{1}{p}$ for $n \ge 4$. It is open whether there are local smoothing estimates at the endpoint $\sigma = \frac{1}{p}$ for dimensions n = 2, 3.

To study Conjecture 1.1, Wolff [23] initiated a program based on decoupling inequalities. The investigation of decoupling inequalities culminated in Bourgain-Demeter's resolution on the ℓ^2 -decoupling theorem [4]. As a direct corollary, they showed that Conjecture 1.1 is true when $p > 2 + \frac{4}{n-1}$, where $2 + \frac{4}{n-1}$ is the decoupling exponent for the light cone. The approach was adapted to general compact Riemannian manifolds by Beltran-Hickman-Sogge [1], where they proved the same range of p for Conjecture 1.2. We refer to [2] for more historical information.

Conjecture 1.1 was verified when n = 2 by Guth-Wang-Zhang [12], however, via a different approach. The authors there proved a sharp reverse square function estimate, which implies Conjecture 1.1. This approach was modified by Gao-Liu-Miao-Xi [8] to prove Conjecture 1.2 in the same dimensions.

In this paper, we follow Wolff's footsteps and use decoupling inequalities to study the local smoothing problem. Prior to this work, the decoupling range $p > 2 + \frac{4}{n-1}$ is the best-known result for both Conjecture 1.1 and 1.2 when $n \ge 3$. Further discussion of our methods will be provided after an introduction to the Fourier integral operator, a standard tool for the local smoothing problem on manifolds.

1.2. Fourier integral operator (FIO). Through a standard parametrix construction of the half wave operator $e^{it\sqrt{-\Delta_g}}$ (see [2] for details), the study of Conjecture 1.2 boils down to the analysis of certain Fourier integral operators:

(1.5)
$$\mathcal{F}f(x,t) := \int_{\mathbb{R}^n} e^{i\phi(x,t;\xi)} a(x,t;\xi) \widehat{f}(\xi) \mathrm{d}\xi.$$

Here, the symbol $a \in S^0(\mathbb{R}^{n+1} \times \mathbb{R}^n)$ is supported in $\{(x,t;\xi) : |x|^2 + t^2 \leq 1\}$. The phase function $\phi(x,t;\xi)$ is homogeneous of degree 1 in ξ and smooth away from $\xi = 0$. Also, ϕ satisfies

• (H1) Non-degeneracy condition.

rank $\partial_{x\xi}^2 \phi(x,t;\xi) = n$ for all $(x,t;\xi) \in \text{supp } a$.

• (H2) Positive definiteness condition. Consider the Gauss map G: supp $a \to S^n$ by $G(z;\xi) := \frac{G_0(z;\xi)}{|G_0(z;\xi)|}$, where

$$G_0(z;\xi) := \bigwedge_{j=1}^n \partial_{\xi_j} \partial_z \phi(z;\xi).$$

Then for all $(z, \xi_0) \in \text{supp } a$,

$$\partial_{\xi\xi}^2 \langle \partial_z \phi(z,\xi), G(z,\xi_0) \rangle |_{\xi=\xi_0}$$

has rank n-1 with n-1 positive eigenvalues.

A prototypical example for the phase function is given by $\phi(x,t;\xi) = \langle x,\xi \rangle + t|\xi|$. The Fourier integral operator defined using this phase function corresponds to the classical half-wave propagator $e^{it\sqrt{-\Delta}}f$. One may also keep in mind another example $\phi(x,t;\xi) = \langle x,\xi \rangle + t|\xi'|^2/\xi_n$. Here $\xi' = (\xi_1,\ldots,\xi_{n-1})$.

As shown in [2], to prove Conjecture 1.2, it suffices to prove the following conjecture for FIOs satisfying the two conditions (H1) and (H2).

Conjecture 1.5. Let \mathcal{F} be defined in (1.5). Suppose \mathcal{F} satisfies conditions (H1) and (H2). Then for $p \ge p_n^+$ and $\sigma > (n-1)(\frac{1}{2} - \frac{1}{p}) - \frac{1}{p}$,

(1.6)
$$\|\mathcal{F}f\|_{L^p(\mathbb{R}^n \times [1,2])} \lesssim \|f\|_{L^p_{\sigma}(\mathbb{R}^n)}.$$

Remark 1.6. Via a similar argument, to prove Conjecture 1.1, it suffices to prove (1.6) with $\mathcal{F} = e^{it\sqrt{-\Delta}}$ for the given range of p.

Similarly, to establish Theorem 1.3, we only need to prove

Theorem 1.7. Conjecture 1.5 is true when $p \ge p(n)$, where p(n) is given in (1.4).

1.3. Main ideas and outline. To explain our ideas, we take $\mathcal{F} = e^{it\sqrt{-\Delta}}$ in Conjecture 1.5. The same ideas apply to general FIOs.

Since the kernel of the operator $e^{it\sqrt{-\Delta}}$ is essentially the Fourier inversion of the surface measure of the cone $\{(\xi, |\xi|) : \xi \in \mathbb{R}^n\}$, it is natural to connect the local smoothing problem with Fourier restriction theory. This connection was shown to be successful by Rogers [18] in the study of the Schrödinger equations (with the half-wave operator $e^{it\sqrt{-\Delta}}$ being replaced by the Schrödinger operator $e^{it\Delta}$). In fact, Rogers showed that the local smoothing conjecture for Schrödinger equations is equivalent to the Fourier restriction conjecture for paraboloid. Moreover, since the tools developed for the parabolic restriction conjecture are versatile and apply equally well to hypersurfaces with a positive definite second fundamental form, it is reasonable to believe that the same ideas will extend to the local smoothing problem of the same type. Successful examples can be found in [6], where fractional Schrödinger equations with the operators $e^{it(-\Delta)^{\alpha/2}}$, $\alpha > 1$, are investigated.

However, there is a significant difference when one attempts to relate the local smoothing problem for wave equations to the conic restriction problem. Indeed, the local smoothing problem is generally believed to be strictly harder than the conic and parabolic restriction problem (see [21]). For example, the exponent $2 + \frac{4}{n-1}$, also known as the Stein-Tomas exponent, was established for the restriction problem back in the 1970s. In contrast, for the local smoothing problem, this exponent was only obtained much later by Bourgain and Demeter in 2014.

One key difference between paraboloid and light cone is that the former possesses a better symmetry: the natural rescaling for paraboloid, the parabolic rescaling (see [9]), is an isotropic rescaling when restricting to the initial data, whereas the natural rescaling for the light cone, the Lorentz rescaling (see Section 5.1), is nonisotropic. As a consequence, although induction is well-established in the study of the parabolic restriction problem and hence the related local smoothing problem for Schrödinger equations, there seems to be no satisfactory induction on scales scheme for wave equations.

In this paper, we propose a possible candidate for induction on scales for the wave equations. To better explain our idea, we will compare the half-wave operator $e^{it\sqrt{-\Delta}}$ with the Schrödinger operator $e^{it\Delta}$. After standard reductions, the desired local smoothing estimates for the two operators and the parabolic restriction problem are reduced to the following local forms:

Local smoothing estimate for wave equations: For supp $\widehat{f} \subset B_1^n \setminus B_{1/2}^n$,

(1.7)
$$\|e^{it\sqrt{-\Delta}}f\|_{L^p(B^{n+1}_R)} \lesssim R^{(n-1)(\frac{1}{2}-\frac{1}{p})}\|f\|_p.$$

Local smoothing estimate for Schrödinger equations: For supp $\widehat{f} \subset B_1^n$,

(1.8)
$$\|e^{it\Delta}f\|_{L^p(B_R^{n+1})} \lesssim R^{n(\frac{1}{2}-\frac{1}{p})} \|f\|_p$$

Fourier restriction estimate for paraboloid: For supp $\widehat{f} \subset B_1^n$,

(1.9)
$$\|e^{it\Delta}f\|_{L^p(B^{n+1}_R)} \lesssim \|\hat{f}\|_p.$$

To study the restriction estimate (1.9) via induction on scales, Guth [9] introduced the mixed norm

(1.10)
$$\|\widehat{f}\|_{2}^{\frac{2}{p}} \sup_{\theta} \|\widehat{f}_{\theta}\|_{L^{2}_{avg}(\theta)}^{1-\frac{2}{p}}$$

in place of the L^p -norm $\|\widehat{f}\|_{L^p(B_1^n)}$, where $\|\widehat{f}_{\theta}\|_{L^2_{avg}(\theta)}^2 = |\theta|^{-1} \|\widehat{f}_{\theta}\|_{L^2(\theta)}^2$, and θ ranges over $R^{-1/2}$ -balls in B_1^n . The mixed norm (1.10) can be viewed as the L^p -norm by interpolation, in which the quantity $\sup_{\theta} \|\widehat{f}_{\theta}\|_{L^2_{avg}(\theta)}$ plays the roll of $\|\widehat{f}\|_{L^{\infty}}$. It is crucial that $\sup_{\theta} \|\widehat{f}_{\theta}\|_{L^2_{avg}(\theta)}$ is an L^2 -average. If this is simply replaced by $\|\widehat{f}\|_{L^{\infty}}$, inductions will generally fail.

Given the connection between the parabolic restriction conjecture and the local smoothing conjecture for Schrödinger equations, it is thus natural to consider a suitable mixed norm similar to (1.10) to investigate (1.8). One subtlety is that the right-hand side of (1.8) lies in the physical space, while the right-hand side of (1.9) lies in the frequency space. This leads to a different mixed norm, which may be chosen as follows:

$$R^{n(\frac{1}{2}-\frac{1}{p})} \|f\|_{2}^{\frac{2}{p}} \sup_{B} \|f\|_{L^{2}_{avg}(B)}^{1-\frac{2}{p}}$$

where B ranges over $R^{1/2}$ -balls in B_R^n . Using this mixed norm, it is possible that the recent framework developed by Wang and the second author in [22] for the restriction conjecture can lead to parallel results for local smoothing estimate for Schrödinger equations (1.8).

Based on the above discussion, it seems natural to study the local smoothing estimate for wave equations (1.7) by considering a similar mixed norm:

(1.11)
$$R^{(n-1)(\frac{1}{2}-\frac{1}{p})} \|f\|_{2}^{\frac{2}{p}} \sup_{B} \|f\|_{L^{2}_{avg}(B)}^{1-\frac{2}{p}},$$

where B ranges over $R^{1/2}$ -balls in B_R^n . However, this naive choice of the mixed norm does not work, since the form in (1.11) is not invariant under Lorentz rescaling, a non-isotropic one.

Inspired by the work [12], we introduce a mixed norm

(1.12)
$$R^{(n-1)(\frac{1}{2}-\frac{1}{p})} \|f\|_2^{\frac{1}{p}} \mathcal{W}(f, B_R^{n+1})^{1-\frac{2}{p}},$$

where the wave packet density $\mathcal{W}(f, B_R^{n+1})$ is a rescaling-invariant form of an L^2 average that serves as an analogue of $||f||_{\infty}$. The precise definition of $\mathcal{W}(f, B_R^{n+1})$ is given in Definition 4.1. Roughly speaking, it is the supreme of all the L^2 -average over conic wave packets and their rescaled counterparts under Lorentz rescaling.

The introduction of the wave packet density is one of the main ideas of this work. With the mixed norm (1.12) in hand, induction on scales becomes possible, and we would like to adapt the framework developed in [22] to study (1.7). After the standard wave packet decomposition and several steps of dyadic pigeonholing, our goal is to establish the following: Suppose $f = \sum_{T \in \mathbb{T}} f_T$ is a wave packet decomposition of f, where \mathbb{T} is a family of $1 \times R^{1/2} \times \cdots \times R^{1/2} \times R$ -planks. Given a union of unit balls $X \subset B_R^{n+1}$ such that each unit ball $B \subset X$ intersects with $\lesssim \mu$ planks in \mathbb{T} , prove that when p = p(n),

(1.13)
$$\|e^{it\sqrt{-\Delta}}f\|_{L^p(X)} \lesssim R^{(n-1)(\frac{1}{2}-\frac{1}{p})} \|f\|_2^{\frac{2}{p}} \mathcal{W}(f, B_R^{n+1})^{1-\frac{2}{p}}$$

As in [22], we aim to

- (1) Prove a refined decoupling inequality for the cone to bound $||e^{it\sqrt{-\Delta}}f||_{L^{q_n}(X)}$ with $q_n = \frac{2(n+1)}{n-1}$ being the decoupling exponent.
- (2) Explore orthogonality in the L^2 space. That is, estimate $||e^{it\sqrt{-\Delta}}f||_{L^2(X)}$.

It is fairly standard to establish item (1), and we prove the refined decoupling inequality for general FIOs in the appendix. However, it is not clear what the optimal L^2 -orthogonality for $||e^{it\sqrt{-\Delta}}f||_{L^2(X)}$ is, as X is a union of unit balls. One simple approach is to enlarge X to its $R^{1/2}$ -neighborhood and then use L^2 orthogonality on the $R^{1/2}$ -balls. As we will see below, this already allows us to obtain some useful L^2 estimates. A refined approach to the L^2 estimate will be discussed later in Subsection 1.4. This refined L^2 argument is another key idea of the paper, ultimately allowing us to obtain the sharp estimate in dimensions 3+1.

We introduce a parameter l that measures how many $R^{1/2}$ -balls interact with a plank T. Specifically, for each plank $T \in \mathbb{T}$, define a shading $Y(T) = X \cap T$, and assume that Y(T) intersects with $\sim l$ many $R^{1/2}$ -balls in \mathbb{R}^{n+1} . Thus, by using L^2 -orthogonality inside each $R^{1/2}$ -ball in $N_{R^{1/2}}(X)$, we have the following L^2 -estimate

(1.14)
$$\|e^{it\sqrt{-\Delta}}f\|_{L^2(X)}^2 \lesssim \|e^{it\sqrt{-\Delta}}f\|_{L^2(N_{R^{1/2}}(X))}^2 \lesssim (lR^{1/2})\|f\|_2^2.$$

Next, at the decoupling endpoint $p = \frac{2(n+1)}{n-1}$, we use the refined decoupling inequality for the cone to have

$$\|e^{it\sqrt{-\Delta}}f\|_{L^p(X)}^p \lesssim \mu^{\frac{2}{n-1}} \sum_{T \in \mathbb{T}} \|e^{it\sqrt{-\Delta}}f_T\|_p^p,$$

After normalization, we assume that $|e^{it\sqrt{-\Delta}}f_T|$ is essentially the characteristic function of the $1 \times R^{1/2} \times \cdots \times R^{1/2} \times R$ -plank *T*. Thus, when $p = \frac{2(n+1)}{n-1}$,

(1.15)
$$\|e^{it\sqrt{-\Delta}}f\|_{L^p(X)}^p \lesssim \mu^{\frac{2}{n-1}}(R^{\frac{n+1}{2}}\#\mathbb{T})$$

It remains to find an upper bound for $\mu^{\frac{2}{n-1}} \#\mathbb{T}$, which is an incidence problem between the unit balls in X and the $1 \times R^{1/2} \times \cdots \times R^{1/2} \times R$ -planks in \mathbb{T} .

Now we are going to use the power of induction on scales. Broadly speaking, induction on scales allows us to impose a two-ends condition (which is a non-concentration condition) on the shading Y(T). Also, the wave packet density $\mathcal{W}(f, B_R^n)$ imposes a non-concentration condition on the collection of planks \mathbb{T} (see Remark 4.8), which, in particular, implies $\mathcal{W}(f, B_R^{n+1})^2 \gtrsim R^{-\frac{n+1}{4}} (\#\mathbb{T})^{1/2}$. Therefore, by using the hairbrush structure (see Figure 5.5) and a "2-broad" assumption on each unit ball in X, we have

$$\mu \lesssim l^{-1/2} (\#\mathbb{T})^{1/2} \lesssim l^{-1/2} R^{\frac{n+1}{4}} \mathcal{W}(f, B_R^{n+1})^2.$$

We remark that the hairbrush structure for the cone has the following special property: all the light rays intersecting a fixed line are essentially disjoint.

Finally, we plug the incidence bound back to (1.15) to get when $p = \frac{2(n+1)}{n-1}$,

$$\|e^{it\sqrt{-\Delta}}f\|_{L^p(X)}^p \lesssim R^2 (R^{-\frac{n-3}{2(n-1)}}l^{-\frac{1}{n-1}}) \|f\|_2^2 \mathcal{W}(f, B_R^{n+1})^{p-2}.$$

Interpolate this with (1.14) and use the fact that $l \leq R^{1/2}$ to conclude (1.7) for $p = 2 + \frac{8}{3n-4}$. We refer to Section 2.1 for a sketch of the proof. This concludes Theorem 1.7 when $n \geq 4$.

Remark 1.8. The wave packet density $\mathcal{W}(f, B_R^n)$ is quite versatile. Modifications on its definition will allow us to impose stronger non-concentration conditions on the set of planks \mathbb{T} . We refer the readers to Remark 4.9 for further discussions.

1.4. A refined L^2 estimate. Finally, let us focus on n = 3 and further explore the L^2 -orthogonality. The approach outlined in the previous subsection already gives strong local smoothing estimates for wave equations (note that $2 + \frac{8}{3n-4} < 2 + \frac{4}{n-1}$ when $n \ge 3$) in this dimension. Nonetheless, it does not make full use of the local information of the set X in the L^2 estimate (1.14). This is because in (1.14), we bound $\|e^{it\sqrt{-\Delta}}f\|_{L^2(X)}^2$ simply by $\|e^{it\sqrt{-\Delta}}f\|_{L^2(N_{R^{1/2}}(X))}^2$ and give up the information of X on each $R^{1/2}$ -ball.

To capture this information, we use the broad-narrow argument and the approach developed in [15] to prove a refined L^2 estimate in place of (1.14). For each $R^{1/2}$ -ball Q, we assume $|X \cap Q| \sim \sigma$ for some positive integer σ . Via the covering lemma, Lemma 5.7, there is a set of $1 \times R^{1/2} \times R^{1/2} \times R^{1/2}$ -slabs $\mathbb{S}(Q)$ such that the following is true:

- (1) $\mathbb{S}(Q)$ essentially forms a cover of $X \cap Q$.
- (2) $\#\mathbb{S}(Q) \sim \sigma/\rho$.
- (3) For any arbitrary $1 \times R^{1/2} \times R^{1/2} \times R^{1/2}$ -slab $S \subset Q, |S \cap X| \lesssim \rho$.

Let f_Q be the sum of wave packets f_T that $T \cap Q \neq \emptyset$. The key observation is that if each $B \subset X \cap Q$ is "4-broad", then we have the L^2 estimate

$$\|e^{it\sqrt{-\Delta}}f\|_{L^{2}(X\cap Q)}^{2} \lesssim \sum_{S\in\mathbb{S}(Q)} \|e^{it\sqrt{-\Delta}}f\|_{L^{2}(X\cap S)}^{2} \lesssim \#\mathbb{S}(Q)\|f_{Q}\|_{2}^{2} \sim (\sigma/\rho)\|f_{Q}\|_{2}^{2}.$$

Sum up all $R^{1/2}$ -balls to obtain the refined L^2 estimate

(1.16)
$$\|e^{it\sqrt{-\Delta}}f\|_{L^{2}(X)}^{2} \lesssim l(\sigma/\rho)\|f\|_{2}^{2}.$$

Note that (1.16) is stronger than (1.14) when σ/ρ is smaller than $R^{1/2}$.

On the other hand, let $\mathbb{T}(Q)$ be the set of planks in \mathbb{T} that intersect Q. Since ${T \cap Q}_{T \in \mathbb{T}(Q)}$ are distinct $1 \times R^{1/2} \times R^{1/2} \times R^{1/2}$ -slabs and since $|S \cap X| \lesssim \rho$ for any arbitrary $1 \times R^{1/2} \times R^{1/2} \times R^{1/2}$ -slab $S \subset Q$, by double counting the unit balls in $X \cap Q$ and planks in $\mathbb{T}(Q)$, we have $\#\mathbb{T}(Q) \gtrsim \mu(\sigma/\rho)$. Using the hairbrush structure, this estimate gives an improved incidence estimate

$$\mu \lesssim (\#\mathbb{T})^{1/2} \rho^{1/2} (l\sigma)^{-1/2} \lesssim \rho^{1/2} (l\sigma)^{-1/2} R \cdot \mathcal{W}(f, B_R^4)^2.$$

Plugging this back to (1.15) and interpolating with (1.16), we prove (1.6) when n = 3 and $p = \frac{10}{3}$, conditioned on the assumption that each $B \subset X \cap Q$ is 4-broad. If each $B \subset X \cap Q$ is not 4-broad, then we use induction and decoupling to conclude (1.6) at the same exponent $p = \frac{10}{3}$, thereby completing the proof of Theorem 1.7. A sketch of our argument will be given in Section 2.2. The idea developed for n = 3 might lead to another improvement over Theorem 1.7 when n > 4. However, it is unclear to us whether this is enough to solve Conjecture 1.5 for all higher dimensions.

1.5. A broad & two-ends algorithm. We conclude the introduction with a comment on a technical difficulty encountered during the proof. Let $e^{it\sqrt{-\Delta}}f =$ $\sum_{T \in \mathbb{T}} e^{it\sqrt{-\Delta}} f_T$ be the wave packet decomposition. Via a standard broad-narrow reduction, it suffices to consider the "broad" norm $\|\sum_{T\in\mathbb{T}}e^{it\sqrt{-\Delta}}f_T\|_{\mathrm{BL}^p(X)}$. That is, for each unit ball $B \subset X$, $|\sum_{T \in \mathbb{T}} e^{it\sqrt{-\Delta}} f_T| \mathbf{1}_B$ is "broad", which essentially means that its contribution comes from groups of wave packets with quantitatively linearly independent directions. To utilize the "broad" information and the twoends condition on the shading $Y(T) = X \cap T$, we need the following two conditions to hold simultaneously:

- (1) For each unit ball $B \subset X$, $|\sum_{T \in \mathbb{T}} e^{it\sqrt{-\Delta}} f_T| \mathbf{1}_B$ is broad. (2) For each plank $T \in \mathbb{T}$, $Y(T) = X \cap T$ satisfies a two-ends condition.

It turns out that achieving the two conditions requires considerable effort, mostly because the sum in $|\sum_{T \in \mathbb{T}} e^{it\sqrt{-\Delta}} f_T| \mathbf{1}_B$ is a sum of oscillatory functions. To compare, if we were to consider $\sum_{T \in \mathbb{T}} |e^{it\sqrt{-\Delta}} f_T| \mathbf{1}_B$, where oscillation is removed, then several steps of dyadic pigeonholing would suffice to ensure the two required conditions hold simultaneously. When there is oscillation, as is the case here, we essentially prove that if the two conditions do not hold simultaneously, then the total incidence $\sum_{B \subset X} \#\{T \in \mathbb{T} : T \cap B \neq \emptyset\}$ must decrease by a factor of R^{κ} for some absolute $\kappa > 0$. As a result, we develop an algorithm to show that the two conditions must hold simultaneously after finite steps of refinement. See Section 6, Remark 6.6, and Section 7 for details.

1.6. Structure of the paper. In Section 2, we give a sketch of the proof of our main result. Section 3 discusses preliminaries, followed by the introduction of wave packet density in Section 4. Section 5 contains some geometric results, and we introduce the algorithm in Section 6 that helps us realize the broad & two-ends reduction on our operator. Finally, we prove the main theorem in Sections 7 and 8. The appendix includes a proof of the refined decoupling theorem for the cone in the variable coefficient setting.

Notation: Throughout the paper, we use #E to denote the cardinality of a finite set. If \mathcal{E} is a family of sets in \mathbb{R}^n , we use $\cup_{\mathcal{E}}$ to denote $\cup_{E \in \mathcal{E}} E$. For $A, B \ge 0$, we use $A \lesssim B$ to mean $A \le CB$ for an absolute (big) constant C, and use $A \sim B$ to mean $A \lesssim B$ and $B \lesssim A$. For a given R > 1, we use $A \lesssim B$ to denote $A \le c_{\eta} R^{\eta} B$ for all $\eta > 0$. For two finite sets E, F, we say E is a **refinement** of F, if $E \subset F$ and $\#E \gtrsim \#F$. We use RapDec(R) to denote the quantity such that for all $R \ge 1$ and all $N \ge 1$,

$$\operatorname{RapDec}(R) \leq C_N R^{-N}$$

We use $B_r^m(z)$ to denote the ball of radius r centered at z in \mathbb{R}^m . When the center is not specified, B_r^m typically refers to the ball of radius r centered at the origin, unless stated otherwise. We use $N_r(X)$ to denote the r-neighborhood of X.

Choice of parameters: $0 < \varepsilon < 1$, $R \in [1, \lambda^{1-\varepsilon}]$, $K = R^{\varepsilon^{50}}$, $K_{\circ} = R^{\varepsilon^{100}}$, $\kappa = R^{\varepsilon^{500}}$, $\delta = \varepsilon^{1000}$. Under such choice, we have

$$R^{\delta} \ll \kappa \ll K_{\circ} \ll K \ll R \ll \lambda.$$

2. A sketch of the proof for Theorem 1.7

In this section, we give a sketch of our proof of Theorem 1.7, mostly focusing on the numerology. For simplicity, we take $\mathcal{F} = e^{it\sqrt{-\Delta}}$ in (1.6). After standard global-to-local reductions, it suffices to prove that for p = p(n) and a function fdefined on \mathbb{R}^n with $\operatorname{supp} \widehat{f} \subset B_1^n \setminus B_{1/2}^n$,

$$||e^{it\sqrt{-\Delta}}f||_{L^p(B_R^{n+1})} \lesssim R^{(n-1)(\frac{1}{2}-\frac{1}{p})}||f||_p.$$

Let $f = \sum_{T \in \mathbb{T}} f_T$ be the wave packet decomposition, where \mathbb{T} is a family of $1 \times R^{1/2} \times \cdots \times R^{1/2} \times R$ -planks. Let $q_n = \frac{2(n+1)}{n-1}$ be the decoupling exponent. By dyadic pigeonholing and rescaling, we can assume each wave packet $e^{it\sqrt{-\Delta}}f_T$ has amplitude ~ 1 , and hence $\|e^{it\sqrt{-\Delta}}f_T\|_{L^{q_n}(B_R)}^{q_n} \sim R^{\frac{n+1}{2}}$ for all $T \in \mathbb{T}$. For the purpose of induction, we replace the L^p -norm $\|f\|_p$ by a mixed norm $\|f\|_2^{\frac{2}{p}}\mathcal{W}(f, B_R^{n+1})^{1-\frac{2}{p}}$. Roughly speaking, the wave packet density $\mathcal{W}(f, B_R^{n+1})$ is defined by

$$\mathcal{W}(f, B_R^{n+1}) := \sup_U (\frac{1}{|U|} \int_{\mathbb{R}^n} \sum_{T \in \mathbb{T}[U]} |e^{it\sqrt{-\Delta}} f_T|^2)^{1/2}.$$

Here, U ranges over all $Rs^2 \times Rs \times \cdots \times Rs \times R$ -planks where s ranges over dyadic numbers in $[R^{-1/2}, 1]$. We also require the longest direction of U points to a light ray direction, and the shortest direction of U points to the normal direction of the cone at that light ray. $\mathbb{T}[U]$ is the set of planks contained in U that share the same direction as U. We remark that the U defined here is a higher-dimensional generalization of the one in [12].

It suffices to show that when p = p(n),

(2.1)
$$\|\sum_{T\in\mathbb{T}}e^{it\sqrt{-\Delta}}f_T\|_{L^p(B_R^{n+1})}^p \lesssim R^{(n-1)(\frac{p}{2}-1)}\|f\|_2^2 \mathcal{W}(f, B_R^{n+1})^{p-2}$$

By testing with U = T and $U = B_R^{n+1}$, we have the following lower bound for $\mathcal{W}(f, B_R^{n+1})$:

(2.2)
$$\mathcal{W}(f, B_R^{n+1})^2 \gtrsim \max\{1, R^{-\frac{n+1}{2}}(\#\mathbb{T})\} \gtrsim R^{-\frac{n+1}{4}}(\#\mathbb{T})^{1/2}.$$

We also record the refined decoupling inequality for cones, whose variable coefficient version is proved in Theorem A.3.

Theorem 2.1. Suppose we have the wave packet decomposition for $e^{it\sqrt{-\Delta}}f$ in B_{R}^{n+1} :

$$e^{it\sqrt{-\Delta}}f = \sum_{T \in \mathbb{W}} e^{it\sqrt{-\Delta}}f_T.$$

Assume that $||f_T||_{L^p}$ are about the same for all $T \in \mathbb{W}$. Let Y be a disjoint union of unit balls in B_R^{n+1} , each of which intersects $\leq M$ many $T \in \mathbb{W}$. Then for $2 \leq p \leq \frac{2(n+1)}{n-1}$ and any $\varepsilon > 0$,

(2.3)
$$\|e^{it\sqrt{-\Delta}}f\|_{L^p(Y)} \lesssim R^{\varepsilon} M^{\frac{1}{2}-\frac{1}{p}} \Big(\sum_{T\in\mathbb{W}} \|e^{it\sqrt{-\Delta}}f_T\|_{L^p(w_{B_R^{n+1}})}^p\Big)^{1/p}.$$




FIGURE 1.

- 2.1. Sketch for $n \ge 4$. By several steps of dyadic pigeonholing, we can obtain
 - (a) a union of unit balls $X \subset B_R^{n+1}$, (b) a number $\mu \in [1, R^{(n-1)/2}]$,

 - (c) a number $l \in [1, R^{1/2}]$,
 - (d) a number $\lambda \in [1, R^{(n+1)/2}]$

such that the following is true (see Figure 1):

(1)
$$\|\sum_{T\in\mathbb{T}} e^{it\sqrt{-\Delta}} f_T\|_{L^p(B)}$$
 are about the same for all $B \subset X$, and

(2.4)
$$\|\sum_{T\in\mathbb{T}}e^{it\sqrt{-\Delta}}f_T\|_{L^p(B_R^{n+1})}^p \lesssim \|\sum_{T\in\mathbb{T}}e^{it\sqrt{-\Delta}}f_T\|_{L^p(X)}^p.$$

- (2) Each $B \subset X$ intersects with $\sim \mu$ many planks in \mathbb{T} .
- (3) For each plank $T \in \mathbb{T}$, the shading $Y(T) := X \cap T$ intersects with $\sim l$ many $R^{1/2}$ -balls.
- (4) For each $T \in \mathbb{T}$, $|Y(T)| \sim \lambda$.

With the help of the broad & two-ends algorithm discussed in Sections 6 and 7, we can further assume that

- (5) The shading Y(T) satisfies a two-ends condition for all $T \in \mathbb{T}$.
- (6) For each unit ball $B \subset X$, $|\sum_{T \in \mathbb{T}} e^{it\sqrt{-\Delta}} f_T| \mathbf{1}_B$ is "2-broad", which implies the following: There exist two refinements $\mathbb{T}_1(B)$, $\mathbb{T}_2(B)$ of $\mathbb{T}(B)$ such that T_1 and T_2 are quantitatively transverse for all $T_1 \in \mathbb{T}_1(B), T_2 \in \mathbb{T}_2(B)$. Here $\mathbb{T}(B) := \{T \in \mathbb{T} : Y(T) \cap B \neq \emptyset\}.$

Next, apply the refined decoupling theorem, Theorem 2.1, to the right-hand side of (2.4) so that, recalling $q_n = \frac{2(n+1)}{n-1}$,

$$\|\sum_{T\in\mathbb{T}}e^{it\sqrt{-\Delta}}f_T\|_{L^{q_n}(X)}^{q_n} \lesssim \mu^{\frac{2}{n-1}}\sum_{T\in\mathbb{T}}\|e^{it\sqrt{-\Delta}}f_T\|_{L^{q_n}(B_R^{n+1})}^{q_n}.$$

Since $R^{\frac{n+1}{2}} \sim \|e^{it\sqrt{-\Delta}} f_T\|_{L^{q_n}(B_R^{n+1})}^{q_n} \lesssim R^{-\frac{n+1}{n-1}} \|e^{it\sqrt{-\Delta}} f_T\|_{L^2(B_R^{n+1})}^{q_n} \lesssim \|f_T\|_2^{q_n}$, we get

(2.5)
$$\|\sum_{T\in\mathbb{T}}e^{it\sqrt{-\Delta}}f_T\|_{L^{q_n}(X)}^{q_n} \lesssim \mu^{\frac{2}{n-1}}(R^{\frac{n+1}{2}}\#\mathbb{T}) \lesssim R^2(R^{-1}\mu^{\frac{2}{n-1}})\|f\|_2^2.$$

To obtain an upper on μ , we will use the hairbrush structure (see Figure 5). On the one hand, by double-counting, we have

(2.6)
$$|X| \sim \mu^{-1} \lambda(\#\mathbb{T}).$$

On the other hand, pick a plank $T_0 \in \mathbb{T}$ and define a hairbrush

$$\mathbb{H}(T_0) = \{T \in \mathbb{T} : Y(T) \cap Y(T_0) \neq \emptyset, T \text{ and } T_0 \text{ are quantitatively transverse} \}.$$

One sees that the planks in $\mathbb{H}(T_0)$ are morally disjoint. Therefore,

(2.7)
$$|X| \ge |\bigcup_{T \in \mathbb{H}(T_0)} Y(T)| \gtrsim \sum_{T \in \mathbb{H}(T_0)} |Y(T)| \gtrsim l\mu\lambda,$$

where we used $|Y(T)| \sim \lambda$ and $\#\mathbb{H}(T_0) \sim l\mu$.

Combining (2.6) and (2.7), we get

$$\mu \lesssim l^{-1/2} (\#\mathbb{T})^{1/2}.$$

Plug this incidence bound back to (2.5) and use (2.2) to have

$$\begin{aligned} \|\sum_{T\in\mathbb{T}} e^{it\sqrt{-\Delta}} f_T \|_{L^{q_n}(X)}^{q_n} &\lesssim R^2 (R^{-1} (l^{-1} \# \mathbb{T})^{\frac{1}{n-1}}) \|f\|_2^2 \\ &\lesssim R^2 (R^{-\frac{n-3}{2(n-1)}} l^{-\frac{1}{n-1}}) \|f\|_2^2 \mathcal{W}(f, B_R^{n+1})^{q_n-2} \\ &= R^{(n-1)(\frac{1}{2} - \frac{1}{q_n})} (R^{-\frac{n-3}{2(n-1)}} l^{-\frac{1}{n-1}}) \|f\|_2^2 \mathcal{W}(f, B_R^{n+1})^{q_n-2}. \end{aligned}$$

$$(2.8)$$

10

Since Y(T) intersects with $\leq l \mod R^{1/2}$ -balls for each plank $T \in \mathbb{T}$, by using L^2 -orthogonality inside each $R^{1/2}$ -ball in $N_{R^{1/2}}(X)$,

(2.9)
$$\|\sum_{T\in\mathbb{T}}e^{it\sqrt{-\Delta}}f_T\|_{L^2(X)}^2 \lesssim \|\sum_{T\in\mathbb{T}}e^{it\sqrt{-\Delta}}f_T\|_{L^2(N_{R^{1/2}}(X))}^2 \lesssim lR^{1/2}\|f\|_2^2.$$

Interpolating (2.8) with (2.9) and noting that $l \lesssim R^{1/2}$, we obtain (2.1) when $p = 2 + \frac{8}{3n-4}$.

2.2. Sketch for n = 3. As outlined in the introduction, the key new ingredient in this case is a refined L^2 estimate. This leads to an enhanced hairbrush estimate, yielding an improved incidence result.

Recall (2.1) that our goal is to prove

(2.10)
$$\|e^{it\sqrt{-\Delta}}f\|_{L^p(B_R^4)}^p \lesssim R^{2(\frac{1}{2}-\frac{1}{p})} \|f\|_2^2 \mathcal{W}(f, B_R^4)^{p-2}$$

for $p = \frac{10}{3}$. Similar to the previous subsection, we can obtain

- (a) a union of unit balls $X \subset B_R^4$,
- (b) a number $\mu \in [1, R]$,
- (c) a number $l \in [1, R^{1/2}]$,
- (d) a number $\lambda \in [1, R^2]$

such that the following is true:

(1) $\|\sum_{T \in \mathbb{T}} e^{it\sqrt{-\Delta}} f_T\|_{L^p(B)}$ are about the same for all $B \subset X$, and

(2.11)
$$\|\sum_{T\in\mathbb{T}}e^{it\sqrt{-\Delta}}f_T\|_{L^p(B_R^4)}^p \lesssim \|\sum_{T\in\mathbb{T}}e^{it\sqrt{-\Delta}}f_T\|_{L^p(X)}^p.$$

- (2) Each $B \subset X$ intersects with $\sim \mu$ many planks in \mathbb{T} .
- (3) For each plank $T \in \mathbb{T}$, the shading $Y(T) := X \cap T$ intersects with $\sim l$ many $R^{1/2}$ -balls.
- (4) For each $T \in \mathbb{T}$, $|Y(T)| \sim \lambda$.
- (5) The shading Y(T) satisfies a two-ends condition for all $T \in \mathbb{T}$.
- (6) For each unit ball $B \subset X$, $|\sum_{T \in \mathbb{T}} e^{it\sqrt{-\Delta}} f_T | \mathbf{1}_B$ is "4-broad".

In addition, by several more steps of dyadic pigeonholing and a covering lemma, Lemma 5.7, we can obtain

- (e) a number $\sigma \in [1, R^2]$,
- (f) a number $\rho \in [1, R^{3/2}],$
- (g) a number $m \ge 1$,

such that the following is true:

(7) There is a set of $R^{1/2}$ -balls $\mathcal{Q} = \{Q\}$ that cover X. For each $Q \in \mathcal{Q}$,

$$|X \cap Q| \sim \sigma.$$

(8) For any $Q \in Q$, $X \cap Q$ is ρ -regular with respect to $1 \times R^{1/2} \times R^{1/2} \times R^{1/2}$ slabs, in the following sense: For any $1 \times R^{1/2} \times R^{1/2} \times R^{1/2}$ -slab $S \subset Q$,

$$|X \cap S| \le \rho$$

and there are

$$\sim |X \cap Q|/\rho \sim \sigma/\rho$$

many $1 \times R^{1/2} \times R^{1/2} \times R^{1/2}$ -slabs that cover $X \cap Q$. Denote by $\mathbb{S}(Q) = \{S\}$ these slabs that cover $X \cap Q$, so $\#\mathbb{S}(Q) \sim \sigma/\rho$

(9) For each $Q \in \mathcal{Q}$ that intersects Y(T),

$$|Q \cap Y(T)| \sim \lambda/l,$$

which yields

$$ho \gtrsim \lambda/l$$

by the definition of ρ , since $Q \cap T$ is a $1 \times R^{1/2} \times R^{1/2} \times R^{1/2}$ -slab and since $Q \cap Y(T) = X \cap (Q \cap T)$.

(10) For each $Q \in \mathcal{Q}$,

$$\#\mathbb{T}(Q) \sim m,$$

where $\mathbb{T}(Q) = \{T \in \mathbb{T} : Q \cap T \neq \emptyset\}$. Note that the $1 \times R^{1/2} \times R^{1/2} \times R^{1/2}$ -slabs $\{Q \cap T : T \in \mathbb{T}(Q)\}$ are essentially distinct.

Let us see how these data lead to the proof of (2.10). Similar to (2.5), via the refined decoupling theorem, Theorem 2.1, we have

(2.12)
$$\|\sum_{T\in\mathbb{T}}e^{it\sqrt{-\Delta}}f_T\|_{L^4(X)}^4 \lesssim \mu(\#\mathbb{T}) \lesssim R^2(R^{-1}\mu)\|f\|_2^2.$$

Next, we will obtain an upper bound for μ . Inside an $R^{1/2}$ -ball $Q \in \mathcal{Q}$, by doublecounting the incidence between the unit balls in $X \cap Q$ and the $1 \times R^{1/2} \times R^{1/2} \times R^{1/2}$ slabs $\{Q \cap T : T \in \mathbb{T}(Q)\}$ and noting that $\lambda/l \leq \rho$, we obtain

(2.13)
$$\mu\sigma \lesssim \rho m.$$

What follows is a hairbrush argument. Fix a plank $T_0 \in \mathbb{T}$ and let $\overline{T}_0 = N_{R^{1/2}}(T_0)$ be the *R*-tube containing T_0 . For each unit ball $B \subset X$, let $\mathbb{T}(B) := \{T \in \mathbb{T} : Y(T) \cap B \neq \emptyset\}$. From the outcome that $|\sum_{T \in \mathbb{T}} e^{it\sqrt{-\Delta}} f_T| \mathbf{1}_B$ is broad, we know that for each $B \subset X \cap \overline{T}_0$, there is a refinement $\mathbb{T}_1(B)$ of $\mathbb{T}(B)$ such that planks in $\mathbb{T}_1(B)$ are quantitatively transverse to \overline{T}_0 . Since there are $\sim l$ many $R^{1/2}$ -balls $Q \in Q$ intersecting T_0 , we have $\#\mathbb{H}(T_0) \gtrsim lm$, where

 $\mathbb{H}(T_0) = \{T \in \mathbb{T} : Y(T) \cap (X \cap \overline{T}_0) \neq \emptyset, T \text{ and } T_0 \text{ are quantitatively transverse} \}.$ Therefore,

(2.14)
$$|X| \gtrsim \sum_{T \in \mathbb{H}(T_0)} |Y(T)| \gtrsim \lambda lm.$$

On the other hand, we have by double-counting that

$$(2.15) |X| \sim \mu^{-1}(\#\mathbb{T})\lambda.$$

Combining (2.14) and (2.15), we get

$$\mu \lesssim \# \mathbb{T}(lm)^{-1},$$

which, via (2.13), implies

$$\mu \lesssim (\#\mathbb{T})^{1/2} \rho^{1/2} (l\sigma)^{-1/2}.$$

Plug this incidence estimate back to (2.12) and use (2.2) to have

(2.16)
$$\begin{aligned} \|\sum_{T\in\mathbb{T}} e^{it\sqrt{-\Delta}} f_T\|_{L^4(X)}^4 &\lesssim R^2 (R^{-1}(\#\mathbb{T})^{1/2} \rho^{1/2} (l\sigma)^{-1/2}) \|f\|_2^2 \\ &\lesssim R^2 (\rho^{1/2} (l\sigma)^{-1/2}) \|f\|_2^2 \mathcal{W}(f, B_R^4)^2. \end{aligned}$$

This is the L^4 estimate with the improved incidence result. Next, we will prove a refined L^2 estimate.

12

Recall that for each unit ball $B \subset X$, $|\sum_{T \in \mathbb{T}} e^{it\sqrt{-\Delta}} f_T | \mathbf{1}_B$ is broad. Since the planks intersecting B are essentially from a 3-dimensional family of directions, there are only two separate cases to consider:

- (1) For all $B \subset X$, $|\sum_{T \in \mathbb{T}} e^{it\sqrt{-\Delta}} f_T | \mathbf{1}_B$ is not "4-broad".
- (2) For all $B \subset X$, $|\sum_{T \in \mathbb{T}} e^{it\sqrt{-\Delta}} f_T| \mathbf{1}_B$ is "4-broad"

If the first case happens, then via decoupling and induction on scales in a somewhat

standard manner, we can conclude (2.10) when $p = \frac{10}{3}$. Now let us assume that for all $B \subset X$, $|\sum_{T \in \mathbb{T}} e^{it\sqrt{-\Delta}} f_T| \mathbf{1}_B$ is "4-broad". That is, there exist 4 quantitatively linearly independent directional caps $\{\tau_j\}_{j=1,\ldots,4}$ such that for all $j = 1, \ldots, 4$,

$$|\sum_{T\in\mathbb{T}}e^{it\sqrt{-\Delta}}f_T|\mathbf{1}_B\lesssim|\sum_{T\in\mathbb{T}_{\tau_j}}e^{it\sqrt{-\Delta}}f_T|\mathbf{1}_B.$$

Here $\mathbb{T}_{\tau_j} := \{T \in \mathbb{T} : \text{ the direction of } T \text{ is contained in } \tau_j \}.$

For each $R^{1/2}$ -ball $Q \subset \mathcal{Q}$, recall that $X \cap Q$ has a cover $\mathbb{S}(Q)$ by $1 \times R^{1/2} \times \mathbb{S}(Q)$ $R^{1/2} \times R^{1/2}$ -slabs and $\#\mathbb{S}(Q) \sim \sigma/\rho$. We claim that for all $S \in \mathbb{S}(Q)$

(2.17)
$$\|\sum_{T\in\mathbb{T}}e^{it\sqrt{-\Delta}}f_T\|_{L^2(S\cap X)}^2 \lesssim \sum_{T\in\mathbb{T}(Q)}\|f\|_2^2.$$

(2.17) is analogous to equation (2.8) in [15]. The proof of (2.17) involves several steps of pigeonholing. Intuitively, it is based on the following observation: Since $\sum_{T \in \mathbb{T}} e^{it\sqrt{-\Delta}} f_T |\mathbf{1}_B|$ is "4-broad" for each unit ball $B \subset S \cap X$, we can choose a directional cap τ_j such that τ_j is quantitatively transverse to the slab S. Let us assume that the cap τ_j is uniform for all $B \subset S \cap X$. Therefore, since the planks in \mathbb{T}_{τ_i} are all quantitatively transverse to S (see Figure 2), by L²-orthogonality, we have

$$\begin{split} &\|\sum_{T\in\mathbb{T}}e^{it\sqrt{-\Delta}}f_T\|_{L^2(S\cap X)}^2 \lesssim \|\sum_{T\in\mathbb{T}(Q)}e^{it\sqrt{-\Delta}}f_T\|_{L^2(S\cap X)}^2 \\ &\lesssim \|\sum_{T\in\mathbb{T}_{\tau_j}\cap\mathbb{T}(Q)}e^{it\sqrt{-\Delta}}f_T\|_{L^2(S\cap X)}^2 \lesssim \|\sum_{T\in\mathbb{T}_{\tau_j}\cap\mathbb{T}(Q)}e^{it\sqrt{-\Delta}}f_T\|_{L^2(S)}^2 \\ &\lesssim \sum_{T\in\mathbb{T}_{\tau_j}\cap\mathbb{T}(Q)}\|f_T\|_2^2 \lesssim \sum_{T\in\mathbb{T}(Q)}\|f_T\|_2^2. \end{split}$$

This proves (2.17).

Apply (2.17) to all $S \in \mathbb{S}(Q)$ and all $Q \in Q$. Since $\#\mathbb{S}(Q) \sim \sigma/\rho$ and since each $T \in \mathbb{T}$ intersects $\leq l$ many $R^{1/2}$ -balls in \mathcal{Q} , we have

$$\|e^{it\sqrt{-\Delta}}f\|_{L^{2}(X)}^{2} \lesssim \sum_{Q \in \mathcal{Q}} \sum_{S \in \mathbb{S}(Q)} \|e^{it\sqrt{-\Delta}}f\|_{L^{2}(S \cap X)}^{2} \lesssim l(\sigma/\rho) \sum_{T \in \mathbb{T}} \|f_{T}\|_{2}^{2},$$

which implies the L^2 estimate

(2.18)
$$\|e^{it\sqrt{-\Delta}}f\|_{L^2(X)}^2 \lesssim l(\sigma/\rho)\|f\|_2^2$$

Interpolate (2.12) and (2.18) to obtain (2.10) for $p = \frac{10}{3}$.




FIGURE 2.

3. Preliminaries

In this section, we set up some notations and introduce basic properties for Fourier integral operators. We denote the variables as $z = (x,t) = (x', x_n, t) \in \mathbb{R}^{n-1} \times \mathbb{R} \times \mathbb{R}$ and $\xi = (\xi', \xi_n) \in \mathbb{R}^{n-1} \times \mathbb{R}$ in the rest of the paper.

3.1. Reduction to a local estimate. Recall the Fourier integral operator defined in (1.5) and the associated assumptions (H1) and (H2). By considering the new phase function $\tilde{\phi}(z;\xi) = \phi(z;\xi) - \phi(0;\xi)$, we can assume that for all α ,

(3.1)
$$\partial_{\xi}^{\alpha}\phi(0;\xi) \equiv 0.$$

Let $\lambda \geq 1$. Given a Fourier integral operator \mathcal{F} defined in (1.5), let

(3.2)
$$\mathcal{F}^{\lambda}f(x,t) := \int_{\mathbb{R}^n} e^{i\phi^{\lambda}(x,t;\xi)} a^{\lambda}(x,t;\xi) \widehat{f}(\xi) \mathrm{d}\xi$$

where ϕ^{λ} and a^{λ} are defined as

$$\phi^{\lambda}(x,t;\xi) := \lambda \phi(x/\lambda,t/\lambda;\xi), \quad a^{\lambda}(x,t;\xi) := a(x/\lambda,t/\lambda;\xi).$$

By a standard Littlewood-Paley decomposition (see, for example, [8, Section 1.3]) and by a smooth partition on the unit sphere S^{n-1} , we can assume the amplitude function a obeys that $\operatorname{supp}_{\xi} a \subset \mathbb{A}^n(1)^\circ$, the interior of $\mathbb{A}^n(1)$, where

$$\mathbb{A}^{n}(1) := \{ (\xi', \xi_n) \in \mathbb{R}^{n} : 1/2 \le \xi_n \le 2, |\xi'| \le \xi_n \}.$$

Therefore, Theorem 1.7 reduces to the following result:

Proposition 3.1. Let \mathcal{F} be a Fourier integral operator given by (1.5). Suppose the phase function ϕ satisfies (3.1), and the amplitude function a obeys $\operatorname{supp}_{\xi} a \subset \mathbb{A}^n(1)^\circ$. Let \mathcal{F}^{λ} be given by (3.2). Then for all $\varepsilon > 0$ and all p > p(n),

(3.3)
$$\|\mathcal{F}^{\lambda}f\|_{L^{p}(B^{n+1}_{\lambda})} \leq C_{\varepsilon,p}\lambda^{(n-1)(\frac{1}{2}-\frac{1}{p})+\varepsilon}\|f\|_{L^{p}(\mathbb{R}^{n})}.$$

3.2. Quantitative conditions. We need to strengthen Proposition 3.1 to obtain uniform estimates for a class of Fourier integral operators (FIOs). This is necessary because we aim to use induction on scales, but an individual operator \mathcal{F} is not invariant under rescaling. Fortunately, rescaling preserves certain properties of the FIOs, enabling us to perform induction for a broader class of operators simultaneously. We will adopt the framework established in [1, Section 2.3].

Let $c_{\text{par}} > 0$ be a small fixed constant. For $\mathbf{A} = (A_1, A_2, A_3) \in [1, \infty)^3$, consider the following conditions on the phase function:

- $\begin{array}{l} (\mathrm{H1}_{\mathbf{A}}) \ |\partial_{\xi x}^2 \phi(z;\xi) I_n| \leq c_{\mathrm{par}} A_1 \text{ for all } (z;\xi) \in B_1^{n+1} \times \mathbb{A}^n(1). \\ (\mathrm{H2}_{\mathbf{A}}) \ |\partial_{\xi' \xi'}^2 \partial_t \phi(z;\xi) \frac{1}{\xi_n} I_{n-1}| \leq c_{\mathrm{par}} A_2 \text{ for all } (z;\xi) \in B_1^{n+1} \times \mathbb{A}^n(1). \end{array}$
- $(\mathrm{D1}_{\mathbf{A}}) \ \|\partial_{\xi}^{\beta}\partial_{x_{k}}\phi\|_{L^{\infty}(B_{1}^{n+1}\times\mathbb{A}^{n}(1))} \leq c_{\mathrm{par}}A_{1} \text{ for all } 1 \leq k \leq n \text{ and } \beta \in \mathbb{N}_{0}^{n} \text{ with } \mathbb{A}^{n}(\mathbb{A}^{n})$ $2 \leq |\beta| \leq 3$ satisfying $|\beta'| \geq 2$;

 $\|\partial_{\xi'}^{\beta'}\partial_{x_k}\phi\|_{L^{\infty}(B_1^{n+1}\times\mathbb{A}^n(1))} \leq \frac{c_{\text{par}}}{2n}A_1 \text{ for all } \beta' \in \mathbb{N}_0^{n-1} \text{ with } |\beta'| = 3.$ (D2_A) For some large integer $N = N_{\varepsilon,M,p} \in \mathbb{N}$ depending only on the dimension n and the fixed choice of ε , M and p, one has

$$\|\partial_{\xi}^{\beta}\partial_{z}^{\alpha}\phi\|_{L^{\infty}(B_{1}^{n+1}\times\mathbb{A}^{n}(1))}\leq\frac{c_{\mathrm{par}}}{2n}A_{3}$$

for all $(\alpha, \beta) \in \mathbb{N}_0^{n+1} \times \mathbb{N}_0^n$ with $2 \leq |\alpha| \leq 4N$ and $1 \leq |\beta| \leq 4N + 2$ satisfying $1 \leq |\beta| \leq 4N$ or $|\beta'| \geq 2$.

Finally, it is useful to assume a margin condition on the spatial support of the amplitude a:

(M_A) dist(supp_z a, $\mathbb{R}^{n+1} \setminus B_1^{n+1}$) $\geq \frac{1}{4A_2}$.

Datum (ϕ, a) satisfying $(H1_A), (H2_A), (D1_A), (D2_A)$ and (M_A) (in addition to (H1) and (H2) is said to be of type A. By cutting the amplitude function into pieces and various rescaling arguments, it is possible to reduce to the case where $\mathbf{A} = \mathbf{1} := (1, 1, 1)$. We refer to [1, Section 2.5] for more details.

Throughout the paper, we will assume the FIOs are of type 1. Proposition 3.1 can be strengthened as

Proposition 3.2. Let \mathcal{F} be a Fourier integral operator of type 1 given by (1.5). Suppose the phase function ϕ satisfies (3.1), and the amplitude function a obeys $\operatorname{supp}_{\varepsilon} a \subset \mathbb{A}^n(1)^{\circ}$. Let \mathcal{F}^{λ} be given by (3.2). Then for all $\varepsilon > 0$ and all p > p(n), there exists a constant $C_{\varepsilon,p}$ independent to \mathcal{F} , such that

(3.4)
$$\|\mathcal{F}^{\lambda}f\|_{L^{p}(B^{n+1}_{\lambda})} \leq C_{\varepsilon,p}\lambda^{(n-1)(\frac{1}{2}-\frac{1}{p})+\varepsilon}\|f\|_{L^{p}(\mathbb{R}^{n})}.$$

Here, $C_{\varepsilon,p} > 0$ is uniform for all \mathcal{F} of type 1.

3.3. Wave packet decomposition. Fix $0 < \varepsilon < 1/100$. We will perform the wave packet decomposition for a function f with $\operatorname{supp} \widehat{f} \subset \mathbb{A}^n(1)$ at a scale $R \in [1, \lambda^{1-\varepsilon}]$.

By condition (H1), we see that $\partial_{\xi} \phi(\cdot, t; \xi)$ is a local diffeomorphism. Indeed, for fixed t, ξ , the map $x \mapsto \partial_{\xi} \phi(x, t; \xi)$ has non-vanishing Jacobian. By rescaling, we may assume $\partial_{\xi} \phi(\cdot, t; \xi)$ is a diffeomorphism on the domain B_1^n for any $(t, \xi) \in$ $[-1,1] \times \mathbb{A}^n(1)$. Therefore, it has an inverse function which we defined as follows.

Definition 3.3. For $(t,\xi) \in [-1,1] \times \mathbb{A}^n(1)$, let $\Phi(u,t;\xi)$ be the unique solution to

(3.5)
$$\partial_{\xi}\phi(\Phi(u,t;\xi),t;\xi) = u$$

Here, we may assume the equation holds for u in a small neighborhood of the origin.

Since ϕ is 1-homogeneous in ξ , we have $\partial_{\xi}\phi(z;\xi) = \partial_{\xi}\phi(z;s\xi)$. Hence,

(3.6)
$$\Phi(u,t;\xi) = \Phi(u,t;s\xi)$$

By the inverse function theorem and conditions $(D1_{\mathbf{A}}), (D2_{\mathbf{A}})$, we obtain favorable upper bounds on the L^{∞} norm of Φ and its derivatives. Taking the derivative in u in (3.5), we get $\partial_{\varepsilon_x}^2 \phi \cdot \partial_u \Phi = I_n$. Thus,

$$\det \partial_u \Phi(u, t; \xi) \gtrsim 1.$$

In other words, $\partial_u \Phi(u,t;\xi)$ is a quantitatively non-singular matrix.

We define

$$\gamma^{\lambda}(u,t;\xi) := \lambda \Phi(u/\lambda,t/\lambda;\xi).$$

Then γ^{λ} satisfies

(3.7)
$$\partial_{\xi}\phi^{\lambda}(\gamma^{\lambda}(u,t;\xi),t;\xi) = u.$$

For fixed $(u;\xi)$, $t \mapsto (\gamma^{\lambda}(u,t;\xi),t)$ is a curve, with t ranging over [-R,R]. Our wave packets will be certain non-isotropic neighborhoods of such curves.

Now we present the wave packet decomposition for $\mathcal{F}^{\lambda}f$ in B_R^{n+1} . In fact, we will perform the wave packet decomposition for $f: f = \sum_T f_T$ first. Then, by applying the linear operator \mathcal{F}^{λ} , we get $\mathcal{F}^{\lambda}f = \sum_T \mathcal{F}^{\lambda}f_T$.

We begin by decomposing in frequency space. Fix a maximally $R^{-1/2}$ -separated subset of $B_1^{n-1} \times \{1\}$. For each ξ_{θ} belonging to this subset, define

(3.8)
$$\theta = \{ (\xi', \xi_n) \in \mathbb{A}^n(1) : |\xi'/\xi_n - \xi_\theta| \le R^{-1/2} \}.$$

Then θ is roughly a $1 \times R^{-1/2} \times \cdots \times R^{-1/2}$ -box, and ξ_{θ} is the center of θ . We call such θ an $R^{-1/2}$ -**cap** and denote the collection of these θ by $\Theta_{R^{-1/2}}$. Note that $\Theta_{R^{-1/2}}$ form a finitely overlapping covering of $\mathbb{A}^n(1)$. Choose $\{\psi_{\theta}(\xi)\}_{\theta \in \Theta_{R^{-1/2}}}$ to be a smooth partition of unity adapted to $\Theta_{R^{-1/2}}$, so that $\sum_{\theta \in \Theta_{R^{-1/2}}} \psi_{\theta}(\xi) = 1$ for $\xi \in \mathbb{A}^n(1)$. Since $\operatorname{supp} \widehat{f} \subset \mathbb{A}^n(1)$, we have the frequency decomposition

$$\widehat{f} = \sum_{\theta \in \Theta_{R^{-1/2}}} \psi_{\theta} \widehat{f}$$

We denote $f_{\theta} := (\psi_{\theta} \widehat{f})^{\vee}$, and hence

$$f = \sum_{\theta \in \Theta_{R^{-1/2}}} f_{\theta}.$$

Before performing decomposition in the physical space, we introduce a notation.

Definition 3.4. Let $U \subset \mathbb{R}^n$ be a box of dimensions $a_1 \times \cdots \times a_n$. Define U^* to be the **dual box** of U, which is a box centered at the origin of dimensions $a_1^{-1} \times \cdots \times a_n^{-1}$, and with edges parallel to the corresponding edges of U.

Now for a fixed $\theta \in \Theta_{R^{-1/2}}$, we cover B_R^n by $1 \times R^{1/2} \times \cdots \times R^{1/2}$ -slabs that are parallel to θ^* . Denote this collection of slabs by

(3.9)
$$\mathbb{T}^{\flat}_{\theta} = \{T^{\flat}\}.$$

Choose a set of non-negative smooth functions $\{\eta_{T^{\flat}}\}$ adapted to $\mathbb{T}^{\flat}_{\theta}$, so that $\eta_{T^{\flat}}$ decays rapidly outside T^{\flat} , $\sum \eta_{T^{\flat}} = 1$ in B^{n}_{R} and $\operatorname{supp} \widehat{\eta}_{T^{\flat}} \subset \theta$. Thus, inside B^{n}_{R} , we have

$$\widehat{f} = \sum_{\theta \in \Theta_{R^{-1/2}}} \sum_{T^{\flat} \in \mathbb{T}_{\theta}^{\flat}} \widehat{\eta}_{T^{\flat}} * (\psi_{\theta} \widehat{f}) + \operatorname{RapDec}(R) \|f\|_{2}$$




FIGURE 3. Wave packet

Let $\delta = \varepsilon^{1000}$, which is a small number to handle rapidly decaying tails. For each T^{\flat} , define

$$T := \{ (\gamma^{\lambda}(u, t; \xi_{\theta}), t) : u \in R^{\delta} T^{\flat}, |t| \le R \}.$$

Here, $R^{\delta}T^{\flat}$ is a non-isotropic dilate of T^{\flat} of dimensions $R^{2\delta} \times R^{1/2+\delta} \times \cdots \times R^{1/2+\delta}$. Each T is a "curved plank" (see Figure 3). We denote $\mathbb{T}_{\theta} := \{T : T^{\flat} \in \mathbb{T}_{\theta}^{\flat}\}$ and

$$f_T := \eta_{T^\flat}(\psi_\theta^{\lor} * f)$$

This gives the wave packet decomposition in B_R^n

$$f = \sum_{\theta \in \Theta_{R^{-1/2}}} \sum_{T \in \mathbb{T}_{\theta}} f_T + \operatorname{RapDec}(R) \|f\|_2.$$

Therefore, in B_R^{n+1} , we obtain

$$\mathcal{F}^{\lambda}f = \sum_{\theta \in \Theta_{R^{-1/2}}} \sum_{T \in \mathbb{T}_{\theta}} \mathcal{F}^{\lambda}f_{T} + \operatorname{RapDec}(R) \|f\|_{2}.$$

Definition 3.5 (Curved plank). Given $\xi_0 \in \mathbb{A}^n(1)$ and a box $V \subset B_R^n$, define

(3.10)
$$\Gamma_V(\xi_0; R) := \{ (\gamma^\lambda(u, t; \xi_0), t) : u \in V, |t| \le R \}$$

 $\Gamma_V(\xi_0; R)$ is called a curved plank with base V and direction ξ_0 .

We primarily focus on the case where V is a slab with dimensions $Rs^2 \times Rs \times \cdots \times Rs$ for $s \in [R^{-1/2}, 1]$. Next, we define the non-isotropic dilation for such slabs.

Definition 3.6. For a slab $U \subset \mathbb{R}^n$ of dimensions $Rs^2 \times Rs \times \cdots \times Rs$ with $s \in [R^{-1/2}, 1]$ and a constant C > 0, define CU, the non-isotropic C-dilation of U, to be a slab of dimensions $C^2Rs^2 \times CRs \times \cdots \times CRs$ with the same center as U.

For two slabs $U_1, U_2 \subset \mathbb{R}^n$, we say they are **comparable** if

$$\frac{1}{C}U_1 \subset U_2 \subset CU_1.$$

Here, $C = C_{\phi}$ is a constant that may vary from line to line, but eventually only depends on ϕ .

Remark 3.7. We will also need the isotropic dilation. Namely, for a $a_1 \times \cdots \times a_n$ box U, the isotropic C-dilation of U will be the box of dimensions $Ca_1 \times \cdots \times Ca_n$ with the same center as U. We will always specify when doing isotropic dilation; otherwise, the notation CU always refer to the non-isotropic dilation.

We also define the dilation and comparability for curved planks. Since a curved plank is not convex, we will define these notions in terms of its base.

Definition 3.8. For a curved plank $\Gamma_V(\xi_0; R)$ and a constant C > 0, we define the (non-isotropic) C-dilation of $\Gamma_V(\xi_0; R)$ to be

 $C\Gamma_V(\xi_0; R) := \{ (\gamma^{\lambda}(u, t; \xi_0), t) : u \in CV, |t| \le R \}.$

For two curved planks $\Gamma_{V_1}(\xi_1; R), \Gamma_{V_2}(\xi_2; R)$, we say they are **comparable** if

$$\frac{1}{C}\Gamma_{V_1}(\xi_1; R) \subset \Gamma_{V_2}(\xi_2; R) \subset C\Gamma_{V_1}(\xi_1; R)$$

for some constant C only depending on ϕ . If $\Gamma_{V_1}(\xi_1; R), \Gamma_{V_2}(\xi_2; R)$ are not comparable, we say they are **distinct**. We may identify curved planks that are comparable.

The next lemma concerns the geometry of curved planks.

Lemma 3.9. Let $s \in [R^{-1/2}, 1]$. Let $V \subset B_R^n$ be a slab of dimensions $Rs^2 \times Rs \times$ $\cdots \times Rs$. Then for $|t_0| \leq R$, each t_0 -slice $\Gamma_V(\xi_0; R) \cap \{t = t_0\}$ is comparable to an $Rs^2 \times Rs \times \cdots \times Rs$ -slab.

Proof. Suppose V is centered at u_0 . We express each point in V as $u_0 + u$, where u ranges over $V - u_0$, an $Rs^2 \times Rs \times \cdots \times Rs$ -box centered at the origin.

For fixed (t_0, ξ_0) , by applying Taylor's expansion in the *u* variable, we have (3.11)

 $\gamma^{\lambda}(u_{0} + u, t_{0}; \xi_{0}) = \gamma^{\lambda}(u_{0}, t_{0}; \xi_{0}) + \partial_{u}\gamma^{\lambda}(u_{0}, t_{0}; \xi_{0}) \cdot u + O(\|\partial_{uu}^{2}\gamma^{\lambda}(\cdot, t_{0}; \xi_{0})\|_{\infty})|u|^{2}.$ Since $\partial_{uu}^2 \gamma^{\lambda}(u_0, t_0; \xi_0) = \lambda^{-1} \partial_{uu}^2 \Phi(u_0/\lambda, t_0/\lambda; \xi_0)$, whose L^{∞} norm is $O(\lambda^{-1})$,

 $O(\|\partial_{uu}^2 \gamma^{\lambda}(\cdot, t_0; \xi_0)\|_{\infty})|u|^2 \lesssim \lambda^{-1} (Rs)^2 \le Rs^2.$

Therefore, the sets $\Gamma_V(\xi_0; R) \cap \{t = t_0\}$ and

 $\gamma^{\lambda}(u_0, t_0; \xi_0) + \{\partial_u \gamma^{\lambda}(u_0, t_0; \xi_0) \cdot x : x \in V - u_0\}$

are within a distance $\leq Rs^2$ from each other. Since $\partial_u \gamma^\lambda(u_0, t_0; \xi_0)$ is quantitative non-singular, the latter is comparable to $V - u_0$ which is a $Rs^2 \times Rs \times \cdots \times Rs$ -slab. Therefore, $\Gamma_V(\xi_0; R) \cap \{t = t_0\}$ is also comparable to such a slab.

Remark 3.10. From the above lemma, we see that in Figure 3, each horizontal slice of T is a slab of dimensions $R^{2\delta} \times R^{1/2+\delta} \times \cdots \times R^{1/2+\delta}$. Though, they may not be parallel.

The next lemma concerns the perturbation of a curved plank resulting from choosing different directions. It was essentially proved in [8].

Lemma 3.11. Let $s \in [R^{-1/2}, 1]$, and let $\tau \in \Theta_s$. Define $V_{\tau,R} := (Rs^2)\tau^*$ to be an isotropic dilate of τ^* , which is an $Rs^2 \times Rs \times \cdots \times Rs$ -box. Let $V \subset B_R^n$ be a translated copy of $V_{\tau,R}$. Then for any $\xi_1, \xi_2 \in \tau$, the two curved planks $\Gamma_V(\xi_1; R)$ and $\Gamma_V(\xi_2; R)$ are comparable. In other words, $\Gamma_V(\xi_1; R) \subset \Gamma_{CV}(\xi_2; R)$ for some constant C > 0 and vice versa.

Proof. It suffices to show that there exists a large constant C > 0 such that

(3.12)
$$\{\gamma^{\lambda}(u,t;\xi_2): u \in V\} \subset \{\gamma^{\lambda}(u,t;\xi_1): u \in CV\}.$$

Let $u^{\lambda}(z;\xi) := \partial_{\xi} \phi^{\lambda}(z;\xi)$, so $u^{\lambda}(\gamma^{\lambda}(u,t;\xi),t;\xi) = u$ (recall (3.7)). To obtain (3.12), we only need to show that for any $u \in V$,

$$u^{\lambda}(\gamma^{\lambda}(u,t;\xi_2),t;\xi_1) \subset CV.$$

Since $u^{\lambda}(\gamma^{\lambda}(u,t;\xi_2),t;\xi_2) = u \in V$, it suffices to show

(3.13)
$$u^{\lambda}(z;\xi_1) - u^{\lambda}(z;\xi_2) \subset 4V_{\tau,R}.$$

for all $z \in B_R^{n+1}$ and all $\xi_1, \xi_2 \in \tau$. This is true by Lemma 4.3 in [8].

Using the notation of curved plank, each wave packet $T \in \mathbb{T}_{\theta}$ can be written as

$$T = \Gamma_{R^{\delta}T^{\flat}}(\xi_{\theta}; R),$$

a curved plank with base $R^{\delta}T^{\flat}$ and direction ξ_{θ} . We call each such T an R-plank. Note that each t-slice $T \cap \{t = t_0\}$ of T is morally an $R^{2\delta} \times R^{1/2+\delta} \times \cdots \times R^{1/2+\delta}$ -slab. Moreover, $\Gamma_{R^{\delta}T^{\flat}}(\xi_{\theta}; R)$ and $\Gamma_{R^{\delta}T^{\flat}}(\xi; R)$ are comparable for all $\xi \in \theta$.

Lemma 3.12. $(\mathcal{F}^{\lambda}f_T)\mathbf{1}_{B_{p}^{n+1}}$ is essentially supported in T. In other words,

$$(\mathcal{F}^{\lambda}f_T)\mathbf{1}_{B_R^{n+1}\setminus T} = \operatorname{RapDec}(R)||f||_2$$

Proof. Note that

$$\mathcal{F}^{\lambda}f_{T}(x,t) = \int_{\mathbb{R}^{n}} e^{i\phi^{\lambda}(x,t;\xi)} a^{\lambda}(x,t;\xi)\widehat{\eta}_{T^{\flat}} * (\psi_{\theta}\widehat{f}) \mathrm{d}\xi.$$

Since a^{λ} is a smoothing symbol and \hat{f}_T is supported in 2θ , we have

$$\mathcal{F}^{\lambda}f_{T}(x,t) = \int_{\mathbb{R}^{n}} \left(e^{i\phi^{\lambda}(x,t;\xi)} a^{\lambda} \widetilde{\psi}_{\theta} \right) \widehat{\eta}_{T^{\flat}} * (\psi_{\theta}\widehat{f}) \mathrm{d}\xi + \mathrm{RapDec}(R) \|f\|_{2}$$

where $\widetilde{\psi}_{\theta}$ is a bump function on 2θ . Let $G_{x,t}(\xi)$ be so that $\overline{G_{x,t}(\xi)} = e^{i\phi^{\lambda}(x,t;\xi)}a^{\lambda}\widetilde{\psi}_{\theta}$. By Plancherel,

(3.14)
$$\mathcal{F}^{\lambda}f_{T}(x,t) = \int_{\mathbb{R}^{n}} \overline{\check{G}_{x,t}(y)} \Big(\eta_{T^{\flat}}(\psi_{\theta}^{\vee} * f) \Big)(y) \mathrm{d}y + \mathrm{RapDec}(R) \|f\|_{2}.$$

Let us compute

$$\overline{\check{G}_{x,t}(y)} = \int_{\mathbb{R}^n} e^{i(\langle y,\xi\rangle - \phi^\lambda(x,t;\xi))} \overline{a^\lambda \widetilde{\psi}_\theta} \mathrm{d}\xi.$$

Via the change of variable $\xi = \xi_{\theta} + \eta$, we have

$$\overline{\check{G}_{x,t}(y)} = \int_{\mathbb{R}^n} e^{i(\langle y,\xi_\theta + \eta \rangle - \phi^\lambda(x,t;\xi_\theta + \eta))} \overline{a^\lambda(z;\xi_\theta + \eta)} \widetilde{\psi}_\theta(\xi_\theta + \eta) \mathrm{d}\xi.$$

By the method of non-stationary phase (see [10, Lemma 5.4] for a similar argument), $|\check{G}_{x,t}(y)| = \operatorname{RapDec}(R)$ if

$$\partial_{\xi} \phi^{\lambda}(x,t;\xi_{\theta}+\eta) - y \notin \theta^*.$$

Recall (3.13), which implies that $\partial_{\xi}\phi^{\lambda}(x,t;\xi_{\theta}+\eta) - \partial_{\xi}\phi^{\lambda}(x,t;\xi_{\theta}) \in 4\theta^{*}$. Therefore, $|\check{G}_{x,t}(y)| = \operatorname{RapDec}(R)$ if

(3.15)
$$\partial_{\xi}\phi^{\lambda}(x,t;\xi_{\theta}) - y \notin 5\theta^*$$

Now we are ready to prove the lemma. If $(x_0, t_0) \in B_R^{n+1} \setminus T$, then

$$[x_0,t_0) \notin (T \cap \{t=t_0\}) = \{\gamma^{\lambda}(u,t_0;\xi_{\theta}) : u \in R^{\delta}T^{\flat}\}.$$

By applying $\partial_{\xi}\phi^{\lambda}(\cdot, \cdot; \xi_{\theta})$ to both sides (recall (3.7)), we see that the above is a consequence of

(3.16)
$$\partial_{\xi}\phi^{\lambda}(x_0, t_0; \xi_{\theta}) \notin R^{\delta}T^{\flat}$$

We use (3.14) to show $\mathcal{F}^{\lambda}f_T(x_0, t_0) = \operatorname{RapDec}(R) ||f||_2$: Since $\eta_{T^{\flat}}$ decreases rapidly outside $R^{\delta/2}T^{\flat}$, it suffices to show $|\check{G}_{x_0,t_0}(y)| = \operatorname{RapDec}(R)$ for $y \in R^{\delta/2}T^{\flat}$. This boils down to verifying (3.15), or equivalently, $\partial_{\xi}\phi^{\lambda}(x_0, t_0; \xi_{\theta}) \notin y + 5\theta^* \subset 5R^{\delta/2}T^{\flat}$, since T^{\flat} is a translated copy of θ^* and since $y \in T^{\flat}$. However, this is given by (3.16).

3.4. Comparing wave packets at different scales. Next, we discuss the wave packet decomposition for $\mathcal{F}^{\lambda}f$ inside a ball not necessarily centered at the origin. Fix $z_0 = (x_0, t_0) \in B_{\lambda}^{n+1}$ and consider the ball $B_R^{n+1}(z_0)$. Define \tilde{f} so that

$$(\widetilde{f})^{\wedge}(\xi) := e^{i\phi^{\lambda}(z_0;\xi)}\widehat{f}(\xi).$$

As a result,

$$\mathcal{F}^{\lambda}f(z) = \widetilde{\mathcal{F}}^{\lambda}\widetilde{f}(\widetilde{z}) \text{ for } \widetilde{z} = z - z_0,$$

where $\widetilde{\mathcal{F}}^{\lambda}$ is the Fourier integral operator whose phase $\widetilde{\phi}^{\lambda}$ and amplitude \widetilde{a}^{λ} are given by

$$\widetilde{\phi}(z;\xi) := \phi(z + \frac{z_0}{\lambda};\xi) - \phi(\frac{z_0}{\lambda};\xi) \text{ and } \widetilde{a}(z;\xi) := a(z + \frac{z_0}{\lambda};\xi)$$

If $z \in B_R^{n+1}(z_0)$, then $\tilde{z} \in B_R^{n+1}$, and we can therefore apply the wave packet decomposition for $\tilde{\mathcal{F}}^{\lambda}\tilde{f}$ inside B_R^{n+1} :

$$\widetilde{\mathcal{F}}^{\lambda}\widetilde{f} = \sum_{\widetilde{T}} \widetilde{\mathcal{F}}^{\lambda}\widetilde{f}_{\widetilde{T}} + \operatorname{RapDec}(R) \|f\|_{2}.$$

Each \widetilde{T} is given by the collection of $\widetilde{z} \in B_R^{n+1}$ satisfying

$$\partial_{\xi}\phi(\frac{\widetilde{z}}{\lambda}+\frac{z_0}{\lambda};\xi_{\theta})=\frac{v}{\lambda}+\partial_{\xi}\phi(\frac{z_0}{\lambda};\xi_{\theta}), \text{ for some } v\in R^{\delta}T^{\flat}.$$

Recall (3.9) that $\{T^{\flat}\}$ are translated copies of θ^* that cover B_R^n .

For a given z_0 , define

(3.17)
$$T(z_0) := \left\{ \left(\gamma^{\lambda} (v + \partial_{\xi} \phi^{\lambda}(z_0; \xi_{\theta}), t; \xi_{\theta}), t \right) : v \in R^{\delta} T^{\flat}, |t - t_0| \le R \right\}.$$

20

Remark 3.13. If we replace ξ_{θ} by any $\xi \in \theta$ in the definition of $T(z_0)$ in (3.17), then the resulting curved plank is comparable to the original curved plank defined using ξ_{θ} . This is a consequence of Lemma 3.11. We will use this fact later when comparing wave packets in different scales.

Thus, the function

$$\widetilde{\mathcal{F}}^{\lambda}\widetilde{f}_{\widetilde{T}}(z-z_0)$$

is essentially supported in $T(z_0)$ when restricted to $B_R^{n+1}(z_0)$. We denote the collection of planks $\{T(z_0): T^{\flat} \in \mathbb{T}_{\theta}^{\flat}\}$ as

(3.18)
$$\mathbb{T}_{\theta}(z_0) := \{ T(z_0) : T^{\flat} \in \mathbb{T}_{\theta}^{\flat} \}.$$

For each $T \in \bigcup_{\theta} \mathbb{T}_{\theta}(z_0)$, we also define

(3.19)
$$\widehat{f}_T(\xi) := e^{-i\phi^{\lambda}(z_0;\xi)} (\widetilde{f}_{\widetilde{T}})^{\wedge}(\xi),$$

where $T^{\flat} = \widetilde{T}^{\flat}$. Under this notation,

$$\mathcal{F}^{\lambda}f = \sum_{T \in \bigcup_{\theta} \mathbb{T}_{\theta}(z_0)} \mathcal{F}^{\lambda}f_T + \operatorname{RapDec}(R) \|f\|_2$$

in $B_R^{n+1}(z_0)$. For each $T \in \mathbb{T}_{\theta}(z_0)$, we remark that $(\mathcal{F}^{\lambda}f_T)\mathbf{1}_{B_R^{n+1}(z_0)}$ is essentially supported in T^{\flat} , and \hat{f}_T is supported in 2θ .

We have the following lemma regarding the comparison of wave packets at different scales. Its counterpart can be found in [10, Lemma 9.1].

Lemma 3.14. Let $r \in [1, R)$, and let $\theta \in \Theta_{R^{-1/2}}$, $\tilde{\theta} \in \Theta_{r^{-1/2}}$. Suppose $B_r^{n+1}(z_0) \subset B_R^{n+1}$, and suppose $T \in \mathbb{T}_{\theta}$, $\tilde{T} \in \mathbb{T}_{\tilde{\theta}}(z_0)$. If either $\theta \not\subset 4\tilde{\theta}$, or $100T \cap 100\tilde{T} = \emptyset$, then

$$(f_T)_{\widetilde{T}} = \operatorname{RapDec}(R) ||f||_2.$$

Proof. For simplicity, we denote $e^{i\phi^{\lambda}(z_0;D)}g := (e^{i\phi^{\lambda}(z_0;\cdot)}\widehat{g})^{\vee}$. Recall (3.19) for the definition of the wave packet decomposition inside $B_r^{n+1}(z_0)$, we have

$$e^{i\phi^{\lambda}(z_0;D)}\Big(f_T\Big)_{\widetilde{T}} = \eta_{\widetilde{T}^{\flat}}\Big(\psi_{\widetilde{\theta}}^{\vee} * (e^{i\phi^{\lambda}(z_0;D)}f_T)\Big).$$

Note that $\operatorname{supp}(e^{i\phi^{\lambda}(z_0;D)}f_T)^{\wedge} = \operatorname{supp}\widehat{f}_T$, which is essentially contained in 2θ . Since $\operatorname{supp}\psi_{\widetilde{\theta}} \subset \widetilde{\theta}$, if $\theta \not\subset 4\widetilde{\theta}$, we have $e^{i\phi^{\lambda}(z_0;D)} (f_T)_{\widetilde{T}} = \operatorname{RapDec}(R) ||f||_2$, which implies $(f_T)_{\widetilde{T}} = \operatorname{RapDec}(R) ||f||_2$.

Next, we assume $\theta \subset 4\widetilde{\theta}$ and $10T \cap \widetilde{T} = \emptyset$. Compute

$$e^{i\phi^{\lambda}(z_{0};D)}f_{T}(x) = \int_{\mathbb{R}^{n}} e^{i(\phi^{\lambda}(z_{0};\xi) + \langle x,\xi\rangle)}\widehat{\eta}_{T^{\flat}} * (\psi_{\theta}\widehat{f})d\xi$$
$$= \int_{\mathbb{R}^{n}} \left(e^{i(\phi^{\lambda}(z_{0};\xi) + \langle x,\xi\rangle)}\widetilde{\psi}_{\theta} \right) \widehat{\eta}_{T^{\flat}} * (\psi_{\theta}\widehat{f})d\xi + \operatorname{RapDec}(R) \|f\|_{2},$$

where $\widetilde{\psi}_{\theta}$ is a bump function on 2θ . Introduce $G_{z_0,x}(\xi)$ such that $\overline{G_{z_0,x}(\xi)} = e^{i(\phi^{\lambda}(z_0;\xi) + \langle x,\xi \rangle)}\widetilde{\psi}_{\theta}$. By Plancherel,

(3.20)
$$e^{i\phi^{\lambda}(z_0;D)}f_T(x) = \int_{\mathbb{R}^n} \overline{\check{G}_{z_0,x}(y)} \Big(\eta_{T^{\flat}}(\psi_{\theta}^{\vee} * f)\Big)(y) \mathrm{d}y + \operatorname{RapDec}(R) \|f\|_2.$$

Let us compute

$$\check{G}_{z_0,x}(y) = \int_{\mathbb{R}^n} e^{i(-\phi^{\lambda}(z_0;\xi) + \langle y - x,\xi \rangle)} \widetilde{\psi}_{\theta} \mathrm{d}\xi.$$

By the same argument as in the proof of Lemma 3.12, $|\check{G}_{z_0,x}(y)| = \operatorname{RapDec}(R)$ if

(3.21)
$$\partial_{\xi}\phi^{\lambda}(z_0;\xi_{\theta}) + x - y \notin 5\theta^*.$$

To show $\eta_{\widetilde{T}^{\flat}}\left(\psi_{\widetilde{\theta}}^{\vee} * (e^{i\phi^{\lambda}(z_0;D)}f_T)\right) = \operatorname{RapDec}(R) \|f\|_2$, we just need to show (3.20) = $\operatorname{RapDec}(R) \|f\|_2$ for $x \in r^{\delta/2} \widetilde{T}^{\flat}$. This boils down to show $|\check{G}_{z_0,x}(y)| = \operatorname{RapDec}(R)$ for $x \in r^{\delta/2} \widetilde{T}^{\flat}, y \in R^{\delta/2} T^{\flat}$. By (3.21), it suffices to show

(3.22)
$$\partial_{\xi}\phi^{\lambda}(z_0;\xi_{\theta}) + x - y \notin \theta^* \text{ for all } x \in r^{\delta/2}\widetilde{T}^{\flat}, y \in R^{\delta/2}T^{\flat}.$$

Since T^{\flat} is a translated copy of θ^* , (3.22) follows from

$$2R^{\delta/2}T^{\flat} \cap (\partial_{\xi}\phi^{\lambda}(z_0;\xi_{\theta}) + r^{\delta/2}\widetilde{T}^{\flat}) = \emptyset.$$

By applying the transformation $\partial_{\xi}(\cdot, t; \xi_{\theta})$ (recall (3.7)), the above is a consequence of the fact that 4T and

$$\{\left(\partial_{\xi}\phi^{\lambda}(v+\partial_{\xi}\phi^{\lambda}(z_{0};\xi_{\theta}),t;\xi_{\theta}),t\right):v\in r^{\delta/2}\widetilde{T}^{\flat}\}$$

are disjoint. However, by Remark 3.13, the latter one is contained in $100\tilde{T}$. Thus, if $4T \cap 100T = \emptyset$, then the aforementioned two planks are disjoint, yielding (3.22).

3.5. Broad norm. We introduce the broad norm in the variable coefficient setting. A similar setup can be found in [10, Section 1.5].

Let $K = R^{\varepsilon^{50}}$. Recall (3.8) that $\mathbb{A}^n(1)$ can be partitioned into caps $\Theta_{K^{-1}} = \{\tau\}$, where each τ has dimensions $1 \times K^{-1} \times \cdots \times K^{-1}$. In view of the rescaling ϕ^{λ} of the phase function, define the rescaled generalized Gauss map

$$G^{\lambda}(z;\xi) := G(z/\lambda;\xi) \text{ for } (z;\xi) \in \text{supp } a^{\lambda}.$$

For each $z \in B_{\lambda}^{n+1}$, there is a range of normal directions associated with the cap τ , as given by

$$G^{\lambda}(z;\tau) := \{ G^{\lambda}(z;\xi) : \xi \in \tau, (z;\xi) \in \text{supp } a^{\lambda} \}.$$

For any subspace $V \subset \mathbb{R}^{n+1}$, define $\measuredangle(G^{\lambda}(z;\tau), V)$ to be the smallest angle between non-zero vectors $v \in V$ and $v' \in G^{\lambda}(z; \tau)$. Fix $A \ge 1$ and $1 \le k \le n+1$. For a K^2 -ball $B_{K^2}^{n+1} \subset B_{\lambda}^{n+1}$ centered at z_0 , define

$$\mu_{\mathcal{F}^{\lambda}f}(B_{K^{2}}^{n+1}) := \min_{V_{1},...,V_{A}} \max_{\tau \notin V_{i}} \int_{B_{K^{2}}^{n+1}} |\mathcal{F}^{\lambda}f_{\tau}|^{p}$$

Here, V_1, \ldots, V_A are taken over all (k-1)-dimensional subspaces, and $\tau \notin V_i$ refers to those $\tau \in \Theta_{K^{-1}}$ such that $\measuredangle(G^{\lambda}(z_0;\tau), V_i) > K^{-2}$ for all $1 \le i \le A$. If $U \subset B_{\lambda}^{n+1}$ is a disjoint union of K^2 -balls, then we define the broad norm as

$$\|\mathcal{F}^{\lambda}f\|_{\mathrm{BL}^p_{k,A}(U)}^p := \sum_{\substack{B^{n+1}_{K^2} \subset U \\ K^2}} \mu_{\mathcal{F}^{\lambda}f}(B^{n+1}_{K^2}).$$

The broad norm enjoys the following properties. See [10] for their proofs.

Lemma 3.15. Suppose $1 \le p < \infty$, $A_1, A_2 \ge 1$ and $A = A_1 + A_2$, then $\|\mathcal{F}^{\lambda}(f_{1}+f_{2})\|_{\mathrm{BL}_{k-4}^{p}(U)} \lesssim \|\mathcal{F}^{\lambda}f_{1}\|_{\mathrm{BL}_{k-4}^{p}(U)} + \|\mathcal{F}^{\lambda}f_{2}\|_{\mathrm{BL}_{k-4}^{p}(U)}.$

22

Lemma 3.16. Suppose $1 \le p, p_1, p_2 < \infty$, $0 \le \alpha_1, \alpha_2 \le 1$ satisfy $\alpha_1 + \alpha_2 = 1$ and

$$\frac{1}{p} = \frac{\alpha_1}{p_1} + \frac{\alpha_2}{p_2}$$

Suppose also $A_1, A_2 \ge 1$ and $A = A_1 + A_2$. Then,

$$\|\mathcal{F}^{\lambda}f\|_{\mathrm{BL}^{p}_{k,A}(U)} \lesssim \|\mathcal{F}^{\lambda}f\|_{\mathrm{BL}^{p_{1}}_{k,A_{1}}(U)}^{\alpha_{1}}\|\mathcal{F}^{\lambda}f\|_{\mathrm{BL}^{p_{2}}_{k,A_{2}}(U)}^{\alpha_{2}}.$$

3.6. Some L^2 results. The first one is a local L^2 estimate for a sum of wave packets whose directions are transverse to the integration domain.

Lemma 3.17. Let $1 \leq K^{1000} \leq R$. Let S be a $1 \times R^{1/2} \times \cdots \times R^{1/2}$ -slab in \mathbb{R}^{n+1} parallel to the n-dimensional subspace V. Let $\tau \in \Theta_{K^{-1}}$. Suppose there exists $z_0 \in S$ such that

$$\measuredangle(G^{\lambda}(z_0;\tau),V) > K^{-2}$$

Then for any set of distinct R-planks \mathbf{T} that $\theta(T) \subset \tau$ for all $T \in \mathbf{T}$, we have

$$\|\sum_{T\in\mathbf{T}} \mathcal{F}^{\lambda} f_{T}\|_{L^{2}(S)}^{2} \lesssim K^{O(1)} \sum_{T\in\mathbf{T}} \|f_{T}\|_{2}^{2}.$$

Proof. We refer the reader to Figure 2, where the planks are curved rather than straight. Since we are allowed to lose a $K^{O(1)}$ factor, by the triangle inequality, we assume the direction caps of all the $T \in \mathbf{T}'$ lie in a K^{-10n} -cap. Let $\mathbf{T} = \bigcup_{\theta} \mathbf{T}_{\theta}$, where the directional cap for the planks in \mathbf{T}_{θ} is θ . Let $f_{\theta} = \sum_{T \in \mathbf{T}_{\theta}} f_{T}$, so that

$$\sum_{T \in \mathbf{T}} \mathcal{F}^{\lambda} f_T = \sum_{\theta} \mathcal{F}^{\lambda} f_{\theta}.$$

We will show

(3.23)
$$\|\sum_{\theta} \mathcal{F}^{\lambda} f_{\theta}\|_{L^{2}(S)}^{2} \lesssim K^{O(1)} \sum_{\theta} \|f_{\theta}\|_{2}^{2}.$$

Then the lemma follows from the L^2 -orthogonality of $\sum_{T \in \mathbf{T}_{\theta}} f_T$.

Let $\mathbf{1}_{S}^{*}$ be a bump function that equals 1 on S and is supported on 2S. By expanding the L^{2} norm, the left-hand side of (3.23) is bounded above by

(3.24)
$$\sum_{\theta_1,\theta_2} \int \mathbf{1}_S^* e^{i(\phi^{\lambda}(z;\xi_1) - \phi^{\lambda}(z;\xi_2))} \mathrm{d}z \int a^{\lambda}(z;\xi_1) \overline{a^{\lambda}(z;\xi_2)} \widehat{f}_{\theta_1}(\xi_1) \overline{\widehat{f}_{\theta_2}(\xi_2)} \mathrm{d}\xi_1 \mathrm{d}\xi_2.$$

Note that (3.23) follows from the estimate

(3.25)
$$\left|\int \mathbf{1}_{S}^{*} e^{i(\phi^{\lambda}(z;\xi_{1})-\phi^{\lambda}(z;\xi_{2}))} \mathrm{d}z\right| = R^{n/2} (1+R^{1/2}K^{-O(1)}|\xi_{1}-\xi_{2}|)^{-10n}$$

In fact, for each θ , partition θ into $R^{-1/2}$ -balls ω , and by let $\hat{f}_{\omega} = \hat{f}_{\theta} \mathbf{1}_{\omega}$. Then \hat{f}_{ω} is supported in an $R^{1/2}$ -ball, and we have

$$(3.24) = \sum_{\omega_1,\omega_2} \int \mathbf{1}_S^* e^{i(\phi^{\lambda}(z;\xi_1) - \phi^{\lambda}(z;\xi_2))} \mathrm{d}z \int a^{\lambda}(z;\xi_1) \overline{a^{\lambda}(z;\xi_2)} \widehat{f}_{\omega_1}(\xi_1) \overline{\widehat{f}_{\omega_2}(\xi_2)} \mathrm{d}\xi_1 \mathrm{d}\xi_2,$$

which, via (3.25) and the estimate $\|\widehat{f}_{\omega}\|_1 \lesssim R^{-n/4} \|\widehat{f}_{\omega}\|_2$, is bounded above by

$$\sum_{\omega_{1},\omega_{2}} \int |f_{\omega_{1}}(\xi_{1})f_{\omega_{2}}(\xi_{2})|R^{n/2}(1+R^{1/2}K^{-O(1)}|\xi_{1}-\xi_{2}|)^{-10n}d\xi_{1}d\xi_{2}$$

$$\lesssim \sum_{\omega_{1},\omega_{2}} (1+R^{1/2}K^{-O(1)}|c(\omega_{1})-c(\omega_{2})|)^{-10n}\|\widehat{f}_{\omega_{1}}\|_{2}\|\widehat{f}_{\omega_{2}}\|_{2}$$

$$\lesssim K^{O(1)}\sum_{\omega} \|\widehat{f}_{\omega}\|_{2}^{2} = K^{O(1)}\sum_{\theta} \|\widehat{f}_{\theta}\|_{2}^{2}.$$

Here $c(\omega)$ is the center of ω . This proves (3.23).

By the method of stationary phase, to prove (3.25), it suffices to show that there exists $\vec{v} \parallel V$ such that for all $z \in 2S$,

(3.26)
$$|\partial_z(\phi^{\lambda}(z;\xi_1) - \phi^{\lambda}(z;\xi_2)) \cdot \vec{v}| \gtrsim K^{-O(1)} |\xi_1 - \xi_2|$$

Via Taylor's theorem, $\partial_z \phi^{\lambda}(z;\xi_1) - \partial_z \phi^{\lambda}(z;\xi_2) = \partial_{z\xi}^2 \phi^{\lambda}(z;\xi_2) \cdot (\xi_1 - \xi_2) + O(|\xi_1 - \xi_2|^2)$. Since $|\partial_{z\xi}^2 \phi^{\lambda}(z;\xi_2) \cdot (\xi_1 - \xi_2)| \gtrsim |\xi_1 - \xi_2|$ and since $|\xi_1 - \xi_2| \lesssim K^{-100}$, the following is true:

- (1) $|\partial_z(\phi^{\lambda}(z;\xi_1) \phi^{\lambda}(z;\xi_2))| \sim |\xi_1 \xi_2|.$ (2) $\angle (\partial_z(\phi^{\lambda}(z;\xi_1) \phi^{\lambda}(z;\xi_2)), \partial^2_{z\xi}\phi^{\lambda}(z;\xi_2) \cdot (\xi_1 \xi_2)) \lesssim K^{-100}.$

Thus, (3.26) boils down to showing that for all $z \in 2S$,

(3.27)
$$\mathcal{L}(\partial_{z\xi}^2 \phi^{\lambda}(z;\xi_2) \cdot (\xi_1 - \xi_2), V^{\perp}) \gtrsim K^{-50}.$$

Note that when $|z - z_0| \lesssim R^{1/2}$,

$$|\partial_{z\xi}^2 \phi^{\lambda}(z;\xi_2) - \partial_{z\xi}^2 \phi^{\lambda}(z_0;\xi_2)| \lesssim |z - z_0| \cdot \|\partial_{zz\xi}^3 \phi^{\lambda}\|_{\infty} \lesssim R^{1/2} \lambda^{-1} < K^{-100}.$$

Hence, to show (3.27), we just need to show

(3.28)
$$\measuredangle (\partial_{z\xi}^2 \phi^{\lambda}(z_0;\xi_2) \cdot (\xi_1 - \xi_2), V^{\perp}) > K^{-30}.$$

However, by the assumption in the lemma, we have

$$\measuredangle(G^{\lambda}(z_0;\xi_2),V) > K^{-2},$$

which, since $G^{\lambda}(z_0;\xi_2)$ is orthogonal to $\partial_{z\xi}^2 \phi^{\lambda}(z_0;\xi_2) \cdot (\xi_1 - \xi_2)$, implies (3.28). \Box

As a direct corollary, we have

Lemma 3.18. Let Q be an $\mathbb{R}^{1/2}$ -ball and \mathbf{T} is a set of wave packets. Then

$$\|\sum_{T \in \mathbf{T}} \mathcal{F}^{\lambda} f_{T}\|_{L^{2}(Q)}^{2} \lesssim R^{1/2} \sum_{T \in \mathbf{T}} \|f_{T}\|_{2}^{2}.$$

4. WAVE PACKET DENSITY

In this section, we introduce the wave packet density $\mathcal{W}(f, B_R^{n+1}(z_0))$. Given a s-cap $\tau \in \Theta_s$ with $R^{-1/2} \leq s \leq 1$, recall that

$$V_{\tau,R} = (Rs^2)\tau$$

is an isotropic dilate of τ^* , a box centered at the origin of dimensions $Rs^2 \times Rs \times$ $\cdots \times Rs.$

Definition 4.1 (Wave packet density). Suppose g is a function defined in \mathbb{R}^n such that $\operatorname{supp} \widehat{g} \subset \mathbb{A}^n(1)$, and suppose $B_R^{n+1}(z_0) = B_R^{n+1}(x_0, t_0) \subset B_\lambda^{n+1}$. Define

 $\mathcal{W}(g, B_R^{n+1}(z_0)) :=$

(4.1)
$$\sup_{R^{-1/2} \le s \le 1} \sup_{\tau \in \Theta_s} \sup_{V \mid |V_{\tau,R}} \left(\frac{1}{|V|} \int \sum_{\substack{\theta \subset \tau \\ \theta \in \Theta_{R^{-1/2}}}} \sum_{\substack{T \in \mathbb{T}_{\theta}(z_0) \\ T^{\flat} \subset V}} |g_T|^2 \right)^{1/2}$$

Here $V \parallel V_{\tau,R}$ means V is a translated copy of $V_{\tau,R}$, and recall (3.18) for $\mathbb{T}_{\theta}(z_0)$.

Remark 4.2. We give some intuitive explanations to the set of planks $\{T \in \bigcup_{\theta \subset \tau} \mathbb{T}_{\theta}(z_0) : T^{\flat} \subset V\}$ that appeared in the above summation. For each $V \parallel V_{\tau,R}$, mimicking Definition 3.5, we define

(4.2)
$$\Gamma_V(z_0;\xi_\tau;R) := \{ (\gamma^\lambda(v+\partial_\xi \phi^\lambda(z_0;\xi_\tau),t;\xi_\tau) : v \in V, |t-t_0| \le R \}.$$

When $z_0 = z$, it recovers Definition 3.5. From (3.17), one sees that for $T \in \mathbb{T}(z_0)$, $T = \Gamma_{R^{\delta}T^{\flat}}(z_0;\xi_{\theta};R)$. We call $\Gamma_V(z_0;\xi_{\tau};R)$ a curved plank in a ball centered at z_0 . Note that Lemma 3.9 and Lemma 3.12 still hold in this setting when we move the center from the origin to z_0 . By Lemma 3.12, $\Gamma_V(z_0;\xi_{\tau};R)$ is comparable to $\Gamma_V(z_0;\xi_{\theta};R)$ for $\theta \subset \tau$. Hence, we may identify $\Gamma_V(z_0;\xi_{\tau};R)$ and $\Gamma_V(z_0;\xi_{\theta};R)$, and the condition $T^{\flat} \subset V$ implies $T \subset \Gamma_V(z_0;\xi_{\tau};R)$. Therefore, the wave packets summed in (4.1) are morally those T that are contained in the fat curved plank $\Gamma_V(z_0;\xi_{\tau};R)$ with directional cap contained in τ . In this regard, $\mathcal{W}(g, B_R^{n+1}(z_0))$ can be viewed as the density of wave packets among all the fat curved planks in $B_R^{n+1}(z_0)$.

Lemma 4.3. Suppose g is a function defined in \mathbb{R}^n with $\operatorname{supp} \widehat{g} \subset \mathbb{A}^n(1)$. Let $g_{\mathbb{T}_1} = \sum_{T \in \mathbb{T}_1} g_T$ and $g_{\mathbb{T}_2} = \sum_{T \in \mathbb{T}_2} g_T$ be the sums of wave packets at scale R in $B_R^{n+1}(z_0)$. If $\mathbb{T}_1 \subset \mathbb{T}_2$, then

$$\mathcal{W}(g_{\mathbb{T}_1}, B_R^{n+1}(z_0))^2 \le \mathcal{W}(g_{\mathbb{T}_2}, B_R^{n+1}(z_0))^2$$

Proof. The proof is straightforward from the definition.

Lemma 4.4. Suppose a function g defined in \mathbb{R}^n is a sum of wave packets $g = \sum_{T \in \mathbb{T}} g_T$. Suppose $\|g_T\|_2^2 \gtrsim R^{\frac{n-1}{2}}$ for all $T \in \mathbb{T}$. Then

$$\mathcal{W}(g, B_R^{n+1})^2 \gtrsim \max\{1, R^{-\frac{n+1}{2}}(\#\mathbb{T})\} \gtrsim R^{-\frac{n+1}{4}}(\#\mathbb{T})^{1/2}.$$

Proof. In the first inequality, the lower bound 1 is obtained by testing with $V = T^{\flat}$ in (4.1), and the lower bound $R^{-\frac{n+1}{2}}(\#\mathbb{T})$ is obtained by testing with $V = B_R^n$. \Box

The next two lemmas are the main reasons why the wave packet density is appropriate for induction on scales. The first lemma suggests the wave packet density is an appropriate substitute for the L^{∞} norm.

Lemma 4.5. Suppose g is a function defined in \mathbb{R}^n with $\operatorname{supp} \widehat{g} \subset \mathbb{A}^n(1)$. Then

 $\mathcal{W}(g, B_R^{n+1}) \lesssim \|g\|_{\infty}.$

Proof. For $\tau, V \parallel V_{\tau,R}$, denote $\mathbb{T} = \{T \in \bigcup_{\theta \subset \tau} \mathbb{T}_{\theta}(z_0) : T^{\flat} \subset V\}$. We just need to prove

$$\frac{1}{|V|} \int \sum_{T \in \mathbb{T}} |g_T|^2 \lesssim ||g||_{\infty}^2.$$

Recall that both $\{\psi_{\theta}\}$ and $\{\eta_{T^{\flat}}\}$ are families of non-negative functions, and that $\{\operatorname{supp}(\psi_{\theta})\}_{\theta}$ are finitely overlapping. By pigeonholing, we can choose $\mathbb{T}' \subset \mathbb{T}$ so that the wave packets $\{g_T : T \in \mathbb{T}'\}$ are essentially orthogonal, and we have

(4.3)
$$\frac{1}{|V|} \int \sum_{T \in \mathbb{T}} |g_T|^2 \lesssim \frac{1}{|V|} \int \sum_{T \in \mathbb{T}'} |g_T|^2 \lesssim \frac{1}{|V|} \int |\sum_{T \in \mathbb{T}'} g_T|^2.$$

Decompose

$$\sum_{T \in \mathbb{T}'} g_T = \sum_{T \in \mathbb{T}'} (g \mathbf{1}_{100V})_T + \sum_{T \in \mathbb{T}'} (g \mathbf{1}_{(100V)^c})_T.$$

We claim that $(g\mathbf{1}_{(100V)^c})_T = \operatorname{RapDec}(R) ||g||_{\infty}$. Note that,

$$(g\mathbf{1}_{(100V)^c})_T = \eta_{T^\flat} \big(\psi_\theta^{\lor} * (g\mathbf{1}_{(100V)^c})\big).$$

Since ψ_{θ}^{\vee} is essentially supported in $\theta^* \subset V_{\tau,R}$, $\psi_{\theta}^{\vee} * (g\mathbf{1}_{(100V)^c})$ is essentially supported in $(100V)^c + V_{\tau,R} \subset (90V)^c$. The claim follows from the fact that $\eta_{T^{\flat}}$ is essentially supported in $T^{\flat} \subset V$. Therefore,

$$(4.3) \lesssim \frac{1}{|V|} \int_{V} |\sum_{T \in \mathbb{T}'} (g\mathbf{1}_{100V})_{T}|^{2} + \operatorname{RapDec}(R) ||g||_{\infty}^{2}$$
$$\lesssim \frac{1}{|V|} \int |g\mathbf{1}_{100V}|^{2} + \operatorname{RapDec}(R) ||g||_{\infty}^{2}$$
$$\lesssim ||g||_{\infty}^{2} + \operatorname{RapDec}(R) ||g||_{\infty}^{2}.$$

The second lemma establishes a connection between wave packet densities at different scales.

Lemma 4.6. Suppose $1 \le r \le R^{1-10\delta}$ and $B_r^{n+1}(z_0) \subset B_R^{n+1}$. Then $\mathcal{W}(f, B_r^{n+1}(z_0)) \lesssim R^{O(\delta)}(\frac{R}{r})^{\frac{n-1}{2}} \mathcal{W}(f, B_R^{n+1}) + \operatorname{RapDec}(R) ||f||_2.$

Proof. By definition, we can assume $\mathcal{W}(f, B_r^{n+1}(z_0))$ is attained at the pair $(\widetilde{V}, \widetilde{\tau})$. In other words, there are $\widetilde{s} \in [r^{-1/2}, 1], \ \widetilde{\tau} \in \Theta_{\widetilde{s}}, \ \widetilde{V} \parallel V_{\widetilde{\tau},r}$ such that

(4.4)
$$\mathcal{W}(f, B_r^{n+1}(x_0))^2 \lesssim \frac{1}{|\widetilde{V}|} \int \sum_{\widetilde{T} \in \widetilde{\mathbf{T}}} |f_{\widetilde{T}}|^2 \lesssim \frac{1}{|\widetilde{V}|} \int \left| \sum_{\widetilde{T} \in \widetilde{\mathbf{T}}'} f_{\widetilde{T}} \right|^2,$$

where $\widetilde{\mathbf{T}}' \subset \widetilde{\mathbf{T}}$ is an appropriate subset, and $\widetilde{\mathbf{T}}$ is defined as

$$\widetilde{\mathbf{T}}:=\{\widetilde{T}\in\bigcup_{\widetilde{\theta}\in\Theta_{r^{-1/2}},\widetilde{\theta}\subset\widetilde{\tau}}\mathbb{T}_{\widetilde{\theta}}:\widetilde{T}^{\flat}\subset\widetilde{V}\}.$$

Next, we analyze how the wave packets at scale r interact with the wave packets at scale R. Recall the wave packet decomposition at scale R in B_R^n :

$$f = \sum_{\theta \in \Theta_{R^{-1/2}}} \sum_{T \in \mathbb{T}_{\theta}} f_T + \operatorname{RapDec}(R) \|f\|_2.$$

We define a relation \prec between wave packets of scales r and R. For two planks $\widetilde{T} \in \mathbb{T}_{\widetilde{\theta}}(z_0), T \in \mathbb{T}_{\theta}$, if $\theta \subset 4\widetilde{\theta}$ and $100T \cap 100\widetilde{T} \neq \emptyset$, we write

$$T \prec T.$$

By Lemma 3.14, $(f_T)_{\widetilde{T}} = \operatorname{RapDec}(R) ||f||_2$ unless $\widetilde{T} \prec T$.


Let $s = \widetilde{s}(\frac{r}{R})^{1/2} R^{5\delta}$, so $s \in [R^{-1/2+5\delta}, 1]$. Let $\Theta_s(4\widetilde{\tau}) \subset \Theta_s$ be a subset with $\#\Theta_s(4\widetilde{\tau}) \sim (\frac{R}{r})^{\frac{n-1}{2}}$ such that

$$4\widetilde{\tau} \subset \bigcup_{\tau \in \Theta_s(4\widetilde{\tau})} \tau.$$

For each $\tau \in \Theta_s(4\tilde{\tau})$, let $V_{\tau} \parallel V_{\tau,R}$ be such that $V_{\tau} \supset \tilde{V} + \partial_{\xi} \phi^{\lambda}(z_0;\xi_{\tau})$, and define

$$\mathbf{T}_{\tau} := \{ T \in \bigcup_{\theta \in \Theta_{R^{-1/2}}, \theta \subset \tau} \mathbb{T}_{\theta} : T^{\flat} \subset V_{\tau} \}.$$

By remark 4.2, $\widetilde{\mathbf{T}}$ is morally the set of *r*-planks contained in $\Gamma_{\widetilde{V}}(z_0; \xi_{\widetilde{\tau}}; r)$ with directional cap contained in $\widetilde{\tau}$; \mathbf{T}_{τ} is morally the set of *R*-planks contained in $\Gamma_{V_{\tau}}(\xi_{\tau}; R)$ with directional cap contained in τ . See Figure 4 for the geometry of $\Gamma_{\widetilde{V}}(z_0; \xi_{\widetilde{\tau}}; r)$ and $\{\Gamma_{V_{\tau}}(\xi_{\tau}; R) : \tau \in \Theta_s(4\widetilde{\tau})\}$. Let





We claim that if $\widetilde{T} \prec T$ for some $\widetilde{T} \in \widetilde{\mathbf{T}}$, then $T \in \mathbf{T}$. Indeed, since $\widetilde{T} \prec T$, we have $T \in \mathbb{T}_{\theta}$ for some $\theta \subset 4\widetilde{\tau}$. Thus, we can find $\tau \in \Theta_s(4\widetilde{\tau})$ such that $\theta \subset \tau$. Consider the curved plank $\Gamma_{\widetilde{V}}(z_0;\xi_{\widetilde{\tau}};r)$ as in (4.2). By Remark 4.2, planks in $\widetilde{\mathbf{T}}$ are all contained in $\Gamma_{C\widetilde{V}}(z_0;\xi_{\widetilde{\tau}};r)$ for some large C > 0. By Lemma 3.11, $\Gamma_{C\widetilde{V}}(z_0;\xi_{\widetilde{\tau}};r) \subset \Gamma_{C^2\widetilde{V}}(z_0;\xi_{\tau};r)$.

Since T intersects the dilate of some \widetilde{T} in $\widetilde{\mathbf{T}}$, we know that $T \cap \Gamma_{C^3 \widetilde{V}}(z_0; \xi_\tau; r) \neq \emptyset$, which implies $R^{\delta} T^{\flat} \cap \left(C^3 \widetilde{V} + \partial_{\xi} \phi^{\lambda}(z_0; \xi_\tau) \right) \neq \emptyset$ via (4.2). This would imply

$$T^{\flat} \subset V_{\tau}.$$

This shows $T \in \mathbf{T}_{\tau} \subset \mathbf{T}$ and hence the claim.

Consequently, from (4.4), we have

$$\begin{split} \mathcal{W}(f, B_R^{n+1}(z_0))^2 &\lesssim \frac{1}{|\widetilde{V}|} \int \bigg| \sum_{\widetilde{T} \in \widetilde{\mathbf{T}}'} \bigg(\sum_{T \in \mathbf{T}} f_T \bigg)_{\widetilde{T}} \bigg|^2 + \operatorname{RapDec}(R) \|f\|_2^2 \\ &\lesssim \frac{1}{|\widetilde{V}|} \int \bigg| \sum_{T \in \mathbf{T}} f_T \bigg|^2 + \operatorname{RapDec}(R) \|f\|_2^2 \\ &\lesssim \frac{1}{|\widetilde{V}|} \int \sum_{T \in \mathbf{T}} |f_T|^2 + \operatorname{RapDec}(R) \|f\|_2^2 \\ &= \sum_{\tau \in \Theta_s(4\widetilde{\tau})} \frac{1}{|\widetilde{V}|} \int \sum_{T \in \mathbf{T}_\tau} |f_T|^2 + \operatorname{RapDec}(R) \|f\|_2^2. \end{split}$$

Now we finish the proof of the lemma. By the definition of \mathbf{T}_{τ} , we have

$$\mathcal{W}(f, B_r^{n+1}(z_0))^2 \lesssim \sum_{\tau \in \Theta_s(4\tilde{\tau})} \frac{|V_\tau|}{|\widetilde{V}|} \left(\frac{1}{|V_\tau|} \int \sum_{\substack{\theta \subset \tau \\ \theta \in \Theta_{R^{-1/2}}}} \sum_{\substack{T \in \mathbb{T}_\theta \\ T^\flat \subset V_\tau}} |f_T|^2 \right) + \operatorname{RapDec}(R) ||f||_2^2.$$

Note that each V_{τ} is of dimensions $R^{10\delta}r\tilde{s}^2 \times R^{5\delta}(Rr)^{1/2}\tilde{s} \times \cdots \times R^{5\delta}(Rr)^{1/2}\tilde{s}$. We have

$$\frac{|V_{\tau}|}{|\widetilde{V}|} \lesssim R^{O(\delta)} (\frac{R}{r})^{\frac{n-1}{2}}$$

Also note

$$\#\Theta_s(4\widetilde{\tau}) \lesssim R^{O(\delta)}(\frac{R}{r})^{\frac{n-1}{2}}.$$

We obtain

$$\mathcal{W}(f, B_r^{n+1}(z_0))^2 \lesssim R^{O(\delta)}(\frac{R}{r})^{n-1} \mathcal{W}(f, B_R^{n+1})^2 + \text{RapDec}(R) \|f\|_2^2.$$

Remark 4.7. The $R^{O(\delta)}$ factor appears here since the wave packet T has size $R^{2\delta} \times R^{1/2+\delta} \times \cdots \times R^{1/2+\delta}$, rather than $1 \times R^{1/2} \times \cdots \times R^{1/2}$. Though, the lemma still works for the induction step since δ is very small compared to ε .

Remark 4.8. The wave packet density introduced in Definition 4.1 naturally imposes a non-concentration condition on the set of planks \mathbb{T} , where $g = \sum_{T \in \mathbb{T}} g_T$ is the wave packet decomposition of g. Indeed, if $||g_T||_2$ are about the same for all $T \in \mathbb{T}$, then the assumption $\mathcal{W}(g, B_R^{n+1}(z_0)) \lesssim 1$ is the necessary condition for the following Kakeya estimate to hold: $|\bigcup_{\mathbb{T}}| \gtrsim \sum_{T \in \mathbb{T}} |T|$. However, it is likely that this is not a sufficient condition. Under the current definition of wave packet density, the assumption $\mathcal{W}(g, B_R^{n+1}(z_0)) \lesssim 1$ is analogous to the classical "(n-1)-ball condition" for a fractal set (see [14, (1.1)] for an example with n = 3 and \mathbb{T} replaced by a family of tubes).

Remark 4.9 (Continuation of Remark 4.8). Nevertheless, wave packet density is a versatile concept. One may modify its definition to impose a stronger nonconcentration condition on the planks \mathbb{T} . For instance, the next modification of Definition 4.1 will enable us to impose a non-concentration condition on the planks \mathbb{T} , similar to the *Convex Wolff Axiom* (see [22, Definition 0.12]) for tubes:

28

For dyadic numbers $s_1, s_2 \ldots, s_{n-1} \in [R^{-1/2}, 1]$, and τ_{\circ} being a convex subset of $[-1, 1]^{n-1} \times \{1\}$ of dimensions $s_1 \times \cdots \times s_{n-1}$, define $\tau := \{\xi \in \mathbb{A}^n(1) : \xi/|\xi| \in \tau_{\circ}\}$. We say τ is an (s_1, \ldots, s_{n-1}) -cap. Then τ is a convex set and can be roughly viewed as a box of dimensions $1 \times s_1 \times \cdots \times s_{n-1}$. For example, a cap $\tau \in \Theta_s$ is an (s, \ldots, s) -cap. Given an (s_1, \ldots, s_{n-1}) -cap τ , we define its dual box $V_{\tau,R}$ in the physical space \mathbb{R}^n as follows: Let $s_{\max} := \max_{1 \le i \le n-1} s_i$. Define $V_{\tau,R}$ to be the box centered at the origin, of dimensions $Rs_{\max}^2 \times Rs_1 \times \cdots \times Rs_{n-1}$, whose edges are parallel to the corresponding edges of the $1 \times s_1 \times \cdots \times s_{n-1}$ -box τ .

Now, we give a generalization of Definition 4.1. One can check that the lemmas corresponding to Lemmas 4.3, 4.5, and 4.6 are also true.

Generalization of Definition 4.1: Suppose g is a function in \mathbb{R}^n with $\operatorname{supp} \widehat{g} \subset \mathbb{A}(1)$, and $B^n_B(x_0)$ is a ball. Define

$$\widetilde{\mathcal{W}}(g, B_R^{n+1}(z_0)) := \sup_{\tau} \sup_{\substack{V \mid V_{\tau,R} \\ V \subset B_R^n}} \left(\frac{1}{|V|} \int \sum_{\substack{\theta \subset \tau \\ \theta \in \Theta_R}} \sum_{\substack{T^\flat \subset V \\ T \in \mathbb{T}_\theta(z_0)}} |g_T|^2 \right)^{1/2}.$$

Here, \sup_{τ} is taken over all caps (s_1, \ldots, s_{n-1}) -caps τ for all dyadic numbers $s_1, s_2, \ldots, s_{n-1} \in [R^{-1/2}, 1]$.

Finally, we reduce Proposition 3.2 to the following theorem, which is formulated in terms of a mixed norm defined via the wave packet density.

Theorem 4.10. Let \mathcal{F} be a type 1 Fourier integral operator given by (1.5). Suppose the phase function ϕ satisfies (3.1), and the amplitude function a obeys $\operatorname{supp}_{\xi} a \subset \mathbb{A}^n(1)^\circ$. Let \mathcal{F}^{λ} be given by (3.2). Then for all $\varepsilon > 0$ and when p = p(n), there exists a constant C_{ε} that is independent to \mathcal{F} , such that

(4.5)
$$\|\mathcal{F}^{\lambda}f\|_{L^{p}(B_{R}^{n+1})} \leq C_{\varepsilon}R^{(n-1)(\frac{1}{2}-\frac{1}{p})+\varepsilon}\|f\|_{2}^{\frac{2}{p}}\mathcal{W}(f,B_{R}^{n+1})^{1-\frac{2}{p}}$$

for all f with $\operatorname{supp} \widehat{f} \subset \mathbb{A}^n(1)$ and all $R \in [1, \lambda^{1-\varepsilon}]$.

Proof that Theorem 4.10 implies Proposition 3.2. Since $\operatorname{supp}_{\xi} a \subset \mathbb{A}^n(1)^\circ$, it follows that $\mathcal{F}^{\lambda}f = \mathcal{F}^{\lambda}(\mathbf{1}_{\mathbb{A}^n(1)}^*\widehat{f})^{\vee}$, where $\mathbf{1}_{\mathbb{A}^n(1)}^*$ is a bump that equals to 1 on $\operatorname{supp}_{\xi} a$ and is supported in $\mathbb{A}^n(1)$. By summing up all balls B_R^{n+1} inside B_{λ}^{n+1} , Theorem 4.10 and Lemma 4.5 imply that for all $\varepsilon > 0$ and when p = p(n),

(4.6)
$$\|\mathcal{F}^{\lambda}f\|_{L^{p}(B^{n+1}_{\lambda})} \leq C_{\varepsilon}\lambda^{(n-1)(\frac{1}{2}-\frac{1}{p})+\varepsilon}\|f\|_{2}^{\frac{2}{p}}\|f\|_{\infty}^{1-\frac{2}{p}}$$

We see that (4.6) implies (3.4) when f is a characteristic function and when p = p(n). Therefore, the range p > p(n) for (3.4) follows from a real interpolation between the restrict-type estimate when p = p(n) and the trivial bound $\|\mathcal{F}^{\lambda}f\|_{\infty} \lesssim \lambda^{\frac{n-1}{2}} \|f\|_{\infty}$. This proves Proposition 3.2.

5. Some geometric results

In this section, we establish several geometric results that will be used to prove Theorem 4.10 in subsequent sections. 5.1. Lorentz rescaling. Suppose $\tau \in \Theta_{K^{-1}}$ and $\operatorname{supp} \widehat{f} \subset \tau$. We want to find an upper bound for $\|\mathcal{F}^{\lambda}f\|_{L^{p}(B_{R}^{n+1})}$ by induction. The idea is to perform a rescaling so that $\mathcal{F}^{\lambda}f$ becomes $\widetilde{\mathcal{F}}^{\lambda/K^{2}}g$, where $\widetilde{\mathcal{F}}$ is a new Fourier integral operator, and g is a function with $\operatorname{supp} \widehat{g} \subset \mathbb{A}^{n}(1)$. At the same time, the scale of the integration domain drops from R to R/K^{2} . Moreover, when \mathcal{F} is of type 1, the new operator $\widetilde{\mathcal{F}}$ is of type (1, 1, C) for some constant C (see [1, Section 2.5]). Therefore, by partitioning the new operator $\widetilde{\mathcal{F}}$ into O(1) parts, we can use an induction hypothesis at the scale $(\lambda/K^{2}, R/K^{2})$ to bound $\|\mathcal{F}^{\lambda}f\|_{L^{p}(R^{n+1})}$.

scale $(\lambda/K^2, R/K^2)$ to bound $\|\mathcal{F}^{\lambda}f\|_{L^p(B_R^{n+1})}$. Next, we describe the rescaling and the relevant geometry. Given a $\tau \in \Theta_{K^{-1}}$, let $V_{\tau,R} \subset B_R^n$ be a dual $RK^{-2} \times RK^{-1} \times \cdots \times RK^{-1}$ -box, as defined in Lemma 3.11. Tile B_R^n by translated copies of $V_{\tau,R}$ and denote them by $\mathbb{B}_{\tau}^{\flat} = \{\Box^{\flat}\}$. For each $\Box^{\flat} \in \mathbb{B}_{\tau}^{\flat}$, define a curved plank

(5.1)
$$\Box := \Gamma_{\Box^{\flat}}(\xi_{\tau}; R) = \{ (\gamma^{\lambda}(v, t; \xi_{\tau}), t; \xi_{\tau}) : v \in \Box^{\flat}, |t| \le R \}.$$

Denote $\mathbb{B}_{\tau} = \{ \Box : \Box^{\flat} \in \mathbb{B}_{\tau}^{\flat} \}$. Observe that \mathbb{B}_{τ} form a covering of B_R^{n+1} , and each $T \in \bigcup_{\theta \subset \tau} \mathbb{T}_{\theta}$ belongs to O(1) many $\Box \in \mathbb{B}_{\tau}$. We assign each $T \in \bigcup_{\theta \subset \tau} \mathbb{T}_{\theta}$ to a single \Box to which T belongs, and denote the collection of T assigned to \Box by \mathbb{T}_{\Box} . This gives a partition

(5.2)
$$\bigcup_{\theta \subset \tau} \mathbb{T}_{\theta} = \bigcup_{\Box} \mathbb{T}_{\Box}.$$

Given a $\tau \in \Theta_{K^{-1}}$, write $\xi_{\tau} = (\omega, 1)$, and define $\Upsilon_{\omega}(y, t) := (\gamma(y, t; \omega, 1), t)$ and $\Upsilon_{\omega}^{\lambda}(y, t) := \lambda \Upsilon_{\omega}(y/\lambda, t/\lambda)$. Introduce two non-isotropic dilations $D_K(y', y_n, t) := (Ky', y_n, K^2t)$ and $D'_{K^{-1}}(y', y_n) := (K^{-1}u', K^{-2}u_n)$. Then, as shown in [1, Proof of Lemma 2.3], we have

(5.3)
$$\mathcal{F}^{\lambda}g \circ \Upsilon^{\lambda}_{\omega} \circ D_K = \widetilde{\mathcal{F}}^{\lambda/K^2}\widetilde{g},$$

where

$$\widetilde{g}(\eta) := K^{-(n-1)}g(\eta_n\omega + K^{-1}\eta', \eta_n),$$

(5.4)
$$\widetilde{\mathcal{F}}^{\lambda/K^2}\widetilde{g}(y,t) := \int_{\mathbb{R}^n} e^{i\widetilde{\phi}^{\lambda/K^2}(y,t;\eta)} \widetilde{a}^{\lambda}(y,t;\eta)\widetilde{g}(\eta) \mathrm{d}\eta$$

Here the phase $\phi(y, t; \eta)$ is given by

$$\langle y,\eta\rangle + \int_0^1 (1-r) \langle \partial_{\xi'\xi'}^2 \phi(\Upsilon_\omega(D'_{K^{-1}}y,t);\eta_n\omega + rK^{-1}\eta',\eta_n)\eta',\eta'\rangle \mathrm{d}r,$$

and the amplitude function $\tilde{a}(y,t;\eta)$ is given by

$$\widetilde{a}(y,t;\eta) := a(\Upsilon_{\omega}(D'_{K^{-1}}y;t);\eta_n\omega + K^{-1}\eta',\eta_n).$$

We remark that for each $\Box \in \mathbb{B}_{\tau}$, $(\Upsilon^{\lambda}_{\omega} \circ D_K)^{-1}(\Box)$ is morally a R/K^2 -ball.

Now we state and prove the following lemma.

Lemma 5.1. Let $1 \leq K \leq R^{\varepsilon}$. Suppose there is a constant C > 0 such that for all function g defined in \mathbb{R}^n with $\operatorname{supp} \widehat{g} \subset \mathbb{A}^n(1)$ and all type 1 FIO $\widetilde{\mathcal{F}}$,

(5.5)
$$\|\widetilde{\mathcal{F}}^{\lambda/K^2}g\|_{L^p(B^{n+1}_{R/K^2})} \le C\|g\|_2^{\frac{2}{p}} \mathcal{W}(g, B^{n+1}_{R/K^2})^{1-\frac{2}{p}}$$

Let $\tau \in \Theta_{K^{-1}}$. Then for all function f defined in \mathbb{R}^n with $\operatorname{supp} \widehat{f} \subset \tau$ and all type **1** FIO $\widetilde{\mathcal{F}}$, we have

$$\|\mathcal{F}^{\lambda}f\|_{L^{p}(B^{n+1}_{R})} \lesssim CK^{\frac{2}{p}} \|f\|_{2}^{\frac{2}{p}} \mathcal{W}(f, B^{n+1}_{R})^{1-\frac{2}{p}}.$$

Proof. Write $\tau = \bigcup_{\theta \subset \tau} \theta$, where $\theta \in \Theta_{R^{-1/2}}$. Recall the definitions of $\mathbb{B}_{\tau} = \{\Box\}$ in (5.1) and \mathbb{T}_{\Box} in (5.2). Perform the wave packet decomposition for $\mathcal{F}^{\lambda}f$ inside B_{R}^{n+1} to have

$$\mathcal{F}^{\lambda}f = \sum_{\theta \subset \tau} \sum_{T \in \mathbb{T}_{\theta}} \mathcal{F}^{\lambda}f_{T} = \sum_{\Box} \sum_{T \in \mathbb{T}_{\Box}} \mathcal{F}^{\lambda}f_{T}.$$

Recall the rescaling $\Upsilon^{\lambda}_{\omega} \circ D_{K}$ in (5.3). For each $\Box \in \mathbb{B}_{\tau}$, let $B^{n+1}_{R/K^{2},\Box}$ be the R/K^{2} ball containing $(\Upsilon^{\lambda}_{\omega} \circ D_{K})^{-1}(\Box)$. Note that, under this rescaling, each $T \in \mathbb{T}_{\Box}$ becomes a R/K^{2} -plank \widetilde{T} in $B^{n+1}_{R/K^{2},\Box}$. We denote $\widetilde{\mathbb{T}}_{\Box} = {\widetilde{T} : T \in \mathbb{T}_{\Box}}$. Thus, under the same rescaling, the function $\mathcal{F}^{\lambda}f$ becomes

$$\sum_{\Box} \sum_{\widetilde{T} \in \widetilde{\mathbb{T}}_{\Box}} \widetilde{\mathcal{F}}^{\lambda/K^2} \widetilde{g}_{\widetilde{T}}$$

where $\widetilde{\mathcal{F}}$ is an FIO of type (1, 1, C) for some constant C.

By the essential disjointness of \Box , we have

(5.6)
$$\|\mathcal{F}^{\lambda}f\|_{L^{p}(B_{R}^{n+1})}^{p} \lesssim \sum_{\Box} \|\sum_{T \in \mathbb{T}_{\Box}} \mathcal{F}^{\lambda}f_{T}\|_{L^{p}(\Box)}^{p}.$$

For each \Box , since $\Upsilon^{\lambda}_{\omega}$ is a diffeomorphism with determinant ~ 1 ,

$$\|\sum_{T\in\mathbb{T}_{\square}}\mathcal{F}^{\lambda}f_{T}\|_{L^{p}(\square)}\sim K^{\frac{n+1}{p}}\|\sum_{\widetilde{T}\in\widetilde{\mathbb{T}}_{\square}}\widetilde{\mathcal{F}}^{\lambda/K^{2}}g_{\widetilde{T}}\|_{L^{p}(B^{n+1}_{R/K^{2},\square})},$$

which, by partitioning $\widetilde{\mathcal{F}}$ into O(1) parts and by (5.5), is bounded by

$$\lesssim K^{\frac{n+1}{p}} C \| \sum_{\widetilde{T} \in \widetilde{\mathbb{T}}_{\square}} g_{\widetilde{T}} \|_2^{\frac{2}{p}} \mathcal{W}(\sum_{\widetilde{T} \in \widetilde{\mathbb{T}}_{\square}} g_{\widetilde{T}}, B^{n+1}_{R/K^2,\square})^{1-\frac{2}{p}}.$$

Again, since $\Upsilon^{\lambda}_{\omega}$ is a diffeomorphism with determinant $\sim 1,$ we have

$$\|\sum_{\widetilde{T}\in\widetilde{\mathbb{T}}_{\square}}g_{\widetilde{T}}\|_2 \sim K^{-\frac{n-1}{2}}\|\sum_{T\in\mathbb{T}_{\square}}f_T\|_2.$$

Finally, notice that

$$\mathcal{W}(\sum_{\widetilde{T}\in\widetilde{\mathbb{T}}_{\square}}g_{\widetilde{T}}, B_{R/K^{2},\square}^{n+1}) \sim \mathcal{W}(\sum_{T\in\mathbb{T}_{\square}}f_{T}, B_{R}^{n+1}) \lesssim \mathcal{W}(f, B_{R}^{n+1}).$$

Plugging the aforementioned information into (5.6), we obtain

$$\begin{split} \|\mathcal{F}^{\lambda}f\|_{L^{p}(B^{n+1}_{R})}^{p} &\lesssim C^{p}K^{2}\sum_{\Box}\|\sum_{T\in\mathbb{T}_{\Box}}f_{T}\|_{2}^{2}\mathcal{W}(f,B^{n+1}_{R})^{p-2} \\ &\lesssim C^{p}K^{2}\|f\|_{2}^{2}\mathcal{W}(f,B^{n+1}_{R})^{p-2}. \end{split}$$

5.2. Hairbrush. We call $\mathcal{T} \subset B_R^{n+1}$ an *R*-tube if \mathcal{T} has the form

 $\mathcal{T} = \Gamma_B(\xi; R)$

for some $R^{1/2}$ -ball $B = B^n_{R^{1/2}}(v)$ and $\xi \in \mathbb{A}^n(1)$ (recall Definition 3.5). Since $\Gamma_B(\xi; R) = \Gamma_B(s\xi; R)$ (recall (3.6)), we may assume $\xi = (\xi', 1)$ by rescaling. We call ξ the direction of the tube and v the position of the tube. Notice that the core curve of the tube is given by

(5.7)
$$\{(\gamma^{\lambda}(v,t;\xi),t): |t| \le R\}.$$

Similar to Definition 3.8, we make the following definition.

Definition 5.2. Two *R*-tubes $\Gamma_{B_1}(\xi_1; R)$ and $\Gamma_{B_2}(\xi_2; R)$ are distinct if

 $\Gamma_{B_i}(\xi_i; R) \not\subset \Gamma_{CB_k}(\xi_k; R)$

for $j, k \in \{1, 2\}, j \neq k$, and some absolute big constant C.

Also, similar to Lemma 3.11, we have

Lemma 5.3. Two R-tubes $\Gamma_{B_1}(\xi_1; R)$ and $\Gamma_{B_2}(\xi_2; R)$ are distinct if one of the following is true:

- (1) dist $(\xi'_1, \xi'_2) \gtrsim R^{-1/2}$. (2) dist $(B_1, B_2) \gtrsim R^{1/2}$.

Recall $K = R^{\varepsilon^{50}}$ and let $K_0 = R^{\varepsilon^{100}}$

Lemma 5.4. Let $\mathcal{T} = \{\mathcal{T}\}$ be a family of distinct *R*-tubes and let \mathcal{T}_0 be an *R*-tube in B_R^{n+1} . Let $\omega_1 = \mathbb{R}^n \times [i_1 R K_{\circ}^{-1}, (i_1 + 1) R K_{\circ}^{-1}]$ and $\omega_2 = \mathbb{R}^n \times [i_2 R K_{\circ}^{-1}, (i_2 + 1) R K_{\circ}^{-1}]$ be two horizontal regions with $i_1, i_2 \in \mathbb{Z}, |i_1|, |i_2| \leq K_{\circ}$ and $|i_1 - i_2| \geq 2$. Suppose for any $\mathcal{T} \in \mathcal{T}$, the direction of \mathcal{T}_0 and the direction of \mathcal{T} are K^{-1} separated, $\mathcal{T}_0 \cap \mathcal{T} \neq \emptyset$, and $\mathcal{T}_0 \cap \mathcal{T} \subset \omega_1$. Then for $z \in \omega_2$,

(5.8)
$$\sum_{\mathcal{T}\in\boldsymbol{\mathcal{T}}} \mathbf{1}_{\mathcal{T}}(z) \lesssim (KK_{\circ})^{O(1)}$$

Before proving the lemma, we would like to discuss the proof idea when \mathcal{F} is a constant coefficient FIO (note that \mathcal{T} depends on the operator \mathcal{F}). The variable coefficient case follows similarly.

We refer to Figure 5 for the key geometric observation. When \mathcal{F} is a constant coefficient FIO, each R-tube \mathcal{T} is a straight tube pointing to a light ray direction. Fix $t \in [i_2 R K_{\circ}^{-1}, (i_2 + 1) R K_{\circ}^{-1}]$, and let $\Pi = \mathbb{R}^n \times \{t\}$ be a horizontal hyperplane in ω_2 . It suffices to prove (5.8) for $z \in \Pi$. Let $\{Q\}$ be a finite-overlapping cover of $\mathcal{T}_0 \cap \omega_1$ by $R^{1/2}$ -balls. Since $\mathcal{T} \cap \mathcal{T}_0 \subset \omega_1$, we assign each $\mathcal{T} \in \mathcal{T}$ to a Q such that $\mathcal{T}_0 \cap \mathcal{T} \cap Q \neq \emptyset$. Denote the set of tubes assigned to Q by $\mathcal{T}(Q)$ so that

$$\mathcal{T} = \bigsqcup_{Q} \mathcal{T}(Q)$$

Let $C(Q) := \bigcup_{\mathcal{T} \in \mathcal{T}(Q)} \mathcal{T} \cap \Pi$, which is contained in the $R^{1/2}$ -neighborhood of a (n-1)-sphere in Π . Since dist $(\omega_1, \omega_2) \ge RK_{\circ}^{-1}$, the spheres C(Q) for different Qhave $R^{1/2}K_{\circ}^{-1}$ -separated radius, and all of them are tangent at $\mathcal{T}_0 \cap \Pi$. Also, since each \mathcal{T} and \mathcal{T}_0 are K^{-1} -separated in direction, $C(Q) \cap N_{100(KK_0)^{-1}R}(\mathcal{T}_0 \cap \Pi) = \emptyset$. Therefore, the sets $\{C(Q)\}_Q$ are $\lesssim (KK_\circ)^{O(1)}$ -overlapping, and hence $\{\mathcal{T} \cap \Pi\}_{\mathcal{T} \in \mathcal{T}}$ are $(KK_{\circ})^{O(1)}$ -overlapping. We remark that a O(1) loss in exponent is acceptable, as $(KK_{\circ})^{O(1)} \leq R^{10\varepsilon}$.




FIGURE 5. Hairbrush for light rays

The variable coefficient case will be clear once we understand the geometry of each corresponding C(Q). As we will see, each C(Q) is contained in the $R^{1/2}$ -neighborhood of a certain surface with a positive definite second fundamental form.

Proof of Lemma 5.4. After affine transformation, we assume \mathcal{T}_0 has position v = 0 and direction $(\xi', \xi_n) = (0, \ldots, 0, 1)$. Also, we want to simplify the phase function. We claim that, by applying a certain change of variables, we may assume

$$\phi(x,t;\xi) = \langle x,\xi \rangle + t \frac{h(\xi')}{\xi_n} + \mathcal{E}(x,t;\xi'/\xi_n)\xi_n,$$

where h is quadratic and $|\mathcal{E}(x,t;\xi')| = O(|t||\xi'|^3 + |z|^2|\xi'|^2)$ is the higher order term. The reader may compare it with [3, (1.11)].

Now, we prove the claim. Recall the notations z = (x, t), $x = (x', x_n)$, $\xi = (\xi', \xi_n)$. For $i \leq n$, note that $z_i = x_i$. Since ϕ is 1-homogeneous in ξ , we focus on $\phi(x, t; \xi', 1)$. Write

$$\phi(z;\xi',1) = \phi(z;0,1) + g(z;\xi'),$$

with $g(z;0) \equiv 0$. By the **(H1)** condition of ϕ , we have $\partial_x \phi(z;0,1) \neq 0$. We may assume $\partial_{z_n} \phi(z;0,1) \neq 0$, so that we can perform the change of variables $\tilde{z}_n = \phi(z;0,1)$ and $\tilde{z}_i = z_i$ $(i \neq n)$. In the new coordinate (we still use z to denote \tilde{z}), we have

$$\phi(z;\xi',1) = z_n + g(z;\xi').$$

Apply Taylor's expansion on g to have

$$\phi(z;\xi',1) = z_n + \sum_{i=1}^{n+1} \sum_{j=1}^{n-1} c_{ij} z_i \xi_j + g_1(z;\xi'),$$

where $\partial_{z\xi'}^2 g_1(0,0) = 0$. There is no ξ -term in the above expansion since $\partial_{\xi}^{\alpha} \phi(0;0) = 0$ by (3.1). Since ϕ is 1-homogeneous in ξ , we have

$$\phi(z;\xi',\xi_n) = z_n\xi_n + \sum_{i=1}^{n+1}\sum_{j=1}^{n-1}c_{ij}z_i\xi_j + g_1(z;\xi'/\xi_n)\xi_n.$$

By condition **(H1)** that $\partial_{x\xi}^2 \phi(0,0)$ has rank *n*, we see that

$$\partial_{x\xi}^2(z_n\xi_n + \sum_{i=1}^{n+1}\sum_{j=1}^{n-1}c_{ij}z_i\xi_j)$$

has rank n. Hence, a linear change of variables in z gives

$$\phi(z;\xi',\xi_n) = \sum_{i=1}^n z_i \xi_i + g_2(z;\xi'/\xi_n)\xi_n.$$

Thus, in appropriate coordinates,

$$\phi(z;\xi',1) = z_n + \sum_{i=1}^{n-1} z_i \xi_i + g_2(z;\xi').$$

By Taylor's expansion, we can further write

$$\phi(z;\xi',1) = z_n + \sum_{i=1}^{n-1} z_i \xi_i + \sum_{i=1}^{n-1} a_i(z)\xi_i + \sum_{i=1}^{n+1} z_i b_i(\xi') + g_3(z;\xi'),$$

where a_i, b_i are quadratic and $|g_3(z;\xi')| = O(|z|^3|\xi'| + |x||\xi'|^3))$. Via the changes of variables $\tilde{z}_i = z_i + a_i(z), \tilde{\xi}_i = \xi_i + b_i(\xi')$ for $i \leq n-1$, we have

$$\phi(z;\xi',1) = z_n + \sum_{i=1}^{n-1} z_i \xi_i + z_n b_n(\xi') + z_{n+1} b_{n+1}(\xi') + g_4(z;\xi'),$$

where g_4 satisfies the same condition as g_3 . Again, since ϕ is 1-homogeneous in ξ ,

$$\phi(z;\xi',\xi_n) = z_n(\xi_n + b_n(\xi'/\xi_n)\xi_n) + \sum_{i=1}^{n-1} z_i\xi_i + z_{n+1}b_{n+1}(\xi'/\xi_n)\xi_n + g_4(z;\xi'/\xi_n)\xi_n.$$

Perform the 1-homogeneous change of variables $\tilde{\xi}_n = \xi_n + b_n(\xi'/\xi_n)\xi_n$. Noting that $\xi_n \sim 1$ and since b_n is quadratic, when $|\xi'|$ is sufficiently small, $\tilde{\xi}_n \sim 1$. Thus, we reduce to the form

$$\phi(z;\xi',\xi_n) = \sum_{i=1}^n z_i \xi_i + z_{n+1} \frac{h(\xi')}{\xi_n} + g_5(z;\xi'/\xi_n)\xi_n,$$

where h is quadratic and $g_5(z;\xi') = O(|z|^3|\xi'| + |z||\xi'|^3)$. Finally, as before, we perform a change of variables $\tilde{z}_i = z_i + O(|z|^3)$ $(i \le n)$ to eliminate the $|z|^3|\xi'|$ term in g_5 ; then perform $\tilde{\xi}_i = \xi_i + O(|\xi|^3)$ $(i \le n-1)$ to eliminate the $|x||\xi'|^3$ term in g_5 . Eventually, we can reduce ϕ to the following form (note that $t = z_{n+1}$)

$$\phi(x,t;\xi) = \langle x,\xi \rangle + t \frac{h(\xi')}{\xi_n} + \mathcal{E}(x,t;\xi'/\xi_n)\xi_n,$$

where $\mathcal{E}(x,t;\xi') = O(|z|^2|\xi'|^2 + |t||\xi'|^3)$. This finishes the proof of the claim. Moreover, by condition **(H2)**, we know that $\operatorname{Hess}(h)$ is positive definite. Thus, under appropriate coordinates of the physical variable z = (x, t), by another linear change of variable on ξ' , we can further reduce ϕ to

(5.9)
$$\phi(x,t;\xi) = \langle x,\xi \rangle + t \frac{|\xi'|^2}{\xi_n} + \mathcal{E}(x,t;\xi'/\xi_n)\xi_n.$$

By partitioning the amplitude function in the operator \mathcal{F} a priori, we assume that $|\partial_{t,\xi'}\mathcal{E}(x,t;\xi'/\xi_n)| \leq (10n)^{-10}$ for $\xi \in \operatorname{supp}_{\xi} a$, the ξ -support of the the amplitude function $a(\cdot;\cdot)$. Denote $\widetilde{\mathcal{E}}(x,t;\xi',\xi_n) = \mathcal{E}(x,t;\xi'/\xi_n)\xi_n$, and $\widetilde{\mathcal{E}}^{\lambda}(x,t;\xi',\xi_n) = \lambda \widetilde{\mathcal{E}}(x/\lambda,t/\lambda;\xi',\xi_n)$.

We return to the proof of Lemma 5.4. One can compute

$$\partial_{\xi}\phi(x,t;\xi) = (x' + 2t\frac{\xi'}{\xi_n} + \partial_{\xi'}\widetilde{\mathcal{E}}, x_n - t\frac{|\xi'|^2}{\xi_n^2} + \partial_{\xi_n}\widetilde{\mathcal{E}}).$$

Hence,

$$\partial_{\xi}\phi^{\lambda}(x,t;\xi) = (x' + 2t\frac{\xi'}{\xi_n} + \partial_{\xi'}\widetilde{\mathcal{E}}^{\lambda}, x_n - t\frac{|\xi'|^2}{\xi_n^2} + \partial_{\xi_n}\widetilde{\mathcal{E}}^{\lambda}).$$

Recall (3.7). To find out the core curve of \mathcal{T}_0 , let $(\xi', \xi_n) = (0, 1)$ and solve the equation $\partial_{\xi} \phi^{\lambda}(x, t; 0, 1) = 0$. Since *h* is quadratic and since $\mathcal{E}(x, t; \xi') = O(|z|^2 |\xi'|^2 + |t||\xi'|^3)$, the solution is x = 0. Hence, the core curve of \mathcal{T}_0 is $\{(0, t) : |t| \leq R\}$.

Suppose \mathcal{T} is a tube whose core curve intersects \mathcal{T}_0 at $z_0 = (0, t_0)$. Then there exists ξ' so that the points (x, t) on the core curve of \mathcal{T} satisfy

$$\partial_{\xi}\phi^{\lambda}(x,t;\xi',1) = \partial_{\xi}\phi^{\lambda}(0,t_0;\xi',1)$$

Solve this equation and obtain

(5.10)
$$\begin{aligned} x' &= x'(t, t_0, \xi') = -2(t - t_0)\xi' + \partial_{\xi'} \Big(\widetilde{\mathcal{E}}^{\lambda}(x, t; \xi', 1) - \widetilde{\mathcal{E}}^{\lambda}(x, t_0; \xi', 1) \Big), \\ x_n &= x_n(t, t_0, \xi') = (t - t_0)|\xi'|^2 + \partial_{\xi_n} \Big(\widetilde{\mathcal{E}}^{\lambda}(x, t; \xi', 1) - \widetilde{\mathcal{E}}^{\lambda}(x, t_0; \xi', 1) \Big). \end{aligned}$$

Let \mathcal{Q} be a finite-overlapping cover of $\mathcal{T} \cap \omega_1$ by $\mathbb{R}^{1/2}$ -balls. We may assume the center of each $Q \in \mathcal{Q}$ is of form $(0, \ldots, 0, t_Q)$. Since $\mathcal{T} \cap \mathcal{T}_0 \subset \omega_1$, we assign each $\mathcal{T} \in \mathcal{T}$ to a $Q \in \mathcal{Q}$ such that $\mathcal{T}_0 \cap \mathcal{T} \cap Q \neq \emptyset$. Denote the set of tubes assigned to Q by $\mathcal{T}(Q)$ so that

$$\mathcal{T} = \bigsqcup_{Q} \mathcal{T}(Q).$$

For each tube $\Gamma_B(\xi; R) \in \mathcal{T}(Q)$, we may assume that the core curve of $\Gamma_B(\xi; R)$ intersects the core curve of \mathcal{T}_0 at the center of Q. Fix an arbitrary t_1 such that $\mathbb{R}^n \times \{t_1\} \subset \omega_2$. For each $\mathcal{T} \in \mathcal{T}$, let $x(\mathcal{T}; t_1)$ be the intersection of the core curve of \mathcal{T} with the horizontal hyperplane $\mathbb{R}^n \times \{t_1\}$. By (5.10),

(5.11)
$$x(\mathcal{T};t_1) = (x'(t_1, t_Q, \xi'), x_n(t_1, t_Q, \xi')).$$

Let $B(\mathcal{T};t_1) = B_{R^{1/2}}^n(x(\mathcal{T};t_1))$. It suffices to show that the balls $\{B(\mathcal{T};t_1)\}_{\mathcal{T}\in\mathcal{T}}$ are $(KK_{\circ})^{O(1)}$ -overlapping.

Note that $|\xi| \gtrsim K^{-1}$, since the directions of the tubes in \mathcal{T} and the direction of \mathcal{T}_0 , which is 0, are K^{-1} -separated. Since tubes in \mathcal{T} are distinct, by Lemma 5.3, the points $(R\xi', t_Q)$ corresponding to $x(\mathcal{T}; t_1)$ (as in (5.11)) are $R^{1/2}$ -separated for all $\mathcal{T} \in \mathcal{T}$. Now consider (x', x_n) as a function of (ξ', t_0) as in (5.10). Calculate

(5.12)
$$\partial_{\xi',t_0}(x',x_n) = \begin{pmatrix} -2(t_1-t_0)I_{n-1} & 2(t_1-t_0)(\xi')^T \\ 2\xi' & -|\xi'|^2 \end{pmatrix} + E,$$

where the error term E is a matrix such that

$$E = (10n)^{-10} \begin{pmatrix} O(|t_1 - t_0|) \cdot I_{n-1} & O(|t_1 - t_0| \cdot |\xi|) \cdot (\tilde{I}_{n-1})^T \\ O(|\xi'|^2) \tilde{I}_{n-1} & O(|\xi'|^2) \end{pmatrix}.$$

Here $\vec{1}_{n-1}$ is the vector $(1, \ldots, 1) \in \mathbb{R}^{n-1}$.

Therefore, since $|t_1 - t_Q| \gtrsim RK_{\circ}^{-1}$ and since $|\xi'| \gtrsim K^{-1}$, the Jacobian (5.12) shows that the points $\{x(\mathcal{T};t_1)\}_{\mathcal{T}\in\mathcal{T}}$ are $R^{1/2}(K_{\circ}K)^{-O(1)}$ -separated, which implies that $\{B(\mathcal{T};t_1)\}_{\mathcal{T}\in\mathcal{T}}$ are $(KK_{\circ})^{O(1)}$ -overlapping.

As a corollary, we have the result for planks.

Lemma 5.5. Suppose \mathcal{T}_0 is the $R^{1/2}$ -neighborhood of a R-plank T_0 and $\mathbf{T} = \{T\}$ is a set of distinct R-planks contained in B_R^{n+1} . Let

$$\omega_1 = \mathbb{R}^n \times [i_1 R K_{\circ}^{-1}, (i_1 + 1) R K_{\circ}^{-1}], \ \omega_2 = \mathbb{R}^n \times [i_2 R K_{\circ}^{-1}, (i_2 + 1) R K_{\circ}^{-1}]$$

be two horizontal regions with $i_1, i_2 \in \mathbb{Z}, |i_1|, |i_2| \leq K_{\circ}$ and $|i_1 - i_2| \geq 2$. Suppose for all $T \in \mathbf{T}$, the direction of \mathcal{T}_0 and the direction of T are K^{-1} -separated, and $\mathcal{T}_0 \cap T \neq \emptyset$ with $\mathcal{T}_0 \cap T \subset \omega_1$. Then for all $x \in \omega_2$,

(5.13)
$$\sum_{T \in \mathbf{T}} \mathbf{1}_T(x) \lesssim (KK_\circ)^{O(1)}$$

Proof. Note that we can cover $\cup_{\mathbb{T}}$ by a family of distinct R-tubes such that

- (1) Each *R*-plank *T* is contained in O(1) many R^{δ} -dilate of *R*-tubes.
- (2) For each R-tube, the R-planks inside the R^{δ} -dilate of it are finitely overlapping.

By partitioning \mathcal{T}_0 into $R^{O(\delta)}$ many *R*-tubes, we then apply Lemma 5.4 to conclude the proof.

5.3. A covering lemma. In this subsection, we focus on the geometry inside each $R^{1/2}$ -ball. Although the main result is stated for all dimensions, we will only apply it in the context of 3+1 dimensions.

Suppose T is an R-plank and B is an $R^{1/2}$ -ball. By Taylor's expansion of the core curve of T inside B, one sees that $T \cap B$ is contained in a $R^{2\delta} \times R^{1/2} \times \cdots \times R^{1/2}$ -slab. We will prove a covering lemma for $K^2 \times R^{1/2} \times \cdots \times R^{1/2}$ -slabs inside an $R^{1/2}$ -ball. As before, $K = R^{\varepsilon^{50}}$.

Definition 5.6. Let $R \ge 1$ and let ρ be such that $1 \le \rho \le R^{O(1)}$. Let **B** be a set of disjoint K^2 -balls inside an $R^{1/2}$ -ball $Q \subset \mathbb{R}^{n+1}$. We say **B** is ρ -regular with respect to $K^2 \times R^{1/2} \times \cdots \times R^{1/2}$ -slabs in Q if

(1) For any $K^2 \times R^{1/2} \times \cdots \times R^{1/2}$ -slab S in Q,

 $\#\{B \in \mathbf{B} : B \cap S \neq \emptyset\} \le 2\rho.$

(2) There exists a collection \mathbb{S} of $K^2 \times R^{1/2} \times \cdots \times R^{1/2}$ -slabs with $\#\mathbb{S} \lesssim \#\mathbf{B}/\rho$ so that the balls in \mathbf{B} all intersect with $\bigcup_{S \in \mathbb{S}} S$, and hence are all contained in $\bigcup_{S \in \mathbb{S}} SS$.

When Q is clear and K, R are fixed, we simply call **B** ρ -regular.

Definition 5.6 is similar to Definition 1.9 in [15]. The next Lemma 5.7 is similar to Lemma 2.1 in the same paper.

Lemma 5.7. Suppose **B** is a disjoint union of K^2 -balls inside an $R^{1/2}$ -ball $Q \subset \mathbb{R}^4$. Then there exists a number ρ and a subset $\mathbf{B}' \subset \mathbf{B}$ with $\#\mathbf{B}' \gtrsim (\log R)^{-1} \#\mathbf{B}$, such that \mathbf{B}' is ρ -regular.

Proof. The proof depends on a greedy algorithm. The algorithm is designed to construct a partition of **B** into $O(\log R)$ subsets $\{\mathbf{B}_{\rho}\}_{\rho}$ where ρ are dyadic numbers in $[1, R^2]$, and \mathbf{B}_{ρ} is ρ -regular.

Let $\mathbf{B}_1 = \mathbf{B}$. We will construct a sequence S_1, S_2, \ldots , along with ρ_1, ρ_2, \ldots in the following way. Let $\mathbf{B}_j = \mathbf{B} \setminus \left(\bigcup_{i \leq j-1} \{ B \in \mathbf{B} : B \cap S_i \neq \emptyset \} \right)$. One sees that $\mathbf{B}_j = \mathbf{B}_{j+1} \sqcup \{ B \in \mathbf{B}_j : B \cap S_j \neq \emptyset \}$. Define

$$\rho_j = \max_{\substack{S \text{ is a } K^2 \times R^{1/2} \times \cdots \times R^{1/2} \text{-slab}} \#\{B \in \mathbf{B}_j : B \cap S \neq \emptyset\}.$$

We use S_j to denote the $K^2 \times R^{1/2} \times \cdots \times R^{1/2}$ -slab that attains the maximum.

We keep doing until $\mathbf{B}_{m+1} = \emptyset$ at certain step. By the definition, we have the monotonicity $\rho_i \geq \rho_{i+1}$ for any *i*. For each dyadic $\rho \in [1, \mathbb{R}^{O(1)}]$, we let $I_{\rho} \subset [1, m] \cap \mathbb{N}$ be the indices such that $\rho_j \in [\rho, 2\rho)$ for any $j \in I_{\rho}$. We obtain a partition of indices

$$[1,m] \cap \mathbb{N} = \bigsqcup_{\rho} I_{\rho}.$$

For each dyadic ρ , we inductively define

$$\mathbf{B}_{\rho} = \{ B \in \mathbf{B} : B \cap \bigcup_{i \in I_{\rho}} S_i \neq \emptyset \} \setminus \Big(\bigcup_{\rho' > \rho} \mathbf{B}_{\rho'} \Big).$$

Here, $\bigcup_{\rho'>\rho}$ ranges over those dyadic numbers ρ' bigger that ρ . Thus,

$$\mathbf{B} = \bigsqcup_{\rho} \mathbf{B}_{\rho}.$$

By pigeonholing, there exists ρ such that $\#\mathbf{B}_{\rho} \gtrsim (\log R)^{-1} \#\mathbf{B}$. We claim that \mathbf{B}_{ρ} is ρ -regular. By the definition of ρ_j , we see that for any $K^2 \times R^{1/2} \times \cdots \times R^{1/2}$ -slab $S, \#\{B \in \mathbf{B}_{\rho} : B \cap S \neq \emptyset\} \leq \sup_{i \in I_{\rho}} \rho_i \leq 2\rho$. Note that

$$\mathbf{B}_{\rho} = \bigcup_{j \in I_{\rho}} \mathbf{B}_j \setminus \mathbf{B}_{j+1} = \bigsqcup_{j \in I_{\rho}} \{ B \in \mathbf{B}_j : B \cap S_j \neq \emptyset \}.$$

We have

$$\sum_{i \in I_{\rho}} \rho_i = \sum_{j \in I_{\rho}} \{ B \in \mathbf{B}_j : B \cap S_j \neq \emptyset \} = \# \mathbf{B}_{\rho}.$$

Therefore, $\#I_{\rho} \lesssim \#\mathbf{B}_{\rho}/\rho$, and all the balls in \mathbf{B}_{ρ} intersect with $\bigcup_{i \in I_{\rho}} S_i$.

Remark 5.8. Note that the proof of Lemma 5.7 does not rely on any geometric properties of the $K^2 \times R^{1/2} \times \cdots \times R^{1/2}$ -slabs. That is to say, a similar covering lemma is expected if the family of $K^2 \times R^{1/2} \times \cdots \times R^{1/2}$ -boxes is replaced by another family of geometric objects (for example, a family of tubes, which was used in [15]).

6. An Algorithm

In this section, we present (a single step of) an algorithm designed to establish the broad & two-ends structure on the sum of wave packets $\sum_T \mathcal{F}^{\lambda} f_T$ mentioned in the introduction. Fix $R \geq 1$ and recall our choice of parameters $K = R^{\varepsilon^{50}}, K_{\circ} = R^{\varepsilon^{100}}$. We may assume $R^{1/2}, K, K_{\circ} \in \mathbb{N}$. Let **T** be a family of *R*-planks and **B** be a family of K^2 -balls in B_R^{n+1} . By slightly modify the parameters (up to a constant multiple), we may assume $K^2 |R^{1/2}| R K_{\circ}^{-1}$.

Let $\Omega = \{\omega\}$ be a collection of horizontal regions where each ω is of the form

$$\mathbb{R}^n \times [iRK_{\circ}^{-1}, (i+1)RK_{\circ}^{-1})]$$

where $i \in \mathbb{Z} \cap [-K_{\circ}, K_{\circ} - 1]$), so Ω forms a disjoint cover of B_R^{n+1} . In the algorithm, we will also work with $\{Q\}$, a set of $R^{1/2}$ -balls. We assume each Q is of form

$$\prod_{j=1}^{n+1} [i_j R^{1/2}, (i_j+1)R^{1/2}],$$

where $i_j \in \mathbb{Z} \cap [-R^{1/2}, R^{1/2} - 1]$. We may assume each $B \in \mathbf{B}$ is of form

$$\prod_{j=1}^{n+1} [i_j K^2, (i_j+1)K^2],$$

where $i_j \in \mathbb{Z} \cap [-RK^{-2}, RK^{-2} - 1]$. By our assumption, each B is contained in one Q, and each Q is contained in one ω . Given $T \in \mathbf{T}$, we have

$$\{B \in \mathbf{B} : B \cap T \neq \emptyset\} = \bigsqcup_{\omega \in \Omega} \{B \in \mathbf{B} : B \cap T \neq \emptyset, B \subset \omega\}.$$

Since the two generic regions in Ω are RK_{\circ}^{-1} -separated, the following two statements are conceptually equivalent:

- (1) The K^2 -balls in **B** intersecting T are from many different regions in Ω .
- (2) The shading $Y(T) = \bigcup_{\mathbf{B}} \cap T$ satisfies a two-ends condition.

To formulate the two-ends property, we introduce the following notion.

Definition 6.1 (Shaded incidence triple). Let $R \ge 1$, let $K = R^{\varepsilon^{50}}$, and let Ω be the set of horizontal regions introduced above. A shaded incidence triple (or simply triple) (**B**, **T**; \mathcal{G}) is the following:

- (1) $\mathbf{B} = \{B\}$ is a set of K^2 -balls in B_R^{n+1} ;
- (2) $\mathbf{T} = \{T\}$ is a set of *R*-planks in B_R^{n+1} ;
- (3) \mathcal{G} is the **shading map** that $\mathcal{G} : \mathbf{T} \to P(\Omega)$. Here P(X) is the collection of subsets of X, which is also called the power set of X.

We remark that for a given plank T, $\mathcal{G}(T)$ is a subset of Ω . For a triple, we will be interested in the following incidence:

$$\mathcal{I}(\mathbf{B},\mathbf{T};\mathcal{G}) := \#\{(B,T) \in \mathbf{B} \times \mathbf{T} : B \cap T \neq \emptyset, B \subset \cup_{\mathcal{G}(T)}\}.$$

We also define for any $B \in \mathbf{B}$,

$$\mathbf{T}(B;\mathcal{G}) := \{ T \in \mathbf{T} : B \cap T \neq \emptyset, B \subset \cup_{\mathcal{G}(T)} \},\$$

and for any $T \in \mathbf{T}$,

$$\mathbf{B}(T;\mathcal{G}) := \{ B \in \mathbf{B} : B \cap T \neq \emptyset, B \subset \bigcup_{\mathcal{G}(T)} \}.$$

By a double-counting argument, one has

(6.1)
$$\mathcal{I}(\mathbf{B},\mathbf{T};\mathcal{G}) = \sum_{B \in \mathbf{B}} \#\mathbf{T}(B;\mathcal{G}) = \sum_{T \in \mathbf{T}} \#\mathbf{B}(T;\mathcal{G}).$$

To formulate the broad property, we introduce the following notions.

Definition 6.2. For an *R*-plank *T*, we use $\theta(T) \in \Theta_{R^{-1/2}}$ to denote its directional cap. Let *S* be a union of $\theta \in \Theta_{R^{-1/2}}$ and let **T** be a set of *R*-planks. We define

$$\mathbf{T}[S] := \{ T \in \mathbf{T} : \theta(T) \subset S \}.$$

Here are two examples:

(1) $\mathbf{T}[\theta] = \{T \in \mathbf{T} : \theta(T) = \theta\}.$ (2) For $\tau \in \Theta_{K^{-1}}, \mathbf{T}[(3\tau)^c] = \{T \in \mathbf{T} : \theta(T) \not\subset 3\tau\}.$

Definition 6.3 (Broad incidence). Given a triple $(\mathbf{B}, \mathbf{T}; \mathcal{G})$, define

$$\mathcal{I}_{\mathrm{br}}(\mathbf{B},\mathbf{T};\mathcal{G}):=\inf_{\tau\in\Theta_{K^{-1}}}\mathcal{I}(\mathbf{B},\mathbf{T}[(3\tau)^{c}];\mathcal{G}).$$

For a set of planks \mathbf{T} , we define the broad cardinality as

$$\#_{\operatorname{br}}\mathbf{T} := \min_{\tau \in \Theta_{K^{-1}}} \#\mathbf{T}[(3\tau)^c].$$

Algorithm:

Now we can present the algorithm. It allows us to refine a given triple $(\mathbf{B}, \mathbf{T}, \mathcal{G})$, where \mathbf{T} corresponds to the set of wave packets and \mathbf{B} corresponds to the integration domain. At the end of the algorithm, we will obtain a "refinement" of $(\mathbf{B}, \mathbf{T}, \mathcal{G})$ along with several associated parameters. This will be summarized in Proposition 6.5. We will use the notation: for two finite sets $E \subset F$, we say E is a **refinement** of F if $\#E \gtrsim \#F$.

Initial stage: Given an incidence triple $(\mathbf{B}, \mathbf{T}; \mathcal{G})$ and a $\nu > 0$ such that for $B \in \mathbf{B}$,

(6.2)
$$\| \sum_{T \in \mathbf{T}(B; \Sigma, \mathcal{G})} \mathcal{F}^{\lambda} f_{T} \|_{L^{p}(B)} \sim \nu$$

Step 1: Broad multiplicity in each K^2 -ball $B \in \mathbf{B}$.

By dyadic pigeonholing on $\{\#_{br}\mathbf{T}(B;\mathcal{G}): B \in \mathbf{B}\}$, we can find a refinement \mathbf{B}_1 of **B** and a $\mu \in [1, R^{O(1)}] \cup \{0\}$ such that

(6.3)
$$\begin{aligned} \#_{\mathrm{br}}\mathbf{T}(B;\mathcal{G}) \in [\mu, 2\mu) \text{ when } \mu \geq 1, \\ \#_{\mathrm{br}}\mathbf{T}(B;\mathcal{G}) = 0 \text{ when } \mu = 0, \end{aligned}$$

for all $B \in \mathbf{B}_1$. Note that

(6.4)
$$\mathcal{I}(\mathbf{B}_1, \mathbf{T}; \mathcal{G}) = \sum_{B \in \mathbf{B}_1} \# \mathbf{T}(B; \mathcal{G}) \ge \# \mathbf{B}_1 \mu$$

Step 2: Two-ends reduction

From Step 1, we obtain a triple $(\mathbf{B}_1, \mathbf{T}; \mathcal{G})$. For a plank $T \in \mathbf{T}$ and a dyadic number $1 \leq \lambda \leq R^{O(1)}$, we define

(6.5)
$$\mathcal{G}_{\lambda}(T) := \{ \omega \in \mathcal{G}(T) : \#\{B \in \mathbf{B}_1(T; \mathcal{G}) : B \subset \omega\} \in [\lambda, 2\lambda) \}.$$

Note that \mathcal{G}_{λ} is a new shading map. Since $\mathcal{G}(T) = \bigsqcup_{\lambda} \mathcal{G}_{\lambda}(T)$, we have for each B,

$$\mathbf{T}(B;\mathcal{G}) = \bigsqcup_{\lambda} \mathbf{T}(B;\mathcal{G}_{\lambda}),$$

and hence

$$\sum_{T \in \mathbf{T}(B;\mathcal{G})} \mathcal{F}^{\lambda} f_T = \sum_{\lambda} \sum_{T \in \mathbf{T}(B;\mathcal{G}_{\lambda})} \mathcal{F}^{\lambda} f_T.$$

By pigeonholing, there is a $\lambda(B)$ such that

$$\nu \sim \big\| \sum_{T \in \mathbf{T}(B;\mathcal{G})} \mathcal{F}^{\lambda} f_T \big\|_{L^p(B)} \lesssim \big\| \sum_{T \in \mathbf{T}(B;\mathcal{G}_{\lambda(B)})} \mathcal{F}^{\lambda} f_T \big\|_{L^p(B)}.$$

By dyadic pigeonholing on the tuples

$$\left\{ \left(\lambda(B), \left\| \sum_{T \in \mathbf{T}(B; \mathcal{G}_{\lambda(B)})} \mathcal{F}^{\lambda} f_{T} \right\|_{L^{p}(B)} \right) \right\}_{B \in \mathbf{B}_{1}},$$

there exist \mathbf{B}_2 , λ , and ν_1 , such that the following is true:

- (1) \mathbf{B}_2 is a refinement of \mathbf{B}_1 .
- (2) $\lambda(B) = \lambda$ for all $B \in \mathbf{B}_2$.
- (3) For all $B \in \mathbf{B}_2$, $\|\sum_{T \in \mathbf{T}(B;\mathcal{G}_\lambda)} \mathcal{F}^\lambda f_T\|_{L^p(B)} \sim \nu_1 \gtrsim \nu$.

Next, we will prune the wave packets **T**. Consider the partition $\mathbf{T} = \bigsqcup_{\beta} \mathbf{T}_{\beta}$, where $\beta \in [1, 2K_{\circ}]$ ranges over dyadic numbers and

$$\mathbf{T}_{\beta} = \{T \in \mathbf{T} : \beta \le \# \mathcal{G}_{\lambda}(T) < 2\beta\}$$

For each $B \in \mathbf{B}_2$, we have

(6.6)

$$\mathbf{T}(B;\mathcal{G}_{\lambda}) = \bigsqcup_{\beta} \mathbf{T}_{\beta}(B;\mathcal{G}_{\lambda}).$$

As a result, for each $B \in \mathbf{B}_2$,

$$\sum_{T \in \mathbf{T}(B;\mathcal{G}_{\lambda})} \mathcal{F}^{\lambda} f_{T} = \sum_{\beta} \sum_{T \in \mathbf{T}_{\beta}(B;\mathcal{G}_{\lambda})} \mathcal{F}^{\lambda} f_{T}$$

By pigeonholing, there is a $\beta(B)$ such that

$$\Big\|\sum_{T\in\mathbf{T}(B;\mathcal{G}_{\lambda})}\mathcal{F}^{\lambda}f_{T}\Big\|_{L^{p}(B)} \lesssim \Big\|\sum_{T\in\mathbf{T}_{\beta(B)}(B;\mathcal{G}_{\lambda})}\mathcal{F}^{\lambda}f_{T}\Big\|_{L^{p}(B)}.$$

Since $\|\sum_{T \in \mathbf{T}(B;\mathcal{G}_{\lambda})} \mathcal{F}^{\lambda} f_{T}\|_{L^{p}(B)} \sim \nu_{1}$ for all $B \in \mathbf{B}_{2}$, by a similar dyadic pigeonholing argument as before, there exist \mathbf{B}_{3} , β , and ν_{2} , such that:

- (1) \mathbf{B}_3 is a refinement of \mathbf{B}_2 .
- (2) $\beta(B) = \beta$ for all $B \in \mathbf{B}_3$;
- (3) For all $B \in \mathbf{B}_3$, $\|\sum_{T \in \mathbf{T}_{\beta}(B;\mathcal{G}_{\lambda})} \mathcal{F}^{\lambda} f_T\|_{L^p(B)} \sim \nu_2 \gtrsim \nu_1$.

Note that for each $T \in \mathbf{T}_{\beta}$, one has

$$#\mathbf{B}_{1}(T;\mathcal{G}_{\lambda}) = \#\{B \in \mathbf{B}_{1} : B \cap T \neq \emptyset, B \subset \cup_{\mathcal{G}_{\lambda}(T)}\}\$$
$$= \sum_{\omega \in \mathcal{G}_{\lambda}(T)} \#\{B \in \mathbf{B}_{1} : B \cap T \neq \emptyset, B \subset \omega\}\$$
$$\leq \sum_{\omega \in \mathcal{G}_{\lambda}(T)} \#\{B \in \mathbf{B}_{1}(T;\mathcal{G}) : B \subset \omega\}.$$

By (6.5) and (6.6), $\#\{B \in \mathbf{B}_1(T; \mathcal{G}) : B \subset \omega\} < 2\lambda$ when $\omega \in \mathcal{G}_\lambda(T); \#\mathcal{G}_\lambda(T) < 2\beta$ when $T \in \mathbf{T}_\beta$. Therefore, $\#\mathbf{B}_1(T; \mathcal{G}_\lambda) \lesssim \beta\lambda$ for $T \in \mathbf{T}_\beta$, which gives

(6.7)
$$\mathcal{I}(\mathbf{B}_1, \mathbf{T}_\beta; \mathcal{G}_\lambda) = \sum_{T \in \mathbf{T}_\beta} \# \mathbf{B}_1(T; \mathcal{G}_\lambda) \lesssim \# \mathbf{T}_\beta \cdot \beta \lambda.$$

Step 3: Uniformization on each $R^{1/2}$ -ball

This step is designed for our argument in 3+1 dimensions. The information from this step will not be used in the proof for higher dimensions

Let $\{Q\}$ be a set of $R^{1/2}$ -balls that form a partition of B_R^{n+1} . For each $R^{1/2}$ ball Q, consider the set $\mathbf{B}_3|_Q = \{B \in \mathbf{B}_3 : B \subset Q\}$. By Lemma 5.7, there is a refinement $\mathbf{B}'_3|_Q$ of $\mathbf{B}_3|_Q$ and a number $\rho(Q) \ge 1$ such that $\mathbf{B}'_3|_Q$ is ρ -regular. Let $\mathbf{B}'_3 = \bigcup_Q \mathbf{B}'_3|_Q$, so \mathbf{B}'_3 is a refinement of \mathbf{B}_3 . For each $R^{1/2}$ -ball Q, we consider the following quantities:

- (1) $\rho(Q);$
- (2) $\# \mathbf{B}'_3|_Q;$
- (3) $\#_{\mathrm{br}} \bigcup_{B \in \mathbf{B}_{4}|_{O}} \mathbf{T}_{\beta}(B; \mathcal{G}_{\lambda})$. (Recall $\#_{\mathrm{br}}$ in Definition 6.3.)

By pigeonholing, we can find a set of $R^{1/2}$ -balls $\mathcal{Q} = \{Q\}$ and numbers ρ, σ, m, ι , such that for all $Q \in \mathcal{Q}$ the following is true:

(1) $\rho(Q) \sim \rho$. (2) $\#\mathbf{B}'_{3}|_{Q} \sim \sigma$ (3) $\#_{\mathrm{br}} \bigcup_{B \in \mathbf{B}'_{3}|_{Q}} \mathbf{T}_{\beta}(B; \mathcal{G}_{\lambda}) \sim m$. (4) $\mathcal{I}_{\mathrm{br}}(\mathbf{B}'_{3}|_{Q}, \mathbf{T}; \mathcal{G}) \sim \iota$ (5) $\# \bigcup_{Q \in \mathcal{Q}} \mathbf{B}'_{3}|_{Q} \gtrsim \#\mathbf{B}'_{3}$.

Let \mathbf{B}_4 be a refinement of \mathbf{B}'_3 defined as

$$\mathbf{B}_4 := \bigcup_{Q \in \mathcal{Q}} \mathbf{B}_3'|_Q.$$

We also define an important quantity:

(6.8)
$$l := \sup_{T \in \mathbf{T}_{\beta}} \# \{ Q \in \mathcal{Q} : Q \cap T \neq \emptyset \}.$$

What follows is some numerology regarding the parameters we have introduced. Recall (6.3). Hence, for any $B \in \mathbf{B}_1$ and any $\tau \in \Theta_{K^{-1}}$,

$$\#\mathbf{T}[(3\tau)^c](B;\mathcal{G}) \ge \#_{\mathrm{br}}\mathbf{T}(B;\mathcal{G}) \ge \mu.$$

Note that $\mathbf{B}_4 \subset \mathbf{B}_1$ and $\#\mathbf{B}_4|_Q \sim \sigma$ for all $Q \in \mathcal{Q}$ (here $\mathbf{B}_4|_Q = \{B \in \mathbf{B}_4 : B \subset Q\}$). Therefore, for all $Q \in \mathcal{Q}$,

(6.9)
$$\mathcal{I}_{\mathrm{br}}(\mathbf{B}_{4}|_{Q},\mathbf{T};\mathcal{G}) = \inf_{\tau \in \Theta_{K^{-1}}} \mathcal{I}(\mathbf{B}_{4}|_{Q},\mathbf{T}[(3\tau)^{c}];\mathcal{G})$$
$$= \inf_{\tau \in \Theta_{K^{-1}}} \sum_{B \in \mathbf{B}_{4}|_{Q}} \#\mathbf{T}[(3\tau)^{c}](B;\mathcal{G}) \gtrsim \sigma \mu.$$

(6.9) is our first estimate. To establish our second estimate, note that by (6.1),

$$\mathcal{I}_{\mathrm{br}}(\mathbf{B}_4|_Q, \mathbf{T}_\beta; \mathcal{G}_\lambda) = \inf_{\tau \in \Theta_{K^{-1}}} \sum_{T \in \mathbf{T}_\beta[(3\tau)^c]} \#(\mathbf{B}_4|_Q)(T; \mathcal{G}_\lambda).$$

Thus, for any $\tau \in \Theta_{K^{-1}}$, we have

$$\mathcal{I}_{\mathrm{br}}(\mathbf{B}_{4}|_{Q},\mathbf{T}_{\beta};\mathcal{G}_{\lambda}) \leq \sum_{T\in\mathbf{T}_{\beta}[(3\tau)^{c}]} \#(\mathbf{B}_{4}|_{Q})(T;\mathcal{G}_{\lambda})$$
$$= \sum_{T\in\bigcup_{B\in\mathbf{B}_{4}|_{Q}}\mathbf{T}_{\beta}[(3\tau)^{c}](B;\mathcal{G}_{\lambda})} \#(\mathbf{B}_{4}|_{Q})(T;\mathcal{G}_{\lambda}),$$

as $(\mathbf{B}_4|_Q)(T;\mathcal{G}_\lambda) = \emptyset$ when $T \notin \bigcup_{B \in \mathbf{B}_4|_Q} \mathbf{T}_\beta[(3\tau)^c](B;\mathcal{G}_\lambda)$. Choose τ so that

(6.10)
$$\# \bigcup_{B \in \mathbf{B}_4|_Q} \mathbf{T}_\beta[(3\tau)^c](B; \mathcal{G}_\lambda) = \#_{\mathrm{br}} \bigcup_{B \in \mathbf{B}_4|_Q} \mathbf{T}_\beta(B; \mathcal{G}_\lambda) \sim m.$$

Note that $\#(\mathbf{B}_4|_Q)(T;\mathcal{G}_\lambda) \leq \#\{B \in \mathbf{B}_4|_Q : B \cap T \neq \emptyset\}$. Since $\mathbf{B}_4|_Q$ is ρ -regular,

$$\#(\mathbf{B}_4|_Q)(T;\mathcal{G}_\lambda) \lesssim \rho$$

Therefore, we obtain our second estimate

(6.11)
$$\mathcal{I}_{\mathrm{br}}(\mathbf{B}_4|_Q, \mathbf{T}_\beta; \mathcal{G}_\lambda) \lesssim \rho m.$$

At this point, the algorithm stops.

Before concluding the algorithm as a theorem, we present an incidence result based on the parameters introduced in the algorithm.

Lemma 6.4. Suppose $\beta > 100$. Then

(6.12)
$$\#\mathbf{B}_1 \gtrsim (KK_\circ)^{-O(1)} lm\lambda$$

Proof. By the definition of l in (6.8), there exists $T_0 \in \mathbf{T}_{\beta}$ such that

$$l = \#\{Q \in \mathcal{Q} : Q \cap T_0 \neq \emptyset\}$$

By pigeonholing, there exists $\omega_1 \in \Omega$ such that

$$\#\{Q \in \mathcal{Q} : Q \cap T \neq \emptyset, Q \subset \omega_1\} \gtrsim lK_{\circ}^{-1}.$$

Denote $\mathcal{Q}' = \{Q \in \mathcal{Q} : Q \cap T \neq \emptyset, Q \subset \omega_1\}$. Let \mathcal{T}_0 be the $R^{1/2}$ -neighborhood of the *R*-plank T_0 . Recall (6.10) that $\#_{\mathrm{br}} \bigcup_{B \in \mathbf{B}_4|_Q} \mathbf{T}_{\beta}(B; \mathcal{G}_{\lambda}) \sim m$ for each $Q \in \mathcal{Q}$. Let $\tau \in \Theta_{K^{-1}}$ be the cap containing the direction of \mathcal{T}_0 , so

$$\# \bigcup_{B \in \mathbf{B}_4|_Q} \mathbf{T}_\beta[(3\tau)^c](B; \mathcal{G}_\lambda) \gtrsim m.$$

Now we define

$$\mathbf{T}' = \bigcup_{Q \in \mathcal{Q}'} \bigcup_{B \in \mathbf{B}_4|_Q} \mathbf{T}_\beta[(3\tau)^c](B; \mathcal{G}_\lambda),$$

which can be viewed as a hairbrush with stem \mathcal{T}_0 . Since the directions of planks $T \in \mathbf{T}'$ are K^{-1} -separated with the direction of \mathcal{T}_0 , each $T \in \mathbf{T}'$ belongs to $\leq K^{O(1)}$ different $\{\bigcup_{B \in \mathbf{B}_4|_{\mathcal{O}}} \mathbf{T}_{\beta}[(3\tau)^c](B; \mathcal{G}_{\lambda}) : Q \in \mathcal{Q}'\}$. Therefore,

$$\#\mathbf{T}' \gtrsim K^{-O(1)} \sum_{Q \in \mathcal{Q}'} \# \bigcup_{B \in \mathbf{B}_4|_Q} \mathbf{T}_\beta[(3\tau)^c](B; \mathcal{G}_\lambda) \gtrsim (KK_\circ)^{-1} lm.$$

Next, we consider the set

$$\mathbf{B}' = \bigcup_{T \in \mathbf{T}'} \mathbf{B}_1(T; \mathcal{G}_\lambda).$$

Note that $\mathbf{T}' \subset \mathbf{T}_{\beta}$ and for each $T \in \mathbf{T}_{\beta}, \# \mathcal{G}_{\lambda}(T) \geq \beta > 100$. Recall (6.5). By pigeonholing on the horizontal regions in Ω , we can find a uniform $\omega_2 \in \Omega$ with $\operatorname{dist}(\omega_1,\omega_2) > RK_{\circ}^{-1}$, so that

$$\#\{B \in \mathbf{B}_1(T; \mathcal{G}_\lambda) : B \subset \omega_2\} \ge \lambda$$

holds for $\geq (2K_{\circ})^{-1} \# \mathbf{T}'$ many $T \in \mathbf{T}'$. By Lemma 5.5, $\{T\}_{T \in \mathbf{T}'}$ are $(KK_{\circ})^{O(1)}$ overlapping in ω_2 . As a result, we have

$$#\mathbf{B}' \gtrsim (2K_{\circ})^{-1} #\mathbf{T}'(KK_{\circ})^{-O(1)} \lambda \gtrsim (KK_{\circ})^{-O(1)} lm\lambda.$$

Therefore, as $\mathbf{B}' \subset \mathbf{B}_1$, we have

$$\#\mathbf{B}_1 \gtrsim (KK_\circ)^{-O(1)} lm\lambda.$$

We summarize the algorithm as a Proposition.

Proposition 6.5. Let **T** be a set of R-planks, **B** be a set of K^2 -balls in B_R^{n+1} , and \mathcal{G} be a shading map on **T**. Suppose $\{f_T\}_{T \in \mathbf{T}}$ is a set of wave packets and there exist $\nu > 0$ such that for all $B \in \mathbf{B}$,

$$\Big\|\sum_{T\in\mathbf{T}(\mathbf{B};\mathcal{G})}\mathcal{F}^{\lambda}f_{T}\Big\|_{L^{p}(B)}\sim\nu.$$

Then there exists

- (1) parameters $\mu, \lambda, \beta, \rho, \sigma, m, \iota, l, \nu'$,
- (2) a sequence of refinements $\mathbf{B}_4 \subset \mathbf{B}_3 \subset \mathbf{B}_2 \subset \mathbf{B}_1 \subset \mathbf{B}$,
- (3) a subset of planks $\mathbf{T}_{\beta} \subset \mathbf{T}$, a shading map \mathcal{G}_{λ} , and a set of $\mathbb{R}^{1/2}$ -balls \mathcal{Q} ,

such that $\mathcal{G}_{\lambda}(T) \subset \mathcal{G}(T)$ for all $T \in \mathbf{T}$, and the following is true:

- (a) $\|\sum_{T \in \mathbf{T}_{\beta}(B;\mathcal{G}_{\lambda})} \mathcal{F}^{\lambda} f_{T}\|_{L^{p}(B)} \sim \nu' \gtrsim \nu$ for all $B \in \mathbf{B}_{4}$. (b) $\#\mathbf{B}_{4}|_{Q} \sim \sigma$, and $\mathbf{B}_{4}|_{Q}$ is ρ -regular for all $Q \in \mathcal{Q}$. Here, $\mathbf{B}_{4}|_{Q}$ denotes $\{B \in \mathbf{B}_{4}, B_{4}|_{Q}\}$ $\mathbf{B}_4: B \subset Q\}.$
- (c) $l = \sup_{T \in \mathbf{T}_{\beta}} \# \{ Q \in \mathcal{Q} : Q \cap T \neq \emptyset \}.$

Moreover, we have the following estimates of these parameters:

- (i) $\mathcal{I}(\mathbf{B}_1, \mathbf{T}; \mathcal{G}) \geq \# \mathbf{B}_1 \mu$.
- (*ii*) $\mathcal{I}(\mathbf{B}_1, \mathbf{T}_\beta; \mathcal{G}_\lambda) \lesssim \# \mathbf{T}_\beta \beta \lambda.$
- (iii) $\iota \sim \mathcal{I}_{\mathrm{br}}(\mathbf{B}_4|_Q, \mathbf{T}; \mathcal{G}) \gtrsim \sigma \mu$ for all $Q \in \mathcal{Q}$.
- (iv) $\mathcal{I}_{\mathrm{br}}(\mathbf{B}_4|_Q, \mathbf{T}_{\beta}; \mathcal{G}_{\lambda}) \lesssim \rho m \text{ for each } Q \in \mathcal{Q}.$
- (v) $\beta \leq K_{\circ}$, and if $\beta > 100$, then $\#\mathbf{B}_1 \gtrsim (KK_{\circ})^{-O(1)} lm\lambda$.
- (vi) For all $B \in \mathbf{B}_1$, $\#_{\mathrm{br}}\mathbf{T}(B;\mathcal{G}) \lesssim \mu$.
- (vii) For all $T \in \mathbf{T}_{\beta}$, $\# \mathcal{G}_{\lambda}(T) \lesssim \beta$.

Proof. Item (i) follows from (6.4). Item (ii) follows from (6.7). Item (iii) follows from (6.9). Item (iv) follows from (6.11). Item (v) follows from (6.12). Item (vi) follows from (6.3). Item (vii) follows from (6.6). \square

Remark 6.6. The incidence estimates in Proposition 6.5 are only useful if we can establish the two bounds

- (1) $\mathcal{I}(\mathbf{B}_1, \mathbf{T}_{\beta}; \mathcal{G}_{\lambda}) \gtrsim \mathcal{I}(\mathbf{B}_1, \mathbf{T}; \mathcal{G}).$ (2) $\mathcal{I}_{\mathrm{br}}(\mathbf{B}_4|_Q, \mathbf{T}_{\beta}; \mathcal{G}_{\lambda}) \gtrsim \mathcal{I}_{\mathrm{br}}(\mathbf{B}_4|_Q, \mathbf{T}; \mathcal{G})$ for all $Q \in \mathcal{Q}$.

However, this is generally not true, mostly because the sum in (6.2) involves oscillatory functions rather than positive functions. To achieve these bounds, we will repeatedly run the algorithm. As shown in the next section, after a finite number of iterations, we can essentially realize these two bounds.

7. Finish the proof: iteration of Proposition 6.5

We will prove Theorem 4.10 in this and the next section. For the reader's convenience, we restate it below.

Theorem 7.1. Let \mathcal{F} be a type 1 Fourier integral operator given by (1.5). Suppose the phase function ϕ satisfies (3.1), and the amplitude function a obeys $\operatorname{supp}_{\xi} a \subset \mathbb{A}^n(1)^\circ$. Let \mathcal{F}^{λ} be given by (3.2). Then for all $\varepsilon > 0$ and when p = p(n), there exists a constant C_{ε} that is independent to \mathcal{F} , such that

(7.1)
$$\|\mathcal{F}^{\lambda}f\|_{L^{p}(B^{n+1}_{R})} \leq C_{\varepsilon}R^{(n-1)(\frac{1}{2}-\frac{1}{p})+\varepsilon}\|f\|_{2}^{\frac{2}{p}}\mathcal{W}(f,B^{n+1}_{R})^{1-\frac{2}{p}}.$$

for all f with $\operatorname{supp} \widehat{f} \subset \mathbb{A}^n(1)$ and all $R \in [1, \lambda^{1-\varepsilon}]$.

We will perform a two-parameter inductions on (λ, R) with $1 \leq R \leq \lambda^{1-\varepsilon}$. Our base case is $R \leq 100$, which is not hard to verify. Suppose Theorem 7.1 has been proved for (λ', R') satisfying $\lambda' \leq \lambda, R' \leq R/100$. We now prove for the scale (λ, R) .

Consider the wave packet decomposition for $\mathcal{F}^{\lambda}f$ in B_{R}^{n+1} at scale R:

$$\mathcal{F}^{\lambda}f = \sum_{\theta \in \Theta_{R}^{-1/2}} \sum_{T \in \mathbb{T}_{\theta}} \mathcal{F}^{\lambda}f_{T}$$

Let $w_{B_R^{n+1}}$ be a weight that is ~ 1 on B_R^{n+1} and decreases rapidly outside B_R^{n+1} . By a standard dyadic pigeonholing argument (see, for instance, [22, Section 5]) and homogeneity, we assume that there exists $\mathbb{T} \subset \bigcup_{\theta \in \Theta_{n-1}/2} \mathbb{T}_{\theta}$ such that

(7.2)
$$\|\mathcal{F}^{\lambda}f\|_{L^{p}(B_{R}^{n+1})} \lesssim \|\sum_{T \in \mathbb{T}} \mathcal{F}^{\lambda}f_{T}\|_{L^{p}(B_{R}^{n+1})},$$

and for all $T \in \mathbb{T}$ and $q_n = \frac{2(n+1)}{n-1}$,

(7.3)
$$\|\mathcal{F}^{\lambda}f_{T}\|_{L^{q_{n}}(w_{B_{R}^{n+1}})} \sim R^{\frac{n-1}{4}}.$$

The normalization (7.3) is set because $|T|^{1/q_n} \approx R^{\frac{n+1}{2}\frac{1}{q_n}} = R^{\frac{n-1}{4}}$.

By dyadic pigeonholing, there is a family of disjoint K^2 -balls $\mathcal{B} = \{B\}$ such that $\|\sum_{T \in \mathbb{T}} \mathcal{F}^{\lambda} f_T\|_{L^p(B)}$ are about the same for all $B \in \mathcal{B}$ and

$$\left\|\sum_{T\in\mathbb{T}}\mathcal{F}^{\lambda}f_{T}\right\|_{L^{p}(B_{R}^{n+1})}^{p} \lessapprox \sum_{B\in\mathcal{B}}\left\|\sum_{T\in\mathbb{T}}\mathcal{F}^{\lambda}f_{T}\right\|_{L^{p}(B)}^{p}.$$

Since $\mathcal{F}^{\lambda} f_T$ is essentially supported in T, for each $B \in \mathcal{B}$, we have

$$\|\sum_{T\in\mathbb{T}}\mathcal{F}^{\lambda}f_{T}\|_{L^{p}(B)}\sim\|\sum_{T\in\mathbb{T},T\cap B\neq\emptyset}\mathcal{F}^{\lambda}f_{T}\|_{L^{p}(B)}.$$

Thus, $\|\sum_{T \in \mathbb{T}, T \cap B \neq \emptyset} \mathcal{F}^{\lambda} f_T \|_{L^p(B)}$ are about the same for all $B \in \mathcal{B}$, and

(7.4)
$$\|\sum_{T\in\mathbb{T}}\mathcal{F}^{\lambda}f_{T}\|_{L^{p}(B_{R}^{n+1})}^{p} \lesssim \sum_{B\in\mathcal{B}}\|\sum_{T\in\mathbb{T},T\cap B\neq\emptyset}\mathcal{F}^{\lambda}f_{T}\|_{L^{p}(B)}^{p}$$

For simplicity, we introduce the following notation.

Definition 7.2. Given a triple $(\mathbf{B}, \mathbf{T}; \mathcal{G})$, we denote

$$L^{p}(\mathbf{B},\mathbf{T};\mathcal{G};f) := \sum_{B \in \mathbf{B}} \|\sum_{T \in \mathbf{T}(B;\mathcal{G})} \mathcal{F}^{\lambda} f_{T}\|_{L^{p}(B)}^{p}$$

Let $\mathcal{G}^{(0)}$ be the trivial shading map: $\mathcal{G}^{(0)}(T) = \Omega$. Therefore, $\mathbb{T}(B; \mathcal{G}^{(0)}) = \{T \in \mathbb{C}\}$ $\mathbb{T}: T \cap B \neq \emptyset$. Denote $\mathbb{T}^{(0)} = \mathbb{T}, \mathcal{B}^{(0)} = \mathcal{B}$, so

(7.5) R.H.S. of (7.4) =
$$L^p(\mathcal{B}^{(0)}, \mathbb{T}^{(0)}; \mathcal{G}^{(0)}; f)$$
.

We also introduce the following notion.

Definition 7.3. For two shading maps $\mathcal{G}, \mathcal{G}'$, we write $\mathcal{G} \subset \mathcal{G}'$ if the domain of \mathcal{G} is a subset of the domain of \mathcal{G}' , and $\mathcal{G}(T) \subset \mathcal{G}'(T)$ for any T in the domain of \mathcal{G} . For two triples, we write

$$(\mathbf{B},\mathbf{T};\mathcal{G})\subset (\mathbf{B}',\mathbf{T}';\mathcal{G}')$$

if
$$\mathbf{B} \subset \mathbf{B}', \mathbf{T} \subset \mathbf{T}'$$
, and $\mathcal{G} \subset \mathcal{G}'$.

The triple $(\mathcal{B}^{(0)}, \mathbb{T}^{(0)}; \mathcal{G}^{(0)})$ will be the input of the algorithm given in Section 6. Next, we will repeatedly use Proposition 6.5 to realize the scenario described in Remark 6.6.

Iteration of Proposition 6.5.

At the initial stage (step 0), we are given a $\nu^{(0)} > 0$ and a triple $(\mathcal{B}^{(0)}, \mathbb{T}^{(0)}; \mathcal{G}^{(0)})$ such that for all $B \in \mathcal{B}^{(0)}$,

$$\|\sum_{T\in\mathbb{T}^{(0)}(B;\mathcal{G}^{(0)})}\mathcal{F}^{\lambda}f_{T}\|_{L^{p}(B)}\sim\nu^{(0)}$$

At step i, we apply Proposition 6.5 to the triple $(\mathcal{B}^{(i)}, \mathbb{T}^{(i)}; \mathcal{G}^{(i)})$ and the number $\nu^{(i)}$, which gives the following restatement of Proposition 6.5 in the new notation:

List 1. There exists

- (1) parameters $\mu^{(i)}, \lambda^{(i)}, \beta^{(i)}, \rho^{(i)}, \sigma^{(i)}, m^{(i)}, \iota^{(i)}, l^{(i)}, \nu^{(i+1)}$.

(2) a sequence of refinements $\mathcal{B}_4^{(i)} \subset \mathcal{B}_3^{(i)} \subset \mathcal{B}_2^{(i)} \subset \mathcal{B}_1^{(i)} \subset \mathcal{B}^{(i)}$. (3) a set $\mathbb{T}_{\beta^{(i)}}^{(i)} \subset \mathbb{T}^{(i)}$, a shading map $\mathcal{G}_{\lambda^{(i)}}^{(i)} \subset \mathcal{G}^{(i)}$, and a set of $R^{1/2}$ -balls $\mathcal{Q}^{(i)}$,

such that, by defining

$$\mathcal{B}^{(i+1)} := \mathcal{B}_4^{(i)}, \ \mathbb{T}^{(i+1)} := \mathbb{T}_{\beta^{(i)}}^{(i)}, \ \mathcal{G}^{(i+1)} := \mathcal{G}_{\lambda^{(i)}}^{(i)}$$

which will serve as the input for step i + 1, the following is true:

- (a) $\|\sum_{T \in \mathbb{T}^{(i+1)}(B;\mathcal{G}^{(i+1)})} \mathcal{F}^{\lambda} f_T\|_{L^p(B)} \sim \nu^{(i+1)} \gtrsim \nu^{(i)}$ for all $B \in \mathcal{B}^{(i+1)}$.
- (b) $\#\mathcal{B}^{(i+1)}|_Q \sim \sigma^{(i)}$, and $\mathcal{B}^{(i+1)}|_Q$ is $\rho^{(i)}$ -regular for all $Q \in \mathcal{Q}^{(i)}$. Here $\mathcal{B}^{(i+1)}|_Q = \{B \in \mathcal{B}^{(i+1)} : B \subset Q\}.$
- (c) $\tilde{l}^{(i)} = \sup_{T \in \mathbb{T}^{(i+1)}} \# \{ Q \in \mathcal{Q}^{(i)} : Q \cap T \neq \emptyset \}.$

Moreover, we have the following estimates of these parameters:

- (i) $\mathcal{I}(\mathcal{B}_1^{(i)}, \mathbb{T}^{(i)}; \mathcal{G}^{(i)}) \ge \# \mathcal{B}_1^{(i)} \mu^{(i)}.$

- (i) $\mathcal{I}(\mathcal{D}_1^{(i)}, \mathbb{T}^{(i+1)}; \mathcal{G}^{(i+1)}) \leq \#\mathbb{T}^{(i+1)}\beta^{(i)}\lambda^{(i)}.$ (ii) $\iota^{(i)} \sim \mathcal{I}_{\mathrm{br}}(\mathcal{B}^{(i+1)}|_Q, \mathbb{T}^{(i)}; \mathcal{G}^{(i)}) \gtrsim \sigma^{(i)}\mu^{(i)}$ for all $Q \in \mathcal{Q}^{(i)}.$ (iv) $\mathcal{I}_{\mathrm{br}}(\mathcal{B}^{(i+1)}|_Q, \mathbb{T}^{(i+1)}; \mathcal{G}^{(i+1)}) \leq \rho^{(i)}m^{(i)}$ for all $Q \in \mathcal{Q}^{(i)}.$

- (v) $\beta^{(i)} \leq K_{\circ}$, and if $\beta^{(i)} > 100$, then $\# \mathcal{B}_{1}^{(i)} \gtrsim (KK_{\circ})^{-O(1)} l^{(i)} m^{(i)} \lambda^{(i)}$.
- (vi) For all $B \in \mathbf{B}_1^{(i)}, \#_{\mathrm{br}} \mathbb{T}^{(i)}(B; \mathcal{G}^{(i)}) \lesssim \mu^{(i)}$
- (vii) For all $T \in \mathbb{T}^{(i+1)}$, $\#\mathcal{G}^{(i+1)}(T) \leq \beta^{(i)}$.

The sets and parameters obtained in the above iteration obey certain monotonicity properties. We state some of them that will be used later.

List 2. We have the following monotonicity on parameters:

 $\begin{array}{ll} (\mathrm{i}) \ \ \mu^{(i)} \geq \mu^{(i+1)}, \lambda^{(i)} \geq \lambda^{(i+1)}, \beta^{(i)} \geq \beta^{(i+1)}, \rho^{(i)} \geq \rho^{(i+1)}, \sigma^{(i)} \geq \sigma^{(i+1)}, m^{(i)} \geq m^{(i+1)}, \mu^{(i)} \geq l^{(i+1)}. \end{array}$ (ii) $\nu^{(i+1)} \gtrsim \nu^{(i)}$.

Also, we have the following monotonicity for sets and incidence:

- (1) $(\mathcal{B}^{(i)}, \mathbb{T}^{(i)}; \mathcal{G}^{(i)}) \supset (\mathcal{B}^{(i+1)}, \mathbb{T}^{(i+1)}; \mathcal{G}^{(i+1)}).$
- (2) $\mathcal{Q}^{(i)} \supset \mathcal{Q}^{(i+1)}$.
- (3) $\mathcal{I}(\mathcal{B}^{(i)}, \mathbb{T}^{(i)}; \mathcal{G}^{(i)}) \ge \mathcal{I}(\mathcal{B}^{(i+1)}, \mathbb{T}^{(i+1)}; \mathcal{G}^{(i+1)})$
- $\begin{array}{l} (4) \quad \mathcal{I}_{\mathrm{br}}(\mathcal{B}^{(i)}, \mathbb{T}^{(i)}; \mathcal{G}^{(i)}) \geq \mathcal{I}_{\mathrm{br}}(\mathcal{B}^{(i+1)}, \mathbb{T}^{(i+1)}; \mathcal{G}^{(i+1)}). \\ (5) \quad \#\mathcal{B}^{(i)} \lessapprox \#\mathcal{B}^{(i+1)} \leq \#\mathcal{B}^{(i)}. \end{array}$

As a consequence of (ii) and (5), we have

- (a) $\|\sum_{T \in \mathbb{T}^{(i)}(B;\mathcal{G}^{(i)})} \mathcal{F}^{\lambda} f_T\|_{L^p(B)} \lesssim \|\sum_{T \in \mathbb{T}^{(i+1)}(B;\mathcal{G}^{(i+1)})} \mathcal{F}^{\lambda} f_T\|_{L^p(B)}$ for all $B \in \mathbb{C}$ $\mathcal{B}^{(i+1)}$
- (b) $L^p(\mathcal{B}^{(i)}, \mathbb{T}^{(i)}; \mathcal{G}^{(i)}; f) \lesssim L^p(\mathcal{B}^{(i+1)}, \mathbb{T}^{(i+1)}; \mathcal{G}^{(i+1)}; f).$

At this point, we have completed the description of the iteration.

Now we are going to a uniformization for the parameters obtained in the iteration. For this, we introduce a new parameter $\kappa = R^{\varepsilon^{500}}$. One may compare it with other parameters: $R^{\varepsilon} \gg K \gg K_{\circ} \gg \kappa \gg 1$.

Note that the following 11 factors $\mu^{(i)}, \lambda^{(i)}, \beta^{(i)}, \rho^{(i)}, \sigma^{(i)}, m^{(i)}, \iota^{(i)}, l^{(i)}, \#\mathcal{B}^{(i)}, \mu^{(i)}, \mu^{(i)}$ $\mathbb{T}^{(i)}, \mathcal{I}(\mathcal{B}_1^{(i)}, \mathbb{T}^{(i)}; \mathcal{G}^{(i)})$ are all natural numbers and all $\leq R^{10n}$. Partition $[1, R^{10n}]^{11}$ into $O_{\varepsilon}(1)$ balls of radius κ . Since these factors are monotone decreasing, by pigeonholing, we can find an integer $N \leq \varepsilon^{-O(1)} \lesssim_{\varepsilon} 1$ so that the following is true:

List 3. For i = N - 2, N - 1, N, N + 1, N + 2, we have the reverse control on the parameters:

 $(1) \ \mu^{(i)} \le \kappa \mu^{(i+1)}, \lambda^{(i)} \le \kappa \lambda^{(i+1)}, \beta^{(i)} \le \kappa \beta^{(i+1)}, \rho^{(i)} \le \kappa \rho^{(i+1)}, \sigma^{(i)} \le \kappa \sigma^{(i+1)}, \beta^{(i)} \le \kappa \sigma^{(i)} \le \kappa \sigma^{(i+1)}, \beta^{(i)} \le \kappa \sigma^{(i+1)}, \beta^{(i)} \le \kappa \sigma^{(i+1)}, \beta^$ $\iota^{(i)} \leq \kappa \iota^{(i+1)}, \ m^{(i)} \leq \kappa m^{(i+1)}, \ l^{(i)} \leq \kappa l^{(i+1)}, \ \# \mathcal{B}^{(i)} \leq \kappa \# \mathcal{B}^{(i+1)}, \ \# \mathbb{T}^{(i)} \leq \kappa \# \mathcal{B}^{(i+1)}, \ \# \mathbb{T}^{(i)} \leq \kappa \# \mathcal{B}^{(i+1)}, \ \# \mathbb{T}^{(i)} \leq \kappa \# \mathcal{B}^{(i)} \leq \kappa \# \mathcal{B}^{(i+1)}, \ \# \mathbb{T}^{(i)} \leq \kappa \# \mathcal{B}^{(i)} \leq \kappa \# \mathcal{B}^{(i+1)}, \ \# \mathbb{T}^{(i)} \leq \kappa \# \mathcal{B}^{(i)} \leq \kappa \# \mathcal{B}^{($ $\kappa \mathbb{T}^{(i+1)}, \mathcal{I}(\mathcal{B}_1^{(i)}, \mathbb{T}^{(i)}; \mathcal{G}^{(i)}) \leq \kappa \mathcal{I}(\mathcal{B}_1^{(i+1)}, \mathbb{T}^{(i+1)}; \mathcal{G}^{(i+1)}).$

As a consequence of this and items (i)-(iv) in List 1, when i = N, we have

- (2) $#\mathcal{B}_1^{(i)}\mu^{(i)} \lesssim \kappa^2 #\mathbb{T}^{(i)}\beta^{(i)}\lambda^{(i)}.$
- (3) For all $Q \in Q^{(i)}, \sigma^{(i)}\mu^{(i)} \leq \kappa \rho^{(i)}m^{(i)}$.

This, in particular, implies the following lemma:

Lemma 7.4. Let N be such that List 3 is true for i = N. Then when $\beta^{(N)} > 100$,

$$\mu^{(N)} \lesssim R^{\varepsilon^2} (\#\mathbb{T}^{(N)})^{1/2} (l^{(N)} \sigma^{(N)})^{-1/2} (\rho^{(N)})^{1/2}.$$

Proof. By item (v) in List 1 and item (2) in List 3,

$$(KK_{\circ})^{-O(1)}l^{(N)}m^{(N)}\lambda^{(N)}\mu^{(N)} \lesssim \kappa^2 \# \mathbb{T}^{(N)}\beta^{(N)}\lambda^{(N)}.$$

This concludes the Lemma by item (3) in List 3 and since $(\kappa K K_{\circ})^{O(1)} \lesssim R^{\varepsilon^2}$. \Box

Let us summarize what we have into a theorem.

Theorem 7.5. There exist

(1) parameters $\mu, \beta, \rho, \sigma, l$,

- (2) a triple $(\mathbf{B}, \mathbf{T}, \mathcal{G})$,
- (3) a collection of $\mathbb{R}^{1/2}$ -balls \mathcal{Q} ,

such that the following is true:

(a) $\cup_{\mathbf{B}} \subset \cup_{\mathcal{Q}}$. (b) $\#\mathbf{B}|_Q \sim \sigma$, and $\mathbf{B}|_Q$ is ρ -regular for all $Q \in \mathcal{Q}$. Here $\mathbf{B}|_Q = \{B \in \mathbf{B} : B \subset Q\}$. (c) $l = \sup_{T \in \mathbf{T}} \#\{Q \in \mathcal{Q} : Q \cap T \neq \emptyset\}$.

Moreover, we have the following estimates:

- (i) $\#_{\mathrm{br}} \mathbf{T}(B; \mathcal{G}) \lesssim \mu \text{ for all } B \in \mathbf{B}.$ (ii) $L^p(\mathcal{B}^{(0)}, \mathbb{T}^{(0)}; \mathcal{G}^{(0)}; f) \lesssim L^p(\mathbf{B}, \mathbf{T}; \mathcal{G}; f) \text{ (recall (7.5)).}$
- (iii) $\mathcal{G}(T) \lesssim \beta$ for all $T \in \mathbf{T}$, and if $\beta \ge 100$, $\mu \lesssim R^{\varepsilon^2} (\#\mathbb{T})^{1/2} (l\sigma)^{-1/2} \rho^{1/2}$.

Proof. Let $N \leq_{\varepsilon} 1$ be the natural number obtained in List (3). Take $(\mathbf{B}, \mathbf{T}, \mathcal{G}) = (\mathcal{B}^{(N+1)}, \mathbb{T}^{(N+1)}; \mathcal{G}^{(N+1)})$ and $(\mu, \beta, \rho, \sigma, l, \mathcal{Q}) = (\mu^{(N)}, \beta^{(N)}, \rho^{(N)}, \sigma^{(N)}, l^{(N)}, \mathcal{Q}^{(N)})$. Then item (a) follows by definition, and items (b) and (c) follow from items (b) and (c) in List 1. Items (i), (iii) follow from item (vi), (vii) in List 1 and Lemma 7.4. Finally, item (ii) follows from item (b) in List 2, and the fact that $N \leq_{\varepsilon} 1$. \Box

8. FINISH THE PROOF: CASE STUDY

Let the triple $(\mathbf{B}, \mathbf{T}, \mathcal{G})$, the set of $R^{1/2}$ -balls, and the parameters $(\mu, \beta, \rho, \sigma, l)$ be given by Theorem 7.5. As a consequence of item (ii) in Theorem 7.5 and recall (7.5), (7.4), and (7.2), we have

(8.1)
$$\|\mathcal{F}^{\lambda}f\|_{L^{p}(B_{R}^{n+1})}^{p} \lessapprox \sum_{B \in \mathbf{B}} \|\sum_{T \in \mathbf{T}(B;\mathcal{G})} \mathcal{F}^{\lambda}f_{T}\|_{L^{p}(B)}^{p}$$

We partition each horizontal region $\omega \cap B_R^{n+1}$ ($\omega \in \Omega$) into a collection of R/K_{\circ} balls, resulting in a partition of B_R^{n+1} into R/K_{\circ} -balls, which we denote by $\{B_k\}$. Using this partition, we have

R.H.S. of (8.1) =
$$\sum_{k} \sum_{B \in \mathbf{B}, B \subset B_{k}} \| \sum_{T \in \mathbf{T}(B; \mathcal{G})} \mathcal{F}^{\lambda} f_{T} \|_{L^{p}(B)}^{p}$$
.

Denote

(8.2)
$$\mathbf{T}_k = \{T \in \mathbf{T} : \mathcal{G}(T) \cap B_k \neq \emptyset\},\$$

and let

(8.3)
$$f_k := \sum_{T \in \mathbf{T}_k} f_T.$$

Since $\mathcal{F}^{\lambda} f_T$ is essentially supported in T, when $B \subset B_k$, we have

$$\|\sum_{T\in\mathbf{T}(B;\mathcal{G})}\mathcal{F}^{\lambda}f_{T}\|_{L^{p}(B)} \lesssim \|\mathcal{F}^{\lambda}f_{k}\|_{L^{p}(B)}.$$

Therefore, we obtain

(8.4) R.H.S. of (8.1)
$$\lesssim \sum_{k} \sum_{B \in \mathbf{B}, B \subset B_{k}} \|\mathcal{F}^{\lambda} f_{k}\|_{L^{p}(B)}^{p} \lesssim \sum_{k} \|\mathcal{F}^{\lambda} f_{k}\|_{L^{p}(B_{k})}^{p}.$$

Denote by $\beta = \beta^{(N)}$. We will discuss the following two cases separately:

- (1) **One-end**: $\beta \le 100$.
- (2) **Two-ends**: $\beta > 100$.

In the rest of the section, we prove Theorem 7.1. The numerology of the two cases, n = 3 and $n \ge 4$, differs slightly but follows the same strategy. Therefore, we will address them simultaneously.

8.1. **One-end.** In this subsection, we prove Theorem 7.1 assuming $\beta \leq 100$. Recall Theorem 7.5 item (iii) that $\mathcal{G}(T) \leq \beta \leq 100$ for all $T \in \mathbf{T}$. Since $\beta \leq 100$, each wave packet f_T is included in the summation of at most O(1) different f_k . Thus, by L^2 -orthogonality,

(8.5)
$$\sum_{k} \|f_k\|_2^2 \lesssim \|f\|_2^2.$$

Next, we use the induction hypothesis at the scales $(\lambda, R/K_{\circ})$ and apply (7.1) to each B_k . We note that (7.1) is stated for balls centered at the origin, but a change of coordinates allows the result to hold for balls not necessarily centered at the origin. Suppose $B_k = B_{R/K_{\circ}}^{n+1}(z_k)$. Apply (7.1) to obtain

$$\|\mathcal{F}^{\lambda}f_{k}\|_{L^{p}(B_{k})}^{p} \leq C_{\varepsilon}(\frac{R}{K_{\circ}})^{(n-1)(\frac{p}{2}-1)+p\varepsilon}\|f_{k}\|_{2}^{2}\mathcal{W}(f_{k},B_{R/K_{\circ}}^{n+1}(z_{k}))^{p-2}$$

By Lemma 4.6 in which $r = RK_0^{-1}$, we have

$$\mathcal{W}(f_k, B_{R/K_{\circ}}^{n+1}(z_k)) \lesssim R^{O(\delta)} K_{\circ}^{\frac{n-1}{2}} \mathcal{W}(f_k, B_R^{n+1}).$$

Since the wave packets summed in f_k are a subset of those summed in f, it follows from Lemma 4.3 that

$$\mathcal{W}(f_k, B_R^{n+1}) \lesssim \mathcal{W}(f, B_R^{n+1}).$$

Combining the calculations above and recalling (8.4), we obtain

$$\begin{aligned} \|\mathcal{F}^{\lambda}f\|_{L^{p}(B^{n+1}_{R})}^{p} &\lesssim \sum_{k} C_{\varepsilon}(\frac{R}{K_{\circ}})^{(n-1)(\frac{p}{2}-1)+p\varepsilon} R^{O(\delta)} K_{\circ}^{\frac{n-1}{2}(p-2)} \|f_{k}\|_{2}^{2} \mathcal{W}(f, B^{n+1}_{R})^{p-2} \\ &\lesssim C_{\varepsilon} R^{O(\delta)} K_{\circ}^{-p\varepsilon} R^{(n-1)(\frac{p}{2}-1)+p\varepsilon} \|f\|_{2}^{2} \mathcal{W}(f, B^{n+1}_{R})^{p-2}. \end{aligned}$$

The second inequality follows from (8.5). This closes the induction since $K_{\circ} = R^{\varepsilon^{100}}$ is large enough and hence proves Theorem 7.1.

8.2. Two-ends: broad narrow reduction. By pigeonholing on the R/K_{\circ} balls summed in (8.4), there exists a B_k so that

(8.6)
$$\|\mathcal{F}^{\lambda}f\|_{L^{p}(B_{R}^{n+1})}^{p} \lesssim K_{\circ}^{O(1)} \sum_{B \in \mathbf{B}, B \subset B_{k}} \|\mathcal{F}^{\lambda}f_{k}\|_{L^{p}(B)}^{p}.$$

What follows is a somewhat standard broad-narrow method. Recall the broad norm $\|\cdot\|_{BL_{k-4}^p}^p$ defined in Subsection 3.5. Let

 $k_n = 4$ when n = 3 and $k_n = 2$ when $n \ge 4$.

We remark that the k in the broad norm is irrelevant to the k in B_k .

We choose A = 10000. For any K^2 -ball $B \subset B_k$, let V_1, \ldots, V_A be the set of subspaces of dimension k_n of dimension where the broad norm is attained:

$$\|\mathcal{F}^{\lambda}f_{k}\|_{\mathrm{BL}_{k_{n},A}^{p}(B)}^{p} = \max_{\tau \notin V_{i}, 1 \leq i \leq A} \int_{B} |\mathcal{F}^{\lambda}f_{k,\tau}|^{p}$$

Here " $\tau \notin V_i, 1 \leq i \leq A$ " means " τ ranges over $\tau \in \Theta_{K^{-1}}$ with $\measuredangle(G(\tau), V_i) > K^{-2}$ for all $1 \leq i \leq A$ ". Also, $f_{k,\tau} = \sum_{T \in \mathbf{T}_k[\tau]} f_T$ (recall Definition 6.2).

Therefore, we have

(8.7)
$$\|\mathcal{F}^{\lambda}f_{k}\|_{L^{p}(B)}^{p} \lesssim K^{O(1)}\|\mathcal{F}^{\lambda}f_{k}\|_{\mathrm{BL}_{k_{n},A}^{p}(B)}^{p} + \sum_{i=1}^{A}\int_{B}|\sum_{\tau\in V_{i}}\mathcal{F}^{\lambda}f_{k,\tau}|^{p}.$$

Here " $\tau \in V_i$ " means $\measuredangle(G^\lambda(z_B, \tau), V_i) \le K^{-2}$, where z_B is the center of B.

The following standard lemma provides an upper bound on $\#\{\tau : \tau \in V_i\}$. It appears in various references, for example, in the display below (7.6) in [7]. Nevertheless, we include the proof here for completeness.

Lemma 8.1. For any (k-1)-dimensional subspace $V \subset \mathbb{R}^{n+1}$, then

(8.8)
$$\#\{\tau \in \Theta_{K^{-1}} : \measuredangle(G^{\lambda}(z_B, \tau), V) \le K^{-2}\} \lesssim \max\{1, K^{k-3}\}$$

Proof. Without loss of generality, we assume $z_B = 0$. Via change of variables, we also assume ϕ is of reduced form as in the proof of Lemma 5.4, equation (5.9):

$$\phi(x,t;\xi) = \langle x,\xi \rangle + t \frac{|\xi'|^2}{\xi_n} + \widetilde{\mathcal{E}}(x,t;\xi),$$

where $|\widetilde{\mathcal{E}}(x,t;\xi)| = O(|t||\xi'|^3 + |z|^2|\xi'|^2)$. Applying $\frac{d}{dt}$ to $\partial_{\xi}\phi^{\lambda}(\gamma^{\lambda}(u,t;\xi),t;\xi) = u$, one obtains

$$\partial_{z\xi}^2 \phi^{\lambda} \cdot \left(\frac{d}{dt}\gamma^{\lambda}, 1\right) = 0.$$

Hence, $(\frac{d}{dt}\gamma^{\lambda}(z;\xi),1)$ is parallel to $G^{\lambda}(z;\xi)$. One can compute as in (5.10) that

$$\frac{d}{dt}\gamma^{\lambda}(z_B;\xi',1) = \frac{d}{dt}\gamma^{\lambda}(0;\xi',1) = (-2\xi',|\xi'|^2).$$

Since $(-2\xi', |\xi'|^2, 1)$ is parallel to $G^{\lambda}(z_B; \xi)$, the left-hand side of (8.8) is bounded above by

$$\#\{\tau \in \Theta_{K^{-1}} : (-2\xi'_{\tau}, |\xi'_{\tau}|^2, 1) \in N_{CK^{-2}}(V)\}.$$

Let $\mathcal{S}^{n-1} := \{(-2\xi', |\xi'|^2, 1) : |\xi'| \le 100^{-1}\}.$ Since \mathcal{S}^{n-1} is aways from 0,
 $N_{CK^{-2}}(V) \cap \mathcal{S}^{n-1} \subset N_{C^2K^{-1}}(V \cap \mathcal{S}^{n-1}).$

Since $V \cap S^{n-1}$ is at most k-3 dimensional and since $\{(-2\xi'_{\tau}, |\xi'_{\tau}|^2, 1)\}_{\tau \in \Theta_{K^{-1}}}$ are K^{-1} -separated, we have

$$\#\{\tau \in \Theta_{K^{-1}} : (\xi_{\tau}', |\xi_{\tau}'|^2, 1) \in N_{CK^{-2}}(V)\} \lesssim \max\{1, K^{k-3}\}.$$

By using Theorem A.2 at $R = K^2$, p = p(n), Lemma 8.1 and Hölder's inequality, we can bound the right-hand side of (8.7) by (the rapidly decreasing term can be discarded, as we have done the normalization in (7.3))

$$\sum_{i=1}^{A} \int_{B} \left| \sum_{\tau \in V_{i}} \mathcal{F}^{\lambda} f_{k,\tau} \right|^{p} \lesssim K^{\left(\frac{1}{2} - \frac{1}{p}\right)p} \sum_{\tau \in \Theta_{K^{-1}}} \int_{w_{B}} |\mathcal{F}^{\lambda} f_{k,\tau}|^{p} \text{ when } n = 3,$$

where w_B is a weight that is ~ 1 on B and decreases rapidly outside B. Since $\#\{\tau \in V\} = O(1)$ when V is a one-dimensional subspace, we have

$$\sum_{i=1}^{A} \int_{B} |\sum_{\tau \in V_{i}} \mathcal{F}^{\lambda} f_{k,\tau}|^{p} \lesssim \sum_{\tau \in \Theta_{K^{-1}}} \int_{B} |\mathcal{F}^{\lambda} f_{k,\tau}|^{p} \text{ when } n \ge 4.$$

Denote $X := \bigcup_{B \in \mathbf{B}: B \subset B_k} B$. Summing up all $\{B \in \mathbf{B}: B \subset B_k\}$ in (8.7) recalling (8.6), we have when n = 3,

(8.9)
$$\|\mathcal{F}^{\lambda}f\|_{L^{p}(B_{R}^{4})}^{p} \lesssim (KK_{\circ})^{O(1)} \|\mathcal{F}^{\lambda}f_{k}\|_{\mathrm{BL}_{4,A}^{p}(X)}^{p} + K_{\circ}^{O(1)}K^{(\frac{1}{2}-\frac{1}{p})p} \sum_{\tau} \int_{B_{R}^{4}} |\mathcal{F}^{\lambda}f_{k,\tau}|^{p}$$

and when $n \ge 4$,

(8.10)
$$\|\mathcal{F}^{\lambda}f\|_{L^{p}(B_{R}^{n+1})}^{p} \lesssim (KK_{\circ})^{O(1)} \|\mathcal{F}^{\lambda}f_{k}\|_{\mathrm{BL}_{2,A}^{p}(X)}^{p} + K_{\circ}^{O(1)} \sum_{\tau} \int_{B_{R}^{n+1}} |\mathcal{F}^{\lambda}f_{k,\tau}|^{p}.$$

If the first term on the right-hand side of (8.9) dominates when n = 3 or the first term of (8.10) dominates when $n \ge 4$, we say we are in the **broad case**. If the second term on the right-hand side of (8.9) dominates when n = 3 or the second term of (8.10) dominates when $n \ge 4$, we say we are in the **narrow case**.

8.3. Two-ends: narrow case. Recall (8.9) and (8.10). In this case, we have

$$\begin{aligned} \|\mathcal{F}^{\lambda}f\|_{L^{p}(B_{R}^{4})}^{p} &\lesssim K_{\circ}^{O(1)}K^{(\frac{1}{2}-\frac{1}{p})p}\sum_{\tau}\int_{B_{R}^{4}}|\mathcal{F}^{\lambda}f_{k,\tau}|^{p} \quad \text{when } n=3;\\ \|\mathcal{F}^{\lambda}f\|_{L^{p}(B_{R}^{n+1})}^{p} &\lesssim K_{\circ}^{O(1)}\sum_{\tau}\int_{B_{R}^{n+1}}|\mathcal{F}^{\lambda}f_{k,\tau}|^{p} \quad \text{when } n\geq 4. \end{aligned}$$

For each τ , by Lemma 5.1 and by using the induction hypothesis of Theorem 7.1 at scale R/K^2 , we have

$$\|\mathcal{F}^{\lambda}f_{k,\tau}\|_{L^{p}(B^{n+1}_{R})} \lesssim K^{\frac{2}{p}}(\frac{R}{K^{2}})^{(n-1)(\frac{1}{2}-\frac{1}{p})+\varepsilon} \|f_{k,\tau}\|_{2}^{\frac{2}{p}} \mathcal{W}(f_{k,\tau}, B^{n+1}_{R})^{1-\frac{2}{p}}$$

By Lemma 4.3, we get

$$\mathcal{W}(f_{k,\tau}, B_R^{n+1}) \lesssim \mathcal{W}(f, B_R^{n+1}).$$

Thus, when n = 3, $\|\mathcal{F}^{\lambda}f\|_{L^{p}(B^{4}_{\mathcal{D}})}^{p}$ is

$$\lesssim K_{\circ}^{O(1)} R^{2(\frac{1}{2} - \frac{1}{p})p + p\varepsilon} K^{(\frac{1}{2} - \frac{1}{p})p + 2 - 4(\frac{1}{2} - \frac{1}{p})p - 2p\varepsilon} \sum_{\tau} \|f_{k,\tau}\|_{2}^{2} \mathcal{W}(f, B_{R}^{4})^{p-2}$$

and when $n \geq 4$, $\|\mathcal{F}^{\lambda}f\|_{L^{p}(B^{n+1}_{R})}^{p}$ is

$$\lesssim K_{\circ}^{O(1)} R^{(n-1)(\frac{1}{2} - \frac{1}{p})p + p\varepsilon} K^{2-2(n-1)(\frac{1}{2} - \frac{1}{p})p - 2p\varepsilon} \sum_{\tau} \|f_{k,\tau}\|_{2}^{2} \mathcal{W}(f, B_{R}^{n+1})^{p-2}.$$

Note that when n = 3, we have $(\frac{1}{2} - \frac{1}{p})p + 2 - 4(\frac{1}{2} - \frac{1}{p})p = 0$, as $p = p(3) = \frac{10}{3}$, and when $n \ge 4$, we have $2 - 4(\frac{1}{2} - \frac{1}{p})p \le 0$, as $p = p(n) > 2 + \frac{2}{n-1}$.

Therefore, in both cases,

$$\|\mathcal{F}^{\lambda}f\|_{L^{p}(B^{n+1}_{R})}^{p} \lesssim K_{\circ}^{O(1)}R^{(n-1)(\frac{1}{2}-\frac{1}{p})p+p\varepsilon}K^{-2p\varepsilon}\|f\|_{2}^{2}\mathcal{W}(f,B^{n+1}_{R})^{p-2}.$$

Since $K = R^{\varepsilon^{50}}$ and since $K_{\circ} = R^{\varepsilon^{100}}$, we have $K^{-2p\varepsilon}K_{\circ}^{O(1)} \leq R^{-\varepsilon^{100}}$. Thus, the induction is closed, and we finish the proof of Theorem 7.1.

8.4. Two-ends: broad case. Recall (8.9) and (8.10). In this case, we have

(8.11)
$$\|\mathcal{F}^{\lambda}f\|_{L^{p}(B_{R}^{n+1})}^{p} \lesssim (KK_{\circ})^{O(1)} \|\mathcal{F}^{\lambda}f_{k}\|_{\mathrm{BL}_{k_{n},A}^{p}(X)}^{p},$$

where $k_n = 4$ when n = 3 and $k_n = 2$ when $n \ge 4$.

Let $q_n = \frac{2(n+1)}{n-1}$ be the decoupling exponent. Write $\frac{1}{p(n)} = \alpha(n)\frac{1}{2} + (1-\alpha(n))\frac{1}{q_n}$, so that $\alpha(3) = \frac{1}{5}$ and $\alpha(n) = \frac{n-2}{3n}$ for $n \ge 4$.

By Lemma 3.16, we have

(8.12)
$$\|\mathcal{F}^{\lambda}f_{k}\|_{\mathrm{BL}^{p}_{4,A}(X)} \lesssim \|\mathcal{F}^{\lambda}f_{k}\|_{\mathrm{BL}^{2}_{4,A/2}(X)}^{\frac{1}{5}} \|\mathcal{F}^{\lambda}f_{k}\|_{\mathrm{BL}^{4}_{4,A/2}(X)}^{\frac{4}{5}}$$
 when $n = 3;$

(8.13)
$$\|\mathcal{F}^{\lambda}f_{k}\|_{\mathrm{BL}^{p}_{2,A}(X)} \lesssim \|\mathcal{F}^{\lambda}f_{k}\|_{\mathrm{BL}^{2}_{2,A/2}(X)}^{\alpha(n)} \|\mathcal{F}^{\lambda}f_{k}\|_{\mathrm{BL}^{q_{n}}_{2,A/2}(X)}^{1-\alpha(n)}$$
 when $n \ge 4$.

It remains to estimate $\|\mathcal{F}^{\lambda}f_{k}\|_{\mathrm{BL}^{2}_{4,A/2}(X)}$ and $\|\mathcal{F}^{\lambda}f_{k}\|_{\mathrm{BL}^{4}_{4,A/2}(X)}$ when n = 3 (or $\|\mathcal{F}^{\lambda}f_{k}\|_{\mathrm{BL}^{2}_{2,A/2}(X)}$ and $\|\mathcal{F}^{\lambda}f_{k}\|_{\mathrm{BL}^{q_{n}}_{2,A/2}(X)}$ when $n \geq 4$). Recall (8.3), (8.2), (8.1), and Theorem 7.5.

8.4.1. An L^{q_n} -estimate using refined decoupling. Note that $X = (\cup_{\mathbf{B}}) \cap B_k$. Thus, for each K^2 -ball $B \subset X$, by item (i) in Theorem 7.5, we have

$$\#_{\mathrm{br}}\{T \in \mathbf{T}_k : T \cap B \neq \emptyset\} \le \#_{\mathrm{br}}\mathbf{T}(B;\mathcal{G}) \lesssim \mu$$

Therefore, we can find $\tau' \in \Theta_{K^{-1}}$ such that

$$\#\{T \in \mathbf{T}_k : T \cap B \neq \emptyset, \, \theta(T) \not\subset 3\tau'\} \lesssim \mu$$

Recall $k_n = 4$ when n = 3, $k_n = 2$ when $n \ge 4$. By the definition of broad norm and noting A/2 = 500, we have

$$\|\mathcal{F}^{\lambda}f_k\|_{\mathrm{BL}^{q_n}_{k_n,A/2}(B)}^{q_n} \lesssim K^{O(1)} \max_{\tau \not\subset 3\tau'} \int_B |\mathcal{F}^{\lambda}f_{k,\tau}|^{q_n},$$

where, recall that $f_{k,\tau} = \sum_{T \in \mathbf{T}_k[\tau]} f_T$. Hence, there is a $\tau(B) \not\subset 3\tau'$ such that

(8.14)
$$\|\mathcal{F}^{\lambda}f_{k}\|_{\mathrm{BL}^{q_{n}}_{k_{n},A/2}(B)}^{q_{n}} \lesssim K^{O(1)} \int_{B} |\mathcal{F}^{\lambda}f_{k,\tau(B)}|^{q_{n}},$$

and since $\tau(B) \not\subset 3\tau'$, we have

(8.15)
$$\#\{\mathbf{T}_k: T \cap B \neq \emptyset, \, \theta(T) \subset \tau(B)\} \lesssim \mu$$

By pigeonholing on the set $\{\tau(B)\}_{B \subset X}$, we can find a uniform $\tau \in \Theta_{K^{-1}}$, a union of sub-collection of K^2 -balls $X' \subset X$, such that $\tau(B) = \tau$ for $B \subset X'$ and

$$\|\mathcal{F}^{\lambda}f_{k}\|_{\mathrm{BL}^{q_{n}}_{q_{n},A/2}(X)}^{q_{n}} \lesssim K^{O(1)}\|\mathcal{F}^{\lambda}f_{k}\|_{\mathrm{BL}^{q_{n}}_{q_{n},A/2}(X')}^{q_{n}}$$

Hence, by (8.14), we get

$$\|\mathcal{F}^{\lambda}f_{k}\|_{\mathrm{BL}^{q_{n}}_{q_{n},A/2}(X)}^{q_{n}} \lesssim K^{O(1)} \sum_{B \subset X'} \int_{B} |\mathcal{F}^{\lambda}f_{k,\tau}|^{q_{n}} = K^{O(1)} \|\mathcal{F}^{\lambda}f_{k,\tau}\|_{L^{q_{n}}(X')}^{q_{n}}$$

Recall (7.3) and Lemma 3.18. Since $\mathcal{F}^{\lambda} f_T$ is essentially constant on T, we have

$$R^{\frac{n+1}{2}} \sim \|\mathcal{F}^{\lambda} f_{T}\|_{L^{q_{n}}(w_{B_{R}^{n+1}})}^{q_{n}} \lesssim R^{\frac{n+1}{n-1}} \|\mathcal{F}^{\lambda} f_{T}\|_{L^{2}(w_{B_{R}^{n+1}})}^{q_{n}} \lesssim \|f_{T}\|_{2}^{q_{n}}.$$

Also, by Lemma 4.4, we have

$$\mathcal{W}(f, B_R^{n+1})^2 \gtrsim \max\{1, R^{-\frac{n+1}{2}}(\#\mathbf{T})\} \gtrsim R^{-\frac{n+1}{4}}(\#\mathbf{T})^{1/2}.$$

Therefore, by (8.15) and recalling (7.3), we can apply Theorem A.3 to get

(8.16)
$$\|\mathcal{F}^{\lambda}f_{k}\|_{\mathrm{BL}^{4}_{4,A/2}(X)}^{4} \lesssim K^{O(1)}\|\mathcal{F}^{\lambda}f_{k,\tau}\|_{L^{4}(X')}^{4}$$

$$\lesssim K^{O(1)} \mu \sum_{T \in \mathbf{T}_{k}[\tau]} \| \mathcal{F}^{\lambda} f_{T} \|_{L^{4}(w_{B_{R}^{4}})}^{4} \lesssim K^{O(1)} \mu(\#\mathbf{T} \cdot R^{2})$$
$$\lesssim K^{O(1)} R^{2} \cdot (\mu(\#\mathbf{T})^{-1/2}) \| f \|_{2}^{2} W(f, B_{R}^{4})^{2} \text{ when } n = 3;$$

and also

$$(8.17) \quad \|\mathcal{F}^{\lambda}f_{k}\|_{\mathrm{BL}_{2,A/2}^{q_{n}}(X)}^{q_{n}} \lessapprox K^{O(1)}\|\mathcal{F}^{\lambda}f_{k,\tau}\|_{L^{q_{n}}(X')}^{q_{n}}$$
$$\lesssim K^{O(1)}\mu^{\frac{2}{n-1}}\sum_{T\in\mathbf{T}_{k}[\tau]}\|\mathcal{F}^{\lambda}f_{T}\|_{L^{q_{n}}(w_{B_{R}^{n+1}})}^{q_{n}} \lesssim K^{O(1)}\mu^{\frac{2}{n-1}}(\#\mathbf{T}\cdot R^{\frac{n+1}{2}})$$
$$\lesssim K^{O(1)}R^{2} \cdot (R^{-\frac{n-3}{2(n-1)}}\mu^{\frac{2}{n-1}}(\#\mathbf{T})^{-\frac{1}{n-1}})\|f\|_{2}^{2}W(f, B_{R}^{n+1})^{p_{n}-2} \quad \text{when } n \ge 4$$

Finally, we will finish the proof for n = 3 and $n \ge 4$ separately.

8.4.2. Proof for the case n = 3. We are going to estimate $\|\mathcal{F}^{\lambda}f_k\|_{\mathrm{BL}^2_{4,A/2}(X)}$ inside each $R^{1/2}$ -ball $Q \in \mathcal{Q}$ separately and then sum them up. Recall that $X = \cup_{\mathbf{B}} \cap B_k$. By Theorem 7.5, $X|_Q$ is ρ -regular for all $Q \in \mathcal{Q}$. Thus, for each $Q \in \mathcal{Q}$, there exists a set of $10K^2 \times R^{1/2} \times R^{1/2} \times R^{1/2}$ -slabs $\mathbb{S}(Q)$ with $\#\mathbb{S}(Q) \lesssim \sigma/\rho$ such that $X \cap Q \subset \bigcup_{\mathbb{S}(Q)}$. Thus,

(8.18)
$$\|\mathcal{F}^{\lambda}f_{k}\|_{\mathrm{BL}^{2}_{4,A/2}(X\cap Q)}^{2} \leq \sum_{S\in\mathbb{S}(Q)} \|\mathcal{F}^{\lambda}f_{k}\|_{\mathrm{BL}^{2}_{4,A/2}(S)}^{2}.$$

Fix an $S \in S(Q)$ and let V be the 3-dimensional subspace that is parallel to S. By the definition of the broad norm, we have

$$\|\mathcal{F}^{\lambda}f_k\|_{\mathrm{BL}^2_{4,A/2}(S)}^2 \leq \sum_{B \subset S} \max_{\tau \notin V} \int_B |\mathcal{F}^{\lambda}f_{k,\tau}|^2.$$

By pigeonholing on the caps $\{\tau\}$, there exists a uniform $\tau \notin V$ such that

$$\|\mathcal{F}^{\lambda}f_{k}\|_{\mathrm{BL}^{2}_{4,A/2}(S)}^{2} \lesssim K^{O(1)} \int_{S} |\mathcal{F}^{\lambda}f_{k,\tau}|^{2}.$$

Here $f_{k,\tau} = \sum_{T \in \mathbf{T}_k[\tau]} f_T$. Apply Lemma 3.17 and note that $\mathbf{T}_k[\tau] \subset \mathbf{T}_k \subset \mathbf{T}$. We get

$$\int_{S} |\mathcal{F}^{\lambda} f_{k,\tau}|^2 \lesssim K^{O(1)} \sum_{T \in \mathbf{T}_k[\tau], T \cap Q \neq \emptyset} ||f_T||_2^2 \lesssim K^{O(1)} \sum_{T \in \mathbf{T}, T \cap Q \neq \emptyset} ||f_T||_2^2.$$

Plugging it back to (8.18) and summing up all $Q \in \mathcal{Q}$, we get

$$\begin{aligned} \|\mathcal{F}^{\lambda}f_{k}\|_{\mathrm{BL}^{2}_{4,A/2}(X)}^{2} &\lesssim K^{O(1)}\sum_{Q\in\mathcal{Q}}\sum_{S\in\mathbb{S}(Q)}\sum_{T\in\mathbf{T}:T\cap Q\neq\emptyset}\|f_{T}\|_{2}^{2}\\ &\lesssim K^{O(1)}\#\mathbb{S}(Q)\sum_{T\in\mathbf{T}}\sum_{Q\in\mathcal{Q}:T\cap Q\neq\emptyset}\|f_{T}\|_{2}^{2} \lesssim K^{O(1)}l(\sigma/\rho)\|f\|_{2}^{2}.\end{aligned}$$

In the last inequality, we use item (c) in Theorem 7.5 and L^2 -orthogonality.

Now we can finish the proof. By (8.11), (8.12), (8.16) and (8.18), we get

$$\|\mathcal{F}^{\lambda}f\|_{L^{p}(B_{R}^{4})} \lesssim (KK_{\circ})^{O(1)} R^{2(\frac{1}{2}-\frac{1}{p})} (\mu(\#\mathbf{T})^{-1/2})^{1/5} (l\sigma/\rho)^{1/10} \|f\|_{2}^{\frac{2}{p}} W(f, B_{R}^{4})^{1-\frac{2}{p}}.$$

This implies (7.1) since $(KK_{\circ})^{O(1)} \lesssim R^{\varepsilon^{2}}$ and since $\mu(\#\mathbf{T})^{-1/2} (l\sigma/\rho)^{1/2} \lesssim R^{\varepsilon^{2}},$
which follows from item (iii) in Theorem 7.5.

8.4.3. Proof for the case $n \ge 4$. To estimate $\|\mathcal{F}^{\lambda}f_k\|_{\mathrm{BL}^2_{2,A/2}(X)}^2$, we will simply use the orthogonality on each $R^{1/2}$ -ball Q. We proceed as follows.

(8.19)
$$\|\mathcal{F}^{\lambda}f_{k}\|_{\mathrm{BL}^{2}_{2,A/2}(X)}^{2} = \sum_{Q \in \mathcal{Q}} \|\mathcal{F}^{\lambda}f_{k}\|_{\mathrm{BL}^{2}_{2,A/2}(X \cap Q)}^{2} \leq \sum_{Q \in \mathcal{Q}} \|\mathcal{F}^{\lambda}f_{k}\|_{L^{2}(Q)}^{2}.$$

Note that

$$\int_{Q} |\mathcal{F}^{\lambda} f_{k}|^{2} \lesssim \int_{Q} |\sum_{T \in \mathbf{T}_{k}, T \cap Q \neq \varnothing} f_{T}|^{2}$$

Since $\mathbf{T}_k \subset \mathbf{T}$, apply Lemma 3.18 so that

$$\int_{Q} |\mathcal{F}^{\lambda} f_{k}|^{2} \lesssim R^{1/2} \sum_{T \in \mathbf{T}_{k}, T \cap Q \neq \varnothing} \|f_{T}\|_{2}^{2} \lesssim R^{1/2} \sum_{T \in \mathbf{T}, T \cap Q \neq \varnothing} \|f_{T}\|_{2}^{2}.$$

Plugging it back to (8.19), we get

(8.20)
$$\begin{aligned} \|\mathcal{F}^{\lambda}f_{k}\|_{\mathrm{BL}^{2}_{2,A/2}(X)}^{2} \lesssim R^{1/2} \sum_{Q \in \mathcal{Q}} \sum_{T \in \mathbf{T}: T \cap Q \neq \emptyset} \|f_{T}\|_{2}^{2} \\ \lesssim R^{1/2} \sum_{T \in \mathbf{T}} \sum_{Q \in \mathcal{Q}: T \cap Q \neq \emptyset} \|f_{T}\|_{2}^{2} \\ \lesssim (lR^{1/2}) \|f\|_{2}^{2}. \end{aligned}$$

In the last inequality, we use item (c) in Theorem 7.5 and L^2 -orthogonality.

Now we can finish the proof. By (8.11), (8.13), (8.17), and (8.19), we get (8.21)

$$\|\mathcal{F}^{\lambda}f\|_{L^{p}(B_{R}^{n+1})} \lesssim (KK_{\circ})^{O(1)} R^{(n-1)(\frac{1}{2}-\frac{1}{p})} C(\mu,\mathbf{T},l,R) \|f\|_{2}^{\frac{2}{p}} W(f,B_{R}^{n+1})^{1-\frac{2}{p}},$$

where $C(\mu, \mathbf{T}, l, R)$ equals to

$$(l^{1/2}R^{1/4})^{\alpha(n)}(R^{-\frac{n-3}{2(n-1)}}\mu^{\frac{2}{n-1}}(\#\mathbf{T})^{-\frac{1}{n-1}})^{\frac{1-\alpha(n)}{q_n}}$$

By item (iii) in Theorem 7.5, we have

$$\mu \lesssim R^{\varepsilon^2} (\#\mathbf{T})^{1/2} (l\sigma)^{-1/2} \rho^{1/2} \le R^{\varepsilon^2} (\#\mathbf{T})^{1/2} l^{-1/2},$$

which implies

$$C(\mu, \mathbf{T}, l, R) \lesssim R^{\varepsilon^2} (l^{1/2} R^{1/4})^{\alpha(n)} (R^{-\frac{n-3}{2(n-1)}} l^{-\frac{1}{n-1}})^{\frac{1-\alpha(n)}{q_n}}$$

To conclude (7.1) from (8.21), it remains to show

$$(l^{1/2}R^{1/4})^{\alpha(n)} (R^{-\frac{n-3}{2(n-1)}}l^{-\frac{1}{n-1}})^{\frac{1-\alpha(n)}{q_n}} \leq 1.$$

However, since the power of ℓ is $\frac{\alpha(n)}{2} - \frac{1-\alpha(n)}{2(n+1)} > 0$ with our choice $\alpha(n) = \frac{n-2}{3n}$, this is true as $l \leq R^{1/2}$.

APPENDIX A. DECOUPLING INEQUALITIES

Suppose f(x) is a function in \mathbb{R}^n with $\operatorname{supp} \widehat{f} \subset \mathbb{A}^n(1)$ with the wave packet decomposition inside B_R :

$$\mathcal{F}^{\lambda}f = \sum_{\theta \in \Theta_{R^{-1/2}}} \sum_{T \in \mathbb{T}_{\theta}} \mathcal{F}^{\lambda}f_{T}.$$

Beltran, Hickman, and Sogge [1] proved the following variable coefficient version of $\ell^p L^p$ -decoupling. We remark that $\mathcal{F}^{\lambda} f = T^{\lambda} \hat{f}$, where T^{λ} is the oscillatory integral operator as in [1]. Also, as we will focus exclusively on (n + 1) dimensions in the appendix, we simplify the notation by writing $B_r^{n+1}(z)$ as $B_r(z)$.

Theorem A.1 (ℓ^p -decoupling). Let $2 \leq p \leq \frac{2(n+1)}{n-1}$. For any $\varepsilon > 0$, one has

$$\|\mathcal{F}^{\lambda}f\|_{L^{p}(B_{R})} \lesssim_{\varepsilon, M, \phi, a} R^{\frac{n-1}{2}(\frac{1}{2}-\frac{1}{p})+\varepsilon} (\sum_{\theta \in \Theta_{R}^{-1/2}} \|\mathcal{F}^{\lambda}f_{\theta}\|_{L^{p}(w_{B_{R}})}^{p})^{1/p} + R^{-M} \|f\|_{2}.$$

Here w_{B_R} is a weight that is ~ 1 on B_R and decreases rapidly outside B_R .

A slight modification of the proof in [1] also gives the ℓ^2 -decoupling inequality.

Theorem A.2 (ℓ^2 -decoupling). Let $2 \le p \le \frac{2(n+1)}{n-1}$. For any $\varepsilon > 0$, one has

$$\|\mathcal{F}^{\lambda}f\|_{L^{p}(B_{R})} \lesssim_{\varepsilon, M, \phi, a} R^{\varepsilon} (\sum_{\theta \in \Theta_{R}^{-1/2}} \|\mathcal{F}^{\lambda}f_{\theta}\|_{L^{p}(w_{B_{R}})}^{p})^{1/p} + R^{-M} \|f\|_{2}.$$

In the rest of the appendix, we prove the following refined decoupling inequality, which also implies Theorem A.2. The implication can be found below Theorem 4.2 in [11]. The refined decoupling inequality for the cone was proved in [5, Appendix A], when \mathcal{F}^{λ} is of constant coefficient. Similarly, the proof for the variable coefficient case is based on the ℓ^2 -decoupling theorem and an induction on scales argument.

Theorem A.3. Fix $\nu = \varepsilon^{50}$ and let $K = R^{\nu}$. Suppose we have the wave packet decomposition at scale R in B_B^{n+1} :

$$\mathcal{F}^{\lambda}f = \sum_{T \in \mathbb{W}} \mathcal{F}^{\lambda}f_{T}.$$

Assume that $\|\mathcal{F}^{\lambda}f_{T}\|_{L^{p}(w_{B_{R}})}$ are about the same for all $T \in \mathbb{W}$. Let Y be a disjoint union of K^{2} -balls in B_{R} , each of which intersects $\leq M$ many $T \in \mathbb{W}$. Then for $2 \leq p \leq \frac{2(n+1)}{n-1}$ and any $\varepsilon > 0$,

$$\|\mathcal{F}^{\lambda}f\|_{L^{p}(Y)} \leq C_{\varepsilon,\nu}R^{\varepsilon}\left(\frac{M}{|\mathbb{W}|}\right)^{\frac{1}{2}-\frac{1}{p}}\left(\sum_{T\in\mathbb{W}}\|\mathcal{F}^{\lambda}f_{T}\|_{L^{p}(w_{B_{R}})}^{2}\right)^{1/2} + \operatorname{RapDec}(R)\|f\|_{2}.$$

Since $\|\mathcal{F}^{\lambda}f_{T}\|_{L^{p}(w_{B_{R}})}$ are about the same for all $T \in \mathbb{W}$, we equivalently have

(A.1)
$$\|\mathcal{F}^{\lambda}f\|_{L^{p}(Y)} \leq C_{\varepsilon,\nu}R^{\varepsilon}M^{\frac{1}{2}-\frac{1}{p}}(\sum_{T\in\mathbb{W}}\|\mathcal{F}^{\lambda}f_{T}\|_{L^{p}(w_{B_{R}})}^{p})^{1/2} + \operatorname{RapDec}(R)\|f\|_{2}.$$

Proof. We will prove (A.1) by induction on the scale R. Suppose (A.1) holds for scales $\lambda' \leq \lambda/K^2, R' \leq R/K^2$. Let

$$R_1 = R/K^2 = R^{1-2\nu}$$
 and $K_1 = R_1^{\nu} = R^{\nu-2\nu^2}$.

For $\tau \in \Theta_{K^2}$, we define $\Box_{\tau}^{\flat} = (R/K^{-2})\tau^*$ to be the isotropic dilate of τ^* , a box centered at the origin in \mathbb{R}^n of dimensions $RK^{-2} \times RK^{-1} \times \cdots \times RK^{-1}$. Cover B_R^n by translated copies of \Box_{τ}^{\flat} and denote this cover by $\mathbb{B}_{\tau}^{\flat} = \{\Box^{\flat}\}$. For each \Box^{\flat} , as in (3.10), we define

$$\Box = \Gamma_{\Box^{\flat}}(\xi_{\tau}; R).$$

Let $\mathbb{B}_{\tau} = \{\Box : \Box^{\flat} \in \mathbb{B}_{\tau}^{\flat}\}$, so the fat curved planks in \mathbb{B}_{τ} form a finitely overlapping covering of the n+1 dimensional ball B_R . This induces a partition of wave packets $\mathbb{W} = \bigsqcup_{\tau \in \Theta_{K^2}} \bigsqcup_{\Box \in \mathbb{B}_{\tau}} \mathbb{W}_{\Box}$. Here for each $T \in \mathbb{W}$, we assign T to \mathbb{W}_{\Box} if $T \subset \Box$. In cases where there are multiple valid choices for \Box , we simply choose one.




FIGURE 6. Rescaling

For each $\Box \in \mathbb{B}_{\tau}$, let $\mathcal{Q}_{\Box} = \{Q_{\Box}\}$ be a set of planks that form a finitely overlapping covering of 100 \Box , where each Q_{\Box} is of form

(A.2)
$$\Gamma_W(\bar{z};\xi_\tau;K^2K_1^2)$$

(see (4.2)) for some point $\overline{z} \in 100\square$ and some $K_1^2 \times KK_1^2 \times \cdots \times KK_1^2$ -slab W parallel to $K_1^2\tau^*$, an isotropic dilate of τ^* . The size of Q_{\square} is chosen so that when \square is rescaled to become B_{R_1} , each Q_{\square} becomes a K_1^2 -ball, enabling us to perform

induction. See Figure 6. Let $\{\eta_{Q_{\square}}\}_{Q_{\square} \in Q_{\square}}$ be a partition of unity adapted to $Q_{\square} = \{Q_{\square}\}.$

Write $\mathcal{F}^{\lambda} f$ as

$$\mathcal{F}^{\lambda}f = \sum_{\mathbb{W}} \mathcal{F}^{\lambda}f_{T} = \sum_{\Box} \sum_{T \in \mathbb{W}_{\Box}} \mathcal{F}^{\lambda}f_{T}$$

For each \Box , by dyadic pigeonholing on $\{\#\{T \in \mathbb{W}_{\Box} : T \cap Q_{\Box} \neq \emptyset\} : Q_{\Box} \in \mathcal{Q}_{\Box}\}$, there exists a number $M'(\Box)$ and a set Y_{\Box} , which is a union of Q_{\Box} , such that

(1) $\#\{T \in \mathbb{W}_{\Box} : T \cap Q_{\Box} \neq \emptyset\} \sim M'(\Box) \text{ for all } Q_{\Box} \subset Y_{\Box}.$

(2) We have

(A.3)
$$\|\mathcal{F}^{\lambda}f\|_{L^{p}(Y)} \lesssim (\log R) \left\|\sum_{\Box} \sum_{T \in \mathbb{W}_{\Box}} \eta_{Y_{\Box}} \mathcal{F}^{\lambda}f_{T}\right\|_{L^{p}(Y)}$$

Here, $\eta_{Y_{\square}} = \sum_{Q_{\square} \subset Y_{\square}} \eta_{Q_{\square}}$. By dyadic pigeonholing on $\{M'(\square)\}_{\square}$, there exists a set \mathbb{B} and a uniform number M' such that

(1) $M'(\Box) \sim M'$ for all $\Box \in \mathbb{B}$. (2) We have

(A.4)
$$\left\|\sum_{\Box}\sum_{T\in\mathbb{W}_{\Box}}\eta_{Y_{\Box}}\mathcal{F}^{\lambda}f_{T}\right\|_{L^{p}(Y)} \lesssim (\log R)^{2}\left\|\sum_{\Box\in\mathbb{B}}\sum_{T\in\mathbb{W}_{\Box}}\eta_{Y_{\Box}}\mathcal{F}^{\lambda}f_{T}\right\|_{L^{p}(Y)}.$$

Finally, by dyadic pigeonholing on $\{\#\{Y_{\Box}: Y_{\Box} \cap Q \neq \emptyset, \Box \in \mathbb{B}\} : Q \subset Y\}$, there exists a number M'' and a subset $Y' \subset Y$ such that

(1) $\#\{Y_{\Box}: Y_{\Box} \cap Q \neq \emptyset, \Box \in \mathbb{B}\} \sim M''$ for all $Q \subset Y'$. (2) We have

(A.5)
$$\left\|\sum_{\Box\in\mathbb{B}}\sum_{T\in\mathbb{W}_{\Box}}\eta_{Y_{\Box}}\mathcal{F}^{\lambda}f_{T}\right\|_{L^{p}(Y)}\lesssim (\log R)\right\|\sum_{\Box\in\mathbb{B}}\sum_{T\in\mathbb{W}_{\Box}}\eta_{Y_{\Box}}\mathcal{F}^{\lambda}f_{T}\right\|_{L^{p}(Y')}.$$

We want to estimate the right-hand side of (A.5) using Bourgain-Demeter's ℓ^2 decoupling theorem ([4, Theorem 1.2]), that is, the ℓ^2 decoupling theorem in the constant coefficient setting. This requires us to study the Fourier transform of the function in the right-hand side of (A.5).

For each K^2 -ball $Q \subset Y'$, let $\mathbf{1}_Q^*$ be a smooth bump function adapted to Q, so

(A.6)
$$\left\|\sum_{\square\in\mathbb{B}}\sum_{T\in\mathbb{W}_{\square}}\eta_{Y_{\square}}\mathcal{F}^{\lambda}f_{T}\right\|_{L^{p}(Q)}\lesssim \left\|\mathbf{1}_{Q}^{*}\sum_{\square\in\mathbb{B}}\sum_{T\in\mathbb{W}_{\square}}\eta_{Y_{\square}}\mathcal{F}^{\lambda}f_{T}\right\|_{p}.$$

Suppose $Q = B_{K^2}(z_0)$ is centered at z_0 . For each $\tau \in \Theta_{K^{-1}}$, define the cap

 $P(z_0;\tau):=N_{K^{-2+\delta}}(\{\partial_z\phi^\lambda(z_0;\xi):\xi\in 2\tau\}).$

Here, $\delta = \varepsilon^{1000}$. Note that $\{P(z_0; \tau)\}_{\tau \in \Theta_{K^{-1}}}$ is a set of $1 \times K^{-1} \times \cdots \times K^{-1} \times K^{-2+\delta}$ -caps that form a finite-overlapping cover of the $K^{-2+\delta}$ -neighborhood of the conical surface

$$\Sigma_{z_0} := \{ \partial_z \phi^\lambda(z_0; \xi) : \xi \in \mathbb{A}^n(1) \}.$$

One may view $P(z_0; \tau)$ as the $K^{-2+\delta}$ -neighborhood of the tangent plane of Σ_{z_0} at ξ_{τ} . Note that the normal direction of the tangent plane is $(\partial_t \gamma^{\lambda}(u_0, t_0; \xi_{\tau}, t_0), 1)$,

where $u_0 = \partial_{\xi} \phi^{\lambda}(z_0; \xi_{\tau})$. Thus, we can think of $P(z_0; \tau)$ as

$$\partial_z \phi^{\lambda}(z_0; \xi_{\tau}) + \left(\left\{ \partial_{\xi} \partial_z \phi^{\lambda}(z_0; \xi_{\tau}) \cdot (\xi - \xi_{\tau}) : \xi \in \tau \right\} \\ \oplus \left\{ r(\partial_t \gamma^{\lambda}(u_0, t_0; \xi_{\tau}, t_0), 1) : -K^{-2+\delta} \le r \le K^{-2+\delta} \right\} \right).$$

Here \oplus denotes the orthogonal sum. This shows that a dual box of $P(z_0; \tau)$ is parameterized by

$$\left(\operatorname{Proj}_{\mathbb{R}^n}\left(\left\{\partial_{\xi}\partial_{z}\phi^{\lambda}(z_0;\xi_{\tau})\cdot(\xi-\xi_{\tau}):\xi\in\tau\right\}\right)\right)^* + \left\{r(\partial_{t}\gamma^{\lambda}(u_0,t_0;\xi_{\tau},t_0),1):-K^{2-\delta}\leq r\leq K^{2-\delta}\right\}.$$

Since $\partial_{\xi}\partial_x \phi^{\lambda}(z_0;\xi_{\tau}) \cdot \partial_u \gamma^{\lambda}(u_0,t_0;\xi_{\tau}) = I_n$ and since ϕ obeys the quantitative condition (H1_A), we have

$$\left(\operatorname{Proj}_{\mathbb{R}^n}\left(\left\{\partial_{\xi}\partial_{z}\phi^{\lambda}(z_0;\xi_{\tau})\cdot(\xi-\xi_{\tau}):\xi\in\tau\right\}\right)\right)^* = \left(\left\{\partial_{\xi}\partial_{x}\phi^{\lambda}(z_0;\xi_{\tau})\cdot(\xi-\xi_{\tau}):\xi\in\tau\right\}\right)^* \sim \left\{\partial_{u}\gamma^{\lambda}(u_0,t_0;\xi_{\tau})\cdot u:u\in\tau^*\right\}.$$

Hence, we get

(A.7)
$$P^*(z_0;\tau) \sim \left\{ \partial_u \gamma^{\lambda}(u_0, t_0; \xi_{\tau}) \cdot u : u \in \tau^* \right\} + \left\{ r(\partial_t \gamma^{\lambda}(u_0, t_0; \xi_{\tau}), 1) : -K^{2-\delta} \le r \le K^{2-\delta} \right\}.$$

Lemma A.4. Given $\tau \in \Theta_{K^{-1}}$ and $\Box \in \mathbb{B}_{\tau}$,

$$\mathbf{1}_Q^* \sum_{T \in \mathbb{W}_{\square}} \eta_{Y_{\square}} \mathcal{F}^{\lambda} f_T,$$

as a function in \mathbb{R}^{n+1} , has Fourier transform essentially supported in $2P(z_0; \tau)$.

Proof. Note that, up to a rapidly decreasing term,

$$\mathbf{1}_{Q}^{*}\sum_{T\in\mathbb{W}_{\Box}}\eta_{Y_{\Box}}\mathcal{F}^{\lambda}f_{T}=\mathbf{1}_{Q}^{*}\sum_{Q_{\Box}\subset Y_{\Box},Q_{\Box}\cap 2Q\neq\emptyset}\eta_{Q_{\Box}}\sum_{T\in\mathbb{W}_{\Box}}\mathcal{F}^{\lambda}f_{T}$$

Let $P_0(z_0;\tau)$ be a translated copy of $P(z_0;\tau)$ that is centered at the origin. Since $(\mathbf{1}_Q^*\eta_{Q_{\square}}\mathcal{F}^{\lambda}f_T)^{\wedge} = \widehat{\eta}_{Q_{\square}} * (\mathbf{1}_Q^*\mathcal{F}^{\lambda}f_T)^{\wedge}$, it suffices to show that

(1) $(\mathbf{1}_Q^* \mathcal{F}^{\lambda} f_T)^{\wedge}$ is essentially supported in $P(z_0; \tau)$.

(2) $\hat{\eta}_{Q_{\Box}}$ is essentially supported in $P_0(z_0; \tau)$.

We write

(A.8)
$$(\mathbf{1}_Q^* \mathcal{F}^{\lambda} f_T)^{\wedge}(\eta) = \int \mathbf{1}_Q^* e^{i(\phi^{\lambda}(z;\xi) - \langle z,\eta \rangle)} \mathrm{d}z \int a^{\lambda}(z;\xi) \widehat{f}_T \mathrm{d}\xi$$

Note that \hat{f}_T is supported in $2\theta \subset 2\tau$. Thus, to show item (1), we just need to show that for all $\xi \in 2\tau$,

(A.9)
$$\int \mathbf{1}_Q^* e^{i(\phi^{\lambda}(z;\xi) - \langle z,\eta \rangle)} \mathrm{d}z$$

decays rapidly outside $P(z_0; \tau)$. By stationary phase method, this is true if $|\partial_z \phi^{\lambda}(z; \xi) - \eta| \ge K^{-2+\delta}$ for all $z \in 2Q$. Note that $|\partial_z \phi^{\lambda}(z;\xi) - \partial_z \phi^{\lambda}(z_0;\xi)| \leq |z-z_0|\lambda^{-1} \leq K^{-3}$, as K is a small power of λ . Hence, (A.9) is negligible if

$$|\partial_z \phi^\lambda(z_0;\xi) - \eta| \ge K^{-2+\delta}.$$

Therefore, (A.8) is essentially supported in

$$N_{K^{-2+\delta}}(\{\partial_z\phi^\lambda(z_0;\xi):\xi\in 2\tau\})=P(z_0;\tau).$$

This proves item (1).

Regarding item (2), we study $\eta_{Q_{\square}}$ for $Q_{\square} \cap 2Q \neq \emptyset$. By (A.2), for some slab W,

$$Q_{\Box} = \{ (\gamma^{\lambda}(u + \partial_z \phi^{\lambda}(\bar{z}; \xi_{\tau}), t; \xi_{\tau}), t) : u \in W, |t - \bar{t}| \le K^2 K_1^2 \},\$$

where $\bar{z} = (\bar{x}, \bar{t})$. We first show that Q_{\Box} is morally a box. Let \bar{u} be the center of W. By Taylor's expansion, we have

$$\gamma^{\lambda}(u + \partial_{z}\phi^{\lambda}(\bar{z};\xi_{\tau}),t;\xi_{\tau}) = \gamma^{\lambda}(\bar{u} + \partial_{z}\phi^{\lambda}(\bar{z};\xi_{\tau}),\bar{t};\xi_{\tau})$$
(A.10)

$$+\partial_{u}\gamma^{\lambda}(\bar{u} + \partial_{z}\phi^{\lambda}(\bar{z};\xi_{\tau}),\bar{t};\xi_{\tau}) \cdot (u - \bar{u})$$

$$+\partial_{t}\gamma^{\lambda}(\bar{u} + \partial_{z}\phi^{\lambda}(\bar{z};\xi_{\tau}),\bar{t};\xi_{\tau})(t - \bar{t}) + O(\lambda^{-1}(KK_{1})^{2}).$$

Since $K, K_1 \leq R^{\nu} \leq \lambda^{\nu}, O(\lambda^{-1}(KK_1)^2) = O(1)$. Hence, Q_{\Box} can be approximated by the linear term in (A.10), yielding that Q_{\Box} is morally a box. As a result, the translation of Q_{\Box} to the origin can be essentially parameterized by

$$\begin{split} \bigg\{ \Bigg(\partial_u \gamma^\lambda (\bar{u} + \partial_z \phi^\lambda(\bar{z}; \xi_\tau), \bar{t}; \xi_\tau) \cdot u + \partial_t \gamma^\lambda (\bar{u} + \partial_z \phi^\lambda(\bar{z}; \xi_\tau), \bar{t}; \xi_\tau) t, t \Bigg) \\ &: u \in K_1^2 \tau^*, |t| \le K^2 K_1^2 \bigg\}. \end{split}$$

Again, since $|\bar{u}| \leq K^2$ and $|\bar{z} - z_0| \leq K^2$, by using Taylor's expansion to get rid of the second order term, the above rectangle can be essentially parameterized by

(A.11)
$$\left\{ \left(\partial_u \gamma^{\lambda}(u_0, t_0; \xi_{\tau}) \cdot u + \partial_t \gamma^{\lambda}(u_0, t_0; \xi_{\tau}) t, t \right) : u \in K_1^2 \tau^*, |t| \le K^2 K_1^2 \right\},$$

where replace \bar{u} by 0 and replace \bar{z} by z_0 (recall that $u_0 = \partial_{\xi} \phi^{\lambda}(z_0; \xi_{\tau})$).

To show that $\hat{\eta}_{Q_{\square}}$ is essentially supported in $P_0(z_0; \tau)$, we just need to show that Q_{\square} contains a translated copy of $P_0^*(z_0; \tau)$. In other words, we just need to show (A.11) contains (A.7), which is true.

Now we can use Bourgain-Demeter's ℓ^2 decoupling theorem in Q to have

R.H.S of (A.6)
$$\leq C_{\varepsilon} K^{\varepsilon/100} (M'')^{\frac{1}{2} - \frac{1}{p}} \left(\sum_{\Box \in \mathbb{B}} \left\| \sum_{T \in \mathbb{W}_{\Box}} \eta_{Y_{\Box}} \mathcal{F}^{\lambda} f_{T} \right\|_{L^{p}(\omega_{Q})}^{p} \right)^{1/p}$$
.

Recall (A.3), (A.4), and (A.5). Sum up $Q \subset Y'$ so that

$$\|\mathcal{F}^{\lambda}f\|_{L^{p}(Y)} \lesssim C_{\varepsilon}(\log R)^{100} K^{\varepsilon/100}(M'')^{\frac{1}{2}-\frac{1}{p}} \left(\sum_{\square \in \mathbb{B}} \left\|\sum_{T \in \mathbb{W}_{\square}} \mathcal{F}^{\lambda}f_{T}\right\|_{L^{p}(Y_{\square})}^{p}\right)^{1/p}.$$

Similar to Lemma 5.1, use (A.1) at scale $R_1 = R/K^2$ as an induction hypothesis (see Figure 6) to get

$$\left\|\sum_{T\in\mathbb{W}_{\square}}\mathcal{F}^{\lambda}f_{T}\right\|_{L^{p}(Y_{\square})}^{p} \lesssim C_{\varepsilon,\delta}R^{\varepsilon}K^{-2\varepsilon}\left(\frac{M'}{|\mathbb{W}_{\square}|}\right)^{\frac{1}{2}-\frac{1}{p}}\left(\sum_{T\in\mathbb{W}_{\square}}\|\mathcal{F}^{\lambda}f_{T}\|_{L^{p}(\omega_{B_{R}})}^{2}\right)^{1/2}$$

Therefore, $\|\mathcal{F}^{\lambda}f\|_{L^{p}(Y)}$ is bounded above by

$$C_{\varepsilon}(\log R)^{100}K^{\varepsilon/100}C_{\varepsilon,\delta}R^{\varepsilon}K^{-2\varepsilon}\left(\frac{M'M''}{|\mathbb{W}_{\square}|}\right)^{\frac{1}{2}-\frac{1}{p}}\left(\sum_{\square\in\mathbb{B}}\left(\sum_{T\in\mathbb{W}_{\square}}\|\mathcal{F}^{\lambda}f_{T}\|_{L^{p}(\omega_{B_{R}})}^{2}\right)^{p/2}\right)^{1/p}.$$

Since $\|\mathcal{F}^{\lambda}f_{T}\|_{p}$ are comparable for all $T \in \mathbb{W}$, we thus have that $\|\mathcal{F}^{\lambda}f\|_{L^{p}(Y)}$ is bounded above by

$$C_{\varepsilon}(\log R)^{100}K^{\varepsilon/100}C_{\varepsilon,\delta}R^{\varepsilon}K^{-2\varepsilon}\left(\frac{M'M''}{|\mathbb{W}|}\right)^{\frac{1}{2}-\frac{1}{p}}\left(\frac{|\mathbb{B}||\mathbb{W}_{\square}|}{|\mathbb{W}|}\right)^{\frac{1}{p}}\left(\sum_{T\in\mathbb{W}}\|\mathcal{F}^{\lambda}f_{T}\|_{L^{p}(\omega_{B_{R}})}^{2}\right)^{1/p}$$

Recall $K = R^{\varepsilon^{50}}$. Thus, when R is large enough, $C_{\varepsilon}(\log R)^{100}K^{\varepsilon/100}K^{-2\varepsilon} \leq 1/1000$. Since $|\mathbb{B}||\mathbb{W}_{\Box}| \leq 2|\mathbb{W}|$ and since $M'M'' \leq 4M$, we therefore obtain

$$\|\mathcal{F}^{\lambda}f\|_{L^{p}(Y)} \leq C_{\varepsilon,\delta}R^{\varepsilon}(\frac{M}{|\mathbb{W}|})^{\frac{1}{2}-\frac{1}{p}}(\sum_{T\in\mathbb{W}}\|\mathcal{F}^{\lambda}f_{T}\|_{L^{p}(\omega_{B_{R}})}^{2})^{1/2}.$$

References

- D. Beltran, J. Hickman, and C. D. Sogge. Variable coefficient Wolff-type inequalities and sharp local smoothing estimates for wave equations on manifolds. *Anal. PDE*, 13(2):403–433, 2020.
- [2] D. Beltran, J. Hickman, and C. D. Sogge. Sharp local smoothing estimates for Fourier integral operators. In *Geometric aspects of harmonic analysis*, volume 45 of *Springer INdAM Ser.*, pages 29–105. Springer, Cham, [2021] ©2021.
- [3] J. Bourgain. L^p-estimates for oscillatory integrals in several variables. Geom. Funct. Anal., 1(4):321–374, 1991.
- [4] J. Bourgain and C. Demeter. The proof of the l² decoupling conjecture. Ann. of Math. (2), 182(1):351–389, 2015.
- [5] S. Gan, S. Guo, L. Guth, T. L. J. Harris, D. Maldague, and H. Wang. On restricted projections to planes in R³. To appear in American Journal of Mathematics, 2022.
- [6] S. Gan, C. Oh, and S. Wu. A note on local smoothing estimates for fractional Schrödinger equations. J. Funct. Anal., 283(5):Paper No. 109558, 36, 2022.
- [7] C. Gao, B. Liu, C. Miao, and Y. Xi. Improved local smoothing estimate for the wave equation in higher dimensions. J. Funct. Anal., 284(9):Paper No. 109879, 48, 2023.
- [8] C. Gao, B. Liu, C. Miao, and Y. Xi. Square function estimates and local smoothing for Fourier integral operators. Proc. Lond. Math. Soc. (3), 126(6):1923–1960, 2023.
- [9] L. Guth. A restriction estimate using polynomial partitioning. J. Amer. Math. Soc., 29(2):371-413, 2016.
- [10] L. Guth, J. Hickman, and M. Iliopoulou. Sharp estimates for oscillatory integral operators via polynomial partitioning. Acta Math., 223(2):251–376, 2019.
- [11] L. Guth, A. Iosevich, Y. Ou, and H. Wang. On Falconer's distance set problem in the plane. *Invent. Math.*, 219(3):779–830, 2020.
- [12] L. Guth, H. Wang, and R. Zhang. A sharp square function estimate for the cone in ℝ³. Ann. of Math. (2), 192(2):551–581, 2020.
- [13] Y. Heo, F. Nazarov, and A. Seeger. Radial Fourier multipliers in high dimensions. Acta Math., 206(1):55–92, 2011.
- [14] N. H. Katz, S. Wu, and J. Zahl. Kakeya sets from lines in SL₂. Ars Inven. Anal., pages Paper No. 6, 23, 2023.

- [15] X. Li and S. Wu. On almost everywhere convergence of planar Bochner-Riesz mean. arXiv:2407.20887, 2024.
- [16] W. P. Minicozzi, II and C. D. Sogge. Negative results for Nikodym maximal functions and related oscillatory integrals in curved space. *Math. Res. Lett.*, 4(2-3):221–237, 1997.
- [17] J. C. Peral. L^p estimates for the wave equation. J. Functional Analysis, 36(1):114–145, 1980.
- [18] K. M. Rogers. A local smoothing estimate for the Schrödinger equation. Adv. Math., 219(6):2105–2122, 2008.
- [19] A. Seeger, C. D. Sogge, and E. M. Stein. Regularity properties of Fourier integral operators. Ann. of Math. (2), 134(2):231–251, 1991.
- [20] C. D. Sogge. Propagation of singularities and maximal functions in the plane. Invent. Math., 104(2):349–376, 1991.
- [21] T. Tao. The Bochner-Riesz conjecture implies the restriction conjecture. Duke Math. J., 96(2):363–375, 1999.
- [22] H. Wang and S. Wu. Restriction estimates using decoupling theorems and two-ends Furstenberg inequalities. arXiv:2411.08871, 2024.
- [23] T. Wolff. Local smoothing type estimates on L^p for large p. Geom. Funct. Anal., 10(5):1237–1288, 2000.

Shengwen Gan, Department of Mathematics, University of Wisconsin-Madison, USA $\mathit{Email}\ address: \tt sgan7@wisc.edu$

SHUKUN WU, DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY BLOOMINGTON, USA *Email address:* shukwu@iu.edu