# BOUNDARY FRAMINGS FOR lcs FOUR-MANIFOLDS

#### J MORAVA

ABSTRACT. In work with Urs Schreiber [§1] we construct a rational homotopytheoretic model for a classifying space of locally conformally symplectic (lcs) structures on four-manifolds. This leads to the definition of a cobordism category of three-manifolds 'anchored' by principal  $\Omega^2 S^2$  - bundles (§2, generalizing contact structures). Powerful  $\mathfrak{sl}_2\mathbb{R}$  - representation-valued Hodge-Lefschetz cohomology [3, 31, 32], going back to Chern and Weil), taking values in the  $\mathbb{Z}$ -graded bidifferential category  $\mathfrak{B}^*$  of [1](lemma 1.3), is available for its study.

§3 observes that David Ebin's [4] configuration groupoid [Metrics//D] for Euclidean general relativity can be represented (cohomologically over  $\mathbb{Q}$ ) in terms of a category

$$\mathbb{A}(X) := \int_{K \in [\mathbb{C}\mathrm{cpct}_0 / / \mathbb{D}]} [X / / K] \to \mathcal{B}^*$$

of objects (or 'aspects') of X (as a space, under the action of its group  $\mathbb{D}$  of (pointed, etc) diffeomorphisms). This provides a notational system for the higher category of actions of compact connected groups K on X, which is classically accessible by Atiyah-Bott/Duistermaat-Heckmann [2] ...

## §I A rational homotopy-theoretic classifying space for lcs four-manifolds

**1.1** In the following,  $X \supset \partial X = Y$  will be a connected compact oriented fourdimensional manifold with (for now) connected boundary, together with the locally conformally symplectic structure [5, 15, 27] defined by a two-form  $\omega \in \Omega^2_{\mathbb{R}}(X)$ , which is **effective** in that

★  $(\forall x \in X) (\omega \land \omega)(x) \neq 0$ , and moreover there is a (necessarily unique) [18] one-form  $\vartheta$  such that

•  $d\vartheta = 0$  with  $d\omega = \vartheta \wedge \omega$ .

Working in a closed model category of differential commutative graded  $\mathbb{Q}$ -algebras, we construct a commutative diagram ( $\omega \mapsto \omega_2, \vartheta \to \omega_1$ )

with dotted arrows signifying cofibrant (minimal model) replacements, following Neisendorfer  $[26](\S5.6a)$  and Meier  $[19](\S2.3)$ .

J MORAVA

I am indebted to Urs Schreiber for this – these dcas are **not** nilpotent [16][Def 1.10], but  $\pi_1 = \mathbb{Z}$  has cohomological dimension one, which suffices for this construction – and, more generally, for drawing my attention to the utility of rational-homotopytheoretic methods, eg [10](ex 2.41), [12](§4.6, open M5 branes), [34] in mathematical physics. I also owe him thanks for his help and patience, but the mistakes here are my responsibility.

 $1.2~\mathrm{An}\ \mathsf{lcs}\ \mathrm{structure}\ \mathrm{on}\ X$  defines a homomorphism

$$\Omega^* \operatorname{Blcs} \to \Omega^*_{\mathbb{R}}(X)$$

making the real differential forms on X into an effective  $\Omega^*$ Blcs algebra.

 $\mathbf{Prop}\ A\ collar\ (0,1]\times Y\subset X$  for the boundary  $\partial X=Y$  defines a commutative diagram



presenting an  $\Omega^* \Omega S^2$  - algebra structure on  $\Omega^* Y$  as boundary data for a  $\Omega^* Blcs$  structure on  $\Omega^* X$ .

[The right vertical map is defined by restriction. Note that  $\Omega^*$  signifies an algebra of differential forms, whereas  $\Omega$  indicates a based loop space.]

\* **1.3** Recall [5, 21] that a contact form ( $\alpha \in \Omega^1 Y$  such that  $(\forall x) \ \alpha \land d\alpha \neq 0$ , *cf* **a3**) defines an effective  $\Omega^*$ Blcs - algebra structure on  $\Omega^*(I \times Y)$ , by

$$\omega_2 = -t^{-1}d(t\alpha), \ \omega_1 = t^{-1}dt = \vartheta \in \mathbf{\Omega}^1 \mathbf{I};$$

indeed

$$d\omega_2 = \vartheta \wedge \omega_2, \text{ and } \omega_2 \wedge \omega_2 = -2\alpha \wedge d\alpha \wedge \vartheta \neq 0$$

**1.4** We will return to contact forms in  $[\mathbf{a2}]$  below, but for the moment recall [29] that a contact form has both an associated real  $[\xi] \in \operatorname{Pic}_{\mathbb{R}} Y \cong H^{1}(Y, \mathbb{Z}_{2})$  line bundle as well as a complex  $[\eta] \in H^{2}(Y, \mathbb{Z}_{2}) \cong \operatorname{Pic}_{\mathbb{C}}(Y)$  (related by  $[\xi] \otimes_{\mathbb{R}} [\eta] = [\eta]$ ), and that an almost contact structure is a reduction of the structure group of Y to

$$\mathbb{T} \rtimes \mathbb{Z}_2 \subset \mathrm{SU}(2) = \mathrm{Spin}(3)$$
.

The class  $[\vartheta] \in H^1(Y, \mathbb{R})$  can perhaps be regarded as a lift of  $[\xi]$ .

### **§II** Spin, contact, and line structures

2.0 An almost contact structure on Y as above is rationally indistinguishable from an  $\Omega^*\Omega S^2$  - algebra structure, but this loses two-torsion information. We argue here for working in a topos

 $\{X / |\operatorname{Fin}_{\mathbb{C}}^{\otimes}|^+\}$ 

of spaces with enough basepoints – alternatively, spaces endowed with a principal line or  $\Omega^2 S^2 \simeq \Omega^2 SU(2) \rtimes \mathbb{Z}$  bundle<sup>1</sup> – as a rationally equivalent alternative. Such

<sup>&</sup>lt;sup>1</sup>a noncommutative harmonic oscillator [23]

framings/orientations/basepoints were proposed in [22], as useful in renormalization theory.

2.1 Recall that under Pontrjagin multiplication, the integral homology

$$\mathsf{H}_*(\Omega S^2, \mathbb{Z}) \cong \mathbb{Z}[\mathbf{x}]$$

is polynomial on a single generator  $\mathbf{x}$  of degree **one**, and is therefore **not** a (graded-) commutative algebra. The cohomology

$$\mathsf{H}^*(\Omega \mathsf{S}^2, \mathbb{Z}) \cong \mathsf{E}([\omega_1]) \otimes \mathsf{P}([\omega_2])$$

is however graded - commutative, with  $[\omega_2]$  Kronecker dual to  $\mathbf{x}^2$ . There is further background in appendix  $[\mathbf{a1}]$  below.]

**Definition** The map

$$\mathbb{C}^{\times} \ni z \mapsto [\mathsf{q} \to z\mathsf{q} z^{-1}] \in \mathrm{Maps}_{\{1\}}(\mathbb{H}^{\times}, \mathbb{H}^{\times})$$

extends by collapse maps when  $z \in \{0, \infty\}$  to define

$$\mathbb{P}^1(\mathbb{C}) = \mathbb{C}^{\times} \cup \{0, \infty\} \to \operatorname{Maps}_{\operatorname{deg}=1}(S^3, S^3) ,$$

(*ie* if  $z = \operatorname{rexp}(i\alpha)$  with  $r \in \mathbb{R}^{\times}$  then  $z \to [q \to e^{i\alpha}qe^{-i\alpha}]$ , but if  $r = 0, \infty$  it defines a Mercator-like compactification, with fixed points at the north and south poles).

Taking adjoints gives a loop map [a3]

$$[S^3 \to \operatorname{Maps}(\mathbb{P}^1(\mathbb{C}), S^3) \cong \Omega^2 S^3] = \eta^2 \in \pi_5 S^3 \cong \mathbb{Z}_2$$

and thus a composition

$$B\eta^2 : BSU(2) \to \Omega S^3 \to \Omega S^2$$

(the last map being the universal cover).

IIUC the composition  $\mathbb{T} \rtimes \mathbb{Z}_2 \to \mathrm{SU}(2) \to \Omega^2 S^2$  therefore assigns a line bundle to an almost contact structure via the remarkable relation

$$\eta^3 = 4\nu$$

of Ravenel [28](1.1.14). Note that  $\eta^2$  is the first nontrivial element in the cokernel of stable J.

2.2 A spin structure on X thus has compatible associated line bundles

on both X and its boundary.

More generally, our topos

$$\{X \mid |\operatorname{Fin}_{\mathbb{C}}^{\otimes}|^+\} \to \{X \mid \operatorname{BSpin}\}$$

lies behind that of manifolds with spin structures, which already has a rich deformation theory [17].

#### J MORAVA

**2.3 Proposition** (Compact) four-manifolds with effective  $\Omega^*$ Blcs - algebra structures as morphisms, and three-manifolds with compatible (cf §1.2-3) boundary line bundles as objects, form a cobordism (topological 3-) category.

[More precisely, its morphism objects  $\operatorname{Mor}_*(Y', Y)$  are the topological groupoids with lcs cobordisms  $W \supset \partial W = Y'_{\operatorname{op}} \sqcup Y \to \Omega S^2$  as objects, and diffeomorphisms preserving the line and lcs structure [15](Theorem 3.7) as morphisms. In particular, its morphisms compose under glueing along compatible boundaries.]

The effectiveness of an  $\Omega^*$ Blcs-structure permits the construction of a symplectic Hodge duality operator and so [3, 31, 32] leads to an  $\mathfrak{sl}_2\mathbb{R}$  - representation valued Hodge-Lefschetz bidifferential cohomology theory [1] with good finiteness properties.

### **§III** Aspects of manifolds

**3.0** Recall that a presheaf  $\Phi : \mathbb{C}^{\mathrm{op}} \to \mathcal{D}$  defines Grothendieck's category  $\int_{\Phi} \mathcal{D} \to \mathbb{C}$  (with objects  $(C \in \mathbb{C}, c \in C)$  and morphisms  $\mathfrak{u} : (C, c) \to (C', c')$  defined by functors  $\mathfrak{u} : C' \to C$  such that  $\Phi(\mathfrak{u})(c) = c'$ ).

If, for example, X is a compact connected manifold and  $\mathbb{D}$  is the topological group [11, 15] of its smooth based diffeomorphisms (respecting the boundary  $\partial X \to \pi_0 \partial X$ ), let

$$\operatorname{Cpct}_0/\mathbb{D} := [\operatorname{Cpct}_0 \subset \mathbb{D}\}//\mathbb{D}^{\operatorname{conj}}]$$

be its category of compact connected subgroups under conjugation by elements of  $\mathbb D.$  Integration

$$\begin{split} \mathbb{A}(X) &:= \int_{K \in \mathfrak{C}\mathrm{pct}_0/\mathbb{D}} [X / \! / K] \\ & \bigvee \\ \mathfrak{C}\mathrm{pct}_0 / \mathbb{D} \end{split}$$

along the presheaf  $K \to [X/\!/K]$  of groupoids then defines a (higher, fibered) category of 'aspects' of X.

Such a thing is an adéle-like element - for example, a fixed point - of X, but with X indexed by or regarded as a space with (the conjugacy class of) an action of a compact connected group K upon it.

But actions of compact groups on smooth manifolds is a classical subject accessible via localization [2] in characteristic zero, and via tom Dieck/Elmendorf in (Spectra). It can be topologized [32] and equivariant cohomology functors extend to functors on such categories of objects.

**3.1 re topological gravity** The Riemannian metrics on a closed compact smooth manifold X, regarded as the object of an action of the group  $\mathbb{D}$  of diffeomorphisms which frame a basepoint ([11] see also [15]) define a topological groupoid

[Metrics //D]

which models the space of states or configurations in Euclidean general relativity. Myers and Steenrod [25] show that the isotropy group of a Riemannian metric g on a compact manifold X is compact. As a corollary, we have a commutative diagram

## Proposition



where the fibers of the left diagonal map are spaces of metrics g with  $iso_0(g) = K$ : which, being cones, are contractible.

Arguing from a (hypothetical) Leray sseq

 $H^*(|[\operatorname{\mathbb{C}pct}_0/\mathbb{D}]|; \operatorname{\mathcal{H}}^*(\operatorname{iso})) \Rightarrow H^*|[\operatorname{\mathbb{M}etrics}(X)/\!/\mathbb{D}]|$ 

suggests that a natural way to think of the target of this sseq is as an element

$$[\mathrm{Rt}_0^!]:\mathbb{A}(X)\to \int_{K\in \mathfrak{C}\mathrm{pct}_0/\mathbb{D}}H^*_{\mathfrak{B}}|[X/\!/K]|$$

of a category of natural transformations of functors (defined over a topos of conjugacy classes of connected compact subgroups of  $\mathbb{D}$ ).

**3.2** Let us then consider the cobordism category of compact **Riemannian** fourmanifolds with **effective**  $\Omega^*$ Blcs - algebra structures as morphisms, and compatible (Riemannian) three-manifolds, anchored by line bundles, as objects, as a category fibered over the construction in §2.3 above.

This provides a reasonable background for algebraic models of quantum gravity adapted to topology change. Considered as a generalized Morse theory, it weighs symmetric objects and states very heavily.

### appendices

 $\bullet$  a1 This is a remark about a rough circle of ideas, a homotopy-theoretic version of Wess-Zumino-Witten theory, where

A principal  $\Omega^2 S^2$  (or line) bundle on X is homotopically equivalent to a field  $X \to |\operatorname{Fin}_{\mathbb{C}}^{\otimes}|^+$  of (virtual) approximations of the cx line by finite subsets.

The set of such equivalence classes has group structure, defining the homomorphism

$$\pi^2(S^1 \wedge X) := \operatorname{Pic}_{wzw}(X) \to \operatorname{Pic}_{\mathbb{C}}(X) \cong H^2(X, \mathbb{Z})$$

induced by a generator of  $H^2(\Omega S^2, \mathbb{Z})$ .

A line bundle has an underlying (cx) line bundle; they are candidates for homotopytheoretic gauge fields, in particular on 3-manifolds. They can be interpreted as fields of WZW models, as follows: Taking loopspaces sends fibrations to fibrations; applied to Hopf

 $\mathbb{T} = S^1 \to SU(2) = S^3 \to \mathbb{C}P^1 = S^2$ 

we get the universal cover

 $\Omega \mathbb{T} \simeq \mathbb{Z} \to \Omega SU(2) \to \Omega S^2$ 

so looping once again defines a homotopy equivalence

$$\Omega^2 \mathrm{SU}(2) \ (\cong \Omega^2 \mathbb{R}^3_+) \to \Omega^2_e \mathbb{S}^2$$

with the identity component of a semi-direct product

$$\Omega^2 \mathbb{S}^2 \simeq \Omega^2 \mathrm{SU}(2) \rtimes \mathbb{Z} \sim [\Omega^2 \mathrm{SU}(2) / / \mathbb{Z}]$$

though one must be careful because group-like spaces are not necessarily groups.

Note that  $\mathbb{Z} \cong \pi_0(\Omega^2 S^2)$  builds the writhe or determinant of a braid (as in §8 of [24]: a map  $X \to Y$  of Poincaré duality spaces, with a lift of its normal spherical bundle to the classifying space<sup>2</sup> of (degree one self-maps/homotopy automorphisms of)  $S^3$ ) is Steenrod representable, by work of Hopkins and Mahowald) into geometry. In this language there's a topos of (Poincaré duality) spaces over B $\mathcal{H}Aut(S^3)$  which is where solutions of variational geometric problems live, while this note argues/claims that the topos of (4D) spaces over B $\mathcal{H}Aut(S^2)$  is a good place for renormalizeable 4D Riemannian geometry/qEuclidean GR.

• There is a nice 3-form

## $\mathfrak{a}\mathfrak{d}\mathfrak{a} + \lambda \cdot \kappa(\mathfrak{a},\mathfrak{a},\mathfrak{a})$

on the space of  $\mathfrak{su}(2)$  - connections, where  $\kappa$  is related to the Killing form of Lie theory and to the generator of  $\pi_3$  of a (...) simple Lie group; it leads to invariants of three-manifolds defined without reference to Riemannian geometry. The Bott-Cartan element underlying  $\kappa$  is the class of origin for  $4\nu_{(2)} = \eta^3 \in \pi_3^S$  in the stable homotopy groups of spheres; note that both  $\eta$  and  $\eta^3$  are in the image of the J-homomorphism, but  $\eta^2$  is not.

• a2 In these notes  $Y \cong Y(\mathcal{B})$  is in principle a compact connected closed oriented three-manifold, presented as an open book with binding  $\mathcal{B} \subset Y$  [7](§2) [14], eg the mapping torus of a two-manifold. Any  $(\ldots)$  three-manifold has such presentations.

• [Thurston-Winkelnkemper] [28] There is a contact form  $\alpha$ , *ie*  $\alpha \wedge d\alpha \neq 0$  supported by  $\mathcal{B}$ .

• [Giroux] [13] : The space of contact forms supported by a binding  $\mathcal{B}$  is connected (under isotopy = smooth deformation of the contact form  $\alpha^3$ ).

• [19] Poincaré's lemma in link calculus [14] (Prop 4.5.11, Exercises 5.3.13] associates to (the link  $\mathcal{B}$  binding)  $Y(\mathcal{B})$ , a four-manifold  $X \supset \partial X \cong Y$  with intersection form  $H^2(X/Y;\mathbb{Z})$  defined on  $H_2(X,\mathbb{Z})$  as in [19] (§5). The coboundary

$$H^{1}Y \rightarrow H^{2}X/Y : \operatorname{Pic}_{wzw}Y \rightarrow \operatorname{Pic}_{\mathbb{C}}X/Y$$

<sup>&</sup>lt;sup>2</sup>which is the (to me mysterious) universal cover  $\Omega^2 SU(2)$ 

<sup>&</sup>lt;sup>3</sup>accomplished by stabilizing [7](§4.31 Fig 12] under certain (? 2-categorical) moves

sends lines on the boundary to complex lines on the interior of X..

• **a3** Check of §2.1: The diagram

$$S^{3} \times S^{3} \xrightarrow{\eta^{2} \times \eta^{2}} \Omega^{2} S^{3} \times \Omega^{2} S^{3}$$

$$\downarrow^{\mu_{S^{3}}} \qquad \qquad \downarrow^{\mu_{\Omega^{2} S^{3}}} \Omega^{2} S^{3}$$

commutes, because across the top we have

$$q,k \rightarrow [z \rightarrow z q z^{-1}], [w \rightarrow w k w^{-1}] \;,$$

then down along the right we have

$$[z \rightarrow zqz^{-1}], [w \rightarrow wkw^{-1}] \mapsto [u \rightarrow u \times u \rightarrow uqu^{-1} \cdot uku^{-1} = ukqu^{-1}$$

while around the left side we have  $q, k \mapsto [u \to u q k u^{-1}]$  .

To be more precise about basepoints: in somewhat mixed notation, we have a commutative diagram

in which  $\rho(r, \omega, q) = \omega q \omega^{-1}$  for  $r \in \mathbb{R}^{\times}$  and  $\rho(0, -, -) = 0, \rho(\infty, -, -) = \infty$ . The downward left collapse  $\{0, \infty\} \times \mathbb{T} \to [0, \infty] \times \mathbb{T} \to S^2$  is a cofibration but the right side is just an inclusion. The assertion is not that  $\rho$  is continuous, but that collapsing the two endpoints of a cylinder does define a continuous if not smooth  $\rho$ .

#### References

- 1. D Angella, A Otiman, N Tardini Cohomologies of locally conformally symplectic manifolds and solvmanifolds https://arxiv.org/abs/1703.05512
- 2. M Atiyah, R Bott, The moment map and equivariant cohomology, Topology 23, 1-28 (1984)
- 3. Dong Yan, Hodge structures on symplectic manifolds, Adv Math 120,143 –154 (1996)
- D Ebin The manifold of Riemannian metrics. Global Analysis (Proc. Sympos. Pure Math (1968) 11–40 AMS (1970)
- 5. Y Eliashberg On symplectic manifolds with some contact properties. J. Differential Geom. 33 (1991) 233 238
- 6. A few remarks about symplectic filling, https://arxiv.org/abs/math/0311459
- J Etnyre Lectures on open book decompositions and contact structures, https://arxiv.org/abs/math/0409402
- Lectures on contact geometry in low-dimensional topology https://arxiv.org/abs/math/0610798
- 9. K Honda On symplectic cobordisms, https://arxiv.org/abs/math/0102162
- 10. D Fiorenza, H Sati, U Schreiber, The character map in (twisted differential) non-abelian cohomology https://arxiv.org/abs/2009.11909

#### J MORAVA

- 11. M Freedman, Controlled Mather-Thurston theorems, https://arxiv.org/abs/2006.00374
- G Giotopoulos, H Sati, U Schreiber, Flux Quantization on M5-branes https://arxiv.org/abs/2406.11304
- E Giroux, Géométrie de contact: de la dimension trois vers les dimensions supérieures. Proceedings of the Beijing ICM Vol. II 405–414, Higher Ed. Press, Beijing 2002
- 14. R Gompf, A Stipsicz, 4-manifolds and Kirby calculus. AMS Grad Studies 20 (1999)
- S Haller, T Rybicki, On the group of diffeomorphisms preserving a locally conformal symplectic structure. Ann. Global Anal. Geom. 17 (1999) 475 – 502
- 16. K Hess, Rational homotopy theory https://arxiv.org/abs/math/0604626
- M Hopkins, I M Singer, Quadratic functions in geometry, topology, and -theory. J. Differential Geom. 70 (2005) 329–452.
- Haw-Chung Lee, A kind of even-dimensional differential geometry and its application to exterior calculus. Amer. J. Math. 65 (1943) 433 – 438
- W Meier, Minimal models for nonnilpotent spaces, Comment. Math. Helv. 55 (1980) 622 633
- 20. J Morava, At the boundary of Minkowski space, https://arxiv.org/abs/2111.08053
- —-, A locally conformally symplectic structure on Kerr space-time https://arxiv.org/abs/2212.08210
- 22. —, A homotopy-theoretic context for CKM/Birkhoff renormalization https://arxiv.org/abs/2307.10148
- 23. —, Some very low-dimensional algebraic topology https://arxiv.org/abs/2411.15885
- —, D Rolfsen, Toward the group completion of the Burau representation. Trans. AMS 376 (2023) 1845 - 1865 https://arxiv.org/abs/1809.01994
- S B Myers, N Steenrod, The group of isometries of a Riemannian manifold. Ann. of Math. 40 (1939) 400 –416.
- J Neisendorfer, Lie algebras, coalgebras and rational homotopy theory for nilpotent spaces, Pacific J. Math. 74 (1978) 429 – 460
- 27. A Otiman, Locally conformally symplectic bundles, https://arxiv.org/abs/1510.02770
- D Ravenel, Complex cobordism and stable homotopy groups of spheres Academic Press Pure and Applied 121 (1986)
- 29. R E Stong, Contact manifolds, J. Differential Geometry 9 (1974) 219 238
- 30. W Thurston, H Winkelnkemper On the existence of contact forms, Proc. AMS 52 (1975) 345-347
- Tseng Li-Sheng, Yau Shing-Tung, Cohomology and Hodge theory on symplectic manifolds I, J Diff Geo 91 (2012) 383 – 416 ; II, J. Diff Geo 91 (2012) 417 – 443
- 32. —, —, Tsai Chung-Jun, Hodge theory on symplectic manifolds: III. J. Diff Geo103 (2016) 83 143
- 33. https://en.wikipedia.org/wiki/Chabauty\_topology
- 34. https://en.wikipedia.org/wiki/Misalignment\_mechanism

DEPARTMENT OF MATHEMATICS, THE JOHNS HOPKINS UNIVERSITY, BALTIMORE, MARYLAND

Email address: jmorava1@jhu.edu