# Lipschitz-Driven Inference: Bias-corrected Confidence Intervals for Spatial Linear Models

David R. Burt<sup>\*1</sup> Renato Berlinghieri<sup>\*1</sup> Stephen Bates<sup>1</sup> Tamara Broderick<sup>1</sup>

### Abstract

Linear models remain ubiquitous in modern spatial applications — including climate science, public health, and economics - due to their interpretability, speed, and reproducibility. While practitioners generally report a form of uncertainty, popular spatial uncertainty quantification methods do not jointly handle model misspecification and distribution shift — despite both being essentially always present in spatial problems. In the present paper, we show that existing methods for constructing confidence (or credible) intervals in spatial linear models fail to provide correct coverage due to unaccountedfor bias. In contrast to classical methods that rely on an i.i.d. assumption that is inappropriate in spatial problems, in the present work we instead make a spatial smoothness (Lipschitz) assumption. We are then able to propose a new confidence-interval construction that accounts for bias in the estimation procedure. We demonstrate that our new method achieves nominal coverage via both theory and experiments. Code to reproduce experiments is available at https://github. com/DavidRBurt/Lipschitz-Driven-Inference.

# 1. Introduction

Preprint.

Scientists often want to understand the sign and magnitude of an association in cases where covariates vary spatially. Examples include the relationship between aerosol concentrations and regional precipitation changes (Westervelt et al., 2018), the association between proximity to major highways and the prevalence of dementia (Li et al., 2023), and the link between air pollution exposure and birth weight (Lee et al., 2022). Here, the covariates (aerosol concentrations, proximity to highways, and air pollution) may all be thought of as a function of spatial location. Moreover, scientists often have access to data at some spatial locations but want to infer the association of interest at other locations. For instance, a country might measure birth weight and air pollution at the municipal level for some municipalities — and be interested in understanding the association between air pollution and birth weight in unobserved municipalities.

A priori we expect birth weight as a response to have a potentially nonlinear relationship to air pollution as a covariate. So a natural idea is to fit a flexible, nonparametric model, such as a neural network, Gaussian process, or random forest. While these methods can be expected to achieve high predictive accuracy, they often lack interpretability, and it is not clear how to draw conclusions about the specific relationships between covariates and the response variable (Rudin, 2019; Doshi-Velez and Kim, 2017).

Another option, made precise by Buja et al. (2019), is to recognize the potentially nonlinear relationship between covariates and response, but fit a linear model to the data essentially to summarize any trend. Namely, one can use the parameter estimate as a best guess of the association trend between covariates and response, and a confidence interval captures uncertainty in this trend. And indeed, linear models remain widespread to the present day despite an understanding that they will generally be misspecified. All of the applied papers above use linear regression. Moreover, Castro Torres and Akbaritabar (2024, Figure 1) conducted a survey of publications up to 2022 that report quantitative methods; they found that over half of all such papers in various areas (agricultural science, social sciences, and medical and health sciences) reported results from a linear model.

Providing a valid confidence interval in this setting remains, however, an open problem. Namely, we are interested in the common setting where the linear model is misspecified, and we would like to understand associations at spatial locations that may be different from those in the observed data. Under misspecification, Buja et al. (2019) observes that the ordinary least squares (OLS) estimator now depends on the covariate values, and classic confidence intervals are valid only at exactly the set of observed (training) covariate values. The heteroskedasticity-consistent sandwich estimator (Huber, 1967; White, 1980a;b; Buja et al., 2019) provides valid confidence intervals under misspecification, but then

arXiv:2502.06067v1 [stat.ML] 9 Feb 2025

 $<sup>^{*}</sup>$ Equal contribution  $^{1}$ MIT LIDS. Correspondence to: David R. Burt <dburt@mit.edu>, Renato Berlinghieri <renb@mit.edu>.

we need to assume that all covariates are samples from a single population. Generalized least squares linear regression (GLS, Aitken 1936) is designed to handle spatial correlation in residuals (Cressie, 2015, pages 22-24). But there is no reason to expect that GLS solves the bias issues that arise in OLS under a combination of misspecification and distribution shift. Spatial data analysts often report Bayesian credible intervals, e.g. with a Gaussian process model, but these are known to underestimate uncertainty in the simultaneous presence of misspecification and distribution shift (Walker, 2013; Müller, 2013). Finally, importance weighting methods (Shimodaira, 2000) have two drawbacks. (1) They assume one is able to de-bias the estimator, which need not be true. (2) Confidence interval validity relies on exact estimation of importance weights, which will never be perfectly achieved in practice. We discuss related work in more detail in Appendix A.

In what follows, we show in both real and simulated experiments (Section 4) that all of the confidence-interval constructions above can provide (much) less than nominal coverage. **Our principal contribution** is to provide the first confidence intervals for spatial linear regression that are guaranteed to achieve frequentist coverage at the nominal level (a) under nonparametric assumptions about the data-generating distribution and (b) when the target locations need not match the observed data. Along the way, we formalize the problem of estimation in this setting. In our experiments, our method is (by far) the only method to consistently achieve nominal coverage.

### 2. Problem Setup

We start by describing the available data. After briefly reviewing classic well-specified linear regression, we set up the misspecified case with different target and source data, and we establish our estimand in this case.

**Data.** Following the covariate-shift literature (Ben-David et al., 2006; Pan and Yang, 2010; Csurka, 2017), we refer to our fully observed (training) data as the *source* data; likewise, we let *target* data denote the (test) locations and covariates where we do not observe the response but would like to understand the association between covariates and response. In particular, the source data consists of N triplets  $(S_n, X_n, Y_n)_{n=1}^N$ , with spatial location  $S_n \in S$ , covariate  $X_n \in \mathbb{R}^P$ , and response  $Y_n \in \mathbb{R}$ . Here  $P \ge 1$ , and S represents geographic space; we assume S is a metric space with metric  $d_S$ . We collect the source covariates in the matrix  $X \in \mathbb{R}^{N \times P}$  and the source responses in the N-long column vector Y. The target data consists of M pairs  $(S_m^*, X_m^*)_{m=1}^M$ , with  $S_m^* \in S$ ,  $X_m^* \in \mathbb{R}^P$ . The corresponding responses  $Y_m^* \in \mathbb{R}$ ,  $1 \le m \le M$  are unobserved. We collect the target covariates in  $X^* \in \mathbb{R}^{M \times P}$  and responses in column vector  $Y^* \in \mathbb{R}^M$ .

**Review: Well-specified Linear Model.** Though we will focus on the misspecified (spatial) case, we start by reviewing the classic well-specified case for comparison purposes.

In the classic well-specified setup, we have

$$Y_n = \theta_{\text{OLS}}^{\text{T}} X_n + \epsilon_n, \quad Y_m^{\star} = \theta_{\text{OLS}}^{\text{T}} X_m^{\star} + \epsilon_m^{\star} \qquad (1)$$

with column-vector parameter  $\theta_{OLS} \in \mathbb{R}^P$  and  $\epsilon_n, \epsilon_m^{\star} \stackrel{iid}{\sim} \mathcal{N}(0, \sigma^2)$  for some (unknown)  $\sigma^2 > 0$ . For any fixed set of source data points (and assuming invertibility holds as needed), we can recover the parameter exactly as

$$\theta_{\text{OLS}} = \arg\min_{\theta \in \mathbb{R}^P} \mathbb{E} \Big[ \frac{1}{N} \sum_{n=1}^{N} (Y_n - \theta^{\mathrm{T}} X_n) \Big]$$
(2)

$$= \mathbb{E}[X^{\mathrm{T}}X]^{-1}\mathbb{E}[X^{\mathrm{T}}Y] = (X^{\mathrm{T}}X)^{-1}X^{\mathrm{T}}\mathbb{E}[Y].$$
(3)

An analogous formula holds for target points;  $\theta_{OLS}$  is constant across covariate values in any case. Since the population expectation is unknown, analysts typically estimate  $\theta_{OLS}$  via maximum likelihood. The standard estimator and confidence interval at level  $\alpha$  for the *p*th coefficient,  $\theta_{OLS,p}$ , appear below in Equation (4). Under correct specification, the confidence interval is valid: that is, it provides nominal coverage. E.g., a 95% confidence interval contains the true parameter 95% of the time under resampling.

$$\hat{\theta}_{\text{OLS},p} = e_p^{\text{T}} (X^{\text{T}} X)^{-1} X^{\text{T}} Y, I_{\text{OLS},p} = \hat{\theta}_{\text{OLS},p} \pm z_\alpha \rho,$$
(4)

where  $\rho^2 = \sigma^2 e_p^{\rm T} (X^{\rm T} X)^{-1} e_p$ ,  $z_{\alpha}$  is the  $\alpha$  quantile of a standard normal distribution, and  $e_p$  denotes the *P*dimensional vector with a 1 in entry *p* and 0s elsewhere. The noise variance  $\sigma^2$  in Equation (4) is typically replaced with an estimate  $\hat{\sigma}^2$ .

**Misspecified Spatial Setup: Data-Generating Process.** In what follows, we assume the data is generated as

$$Y_n = f(X_n, S_n) + \epsilon_n, Y_m^{\star} = f(X_m^{\star}, S_m^{\star}) + \epsilon_m^{\star}, \quad (5)$$

for a function f that need not have a parametric form and with  $\epsilon_n, \epsilon_m^{\star} \stackrel{iid}{\sim} \mathcal{N}(0, \sigma^2)$  for some (unknown)  $\sigma^2 > 0$ .

For ease of development, we will next make the simplifying assumption that source and target covariates are fixed functions of spatial location. Recall that all of the covariates in our examples (aerosol concentrations, proximity to highways, and air pollution) can be expected to vary spatially and be measured with minimal error. We similarly expect meteorological variables such as precipitation, humidity, and temperature to be reasonably captured by this assumption.<sup>1</sup>

<sup>&</sup>lt;sup>1</sup>Conversely demographic covariates may more reasonably be thought of as noisy functions of space, and further work is needed to handle the noisy case.

**Assumption 1.** There exists a function  $\chi : S \to \mathbb{R}^P$  such that  $X_m^* = \chi(S_m^*)$  for  $1 \le m \le M$  and  $X_n = \chi(S_n)$  for  $1 \le n \le N$ .

Under assumption 1, our data-generating process simplifies. **Assumption 2.** There exists a function  $f : S \to \mathbb{R}$  such that  $\forall m \in \{1, ..., M\}, Y_m^{\star} = f(S_m^{\star}) + \epsilon_m^{\star}$  and  $\forall n \in \{1, ..., N\}, Y_n = f(S_n) + \epsilon_n$ , where  $\epsilon_m^{\star}, \epsilon_n \overset{iid}{\sim} \mathcal{N}(0, \sigma^2)$ .

**Our Estimand.** At a high level, our goal is to capture the relationship between a covariate and the response variable at target locations using data from source locations, while taking into account that these two sets of locations may differ. From this perspective, we can define our estimand as the parameter of the best linear approximation to the response, where "best" is defined by minimizing squared error over the target locations.

$$\theta_{\text{OLS}}^{\star} = \arg\min_{\theta \in \mathbb{R}^{P}} \mathbb{E}\Big[\sum_{m=1}^{M} (Y_{m}^{\star} - \theta^{\top} X_{m}^{\star})^{2} |S_{m}^{\star}\Big].$$
(6)

As in White (1980b); Buja et al. (2019), since the datagenerating process may be non-linear,  $\theta_{OLS}^{\star}$  is no longer constant like the well-specified case, but rather we expect it to vary as a function of the target locations.

As an aside, we note that an alternative estimand that is similar in spirit would treat the source and target locations as random, each with their own respective distribution; in this case, the estimand in Equation (6) would lose the conditioning on  $S_m^{\star}$ , and covariate-shift methods could apply. First, we note that we are not aware of approaches for constructing finite-sample confidence intervals using current covariateshift methodology, so the question of valid confidence intervals remains open in any case. That being said, we choose to fix the source and target locations in our present work because, in the spatial applications of interest here, these locations might not each plausibly be sampled i.i.d. from a population. For instance, if source data arises from the United States Environmental Protection Agency (EPA), we know that EPA sensors are sited with other sensors in mind and thus are known to not be i.i.d. Also particular target locations, such as certain municipalities missing data, may be of interest; especially when there may be only a handful or even just one such target location, it's not clear that there is a meaningful population representation. So we henceforth use the conditional form of Equation (6).

To solve the minimization in Equation (6), it will be convenient to assume invertibility, as for OLS.

**Assumption 3.**  $X^{\star T}X^{\star}$  is invertible.

With assumptions 1 and 3, we can solve the minimization in Equation (6) to find

$$\theta_{\text{OLS}}^{\star} = (X^{\star \text{T}} X^{\star})^{-1} X^{\star \text{T}} \mathbb{E}[Y^{\star} | S^{\star}].$$
(7)

See Appendix E.1 for a derivation. To the best of our knowledge, the target-conditional estimand in Equation (7) has not been previously proposed or analyzed in spatial linear regression.

To estimate  $\theta_{OLS}^{\star}$  well, we need to estimate the conditional expectation  $\mathbb{E}[Y^{\star} \mid S^{\star}]$  well. To hope to do so, we need to make regularity assumptions that make accurate estimation feasible. In classical and covariate-shift settings, these take the form of i.i.d. and bounded-density-ratio assumptions. Since we've seen that the classic assumptions are inappropriate in this spatial setting, we instead assume f is shared across source and target (assumption 2) and not varying so quickly in space as to be difficult to learn from limited data.

**Assumption 4.** The conditional expectation, f, is *L*-Lipschitz as a function from  $(S, d_S) \rightarrow (\mathbb{R}, |\cdot|)$ . That is, for any  $s, s' \in S$ ,  $|f(s) - f(s')| \leq Ld_S(s, s')$ .

As an illustrative example of how this assumption might be satisfied, consider  $\chi$  an  $L_1$ -Lipschitz function of the spatial domain, and  $f(s) = \beta^T \chi(s) + g(s)$  with g a fixed,  $L_2$ -Lipschitz function of S. Then f is a  $(||\beta||_2 L_1 + L_2)$ -Lipschitz function of the spatial domain.

### 3. Lipschitz-Driven Inference

We next provide a confidence interval for  $\theta^{\star}_{OLS,p}$ , the *p*th coefficient of the target-conditional least squares estimand (Equation (7)). We support its validity with theory (in the present section) and experiments (in Section 4). To that end, we start by providing an efficiently-computable point estimate. We end by discussing the role and choice of the Lipschitz constant *L*.

**Lipschitz-Driven Point Estimation.** Since the target covariates are known, the key challenge in estimating Equation (7) is estimating the unknown quantity  $\mathbb{E}[Y^*|S^*]$ .

For our first approximation, recall that, by assumption 4, f varies smoothly in space. Since the conditional distribution of the responses given the spatial locations is the same function for both target and source data (assumption 2), we can approximate  $\mathbb{E}[Y^*|S^*]$  by a weighted average of  $\mathbb{E}[Y|S]$  values for locations S near  $S^*$ . Concretely, let  $\Psi \in \mathbb{R}^{M \times N}$  be a (non-negative) matrix of weights. If  $\Psi$  assigns weight mostly to source locations near each corresponding target location, then by the Lipschitz assumption (assumption 4),

$$\mathbb{E}[Y^{\star}|S^{\star}] \approx \Psi \mathbb{E}[Y|S]. \tag{8}$$

 $\mathbb{E}[Y|S]$  is also unobserved, so we next approximate it by observed values of Y (at each source location in S):

$$\Psi \mathbb{E}[Y|S] \approx \Psi Y. \tag{9}$$

Together, these two approximations yield the estimator:

$$\hat{\theta}_p^{\Psi} = e_p^{\mathrm{T}} \left( X^{\star \mathrm{T}} X^{\star} \right)^{-1} X^{\star \mathrm{T}} \Psi Y.$$
(10)

**Algorithm 1** Lipschitz-Driven CI for  $\theta_{OLS,n}^{\star}$ 

input  $\{(S_n, X_n, Y_n)\}_{n=1}^N, \{(S_m^{\star}, X_m^{\star})\}_{m=1}^M$ , Lipschitz constant L, confidence level  $1 - \alpha, \sigma^2$  (optional)

- **output** A  $(1 \alpha)$ -confidence interval  $I^{\Psi}$  for  $\theta^{\star}_{OLS,p}$ 1:  $\Psi \leftarrow 1$ -NN $(\{S_n\}_{n=1}^N, \{S_m^{\star}\}_{m=1}^M)$  (Definition 5) 2:  $\hat{\theta}_p^{\Psi} \leftarrow e_p^{\top} (X^{\star \top} X^{\star})^{-1} X^{\star \top} \Psi Y$ 3:  $w \leftarrow e_p^{\top} (X^{\star \top} X^{\star})^{-1} X^{\star \top}, v^{\Psi} \leftarrow w \Psi$
- 3:  $w \leftarrow e_p^+(X^{\star+}X^{\star}) \stackrel{T}{\to} X^{\star+}, v^{\Psi} \leftarrow w \Psi$ 4:  $B \leftarrow \sup_{g \in \mathcal{F}_L} \left| \sum_{m=1}^M w_m g(S_m^{\star}) - \sum_{n=1}^N v_n^{\Psi} g(S_n) \right|$ . Compute with linear program (Appendix B.2).
- 5: If  $\sigma^2$  unknown:

$$\sigma^2 := \hat{\sigma}_N^2 \leftarrow \inf_{g \in \mathcal{F}_L} \frac{1}{N} \sum_{n=1}^N (Y_n - g(S_n))^2 \quad (12)$$

by quadratic program (Appendix B.5).

- 6:  $c \leftarrow \sigma \|v^{\Psi}\|_2$
- 7: Find  $\Delta(\alpha)$  satisfying

$$\Phi(\Delta(\alpha)) - \Phi(\frac{-2B}{c} - \Delta(\alpha)) = 1 - \alpha,$$

by root finding algorithm (Appendix B.4).

8:  $I^{\Psi} \leftarrow \left[\hat{\theta}_p^{\Psi} - B - c\,\Delta(\alpha) , \,\hat{\theta}_p^{\Psi} + B + c\,\Delta(\alpha)\right].$ 

In our experiments, we construct  $\Psi$  as follows.

**Definition 5** (Nearest-Neighbor Weight Matrix). *Define the 1-nearest neighbor weight matrix by* 

$$\Psi_{mn} = \begin{cases} 1 & S_n = \text{closest source location to } S_m^{\star} \\ 0 & \text{otherwise} \end{cases}$$
(11)

We break ties uniformly at random.

While this simple construction works well in our present experiments, we discuss the potential benefits of other constructions in Appendix B.1.

**Lipschitz-Driven Confidence Intervals.** We detail how to efficiently compute our proposed confidence interval for  $\theta^*_{OLS,p}$  in Algorithm 1. We prove its validity in Theorem 7 below. Before stating our result, we establish relevant notation and intuition for how our method works. First, we show the difference between our estimand and the estimator is normally distributed. Toward that goal, we start by writing

$$\theta_{\text{OLS},p}^{\star} - \hat{\theta}_{p}^{\Psi} = \sum_{m=1}^{M} w_{m} f(S_{m}^{\star}) - \sum_{n=1}^{N} v_{n}^{\Psi} Y_{n},$$
(13)

for  $w := e_p^{\mathrm{T}} (X^{\star \mathrm{T}} X^{\star})^{-1} X^{\star \mathrm{T}} \in \mathbb{R}^M$  and  $v^{\Psi} := w \Psi \in \mathbb{R}^N$ . By assumption 2, the righthand side of Equation (13) can be written

$$\underbrace{\sum_{m=1}^{M} w_m f(S_m^{\star}) - \sum_{n=1}^{N} v_n^{\Psi} f(S_n)}_{\text{bias}} - \underbrace{\sum_{n=1}^{N} v_n^{\Psi} \epsilon_n}_{\text{randomness}}.$$
 (14)

That is, Equation (14) expresses  $\theta^{\star}_{OLS,p} - \hat{\theta}^{\Psi}_{p}$  as the sum of (i) a *bias* term due to differing locations between the source and target data and (ii) a a mean-zero Gaussian *randomness* term due to observation noise.

Since the spatial locations are fixed, the bias term is not random and can be written as  $b \in \mathbb{R}$ . It follows that

$$\theta^{\star}_{\text{OLS},p} - \hat{\theta}^{\Psi}_{p} \sim \mathcal{N}(b, \sigma^{2} \| v^{\Psi} \|_{2}^{2})$$
(15)

since the variance of the randomness term is the sum of the variances of its (independent) summands.

Our strategy from here will be to (1) bound b, (2) establish a valid confidence interval using our bound on b while assuming fixed  $\sigma^2$ , and (3) estimate  $\sigma^2$  consistently (as  $N \to \infty$ ).

To bound the bias b, we use assumption 4 to write

$$|b| \le \sup_{g \in \mathcal{F}_L} \left| \sum_{m=1}^M w_m g(S_m^\star) - \sum_{n=1}^N v_n^\Psi g(S_n) \right|, \quad (16)$$

where  $\mathcal{F}_L$  is the space of *L*-Lipschitz functions from  $S \rightarrow \mathbb{R}$ . In Appendix E.3, we show that it is possible to use Kantorovich-Rubinstein duality to restate the righthand side of Equation (16) as a Wasserstein-1 distance between discrete measures. This alternative formulation is useful since it can be cast as a linear program (Peyré and Cuturi, 2019, Chapter 3); see Appendix E.3. Let *B* denote the output of this linear program.

Given  $B \geq |b|$ , the following lemma (with proof in Appendix E.4) allows us to construct a confidence interval for  $\theta^{\star}_{\text{OLS},p}$  centered on  $\hat{\theta}^{\Psi}_{p}$ . We discuss the benefits of this construction over alternative approaches in Appendix B.3.

**Lemma 6.** Let  $b \in [-B, B]$ ,  $\tilde{c} > 0$ , and  $\alpha \in (0, 1)$ . Then the narrowest  $1 - \alpha$  confidence interval that is symmetric and valid for all  $\mathcal{N}(b, \tilde{c}^2)$  is of the form  $[-B - \tilde{c}\Delta, B + \tilde{c}\Delta]$ where  $\Delta$  is the solution of:

$$\Phi(\Delta) - \Phi\left(\frac{-2B}{\tilde{c}} - \Delta\right) = 1 - \alpha \tag{17}$$

with  $\Phi$  the cumulative density function of a standard normal distribution. Also, the  $\Delta$  satisfying this inequality is  $\Delta \in [\Phi^{-1}(1-\alpha), \Phi^{-1}(1-\frac{\alpha}{2})].$ 

The resulting confidence interval appears in Algorithm 1. We next establish its validity. So far, we have covered only the known  $\sigma^2$  case. We handle the unknown  $\sigma^2$  case after the following theorem.

**Theorem 7.** Suppose  $(S_m^{\star}, X_m^{\star}, Y_m^{\star})_{m=1}^M$  and  $(S_n, X_n, Y_n)_{n=1}^N$  satisfy assumptions 1 to 4 with known  $\sigma^2$ . Define the (random) interval  $I^{\Psi}$  as in Algorithm 1 using the known value of  $\sigma^2$ . Then with probability at least  $1 - \alpha$ ,  $\theta_{OLS,p}^{\bullet} \in I^{\Psi}$ . That is,  $I^{\Psi}$  has coverage (conditional on the test locations) at least  $1 - \alpha$ .

**Bias-corrected Confidence Intervals for Spatial Linear Models** 



*Figure 1.* Coverages (left) and confidence interval widths (right) for our method as well as 5 other methods. Only our method and GP BCIs consistently achieve the nominal coverage rate (95%) in this example; the GP BCIs line (dashed blue) overlaps with ours, in solid black, for most of the shifts. Of the two methods with correct coverage, our method yields much narrower intervals.

In Appendix E.2, we prove validity of our confidence interval for a generic choice of weight matrix  $\Psi$ . Theorem 7 is an immediate corollary of that result.

**Consistent Estimation of the Noise Variance**  $\sigma^2$ . Generally, the noise variance  $\sigma^2$  is unknown, so we will substitute an estimate for  $\sigma^2$  in the calculation of the confidence interval  $I^{\Psi}$  (Step 5 in Algorithm 1). In Corollary 9 below, we show that the resulting confidence interval has asymptotically valid coverage. To that end, we first show that the estimator in Equation (12) is consistent for  $\sigma^2$ .

**Proposition 8.** Suppose the spatial domain  $S = [-A, A]^D$ for some  $A > 0, D \in \mathbb{N}$ . Take assumptions 2 and 4. For any sequence of source spatial locations  $(S_n)_{n=1}^{\infty}$ , take  $\hat{\sigma}_N^2$  as in Equation (12). Then  $\hat{\sigma}_N^2 \to \sigma^2$  in probability as  $N \to \infty$ .

See Appendix E.5 for a proof. For intuition, recall that the conditional expectation minimizes expected squared error over all functions. Since the conditional expectation is *L*-Lipschitz (assumption 4), we have

$$\sigma^2 = \inf_{g \in \mathcal{F}_L} \mathbb{E}\left[\frac{1}{N} \sum_{n=1}^N (Y - g(S_n))^2 \middle| S\right].$$
(18)

The empirical version of Equation (18) is Equation (12), which should be a good estimate for  $\sigma^2$  if N is large.

When S is a subset of Euclidean space or a subset of the sphere with spherical distance, the minimization in Equation (12) yields a quadratic program for computing  $\hat{\sigma}_N^2$ . We provide implementation details in Appendix B.5.

Given Proposition 8, it follows from Slutsky's Lemma that the resulting confidence interval is asymptotically valid.

**Corollary 9.** For S as in Proposition 8, with the assumptions and notation of Theorem 7, but with  $\sigma^2$  unknown, the confidence interval  $I^{\Psi}$  from Algorithm 1 has asymptotic (with fixed M and as  $N \to \infty$ ) coverage at level  $1 - \alpha$ .

See Appendix E.5 for a proof.

**Choice of the Lipschitz Constant** *L*. The Lipschitz assumption allows us to make inferences about the target data from the source data. The Lipschitz constant subsequently enters our intervals in two principal ways: via *B* and via  $\hat{\sigma}_N^2$ . Intuitively, larger values of *L* (allowing for less smooth responses) lead to our algorithm constructing confidence intervals with larger bounds (*B*) on the bias but smaller estimated residual variance ( $\hat{\sigma}_N^2$ ). We depict this trade-off in a concrete example in Section 4 (Figure 3).

Ultimately the choice of Lipschitz constant must be guided by domain knowledge. We give a concrete example describing our choice of the Lipschitz constant in our real-data experiment on tree cover (Section 4). In our simulated experiments, we know the minimum value for which assumption 4 holds; call it  $L_0$ . So we first choose  $L = L_0$ . Then we perform ablation studies in both simulated and real data showing that we essentially maintain coverage while varying L over an order of magnitude around our initial choices. We show that further decreasing L can decrease coverage and discuss why it is useful to err on the side of conservatism (i.e., a larger L).

### 4. Experiments

In simulated and real data experiments, we find that our method consistently achieves nominal coverage, whereas all the alternatives dramatically fail to do so. We also provide ablation studies to evaluate the effect of varying the Lipschitz constant in both simulated and real settings.

Baselines. We compare to five alternative constructions.

*Ordinary Least Squares (OLS).* We treat the noise variance as unknown and estimate it as the average squared residual, with a correction for the number of degrees of freedom and a



*Figure 2.* Coverages (left) and confidence intervals (right) for our method as well as the 3 other methods that could efficiently run on this data. Only our method achieves the nominal coverage rate.

t-statistic instead of a z-statistic (Greene, 2012, pp. 50–52).

Sandwich Estimator. The sandwich estimator (Huber, 1967; White, 1980a;b) uses the same point estimate as OLS but a different variance estimate. We take the variance estimate from MacKinnon and White (1985, Equation 6):  $\frac{1}{N-P}e_p^{\rm T}(X^{\rm T}X)^{-1}(X^{\rm T}RX)(X^{\rm T}X)^{-1}e_p$ , where *R* is a diagonal matrix with the squared residuals as entries.

*Importance Weighted Least Squares (KDEIW).* As suggested by Shimodaira (2000, Section 9), we calculate importance weights via kernel density estimation (KDE). We select the bandwidth parameter with 5-fold cross validation; see Appendix B.7. Given the KDE weights, we use the point estimate and confidence interval from weighted least squares.

Generalized Least Squares (GLS). We maximize the likelihood of  $Y \sim \mathcal{N}(\theta^T X, \Sigma)$ , with  $\Sigma$  specified by an isotropic Matérn 3/2 covariance function and a nugget, to select the parameters of the covariance function and nugget variance. Then we use the restricted spatial-regression framework (Hodges and Reich, 2010); since we project the spatially-correlated error term onto the orthogonal complement of the covariates, the point estimate coincides with OLS.

Gaussian Process Bayesian Credible Intervals (GP BCIs). We use the model  $Y(S) = \theta^T X(S) + g(S) + \epsilon$ , with  $\theta^T \sim \mathcal{N}(0, \lambda^2 I_P), g \sim \mathcal{GP}(0, k_{\gamma}), k_{\gamma}$  an isotropic Matérn 3/2 kernel function with hyperparameters  $\gamma$ , and  $\epsilon \sim \mathcal{N}(0, \delta^2)$ . We select  $\{\lambda, \gamma, \delta\}$  by maximum likelihood. We report posterior credible intervals for  $\theta_p$ .

**Single Covariate Simulation.** In our first simulation, the source locations are uniform on  $S = [-1, 1]^2$  (blue points in Figure 5 left plots). The target locations are uniform on  $[\frac{-1+\mathrm{shift}}{1+|\mathrm{shift}|}, \frac{1+\mathrm{shift}}{1+|\mathrm{shift}|}]$ , where shift controls the degree of distribution shift between source and target (orange points in Figure 5 left plots). In this experiment, the single covariate  $X = \chi(S) = S^{(1)} + S^{(2)}$  (Figure 5, third plot). And

the response is  $Y = X + \frac{1}{2}((S^{(1)})^2 + (S^{(2)})^2) + \epsilon$ , with  $\epsilon \sim \mathcal{N}(0, 0.1^2)$ . Figure 5, fourth plot, shows the conditional expectation of the response given location. We can compute the ground truth parameter in closed form because we have access to the conditional expectation of the response (Equation (7)). We vary shift  $\in [0, \pm 0.2, \pm 0.4, \pm 0.6, \pm 0.8]$  and run 250 seeds for each shift.

Figure 1, left, shows that only our method and the GP consistently achieve nominal coverage. Given correct coverage, narrower (i.e., more precise) confidence intervals are desirable; Figure 1, right, shows that our method yields narrower intervals than the GP. KDEIW comes close to achieving nominal coverage when there is no shift. But under any shift, it is not able to fully debias the estimate, and coverage drops. For large M, we expect the sandwich estimator to achieve nominal coverage at shift = 0 since it is guaranteed to cover the population under first-order misspecification without distribution shift (Huber, 1967; White, 1980b). But under any of the depicted non-zero shifts, the sandwich, OLS, and GLS achieve zero coverage.

The extreme narrowness of the OLS, sandwich, KDEIW, and GLS intervals (Figure 1, right) suggests that the problem with these methods is exactly their overconfidence. Because these approaches assume that the estimator is unbiased (or can be debiased), and that errors are independent and Gaussian, their intervals contract far too quickly, even with small amounts of data (here, N = 300). Essentially, these methods' reliance on strong modeling assumptions leads to non-robust coverage.

We note that the confidence intervals and coverages for each method have (approximately) the same values for either  $\pm$ shift in this experiment (Figure 1, right) because the covariate is symmetric around the line  $S^{(1)} = S^{(2)}$ ; see the middle right plot of Figure 5 in Appendix C. So positive and negative shift values move the source and target loca-



*Figure 3.* Left: the confidence interval width of our method as a function of shift for each Lipschitz constant *L*. All *L* yield coverage of 1.0. Middle and right: the confidence interval width (solid line with dot marker) as a function of the Lipschitz constant for shift = 0 (middle) and shift = 0.8 (right). The vertical axis is shared across all three plots. The bias contribution to the width (dashed line, x marker) is monotonically increasing in *L*. The randomness contribution (dashed line, square marker) is monotonically decreasing.

tions symmetrically across the  $S^{(1)} = S^{(2)}$  line. We do not expect such a symmetry for general covariates and will not see it in our next simulation (Figure 2, right).

Simulation with Several Covariates. Our second simulation generates locations as in the previous experiment. Now we use N = 10,000 and M = 100. And we generate 3 covariates,  $X^{(1)} = \sin(S^{(1)}) + \cos(S^{(2)})$ ,  $X^{(2)} = \cos(S^{(1)}) - \sin(S^{(2)})$  and  $X^{(3)} = S^{(1)} + S_2$ . The response is  $Y = X^{(1)}X^{(2)} + \frac{1}{2}((S^{(1)})^2 + (S^{(2)})^2) + \epsilon$  with  $\epsilon \sim \mathcal{N}(0, 0.1^2)$ . We focus on inference for the first coefficient. Calculation of  $\hat{\sigma}_N^2$  scales poorly with N due to needing to solve a quadratic program. So here we instead estimate  $\sigma^2$  using the squared error of leave-one-out 1-nearest neighbor regression fit on the source data. See Appendix B.6 for details. We compare against the same set of methods except we do not include GLS or GP BCIs since these require further approximations to scale for this N.

We again find that our method achieves coverage while the other methods do not (Figure 2, left). In fact, in this experiment, no other method achieves coverage over 30% across any shift value (even 0). As before, we see that competing methods are overconfident, with very small CI widths (Figure 2, right).

For methods besides our own, coverage levels at 0 are generally lower in this experiment at shift = 0 than in the previous experiment. The difference is that N is much larger here (making confidence intervals narrower and exacerbating overconfidence) while M is the same. The sandwich estimator covers the analogue of  $\theta^*_{OLS,p}$  where spatial locations are treated as random. So it has good coverage at shift = 0 when  $M \gg N$ , but not when  $M \ll N$ .

Effect of Lipschitz Constant on Confidence Intervals in Simulation Experiment. In the experiments above, we know the minimum value ( $L_0$ ) for which assumption 4 holds. Above we fit our method with  $L = L_0$  ( $L_0 = 2\sqrt{2}$  and  $3\sqrt{2}$ , respectively). Now we repeat the first simulation but vary  $L \in \{0.1, 0.5, 1.0, 2.0, 3.5, 5, 7.5, 10\}$ . All of these L values yield coverage of 1.0 for our method, above the nominal coverage of 0.95, even though coverage is not guaranteed by our theory for  $L \le 2\sqrt{2} \approx 2.8$ .

We next demonstrate how the confidence interval width reflects a bias-randomness trade-off as L varies. If the noise were known, the confidence interval would be monotonically increasing in L. Since it is unknown, only the contribution of the bias to the interval width (2B, Step 4 in Algorithm 1) increases (Figure 3, middle and left, x marker). Conversely, smaller values of L yield larger values for  $\hat{\sigma}_N^2$ , so the randomness contribution to the interval width (2c, Step 5 in Algorithm 1, square marker) increases (Figure 3, middle and left). The full confidence interval width,  $2B + 2c\Delta(\alpha)$ , is non-monotonic in L.

Tree Cover Linear Regression. We use a linear regression model  $Y_{\text{Tree Cover }\%} = \sum_{p \in \mathcal{P}} \theta_p X_p$  to quantify how tree cover percentage in the contiguous United States (CONUS) in the year 2021 relates to three variables,  $\mathcal{P} =$ {Aridity\_Index, Elevation, Slope}. We use the 983 data points from Lu et al. (2024), who in turn draw on (USFS, 2023; Trabucco, 2019; NASA Jet Propulsion Lab, 2020). We define our target region in the West portion of CONUS as locations with latitude in the range (25, 50) and longitude in the range (-125, -110). Out of all points in this region, we designate 50% — totaling 133 sites — as target data. Next, we select the source data by taking a uniform random sample of 20% of the remaining spatial locations, repeated over 250 random seeds to assess coverage performance. Each seed yields 170 source locations. Figure 8 illustrates the spatial split between source and target data for a representative seed. We discuss the data and our pre-processing in detail in Appendix D.

We compare coverages for confidence intervals of the three parameters,  $\theta_{\text{Aridity_Index}}$ ,  $\theta_{\text{Elevation}}$ , and  $\theta_{\text{Slope}}$ . We discuss how we evaluate coverage in Appendix D.3. In the top



*Figure 4.* Coverages (upper) and confidence interval widths (lower) for our method as well as 5 other methods. Each column represents a parameter in the tree cover experiment. Only our method consistently achieves the nominal coverage.

row of Figure 4, we see that our method is the only one to achieve the 95% nominal coverage for all three parameters. Conversely, for the *Slope* parameter, every other method achieves coverage at most 54%. In the bottom row of Figure 11, we again see that alternative methods fail to provide coverage due to their overconfidence (small widths). To see the confidence intervals themselves (rather than just the summary plots here), in Figure 11, we depict all methods' constructed CIs across all three parameters for 7 of the 250 seeds. In Appendix D.5, we conduct a similar analysis but with target locations in the Southeast, rather than West, of CONUS. The results align with our discussion here.

Choice of Lipschitz Constant in the Tree Cover Experiment. For the tree cover experiment, we leverage domain knowledge to set the Lipschitz constant to L = 0.2, in units of percent tree cover per kilometer (km). This choice implies that a 1% change in tree cover corresponds to moving 1/0.2 = 5 km. To arrive at this choice, we observe that in certain regions of the U.S., such as the Midwest, tree coverage remains relatively uniform over several kilometers, so smaller Lipschitz constants (e.g., L = 0.02, corresponding to a 1% change over 50 km) would be appropriate. However, in other regions - such as the western U.S., where elevation changes are more pronounced (e.g., the Rockies, California, and the Pacific Northwest) - tree cover can change sharply over short distances. To account for such variations conservatively, we choose L = 0.2. More generally, for real-world applications, we recommend the following strategy: (i) use domain knowledge to select a reasonable Lipschitz constant for the response variable, and (ii) inflate the Lipschitz constant to ensure a conservative estimate (which is more likely to satisfy assumption 4).

Effect of Lipschitz Constant on Confidence Intervals in Tree Cover Experiment. In Figure 10, we show the coverage and width of our confidence intervals across three orders of magnitude of Lipschitz constants (L from 0.001 to 1). For L varying between 0.1 and 1 (a single order of magnitude variation around our chosen value of 0.2), we find that coverage is always met except in one case (Aridity Index with L = 0.1), where it is very close to nominal (89% instead of 95%). For Slope and Elevation, coverage is met or very nearly met for all L values. For Aridity Index, coverage is low for  $L \le 0.1$ . Meanwhile, confidence intervals become noticeably wider for L > 0.5while remaining relatively stable for smaller values. These results support our intuition to err on the side of larger Lvalues to be conservative and maintain coverage.

### 5. Discussion

We propose a new method for constructing confidence intervals for spatial linear models. We show via theory and experiments that our intervals accurately quantify uncertainty under misspecification and covariate shift. In experiments, our method is the only method that consistently achieves (or even comes close to) nominal coverage. We observe that, very commonly in spatial data analyses, covariates and responses may be observed at different locations in space. Since our method does not actually use the source covariate values in inference for  $\theta_{OLS}^*$ , it can be applied in this common missing-data scenario. Though it requires additional work, we believe the ideas here will extend naturally to the widely used class of generalized linear models.

### Acknowledgments

This work was supported in part by an Office of Naval Research Early Career Grant, Generali, and a Microsoft Trustworthy AI Grant.

# References

- Agrawal, A., Verschueren, R., Diamond, S., and Boyd, S. (2018). A rewriting system for convex optimization problems. *Journal of Control and Decision*, 5(1):42–60.
- Aitken, A. C. (1936). IV.—on least squares and linear combination of observations. *Proceedings of the Royal Society of Edinburgh*, 55:42–48.
- Ben-David, S., Blitzer, J., Crammer, K., and Pereira, F. (2006). Analysis of representations for domain adaptation. In Advances in Neural Information Processing Systems.
- Brent, R. P. (1971). An algorithm with guaranteed convergence for finding a zero of a function. *The Computer Journal*, 14(4):422–425.
- Brunsdon, C., Fotheringham, A. S., and Charlton, M. E. (1996). Geographically weighted regression: a method for exploring spatial nonstationarity. *Geographical analysis*, 28(4):281–298.
- Buja, A., Brown, L., Berk, R., George, E., Pitkin, E., Traskin, M., Zhang, K., and Zhao, L. (2019). Models as Approximations I: Consequences Illustrated with Linear Regression. *Statistical Science*, 34(4):523–544.
- Castro Torres, A. F. and Akbaritabar, A. (2024). The use of linear models in quantitative research. *Quantitative Science Studies*, 5(2):426–446.
- Cressie, N. (2015). *Statistics for spatial data*. John Wiley & Sons.
- Csurka, G. (2017). A Comprehensive Survey on Domain Adaptation for Visual Applications, pages 1–35. Springer International Publishing.
- Diamond, S. and Boyd, S. (2016). CVXPY: A Pythonembedded modeling language for convex optimization. *Journal of Machine Learning Research*, 17(83):1–5.
- Doshi-Velez, F. and Kim, B. (2017). Towards a rigorous science of interpretable machine learning. *arXiv preprint arXiv:1702.08608*.
- Flamary, R., Courty, N., Gramfort, A., Alaya, M. Z., Boisbunon, A., Chambon, S., Chapel, L., Corenflos, A., Fatras, K., Fournier, N., Gautheron, L., Gayraud, N. T., Janati, H., Rakotomamonjy, A., Redko, I., Rolet, A., Schutz, A., Seguy, V., Sutherland, D. J., Tavenard, R., Tong, A., and Vayer, T. (2021). POT: Python Optimal Transport. *Journal of Machine Learning Research*, 22(78):1–8.

- Fotheringham, A. S., Brunsdon, C., and Charlton, M. (2002). Geographically Weighted Regression: The Analysis of Spatially Varying Relationships. Wiley.
- Gilbert, B., Ogburn, E. L., and Datta, A. (2024). Consistency of common spatial estimators under spatial confounding. *Biometrika*.
- Goulart, P. J. and Chen, Y. (2024). Clarabel: An interiorpoint solver for conic programs with quadratic objectives. *arXiv preprint arXiv:2405.12762*.
- Gramacy, R. B. (2020). Surrogates: Gaussian Process Modeling, Design, and Optimization for the Applied Sciences. Chapman and Hall/CRC.
- Greene, W. H. (2012). *Econometric Analysis: 7th Edition*. USA: Pearson Prentice Hall.
- Gretton, A., Smola, A., Huang, J., Schmittfull, M., Borgwardt, K., and Schölkopf, B. (2008). Covariate shift by kernel mean matching. In *Dataset Shift in Machine Learning*. MIT Press.
- Heaton, M. J., Datta, A., Finley, A. O., Furrer, R., Guinness, J., Guhaniyogi, R., Gerber, F., Gramacy, R. B., Hammerling, D., Katzfuss, M., et al. (2019). A case study competition among methods for analyzing large spatial data. *Journal of Agricultural, Biological and Environmental Statistics*, 24:398–425.
- Hinkley, D. V. (1977). Jackknifing in unbalanced situations. *Technometrics*, 19(3):285–292.
- Hodges, J. S. and Reich, B. J. (2010). Adding spatiallycorrelated errors can mess up the fixed effect you love. *The American Statistician*, 64(4):325–334.
- Huber, P. J. (1967). The behavior of maximum likelihood estimates under nonstandard conditions. In *Proceedings* of the Fifth Berkeley Symposium on Mathematical Statistics and Probability, Volume 1: Statistics, pages 221–234. University of California Press.
- Kanamori, T., Suzuki, T., and Sugiyama, M. (2012). Statistical analysis of kernel-based least-squares density-ratio estimation. *Machine Learning*, 86:335–367.
- Kirszbraun, M. D. (1934). Über die zusammenziehende und lipschitzsche transformationen. Fundamenta Mathematicae, 22:77–108.
- Lee, J., Costello, S., Balmes, J. R., and Holm, S. M. (2022). The Association between Ambient PM2.5 and Low Birth Weight in California. *International Journal of Environmental Research and Public Health*, 19(20):13554.

- Li, C., Gao, D., Cai, Y. S., Liang, J., Wang, Y., Pan, Y., Zhang, W., Zheng, F., and Xie, W. (2023). Relationships of Residential Distance to Major Traffic Roads with Dementia Incidence and Brain Structure Measures: Mediation Role of Air Pollution. *Health Data Science*, 3:0091.
- Liu, D. C. and Nocedal, J. (1989). On the limited memory BFGS method for large scale optimization. *Math. Program.*, 45(1–3):503–528.
- Lu, K., Bates, S., and Wang, S. (2024). Quantifying uncertainty in area and regression coefficient estimation from remote sensing maps. arXiv preprint arXiv:2407.13659.
- MacKinnon, J. G. and White, H. (1985). Some heteroskedasticity-consistent covariance matrix estimators with improved finite sample properties. *Journal of Econometrics*, 29(3):305–325.
- Matthews, A. G. d. G., van der Wilk, M., Nickson, T., Fujii, K., Boukouvalas, A., León-Villagrá, P., Ghahramani, Z., and Hensman, J. (2017). GPflow: A Gaussian process library using TensorFlow. *Journal of Machine Learning Research*, 18(40):1–6.
- Mayor, J. R., Sanders, N. J., Classen, A. T., Bardgett, R. D., Clement, J.-C., Fajardo, A., Lavorel, S., Sundqvist, M. K., Bahn, M., Chisholm, C., et al. (2017). Elevation alters ecosystem properties across temperate treelines globally. *Nature*, 542(7639):91–95.
- Müller, U. K. (2013). Risk of Bayesian inference in misspecified models, and the sandwich covariance matrix. *Econometrica*, 81(5):1805–1849.
- NASA Jet Propulsion Lab (2020). NASADEM merged DEM global 1 arc second v001 [data set]. NASA EOSDIS Land Processes DAAC.
- Nobre, W. S., Schmidt, A. M., and Pereira, J. B. M. (2021). On the Effects of Spatial Confounding in Hierarchical Models. *International Statistical Review*, 89(2):302–322.
- Paciorek, C. J. (2010). The Importance of Scale for Spatial-Confounding Bias and Precision of Spatial Regression Estimators. *Statistical Science*, 25(1):107–125.
- Page, G. L., Liu, Y., He, Z., and Sun, D. (2017). Estimation and Prediction in the Presence of Spatial Confounding for Spatial Linear Models. *Scandinavian Journal of Statistics*, 44(3):780–797.
- Pan, S. J. and Yang, Q. (2010). A survey on transfer learning. *IEEE Transactions on Knowledge and Data Engineering*, 22(10):1345–1359.
- Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M., Prettenhofer, P.,

Weiss, R., Dubourg, V., Vanderplas, J., Passos, A., Cournapeau, D., Brucher, M., Perrot, M., and Duchesnay, E. (2011). Scikit-learn: Machine learning in Python. *Journal* of Machine Learning Research, 12:2825–2830.

- Peyré, G. and Cuturi, M. (2019). Computational Optimal Transport: With Applications to Data Science. *Foundations and Trends*® *in Machine Learning*, 11(5-6):355– 607.
- Portier, F., Truquet, L., and Yamane, I. (2024). Nearest neighbor sampling for covariate shift adaptation. *arXiv* preprint 2312.09969.v2.
- Que, Q. and Belkin, M. (2013). Inverse density as an inverse problem: the fredholm equation approach. In *Advances in Neural Information Processing Systems*, pages 1484–1492.
- Rasmussen, C. E. and Williams, C. K. I. (2006). *Gaussian Processes for Machine Learning*. The MIT Press.
- Rolf, E., Klemmer, K., Robinson, C., and Kerner, H. (2024). Position: Mission critical – satellite data is a distinct modality in machine learning. In *Proceedings of the 41st International Conference on Machine Learning*.
- Rudin, C. (2019). Stop explaining black box machine learning models for high stakes decisions and use interpretable models instead. *Nature machine intelligence*, 1(5):206– 215.
- Rudin, W. (1976). Principles of Mathematical Analysis. International series in pure and applied mathematics. McGraw-Hill.
- Sandel, B. and Svenning, J.-C. (2013). Human impacts drive a global topographic signature in tree cover. *Nature Communications*, 4(1):2474.
- Seabold, S. and Perktold, J. (2010). statsmodels: Econometric and statistical modeling with python. In *9th Python in Science Conference*.
- Shimodaira, H. (2000). Improving predictive inference under covariate shift by weighting the log-likelihood function. *Journal of Statistical Planning and Inference*, 90(2):227–244.
- Sugiyama, M., Suzuki, T., and Kanamori, T. (2012). *Density Ratio Estimation in Machine Learning*. Cambridge University Press.
- Sugiyama, M., Suzuki, T., Nakajima, S., Kashima, H., von Bünau, P., and Kawanabe, M. (2008). Direct importance estimation for covariate shift adaptation. *Annals of the Institute of Statistical Mathematics*, 60(4):699–746.

- Trabucco, A. (2019). Global aridity index and potential evapotranspiration (ET0) climate database v2. *CGIAR Consortium for Spatial Information*.
- Tuia, D., Persello, C., and Bruzzone, L. (2016). Domain adaptation for the classification of remote sensing data: An overview of recent advances. *IEEE Geoscience and Remote Sensing Magazine*, 4(2):41–57.
- USFS (2023). USFS tree canopy cover v2021.4 (Conterminous United States and Southeastern Alaska).
- Valentine, F. A. (1945). A Lipschitz condition preserving extension for a vector function. *American Journal of Mathematics*, 67(1):83–93.
- Villani, C. (2009). *Optimal Transport: Old and New*. Springer.
- Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D., Burovski, E., Peterson, P., Weckesser, W., Bright, J., van der Walt, S. J., Brett, M., Wilson, J., Millman, K. J., Mayorov, N., Nelson, A. R. J., Jones, E., Kern, R., Larson, E., Carey, C. J., Polat, İ., Feng, Y., Moore, E. W., VanderPlas, J., Laxalde, D., Perktold, J., Cimrman, R., Henriksen, I., Quintero, E. A., Harris, C. R., Archibald, A. M., Ribeiro, A. H., Pedregosa, F., van Mulbregt, P., and SciPy 1.0 Contributors (2020). SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python. *Nature Methods*, 17:261–272.
- Wainwright, M. J. (2019). *High-Dimensional Statistics: A Non-Asymptotic Viewpoint*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, Cambridge.
- Walker, S. G. (2013). Bayesian inference with misspecified models. *Journal of Statistical Planning and Inference*, 143(10):1621–1633.
- Weber, L. M., Saha, A., Datta, A., Hansen, K. D., and Hicks, S. C. (2023). nnSVG for the scalable identification of spatially variable genes using nearest-neighbor Gaussian processes. *Nature Communications*, 14(1):1–12.
- Westervelt, D. M., Conley, A. J., Fiore, A. M., Lamarque, J.-F., Shindell, D. T., Previdi, M., Mascioli, N. R., Faluvegi, G., Correa, G., and Horowitz, L. W. (2018). Connecting regional aerosol emissions reductions to local and remote precipitation responses. *Atmospheric Chemistry and Physics*, 18(16):12461–12475.
- White, H. (1980a). A heteroskedasticity-consistent covariance matrix estimator and a direct test for heteroskedasticity. *Econometrica*, 48(4):817–838.
- White, H. (1980b). Using least squares to approximate unknown regression functions. *International Economic Review*.

Yu, H., Fotheringham, A. S., Li, Z., Oshan, T., and Wolf, L. J. (2020). On the measurement of bias in geographically weighted regression models. *Spatial Statistics*, 38:100453.

# **A. Extended Related Work**

In this section, we discuss related work on bias in linear regression in spatial settings, local approaches to regression that often rely on data-borrowing strategies similar to our nearest neighbor approach, and covariate shift.

Bias in Spatial Regression Linear models with a Gaussian process random effect — that is, models of the form

$$Y(S) = \theta^{\mathrm{T}} X(S) + g(S) + \epsilon, \tag{19}$$

with g a Gaussian process and  $\epsilon$  independent and identically distributed noise — are a classic tool in spatial regression and remain widely used in applications (Weber et al., 2023; Gramacy, 2020; Heaton et al., 2019). Whether to treat the covariates as fixed functions of spatial location or random is a topic of significant debate. In order for the model to be identifiable, it is generally necessary for the covariates to be thought of as random (Gilbert et al., 2024, Proposition 1). However, Paciorek (2010) observed it may be more reasonable to think of covariates as fixed (the perspective we also take). If X and g are not independent (in the random case) or close to orthogonal in the fixed case, Paciorek (2010) illustrates that bias is introduced into the estimation of  $\theta$ . And that the degree of this bias depends on the relative scales over which X and g vary. He gives a closed form for this bias under strong linear-Gaussian assumptions that depend on parameters that would generally be unknown. Page et al. (2017) builds on the result of Paciorek (2010), and shows that bias in predictions made by the spatial model due to confounding may be small, despite biases in parameter estimation. Hodges and Reich (2010) proposed using restricted spatial regression, essentially explaining as much of the variation in the response as possible by the covariates to reduce bias in estimation due to spatially-correlated error. We take this approach in the GLS baseline used in our experiments. Nobre et al. (2021) extended earlier work about bias due to spatial confounding by considering the case when multiple observations are available at the same spatial location with independent noise. They showed that spatial regression models can still be biased with repeated observations due to confounding.

In our work, we focus on bias in estimation of the target conditional OLS. Because we do not assume the model is wellspecified, there is no 'true' value of the parameter to estimate. Instead, our goal is to estimate the 'best' (in a least squares sense) linear approximation to the response. We focus on the case when X is a fixed function of location (assumption 1) and make weak non-parametric, smoothness assumptions (assumption 4). This contrasts with prior works that generally assume the covariates are random and make linear-Gaussian assumptions to calculate the bias (Paciorek, 2010; Page et al., 2017). Our work shows how to directly incorporate bias into confidence intervals for  $\theta$  under our weak assumptions, but does not address issues of (non-)identifiability in the estimation problem.

Gilbert et al. (2024) showed that despite finite-sample bias in estimation of  $\theta$  under spatial confounding, consistent estimation of  $\theta$  is possible in the identifiable case when X is random under infill asymptotics. This is essentially orthogonal to our work, as we focus on (finite-sample valid) confidence intervals in the case when X is fixed.

### Local Versus Global Regression Estimates

An alternative perspective on regression in spatial settings, referred to as *geographically weighted regression* (GWR), focuses on estimating the association between the response and covariates at each location in space (Brunsdon et al., 1996; Fotheringham et al., 2002). For this regression problem to be well-defined, the covariates at each spatial location must be treated as random variables. Like our method, this approach uses nearest neighbors or local smoothing approaches to estimate the local coefficients at each location in space. And — as in our work — because of this "data-borrowing" from nearby locations, bias is introduced into the estimation of these local regression coefficients. The confidence intervals reported for GWR typically do not account for this bias (Yu et al., 2020). While Yu et al. (2020) provide a formula for the bias introduced by "data-borrowing", their formula relies on the assumption that the model is well-specified and depends on unobservable parameters.

We focus on the estimation bias in the estimation of global parameters. Although local parameter estimates are often of interest in spatial analyses (Fotheringham et al., 2002, Chapter 1), global estimates are also useful as summary statistics for making decisions that impact regions or multiple localities. And we characterize the bias in these estimates under weak, non-parametric assumptions and incorporate it into our confidence intervals. Extensions of our approach to GWR coefficients are a promising direction for future research. However, this extension is not simple, as it involves accounting for randomness in the covariates.

**Covariate Shift and Importance Weighting.** Differences between the source locations (with observed responses) and the target locations (at which we want to study the relationship between variables) can be seen as an instance of covariate

shift (Rolf et al., 2024; Tuia et al., 2016; Weber et al., 2023). Covariate shift is often dealt with by importance weighting, reweighting each training example to account for differences in the density of the source and target distributions. Prior work in spatial machine learning has generally focused on addressing covariate shift in the context of estimating a method's prediction error. However, these approaches can, in principle, be extended to parameter estimation in OLS, mirroring the broader covariate shift literature (Shimodaira, 2000).

While in the main text we focused on density ratio estimation using kernel density estimation (Shimodaira, 2000), many other approaches for density ratio estimation can be used (Sugiyama et al., 2012; Kanamori et al., 2012; Gretton et al., 2008; Que and Belkin, 2013; Sugiyama et al., 2008). Importance weighting de-biases the estimator, but can only be used if the distribution of the source data is supported on a region containing the test data. And the confidence intervals obtained by importance weighting with estimated weights do not account for errors in estimation of the density ratio (if this exists). These confidence intervals are therefore not guaranteed to achieve the nominal coverage rate. And they are not applicable in cases where extrapolation is required, as the estimator cannot necessarily be de-biased in these cases.

In contrast, our approach ensures nominal coverage even when de-biasing the estimate via importance weighting is not possible. This advantage comes at the cost of an additional regularity assumption (assumption 4), as well as often empirically wider confidence intervals.

# **B.** Implementation Details

In this section, we describe the implementation details of our method. Particularly, this involves computing the upper bound on the bias, computation of  $\Delta$  in Lemma 6 and computation of  $\hat{\sigma}_N^2$  from Equation (12).

#### **B.1.** Selecting the Matrix $\Psi$ .

In the main text we selected  $\Psi$  by using the nearest source location to each target location. We now provide additional discussion about alternative choices of  $\Psi$  and possible trade-offs.

A generalization of the 1-nearest neighbor approach is to consider a  $\Psi$  matrix determined by K-nearest neighbors.

Definition 10 (Nearest-Neighbor Weight Matrix). Define the K-nearest neighbor weight matrix by

$$\Psi_{mn} = \begin{cases} \frac{1}{K} & S_n \in \{K \text{ closest source locations to } S_m^* \}\\ 0 & \text{otherwise} \end{cases}.$$
(20)

For definiteness, we assume that, if multiple sources are the same distance from a target, ties are broken uniformly at random.

When using a K-nearest neighbor matrix, there is an inherent bias–variance trade-off in choosing K. Increasing K broadens the geographic range of source observations used, hence the bias will increase; however, it also lowers the variance by averaging over more responses. We do not investigate this trade-off in detail. In the experiments we ran, we found 1-nearest neighbor to work well. This is consistent with results in the mean estimation literature suggesting that if source and target distributions overlap substantially, the variance term remains manageable, even when using 1-nearest neighbor approaches (Portier et al., 2024).

### **B.2. Implementation of the Wasserstein Bound Calculation**

We show in Appendix E.3 how Equation (16) reduces to computing a Wasserstein-1 distance between empirical measures. To implement the Wasserstein distance calculation, we rely on the Python Optimal Transport library (Flamary et al., 2021). For the simulated experiments, we compute the cost matrix using Euclidean distances between spatial locations. For the real-world experiment, we use the Haversine distance to account for Earth's curvature.

#### B.3. Use of Confidence Interval Lemma 6

In all experiments, we construct confidence intervals as in Lemma 6. A simpler alternative, which guarantees  $1 - \alpha$  coverage for all Gaussian distributions  $\mathcal{N}(b, c^2)$  with  $b \in [-B, B]$ , is to form two-sided confidence intervals for each  $b \in [-B, B]$  and then take their union. This produces an interval of the form

$$[-B - c \Phi^{-1}(1 - \frac{\alpha}{2}), B + c \Phi^{-1}(1 - \frac{\alpha}{2})]$$

However, the confidence intervals from Lemma 6 are never longer than this union-based approach, and the root-finding step required to compute them adds negligible overhead. Consequently, we opt for the intervals in Lemma 6 in all our experiments.

### **B.4.** Computation of $\Delta$ for Lemma 6

Define  $g(\Delta) = \Phi(\Delta) - \Phi(-\frac{2B}{c} - \Delta) - 1 + \alpha$ . Our goal is to find a root of g in the interval  $[\Phi^{-1}(1 - \alpha), \Phi^{-1}(1 - \alpha/2)]$ . We first show that g is monotonic, that a root exists in this interval, and that the root is unique.

Differentiating, we see  $g'(\Delta) = \phi(\Delta) + \phi(-\frac{2B}{c} - \Delta)$ , where  $\phi$  denotes the Gaussian probability density function. This is strictly positive, and so g is strictly monotone increasing.

Also,

$$g(\Phi^{-1}(1-\alpha)) = 1 - \alpha - \Phi\left(-\frac{2B}{c} - \Phi^{-1}(1-\alpha)\right) - 1 + \alpha < 1 - \alpha - 1 + \alpha = 0.$$
 (21)

by non-negativity of the CDF. And,

$$g(\Phi^{-1}(1-\alpha/2)) = 1 - \alpha/2 - \Phi\left(-\frac{2B}{c} - \Phi^{-1}(1-\alpha/2)\right) - 1 + \alpha$$
(22)

$$= \alpha/2 - \left(1 - \Phi\left(\frac{2B}{c} + \Phi^{-1}(1 - \alpha/2)\right)\right).$$
(23)

We used symmetry of the Gaussian in the second equality. Then,

$$\Phi\left(\frac{2B}{c} + \Phi^{-1}(1 - \alpha/2)\right) \ge 1 - \alpha/2,\tag{24}$$

and so  $g(\Phi^{-1}(1 - \alpha/2)) \le 0$ . We conclude that g has a root in the interval. By strict monotonicity of g, this root is unique. We use Brent's method (Brent, 1971) as implemented in Scipy (Virtanen et al., 2020) to compute this root numerically.

### **B.5.** Computation of $\hat{\sigma}_N^2$ via quadratic programming

To estimate the noise parameter, we need to solve the minimization problem

$$\hat{\sigma}_N^2 = \inf_{g \in \mathcal{F}_L} \frac{1}{N} \sum_{n=1}^N (Y_n - g(S_n))^2.$$
(25)

The first obstacle is that the infimum is taken over an infinite-dimensional space. However, by the Kirszbraun theorem (Kirszbraun, 1934; Valentine, 1945), every *L*-Lipschitz function defined on a subset of  $\mathbb{R}^D$  or  $\mathbb{S}^D$  can be extended to an *L*-Lipschitz function on the whole domain. Since our objective function depends only on the values of g at the source spatial locations, we only need to enforce the Lipschitz condition between all pairs of source spatial locations, not over the entire domain. Enforcing the Lipschitz condition at these N source locations amounts to N(N-1)/2 linear inequality constraints.

Particularly, if we define  $G = (g(S_1), g(S_2), \dots, g(S_n))$ , then we have the constraints

$$AG \le L \operatorname{vec}(\Gamma) \tag{26}$$

where L is the Lipschitz constant,  $\Gamma \in \mathbb{R}^{N^2-N}$  is the matrix of pairwise distances between distinct points in S, and  $A \in \mathbb{R}^{(N^2-N)\times N}$  is a sparse matrix with exactly one 1 and one -1 in distinct rows of each column, representing all such pairs in its rows.

The objective is a symmetric, positive-definite quadratic form since it is a sum of squares. Therefore, the optimization problem is a quadratic program.

In practice, we use the Scipy sparse matrix algebra (Virtanen et al., 2020) and the CLARABEL solver (Goulart and Chen, 2024) through the CVXPY optimization interface (Diamond and Boyd, 2016; Agrawal et al., 2018) to solve this quadratic program.

# **B.6. Scalable Estimation of** $\sigma^2$

The quadratic programming approach for estimating  $\sigma^2$  outlined in the main text (Equation (12)) and described in detail in the previous section does not scale well to large numbers of source locations. Therefore, for our synthetic experiment with N = 10,000, we take a different approach.

For  $1 \le n \le N$ , let  $\eta(n) = \arg \min_{n' \ne n} d_{\mathcal{S}}(S_n, S'_n)$ . That is  $\eta(n)$  is the index of the nearest point to  $S_n$ . Define the estimator,

$$\tilde{\sigma}_N^2 = \frac{1}{2N} \sum_{n=1} (Y_n - Y_{\eta(n)})^2.$$
(27)

Then as long as  $d_{\mathcal{S}}(S_n, S'_n) \approx 0$ , by the Lipschitz assumption,  $Y_n - Y_{\eta(n)} \approx \epsilon_n - \epsilon_{\eta(n)}$ , and so

$$\tilde{\sigma}_N^2 \approx \frac{1}{2N} \sum_{n=1}^N (\epsilon_n - \epsilon_{\eta(n)})^2 = \frac{1}{2N} \left( \sum_{n=1}^N \epsilon_n - \sum_{n=1}^N \epsilon_n \epsilon_{\eta(n)} + \sum_{n=1}^N \epsilon_{\eta(n)}^2 \right).$$
(28)

Then,

$$\mathbb{E}[\tilde{\sigma}_N^2] = \sigma^2. \tag{29}$$

In general, we expect the estimate to concentrate around its expectation, provided that no single point in the source data is the nearest neighbor of too many other points in the source data.

### **B.7. Implementation of Baseline Methods**

We now describe the details of the implementation of the baseline methods.

**Ordinary Least Squares.** We use the ordinary least squares implementation from statsmodel (Seabold and Perktold, 2010). We use the default implementation, which calculates the variance as the average sum of squared residuals with a degrees-of-freedom correction, as in Equation (4). We use a t-statistic to compute the corresponding confidence interval, which is again the default in statsmodel (Seabold and Perktold, 2010).

Sandwich Estimator. We use the sandwich estimation procedure included in ordinary least squares in statsmodels (Seabold and Perktold, 2010). We use the HC1 function, which implements the sandwich estimator with the degrees-of-freedom correction from Hinkley (1977); MacKinnon and White (1985). That is, the variance is estimated as  $\frac{1}{N-P}e_p^{\rm T}(X^{\rm T}X)^{-1}(X^{\rm T}RX)(X^{\rm T}X)^{-1}e_p$ . We use the default settings in statsmodels, which use a z-statistic with the sandwich estimator to compute the corresponding confidence interval.

**Importance Weighted Least Squares.** We use the scikit-learn (Pedregosa et al., 2011) implementation of kernel density estimation to estimate the density of test and train point separately. We use a Gaussian kernel (the default) and perform 5-fold cross validation to select the bandwidth parameter, maximizing the average log likelihood of held-out points in each fold. For the simulation experiments, we performed cross-validation to select the bandwidth parameters over the set  $\{0.01, 0.025, 0.05, 0.1, 0.25, 0.5\}$ . We selected this set of possible bandwidths to span several orders of magnitude from very short bandwidths, to bandwidths on the same order as the entire domain. We select the bandwidths for the source and target density estimation problems separately. For the real data experiments, the bandwidths was selected from the set  $\{0.002, 0.005, 0.01, 0.02, 0.05, 0.1, 0.2, 0.5, 1\}$ . This range was selected since the maximum Haversine distance between points in the spatial domain of interest is approximately 1. Once density estimates are obtained, we evaluate the ratio of the density function on the training locations, and use these as weights to perform weighted least squares. Weighted least squares is performed using the default settings in statsmodels (Seabold and Perktold, 2010).

**Generalized Least Squares.** We perform generalized least squares in a two-stage manner. We first approximate the covariance structure with a Gaussian process regression model. Then we use this approximation to fit a generalized least squares model with restricted spatial regression (Hodges and Reich, 2010).

More precisely, first we optimize the maximum likelihood of a Gaussian process regression model with a linear mean function depending on the covariates including an intercept and Matérn 3/2 kernel defined on the spatial source locations. We use the GPFlow (Matthews et al., 2017) implementation of the likelihood, and L-BFGS (Liu and Nocedal, 1989) for

numerical optimization of the likelihood. The optimization is initialized using the GPFlow default parameters for the mean and covariance functions.

Once the maximum likelihood parameters have been found, we use the found prior covariance function for defining the covariance between datapoints to be used in the generalized least squares routine. We use restricted spatial regression (Hodges and Reich, 2010), and so the covariance matrix is defined as  $PKP + \lambda^2 I_N$ , where  $\lambda^2$  is the noise variance selected by maximum likelihood in the GP model, K is the  $N \times N$  matrix formed by evaluating the Matérn 3/2 kernel with the maximum likelihood kernel parameters on the source locations, and P is the orthogonal projection onto the orthogonal complement of the covariates (including intercept), i.e.  $P = I_N - X(X^TX)^{-1}X^T$ . This covariance matrix  $PKP + \lambda^2 I_N$  is passed to the GLS method in statsmodel, and confidence intervals as well as point estimates are computed using the default settings.

#### Gaussian Process Bayesian Credible Intervals.

We first optimize the maximum likelihood of a Gaussian process regression model with Matérn 3/2 kernel defined on the spatial source locations summed with a linear kernel defined on the covariates. This has the same likelihood as having a linear mean function in the covariates with a Gaussian prior over the weights, see Rasmussen and Williams (2006, Page 28). We use the GPFlow (Matthews et al., 2017) implementation of the likelihood, and L-BFGS (Liu and Nocedal, 1989) for numerical optimization of the likelihood. The optimization is initialized using the GPFlow default parameters for the mean and covariance functions. Once we have calculated the maximum likelihood parameters, we compute the posterior credible interval for  $\theta$ . The posterior over  $\theta$  is Gaussian and has the closed form,

$$\theta_{\text{post}} \sim \mathcal{N}(\Sigma_{\text{post}}^{-1} X^{\mathrm{T}} \Sigma^{-1} Y, \Sigma_{\text{post}})$$
 (30)

where

$$\Sigma_{\text{post}} = \left( X^{\text{T}} \Sigma^{-1} X + \frac{1}{\lambda^2} I_P \right), \tag{31}$$

where  $\lambda^2$  is the prior variance of  $\theta$ ,  $\Sigma = K + \delta^2 I_N$ , where  $\delta^2$  is the variance of the noise, and K is the  $N \times N$  kernel matrix formed by evaluating the Matérn 3/2 kernel on the source spatial locations. The posterior for  $e_p^{\rm T} \theta_{\rm post}$  is therefore also Gaussian,

$$e_p^{\mathrm{T}}\theta_{\mathrm{post}} \sim \mathcal{N}(e_p^{\mathrm{T}}\Sigma_{\mathrm{post}}^{-1}X^{\mathrm{T}}\Sigma^{-1}Y, e_p^{\mathrm{T}}\Sigma_{\mathrm{post}}e_p).$$
(32)

Credible intervals are then computed using z-scores together with this mean and variance.

### **C. Simulations Additional Details**

In this section, we present figures to visualize the data generating process for simulation experiments. In all experiments, an intercept is included in the regression as well as the covariates described.

#### C.1. Data Generation for the Single Covariate Experiment

We show example datasets used in the simulation experiment with a single covariate and the simulation experiment in which we investigated the impact of Lipschitz constant on confidence interval width and coverage in Figure 5. The left most two panels show the distribution of source (blue) and target (orange) locations for two values of the shift parameter. Large positive values of the shift parameter lead to target distributions clustered to the top right, values close to zero lead to the target locations being approximately uniformly distributed and large negative values lead to target locations clustered to the bottom left. The third panel from the left in Figure 5 shows the covariate plotted as a function of spatial location,

$$\chi(S^{(1)}, S^{(2)}) = S^{(1)} + S^{(2)}.$$
(33)

The right most panel shows the conditional expectation of the response plotted as a function of spatial location,

$$\mathbb{E}[Y^{\star}|S^{\star} = (S^{(1)}, S^{(2)})] = S^{(1)} + S^{(2)} + \frac{1}{2}((S^{(1)})^2 + (S^{(2)})^2).$$
(34)



*Figure 5.* Spatial sites for the source (blue) and target (orange) data are shown in the left most plots for different values of shift used in generating the data. More extreme values of the shift parameter lead to larger biases in parameter estimation from the training data without adjustment. The third plot from the left shows the covariate surface, while the fourth shows the expected response at each spatial location.

The gradient of this conditional expectation is,

$$\begin{bmatrix} 1 + S^{(1)} \\ 1 + S^{(2)} \end{bmatrix}$$
(35)

which has norm

$$\sqrt{(1+S^{(1)})^2 + (1+S^{(2)})^2}.$$
(36)

This obtains a maximum of  $2\sqrt{2}$  on  $[-1, 1]^2$ , and so this is the Lipschitz constant of  $f(S) = \mathbb{E}[Y|S]$ .

### C.2. Data Generation for the Three Covariate Experiment

In Figure 6 we show data generated for the three covariate shift experiment. Source and target locations are generated as in the one covariate simulation, but with N = 10,000 (M = 100 is still used). These are not shown in Figure 6.

The covariates are, from left to right in Figure 6

$$X^{(1)} = \sin(S^{(1)}) + \cos(S^{(2)}) \tag{37}$$

$$X^{(2)} = \cos(S^{(1)}) - \sin(S^{(2)}) \tag{38}$$

$$X^{(3)} = S^{(1)} + S^{(2)}.$$
(39)

The conditional expectation of the response (right most panel in Figure 6) is

$$\mathbb{E}[Y|S = (S^{(1)}, S^{(2)}] = X^{(1)}X^{(2)} + \frac{1}{2}((S^{(1)})^2 + (S^{(2)})^2)$$
(40)

$$= (\sin(S^{(1)}) + \cos(S^{(2)}))(\cos(S^{(1)}) - \sin(S^{(2)})) + \frac{1}{2}((S^{(1)})^2 + (S^{(2)})^2).$$
(41)

The gradient of this conditional expectation is

$$\begin{bmatrix} \cos(2S^{(1)}) - \sin(S^{(1)} + S^{(2)}) + S^{(1)} \\ -\cos(2S^{(1)}) + \sin(S^{(1)} - S^{(2)}) + S^{(2)} \end{bmatrix}.$$
(42)

We see that both arguments of this gradient are less than 3 in absolute value, and therefore the norm of this Lipschitz constant is less than or equal to  $\sqrt{3^2 + 3^2} = 3\sqrt{2}$ .

# **D.** Tree Cover Experiment Additional Details

In this section we provide additional details for the real data tree cover experiment, as well as figures to visualize the data and additional experimental results.

**Bias-corrected Confidence Intervals for Spatial Linear Models** 



*Figure 6.* The first 3 plots from the left show the covariate surfaces, while the fourth shows the expected response at each spatial location for the second simulated experiment. The source and target locations (not shown) are the same as in Figure 5, though with N = 10,000.

# D.1. Tree Cover Data

Our analysis draws on data from Lu et al. (2024), who manually labeled 983 high-resolution Google Maps satellite images for tree cover percentage. These images were selected from random locations within the 2021 USFS Tree Canopy Cover (TCC) product (USFS, 2023). We follow Lu et al. (2024) and use three covariates:

- 1. *Global Aridity Index (1970–2000)*: Averaged at a 30 arc-seconds resolution (Trabucco, 2019). This index is calculated as the ratio of precipitation to evapotranspiration, with lower values indicating more arid conditions.
- 2. Elevation: Provided by NASA's 30-meter resolution dataset (NASA Jet Propulsion Lab, 2020).
- 3. *Slope*: NASA's 30 m Digital Elevation Model. Also provided by NASA's 30-meter resolution dataset (NASA Jet Propulsion Lab, 2020).

Figure 7 provides a visual overview of both the tree cover and covariates. As a preprocessing step, we convert the (latitude, longitude) coordinates of each data point into radians. This allows us to use the Haversine formula to compute distances in kilometers for the Wasserstein-1 cost and the nearest neighbor weighting procedure.

Elevation and slope are important factors influencing tree cover worldwide. Generally, areas at lower elevations tend to have more tree cover, often due to higher temperatures (Mayor et al., 2017), and sloped terrains also support greater tree coverage (Sandel and Svenning, 2013). As done in Lu et al. (2024), we focus on these three covariates and did not include additional factors that might affect tree cover. This decision was made because our primary objective is to demonstrate uncertainty quantification rather than to provide a comprehensive explanation of tree cover dynamics.

Figure 7 provides a visual overview of the distribution for both the tree cover and covariates.

### D.2. Source and Target Data Split and Spatial Preprocessing

We define our target region to be the Western portion of the Continental United States. In particular, we consider locations that have latitudes between  $25^{\circ}$  and  $50^{\circ}$  and longitudes between  $-125^{\circ}$  and  $-110^{\circ}$ . Within spatial locations in this defined region, we pick 50% of all spatial locations — totaling 133 sites — as *target* data.

To select the *source* data, we perform the following steps:

- 1. Consider all the remaining spatial locations, i.e. exclude the 133 target points from the pool of 983 spatial points. This leaves us with 850 points.
- 2. Since we are interested in evaluating whether our method and the baselines achieve nominal coverage, uniformly randomly sample 20% of the remaining locations across 250 different random seeds. By doing this, for each random seed we have 170 source locations.

Figure 8 visually represents the spatial distribution of the source and target locations for one representative random seed.

As a preprocessing step, we also convert the geographical coordinates (latitude and longitude) of each data point from degrees to radians. This conversion is essential because it allows us to apply the Haversine formula, which calculates the



*Figure 7.* Tree cover response and covariates. The dots represent the 983 locations considered. Top left: distribution of tree cover percentage. Top right: Average Aridity Index, measured as the ratio of precipitation to evapotranspiration. Bottom left: Elevation, measured in meters. Bottom right: Slope, measured in degrees.

great-circle distance between two points on the Earth's surface in kilometers, to compute distances in kilometers for the Wasserstein-1 cost and the nearest neighbor weighting procedure.

#### D.3. Estimating the Ground Truth Parameters to Evaluate Coverage

In order to evaluate coverage, we repeat the data subsampling process described above 250 times. Ideally, we would estimate the coverage as the proportion of these seeds in which the estimand  $\theta^*_{OLS,p}$  falls inside the confidence interval we construct for each method. However, we cannot evaluate  $\theta^*_{OLS,p}$  directly, even though we have access to the target responses. To account for sampling variability, we use our method and each baseline to construct a confidence interval for the difference,

$$\hat{\theta}_p^{\star} - \hat{\theta}_p \tag{43}$$

where  $\hat{\theta}_p^{\star} = e_p^{\mathrm{T}} (X^{\star \mathrm{T}} X^{\star})^{-1} X^{\star \mathrm{T}} Y^{\star}$  is the estimated parameter using the target data (which our method and baselines don't have access to) and  $\hat{\theta}_p$  is the estimated parameter we compute with each method using the source responses. If this confidence interval contains 0, we count the method as having covered the true parameter, while if it doesn't we count the method as not having covered the true parameter. We estimate the variance of  $\hat{\theta}_p^{\star}$  using the model-trusting standard errors  $\hat{\sigma}^2 = \frac{1}{N-P} \sum_{n=1}^{N} r_n^2$  where  $r_n$  are the residuals of the model fit on the training data. We expect this to inflate our estimate of the standard variance of the target OLS estimate if the response surface is nonlinear, as the residuals will be larger due to bias. By possibly overestimating and incorporating this sampling variability into confidence intervals, we expect the calculated coverages to overestimate the true coverages. The resulting coverages are shown in the top row of Figure 4.

In Figure 9 we also report confidence intervals by calculating the proportion of times that  $\hat{\theta}_p^*$  is contained in each confidence interval.  $\hat{\theta}_p$  is an unbiased estimate for  $\theta_{\text{OLS},p}^*$  whether or not the model is well-specified. However, we might expect that the coverages reported with this approach underestimate the actual coverage of each method's confidence intervals due to not accounting for sampling variability in  $\hat{\theta}_p^*$ .



Figure 8. Split of the tree cover dataset in a target distribution in the West of the United States. Target locations are shown in orange, while source locations are shown in blue.

### **D.4. Experiment with Varying Lipschitz Constant**

In this section, we provide an additional experiment with these real data where the focus is to assess how varying the underlying Lipschitz constant in assumption 4 changes coverage and interval width. For this experiment we consider 9 different values for the Lipschitz constant,  $L \in \{0.001, 0.005, 0.01, 0.02, 0.05, 0.1, 0.2, 0.5, 1\}$ . This range of values is a reasonable range of values that a practitioner with domain knowledge in the tree cover field might be willing to specify. Indeed, the smallest Lipschitz constant considered, 0.001, corresponds to assuming that to have a 1% increase in tree cover we need to move 1/0.001 = 1000 kilometers. This is quite an extreme value, but in some parts of arid regions such as New Mexico and Arizona it can be true. On the other hand, the largest Lipschitz constant considered, 1, corresponds to assuming that to have 1% increase in tree cover we need to move 1 kilometer. This is also extreme, but in regions in the US where elevation changes are very pronounced (e.g. in the Colorado Rockies), tree cover can vary sharply over very short distances.

In Figure 10, we present the results of this experiment. We find that varying the Lipschitz constant within this range does not significantly impact coverage. Specifically, for *slope* and *elevation*, coverage remains consistent across all constants except L = 0.001. For *aridity index*, coverage is low for  $L \le 0.1$  but exceeds 95% for  $L \ge 0.2$ . Meanwhile, confidence intervals become noticeably wider for L > 0.5 while remaining relatively stable for smaller values.

### **D.5. Additional Experiment: Target South-East US**

In this experiment, we define our target region in the Southeastern portion of CONUS at locations with latitude in the range (25, 38) and longitude in the range (-100, -75). Out of all spatial points in this region, 50% — totaling 118 sites — are designated as target data. Next, we select the source data by taking a uniform random sample of 20% of the remaining spatial locations, repeated over 250 random seeds to assess coverage performance. Each seed yields 173 source locations. Figure 12 illustrates the spatial split between source and target data for a representative seed. As before, as a preprocessing step we convert the (latitude, longitude) coordinates of each data point into radians.

**Results.** We report the results for the confidence interval coverage and width in Figure 13. As before, our method consistently achieves or exceeds 95% nominal coverage for all parameters. All competing methods but KDEIW here achieves 95% nominal coverage (or close to 95% nominal coverage) for the aridity index and slope parameter when considering the coverage for the  $\hat{\theta}_p^* - \hat{\theta}_p$  (top row). All competing methods fall short of the nominal threshold for the elevation parameter. The wider intervals produced by our method (last row) reflect the trade-off between achieving reliable coverage and maintaining narrower intervals.

In the middle row we show the coverage for the point estimate  $\hat{\theta}_p^{\star}$ . Here we see how our method is the only one that achieves nominal coverage for all the parameters. In particular, all the other methods do not achieve nominal coverage for any of the parameters.



*Figure 9.* Coverages for the difference (upper), coverages for the point estimate (middle), and confidence interval widths (lower) for our method as well as 5 other methods for the West US data. Each column represents a parameter in the tree cover experiment. Only our method consistently achieves the nominal coverage.

**Varying Lipschitz constant.** Finally, we report also for this experiment results when varying the assumed Lipschitz constant. As explained in Appendix D.4, we consider 9 different values for the Lipschitz constant,  $L \in \{0.001, 0.005, 0.01, 0.02, 0.05, 0.1, 0.2, 0.5, 1\}$ . We report the results in Figure 15. As before, we find that varying the Lipschitz constant within this range does not significantly impact coverage. Specifically, here we see that for *aridity index* and *elevation*, coverage remains consistent across all constants above L = 0.01. For *slope*, 95% nominal coverage is achieved for  $L \ge 0.05$ . And — as before — confidence intervals become noticeably wider for L > 0.5 while remaining relatively stable for smaller values.



Figure 10. Coverage and average confidence interval widths over 250 seeds for 9 different values for the Lipschitz constant  $L \in \{0.001, 0.005, 0.01, 0.02, 0.05, 0.1, 0.2, 0.5, 1\}$ . The horizontal axis is shared across all plots.



*Figure 11.* Confidence intervals for different seeds for the West US. Each row shows confidence intervals for the various methods over the three parameters for a given seed. The dashed vertical lines represent the true parameters (black is the point estimate, orange is a 95% confidence interval). The blue dots the point estimates for the different methods, and the blue lines are the confidence intervals.



*Figure 12.* Spatial sites for the source (blue) and target (orange) data. The target data are chosen from the south-eastern part of the CONUS, whereas the source data cover the whole region.



*Figure 13.* Coverages for the difference (upper), coverages for the point estimate (middle), and confidence interval widths (lower) for our method as well as 5 other methods for the Southeast US data. Each column represents a parameter in the tree cover experiment. Only our method consistently achieves the nominal coverage.



*Figure 14.* Confidence intervals for different seeds for the South East US. Each row shows confidence intervals for the various methods over the three parameters for a given seed. The dashed vertical lines represent the true parameters (black is the point estimate, orange is a 95% confidence interval). The blue dots the point estimates for the different methods, and the blue lines are the confidence intervals.



Figure 15. Coverage and average confidence interval widths over 250 seeds for 9 different values for the Lipschitz constant  $L \in \{0.001, 0.005, 0.01, 0.02, 0.05, 0.1, 0.2, 0.5, 1\}$ .

# **E. Proofs**

# E.1. Derivation of Target-Conditional Ordinary Least Squares

We derive the OLS (Ordinary Least Squares) estimand for the given optimization problem:

$$\theta_{\text{OLS}}^{\star} = \arg\min_{\theta \in \mathbb{R}^{P}} \mathbb{E}\left[\sum_{m=1}^{M} (Y_{m}^{\star} - \theta^{\top} X_{m}^{\star})^{2} \middle| S_{m}^{\star} \right],$$
(44)

where  $Y_m^{\star}$  is unobserved, and  $X_m^{\star}$  is observed and a fixed function of  $S_m^{\star}$ .

First, expand the squared term inside the expectation:

$$(Y_m^{\star} - \theta^{\top} X_m^{\star})^2 = (Y_m^{\star})^2 - 2Y_m^{\star} \theta^{\top} X_m^{\star} + (\theta^{\top} X_m^{\star})^2.$$
(45)

Substitute this back into the expectation:

$$\mathbb{E}\left[\sum_{m=1}^{M} (Y_m^{\star} - \theta^{\top} X_m^{\star})^2 \middle| S_m^{\star} \right] = \mathbb{E}\left[\sum_{m=1}^{M} (Y_m^{\star})^2 - 2Y_m^{\star} \theta^{\top} X_m^{\star} + (\theta^{\top} X_m^{\star})^2 \middle| S_m^{\star} \right]$$
(46)

Since the expectation is linear, we can separate the terms:

$$\mathbb{E}\left[\sum_{m=1}^{M} (Y_m^{\star})^2 \Big| S_m^{\star}\right] - 2\mathbb{E}\left[\sum_{m=1}^{M} Y_m^{\star} \theta^{\top} X_m^{\star} \Big| S_m^{\star}\right] + \mathbb{E}\left[\sum_{m=1}^{M} (\theta^{\top} X_m^{\star})^2 \Big| S_m^{\star}\right]$$
(47)

Now we can simplify each of the terms as follows:

- The first term,  $\sum_{m=1}^{M} \mathbb{E}\left[ (Y_m^{\star})^2 \middle| S_m^{\star} \right]$ , does not depend on  $\theta$ , so it can be treated as a constant with respect to the optimization problem
- The second term,  $-2\mathbb{E}\left[\sum_{m=1}^{M} Y_m^{\star} \theta^{\top} X_m^{\star} \middle| S_m^{\star}\right]$ , can be rewritten using the linearity of expectation, the fact that  $\theta$  is not random, and assumption 1:

$$-2\theta^{\top} \sum_{m=1}^{M} X_m^{\star} \mathbb{E} \left[ Y_m^{\star} \middle| S_m^{\star} \right]$$
(48)

• The third term,  $\mathbb{E}\left[\sum_{m=1}^{M} (\theta^{\top} X_m^{\star})^2 \middle| S_m^{\star}\right]$ , is non-random by assumption 1 and can be rewritten as:

$$\sum_{m=1}^{M} (\theta^{\top} X_m^{\star})^2 \tag{49}$$

The optimization problem then becomes

$$\theta_{\text{OLS}}^{\star} = \arg\min_{\theta \in \mathbb{R}^{P}} \left\{ \text{constant} - 2\theta^{\top} \sum_{m=1}^{M} X_{m}^{\star} \mathbb{E} \left[ Y_{m}^{\star} \middle| S_{m}^{\star} \right] + \sum_{m=1}^{M} (\theta^{\top} X_{m}^{\star})^{2} \right\}.$$
(50)

And since the constant term does not affect the optimization, we can drop it to get

$$\theta_{\text{OLS}}^{\star} = \arg\min_{\theta \in \mathbb{R}^{P}} \left\{ -2\theta^{\top} \sum_{m=1}^{M} X_{m}^{\star} \mathbb{E} \left[ Y_{m}^{\star} \middle| S_{m}^{\star} \right] + \sum_{m=1}^{M} (\theta^{\top} X_{m}^{\star})^{2} \right\}$$
(51)

To find the minimizer, we take the derivative of the objective function with respect to  $\theta$  and set it to zero:

$$\frac{\partial}{\partial \theta} \left\{ -2\theta^{\top} \sum_{m=1}^{M} X_m^{\star} \mathbb{E} \left[ Y_m^{\star} \middle| S_m^{\star} \right] + \sum_{m=1}^{M} (\theta^{\top} X_m^{\star})^2 \right\} = 0$$
(52)

$$-2\sum_{m=1}^{M} X_m^{\star} \mathbb{E}\left[Y_m^{\star} \middle| S_m^{\star}\right] + 2\sum_{m=1}^{M} X_m^{\star} (X_m^{\star})^{\top} \theta = 0$$
<sup>(53)</sup>

And by inverting this and using the fact that we assumed that  $X^{\star T}X^{\star}$  is invertible

$$\theta_{\text{OLS}}^{\star} = \left(\sum_{m=1}^{M} X_m^{\star} (X_m^{\star})^{\top} \theta\right)^{-1} \sum_{m=1}^{M} X_m^{\star} \mathbb{E}\left[Y_m^{\star} \middle| S_m^{\star}\right]$$
(54)

which in matrix form can be written as

$$\theta_{\text{OLS}}^{\star} = (X^{\star T} X^{\star})^{-1} X^{\star T} \mathbb{E}[Y^{\star} | S^{\star}].$$
(55)

as in Equation (7).

#### E.2. Proof of Theorem 7

In the main text, we stated our confidence interval for the 1NN choice of  $\Psi$ . We now state version for general non-negative matrices with columns summing to 1.

**Theorem 11.** Suppose  $(S_m^{\star}, X_m^{\star}, Y_m^{\star})_{m=1}^M$  and  $(S_n, X_n, Y_n)_{n=1}^N$  satisfy assumptions 1 to 4. Define w as in Algorithm 1. For any  $\Psi \in \mathbb{R}^{M \times N}$ , a matrix with non-negative entries with columns summing to 1, define  $v^{\Psi} = w\Psi$ . Take  $\hat{\theta}_p^{\Psi}$  as in Equation (10). Define  $c = \sigma ||v^{\Psi}||_2$ , with  $\sigma^2$  the variance of the additive noise from assumption 2. Define the (random) interval  $I^{\Psi}$  as in Algorithm 1 with known  $\sigma^2$ . Then with probability at least  $1 - \alpha$ ,  $\theta_{OLS,p}^{\star} \in I^{\Psi}$ . That is,  $I^{\Psi}$  has coverage (conditional on the test locations) at least  $1 - \alpha$ .

*Proof.* We being as in the main text. We show the difference between our estimand and estimator is normally distributed. To that end, we decompose the difference between our estimand and estimator into a bias term and a mean-zero noise term.

$$\theta_{\text{OLS},p}^{\star} - \hat{\theta}_{p}^{\Psi} = \sum_{m=1}^{M} w_{m} f(S_{m}^{\star}) - \sum_{n=1}^{N} v_{n}^{\Psi} Y_{n},$$
(56)

for  $w := e_p^T (X^{\star T} X^{\star})^{-1} X^{\star T} \in \mathbb{R}^M$  and  $v^{\Psi} := w \Psi \in \mathbb{R}^N$ . By assumption 2, the righthand side of Equation (13) can be written

$$\underbrace{\sum_{m=1}^{M} w_m f(S_m^{\star}) - \sum_{n=1}^{N} v_n^{\Psi} f(S_n)}_{\text{bias}} - \underbrace{\sum_{n=1}^{N} v_n^{\Psi} \epsilon_n}_{\text{randomness}}.$$
(57)

Since the spatial locations are fixed, the bias term is not random and can be written as  $b \in \mathbb{R}$ . We can calculate the variance directly,

$$\mathbb{V}[\theta_{\text{OLS},p}^{\star} - \hat{\theta}_{p}^{\Psi}] = \mathbb{V}[\sum_{n=1}^{N} v_{n}^{\Psi} \epsilon_{n}] = \sum_{n=1}^{N} (v_{n}^{\Psi})^{2} \mathbb{V}[\epsilon_{n}] = \sigma^{2} \|v^{\Psi}\|_{2}^{2}.$$
(58)

We used the  $\epsilon_n$  are independent and identically distributed with variance  $\sigma^2$  (assumption 2).

It follows that

$$\theta^{\star}_{\text{OLS},p} - \hat{\theta}^{\Psi}_{p} \sim \mathcal{N}(b, \sigma^{2} \| v^{\Psi} \|_{2}^{2})$$
(59)

To bound the bias b, we use assumption 4 to write

$$|b| \le \sup_{g \in \mathcal{F}_L} \left| \sum_{m=1}^M w_m g(S_m^{\star}) - \sum_{n=1}^N v_n^{\Psi} g(S_n) \right|.$$
(60)

We can therefore apply Lemma 6 to complete the proof.

### E.3. Computing an Upper Bound on the Bias of Our Estimand with Wasserstein-1 Distance

First of all observe that if  $\sum_{m=1}^{M} w_m = 1$ ,  $\sum_{n=1}^{N} v_n^{\Psi} = 1$ ,  $v_n^{\Psi} \ge 0$  for  $1 \le n \le N$ , and  $w_m \ge 0$  for  $1 \le m \le M$  then the supremum in Equation (16) would be equal to a Wasserstein-1 distance by Kantorovich-Rubinstein duality (Villani, 2009, Theorem 5.10, Case 5.16).

Next, consider what happens if  $\sum_{m=1}^{M} w_m - \sum_{n=1}^{N} v_n^{\Psi} \neq 0$ . We show that the right-hand side of Equation (16) is infinite. Assume by contradiction that there exists a C > 0 that upper bounds this supremum. Because  $\sum_{m=1}^{M} w_m - \sum_{n=1}^{N} v_n^{\Psi} \neq 0$  and all constant functions are *L*-Lipschitz, for any  $\gamma > 0$ 

$$\sup_{g \in \mathcal{F}_{L}} \left| \sum_{m=1}^{M} w_{m} g(S_{m}^{\star}) - \sum_{n=1}^{N} v_{n}^{\Psi} g(S_{n}) \right| \geq \left| \sum_{m=1}^{M} w_{m} \frac{C + \gamma}{\sum_{m=1}^{M} w_{m} - \sum_{n=1}^{N} v_{n}^{\Psi}} - \sum_{n=1}^{N} v_{n}^{\Psi} \frac{C + \gamma}{\sum_{m=1}^{M} w_{m} - \sum_{n=1}^{N} v_{n}^{\Psi}} \right| \quad (61)$$
$$= C + \gamma.$$

This contradicts the assumption that C is an upper bound on the supremum. Because C was arbitrary, the right hand side of Equation (16) is infinite, as desired.

Our assumption that  $\Psi$  is a non-negative matrix whose columns sum to 1 avoids this situation. We formalize our upper bound on the bias in the following proposition.

**Proposition 12.** Suppose that  $\sum_{n=1}^{N} \Psi_{m,n} = 1$  for all m. Let  $w \in \mathbb{R}^{M}$  and  $v^{\Psi} = (w\Psi)^{\mathrm{T}} \in \mathbb{R}^{N}$ .

$$\sup_{g \in \mathcal{F}_L} \left| \sum_{m=1}^M w_m g(S_m^\star) - \sum_{n=1}^N v_n^\Psi g(S_n) \right| = ALW_1 \left( \sum_{m \in I} \frac{w_m}{A} \delta_{S_m^\star} + \sum_{n \in I'} \frac{-v_n^\Psi}{A} \delta_{S_n}, \sum_{m \in J} \frac{-w_m}{A} \delta_{S_m^\star} + \sum_{n \in J'} \frac{v_n^\Psi}{A} \delta_{S_n} \right),$$
(63)

where  $I = \{1 \le i \le M : w_i \ge 0\}$ ,  $I' = \{1 \le i \le N : v_i^{\Psi} < 0\}$ ,  $J = \{1 \le j \le M : w_j < 0\}$  and  $J' = \{1 \le j \le N : v_j^{\Psi} \ge 0\}$  and  $A = \frac{1}{2} \left( \sum_{m=1}^M |w_m| + \sum_{n=1}^N |v_n^{\Psi}| \right)$ .

*Proof.* First, observe that  $\sum_{n=1}^{N} v_n^{\Psi} = \sum_{n=1}^{N} (w\Psi)_n = \sum_{n=1}^{N} \sum_{m=1}^{M} w_m \Psi_{m,n} = \sum_{m=1}^{M} w_m \sum_{n=1}^{N} \Psi_{m,n} = \sum_{m=1}^{M} w_m$ . Next, we normalize the weights to sum in absolute value to 2, and rescale the function class to consist of 1-Lipschitz function. Define  $A = \frac{1}{2} \left( \sum_{m=1}^{M} |w_m| + \sum_{n=1}^{N} |v_n^{\Psi}| \right)$ . Then

$$\sup_{g \in \mathcal{F}_L} \left| \sum_{m=1}^M w_m g(S_m^{\star}) - \sum_{n=1}^N v_n^{\Psi} g(S_n) \right| = \sup_{g \in \mathcal{F}_1} \left| \sum_{m=1}^M w_m Lg(S_m^{\star}) - \sum_{n=1}^N v_n^{\Psi} Lg(S_n) \right|$$
(64)

$$= AL \sup_{g \in \mathcal{F}_1} \left| \sum_{m=1}^M \frac{w_m}{A} g(S_m^\star) - \sum_{n=1}^N \frac{v_n^\Psi}{A} g(S_n) \right|$$
(65)

where we have used that a function is L-Lipschitz if and only if it can be written by scaling a 1-Lipschitz function by L.

Define  $I = \{1 \le i \le M : w_i \ge 0\}, I' = \{1 \le i \le N : v_i^{\Psi} < 0\}, J = \{1 \le j \le M : w_j < 0\}$  and  $J' = \{1 \le j \le N : v_j^{\Psi} \ge 0\}$ . Then

$$AL \sup_{g \in \mathcal{F}_1} \left| \sum_{m=1}^M \frac{w_m}{A} g(S_m^\star) - \sum_{n=1}^N \frac{v_n^\Psi}{A} g(S_n) \right|$$
(66)

$$= AL \sup_{g \in \mathcal{F}_1} \left| \sum_{m \in I} \frac{w_m}{A} g(S_m^\star) + \sum_{n \in I'} \frac{-v_n^\Psi}{A} g(S_n) - \left( \sum_{m \in J} \frac{-w_m}{A} g(S_m^\star) + \sum_{n \in J'} \frac{v_n^\Psi}{A} g(S_n) \right) \right|$$
(67)

Because  $\sum_{n=1}^{N} v_n^{\Psi} = \sum_{m=1}^{M} w_m$  and I and J partition the index sets,

$$\sum_{m \in I} w_m + \sum_{n \in I'} -v_n^{\Psi} = \sum_{m \in J} -w_m + \sum_{n \in J'} v_n^{\Psi}.$$
(68)

And because the set I, I', J, J' sort the indices into positive and negative parts

$$\sum_{m \in I} w_m + \sum_{m \in J} -w_m + \sum_{n \in I'} -v_n^{\Psi} + \sum_{n \in J'} v_n^{\Psi} = \sum_{m=1}^M |w_m| + \sum_{n=1}^N |v_n^{\Psi}| = 2A.$$
 (69)

Therefore,

$$\sum_{m \in I} \frac{w_m}{A} \delta_{S_m^\star} + \sum_{n \in I'} \frac{-v_n^\Psi}{A} \delta_{S_n} \text{ and } \sum_{m \in J} \frac{-w_m}{A} \delta_{S_m^\star} + \sum_{n \in J'} \frac{v_n^\Psi}{A} \delta_{S_n}$$
(70)

are probability measures. We can apply Kantorovich-Rubinstein duality to write,

$$B \le ALW_1 \left( \sum_{m \in I} \frac{w_m}{A} \delta_{S_m^\star} + \sum_{n \in I'} \frac{-v_n^\Psi}{A} \delta_{S_n}, \sum_{m \in J} \frac{-w_m}{A} \delta_{S_m^\star} + \sum_{n \in J'} \frac{v_n^\Psi}{A} \delta_{S_n} \right).$$
(71)

where  $W_1$  denotes the 1-Wasserstein distance.

We compute the Wasserstein-1 distance by linear programming, see discussion in Appendix B.2. Scalable upper bounds could also be computed by exhibiting a coupling between the measures (for example by solving an entropy regularized optimal transport problem). See (Peyré and Cuturi, 2019, Chapters 3 and 4) for details on computation of Wasserstein distances.

### E.4. Proof of Lemma 6

*Proof of Lemma 6.* We aim to prove that the interval  $[-B - \Delta, B + \Delta]$  is the narrowest  $1 - \alpha$  confidence interval that is valid for all  $b \in [-B, B]$  where  $\Delta$  is the solution of Equation (17).

**Ensuring Coverage Probability.** Suppose that  $X_b \sim N(b, \tilde{c}^2)$  for  $b \in [-B - \Delta, B + \Delta]$ . Then,

$$\Pr(X_b \in [-B - \Delta, B + \Delta]) = \Phi\left(\frac{B + \Delta - b}{\tilde{c}}\right) - \Phi\left(\frac{-B - \Delta - b}{\tilde{c}}\right),$$

To construct a valid confidence interval for any  $b \in [-B, B]$ , we require that

$$\Pr(X_b \in [-B - \Delta, B + \Delta]) \ge 1 - \alpha, \quad \forall b \in [-B, B].$$

This ensures  $1 - \alpha$  coverage over all possible values of b in [-B, B].

**Reduce the problem to Worst-Case Coverage.** To find the narrowest interval, we identify the worst-case value of *b* that minimizes the coverage probability. Let

$$C(b;\Delta) = \Phi\left(\frac{B+\Delta-b}{\tilde{c}}\right) - \Phi\left(\frac{-B-\Delta-b}{\tilde{c}}\right),$$

denote the coverage probability of the interval  $[-B - \Delta, B + \Delta]$  for  $X_b \sim \mathcal{N}(b, \tilde{c}^2)$ . In order to ensure the interval is valid for all b coverage, we want to bound below.

$$\inf_{b \in [-B,B]} C(b;\Delta)$$

The interval  $[-B - \Delta, B + \Delta]$  is symmetric about 0, and the Probability Density Function for a Gaussian of mean b is symmetric about b. Thus, the coverage probabilities at b = -B and b = B are equal. Consequently, it suffices to consider  $b \in [0, B]$ .

Moreover, observe that  $C(b; \Delta)$  is a strictly decreasing function of b on [0, B] since (i)  $\Phi\left(\frac{B+\Delta-b}{\bar{c}}\right)$  decreases as b increases (because  $B + \Delta - b$  decreases and  $\Phi(z)$  is monotonic) and (ii)  $\Phi\left(\frac{-B-\Delta-b}{\bar{c}}\right)$  also decreases as b increases (because  $-B - \Delta - b$  becomes more negative). Thus,  $C(b; \Delta)$  is a strictly decreasing function of b on [0, B]. The minimum value of  $C(b; \Delta)$  occurs at b = B.

**Ensuring coverage in the worst case.** At the worst-case value b = B, the coverage probability is:

$$C(B;\Delta) = \Phi\left(\frac{\Delta}{\tilde{c}}\right) - \Phi\left(\frac{-2B - \Delta}{\tilde{c}}\right)$$

To ensure that the interval  $[-B - \Delta, B + \Delta]$  achieves at least  $1 - \alpha$  coverage for all  $b \in [-B, B]$ , we solve:

$$\Phi\left(\frac{\Delta}{\tilde{c}}\right) - \Phi\left(\frac{-2B - \Delta}{\tilde{c}}\right) = 1 - \alpha.$$
(72)

This guarantees that the interval is valid for all b and achieves the desired coverage level.

**Narrowest interval.** The narrowest interval corresponds to the smallest  $\Delta$  that satisfies the Equation (72). By construction, any smaller  $\Delta$  would fail to achieve the required coverage for  $b = \pm B$ , violating the validity condition.

#### E.5. Proof of Proposition 8

In this section, we prove Proposition 8. For simplicity of exposition, we prove the result for  $S = [0, 1]^D$ . The result generalizes to any spatial domain which is a compact metric space.

*Proof of Proposition 8.* Using assumption 2 and expanding the quadratic form  $(\epsilon_n + (f(S_n) - g(S_n)))$ , we have

$$\hat{\sigma}_N^2 - \sigma^2 = Z_N + \zeta_N,\tag{73}$$

where  $\zeta_N = \inf_{g \in \mathcal{F}_L} \frac{1}{N} \sum_{n=1}^N (f(S_n) - g(S_n))^2 + \frac{1}{N} \sum_{n=1}^N \epsilon_n (f(S_n) - g(S_n))$  and  $Z_N = \frac{1}{N} \sum_{n=1}^N \epsilon_n^2 - \sigma^2$ . Since  $Z_N$  is an average of independent and identically distributed variable, and since  $\mathbb{E}[Z_N] = 0$ , the law of large numbers (LLN) implies  $Z_N \to 0$  in probability. Because  $Z_N \to 0$ , if  $\zeta_N \to 0$  in probability we can conclude by Slutsky's Lemma that  $\hat{\sigma}_N^2 \to \sigma^2$  in probability. Therefore, the remainder of the proof involves showing  $\zeta_N \to 0$  in probability.

Define  $f_N$  to be the empirically centered version of f, that is  $f_N = f - \frac{1}{N} \sum_{n=1}^N f(S_n)$ . Then since the space of Lipschitz functions is invariant to shifts by constant functions

$$\zeta_N = \inf_{g \in \mathcal{F}_L} \frac{1}{N} \sum_{n=1}^N (f_N(S_n) - g(S_n))^2 + \frac{1}{N} \sum_{n=1}^N \epsilon_n (f_N(S_n) - g(S_n)).$$
(74)

Define the process,

$$\tau_N(g) = \frac{1}{N} \sum_{n=1}^N (f_N(S_n) - g(S_n))^2 + \frac{1}{N} \sum_{n=1}^N \epsilon_n (f_N(S_n) - g(S_n)),$$
(75)

so that  $\zeta_N$  is the infimum of  $\tau_N$ .  $\tau_N(f_N) = 0$ . Therefore, for  $\zeta_N \leq 0$  almost surely. It remains to show that for any  $\delta < 0$ ,  $\lim_{N \to \infty} \Pr(\zeta_N < \delta) \to 0$ .

The essential challenge to showing that for any  $\delta < 0$ ,  $\lim_{N\to\infty} \Pr(\zeta_N < \delta) \to 0$  is the infimum over the space of Lipschitz functions. Our proof has three steps. First, we show that it suffices to consider a subset of the space of Lipschitz functions with bounded infinity norm. Second, we show that this space is compact as a subset of  $L^{\infty}$ . Third, we show that because the infimum is then over a compact set, it can be well-approximated by a minimum over a finite set of functions. And then a union bound suffices to prove the claim.

#### Step 1: It's Enough to Consider a Bounded Subset of Lipschitz Functions.

Because  $f_N$  has empirical mean 0 and is continuous, by the intermediate value theorem it takes on the value 0 somewhere on  $[0,1]^D$ . Because  $f_N$  is L-Lipschitz and defined on a set of diameter  $\sqrt{D}$ , and 0 somewhere inside this set,  $||f_N||_{\infty} \leq L\sqrt{D}$ .

Define the set  $\overline{\mathcal{F}_L} = \mathcal{F}_L \cap B_{\infty}(2L\sqrt{D} + 2\sigma^2)$ , where  $B_{\infty}(r)$  denotes the space of functions uniformly bounded by constant r on  $[0, 1]^D$ . By subadditivity of measure, for any  $\delta < 0$ 

$$\Pr(\zeta_N < \delta) \le \Pr\left(\inf_{g \in \overline{\mathcal{F}_L}} \tau_N(g) < \delta\right) + \Pr\left(\inf_{g \in \mathcal{F}_L \setminus \overline{\mathcal{F}_L}} \tau_N(g) < \delta\right)$$
(76)

$$\leq \Pr\left(\inf_{g\in\overline{\mathcal{F}_L}}\tau_N(g)<\delta\right) + \Pr\left(\inf_{g\in\mathcal{F}_L\setminus\overline{\mathcal{F}_L}}\tau_N(g)<0\right).$$
(77)

We first consider the second term in this sum and show it tends to 0. We apply a crude Cauchy-Schwarz bound to the second term in Equation (75) so that for any g,

$$\tau_N(g) \ge \sqrt{\frac{1}{N} \sum_{n=1}^N (f(S_n) - g(S_n))^2} \left( \sqrt{\frac{1}{N} \sum_{n=1}^N (f(S_n) - g(S_n))^2} - \sqrt{\frac{1}{N} \sum_{n=1}^N \epsilon_n^2} \right).$$
(78)

Therefore,  $\tau(g) < 0$  implies

$$\frac{1}{N}\sum_{n=1}^{N}\epsilon_n^2 \le \frac{1}{N}\sum_{n=1}^{N}(f_N(S_n) - g(S_n))^2.$$
(79)

For any  $g \in \mathcal{F}_L \setminus \overline{\mathcal{F}_L}$  because g takes on a value at least  $2L\sqrt{D} + 2\sigma^2$  and is L-Lipschitz, g is larger than  $L\sqrt{D} + 2\sigma^2$  over the entire unit cube. And because  $||f_N|| \leq L\sqrt{D}$ ,  $f_N(S_n) - g(S_n) \geq 2\sigma^2$  for all n. Therefore for all  $g \in \mathcal{F}_L \setminus \overline{\mathcal{F}_L}$ 

$$\frac{1}{N}\sum_{n=1}^{N}(f_N(S_n) - g(S_n))^2 \ge 2\sigma^2.$$
(80)

We conclude that

$$\lim_{N \to \infty} \Pr\left(\inf_{g \in \mathcal{F}_L \setminus \overline{\mathcal{F}_L}} \tau_N(g) < 0\right) \le \lim_{N \to \infty} \Pr\left(\frac{1}{N} \sum_{n=1}^N \epsilon_n^2 \ge 2\sigma^2\right) = 0.$$
(81)

where the final inequality is the law of large numbers.

 $\overline{\mathcal{F}_L}$  is a Compact Subset of the Space of Bounded Functions with Sup Norm. All that remains to show is that for any  $\delta < 0$ ,

$$\Pr\left(\inf_{g\in\overline{\mathcal{F}_L}}\tau_N(g)<\delta\right)\to 0.$$
(82)

The idea (following standard arguments made with empirical processes) is that we can take a cover of  $\overline{\mathcal{F}_L}$  of finite size, such that each element of  $\tau_N(g)$  is almost constant over elements of this cover. This essentially lets us approximate the infimum with a minimum over a finite set, up to small error. And once the problem is reduced to a minimum we can apply a union bound and the law of large number. We will formalize this in the next paragraph, but we first show that  $\overline{\mathcal{F}_L}$  is compact.

Because every Lipschitz continuous function is equicontinuous, functions in  $\overline{\mathcal{F}_L}$  are pointwise bounded by  $2L\sqrt{D} + 2\sigma^2$ , and  $[0,1]^D$  is compact, we may apply Arzela-Ascoli (Rudin, 1976, Theorem 7.25) to conclude that  $\overline{\mathcal{F}_L}$  with the sup norm is sequentially compact. It is therefore compact, as a sequentially compact metric space is compact.

#### Step 3: Reduction to a Minimum over a Finite Set and a Union Bound.

For any  $\delta' < 0$  and any  $g \in \overline{\mathcal{F}_L}$ ,

$$\lim_{N \to \infty} \Pr(\tau_N(g) < \delta') \le \lim_{N \to \infty} \Pr\left(\frac{1}{N} \sum_{n=1}^N \epsilon_n g(S_n) < \delta'\right) = 0$$
(83)

where the final equality is the law of large numbers.

Therefore, for any finite set  $C \subset \overline{\mathcal{F}_L}$ ,

$$\lim_{N \to \infty} \Pr(\min_{g \in C} \tau_N(g) < \delta') \le \lim_{N \to \infty} \sum_{g \in C} \Pr\left(\frac{1}{N} \sum_{n=1}^N \epsilon_n g(S_n) < \delta'\right)$$
(84)

$$= \sum_{g \in C} \lim_{N \to \infty} \Pr\left(\frac{1}{N} \sum_{n=1}^{N} \epsilon_n g(S_n) < \delta'\right) = 0.$$
(85)

We used countable subadditivity in the inequality.

Now for any  $\gamma > 0$ , there exists a finite set of functions  $C_{\gamma} \subset \overline{\mathcal{F}_L}$  such that for any  $g \in \overline{\mathcal{F}_L}$ , there exists a  $g' \in C_{\gamma}$  with  $\|g - g'\|_{\infty} \leq \gamma$ . And since  $\tau_N(g)$  is pathwise uniformly (in N) continuous on  $\overline{\mathcal{F}_L}$  equipped with sup norm,

$$\inf_{g \in \mathcal{F}_L} \tau_N(g) \ge \min_{g \in C_\gamma} \tau_N(g) - \rho(\gamma)$$
(86)

where is a nonnegative function such that  $\lim_{\gamma \to 0} \rho(\gamma) = 0$ . Therefore, for any  $\delta < 0$ , we can find a  $\gamma$  such that  $\rho(\gamma) \le -\frac{\delta}{2}$ . For this  $\gamma$ , applying Equation (84) with  $\delta' = \frac{\delta}{2}$  allows us to conclude that

$$\lim_{N \to \infty} \Pr(\inf_{g \in \overline{\mathcal{F}}_L} \tau_N(g) \le \delta) = 0.$$
(87)

This is a uniform law of large number for the class of Lipschitz and bounded functions. More quantitative results are likely possible using empirical process theory, see Wainwright (2019, Chapter 5).

**Proof of Asymptotic Coverage (Corollary 9).** We now prove Corollary 9. We begin by recalling the definition of an asymptotically valid confidence interval.

**Definition 13.** We say a sequence of (random) intervals  $(I_N)_{N=1}^{\infty}$  has asymptotically valid coverage of  $\theta$  at level  $(1 - \alpha)$  if

$$\lim_{N \to \infty} \Pr(\theta \in I_N) = 1 - \alpha \tag{88}$$

Corollary 9 follows from Theorem 7 and Proposition 8 by the following lemma, which is a special case of Slutsky's lemma. **Lemma 14.** Let  $\sigma^2 > 0$ . Suppose that  $\theta_{OLS,p}^{\star} - \hat{\theta}_p^{\Psi} - b_N \sim \mathcal{N}(0, \sigma^2)$  where  $(b_N)_{N=1}^N$  is a fixed sequence. Suppose  $\hat{\sigma}_N^2$  converges in probability to  $\sigma^2$ . Then,

$$\frac{1}{\hat{\sigma}_N^2} \left( \theta^*_{OLS,p} - \hat{\theta}_p^\Psi - b_N \right) \to \mathcal{N}(0,1)$$
(89)

where convergence is in probability.

*Proof.* The result is a special case of Slutsky's lemma, using  $\hat{\sigma}_N^2 \rightarrow \sigma^2 > 0$ .

Proof of Corollary 9. Define  $b_N = \sum_{m=1}^M w_m f(S_m^{\star}) - \sum_{n=1}^N v_n^{\Psi} f(S_n^{\star})$ . By Equation (15)

$$(\theta_{\text{OLS}}^{\star} - \hat{\theta}_p^{\Psi} - b_N) \sim \mathcal{N}(0, \sigma^2).$$
(90)

If  $\sigma^2 = 0$ , then  $\hat{\sigma}_N^2 = 0$  for all N, because the conditional expectation is an L-Lipschitz function leading to 0 squared error in the minimization algorithm used to calculate  $\hat{\sigma}_N^2$ . Therefore, the resulting confidence interval has coverage  $(1 - \alpha)$  for all N by Theorem 7.

For  $\sigma^2 > 0$  we apply Lemma 14 to conclude that

$$\frac{1}{\hat{\sigma}_N^2} \left( \theta^{\star}_{\text{OLS},p} - \hat{\theta}_p^{\Psi} - b_N \right) \to \mathcal{N}(0,1) \tag{91}$$

where convergence is in probability.

Convergence in probability implies convergence in distribution, and therefore convergence of quantiles at continuity points. The Gaussian CDF is continuous. Therefore, the quantile computation in Lemma 6 using  $\hat{\sigma}_N^2$  in place of  $\sigma^2$  produces an asymptotically valid confidence interval in the limit as  $N \to \infty$ .