

THE BASIC LOCUS OF RAMIFIED UNITARY RAPOPORT-ZINK SPACE AT MAXIMAL VERTEX LEVEL

QIAO HE, YU LUO, AND YOUSHENG SHI

ABSTRACT. We construct the Bruhat-Tits stratification of the ramified unitary Rapoport-Zink space, with the level being the stabilizer of a vertex lattice. We develop the local model theory for Bruhat-Tits strata, proving their normality and Cohen-Macaulayness, and provide precise dimension formulas. Additionally, we establish an explicit isomorphism between Bruhat-Tits strata and Deligne-Lusztig varieties, revealing new phenomena beyond the previously studied Coxeter-type cases.

CONTENTS

1. Introduction	1
2. Statements of the main results	5
3. Rapoport-Zink spaces	11
4. Local models of basic locus strata	16
5. Bruhat-Tits stratification	29
6. Deligne-Lusztig varieties	34
7. Relation between Bruhat-Tits strata and Deligne-Lusztig varieties	45
References	52

1. INTRODUCTION

In this paper, we study the reduced locus of basic unitary Rapoport-Zink spaces. This study contributes to the broader theory of reduction of integral models of Shimura varieties. For historical context and background on this topic, we refer to [Vol10].

Basic unitary Rapoport-Zink spaces have been proven important in arithmetic intersection theory, particularly in the Kudla-Rapoport conjecture [LZ22a, LZ22b, HLSY23] and the arithmetic fundamental lemma and the arithmetic transfer conjecture [Zha21, Zha24].

The reduced locus of basic unitary Rapoport-Zink spaces was first studied by Vollaard and Wedhorn [Vol10, VW11] for the unramified unitary group with signature $(1, n - 1)$ and hyperspecial level. Their work revealed a fundamental structure: a Bruhat-Tits stratification indexed by the Bruhat-Tits building of the unitary group. In this stratification, each stratum admits an explicit geometric description as disjoint union of Deligne-Lusztig varieties. This elegant structural characterization has since proven to be more universal, extending to several related settings.

When the prime is ramified, which is the case we are interested in, the reduced locus was studied by Rapoport-Terstiege-Wilson [RTW14] for self-dual levels and by Wu [Wu16] for exotic smooth levels. These level structure are of ‘‘Coxeter type’’ level structure, as defined in [GHN24]. We refer to loc. cit. for a group-theoretical study of Coxeter type ADLV. In this paper, we extend the study of the Bruhat-Tits stratification to all maximal vertex levels, where the level is constructed as the stabilizer of a vertex lattice. These cases generally fall outside the Coxeter type framework. Our treatment is moduli-theoretic, not group-theoretic, offering a more geometrically intuitive perspective on the stratification structure.

Let F/F_0 be a ramified quadratic extension of p -adic fields ($p \geq 3$) with uniformizers π and π_0 respectively, satisfying $\pi^2 = \pi_0$. Let V be a hermitian space over F of dimension n with Hasse invariant ε . The basic Rapoport-Zink space $\mathcal{N}_{n,\varepsilon}^{[h]}(1, n-1)$ is a certain formal scheme that relates to the unitary group $U(V)$ and some vertex lattice $L \subseteq V$ of type h (along with the minimal element $b \in B(U(V))$ and a conjugacy class of geometric cocharacter $\mu_{1,n-1}$), see §3.3. The main result of this paper is a description of the reduced locus $\mathcal{N}_{n,\varepsilon,\text{red}}^{[h]}$ in terms of the Bruhat-Tits building of the inner twist $U(\mathbb{V})$, where \mathbb{V} is a hermitian space over F of dimension n and Hasse invariant $-\varepsilon$.

We denote by $\mathcal{L}_{\mathcal{Z}}$ (resp. $\mathcal{L}_{\mathcal{Y}}$) the set of vertex lattices $\Lambda \subset \mathbb{V}$ of type $\geq h$ (resp. $\leq h$). For each $\Lambda_1 \in \mathcal{L}_{\mathcal{Z}}$ and $\Lambda_2 \in \mathcal{L}_{\mathcal{Y}}$, we define closed subschemes $\mathcal{Z}(\Lambda_1)$ and $\mathcal{Y}(\Lambda_2^\sharp)$ of the special fiber of the RZ space $\overline{\mathcal{M}}_{n,\varepsilon}^{[h]}$ (see Definition 2.2). The main result of the paper is the decomposition of the underlying reduced scheme:

$$\mathcal{N}_{n,\varepsilon,\text{red}}^{[h]} = \left(\bigcup_{\Lambda_1 \in \mathcal{L}_{\mathcal{Z}}} \mathcal{Z}(\Lambda_1) \right) \cup \left(\bigcup_{\Lambda_2 \in \mathcal{L}_{\mathcal{Y}}} \mathcal{Y}(\Lambda_2^\sharp) \right).$$

We also show that those subschemes satisfy nice inclusions relations corresponding to lattice inclusion (Theorem 2.4). These relations enable us to construct a stratification, the *Bruhat-Tits stratification*. The proof relies on Dieudonné theory to transform the problem into a semi-linear algebra problem, then it boils down to the crucial lemma, which is originally established in [VW11, RTW14] and later generalized in [KR12, HZ25].

We note that by definition, $\mathcal{Z}(\Lambda_1)$ and $\mathcal{Y}(\Lambda_2^\sharp)$ are closed subschemes of the special fiber. We prove that these subschemes are reduced. The proof proceeds by constructing a local model diagram

$$\begin{array}{ccc} & \tilde{\mathcal{Z}}(\Lambda_1) & \\ \varphi \swarrow & & \searrow \pi \\ \mathcal{Z}(\Lambda_1) & & \overline{\mathcal{M}}_n^{[h]}(t), \end{array}$$

where $\overline{\mathcal{M}}_n^{[2h]}(t)$ denotes the *strata local model* of type $t = t(\Lambda_1)$ (Definition 4.1). The morphisms φ and π are smooth of equal dimension. In particular, the \mathcal{Z} -stratum $\mathcal{Z}(\Lambda_1)$ shares many common geometric properties with its strata local model. Similar local model diagrams exist for both the \mathcal{Y} -stratum $\mathcal{Y}(\Lambda_2^\sharp)$ and the intersection stratum $\mathcal{Z}(\Lambda_1) \cap \mathcal{Y}(\Lambda_2^\sharp)$.

Similar to the local models associated to Shimura varieties and Rapoport-Zink spaces, strata local models are defined by purely linear algebraic data, enabling simpler analysis than Bruhat-Tits strata. By embedding strata local models into partial affine flag varieties and relating them

to Schubert varieties, we obtain the reducedness of strata local models and, consequently, of the corresponding Bruhat-Tits strata. It remains an interesting question to characterize strata local models and deduce more geometric properties using group-theoretical methods.

By explicit computations, we establish numerous geometric properties of the Bruhat-Tits strata, including normality, Cohen-Macaulayness, and precise dimension formulas. A complete list of these properties is presented in Theorem 2.3.

Prior to our work, the reducedness of Bruhat-Tits strata was established only in specific cases with smooth structure: see Vollaard-Wedhron [VW11, Thm. 3.10] and Li-Zhu [LZ17, Cor. 3.2.3] for unramified hyperspecial level and He-Li-Shi-Yang [HLSY23, §3] for Krämer models. It turns out that local model diagrams can be constructed whenever a version of the crucial lemma holds, for example, in unramified cases [Vol10, VW11, Cho18], ramified cases [RTW14, Wu16], higher signatures [FHI24, Tre23], and Krämer models [HLSY23] and various splitting models [HLS24, ZZ23, ZZ24]. Our technique for proving reducedness is likely to generalize to all these cases.

Finally, we establish an explicit isomorphism between Bruhat-Tits strata and Deligne-Lusztig varieties, extending similar results from [VW11, RTW14, Wu16]. This geometric connection not only illuminates the structure of Bruhat-Tits strata but has also proven important in arithmetic intersection theory, see, for example, [LZ22a, RTZ13, HLZ19].

The relationship between reduced locus of Bruhat-Tits strata and Deligne-Lusztig varieties has been studied group-theoretically in [GHN19]: the perfection of the special fiber of the Rapoport-Zink space is isomorphic to an ADLV. In our case, the ADLV is fully Newton-Hodge decomposable ([GHN19, §1.3]), and the EKOR strata decompose into weak Bruhat-Tits strata ([GHN24, §2.4]) where each stratum is isomorphic to the perfection of a Deligne-Lusztig variety. For Coxeter-type parahoric levels in the unitary group of signature $(n - 1, 1)$ (studied in [VW11, RTW14, Wu16]), [GHN24] shows that the weak Bruhat-Tits stratification coincides with the classical Bruhat-Tits stratification.

While the group-theoretical approach provides a comprehensive framework, it has some limitations: it is purely topological, and hence may lose geometric and arithmetic information. Moreover, it turns out that the lattice descriptions of Bruhat-Tits stratification are more useful in some arithmetic applications, such as describing special cycles.

We establish a scheme-theoretic relationship between Bruhat-Tits strata and Deligne-Lusztig varieties. Compared to the Coxeter type cases, several new phenomena emerge. Due to the extensive notation and results involved, we refer readers to §2.3 for a complete result. We want to point out a new phenomenon in the non-Coxeter type case. In [RTW14, Prop. 5.3], one of the important steps is studying the space

$$S_\Lambda = \{\mathcal{V} \subset (\Lambda^\sharp/\Lambda) \otimes_{\mathbb{F}_q} \mathbb{F} \mid \mathcal{V} \text{ is Lagrangian, and } \mathcal{V} \cap \Phi(\mathcal{V}) \stackrel{\leq 1}{\subseteq} \mathcal{V}\}.$$

Here Λ^\sharp/Λ is equipped with a natural symplectic structure (see §1.2) and extends to \mathbb{F} . The map Φ denotes the Frobenius action on this space. In loc. cit, it is shown that the space S_Λ admits a first-step decomposition:

$$S_\Lambda = X_P(\text{id}) \amalg X_P(w),$$

where $X_P(\text{id})$ corresponds to Φ -stable subspaces and $X_P(w)$ to the non- Φ -stable ones. We refer the reader to [RTW14, Prop. 5.3] for detailed definitions of the notation.

In the non-Coxeter type case, the subspaces \mathcal{V} parameterized by S_Λ are totally isotropic but not Lagrangian. The first-step decomposition consists of three components:

$$S_\Lambda = X_P(\text{id}) \amalg X_P(w) \amalg X_P(w'),$$

where $X_P(\text{id})$ corresponds to Φ -stable subspaces, while $X_P(w)$ (resp. $X_P(w')$) corresponds to non- Φ -stable subspaces \mathcal{V} such that $\mathcal{V} + \Phi(\mathcal{V})$ is isotropic (resp. non-isotropic). This decomposition relates to the Kottwitz-Rapoport strata; see Remark 7.11. Due to the additional pieces $X_P(w')$, the structure of S_Λ becomes more complicated than the Coxeter type. Nevertheless, we managed to give a complete description of the stratification in terms of Deligne-Lusztig varieties for any maximal vertex level, see §2.3 for a list of results.

1.1. Acknowledgement. Y. Luo would like to thank Michael Rapoport for his encouragement and comments on this paper. Q. He would like to thank Rong Zhou for collaboration and various discussions that were very helpful for the current project. Q. He and Y. Luo would like to thank the Institute for Advanced Study in Mathematics at Zhejiang University and the Morningside Center of Mathematics at Chinese Academy of Sciences for their hospitality during the Summer 2024 when part of this work was done. Y. Shi would like to thank Zhejiang University for its start-up grant.

1.2. Notation.

- Let F/F_0 be a ramified quadratic extension of a p -adic field, with $p \geq 3$. For any element $a \in F$, we denote its Galois conjugate over F_0 by \bar{a} . Let $\pi \in F$ and $\pi_0 \in F_0$ be uniformizers satisfying $\pi^2 = \pi_0$. We denote by \mathbb{F}_q their common residue field and by \mathbb{F} its algebraic closure.
- For any local field K , we denote by \check{K} the completion of the maximal unramified extension K . Let $\sigma \in \text{Gal}(\check{F}_0/F_0)$ be the Frobenius element. We fix an embedding of rings $i_0 : O_{F_0} \hookrightarrow O_{\check{F}_0}$ and an embedding $i : O_F \hookrightarrow O_{\check{F}}$ extending i_0 . Finally, we define $\bar{i} : O_F \rightarrow O_{\check{F}}$ by $a \mapsto i(\bar{a})$.
- Let (V, h) be an F/F_0 -hermitian space and $\Lambda \subset V$ be a lattice over F/F_0 . The hermitian dual of Λ is defined as

$$\Lambda^\sharp := \{v \in V \mid h(x, \Lambda) \in O_F\}.$$

We call Λ a *vertex lattice* if it satisfies

$$\pi\Lambda^\sharp \subseteq \Lambda \subseteq \Lambda^\sharp.$$

The *type* of a vertex lattice Λ , denoted $t(\Lambda)$, is defined as $\dim(\Lambda^\sharp/\Lambda)$.

- For any vertex lattice $\Lambda \subset V$, we define its base change to \check{F} as $\check{\Lambda} := \Lambda \otimes_{O_F} O_{\check{F}}$.
- Given a vertex lattice $\Lambda \subset V$, we define two \mathbb{F}_q -vector spaces:

$$V_\Lambda := \Lambda^\sharp/\Lambda, \quad \text{and} \quad V_{\Lambda^\sharp} = \Lambda/\pi\Lambda^\sharp.$$

The space V_Λ carries a natural symplectic structure given by

$$\langle \cdot, \cdot \rangle : V_\Lambda \times V_\Lambda \rightarrow \mathbb{F}_q, \quad \langle x, y \rangle \mapsto \pi h(\tilde{x}, \tilde{y}) \pmod{\pi},$$

where \tilde{x} and \tilde{y} denote arbitrary lifts of x and y to Λ^\sharp , resp.

The space V_{Λ^\sharp} carries a natural symmetric structure given by

$$(\cdot, \cdot) : V_{\Lambda^\sharp} \times V_{\Lambda^\sharp} \rightarrow \mathbb{F}_q, \quad (x, y) \mapsto h(\tilde{x}, \tilde{y}) \pmod{\pi}$$

where \tilde{x} and \tilde{y} denote arbitrary lifts of x and y to Λ , resp. Standard facts tell us that these two forms are non-degenerate.

- Let $\text{Nilp}_{O_{\check{F}}}$ denote the category of $O_{\check{F}}$ -schemes S where π is locally nilpotent on S . For any such scheme S , we denote its special fiber $S \times_{\text{Spf } O_{\check{F}}} \mathbb{F}$ by \bar{S} .

2. STATEMENTS OF THE MAIN RESULTS

Let F/F_0 be a ramified quadratic extension of p -adic fields ($p \geq 3$) with uniformizers π and π_0 respectively, satisfying the relation $\pi^2 = \pi_0$. Denote by \mathbb{F}_q the residue field of F and let $\mathbb{F} = \overline{\mathbb{F}_q}$ be its algebraic closure. Let \check{F} be the completion of the maximal unramified extension of F .

2.1. Bruhat-Tits strata. We will fix a framing object and define Rapoport-Zink (RZ) spaces, see §3.3 for details. Let $(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ be a fixed supersingular unitary O_{F_0} -module of rank n and type h with signature $(n-1, 1)$ (Definition 3.2). Here, supersingular means that the relative isocrystal has all relative slopes $1/2$. To this framing object we associate a hermitian space \mathbb{V} of dimension n over F (see §3.2). Let $\varepsilon = \varepsilon(\mathbb{V}) := -\text{Hasse}(\mathbb{V})$ denote the negative of its Hasse invariant.

The wedge RZ space $\mathcal{N}_{n, \varepsilon}^{[h], \wedge}$ associated to this framing object is a formal scheme over $\text{Spf } O_{\check{F}}$. It represents the moduli functor that assigns to each S in $\text{Nilp } O_{\check{F}}$ the set of isomorphism classes of tuples $(X, \iota, \lambda; \rho)$, where (X, ι, λ) is a unitary O_F -module over S and

$$\rho : X \times_S \bar{S} \rightarrow \mathbb{X} \times_{\mathbb{F}} \bar{S}$$

is an O_F -linear quasi-isogeny of height 0 over the special fiber \bar{S} . The RZ space $\mathcal{N}_{n, \varepsilon}^{[h]}$ is defined as the flat closure of $\mathcal{N}_{n, \varepsilon}^{[h], \wedge}$. We denote by $\bar{\mathcal{N}}_{n, \varepsilon}^{[h]}$ the special fiber of $\mathcal{N}_{n, \varepsilon}^{[h]}$ and by $\mathcal{N}_{n, \varepsilon, \text{red}}^{[h]}$ its reduced subscheme.

Recall the following lemma:

Lemma 2.1 ([RTW14, Lem. 6.1]). *Let κ be any perfect field over \mathbb{F} and let $M \subset \Lambda \otimes W_O(\kappa)$ be a $O_F \otimes W_O(\kappa)$ -lattice such that $M \subseteq M^\sharp$. Then M and M^\sharp are stable under F, V and Π . \square*

Let $\Lambda \subset \mathbb{V}$ be a vertex lattice of type t . By Lemma 2.1, both $\check{\Lambda}$ and $\check{\Lambda}^\sharp$ are stable under F, V and Π . By (relative) Dieudonné theory, the lattices $\check{\Lambda}$ and $\check{\Lambda}^\sharp$ correspond to strict $O_{\check{F}_0}$ -modules over \mathbb{F} , denoted X_Λ and X_{Λ^\sharp} resp, together with quasi-isogenies $\rho_\Lambda : X_\Lambda \rightarrow \mathbb{X}$ and $\rho_{\Lambda^\sharp} : X_{\Lambda^\sharp} \rightarrow \mathbb{X}$, resp.

Definition 2.2. Let $\mathcal{L}_{\mathcal{Z}}$ denote the set of all vertex lattices in \mathbb{V} of type $\geq h$, and let $\mathcal{L}_{\mathcal{Y}}$ denote the set of all vertex lattices in \mathbb{V} of type $\leq h$. We define the following two kinds of Bruhat-Tits (BT) strata:

(1) For any $\Lambda \in \mathcal{L}_{\mathcal{Z}}$, the \mathcal{Z} -stratum $\mathcal{Z}(\Lambda)$ is the subfunctor of $\bar{\mathcal{N}}_{n, \varepsilon}^{[h]}$ that assigns to each \mathbb{F} -scheme S the set of tuples $(X, \iota, \lambda, \rho)$ such that the composition $\rho_{\Lambda, X} := \rho^{-1} \circ (\rho_\Lambda)_S$ is an isogeny.

(2) For any $\Lambda \in \mathcal{L}_{\mathcal{Y}}$, the \mathcal{Y} -stratum $\mathcal{Y}(\Lambda^\sharp)$ is the subfunctor of $\bar{\mathcal{N}}_{n, \varepsilon}^{[h]}$ that assigns to each \mathbb{F} -scheme S the set of tuples $(X, \iota, \lambda, \rho)$ such that the composition $\rho_{\Lambda^\sharp, X^\vee} := \rho^\vee \circ \lambda_{\mathbb{X}} \circ \rho_{\Lambda^\sharp}$ is an isogeny, where $\rho_{\Lambda^\sharp} = \rho_\Lambda \circ \lambda_\Lambda^{-1}$.

By [RZ96, Lem. 2.10], $\mathcal{Z}(\Lambda)$ and $\mathcal{Y}(\Lambda^\sharp)$ are closed subschemes of $\overline{\mathcal{N}}_{n,\varepsilon}^{[h]}$. Following the arguments of [VW11, Lem. 4.2], both strata are representable by projective schemes over \mathbb{F} .

When $t_1 = t_2 = h$, the strata $\mathcal{Z}(\Lambda_1)$ and $\mathcal{Y}(\Lambda_2^\sharp)$ are each a single geometric point. In §4, we will prove the following results:

Theorem 2.3. *Let Λ_1 and Λ_2 be vertex lattices of type $t_i = t(\Lambda_i)$ with $t_2 < h < t_1$.*

(1) *The Bruhat-Tits strata $\mathcal{Z}(\Lambda_1)$, $\mathcal{Y}(\Lambda_2^\sharp)$, and their intersection $\mathcal{Z}(\Lambda_1) \cap \mathcal{Y}(\Lambda_2^\sharp)$ are normal and Cohen-Macaulay. In particular, they lie in the reduced subscheme $\mathcal{N}_{n,\varepsilon,\text{red}}^{[h]} \subset \overline{\mathcal{N}}_{n,\varepsilon}^{[h]}$.*

(2) *The dimensions are:*

- (i) $\dim \mathcal{Z}(\Lambda_1) = \frac{1}{2}(t_1 + h)$
- (ii) $\dim \mathcal{Y}(\Lambda_2^\sharp) = n - \frac{1}{2}(h - t_2) - 1$
- (iii) $\dim(\mathcal{Z}(\Lambda_1) \cap \mathcal{Y}(\Lambda_2^\sharp)) = \frac{1}{2}(t_1 - t_2) - 1$

(3) *When $h = 2\lfloor n/2 \rfloor$, all Bruhat-Tits strata and their intersections are smooth.*

(4) *When $h \neq 2\lfloor n/2 \rfloor$,*

- (i) $\mathcal{Z}(\Lambda_1)$ *is smooth if and only if $t_1 - h = 2$;*
- (ii) $\mathcal{Y}(\Lambda_2^\sharp)$ *is smooth if and only if $h - t_2 = 2$;*
- (iii) $\mathcal{Z}(\Lambda_1) \cap \mathcal{Y}(\Lambda_2^\sharp)$ *is smooth if and only if either $\mathcal{Z}(\Lambda_1)$ or $\mathcal{Y}(\Lambda_2^\sharp)$ is smooth.*

(5) *When $h \neq 2\lfloor n/2 \rfloor$ and $|h - t_i| > 2$:*

- (i) *The \mathcal{Z} -stratum is Gorenstein if and only if $t_1 = 3h + 4$*
- (ii) *The \mathcal{Y} -stratum is Gorenstein if and only if $t_2 = 3h - 2n$*
- (iii) $\mathcal{Z}(\Lambda_1) \cap \mathcal{Y}(\Lambda_2^\sharp)$ *is Gorenstein if and only if $2h = t_1 + t_2$*

The key ingredient in our proof is the local model diagram relating the Bruhat-Tits strata to their corresponding local models. For a vertex lattice $\Lambda \in \mathcal{L}_{\mathcal{Z}}$ of type $t \geq h$, we construct the following local model diagram:

$$\begin{array}{ccc}
 & \tilde{\mathcal{Z}}(\Lambda) & \\
 \varphi \swarrow & & \searrow \pi \\
 \mathcal{Z}(\Lambda) & & \overline{\mathcal{M}}_n^{[h]}(t),
 \end{array} \tag{2.1}$$

where $\overline{\mathcal{M}}_n^{[h]}(t)$ is the strata local model of type t (Definition 4.1). The construction and properties of this diagram are studied in §4.4. We show that the morphisms φ and π are smooth of equal relative dimension. Consequently, the proof of Theorem 2.3 reduces to verifying the corresponding properties for the strata local model $\overline{\mathcal{M}}_n^{[h]}(t)$, which we establish through explicit local chart computations in §4.5.

We want to note that the local model approach only allows us to determine certain local properties of Bruhat-Tits strata, such as reducedness and dimensions. To investigate global properties like connectedness, we must rely on the classical approach, which identifies Bruhat-Tits strata with Deligne-Lusztig varieties, see §2.3.

2.2. Bruhat-Tits stratification. The next main result of our paper is the following:

Theorem 2.4. *Recall that $\mathcal{L}_{\mathcal{Z}}$ (resp. $\mathcal{L}_{\mathcal{Y}}$) is the collections of vertex lattices in \mathbb{V} of type $\geq h$ (resp. $\leq h$). The reduced locus decomposes as:*

$$\mathcal{N}_{n,\varepsilon,\text{red}}^{[h]} = \left(\bigcup_{\Lambda_1 \in \mathcal{L}_{\mathcal{Z}}} \mathcal{Z}(\Lambda_1) \right) \cup \left(\bigcup_{\Lambda_2 \in \mathcal{L}_{\mathcal{Y}}} \mathcal{Y}(\Lambda_2^\sharp) \right),$$

with inclusions characterized by:

- $\mathcal{Z}(\Lambda_1) \subseteq \mathcal{Z}(\Lambda_1')$ if and only if $\Lambda_1 \supseteq \Lambda_1'$;
- $\mathcal{Y}(\Lambda_2^\sharp) \subseteq \mathcal{Y}(\Lambda_2'^\sharp)$ if and only if $\Lambda_2 \subseteq \Lambda_2'$;
- $\mathcal{Z}(\Lambda_1) \cap \mathcal{Y}(\Lambda_2^\sharp) \neq \emptyset$ if and only if $\Lambda_1 \subseteq \Lambda_2$;
- $\mathcal{Z}(\Lambda_1) \subseteq \mathcal{Y}(\Lambda_2^\sharp)$ if and only if $t(\Lambda_1) = h$ and $\Lambda_1 \supseteq \Lambda_2$;
- $\mathcal{Y}(\Lambda_2^\sharp) \subseteq \mathcal{Z}(\Lambda_1)$ if and only if $t(\Lambda_2) = h$ and $\Lambda_2 \subseteq \Lambda_1$.

Moreover, for any vertex lattice $\Lambda \in \mathcal{L}_{\mathcal{Z}} \cap \mathcal{L}_{\mathcal{Y}}$ (i.e., of type h), the stratum $\mathcal{Z}(\Lambda) = \mathcal{Y}(\Lambda^\sharp)$ is a discrete geometric point, called the worst point.

We refer the reader to Theorem 2.8 and Corollary 2.9 for the connectedness and irreducibility of the BT-strata and of the reduced locus of the RZ space.

Corollary 2.5. *Combining with the dimension formula in Theorem 2.3(2), we obtain:*

(1) *For special cases:*

- (i) ([RTW14, Thm. 1.1]) *When n is odd and $h = 0$, $\dim \mathcal{N}_{n,\varepsilon,\text{red}}^{[h]} = \frac{1}{2}(n - 1)$.*
- (ii) ([RTW14, Thm. 1.1]) *When n is even and $h = 0$ and $\varepsilon = 1$, $\dim \mathcal{N}_{n,\varepsilon,\text{red}}^{[h]} = \frac{1}{2}n - 1$.*
- (iii) ([RTW14, Thm. 1.1]) *When n is even and $h = 0$ and $\varepsilon = -1$, $\dim \mathcal{N}_{n,\varepsilon,\text{red}}^{[h]} = \frac{1}{2}n$.*
- (iv) ([Wu16, Thm. 5.18]) *When $h = n$, $\dim \mathcal{N}_{n,\varepsilon,\text{red}}^{[h]} = \frac{1}{2}n - 1$.*
- (v) ([Wu16, Thm. 5.18]) *When $h = n - 1$, $\dim \mathcal{N}_{n,\varepsilon,\text{red}}^{[h]} = \frac{1}{2}(n - 1)$.*
- (vi) *When $h = n - 2$ and $\varepsilon = -1$, $\dim \mathcal{N}_{n,\varepsilon,\text{red}}^{[h]} = \frac{1}{2}n - 2$.*

(2) *For the other cases*

$$\dim \mathcal{N}_{n,\varepsilon,\text{red}}^{[h]} = \begin{cases} \max \left\{ \frac{1}{2}(n + h - 1), n - \frac{1}{2}h + 1 \right\} & \text{if } n \text{ is odd;} \\ \max \left\{ \frac{1}{2}(n + h) - 1, n - \frac{1}{2}h + 1 \right\} & \text{if } n \text{ is even and } \varepsilon = 1; \\ \max \left\{ \frac{1}{2}(n + h), n - \frac{1}{2}h + 1 \right\} & \text{if } n \text{ is even and } \varepsilon = -1. \end{cases}$$

Example 2.6. (1) When $h = 0$, the reduced locus $\mathcal{N}_{n,\varepsilon,\text{red}}^{[0]}$ is connected and only consists of \mathcal{Z} -strata (note that while \mathcal{Y} -strata appear only at worst points, these can be replaced by \mathcal{Z} -strata):

$$\mathcal{N}_{n,\varepsilon,\text{red}}^{[0]} = \bigcup_{\Lambda \in \mathcal{L}_{\mathcal{Z}}} \mathcal{Z}(\Lambda) \quad \text{such that} \quad \mathcal{Z}(\Lambda) \subseteq \mathcal{Z}(\Lambda') \quad \text{if and only if} \quad \Lambda \supseteq \Lambda'.$$

Defining the open stratum $\mathcal{Z}^\circ(\Lambda) := \mathcal{Z}(\Lambda) \setminus \bigcup_{\Lambda' \subsetneq \Lambda} \mathcal{Z}(\Lambda')$, we have

$$\mathcal{N}_{n,\varepsilon,\text{red}}^{[0]} = \coprod_{\Lambda \in \mathcal{L}_{\mathcal{Z}}} \mathcal{Z}^\circ(\Lambda) \quad \text{with closure relations} \quad \overline{\mathcal{Z}^\circ(\Lambda)} = \mathcal{Z}(\Lambda) = \coprod_{\substack{\Lambda' \in \mathcal{L}_{\mathcal{Z}} \\ \Lambda' \subsetneq \Lambda}} \mathcal{Z}^\circ(\Lambda').$$

This recovers the Bruhat-Tits stratification described in [RTW14].

(2) Similarly, when $h = 2\lfloor n/2 \rfloor$, the reduced locus only consists of \mathcal{Y} -strata, recovering the results of [Wu16].

When $h \neq 0$ or $2\lfloor n/2 \rfloor$, both \mathcal{Z} -strata and \mathcal{Y} -strata can appear simultaneously. This complicates both the definition of open strata and their closure relations, this is the reason we stick to closed strata.

Example 2.7. Consider the case where $n = 2m$ is even and $h = n - 2$.

(1) When $\varepsilon = 1$, the hermitian space \mathbb{V} is non-split, and $\mathcal{L}_{\mathcal{Z}}$ consists of vertex lattices $\Lambda \subset C$ of type $n - 2 = h$. By Theorem 2.4(3), we obtain a Bruhat-Tits stratification involving only \mathcal{Y} -strata.

(2) When $h = n - 2 > 0$ and $\varepsilon = -1$, the hermitian space \mathbb{V} is split, and $\mathcal{L}_{\mathcal{Z}}$ consists of vertex lattices of types $n - 2$ and n . The reduced locus $\mathcal{N}_{n,-1,\text{red}}^{[n-2]}$ decomposes as a union of irreducible components:

$$\mathcal{N}_{n,-1,\text{red}}^{[n-2]} = \bigcup_{\substack{\Lambda_n \subset C \\ t(\Lambda_n)=n}} \mathcal{Z}(\Lambda_n) \cup \bigcup_{\substack{\Lambda_0 \subset C \\ t(\Lambda_0)=0}} \mathcal{Y}(\Lambda_0^\sharp),$$

where $\dim \mathcal{Z}(\Lambda_n) = n - 1$ and $\dim \mathcal{Y}(\Lambda_0^\sharp) = n/2$. The intersection $\mathcal{Z}(\Lambda_n) \cap \mathcal{Y}(\Lambda_0^\sharp)$ is nonempty if and only if $\Lambda_0 \subset \Lambda_n$, in which case $\dim \mathcal{Z}(\Lambda_n) \cap \mathcal{Y}(\Lambda_0^\sharp) = n/2 - 1$. This structure is parallel to the Balloon-Ground stratification in the unramified case studied in [KR12]; see also [LRZ24, §14.1].

2.3. Relation to Deligne-Lusztig variety. In this subsection, we address the relationship between Bruhat-Tits strata and Deligne-Lusztig varieties. This section summarizes the key results presented in §6 and §7. Before proceeding, we introduce the following notation:

- We refer the reader to Proposition 5.4 for the lattice description of Bruhat-Tits strata. For instance, for any $\Lambda \in \mathcal{L}_{\mathcal{Z}}$ and any perfect field κ over \mathbb{F} , we have

$$\mathcal{Z}(\Lambda)(\kappa) = \{(X, \iota_X, \lambda_X, \rho_X) \in \mathcal{N}_{n,\varepsilon}^{[h]}(\kappa) \mid \Lambda \otimes W_O(\kappa) \subseteq M(X) \subseteq M(X)^\sharp \subseteq \Lambda^\sharp \otimes W_O(\kappa)\},$$

where $W_O(\kappa)$ is the ring of ramified Witt vectors, see §3.1.

- The definitions of S_Λ , R_{Λ^\sharp} , and $S_{[\Lambda_1, \Lambda_2]}$ are given in equations (6.3), (6.12), and (6.13), resp. For instance, for any $\Lambda \in \mathcal{L}_{\mathcal{Z}}$ and any field k over \mathbb{F} , we have

$$S_\Lambda(k) = \{\mathcal{V} \subset (\Lambda^\sharp/\Lambda) \otimes_{\mathbb{F}_q} k \mid \mathcal{V} \text{ is isotropic, and } \dim \mathcal{V} = \frac{t-h}{2}, \text{ and } \dim(\mathcal{V} \cap \Phi(\mathcal{V})) \geq \frac{t-h}{2} - 1\}.$$

We now summarize our results.

- For any vertex lattice $\Lambda \in \mathcal{L}_{\mathcal{Y}}$ of type $t = t(\Lambda)$, since the types are always even integers, we denote $\mathbf{t} = t(\Lambda)/2$ and $\mathbf{h} = h/2$ for more convenient indexing. Theorem 6.3 yields the decomposition:

$$S_\Lambda = \left(\prod_{0 \leq j \leq \mathbf{h} < i \leq \mathbf{t}} X_{P_{ij}}(w_{ij}) \right) \amalg \left(\prod_{0 \leq j < \mathbf{h} < i \leq \mathbf{t}} X_{P_{ij}}(w'_{ij}) \right) \amalg X_{P_{\mathbf{h}\mathbf{h}}}(\text{id}).$$

This decomposition implies that S_Λ is irreducible. All group-theoretic data are related to the symplectic group. For the definitions of the parabolic subgroups P_{ij} and Weyl group elements w_{ij} and w'_{ij} , we refer to the discussion preceding Theorem 6.3.

- For any vertex lattice $\Lambda \in \mathcal{L}_y$ of type $t = t(\Lambda)$. We denote by $m = \lfloor n/2 \rfloor$, $\mathbf{h}' = m - \mathbf{h}$, and $\mathbf{t}' = m - t/2$. Theorem 6.10 yields the decomposition:

$$R_{\Lambda^\sharp} = \begin{cases} \prod_{\substack{0 \leq j \leq \mathbf{h}' < i \leq \mathbf{t}' \\ \delta \in \{\pm\}}} X_{P_{ij}}(w_{ij}^\delta) & \text{if } h = n, \\ \left(\prod_{0 \leq j \leq \mathbf{h}' < i \leq \mathbf{t}'} X_{P_{ij}}(w_{ij}) \right) \amalg \left(\prod_{\substack{0 \leq j < \mathbf{h}' < i \leq \mathbf{t}' \\ \delta \in \{\pm\}}} X_{P_{ij}}(w_{ij}^{\delta'}) \right) \amalg X_{P_{\mathbf{h}'\mathbf{h}'}}(\text{id}) & \text{if } h = n - 2, \\ \left(\prod_{0 \leq j \leq \mathbf{h}' < i \leq \mathbf{t}'} X_{P_{ij}}(w_{ij}) \right) \amalg \left(\prod_{0 \leq j < \mathbf{h}' < i \leq \mathbf{t}'} X_{P_{ij}}(w_{ij}') \right) \amalg X_{P_{\mathbf{h}'\mathbf{h}'}}(\text{id}) & \text{otherwise.} \end{cases}$$

This decomposition implies that R_{Λ^\sharp} has two disjoint irreducible components when $n = 2m$ is even and $h = n$; in all other cases, R_{Λ^\sharp} are irreducible. All group-theoretic data are related to the orthogonal group. For the definitions of the parabolic subgroups P_{ij} and Weyl group elements w_{ij} (w_{ij}^δ) and w_{ij}' ($w_{ij}^{\delta'}$), we refer to the discussion preceding Theorem 6.10.

- Let $\Lambda_1 \in \mathcal{L}_Z$ (resp. $\Lambda_2 \in \mathcal{L}_y$) be a vertex lattice of type $t_1 = t(\Lambda_1)$ (resp. $t_2 = t(\Lambda_2)$) such that $\Lambda_1 \subseteq \Lambda_2$. For more convenient indexing, we denote by $\mathbf{h} = h/2$ and $\mathbf{t}_1 = t_1/2$ and $\mathbf{t}_2 = t_2/2$. Proposition 6.11 yields the decomposition:

$$S_{[\Lambda_1, \Lambda_2]} = \prod_{\mathbf{t}_2 \leq j \leq \mathbf{h} \leq i \leq \mathbf{t}_1} X_{P_{ij}}(w_{ij}),$$

This decomposition implies that $S_{[\Lambda_1, \Lambda_2]}$ is irreducible. All group-theoretic data are related to the general linear group. For definitions of the parabolic subgroups P_{ij} and Weyl group elements w_{ij} , we refer to the discussion before Proposition 6.11.

Theorem 2.8. *Let $\Lambda_1 \in \mathcal{L}_Z$ and $\Lambda_2 \in \mathcal{L}_y$ be vertex lattices.*

- (1) *There are canonical isomorphisms of schemes over \mathbb{F} ,*

$$\Psi_Z : \mathcal{Z}(\Lambda_1) \cong S_{\Lambda_1} \quad \text{and} \quad \Psi_Y : \mathcal{Y}(\Lambda_2^\sharp) \cong R_{\Lambda_2^\sharp}. \quad (2.2)$$

Consequently, all \mathcal{Z} -strata $\mathcal{Z}(\Lambda_1)$ are irreducible. The \mathcal{Y} -stratum $\mathcal{Y}(\Lambda_2^\sharp)$ consists of two disjoint irreducible components when $n = 2m$ is even and $h = n$, and is irreducible otherwise.

- (2) *For $\Lambda_1 \subseteq \Lambda_2$, there is an isomorphism*

$$\mathcal{Z}(\Lambda_1) \cap \mathcal{Y}(\Lambda_2^\sharp) \cong S_{[\Lambda_1, \Lambda_2]}.$$

Consequently, the intersection $\mathcal{Z}(\Lambda_1) \cap \mathcal{Y}(\Lambda_2^\sharp)$ is irreducible when $t(\Lambda_2) < h < t(\Lambda_1)$. It is a discrete geometric point when $t(\Lambda_2) = h$ or $t(\Lambda_1) = h$.

The isomorphisms Ψ_Z and Ψ_Y are established in Theorems 7.3 and 7.9, respectively. Their geometric properties follow from Theorems 6.3 and 6.10, respectively. The intersection isomorphism is proven in Proposition 7.10, while its geometric properties are derived in Proposition 6.11.

Note that the geometric properties of Bruhat-Tits strata, including their dimension and normality, can also be derived from properties of the corresponding Deligne-Lusztig varieties.

Corollary 2.9. *The reduced locus $\mathcal{N}_{n, \varepsilon, \text{red}}^{[h]}$ is connected except when $h = n$, in which case it has exactly two connected components.*

Remark 2.10. (1) Theorem 6.3, Theorem 6.10, and Proposition 6.11 provide moduli descriptions for all Deligne-Lusztig varieties involved. Via the isomorphisms in (2.2), we can explicitly identify the corresponding strata in the reduced locus.

(2) An analogous result exists for the orthogonal case with vertex level. The reader may compare our Theorem 2.8 with [HZ25, Thm. 7.26].

(3) See also [LRZ25, §5] for a proof of Corollary 2.9 using the group-theoretical method.

2.4. Kottwitz-Rapoport stratification. To conclude this section, we discuss the relationship between the Kottwitz-Rapoport (KR) stratification and the Bruhat-Tits (BT) stratification. While this material is not essential for the main results of our paper, it presents several interesting connections. We begin by defining the (closed) Kottwitz-Rapoport strata in the reduced locus of the Rapoport-Zink space.

Definition 2.11. Let us define two \mathbb{F} -schemes associated with the reduced locus of $\mathcal{N}_{n,\varepsilon,\text{red}}^{[h]}$:

(1) The \mathbb{F} -scheme \mathcal{Z} is the reduced locus of the subfunctor of $\overline{\mathcal{N}}_{n,\varepsilon}^{[h]}$ consisting of tuples $(X, \iota, \lambda, \rho)$ over an \mathbb{F} -scheme S that satisfy: $\lambda^\vee(\text{Fil}(X^\vee)) \subseteq \iota(\pi)D(X)$.

(2) The \mathbb{F} -scheme \mathcal{Y} is the reduced locus of the subfunctor of $\overline{\mathcal{N}}_{n,\varepsilon}^{[h]}$ consisting of tuples $(X, \iota, \lambda, \rho)$ over an \mathbb{F} -scheme S that satisfy: $\lambda(\text{Fil}(X)) \subseteq \iota(\pi)D(X^\vee)$.

Remark 2.12. For strongly non-special h , i.e., when $h \neq 0, n - 2$ and n , the genuine Kottwitz-Rapoport stratification (see [He16, §1.6]) decomposes as:

$$\mathcal{N}_{n,\varepsilon,\text{red}}^{[h]} = (\mathcal{Z} \setminus \mathcal{Y}) \amalg (\mathcal{Y} \setminus \mathcal{Z}) \amalg (\mathcal{Z} \cap \mathcal{Y} \setminus \text{WT}) \amalg \text{WT}$$

where WT, called the *worst points*, is the disjoint union of all $\mathcal{Z}(\Lambda)$ satisfying $t(\Lambda) = h$ (see Theorem 2.4). This stratification is intimately related to the Kottwitz-Rapoport stratification of the special fiber of the corresponding local model of $\mathcal{N}_{n,\varepsilon}^{[h]}$. In the local model setting, the Kottwitz-Rapoport strata are indexed by admissible sets (see [PR08, §11] or [Zhu14, Thm. 8.1]), and when h is strongly non-special, a direct computation shows that admissible sets consists of four elements:

$$\begin{array}{ccc} w_1 & \searrow & \\ & & w_{12} \longrightarrow \text{id}, \\ w_2 & \nearrow & \end{array}$$

where the arrows indicate the Bruhat order. The extreme elements w_1 and w_2 correspond to $\mathcal{Z} \setminus \mathcal{Y}$ and $\mathcal{Y} \setminus \mathcal{Z}$ respectively, while w_{12} corresponds to $\mathcal{Z} \cap \mathcal{Y} \setminus \text{WT}$ and id corresponds to the worst point. In particular, the subvarieties \mathcal{Z} and \mathcal{Y} in our definition correspond to the irreducible components of the special fiber of the local model. In the special case when $n = h - 2$, the variety \mathcal{Z} further decomposes into two components; we will address this case in Remark 7.12.

We show in §5.3 that:

Proposition 2.13. (1) *The reduced locus decomposes as:*

$$\mathcal{N}_{n,\varepsilon,\text{red}}^{[h]} = \mathcal{Z} \cup \mathcal{Y}.$$

(2) For all lattices $\Lambda_1 \in \mathcal{L}_{\mathcal{Z}}$ and $\Lambda_2 \in \mathcal{L}_{\mathcal{Y}}$, we have $\mathcal{Z}(\Lambda_1) \subset \mathcal{Z}$ and $\mathcal{Y}(\Lambda_2^\sharp) \subset \mathcal{Y}$.

However, these inclusions are strictly proper. Indeed, in the strongly non-special case, as shown in Proposition 5.3, we have:

$$\bigcup_{\Lambda_1 \in \mathcal{L}_{\mathcal{Z}}} \mathcal{Z}(\Lambda_1) \subsetneq \mathcal{Z} \quad \text{and} \quad \bigcup_{\Lambda_2 \in \mathcal{L}_{\mathcal{Y}}} \mathcal{Y}(\Lambda_2^\sharp) \subsetneq \mathcal{Y}.$$

This reveals a non-trivial interaction between the BT-stratification and KR-stratification: the \mathcal{Z} -KR-stratum receives contributions from certain \mathcal{Y} -BT-strata, and vice versa (except if no $\mathcal{Z}(\Lambda)$, resp. $\mathcal{Y}(\Lambda^\sharp)$). The precise nature of these contributions can be determined explicitly from the moduli description.

In §7.4, we prove the following result:

Proposition 2.14. (1) The isomorphism $\Psi_{\mathcal{Z}}$ restricts to give:

$$\begin{aligned} \text{(i)} \quad & \mathcal{Z}(\Lambda_1) \setminus \mathcal{Y} \cong \prod_{i=0}^{\mathfrak{h}-1} X_{P_{[i, \mathfrak{h}]}}(w_{i, \mathfrak{h}}); \\ \text{(ii)} \quad & \mathcal{Z}(\Lambda_1) \cap \mathcal{Y} \cong \left(\prod_{0 \leq j < \mathfrak{h} < i \leq \mathfrak{t}} X_{P_{ij}}(w_{ij}) \right) \amalg \left(\prod_{0 \leq j < \mathfrak{h} < i \leq \mathfrak{t}} X_{P_{ij}}(w'_{ij}) \right) \amalg X_{P_{\mathfrak{h}\mathfrak{h}}}(\text{id}). \end{aligned}$$

(2) The isomorphism $\Psi_{\mathcal{Y}}$ similarly restricts to give:

$$\begin{aligned} \text{(i)} \quad & \mathcal{Y}(\Lambda_2^\sharp) \setminus \mathcal{Z} \cong \prod_{i=0}^{\mathfrak{h}'-1} X_{P_{i, \mathfrak{h}'}}(w_{i, \mathfrak{h}'}); \\ \text{(ii)} \quad & \mathcal{Y}(\Lambda_2^\sharp) \cap \mathcal{Z} \cong \begin{cases} \emptyset & \text{if } h = n, \\ \left(\prod_{0 \leq j < \mathfrak{h}' < i \leq \mathfrak{t}'} X_{P_{ij}}(w_{ij}) \right) \amalg \left(\prod_{\substack{0 \leq j < \mathfrak{h}' < i \leq \mathfrak{t}' \\ \delta \in \{\pm\}}} X_{P_{ij}}(w'_{ij}{}^\delta) \right) \amalg X_{P_{\mathfrak{h}'\mathfrak{h}'}}(\text{id}) & \text{if } h = n - 2, \\ \left(\prod_{0 \leq j < \mathfrak{h}' < i \leq \mathfrak{t}'} X_{P_{ij}}(w_{ij}) \right) \amalg \left(\prod_{0 \leq j < \mathfrak{h}' < i \leq \mathfrak{t}'} X_{P_{ij}}(w'_{ij}) \right) \amalg X_{P_{\mathfrak{h}'\mathfrak{h}'}}(\text{id}) & \text{otherwise.} \end{cases} \end{aligned}$$

These decompositions play a key role in refining KR strata to EKOR-strata (cf. [GHN24, §2.4]).

3. RAPOPORT-ZINK SPACES

In this section, we introduce the theory of ramified unitary Rapoport-Zink spaces with vertex-level structures.

3.1. Review of strict O_{F_0} -modules. We begin with a review of O_{F_0} -strict modules. For a comprehensive treatment, we refer to [Mih22, KRZ23, MLZ25]. Throughout this work, we assume $p \neq 2$.

Let F_0/\mathbb{Q}_p be an extension of p -adic fields with a fixed uniformizer $\pi_0 \in O_{F_0}$. We assume that the residue field $O_{F_0}/(\pi_0)$ is finite of order q . For an O_{F_0} -scheme S , a *strict O_{F_0} -module* over S is a pair (X, ι) consisting of a pair (X, ι) where X is a p -divisible group over S and $\iota : O_{F_0} \rightarrow \text{End}(X)$ is an action such that O_{F_0} acts on $\text{Lie}(X)$ via the structure morphism $O_{F_0} \rightarrow \mathcal{O}_S$. A strict O_{F_0} -module (X, ι) is called *formal* if its underlying p -divisible group X is formal. The dimension of a strict O_{F_0} -module is the dimension of its underlying p -divisible group. For the notion of (relative) *height* of strict O_{F_0} -modules, we direct the reader to the aforementioned references.

For any π_0 -adic algebra R over O_{F_0} , the *ring of ramified Witt vectors* is an O_{F_0} -algebra $W_{O_{F_0}}(R)$ which is part of a quadruple:

$$(W_O(R), I(R), \sigma, \dot{\sigma}) = (W_{O_{F_0}}(R), I_{O_{F_0}}(R), \sigma, \dot{\sigma}),$$

where $I(R) \subset W_O(R)$ is an ideal. $\sigma : W_O(R) \rightarrow W_O(R)$ and $\dot{\sigma} : I(R) \rightarrow W_O(R)$ are $W_O(R)$ -linear maps. See [ACZ16, §1.2.1] for the precise definition.

Definition 3.1 (*O*-displays, [ACZ16, §2.1]). Let R be a π_0 -adic O_{F_0} -algebra. An *O_{F_0} -display over R* is a quadruple $\mathcal{P} = (P, Q, \mathbf{F}, \dot{\mathbf{F}})$ whose entries are of the following kind:

- P is a finite projective $W_O(R)$ -module;
- $Q \subseteq P$ is a submodule with $I(R)P \subseteq Q$ and such that P/Q is a projective R -module;
- $\mathbf{F} : P \rightarrow P$ and $\dot{\mathbf{F}} : Q \rightarrow P$ are two σ -linear maps in the sense that

$$\mathbf{F}(\xi x) = \sigma(\xi)\mathbf{F}(x), \quad \dot{\mathbf{F}}(\xi y) = \sigma(\xi)\dot{\mathbf{F}}(y) \quad \text{for } \xi \in W_O(R), x \in P \text{ and } y \in Q.$$

We require that these data satisfy the following two axioms:

- (1) $\dot{\mathbf{F}}(Q)$ generates P as a $W_O(R)$ -module;
- (2) For $x \in P$ and $\xi \in I(R)$, we have

$$\dot{\mathbf{F}}(\xi x) = \dot{\sigma}(\xi)\mathbf{F}(x). \tag{3.1}$$

Let S be a formal scheme over $\mathrm{Spf} O_{F_0}$. By the result of Zink and Lau, which is generalized by Ahsendorf-Chen-Zink [ACZ16], we have a equivalence of categories:

$$\mathrm{BT} : \{\text{nilpotent } O_{F_0}\text{-displays}\} \xrightarrow{\sim} \{\text{strict formal } O_{F_0}\text{-modules}\}, \tag{3.2}$$

which is compatible with base change. We denote a quasi-inverse of this equivalence by

$$X \mapsto \mathcal{P}(X) = (P(X), Q(X), \mathbf{F}(X), \dot{\mathbf{F}}(X)).$$

This functor naturally induces a relative crystal $\mathbb{D}(X)$ valued in the category of O_{F_0} -pd-thickenings.

Let $D(X)$ denote the (covariant relative) de Rham realization of X , defined as

$$D(X) := P(X)/I(R)P(X) = \mathbb{D}(X)(R),$$

This is a locally free \mathcal{O}_S -module whose rank equals the height of X . The Hodge filtration $\mathrm{Fil}(X) = Q(X)/I(R)P(X) \subset D(X)$ fits into a canonical short exact sequence of \mathcal{O}_S -modules:

$$0 \rightarrow \mathrm{Fil}(X) \rightarrow D(X) \rightarrow \mathrm{Lie}(X) \rightarrow 0.$$

By (relative) Grothendieck-Messing theory, deformations of X along O_{F_0} -pd-thickenings are in canonical bijection with liftings of the Hodge filtration.

We now specialize to the case where $X = (X, \iota)$ is *biformal*, a notion introduced in [Mih22, Definition 11.9]. For such biformal strict O_{F_0} -modules, we can define the *relative dual* X^\vee and consequently the notion of *relative polarization* (see [Mih22, Definition 11.9]). The duality structure induces a perfect pairing

$$D(X) \times D(X^\vee) \rightarrow \mathcal{O}_S$$

such that $\mathrm{Fil}(X) \subset D(X)$ and $\mathrm{Fil}(X^\vee)$ are orthogonal complements to each other.

When $S = \text{Spec } R$ is perfect, the nilpotent O_{F_0} -display admits an alternative description via the relative Dieudonné module $M(X) := P(X)$ over $W_O(R)$. This module is equipped with a σ -linear operator F and a σ^{-1} -linear operator V satisfying the relation $FV = VF = \pi \text{id}$. This description yields an equivalence of categories.

3.2. Framing object.

Definition 3.2. Let h, n be integers with $0 \leq h \leq n$ and h even. For any $S \in \text{Nilp } O_{\check{F}}$, a *unitary O_F -module* of rank n and type h (with signature $(n-1, 1)$) over S is a triple (X, ι, λ) satisfying:

- (1) X is a strict biformal O_{F_0} -module over S of height $2n$ and dimension n .
- (2) $\iota : O_F \rightarrow \text{End}(X)$ is an action of O_F on X extending the O_{F_0} -action. The induced action of O_F on $\text{Fil}(X)$ satisfies:
 - (Kottwitz condition) The characteristic polynomial satisfies

$$\text{char}(\iota(\pi) \mid \text{Fil}(X)) = (T - \pi)(T + \pi)^{n-1};$$

- (Wedge condition) The following relations hold:

$$\bigwedge^2 (\iota(\pi) - \pi \mid \text{Fil}(X)) = 0; \quad \bigwedge^n (\iota(\pi) + \pi \mid \text{Fil}(X)) = 0.$$

- (Spin condition) When n is even and $h = n$, we require that $\iota(\pi) - \pi$ is non-vanishing on $\text{Fil}(X)$
- (3) λ is a (relative) polarization of X that is O_F/O_{F_0} -semilinear in the sense that the Rosati involution Ros_λ induces the non-trivial involution $\overline{(-)} \in \text{Gal}(F/F_0)$ on $\iota : O_F \rightarrow \text{End}(X)$.
- (4) We require that $\ker[\lambda] \subseteq X[\iota(\pi)]$ with order q^h .

By the kernel condition in (4), there exists a unique isogeny λ^\vee such that the composition

$$X \xrightarrow{\lambda} X^\vee \xrightarrow{\lambda^\vee} X$$

equals $\iota(\pi)$.

An isomorphism between two such triples $(X_1, \iota_1, \lambda_1) \xrightarrow{\sim} (X_2, \iota_2, \lambda_2)$ is an O_F -linear isomorphism $\varphi : X_1 \xrightarrow{\sim} X_2$ satisfying $\varphi^*(\lambda_2) = \lambda_1$.

The signature conditions imposed on unitary O_F -modules ensures that the associated Rapoport-Zink space has the property of “topological flatness” (see Proposition 3.4). In the special case where $h = n$, the spin condition is essential, see [RSZ17, Remark 3.11].

Let $(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ be a unitary O_F -module of dimension n over $\text{Spec } \mathbb{F}$. The associated rational Dieudonné module $N = M(\mathbb{X})[1/\pi_0]$ is a $2n$ -dimensional \check{F}_0 -vector space equipped with a σ -linear operator F and a σ^{-1} -linear operator V . Throughout this paper, we restrict our attention to the *supersingular* case, which means the rational Dieudonné module $N = M(\mathbb{X})[1/\pi_0]$ has all relative slopes $\frac{1}{2}$.

The O_F -action $\iota_{\mathbb{X}} : O_F \rightarrow \text{End}(\mathbb{X})$ induces on N an action that commutes with both operators F and V . The polarization of \mathbb{X} induces a skew-symmetric \check{F}_0 -bilinear form $\langle \cdot, \cdot \rangle$ on N satisfying

$$\langle Fx, y \rangle = \langle x, Vy \rangle^\sigma, \quad \langle \iota(a)x, y \rangle = \langle x, \iota(\bar{a})y \rangle, \quad \text{for any } x, y \in N, a \in O_F.$$

Furthermore, N is an n -dimensional \check{F} -vector space equipped with the \check{F}/\check{F}_0 -hermitian form $h(\cdot, \cdot)$ defined by:

$$h(x, y) := \delta(\langle \pi x, y \rangle + \pi \langle x, y \rangle),$$

where δ is a fixed element in $O_{\check{F}_0}^\times$ satisfying $\sigma(\delta) = -\delta$. The bilinear form $\langle \cdot, \cdot \rangle$ can be recovered from $h(\cdot, \cdot)$ via the relation:

$$\langle x, y \rangle = \frac{1}{2\delta} \operatorname{Tr}_{\check{F}/\check{F}_0}(\pi^{-1}h(x, y)).$$

Let $\tau := \Pi V^{-1}$ and define $\mathbb{V} := N^{\tau=1}$. Then \mathbb{V} is an F -vector space of dimension n , and we have $N = \mathbb{V} \otimes_{F_0} \check{F}_0$. The F/F_0 -hermitian form $h(\cdot, \cdot)$ restricts to \mathbb{V} , and we maintain this notation for the restricted form.

We define the *sign* of \mathbb{X} , denoted $\varepsilon = \varepsilon(\mathbb{X})$, as $-\operatorname{Hasse}(\mathbb{V})$, where $\operatorname{Hasse}(-)$ is the Hasse invariant of the hermitian space taking values in ± 1 . For any dimension n , type h , and sign ε , we denote by $(\mathbb{X}_{n,\varepsilon}^{[h]}, \iota_{\mathbb{X}_{n,\varepsilon}^{[h]}}, \lambda_{\mathbb{X}_{n,\varepsilon}^{[h]}})$ the corresponding *framing object*, which is a unitary O_F -module over $\operatorname{Spec} \mathbb{F}$. For the existence and uniqueness of these framing objects, we refer to [LRZ25, §5].

3.3. Two Rapoport-Zink spaces. In this subsection, we introduce two different Rapoport-Zink (RZ) spaces.

Definition 3.3. Let $(\mathbb{X}_{n,\varepsilon}^{[h]}, \iota_{\mathbb{X}_{n,\varepsilon}^{[h]}}, \lambda_{\mathbb{X}_{n,\varepsilon}^{[h]}})$ be a framing object over \mathbb{F} of rank n and type h .

(1) The wedge (relative) RZ space $\mathcal{N}_{n,\varepsilon}^{[h],\wedge}$ is the functor

$$\mathcal{N}_{n,\varepsilon}^{[h],\wedge} \longrightarrow \operatorname{Spf} O_{\check{F}}$$

that assigns to each scheme S the set of isomorphism classes of tuples $(X, \iota, \lambda, \rho)$, where

- (X, ι, λ) is a unitary O_F -module over S of dimension n and type h .
- $\rho : X \times_S \overline{S} \rightarrow \mathbb{X}_{n,\varepsilon}^{[h]} \times_{\mathbb{F}} \overline{S}$ is an O_F -linear quasi-isogeny of height 0 over the reduction $\overline{S} := S \times_{\operatorname{Spf} O_{\check{F}}} \operatorname{Spec} \mathbb{F}$ such that $\rho^*(\lambda_{\mathbb{X}_{n,\varepsilon}^{[h]}}|_{\overline{S}}) = \lambda_{\overline{S}}$.

(2) The (relative) RZ space $\mathcal{N}_{n,\varepsilon}^{[h]}$ is defined as the closed formal subscheme of $\mathcal{N}_{n,\varepsilon}^{[h],\wedge}$ cut out by the ideal sheaf $\mathcal{O}_{\mathcal{N}_{n,\varepsilon}^{[h],\wedge}}[\pi_0^\infty] \subset \mathcal{O}_{\mathcal{N}_{n,\varepsilon}^{[h],\wedge}}$. This is the maximal flat closed formal subscheme of $\mathcal{N}_{n,\varepsilon}^{[h],\wedge}$.

By [RZ96], the RZ spaces $\mathcal{N}_{n,\varepsilon}^{[h],\wedge}$ and $\mathcal{N}_{n,\varepsilon}^{[h]}$ are representable by formal schemes locally of finite type over $\operatorname{Spf} O_{\check{F}}$. Both spaces have relative dimension $n - 1$ (see Proposition 3.4). In two special cases, these spaces coincide: the self-dual case (when $h = 0$) and the π -modular case (when n is even and $h = n$), as shown in [Pap00, RSZ17]. In all other cases, the spaces are distinct. While there exists a moduli description of the RZ space $\mathcal{N}_{n,\varepsilon}^{[h]}$ using the strengthened spin condition [Luo24], we do not require it for our purposes.

Proposition 3.4. *For any field k over \mathbb{F} , the closed immersion between formal closed subschemes yields an equality of geometric points $\mathcal{N}_{n,\varepsilon}^{[h]}(k) = \mathcal{N}_{n,\varepsilon}^{[h],\wedge}(k)$. Consequently, these spaces share identical reduced loci.*

Proof. In the π -modular case, this proposition is established in [Wu16, Prop. 3.4]. We therefore assume $n \neq h$ and set $\mathfrak{h} = h/2$. Following [RZ96], with linear modification to $\mathcal{N}_{n,\varepsilon}^{[h]}$ as in [Pap00, Prop. 2.3], we have the local model diagram for $\mathcal{N}_{n,\varepsilon}^{[h],\wedge}$:

$$\begin{array}{ccccc}
 & & \tilde{\mathcal{N}}_{n,\varepsilon}^{[h]} & & \\
 & \swarrow & \downarrow & \searrow & \\
 \mathcal{N}_{n,\varepsilon}^{[h]} & & \tilde{\mathcal{N}}_{n,\varepsilon}^{[h],\wedge} & & \mathcal{M}_n^{[h]} \\
 \downarrow & \swarrow & \searrow & \downarrow & \\
 \mathcal{N}_{n,\varepsilon}^{[h],\wedge} & & & & \mathcal{M}_n^{[h],\wedge}
 \end{array}$$

where all diagonal arrows are smooth of equal relative dimension. The right-hand-side spaces are defined in Definition 4.1.

Using this diagram, our problem reduces to showing that the local model $\mathcal{M}_n^{[h],\wedge}$ is “topologically flat”, that is, the closed immersion $\mathcal{M}_n^{[h]} \hookrightarrow \mathcal{M}_n^{[h],\wedge}$ is defined by a nilpotent ideal.

To be more precise, for any point $(X, \iota, \lambda, \rho) \in \mathcal{N}_{n,\varepsilon}^{[h],\wedge}(k)$, [RZ96, Appendix] ensures the existence of an étale extension $\text{Spec } R \rightarrow \text{Spec } k$ with trivialization:

$$[\cdots \rightarrow D(X)_R \rightarrow D(X^\vee)_R \rightarrow \cdots] \simeq [\cdots \rightarrow \mathbf{L}_{-\mathfrak{h}} \otimes_{O_{F_0}} R \rightarrow \mathbf{L}_{\mathfrak{h}} \otimes_{O_{F_0}} R \rightarrow \cdots].$$

where \mathbf{L}_i denotes the standard lattice chain defined in (4.1). Under this trivialization, the Hodge filtration determines an R -point $(\mathcal{F}_i \subset \mathbf{L}_i \otimes_{O_{F_0}} R)$ of $\mathcal{M}_n^{[h],\wedge}$ (see Definition 4.1).

By the linear modification of the local model ([Pap00, Prop. 2.3]), a k -point $(X, \iota, \lambda, \rho) \in \mathcal{N}_{n,\varepsilon}^{[h],\wedge}(k)$ lies in $\mathcal{N}_{n,\varepsilon}^{[h]}(k)$ if and only if its Hodge filtration defines an R -point $(\mathcal{F}_i \subset \mathbf{L}_{i,R}) \in \mathcal{M}_n^{[h]}(R)$ after trivialization. Thus, our problem reduces to showing that $\mathcal{M}_n^{[h]}(R) = \mathcal{M}_n^{[h],\wedge}(R)$ for any étale k -algebra R , which would follow from the topological flatness of the wedge local model.

Now we establish the topological flatness. For odd $n = 2m + 1$, this follows directly from [Smi11]. For even $n = 2m$, by [Smi14, Thm. 9.6.1], we only need to verify that both local models are indexed by the same permissible elements in the double quotient $W_{\mathfrak{h}} \backslash \widetilde{W} / W_{\mathfrak{h}}$ of the affine Weyl group.

In [Smi14], it is shown that spin-permissibility (distinct from the spin condition in Definition 3.2) is equivalent to permissibility. Thus, we only need to prove that wedge-permissibility implies spin-permissibility in our case. For the relevant terminology, see [Smi14, §7].

By [Smi14, Proof of Prop. 7.2.2.], if an element $\tilde{w} \in W/W_{\mathfrak{h}}$ is naive-permissible, then it is a 2-face. From [Smi14, Prop. 7.2.2.], we have $0 \leq \mu_{\mathfrak{h}}^{\tilde{w}}(j) \leq 2$ for all j . Moreover, by [Smi14, Lem. 4.3.7]), we have the following basic inequality:

$$1 \leq \mu_{\mathfrak{h}}^{\tilde{w}}(j) + \mu_{\mathfrak{h}}^{\tilde{w}}(n+1-j) \leq 2 \quad \text{for all } j \in \{\mathfrak{h} + 1, \dots, n - \mathfrak{h}\}. \quad (3.3)$$

Consequently, for any $j \in \{\mathfrak{h} + 1, \dots, n - \mathfrak{h}\}$, if $\mu_{\mathfrak{h}}^{\tilde{w}}(j) \neq 0$, then either $\mu_{\mathfrak{h}}^{\tilde{w}}(j) = \mu_{\mathfrak{h}}^{\tilde{w}}(n+1-j) = 1$, or $\mu_{\mathfrak{h}}^{\tilde{w}}(j) = 2, \mu_{\mathfrak{h}}^{\tilde{w}}(n+1-j) = 0$.

For wedge-permissible elements \tilde{w} , [Smi14, Prop. 7.3.2] yields

$$\#\{j \mid \mu_{\mathfrak{h}}^{\tilde{w}}(j) = 0\} \leq 1 \quad (3.4)$$

According to [Smi14, Prop. 7.4.7], spin-permissibility is equivalent to wedge-permissibility together with the (P3) condition:

- (1) If $\mu_{\mathfrak{h}}^{\tilde{w}}$ is self-dual, i.e. if $\mu_{\mathfrak{h}}^{\tilde{w}} = \mu_{-\mathfrak{h}}^{\tilde{w}}$ ([Smi14, Def. 4.3.9]), then $\#\{j \mid \mu_{\mathfrak{h}}^{\tilde{w}}(j) = 0\} \equiv 1 \pmod{2}$;
- (2) In any case¹, there exists $j \in \{\mathfrak{h} + 1, \dots, n - \mathfrak{h}\}$ such that $\mu_{\mathfrak{h}}^{\tilde{w}}(j) = 1$.

When $\mathfrak{h} \leq m - 2$, we have $\#\{\mathfrak{h} + 1, \dots, n - \mathfrak{h}\} \geq 4$. In this case, wedge-permissibility (3.4) ensures the existence of at least one $j \in \mathfrak{h} + 1, \dots, n - \mathfrak{h}$ satisfying $\mu_{\mathfrak{h}}^{\tilde{w}}(j) = 1$.

When $\mathfrak{h} = m - 1$, if there exists no $j \in \{\mathfrak{h} + 1, \dots, n - \mathfrak{h}\} = \{m, m + 1\}$ such that $\mu_{\mathfrak{h}}^{\tilde{w}}(j) = 1$, then by [Smi14, Lem. 7.4.5], $\mu_{\mathfrak{h}}^{\tilde{w}}$ is self-dual. Therefore, it satisfies (P3)(1) and hence the spin condition. \square

Let $N = D(\mathbb{X}^{[h]}_{n, \varepsilon})[\frac{1}{\pi_0}]$ denote the rational Dieudonné module of the framing object.

Proposition 3.5. *Let κ be a perfect field over \mathbb{F} . There is a bijection between $\mathcal{N}_{n, \varepsilon, \text{red}}^{[h]}(\kappa)$ and the set of $W_O(\kappa)$ -lattices*

$$\left\{ M \subset N \otimes W_O(\kappa) \mid \pi M^\sharp \subseteq M \stackrel{h}{\subset} M^\sharp, \quad \Pi M \subseteq \tau^{-1}(M) \subseteq \Pi^{-1}M, \quad M \stackrel{\leq 1}{\subseteq} (M + \tau(M)) \right\}.$$

When n is even and $h = n$, the last relation is replaced by $M \stackrel{1}{\subset} (M + \tau(M))$.

Proof. Since the π -modular case is proved in [Wu16], we focus on the remaining cases. For any point in $\mathcal{N}_{n, \varepsilon, \text{red}}^{[h]}(\kappa) = \mathcal{N}_{n, \varepsilon, \text{red}}^{[h], \wedge}(\kappa)$, its Dieudonné module establishes a bijection between $\overline{\mathcal{N}}_{n, \varepsilon}^{[h]}(\kappa)$ and the set of $W_O(\kappa)$ -lattices:

$$\left\{ M \subset N \otimes W_O(\kappa) \mid \pi M^\sharp \subseteq M \stackrel{h}{\subset} M^\sharp, \quad \Pi M \subseteq M, \quad \pi_0 M \subset VM \stackrel{n}{\subset} M, \quad VM \stackrel{\leq 1}{\subseteq} VM + \Pi M \right\}.$$

where $M \stackrel{h}{\subset} M^\sharp$ comes from the polarization; $\Pi M \subset M$ due to the O_F -stability; $VM \stackrel{\leq 1}{\subset} VM + \Pi M$ follows from the Kottwitz and wedge conditions, where $VM/\pi_0 M \subset M/\pi_0 M$ is identified with the Hodge filtration. Moreover, we have the following equivalences: (1) $\pi_0 M \subset VM \subset M$ is equivalent to $\Pi M \subset \tau^{-1}(M) \subset \Pi^{-1}M$; (2) $VM \stackrel{\leq 1}{\subset} VM + \Pi M$ is equivalent to $M \stackrel{\leq 1}{\subset} (M + \tau(M))$. This completes the proof. \square

4. LOCAL MODELS OF BASIC LOCUS STRATA

In this section, we introduce local models of Bruhat-Tits strata.

4.1. Standard polarized lattice chain. In this subsection, we recall the basic setup for local models.

Consider the vector space F^n with the standard F -basis e_1, \dots, e_n . We equip it with a split F/F_0 -Hermitian form

$$h : F^n \times F^n \rightarrow F, \quad h(ae_i, be_j) = \bar{a}b\delta_{i, n+1-j}, \quad a, b \in F.$$

¹While the statement of [Smi14, Prop. 7.4.7] may be ambiguous, note that the existence of $j \in \{\mathfrak{h} + 1, \dots, n - \mathfrak{h}\}$ such that $\mu_{\mathfrak{h}}^{\tilde{w}}(j) = 1$ is equivalent to $a_{\mathfrak{h}} > a_{\mathfrak{h}}^\perp$, see the bottom of page 333 (resp. 334) for the definition of $a_{\mathfrak{h}}$ (resp. $a_{\mathfrak{h}}^\perp$) and the middle of page 336 for the proof that this implies the spin condition.

Attached to ϕ are the respective alternating and symmetric F_0 -bilinear forms $F^n \times F^n \rightarrow F_0$ given by

$$\langle x, y \rangle := \frac{1}{2} \operatorname{Tr}_{F/F_0}(\pi^{-1}h(x, y)) \quad \text{and} \quad (x, y) := \frac{1}{2} \operatorname{Tr}_{F/F_0}(h(x, y)).$$

For each integer $i = bn + c$ with $0 \leq c < n$, define the standard O_F -lattices

$$\mathbf{L}_i := \sum_{j=1}^c \pi^{-b-1} O_F e_j + \sum_{j=c+1}^n \pi^{-b} O_F e_j \subset F^n. \quad (4.1)$$

For all i , the \langle, \rangle -dual \mathbf{L}^\vee of \mathbf{L}_i in F^n is \mathbf{L}_{-i} , by which we mean that

$$\{x \in F^n \mid \langle \mathbf{L}_i, x \rangle \subset O_{F_0}\} = \mathbf{L}_{-i}.$$

By restriction, we have a perfect O_{F_0} -bilinear pair:

$$\mathbf{L}_i \times \mathbf{L}_{-i} \xrightarrow{\langle, \rangle} O_{F_0}$$

Similarly, \mathbf{L}_{n-i} is the $(,)$ -dual of \mathbf{L}_i in F^n . The \mathbf{L}_i 's forms a complete, periodic, self-dual lattice chain

$$\cdots \subset \mathbf{L}_{-2} \subset \mathbf{L}_{-1} \subset \mathbf{L}_0 \subset \mathbf{L}_1 \subset \mathbf{L}_2 \subset \cdots.$$

For any subset $I \subset \{1, \dots, m\}$ with $m = \lfloor \frac{n}{2} \rfloor$, we define the standard polarized chain \mathbf{L}_I as a sub-lattice chain with indices $i \in \pm I + n\mathbb{Z}$.

For even integers r, s, t , we define the following index sets: $[s] := \{\pm \mathfrak{s}\} + n\mathbb{Z}$ and $[s, t] := \{\pm \mathfrak{s}, \pm \mathfrak{t}\} + n\mathbb{Z}$ and $[r, s, t] := \{\pm \mathfrak{r}, \pm \mathfrak{s}, \pm \mathfrak{t}\} + n\mathbb{Z}$. For any such index set I , we define the standard lattice chain $\mathbf{L}_I := \{\mathbf{L}_i\}_{i \in I}$. This chain is self-dual in the sense of [RZ96]. We use these half-integral indices to maintain consistency with the local model indexing conventions used in [PR09, Luo24].

4.2. Local models. We first recall the definition of the relative local model. For any O_F -algebra R , let $\mathbf{L}_{i,R}$ denote the tensor product $\mathbf{L}_i \otimes_{O_{F_0}} R$. Let $\Pi := \pi \otimes 1$ and $\pi := 1 \otimes \pi$.

Definition 4.1. (1) The *wedge local model* $\mathcal{M}_n^{[h], \wedge}$ is a projective scheme over $\operatorname{Spec} O_F$. It represents the moduli problem that assigns to each O_F -algebra R the set of all families $(\mathcal{F}_i \subset \mathbf{L}_{i,R})_{i \in [h]}$ such that:

LM1. for all $i \in [h]$, \mathcal{F}_i is an $O_F \otimes_{O_{F_0}} R$ -submodule of $\mathbf{L}_{i,R}$, and an R -direct summand of rank n ;

LM2. for all $i, j \in [h]$ with $i < j$, the natural arrow $\mathbf{L}_{i,R} \rightarrow \mathbf{L}_{j,R}$ carries \mathcal{F}_i into \mathcal{F}_j ;

LM3. for all $i \in [h]$, the isomorphism $\mathbf{L}_{i,R} \xrightarrow{\Pi} \mathbf{L}_{i-n,R}$ identifies

$$\mathcal{F}_i \xrightarrow{\sim} \mathcal{F}_{i-n};$$

LM4. for all $i \in [h]$, the perfect R -bilinear pairing

$$\mathbf{L}_{i,R} \times \mathbf{L}_{-i,R} \xrightarrow{\langle -, - \rangle \otimes R} R$$

identifies \mathcal{F}_i^\perp with \mathcal{F}_{-i} inside $\mathbf{L}_{-i,R}$; and

LM5. For all $i \in [h]$, the action of $\Pi = \pi \otimes 1 \in O_F \otimes_{O_{F_0}} R$ on \mathcal{F}_i satisfies the following signature conditions:

- (Kottwitz condition) The characteristic polynomial satisfies:

$$\operatorname{char}(\Pi \mid \mathcal{F}_i) = (T - \pi)(T + \pi)^{n-1};$$

- (Wedge condition) The following operators vanish:

$$\bigwedge^2(\Pi - \pi | \mathcal{F}_i) = 0; \quad \bigwedge^n(\Pi + \pi | \mathcal{F}_i) = 0.$$

- (Spin condition) For even $n = 2m$ and $h = n$, the operator $\Pi - \pi$ is non-zero on \mathcal{F}_m

(2) We define the *local model* $\mathcal{M}_n^{[h]}$ as the flat closure of $\mathcal{M}_n^{[h],\wedge}$. To be more precise, it is the scheme-theoretic closure of the generic fiber of the wedge local model:

$$\begin{array}{ccc} \mathcal{M}_n^{[h]} & \hookrightarrow & \mathcal{M}_n^{[h],\wedge} \\ \uparrow & \nearrow & \\ \mathcal{M}_{n,\eta}^{[h],\wedge} & & \end{array}$$

4.3. Strata local model. In this subsection, we define the strata local model and study its basic properties.

Definition 4.2. Let R be an \mathbb{F}_q -algebra and let $\mathbf{L}_{[h]}$ and $\mathbf{L}_{[h,t]}$ be two standard lattice chains.

(1) The *pivoting filtration*² of type t is the fixed filtration with indices in $[t]$:

$$(\mathcal{F}_i := \prod_{i,R} \mathbf{L}_{i,R})_{i \in [t]}.$$

(2) For any R -point $(\mathcal{F}_i \subset \mathbf{L}_{i,R})_{i \in [h]}$ in $\mathcal{M}_{n,\mathbb{F}}^{[h]}$, we say it is *pinned by the pivoting filtration* of type t if for any $i < j$ with either $i \in [h], j \in [t]$ or $i \in [t], j \in [h]$, the natural morphism $\mathbf{L}_{i,R} \rightarrow \mathbf{L}_{j,R}$ maps \mathcal{F}_i into \mathcal{F}_j . Here, when $i \in [t]$, \mathcal{F}_i refers to the pivoting filtration.

(3) The *strata local model* $\overline{\mathcal{M}}_n^{[h]}(t)$ is defined as the closed subscheme of the special fiber $\overline{\mathcal{M}}_n^{[h]}$ that parameterizes all points pinned by the pivoting filtration of type t .

We will relate these to the Bruhat-Tits strata in the next subsection.

Theorem 4.3. *The strata local model $\overline{\mathcal{M}}_n^{[h]}(t)$ is reduced.*

Proof. We use the theory of unitary affine flag varieties $\mathcal{F}\ell_I$ as developed in [PR08]. Let $\mathbb{F}((u))/\mathbb{F}((t))$ be the ramified quadratic extension of function fields with $u^2 = t$. The standard polarized lattice chains \mathbb{L}_i for $\mathbb{F}((u))/\mathbb{F}((t))$ are defined as in §4.1. Via the isomorphism $\mathbf{L}_i/\pi_0\mathbf{L}_i \simeq u^{-1}\mathbb{L}_i/u\mathbb{L}_i$, one can embed the special fiber of the local model $\overline{\mathcal{M}}_n^{[h]}$ into the affine flag varieties. The pivoting filtration $(\prod_{i,\mathbb{F}} \mathbf{L}_{i,\mathbb{F}} \subset \mathbf{L}_{i,\mathbb{F}})_{i \in [t]}$ lifts to the geometric point $(u\mathbb{L}_i \subset \mathbb{L}_i \subset u^{-1}\mathbb{L}_i)_{i \in [t]} =: *_{[t]} \in \mathcal{F}\ell_{[t]}(\mathbb{F})$, which is an $L^+P_{[t]}$ -invariant point. Consider the following diagram:

$$\begin{array}{ccc} & \mathcal{F}\ell_{[h,t]} & \\ p_h \swarrow & & \searrow p_t \\ \overline{\mathcal{M}}_n^{[h]} \subset \mathcal{F}\ell_{[h]} & & \mathcal{F}\ell_{[t]} \ni *_{[t]} \end{array}$$

From the moduli description, we have the equality:

$$\overline{\mathcal{M}}_n^{[h]}(t) = p_h^{-1}(\overline{\mathcal{M}}_n^{[h]}) \cap p_t^{-1}(*_{[t]}).$$

²This terminology is due to S. Kudla.

To be more precise, an R -point of $p_h^{-1}(\overline{\mathcal{M}}_n^{[h]})$ is a filtration $(\mathcal{F}_i \subset \mathbf{L}_{i,R})_{i \in [h,t]}$ such that the subfamily $(\mathcal{F}_i \subset \mathbf{L}_{i,R})_{i \in [h]}$ defines a point of $\overline{\mathcal{M}}_{[h]}^{\text{loc}}(R)$. The intersection $p_h^{-1}(\overline{\mathcal{M}}_n^{[h]}) \cap p_t^{-1}(*_{[t]})$ parameterizes those filtrations $(\mathcal{F}_i \subset \mathbf{L}_{i,R})_{i \in [h,t]}$ where $(\mathcal{F}_i \subset \mathbf{L}_{i,R})_{i \in [t]}$ is the pivoting filtration.

Since $*_{[t]}$ is $L^+P_{[t]}$ -invariant, it is a Schubert variety. Thus, its preimage $p_t^{-1}(*_{[t]})$ is a union of Schubert varieties. By the Coherence conjecture, the special fiber $\overline{\mathcal{M}}_n^{[h]} \subset \mathcal{F}\ell_{[h]}$ of the local model is a union of Schubert varieties (see [PR09, Thm. 4.1]). Therefore, $\overline{\mathcal{M}}_n^{[h]}(t)$ is an intersection of unions of Schubert varieties. It is compatibly Frobenius split in a sufficiently large Schubert variety and hence reduced, see [GÖ1, §2]. \square

One can also define the strata local model $\overline{\mathcal{M}}_n^{[h]}(t_1, t_2)$ with respect to two classes of pivoting filtrations. The cases where $t_1, t_2 < h$ or $t_1, t_2 > h$ are degenerate. For $t_1 < h < t_2$, we obtain a closed subscheme of the special fiber of the local model that fits into the following diagram:

$$\begin{array}{ccccccccc} \mathbf{L}_{-t_1,R} & \longrightarrow & \mathbf{L}_{-h,R} & \longrightarrow & \mathbf{L}_{-t_2,R} & \longrightarrow & \mathbf{L}_{t_2,R} & \longrightarrow & \mathbf{L}_{h,R} & \longrightarrow & \mathbf{L}_{t_1,R} \\ \uparrow & & \uparrow & & \uparrow & & \uparrow & & \uparrow & & \uparrow \\ \Pi\mathbf{L}_{-t_1,R} & \longrightarrow & \mathcal{F}_{-h,R} & \longrightarrow & \Pi\mathbf{L}_{-t_2,R} & \longrightarrow & \Pi\mathbf{L}_{t_2,R} & \longrightarrow & \mathcal{F}_{h,R} & \longrightarrow & \Pi\mathbf{L}_{t_1,R} \end{array} .$$

The strata local model $\overline{\mathcal{M}}_n^{[h]}(t_1, t_2)$ is again reduced. This follows from the same strategy as in Theorem 4.3: considering projections of affine flag varieties

$$\begin{array}{ccc} & \mathcal{F}\ell_{[h,t_1,t_2]} & \\ p_{t_1} \swarrow & \downarrow p_h & \searrow p_{t_2} \\ \mathcal{F}\ell_{[t_1]} & \mathcal{F}\ell_{[h]} & \mathcal{F}\ell_{[t_2]} \end{array} .$$

The strata local model now is the intersection

$$\overline{\mathcal{M}}_n^{[h]}(t_1, t_2) = p_h^{-1}(\overline{\mathcal{M}}_n^{[h]}) \cap p_{t_1}^{-1}(*_{[t_1]}) \cap p_{t_2}^{-1}(*_{[t_2]}).$$

Remark 4.4. The strata local model can also be defined when F/F_0 is unramified. In this case, the pivoting filtration is the filtration $(V\Lambda/\pi\Lambda \subset \Lambda/\pi\Lambda)$, where V is the Verschiebung of the rational Dieudonné module of the framing object. This lifts to a point $(\pi\Lambda \subset V\Lambda \subset \Lambda)$ in the unramified unitary affine flag variety as a closed Schubert cell.

Since the unramified unitary group splits over $W_{O_{F_0}}(\mathbb{F})[\frac{1}{\pi_0}]$, we can reduce to the local model and affine Grassmannian of GL_n . In this case, the vertex lattices correspond to the “vertex lattices” \mathcal{L}_i^\pm studied in [Cho18].

For example, the strata local model $\overline{\mathcal{M}}_n^{[0]}(t)$ for $t = 2k + 1$ parameterizes one dimensional subspaces $\mathcal{F}_0 \subset \Lambda_{0,R}$ with the following factorization:

$$\begin{array}{ccccc} \mathbf{L}_{-k-1,R} & \longrightarrow & \mathbf{L}_{0,R} & \longrightarrow & \mathbf{L}_{k,R} \\ \uparrow & & \uparrow & & \uparrow \\ (0) & \longrightarrow & \mathcal{F}_{0,R} & \longrightarrow & (0) \end{array}$$

While this does not correspond to any classical local model, the proof of Theorem 4.3 still applies. A direct computation shows that $\overline{\mathcal{M}}_n^{[0]}(t)$ is smooth.

4.4. Local model diagram. In this subsection, we establish the local model diagram connecting the Bruhat-Tits strata to the strata local model. This construction is conceptual in nature, and does not require a moduli-theoretic description.

Definition 4.5. Let Λ be a vertex lattice.

(1) For $t(\Lambda) \geq h$, define $\tilde{\mathcal{Z}}(\Lambda)$ to be a projective formal scheme over \mathbb{F} that represents the functor sending each \mathbb{F} -algebra R to the set of tuples $(X, \iota, \lambda, \rho; f)$, where:

- $(X, \iota, \lambda, \rho) \in \mathcal{Z}(\Lambda)(R)$ is an R -point of the \mathcal{Z} -stratum $\mathcal{Z}(\Lambda)$;
- f is an isomorphism between the standard lattice chain $\mathcal{L}_{[h,t],R}$ and the lattice chain of de Rham realizations:

$$f : \begin{array}{ccccccc} \mathbf{L}_{-t,R} & \longrightarrow & \mathbf{L}_{-h,R} & \longrightarrow & \mathbf{L}_{h,R} & \longrightarrow & \mathbf{L}_{t,R} \\ \downarrow \sim & & \downarrow \sim & & \downarrow \sim & & \downarrow \sim \\ D(X_\Lambda) & \xrightarrow{\rho_{\Lambda,*}} & D(X) & \xrightarrow{\lambda_*} & D(X^\vee) & \xrightarrow{\rho_{\Lambda^\sharp}^\vee} & D(X_{\Lambda^\sharp}). \end{array}$$

(2) For $t(\Lambda) \leq h$, define $\tilde{\mathcal{Y}}(\Lambda^\sharp)$ to be a projective formal scheme over \mathbb{F} that represents the functor sending each \mathbb{F} -algebra R to the set of tuples $(X, \iota, \lambda, \rho; f)$, where:

- $(X, \iota, \lambda, \rho) \in \mathcal{Y}(\Lambda^\sharp)(R)$ is a R -point of the \mathcal{Y} -stratum $\mathcal{Y}(\Lambda^\sharp)$;
- f is an isomorphism between the standard lattice chain $\mathcal{L}_{[h,t],R}$ and the lattice chain of de Rham realizations:

$$f : \begin{array}{ccccccc} \mathbf{L}_{t,R} & \longrightarrow & \mathbf{L}_{h,R} & \longrightarrow & \mathbf{L}_{n-h,R} & \longrightarrow & \mathbf{L}_{n-t,R} \\ \downarrow \sim & & \downarrow \sim & & \downarrow \sim & & \downarrow \sim \\ D(X_{\Lambda^\sharp}) & \longrightarrow & D(X^\vee) & \xrightarrow{\lambda_*^\vee} & D(X) & \longrightarrow & D(X_\Lambda). \end{array}$$

We now construct the local model diagram for the \mathcal{Z} -strata. The construction and properties for the \mathcal{Y} -strata follow analogously. For a vertex lattice Λ of type $t(\Lambda) \geq h$, we construct the following local model diagram:

$$\begin{array}{ccc} & \tilde{\mathcal{Z}}(\Lambda) & \\ \varphi \swarrow & & \searrow \pi \\ \mathcal{Z}(\Lambda) & & \overline{\mathcal{M}}_n^{[h]}(t). \end{array} \tag{4.2}$$

The map φ is defined by forgetting the trivialization f :

$$\varphi : \tilde{\mathcal{Z}}(\Lambda) \rightarrow \mathcal{Z}(\Lambda), \quad (X, \iota, \lambda, \rho; f) \mapsto (X, \iota, \lambda, \rho).$$

Let $\mathcal{G} := \text{Aut}(\mathbf{L}_{[h,t]})$ be the group scheme over O_{F_0} consisting of automorphisms of $\mathbf{L}_{[h,t]}$ that preserves the PEL data. By [RZ96, Thm. 3.16], $\mathcal{G} \otimes_{O_{F_0}} O_F$ is a smooth group scheme acting on $\tilde{\mathcal{Z}}(\Lambda)$. Moreover:

Theorem 4.6 ([RZ96, Thm. 3.16]). *The morphism φ is a $\mathcal{G}_{\mathbb{F}}$ -torsor. In particular, φ is a smooth morphism of relative dimension $\dim \mathcal{G}_{\mathbb{F}}$. \square*

The morphism π is defined by attaching the de Rham realization of the strict O_{F_0} -modules to the lattice chain via f . To be more precise, for an R -point of $\tilde{\mathcal{Z}}(\Lambda)$, we have morphisms

$$X_{\Lambda,R} \xrightarrow{\rho_\Lambda} X \xrightarrow{\lambda} X^\vee \xrightarrow{\rho_{\Lambda^\#}} X_{\Lambda^\#,R}.$$

By taking the Hodge filtration, we obtain a chain of filtrations:

$$\begin{array}{ccccccc} D(X_\Lambda) & \longrightarrow & D(X) & \longrightarrow & D(X^\vee) & \longrightarrow & D(X_{\Lambda^\#}) \\ \uparrow & & \uparrow & & \uparrow & & \uparrow \\ \text{Fil}(X_\Lambda) & \longrightarrow & \text{Fil}(X) & \longrightarrow & \text{Fil}(X^\vee) & \longrightarrow & \text{Fil}(X_{\Lambda^\#}) \end{array}$$

Recall that we define $X_{\Lambda,R} = X_\Lambda \times_{\mathbb{F}} R$ and $X_{\Lambda^\#,R} = X_{\Lambda^\#} \times_{\mathbb{F}} R$. Then we have

$$\text{Fil}(X_\Lambda) = \Pi D(X_\Lambda) \otimes_{\mathbb{F}} R, \quad \text{Fil}(X_{\Lambda^\#}) = \Pi D(X_{\Lambda^\#}) \otimes_{\mathbb{F}} R.$$

Applying f , we obtain the desired filtration in $\overline{\mathcal{M}}_n^{[h]}(t)$:

$$\begin{array}{ccccccc} \mathbf{L}_{-\mathbf{t},R} & \longrightarrow & \mathbf{L}_{-\mathbf{n},R} & \longrightarrow & \mathbf{L}_{\mathbf{n},R} & \longrightarrow & \mathbf{L}_{\mathbf{t},R} \\ \uparrow & & \uparrow & & \uparrow & & \uparrow \\ \Pi \mathbf{L}_{-\mathbf{t},R} & \longrightarrow & f(\text{Fil}(X)) & \longrightarrow & f(\text{Fil}(X^\vee)) & \longrightarrow & \Pi \mathbf{L}_{\mathbf{t},R} \end{array}$$

Proposition 4.7 (Grothendieck-Messing). *The morphism π is smooth of relative dimension equal to $\dim \mathcal{G}_{\mathbb{F}}$.*

Proof. Let R_0 an \mathbb{F} -algebra and $\text{Spec } R_0 \hookrightarrow \text{Spec } R$ be a first-order thickening. Let $x \in \overline{\mathcal{M}}_n^{[h]}(t)(R)$ be an R -point with reduction $\bar{x} \in \overline{\mathcal{M}}_n^{[h]}(t)(R_0)$. Let $f_0 \in \tilde{\mathcal{Z}}(\Lambda)(R_0)$ be an R_0 -point that lifts \bar{x} :

$$\begin{array}{ccc} & & \tilde{\mathcal{Z}}(\Lambda) \\ & \nearrow f_0 & \downarrow \pi \\ \text{Spec } R_0 & \hookrightarrow \text{Spec } R & \xrightarrow{x} \overline{\mathcal{M}}_n^{[h]}(t). \end{array}$$

To show formal smoothness, we need to construct a lift f as shown in the diagram. The existence of such a lift is equivalent to finding compatible lifts of the following data:

$$\begin{array}{ccccccc} X_{\Lambda,R} \rightarrow X \rightarrow X^\vee \rightarrow X_{\Lambda^\#,R} & \rightsquigarrow & X_{\Lambda,R_0} \rightarrow \overline{X} \rightarrow \overline{X}^\vee \rightarrow X_{\Lambda^\#,R_0}. & (4.3) \\ \mathbf{L}_{-\mathbf{t},R} \rightarrow \mathbf{L}_{-\mathbf{n},R} \rightarrow \mathbf{L}_{\mathbf{n},R} \rightarrow \mathbf{L}_{\mathbf{t},R} & \rightsquigarrow & \mathbf{L}_{-\mathbf{t},R_0} \rightarrow \mathbf{L}_{-\mathbf{n},R_0} \rightarrow \mathbf{L}_{\mathbf{n},R_0} \rightarrow \mathbf{L}_{\mathbf{t},R_0} \\ \downarrow \sim & & \downarrow \sim & & \downarrow \sim & & \downarrow \sim \\ \mathbb{D}(X_\Lambda)_R \rightarrow \mathbb{D}(\overline{X})_R \rightarrow \mathbb{D}(\overline{X}^\vee)_R \rightarrow \mathbb{D}(X_{\Lambda^\#})_R & \rightsquigarrow & \mathbb{D}(X_\Lambda)_{R_0} \rightarrow \mathbb{D}(\overline{X})_{R_0} \rightarrow \mathbb{D}(\overline{X}^\vee)_{R_0} \rightarrow \mathbb{D}(X_{\Lambda^\#})_{R_0}. & (4.4) \end{array}$$

By the (relative) Grothendieck-Messing theorem, lifting the diagram (4.3) is equivalent to lifting the Hodge filtration:

$$\begin{array}{ccccccc}
\mathbb{D}(X_\Lambda)_R & \rightarrow & \mathbb{D}(\overline{X})_R & \rightarrow & \mathbb{D}(\overline{X}^\vee)_R & \rightarrow & \mathbb{D}(X_{\Lambda^\sharp})_R \\
\uparrow & & \uparrow & & \uparrow & & \uparrow \\
\Pi\mathbb{D}(X_\Lambda)_R & \rightarrow & \mathcal{F}_X & \longrightarrow & \mathcal{F}_{X^\vee} & \rightarrow & \Pi\mathbb{D}(X_{\Lambda^\sharp})_R
\end{array}
\rightsquigarrow
\begin{array}{ccccccc}
\mathbb{D}(X_\Lambda)_{R_0} & \rightarrow & D(\overline{X}) & \rightarrow & D(\overline{X}^\vee) & \rightarrow & \mathbb{D}(X_{\Lambda^\sharp})_{R_0} \\
\uparrow & & \uparrow & & \uparrow & & \uparrow \\
\Pi\mathbb{D}(X_\Lambda)_{R_0} & \rightarrow & \text{Fil}(\overline{X}) & \rightarrow & \text{Fil}(\overline{X}^\vee) & \rightarrow & \Pi\mathbb{D}(X_{\Lambda^\sharp})_{R_0}
\end{array}.
\tag{4.5}$$

Combining (4.5) with (4.4), we reduce the problem to finding a lifting:

$$\begin{array}{ccccccc}
\mathbf{L}_{-t,R} & \rightarrow & \mathbf{L}_{-\mathfrak{h},R} & \rightarrow & \mathbf{L}_{\mathfrak{h},R} & \rightarrow & \mathbf{L}_{t,R} \\
\uparrow & & \uparrow & & \uparrow & & \uparrow \\
\Pi\mathbf{L}_{-t,R} & \rightarrow & \widetilde{\mathcal{F}}_{-\mathfrak{h}} & \longrightarrow & \widetilde{\mathcal{F}}_{\mathfrak{h}} & \rightarrow & \Pi\mathbf{L}_{t,R}
\end{array}
\rightsquigarrow
\begin{array}{ccccccc}
\mathbf{L}_{-t,R_0} & \rightarrow & \mathbf{L}_{-\mathfrak{h},R_0} & \rightarrow & \mathbf{L}_{\mathfrak{h},R_0} & \rightarrow & \mathbf{L}_{t,R_0} \\
\uparrow & & \uparrow & & \uparrow & & \uparrow \\
\Pi\mathbf{L}_{-t,R_0} & \rightarrow & \mathcal{F}_{-\mathfrak{h}} & \longrightarrow & \mathcal{F}_{\mathfrak{h}} & \rightarrow & \Pi\mathbf{L}_{t,R_0}
\end{array}.
\tag{4.6}$$

Such a lifting can be obtained from the map $x \rightsquigarrow \bar{x}$ by compatibility. Moreover, the uniqueness of the lifting f follows directly from the construction. Furthermore, we observe that there are no additional constraints on the isomorphism (4.4), which implies that the map π has relative dimension $\dim \mathcal{G}_{\mathbb{F}}$. \square

Corollary 4.8. *The moduli functors $\mathcal{Z}(\Lambda)$ and $\mathcal{Y}(\Lambda^\sharp)$ are reduced.*

Proof. Consider the local model diagram (4.2). Since the morphisms φ and π are smooth of equal dimension, each point in the BT-strata is étale locally isomorphic to a point in the corresponding stratum of the local model. The reducedness then follows from Theorem 4.3. \square

Remark 4.9. (1) The local model diagram between $\mathcal{Z}(\Lambda_1) \cap \mathcal{Y}(\Lambda_2^\sharp)$ and $\mathcal{M}_{n,\mathbb{F}}^{[h]}(t_1, t_2)$ can be constructed and proved using analogous arguments.

(2) When F/F_0 is unramified, the local model diagram can be constructed in the same manner, and the proofs follow the same line of reasoning.

4.5. Local chart computations. The strata local model of BT-strata provides a powerful tool for studying the geometry of BT-strata without reference to Deligne-Lusztig varieties. In this subsection, we compute the local charts of the strata local models and establish Theorem 2.3.

4.5.1. Strata local model of \mathcal{Z} -strata. We begin by computing the strata local model associated to $\mathcal{Z}(\Lambda_1)$. By an unramified base change, we can reduce to the case where the hermitian form is split. Since the strata local model is equivariant under the action of the loop group, it suffices to compute an affine chart at the worst point. In this case, the strata local model parameterizes lattice chains:

$$\begin{array}{ccccccc}
\mathbf{L}_{-t_1,R} & \xrightarrow{\lambda_1} & \mathbf{L}_{-\mathfrak{h},R} & \xrightarrow{\lambda} & \mathbf{L}_{\mathfrak{h},R} & \xrightarrow{\lambda_2} & \mathbf{L}_{t_1,R} \\
\uparrow & & \uparrow & & \uparrow & & \uparrow \\
\Pi\mathbf{L}_{-t_1,R} & \longrightarrow & \mathcal{F}_{-\mathfrak{h}} & \longrightarrow & \mathcal{F}_{\mathfrak{h}} & \longrightarrow & \Pi\mathbf{L}_{t_1,R}
\end{array}$$

We select the same affine charts U as those given in [Luo24, §3.1]:

$$\mathcal{F}_{\mathfrak{h}} = \text{colspan} \begin{pmatrix} & h/2 & n-h & h/2 \\ D & 0 & C & \\ F & X_4 & E & \\ B & 0 & A & \\ I & & & \\ & I & & \\ & & I & \end{pmatrix} \begin{matrix} h/2 \\ n-h \\ h/2 \end{matrix}, \quad \mathcal{F}_{-\mathfrak{h}} = \text{colspan} \begin{pmatrix} h/2 & n-h & h/2 \\ -D & E^{\text{ad}} & -C \\ 0 & X_4 & 0 \\ -B & -F^{\text{ad}} & -A \\ I & & \\ & I & \\ & & I \end{pmatrix} \begin{matrix} h/2 \\ n-h \\ h/2 \end{matrix}. \quad (4.7)$$

For notational convenience, we denote by \mathcal{F}_{\bullet} the matrix whose columns span \mathcal{F}_{\bullet} . Let $X^{\text{ad}} = HX^tH$, where H is the antidiagonal identity matrix. The transition map λ^{\vee} factors as the composition:

$$\lambda^{\vee} : \mathbf{L}_{\mathfrak{h},R} \longrightarrow \mathbf{L}_{t_1,R} \longrightarrow \mathbf{L}_{n-t_1,R} \longrightarrow \mathbf{L}_{n-\mathfrak{h},R}. \quad (4.8)$$

We have $\lambda^{\vee}(\mathcal{F}_{\mathfrak{h}}) = \text{III}_{n-\mathfrak{h},R}$. By [Luo24, Thm. 6.3.2], we obtain $A = B = C = D = 0$. The coordinates can be further refined as follows:

$$\mathcal{F}_{\mathfrak{h}} = \begin{pmatrix} & h/2 & \frac{t_1-h}{2} & n-t_1 & \frac{t_1-h}{2} & h/2 \\ 0 & 0 & 0 & 0 & 0 \\ f_1 & Z_{11} & Z_{12} & Z_{13} & e_1 \\ f_2 & Z_{21} & Z_{22} & Z_{23} & e_2 \\ f_3 & Z_{31} & Z_{32} & Z_{33} & e_3 \\ 0 & 0 & 0 & 0 & 0 \\ I & & & & \\ & I & & & \\ & & I & & \\ & & & I & \\ & & & & I \end{pmatrix} \begin{matrix} h/2 \\ \frac{t_1-h}{2} \\ n-t_1 \\ \frac{t_1-h}{2} \\ h/2 \end{matrix}, \quad \mathcal{F}_{-\mathfrak{h}} = \begin{pmatrix} h/2 & \frac{t_1-h}{2} & n-t_1 & \frac{t_1-h}{2} & h/2 \\ 0 & e_3^{\text{ad}} & e_2^{\text{ad}} & e_1^{\text{ad}} & 0 \\ 0 & Z_{11} & Z_{12} & Z_{13} & 0 \\ 0 & Z_{21} & Z_{22} & Z_{23} & 0 \\ 0 & Z_{31} & Z_{32} & Z_{33} & 0 \\ 0 & -f_3^{\text{ad}} & -f_2^{\text{ad}} & -f_1^{\text{ad}} & 0 \\ I & & & & \\ & I & & & \\ & & I & & \\ & & & I & \\ & & & & I \end{pmatrix} \begin{matrix} h/2 \\ \frac{t_1-h}{2} \\ n-t_1 \\ \frac{t_1-h}{2} \\ h/2 \end{matrix}. \quad (4.9)$$

With respect to this partition, we can express the transition matrices as:

$$\lambda_1 = \begin{pmatrix} I_{n-t_1} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & I_h & 0 & 0 & 0 \\ 0 & 0 & 0 & I_{n-t_1} & 0 & 0 \\ 0 & I_{t_1-\mathfrak{h}} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & I_{\mathfrak{h}} \end{pmatrix}, \quad \lambda_2 = \begin{pmatrix} I_{\mathfrak{h}} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & I_{n-t_1} & 0 & 0 & 0 \\ 0 & 0 & 0 & I_h & 0 & 0 \\ 0 & I_{t_1-\mathfrak{h}} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & I_{n-t_1} \end{pmatrix}.$$

Therefore, $\lambda_1(\text{III}_{-t_1}) \subset \mathcal{F}_{-\mathfrak{h}}$ implies that

$$\begin{aligned} Z_{11} = 0, Z_{12} = 0, Z_{21} = 0, Z_{22} = 0, Z_{31} = 0, Z_{32} = 0, \\ e_3^{\text{ad}} = 0, e_2^{\text{ad}} = 0, -f_3^{\text{ad}} = 0, -f_2^{\text{ad}} = 0. \end{aligned}$$

The inclusion $\lambda_2(\mathcal{F}_{\mathbf{k}}) \subset \text{III}_{\mathbf{t}_1}$ implies that

$$\begin{aligned} Z_{21} = 0, Z_{22} = 0, Z_{23} = 0, Z_{31} = 0, Z_{32} = 0, Z_{33} = 0, \\ f_2 = 0, f_3 = 0, e_2 = 0, e_3 = 0. \end{aligned}$$

Therefore, the only nonzero matrices are Z_{13}, f_1 and e_1 . By the local model axioms, we have $X_4 = X_4^{\text{ad}}, \text{Tr}(X_4) = 0$, and $\wedge^2 X = 0$ ([Luo24, Prop. 4.1.1]). After simplification, the affine coordinate ring of the open affine chart U is isomorphic to:

$$\frac{\mathbb{F}[Z_{13}, e_1, f_1]}{Z_{13} - Z_{13}^{\text{ad}}, \wedge^2(e_1, f_1, Z_{13})}.$$

This ring defines a symmetric determinantal variety corresponding to a $(\mathbf{t}_1 + \mathbf{k}) \times (\mathbf{t}_1 - \mathbf{k})$ matrix. By [Luo24, Thm. 4.2.2], this variety is normal and Cohen-Macaulay of dimension $\mathbf{t}_1 + \mathbf{k}$. Moreover, it is smooth when $\mathbf{t}_1 - \mathbf{k} = 1$ and singular otherwise.

Recall the following result of Conca:

Proposition 4.10 ([Con94]). *Let $X = (Y, Z)$ be a matrix of indeterminates, where Y is an $m \times m$ matrix and Z is an $m \times (n - m)$ matrix. The ring*

$$\frac{k[X]}{\wedge^2 X, Y - Y^t}$$

is Gorenstein if and only if $2m = n + 2$.

As a consequence, our stratum local model is Gorenstein if and only if $2(\mathbf{t}_1 - \mathbf{k}) = (\mathbf{t}_1 + \mathbf{k}) + 2$, i.e. $\mathbf{t}_1 = 3\mathbf{k} + 2$.

4.5.2. *Strata local model of \mathcal{Y} -strata.* We exclude the case where n is even and $h = n$, as in this situation the local model does not have a worst point. The strata local model then parameterizes lattice chains

$$\begin{array}{ccccccc} \mathbf{L}_{\mathbf{t}_2, R} & \xrightarrow{\lambda_2} & \mathbf{L}_{\mathbf{k}, R} & \xrightarrow{\lambda^\vee} & \mathbf{L}_{n-\mathbf{k}, R} & \xrightarrow{\lambda_1} & \mathbf{L}_{n-\mathbf{t}_2, R} \\ \uparrow & & \uparrow & & \uparrow & & \uparrow \\ \text{III}_{\mathbf{t}_2, R} & \longrightarrow & \mathcal{F}_{\mathbf{k}} & \longrightarrow & \mathcal{F}_{n-\mathbf{k}} & \longrightarrow & \text{III}_{n-\mathbf{t}_2, R} \end{array}$$

We choose the same affine charts as in the previous case. In particular, the formulas for $\mathcal{F}_{\mathbf{k}}$ and $\mathcal{F}_{n-\mathbf{k}}$ remain as in equation (4.7) (note that $\mathcal{F}_{-\mathbf{k}}$ has the same form as $\mathcal{F}_{\mathbf{k}}$ by the local model axiom).

For reasons similar to those in equation (4.8), the transition map λ carries $\mathcal{F}_{-\mathfrak{n}}$ to $\Pi\Lambda_{\mathfrak{n}}$. By [Luo24, Thm. 6.3.2], this implies $X_4 = 0$. We can further refine the coordinates as follows

$$\mathcal{F}_{\mathfrak{n}} = \begin{pmatrix} \frac{t_2}{2} & \frac{h-t_2}{2} & n-h & \frac{h-t_2}{2} & \frac{t_2}{2} \\ D_1 & D_2 & 0 & C_1 & C_2 \\ D_3 & D_4 & 0 & C_3 & C_4 \\ F_1 & F_2 & 0 & E_1 & E_2 \\ B_1 & B_2 & 0 & A_1 & A_2 \\ B_3 & B_4 & 0 & A_3 & A_4 \\ I & & & & \\ & I & & & \\ & & I & & \\ & & & I & \\ & & & & I \end{pmatrix} \begin{matrix} t_2/2 \\ \frac{h-t_2}{2} \\ n-h \\ \frac{h-t_2}{2} \\ t_2/2 \end{matrix}, \quad \mathcal{F}_{n-\mathfrak{n}} = \begin{pmatrix} t_2/2 & \frac{h-t_2}{2} & n-h & \frac{h-t_2}{2} & t_2/2 \\ -D_1 & -D_2 & E_2^{\text{ad}} & -C_1 & -C_2 \\ -D_3 & -D_4 & E_1^{\text{ad}} & -C_3 & -C_4 \\ 0 & 0 & 0 & 0 & 0 \\ -B_1 & -B_2 & -F_2^{\text{ad}} & -A_1 & -A_2 \\ -B_3 & -B_4 & -F_1^{\text{ad}} & -A_3 & -A_4 \\ I & & & & \\ & I & & & \\ & & I & & \\ & & & I & \\ & & & & I \end{pmatrix} \begin{matrix} t_2/2 \\ \frac{h-t_2}{2} \\ n-h \\ \frac{h-t_2}{2} \\ t_2/2 \end{matrix}. \quad (4.10)$$

With respect to this partition, the transition matrices can be represented as

$$\lambda_1 = \begin{pmatrix} I_{n-\mathfrak{n}} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & I_{t_2} & 0 & 0 & 0 \\ 0 & 0 & 0 & I_{n-h} & 0 & 0 \\ 0 & I_{\mathfrak{n}-t_2} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & I_{t_2} \end{pmatrix}, \quad \lambda_2 = \begin{pmatrix} I_{t_2} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & I_{n-\mathfrak{n}} & 0 & 0 & 0 \\ 0 & 0 & 0 & I_{t_2} & 0 & 0 \\ 0 & I_{\mathfrak{n}-t_2} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & I_{n-\mathfrak{n}} \end{pmatrix}$$

Now $\lambda_1(\mathcal{F}_{n-\mathfrak{n}}) \subset \Pi\Lambda_{n-t_2}$ implies that

$$C = 0, D = 0, E = 0, A_3 = 0, A_4 = 0, B_3 = 0, B_4 = 0, -F_1^{\text{ad}} = 0.$$

The inclusion $\lambda_2(\Pi\Lambda_{\ell}) \subset \mathcal{F}_h$ implies that

$$A = 0, C = 0, E = 0, B_1 = 0, B_3 = 0, D_1 = 0, D_3 = 0, F_1 = 0.$$

The local model relations [Luo24, Prop. 4.1.5] imply the following relations

$$\bigwedge^2 (B_2, F_2) = 0, B = -\frac{1}{2}F^{\text{ad}}F.$$

By the proof of Theorem 4.1.9 in loc.cit, we can further simplify the affine ring into

$$\frac{\mathbb{F}[F_2]}{\bigwedge^2 F_2}.$$

This defines a determinantal variety corresponding to a $(\mathfrak{n} - t_2) \times (n - 2\mathfrak{n})$ matrix. The variety is normal and Cohen-Macaulay of dimension $n - \mathfrak{n} - t_2 - 1$. In the almost π -modular case, where $n - 2\mathfrak{n} = 1$, the matrix F_2 reduces to a column vector, and the affine ring is smooth. Similarly, when $h - t_2 = 1$, F_2 becomes a row vector, and the variety is again smooth. In all other cases, the variety is singular. Moreover, it is Gorenstein if and only if $(\mathfrak{n} - t_2) = (n - 2\mathfrak{n})$, i.e. $t_2 = 3\mathfrak{n} - n$.

4.5.3. *Intersection of \mathcal{Y} -strata and \mathcal{Z} -strata.* We consider the intersection $\mathcal{Z}(\Lambda_1) \cap \mathcal{Y}(\Lambda_2^\sharp)$ where $t(\Lambda_1) = t_1 = 2\mathbf{t}_1$ and $t(\Lambda_2) = t_2 = 2\mathbf{t}_2$. This intersection is related to the strata local model $\overline{\mathcal{M}}_n^{[h]}(t_1, t_2)$. Maintaining the notation from equation (4.7), we further decompose:

$$f_1 = (f_{11}, f_{12}), \quad f_{11} : (\mathbf{t}_1 - \mathbf{h}) \times \mathbf{h}, \quad f_{12} : (\mathbf{t}_1 - \mathbf{h}) \times (\mathbf{h} - \mathbf{t}_2).$$

Combining the computations from the previous two subsections, we find that the affine chart of the stratum local model is isomorphic to the spectrum of the ring:

$$\frac{\mathbb{F}[f_{12}]}{\wedge^2 f_{12}}.$$

This ring defines a variety that is normal and Cohen-Macaulay of dimension $\mathbf{t}_1 - \mathbf{t}_2 - 1$. The variety is smooth if and only if $\mathbf{t}_1 - \mathbf{h} = 1$ or $\mathbf{h} - \mathbf{t}_2 = 1$. In all other cases, it is singular and is Gorenstein if and only if $\mathbf{t}_1 - \mathbf{h} = \mathbf{h} - \mathbf{t}_2$.

4.5.4. *Strata local model of \mathcal{Y} -strata: π -modular case.* We now compute the affine charts of the π -modular local model. In this case, the local model does not contain a worst point, cf. [PR09, Rem. 5.3]. We note that the affine chart chosen in their work does not apply to our situation. The strata local model in this case parameterizes lattice chains

$$\begin{array}{ccccc} \mathbf{L}_{\mathbf{t}_2, R} & \xrightarrow{\lambda_2} & \mathbf{L}_{m, R} & \xrightarrow{\lambda_1} & \mathbf{L}_{n-\mathbf{t}_2, R} \\ \uparrow & & \uparrow & & \uparrow \\ \Pi\mathbf{L}_{\mathbf{t}_2, R} & \longrightarrow & \mathcal{F}_m & \longrightarrow & \Pi\mathbf{L}_{n-\mathbf{t}_2, R} \end{array}$$

We choose the standard basis as given in [Luo24, (3.1.1)]. Over the special fiber, the two transition maps are represented by the following matrices:

$$\lambda_2 = \begin{pmatrix} I_{\mathbf{t}_2} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & I_m & 0 & 0 & 0 \\ 0 & 0 & 0 & I_{\mathbf{t}_2} & 0 & 0 \\ 0 & I_{m-\mathbf{t}_2} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & I_m \end{pmatrix}, \quad \lambda_1 = \begin{pmatrix} I_m & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & I_{\mathbf{t}_2} & 0 & 0 & 0 \\ 0 & 0 & 0 & I_m & 0 & 0 \\ 0 & I_{m-\mathbf{t}_2} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & I_{\mathbf{t}_2} \end{pmatrix}.$$

Consider the \mathbb{F} -span of the following vectors

$$e_1, \dots, e_{m-1}, e_{m+1}, \pi e_{m+1}, \dots, \pi e_n, \quad (4.11)$$

It defines a \mathbb{F} -point of $\overline{\mathcal{M}}_n^{[n]}$, as can be checked using the spin condition in [RSZ18, Def. 3.9]. We can embed the special fiber $\overline{\mathcal{M}}_n^{[n]}$ into the Grassmannian $\mathrm{Gr}(n, \Lambda_{m, \mathbb{F}})$. Choose the affine chart U of

the point (4.11) in the Grassmannian as

$$\mathcal{F} = \begin{pmatrix} t_2/2 & \frac{n-t_2-2}{2} & 1 & 1 & \frac{n-t_2-2}{2} & t_2/2 \\ X_{11} & X_{12} & X_{13} & X_{14} & X_{15} & X_{16} \\ X_{21} & X_{22} & X_{23} & X_{24} & X_{25} & X_{26} \\ Y_1 & Y_2 & Y_3 & Y_4 & Y_5 & Y_6 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ X_{31} & X_{32} & X_{33} & X_{34} & X_{35} & X_{36} \\ X_{41} & X_{42} & X_{43} & X_{44} & X_{45} & X_{46} \\ I_{t_2} & & & & & \\ & I_{m-t_2-1} & & & & \\ Z_1 & Z_2 & Z_3 & Z_4 & Z_5 & Z_6 \\ & & & 1 & & \\ & & & & I_{m-t_2-1} & \\ & & & & & I_{t_2} \end{pmatrix}.$$

Now two subspaces $\lambda_2(\mathbf{III}_{\ell,R})$ and $\lambda_1(\mathcal{F})$ are spanned by the following matrices respectively:

$$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ I_\ell & 0 & 0 & 0 & 0 & 0 \\ Z_1 & Z_2 & Z_3 & Z_4 & Z_5 & Z_6 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ X_{31} & X_{32} & X_{33} & X_{34} & X_{35} & X_{36} \\ 0 & 0 & 0 & 0 & 0 & I_\ell \end{pmatrix}, \begin{pmatrix} X_{11} & X_{12} & X_{13} & X_{14} & X_{15} & X_{16} \\ X_{21} & X_{22} & X_{23} & X_{24} & X_{25} & X_{26} \\ Y_1 & Y_2 & Y_3 & Y_4 & Y_5 & Y_6 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ X_{41} & X_{42} & X_{43} & X_{44} & X_{45} & X_{46} \\ I_{t_2} & 0 & 0 & 0 & 0 & 0 \\ 0 & I_{m-t_2-1} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & I_{m-t_2-1} & 0 \\ 0 & 0 & 0 & 0 & 0 & I_{t_2} \end{pmatrix}.$$

Now $\lambda_2(\mathbf{III}_{t_2}) \subset \mathcal{F}_m$ implies

$$\begin{aligned} X_{11} &= 0, X_{21} = 0, X_{31} = 0, X_{41} = 0, Y_1 = 0, Z_1 = 0, \\ X_{14} &= 0, X_{24} = 0, X_{34} = 0, X_{44} = 0, Y_4 = 0, Z_4 = 0, \\ X_{15} &= 0, X_{25} = 0, X_{35} = 0, X_{45} = 0, Y_5 = 0, Z_5 = 0, \\ X_{16} &= 0, X_{26} = 0, X_{36} = 0, X_{46} = 0, Y_6 = 0, Z_6 = 0. \end{aligned}$$

The inclusion $\lambda_1(\mathcal{F}_m) \subset \text{III}_{n-\mathbf{t}_2}$ implies

$$\begin{aligned} X_{11} &= 0, X_{12} = 0, X_{13} = 0, X_{14} = 0, X_{15} = 0, X_{16} = 0, \\ X_{21} &= 0, X_{22} = 0, X_{23} = 0, X_{24} = 0, X_{25} = 0, X_{26} = 0, \\ Y_1 &= 0, Y_2 = 0, Y_3 = 0, Y_4 = 0, Y_5 = 0, Y_6 = 0, \\ X_{41} &= 0, X_{42} = 0, X_{43} = 0, X_{44} = 0, X_{45} = 0, X_{46} = 0. \end{aligned}$$

By simplification, the subspace \mathcal{F}_m is spanned by the matrix of the form

$$\mathcal{F}_k = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & X_{32} & X_{33} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ I_{\mathbf{t}_2} & & & & & \\ & I_{m-\mathbf{t}_2-1} & & & & \\ 0 & Z_2 & Z_3 & 0 & 0 & 0 \\ & & & 1 & & \\ & & & & I_{m-\mathbf{t}_2-1} & \\ & & & & & I_{\mathbf{t}_2} \end{pmatrix}.$$

Now we apply the remaining local model axioms. Due to our choice of affine chart, the spin condition is automatically satisfied. The Kottwitz condition holds since the upper half of the matrix is nilpotent. The wedge condition implies that $X_{32} = 0$. It remains to interpret the isotropic condition. With respect to our chosen standard basis, the symmetric form $(,)$ is represented by the matrix

$$M = \begin{pmatrix} & & -H \\ & H & \\ -H & H & \end{pmatrix},$$

where H is the $m \times m$ antidiagonal unit matrix. The isotropic condition implies that $\mathcal{F}_m^t M \mathcal{F}_m = 0$, which boils down to

$$A + A^t = 0, \quad A = \begin{pmatrix} I_{\mathbf{t}_2} & & \\ & I_{m-\mathbf{t}_2-1} & Z_2^t \\ & & Z_3 \end{pmatrix} H \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & X_{33} \\ 0 & 0 & 0 \end{pmatrix}.$$

This implies that $Z_3 = 0$ and $HX_{33} + Z_2^t = 0$. Therefore, the affine chart of the strata local model is isomorphic to:

$$\text{Spec } \mathbb{F}[X_{33}].$$

This defines a smooth variety of dimension $m - \mathbf{t}_2 - 1$. Since $\overline{\mathcal{M}}_n^{[n]}(t)$ is contained in the preimage of the worst point $(\text{III}_{\mathbf{t}_2, \mathbb{F}} \subset \mathbf{L}_{\mathbf{t}_2, \mathbb{F}}, \text{III}_{n-\mathbf{t}_2, \mathbb{F}} \subset \mathbf{L}_{n-\mathbf{t}_2, \mathbb{F}})$, the affine chart U intersects all irreducible

components of the strata local model. Therefore, the strata local model is smooth and irreducible of dimension $m - \mathfrak{t}_2 - 1$.

5. BRUHAT-TITS STRATIFICATION

In this section, we begin by proving a crucial lemma, then proceed to define the Bruhat-Tits stratification of the reduced locus $\mathcal{N}_{n,\varepsilon,\text{red}}^{[h]}$. In the final subsection, we establish its relationship with the Kottwitz-Rapoport strata.

5.1. The crucial lemma. Let κ be any perfect field over \mathbb{F} . We denote by $M = M(X)$ the Dieudonné module of $(X, \iota_X, \lambda_X, \rho_X) \in \mathcal{N}_{n,\varepsilon}^{[h]}(\kappa)$. By Proposition 3.5, we have

$$\pi M^\sharp \stackrel{n-h}{\subseteq} M \stackrel{h}{\subseteq} M^\sharp, \quad \text{and} \quad M \stackrel{\leq 1}{\subseteq} M + \tau(M).$$

The following discussion is inspired by and is essentially the same as [HZ25, §4.3] once we replace the Φ in [HZ25] by τ . See also [KR12, §3].

We denote by $T_i(M)$ the summation $M + \tau(M) + \cdots + \tau^i(M)$. Recall that we have $(M^\sharp)^\sharp = M$ and $\tau(M^\sharp) = (\tau(M))^\sharp$. **From now on, we always assume c (resp. d) is the smallest non-negative integer such that $T_c(M)$ (resp. $T_d(M^\sharp)$) is τ -invariant.** Note that c and d exists by [RZ96, Prop. 2.17]. It is not hard to show that

$$M \stackrel{1}{\subset} T_1(M) \stackrel{1}{\subset} T_2(M) \cdots \stackrel{1}{\subset} T_c(M), \quad \text{and} \quad M^\sharp \stackrel{1}{\subset} T_1(M^\sharp) \stackrel{1}{\subset} T_2(M^\sharp) \cdots \stackrel{1}{\subset} T_d(M^\sharp).$$

Lemma 5.1. $[M + \tau(M) : M] = 1$ implies that $[M^\sharp + \tau(M^\sharp) : M^\sharp] = 1$.

Proof. By taking the dual, we have $[M^\sharp + \tau(M^\sharp) : M^\sharp] = [M : M \cap \tau(M)]$. Moreover, $[M : M \cap \tau(M)] = [M + \tau(M) : M] = 1$. \square

Lemma 5.2 ([HZ25, Lem. 4.7]). *We have*

$$(1) \quad T_c(M) \subseteq \bigcap_{i \in \mathbb{Z}_{\geq 0}} \tau^i(\pi^{-1}M) \quad \text{and} \quad T_d(M^\sharp) \subseteq \bigcap_{i \in \mathbb{Z}_{\geq 0}} \tau^i(\pi^{-1}M^\sharp).$$

(2) *If $\tau(M) \subseteq M^\sharp$ and $c \leq d$, then*

$$T_c(M) \subseteq \bigcap_{i \in \mathbb{Z}_{\geq 0}} \tau^i(M^\sharp).$$

(3) *If $\tau(\pi M^\sharp) \subseteq M$ and $d \leq c$, then*

$$T_d(\pi M^\sharp) \subseteq \bigcap_{i \in \mathbb{Z}_{\geq 0}} \tau^i(M).$$

Proof. For (1), we only prove the statement for $T_c(M)$ since the proof for $T_d(M^\sharp)$ is literally the same. First, note that

$$\tau(M) \subseteq \pi^{-1}M.$$

Hence

$$T_c(M) \subseteq \pi^{-1}T_{c-1}(M).$$

Since $T_c(M)$ is τ -invariant, we have

$$T_c(M) \subseteq \pi^{-1}\tau(T_{c-1}(M)) \cap \pi^{-1}T_{c-1}(M) = \pi^{-1}\tau(T_{c-2}(M)).$$

Here, for the last equality, we use the facts that:

- (i) $\tau(T_{c-1}(M)) \neq T_{c-1}(M)$,
- (ii) $\tau(T_{c-2}(M)) \subseteq \tau(T_{c-1}(M)) \cap T_{c-1}(M)$, and
- (iii) $[T_{c-1}(M) : T_{c-2}(M)] = [\tau(T_{c-1}(M)) : \tau(T_{c-2}(M))] = 1$.

Since $T_c(M)$ is τ -invariant, we get

$$T_c(M) \subseteq \pi^{-1}T_{c-2}(M).$$

Inductively, we have $T_c(M) \subseteq \pi^{-1}M$. Since $T_c(M)$ is τ -invariant, we have

$$T_c(M) \subseteq \pi^{-1} \bigcap_{i \in \mathbb{Z}_{\geq 0}} \tau^i(M) \subseteq \pi^{-1} \bigcap_{0 \leq i \leq f} \tau^i(M)$$

for any $f \in \mathbb{Z}_{\geq 0}$.

Part (2) and (3) are essentially the same as (1), once we observe that the assumption $c \leq d$ implies that $\tau(T_{c-1}(M^\sharp)) \neq T_{c-1}(M^\sharp)$ and $\tau(T_{c-1}(M^\sharp)) \cap T_{c-1}(M^\sharp) = \tau(T_{c-2}(M^\sharp))$ hold. Similarly, $d \leq c$ implies that $\tau(T_{d-1}(M)) \cap T_{d-1}(M) = \tau(T_{d-2}(M))$ \square

Proposition 5.3 ([HZ25, Prop. 4.8]). *Assume $X \in \mathcal{N}_{n,\varepsilon}^{[h]}(\kappa)$ and $M = M(X)$. Then one of the following will happen:*

(Case \mathcal{Y}): $\pi(T_c(M))^\sharp \subseteq \pi M^\sharp \subseteq M \subseteq T_c(M) \subseteq (T_c(M))^\sharp \subseteq M^\sharp$. In particular, $T_c(M)$ is a vertex lattice of type $t \leq h$.

(Case \mathcal{Z}): $\pi M^\sharp \subseteq \pi T_d(M^\sharp) \subseteq T_d(M^\sharp)^\sharp \subseteq M \subseteq M^\sharp \subseteq T_d(M^\sharp)$. In particular, $T_d(M^\sharp)^\sharp$ is a vertex lattice of type $t \geq h$.

More precisely, we have:

- (1) If $\tau(\pi M^\sharp) \not\subseteq M$, then M satisfies (Case \mathcal{Y}).
- (2) If $\tau(M) \not\subseteq M^\sharp$, then M satisfies (Case \mathcal{Z}).
- (3) If $\tau(\pi M^\sharp) \subseteq M$ and $\tau(M) \subseteq M^\sharp$, then M satisfies (Case \mathcal{Y}) if $c \leq d$.
- (4) If $\tau(\pi M^\sharp) \subseteq M$ and $\tau(M) \subseteq M^\sharp$, then M satisfies (Case \mathcal{Z}) if $d \leq c$.

Proof. (1) Assume $\tau(M^\sharp) \not\subseteq \pi^{-1}M$. Since $M \subseteq T_c(M)$, it suffices to show $T_c(M) \subseteq T_c(M)^\sharp$. Since $M \stackrel{\leq 1}{\subset} M + \tau(M)$, $\tau(\pi M^\sharp) \not\subseteq M$ and $\tau(\pi M^\sharp) \subset \tau(M)$, we have $M + \tau(M) = M + \tau(\pi M^\sharp)$. In fact, an inductive argument on $T_i(M)$ shows that

$$T_c(M) = M + \tau(\pi M^\sharp) + \cdots + \tau^c(\pi M^\sharp). \quad (5.1)$$

Equivalently,

$$T_c(M)^\sharp = M^\sharp \cap \left(\bigcap_{1 \leq i \leq c} \tau^i(\pi^{-1}M) \right). \quad (5.2)$$

According to Lemma 5.2, we have

$$\begin{aligned} T_c(\pi M^\sharp) &\subseteq \bigcap_{0 \leq i \leq c} \tau^i(M^\sharp) \subseteq M^\sharp \cap \left(\bigcap_{1 \leq i \leq c} \tau^i(\pi^{-1}M) \right), \\ M &\subseteq M^\sharp \cap T_c(M) \subseteq M^\sharp \cap \left(\bigcap_{1 \leq i \leq c} \tau^i(\pi^{-1}M) \right). \end{aligned} \quad (5.3)$$

Therefore,

$$T_c(M) \stackrel{(5.1)}{\subseteq} M + T_c(\pi M^\sharp) \stackrel{(5.3)}{\subseteq} M^\sharp \cap \left(\bigcap_{1 \leq i \leq c} \tau^i(\pi^{-1}M) \right) \stackrel{(5.2)}{=} T_c(M)^\sharp.$$

(2) Now we assume $\tau(M) \not\subseteq M^\sharp$. By construction, we have $T_d(M^\sharp)^\sharp \subseteq M \subseteq M^\sharp \subseteq T_d(M^\sharp)$. Therefore, we only need to show $\pi T_d(M^\sharp) \subseteq T_d(M^\sharp)^\sharp$. Since $M^\sharp \stackrel{\leq 1}{\subset} M^\sharp + \tau(M^\sharp)$ and $\tau(M) \not\subseteq M^\sharp$, we have $M^\sharp + \tau(M^\sharp) = M^\sharp + \tau(M)$. In fact, an inductive argument shows that

$$T_d(M^\sharp) = M^\sharp + \tau(M) + \cdots + \tau^d(M). \quad (5.4)$$

Equivalently,

$$T_d(M^\sharp)^\sharp = M \cap \left(\bigcap_{1 \leq i \leq d} \tau^i(M^\sharp) \right). \quad (5.5)$$

According to Lemma 5.2, we have

$$\begin{aligned} T_d(\pi M) &\subseteq \bigcap_{0 \leq i \leq d} \tau^i(M) \subseteq M \cap \left(\bigcap_{1 \leq i \leq d} \tau^i(M^\sharp) \right), \\ \pi M^\sharp &\subseteq M \cap T_d(\pi M^\sharp) \subseteq M \cap \left(\bigcap_{1 \leq i \leq d} \tau^i(M^\sharp) \right). \end{aligned} \quad (5.6)$$

Therefore,

$$\pi T_d(M^\sharp) \stackrel{(5.4)}{\subseteq} \pi M^\sharp + T_d(\pi M) \stackrel{(5.6)}{\subseteq} M \cap \left(\bigcap_{1 \leq i \leq d} \tau^i(M^\sharp) \right) \stackrel{(5.5)}{=} T_d(M^\sharp)^\sharp.$$

(3) Assuming $\tau(M) \subseteq M^\sharp$ and $c \leq d$, we will show that M satisfies (Case \mathcal{Y}) (a stronger statement than (3) in the assertion). Note that $M + \tau(M) \subseteq M^\sharp$. Inductively, we see that $T_c(M) \subseteq T_{c-1}(M^\sharp)$. Since $T_c(M)$ is τ -invariant, we have $T_c(M) \subseteq \bigcap_i \tau^i(M^\sharp) \subseteq \bigcap_{0 \leq i \leq c} \tau^i(M^\sharp) \subseteq (T_c(M))^\sharp$ by Lemma 5.2. Hence, we have

$$M \subseteq T_c(M) \subseteq (T_c(M))^\sharp \subseteq M^\sharp.$$

Since $\pi M^\sharp \subseteq M \subseteq M^\sharp$, we have

$$\pi(T_c(M))^\sharp \subseteq \pi M^\sharp \subseteq M \subseteq T_c(M) \subseteq (T_c(M))^\sharp \subseteq M^\sharp.$$

The proof of (4) follows from the same argument as the proof of (3). \square

5.2. Bruhat-Tits stratification. In this subsection, we define the Bruhat-Tits stratification. Let $\Lambda \subset V$ be a vertex lattice of type t . Since t is an even integer, we write $t = 2\mathbf{t}$. Recall from Section 1.2 that for any lattice Λ , we let $\check{\Lambda} := \Lambda \otimes_{O_F} O_{\check{F}}$. We define two important quotient spaces: $V_{\Lambda} := \Lambda^{\sharp}/\Lambda$ equipped with an alternating form and $V_{\Lambda^{\sharp}} := \Lambda/\pi\Lambda^{\sharp}$ equipped with a symmetric form. By extending scalars to \mathbb{F} , we obtain $\Omega_{\Lambda} := V_{\Lambda} \otimes_{\mathbb{F}_q} \mathbb{F}$ and $\Omega_{\Lambda^{\sharp}} := V_{\Lambda^{\sharp}} \otimes_{\mathbb{F}_q} \mathbb{F}$. The alternating form on Ω_{Λ} and the symmetric form on $\Omega_{\Lambda^{\sharp}}$ are defined by extending the corresponding forms on V_{Λ} and $V_{\Lambda^{\sharp}}$ via scalar extension.

From Proposition 3.5, we have a lattice-theoretic description of $\mathcal{N}_{n,\varepsilon,\text{red}}^{[h]}(\kappa)$, where κ is a perfect field over \mathbb{F} . These descriptions naturally extend to give a lattice-theoretic characterization of Bruhat-Tits strata on κ -points:

Proposition 5.4. *Let κ be any perfect field over \mathbb{F} . The κ -points of the Bruhat-Tits strata can be described as follows:*

(1) *Assume Λ is a vertex lattice of type $t \geq h$. Then*

$$\mathcal{Z}(\Lambda)(\kappa) = \{(X, \iota_X, \lambda_X, \rho_X) \in \mathcal{N}_{n,\varepsilon}^{[h]}(\kappa) \mid \Lambda \otimes W_O(\kappa) \subseteq M(X) \subseteq M(X)^{\sharp} \subseteq \Lambda^{\sharp} \otimes W_O(\kappa)\}.$$

(2) *Assume Λ is a vertex lattice of type $t \leq h$. Then*

$$\mathcal{Y}(\Lambda^{\sharp})(\kappa) = \{(X, \iota_X, \lambda_X, \rho_X) \in \mathcal{N}_{n,\varepsilon}^{[h]}(\kappa) \mid \pi\Lambda^{\sharp} \otimes W_O(\kappa) \subseteq \pi M(X)^{\sharp} \subseteq M(X) \subseteq \Lambda \otimes W_O(\kappa)\}.$$

□

Theorem 5.5. *Recall that $\mathcal{L}_{\mathcal{Z}}$ (resp. $\mathcal{L}_{\mathcal{Y}}$) denotes the set of all vertex lattices in \mathbb{V} of type $\geq h$ (resp. $\leq h$). Let κ be any perfect field over \mathbb{F} , then we have following:*

(1) *The reduced locus $\mathcal{N}_{n,\varepsilon,\text{red}}^{[h]}$ is the union of closed subvarieties:*

$$\mathcal{N}_{n,\varepsilon}^{[h]}(\kappa) = \left(\bigcup_{\Lambda_1 \in \mathcal{L}_{\mathcal{Z}}} \mathcal{Z}(\Lambda_1)(\kappa) \right) \cup \left(\bigcup_{\Lambda_2 \in \mathcal{L}_{\mathcal{Y}}} \mathcal{Y}(\Lambda_2^{\sharp})(\kappa) \right),$$

Moreover, these strata satisfy the following inclusion relations:

(i) *For any Λ_1 and Λ'_1 in $\mathcal{L}_{\mathcal{Z}}$, $\mathcal{Z}(\Lambda_1)(\kappa) \subseteq \mathcal{Z}(\Lambda'_1)(\kappa)$ if and only if $\Lambda_1 \supseteq \Lambda'_1$,*

(ii) *For any Λ_2 and Λ'_2 in $\mathcal{L}_{\mathcal{Y}}$, $\mathcal{Y}(\Lambda_2^{\sharp})(\kappa) \subseteq \mathcal{Y}(\Lambda'^{\sharp}_2)(\kappa)$ if and only if $\Lambda_2 \subseteq \Lambda'_2$.*

(2) *For any $\Lambda_1 \in \mathcal{L}_{\mathcal{Z}}$ and $\Lambda_2 \in \mathcal{L}_{\mathcal{Y}}$, the intersection $\mathcal{Z}(\Lambda_1)(\kappa) \cap \mathcal{Y}(\Lambda_2^{\sharp})(\kappa)$ is non-empty if and only if $\Lambda_1 \subseteq \Lambda_2$.*

(3) *For $\Lambda \in \mathcal{L}_{\mathcal{Z}} \cap \mathcal{L}_{\mathcal{Y}}$ (i.e., Λ is a vertex lattice of type h), the set $\mathcal{Z}(\Lambda) = \mathcal{Y}(\Lambda^{\sharp})(\kappa)$ is a singleton, corresponding to a discrete point in the RZ space called the worst point.*

Proof. (1) The decomposition follows from the crucial lemma (Proposition 5.3), combined with the characterization of κ -points given in Propositions 3.5 and 5.4.

(2) By Proposition 5.4, a κ -point in $\mathcal{Y}(\Lambda_2^{\sharp})$ corresponds to a lattice M such that:

$$\pi\Lambda_2^{\sharp} \otimes W_O(\kappa) \subseteq \pi M^{\sharp} \subseteq M \subseteq \Lambda_2 \otimes W_O(\kappa) \subseteq \Lambda_2^{\sharp} \otimes W_O(\kappa) \subseteq M^{\sharp}.$$

If $\mathcal{Z}(\Lambda_1) \cap \mathcal{Y}(\Lambda_2^{\sharp}) \neq \emptyset$, then there exists a lattice $M \subset \mathbb{V} \otimes W_O(\kappa)$ of type h such that

$$\Lambda_1 \otimes W_O(\kappa) \subseteq M \subseteq \Lambda_2 \otimes W_O(\kappa).$$

This implies $\Lambda_1 \subseteq \Lambda_2$. Conversely, suppose $\Lambda_1 \subseteq \Lambda_2$, then we can find a vertex lattice of type h such that $\Lambda_1 \subseteq \Lambda \subseteq \Lambda_2$. Let $M := \Lambda \otimes W_O(\kappa) \subset \Lambda^\sharp \otimes W_O(\kappa)$. By Lemma 2.1, M stable under F, V and Π . Since $\Lambda \subset \mathbb{V}$, we have $\tau(M) = M$, hence $M \in \mathcal{Z}(\Lambda_1) \cap \mathcal{Y}(\Lambda_2^\sharp)(\kappa)$.

(3) For any vertex lattice $\Lambda \subset \mathbb{V}$ of type h , we have by definition $\mathcal{Z}(\Lambda)(\kappa) = \mathcal{Y}(\Lambda)(\kappa) = \{\Lambda \otimes W_O(\kappa)\}$ which is a single point (corresponds to the worst point in the local model). \square

5.3. Kottwitz-Rapoport strata. In this section, we examine the relationship between two fundamental types of strata: the (closed) BT strata and the (closed) Kottwitz-Rapoport (KR) strata. Our discussion relies on notation and concepts from local model theory, which is developed in Section 4.

Proposition 5.6. *Let $M \subset N \otimes W_O(\kappa)$ represents a point in $\mathcal{N}_{n,\varepsilon}^{[h]}(\kappa)$. Then one of the following inclusions must hold:*

$$\tau(M) \subseteq M^\sharp, \quad \text{or} \quad \tau(\Pi M^\sharp) \subseteq M.$$

We present two independent proofs of this result: the first proof uses the crucial lemma, the second proof uses local model theory.

First proof. Assume $(X, \iota_X, \lambda_X, \rho_X) \in \mathcal{Z}(\Lambda)(\kappa)$ corresponds to the lattices chain $\Lambda \otimes W_O(\kappa) \subseteq M \subseteq M^\sharp \subseteq \Lambda^\sharp \otimes W_O(\kappa)$, we have

$$\pi M^\sharp \subseteq \pi \Lambda^\sharp \otimes W_O(\kappa) \subseteq \Lambda \otimes W_O(\kappa) \subseteq M.$$

Then we have

$$\tau(\pi M^\sharp) \subseteq \tau(\Lambda \otimes W_O(\kappa)) = \Lambda \otimes W_O(\kappa) \subseteq M.$$

By definition, we have $(X, \iota_X, \lambda_X, \rho_X) \in \mathcal{Z}$. The same argument holds for \mathcal{Y} -strata. \square

Second proof. For a strict O_{F_0} -module X over a perfect field κ , recall its associated Dieudonné module

$$\pi_0 M \subset VM \subset M.$$

There exists an identification between short exact sequences:

$$\begin{aligned} 0 \rightarrow \text{Fil}(X) \rightarrow D(X) \rightarrow \text{Lie}(X) \rightarrow 0 \\ \cong \quad 0 \rightarrow VM/\pi_0 M \rightarrow M/\pi_0 M \rightarrow M/VM \rightarrow 0. \end{aligned}$$

We have two transition maps $\lambda : X \rightarrow X^\vee$ and $\lambda^\vee : X^\vee \rightarrow X$. By [Luo24, Thm. 6.2.2], for the Hodge filtrations

$$\text{Fil}(X) \subset D(X), \quad \text{Fil}(X^\vee) \subset D(X^\vee),$$

we have either

$$\lambda(\text{Fil}(X)) \subseteq \iota(\pi)D(X^\vee), \quad \text{or} \quad \lambda^\vee(\text{Fil}(X^\vee)) \subseteq \iota(\pi)D(X).$$

Translating into the Dieudonné modules, this means:

$$\lambda(VM/\pi_0 M) \subseteq \Pi M^\sharp/\pi_0 M^\sharp, \quad \text{or} \quad \lambda^\vee(VM^\sharp/\pi_0 M^\sharp) \subseteq \Pi^{-1}(\Pi M/\pi_0 M).$$

This is equivalent to having either

$$VM \subseteq \Pi M^\sharp, \quad \text{or} \quad VM^\sharp \subseteq M.$$

The assertion now follows from the definition of τ . \square

We define (closed) Kottwitz-Rapoport strata in the reduced locus of the RZ space.

Definition 5.7. (1) The \mathbb{F} -scheme \mathcal{Z} is defined as the reduced locus of the subfunctor of $\overline{\mathcal{N}}_{n,\varepsilon}^{[h]}$ consisting of tuples $(X, \iota, \lambda, \rho)$ over an \mathbb{F} -scheme S such that $\lambda^\vee(\mathrm{Fil}(X^\vee)) \subseteq \iota(\pi)D(X)$.

(2) The \mathbb{F} -scheme \mathcal{Y} is defined as the reduced locus of the subfunctor of $\overline{\mathcal{N}}_{n,\varepsilon}^{[h]}$ consisting of tuples $(X, \iota, \lambda, \rho)$ over an \mathbb{F} -scheme S such that $\lambda(\mathrm{Fil}(X)) \subseteq \iota(\pi)D(X^\vee)$.

Proposition 5.8. (1) We have the decomposition of the reduced locus,

$$\mathcal{N}_{n,\varepsilon,\mathrm{red}}^{[h]} = \mathcal{Z} \cup \mathcal{Y}.$$

Furthermore, for any perfect field κ over \mathbb{F} , we have lattice descriptions:

$$\mathcal{Z}(\kappa) = \left\{ M \in \mathcal{N}_{n,\varepsilon,\mathrm{red}}^{[h]}(\kappa) \mid \tau(\Pi M^\sharp) \subseteq M \right\}, \quad \text{and} \quad \mathcal{Y}(\kappa) = \left\{ M \in \mathcal{N}_{n,\varepsilon,\mathrm{red}}^{[h]}(\kappa) \mid \tau(M) \subseteq M^\sharp \right\}.$$

(2) For any $\Lambda_1 \in \mathcal{L}_{\mathcal{Z}}$ and $\Lambda_2 \in \mathcal{L}_{\mathcal{Y}}$, we have $\mathcal{Z}(\Lambda_1) \subset \mathcal{Z}$ and $\mathcal{Y}(\Lambda_2^\sharp) \subset \mathcal{Y}$.

Proof. Part (1) follows directly from Proposition 5.6 and the definition. For part (2), assume $(X, \iota_X, \lambda_X, \rho_X) \in \mathcal{Z}(\Lambda)(\kappa)$ corresponds to the lattice chain $\Lambda \otimes W_O(\kappa) \subseteq M \subseteq M^\sharp \subseteq \Lambda^\sharp \otimes W_O(\kappa)$. Then

$$\pi M^\sharp \subseteq \pi \Lambda^\sharp \otimes W_O(\kappa) \subseteq \Lambda \otimes W_O(\kappa) \subseteq M.$$

Then we have

$$\tau(\pi M^\sharp) \subseteq \tau(\Lambda \otimes W_O(\kappa)) = \Lambda \otimes W_O(\kappa) \subseteq M.$$

By definition, we have $(X, \iota_X, \lambda_X, \rho_X) \in \mathcal{Z}$. The same argument holds for \mathcal{Y} -strata. \square

Remark 5.9. (1) Proposition 5.4(2) can also be derived from the local model computation in §4.

(2) For the converse of Proposition 5.8(2), we have the following proper inclusions:

$$\bigcup_{\Lambda_1 \in \mathcal{L}_{\mathcal{Z}}} \mathcal{Z}(\Lambda_1) \subsetneq \mathcal{Z} \quad \text{and} \quad \bigcup_{\Lambda_2 \in \mathcal{L}_{\mathcal{Y}}} \mathcal{Y}(\Lambda_2^\sharp) \subsetneq \mathcal{Y}.$$

This follows from Proposition 5.3.

6. DELIGNE-LUSZTIG VARIETIES

In this section, we study a class of generalized Deligne-Lusztig varieties associated with symplectic, orthogonal, and general linear groups. These algebraic varieties are then used to derived global geometric properties of Bruhat-Tits strata of Rapoport-Zink spaces.

6.1. Deligne-Lusztig varieties. Let G_0 be a reductive group over \mathbb{F}_q . We fix a maximal torus T_0 and Borel subgroup B_0 over \mathbb{F}_q . Let G be the reductive group $G_0 \otimes_{\mathbb{F}_q} \mathbb{F}$ over \mathbb{F} and $T := T_0 \otimes_{\mathbb{F}_q} \mathbb{F}$ and $B := B_0 \otimes_{\mathbb{F}_q} \mathbb{F}$, with a Frobenius action Φ . Let $W = W_G$ be the Weyl group of G , let W_I be the subgroup of W generated by simple reflections in I , where I is a subset of the set of simple reflections S of W .

Let $I, J \subset S$ be two non-empty subsets. Every double coset in $W_I \backslash W / W_J$ contains a unique element of minimal length. Let ${}^I W^J \subset W$ be the set of such elements. Then ${}^I W^J \rightarrow W_I \backslash W / W_J$ is a bijection, which we regard as an identification.

Definition 6.1. For each $w \in {}^I W^{\Phi(I)}$, the (generalized) Deligne-Lusztig variety $X_{P_I}(w)$ is defined as

$$X_{P_I}(w) := \{g \in G/P_I : g^{-1}\Phi(g) \in P_I w P_{\Phi(I)}\}.$$

In the remaining part of the paper, we will write DL variety for Deligne-Lusztig variety.

We recall and generalize the *Görtz local model diagram* introduced in [GY10, §5.2]:

$$\begin{array}{ccc} & G & \\ \pi \swarrow & & \searrow L \\ G/P & & G/P. \end{array} \quad (6.1)$$

Here, π denotes the natural projection and L is the Lang map, which sends an element g to $g^{-1}\Phi(g)P$. Both maps π and L are smooth of relative dimension $\dim P$, and we have the equality

$$\pi^{-1}(C_P(w)) = L^{-1}(X_P(w)),$$

where $C_P(w) = PwP/P$ is the (generalized) Schubert cell in the partial flag variety G/P . This correspondence allows us to deduce properties of DL varieties from their corresponding Schubert cells.

Proposition 6.2. *Let $I \subset S$ be a non-empty subset and let $w \in {}^I W^{\Phi(I)}$.*

- (1) *The DL variety $X_{P_I}(w)$ is smooth of dimension $\ell(w) + \ell(W_{\Phi(I)}) - \ell(W_{I \cap \Phi(I)})$. Here $\ell(W_I)$ is the length of the longest element in the Weyl group W_I , and ${}^w I := wIw^{-1}$.*
- (2) *The DL variety $X_{P_I}(w)$ is irreducible if and only if $W_I w$ is not contained in a proper Φ -stable standard parabolic subgroup of W .*

Proof. Part (1) follows from [Hoe10, Lem. 2.1.3] and part (2) follows from [BR06]. □

6.2. Symplectic case. In this subsection, we study DL varieties for symplectic groups. In contrast to [RTW14], the spaces we consider here are not of Coxeter type. Recall the notation convention that $h = 2\mathfrak{t}$. Let $\Lambda \subset C$ be a vertex lattice of type $t(\Lambda) = t > h$. Suppose from now on that $t \neq 0$. We set

$$V = V_\Lambda := \Lambda^\sharp / \Lambda$$

with induced symplectic form $\langle \cdot, \cdot \rangle$. We fix a basis $V = \text{Span}_{\mathbb{F}_q}(e_1, \dots, e_{\mathfrak{t}}, f_1, \dots, f_{\mathfrak{t}})$ such that $\langle e_i, f_j \rangle = \delta_{ij}$, and $\langle e_i, e_j \rangle = \langle f_i, f_j \rangle = 0$ for any $1 \leq i, j \leq \mathfrak{t}$.

Let \mathbb{F} be the algebraic closure of \mathbb{F}_q with Frobenius Φ . We denote by $\Omega = \Omega_\Lambda = V_\Lambda \otimes_{\mathbb{F}_q} \mathbb{F}$ the symplectic space over \mathbb{F} . Consider the standard isotropic flags \mathcal{F}_\bullet in Ω defined by

$$\mathcal{F}_i = \text{Span}_{\mathbb{F}}\{e_1, \dots, e_i\} \quad \text{for } 1 \leq i \leq \mathfrak{t}. \quad (6.2)$$

This pins the choice of maximal torus and Borel $T \subset B \subset G := \text{Sp}(\Omega)$ which is stable under the Φ -action. We use $\Delta^* = \{s_1, \dots, s_{\mathfrak{t}}\}$ to denote the set of corresponding simple reflections in the Weyl group $W = N(T)/T$, where:

- for $1 \leq i \leq \mathfrak{t} - 1$, the reflection s_i interchanges $e_i \leftrightarrow e_{i+1}$ and $f_i \leftrightarrow f_{i+1}$, and fixes the other basis elements;

- The element $s_{\mathbf{t}}$ interchanges $e_{\mathbf{t}} \leftrightarrow f_{\mathbf{t}}$.

For each $0 \leq \mathfrak{s} \leq \mathfrak{h} \leq \mathfrak{r} \leq \mathbf{t} - 1$, we denote by:

$$I_{\mathfrak{r}\mathfrak{s}} := \{s_1, \dots, s_{\mathbf{t}-\mathfrak{r}-1}, s_{\mathbf{t}-\mathfrak{s}+1}, \dots, s_{\mathbf{t}}\} = \{s_1, \dots, s_{\mathbf{t}}\} \setminus \{s_{\mathbf{t}-\mathfrak{r}}, \dots, s_{\mathbf{t}-\mathfrak{s}}\}.$$

Note that a lot of objects we consider in the remaining of the section will depend on h (e.g. $I_{\mathfrak{r}\mathfrak{s}}$). To save notations, we drop h in the notations. We denote by $W_{\mathfrak{r}\mathfrak{s}}$ the subgroup of the Weyl group generated by elements in $I_{\mathfrak{r}\mathfrak{s}}$ and denote by $P_{\mathfrak{r}\mathfrak{s}}$ the corresponding standard parabolic subgroup.

Consider now the parabolic subgroup $P_{\mathfrak{h}\mathfrak{h}}$. The space $G/P_{\mathfrak{h}\mathfrak{h}}$ parametrizes isotropic subspaces in Ω of dimension $\mathbf{t} - \mathfrak{h}$. We consider the subvariety S_{Λ} of $G/P_{\mathfrak{h}\mathfrak{h}}$ such that for any field k over \mathbb{F} , its k -points are

$$S_{\Lambda}(k) = \{\mathcal{V} \subset \Omega_{\Lambda, k} \mid \mathcal{V} \text{ is isotropic, and } \dim \mathcal{V} = \mathbf{t} - \mathfrak{h}, \text{ and } \dim(\mathcal{V} \cap \Phi(\mathcal{V})) \geq \mathbf{t} - \mathfrak{h} - 1\}. \quad (6.3)$$

For any $0 \leq \mathfrak{s} \leq \mathfrak{h} < \mathfrak{r} \leq \mathbf{t}$, we let:

- $g_i := s_i s_{i+1} \cdots s_{\mathbf{t}-1} s_{\mathbf{t}} s_{\mathbf{t}-1} \cdots s_{i+1} s_i$.
- $w_{\mathfrak{r}\mathfrak{s}} := (s_{\mathbf{t}-\mathfrak{h}} s_{\mathbf{t}-\mathfrak{h}+1} \cdots s_{\mathbf{t}-\mathfrak{s}-1} g_{\mathbf{t}-\mathfrak{s}}) \cdot (s_{\mathbf{t}-\mathfrak{h}-1} s_{\mathbf{t}-\mathfrak{h}-2} \cdots s_{\mathbf{t}-\mathfrak{r}+1})$.
- $w'_{\mathfrak{r}\mathfrak{s}} := (s_{\mathbf{t}-\mathfrak{h}} s_{\mathbf{t}-\mathfrak{h}-1} \cdots s_{\mathbf{t}-\mathfrak{r}+1}) \cdot (s_{\mathbf{t}-\mathfrak{h}+1} s_{\mathbf{t}-\mathfrak{h}+2} \cdots s_{\mathbf{t}-\mathfrak{s}-1})$.

Hence $w_{\mathfrak{h}+1, \mathfrak{s}} = s_{\mathbf{t}-\mathfrak{h}} s_{\mathbf{t}-\mathfrak{h}+1} \cdots s_{\mathbf{t}-\mathfrak{s}-1} g_{\mathbf{t}-\mathfrak{s}}$, and $w_{\mathfrak{r}\mathfrak{h}} = g_{\mathbf{t}-\mathfrak{h}} s_{\mathbf{t}-\mathfrak{h}-1} s_{\mathbf{t}-\mathfrak{h}-2} \cdots s_{\mathbf{t}-\mathfrak{r}+1}$, and $w'_{\mathfrak{r}, \mathfrak{h}-1} = s_{\mathbf{t}-\mathfrak{h}} s_{\mathbf{t}-\mathfrak{h}-1} \cdots s_{\mathbf{t}-\mathfrak{r}+1}$. Note that the element g_i interchanges $e_i \leftrightarrow f_i$ and fixes all other elements in the basis.

The main result of this subsection is the following:

Theorem 6.3. *We have the following stratification:*

$$S_{\Lambda} = \left(\prod_{0 \leq j \leq \mathfrak{h} < i \leq \mathbf{t}} X_{P_{ij}}(w_{ij}) \right) \amalg \left(\prod_{0 \leq j < \mathfrak{h} < i \leq \mathbf{t}} X_{P_{ij}}(w'_{ij}) \right) \amalg X_{P_{\mathfrak{h}\mathfrak{h}}}(\text{id}), \quad (6.4)$$

such that for each $0 \leq \mathfrak{s} \leq \mathfrak{h} \leq \mathfrak{r} \leq \mathbf{t}$, we have the following closure relations:

$$\overline{X_{P_{\mathfrak{r}\mathfrak{s}}}(w_{\mathfrak{r}\mathfrak{s}})} = \left(\prod_{\mathfrak{s} \leq j \leq \mathfrak{h} < i \leq \mathfrak{r}} X_{P_{ij}}(w_{ij}) \right) \amalg \left(\prod_{0 \leq j < \mathfrak{h} < i \leq \mathfrak{r}} X_{P_{ij}}(w'_{ij}) \right) \amalg X_{P_{\mathfrak{h}\mathfrak{h}}}(\text{id}); \quad (6.5)$$

$$\overline{X_{P_{\mathfrak{r}\mathfrak{s}}}(w'_{\mathfrak{r}\mathfrak{s}})} = \left(\prod_{\mathfrak{s} \leq j < \mathfrak{h} < i \leq \mathfrak{r}} X_{P_{ij}}(w'_{ij}) \right) \amalg X_{P_{\mathfrak{h}\mathfrak{h}}}(\text{id}). \quad (6.6)$$

Moreover, S_{Λ} is irreducible and normal of dimension $\mathbf{t} + \mathfrak{h}$.

Proof. For $\mathfrak{s} \leq \mathfrak{h} < \mathfrak{r}$, define a locally closed subvariety $S_{\mathfrak{r}\mathfrak{s}}$ of $G/P_{\mathfrak{r}\mathfrak{s}}$ by specifying its k -points for any field k over \mathbb{F}

$$S_{\mathfrak{r}\mathfrak{s}}(k) := \left\{ \begin{array}{l} (\mathcal{F}_{\mathbf{t}-\mathfrak{r}} \subset \dots \subset \mathcal{F}_{\mathbf{t}-\mathfrak{s}}) \\ \in G/P_{\mathfrak{r}\mathfrak{s}}(k) \end{array} \left| \begin{array}{l} \mathcal{F}_i = \mathcal{F}_{i+1} \cap \Phi(\mathcal{F}_{i+1}) \text{ for } \mathbf{t} - \mathfrak{r} \leq i \leq \mathbf{t} - \mathfrak{h} - 1; \\ \mathcal{F}_{j+1} = \mathcal{F}_j + \Phi(\mathcal{F}_j) \text{ for } \mathbf{t} - \mathfrak{h} \leq j \leq \mathbf{t} - \mathfrak{s} - 1; \\ \mathcal{F}_{\mathbf{t}-\mathfrak{r}} = \Phi(\mathcal{F}_{\mathbf{t}-\mathfrak{r}}) \text{ and } \mathcal{F}_{\mathbf{t}-\mathfrak{s}} + \Phi(\mathcal{F}_{\mathbf{t}-\mathfrak{s}}) \text{ is non-isotropic.} \end{array} \right. \right\}.$$

For $\mathfrak{s} < \mathfrak{h} < \mathfrak{r}$, define a locally closed subvariety $S'_{\mathfrak{r}\mathfrak{s}}$ of $G/P_{\mathfrak{r}\mathfrak{s}}$ by specifying its k -points for any field k over \mathbb{F}

$$S'_{\mathfrak{r}\mathfrak{s}}(k) := \left\{ \begin{array}{l} (\mathcal{F}_{\mathbf{t}-\mathfrak{r}} \subset \dots \subset \mathcal{F}_{\mathbf{t}-\mathfrak{s}}) \\ \in G/P_{\mathfrak{r}\mathfrak{s}}(k) \end{array} \left| \begin{array}{l} \mathcal{F}_i = \mathcal{F}_{i+1} \cap \Phi(\mathcal{F}_{i+1}) \text{ for } \mathbf{t} - \mathfrak{r} \leq i \leq \mathbf{t} - \mathfrak{h} - 1; \\ \mathcal{F}_{j+1} = \mathcal{F}_j + \Phi(\mathcal{F}_j) \text{ for } \mathbf{t} - \mathfrak{h} \leq j \leq \mathbf{t} - \mathfrak{s} - 1; \\ \mathcal{F}_{\mathbf{t}-\mathfrak{r}} = \Phi(\mathcal{F}_{\mathbf{t}-\mathfrak{r}}) \text{ and } \mathcal{F}_{\mathbf{t}-\mathfrak{s}} = \Phi(\mathcal{F}_{\mathbf{t}-\mathfrak{s}}). \end{array} \right. \right\}.$$

Finally it is clear that $X_{P_{\mathbf{n}}}(\text{id})$ is the subvariety of $G/P_{\mathbf{n}}$ whose k -points is the finite set

$$X_{P_{\mathbf{n}}}(\text{id})(k) = \{\mathcal{F}_{\mathbf{t}-\mathbf{n}} \in G/P_{\mathbf{n}}(k) \mid \mathcal{F}_{\mathbf{t}-\mathbf{n}} = \Phi(\mathcal{F}_{\mathbf{t}-\mathbf{n}})\}.$$

We claim that:

$$S_{\Lambda} = \left(\prod_{0 \leq j \leq \mathbf{n} < i \leq \mathbf{t}} S_{\mathbf{r}\mathfrak{s}} \right) \amalg \left(\prod_{0 \leq j < \mathbf{n} < i \leq \mathbf{t}} S'_{\mathbf{r}\mathfrak{s}} \right) \amalg X_{P_{\mathbf{n}}}(\text{id}). \quad (6.7)$$

Assuming (6.7), the decomposition (6.4) now follow from Proposition 6.7, and Proposition 6.8 below. The closure relations (6.5) and (6.6) now follows from the moduli interpretations of $S_{\mathbf{r}\mathfrak{s}}$ and $S'_{\mathbf{r}\mathfrak{s}}$. In particular we see that $X_{P_{\mathbf{n}}}(w_{\mathbf{t}\mathbf{n}})$ is an open dense subvariety of S_{Λ} , hence the irreducibility and dimension of S_{Λ} follows from Proposition 6.2 and Proposition 6.7. The normality of S_{Λ} follows from the normality of Schubert varieties and the Görtz local model diagram (6.1).

In the remaining part of the proof, we prove (6.7). For any point $z = (\mathcal{F}_{\mathbf{t}-\mathbf{n}}) \in S_{\Lambda}(k)$, consider the flag

$$(\mathcal{F}_i)_{i \in \mathbb{Z}} := (\dots \subseteq \mathcal{F}_{\mathbf{t}-\mathbf{n}-1} \subseteq \mathcal{F}_{\mathbf{t}-\mathbf{n}} \subseteq \mathcal{F}_{\mathbf{t}-\mathbf{n}+1} \subseteq \dots),$$

such that for $i < \mathbf{t} - \mathbf{n}$, $\mathcal{F}_i = \mathcal{F}_{i+1} \cap \Phi(\mathcal{F}_{i+1})$ and for $i > \mathbf{t} - \mathbf{n}$, $\mathcal{F}_i = \mathcal{F}_{i-1} + \Phi(\mathcal{F}_{i-1})$. Since Ω_{Λ} is a finite dimensional vector space, the flag $(\mathcal{F}_i)_{i \in \mathbb{Z}}$ stabilizes at both ends.

Let $\mathbf{r} \geq \mathbf{n}$ be the unique integer such that $\mathcal{F}_{\mathbf{t}-\mathbf{r}-1} = \mathcal{F}_{\mathbf{t}-\mathbf{r}} \subsetneq \mathcal{F}_{\mathbf{t}-\mathbf{r}+1}$. Let $\mathfrak{s} \leq \mathbf{n}$ be the unique integer such that one of the following two situations occurs:

- (a) $\mathcal{F}_{\mathbf{t}-\mathfrak{s}-1} \subsetneq \mathcal{F}_{\mathbf{t}-\mathfrak{s}}$, $\mathcal{F}_{\mathbf{t}-\mathfrak{s}}$ is isotropic, and $\mathcal{F}_{\mathbf{t}-\mathfrak{s}+1}$ is anisotropic.
- (b) $\mathcal{F}_{\mathbf{t}-\mathfrak{s}-1} \subsetneq \mathcal{F}_{\mathbf{t}-\mathfrak{s}} = \mathcal{F}_{\mathbf{t}-\mathfrak{s}+1}$ and $\mathcal{F}_{\mathbf{t}-\mathfrak{s}}$ is isotropic.

Depending on whether the flag $(\mathcal{F}_i)_{i \in \mathbb{Z}}$ stabilizes first or becomes anisotropic first at the right end, exactly one of the above situations will occur.

Consider the isotropic flag $(\mathcal{F}_{\mathbf{t}-\mathbf{r}} \subset \dots \subset \mathcal{F}_{\mathbf{t}-\mathfrak{s}})$, we claim that

$$\dim_k \mathcal{F}_i = i \quad \text{for} \quad \mathbf{t} - \mathbf{r} \leq i \leq \mathbf{t} - \mathfrak{s}.$$

Indeed, for any $\mathbf{t} - \mathbf{r} < i < \mathbf{t} - \mathbf{n}$, we have

$$[\mathcal{F}_i : \mathcal{F}_{i-1}] = [\mathcal{F}_i : \mathcal{F}_i \cap \Phi(\mathcal{F}_i)] = [\Phi(\mathcal{F}_i) + \mathcal{F}_i : \Phi(\mathcal{F}_i)] = [\mathcal{F}_i + \Phi^{-1}(\mathcal{F}_i) : \mathcal{F}_i],$$

where for any $\mathcal{F} \subset \mathcal{G}$, we denote by $[\mathcal{F} : \mathcal{G}] := \dim_k \mathcal{F}/\mathcal{G}$. Since $\mathcal{F}_i = \mathcal{F}_{i+1} \cap \Phi(\mathcal{F}_{i+1})$, we have $\mathcal{F}_i + \Phi^{-1}(\mathcal{F}_i) \subseteq \mathcal{F}_{i+1}$. Hence we obtain

$$[\mathcal{F}_i : \mathcal{F}_{i-1}] \leq [\mathcal{F}_{i+1} : \mathcal{F}_i] \leq \dots \leq [\mathcal{F}_{\mathbf{t}-\mathbf{n}} : \mathcal{F}_{\mathbf{t}-\mathbf{n}-1}] \leq 1,$$

where the last inequality follows from the definition of S_{Λ} .

Since by assumption $\mathcal{F}_i \neq \mathcal{F}_{i-1}$ for any $\mathbf{t} - \mathbf{r} \leq i \leq \mathbf{t} - \mathbf{n}$, we know $[\mathcal{F}_i : \mathcal{F}_{i-1}] = 1$, hence $\dim_k \mathcal{F}_{i-1} = i - 1$. Similarly $\dim_k \mathcal{F}_i = i$ for $\mathbf{t} - \mathbf{n} < i \leq \mathbf{t} - \mathfrak{s}$. In other words the isotropic flag $(\mathcal{F}_{\mathbf{t}-\mathbf{r}} \subset \dots \subset \mathcal{F}_{\mathbf{t}-\mathfrak{s}})$ is a point in $G/P_{\mathbf{r}\mathfrak{s}}(k)$.

Now if situation (a) occurs for $\mathcal{F}_{\mathbf{t}-\mathfrak{s}}$, then $(\mathcal{F}_{\mathbf{t}-\mathbf{r}} \subset \dots \subset \mathcal{F}_{\mathbf{t}-\mathfrak{s}})$ lies in $S_{\mathbf{r}\mathfrak{s}}(k)$. If situation (b) occurs, then $(\mathcal{F}_{\mathbf{t}-\mathbf{r}} \subset \dots \subset \mathcal{F}_{\mathbf{t}-\mathfrak{s}})$ lies in $S'_{\mathbf{r}\mathfrak{s}}(k)$, resp. in $X_{P_{\mathbf{n}}}(\text{id})(k)$, if in addition $\mathbf{r} = \mathfrak{s} = \mathbf{n}$. Conversely, any point $z = (\mathcal{F}_{\mathbf{t}-\mathbf{r}} \subset \dots \subset \mathcal{F}_{\mathbf{t}-\mathfrak{s}})$ in $S_{\mathbf{r}\mathfrak{s}}(k)$ or $S'_{\mathbf{r}\mathfrak{s}}(k)$ or $X_{P_{\mathbf{n}}}(\text{id})(k)$ gives rise to a point $\mathcal{F}_{\mathbf{t}-\mathbf{n}} \in S_{\Lambda}$, and the point z can be recovered from $\mathcal{F}_{\mathbf{t}-\mathbf{n}}$ by the above procedure. This proves (6.7). \square

Remark 6.4. The DL varieties on the right hand side of (6.4) are all fine DL varieties in the sense of [Hoe10, §2.1.2] and [HLZ19, §2.3]. Theorem 6.3 is a special example of stratification into fine Deligne–Lusztig varieties, see [HLZ19, Thm. 2.3.1].

Remark 6.5. Let

$$w_\Lambda := s_{t-\mathfrak{n}} s_{t-\mathfrak{n}+1} \cdots s_{t-1} s_t s_{t-1} \cdots s_{t-\mathfrak{n}+1} s_{t-\mathfrak{n}} \quad \text{and} \quad w'_\Lambda := s_{t-\mathfrak{n}}.$$

One can verify that (cf. the proof of Proposition 6.7)

$$S_\Lambda = X_{P_{\mathfrak{n}\mathfrak{n}}}(w_\Lambda) \amalg X_{P_{\mathfrak{n}\mathfrak{n}}}(w'_\Lambda) \amalg X_{P_{\mathfrak{n}\mathfrak{n}}}(\text{id}), \quad (6.8)$$

where we can describe these DL varieties as follows:

$$X_{P_{\mathfrak{n}\mathfrak{n}}}(w_\Lambda) = \{\mathcal{V} \in S_\Lambda \mid \mathcal{V} \neq \Phi(\mathcal{V}) \text{ and } \mathcal{V} + \Phi(\mathcal{V}) \text{ is not isotropic}\};$$

$$X_{P_{\mathfrak{n}\mathfrak{n}}}(w'_\Lambda) = \{\mathcal{V} \in S_\Lambda \mid \mathcal{V} \neq \Phi(\mathcal{V}) \text{ and } \mathcal{V} + \Phi(\mathcal{V}) \text{ is isotropic}\};$$

$$X_{P_{\mathfrak{n}\mathfrak{n}}}(\text{id}) = \{\mathcal{V} \in S_\Lambda \mid \mathcal{V} = \Phi(\mathcal{V})\}.$$

We have the following decomposition, which refines the stratification (6.4):

$$\begin{aligned} \bullet X_{P_{\mathfrak{n}\mathfrak{n}}}(w_\Lambda) &= \prod_{i=h+1}^t X_{P_{i\mathfrak{n}}}(w_{i\mathfrak{n}}); \\ \bullet X_{P_{\mathfrak{n}\mathfrak{n}}}(w'_\Lambda) &= \left(\prod_{0 \leq j < \mathfrak{n} < i \leq t} X_{P_{ij}}(w_{ij}) \right) \amalg \left(\prod_{0 \leq j < \mathfrak{n} < i \leq t} X_{P_{ij}}(w'_{ij}) \right). \end{aligned}$$

This result can be established through two approaches: either via a group-theoretical argument as shown in [HZ25, Thm. 5.4], or through the moduli descriptions given in Propositions 6.7 and 6.8.

This decomposition corresponds to the stratification of *fine DL varieties*, as discussed in [HLZ19, §2.3]. In our context, these strata themselves are Deligne–Lusztig varieties, as demonstrated in [Hoe10, Thm. 2.1.7(1)]. This provides an alternative proof of Theorem 6.3, though we will not elaborate on this approach here. When $\mathfrak{n} = 0$, the component $X_{P_{\mathfrak{n}\mathfrak{n}}}(w'_\Lambda)$ is absent, which is evident both from the indices in (6.5) and from the moduli description in Proposition 6.8.

To complete the proof of Theorem 6.3, it remains to state and prove Proposition 6.7 and Proposition 6.8 below. First of all, it is straightforward to verify the following:

Lemma 6.6. (1) The element $w_{\mathfrak{r}\mathfrak{n}}$ stabilizes e_i and f_i for $1 \leq i \leq t - \mathfrak{r}$ and $t - \mathfrak{n} + 1 \leq i \leq t$. It acts on the remaining elements in the basis as follows:

$$e_{t-\mathfrak{r}+1} \xleftarrow{\quad \cdots \quad} e_{t-\mathfrak{n}} \xleftarrow{\quad f_{t-\mathfrak{r}+1} \quad} \cdots \xleftarrow{\quad} f_{t-\mathfrak{n}}.$$

Moreover, $w_{\mathfrak{r}\mathfrak{n}}$ is the minimal representative element in the double coset $W_{\mathfrak{r}} w_{\mathfrak{r}\mathfrak{n}} W_{\mathfrak{r}}$.

(2) The element $w_{\mathfrak{r}\mathfrak{s}}$ stabilizes e_i and f_i for $1 \leq i \leq t - \mathfrak{r}$ and $t - \mathfrak{s} + 1 \leq i \leq t$. It acts on the remaining e_i 's as follows:

$$\begin{array}{ccccccc} e_{t-\mathfrak{r}+1} & \xleftarrow{\quad \cdots \quad} & e_{t-\mathfrak{n}} & \xleftarrow{\quad} & e_{t-\mathfrak{n}+1} & \xrightarrow{\quad} & \cdots & \xrightarrow{\quad} & e_{t-\mathfrak{s}} \\ & & & & & & & & \\ f_{t-\mathfrak{r}+1} & \xleftarrow{\quad \cdots \quad} & f_{t-\mathfrak{n}} & \xleftarrow{\quad} & f_{t-\mathfrak{n}+1} & \xrightarrow{\quad} & \cdots & \xrightarrow{\quad} & f_{t-\mathfrak{s}} \end{array}$$

Moreover, $w_{\mathfrak{r}\mathfrak{s}}$ is the minimal representative element in the double coset $W_{\mathfrak{r},\mathfrak{s}} w_{\mathfrak{r}\mathfrak{s}} W_{\mathfrak{r},\mathfrak{s}}$.

(3) The element $w'_{\mathbf{r}\mathfrak{s}}$ stabilizes e_i and f_i for $1 \leq i \leq \mathbf{t} - \mathbf{r}$ and $\mathbf{t} - \mathfrak{s} + 1 \leq i \leq \mathbf{t}$. It acts on the remaining e_i 's as follows:

$$e_{\mathbf{t}-\mathbf{r}+1} \xleftarrow{\quad} \cdots \xleftarrow{\quad} e_{\mathbf{t}-\mathbf{n}} \xleftarrow{\quad} e_{\mathbf{t}-\mathbf{n}+1} \xrightarrow{\quad} e_{\mathbf{t}-\mathbf{n}+2} \xrightarrow{\quad} \cdots \xrightarrow{\quad} e_{\mathbf{t}-\mathfrak{s}}$$

We have the same action on the remaining f_i 's. Moreover, $w'_{\mathbf{r},\mathfrak{s}}$ is the minimal representative element in the double coset $W_{\mathbf{r},\mathfrak{s}}w'_{\mathbf{r},\mathfrak{s}}W_{\mathbf{r},\mathfrak{s}}$. \square

Proposition 6.7. For any integers $0 \leq \mathfrak{s} \leq \mathbf{n} < \mathbf{r} \leq \mathbf{t}$, the DL variety $X_{P_{\mathbf{r}\mathfrak{s}}}(w_{\mathbf{r}\mathfrak{s}})$ is the subvariety of $G/P_{\mathbf{r}\mathfrak{s}}$ whose k -points for any field k over \mathbb{F} are characterized by:

$$X_{P_{\mathbf{r}\mathfrak{s}}}(w_{\mathbf{r}\mathfrak{s}})(k) := \left\{ \begin{array}{l} (\mathcal{F}_{\mathbf{t}-\mathbf{r}} \subset \cdots \subset \mathcal{F}_{\mathbf{t}-\mathfrak{s}}) \\ \in G/P_{\mathbf{r}\mathfrak{s}}(k) \end{array} \left| \begin{array}{l} \mathcal{F}_i = \mathcal{F}_{i+1} \cap \Phi(\mathcal{F}_{i+1}) \text{ for } \mathbf{t} - \mathbf{r} \leq i \leq \mathbf{t} - \mathbf{n} - 1; \\ \mathcal{F}_{j+1} = \mathcal{F}_j + \Phi(\mathcal{F}_j) \text{ for } \mathbf{t} - \mathbf{n} \leq j \leq \mathbf{t} - \mathfrak{s} - 1; \\ \mathcal{F}_{\mathbf{t}-\mathbf{r}} = \Phi(\mathcal{F}_{\mathbf{t}-\mathbf{r}}) \text{ and } \mathcal{F}_{\mathbf{t}-\mathfrak{s}} + \Phi(\mathcal{F}_{\mathbf{t}-\mathfrak{s}}) \text{ is non-isotropic.} \end{array} \right. \right\}.$$

Its dimension is $\mathbf{r} + \mathfrak{s}$.

Proof. Let $gP \in X_P(w)$ be any point in the DL variety, we have $g^{-1}\Phi(g) \in Pw\sigma(P)$, in particular, we can find $p_1, p_2 \in P$ such that $\Phi(g) = gp_1w\Phi(p_2)$. Let $g_0 = gp_1$, then we have

$$\Phi(g_0) = g_0w\Phi(p_2)\Phi(p_1) = g_0w\Phi(p_2p_1).$$

Since $g_0P = gP$, we conclude that:

$$gP \in X_P(w) \quad \text{if and only if} \quad \exists g_0 \in gP \text{ such that } \Phi(g_0)\Phi(P) = g_0w\Phi(P). \quad (6.9)$$

In particular, by identifying the right coset gP with partial flags $(\mathcal{F}_{\mathbf{t}-\mathbf{r}} \subset \cdots \subset \mathcal{F}_{\mathbf{t}-\mathfrak{s}})$, we can use Lemma 6.6(i)(ii) and equation (6.9) to show that any partial flag $(\mathcal{F}_i) = g_0P_{\mathbf{r}\mathfrak{s}} \in X_{P_{\mathbf{r}\mathfrak{s}}}(w_{\mathbf{r}\mathfrak{s}})(k)$ belongs to the moduli functor defined in the assertion. Conversely, consider a partial flag $(\mathcal{F}_i) = g_0P_{\mathbf{r}\mathfrak{s}}$ in $\text{Sp}(\Omega_\Lambda)/P_{\mathbf{r}\mathfrak{s}}(k)$ that defines a point in the moduli functor. We will prove that this flag lies in $X_{P_{\mathbf{r}\mathfrak{s}}}(w_{\mathbf{r}\mathfrak{s}})(k)$.

We find a common basis for all \mathcal{F}_i and $\Phi(\mathcal{F}_j)$ in the following three steps:

Step 1. We choose vectors $u_i \in \mathcal{F}_{\mathbf{t}-\mathbf{n}}$ for $1 \leq i \leq \mathbf{t} - \mathbf{n}$ such that $\mathcal{F}_{\mathbf{t}-\mathbf{r}} = \text{Span}_k(u_1, \dots, u_{\mathbf{t}-\mathbf{r}})$ and that

$$\mathcal{F}_i = \mathcal{F}_{i-1} \oplus \langle u_i \rangle \quad \text{for all } \mathbf{t} - \mathbf{r} \leq i \leq \mathbf{t} - \mathbf{n} + 1.$$

Step 2. For each $\mathbf{t} - \mathbf{r} + 1 \leq i \leq \mathbf{t} - \mathbf{n} - 1$, we have $\Phi(\mathcal{F}_i) \stackrel{1}{\subset} \Phi(\mathcal{F}_{i+1})$. For each $\mathbf{t} - \mathbf{r} \leq i \leq \mathbf{t} - \mathbf{n} - 1$, the equality $\mathcal{F}_i = \mathcal{F}_{i+1} \cap \Phi(\mathcal{F}_{i+1})$ implies that

$$\mathcal{F}_i = \mathcal{F}_{i+1} \cap \Phi(\mathcal{F}_{i+1}) \stackrel{1}{\subset} \Phi(\mathcal{F}_{i+1}).$$

These inclusions yield:

$$\Phi(\mathcal{F}_{i+1}) = \mathcal{F}_i + \Phi(\mathcal{F}_i), \quad \text{for all } \mathbf{t} - \mathbf{r} + 1 \leq i \leq \mathbf{t} - \mathbf{n} - 1. \quad (6.10)$$

Therefore, we can find a vector $u_{\mathbf{t}-\mathbf{n}+1} \in \Phi(\mathcal{F}_{\mathbf{t}-\mathbf{n}})$ such that:

$$\Phi(\mathcal{F}_{\mathbf{t}-\mathbf{n}}) = \mathcal{F}_{\mathbf{t}-\mathbf{n}-1} \oplus \langle u_{\mathbf{t}-\mathbf{n}+1} \rangle.$$

Furthermore, by (6.10) and induction, we have $u_{\mathbf{t}-\mathbf{n}+1} \notin \mathcal{F}_i$ for any $\mathbf{t} - \mathbf{r} \leq i \leq \mathbf{t} - \mathbf{n}$, and that:

$$\Phi(\mathcal{F}_{i+1}) = \mathcal{F}_i \oplus \langle u_{\mathbf{t}-\mathbf{n}+1} \rangle, \quad \text{for all } \mathbf{t} - \mathbf{r} \leq i \leq \mathbf{t} - \mathbf{n} - 1.$$

Step 3. Finally, for each $\mathfrak{t} - \mathfrak{h} + 1 \leq j \leq \mathfrak{t} - \mathfrak{s} - 1$ (if $\mathfrak{s} = \mathfrak{h}$, we skip this step), we denote by

$$u_j := \Phi(u_{j-1}) = \Phi^{j-(\mathfrak{t}-\mathfrak{h}+1)}(u_{\mathfrak{t}-\mathfrak{h}+1}).$$

Since $\mathcal{F}_j = \mathcal{F}_{j-1} + \Phi(\mathcal{F}_{j-1})$, by induction, we have

$$\mathcal{F}_j = \text{Span}_k(u_1, \dots, u_j) \quad \text{and} \quad \Phi(\mathcal{F}_j) = \text{Span}_k(u_1, \dots, u_{\mathfrak{t}-\mathfrak{h}-1}, u_{\mathfrak{t}-\mathfrak{h}+1}, \dots, u_{j+1}).$$

To summarize, we have formed a set of vectors u_i such that

$$\mathcal{F}_i = \text{Span}_k(u_1, \dots, u_i), \quad \text{for all } \mathfrak{t} - \mathfrak{r} \leq i \leq \mathfrak{t} - \mathfrak{s},$$

and that for any $\mathfrak{t} - \mathfrak{r} \leq j \leq \mathfrak{t} - \mathfrak{s}$, we have

$$\Phi(\mathcal{F}_j) = \begin{cases} \text{Span}_k(u_1, \dots, u_j, u_{\mathfrak{t}-\mathfrak{h}+1}), & \mathfrak{t} - \mathfrak{r} \leq j \leq \mathfrak{t} - \mathfrak{h}; \\ \text{Span}_k(u_1, \dots, u_{\mathfrak{t}-\mathfrak{h}-1}, u_{\mathfrak{t}-\mathfrak{h}+1}, \dots, u_{j+1}), & \mathfrak{t} - \mathfrak{h} + 1 \leq j \leq \mathfrak{t} - \mathfrak{s}. \end{cases}$$

Notice that:

- Since the subspace $\mathcal{F}_{\mathfrak{t}-\mathfrak{s}}$ is isotropic, we have

$$\langle u_i, u_j \rangle = 0 \quad \text{for all } 1 \leq i, j \leq \mathfrak{t} - \mathfrak{s}.$$

- Since the subspace $\Phi(\mathcal{F}_{\mathfrak{t}-\mathfrak{s}})$ is isotropic, we have

$$\langle u_i, u_{\mathfrak{t}-\mathfrak{s}+1} \rangle = 0 \quad \text{for all } 1 \leq i \neq \mathfrak{t} - \mathfrak{h} \leq \mathfrak{t} - \mathfrak{s}.$$

- Since $\mathcal{F}_{\mathfrak{t}-\mathfrak{s}} + \Phi(\mathcal{F}_{\mathfrak{t}-\mathfrak{s}})$ is anisotropic, we have

$$\langle u_{\mathfrak{t}-\mathfrak{h}}, u_{\mathfrak{t}-\mathfrak{s}+1} \rangle \neq 0. \tag{6.11}$$

By normalizing $u_{\mathfrak{t}-\mathfrak{s}+1}$, we may assume that the pairing equals 1.

Define a linear map between two subspaces of Ω_Λ by sending

$$e_i \mapsto u_i, \quad \text{for } 1 \leq i \leq \mathfrak{t} - \mathfrak{s}; \quad \text{and} \quad f_{\mathfrak{t}-\mathfrak{h}} \mapsto u_{\mathfrak{t}-\mathfrak{s}+1}.$$

This is an isometry, and by Witt's theorem, we can find a $g \in G$ that extends this isometry to an isometry of Ω_Λ . For such g , we have:

$$\left[\mathcal{F}_{\mathfrak{t}-\mathfrak{r}} \subset \dots \subset \mathcal{F}_{\mathfrak{t}-\mathfrak{s}} \right] \xrightarrow{g} \left[\mathcal{F}_{\mathfrak{t}-\mathfrak{r}} \subset \dots \subset \mathcal{F}_{\mathfrak{t}-\mathfrak{s}} \right],$$

and

$$\left[w_{[\mathfrak{r}, \mathfrak{h}]}(\mathcal{F}_{\mathfrak{t}-\mathfrak{r}}) \subset \dots \subset w_{[\mathfrak{r}, \mathfrak{h}]}(\mathcal{F}_{\mathfrak{t}-\mathfrak{h}}) \right] \xrightarrow{g} \left[\Phi(\mathcal{F}_{\mathfrak{t}-\mathfrak{r}}) \subset \dots \subset \Phi(\mathcal{F}_{\mathfrak{t}-\mathfrak{h}}) \right].$$

Recall that \mathcal{F}_i are standard flags defined in (6.2). Equivalently, we have $gP_{\mathfrak{r}\mathfrak{s}} = g_0P_{\mathfrak{r}\mathfrak{s}}$ and that $\Phi(g)P_{\mathfrak{r}\mathfrak{s}} = gw_{\mathfrak{r}\mathfrak{s}}P_{\mathfrak{r}\mathfrak{s}}$. By (6.9), we have $g_0P_{\mathfrak{r}\mathfrak{s}} = gP_{\mathfrak{r}\mathfrak{s}} \in X_{P_{\mathfrak{r}\mathfrak{s}}}(w_{\mathfrak{r}\mathfrak{s}})$.

Finally, by Lemma 6.6(1)(2), we see that $w_{\mathfrak{r}\mathfrak{s}}$ acts trivially on e_i and f_i for $1 \leq i \leq \mathfrak{t} - \mathfrak{r}$ and $\mathfrak{t} - \mathfrak{s} + 1 \leq i \leq \mathfrak{t}$, hence $w_{\mathfrak{r}\mathfrak{s}}I_{\mathfrak{r}\mathfrak{s}} = I_{\mathfrak{r}\mathfrak{s}}w_{\mathfrak{r}\mathfrak{s}}$. By Proposition 6.2(1), the dimension of the DL variety $X_{P_{\mathfrak{r}\mathfrak{s}}}(w_{\mathfrak{r}\mathfrak{s}})$ equals $\ell(w_{\mathfrak{r}\mathfrak{s}}) = \mathfrak{r} + \mathfrak{s}$, as desired. \square

We have similar statements for the other class of DL varieties:

Proposition 6.8. For any integers $0 \leq \mathfrak{s} \leq \mathfrak{h} < \mathfrak{r} \leq \mathfrak{t}$, the DL variety $X_{P_{\mathfrak{r}\mathfrak{s}}}(w'_{\mathfrak{r}\mathfrak{s}})$ is the subvariety of $G/P_{\mathfrak{r}\mathfrak{s}}$ whose k -points for any field k over \mathbb{F} are characterized by:

$$X_{P_{\mathfrak{r}\mathfrak{s}}}(w'_{\mathfrak{r}\mathfrak{s}})(k) := \left\{ \begin{array}{l} (\mathcal{F}_{\mathfrak{t}-\mathfrak{r}} \subset \dots \subset \mathcal{F}_{\mathfrak{t}-\mathfrak{s}}) \\ \in G/P_{\mathfrak{r}\mathfrak{s}}(k) \end{array} \left| \begin{array}{l} \mathcal{F}_i = \mathcal{F}_{i+1} \cap \Phi(\mathcal{F}_{i+1}) \text{ for } \mathfrak{t} - \mathfrak{r} \leq i \leq \mathfrak{t} - \mathfrak{h} - 1; \\ \mathcal{F}_{j+1} = \mathcal{F}_j + \Phi(\mathcal{F}_j) \text{ for } \mathfrak{t} - \mathfrak{h} \leq j \leq \mathfrak{t} - \mathfrak{s} - 1; \\ \mathcal{F}_{\mathfrak{t}-\mathfrak{r}} = \Phi(\mathcal{F}_{\mathfrak{t}-\mathfrak{r}}) \text{ and } \mathcal{F}_{\mathfrak{t}-\mathfrak{s}} = \Phi(\mathcal{F}_{\mathfrak{t}-\mathfrak{s}}). \end{array} \right. \right\}.$$

Its dimension is $\mathfrak{r} - \mathfrak{s}$.

Proof. It is straightforward to verify that $w'_{\mathfrak{r}\mathfrak{s}}I_{\mathfrak{r}\mathfrak{s}} = I_{\mathfrak{r}\mathfrak{s}}w'_{\mathfrak{r}\mathfrak{s}}$, and this concludes the dimension formula. By Lemma 6.6(iii) and (6.9), any partial flag $(\mathcal{F}_i) = g_0P_{\mathfrak{r}\mathfrak{s}} \in X_{P_{\mathfrak{r}\mathfrak{s}}}(w'_{\mathfrak{r}\mathfrak{s}})(k)$ lies in the moduli functor. Conversely, let $(\mathcal{F}_i) = g_0P_{\mathfrak{r}\mathfrak{s}}$ be a partial flag in $G/P_{\mathfrak{r}\mathfrak{s}}(k)$ defines a point in the moduli functor, we show that this flag lies in $X_{P_{\mathfrak{r}\mathfrak{s}}}(w'_{\mathfrak{r}\mathfrak{s}})(k)$.

Following with the same procedure as Proposition 6.7, we can find vectors $u_i \in \Omega_{\Lambda}(k)$ such that

$$\mathcal{F}_i = \text{Span}_k(u_1, \dots, u_i), \quad \text{for all } \mathfrak{t} - \mathfrak{r} \leq i \leq \mathfrak{t} - \mathfrak{s},$$

and that for any $\mathfrak{t} - \mathfrak{r} \leq j \leq \mathfrak{t} - \mathfrak{s} - 1$, we have

$$\Phi(\mathcal{F}_j) = \begin{cases} \text{Span}_k(u_1, \dots, u_j, u_{\mathfrak{t}-\mathfrak{h}+1}), & \mathfrak{t} - \mathfrak{r} \leq j \leq \mathfrak{t} - \mathfrak{h}; \\ \text{Span}_k(u_1, \dots, u_{\mathfrak{t}-\mathfrak{h}-1}, u_{\mathfrak{t}-\mathfrak{h}+1}, \dots, u_{j+1}), & \mathfrak{t} - \mathfrak{h} + 1 \leq j \leq \mathfrak{t} - \mathfrak{s} - 1. \end{cases}$$

Since $\mathcal{F}_{\mathfrak{t}-\mathfrak{s}} = \mathcal{F}_{\mathfrak{t}-\mathfrak{s}-1} + \Phi(\mathcal{F}_{\mathfrak{t}-\mathfrak{s}-1})$ is Φ -stable now, we have

$$\Phi(\mathcal{F}_{\mathfrak{t}-\mathfrak{s}}) = \mathcal{F}_{\mathfrak{t}-\mathfrak{s}} = \text{Span}_k(u_1, \dots, u_j).$$

Since $\mathcal{F}_{\mathfrak{t}-\mathfrak{s}}$ is isotropic, we conclude that

$$\langle u_i, u_j \rangle = 0, \quad \text{for all } 1 \leq i, j \leq \mathfrak{t} - \mathfrak{s}.$$

Define a linear map between two subspaces of Ω_{Λ} by sending $e_i \mapsto u_i$, for $1 \leq i \leq \mathfrak{t} - \mathfrak{s}$. This is an isometry and, by Witt's theorem, we can find a $g \in G$ that extends this isometry to an isometry of Ω_{Λ} . By the same argument as in the proof of Proposition 6.7, we have $g_0P_{\mathfrak{r}\mathfrak{s}} = gP_{\mathfrak{r}\mathfrak{s}} \in X_{P_{\mathfrak{r}\mathfrak{s}}}(w_{\mathfrak{r}\mathfrak{s}})$. \square

Remark 6.9. The DL variety $X_{P_{\mathfrak{r}\mathfrak{s}}}(w'_{\mathfrak{r}\mathfrak{s}}) \subset \text{Sp}(\Omega_{\Lambda})/P_{\mathfrak{r}\mathfrak{s}}$ in Proposition 6.8 is not irreducible in general. Its irreducible components are all isomorphic to the DL variety associated to the general linear group, as described in Proposition 6.12. These components are indexed by pairs of \mathbb{F}_q -rational subspaces $\mathcal{F}_{\mathfrak{t}-\mathfrak{r}}^0 \subset \mathcal{F}_{\mathfrak{t}-\mathfrak{s}}^0 \subset V_{\Lambda}$ of dimensions $\mathfrak{t} - \mathfrak{r}$ and $\mathfrak{t} - \mathfrak{s}$, resp. For further details on this matter, we refer to [RTW14, Prop. 5.7].

6.3. The orthogonal case. First, we introduce the relevant group theoretic notations in the orthogonal case. Let $\Lambda \subset C$ be a vertex lattice of type $t < h$. Since h and t are always even, we still let $\mathfrak{h} = \frac{h}{2}$ and $\mathfrak{t} = \frac{t}{2}$. Let $\mathfrak{n} := \lfloor n/2 \rfloor$ and $\mathfrak{h}' = \mathfrak{n} - \mathfrak{h}$.

6.3.1. *The even dimensional orthogonal case.* Assume n is even so that $n - t = 2(\mathbf{n} - \mathbf{t})$ is even. Let $\Omega = V_\Lambda \otimes_{\mathbb{F}_q} \mathbb{F}$ be the quadratic space over \mathbb{F} with basis $\{e_1, \dots, e_{\mathbf{n}-\mathbf{t}}, f_1, \dots, f_{\mathbf{n}-\mathbf{t}}\}$ such that $\text{Span}_{\mathbb{F}}\{e_1, \dots, e_{\mathbf{n}-\mathbf{t}}\}$ and $\text{Span}_{\mathbb{F}}\{f_1, \dots, f_{\mathbf{n}-\mathbf{t}}\}$ are totally isotropic and $(e_i, f_j) = \delta_{i,j}$. If V_Λ is split, then the Frobenius Φ fixes $e_1, \dots, e_{\mathbf{n}-\mathbf{t}}, f_1, \dots, f_{\mathbf{n}-\mathbf{t}}$. If V_Λ is non-split, then Φ fixes $e_1, \dots, e_{\mathbf{n}-\mathbf{t}-1}, f_1, \dots, f_{\mathbf{n}-\mathbf{t}-1}$ but interchanges $e_{\mathbf{n}-\mathbf{t}} \leftrightarrow f_{\mathbf{n}-\mathbf{t}}$.

Consider the isotropic flags \mathcal{F}_\bullet^\pm and \mathcal{F}_\bullet^\mp in Ω defined by

$$\begin{aligned}\mathcal{F}_i^\pm &= \text{Span}_{\mathbb{F}}\{e_1, \dots, e_i\} \quad \text{for } 1 \leq i \leq \mathbf{n} - \mathbf{t} - 1 \\ \mathcal{F}_{\mathbf{n}-\mathbf{t}}^+ &= \text{Span}_{\mathbb{F}}\{e_1, \dots, e_{\mathbf{n}-\mathbf{t}-1}, e_{\mathbf{n}-\mathbf{t}}\} \\ \mathcal{F}_{\mathbf{n}-\mathbf{t}}^- &= \text{Span}_{\mathbb{F}}\{e_1, \dots, e_{\mathbf{n}-\mathbf{t}-1}, f_{\mathbf{n}-\mathbf{t}}\}.\end{aligned}$$

Hence when V_Λ is split, we have $\Phi \mathcal{F}_\bullet^\pm = \mathcal{F}_\bullet^\pm$. When V_Λ is non-split, we have $\Phi \mathcal{F}_\bullet^\pm = \mathcal{F}_\bullet^\mp$. The stabilizers of \mathcal{F}_\bullet^\pm are the same, which we denote as $B \subset H = \text{SO}(\Omega)$. Hence B is a Φ -stable Borel subgroup containing T where T is a maximal Φ -stable torus $T \subset \text{SO}(\Omega)$. We use $\Delta^* = \{s_1, \dots, s_{\mathbf{n}-\mathbf{t}}\}$ to denote the set of corresponding simple reflections in the Weyl group $W = N(T)/T$ where

- s_i interchanges $e_i \leftrightarrow e_{i+1}$ and $f_i \leftrightarrow f_{i+1}$, and fixes the other basis elements if $i < \mathbf{n} - \mathbf{t}$.
- $s_{\mathbf{n}-\mathbf{t}}$ interchanges $e_{\mathbf{n}-\mathbf{t}-1} \leftrightarrow f_{\mathbf{n}-\mathbf{t}}$ and $f_{\mathbf{n}-\mathbf{t}-1} \leftrightarrow e_{\mathbf{n}-\mathbf{t}}$, and fixes the other basis elements.

We also use t^+ and t^- to denote $s_{\mathbf{n}-\mathbf{t}-1}$ and $s_{\mathbf{n}-\mathbf{t}}$ respectively.

For each $0 \leq \mathfrak{s} \leq \mathfrak{h}' \leq \mathfrak{r} \leq \mathbf{n} - \mathbf{t} - 1$, we let

$$I_{\mathfrak{r}\mathfrak{s}} := \begin{cases} \{s_1, \dots, s_{\mathbf{n}-\mathbf{t}-\mathfrak{r}-1}\} & \text{if } h \geq n - 2, \\ \{s_1, \dots, s_{\mathbf{n}-\mathbf{t}-\mathfrak{r}-1}, s_{\mathbf{n}-\mathbf{t}-\mathfrak{s}+1}, \dots, s_{\mathbf{n}-\mathbf{t}-2}, s_{\mathbf{n}-\mathbf{t}-1}, s_{\mathbf{n}-\mathbf{t}}\} & \text{if } h < n - 2, \end{cases}$$

and let $P_{\mathfrak{r}\mathfrak{s}}$ be the corresponding parabolic subgroup. Next, we define Weyl group elements for DL varieties. For each $0 \leq \mathfrak{s} \leq \mathfrak{h}' < \mathfrak{r} \leq \mathbf{n} - \mathbf{t}$, let

$$\begin{aligned}\bullet \quad g_i^\pm &:= \begin{cases} t^\pm & \text{if } i = \mathbf{n} - \mathbf{t}, \\ t^- t^+ & \text{if } i = \mathbf{n} - \mathbf{t} - 1, \\ s_i s_{i+1} \cdots s_{\mathbf{n}-\mathbf{t}-2} t^- t^+ s_{\mathbf{n}-\mathbf{t}-2} \cdots s_{i+1} s_i & \text{otherwise.} \end{cases} \\ \bullet \quad w_{\mathfrak{r}\mathfrak{s}}^\pm &:= (s_{\mathfrak{h}-\mathbf{t}} s_{\mathfrak{h}-\mathbf{t}+1} \cdots s_{\mathbf{n}-\mathbf{t}-\mathfrak{s}-1} g_{\mathbf{n}-\mathbf{t}-\mathfrak{s}}^\pm) \cdot (s_{\mathfrak{h}-\mathbf{t}-1} s_{\mathfrak{h}-\mathbf{t}-2} \cdots s_{\mathbf{n}-\mathbf{t}-\mathfrak{r}+1}).\end{aligned}$$

Note that when $h \neq n$, then $w_{\mathfrak{r}\mathfrak{s}}^\pm$ (resp. g_i^\pm) is independent of \pm sign and we will simply denote it as $w_{\mathfrak{r}\mathfrak{s}}$ (resp. g_i) in this case.

For each $0 \leq \mathfrak{s} \leq \mathbf{n} - \mathfrak{h} < \mathfrak{r} \leq \mathbf{n} - \mathbf{t}$, let

$$\bullet \quad w'_{\mathfrak{r}\mathfrak{s}} = w_{\mathfrak{r}\mathfrak{s}}^{\prime+} := (s_{\mathfrak{h}-\mathbf{t}} s_{\mathfrak{h}-\mathbf{t}-1} \cdots s_{\mathbf{n}-\mathbf{t}-\mathfrak{r}+1}) \cdot (s_{\mathfrak{h}-\mathbf{t}+1} s_{\mathfrak{h}-\mathbf{t}+2} \cdots s_{\mathbf{n}-\mathbf{t}-\mathfrak{s}-1}).$$

In particular, $w_{\mathbf{n}-\mathfrak{h}+1, \mathfrak{s}} = s_{\mathfrak{h}-\mathbf{t}} s_{\mathfrak{h}-\mathbf{t}+1} \cdots s_{\mathbf{n}-\mathbf{t}-\mathfrak{s}-1} g_{\mathbf{n}-\mathbf{t}-\mathfrak{s}}$, and $w_{\mathfrak{r}, \mathbf{n}-\mathfrak{h}} = g_{\mathfrak{h}-\mathbf{t}} s_{\mathfrak{h}-\mathbf{t}-1} s_{\mathfrak{h}-\mathbf{t}-2} \cdots s_{\mathbf{n}-\mathbf{t}-\mathfrak{r}+1}$, and $w'_{\mathfrak{r}, \mathbf{n}-\mathfrak{h}-1} = s_{\mathfrak{h}-\mathbf{t}} s_{\mathfrak{h}-\mathbf{t}-1} \cdots s_{\mathbf{n}-\mathbf{t}-\mathfrak{r}+1}$. Note that the element g_i interchanges e_i with f_i , $e_{\mathbf{n}-\mathbf{t}}$ with $f_{\mathbf{n}-\mathbf{t}}$ and fixes all the other basis vectors when $i < \mathbf{n} - \mathbf{t}$ and $g_{\mathbf{n}-\mathbf{t}}^\pm = t^\pm$.

When $h = n - 2$, then $\mathfrak{s} = 0$ and $w'_{\mathfrak{r}0} = w_{\mathfrak{r}0}^{\prime+} = t^+ s_{\mathbf{n}-1-\mathbf{t}-1} \cdots s_{\mathbf{n}-\mathbf{t}-\mathfrak{r}+1}$. In this special case, we define another Weyl element:

$$\bullet \quad w_{\mathfrak{r}\mathfrak{s}}^{\prime-} := t^- s_{\mathbf{n}-\mathbf{t}-2} \cdots s_{\mathbf{n}-\mathbf{t}-\mathfrak{r}+1}.$$

6.3.2. *The odd dimensional orthogonal case.* In this subsection, we assume $n = 2\mathbf{n} + 1$ is odd. We set

$$V_{\Lambda^\sharp} := \Lambda/\pi\Lambda^\sharp$$

with quadratic form induced from $(,)$. Let $\Omega = V_{\Lambda^\sharp} \otimes_{\mathbb{F}_q} \mathbb{F}$ be a quadratic space over \mathbb{F} . Since V_{Λ^\sharp} is of odd dimension, we can choose a basis $\{e_1, \dots, e_{\mathbf{n}-\mathbf{t}}, f_1, \dots, f_{\mathbf{n}-\mathbf{t}}, e_{2(\mathbf{n}-\mathbf{t})+1}\}$ such that $\text{Span}_{\mathbb{F}}\{e_1, \dots, e_{\mathbf{n}-\mathbf{t}}\}$ and $\text{Span}_{\mathbb{F}}\{f_1, \dots, f_{\mathbf{n}-\mathbf{t}}\}$ are totally isotropic and $(e_i, f_j) = \delta_{i,j}$ and $(e_{2(\mathbf{n}-\mathbf{t})+1}, e_{2(\mathbf{n}-\mathbf{t})+1}) = 1$ and $e_{2(\mathbf{n}-\mathbf{t})+1}$ is orthogonal to all the other basis vectors and Φ fixes all the basis vectors.

Consider the isotropic flags \mathcal{F}_\bullet in Ω defined by

$$\mathcal{F}_i = \text{Span}_{\mathbb{F}}\{e_1, \dots, e_i\} \text{ for } 1 \leq i \leq \mathbf{n} - \mathbf{t}.$$

The stabilizer of \mathcal{F}_\bullet is denoted as $B \subset H = \text{SO}(\Omega)$. Hence B is a Φ -stable Borel subgroup containing T where T is a maximal Φ -stable torus $T \subset \text{SO}(\Omega)$.

We use $\Delta^* = \{s_1, \dots, s_{\mathbf{n}-\mathbf{t}-2}, s_{\mathbf{n}-\mathbf{t}-1}, s_{\mathbf{n}-\mathbf{t}}\}$ to denote the set of corresponding simple reflections in the Weyl group $W = N(T)/T$, where

- s_i interchanges $e_i \leftrightarrow e_{i+1}$ and $f_i \leftrightarrow f_{i+1}$, and fixes the other basis elements if $i < \mathbf{n} - \mathbf{t}$.
- $s_{\mathbf{n}-\mathbf{t}}$ interchanges $e_{\mathbf{n}-\mathbf{t}} \leftrightarrow f_{\mathbf{n}-\mathbf{t}}$ and $e_{2(\mathbf{n}-\mathbf{t})+1} \leftrightarrow -e_{2(\mathbf{n}-\mathbf{t})+1}$, and fixes the other basis elements.

For each $0 \leq \mathfrak{s} \leq \mathbf{n} - \mathbf{h} \leq \mathbf{r} \leq \mathbf{n} - \mathbf{t} - 1$, we let

$$I_{\mathbf{r}\mathfrak{s}} := \begin{cases} \{s_1, \dots, s_{\mathbf{n}-\mathbf{t}-\mathbf{r}-1}\} & \text{if } h \geq n - 2, \\ \{s_1, \dots, s_{\mathbf{n}-\mathbf{t}-\mathbf{r}-1}, s_{\mathbf{n}-\mathbf{t}-\mathbf{s}+1}, \dots, s_{\mathbf{n}-\mathbf{t}-2}, s_{\mathbf{n}-\mathbf{t}-1}, s_{\mathbf{n}-\mathbf{t}}\} & \text{if } h < n - 2, \end{cases}$$

and $P_{\mathbf{r}\mathfrak{s}}$ be the corresponding parabolic subgroup.

For each $0 \leq \mathfrak{s} \leq \mathbf{n} - \mathbf{h} < \mathbf{r} \leq \mathbf{n} - \mathbf{t}$, let

- $g_i := s_i s_{i+1} \cdots s_{\mathbf{n}-\mathbf{t}-1} s_{\mathbf{n}-\mathbf{t}} s_{\mathbf{n}-\mathbf{t}-1} \cdots s_{i+1} s_i$.
- $w_{\mathbf{r}\mathfrak{s}} := (s_{\mathbf{h}-\mathbf{t}} s_{\mathbf{h}-\mathbf{t}+1} \cdots s_{\mathbf{n}-\mathbf{t}-\mathfrak{s}-1} g_{\mathbf{n}-\mathbf{t}-\mathfrak{s}}) \cdot (s_{\mathbf{h}-\mathbf{t}-1} s_{\mathbf{h}-\mathbf{t}-2} \cdots s_{\mathbf{n}-\mathbf{t}-\mathbf{r}+1})$.
- $w'_{\mathbf{r}\mathfrak{s}} := (s_{\mathbf{h}-\mathbf{t}} s_{\mathbf{h}-\mathbf{t}-1} \cdots s_{\mathbf{n}-\mathbf{t}-\mathbf{r}+1}) \cdot (s_{\mathbf{h}-\mathbf{t}+1} s_{\mathbf{h}-\mathbf{t}+2} \cdots s_{\mathbf{n}-\mathbf{t}-\mathfrak{s}-1})$.

In particular, $w_{\mathbf{n}-\mathbf{h}+1, \mathfrak{s}} = s_{\mathbf{h}-\mathbf{t}} s_{\mathbf{h}-\mathbf{t}+1} \cdots s_{\mathbf{n}-\mathbf{t}-\mathfrak{s}-1} g_{\mathbf{n}-\mathbf{t}-\mathfrak{s}}$, and $w_{\mathbf{r}, \mathbf{n}-\mathbf{h}} = g_{\mathbf{h}-\mathbf{t}} s_{\mathbf{h}-\mathbf{t}-1} s_{\mathbf{h}-\mathbf{t}-2} \cdots s_{\mathbf{n}-\mathbf{t}-\mathbf{r}+1}$, and $w'_{\mathbf{r}, \mathbf{n}-\mathbf{h}-1} = s_{\mathbf{h}-\mathbf{t}} s_{\mathbf{h}-\mathbf{t}-1} \cdots s_{\mathbf{n}-\mathbf{t}-\mathbf{r}+1}$. Note that the element g_i interchanges $e_i \leftrightarrow f_i$, $e_{2d+1} \leftrightarrow -e_{2d+1}$, and fixes all other elements in the basis.

6.3.3. *Orthogonal Deligne-Lusztig varieties.* For a quadratic space Ω over \mathbb{F} , let $R_{\Lambda^\sharp} \subset \text{OGr}(\mathbf{h} - \mathbf{t}, \Omega_\Lambda)$ be the reduced closed subscheme such that for any field k over \mathbb{F} , we have

$$R_{\Lambda^\sharp}(k) = \{\mathcal{V} \subset \Omega_{\Lambda^\sharp, k} \mid \mathcal{V} \text{ is isotropic, and } \dim_k \mathcal{V} = \mathbf{h} - \mathbf{t}, \text{ and } \dim_k(\mathcal{V} + \Phi(\mathcal{V})) \leq \mathbf{h} - \mathbf{t} + 1\}. \quad (6.12)$$

The following result can be proved by essentially the same way as Theorem 6.3 and we will not repeat it here. We also refer the readers to [HZ25] for a group theoretic approach.

Theorem 6.10. Let $\mathfrak{h}' := \mathfrak{n} - \mathfrak{h}$ and $\mathfrak{t}' := \mathfrak{n} - \mathfrak{t}$, where $\mathfrak{n} = \lfloor n/2 \rfloor$ and $\mathfrak{h} = h/2$ and $\mathfrak{t} = t/2$. We have the following stratification:

$$R_{\Lambda^\sharp} = \begin{cases} \prod_{\substack{0 \leq j \leq \mathfrak{h}' < i \leq \mathfrak{t}' \\ \delta \in \{\pm\}}} X_{P_{ij}}(w_{ij}^\delta) & \text{if } h = n, \\ \left(\prod_{0 \leq j \leq \mathfrak{h}' < i \leq \mathfrak{t}'} X_{P_{ij}}(w_{ij}) \right) \amalg \left(\prod_{\substack{0 \leq j < \mathfrak{n} - \mathfrak{h} < i \leq \mathfrak{t}' \\ \delta \in \{\pm\}}} X_{P_{ij}}(w_{ij}^{\prime\delta}) \right) \amalg X_{P_{\mathfrak{h}'\mathfrak{h}'}}(\text{id}) & \text{if } h = n - 2, \\ \left(\prod_{0 \leq j \leq \mathfrak{h}' < i \leq \mathfrak{t}'} X_{P_{ij}}(w_{ij}) \right) \amalg \left(\prod_{0 \leq j < \mathfrak{h}' < i \leq \mathfrak{n} - \mathfrak{t}} X_{P_{ij}}(w'_{ij}) \right) \amalg X_{P_{\mathfrak{h}'\mathfrak{h}'}}(\text{id}) & \text{otherwise,} \end{cases}$$

such that for each $0 \leq \mathfrak{s} \leq \mathfrak{h}' \leq \mathfrak{r} \leq \mathfrak{t}'$, we have the following closure relations:

$$\begin{aligned} \overline{X_{P_{\mathfrak{r}\mathfrak{s}}}(w_{\mathfrak{r}\mathfrak{s}}^\pm)} &= \begin{cases} \prod_{0 \leq \mathfrak{s} \leq \mathfrak{h}' < i \leq \mathfrak{r}} X_{P_{ij}}(w_{ij}^\pm) & \text{if } h = n, \\ \left(\prod_{\mathfrak{s} \leq j \leq \mathfrak{h}' < i \leq \mathfrak{r}} X_{P_{ij}}(w_{ij}) \right) \amalg \left(\prod_{\substack{0 \leq j < \mathfrak{h}' < i \leq \mathfrak{r} \\ \delta \in \{\pm\}}} X_{P_{ij}}(w_{ij}^{\prime\delta}) \right) \amalg X_{P_{\mathfrak{h}'\mathfrak{h}'}}(\text{id}) & \text{if } h = n - 2, \\ \left(\prod_{\mathfrak{s} \leq j \leq \mathfrak{h}' < i \leq \mathfrak{r}} X_{P_{ij}}(w_{ij}) \right) \amalg \left(\prod_{0 \leq j < \mathfrak{h}' < i \leq \mathfrak{r}} X_{P_{ij}}(w'_{ij}) \right) \amalg X_{P_{\mathfrak{h}'\mathfrak{h}'}}(\text{id}) & \text{otherwise.} \end{cases} \\ \overline{X_{P_{\mathfrak{r}\mathfrak{s}}}(w'_{\mathfrak{r}\mathfrak{s}})} &= \left(\prod_{\mathfrak{s} \leq j < \mathfrak{h}' < i \leq \mathfrak{r}} X_{P_{ij}}(w'_{ij}) \right) \amalg X_{P_{\mathfrak{h}'\mathfrak{h}'}}(\text{id}). \end{aligned}$$

Moreover, R_{Λ^\sharp} is irreducible and normal of dimension $n - (\mathfrak{t} + \mathfrak{h}) - 1$.

6.4. General linear case. In this subsection, we present the necessary results concerning generalized DL varieties associated to the general linear group. Let $\Lambda_1 \subset \Lambda_2$ be vertex lattices of type t_1 and t_2 , resp, such that $t_2 \leq h \leq t_1$. Recall that we use the notation $t_i = 2\mathfrak{t}_i$. Let $V = V_{[\Lambda_1, \Lambda_2]} = \Lambda_2/\Lambda_1$ denote the vector space of dimension $t = \mathfrak{t}_1 - \mathfrak{t}_2$ over \mathbb{F}_q and define $\Omega = \Omega_{[\Lambda_1, \Lambda_2]} := V \otimes_{\mathbb{F}_q} \mathbb{F}$, with Frobenius Φ . We choose a basis $V = \text{Span}(e_{\mathfrak{t}_2+1}, \dots, e_{\mathfrak{t}_1})$. Note that the notations here are not standard. Consider the standard flags \mathcal{F}_\bullet defined by

$$\mathcal{F}_i = \text{Span}_{\mathbb{F}}\{e_{\mathfrak{t}_2+1}, \dots, e_i\}, \quad \mathfrak{t}_2 \leq i \leq \mathfrak{t}_1.$$

In particular, we have $\mathcal{F}_{\mathfrak{t}_2} = (0)$ and $\dim_{\mathbb{F}} \mathcal{F}_i = i - \mathfrak{t}_2$. This pins the choice of maximal torus and Borel $T \subset B \subset G := \text{GL}(\Omega_{[\Lambda_1, \Lambda_2]})$. We use $\Delta^* = \{s_{\mathfrak{t}_2+1}, \dots, s_{\mathfrak{t}_1-1}\}$ to denote the set of corresponding simple reflections in the Weyl group W , such that s_i is the permutation between e_i and e_{i+1} . For each $\mathfrak{t}_2 \leq \mathfrak{s} \leq \mathfrak{h} \leq \mathfrak{r} \leq \mathfrak{t}_1$, let

$$I_{\mathfrak{r}\mathfrak{s}} := \{s_{\mathfrak{t}_2}, \dots, s_{\mathfrak{s}-1}, s_{\mathfrak{r}+1}, \dots, s_{\mathfrak{t}_1-1}\}.$$

Denote by $W_{\mathfrak{r}\mathfrak{s}}$ the subgroup of the Weyl group generated by elements in $I_{\mathfrak{r}\mathfrak{s}}$ and by $P_{\mathfrak{r}\mathfrak{s}}$ the corresponding parabolic subgroup.

The partial flag variety $G/P_{\mathfrak{h}\mathfrak{h}}$ parameterizes the subspaces in Ω of dimension $\mathfrak{t}_1 - \mathfrak{h}$. We consider the subvariety $S_{[\Lambda_1, \Lambda_2]}$ of $G/P_{\mathfrak{h}\mathfrak{h}}$ whose k -points for any field k over \mathbb{F} are given by

$$S_{[\Lambda_1, \Lambda_2]}(k) = \{\mathcal{V} \subset \Omega_{[\Lambda_1, \Lambda_2], k} \mid \dim_k \mathcal{V} = \mathfrak{t}_1 - \mathfrak{h} \quad \text{and} \quad \dim_k(\mathcal{V} \cap \Phi(\mathcal{V})) \geq \mathfrak{t}_1 - \mathfrak{h} - 1\}. \quad (6.13)$$

For each $\mathbf{t}_2 \leq \mathfrak{s} \leq \mathbf{h} \leq \mathbf{r} \leq \mathbf{t}_1$, we let

$$w_{\mathbf{r}\mathfrak{s}} := (s_{\mathbf{h}} \cdots s_{\mathbf{r}-2} s_{\mathbf{r}-1}) \cdot (s_{\mathbf{h}-1} \cdots s_{\mathfrak{s}+2} s_{\mathfrak{s}+1}).$$

The main result is the following:

Proposition 6.11. *We have a stratification for $S_{[\Lambda_1, \Lambda_2]}$*

$$S_{[\Lambda_1, \Lambda_2]} = \coprod_{\mathbf{t}_2 \leq j \leq \mathbf{h} \leq i \leq \mathbf{t}_1} X_{P_{ij}}(w_{ij}),$$

such that for each $\mathbf{t}_2 \leq \mathfrak{s} \leq \mathbf{h} \leq \mathbf{r} \leq \mathbf{t}_1$, we have:

$$\overline{X_{\mathbf{r}\mathfrak{s}}(w_{\mathbf{r}\mathfrak{s}})} = \coprod_{\mathfrak{s} \leq j \leq \mathbf{h} \leq i \leq \mathbf{r}} X_{P_{ij}}(w_{ij}).$$

In particular, $S_{[\Lambda_1, \Lambda_2]}$ is irreducible and normal of dimension $\mathbf{t}_1 - \mathbf{t}_2 - 1$.

Proof. This follows from a similar but much simpler argument to the one in the proof of Theorem 6.3, combined with Proposition 6.12. \square

Proposition 6.12. *For any integers $\mathbf{t}_2 \leq \mathfrak{s} \leq \mathbf{h} < \mathbf{r} \leq \mathbf{t}_1$, the DL variety $X_{P_{\mathbf{r}\mathfrak{s}}}(w_{\mathbf{r}\mathfrak{s}})$ is the subvariety of $G/P_{\mathbf{r}\mathfrak{s}}$ whose k -points for any field k over \mathbb{F} are characterized by:*

$$X_{P_{\mathbf{r}\mathfrak{s}}}(w_{\mathbf{r}\mathfrak{s}})(k) := \left\{ \begin{array}{l} (\mathcal{F}_{\mathbf{t}-\mathbf{r}} \subset \cdots \subset \mathcal{F}_{\mathbf{t}-\mathfrak{s}}) \\ \in G/P_{\mathbf{r}\mathfrak{s}}(k) \end{array} \left| \begin{array}{l} \mathcal{F}_i = \mathcal{F}_{i+1} \cap \Phi(\mathcal{F}_{i+1}) \text{ for } \mathfrak{s} \leq i \leq \mathbf{h} - 1; \\ \mathcal{F}_j = \mathcal{F}_{j-1} + \Phi(\mathcal{F}_{j-1}) \text{ for } \mathbf{h} + 1 \leq j \leq \mathbf{r}; \\ \mathcal{F}_{\mathbf{t}} = \Phi(\mathcal{F}_{\mathbf{t}}) \text{ and } \mathcal{F}_{\mathfrak{s}} = \Phi(\mathcal{F}_{\mathfrak{s}}). \end{array} \right. \right\}.$$

It is smooth of dimension $\mathbf{r} - \mathfrak{s} - 1$. Moreover, the DL variety $X_{P_{[\mathbf{t}_1, \mathbf{t}_2]}}(w_{[\mathbf{t}_1, \mathbf{t}_2]})$ is irreducible.

Proof. The element $w_{\mathbf{r}\mathfrak{s}}$ acts on $V_{[\Lambda_1, \Lambda_2]}$ by

$$e_{\mathfrak{s}+1} \begin{array}{c} \xrightarrow{\quad} \cdots \xrightarrow{\quad} e_{\mathbf{h}-1} \xrightarrow{\quad} e_{\mathbf{h}} \xrightarrow{\quad} e_{\mathbf{h}+1} \xrightarrow{\quad} e_{\mathbf{h}+2} \xrightarrow{\quad} \cdots \xrightarrow{\quad} e_{\mathbf{r}} \\ \xleftarrow{\quad} \cdots \xleftarrow{\quad} e_{\mathbf{h}-1} \xleftarrow{\quad} e_{\mathbf{h}} \xleftarrow{\quad} e_{\mathbf{h}+1} \xleftarrow{\quad} e_{\mathbf{h}+2} \xleftarrow{\quad} \cdots \xleftarrow{\quad} e_{\mathbf{r}} \end{array}$$

One can immediately check that:

$$\begin{array}{ll} \mathcal{F}_i = \mathcal{F}_{i+1} \cap w_{\mathbf{r}\mathfrak{s}}(\mathcal{F}_{i+1}), & i = \mathfrak{s} + 1, \dots, \mathbf{h} - 1; \\ \mathcal{F}_j = \mathcal{F}_{j-1} + w_{\mathbf{r}\mathfrak{s}}(\mathcal{F}_{j-1}), & j = \mathbf{h} + 1, \dots, \mathbf{r} - 1; \\ \mathcal{F}_k = w_{\mathbf{r}\mathfrak{s}}(\mathcal{F}_k), & k = \mathbf{r}, \mathfrak{s}. \end{array}$$

The assertions now follow directly. \square

7. RELATION BETWEEN BRUHAT-TITS STRATA AND DELIGNE-LUSZTIG VARIETIES

In this section, we establish the identification between the BT-strata defined in §5 and certain spaces introduced in §6. While one could construct this isomorphism directly using O_{F_0} -displays as indicated in (7.3), we instead follow the p -divisible group construction to maintain consistency with the existing literature [Vol10, VW11, RTW14, Wu16, HLSY23, KR12, Cho18].

7.1. \mathcal{Z} -strata stratification. Throughout the remainder of this paper, we fix a framing object $\mathbb{X}_{n,\varepsilon}^{[h]}$. Recall from §3.2 the hermitian space \mathbb{V} over F of dimension n with $\text{Hasse}(\mathbb{V}) = -\varepsilon$. Let $\Lambda \subset \mathbb{V}$ be any vertex lattice of type $t(\Lambda) \geq h$. We will construct an isomorphism

$$\Psi_{\mathcal{Z}} : \mathcal{Z}(\Lambda) \cong S_{\Lambda},$$

where S_{Λ} is defined in (6.3). For any \mathbb{F} -algebra R and any R -point $(X, \iota, \lambda, \rho) \in \mathcal{Z}(\Lambda)(R)$ in the \mathcal{Z} -stratum (see Definition 2.2), we have chains of isogenies

$$\rho_{\Lambda, \Lambda^{\sharp}} : X_{\Lambda, R} \xrightarrow{\rho_{\Lambda, X}} X \xrightarrow{\lambda} X^{\vee} \xrightarrow{\rho_{X^{\vee}, \Lambda^{\sharp}}} X_{\Lambda^{\sharp}, R}.$$

Let $\rho_{X, \Lambda^{\sharp}}$ be the composition $\rho_{X^{\vee}, \Lambda^{\sharp}} \circ \lambda$. Applying de Rham realization, we have a sequence of R -modules:

$$D(\rho_{\Lambda, \Lambda^{\sharp}}) : D(X_{\Lambda, R}) \xrightarrow{\rho_{\Lambda, X}} D(X) \xrightarrow{\lambda} D(X^{\vee}) \xrightarrow{\rho_{X^{\vee}, \Lambda^{\sharp}}} D(X_{\Lambda^{\sharp}, R}).$$

By definition, the image $\text{Im}(D(\rho_{\Lambda, \Lambda^{\sharp}}))$ is a locally free direct summand of $D(X_{\Lambda^{\sharp}, R}) = \Lambda^{\sharp} \otimes_{\mathbb{F}_q} R$ of corank $t = t(\Lambda) = 2\mathbf{t}$, such that

$$D(X_{\Lambda^{\sharp}, R}) / \text{Im}(D(\rho_{\Lambda, \Lambda^{\sharp}})) \simeq \Lambda^{\sharp} / \Lambda \otimes_{\mathbb{F}_q} R = \Omega_{\Lambda, R}.$$

Since Λ is a vertex lattice, we have $\ker(\rho_{\Lambda, \Lambda^{\sharp}}) \subseteq X_{\Lambda}[\iota(\pi)]$. This implies that the kernel of the composition

$$\rho_{X, \Lambda^{\sharp}} := \rho_{X^{\vee}, \Lambda^{\sharp}} \circ \lambda : X \longrightarrow X_{\Lambda^{\sharp}, R}$$

lies in $X[\iota(\pi)]$. Therefore, there exists an isogeny $\tilde{\rho}_{X, \Lambda^{\sharp}} : X_{\Lambda^{\sharp}} \rightarrow X$ such that $\tilde{\rho}_{X, \Lambda^{\sharp}} \circ \rho_{X, \Lambda^{\sharp}} = \iota(\pi) : X \rightarrow X$.

Lemma 7.1. *For $(X, \iota, \lambda, \rho) \in \mathcal{Z}(\Lambda)(R)$, we have induced filtrations*

$$\begin{array}{ccccc} D(X_{\Lambda, R}) & \xrightarrow{D(\rho_{\Lambda, \Lambda^{\sharp}})} & D(X_{\Lambda^{\sharp}, R}) & \xrightarrow{D(\tilde{\rho}_{X, \Lambda^{\sharp}})} & D(X) \\ \uparrow & & \uparrow & & \uparrow \\ \text{IID}(X_{\Lambda, R}) & \longrightarrow & \text{IID}(X_{\Lambda^{\sharp}, R}) & \longrightarrow & \text{Fil}(X). \end{array}$$

The preimage $D(\tilde{\rho}_{X, \Lambda^{\sharp}})^{-1}(\text{Fil}(X)) \subseteq D(X_{\Lambda^{\sharp}, R})$ is a locally free direct summand that contains $\text{Im}(D(\rho_{\Lambda, \Lambda^{\sharp}}))$. Moreover, the quotient

$$U(X) := D(\tilde{\rho}_{X, \Lambda^{\sharp}})^{-1}(\text{Fil}(X)) / \text{Im}(D(\rho_{\Lambda, \Lambda^{\sharp}})) \subset \Omega_{\Lambda, R} \tag{7.1}$$

is a locally free isotropic direct summand of rank $\frac{1}{2}(t - h)$.

Proof. It suffices to check the condition on κ -points of $\mathcal{Z}(\Lambda)$, where κ is any algebraically closed field over \mathbb{F} . For the remainder of the proof, fix one such κ and denote $\check{\Lambda} := \Lambda \otimes W_O(\kappa)$ and $\check{\Lambda}^{\sharp} := \Lambda^{\sharp} \otimes W_O(\kappa)$, where $W_O(\kappa)$ denotes the ring of ramified Witt vectors (Note that elsewhere in the paper, we used the notation $\check{\Lambda} := \Lambda \otimes O_{F_0}^{\times}$).

By Proposition 5.4, a point $(X, \iota, \lambda, \rho) \in \mathcal{Z}(\Lambda)(\kappa)$ corresponds to a chain of Dieudonné lattices

$$\check{\Lambda} \subseteq M(X) \subseteq M(X)^{\sharp} \subseteq \check{\Lambda}^{\sharp}.$$

The isogeny $\tilde{\rho}_{X,\Lambda^\sharp}$ induces a map between Dieudonné lattices

$$\check{\Lambda}^\sharp \longrightarrow M(X), \quad x \mapsto \Pi \cdot x.$$

Since $\text{Fil}(X) = M(X)/VM(X)$, we have

$$M(\tilde{\rho}_{X,\Lambda^\sharp})^{-1}(\text{Fil}(X)) = \Pi^{-1}VM/\pi_0\check{\Lambda}^\sharp = \tau^{-1}M/\pi_0\check{\Lambda}^\sharp.$$

Moreover, since $\check{\Lambda} \subseteq M(X)$, it follows that $\check{\Lambda} = \tau^{-1}(\check{\Lambda}) \subseteq \tau^{-1}(M(X))$. Therefore

$$\text{Im}(M((\rho_\Lambda)_R)) = \check{\Lambda}/\pi_0\check{\Lambda}^\sharp \subseteq M(\tilde{\rho}_{X,\Lambda^\sharp})^{-1}(\text{Fil}(X)).$$

The quotient

$$U(X) \cong \tau^{-1}(M)/\check{\Lambda} = \Phi^{-1}(M/\check{\Lambda}) \subset \Lambda^\sharp/\Lambda \otimes \kappa = \Omega_{\Lambda,\kappa}$$

has dimension $\frac{1}{2}(t-h)$, where Φ is the Frobenius in $\Omega_{\Lambda,\kappa}$.

For any $x \in M$ and $y \in \check{\Lambda}^\sharp$, we have $\langle \bar{x}, \bar{y} \rangle_{\check{\Lambda}^\sharp/\check{\Lambda}} = 0$ if and only if $\pi\langle x, y \rangle \in (\pi) \subset O_F \otimes_{O_{F_0}} W_O(\kappa)$. This is equivalent to $y \in M^\sharp$, which implies that $M^\sharp/\check{\Lambda}$ is the orthogonal complement of $M/\check{\Lambda}$ in $\Omega_{\Lambda,\kappa}$. In particular, $\Phi^{-1}(M/\check{\Lambda})$ is an isotropic subspace. \square

For the remainder of our discussion, recall that $\Omega_\Lambda = \Lambda^\sharp/\Lambda \otimes \mathbb{F}$, when $t(\Lambda) \neq h$. The partial flag variety $\text{Sp}(\Omega_\Lambda)/P_{\mathbf{n}\mathbf{h}}$ parameterizes isotropic subspaces of dimension $\mathbf{t} - \mathbf{h}$, and S_Λ in §6.2 is defined as a closed subvariety whose k -points for any field k over \mathbb{F} are given by

$$S_\Lambda(k) = \{\mathcal{V} \subset \Omega_{\Lambda,k} \mid \mathcal{V} \text{ is isotropic, and } \dim_k \mathcal{V} = \mathbf{t} - \mathbf{h}, \text{ and } \dim_k(\mathcal{V} \cap \Phi(\mathcal{V})) \geq \mathbf{t} - \mathbf{h} - 1\}.$$

When $t(\Lambda) = h$, the space S_Λ consists of the zero-dimensional subspace. Hence, the stratification in Theorem 6.3 degenerates into $S_\Lambda = X_{P_{\mathbf{n}\mathbf{h}}}(\text{id})$, which is irreducible of dimension 0. By Lemma 7.1, we define the map

$$\Psi_{\mathcal{Z}} : \mathcal{Z}(\Lambda) \longrightarrow \text{Sp}(\Omega_\Lambda)/P_{\mathbf{n}\mathbf{h}}, \quad (X, \iota, \lambda, \rho) \mapsto U(X).$$

Lemma 7.2. *Let κ be any perfect field over \mathbb{F} . The map $\Psi_{\mathcal{Z}}$ defines a bijection between $\mathcal{Z}(\Lambda)(\kappa)$ and $S_\Lambda(\kappa)$, hence it factors through S_Λ , which we will still denote by $\Psi_{\mathcal{Z}}$.*

Proof. By Dieudonné theory, a point $z \in \mathcal{Z}(\Lambda)(\kappa)$ corresponds to a lattice M satisfying

$$\check{\Lambda} \begin{array}{c} \mathbf{n} - \mathbf{h} - \mathbf{t} \\ \subset \\ \end{array} M \begin{array}{c} \mathbf{t} \\ \subset \\ \end{array} M^\sharp \begin{array}{c} \mathbf{n} - \mathbf{h} - \mathbf{t} \\ \subset \\ \end{array} \check{\Lambda}^\sharp,$$

see Proposition 5.4. The wedge condition implies $M \cap \tau(M) \stackrel{\leq 1}{\subset} M$, which is equivalent to

$$\Phi^{-1}(M/\check{\Lambda}) \cap M/\check{\Lambda} \stackrel{\leq 1}{\subset} \Phi^{-1}(M/\check{\Lambda}).$$

Therefore, $\Psi_{\mathcal{Z}}(z) \in S_\Lambda(\kappa)$.

Conversely, let $U \in S_\Lambda(\kappa)$ be any geometric point. The dual space $U^\perp \subset \Omega_\Lambda$ is a subspace of dimension $\mathbf{n} - \mathbf{h} + \mathbf{t}$ containing U . By the definition of S_Λ we have

$$U \cap \Phi(U) \stackrel{\leq 1}{\subset} U. \tag{7.2}$$

We denote by $M \subset \check{\Lambda}^\sharp$ (resp. $M_{\Phi(U)^\sharp} \subset \check{\Lambda}^\sharp$) the preimage of $\Phi(U)$ (resp. $\Phi(U)^\sharp$) under the projection map $\check{\Lambda}^\sharp \rightarrow \check{\Lambda}^\sharp/\check{\Lambda}$.

Let M^\sharp be the hermitian dual of the lattice $M \subset M(\mathbb{X}) \otimes W_O(\kappa)[1/\pi_0]$, which agrees with the symplectic dual in the same space, see §3.2. Since $\check{\Lambda} \subseteq M$, we have $M^\sharp \subseteq \check{\Lambda}^\sharp$. Therefore, we have

$$\begin{aligned} M^\sharp &= \{x \in M(\mathbb{X}) \otimes W_O(\kappa)[1/\pi_0] \mid \langle x, y \rangle \in W_O(\kappa) \text{ for all } y \in M\}, \\ &= \{x \in \check{\Lambda}^\sharp \mid \pi \langle x, y \rangle \equiv 0 \pmod{\pi_0} \text{ for all } y \in M\}, \\ &= \{x \in \check{\Lambda}^\sharp \mid \langle \bar{x}, \bar{y} \rangle = 0, \text{ for all } y \in M\}, \\ &= \{x \in \check{\Lambda}^\sharp \mid \langle \bar{x}, \bar{y} \rangle = 0 \text{ for all } y \in \Phi(U)\}, \\ &= \{x \in \check{\Lambda}^\sharp \mid \bar{x} \in \Phi(U)^\perp \subset \Omega_{\Lambda, \kappa}\} = M_{\Phi(U)^\sharp}. \end{aligned}$$

Since $U \stackrel{h}{\subset} U^\sharp$, we have $M \stackrel{h}{\subset} M^\sharp$. The relation (7.2) implies that $M \cap \tau(M) \stackrel{\leq 1}{\subset} M$. Next, we have

$$\Pi M \subset \Pi \check{\Lambda}^\sharp \subseteq \check{\Lambda} = \tau^{-1} \check{\Lambda} \subseteq \tau^{-1} M \subseteq \tau^{-1} \check{\Lambda}^\sharp = \check{\Lambda}^\sharp \subseteq \Pi^{-1} \Lambda \subseteq \Pi^{-1} M.$$

Hence the lattice M defines a point in $\mathcal{N}_{n, \varepsilon}^{[h]}(\kappa)$. We conclude that $M \in \mathcal{Z}(\Lambda)(\kappa)$. \square

Theorem 7.3. *Let Λ be a vertex lattice of type $t \geq h$ in \mathbb{V} . Then the morphism $\Psi_{\mathcal{Z}}$ defines an isomorphism $\mathcal{Z}(\Lambda) \rightarrow S_\Lambda$.*

Proof. By Lemma 7.2, we know that $\Psi_{\mathcal{Z}}$ is a bijection for any perfect field κ over \mathbb{F} . We will show in Theorem 7.5 that this property extends to any field k over \mathbb{F} . Given this result and the fact that S_Λ is normal (Theorem 6.3) together with the properness of $\Psi_{\mathcal{Z}}$ (as it is a morphism between projective varieties), it follows from Zariski's main theorem that $\Psi_{\mathcal{Z}}$ is an isomorphism. \square

Proposition 7.4. *The isomorphism $\Phi_{\mathcal{Z}}$ is compatible with inclusions of vertex lattices.*

Proof. We can check this on κ -valued points for any perfect field κ over \mathbb{F} , where it is obvious from the definition of the map $\Psi_{\mathcal{Z}}$. \square

To extend Lemma 7.2 to an arbitrary field k over \mathbb{F} , we use O_{F_0} -displays. For any $R \in \text{Nilp}_{O_{\mathbb{F}}}$ and R -point $(X, \iota, \lambda, \rho) \in \mathcal{Z}(\Lambda)(R)$, let $\mathcal{P}(X) = \mathcal{P} = (P, Q, \mathbf{F}, \dot{\mathbf{F}})$ be the associated O_{F_0} -display (see §3.1). Let $\mathcal{P}_\Lambda = (P_\Lambda, Q_\Lambda, \mathbf{F}, \dot{\mathbf{F}})$ and $\mathcal{P}_{\Lambda^\sharp} = (P_{\Lambda^\sharp}, Q_{\Lambda^\sharp}, \mathbf{F}, \dot{\mathbf{F}})$ denote the displays $\mathcal{P}(X_{\Lambda^\sharp})$ and $\mathcal{P}(X_\Lambda)$, resp. The isogeny $X \rightarrow X_{\Lambda^\sharp}$ induces an O_{F_0} -display morphism $\rho_{X, \Lambda^\sharp} : \mathcal{P}(X) \rightarrow \mathcal{P}_{\Lambda^\sharp}$. Define

$$\mathcal{U}(X) := \ker[\Pi : P_{\Lambda^\sharp} \rightarrow P_{\Lambda^\sharp} \twoheadrightarrow P_{\Lambda^\sharp}/\rho_{X, \Lambda^\sharp}(Q)]$$

This defines a $W_O(R)$ -submodule of P_{Λ^\sharp} .

When $R = \kappa$ is an algebraically closed field, by Dieudonné theory, we have $P = M(X)$ and $\text{VM}(X) = Q \cong \rho_{X, \Lambda^\sharp}(Q) \subseteq P_{\Lambda^\sharp} = \Lambda^\sharp \otimes W_O(\kappa)$. Hence

$$\begin{aligned} \mathcal{U}(X) &= \{v \in \Lambda^\sharp \otimes W_O(\kappa) \mid \Pi(v) \in \text{VM}(X)\} \\ &= \{v \in \Lambda^\sharp \otimes W_O(\kappa)[1/\pi] \mid v \in \Pi^{-1} \text{VM}(X) \cap \Lambda^\sharp \otimes W_O(\kappa)\} \\ &= \Pi^{-1} \text{VM}(X) = \tau^{-1} M(X) \subseteq \Lambda^\sharp \otimes W_O(\kappa). \end{aligned}$$

Therefore, we can re-define

$$U(X) := \mathcal{U}(X)/\rho_{\Lambda, \Lambda^\sharp}(P_\Lambda) \subset P_{\Lambda^\sharp}/\rho_{\Lambda, \Lambda^\sharp}(P_\Lambda) = \Lambda^\sharp/\Lambda \otimes W_O(R) = \Lambda^\sharp/\Lambda \otimes R. \quad (7.3)$$

This construction agrees with our previous one, as verified by checking geometric points.

For any field k over \mathbb{F} , the natural map $k \rightarrow \bar{k}$ induces an embedding $W_O(\kappa) \rightarrow W_O(\bar{k})$, making $W_O(\kappa)$ an integral domain as a subring of $W_O(\bar{k})$. Consequently, for any $X \in \mathcal{Z}(\Lambda)(k)$, the induced maps $P_\Lambda \rightarrow P \rightarrow P_{\Lambda^\sharp}$ are injective.

Proposition 7.5. *Let k be any field over \mathbb{F} . The map $\Psi_{\mathcal{Z}}$ defines a bijection between $\mathcal{Z}(\Lambda)(k)$ and $S_\Lambda(k)$.*

Proof. To construct the inverse to $\Psi_{\mathcal{Z}}(k)$, consider a k -point $U \in S_\Lambda(k)$ defining an isotropic subspace $U \stackrel{t-\hbar}{\subset} \Omega_{\Lambda,k}$. Via the isomorphisms $\Omega_{\Lambda,k} = \Lambda^\sharp/\Lambda \otimes k \cong \Lambda^\sharp/\Lambda \otimes W_O(\kappa) = P_{\Lambda^\sharp}/P_\Lambda$, the subspace U lifts to a lattice

$$P_\Lambda \stackrel{t-\hbar}{\subseteq} \mathcal{U} \subseteq P_{\Lambda^\sharp} \quad (7.4)$$

Let $Q := \Pi\mathcal{U} \subset \mathcal{P}_{\Lambda^\sharp}$ and let P be the submodule of $\mathcal{P}_{\Lambda^\sharp}$ generated by $\dot{\mathbf{F}}(Q)$. Then $\mathcal{P} := (P, Q, \mathbf{F}, \dot{\mathbf{F}})$ defines an O_{F_0} -display over k . By (7.4), we have

$$\Pi P_\Lambda = Q_\Lambda \subset Q \subset \Pi P_{\Lambda^\sharp} = Q_{\Lambda^\sharp}$$

which yields natural inclusions $\mathcal{P}_\Lambda \hookrightarrow \mathcal{P} \hookrightarrow \mathcal{P}_{\Lambda^\sharp}$. The definition of S_Λ ensures $\text{BT}(\mathcal{P})$ defines the subspace U in $\mathcal{Z}(\Lambda)(k)$, providing the inverse of $\Psi_{\mathcal{Z}}(k)$. \square

Remark 7.6. An important observation from the proof of Proposition 7.5 is that while $Q \subset P_\Lambda$ uniquely determines $P \subset P_\Lambda$ (since $P \stackrel{=}{=} V^{-1}Q$), the converse fails. Indeed, by [ACZ16, Lem. 2.2], Verschiebung can only be constructed up to a twist

$$V^\sharp : P \rightarrow W_O(R) \otimes_{\sigma, W_O(R)} P.$$

Thus $Q \stackrel{=}{=} VP$ can be recovered from P only up to a Frobenius twist, which fails to be an isomorphism over non-perfect fields k . Consequently, an alternative map via

$$\mathcal{Z}(\Lambda) \rightarrow S_\Lambda \quad X \mapsto \mathcal{U}(X)^{\text{fake}} := \text{Im}[P \rightarrow P_{\Lambda^\sharp}]$$

would not yield a isomorphism $\mathcal{Z}(\Lambda) \cong S_\Lambda$, but differing by a Frobenius twist.

7.2. \mathcal{Y} -strata stratification. Let $\Lambda \subset \mathbb{V}$ be any vertex lattice of type $t(\Lambda) \leq h$. We sketch the construction (similar to §7.1) of the isomorphism

$$\Psi_{\mathcal{Y}} : \mathcal{Y}(\Lambda) \simeq R_{\Lambda^\sharp},$$

where R_{Λ^\sharp} is defined in (6.12). For any \mathbb{F} -algebra R and any R -point $(X, \iota, \lambda, \rho) \in \mathcal{Y}(\Lambda^\sharp)(R)$ in the \mathcal{Y} -stratum, see Definition 2.2, we have an isogeny $\rho_{\Lambda^\sharp, X^\vee} : X_{\Lambda^\sharp} \rightarrow X^\vee$. This implies that the quasi-isogeny $\rho_{X^\vee, \pi^{-1}\Lambda} : X^\vee \rightarrow X_\Lambda$ defined by $\rho_{X^\vee, \pi^{-1}\Lambda} \circ \rho_{\Lambda^\sharp, X^\vee} = \rho_{\Lambda^\sharp, \pi^{-1}\Lambda}$ is an isogeny.

Since Λ is a vertex lattice, we have $\ker[\rho_{X^\vee, \pi^{-1}\Lambda}] \subset X^\vee[\pi]$. Therefore, there exists an isogeny $\tilde{\rho}_{X^\vee, \pi^{-1}\Lambda} : X_{\pi^{-1}\Lambda} \rightarrow X^\vee$ such that $\tilde{\rho}_{X^\vee, \pi^{-1}\Lambda} \circ \rho_{X^\vee, \pi^{-1}\Lambda} = \iota(\pi) : X^\vee \rightarrow X^\vee$.

The proof of the following lemma is the same as that of Lemma 7.1 and we leave the details to the reader.

Lemma 7.7. For $X \in \mathcal{Y}(\Lambda^\sharp)(R)$, we have induced filtrations

$$\begin{array}{ccccc} D(X_{\Lambda^\sharp}) & \xrightarrow{D(\rho_{\Lambda^\sharp, \pi^{-1}\Lambda})} & D(X_\Lambda) & \xrightarrow{D(\tilde{\rho}_{X^\vee, \pi^{-1}\Lambda})} & D(X^\vee) \\ \uparrow & & \uparrow & & \uparrow \\ \Pi D(X_{\Lambda^\sharp}) & \longrightarrow & \Pi D(X_\Lambda) & \longrightarrow & \text{Fil}(X^\vee). \end{array}$$

The preimage $D(\tilde{\rho}_{X^\vee, \pi^{-1}\Lambda})^{-1}(\text{Fil}(X^\vee)) \subseteq D(X_{\pi^{-1}\Lambda, R})$ is a locally free direct summand that contains $\text{Im}(D(\rho_{\Lambda^\sharp, \pi^{-1}\Lambda}))$. Moreover the quotient

$$U(X) := M(\tilde{\rho}_{X^\vee, \pi^{-1}\Lambda})^{-1}(\text{Fil}(X))/\text{Im}M(\rho_{\Lambda^\sharp, \pi^{-1}\Lambda}) \subset \Omega_{\Lambda^\sharp, R} \quad (7.5)$$

is a locally free isotropic direct summand of rank $\frac{1}{2}(h-t)$. \square

For the remainder of our discussion, recall that $\Omega_{\Lambda^\sharp} = \Lambda/\pi\Lambda^\sharp \otimes \mathbb{F}$ when $t(\Lambda) \neq h$. The variety R_{Λ^\sharp} is defined in (6.12) whose k -points for any field k over \mathbb{F} are given by

$$R_{\Lambda^\sharp}(k) = \{\mathcal{V} \subset \Omega_{\Lambda^\sharp, k} \mid \mathcal{V} \text{ is isotropic, and } \dim_k \mathcal{V} = \mathfrak{h} - \mathfrak{t}, \text{ and } \dim_k(\mathcal{V} + \Phi(\mathcal{V})) \leq \mathfrak{h} - \mathfrak{t} + 1\}. \quad (7.6)$$

When $t(\Lambda) = h$, the space R_{Λ^\sharp} consists of the zero-dimensional subspace, hence the stratification in Theorem 6.10 degenerates into $R_{\Lambda^\sharp} = X_{P_{[\mathfrak{h}, \mathfrak{h}]}}(\text{id})$, which is irreducible of dimension 0.

By Lemma 7.7, we define the map $\Psi_{\mathcal{Y}} : \mathcal{Y}(\Lambda^\sharp) \rightarrow R_{\Lambda^\sharp}$.

Lemma 7.8. Let κ be any perfect field over \mathbb{F} . The map $\Psi_{\mathcal{Y}}$ defines a bijection between $\mathcal{Y}(\Lambda^\sharp)(\kappa)$ and $R_{\Lambda^\sharp}(\kappa)$.

Proof. By Dieudonné theory, a point $y \in \mathcal{Y}(\Lambda^\sharp)(\kappa)$ corresponds to a lattice M satisfying

$$\pi\check{\Lambda}^\sharp \overset{\mathfrak{h}-\mathfrak{t}}{\subset} \pi M^\sharp \overset{n-h}{\subset} M \overset{\mathfrak{h}-\mathfrak{t}}{\subset} \check{\Lambda},$$

see Proposition 5.4. The wedge condition implies $M \cap \tau(M) \overset{\leq 1}{\subset} M$, which is equivalent to

$$M^\sharp \cap \tau(M^\sharp) \overset{\leq 1}{\subset} M^\sharp$$

by Lemma 5.1. Hence, we have

$$\Phi^{-1}(\pi M^\sharp / \pi \check{\Lambda}^\sharp) \cap \pi M^\sharp / \pi \check{\Lambda}^\sharp \overset{\leq 1}{\subset} \Phi^{-1}(\pi M^\sharp / \pi \check{\Lambda}^\sharp).$$

Therefore $\Phi_{\mathcal{Y}}(y) \in R_{\Lambda^\sharp}(\kappa)$.

Conversely, let $U \in R_{\Lambda^\sharp}(\kappa)$, $\pi_\Lambda^{-1}(U)$ be any closed point. Let $\pi_\Lambda : \Lambda \rightarrow \Lambda/\pi\Lambda^\sharp$ denote the natural projection map. Then the preimage $\pi_\Lambda^{-1}(U)$ defines a point in $\mathcal{Y}(\Lambda^\sharp)$. Indeed, $U \cap \Phi(U) \overset{\leq 1}{\subset} U$ corresponds to $M^\sharp \cap \tau(M^\sharp) \overset{\leq 1}{\subset} M^\sharp$ and the condition that U is isotropic corresponds to the condition $\pi M^\sharp \subset M$ by similar computations as in the proof of Lemma 7.2. It is straightforward to check that the two maps are inverses to each other. \square

Now the same proof as that of Proposition 7.3 proves the following.

Theorem 7.9. Let $\Lambda \in \mathcal{L}_{\mathcal{Y}}$. Then the morphism $\Psi_{\mathcal{Y}}$ defines an isomorphism $\mathcal{Y}(\Lambda^\sharp) \rightarrow R_{\Lambda^\sharp}$. \square

7.3. Intersection between \mathcal{Y} -strata and \mathcal{Z} -strata. We now discuss the intersection of \mathcal{Y} -strata and \mathcal{Z} -strata. Let $\Lambda_1 \in \mathcal{L}_{\mathcal{Z}}$ and $\Lambda_2 \in \mathcal{L}_{\mathcal{Y}}$ be two vertex lattices such that $\Lambda_1 \subseteq \Lambda_2$. Recall from (6.13) that we defined the variety $S_{[\Lambda_1, \Lambda_2]}$ whose k -points for any field k over \mathbb{F} are given by

$$S_{[\Lambda_1, \Lambda_2]}(k) = \{\mathcal{V} \subset \Omega_{[\Lambda_1, \Lambda_2], k} \mid \dim_k \mathcal{V} = \mathbf{t}_1 - \mathbf{h} \quad \text{and} \quad \dim_k(\mathcal{V} \cap \Phi(\mathcal{V})) \geq \mathbf{t}_1 - \mathbf{h} - 1\}.$$

The main result is:

Proposition 7.10. *Let $\Lambda_1 \in \mathcal{L}_{\mathcal{Z}}$ and let $\Lambda_2 \in \mathcal{L}_{\mathcal{Y}}$ be vertex lattices satisfying $\Lambda_1 \subseteq \Lambda_2$. The restriction of $\Psi_{\mathcal{Z}}$ to the intersection $\mathcal{Z}(\Lambda_1) \cap \mathcal{Y}(\Lambda_2^{\sharp})$ defines an isomorphism:*

$$\Psi_{\mathcal{Z} \cap \mathcal{Y}} : \mathcal{Z}(\Lambda_1) \cap \mathcal{Y}(\Lambda_2^{\sharp}) \cong S_{[\Lambda_1, \Lambda_2]}$$

Proof. By Proposition 7.3, we have a natural isomorphism:

$$\Phi_{\mathcal{Z}} : \mathcal{Z}(\Lambda_1) \xrightarrow{\sim} S_{\Lambda_1}, \quad X \mapsto U(X).$$

The inclusion $\Lambda_1 \subseteq \Lambda_2 \subseteq \Lambda_2^{\sharp} \subseteq \Lambda_1^{\sharp}$ implies that the subspace $(0) \subseteq \Lambda_2/\Lambda_1 \subseteq V_{\Lambda_1}$ is isotropic. Let κ be a fixed algebraically closed field over \mathbb{F} . Let $W := \check{\Lambda}_2/\check{\Lambda}_1 \subset \Omega_{\Lambda_1, \kappa}$. Via the isomorphism $\Psi_{\mathcal{Z}}$, the intersection $\mathcal{Z}(\Lambda_1) \cap \mathcal{Y}(\Lambda_2^{\sharp})$ corresponds to the subvariety of S_{Λ_1} which parameterizes all subspaces

$$\{\mathcal{V} \subset \Omega_{\Lambda_1, \kappa} \mid \mathcal{V} \subseteq W \text{ is isotropic, } \dim \mathcal{V} = \mathbf{t}_1 - \mathbf{h} \quad \text{and} \quad \dim(\mathcal{V} \cap \Phi(\mathcal{V})) \geq \mathbf{t}_1 - \mathbf{h} - 1\}.$$

Since W is isotropic, any subspace $U \subset W$ is automatically isotropic. Moreover, since $\Lambda_2 \subset \mathbb{V}$, is closed under Frobenius, this subvariety is isomorphic to the subvariety of the partial flag variety that parameterizes

$$\{\mathcal{V} \subset \Omega_{\Lambda_1, \kappa} \mid \mathcal{V} \subseteq W \quad \text{with} \quad \dim \mathcal{V} = \mathbf{t}_1 - \mathbf{h} \quad \text{and} \quad \dim(\mathcal{V} \cap \Phi(\mathcal{V})) \geq \mathbf{t}_1 - \mathbf{h} - 1\}.$$

By definition, this is isomorphic to $S_{[\Lambda_1, \Lambda_2]}$. □

7.4. Proof of Proposition 2.14. We recall the statement:

Proposition. (1) *By restriction, the isomorphism $\Psi_{\mathcal{Z}}$ induces isomorphisms:*

- (i) $\mathcal{Z}(\Lambda_1) \setminus \mathcal{Y} \cong \prod_{i=0}^{\mathbf{h}-1} X_{P_{[i, \mathbf{h}]}}(w_{[i, \mathbf{h}]})$;
- (ii) $\mathcal{Z}(\Lambda_1) \cap \mathcal{Y} \cong \left(\prod_{0 \leq j < \mathbf{h} < i \leq \mathbf{t}} X_{P_{ij}}(w_{ij}) \right) \amalg \left(\prod_{0 \leq j < \mathbf{h} < i \leq \mathbf{t}} X_{P_{ij}}(w'_{ij}) \right) \amalg X_{P_{\mathbf{h}\mathbf{h}}}(\text{id})$.

(2) *Similarly, by restriction, the isomorphism $\Psi_{\mathcal{Y}}$ induces isomorphisms:*

- (i) $\mathcal{Y}(\Lambda_2^{\sharp}) \setminus \mathcal{Z} \cong \prod_{i=0}^{\mathbf{h}'-1} X_{P_{i, \mathbf{h}'}}(w_{i, \mathbf{h}'})$;
- (ii) $\mathcal{Y}(\Lambda_2^{\sharp}) \cap \mathcal{Z} \cong \begin{cases} \emptyset & \text{if } h = n, \\ \left(\prod_{0 \leq j < \mathbf{h}' < i \leq \mathbf{t}'} X_{P_{ij}}(w_{ij}) \right) \amalg \left(\prod_{\substack{0 \leq j < \mathbf{h}' < i \leq \mathbf{t}' \\ \delta \in \{\pm\}}} X_{P_{ij}}(w'_{ij}{}^{\delta}) \right) \amalg X_{P_{\mathbf{h}'\mathbf{h}'}}(\text{id}) & \text{if } h = n - 2, \\ \left(\prod_{0 \leq j < \mathbf{h}' < i \leq \mathbf{t}'} X_{P_{ij}}(w_{ij}) \right) \amalg \left(\prod_{0 \leq j < \mathbf{h}' < i \leq \mathbf{t}'} X_{P_{ij}}(w'_{ij}) \right) \amalg X_{P_{\mathbf{h}'\mathbf{h}'}}(\text{id}) & \text{otherwise.} \end{cases}$

Proof of Proposition 2.14. Let κ be a perfect field over \mathbb{F} and let $X \in \mathcal{Z}(\Lambda_1)(\kappa)$. Denote by $M = M(X)$ its associated Dieudonné module.

Via the map $\Psi_{\mathcal{Z}}$, the sum $M + \tau(M)$ induces a subspace $U(X) + \Phi(U(X))$ in $\Omega_{\Lambda} = \check{\Lambda}^{\sharp}/\check{\Lambda}$. This subspace is isotropic if and only if $\Phi(U(X)) \subseteq U(X)^{\perp}$. Equivalently, this occurs if and only if $\tau(M) \subseteq M^{\sharp}$, which in turn holds precisely when $X \in \mathcal{Y}(\kappa)$. The claimed decomposition then follows from the moduli description, see also Remark 6.5, particularly (6.8). Part (2) follows by the same argument as part (1). \square

Remark 7.11. Recall from Remark 6.8 that we have the following first-step decomposition:

$$S_{\Lambda} = X_{P_{\hbar\hbar}}(w_{\Lambda}) \amalg X_{P_{\hbar\hbar}}(w'_{\Lambda}) \amalg X_{P_{\hbar\hbar}}(\text{id}),$$

where we can describe these DL varieties as follows:

$$\begin{aligned} X_{P_{\hbar\hbar}}(w_{\Lambda}) &= \{\mathcal{V} \in S_{\Lambda} \mid \mathcal{V} \neq \Phi(\mathcal{V}) \text{ and } \mathcal{V} + \Phi(\mathcal{V}) \text{ is not isotropic}\}; \\ X_{P_{\hbar\hbar}}(w'_{\Lambda}) &= \{\mathcal{V} \in S_{\Lambda} \mid \mathcal{V} \neq \Phi(\mathcal{V}) \text{ and } \mathcal{V} + \Phi(\mathcal{V}) \text{ is isotropic}\}; \\ X_{P_{\hbar\hbar}}(\text{id}) &= \{\mathcal{V} \in S_{\Lambda} \mid \mathcal{V} = \Phi(\mathcal{V})\}. \end{aligned}$$

From the proof of Proposition 2.14, we established that for any $M(X) \in \mathcal{Z}(\Lambda_1)$ with corresponding isotropic subspace $U(X) \in S_{\Lambda}$, we have $\tau(M) \subseteq M^{\sharp}$ if and only if $U(X) + \Phi(U(X))$ is a totally isotropic subspace. Therefore, via the isomorphism $\Psi_{\mathcal{Z}}$, we have:

$$\mathcal{Z}(\Lambda_1) \setminus \mathcal{Y} \cong X_{P_{\hbar\hbar}}(w_{\Lambda}), \quad \text{and} \quad \mathcal{Z}(\Lambda_1) \cap \mathcal{Y} \cong X_{P_{\hbar\hbar}}(w'_{\Lambda}) \amalg X_{P_{\hbar\hbar}}(\text{id}).$$

Similar results exist for \mathcal{Y} -strata.

Remark 7.12. We remark that when $h = n - 2$, part (2) of Proposition 2.14 agrees with the computation in the local model. More precisely, the special fiber of the local model $\mathcal{M}_n^{[n-2]}$ (see Definition 4.1 for our notation) consists of three irreducible components:

$$\overline{\mathcal{M}_n^{[n-2]}} = \mathcal{Y} \cup \mathcal{Z}^+ \cup \mathcal{Z}^-.$$

See [Yu19, §1.5] for computations. Following the notation in loc. cit, \mathcal{Y} is defined as the vanishing locus of x_1, x_2 , while \mathcal{Z}^+ (resp. \mathcal{Z}^-) is defined by the vanishing locus of the first (resp. second) row of Y , where the matrix Y is defined in p.22 of loc. cit. In particular, the intersection $\mathcal{Z}^+ \cap \mathcal{Z}^-$ is the vanishing locus of the entire matrix Y , which is the worst point.

The closure $\overline{X_{P_{\mathfrak{t}'_0}}(w_{\mathfrak{t}'_0}^{\pm})}$ can be regarded as $\mathcal{Y}(\Lambda_2^{\sharp}) \cap \mathcal{Z}^{\pm}$ (where we use \mathcal{Z}^{\pm} to denote both the KR strata and their corresponding components in the local model). Moreover, we have: $\overline{X_{P_{\mathfrak{t}'_0}}(w_{\mathfrak{t}'_0}^{\pm})} \cap \overline{X_{P_{\mathfrak{t}'_0}}(w_{\mathfrak{t}'_0}^{\mp})} = X_{P_{\hbar\hbar'}}(\text{id})$, which aligns with the fact that $\mathcal{Z}^+ \cap \mathcal{Z}^-$ is the worst point.

REFERENCES

- [ACZ16] Tobias Ahsendorf, Chuangxun Cheng, and Thomas Zink. \mathcal{O} -displays and π -divisible formal \mathcal{O} -modules. *Journal of Algebra*, 457:129–193, 2016.
- [BR06] Cédric Bonnafé and Raphaël Rouquier. On the irreducibility of Deligne–Lusztig varieties. *Comptes Rendus Mathématique*, 343(1):37–39, 2006.
- [Cho18] Sungyoon Cho. The basic locus of the unitary shimura variety with parahoric level structure, and special cycles. *arXiv preprint 1807.09997*, 2018.

- [Con94] Aldo Conca. Divisor class group and canonical class of determinantal rings defined by ideals of minors of a symmetric matrix. *Arch. Math. (Basel)*, 63(3):216–224, 1994.
- [FHI24] Maria Fox, Benjamin Howard, and Naoki Imai. Rapoport-Zink spaces of type $\mathrm{GU}(2, n - 2)$. *arXiv preprint 2308.03816*, 2024.
- [Gö1] Ulrich Görtz. On the flatness of models of certain Shimura varieties of PEL-type. *Math. Ann.*, 321(3):689–727, 2001.
- [GHN19] Ulrich Görtz, Xuhua He, and Sian Nie. Fully Hodge-Newton decomposable Shimura varieties. *Peking Math. J.*, 2(2):99–154, 2019.
- [GHN24] Ulrich Görtz, Xuhua He, and Sian Nie. Basic loci of Coxeter type with arbitrary parahoric level. *Canad. J. Math.*, 76(1):126–172, 2024.
- [GY10] Ulrich Görtz and Chia-Fu Yu. Supersingular Kottwitz–Rapoport strata and Deligne–Lusztig varieties. *Journal of the Institute of Mathematics of Jussieu*, 9(2):357–390, 2010.
- [He16] Xuhua He. Kottwitz-Rapoport conjecture on unions of affine Deligne-Lusztig varieties. *Ann. Sci. Éc. Norm. Supér. (4)*, 49(5):1125–1141, 2016.
- [HLS24] Qiao He, Yu Luo, and Yousheng Shi. Regular models of ramified unitary Shimura varieties at maximal parahoric level. *arXiv preprint arXiv:2410.04500*, 2024.
- [HLSY23] Qiao He, Chao Li, Yousheng Shi, and Tonghai Yang. A proof of the Kudla-Rapoport conjecture for Krämer models. *Invent. Math.*, 234(2):721–817, 2023.
- [HLZ19] Xuhua He, Chao Li, and Yihang Zhu. Fine Deligne-Lusztig varieties and arithmetic fundamental lemmas. *Forum Math. Sigma*, 7:e47, 55, 2019.
- [Hoe10] Maarten Hoeve. Stratifications on moduli spaces of abelian varieties and Deligne-Lusztig varieties. *Thesis, fully internal, Universiteit van Amsterdam*, 2010.
- [HZ25] Qiao He and Rong Zhou. Basic locus of GSpin Rapoport-Zink space with maximal parahoric level. *In preparation*, 2025.
- [KR12] Steven Kudla and Michael Rapoport. Notes on special cycles. *unpublished note*, 2012.
- [KRZ23] Stephen Kudla, Michael Rapoport, and Thomas Zink. On the p -adic uniformization of unitary shimura curves. *arXiv preprint arXiv:2007.05211*, 2023.
- [LRZ24] Chao Li, Michael Rapoport, and Wei Zhang. Quasi-canonical AFL and arithmetic transfer conjectures at parahoric levels. *arXiv preprint 2404.02214*, 2024.
- [LRZ25] Yu Luo, Michael Rapoport, and Wei Zhang. More arithmetic transfer conjectures: the ramified quadratic case. *In preparation*, 2025.
- [Luo24] Yu Luo. On the moduli description of ramified unitary local models of signature $(n - 1, 1)$. *arXiv preprint arXiv:2404.10723*, 2024.
- [LZ17] Chao Li and Yihang Zhu. Remarks on the arithmetic fundamental lemma. *Algebra Number Theory*, 11(10):2425–2445, 2017.
- [LZ22a] Chao Li and Wei Zhang. Kudla–Rapoport cycles and derivatives of local densities. *Journal of the American Mathematical Society*, 35(3):705–797, 2022.
- [LZ22b] Chao Li and Wei Zhang. On the arithmetic Siegel–Weil formula for GSpin Shimura varieties. *Inventiones mathematicae*, 228(3):1353–1460, 2022.
- [Mih22] Andreas Mihatsch. Relative unitary RZ-spaces and the arithmetic fundamental lemma. *Journal of the Institute of Mathematics of Jussieu*, 21(1):241–301, 2022.
- [MLZ25] Andreas Mihatsch, Yu Luo, and Zhiyu Zhang. On unitary Shimura varieties in the ramified case. *In preparation*, 2025.
- [Pap00] George Pappas. On the arithmetic moduli schemes of PEL Shimura varieties. *J. Algebraic Geom.*, 9(3):577–605, 2000.
- [PR08] George Pappas and Michael Rapoport. Twisted loop groups and their affine flag varieties. *Advances in Mathematics*, 219:118–198, 2008.

- [PR09] George Pappas and Michael Rapoport. Local models in the ramified case. III. Unitary groups. *J. Inst. Math. Jussieu*, 8(3):507–564, 2009.
- [RSZ17] Michael Rapoport, Brian Smithling, and Wei Zhang. On the arithmetic transfer conjecture for exotic smooth formal moduli spaces. *Duke Math. J.*, 166(12):2183–2336, 2017.
- [RSZ18] Michael Rapoport, Brian Smithling, and Wei Zhang. Regular formal moduli spaces and arithmetic transfer conjectures. *Mathematische Annalen*, 370(3-4):1079–1175, 2018.
- [RTW14] Michael Rapoport, Ulrich Terstiege, and Sean Wilson. The supersingular locus of the Shimura variety for $\mathrm{GU}(1, n - 1)$ over a ramified prime. *Mathematische Zeitschrift*, 276(3-4):1165–1188, 2014.
- [RTZ13] Michael Rapoport, Ulrich Terstiege, and Wei Zhang. On the arithmetic fundamental lemma in the minuscule case. *Compos. Math.*, 149(10):1631–1666, 2013.
- [RZ96] Michael Rapoport and Thomas Zink. *Period spaces for p -divisible groups*, volume 141 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 1996.
- [Smi11] Brian Smithling. Topological flatness of local models for ramified unitary groups. I. The odd dimensional case. *Advances in Mathematics*, 226(4):3160–3190, 2011.
- [Smi14] Brian Smithling. Topological flatness of local models for ramified unitary groups. II. The even dimensional case. *Journal of the Institute of Mathematics of Jussieu*, 13(2):303–393, 2014.
- [Tre23] Stefania Trentin. On the Rapoport-Zink space for $\mathrm{GU}(2, 4)$ over a ramified prime. *arXiv preprint 2309.11290*, 2023.
- [Vol10] Inken Vollaard. The supersingular locus of the Shimura variety for $\mathrm{GU}(1, s)$. *Canad. J. Math.*, 62(3):668–720, 2010.
- [VW11] Inken Vollaard and Torsten Wedhorn. The supersingular locus of the Shimura variety of $\mathrm{GU}(1, n-1)$ II. *Inventiones mathematicae*, 184(3):519–627, 2011.
- [Wu16] Haifeng Wu. The supersingular locus of unitary Shimura varieties with exotic good reduction. *arXiv preprint arXiv:1609.08775*, 2016.
- [Yu19] Si Yu. *On Moduli Description of Local Models For Ramified Unitary Groups and Resolution of Singularity*. PhD thesis, Johns Hopkins University, 2019.
- [Zha21] Wei Zhang. Weil representation and arithmetic fundamental lemma. *Annals of Mathematics*, 193(3):863–978, 2021.
- [Zha24] Zhiyu Zhang. Maximal parahoric arithmetic transfers, resolutions and modularity. *arXiv preprint 2112.11994*, 2024.
- [Zhu14] Xinwen Zhu. On the coherence conjecture of Pappas and Rapoport. *Ann. of Math. (2)*, 180(1):1–85, 2014.
- [ZZ23] Ioannis Zachos and Zhihao Zhao. Semi-stable and splitting models for unitary Shimura varieties over ramified places. I. *arXiv preprint 2309.16463*, 2023.
- [ZZ24] Ioannis Zachos and Zhihao Zhao. Semi-stable and splitting models for unitary Shimura varieties over ramified places. II. *arXiv preprint 2405.06163*, 2024.

DEPARTMENT OF MATHEMATICS, COLUMBIA UNIVERSITY, 2990 BROADWAY, NEW YORK, NY 10027, USA
Email address: qh2275@columbia.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN MADISON, VAN VLECK HALL, MADISON, WI 53706, USA
Email address: yluo237@wisc.edu

SCHOOL OF MATHEMATICAL SCIENCES, ZHEJIANG UNIVERSITY, 866 YUHANGTANG RD, HANGZHOU, 310058, P.R. CHINA
Email address: 0023140@zju.edu.cn