

CONSTRUCTING EQUIVALENCES BETWEEN FUSION CATEGORIES OF QUANTUM GROUPS AND OF VERTEX OPERATOR ALGEBRAS VIA QUANTUM GAUGE GROUPS

CLAUDIA PINZARI

Dedicated to Sergio Doplicher, with gratitude

ABSTRACT. We summarize our results in [11] on a problem that Sergio Doplicher posed in the 90s in the setting of conformal nets, and on a related problem posed by Yi-Zhi Huang in the setting of vertex operator algebras, see e.g. [53]. The latter asks to find a direct construction of an equivalence due to Finkelberg between the quantum group modular fusion category at certain roots of unity and the fusion categories of affine vertex operator algebras at positive integer levels. The former asks for the construction of a quantum gauge group and field algebra for the category of localized endomorphisms of a conformal net, extending Doplicher-Roberts theory of the compact gauge group and field algebra in high dimensional algebraic QFT. In this summary we mostly focus on the affine VOA aspects (corresponding to the loop group conformal nets).

We solve Huang's problem for all the Lie types for which it is known that the centralizer algebras of the non-negligible tensor powers of a given generating object V for the quantum group fusion category (a list of V for which fusion tensor powers are well defined has been given by Hans Wenzl in [87] for all the Lie types), is generated by the representation of the braid group associated to the R -matrix, and for Huang-Lepowsky ribbon braided tensor structure for the vertex operator algebras. In the classical type A , this property becomes the classical Schur-Weyl duality for the vector representation, and was used by Doplicher and Roberts to construct $SU(d)$ as a special case of their unique compact gauge group and field algebra in algebraic QFT. Presently, known results in the literature on this generating property allow to construct our equivalence for the Lie types A, B, C, D, G_2 . Our approach may be regarded in the setting of Connes noncommutative geometry.

Lack of knowledge of the braid group generation property (i.e. for the Lie types E and F) allows us to show equivalence with Huang-Lepowsky ribbon braided tensor structure for all the ribbon morphisms, and part of those braiding morphisms, and associativity morphisms for which V takes one or two components, respectively.

Our work also gives a unitary modular tensor category structure on the linear category of modules of the Zhu algebra associated to the minimum energy functor (or on the vertex operator algebra) for all the Lie types, equivalent to that from quantum groups. This algebra becomes a unitary weak quasi-Hopf algebra, solving a problem posed by Frenkel and Zhu for all the Lie types [34].

Our main tool is the construction of two semisimple quantum gauge groups and an equivalence of their representation categories: The quantum group fusion category provides a unitary coboundary weak Hopf algebra associated to Wenzl functor in [87]. We then apply a de-quantization continuous curve of Wenzl and a Drinfeld twist method to reach the Zhu algebra. The structures we find in both contexts extend Doplicher-Roberts symmetric functors for rigid symmetric tensor categories with simple unit which play a central role in their duality theory.

1. INTRODUCTION

Moore and Seiberg [63], [64] discovered a connection between their equations and a certain structure of a tensor category arising from rational conformal field theories. The notion of ribbon braided tensor category has been settled in mathematics, and we refer to [21], [66], [27], [26], and, for their relations to compact quantum groups, to [70], and more generally to weak (quasi-) Hopf algebras to [11]. Turaev gave a definition of modular tensor category, based on previous work with Reshetikhin on the construction of knot and 3-manifold invariants see [85]. Several authors contributed to construct modular fusion categories from quantum groups at roots of unity, and a more complete historical account on the quantum group side may be found in [80], [82]. see also the introduction of [11]. We denote by $\mathcal{C}(\mathfrak{g}, q)$ the resulting modular fusion category, with $q = e^{i\pi/d\ell}$, and $d \in \{1, 2, 3\}$ a suitable integer structure constant from Lie theory. Unitary tensor structures have been constructed by Wenzl and Xu [87], [92], and the former paper, as well as more recent papers by Wenzl and his coauthors play a major role in our work, and we try to explain this in this paper.

On the side of CFT, our main interest in this paper is the Wess-Zumino-Witten model. Modular, or more generally ribbon rigid braided tensor category constructions involved many authors among physicists and mathematicians.

Major work was carried out in a series of papers by Kazhdan and Lusztig in the mid 90s. Let \mathfrak{g} be a complex simple Lie algebra. The authors construct rigid braided tensor structures for certain module categories of affine Lie algebras associated to \mathfrak{g} at levels excluding the positive rationals. In their work they consider both generic and non generic values of the levels. In particular they consider the negative integer levels $-k$, $k \in \mathbb{N}$, and proved a braided tensor equivalence of their category \mathcal{O}_{-k} with the non-semisimple category of certain modules of the Drinfeld-Jimbo quantum group $U_q(\mathfrak{g})$ at the same type of roots of unity $q = e^{i\pi/d\ell}$, with the exception of some exceptional cases at low levels [57]. The order ℓ appearing in the root of unity and the positive integer k are related by a well known equality, see e.g. Def. 19.1 and 20.2 in [11] and references therein.

Shortly after, Finkelberg considered a certain semisimple subquotient $\tilde{\mathcal{O}}_{-k}$ of the representation category constructed by Kazhdan and Lusztig, paralleling the subquotient construction in the setting of quantum groups, and described a passage $\tilde{\mathcal{O}}_k \rightarrow \tilde{\mathcal{O}}_{-k}$ claimed as a braided tensor equivalence from the category $\tilde{\mathcal{O}}_k$ of certain integral modules of the affine Lie algebra at the positive integer level k to the subquotient rigid braided tensor category $\tilde{\mathcal{O}}_{-k}$. Combining with the equivalence by Kazhdan and Lusztig, Finkelberg derived a connection between $\tilde{\mathcal{O}}_k \rightarrow \mathcal{C}(\mathfrak{g}, q)$. See the introduction and Theorem 4.3 in [30].

The original work by Finkelberg refers to the cited work in physics by Moore and Seiberg and in algebraic geometry by Beilinson, Feigin, and Mazur [2] for the rigid braided tensor structure on $\tilde{\mathcal{O}}_k$. As I learned from discussions with Huang, rigidity was not proved in [2], and there was no proof of rigidity using this approach.

Remarkably, Huang and Lepowsky introduced and developed a rigorous direct construction of a modular (vertex) tensor category structure in a very general setting for vertex

operator algebras, including as a particular case the affine vertex operator algebras at positive integer levels that we denote by V_{gk} . The construction has been completed by Huang between 2005 and 2008 [47], [51] following a general construction of tensor product theory in the setting of vertex operator algebras by Huang and Lepowsky [42], [43], [44] and by Huang [45], [50]. In particular [45] gives the construction of associativity morphisms which is the first difficulty in the construction of the tensor structure. An historical account may be found in [52]. We also refer to the introductions of [46] and [53]. These works give in particular the first mathematical proof of modularity, that applies moreover to a large class of representation categories of vertex operator algebras.

We note that difficulties in the construction of associativity morphisms in the setting of Drinfeld category were earlier solved by Drinfeld in [23], [24], [25] by the introduction and construction of quasi-Hopf algebras, via differential Knizhnik-Zamolodchikov equation, leading to his remarkable Drinfeld-Kohno theorem connecting via a Drinfeld twist with the quantized universal enveloping algebra. In this case, rigidity of Drinfeld category is derived from the equivalence constructed by Drinfeld in Drinfeld-Kohno theorem.

In particular rigidity and modularity property in the setting of vertex operator algebras was first shown in [47]. To show modularity, and in particular rigidity, Huang first discovered the need of the Verlinde formula. The paper [50] gives a proof of the Verlinde formula in the general case. See also [48] for the proof of convergence and extension property assumed in [45] and [49] for the proof of modular invariance conjecture of Moore and Seiberg needed in the proof of the Verlinde formula.

By a private communication with Huang on the history of this problem, I learned that it finally turned out that the construction by Finkelberg described in [30] connecting $\tilde{\mathcal{O}}_k$ with Kazhdan and Lusztig structure inherited by $\tilde{\mathcal{O}}_{-k}$ could not suffice to derive rigidity of $\tilde{\mathcal{O}}_k$ from that of $\tilde{\mathcal{O}}_{-k}$. This implied that the proof of the equivalence $\tilde{\mathcal{O}}_k \rightarrow \tilde{\mathcal{O}}_{-k}$ was far from complete.

I have been informed by Huang that these doubts were motivated by two facts. On one hand, his earlier direct communications in the 90s had clarified that rigidity was not proved in [2]. On the other, the mentioned property that his proof of rigidity in the setting of vertex operator algebras involved the Verlinde formula indicated that more work was likely needed to complete the proof in [30]. These facts together with early claims of rigidity in [30], motivated Huang to a comparison of the results in [30] with their results on rigidity and this led to note a gap in a proposition of [30] with implications on the main result.

Huang's study had led him to identify the gap in the fact that a suitable constant needed to be shown to be nonzero in a proposition useful to obtain the equivalence between the two local systems considered by Finkelberg and Kazhdan and Lusztig, and this proposition would imply derivation of rigidity in his case from previous work. On the other hand, Huang proof of rigidity in the setting of vertex operator algebras had similarly reduced to show nontriviality of a constant, and to prove this he had used the Verlinde formula as above alluded to, which was a main result in their work. This had led Huang to inform Finkelberg about utility of this formula to show rigidity in his case.

As later reported by Finkelberg in [31], it turned out that the completion of the proof of that proposition leading to rigidity of $\tilde{\mathcal{O}}_k$ and the equivalence $\tilde{\mathcal{O}}_k \rightarrow \tilde{\mathcal{O}}_{-k}$, involved the Verlinde formula as well (which became another main result in Finkelberg's paper). This reflects also on Finkelberg composed equivalence $\tilde{\mathcal{O}}_k \rightarrow \tilde{\mathcal{O}}_{-k} \rightarrow \mathcal{C}(\mathfrak{g}, q)$, with Kazhdan-Lusztig equivalence that moreover still passes through the negative integer levels. This implies in particular that the theorem does not include a few exceptional cases, including $\mathfrak{g} = E_8$ and $k = 2$, which correspond to cases not treated in Kazhdan-Lusztig theory.

Following the introductions in [46], [53], Finkelberg work [30], [31] may be reinterpreted as giving the construction of a braided tensor category structure on $\tilde{\mathcal{O}}_k$. One may also conclude that these papers give a second proof of rigidity and modularity of this category.

An approach to modularity based on algebraic geometry may be found in Theorem 7.0.1 [1] but the proof of rigidity is not given and, as already mentioned it was known to experts that there was no proof based on their approach at the time. The author thanks Y.-Z. Huang for this information. We point out the very recent paper by Etingof and Penneys where the authors show among other things that weak rigidity in the sense of [1] is equivalent to rigidity for certain semisimple braided tensor categories of moderate growth with simple unit, see [28] giving a third proof of rigidity and give applications to a simplification of Huang's proof of rigidity. They also answer positively to a question posed in [1] that for every finite semisimple category over an algebraically closed field the data of a modular functor is essentially equivalent to the structure of a modular fusion category, completing the proof of rigidity in [1].

On the other hand, Finkelberg equivalence theorem is stated as Theorem 7.0.2 in [1] without proof. The braided tensor equivalence between $\tilde{\mathcal{O}}_k \rightarrow \mathcal{C}(\mathfrak{g}, q)$ obtained from the combined works by Finkelberg, Kazhdan and Lusztig is considered indirect by several authors. This leads to the desire of constructing the equivalence in a natural, direct, and explicit way, and this problem is discussed by Y.-Z. Huang in [53] in great detail, where we select only a small part.

Problem 4.4 in [53]. Find a direct construction of this equivalence without using the equivalence given by Kazhdan-Lusztig so that this equivalence covers all the cases, including the important $\mathfrak{g} = E_8$ and $k = 2$ case.

We also mention Gannon's observation in Sect. 6.2.3 of [35] regarding the desire of finding a direct connection between quantum groups and CFT.

A direct construction of the equivalence may be helpful to see how rigidity or modularity in the setting of CFT may be transported from corresponding results in the setting of quantum groups at roots of unity, which seem simpler, or vice versa, by a given explicit equivalence. Or to simplify or give new construction of other structures, e.g. unitary structures, with a quantum group viewpoint, or to see more connections with quantum groups from these fundamental models. The study of how unitarization from the quantum group fusion categories [87] was useful to derive unitarization in representation categories of vertex operator algebras via an explicit equivalence was important for us since the early stages of [11], and further motivated us to study the above problem.

Our approach to Huang problem is based on interpreting Finkelberg-Kazhdan-Lusztig equivalence as an extension to quantum groups of Doplicher-Roberts duality theory for the important class of WZW models that are described by affine Lie algebras at positive integer levels. We shall briefly come back to the previous history of these important models in physics. From the side of mathematical axiomatizations of CFT, we work with respect to Huang and Lepowsky braided tensor structure for the representation category of vertex operator algebras.

Our starting point to approach this problem arose from a problem generously posed by Sergio Doplicher in the 90s to the author in a different setting, that of algebraic quantum field theory. We try to explain this problem.

Algebraic quantum field theory had been initiated by Haag and Kastler in [40] in the 60s, and developed by founding breakthrough works by Doplicher, Haag, and Roberts [15], [16], [17], [18] in the 70s. We refer to the book by Haag [41], for an in-depth introduction to this theory, based on functional analysis, and more specifically on operator algebras, and also for more references.

In this context, I learned from private discussions that for high dimensional quantum field theories categorical structures in their study first emerged from the discovery by Doplicher of the notion of localized endomorphism of the local observable algebra, allowing composition of two of them which was understood as a strict tensor product structure. The mentioned works by Doplicher, Haag, and Roberts led to a deep construction of the structure of a rigid symmetric tensor C^* -category with simple unit, starting from first principles in physics.

Rigidity, called charge conjugation in this setting, is due to particle-antiparticle duality. In their papers rigidity was shown to imply deep and useful properties for the classification of the statistics parameter, and charges of para-Bose or para-Fermi type. Because of the symmetry, their results on the classification statistics parameter, which was shown to be the reciprocal of a positive integer d up to a sign, and C^* -structure, in the 70s it arose the question of whether that was the category of representations of a compact group, because those data arising from a single localised endomorphism were also satisfied by the representation category of a compact Lie subgroup of $SU(d)$, that would be a gauge group of global symmetries. They considered the abelian case in the 70s [17]. In that case, every object had dimension 1, and in modern terminology these categories are referred to as pointed in e.g. [26].

In a series of important papers dated the end of the 80s, Doplicher and Roberts were able to solve the problem in full generality. Their construction gives a unique compact gauge group with representation category equivalent to that of a given symmetric rigid tensor C^* -category with simple unit, yielding the formulation of a new abstract duality theory for compact groups [19], [21],[20], and led to desired applications to high dimensional Algebraic QFT [22].

The term abstract duality refers to the property that the given categories do not have a tensor functor to the category of f.d. vector spaces from the beginning, raising the problem of constructing these structures. It is thus an abstract form of duality in the sense of Tannaka-Krein, where in the latter the tensor structure on the embedding functor

is given from the start. In their theory, an important role is played by the notion of symmetric functor to the category of f.d. vector spaces, for which they show existence and uniqueness. This property allows to find their unique compact gauge group G .

In two or three dimensional quantum field theory, it was noted that the corresponding categories were braided rather than symmetric. This implies that Bose-Fermi alternative does not hold any longer. Models were first described by Streater and Wilde. Doplicher, Haag, Roberts theory of superselection sectors was generalized to braid groups statistics by Fredenhagen, Rheren, Schroer [32], [33]. The representation of the braid group in the tensor C^* -category is unitary.

The discovery of quantum groups first arose in physics literature in the work by Faddeev, introduced as noncommutative function algebras. In the mid 80s breakthrough papers by Drinfeld and Jimbo discovered them as quantization of universal enveloping algebras, and developed their properties. We refer to the introduction in [9] and more historical information at the end of each chapter in their book.

After the discovery of the quantum groups in the mid eighties, Doplicher asked the author to try to extend their theory and consequent applications on the construction of a quantum gauge group and a related field algebra. Of course, several studies were dedicated to extend Doplicher-Roberts duality theory to quantum groups, see Ch. IV.5 in [41] for more references on an early stage of this problem, see also the introduction in [11].

Our approach may be regarded in the setting of Connes noncommutative geometry [13], and follows naturally our previous work in the setting of compact quantum groups, see also the introduction in [12], [14]. In [73], [74], [78], [75], [76], [77], [79] we have developed at the level of tensor categories an analogue of Doplicher-Roberts construction of compact gauge group, although the analogy cannot be performed in a strict way due to new difficulties in the quantum case. We shall come back to this, see the mentioned papers and [11]. One important difference was that for rigid braided tensor categories we do not have a notion of braided tensor functor beyond the symmetric ones, because the category of vector spaces does not admit a braided symmetry. Giving up a compatibility property in the symmetric case implies lack of uniqueness of the functor, and thus of the gauge group. This was noticed early as an easy consequence from their construction.

To treat this difficulty in [10] and [11] we have taken into account Drinfeld notion of coboundary, which finally allows for a generalization. Moreover we have followed both the original choice by Mack and Schomerus [61] of a given dimension function corresponding to non-negligible tensor powers of the vector representation of quantum \mathfrak{sl}_2 at roots of unity and also the study of classification of compact quantum groups corresponding to a given representation category of deformation compact quantum groups by Neshveyev and Yamashita [71], [72]. They both considered a specific positive integer valued dimension function associated to the given category of the quantum group, which was the classical dimension function. This suggested to us to consider in [10] and [11] a specific dimension function on a functor related to classical data, and this was first Wenzl functor [87] for the quantum group fusion category, and then the minimum energy functor for the WZW model [93]. It was particularly important for us that in [87] Wenzl describes an explicit quantization curve associated to his functor that leads to an algebra that identifies with

the Zhu algebra for that model in the setting of vertex operator algebras. Since in [10] we had constructed a natural quantum group structure on the former compatible with the tensor structure, this arose the question of constructing another natural quantum group structure on the latter at the time of [10], following the quantization curve on the unitary structure of tensor products with the aim of making it classical. This involved in [11] the construction of a Drinfeld twist, a notion that Drinfeld had introduced to study equivalence of what is known as the Drinfeld category with the quantum group category in a formal setting.

We refer to Sect. 23 in [11] for an extension of certain results by Neshveyev and Yamashita in the setting of quantum group fusion categories of type A based on Kazhdan-Wenzl theory at roots of unity in our case, and on the weak Hopf algebras constructed in [10], in place of the compact quantum groups. This section is important for us as it suggested that in general the the associativity morphisms in a category arising from CFT may be of a simple nature from the viewpoint of associated quantum gauge groups that need to be constructed, and this finally proved to be the case, we shall come back to this with more precision.

On the other hand, in the setting of compact quantum groups, while extending Doplicher-Roberts theory starting from embedding part of the morphisms of the category into the Hilbert spaces [73], [75], [79], we had already started to fix a dimension function, and from that we constructed a tensor functor from that subcategory. To consider the full category, the analogue Doplicher-Roberts compact gauge group is described by an ergodic action of a larger quantum group on a quantum homogeneous space, rather than a quantum group itself, due to lack of commutativity of the C^* -algebra describing the homogeneous space constructed in [75], and this property is likely to hold also in other models arising from conformal field theory, unless one can show that certain known obstructions already met in [74] for compact quantum groups, vanish. We shall come back to this.

However, for the WZW models arising from affine Lie algebras at positive integer levels of the Lie types mentioned in the abstract, we are able to show in [11] that we obtain genuine quantum groups as quantum gauge groups, described as semisimple weak Hopf algebras with R -matrix, C^* -structure explicitly associated to Drinfeld coboundary matrix, and with unitary representations of the braid group, derived from the quantum group fusion category. This weak Hopf algebra has representation category equivalent to both the starting fusion category and also to that of the corresponding affine vertex operator algebra. Constructing the equivalence from the former to the latter as we do in [11] is our proof of Kazhdan-Lusztig-Finkelberg theorem, see Theorem 2.1.

Among our first references in the context of conformal field theory important for us to start to set up the connection with the work of Wenzl in quantum groups, is the work by Wassermann [86] in the setting of affine Lie algebras, where the key concept is played by the notion of *primary field*, and its initial term, which we use in a fundamental way to connect quantum group fusion category to representations of the Zhu algebra and their special tensor products with the fundamental representation, see Sect 33 in [11]. Most importantly, we describe a connection between Wenzl and Wassermann work via Wenzl quantisation curve in theorem 33.7, to which we refer the reader also for important historical references.

The connection from affine Lie algebra setting to Huang-Lepowsky tensor product theory may be found in Sections 35 and 36.

Our abstract duality methods need new further tools as we shall see from the theory of quantum groups, mainly from the work of Drinfeld and Wenzl, that are not available in the theory of groups. However, we shall find that in both contexts, quantum groups at roots of unity and affine vertex operator algebras at positive integer levels, both representation categories are described by functors extending the symmetric tensor functors in Doplicher-Roberts abstract duality theory, see Sect. 27 in [11]. We call these structures hermitian coboundary weak quasi-Hopf algebras, as the hermitian structure is given by Drinfeld coboundary matrix, which is a modification of the R -matrix with a square root of the ribbon structure.

We notice on the other hand that a stronger property is needed to establish the connection, than the purely categorical characterisation of functors in Sect. 27. This property is expressed in terms of quantum groups, and the first example, albeit in a non-semisimple setting, is the hermitian structure of $U_q(\mathfrak{g})$ at a root of unity by Theorem 29.9. This property says that the $*$ -involution enjoys a precise anticommutation relation (rather than commutation relation as is familiar in Hopf C^* -algebras) with the coproduct and was first noticed in [87]. We axiomatize this property in section 28, as a compatibility property of hermitian coboundary weak quasi-Hopf algebras. Further examples derived from this examples are our weak Hopf and quasi-Hopf algebras, see Sections 29, 30, 31.

We shall derive even more structure than rigidity e.g. unitarity and modularity on the vertex operator algebra representation category from the corresponding quantum group fusion category, via the equivalence, see Sects. 21, 22, 24 in [11]. Moreover our equivalence, which is given by an analogue of the Drinfeld twist in [23], [24], gives an understanding of the unitary structures of representation categories of affine vertex operator algebras as classical compact real forms, or untwisted. We refer to the outlook section for some more information on possible extension of our viewpoint to models beyond the affine vertex operator algebras at positive integer levels, see also [11].

Our main result is Theorem 2.1. In the subsequent sections we outline the plan of our proof. Our proof starts from the work of Wenzl [87] in the setting of quantum group fusion categories. A main idea that we follow is to consider the fundamental representation V of \mathfrak{g} for each Lie type as in [87] where the author explains its generating property and good fusion structure with respect to tensor products with an arbitrary irreducible representation allowing a well-defined truncation procedure to obtain non-negligible objects, with the aim of studying unitarizability of the fusion category. It should also be noted that a main application of that paper is construction of inclusion of subfactors of finite Jones index for all Lie types, extending his previous work.

For all the Lie types we obtain complete identification for the tensor product structure, all the ribbon structure, and those braiding and associativity morphisms where one or two variables are the fundamental representation V , respectively. These are parts (a), (b) of Theorem 2.1. The main method of our proof is the construction of semisimple weak

Hopf or quasi-Hopf algebras in the two settings and an equivalence of their representation categories via the construction of a Drinfeld twist.

The term weak in the setting of quasi-Hopf algebras was first introduced by Mack and Schomerus motivated by the work of Drinfeld for quasi-Hopf algebras and Drinfeld-Kohno theorem [61]. Weakness refers to a non-unital coproduct, needed to exclude the non-physical representations with zero quantum dimension, and with the need of compatibility with the fusion rules of the category arising from a suitable model in CFT, described by a quantized universal enveloping algebra of \mathfrak{sl}_2 at a root of unity.

Mack and Schomerus did not consider the Hopf counterpart in their theory. Weak Hopf algebras were later introduced by Böhm and Szlachanyi motivated by the work of Mack and Schomerus, but with a strictly coassociative coproduct [5]. The definition of weak Hopf algebra has become standard and understood in the sense by Böhm and Szlachanyi.

In [11], and also in this paper, weak Hopf algebras are understood with a different meaning. While a different name would be helpful to avoid confusion, so far we have not been able to find a good name for our algebras to clarify the situation, and we use the term weak Hopf algebras in the sense defined in Sect 9 in [11].

In [11] we develop a general theory of weak Hopf algebras, review and expand the theory of weak quasi-Hopf algebras, mainly Tannaka-Krein aspects, both in the algebraic and in the C^* -algebraic setting, with non-trivial unitary structure, discuss a suitable Drinfeld-Kohno theorem, and its application to the construction of semisimple weak Hopf and quasi-Hopf algebras with structure naturally compatible with that of the involved ribbon braided tensor categories, for quantum groups at roots of unity and affine vertex operator algebras at positive integer levels, including unitary structure as hereditated from the unitary structure of the category due to Wenzl [87]. We follow the strategy of Drinfeld for Drinfeld-Kohno theorem which includes the comparison of braiding morphisms of the two settings, but we do not include the comparison of the associativity morphisms in our analogue, Theorem 28.4 in [11] (more details are given later). We construct a Drinfeld twist for our weak Hopf algebras useful for the braided symmetry, and study associativity morphisms with a uniqueness theorem using the generating representation. We then use this twist to concretely construct unitary structures in the setting of the representation categories of affine vertex operator algebras at positive integer levels with the tensor structure due to Huang and Lepowsky, for all the mentioned Lie types, by transferring the unitary structure from quantum group fusion category constructed in [87].

Our weak Hopf algebras while having a non-unital coproduct, are not in the class of Bohm and Szlachanyi. For example their coproduct is not strictly coassociative. They are not extension of Bohm and Szlachanyi weak Hopf algebras, and we know from results in [11] that the intersection of the two classes are the genuine Hopf algebras.

On the other hand, in our opinion the weak Hopf algebras considered in [11] still deserve to be regarded as Hopf algebras in a weak sense, for various reasons. On one hand they seem to play the role of the quantized universal enveloping algebras in Drinfeld-Kohno theorem, following the original motivation of Mack and Schomerus. Moreover, their representation category is characterized by a tensor functor to the category of Hilbert spaces with non-invertible natural transformations (projections endowed with natural inverse inclusions, in

our case) satisfying tensoriality diagrams, which reduces to the notion of tensor functor when the natural transformations are invertible. Their counit is antimultiplicative and the antipode satisfy the same relations as in the case of Hopf algebras.

We prove the complete identification of braiding and associativity morphisms of the two structures for all the classical Lie types A, B, C, D , and the case G_2 in part (c) of Theorem 2.1, by further refining methods of the previous part but we use more structure in Sect. 8 of [11] that has not emerged in other parts.

For the type A case, the extra structure that we need to use is known as quantum Schur-Weyl duality between the vector representation and the natural representation of the braid group between the tensor powers of the vector representations.

In the classical case, Schur-Weyl duality is a key property for the general linear groups starting with the work of Schur and Weyl. Schur-Weyl duality also takes an important role in Doplicher-Roberts abstract duality theory for compact groups and applications to construction of the field algebra. This classical duality emerges as the starting point for the construction of global gauge symmetry groups in tensor categories arising in [22].

Duality with the braid group representation is a concept at the basis of the origin of quantum groups following V. F. R. Jones [55].

More generally, for Lie types beyond A , we need a corresponding duality between the centralizer algebras of the truncated tensor powers of the fundamental representation V of the Drinfeld-Jimbo quantum group $U_q(\mathfrak{g})$ at the root of unity and the natural representation of the braid group in these algebras, and this leads to the restriction stated in (c). This allows us to apply our abstract uniqueness Theorem 8.19 in [11] for braided tensor structures in a pre-tensor category.

In our work, the need of quantum duality between the braid group representation and the fundamental representation V emerged on the occasion of the proof of our earlier Lemma 7.10, which was not complete in the version 7 of [11], and we worked on completion with the same methods. It was for us a sudden realization of an even greater recognition than we had realized up to that point of the contribution of the authors on extensions of Schur-Weyl duality which has historical interest on his own in representation theory, and more work by Wenzl and his collaborators, and several other authors in the classical and quantum cases for the various Lie types. We shall come back to this in more detail.

In our setting, quantum extension of Schur-Weyl duality to the Lie types has a role to see uniqueness of the associativity morphisms in a tensor category associated to the given braided symmetry with a generating representation V , see Theorem 8.19 in [11]. This applies to both fusion categories associated to quantum groups at roots of unity and of affine VOAs.

Presently it is not known to the author whether the fundamental representation for the exceptional Lie types E and F satisfies braid group duality. The E_8 case is more delicate, because fusion of tensor powers of $U_q(\mathfrak{g})$ at the root of unity is more complicated, as explained in [87], but more work by Wenzl is available for the study of the centralizer algebra [88]. We are grateful to M. Yamashita for pointing out this reference to us.

Among possible applications of our proof are the study of unitary structures of representation categories of vertex operator algebras, due to the clarity offered by a quantum group

viewpoint of the same problem as explained in [87]; the study of equivalences between representation categories of conformal nets and vertex operator algebras via the construction of a common quantum gauge group; to extension of Doplicher-Roberts field algebra construction from the local observable algebra to the conformal field theory setting. Some more discussion may be found in the outlook section 13.

2. THE MAIN RESULT

Theorem 2.1. *Let \mathfrak{g} be a complex simple Lie algebra and let $\text{Rep}(V_{\mathfrak{g}_k})$ be the module category of the corresponding affine vertex operator algebra $V_{\mathfrak{g}_k}$ at level $k \in \mathbb{N}$, endowed with the ribbon braided tensor structure introduced by Huang and Lepowsky and let $A(V_{\mathfrak{g}_k})$ be the Zhu algebra. Then*

- (a) *The representation category $\text{Rep}(A(V_{\mathfrak{g}_k}))$ of the Zhu algebra $A(V_{\mathfrak{g}_k})$ admits a natural structure of ribbon braided tensor category obtained by transporting Huang-Lepowsky structure to $\text{Rep}(A(V_{\mathfrak{g}_k}))$. In this way, Zhu's linear equivalence*

$$Z : \text{Rep}(V_{\mathfrak{g}_k}) \rightarrow \text{Rep}(A(V_{\mathfrak{g}_k}))$$

admits a natural structure of ribbon braided tensor equivalence.

- (b) *Let $A(V_{\mathfrak{g}_k})$ be endowed with the unitary compatible coboundary weak quasi-Hopf structure induced by $\mathcal{C}(\mathfrak{g}, q)$ via the construction of a weak tensor structure on Wenzl functor and of the Drinfeld twist following Theorem 31.1 in [11] (a), and let $\text{Rep}(A(V_{\mathfrak{g}_k}))$ be endowed with the corresponding ribbon braided tensor structure.*

Then $\text{Rep}(A(V_{\mathfrak{g}_k}))$ becomes a unitary coboundary modular tensor category in this way. Moreover the transport in (a) preserves the tensor products, the ribbon for all objects, the braid morphisms for pairs (V_λ, V) and (V, V) , and the associativity morphisms for triples (V_λ, V, V) , (V, V_λ, V) , (V, V, V_λ) with V_λ an arbitrary irreducible of the fusion category.

- (c) *If \mathfrak{g} is of one of the Lie types A, B, C, D, G_2 then the transport preserves the braided tensor structure for all objects. Thus Z is a ribbon braided tensor equivalence. Specifically, the composition*

$$\text{Rep}(V_{\mathfrak{g}_k}) \xrightarrow{Z} \text{Rep}(A(V_{\mathfrak{g}_k})) \xrightarrow{31.1,(a),(b)} \text{Rep}(A_W(\mathfrak{g}, q)) \xrightarrow{\text{TK ribbon equiv}} \mathcal{C}(\mathfrak{g}, q) \quad (2.1)$$

is a ribbon tensor equivalence by application of the indicated theorems.

The definition of unitary coboundary weak Hopf algebra, its restricted version named unitary coboundary weak Hopf algebra, are given in Sects. 29, 30 in [11], by abstracting features of the hermitian structure of $U_q(\mathfrak{g})$ for $|q| = 1$, discovered by Wenzl in [87] in his proof of unitarity of Kirillov inner product, for the associated fusion category as certain roots of unity, based on a complex Drinfeld coboundary matrix. The former more general notion admits a categorical characterization generalizing Doplicher-Roberts symmetric functors.

We would like to notice that the passages described in part (c) of Theorem 2.1 are explicit. The first and the last equivalences are due respectively, to the work by Zhu

combined with that of Huang and Lepowsky. The last is the functor due to Wenzl enriched with our weak tensor structure, preserving Drinfeld coboundary structure derived from the ribbon and R -matrix, see the above cited sections in [11].

The middle equivalence in (2.1) is application of the explicit symmetry provided by our analogue of Drinfeld-Kohno theorem 28.4 in [11]. This symmetry is given by a Drinfeld twist explicitly defined by the action of the R -matrix, more precisely by the braided and ribbon structure in the two settings, that allow to see in the vertex operator algebra setting same structure as that in the quantum group setting.

3. MOTIVATION AND FRAMEWORK OF OUR WORK. CONSTRUCTING BRAIDED TENSOR EQUIVALENCES VIA A QUANTUM GAUGE GROUP GIVEN BY A DISCRETE WEAK HOPF ALGEBRA AND A DRINFELD TWIST.

Before going into the explanation of the statement and outline the main tools of the proof in more detail, we summarize the general framework that motivated and led to our work. In the following, where not indicated the cited sections and results refer to [11], and also for unexplained notation.

We were originally motivated by Doplicher-Roberts abstract duality theory for compact groups and their application to the construction of a global compact gauge group and reconstruction of the net of field algebras from the observable algebras in high dimensional quantum field theories [21], [22].

In the setting of low dimensional quantum field theories, starting with the fusion categories of quantum groups at all the roots of unity that are known to correspond to unitary structures by the work of Wenzl and Xu [87], [92], we in particular construct discrete quantum group-like structures with non-trivial twisted unitary structures for all the Lie types as in [87]. These discrete quantum groups are thought of as duals of compact quantum groups, but their preduals admit no analogy to the case of the compact quantum groups of Woronowicz, because they are non-associative algebras. However, Woronowicz work was strongly inspiring for us. Indeed our first examples in the type A case were first constructed in our previous work [10] via an abstract Tannaka-Krein method in the search of a noncommutative function algebra, using the fusion rules of the corresponding fusion category.

On the other hand, their duals are quite well-behaved discrete C^* -algebras with almost associative coproducts, antimultiplicative antipode and multiplicative counit satisfying the same relation as for Hopf algebras. In this sense they resemble the duals of the compact quantum groups, the main differences being non-unitality of the coproduct and the unitary structure of the tensor product of their representations, which is twisted via a positive operator associated to the R -matrix (Drinfeld coboundary matrix), and makes the representation of the braid group unitary. The need of a twist for the tensor product representation arises from the fact that coproduct and $*$ -involution satisfy an anticommutativity condition, rather than commutativity as it usually is the case for Hopf algebras in the operator algebra setting. This anticommutativity is derived from the same property of the $*$ -involution of the non-semisimple $U_q(\mathfrak{g})$, first pointed out by Wenzl [87].

The weak Hopf algebras constructed in [10] have been extended to all Lie types in Sections 29, 30 of [11] with uniform methods for all the Lie types.

Our algebras are examples of Mack-Schomerus weak quasi-Hopf algebras introduced in [61] as generalization of Drinfeld quasi-Hopf algebras with non-unital coproduct. For $\mathfrak{g} = \mathfrak{sl}_2$ our examples reduce to Mack-Schomerus examples for the algebraic structure. We understand their lack of coassociativity as quite mild, because their representation category is endowed with a weak unitary tensor functor in the category of Hilbert spaces. Thus we regard them as the Hopf counterpart of weak quasi-Hopf algebras, in a position similar to that of Hopf algebras among quasi-Hopf algebras, that in our knowledge has been overlooked.

Here the term weak is understood in the sense first introduced by Mack and Schomerus, the coproduct is non-unital, a feature needed to exclude the non-physical representations with zero quantum dimension, and with the need of compatibility with the tensor structure of the category. The weakness of tensoriality of the embedding functor of their representation category to the category of Hilbert spaces means that the structure natural transformations are surjections onto subspaces of the tensor product spaces endowed with left inverses, rather than isomorphisms, and satisfy two tensoriality diagrams.

Much of Woronowicz theory can be extended to our case. From this via a quantization isomorphism described by Wenzl in [87] and the construction of a Drinfeld twist we construct and describe in some more detail later a quantum group-like structure on the Zhu algebra associated to the affine models described as in the statement of the main result.

These constructions are in some quantum analogy with Doplicher-Roberts construction in the setting of high dimensional algebraic quantum field theory, in that the categories that we study are not given as Tannakian categories. In our case, differently from their case, we have naturally defined linear functors emerging as the minimum energy functor due to Zhu in the setting of vertex operator algebras. Also differently from their case, where the linear functor needed to be constructed but the tensor structure was trivial, in our case we need to construct tensor structures associated to these naturally occurring linear functors to the category of vector spaces.

Our starting point is what we shall refer to as Wenzl functor associated to the quantum group fusion category described in [87]. This paper plays a central role, both on the side of quantum groups and on the side of affine vertex operator algebras, as we next try to explain.

As said, on the the side of quantum groups, our approach has been influenced by results in quantum group theory by Woronowicz but also by the dual viewpoint by Majid for quasi-Hopf algebras, extended to weak quasi-Hopf algebras by Haring-Oldenburg. We start reviewing these Tannakian reconstruction theorems from the viewpoint of Majid and Haring-Oldenburg in Sect. 5 in [11], where the reader may find indication of the original references. Moreover Kazhdan and Wenzl theory is also important to us, and the reader may consult Sect. 23 in [11] and references therein. It is by no means an exaggeration to assert that our paper [11] would have not existed without the paper by Neshveyev, Tuset on Kazhdan-Lusztig theorem for compact quantum groups [68]. This paper was strongly inspiring for our strategy, although methods are different. For similar reasons, we were also

guided by the presentation of Drinfeld-Kohno theorem in the book by Bakalov and Kirillov [1]. These last two references first inspired our work in the search of a unified treatment following a Tannakian approach in the idea of Majid, in our case. An important aspect in our case as far as Majid Tannakian duality for quasi-Hopf algebras, is the search of natural tensor structures for the involved embedding functors into the category of Hilbert spaces (Wenzl functor and Zhu functor in the two settings).

Primary fields first emerged in the physics literature, [3], using previous work by Feigin and Fuks [29] and Kac [56] on the representation theory of the Virasoro algebra. In 1984 Witten introduced what are now known the WZW models, due to Wess, Zumino, Novikov and Witten [90]. In the same year, Knizhnik and Zamolodchikov showed that their correlation functions satisfy a system of partial differential equations [59].

Tsuchiya and Kanie started a rigorous study in [84]. Goddard, Kent and Olive realized the chiral unitary Virasoro minimal models via a coset construction [37].

Moore and Seiberg conjectured some analytic properties of the quantum fields and described the construction of a modular category associated to a general conformal field theory.

The mathematical contribution by Kohno, Tsuchiya and Kanie [84], and the subsequent works by A. Wassermann [86] and Toledano-Laredo [83] on affine Lie algebras, compared with the structure that had previously emphasized by Wenzl in the study of quantum group fusion categories and their unitary structures [87] that led to our approach. The reader may find reference to more of the several original papers in mathematics and physics in Sections 31–36 in [11].

4. THE GENERAL PLAN OF OUR PROOF.

We briefly explain here the plan of our proof of the main theorem for parts (a), (b), and (c). In the following sections we describe the main ideas to prove these parts in greater detail.

Part (a) is an application of a general construction of a natural structure of a (ribbon, braided) tensor equivalence on a given equivalence $\mathcal{C} \rightarrow \text{Rep}(A)$ of semisimple linear categories endowed with an inverse linear equivalence $\text{Rep}(A) \rightarrow \mathcal{C}$, with A a discrete algebra. We outline the main ideas of this tensor structure construction in the next section. The reader may refer to Section 15 in [11] for the detailed construction.

For part (b), we follow a suitable Drinfeld twist method originally used by Drinfeld for Drinfeld-Kohno theorem [23], [24], [25], and in our case for different categories, defined in the statement of the main result. This is the main part of the result. The first main step is the construction of a class of finite dimensional weak Hopf C^* -algebras $A_W(\mathfrak{g}, q)$ for all Lie types and positive integer levels, extending the former examples of [10] in the type A case. They have unitary coboundary structures associated to the fusion category of quantum groups at roots of unity, associated to a specific linear fiber functor $W : \mathcal{C}(\mathfrak{g}, q) \rightarrow \text{Vec}$, that emerges from the work of Wenzl [87]. Complete definitions of unitary coboundary structures on weak quasi-Hopf algebras are given in [11], and are motivated by the work

of several authors, among them Drinfeld, Wenzl, Mack and Schomerus. See Sections 3, 5, 9, 10, 11, 26, 28 in [11].

In Sect. 8 of [11] we prove two main abstract uniqueness results for associativity morphisms and braided symmetries in a semisimple pre-tensor category with a generating object. These are Theorem 8.13 and its extension Theorem 8.19. We explain here briefly how Theorem 8.19 applies to imply part (c) of Theorem 2.1.

The proof of parts (b) and (c) is based on the general theory of semisimple weak quasi-Hopf algebras, weak Hopf algebras, and the theory of Drinfeld twist treated in Sections 3–6, and 8–16 in [11]. This abstract theory splits in two parts. Theory on which part (b) of Theorem 2.1 is based is developed in Sections 3–6, and 9–16. The applicative part of the proof of part (b) is developed in Sections 19–36. The abstract aspects that lead to the proof of part (c) of Theorem 2.1 are the main subject of Section 8.

For part (b), we develop an abstract analogue of Drinfeld-Kohno theorem, Theorem 28.4, and apply it to the setting of fusion categories of quantum groups at suitable roots of unity and of module categories of affine vertex operator algebras at positive integer levels. We construct weak Hopf algebras $A_W(\mathfrak{g}, q)$ associated to the quantum group fusion category for all the Lie types in Section 30, by constructing a weak tensor structure on a linear functor from this category to the category of vector spaces, that emerges in [87] for the special tensor products described in (b). The weak tensor structure is constituted by fusion intertwiners with respect to the action of $U_q(\mathfrak{g})$.

By application of our analogue of Drinfeld-Kohno theorem we obtain the structure of a weak quasi-Hopf algebra on the Zhu algebra $A(V_{\mathfrak{g}_k})$ associated to a corresponding affine vertex operator algebra $V_{\mathfrak{g}_k}$. Thus a natural weak quasitensor structure defined by fusion intertwiners with respect to the action of \mathfrak{g} compatible with the fusion rules of modules of the vertex operator algebra. This induces the structure of a ribbon braided tensor category on the module category of $V_{\mathfrak{g}_k}$, via transport from the the Zhu algebra $A(V_{\mathfrak{g}_k})$.

By construction, this structure is ribbon braided tensor equivalent to the structure arising from the quantum group fusion category. The equivalence is explicitly described by a Drinfeld twist analogous to the Drinfeld twist appearing in Drinfeld's first proof of Drinfeld-Kohno theorem [24]. In our case, the unitary structure of the fusion category of quantum groups at roots of unity in [87] is used in the construction of the Drinfeld twist of $A_W(\mathfrak{g}, q)$.

On the module category of $V_{\mathfrak{g}_k}$ we also have the ribbon braided tensor structure constructed by Huang and Lepowsky, and we need to compare the two structures.

5. OUTLINE OF THE CONSTRUCTION OF A WEAK QUASI-TENSOR STRUCTURE ON THE ZHU ALGEBRA $A(V_{\mathfrak{g}_k})$ VIA WENZL QUANTISATION ISOMORPHISM AND DRINFELD TWIST METHOD.

Let $(\mathcal{C}, \otimes, \iota, \alpha)$ be a semisimple complex tensor category (possibly braided, rigid, ribbon) and let

$$\mathcal{F} : \mathcal{C} \rightarrow \text{Vec}$$

be a faithful linear functor. There are two natural ways to make the representation category $\text{Rep}(A)$ of $A = \text{Nat}_0(\mathcal{F})$ into a tensor category (or a category with the same type of structure as \mathcal{C}) equivalent to \mathcal{C} with all the structure.

The first way is traditionally referred to as Tannaka-Krein duality. One needs a (weak, quasi-) tensor structure (F, G) on \mathcal{F} , and then apply Theorem 6.6 to obtain a (weak, quasi)-bialgebra structure on $A = \text{Nat}_0(\mathcal{F})$ (with extra structure) with the desired equivalence property. The (weak quasi-) tensor structure

$$(\mathcal{F}, F, G) : \mathcal{C} \rightarrow \text{Vec}$$

on a given functor, may be easily identified in some cases, e.g. representation categories of groups, simple complex Lie algebras, or quantum groups or more generally whenever one has a tensor category that naturally admits a tensor functor to Vec , as is the example of the compact quantum groups of Woronowicz. In these cases, Tannakian constructions are ubiquitous.

But in other cases the structure (F, G) needs to be constructed, and in this case the situation is perhaps similar to the problem met by Doplicher and Roberts and Deligne in their abstract duality theories. In Doplicher-Roberts duality for compact groups, the embedding functor itself \mathcal{F} needed to be constructed from an abstract symmetric strict tensor category satisfying suitable conditions, and the tensor structure of \mathcal{F} was trivial.

The Tannakian construction endows A with an associator which depends on both the associativity morphisms α of \mathcal{C} and the structure (F, G) , following Majid version for Drinfeld quasi-Hopf algebras. Thus Tannaka-Krein duality is particularly useful in cases where the associativity morphisms α are not too complicated, in that in this case it puts emphasis on F and G . This may allow to understand most of the associativity property of the category by understanding F and G .

Fusion categories that arise from quantum groups at roots of unity, by the work of many authors, among them Drinfeld, Jimbo, Lusztig, Andersen, Wenzl, are realized in a natural way as strict fusion categories. However, in this case tensor functors to Vec do not exist. A natural linear functor has been naturally constructed in [87] for the roots of unity of interest for the aims of [11], i.e. those that are related to the affine Lie algebras at positive integer levels.

Tensor structures on Wenzl functor \mathcal{F} do not exist in general, and one has to construct more general structures, from the inner structure of the category, in order to reduce to the weak quasi-bialgebra method.

A suitable notion of weak Hopf algebra property is a natural concept to look for, by the analogy with Drinfeld-Kohno theorem that first motivated the work of physicists Mack and Schomerus in [61]. The adjective weak refers to the need of considering truncated fusion rules that respect fusion rules of physical models, and rule out representations of $U_q(\mathfrak{g})$ at roots of unity of vanishing quantum dimension. They accordingly introduced the notion of weak analogues of Drinfeld quasi-Hopf algebras, and called them weak quasi-Hopf algebras, a key concept that we rely on. They constructed a semisimple weak quasi-Hopf algebra associated to a suitable model in conformal field theory for \mathfrak{sl}_2 and truncated fusion rules. However, the weak Hopf property was overlooked by them.

We introduce a notion of weak Hopf algebra in [11]. This property was first noticed in [10] in the type A .

In Sect. 30 we shall construct a weak tensor structure on Wenzl functor, and derive a weak Hopf finite dimensional C^* -algebra $A_W(\mathfrak{g}, q)$ associated to fusion categories of quantum groups at roots of unity for all Lie types, with extra structure, that is the braiding, unitary structures given by a Drinfeld coboundary matrix, also rigid and ribbon, extending the work of [10] for \mathfrak{sl}_N in turn extending and expanding the work by Mack and Schomerus for \mathfrak{sl}_2 [61].

We use the weak Hopf algebra property of $A_W(\mathfrak{g}, q)$ as a replacement of the Hopf algebra property of $U_h(\mathfrak{g})$ in Drinfeld-Kohno theorem. This property indeed plays a crucial role in our work, but in a different way as compared to Drinfeld's proof of Drinfeld-Kohno theorem. In our case, we use the weak Hopf property jointly with fusion and duality properties of a fundamental generating representation V in the application, to study the same aim as Drinfeld, comparison of the braided symmetries and associativity morphisms in the two settings of fusion categories of quantum groups at roots of unity and of module categories of vertex operator algebras.

The unitary aspects of our weak Hopf algebras $A_W(\mathfrak{g}, q)$ are based on the results of [87], on the use of Drinfeld coboundary matrix.

The weak tensor property as compared to the weak quasi-tensor property may prove useful for further constructions of the quantum fields from the observables, with the aim of extending the work of Doplicher and Roberts in high dimensional algebraic quantum field theory, as compared to previous attempts in the literature on low dimensional algebraic quantum field theory, that suffered the difficulties due to a possibly more complicated associator.

6. MAIN TOOLS TO PROVE (A): HOW TO MAKE $\text{Rep}(A(V_{\mathfrak{g}_k}))$ INTO A BRAIDED TENSOR CATEGORY WITH HUANG-LEPOWSKY STRUCTURE.

A second way to construct a braided tensor category structure on $\text{Rep}(A)$ from a braided tensor category \mathcal{C} is useful when \mathcal{C} has a natural faithful linear functor

$$\mathcal{F} : \mathcal{C} \rightarrow \text{Vec}$$

given from the start and a braided tensor category structure $(\mathcal{C}, \otimes, \iota, \alpha, d)$ is constructed as a second step. Then in this case it may be more natural to first consider the discrete algebra $A = \text{Nat}_0(\mathcal{F})$ and then transfer of all the structure of \mathcal{C} to the representation category of A .

We discuss this transfer in Sect. 15 without the need of a weak quasi-tensor structure on \mathcal{F} , in abstract terms. One only needs an inverse linear equivalence $\mathcal{S} : \text{Rep}(A) \rightarrow \mathcal{C}$ of the Tannakian linear equivalence $\mathcal{E} : \mathcal{C} \rightarrow \text{Rep}(A)$. In this situation we may define a braided tensor category structure in $\text{Rep}(A)$ from the braided tensor category structure in \mathcal{C} with the property that $\text{Rep}(A)$ is equivalent to \mathcal{C} as a braided tensor category. The tensor

product, associativity morphisms, braiding morphisms of $\text{Rep}(A)$ are explicitly defined via the equivalence \mathcal{E} and its inverse \mathcal{S} .

Far-reaching historic examples of such \mathcal{F} may be found in conformal field theories (e.g. in the setting of models of conformal nets and in that of vertex operator algebras) as the *minimum energy functor*, before the braided tensor category structure was constructed. In these cases we refer to the braided tensor category structure of modules of vertex operator algebras by Huang and Lepowsky or to the unitary braided tensor category structure of localized endomorphisms of conformal nets.

In the setting of vertex operator algebras, the minimum energy functor is also known as Zhu's functor [93]. Notably, Zhu constructed a semisimple algebra $A(V)$ directly from modules of the vertex operator algebra V , naturally isomorphic to $\text{Nat}_0(\mathcal{F})$. This algebra is well known, and is referred to as the Zhu algebra. He also constructed a linear equivalence $\mathcal{E} : \text{Rep}(V) \rightarrow \text{Rep}(A(V))$ and also a natural linear inverse equivalence $\mathcal{S} : \text{Rep}(A(V)) \rightarrow \text{Rep}(V)$ in the same paper if the vertex operator algebra V satisfies suitable semisimplicity conditions.

Section 15 abstracts Zhu linear equivalence and its inverse, by considering a pair of the Tannakian linear equivalence $\mathcal{E} : \mathcal{C} \rightarrow \text{Rep}(A)$ associated to $\mathcal{F} : \mathcal{C} \rightarrow \text{Vec}$ and a linear inverse equivalence $\mathcal{S} : \text{Rep}(A) \rightarrow \mathcal{C}$ and describes how to make $\text{Rep}(A)$ into a braided tensor category.

Thus, in the case of vertex operator algebras satisfying semisimplicity conditions, one may reduce the analysis of the category of infinite dimensional representations of the vertex operator algebra V to a category of finite dimensional representations of the Zhu algebra $A(V)$ in a natural way. This transfer is the passage described in part (a) of the statement of Theorem 2.1 for the affine vertex operator algebras at a positive integer level k .

We show in [11] that a weak quasi-tensor structure (F_0, G_0) can be associated to the minimum energy functor \mathcal{F} in the case of affine vertex operator algebras at positive integer level. In other words at the end we shall construct natural weak quasi-tensor structures (\mathcal{F}, F_0, G_0) , associated to the weak tensor structure of Wenzl functor of a corresponding fusion category of a quantum group at a root of unity. Since the latter is a rigid and modular tensor category, we in particular directly obtain rigidity and modularity of $\text{Rep}(V_{\mathfrak{g}_k})$ from the corresponding quantum group fusion category. In this way we rediscover a result by Huang for the special case of affine vertex operator algebras at a positive integer level [51].

In the final sections of [11] we discuss in greater detail how Huang and Lepowsky theory together with previous work by Frenkel and Zhu provides \mathcal{F} naturally with the weak quasi-tensor structure (F_0, G_0) , such that F_0 is the 0-grade part of Huang-Lepowsky tensor structure map F .

We shall first identify a quantum group fusion category at a fixed suitable root of unity with that of a corresponding affine VOA fusion category or conformal net fusion category as pre-tensor categories, then we identify them with all the structure on special variables which do not generate all the variables linearly. By analysis of a cohomological kind on any braided tensor category with a generating representation, and further braid group duality properties well studied in the literature for quantum groups, we shall extend this

identification to all variables. We shall give a more detailed overview of our proof of Theorem 2.1 in the following sections.

7. A CENTRAL METHOD IN THE QUANTUM GROUP AND VOA SETTING: THE
FUNDAMENTAL GENERATING REPRESENTATION V FOR ALL LIE TYPES, FOLLOWING
H. WENZL.

A central method of proof for both parts (b) and (c) is the use of the fundamental generating representation V and its properties, following primarily the work of Wenzl. The paper [87] plays a main role especially for the proof of (b). For part (c) we shall later describe results and related works in the literature on which our proof is based.

Part (b) of Theorem 2.1 gives coincidence of the two braided tensor structures for all Lie types (two pairs of associativity morphisms together with their braiding morphisms satisfying the two hexagonal diagrams) on a specific collection of their variables, in the two settings of quantum groups and vertex operator algebras. These collections do not generate additively all the variables of the braiding morphisms or associativity morphisms, because they do not contain all the irreducible objects.

Coincidence of the associativity morphisms and braiding morphisms on the special objects is verified in a direct way, relying on similar structural properties of fusion of two corresponding generating representations in the two cases. In the quantum group case, fusion arises in the setting of non-semisimple representations of $U_q(\mathfrak{g})$ considered in [87], while in the vertex operator algebra case fusion arises in a semisimple setting. To compare the braiding and associativity morphisms on the special variables, we use a Drinfeld twist that we construct trivialising, in a sense, the non-trivial unitary structure in the setting of representations of quantum groups at roots of unity to that of the affine vertex operator algebra. This idea emerges in the construction and analysis of the unitary structure of the fusion category of quantum groups at roots of unity of [87].

The twist takes one unitary structure of a special tensor product arising in the quantum group setting from [87] to the classical unitary structure corresponding to the compact real form of \mathfrak{g} (and this in turn remarkably leads to the construction of a unitary tensor structure on the module category of the vertex operator algebra $V_{\mathfrak{g}_k}$ in a uniform way). This implies that the twist leaves the corresponding structural properties invariant on these special variables. To see this, we work on the side of the vertex operator algebras and we compare with the work by Frenkel and Zhu [34]. An abstract concept describing these common structural properties of the associativity morphisms is introduced in Sect. 8 with the name of a \mathcal{V} -pre-associator of CFT type. The name is motivated by our understanding of the structure of Huang-Lepowsky associativity morphisms.

Direct verification of coincidence of the two braided tensor structures on all the irreducible objects is more complicated due to multiplicity of fusion and non-semisimplicity of $U_q(\mathfrak{g})$. To this aim rather than pursuing a closer analogy with the original Drinfeld-Kohno theorem for all objects, we develop two uniqueness theorems, Theorem 8.13 and its generalization Theorem 8.19. This generalization is made necessary by the actual conclusions reached in our work in the applicative sections on the comparison between special cases

of the two associativity morphisms and their corresponding braided symmetries. Thus the uniqueness theorem 8.19 is formulated with suitable assumptions, where again in the application the abstract notion of \mathcal{V} -pre-associator of CFT-type for the fundamental representation V is used in a central way both on the side of quantum groups and that of vertex operator algebras to verify coincidence of these special braid and associativity morphisms, after application of our Drinfeld-Kohno theorem. But this brief description of Theorems 8.13 and Theorem 8.19 put together with coincidence of the two braided tensor structures on special collections of objects proved in part (b), is still not enough to describe our method of proof to complete the proof of part (c) and with it the proof of Theorem 2.1.

We need more that we describe in the following sections. The first next step is a somewhat complicated cohomological reduction work done in Theorems 8.13 and 8.19, where we use cohomology of the pentagon equation and its interplay with the hexagonal equations defining the braided symmetry for a general semisimple braided tensor category endowed with a generating object.

The cohomological reduction is useful to reduce the problem to the last step, which uncovers a central duality property between the fundamental representation of the quantum group and corresponding representations of the braid group in the intertwining spaces of the fundamental representation, in the various Lie types. The first example of this is Schur-Weyl duality in the type A case, and works in the literature that we shall recall greatly extend this first classical example to other Lie types and to the quantum cases. We shall come back to this, but let us anticipate similarities and differences with the original Drinfeld-Kohno theorem, useful for the proof of (b), that will also be useful to clarify how we were led to uncover the key role of duality of the fundamental representation V , for the proof of the last step (c).

8. MAIN TOOLS TO PROVE (B): ANALOGIES AND DIFFERENCES BETWEEN DRINFELD-KOHN THEOREM AND THEOREM 28.4 IN [11]

We next outline general ideas that make our strategy possible, useful to prove part (b) of Theorem 2.1.

As already explained, the starting point of our work is the paper [87]. In that paper, a fundamental generating representation V is described for each simple Lie algebra \mathfrak{g} , and it is used for the construction of the unitary structure in the associated quantum group fusion category. In our approach we follow the work of Wenzl on the use of the generating representation V .

Secondly, in our approach the notion of coboundary matrix first introduced by Drinfeld in his work on Drinfeld-Kohno theorem for quasi-Hopf algebras in a formal setting, plays a major two-fold role. In our case we adopt the idea of Drinfeld coboundary that we describe for complex weak quasi-Hopf algebras in an analytic setting. This general theory is considered in Sections 26, 27, 28.

On one hand the coboundary matrix in an analytic setting induces a unitary structure which shows striking differences with unitary structures of usual Hopf C^* -algebras, such as for example the compact quantum groups, where the $*$ -involution commutes with the

coproduct. Drinfeld original coboundary is used in [87] in the setting of representations of the quantized universal enveloping algebra $U_q(\mathfrak{g})$ with q a suitable type of root of unity, to construct the unitary braided tensor structure of the associated fusion category. This already gives a striking connection between unitary structures and ribbon braided tensor equivalences established by Drinfeld in Drinfeld-Kohno theorem, for different categories.

Similarities with the original Drinfeld-Kohno theorem regard the treatment of special braiding morphisms, that is based as in Drinfeld proof on a twist given by a square root of the Drinfeld coboundary matrix. This approach is particularly useful for braiding on tensor products of the form $V_\lambda \otimes V$ or $V \otimes V_\lambda$, where multiplicity-free and structural fusion properties of V allow for a direct verification of the equivalence between the two braided symmetries on these objects. Moreover in our case on special triples (or n -tuples) where two (or consecutive $n - 1$ -tuples) coordinates are fixed to V , verification of the equivalence of the two associativity morphisms similarly put in a nutshell reduces to the same properties of V . We do not use the KZ equation to this aim, or in any part of our work, other than the fact that the KZ equation was used by Drinfeld, Kazhdan and Lusztig, and by Huang and Lepowsky in the setting of affine vertex operator algebras, to construct the structure of a braided tensor category.

9. DIFFERENCES BETWEEN THE ORIGINAL DRINFELD-KOHNO THEOREM AND OUR ANALOGUE THEOREM 28.4 IN THE TREATMENT OF THE ASSOCIATIVITY MORPHISMS, TO PROVE (C).

Differences regard mainly the treatment of the associativity morphisms, that in our case as before explained resorts to determining the braiding morphisms via an analogue of Drinfeld square root construction of the coboundary matrix, and to braid group duality for V and to cohomology of a braided tensor structure of a generating representation. Thus in our case the method of proof for coincidence of the associativity morphisms depends on the braid group action primarily and categorical cohomological methods of a kind perhaps different than those used by Drinfeld in the original Drinfeld-Kohno theorem. It would be interesting to study whether the two methods are related.

Differently from the original Drinfeld-Kohno theorem, we do not study the problem of equivalence associativity morphisms in our version version of Drinfeld-Kohno theorem. We do this in Section 8 in [11] with the mentioned different ideas that appeal to duality properties of the fundamental representation V with respect to the representation of the braid group in the tensor powers of V .

More specifically, we first construct semisimple weak Hopf algebras $A_W(\mathfrak{g}, q)$ associated to fusion categories of quantum groups at roots of unity for all Lie types. We show that $A_W(\mathfrak{g}, q)$ satisfy a triviality property of the associator, which makes the defining weak quasi-tensor functor (via a suitable Tannakian duality construction) into a weak tensor functor.

These $A_W(\mathfrak{g}, q)$ play the role of the quantum group over a formal variable in Drinfeld-Kohno theorem. This construction is based on the analysis in [87] and the triviality property of the associator is based on properties observed by Gelfand and Kazhdan for

morphisms between non negligible and negligible objects [36], following the same ideas as in [10] in the type A case.

Then we use a quantization continuous curve that is described in [87] to connect irreducible representations of $A_W(\mathfrak{g}, q)$ with irreducible representations of the Zhu algebra $A(V_{\mathfrak{g}_k})$ associated to a corresponding affine Lie algebra, or vertex operator algebra at a positive integer level.

We apply the natural isomorphism given by extremes of Wenzl continuous curve defining quantization of a suitable finite class of representations of the classical Lie algebra (the Weyl alcove defined by the roots of unity) to representations of $U_q(\mathfrak{g})$. Then we apply our Drinfeld-Kohno theorem to endow the Zhu algebra with the structure of a weak quasi-Hopf algebra with coboundary matrix.

These two passages, in analogy with the original Drinfeld-Kohno theorem, allow to construct the structure of a weak quasi-Hopf algebra in our case on the Zhu algebra $A(V_{\mathfrak{g}_k})$. At this point we can use Zhu's linear equivalence and transport the ribbon braided tensor structure to the module category of the associated affine vertex operator algebra. As already explained we verify that braided symmetry and associativity morphisms correspond to the correct braided symmetry and associativity morphisms in the setting of vertex operator algebras on special pairs or triples of representations of the weak quasi-Hopf algebra.

The unitary structure associated to a coboundary of a semisimple weak quasi-Hopf algebra allows directly unitary representations of the braid group, and, similarly to the case of $U_q(\mathfrak{g})$, anticommutes with the comultiplication. This property resembles the antimultiplicativity property of the $*$ -involution in a noncommutative topological space. The unitary structure enters in the construction of the tensor equivalence in our case via an analytic construction of a Drinfeld twist first performed by Drinfeld for the Drinfeld category.

Drinfeld coboundary is closely related to the ribbon structure, and thus it is relevant for the ribbon preserving property of the KLF equivalence, similarly to the case of Drinfeld-Kohno theorem.

10. MAIN TOOLS TO PROVE (C): COHOMOLOGICAL REDUCTION OF A BRAIDED TENSOR STRUCTURE OF A CATEGORY WITH A GENERATING OBJECT V TO BRAID GROUP DUALITY OF V

Theorem 8.13 and Theorem 8.19 show uniqueness of the associativity morphisms (and also braiding morphisms in the latter case) under some assumptions. These assumptions all aim to determine the associativity morphisms uniquely via data from the braiding, but lead to analysis of a different nature in the application.

The first assumption requires knowledge that in our application, via the methods previously alluded to (i.e. Wenzl isomorphism given by a continuous curve describing quantization of all the irreducible representations of the fusion category [87] and recalled also in

[11], combined with application of our Drinfeld twist) one gets a braided symmetry coinciding, for all pairs of objects, with the braided symmetry associated to Huang-Lepowsky braided tensor category structure.

Moreover in Theorem 8.13 we require that the associativity morphisms coincide on the special triples. These assumptions allow a cohomological analysis on the braided symmetry of a tensor category with a generating object. This analysis does not suffice to complete the proof Theorem 8.13. Before describing the further key property of the fundamental representation that we need, we add some more comments on our methods regarding this part.

In the application on the other hand, while as discussed in the previous parts we reach coincidence of associativity morphisms on special triples thanks to the structure of Huang-Lepowsky associativity morphisms and Drinfeld twist method, regarding the braiding morphisms we limit ourselves to verify the coincidence of the two braided symmetries on the special pairs of objects where one of the variables is fixed to V . These special braiding morphisms generate all the other braiding morphisms, but the generating property of these special morphisms depends also on the associativity morphisms. This poses the need of a more detailed analysis to obtain a uniqueness theorem for both braiding and associativity morphisms based on knowledge of their coincidence on special variables that in our approach we can verify directly. This is the aim of Theorem 8.19, with respect to Theorem 8.13, which on the other hand is useful to present the main ideas of how a full braided symmetry determines the associativity morphisms in case of a generating representation.

The assumptions on coincidence of the associativity morphisms and braided symmetry as required in Theorem 8.19 are verified in the proof of Theorem 2.1 on the side of vertex operator algebras. For the associativity morphisms, methods are similar to those used to verify the assumption of Theorem 8.13. We apply Wenzl isomorphism and Drinfeld twist construction and then we use the structural properties of the Huang-Lepowsky associativity morphisms in Theorem 31.9. For the special braiding morphisms we take into account the formulas given by Toledano-Laredo in Theorem 31.12. These two comparisons first motivated our method of proof for both braided symmetry and associativity morphisms.

The proof of (c) of Theorem 2.1 is reduced to the proof of (b) and application of Theorem 8.19. More in detail, for part (c) we use the fact that the special variables of the associativity morphisms and braided symmetry generate all the triples multiplicatively, in the sense that all the components except one in the variables of the associativity and braiding morphisms are taken by the generating representation in each Lie type. In the proof of Theorem 8.19, and the antecedent Theorems 8.13, we explain the generating property of these special values of the associativity and braiding morphisms with the pentagon equation of associativity morphisms and the hexagonal equations of a braided symmetry in a semisimple tensor category with a generating object but, as anticipated, another important property concerning the braid group representations in the category is needed, to determine the associativity morphisms uniquely.

11. THE ROLE OF BRAID GROUP DUALITY FOR THE GENERATING OBJECT IN DETERMINING THE FULL BRAIDED TENSOR STRUCTURE FOR THE PROOF OF (C).

The uniqueness Theorems 8.13 and 8.19 both apply under a central duality property of the fundamental representation V with respect to the braid group representation in the quantum group tensor category, which is at the core of quantum group representation theory. The first instance of this property is Schur-Weyl duality for the general linear group. This duality says that the algebra of intertwining operators, from a tensor power of the vector representation V of the general linear group to itself (also called centralizer algebras for the tensor powers of the vector representation) is generated by permutation operators [91].

The generalization of Schur-Weyl duality to Lie types beyond A in classical Lie groups and quantum groups is an enormous subject developed by many authors, extending from the classical more familiar cases of type A and vector representation to other Lie types beyond A , with respect to a specified generating representation V , and to the quantum group case.

We need the property that the centralizer algebras (V^n, V^n) with respect to the action of the Drinfeld-Jimbo quantum group $U_q(\mathfrak{g})$ are generated by the natural representation of the braid group induced by the R -matrix. In our case q is a root of unity as usually assumed corresponding to the unitary cases of the associated fusion category, specified in [87], [82], see also Definition 20.2 in [11]. The tensor powers V^n appearing in our setting are canonical choices of truncations of the full tensor powers of the quantum group $U_q(\mathfrak{g})$ obtained by induction over n .

The construction of the iterated truncations V^n of $U_q(\mathfrak{g})$ at the root of unity is explained in [87] and is the starting point to construct our weak Hopf algebras in Sections 29 and 30. The duality property between the truncated tensor powers V^n of the fundamental representation and the representation of the braid group is needed as a further property to complete the proof of FKL theorem on coincidence between associativity morphisms and braiding morphisms on all pairs and triples and is used in our cohomological theorems 8.13 and 8.19. These theorems complete with cohomological arguments what was proved up to that point in our work, coincidence of these structure maps on pairs and triples where one or two coordinated are fixed to the fundamental representation V . To do so, the way how our weak Hopf algebras are described in the application to the quantum group fusion category is also used.

For all the Lie types, the truncated tensor powers V^n are inductively defined over n by a canonical decomposition, which is into irreducible components for $\mathfrak{g} \neq E_8$, of a full tensor product of representations $V_\lambda \otimes V$ of $U_q(\mathfrak{g})$, with V_λ an arbitrary irreducible of the semisimplified fusion category. Then one selects irreducible summands with nonzero quantum dimension. The fusion of the fundamental representation of E_8 is described in [87] as well, and is more delicate.

Returning to the question of whether the braid group representation generates the centralizer algebras (V^n, V^n) for the truncated tensor powers of $U_q(\mathfrak{g})$ at the root of unity,

this question can be studied in the same way as for the case of a corresponding summand with q not a root of unity, as in the truncated summand the root of unity property is not visible. This is explained among other things in Prop. 2.4 in [87], see also Prop 29.2 for the same result with a slightly more detailed proof, where an important role is played by the continuous curve described in that proposition, that passes through generic values of q . See also Section 3.5 in [87] for the special tensor products $V_\lambda \otimes V$ of representations of $U_q(\mathfrak{g})$. Thus it suffices to have the duality property between the full algebras $(V^{\otimes n}, V^{\otimes n})$ of $U_q(\mathfrak{g})$ when q is not a root of unity.

We recall that the fundamental representation V are listed in Sect. 3.5 in [87] for all the Lie types, and in particular are given by

- The vector representation for Lie types A and C ,
- The spinor representation for Lie types B and D . In particular V is the direct sum of the two spinor representations for type D_n with n even.
- The 7-dimensional representation for Lie types G_2 .

The Lie types A and C for the vector representation are well known. If q is not a root of unity, the centralizer algebras $(V^{\otimes n}, V^{\otimes n})$ in the type A case are described by Hecke algebras, a quotient of the complex braid group algebra by well known generators and relations. The first result in this direction is due to Jimbo [54]. For the Lie types B , C , D , when V is the vector representation, Murakami and Birman and Wenzl introduced and studied what are known as the BMW algebras [67], [4]. A nice exposition of the Hecke and BMW algebras may be found in the book by Goodman, de la Harpe and Jones [38]. Duality for the vector representation is of quantum groups was studied by Kirillov and Reshetikhin [58]. A summary of these results and references to the original papers for the vector representation for all the classical Lie types may be found in Theorem 10.2.5 of [9].

The centralizer algebra of the fundamental representation for types B and D is studied in [81], [89].

Type G_2 was shown by Lehrer and Zhang [60], Morrison [65], and Martirosyan and Wenzl [62].

In the application of Theorems 8.13 and 8.19, we work on the side of the quantum group fusion category and by the results above, our assumptions on braid group duality for the fundamental representation of our weak Hopf algebras $A_W(\mathfrak{g}, q)$ constructed in Section 30 are verified for the defining structure arising from Tannakian construction on the quantum group fusion category $\mathcal{C}(\mathfrak{g}, q)$ and Wenzl functor.

12. MORE REMARKS FROM THE VIEWPOINT OF DOPLICHER-ROBERTS RECONSTRUCTION IN AQFT ON THE RELEVANCE OF DUALITY FOR THE FUNDAMENTAL REPRESENTATION IN THE CONSTRUCTION OF QUANTUM GAUGE GROUP

We also remark that Schur-Weyl duality plays an important role in the abstract duality theory for compact groups by Doplicher and Roberts. It is thanks to the fact that the usual permutation symmetry of the category of Hilbert spaces represents all the the intertwining

operators between tensor powers of the defining representation of the unitary groups $SU(d)$, that their construction of the compact gauge group associated to the category of localized endomorphisms of the observable algebras in algebraic quantum field theory takes place, as a closed subgroup $G \subseteq SU(d)$.

If one regards affine vertex operator algebras at positive integer levels as models corresponding to the case of a gauge group given by the simple compact Lie group $SU(d)$ in Doplicher-Roberts theory of quantum gauge fields (i.e. versus the case of gauge group G given by any compact Lie group (a proper closed subgroup of $SU(d)$), then from this viewpoint one may expect that duality properties for the fundamental representation V of a quantum group should play a central role in our approach to FKL theorem. The analogue of permutation symmetry in our case is its generalization to the braid group symmetry, and therefore it is natural to consider all the Lie types for which braid group duality is known to hold for the generating representation V , as a first step towards a Doplicher-Roberts reconstruction of a field algebra from the local observable algebras in low dimensional quantum field theory.

It may therefore be somewhat surprising that the need to use such an important property as generalizations of Schur-Weyl duality to Lie types beyond A and to the quantum case, has not been noted before as a key property to understand the proof of Theorem 2.1, in the literature but also in the previous versions of [11], given that our approach is originally inspired by Doplicher-Roberts duality theory for compact groups with similar applicative intents to the construction of a quantum gauge group that we construct from the emerging categories in low dimensional quantum field theory as an analogue of a gauge group, aimed to the construction of the quantum field algebra from the local observable algebras.

Perhaps similarly surprising is the fact that in our approach we do not invoke modularity. On the other hand by a private discussion with M. Yamashita, we learned that modularity is used in their recent work on classification of fusion categories with fusion rules of type BCD [39].

Thus the fact that braid group duality for the fundamental representation V was the relevant missing observation helpful to complete the proof of Theorem 2.1 in the previous version of [11], which led us to Theorems 8.13 and 8.19 as a completion of Lemma 7.10 in version 7 of [11], was for us a further important recognition of the relevance of the role of quantum groups. We refer the reader to the beautiful historical paper by V. F. R. Jones [55] for a general first introduction to the origin of quantum groups as solutions to the Yang-Baxter equation.

We find it remarkable the deep application that braid group duality has to quantum field theory. This application led us to evaluate even more the relevance of the quantum group viewpoint than as noted up to that point in our study.

We also would like to point out that our results of the categorical characterization of the class of unitary coboundary weak Hopf algebras that we obtain in Sect. 26, 27, 29 in [11] extends the notion of symmetric functors that plays a key role in Doplicher-Roberts theory [21], [22].

13. OUTLOOK

In Doplicher-Roberts theory, a unique gauge compact subgroup of $SU(d)$ is in particular constructed in the case of a rigid symmetric tensor C^* -category with simple unit and a single generating object, to describe the category when the generating property of the permutation group is not satisfied. Their theory has applications to the category of localized endomorphisms arising from the local observable algebras in high dimensional AQFT, and leads to the construction of the field algebra [21], [22].

For the class of the compact quantum groups of Woronowicz there is a construction of their quantum homogeneous spaces and Hilbert bimodule representations developed for non-symmetric rigid tensor categories with simple unit [75], [79]. The construction of the homogeneous space and bimodule representations of the compact quantum group extends to compact quantum groups the idea of closed subgroup construction in Doplicher-Roberts theory. In that quantum case the homogeneous space is an ergodic action of a compact quantum group of a unital C^* -algebra, and may not arise from a quantum subgroup.

In the setting of compact quantum groups, unitary representations of the braid group are not known to arise in a natural way. On the other hand, they arise from fusion categories at roots of unity in [87], and this paper was our first motivation to study these quantum groups and construct weak Hopf C^* -algebras in the sense of [11], with precedents the examples by Mack and Schomerus for \mathfrak{sl}_2 , described as weak quasi-Hopf algebras in those early papers where the study of the unitary structure was not addressed yet [61], and its extension to \mathfrak{sl}_N studied in [10] where the study of the unitary structure was included following the results of [87].

Our weak Hopf C^* -algebras are suitable extensions of the compact quantum groups, and in our experience the failure of strict co-associativity has played a mild role, and has allowed the development of a theory extending that of Hopf algebras, or of compact quantum groups if one allows twisted unitary structure, called Ω -involutions in [11] for which classes of examples arising from ribbon tensor C^* -categories allow unitary representations of the braid group in a natural way, called unitary coboundary weak Hopf C^* -algebras in the same paper.

It is then desirable to search for further generalizations of quantum homogeneous spaces and bimodule representations to these weak Hopf algebras to describe ribbon braided tensor C^* -categories with unitary representations of the braid group, such as the representation categories of models of CFT: conformal nets where the unitary structure is given from the start, or related vertex operator algebras where is constructed. In the affine case that we have studied in our papers, the unitary, rigid, and modular structure in the latter can be transported from that of the quantum group fusion category via the equivalence. This may be useful for those models for which, in analogy to the theory of Doplicher-Roberts for rigid symmetric tensor categories, the generating property of the braid group is not satisfied for the algebras of intertwining morphisms of the category. Or also for models for which weak Hopf algebras can not arise from failure of submultiplicativity of the integer valued dimension function of representations of the Zhu algebra. Natural classes of models to study in this direction are referred to as coset models.

Such an extension may prove useful to construct a field algebra from a braided tensor C^* -category, for which conformal nets provides examples, from the algebra of local observables, taking into consideration the quantum group fusion categories at roots of unity and the associated weak Hopf C^* -algebras constructed in [11], as developments of early works in that direction of the 90s.

Other interesting future developments may regard applications in noncommutative geometry in the sense of Connes [13] of our weak Hopf algebras $A_W(\mathfrak{g}, q)$ associated to the quantum group fusion category $\mathcal{C}(\mathfrak{g}, q)$ or their twisted weak quasi-Hopf algebra structures on the Zhu algebra of an affine vertex operator algebra at a positive integer level $V_{\mathfrak{g}_k}$ to the study of Dirac operators, by expanding an analog in conformal field theory of the groundbreaking work by Neshveyev and Tuset on Dirac operators for compact quantum groups in the setting of Kazhdan-Lusztig theorem for real values of the quantization parameter q [69].

Acknowledgements The author is grateful to Yi-Zhi Huang for several comments on the proof of modularity for fusion categories of affine vertex operator algebras, and the history of Finkelberg-Kazhdan-Lusztig theorem. She is also grateful to Marco V. Giannone for pointing out to her incompleteness of the proof of Lemma 7.10 in version 7 of [11] and to Hans Wenzl for kind email correspondence and for references on duality for tensor powers of the fundamental representation of Drinfeld-Jimbo quantum groups for the Lie types B , D , G_2 , leading to the present paper. Finally she thanks Pavel Etingof for informing her of [28].

REFERENCES

- [1] B. Bakalov, A.A. Kirillov: Lectures on tensor categories and modular functors; Vol. 21. Providence, AMS, 2001.
- [2] A. Beilinson, B. Feigin and B. Mazur, Introduction to algebraic field theory on curves, preprint (1991).
- [3] A. A. Belavin, A. N. Polyakov, and A. B. Zamolodchikov, Infinite conformal symmetries in two-dimensional quantum field theory, *Nuclear Phys. B* 241 (1984), 333–380.
- [4] J. Birman, H. Wenzl: Braid, links and a new algebra, *Tran Amer. Math. Soc.*, **313** (1989), 239–245.
- [5] G. Böhm, K. Szlachanyi: A coassociative C^* -quantum group with nonintegral dimensions, *Lett. Math. Phys.*, **38** (1996), 437–456.
- [6] Carpi et al, *Commun. Math. Phys.*, **390** (2022), 169.
- [7] Carpi et al, *Commun. Math. Phys.* **402** (2023), 169.
- [8] Adamo et al: *Commun. Math. Phys.* **404**, (2023), 1231.

¹The claims, arguments, and reasoning of the recent papers [6], [7], [8] are essentially dependent on the results and in-depth discussions between S.C. and C.P. that took place from 2018 to 2022 on the contents of an entire project including [11] and the topics of the above papers, during a period in which S.C. was a co-author, before breaking the collaboration in early 2023, with a minor contribution in a draft of a single section. In addition, C.P. department digital devices were subject to a serious hacking attack for several months in 2021 during the previous collaboration, which was recognized by authorities, that severely damaged her intellectual property and health.

- [9] V. Chari, A.N. Pressley: A guide to quantum groups. Cambridge University Press, 1995.
- [10] S. Ciamprone, C. Pinzari: Quasi-coassociative C^* -quantum groupoids of type A and modular C^* -categories, *Adv. Math.*, **322** (2017), 971–1032.
- [11] S. Ciamprone, M.V. Giannone, C. Pinzari: Weak quasi-Hopf algebras, C^* -tensor categories and conformal field theory, and the Kazhdan-Lusztig-Finkelberg theorem, arXiv:2101.10016 [math.QA].
- [12] L.S. Cirio, A. D’Andrea, C. Pinzari, S. Rossi, Connected components of compact matrix quantum groups and finiteness conditions. *J. of Funct. Anal.* (2014), **267**, 3154–3204.
- [13] A. Connes: Noncommutative Geometry, Academic Press (1994).
- [14] A. D’Andrea, C. Pinzari, S. Rossi: Polynomial growth of discrete quantum groups, topological dimension of the dual and $*$ -regularity of the Fourier algebra, *Ann. de l’Institut Fourier*, **67**, (2017), 2003–2027.
- [15] S. Doplicher, R. Haag, and J.E. Roberts: Fields, observables and gauge transformations I, *Commun. Math. Phys.*, **13** (1969), 1–23.
- [16] S. Doplicher, R. Haag, and J.E. Roberts: Fields, observables and gauge transformations II, *Commun. Math. Phys.*, **15** (1969), 173–200.
- [17] S. Doplicher, R. Haag, J.E. Roberts: Local Observables and Particle Statistics I, *Comm. Math. Phys.* **28** (1971), 199–230.
- [18] S. Doplicher, R. Haag, J.E. Roberts: Local Observables and Particle Statistics, II, *Comm. Math. Phys.*, **35** (1974), 49–85.
- [19] S. Doplicher, J.E. Roberts: Duals of compact Lie groups realized in the Cuntz algebras and their actions on C^* -algebras, *J. Funct. Anal.*, **74** (1987), 96–120.
- [20] S. Doplicher, J.E. Roberts: Endomorphisms of C^* -algebras, cross products and duality for compact groups, *Ann. Math.*, **130** (1989), 75–119.
- [21] S. Doplicher, J.E. Roberts: A new duality theory for compact groups, *Invent. Math.*, **98** (1989), 157–218.
- [22] S. Doplicher, J. E. Roberts, Why there is a field algebra with a compact gauge group describing the superselection structure in particle physics, *Commun. Math. Phys.*, **131** (1990), 51–107.
- [23] V. G. Drinfeld: On almost cocommutative Hopf algebras, *Algebra Anal.* **1** (1989), n. 2, 30–46; English transl. in *Leningrad Math. J.* **1** (1990).
- [24] V. G. Drinfeld: Quasi-Hopf algebras, *Leningrad Math. J.*, **1** (1990), 1419–1457.
- [25] V. G. Drinfeld: On quasitriangular quasi-Hopf algebras and on a group that is closely connected with $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$, *Leningrad Math. J.*, **2** (1991), n. 4, 829–860.
- [26] P. Etingof, S. Gelaki, D. Nikshych, V. Ostrik: Tensor Categories. Mathematical surveys and monographs, AMS, Volume 205, 2015.
- [27] P. Etingof, D. Nikshych, V. Ostrik: On fusion categories, *Ann. of Math.* , **162** (2005), 581–642.
- [28] P. Etingof, D. Penneys: Rigidity of non-negligible objects of moderate growth in braided categories, arXiv:2412.17681 [math.QA].
- [29] B.L. Feigin and D.B. Fuks: Invariant skewsymmetric differential operators on the line and Verma modules over the Virasoro algebra, *Funct. Anal. Appl.* **16** (1982), 114–126.
- [30] M. Finkelberg: An equivalence of fusion categories, *Geom. Funct. Anal.*, **6** (1996), 249–267.
- [31] M. Finkelberg: Erratum to: an equivalence of fusion categories, *Geom. Funct. Anal.*, **23** (2013), 810–811.
- [32] K. Fredenhagen, K.-H. Rehren, and Bert Schroer: Superselection sectors with braid group statistics and exchange algebras, *Comm. Math. Phys.* **125** (1989) 201–226.
- [33] K. Fredenhagen, K.-H. Rehren, and B. Schroer: Superselection sectors with braid group statistics and exchange algebras II: Geometric aspects and conformal covariance, *Rev. in Math. Phys.* **4** (1992): 113–157.
- [34] I.B. Frenkel, Y.Zhu: Vertex operator algebras associated to representations of affine and Virasoro algebras, **66** *Duke Math. J.*, (1992) 123–168.

- [35] T. Gannon: Moonshine beyond the Monster, the bridge connecting algebra, modular forms and physics, Cambridge University Press, 2006.
- [36] S. Gelfand, D. Kazhdan: Examples of tensor categories, *Inv. Math.* **109** (1992), 595–617.
- [37] P. Goddard, A. Kent, and D.I. Olive: Unitary representations of the Virasoro and super-Virasoro algebras, *Commun. Math. Phys.*, **103** (1986), 105–119.
- [38] F.M. Goodman, P. de la Harpe, and V.F.R. Jones: Coxeter graphs and towers of algebras (Vol. 14). Springer Science and Business Media.
- [39] P. Grossman, S. Neshveyev. M. Yamashita, private communication with M. Yamashita.
- [40] R. Haag and D. Kastler: An algebraic approach to field theory, *J. Math. Phys.*, **5** (1964), 848–861.
- [41] R. Haag: Local quantum physics: Fields, particles, algebras. Springer Science and Business Media, 2012.
- [42] Y.-Z., Huang, J. Lepowsky: A theory of tensor products for module categories for a vertex operator algebra, I, *Selecta Math.*, **1**(1995), 699–756.
- [43] Y.-Z., Huang, J. Lepowsky: A theory of tensor products for module categories for a vertex operator algebra, II, *Selecta Math.* **1** (1995), 757–786.
- [44] Yi-Zhi Huang, James Lepowsky: A theory of tensor products for module categories for a vertex operator algebra, III, *Journal Pure Appl. Alg.*, **100** (1995) 141–171.
- [45] Y.-Z. Huang, A theory of tensor products for module categories for a vertex operator algebra, IV, *J. Pure Appl. Alg.*, **100** (1995) 173–216.
- [46] Yi-Zhi Huang, James Lepowsky: Intertwining operator algebras and vertex tensor categories for affine Lie algebras, *Duke Math. J.*, **99** (1999), 113–134.
- [47] Y. Z. Huang: Vertex operator algebras, the Verlinde conjecture, and modular tensor categories, *Proc. Natl. Acad. Sci. USA*, **102** (2005), 5352–5356.
- [48] Y.-Z. Huang: Differential equations and intertwining operators, *Comm. Contemp. Math.*, **7** (2005), 375–400.
- [49] Y.-Z. Huang: Differential equations, duality and modular invariance *Comm. Contemp. Math.*, **7** (2005), 649–706.
- [50] Y.-Z. Huang: Vertex operator algebras and the Verlinde conjecture, *Comm. Contemp. Math.*, **10** (2008), 103–154.
- [51] Y.-Z. Huang: Rigidity and modularity of vertex tensor categories, *Comm. Contemp. Math.*, **10** (2008), 871–911.
- [52] Y.-Z. Huang, J. Lepowsky: Tensor categories and the mathematics of rational and logarithmic conformal field theory, *J. Phys. A: Math. Theor.* **46** (2013) 494009 (21pp)
- [53] Y.-Z. Huang: Affine Lie algebras and tensor categories, arXiv:1811.05123.
- [54] M. Jimbo: A q -difference analogue of $U(g)$ and the Yang-Baxter equation, *Lett. Math. Phys.*, **10** (1985), 63–69.
- [55] V. F. R. Jones: In and around the origin of quantum groups. arXiv preprint math/0309199.
- [56] V.G. Kac, Contravariant form for infinite-dimensional Lie algebras and Lie superalgebras, Springer Lecture Notes in Physics **94** (1979), 441–445.
- [57] D. Kazhdan, G. Lusztig: Tensor structures arising from affine Lie algebras. I, *J. Amer. Math. Soc.*, **6** (1993), 905–947; II, *J. AMS*, **6** (1993), 949–1011; III, *J. AMS*, **7** (1994), 335–381; IV, *J. AMS*, **7** (1994), 383–453.
- [58] A. N. Kirillov, N. Yu. Reshetikhin: Representation of the algebras $U_q(sl_2)$, q -orthogonal polynomials and invariants of links, in *Infinite dimensional Lie lgebras and groups*, V. G. Kac (Ed), 285–339, World Scientific, Singapore. Yangians and multiplicities of occurrence of the irreducible components of the tensor product of representations of simple Lie algebras *J. Soviet Mth.*, **52**, 3156–3164.
- [59] V. G. Knizhnik and A. B. Zamolodchikov: Current algebra and Wess-Zumino models in two dimensions, *Nuclear Phys.* **B 247** (1984),83–103.
- [60] G.I. Lehrer, R. B.Zhang: Strongly multiplicity free modules for Lie algebras and quantum groups, *J. Alg.*, **306** (2006), 138–174.

- [61] G. Mack, V. Schomerus: Quasi Hopf quantum symmetry in quantum theory, *Nucl. Phys.*, **B 370**, (1992) 185–230.
- [62] L. Martirosyan, H. Wenzl: Affine centralizer algebras, for G_2 , *IMRN*, **12** (2019), 3812–3831.
- [63] G. Moore, N. Seiberg: Polynomial equations for rational conformal field theories, *Phys. Lett.*, **B 212** (1988), 451–460.
- [64] G. Moore and N. Seiberg: Classical and quantum conformal field theory, *Commun. Math. Phys.*, **123** (1989), 177–254.
- [65] S. Morrison: The braid group surjects onto G_2 tensor space, *Pacific J. Math.*, **249** (2011), 189–198.
- [66] M. Müger: Tensor Categories: a selective guided tour, *Rev. Un. Mat. Argentina*, **51 (1)** (2010), 95–163.
- [67] J. Murakami: The Kauffman polynomial of links and representation theory, *Osaka J. Math.*, **24** (1987), 745–758.
- [68] S. Neshveyev, L. Tuset: Notes on the Kazhdan-Lusztig Theorem on Equivalence of the Drinfeld Category and the Category of $U_q(\mathfrak{g})$ -Modules, *Algebr. Represent. Theor.*, (2011) 14: 897.
- [69] S. Neshveyev, L. Tuset: The Dirac operator on compact quantum groups, *J. Reine Angew. Math.*, **641** (2010), 1–20.
- [70] S. Neshveyev, L. Tuset: Compact quantum groups and their representation categories, AMS, 2014.
- [71] S. Neshveyev, M. Yamashita: Classification of non-Kac compact quantum groups of $SU(n)$ type, *IMRN*, **11** (2016), 3356–3391.
- [72] S. Neshveyev, M. Yamashita: Towards a classification of compact quantum groups of Lie type. In *Operator Algebras and Applications: The Abel Symposium 2015* Springer International Publishing, 231–263.
- [73] C. Pinzari: The representation category of the Woronowicz quantum group $S_\mu U(d)$ as a braided tensor C^* -category, *Int. J. Math*, **18** (2007), 113–136.
- [74] C. Pinzari: Embedding ergodic actions of compact quantum groups on C^* -algebras into quotient spaces, *Int. J. Math.*, **18** (2007), 137–164.
- [75] C. Pinzari, J. E. Roberts: A duality theorem for ergodic actions of compact quantum groups on C^* -algebras, *Commun. Math. Phys.*, **277** (2008), 385–421.
- [76] C. Pinzari, J.E. Roberts: Ergodic actions of compact quantum groups from solutions of the conjugate equations, *Kyoto J. Math.* **57**, (2017), 519–552.
- [77] C. Pinzari, J.E. Roberts: Ergodic actions of compact quantum groups from II_1 -subfactors, *J. Geom. Phys.*, **60** (2010), 403–416.
- [78] C. Pinzari, J. E. Roberts: A rigidity result for extensions of braided tensor C^* -categories derived from compact matrix quantum groups, *Comm. Math. Phys*, **306** (2011), 647–662.
- [79] C. Pinzari, J. E. Roberts: A theory of induction and classification of tensor C^* -categories, *J. Noncomm. Geom.* **6** (2012): 665–719.
- [80] E.C. Rowell: From quantum groups to unitary modular tensor categories, *Contemp. Math.*, **413** (2006), 215–230.
- [81] E. Rowell, H. Wenzl: $SO(N)_2$ braid group representations are Gaussians, *Quantum Topol.*, **8** (2017), 1–33.
- [82] S. Sawin: Quantum groups at roots of unity and modularity, *J. Knot Theory*, **15**, (2006) 1245–1277.
- [83] V. Toledano Laredo: Fusion of positive energy representations of $LSpin(2n)$. Ph.D. thesis, St. John’s College, Cambridge, 1997.
- [84] A. Tsuchiya, Y. Kanie: Vertex operators in conformal field theory on \mathbb{P}^1 and monodromy representations of braid group, in *Conformal Field Theory and Solvable Lattice Models*, *Adv. Studies in Pure Math.* **16** (1988), 297–372; Erratum: Vertex Operators in conformal field theory on \mathbb{P}^1 and monodromy representations of the braid group. *Integral systems in quantum field theory and statistical mechanics*, *Advanced studies in pure math.* **19**, Academic Press, Boston 1989, 675–682.
- [85] V. G. Turaev: Quantum invariants of knots and 3-manifolds, *Studies in Mathematics*, Walter de Gruyter, 2010.

- [86] A. Wassermann: Operator algebras and conformal field theory, III. Fusion of positive energy representations of $LSU(N)$ using bounded operators. *Invent. Math.*, **133** (1998), 467–538.
- [87] H. Wenzl: C^* -tensor categories from quantum groups, *J. Amer. Math. Soc.*, **11** (1998), 261–282.
- [88] H. Wenzl: On tensor categories of Lie type E_N , $N \neq 9$, *Adv. Math.*, **177** (2003), 66–104.
- [89] H. Wenzl: Dualities for the spin representations, arXiv:2005.11299 [math.QA].
- [90] E. Witten, Non-abelian bosonization in two dimensions, *Commun. Math. Phys.* **92**, (1984), 455–472.
- [91] H. Weyl: The Classical Groups: Their Invariants and Representations, Princeton University Press, 1946.
- [92] F. Xu: Standard λ -lattices from quantum groups, *Invent. Math.*, **134** (1998), 455–487.
- [93] Y. Zhu: Modular invariance of characters of vertex operator algebras, *J. Amer. Math. Soc.*, **9** (1996), 237–302.

Email address: `pinzari@mat.uniroma1.it`

DIPARTIMENTO DI MATEMATICA, SAPIENZA UNIVERSITÀ DI ROMA, P.LE ALDO MORO, 5 – 00185
ROME, ITALY