Learning Counterfactual Outcomes Under Rank Preservation

Peng Wu¹ Haoxuan Li² Chunyuan Zheng³ Yan Zeng¹ Jiawei Chen⁴ Yang Liu⁵ Ruocheng Guo⁶ Kun Zhang⁷⁸

Abstract

Counterfactual inference aims to estimate the counterfactual outcome at the individual level given knowledge of an observed treatment and the factual outcome, with broad applications in fields such as epidemiology, econometrics, and management science. Previous methods rely on a known structural causal model (SCM) or assume the homogeneity of the exogenous variable and strict monotonicity between the outcome and exogenous variable. In this paper, we propose a principled approach for identifying and estimating the counterfactual outcome. We first introduce a simple and intuitive rank preservation assumption to identify the counterfactual outcome without relying on a known structural causal model. Building on this, we propose a novel ideal loss for theoretically unbiased learning of the counterfactual outcome and further develop a kernel-based estimator for its empirical estimation. Our theoretical analysis shows that the rank preservation assumption is not stronger than the homogeneity and strict monotonicity assumptions, and shows that the proposed ideal loss is convex, and the proposed estimator is unbiased. Extensive semi-synthetic and real-world experiments are conducted to demonstrate the effectiveness of the proposed method.

1. Introduction

Understanding causal relationships is a fundamental goal across various domains, such as epidemiology (Hernán & Robins, 2020), econometrics (Imbens & Rubin, 2015),

Preprint.

and management science (Kallus & Uehara, 2020). Pearl & Mackenzie (2018) define the three-layer causal hierarchy-association, intervention, and counterfactuals-to distinguish three types of queries with increasing complexity and difficulty (Bareinboim et al., 2022). Counterfactual inference, the most challenging level, aims to explore the impact of a treatment on an outcome given knowledge about a different observed treatment and the factual outcome. For example, given a patient who has not taken medication before and now suffers from a headache, we want to know whether the headache would have occurred if the patient had taken the medication initially. Answering such counterfactual queries can provide valuable instructions in scenarios such as credit assignment (Mesnard et al., 2021), root-causal analysis (Budhathoki et al., 2022), and fair decision-making (Imai & Jiang, 2023).

Different from interventional queries, which are prospective and estimate the counterfactual outcome in a hypothetical world via only the observations obtained before treatment (as pre-treatment variables), counterfactual inference is retrospective and further incorporates the factual outcome (as a post-treatment variable) in the observed world. This inherent conflict between the hypothetical and the observed world poses a unique challenge and makes the counterfactual outcome generally unidentifiable, even in randomized controlled experiments (RCTs) (Pearl et al., 2016; Ibeling & Icard, 2020; Bareinboim et al., 2022; Wu et al., 2024).

For counterfactual inference, Pearl et al. (2016) proposed a three-step procedure (abduction, action, and prediction) to estimate counterfactual outcomes. However, it relies on the availability of structural causal models (SCMs) that fully describe the data-generating process (Brouwer, 2022; Xie et al., 2023). In real-world applications, the ground-truth SCM is likely to be unknown, and estimating it requires additional assumptions to ensure identifiability, such as linearity (Shimizu et al., 2006) and additive noise (Hoyer et al., 2008; Peters et al., 2014). Unfortunately, these assumptions are hard to satisfy in practice and restrict the applicability.

To tackle the above problems, several counterfactual learning approaches have been proposed with respect to different identifiability assumptions. For example, Lu et al. (2020), Nasr-Esfahany et al. (2023), and Xie et al. (2023) estab-

¹School of Mathematics and Statistics, Beijing Technology and Business University ²Center for Data Science, Peking University ³School of Mathematical Sciences, Peking University ⁴College of Computer Science and Technology, Zhejiang University ⁵Computer Science and Engineering, University of California, Santa Cruz ⁶ByteDance Research ⁷Department of Philosophy, Carnegie Mellon University ⁸Machine Learning Department, Mohamed bin Zayed University of Artificial Intelligence. Correspondence to: Ruocheng Guo <ruocheng.guo@bytedance.com>.

lished the identifiability of counterfactual outcomes based on homogeneity and strict monotonicity assumptions. The homogeneity assumption posits that the exogenous variable for each individual remains constant across different interventional environments, and the strict monotonicity assumption asserts that the outcome is a strictly monotone function of the exogenous variable given the features. In terms of counterfactual learning, Lu et al. (2020) and Nasr-Esfahany et al. (2023) adopted Pearl's three-step procedure that needs to estimate the SCM initially. In addition, Xie et al. (2023) proposed using quantile regression to estimate counterfactual outcomes that effectively avoid the estimation of SCM. Nevertheless, it relies on a stringent assumption that the conditional quantile functions for different counterfactual outcomes come from the same model and it requires estimating a different quantile value for each individual, leading to a challenging bi-level optimization problem.

In this work, we propose a principled counterfactual learning approach with *intuitive identifiability assumptions and theoretically guaranteed estimation methods*. **On one hand**, we introduce the simple and intuitive rank preservation assumption, positing that an individual's factual and counterfactual outcomes have the same rank in the corresponding distributions of factual and counterfactual outcomes for all individuals. We establish the identifiability of counterfactual outcomes under the rank preservation assumption and show that it is slightly less restrictive than the homogeneity and monotonicity assumptions used in previous studies.

On the other hand, we further propose a theoretically guaranteed method for unbiased estimation of counterfactual outcomes. The proposed estimation method has several desirable merits. First, unlike Pearl's three-step procedure, it does not necessitate a prior estimation of SCMs and thus relies on fewer assumptions than Lu et al. (2020) and Nasr-Esfahany et al. (2023). Second, in contrast to the quantile regression method proposed by Xie et al. (2023), our approach neither restricts conditional quantile functions for different counterfactual outcomes to originate from the same model, nor does it require estimating a different quantile value for each unit. Third, we improve the previous learning approaches by adopting a convex loss for estimating counterfactual outcomes, which leads to a unique solution.

In summary, the main contributions are as follows: (1) We introduce the intuitive rank preservation assumption to identify the counterfactual outcomes with unknown SCM; (2) We propose a novel ideal loss for unbiased learning of the counterfactual outcome and further develop a kernel-based estimator for the ideal loss. In addition, we provide a comprehensive theoretical analysis for the proposed learning approach; (3) We conduct extensive experiments on both semi-synthetic and real-world datasets to demonstrate the effectiveness of the proposed method.

2. Problem Formulation

Throughout, capital letters represent random variables and lowercase letters denote their realizations.

Structural Causal Model (SCM, Pearl, 2009). An SCM \mathcal{M} consists of a causal graph \mathcal{G} and a set of structure equation models $\mathcal{F} = \{f_1, ..., f_p\}$. The nodes in \mathcal{G} are divided into two categories: (a) exogenous variables $\mathbf{U} = (U_1, ..., U_p)$, which represent the environment during data generation, assumed to be mutually independent; (b) endogenous variables $\mathbf{V} = \{V_1, ..., V_p\}$, which denote the relevant features that we need to model in a question of interest. For variable V_j , its value is determined by a structure equation $V_j = f_j(PA_j, U_j), j = 1, ..., p$, where PA_j stands for the set of parents of V_j . SCM provides a formal language for describing how the variables interact and how the resulting distribution would change in response to certain interventions. Based on SCM, we introduce the counterfactual inference problem in the following.

Counterfactual Inference (Pearl, 2009). Suppose that we have three sets of variables denoted by $X, Y, \mathbf{E} \subseteq \mathbf{V}$, counterfactual inference revolves around the question, "given evidence $\mathbf{E} = \mathbf{e}$, what would have happened if we had set X to a different value x'?". Pearl et al. (2016) propose using the three-step procedure to answer the problem: (a) **Abduction**: determine the value of U according to the evidence $\mathbf{E} = \mathbf{e}$; (b) **Action**: modify the model \mathcal{M} by removing the structural equations for X and replacing them with X = x', yielding the modified model $\mathcal{M}_{x'}$; (c) **Prediction**: Use $\mathcal{M}_{x'}$ and the value of U to calculate the counterfactual outcome of Y. In this paper, we focus on estimating the counterfactual outcome for each individual. To illustrate the main ideas, we formulate the common counterfactual inference problem within the context of the backdoor criterion.

Problem Formulation. Let $\mathbf{V} = (Z, X, Y)$, where X causes Y, Z affects both X and Y, and the structure equation of Y is given as

$$Y = f_Y(X, Z, U_X). \tag{1}$$

Let $Y_{x'}$ denotes the potential outcome if we had set X = x'. The counterfactual question, "given evidence (X = x, Z = z, Y = y) of an individual, what would have happened had we set X = x' for this individual", is formally expressed as estimating $y_{x'}$, the realization of $Y_{x'}$ for the individual. Here, we adhere to the deterministic viewpoint of Pearl (2009) and Pearl et al. (2016), treating the value of $Y_{x'}$ for each individual as a fixed constant. According to Pearl's threestep procedure, given the evidence (X = x, Z = z, Y = y)for an individual, the identifiability of its counterfactual value $y_{x'}$ can be achieved by determining the structural equation f_Y and the value of U_X for this individual. This is the key idea underlying most of the existing methods. For clarity, we use $y_{x'}$ to denote the realization of the counterfactual outcome $Y_{x'}$ for a specific individual with observed evidence (X = x, Z = z, Y = y).

3. Analysis of Existing Methods

In this section, we elucidate the challenges of counterfactual inference. Subsequently, we summarize the existing methods and shed light on their limitations.

3.1. Challenges in Counterfactual Inference

The main challenge lies in that the counterfactual value $y_{x'}$ is generally not identifiable, even in randomized controlled experiments (RCTs). By definition, $y_{x'}$ is a quantity involving two "different worlds" at the same time: the observed world with (X = x, Z = z, Y = y) and the hypothetical world where X = x'. We only observe the factual outcome $Y_x = y$ but never observe the counterfactual outcome $Y_{x'}$, which is the fundamental problem in causal inference (Holland, 1986; Morgan & Winship, 2015). This inherent conflict prevents us from simplifying the expression of $y_{x'}$ to a do-calculus expression, making it generally unidentifiable, even in RCTs (Pearl et al., 2016). Therefore, in addition to the widely used assumptions such as conditional exchangeability, overlapping, and consistency (Hernán & Robins, 2020), counterfactual inference requires extra assumptions to ensure identifiability. Essentially, estimating $y_{x'}$ is equivalent to estimating the individual treatment effect $y_{x'} - y_x$, while the conditional average treatment effect (CATE) $\mathbb{E}[Y_{x'} - Y_x | Z = z]$ represents the ATE for a subpopulation with Z = z, overlooking the inherent heterogeneity in this subpopulation caused by the noise terms such as U_X (Albert et al., 2005; Heckman et al., 1997; Djebbari & Smith, 2008; Ding et al., 2019; Lei & Candès, 2021; Ben-Michael et al., 2022; Jin et al., 2023; Wu et al., 2024).

3.2. Summary of Existing Methods

We summarize the existing methods in terms of identifiability assumptions and estimation strategies.

We first present an equivalent expression of Eq. (1) using $(Y_x, Y_{x'})$. Eq. (1) be reformulated as the following system

$$Y_x = f_Y(x, Z, U_x), \ Y_{x'} = f_Y(x', Z, U_{x'})$$

where U_x and $U_{x'}$ denote the values of U_X given X = x and X = x', respectively. The exogenous variable U_X denotes the background and environment information induced by many unmeasured factors (Pearl et al., 2016), and thus U_x and $U_{x'}$ account for the heterogeneity of Y_x and $Y_{x'}$ in the observed and hypothetical worlds, respectively. These two worlds may exhibit different levels of noise due to unmeasured factors (Heckman et al., 1997; Chernozhukov & Hansen, 2005; Ding et al., 2019).

For identification, previous work (Lu et al., 2020; Xie et al., 2023; Nasr-Esfahany et al., 2023) relies on the key homogeneity and strict monotonicity assumptions.

Assumption 3.1 (Homogeneity). $U_x = U_{x'}$.

Assumption 3.2 (Strict Monotonicity). For any given (x, z), $Y_x = f_Y(x, z, U_x)$ is a smooth and strictly monotonic function of U_x ; or it is a bijective mapping from U_x to Y_x .

Assumption 3.1 implies that the value of U_X for each individual remains unchanged across x. Assumption 3.2 implies that Y_x is a strict monotonic function of U_x in the subpopulation of (X = x, Z = z). In Assumption 3.2, the smoothness and strict monotonicity of $f_Y(x, z, U_x)$ are akin to a bijective mapping of Y_x and U_x and serve the same purpose, so we don't distinguish them in detail.

Lemma 3.3. Under Assumptions 3.1-3.2, $y_{x'}$ is identifiable.

For estimation of $y_{x'}$, following Pearl's three-step procedure, Lu et al. (2020) and Nasr-Esfahany et al. (2023) initially estimate f_Y and U_X for each individual. However, estimating f_Y and U_X needs to impose extra assumptions, such as linearity (Shimizu et al., 2006) and additive noise (Peters et al., 2014). In addition, Xie et al. (2023) demonstrate that $y_{x'}$ corresponds to the τ^* -th quantile of the distribution $\mathbb{P}(Y|X = x', Z = z)$, where τ^* is the quantile of y in $\mathbb{P}(Y|X = x, Z = z)$ (See the proof of Lemma 3.3 or Section 4.1 for more details). Based on it, the authors uses quantile regression to estimate $y_{x'}$, which avoids the problem of estimating f_Y and U_X . Nevertheless, this method fits a single model to obtain the conditional quantile functions for both the counterfactual and factual outcomes. Thus, its validity relies on the underlying assumption that the conditional quantile functions of outcomes for different treatment groups stem from the same model. In addition, it involves estimating a distinct quantile value for each individual before deriving the counterfactual outcomes, posing a challenging bi-level optimization problem.

4. Identification through Rank Perservation

In this section, we introduce the rank preservation assumption for identifying $y_{x'}$. From a high-level perspective, identifying $y_{x'}$ essentially involves establishing the relationship between Y_x and $Y_{x'}$ for each individual. Pearl's three-step procedure achieves this by estimating f_Y and U_X , as well as the homogeneity assumption.

4.1. Rank Perservation Assumption

Our identifiability assumption is based on Kendall's rank correlation coefficient defined below.

Definition 4.1 (Kendall, 1938). Let $(x_1, y_1), ..., (x_n, y_n)$ be a set of observations of two random variables (X, Y), such that all the values of x_i and y_i are unique (ties are

neglected for simplicity). Any pair of (x_i, y_i) and (x_j, y_j) , if $(x_j - x_i)(y_j - y_i) > 0$, they are said to be concordant; otherwise they are discordant. The *sample* Kendall rank correlation coefficient is defined as

$$\rho_n(X,Y) = \frac{2}{n(n-1)} \sum_{1 \le i < j \le n} \operatorname{sign}((x_i - x_j)(y_i - y_j)),$$

where sign(t) = -1, 0, 1 for t < 0, t = 0, t > 0, respectively. For any two random variables (X, Y), we define $\rho(X, Y) = 1$, if $\rho_n(X, Y) = 1$ for all integers $n \ge 2$.

The $\rho_n(X, Y)$ also can be written as $2(N_c - N_d)/n(n-1)$, where N_c is the number of concordant pairs, N_d is the number of discordant pairs. It is easy to see that $-1 \leq \rho_n(X, Y) \leq 1$ and if the agreement between the two rankings is perfect (i.e., perfect concordance), $\rho_n(X, Y) = 1$.

Assumption 4.2 (Rank Preservation). $\rho(Y_x, Y_{x'}|Z) = 1$.

For the individual with observation (X = x, Z = z, Y = y), we denote $(y_x = y, y_{x'})$ as its true values of $(Y_x, Y_{x'})$. Assumption 4.2 implies that for this individual, its rankings of y_x and $y_{x'}$ are the same in the distributions of $\mathbb{P}(Y_x|Z = z)$ and $\mathbb{P}(Y_{x'}|Z = z)$, respectively. Therefore, we have

$$\mathbb{P}(Y_x \le y_x | Z = z) = \mathbb{P}(Y_{x'} \le y_{x'} | Z = z).$$
(2)

Since $y_x = y$ is observed and the distributions $\mathbb{P}(Y_x|Z=z)$ and $\mathbb{P}(Y_{x'}|Z=z)$ can be identified as $\mathbb{P}(Y|X=x,Z=z)$ and $\mathbb{P}(Y|X=x',Z=z)$, respectively, by the backdoor criterion (i.e., $(Y_x,Y_{x'}) \perp X|Z$). Therefore, we have the following Proposition 4.3 (see Appendix B for proofs).

Proposition 4.3. Under Assumption 4.2, $y_{x'}$ is identified as the τ^* -th quantile of $\mathbb{P}(Y|X = x', Z = z)$, where τ^* is the quantile of y in the distribution of $\mathbb{P}(Y|X = x, Z = z)$.

Proposition 4.3 shows that Assumption 4.2 can serve as a substitute for Assumptions 3.1-3.2 in identifying $y_{x'}$. Unlike Assumptions 3.1-3.2, Assumption 4.2 is simple and intuitive, as it directly links Y_x and $Y_{x'}$ for each individual. To clarify the relationship between Assumption 4.2 introduced by this work and Assumptions 3.1-3.2 from previous work, we present Proposition 4.4 below.

Proposition 4.4. *The proposed Assumption 4.2 is strictly weaker than Assumptions 3.1-3.2.*

Proposition 4.4 is intuitive, as correlation (Assumption 4.2) does not necessarily imply identity (Assumption 3.1). To illustrate, consider a SCM with $X \in \{0, 1\}, Y_1 = Z + U_1,$ $Y_0 = Z/2 + U_0, U_1 = U_0^3$. In this case, $\rho(Y_0, Y_1|Z) = 1$, but $U_1 \neq U_0$. Nevertheless, Assumption 4.2 is only slightly weaker than Assumptions 3.1-3.2 by allowing $U_{x'} \neq U_x$. Specifically, we can show that if U_x is a strictly monotone increasing function of $U_{x'}$, Assumption 4.2 is equivalent to Assumption 3.2, see Appendix B for proofs.

4.2. Further Relaxation of Strict Monotonicity

In Definition 4.1, we ignore ties for simplicity. However, when the outcome Y is discrete or continuous variables with tied observations, $\rho(Y_x, Y_{x'})$ will always be less than 1. To accommodate such cases, we introduce a modified version of the Kendall rank correlation coefficient given below.

Definition 4.5 (Kendall, 1945). Let $(x_1, y_1), ..., (x_n, y_n)$ be the observations of two random variables (X, Y), the modified Kendall rank correlation coefficient is define as

$$\tilde{\rho}_n(X,Y) = \sum_{1 \le i < j \le n} \frac{\operatorname{sign}((x_i - x_j)(y_i - y_j))}{\sqrt{n(n-1)/2 - T_x} \cdot \sqrt{n(n-1)/2 - T_y}},$$

where T_x is the number of tied pairs in $\{x_1, ..., x_n\}$ and T_y is the number of tied pairs in $\{y_1, ..., y_n\}$. We define $\tilde{\rho}(X, Y) = 1$, if $\tilde{\rho}_n(X, Y) = 1$ for all integers $n \ge 2$.

Compared with Definition 4.1, one can see that $\tilde{\rho}(X, Y)$ adjusts $\rho(X, Y)$ by eliminating the ties in the denominator, and $\tilde{\rho}(X, Y)$ reduces to $\rho(X, Y)$ if there are no ties.

Assumption 4.6 (Rank Preservation). $\tilde{\rho}(Y_x, Y_{x'}|Z) = 1$.

Assumption 4.6 is less restrictive than Assumption 4.2 as it accommodates broader data types of Y. To illustrate, consider a dataset with four individuals where the true values of $(Y_x, Y_{x'})$ are (1, 1), (2, 1.5), (2, 1.5), (3, 2.5). In this scenario, $\sum_{1 \le i < j \le n} \operatorname{sign}((y_{i,x} - y_{j,x})(y_{i,x'} - y_{j,x'}) = 5, T_{Y_x} = 1, T_{Y_{x'}} = 1$, resulting in $\rho(Y_x, Y_{x'}) = 5/6$ and $\tilde{\rho}(Y_x, Y_{x'}) = 5/(\sqrt{6-1} \cdot \sqrt{6-1}) = 1$.

Assumption 4.6 also guarantees the identifiability of $y_{x'}$.

Proposition 4.7. Under Assumption 4.6, the conclusion in *Proposition 4.3 also holds.*

5. Counterfactual Learning

We propose a novel estimation method for counterfactual inference. Suppose that $\{(x_k, z_k, y_k) : k = 1, ..., N\}$ is a sample consisting of N realizations of random variables (X, Z, Y). For an individual, given its evidence (X = x, Z = z, Y = y), we aim to estimate its counterfactual outcome $y_{x'}$, i.e., the realization of $Y_{x'}$ for this individual.

5.1. Rationale and Limitations of Quantile Regression

For estimating $y_{x'}$, Xie et al. (2023) formulate it as the following bi-level optimization problem

$$\tau^* = \arg\min_{\tau} |f_{\tau}(x, z) - y|,$$

$$f_{\tau}^* = \arg\min_{f} \frac{1}{N} \sum_{k=1}^{N} l_{\tau}(y_k - f(x_k, z_k)),$$

where $l_{\tau}(\xi) = \tau \xi \cdot \mathbb{I}(\xi \ge 0) + (\tau - 1)\xi \cdot \mathbb{I}(\xi < 0)$ is the check function (Koenker & Bassett, 1978), the upper

level optimization is to estimate τ^* , the quantile of y in the distribution $\mathbb{P}(Y|X = x, Z = z)$, and the lower level optimization is to estimate the conditional quantile function $q(x, z; \tau) \triangleq \inf_y \{y : \mathbb{P}(Y \le y | X = x, Z = z) \ge \tau\}$ for a given τ . Then $y_{x'}$ can be estimated using $q(x', z; \tau^*)$.

We define two conditional quantile regression functions,

$$q_x(z;\tau) \triangleq \inf_y \{ y : \mathbb{P}(Y_x \le y | Z = z) \ge \tau \},\$$
$$q_{x'}(z;\tau) \triangleq \inf_y \{ y : \mathbb{P}(Y_{x'} \le y | Z = z) \ge \tau \}.$$

By Eq. (2), $y_{x'}$ can be expressed as $q_{x'}(z; \tau^*)$ with τ^* being the quantile of y in the distribution of $\mathbb{P}(Y_x|Z=z)$, i.e., $\mathbb{P}(Y_x \leq y|Z=z) = \tau^*$. Lemma 5.1 (see Appendix C for proofs) shows the rationale behind employing the check function as the loss to estimate conditional quantiles.

Lemma 5.1. We have that

(i) $q_x(Z;\tau) = \arg\min_f \mathbb{E}[l_\tau(Y_x - f(Z))]$ for any given x; (ii) $q(X,Z;\tau) = \arg\min_f \mathbb{E}[l_\tau(Y - f(X,Z))].$

There are two major concerns with the estimation method of Xie et al. (2023). First, it only fits a single quantile regression model for $q(X, Z; \tau)$ to obtain estimates of $q_x(Z; \tau)$ and $q_{x'}(Z; \tau)$. When the two conditional quantile functions $q_x(Z; \tau)$ and $q_{x'}(Z; \tau)$ and $q_{x'}(Z; \tau)$ originate from different models, this method may yield inaccurate estimates. Second, it explicitly requires estimating the quantile τ^* for each individual before estimating the counterfactual outcome $y_{x'}$.

Inspired by Firpo (2007), a simple improvement is to estimate $q_x(z; \tau)$ and $q_{x'}(z; \tau)$ separately. For example, for estimating $q_x(z; \tau)$, the associated loss function is given as

$$R_x(f,\tau) = \frac{1}{N} \sum_{k=1}^{N} \frac{\mathbb{I}(x_k = x) \cdot l_\tau(y_k - f(z_k))}{\hat{p}_x(z_k)}$$

where $p_x(z) = \mathbb{P}(X = x | Z = z)$ is the propensity score, $\hat{p}_x(z)$ is its estimate. Likewise, we could define $R_{x'}(f, \tau)$ by replacing x with x'. Then the estimation procedure for $y_{x'}$ involves four steps: (1) estimating $p_x(z)$; (2) estimating $q_x(z;\tau)$ by minimizing $R_x(f,\tau)$ for a range of candidate values of τ ; (3) identifying the τ^* in the candidate set of τ , that corresponds to the quantile of y in the distribution $\mathbb{P}(Y|X = x, Z = z)$; (4) estimating $y_{x'}$ using $q_{x'}(z;\tau^*)$, where $q_{x'}(z;\tau^*)$ is obtained by minimizing $R_{x'}(f,\tau^*)$. Despite this four-step estimation method that allows $q_x(Z;\tau)$ and $q_{x'}(Z;\tau)$ to come from different models, it still needs to estimate a different τ^* for each individual.

5.2. Enhanced Counterfactual Learning Method

To address the limitations mentioned above in directly applying quantile regression and improve estimation accuracy, we propose a novel loss that produces an unbiased estimator of $y_{x'}$ for the individual with evidence (X = x, Z =

z, Y = y). The proposed ideal loss is constructed as

$$\begin{split} R_{x'}(t|x,z,y) &= \mathbb{E}\left[|Y_{x'} - t| \mid Z = z\right] \\ &+ \mathbb{E}\left[\operatorname{sign}(Y_x - y) \mid Z = z\right] \cdot t, \end{split}$$

which is a function of t and the expectation operator is taken on the random variable of $(Y_x, Y_{x'})$ given Z = z. The proposed estimation method is based on Theorem 5.2.

Theorem 5.2 (Validity of the Proposed Ideal Loss). The loss $R_{x'}(t|x, z, y)$ is convex with respect to t and is minimized uniquely at t^* , where t^* is the solution satisfying

$$\mathbb{P}(Y_{x'} \le t^* | Z = z) = \mathbb{P}(Y_x \le y | Z = z).$$

Theorem 5.2 (see Appendix C for proofs) implies that given the evidence (X = x, Z = z, Y = y) for an individual, the counterfactual outcome $y_{x'}$ satisfies $y_{x'} =$ $\arg \min_t R_{x'}(t|x, z, y)$ under Assumption 4.6. **Importantly**, the loss $R_{x'}(t|x, z, y)$ neither estimates the SCM a priori, nor restricts $q_x(z; \tau)$ and $q_{x'}(z; \tau)$ stem from the same model, and it does not need to estimate a different quantile value for each individual explicitly.

To optimize the ideal loss $R_{x'}(t; x, z, y)$, we first need to estimate it, which presents two significant challenges: (1) $R_{x'}(t|x, z, y)$ involves both Y_x and $Y_{x'}$, but for each unit, we only observe one of them; (2) The terms $\mathbb{E}\left[|Y_{x'} - t| \mid Z = z\right]$ and $\mathbb{E}\left[\operatorname{sign}(Y_x - y) \mid Z = z\right]$ in $R_{x'}(t|x, z, y)$ is conditioned on Z = z, and when Z is a continuous variable with infinite possible values, it cannot be estimated by simply splitting the data based on Z. We employ inverse propensity score and kernel smoothing techniques to overcome these two challenges. Specifically, we propose a kernel-smoothing-based estimator for the ideal loss, which is given as

$$\hat{R}_{x'}(t|x,z,y) = \frac{\sum_{k=1}^{N} K_h(z_k - z) \frac{\mathbb{I}(x_k = x')}{\hat{p}_{x'}(z_k)} |y_k - t|}{\sum_{k=1}^{N} K_h(z_k - z)} + \frac{\sum_{k=1}^{N} K_h(z_k - z) \frac{\mathbb{I}(x_k = x)}{\hat{p}_x(z_k)} \cdot \operatorname{sign}(y_k - y)}{\sum_{k=1}^{N} K_h(z_k - z)} \cdot t$$

where h is a bandwidth/smoothing parameter, $K_h(u) = K(u/h)/h$, and $K(\cdot)$ is a symmetric kernel function (Fan & Gijbels, 1996; Li & Racine, 2007) that satisfies $\int K(u)du = 1$ and $\int uK(u)du = 1$, such as Epanechnikov kernel $K(u) = 3(1 - u^2) \cdot \mathbb{I}(|u| \le 1)/4$ and Gaussian kernel $K(u) = \exp(-u^2/2)/\sqrt{2\pi}$ for $u \in \mathbb{R}$. Then we can estimate $y_{x'}$ by minimizing $\hat{R}_{x'}(t; x, z, y)$ directly.

Proposition 5.3 (Consistency). If $h \to 0$ as $N \to \infty$, $\hat{p}_x(z)$ and $\hat{p}_{x'}(z)$ are consistent estimates of $p_x(z)$ and $p_{x'}(z)$, and the density function of Z is differentiable, then $\hat{R}_{x'}(t|x, z, y)$ converges to $R_{x'}(t|x, z, y)$ in probability.

	Sim-5		Sim-10		Sim-20		Sim-40	
Methods	In-sample	Out-sample	In-sample	Out-sample	In-sample	Out-sample	In-sample	Out-sample
T-learner	2.95 ± 0.02	2.66 ± 0.01	2.99 ± 0.01	3.17 ± 0.01	3.36 ± 0.02	3.19 ± 0.03	5.12 ± 0.02	4.74 ± 0.04
X-learner	2.94 ± 0.01	2.66 ± 0.01	2.98 ± 0.02	3.19 ± 0.02	3.31 ± 0.02	3.21 ± 0.02	5.08 ± 0.04	4.77 ± 0.03
BNN	2.91 ± 0.08	2.64 ± 0.07	2.90 ± 0.11	3.08 ± 0.12	3.21 ± 0.13	3.13 ± 0.16	4.81 ± 0.10	4.54 ± 0.09
TARNet	2.89 ± 0.07	2.64 ± 0.06	2.94 ± 0.07	3.16 ± 0.08	3.18 ± 0.07	3.11 ± 0.07	4.82 ± 0.07	4.56 ± 0.07
CFRNet	2.88 ± 0.07	2.62 ± 0.06	2.94 ± 0.07	3.15 ± 0.08	3.15 ± 0.07	3.08 ± 0.07	4.71 ± 0.12	4.45 ± 0.11
CEVAE	2.92 ± 0.27	2.65 ± 0.21	3.04 ± 0.27	3.11 ± 0.18	3.16 ± 0.17	3.11 ± 0.17	4.88 ± 0.23	4.53 ± 0.20
DragonNet	2.90 ± 0.08	2.63 ± 0.08	3.02 ± 0.07	3.25 ± 0.08	3.16 ± 0.11	3.09 ± 0.10	4.78 ± 0.11	4.50 ± 0.12
DeRCFR	2.88 ± 0.06	2.61 ± 0.06	2.87 ± 0.05	3.07 ± 0.06	3.11 ± 0.07	3.04 ± 0.06	4.77 ± 0.11	4.50 ± 0.10
DESCN	2.93 ± 0.11	2.66 ± 0.09	3.27 ± 0.81	3.46 ± 0.79	3.12 ± 0.20	3.06 ± 0.20	4.91 ± 0.37	4.59 ± 0.35
ESCFR	2.87 ± 0.08	2.62 ± 0.07	2.94 ± 0.08	3.15 ± 0.09	3.03 ± 0.09	3.06 ± 0.09	4.71 ± 0.15	4.43 ± 0.15
CFQP	2.91 ± 0.09	2.67 ± 0.11	3.14 ± 0.30	3.40 ± 0.37	3.21 ± 0.12	3.18 ± 0.11	4.93 ± 0.14	4.55 ± 0.13
Quantile-Reg	2.80 ± 0.06	2.54 ± 0.05	2.78 ± 0.08	3.05 ± 0.09	2.92 ± 0.07	3.01 ± 0.08	4.39 ± 0.13	4.12 ± 0.10
Ours	$\textbf{2.45} \pm \textbf{0.17}$	$\textbf{2.28} \pm \textbf{0.23}$	$\begin{array}{ }\textbf{2.25}\pm\textbf{0.07}\end{array}$	$\textbf{2.33} \pm \textbf{0.07}$	$\textbf{2.51} \pm \textbf{0.07}$	$\textbf{2.46} \pm \textbf{0.06}$	$\textbf{3.74} \pm \textbf{0.26}$	$\textbf{3.66} \pm \textbf{0.21}$

Table 1: $\sqrt{\epsilon_{\text{PEHE}}}$ of individual treatment effect estimation on the simulated Sim-*m* dataset, where *m* is the dimension of *Z*.

Proposition 5.3 indicates that $\hat{R}_{x'}(t|x, z, y)$ is a consistent estimator of $R_{x'}(t|x, z, y)$, demonstrating the validity of the estimated ideal loss. The loss $\hat{R}_{x'}(t|x, z, y)$ is applicable only for discrete treatments due to the terms $\mathbb{I}(x_k = x')$ and $\mathbb{I}(x_k = x)$. However, it can be easily extended to continuous treatments, as detailed in Appendix D.

5.3. Further Theoretical Analysis

We further analyze the properties of the proposed method, including the unbiasedness preservation of the ideal loss, the bias of the estimated loss $\hat{R}_{x'}(t|x, z, y)$ and its impact on the final estimate of the counterfactual outcome $y_{x'}$.

First, we present the property of unbiasedness preservation. Let $R_{x'}^{\text{weight}}(t|x,z,y)$ be the weighted version of $R_{x'}(t|x,z,y)$, defined by

$$\mathbb{E}\left[w(X,Z)|Y_{x'}-t| \left| Z=z\right] + \mathbb{E}\left[w(X,Z)\operatorname{sign}(Y_x-y) \left| Z=z\right] \cdot t,\right]$$

where the weight w(X, Z) is an arbitrary function of (X, Z). The following Theorem 5.4 shows that $R_{x'}^{\text{weight}}(t|x, z, y)$ is also valid for estimating the counterfactual outcome $y_{x'}$.

Theorem 5.4 (Unbiasedness Preservation). The loss $R_{x'}^{\text{weight}}(t|x, z, y)$ is convex in t and is minimized uniquely at t^* that satisfies $\mathbb{P}(Y_{x'} \leq t^*|Z = z) = \mathbb{P}(Y_x \leq y|Z = z)$.

From Theorem 5.4 (see Appendix C for proofs), if we set $w(X, Z) = K_h(Z_k - z)/\{\sum_{k=1}^N K_h(Z_k - z)/N\}$, then the unbiasedness preservation property implicitly indicates that the proposed method is less sensitive to the choice of the kernel function.

Then, we show the bias of the estimated loss $\hat{R}_{x'}(t|x, z, y)$.

Proposition 5.5 (Bias of the Estimated Loss). If $h \to 0$ as $N \to \infty$, $p_x(z)/\hat{p}_x(z)$ and $p_{x'}(z)/\hat{p}_{x'}(z)$ are differentiable with respect to z, and the density function of Z is differentiable, then the bias of $\hat{R}_{x'}(t|x, z, y)$, defined by $\mathbb{E}[\hat{R}_{x'}(t|x, z, y)] - R_{x'}(t|x, z, y)$, is given as

$$\begin{aligned} &\textit{Bias}(\hat{R}_{x'}) = \delta_{p_{x'}} \mathbb{E}\left[|Y_{x'} - t| \mid Z = z\right] \\ &+ \delta_{p_x} \mathbb{E}\left[\textit{sign}(Y_x - y) \mid Z = z\right] \cdot t + O(h^2), \end{aligned}$$

where $\delta_{p_{x'}} = (p_{x'}(z) - \hat{p}_{x'}(z))/\hat{p}_{x'}(z)$ and $\delta_{p_x} = (p_x(z) - \hat{p}_x(z))/\hat{p}_x(z)$ are estimation errors of propensity scores.

From Proposition 5.5, the bias of $R_{x'}(t|x, z, y)$ consists of two components. The first is the estimation error of propensity scores. The second component arises from the kernel smoothing technique and is of order h^2 . In addition, when $\hat{p}_x(z)$ and $\hat{p}_{x'}(z)$ are consistent estimators of $p_x(z)$ and $p_{x'}(z)$ (a weak condition), the bias converges to zero, and Proposition 5.5 simplifies to Proposition 5.3.

Finally, we examine how the estimated $y_{x'}$, denoted as $\hat{y}_{x'} \triangleq \arg \min_t \hat{R}_{x'}(t|x, z, y)$, is influenced by the bias in the estimated loss.

Theorem 5.6 (Bias of the Estimated Counterfactual Outcome). Under the same conditions as in Proposition 5.5, $\hat{y}_{x'}$ converges to $\bar{y}_{x'}$ in probability, where $\bar{y}_{x'}$ satisfies that

$$\mathbb{P}(Y_{x'} \leq \bar{y}_x | Z = z) = \frac{(2 + 2\delta_{p_x})\mathbb{P}(Y_x \leq y | Z = z) + (\delta_{p_{x'}} - \delta_{p_x})}{2 + 2\delta_{p_{x'}}},$$

where $\bar{y}_{x'}$ may not equal to the true value $y_{x'}$ and their difference is the bias.

From Theorem 5.6, one can see that the bias of $\hat{y}_{x'}$ mainly depends on the estimation error of propensity scores. When $\delta_{p_x} = \delta_{p_{x'}}$, the equation in Theorem 5.6 reduces to equation (2), and thus $\bar{y}_{x'} = y_{x'}$, i.e., no bias. Typically, both δ_{p_x} and $\delta_{p_{x'}}$ are small due to the consistency of estimated propensity scores (a weak condition), and thus $\bar{y}_{x'}$ will close to $y_{x'}$.

	Sim-80 ($\rho = 0.3$)		Sim-80 ($\rho = 0.5$)		Sim-40 ($\rho = 0.3$)		Sim-40 ($\rho = 0.5$)	
Methods	In-sample	Out-sample	In-sample	Out-sample	In-sample	Out-sample	In-sample	Out-sample
TARNet	12.63 ± 0.93	12.51 ± 0.90	12.35 ± 1.24	12.68 ± 1.51	8.91 ± 0.97	8.78 ± 0.74	8.76 ± 0.76	8.51 ± 0.68
DragonNet	12.50 ± 0.75	12.36 ± 0.80	12.71 ± 1.29	13.02 ± 1.54	8.83 ± 0.90	8.73 ± 0.72	8.62 ± 0.70	8.39 ± 0.53
ESCFR	12.61 ± 1.09	12.53 ± 1.09	12.56 ± 1.36	12.87 ± 1.64	8.76 ± 1.03	8.65 ± 0.79	8.76 ± 0.78	8.50 ± 0.48
X_learner	12.82 ± 0.91	12.68 ± 0.95	12.74 ± 1.22	12.99 ± 1.43	8.97 ± 0.87	8.81 ± 0.64	8.91 ± 0.75	8.61 ± 0.58
Quantile-Reg	11.59 ± 0.94	11.57 ± 0.97	11.59 ± 1.26	11.91 ± 1.47	8.05 ± 0.73	8.08 ± 0.75	7.74 ± 0.73	7.58 ± 0.73
Ours	$\textbf{9.28} \pm \textbf{0.72}$	$\textbf{9.28} \pm \textbf{0.72}$	$\textbf{9.03} \pm \textbf{1.09}$	$\textbf{9.27} \pm \textbf{0.97}$	$\textbf{7.07} \pm \textbf{0.39}$	$\textbf{7.05} \pm \textbf{0.41}$	$\textbf{7.07} \pm \textbf{1.23}$	$\textbf{6.98} \pm \textbf{1.08}$

Table 2: $\sqrt{\epsilon_{\text{PEHE}}}$ of individual treatment effect estimation on the simulated Sim-*m* dataset, where *m* is the dimension of *Z*.



Figure 1: Estimation performance of individual treatment effects under varying heterogeneity degrees.

6. Experiments

6.1. Synthetic Experiment

Simulation Process. We generate the synthetic dataset by the following process. First, we sample the covariate $Z \sim \mathcal{N}(0, I_m)$ and the treatment $X \sim \text{Bern}(\pi(Z))$, where $\text{Bern}(\cdot)$ is the Bernoulli distribution with probability $\pi(Z) = \mathbb{P}(X = 1 \mid Z) = \sigma(W_x \cdot Z), \sigma(\cdot)$ is the sigmoid function, and $W_x \sim \text{Unif}(-1, 1)^m$, $\text{Unif}(\cdot)$ is the uniform distribution. Then, we sample the noise $U_0 \sim \mathcal{N}(0, 1)$ and $U_1 = \alpha \cdot U_0$ to consider the heterogeneity of the exogenous variables, where α is the hyper-parameter to control the heterogeneity degree. Finally, we simulate $Y_1 = W_y \cdot Z + U_1$ and $Y_0 = W_y \cdot Z/\alpha + U_0$ with $W_y \sim \mathcal{N}(0, I_m)$. We generate 10,000 samples with 63/27/10 train/validation/test split and vary $m \in \{5, 10, 20, 40\}$ in our synthetic experiment.

Baselines and Evaluation Metrics. The competing baselines includes: T-learner (Künzel et al., 2019), X-learner (Künzel et al., 2019), BNN (Johansson et al., 2016), TARNet (Shalit et al., 2017), CFRNet (Shalit et al., 2017), CEVAE (Louizos et al., 2017), DragonNet (Shi et al., 2019), DeRCFR (Wu et al., 2022), DESCN (Zhong et al., 2022), ESCFR (Wang et al., 2023), CFQP (Brouwer, 2022), and Quantile-Reg (Xie et al., 2023). We evaluate the individual treatment effect estimation using the *individual level Precision in Estimation of Heterogeneous Effects* (PEHE):

$$\epsilon_{\text{PEHE}} = \frac{1}{N} \sum_{i=1}^{N} [(\hat{Y}_i(1) - \hat{Y}_i(0)) - (Y_i(1) - Y_i(0))]^2,$$

where $\hat{Y}_i(1)$ and $\hat{Y}_i(0)$ are the predicted values for the corresponding true potential outcomes of unit *i*. It is

noteworthy that ϵ_{PEHE} is tailored for individual-level evaluation and counterfactual estimation, which is different from the common metric (Shalit et al., 2017) given by $\frac{1}{N} \sum_{i=1}^{N} [(\hat{\mu}_1(X_i) - \hat{\mu}_0(X_i)) - (\mu_1(X_i) - \mu_0(X_i))]^2$, where $\mu_1(X_i) - \mu_0(X_i) := \mathbb{E}[Y(1)|X] - \mathbb{E}[Y(0)|X]$ are the true CATE, and $\hat{\mu}_1(X_i) - \hat{\mu}_0(X_i)$ is its estimate. Both in-sample and out-of-sample performances are reported in our experiments. For more implementation details of the proposed method, please refer to Appendix E.

Performance Analysis. The results of estimation performance are shown in Table 1. Our method stably outperforms all baselines with varying covariate dimensions m, demonstrating the effectiveness of the proposed method. In addition, we investigate our method performance with violated assumptions on rank and uncorrelated covariates. Specifically, we modified the data generation process to explore the performance of our method under correlated covariates. Specifically, we sample the covariate $Z \sim \mathcal{N}(0, \Sigma_m)$, where the ρ_{ij} in Σ_m is max $(0.01, \rho^{|i-j|})$. The results are shown in Table 2. The results show that our method still outperforms the baseline methods. See Appendix E for the results with rank assumption violated. Moreover, we further explore the effect of heterogeneity degrees on the performance of the proposed method, as shown in Figure 1, from which one can see that as the heterogeneity degree increases, our method stably outperforms the Quantile-Reg in terms of PEHE. Finally, we examine the effect of different kernels and bandwidths, as shown in Figure 2, our method stably outperforms the Quantile-Reg and ESCFR methods with different kernels and bandwidths.

Learning Counterfactual Outcomes Under Rank Preservation



Figure 2: The estimation performance with different kernels and bandwidths.

Table 3: The experiment results on the IHDP dataset and JOBS dataset. The best result is bolded.

	IHDP				JOBS			
	In-sa	sample Out-sam		ample	In-sample		Out-sample	
Methods	$\sqrt{\epsilon_{\text{PEHE}}}$	ϵ_{ATE}	$\sqrt{\epsilon_{\text{PEHE}}}$	ϵ_{ATE}	R_{Pol}	ϵ_{ATT}	R_{Pol}	ϵ_{ATT}
T-learner	1.49 ± 0.03	0.37 ± 0.05	1.81 ± 0.04	0.49 ± 0.04	0.31 ± 0.06	0.16 ± 0.10	0.27 ± 0.08	0.20 ± 0.07
X-learner	1.50 ± 0.02	0.21 ± 0.05	1.73 ± 0.03	0.36 ± 0.07	0.16 ± 0.04	0.07 ± 0.05	0.16 ± 0.03	0.10 ± 0.09
BNN	2.09 ± 0.16	1.00 ± 0.23	2.37 ± 0.15	1.18 ± 0.19	0.15 ± 0.01	0.08 ± 0.03	0.16 ± 0.02	0.13 ± 0.07
TARNet	1.52 ± 0.07	0.22 ± 0.13	1.78 ± 0.07	0.34 ± 0.18	0.17 ± 0.06	0.06 ± 0.08	0.18 ± 0.09	0.10 ± 0.06
CFRNet	1.46 ± 0.06	0.17 ± 0.15	1.77 ± 0.06	0.32 ± 0.20	0.17 ± 0.03	$\textbf{0.05} \pm \textbf{0.03}$	0.19 ± 0.07	0.10 ± 0.04
CEVAE	4.08 ± 0.88	3.67 ± 1.23	4.12 ± 0.91	3.75 ± 1.23	0.18 ± 0.05	0.09 ± 0.03	0.22 ± 0.08	0.10 ± 0.09
DragonNet	1.49 ± 0.08	0.22 ± 0.14	1.80 ± 0.06	0.29 ± 0.19	0.17 ± 0.06	0.07 ± 0.07	0.20 ± 0.08	0.11 ± 0.09
DeRCFR	1.48 ± 0.06	0.25 ± 0.14	1.69 ± 0.06	0.25 ± 0.14	0.15 ± 0.02	0.14 ± 0.04	0.16 ± 0.04	0.15 ± 0.11
DESCN	2.08 ± 0.98	0.74 ± 1.00	2.67 ± 1.45	1.04 ± 1.46	0.15 ± 0.02	0.21 ± 0.14	0.22 ± 0.16	0.16 ± 0.04
ESCFR	1.46 ± 0.09	0.16 ± 0.16	1.73 ± 0.08	0.27 ± 0.16	0.14 ± 0.02	0.10 ± 0.03	0.15 ± 0.02	0.10 ± 0.08
Quantile-Reg	1.43 ± 0.05	0.14 ± 0.09	1.56 ± 0.03	0.18 ± 0.09	0.14 ± 0.01	0.06 ± 0.01	0.15 ± 0.01	0.07 ± 0.04
CFQP	1.47 ± 0.10	0.18 ± 0.17	1.48 ± 0.05	0.15 ± 0.08	0.15 ± 0.02	0.23 ± 0.15	0.16 ± 0.03	0.15 ± 0.07
Ours	$\textbf{1.41} \pm \textbf{0.02}$	$\textbf{0.11} \pm \textbf{0.10}$	$\textbf{1.50} \pm \textbf{0.06}$	$\textbf{0.13} \pm \textbf{0.08}$	$\textbf{0.08} \pm \textbf{0.04}$	0.06 ± 0.02	$\textbf{0.11} \pm \textbf{0.05}$	$\textbf{0.05} \pm \textbf{0.05}$

6.2. Real-World Experiment

Dataset and Preprocessing. Following previous studies Shalit et al. (2017), Louizos et al. (2017), Yoon et al. (2018), and Yao et al. (2018), we conduct experiments on semi-synthetic dataset IHDP and real-world dataset JOBS. The IHDP dataset (Hill, 2011) is constructed from the Infant Health and Development Program (IHDP) with 747 individuals and 25 covariates. The JOBS dataset (LaLonde, 1986) is based on the National Supported Work program with 3,212 individuals and 17 covariates. We follow Shalit et al. (2017) to split the data into training/validation/testing set with ratios 63/27/10 and 56/24/20 with 100 and 10 repeated times on the IHDP and the JOBS datasets, respectively.

Evaluation Metrics. Following previous studies (Shalit et al., 2017; Louizos et al., 2017; Yao et al., 2018), besides ϵ_{PEHE} , we also use the absolute error in *Average Treatment Effect* (ATE) for evaluation, which is defined as $\epsilon_{\text{ATE}} = \frac{1}{N} |\sum_{i=1}^{N} ((\hat{Y}_i(1) - \hat{Y}_i(0)) - (Y_i(1) - Y_i(0)))|$. We use $\sqrt{\epsilon_{\text{PEHE}}}$ and ϵ_{ATE} to evaluate performance on the IHDP dataset. For the JOBS dataset, since one of the potential outcomes is not available, we evaluate the performance using the absolute error in *Average Treatment effect on the*

Treated (ATT) as $\epsilon_{\text{ATT}} = |\text{ATT} - \frac{1}{|T|} \sum_{i \in T} (\hat{Y}_i(1) - \hat{Y}_i(0)|$ with ATT = $|\frac{1}{|T|} \sum_{i \in T} Y_i - \frac{1}{|C \cap E|} \sum_{i \in C \cap E} Y_i|$. We also use the policy risk $R_{\text{Pol}} = 1 - (\mathbb{E}[Y(1) \mid \hat{Y}(1) - \hat{Y}(0) > 0, X = 1] \cdot \mathbb{P}(\hat{Y}(1) - \hat{Y}(0) > 0) + \mathbb{E}[Y(0) \mid \hat{Y}(1) - \hat{Y}(0) \leq 0, X = 0] \cdot \mathbb{P}(\hat{Y}(1) - \hat{Y}(0) \leq 0))$, where T, C, E are the indexes of treatment sample set, control sample set, and randomized sample set, respectively.

Performance Comparison. The experiment results are shown in Table 3. Similar to the synthetic experiment, the Quantile-Reg method still achieves the most competitive performance compared to the other baselines. Our method stably outperforms all the baselines on both the semi-synthetic dataset IHDP and the real-world dataset JOBS, especially in the out-sample scenario. This provides the empirical evidence of the effectiveness of our method.

7. Conclusion

This work addresses the fundamental challenge of counterfactual inference in the absence of a known SCM and under heterogeneous endogenous variables. We first introduce the rank preservation assumption to identify counterfactual outcomes, showing that it is slightly weaker than the homogeneity and monotonicity assumptions. Then, we propose a novel ideal loss for unbiased learning of counterfactual outcomes and develop a kernel-based estimator for practical implementation. The convexity of the ideal loss and the unbiased nature of the proposed estimator contribute to the robustness and reliability of our method. A potential limitation arises when the propensity score is extremely small in certain data sparsity scenarios, which may cause instability in the estimation method. Further investigation is warranted to address and overcome this challenge.

Impact Statement

This paper proposed a novel counterfactual learning approach that enhances the previous work in terms of relaxed identifiability assumptions and theoretically guaranteed estimation methods. The proposed approach provides valuable tools and insights for researchers and practitioners across various disciplines, facilitating advancements in understanding causal relationships. It guides decision-making processes in complex, real-world scenarios, particularly in estimating individual treatment effects. Applications include credit assignment, root-causal analysis, algorithmic fair decision-making, policy evaluation and improvement, reinforcement learning, counterfactual explanation, counterfactual harm, classification and detection in medical imaging, among others.

References

- Albert, J. M., Gadbury, G. L., and Mascha, E. J. Assessing treatment effect heterogeneity in clinical trialswith blocked binary outcomes. *Biometrical Journal*, 47:662– 673, 2005.
- Anthis, J. R. and Veitch, V. Causal context connects counterfactual fairness to robust prediction and group fairness. *Advances in Neural Information Processing Systems*, 2023.
- Balke, A. and Pearl, J. Counterfactual probabilities: computational methods, bounds and applications. In *Proceedings of the Tenth international conference on Uncertainty in artificial intelligence*, 1994.
- Bareinboim, E., Correa, J. D., Ibeling, D., and Icard, T. On Pearl's Hierarchy and the Foundations of Causal Inference. ACM, 2022.
- Ben-Michael, E., Imai, K., and Jiang, Z. Policy learning with asymmetric utilities. arXiv:2206.10479, 2022.
- Brouwer, E. D. Deep counterfactual estimation with categorical background variables. Advances in Neural Information Processing Systems, 2022.

- Budhathoki, K., Minorics, L., Bloebaum, P., and Janzing, D. Causal structure-based root cause analysis of outliers. In *International Conference on Machine Learning*. PMLR, 2022.
- Chao, P., Blöbaum, P., and Kasiviswanathan, S. P. Interventional and counterfactual inference with diffusion models. *arXiv:2302.00860*, 2023.
- Chen, R., Yang, J., Xiong, H., Bai, J., Hu, T., Hao, J., FENG, Y., Zhou, J. T., Wu, J., and Liu, Z. Fast model debias with machine unlearning. *Advances in Neural Information Processing Systems*, 2023a.
- Chen, X.-H., Yu, Y., Zhu, Z., Yu, Z., Zhenjun, C., Wang, C., Wu, Y., Qin, R.-J., Wu, H., Ding, R., and Fangsheng, H. Adversarial counterfactual environment model learning. Advances in Neural Information Processing Systems, 2023b.
- Chernozhukov, V. and Hansen, C. An IV model of quantile treatment effects. *Econometrica*, 73:245–261, 2005.
- Chipman, H. A., George, E. I., and McCulloch, R. E. Bart: Bayesian additive regression trees. *The Annals of Applied Statistics*, 4(1):266–298, 2010.
- Correa, J. D., Lee, S., and Bareinboim, E. Nested counterfactual identification from arbitrary surrogate experiments. Advances in Neural Information Processing Systems, 2021.
- Ding, P., Feller, A., and Miratrix, L. Decomposing treatment effect variation. *Journal of the American Statistical Association*, 114:304–317, 2019.
- Djebbari, H. and Smith, J. A. Heterogeneous impacts in progresa. *Journal of Econometrics*, 145:64–80, 2008.
- Dudík, M., Langford, J., and Li, L. Doubly robust policy evaluation and learning. In *International Conference on Machine Learning*, pp. 1097–1104. PMLR, 2011.
- Fan, J. and Gijbels, I. *Local Polynomial Modelling and Its Applications*. Chapman and Hall/CRC, 1996.
- Feder, A., Wald, Y., Shi, C., Saria, S., and Blei, D. Data augmentations for improved (large) language model generalization. Advances in Neural Information Processing Systems, 2023.
- Finkelstein, N. and Shpitser, I. Deriving bounds and inequality constraints using logical relations among counterfactuals. In *Proceedings of the 36th Conference on Uncertainty in Artificial Intelligence*, 2020.
- Firpo, S. Efficient semiparametric estimation of quantile treatment effects. *Econometrica*, 75:259–276, 2007.

- Fontanella, A., Antoniou, A., Li, W., Wardlaw, J., Mair, G., Trucco, E., and Storkey, A. Acat: Adversarial counterfactual attention for classification and detection in medical imaging. *International Conference on Machine Learning*, pp. 10153–10169, 2023.
- Hamman, F., Noorani, E., Mishra, S., Magazzeni, D., and Dutta, S. Robust counterfactual explanations for neural networks with probabilistic guarantees. *International Conference on Machine Learning*, pp. 12351–12367, 2023.
- Haugh, M. B. and Singal, R. Counterfactual analysis in dynamic latent state models. *International Conference* on Machine Learning, pp. 12647–12677, 2023.
- Heckman, J. J., Smith, J., and Clements, N. Making the most out of programme evaluations and social experiments: Accounting for heterogeneity in programme impacts. *The Review of Economic Studies*, 64:487–535, 1997.
- Hernán, M. and Robins, J. M. *Causal Inference: What If.* Boca Raton: Chapman and Hall/CRC, 2020.
- Hill, J. L. Bayesian nonparametric modeling for causal inference. *Journal of Computational and Graphical Statistics*, 20(1):217–240, 2011.
- Holland, P. W. Statistics and causal inference. Journal of the American Statistical Association, 81:945–960, 1986.
- Hoyer, P., Janzing, D., Mooij, J. M., Peters, J., and Schölkopf, B. Nonlinear causal discovery with additive noise models. *Advances in Neural Information Process*ing Systems, 21, 2008.
- Ibeling, D. and Icard, T. Probabilistic reasoning across the causal hierarchy. In Proceedings of the Thirty-Fourth AAAI Conference on Artificial Intelligence, 2020.
- Imai, K. and Jiang, Z. Principal fairness for human and algorithmic decision-making. *Statistical Science*, 38(2): 317–328, 2023.
- Imbens, G. W. and Rubin, D. B. Causal Inference For Statistics Social and Biomedical Science. Cambridge University Press, 2015.
- Jin, Y., Ren, Z., and Candès, E. J. Sensitivity analysis of individual treatment effects: A robust conformal inference approach. *Proceedings of the National Academy of Sciences*, 120, 2023.
- Johansson, F., Shalit, U., and Sontag, D. Learning representations for counterfactual inference. In *International Conference on Machine Learning*, pp. 3020–3029. PMLR, 2016.

- Kallus, N. and Uehara, M. Double reinforcement learning for efficient off-policy evaluation in markov decision processes. *Journal of Machine Learning Research*, 21:1–63, 2020.
- Kavouras, L., Tsopelas, K., Giannopoulos, G., Sacharidis, D., Psaroudaki, E., Theologitis, N., Rontogiannis, D., Fotakis, D., and Emiris, I. Fairness aware counterfactuals for subgroups. *Advances in Neural Information Processing Systems*, 2023.
- Kendall, M. G. A new measure of rank correlation. *Biometrika*, 30:81–93, 1938.
- Kendall, M. G. The treatment of ties in ranking problem. *Biometrika*, 33:239–251, 1945.
- Kenny, E. M. and Huang, W. F. The utility of "even if" semifactual explanation to optimise positive outcomes. Advances in Neural Information Processing Systems, 2023.
- Koenker, R. and Bassett, G. Regression quantiles. *Econo*metrica, 46:33–50, 1978.
- Kosorok, M. R. and Laber, E. B. Precision medicine. *Annual Review of Statistics and Its Application*, 6:263–86, 2019.
- Künzel, S. R., Sekhon, J. S., Bickel, P. J., and Yu, B. Metalearners for estimating heterogeneous treatment effects using machine learning. *Proceedings of the national* academy of sciences, 116(10):4156–4165, 2019.
- Kusner, M. J., Loftus, J., Russell, C., and Silva, R. Counterfactual fairness. *Advances in Neural Information Processing Systems*, 30, 2017.
- LaLonde, R. J. Evaluating the econometric evaluations of training programs with experimental data. *The American economic review*, pp. 604–620, 1986.
- Langley, P. Crafting papers on machine learning. In *Interna*tional Conference on Machine Learning, pp. 1207–1216, Stanford, CA, 2000. PMLR.
- Lei, L. and Candès, E. J. Conformal inference of counterfactuals and individual treatment effects. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 83:911–938, 2021.
- Ley, D., Mishra, S., and Magazzeni, D. GLOBE-CE: A translation based approach for global counterfactual explanations. pp. 19315–19342. PMLR, 2023.
- Li, H., Zheng, C., Cao, Y., Geng, Z., Liu, Y., and Wu, P. Trustworthy policy learning under the counterfactual noharm criterion. In *International Conference on Machine Learning*, pp. 20575–20598. PMLR, 2023.

- Li, Q. and Racine, J. S. *Nonparametric econometrics*. Princeton University Press, 2007.
- Liu, Y., Chaudhari, P., and Fakoor, R. Budgeting counterfactual for offline rl. *Advances in Neural Information Processing Systems*, 2023a.
- Liu, Z., Anh, H., Chang, X., Chen, X., and Jia, R. 2d-shapley: A framework for fragmented data valuation. *International Conference on Machine Learning*, pp. 21730–21755, 2023b.
- Louizos, C., Shalit, U., Mooij, J. M., Sontag, D., Zemel, R., and Welling, M. Causal effect inference with deep latent-variable models. *Advances in Neural Information Processing Systems*, 30, 2017.
- Lu, C., Huang, B., Wang, K., Hernández-Lobato, J. M., Zhang, K., and Schölkopf, B. Sample-efficient reinforcement learning viacounterfactual-based data augmentation. In Offline Reinforcement Learning Workshop at Neural Information Processing Systems, 2020.
- Lv, C., Zhang, S., Tian, Y., Qi, M., and Ma, H. Disentangled counterfactual learning for physical audiovisual commonsense reasoning. *Advances in Neural Information Processing Systems*, 2023.
- Melnychuk, V., Frauen, D., and Feuerriegel, S. Partial counterfactual identification of continuous outcomes with a curvature sensitivity model. *Advances in Neural Information Processing Systems*, 2023.
- Mesnard, T., Weber, T., Viola, F., Thakoor, S., Saade, A., Harutyunyan, A., Dabney, W., Stepleton, T., Heess, N., Guez, A., Éric Moulines, Hutter, M., Buesing, L., and Munos, R. Counterfactual credit assignment in model-free reinforcement learning. In *Proceedings of the 38th International Conference on Machine Learning*, pp. 7654–7664. PMLR, 2021.
- Meulemans, A., Schug, S., Kobayashi, S., and Nathaniel Daw, G. W. Would i have gotten that reward? long-term credit assignment by counterfactual contribution analysis. *Advances in Neural Information Processing Systems*, 2023.
- Morgan, S. L. and Winship, C. *Counterfactuals and Causal Inference: Methods and Principles for Social Research.* Cambridge University Press, second edition, 2015.
- Nasr-Esfahany, A., Alizadehand, M., and Shah, D. Counterfactual identifiability of bijective causal models. In *International Conference on Machine Learning*. PMLR, 2023.
- Pearl, J. Causality. Cambridge university press, 2009.

- Pearl, J. and Mackenzie, D. *The Book of Why: The New* Science of Cause and Effect. Hachette Book Group, 2018.
- Pearl, J., Glymour, M., and Jewell, N. P. *Causal Inference in Statistics: A Primer*. John Wiley & Sons, 2016.
- Peters, J., Mooij, J. M., Janzing, D., and Schölkopf, B. Causal discovery with continuous additive noise models. 2014.
- Prabhu, V. U., Yenamandra, S., Chattopadhyay, P., and Hoffman, J. Lance: Stress-testing visual models by generating language-guided counterfactual images. *Advances in Neural Information Processing Systems*, 2023.
- Raman, C., Nonnemaker, A., Villegas-Morcillo, A., Hung, H., and Loog, M. Why did this model forecast this future? information-theoretic saliency for counterfactual explanations of probabilistic regression models. *Advances in Neural Information Processing Systems*, 2023.
- Ribeiro, F. D. S., Xia, T., Monteiro, M., Pawlowski, N., and Glocker, B. High fidelity image counterfactuals with probabilistic causal model. *International Conference on Machine Learning*, pp. 7390–7425, 2023.
- Richens, J. G., Beard, R., and Thompson, D. H. Counterfactual harm. *Advances in Neural Information Processing Systems*, 2022.
- Rosenbaum, P. R. and Rubin, D. B. The central role of the propensity score in observational studies for causal effects. *Biometrika*, 70(1):41–55, 1983.
- Saveski, M., Jecmen, S., Shah, N. B., and Ugander, J. Counterfactual evaluation of peer-review assignment policies. Advances in Neural Information Processing Systems, 2023.
- Schwab, P., Linhardt, L., and Karlen, W. Perfect match: A simple method for learning representations for counterfactual inference with neural networks. *arXiv preprint arXiv:1810.00656*, 2018.
- Shah, A., Dwivedi, R., Shah, D., and Wornell, G. W. On counterfactual inference with unobserved confounding. In *NeurIPS 2022 Workshop on Causality for Real-world Impact*, 2022.
- Shalit, U., Johansson, F. D., and Sontag, D. Estimating individual treatment effect: generalization bounds and algorithms. In *International Conference on Machine Learning*, pp. 3076–3085. PMLR, 2017.
- Shao, J., Qu, Y., Chen, C., Zhang, H., and Ji, X. Counterfactual conservative q learning for offline multi-agent reinforcement learning. *Advances in Neural Information Processing Systems*, 2023.

- Shi, C., Blei, D., and Veitch, V. Adapting neural networks for the estimation of treatment effects. *Advances in Neural Information Processing Systems*, 32, 2019.
- Shimizu, S., Hoyer, P. O., Hyvärinen, A., Kerminen, A., and Jordan, M. A linear non-gaussian acyclic model for causal discovery. *Journal of Machine Learning Research*, 7(10), 2006.
- Shpitser, I. and Pearl, J. What counterfactuals can be tested. In *Proceedings of the Twenty-Third Conference on Uncertainty in Artificial Intelligence*, 2007.
- Sun, Z., He, B., Liu, J., Chen, X., Ma, C., and Zhang, S. Offline imitation learning with variational counterfactual reasoning. *Advances in Neural Information Processing Systems*, 2023.
- Tang, S. and Wiens, J. Counterfactual-augmented importance sampling for semi-offline policy evaluation. Advances in Neural Information Processing Systems, 2023.
- Tian, J. and Pearl, J. Probabilities of causation: Bounds and identification. *Annals of Mathematics and Artificial Intelligence*, 28:287–313, 2000.
- Tsirtsis, S. and Rodriguez, M. G. Finding counterfactually optimal action sequences in continuous state spaces. *Ad*vances in Neural Information Processing Systems, 2023.
- van der Vaart, A. W. and Wellner, J. A. Weak convergence and empirical processes: with application to statistics. Springer, 1996.
- Wager, S. and Athey, S. Estimation and inference of heterogeneous treatment effects using random forests. *Journal* of the American Statistical Association, 113(523):1228– 1242, 2018.
- Wang, H., Fan, J., Chen, Z., Li, H., Liu, W., Liu, T., Dai, Q., Wang, Y., Dong, Z., and Tang, R. Optimal transport for treatment effect estimation. *Advances in Neural Information Processing Systems*, 2023.
- Wu, A., Yuan, J., Kuang, K., Li, B., Wu, R., Zhu, Q., Zhuang, Y., and Wu, F. Learning decomposed representations for treatment effect estimation. *IEEE Transactions* on Knowledge and Data Engineering, 35(5):4989–5001, 2022.
- Wu, P., Ding, P., Geng, Z., and Li, Y. Quantifying individual risk for binary outcome. arXiv:2402.10537, 2024.
- Wu, Z., D'Oosterlinck, K., Geiger, A., Zur, A., and Potts, C. Causal proxy models for concept-based model explanations. *International Conference on Machine Learning*, pp. 37313–37334, 2023.

- Xie, S., Huang, B., Gu, B., Liu, T., and Zhang, K. Advancing counterfactual inference through quantile regression. In *ICML Workshop on Counterfactuals in Minds and Machines*, 2023.
- Yan, H., Kong, L., Gui, L., Chi, Y., Xing, E., He, Y., and Zhang, K. Counterfactual generation with identifiability guarantee. *Advances in Neural Information Processing Systems*, 2023.
- Yan, J. and Wang, H. Self-interpretable time series prediction with counterfactual explanations. *International Conference on Machine Learning*, pp. 39110–39125, 2023.
- Yao, L., Li, S., Li, Y., Huai, M., Gao, J., and Zhang, A. Representation learning for treatment effect estimation from observational data. *Advances in Neural Information Processing Systems*, 31, 2018.
- Yoon, J., Jordon, J., and Van Der Schaar, M. Ganite: Estimation of individualized treatment effects using generative adversarial nets. In *International conference on learning representations*, 2018.
- Zenati, H., Diemert, E., Martin, M., Mairal, J., and Gaillard, P. Sequential counterfactual risk minimization. *International Conference on Machine Learning*, pp. 40681–40706, 2023.
- Zhang, J., Tian, J., and Bareinboim, E. Partial counterfactual identification from observational and experimental data. In *International Conference on Machine Learning*. PMLR, 2022.
- Zhong, K., Xiao, F., Ren, Y., Liang, Y., Yao, W., Yang, X., and Cen, L. Descn: Deep entire space cross networks for individual treatment effect estimation. In *Proceedings* of the 28th ACM SIGKDD Conference on Knowledge Discovery and Data Mining, pp. 4612–4620, 2022.
- Zuo, Z., Khalili, M. M., and Zhang, X. Counterfactually fair representation. *Advances in Neural Information Processing Systems*, 2023.

A. Related Work

Conditional Average Treatment Effect (CATE). CATE also referred to as heterogeneous treatment effect, represents the average treatment effects on subgroups categorized by covariate values, and plays a central role in areas such as precision medicine (Kosorok & Laber, 2019) and policy learning (Dudík et al., 2011). Benefiting from recent advances in machine learning, many methods have been proposed for estimating CATE, including matching methods (Rosenbaum & Rubin, 1983; Schwab et al., 2018; Yao et al., 2018), tree-based methods (Chipman et al., 2010; Wager & Athey, 2018), representation learning methods (Johansson et al., 2016; Shalit et al., 2017; Shi et al., 2019; Wu et al., 2022; Wang et al., 2023), and generative methods (Louizos et al., 2017; Yoon et al., 2018). Unlike the existing work devoted to estimating CATE at the intervention level for subgroups, our work focuses on counterfactual inference at the more challenging and fine-grained individual level.

Counterfactual Inference. Counterfactual inference involves the identification and estimation of counterfactual outcomes. For identification, Shpitser & Pearl (2007) provided an algorithm leveraging counterfactual graphs to identify counterfactual queries. In addition, Correa et al. (2021) discussed the identifiability of nested counterfactuals within a given causal graph. More relevant to our work, Lu et al. (2020) and Xie et al. (2023) studied the identifiability assumptions in the setting of backdoor criterion under homogeneity and strict monotonicity assumptions. Several methods focus on determining its bounds with less stringent assumptions, such as Balke & Pearl (1994), Tian & Pearl (2000), Pearl et al. (2016), Finkelstein & Shpitser (2020), Zhang et al. (2022), and Melnychuk et al. (2023).

For estimation, Pearl et al. (2016) introduced a three-step procedure for counterfactual inference. Many machine learning methods estimate counterfactual outcomes in this framework, such as Lu et al. (2020), Mesnard et al. (2021), Brouwer (2022), Shah et al. (2022), Yan et al. (2023), Nasr-Esfahany et al. (2023) and Chao et al. (2023). Recently, Xie et al. (2023) employed quantile regression to estimate the counterfactual outcomes, effectively circumventing the need for SCM estimation. In our work, we extend the above methods in both identification and estimation.

Recently, counterfactual inference methods have been extensively applied across various application scenarios, such as counterfactual fairness (Kusner et al., 2017; Zuo et al., 2023; Anthis & Veitch, 2023; Kavouras et al., 2023; Chen et al., 2023a), policy evaluation and improvement (Tang & Wiens, 2023; Saveski et al., 2023; Chen et al., 2023b), reinforcement learning (Lu et al., 2020; Tsirtsis & Rodriguez, 2023; Liu et al., 2023a; Shao et al., 2023; Meulemans et al., 2023; Haugh & Singal, 2023; Zenati et al., 2023), imitaion learning (Sun et al., 2023), counterfactual generation (Yan et al., 2023; Prabhu et al., 2023; Feder et al., 2023; Ribeiro et al., 2023), counterfactual explanation (Kenny & Huang, 2023; Raman et al., 2023; Hamman et al., 2023; Wu et al., 2023; Ley et al., 2023), counterfactual harm (Richens et al., 2022; Li et al., 2023), physical audiovisual commonsense reasoning (Lv et al., 2023), interpretable time series prediction (Yan & Wang, 2023), classification and detection in medical imaging (Fontanella et al., 2023), data valuation (Liu et al., 2023b), etc. Therefore, developing novel counterfactual inference methods holds significant practical implications.

B. Proofs in Sections 3 and 4

One can show Lemma 3.3 by a similar argument of the proof of Theorem 1 in Xie et al. (2023). For the sake of self-containedness, we provide a novel proof of it.

Lemma 3.3 Under Assumptions 3.1-3.2, $y_{x'}$ is identifiable.

Proof of Lemma 3.3. First, the distributions $\mathbb{P}(Y_x|Z=z)$ and $\mathbb{P}(Y_{x'}|Z=z)$ can be identified as $\mathbb{P}(Y|X=x,Z=z)$ and $\mathbb{P}(Y|X=x',Z=z)$, respectively, by the backdoor criterion (i.e., $(Y_x,Y_{x'}) \perp X|Z$) of the setting.

Then, according to the model (1), we can equivalently write

$$Y_x = f_Y(x, z, U_x), \ Y_{x'} = f_Y(x', z, U_{x'}),$$

and Y and U_X in model (1) can be expressed as $Y = \sum_{x \in \mathcal{X}} \mathbb{I}(X = x) \cdot Y_x$ and $U_X = \sum_{x \in \mathcal{X}} \mathbb{I}(X = x) \cdot U_x$, where \mathcal{X} is the support set of X and $\mathbb{I}(\cdot)$ is an indicator function. Assumption 3.1 implies that $U_X = U_x = U_{x'}$ conditional on Z, i.e., $Y_x = f_Y(x, z, U_X), \ Y_{x'} = f_Y(x', z, U_X).$

Finally, for the individual with observation (X = x, Z = z, Y = y), we denote $(y_x, y_{x'})$ as the true values of $(Y_x, Y_{x'})$ for this individual. For this individual, we can identify the quantile of y_x in the distribution of $\mathbb{P}(Y_x|Z = z) = \mathbb{P}(Y|X = x, Z = z)$, denoted by τ^* . Let u_{τ^*} be the true value of U_X for this individual, it is the τ^* -quantile in the distribution

 $\mathbb{P}(U_X|Z=z)$, then we have

$$\begin{aligned} \tau^* &= \mathbb{P}(Y_x \leq y_x | Z = z) & \text{(by the definition of } \tau) \\ &= \mathbb{P}(U_x \leq u_\tau | Z = z) & \text{(by Assumption 3.2)} \\ &= \mathbb{P}(U_{x'} \leq u_\tau | Z = z) & \text{(by Assumption 3.1)} \\ &= \mathbb{P}(Y_{x'} \leq f_Y(x', z, u_{\tau^*}) | Z = z) & \text{(by Assumption 3.2)} \\ &= \mathbb{P}(Y_{x'} \leq y_{x'} | Z = z) & \text{(by the definition of } y_{x'}), \end{aligned}$$

which implies that for this individual, its rankings of y_x and $y_{x'}$ are the same in the distributions of $\mathbb{P}(Y_x|Z=z)$ and $\mathbb{P}(Y_{x'}|Z=z)$, resepcctively. Thus, $y_{x'}$ is identified as the τ^* -quantile of the distribution $\mathbb{P}(Y_{x'}|Z=z) = \mathbb{P}(Y|X=x', Z=z)$.

Proposition 4.3 Under Assumption 4.2, $y_{x'}$ is identified as the τ^* -th quantile of $\mathbb{P}(Y|X = x', Z = z)$, where τ^* is the quantile of y in the distribution of $\mathbb{P}(Y|X = x, Z = z)$.

Proof of Proposition 4.3. For the individual with observation (X = x, Z = z, Y = y), we denote $(y_x, y_{x'})$ as the true values of $(Y_x, Y_{x'})$. Assumption 4.2 implies that for this individual, its rankings of y_x and $y_{x'}$ are the same in the distributions of $\mathbb{P}(Y_x|Z=z)$ and $\mathbb{P}(Y_{x'}|Z=z)$, respectively. Therefore,

$$\mathbb{P}(Y_x \le y_x | Z = z) = \mathbb{P}(Y_{x'} \le y_{x'} | Z = z).$$
(3)

Since $y_x = y$ is observed and the distributions $\mathbb{P}(Y_x|Z = z)$ and $\mathbb{P}(Y_{x'}|Z = z)$ can be identified as $\mathbb{P}(Y|X = x, Z = z)$ and $\mathbb{P}(Y|X = x', Z = z)$, respectively, by the backdoor criterion (i.e., $(Y_x, Y_{x'}) \perp X|Z$), we can identify the quantile of y_x in the distribution of $\mathbb{P}(Y|X = x, Z = z)$, denoted by τ^* . Then

$$\mathbb{P}(Y_{x'} \le y_{x'} | Z = z) = \tau^*,$$

which yields that θ is identified as the τ^* -quantile of $\mathbb{P}(Y|X = x', Z = z)$.

The following Proposition 4.4* serves as a complement to Proposition 4.4.

Proposition 4.4^{*} Under Assumption 3.1, or more generally, if U_x is a strictly monotone increasing function of $U_{x'}$, Assumption 4.2 is equivalent to Assumption 3.2.

Proof of Proposition 4.4. According to the model (1), we can equivalently write

$$Y_x = f_Y(x, z, U_x), \ Y_{x'} = f_Y(x', z, U_{x'})$$

Suppose that U_x is a strictly monotone increasing function of $U_{x'}$ (Assumption 3.1, i.e., $U_x = U_{x'}$, is a special case of it). Under this condition, we next prove sufficiency and necessity, respectively.

First, we show that Assumption 3.2 implies Assumption 4.2. If Assumption 3.2 holds, then Y_x is a strictly monotonic function of U_x , and $Y_{x'}$ is a strictly monotonic function of $U_{x'}$. Since U_x is a strictly monotone increasing function of $U_{x'}$, then Y_x is a strictly increasing monotonic function of $Y_{x'}$, which leads to Assumption 4.2.

Second, we show that Assumption 4.2 implies Assumption 3.2. If Assumption 4.2 holds, then given Z = z, Y_x is a strictly increasing function of $Y_{x'}$. When U_x is a strictly monotone increasing function of $U_{x'}$ and note that

$$Y_x = f_Y(x, z, U_X), \ Y_{x'} = f_Y(x', z, U_X),$$

which implies that f_Y is a strictly monotonic function of U_X , i.e., Assumption 3.2 holds.

This finishes the proof.

Proposition 4.7 Under Assumption 4.6, the conclusion in Proposition 4.3 also holds.

Proof of Proposition 4.7. This can be shown through a proof analogous to that of Proposition 4.3.

C. Proofs in Section 5

Recall that $l_{\tau}(\xi) = \tau \xi \cdot \mathbb{I}(\xi \ge 0) + (\tau - 1)\xi \cdot \mathbb{I}(\xi < 0)$, and

$$\begin{split} q(x,z;\tau) &\triangleq \inf_{y} \{ y : \mathbb{P}(Y \leq y | X = x, Z = z) \geq \tau \} \\ q_{0}(z;\tau) &\triangleq \inf_{y} \{ y : \mathbb{P}(Y_{0} \leq y | Z = z) \geq \tau \} \\ q_{1}(z;\tau) &\triangleq \inf_{y} \{ y : \mathbb{P}(Y_{1} \leq y | Z = z) \geq \tau \}. \end{split}$$

Lemma 5.1 We have that

(*i*) $q_x(Z; \tau) = \arg\min_f \mathbb{E}[l_\tau(Y_x - f(Z))]$ for x = 0, 1;(*ii*) $q(X, Z; \tau) = \arg\min_f \mathbb{E}[l_\tau(Y - f(X, Z))].$

Proof of Lemma 5.1. We prove $q_x(Z;\tau) = \arg\min_f \mathbb{E}[l_\tau(Y_x - f(Z))]$, and $q(X,Z;\tau) = \arg\min_f \mathbb{E}[l_\tau(Y - f(X,Z))]$ can be derived by an exactly similar manner. We write

$$\mathbb{E}[l_{\tau}(Y_x - f(Z))] = \mathbb{E}[\mathbb{E}\{l_{\tau}(Y_x - f(Z)) \mid Z\}]$$

To obtain the conclusion, note that $l_{\tau}(Y_x - f(Z))$ is always positive, it suffices to show that

$$q_x(z;\tau) = \arg\min_f \mathbb{E}[l_\tau(Y_x - f(Z)) \mid Z = z]$$
(4)

for any given Z = z. Next, we focus on analyzing the term $\mathbb{E}[l_{\tau}(Y_x - f(Z)) \mid Z = z]$. Given Z = z, f(Z) is a constant and we denote it by c, then

$$\begin{split} & \mathbb{E}[l_{\tau}(Y_{x} - f(Z)) \mid Z = z] \\ &= \mathbb{E}[l_{\tau}(Y_{x} - c) \mid Z = z] \\ &= \mathbb{E}\Big[\tau(Y_{x} - c)\mathbb{I}(Y_{x} \ge c) + (\tau - 1)(Y_{x} - c)\mathbb{I}(Y_{x} < c) \mid Z = z\Big] \\ &= \tau \int_{c}^{\infty} (y_{x} - c)g(y_{x}|z)dy_{x} + (\tau - 1)\int_{-\infty}^{c} (y_{x} - c)g(y_{x}|z)dy_{x}, \end{split}$$

where $g(y_x|z)$ denotes the probability density function of Y_x given Z = z.

Since the check function is a convex function, differentiating $\mathbb{E}[l_{\tau}(Y_x - c) \mid Z = z]$ with respect to c and setting the derivative to zero will yield the solution for the minimum

$$\begin{aligned} &\frac{\partial}{\partial c} \mathbb{E}[l_{\tau}(Y_x - c) \mid Z = z] \\ &= \tau \int_c^{\infty} \frac{\partial}{\partial c} [(y_x - c)g(y_x|z)] dy_x + (\tau - 1) \int_{-\infty}^c \frac{\partial}{\partial c} [(y_x - c)g(y_x|z)] dy_x \\ &= -\tau \Big(1 - \int_{-\infty}^c g(y_x|z) dy_x\Big) + (1 - \tau) \int_{-\infty}^c g(y_x|z) dy_x. \end{aligned}$$

Then let $\frac{\partial}{\partial c}\mathbb{E}[l_{\tau}(Y_x-c)\mid Z=z]=0$ leads to that

$$\int_{-\infty}^{c} g(y_x|z) dy_x = \tau,$$

that is, $c = q_x(z; \tau)$. This completes the proof of Proposition 5.1.

Theorem 5.2. If the probability density function of Y given Z is continuous, then the loss $R_{x'}(t; x, z, y)$ is minimized uniquely at t^* , where t^* is the solution satisfying

$$\mathbb{P}(Y_{x'} \le t^* | Z = z) = \mathbb{P}(Y_x \le y | Z = z).$$

Proof of Theorem 5.2. Recall that

$$R_{x'}(t|x,z,y) = \mathbb{E}\left[|Y_{x'}-t| \mid Z=z\right] + \mathbb{E}\left[\operatorname{sign}(Y_x-y) \mid Z=z\right] \cdot t.$$

Let $g(y_x|z)$ be the probability density function of Y_x given Z = z. By calculation,

$$\mathbb{E}\left[|Y_{x'} - t| \mid Z = z\right] = \int_{t}^{\infty} (y_{x'} - t)g(y_{x'}|z)dy_{x'} + \int_{-\infty}^{t} (t - y_{x'})g(y_{x'}|z)dy_{x'},$$

$$\frac{\partial}{\partial t} \mathbb{E}\left[|Y_{x'} - t| \mid Z = z\right] = -\left(1 - \int_{-\infty}^{t} g(y_{x'}|z)dy_{x'}\right) + \int_{-\infty}^{t} g(y_{x'}|z)dy_{x'} = 2\mathbb{P}(Y_{x'} \le t|Z = z) - 1,$$

and

$$\mathbb{E}\left[\operatorname{sign}(Y_x - y) \mid Z = z\right] = \mathbb{E}\left[-2\mathbb{I}(Y_x \le y) + 1 \mid Z = z\right] = -2\mathbb{P}(Y_x \le y | Z = z) + 1,$$

we have

$$\begin{split} \frac{\partial}{\partial t} R_{x'}(t|x,z,y) &= 2\mathbb{P}(Y_{x'} \leq t|Z=z) - 1 + \mathbb{E}\left[\operatorname{sign}(Y-y) \mid Z=z\right] \\ &= 2\mathbb{P}(Y_{x'} \leq t|Z=z) - 1 - 2\mathbb{P}(Y_x \leq y|Z=z) + 1 \\ &= 2\left\{\mathbb{P}(Y_{x'} \leq t|z) - \mathbb{P}(Y_x \leq y|z)\right\}. \end{split}$$

Since

$$\frac{\partial^2}{\partial t^2} R_{x'}(t|x,z,y) = 2\partial \mathbb{P}(Y_{x'} \le t|z) / \partial t = 2g(y_{x'} = t|z) \ge 0.$$

 $R_{x'}(t|x,z,y)$ is a convex function with respect to t. Letting $\frac{\partial}{\partial t}R_{x'}(t|x,z,y)=0$ yields that

$$\mathbb{P}(Y_{x'} \le t|z) - \mathbb{P}(Y_x \le y|z) = 0.$$

That is, $R_{x'}(t|x, z, y)$ attains its minimum at $t = q_{x'}(z; \tau^*)$, where τ^* is the quantile of y in the distribution $\mathbb{P}(Y_x|Z=z)$.

Proposition 5.3. If $h \to 0$ as $N \to \infty$, $\hat{p}_x(z)$ and $\hat{p}_{x'}(z)$ are consistent estimates of $p_x(z)$ and $p_{x'}(z)$, and the density function of Z is differentiable, then

$$\hat{R}_{x'}(t;x,z,y) \xrightarrow{\mathbb{P}} R_{x'}(t;x,z,y),$$

where $\xrightarrow{\mathbb{P}}$ means convergence in probability.

Proof of Proposition 5.3. For analyzing the theoretical properties of $\hat{R}_{x'}(t; x, z, y)$, we rewritten $\hat{R}_{x'}(t; x, z, y)$ as

$$\hat{R}_{x'}(t;x,z,y) = \frac{\sum_{k=1}^{N} K_h(Z_k - z) \frac{\mathbb{I}(X_k = x')}{\hat{p}_{x'}(Z_k)} |Y_k - t|}{\sum_{k=1}^{N} K_h(Z_k - z)} + \frac{\sum_{k=1}^{N} K_h(Z_k - z) \frac{\mathbb{I}(X_k = x)}{\hat{p}_{x'}(Z_k)} \cdot \operatorname{sign}(Y_k - y)}{\sum_{i=1}^{N} K_h(Z_k - z)} \cdot t,$$

where the capital letters denote random variables and lowercase letters denote their realizations. This is slightly different from that used in the main text.

When $\hat{p}_x(z)$ and $\hat{p}_{x'}(z)$ are consistent estimates of $p_x(z)$ and $p_{x'}(z)$, to show the conclusion, it is sufficient to prove that

$$\frac{\sum_{k=1}^{N} K_h(Z_k - z) \frac{\mathbb{I}(X_k = x')}{p_{x'}(Z_k)} |Y_k - t|}{\sum_{k=1}^{N} K_h(Z_k - z)} \xrightarrow{\mathbb{P}} \mathbb{E}\left[\frac{\mathbb{I}(X = x')}{p_{x'}(z)} |Y - t| \mid Z = z\right] = \mathbb{E}\left[|Y_{x'} - t| \mid Z = z\right],$$
(5)

$$\frac{\sum_{k=1}^{N} K_h(Z_k - z) \frac{\mathbb{I}(X_k = x)}{p_x(Z_k)} \cdot \operatorname{sign}(Y_k - y)}{\sum_{i=1}^{N} K_h(Z_k - z)} \xrightarrow{\mathbb{P}} \mathbb{E}\left[\frac{\mathbb{I}(X = x)}{p_x(z)} \cdot \operatorname{sign}(Y - y) \mid Z = z\right] = \mathbb{E}\left[\operatorname{sign}(Y_x - y) \mid Z = z\right].$$
(6)

We prove equation (5) only, as equation (6) can be addressed similarly.

Note that

$$\frac{\sum_{k=1}^{N} K_h(Z_k-z) \frac{\mathbb{I}(X_k=x')}{p_{x'}(Z_k)} |Y_k-t|}{\sum_{k=1}^{N} K_h(Z_k-z)} = \frac{\frac{1}{N} \sum_{k=1}^{N} K_h(Z_k-z) \frac{\mathbb{I}(X_k=x')}{p_{x'}(Z_k)} |Y_k-t|}{\frac{1}{N} \sum_{k=1}^{N} K_h(Z_k-z)},$$

we analyze the denominator and numerator on the right side of the equation separately. For the denominator, it is an average of N independent random variables and converges to its expectation $\mathbb{E}[K_h(Z_k - z)]$ almost surely. Let $g(z_k)$ be the probability density function of Z_k , and $g^{(1)}(z_k)$ is its first derivative. Since

$$\mathbb{E}[K_h(Z_k - z)] = \int \frac{1}{h} K(\frac{z_k - z}{h}) g(z_k) dz_k$$

$$= \int K(u) g(z + hu) du \quad (\text{let } z_k = z + hu)$$

$$= \int K(u) \cdot \{g(z) + g^{(1)}(z)hu + o(h)\} du \quad (\text{by Taylor Expansion})$$

$$= g(z) \int K(u) du + g^{(1)}(z)h \int K(u) u du + o(h)$$

$$= g(z) + o(h) \quad (\text{by the definition of kernel function}), \quad (7)$$

when $h \to 0$ as $N \to \infty$, the denominator converges to g(z) in probability.

Next, we focus on dealing with the numerator, which also converges to its expectation.

$$\mathbb{E}[K_{h}(Z_{k}-z)\frac{\mathbb{I}(X_{k}=x')}{p_{x'}(Z_{k})}|Y_{k}-t|]$$

$$=\mathbb{E}\Big[K_{h}(Z_{k}-z)\mathbb{E}\Big\{\frac{\mathbb{I}(X_{k}=x')}{p_{x'}(Z_{k})}|Y_{k}-t|\Big|Z_{k}\Big\}\Big] \quad \text{(by the law of iterated expectations)}$$

$$=\mathbb{E}\Big[K_{h}(Z_{k}-z)\mathbb{E}\Big\{\frac{\mathbb{I}(X_{k}=x')}{p_{x'}(Z_{k})}|Y_{x',k}-t|\Big|Z_{k}\Big\}\Big] \quad \text{(write } Y_{k} \text{ as the form of potential outcome)}$$

$$=\mathbb{E}\Big[K_{h}(Z_{k}-z)\mathbb{E}\Big\{|Y_{x',k}-t|\Big|Z_{k}\Big\}\Big] \quad \text{(by backdoor criterion } Y_{x',k} \perp X_{k}|Z_{k}\text{)}. \tag{8}$$

Define $m(Z) = \mathbb{E}[|Y_{x'} - t||Z]$ and $m^{(1)}(Z)$ is its first derivative, then the right side of equation (5) is m(z), and

$$\mathbb{E}\left[K_{h}(Z_{k}-z)\cdot\mathbb{E}\left\{\left|Y_{x',k}-t\right|\left|Z_{k}\right\}\right] = \mathbb{E}\left[K_{h}(Z_{k}-z)\cdot m(Z_{k})\right] \\
= \int \frac{1}{h}K(\frac{z_{k}-z}{h})\cdot m(z_{k})\cdot g(z_{k})dz_{k} \\
= \int K(u)\cdot m(z+hu)\cdot g(z+hu)du \quad (\text{let } z_{k}=z+hu) \\
= \int K(u)\cdot\{m(z)+m^{(1)}(z)hu+o(h)\}\cdot\{g(z)+g^{(1)}(z)hu+o(h)\}du \quad (\text{by Taylor Expansion}) \\
= m(z)g(z)+o(h).$$
(9)

Thus, when $h \to 0$ as $N \to \infty$, the numerator converges to g(z) in probability.

Combining equations (7), (8), and (9) yields the equality (5). This completes the proof.

Theorem 5.4 (Unbiasedness Preservation). The loss $R_{x'}^{\text{weight}}(t|x, z, y)$ is convex in terms of t and is minimized uniquely at t^* , where t^* is the solution satisfying $\mathbb{P}(Y_{x'} \leq t^*|Z=z) = \mathbb{P}(Y_x \leq y|Z=z)$.

Proof of Theorem 5.4. . It is sufficient to show that $R_{x'}^{\text{weight}}(t|x, z, y)$ shares the same minimizer as $R_{x'}(t|x, z, y)$. By the backdoor criterion (i.e., $(Y_x, Y_{x'} \perp X|Z)$), we have that

$$\begin{aligned} R_{x'}^{\text{weight}}(t|x,z,y) \\ &= \mathbb{E}\left[w(X,Z)|Y_{x'}-t| \left| Z=z \right] + \mathbb{E}\left[w(X,Z)\text{sign}(Y_x-y) \left| Z=z \right] \cdot t \right] \\ &= \mathbb{E}[w(X,Z) \mid Z=z] \cdot \mathbb{E}\left[|Y_{x'}-t| \mid Z=z \right] \\ &+ \mathbb{E}[w(X,Z) \mid Z=z] \cdot \mathbb{E}\left[w(X,Z) \cdot \text{sign}(Y_x-y) \mid Z=z \right] \cdot t \\ &= \mathbb{E}[w(X,Z) \mid Z=z] \cdot R_{x'}(t|x,z,y). \end{aligned}$$

Note that $\mathbb{E}[w(X,Z) \mid Z = z]$ is not related to t. Consequently, $R_{x'}^{\text{weight}}(t|x, z, y)$ shares the same unique minimizer as $R_{x'}(t|x, z, y)$. Both minimizers satisfy the condition $\mathbb{P}(Y_{x'} \leq t^* | Z = z) = \mathbb{P}(Y_x \leq y | Z = z)$.

-	-	_

Proposition 5.5 (Bias of the Estimated Loss). If $h \to 0$ as $N \to \infty$, $p_x(z)/\hat{p}_x(z)$ and $p_{x'}(z)/\hat{p}_{x'}(z)$ are differentiable with respect to z, and the density function of Z is differentiable, then the bias of $\hat{R}_{x'}(t|x, z, y)$, defined by $\mathbb{E}[\hat{R}_{x'}(t|x, z, y)] - R_{x'}(t|x, z, y)$, is given as

$$Bias(\hat{R}_{x'}) = \delta_{p_{x'}} \mathbb{E}\left[|Y_{x'} - t| \mid Z = z \right] + \delta_{p_x} \mathbb{E}\left[sign(Y_x - y) \mid Z = z \right] \cdot t + O(h^2).$$

where $\delta_{p_{x'}} = (p_{x'}(z) - \hat{p}_{x'}(z))/\hat{p}_{x'}(z)$ and $\delta_{p_x} = (p_x(z) - \hat{p}_x(z))/\hat{p}_x(z)$ are estimation errors of propensity scores.

Proof of Proposition 5.5. This proof is similar to the proof of Proposition 5.3. Recall that

$$\hat{R}_{x'}(t;x,z,y) = \frac{\sum_{k=1}^{N} K_h(Z_k - z) \frac{\mathbb{I}(X_k = x')}{\hat{p}_{x'}(Z_k)} |Y_k - t|}{\sum_{k=1}^{N} K_h(Z_k - z)} + \frac{\sum_{k=1}^{N} K_h(Z_k - z) \frac{\mathbb{I}(X_k = x)}{\hat{p}_{x}(Z_k)} \cdot \operatorname{sign}(Y_k - y)}{\sum_{i=1}^{N} K_h(Z_k - z)} \cdot t,$$

and

$$R_{x'}(t|x,z,y) = \mathbb{E}\left[|Y_{x'}-t| \mid Z=z\right] + \mathbb{E}\left[\operatorname{sign}(Y_x-y) \mid Z=z\right] \cdot t,$$

Given the estimated propensity scores, to show the conclusion, it suffices to prove that

$$\mathbb{E}\left[\frac{\sum_{k=1}^{N} K_h(Z_k - z) \frac{\mathbb{I}(X_k = x')}{\hat{p}_{x'}(Z_k)} |Y_k - t|}{\sum_{k=1}^{N} K_h(Z_k - z)}\right] = \frac{p_{x'}(z)}{\hat{p}_{x'}(z)} \mathbb{E}\left[|Y_{x'} - t| \mid Z = z\right] + O(h^2).$$
(10)

$$\mathbb{E}\left[\frac{\sum_{k=1}^{N} K_h(Z_k-z) \frac{\mathbb{I}(X_k=x)}{p_x(Z_k)} \cdot \operatorname{sign}(Y_k-y)}{\sum_{i=1}^{N} K_h(Z_k-z)}\right] = \frac{p_x(z)}{\hat{p}_x(z)} \mathbb{E}\left[\operatorname{sign}(Y_x-y) \mid Z=z\right] \cdot t + O(h^2).$$
(11)

We prove the equation (10) only, as remaining equation (11) can be addressed similarly. Let $g(z_k)$ be the probability density

function of Z_k and $m(Z) = \mathbb{E}[|Y_{x'} - t||Z]$. By the proof of Proposition 5.3, $N^{-1} \sum_{k=1}^{N} K_h(Z_k - z) = g(z) + O(h^2)$, and

$$\mathbb{E}\left[\frac{1}{N}\sum_{k=1}^{N}K_{h}(Z_{k}-z)\frac{\mathbb{I}(X_{k}=x')}{\hat{p}_{x'}(Z_{k})}|Y_{k}-t|\right]$$
$$\mathbb{E}[K_{h}(Z_{k}-z)\frac{\mathbb{I}(X_{k}=x')}{\hat{p}_{x'}(Z_{k})}|Y_{k}-t|]$$
$$=\mathbb{E}\left[K_{h}(Z_{k}-z)\cdot\frac{p_{x'}(Z_{k})}{\hat{p}_{x'}(Z_{k})}\mathbb{E}\left\{|Y_{x',k}-t|\Big|Z_{k}\right\}\right]$$
$$=\mathbb{E}\left[K_{h}(Z-z)\cdot\frac{p_{x'}(Z)}{\hat{p}_{x'}(Z)}\cdot m(Z)\right]$$
$$=\frac{p_{x'}(z)}{\hat{p}_{x'}(z)}m(z)g(z)+O(h^{2}).$$

Thus, equation (10) holds using Slutsky's Theorem. This completes the proof.

Theorem 5.6 (Bias of the Estimated Counterfactual Outcome). Under the same conditions as in Proposition 5.5, $\hat{y}_{x'}$ converges to $\bar{y}_{x'}$ in probability, where $\bar{y}_{x'}$ satisfies that

$$\mathbb{P}(Y_{x'} \le \bar{y}_x | Z = z) = \frac{(2 + 2\delta_{p_x})\mathbb{P}(Y_x \le y | Z = z) + (\delta_{p_{x'}} - \delta_{p_x})}{2 + 2\delta_{p_{x'}}},$$
(12)

where $\bar{y}_{x'}$ may not equal to the true value $y_{x'}$ and their difference is the bias.

Proof of Theorem 5.6. By the property of M-estimation (van der Vaart & Wellner, 1996), $\hat{y}_{x'}$ converges to $\bar{y}_{x'}$ in probability, where $\bar{y}_{x'}$ is the minimizer of

$$\bar{R}_{x'}(t;x,z,y),$$

where $\bar{R}_{x'}(t; x, z, y)$ is the probability limit of $\hat{R}_{x'}(t; x, z, y)$. By the proof of Proposition 5.5,

$$\bar{R}_{x'}(t;x,z,y) = \frac{p_{x'}(z)}{\hat{p}_{x'}(z)} \mathbb{E}\left[|Y_{x'} - t| \mid Z = z \right] + \frac{p_x(z)}{\hat{p}_x(z)} \mathbb{E}\left[\operatorname{sign}(Y_x - y) \mid Z = z \right] \cdot t.$$

By a similar proof of Theorem 5.2, as $h \to 0$, $\bar{y}_{x'} = \arg \min_t \bar{R}_{x'}(t; x, z, y)$ satisfies that

$$\mathbb{P}(Y_{x'} \le \bar{y}_x | Z = z) = \frac{(2 + 2\delta_{p_x})\mathbb{P}(Y_x \le y | Z = z) + (\delta_{p_{x'}} - \delta_{p_x})}{2 + 2\delta_{p_{x'}}},$$

D. Extension to Continuous Outcome

When the treatment is continuous, we can estimate the ideal loss with the following estimator

$$\tilde{R}_{x'}(t|x,z,y) = \frac{\sum_{k=1}^{N} K_h(z_k-z) \frac{K_h(x_k-x')}{p_{x'}(z_k)} |y_k-t|}{\sum_{k=1}^{N} K_h(z_k-z)} + \frac{\sum_{k=1}^{N} K_h(z_k-z) \frac{K_h(x_k-x)}{p_{x}(z_k)} \cdot \operatorname{sign}(y_k-y)}{\sum_{k=1}^{N} K_h(z_k-z)} \cdot t,$$

which is a smoothed version of the estimator

$$\hat{R}_{x'}(t|x,z,y) = \frac{\sum_{k=1}^{N} K_h(z_k - z) \frac{\mathbb{I}(x_k = x')}{p_{x'}(z_k)} |y_k - t|}{\sum_{k=1}^{N} K_h(z_k - z)} + \frac{\sum_{k=1}^{N} K_h(z_k - z) \frac{\mathbb{I}(x_k = x)}{p_x(z_k)} \cdot \operatorname{sign}(y_k - y)}{\sum_{k=1}^{N} K_h(z_k - z)} \cdot t,$$

defined in Section 5. In addition, by a proof similar to that of Proposition 5.3, we also can show that $\tilde{R}_{x'}(t;x,z,y) \xrightarrow{\mathbb{P}} R_{x'}(t;x,z,y)$.

Learning Counterfactual Outcomes Under Rank Preservation

	Sim-10 (Rank=0.3)		Sim-10 (Rank=0.5)		Sim-40 (Rank=0.3)		Sim-40 (Rank=0.5)	
Methods	In-sample	Out-sample	In-sample	Out-sample	In-sample	Out-sample	In-sample	Out-sample
TARNet	6.92 ± 0.79	6.96 ± 0.82	4.38 ± 0.59	4.38 ± 0.60	14.91 ± 0.61	14.80 ± 0.82	9.33 ± 0.36	9.33 ± 0.45
DragonNet	6.97 ± 0.83	7.02 ± 0.88	4.43 ± 0.62	4.44 ± 0.62	15.04 ± 0.54	14.94 ± 0.77	9.18 ± 0.30	9.16 ± 0.50
ESCFR	6.91 ± 0.69	6.96 ± 0.76	4.41 ± 0.65	4.41 ± 0.68	15.01 ± 0.58	14.86 ± 0.82	9.25 ± 0.18	9.26 ± 0.31
X_learner	7.01 ± 0.81	7.05 ± 0.85	5.29 ± 0.44	4.98 ± 0.48	15.16 ± 0.64	15.05 ± 0.88	9.43 ± 0.29	9.40 ± 0.42
Quantile-Reg	6.79 ± 0.79	6.63 ± 0.84	4.12 ± 0.63	4.14 ± 0.65	13.12 ± 0.62	13.25 ± 0.87	8.39 ± 0.30	8.43 ± 0.42
Ours	6.00 ± 1.22*	$\textbf{6.01} \pm \textbf{1.29}$	$\textbf{3.88} \pm \textbf{1.25}$	$\textbf{3.90} \pm \textbf{1.30}$	$\textbf{10.76} \pm \textbf{0.52*}$	$\textbf{10.84} \pm \textbf{0.43} \ast$	6.75 ± 0.24*	$\textbf{6.80} \pm \textbf{0.21} \ast$

Table 4: $\sqrt{\epsilon_{\text{PEHE}}}$ of individual treatment effect estimation on the simulated Sim-*m* dataset, where *m* is the dimension of *Z*.

E. Experiment Details and Additional Experiment Results

We run all experiments on the Google Colab platform. For the representation model, we use the MLP for the base model and tune the layers in $\{1, 2, 3\}$. In addition, we adopt the logistic regression model as the propensity model. We tune the learning rate in $\{0.001, 0.005, 0.01, 0.05, 0.1\}$. For the kernel choice, we select the kernel function between the Gaussian kernel function and the Epanechnikov kernel function, and tune the bandwidth in $\{1, 3, 5, 7, 9\}$.

In addition, we investigate the performance of our method when the rank preservation assumption is violated. Specifically, we simulate $Y_1 = (W_y + W_{y1}) \cdot Z + U_1$ and $Y_0 = W_y \cdot Z/\alpha + U_0$ with $W_y \sim \mathcal{N}(0, I_m)$ and $W_{y1} \sim \mathcal{N}(0, \beta I_m)$, where β is the hyper parameter to control the violation degree. In the added experiments, m is in 10, 40, which is consistent with the experiments in our original manuscript, and Kendall's rank correlation coefficient $\rho(Y_0, Y_1)$ is in 0.3, 0.5. We generate 10 different datasets with different seeds and the experiment results are shown in Table 4. The experiment results show that our method still outperforms the baseline methods, even if the rank preservation assumption is violated.