# Existence and Uniqueness of Local Solutions for a Class of Partial Differential-Algebraic Equations

Seyyid Ali Benabdallah and Messaoud Souilah

Dynamic Systems Laboratory (DSL), Department of Mathematics University of Science and Technology Houari Boumediene Algiers, Algeria sbenabdallah@usthb.dz

Dynamic Systems Laboratory (DSL), Department of Mathematics University of Science and Technology Houari Boumediene Algiers, Algeria msouilah@usthb.dz

February 11, 2025

#### Abstract

In this work, we present a result on the local existence and uniqueness of solutions to nonlinear Partial Differential-Algebraic Equations (PDAEs). By applying established theoretical results, we identify the conditions that guarantee the existence of a unique local solution. The analysis relies on techniques from functional analysis, semi-group theory, and the theory of differential-algebraic systems. Additionally, we provide applications to illustrate the effectiveness of this result.

MSC 2020: 47B01, 47F05, 47H20, 47N60.

## 1 Introduction

In this paper, we study a specific type of partial differential-algebraic equations (PDAEs) defined as follows:

$$\begin{cases} V_t(t,x) = F(x,t,U(t,x)), \\ 0 = G(x,t,U(t,x)), \\ V(0,x) = V_0(x), \end{cases}$$
(1)

where U = (V, w) is an unknown function depending on the spatial variable  $x \in \Omega$ and time t, with  $\Omega \subset \mathbb{R}^n$ .

This type of equation models numerous physical and chemical phenomena. For example, the Navier-Stokes equations in fluid mechanics [15] and the transport of ions in

electrolytes [14] fall within this framework. For a broader overview of partial differentialalgebraic equations (PDAEs), we refer the reader to [2], [9], [11], [16], and [17].

In this work, we focus specifically on a class of PDAEs that can be formulated as a problem in a Banach space  $\mathcal{X}$ , enabling us to apply existing results to establish the existence and uniqueness of solutions.

Let  $(\mathcal{E}, \|\cdot\|_{\mathcal{E}})$ ,  $(\mathcal{S}, \|\cdot\|_{\mathcal{S}})$ , and  $(\mathcal{H}, \|\cdot\|_{\mathcal{H}})$  be three Banach spaces, and define  $\mathcal{X} = \mathcal{E} \times \mathcal{S}$  with the norm

$$||U||_{\mathcal{X}} = ||V||_{\mathcal{E}} + ||w||_{\mathcal{S}},$$

where  $U = (V, w), V \in \mathcal{E}$ , and  $w \in \mathcal{S}$ .

If  $G(t, x, U(t, x)) = \mathcal{L}(w(t, x)) - \mathcal{G}(t, x, V(t))$ , we can rewrite the initial-boundaryvalue problem (1) as an abstract problem in Banach space  $\mathcal{X}$ :

$$\begin{cases} V_t(t) = \mathcal{A}(V(t)) + \mathcal{F}(t, U(t)), \\ 0 = \mathcal{L}(w(t)) - \mathcal{G}(t, V(t)), \\ V(0) = V_0, \end{cases} \quad (2)$$

where  $\mathcal{A}: \mathcal{E} \to \mathcal{E}, \ \mathcal{F}: \mathbb{R}_+ \times \mathcal{X} \to \mathcal{E}, \ \mathcal{L}: \mathcal{S} \to \mathcal{H}, \ \text{and} \ \mathcal{G}: \mathbb{R}_+ \times \mathcal{E} \to \mathcal{H}.$ 

By a solution, we mean the functions V(t) and w(t), taking values in  $\mathcal{E}$  and  $\mathcal{S}$ , respectively, such that V(t) and w(t) are continuous for  $t \ge 0$ .

This paper is organized as follows. Section 1 presents the preliminaries necessary for understanding the problem. In Section 2, we establish the existence and uniqueness theorems for the local solution of a class of partial differential-algebraic equations (PDAEs). Finally, Section 3 illustrates our theoretical results through concrete applications.

## 2 Preliminary

To make it easier for the reader, we will introduce the Banach spaces  $\mathcal{X}$  and explain the key concepts and theorems that we will use.

**Theorem 2.1.** [1] Let  $\mathcal{F}: [0, +\infty[\times \mathcal{X} \longrightarrow \mathcal{X}]$  be continuous in t for  $t \geq 0$  and locally Lipchitz continuous in U, uniformly in t on bounded intervals. If  $\mathcal{A}$  is the infinitesimal generator of a  $C_0$  semi-group T(t) on  $\mathcal{X}$  then for every  $U_0 \in \mathcal{X}$  there is a  $t_{max} \leq +\infty$ such that the initial value problem

$$\begin{cases} \frac{dU(t)}{dt} = \mathcal{A}U(t) + \mathcal{F}(U(t)), \\ U(0) = U_0, \end{cases} \qquad t \ge 0, \tag{3}$$

has a unique mild solution U on  $[0, t_{max}]$ . Moreover, if  $t_{max} \leq +\infty$  then

$$\lim_{t \to t_{max}} \|U(t)\|_{\mathcal{X}} = +\infty.$$

**Definition 2.2.** [1] Let T(t) be a  $C_0$  semi-group of contractions on  $\mathcal{X}$ . A continuous function  $U : [0,T] \to \mathcal{X}$  is a mild solution to the problem (3) if it satisfies

$$U(t) = T(t)U_0 + \int_0^t T(t-s)\mathcal{F}(U(s)), ds, \quad 0 \le t \le T,$$

where  $\mathcal{F}$  is locally Lipchitz on  $\mathcal{X}$ .

### 3 Existence and Uniqueness

**Theorem 3.1.** Let  $\mathcal{F} : \mathbb{R}_+ \times \mathcal{X} \to \mathcal{E}$  and  $\mathcal{G} : \mathbb{R}_+ \times \mathcal{E} \to \mathcal{H}$  be continuous in t for  $t \geq 0$ , uniformly in t on bounded intervals, and locally Lipschitz continuous in U and V, respectively. Let  $\mathcal{L} : \mathcal{S} \to \mathcal{H}$  be continuous, linear, and invertible. If  $\mathcal{A}$  is the infinitesimal generator of a  $C_0$ -semigroup T(t),  $t \geq 0$ , on  $\mathcal{E}$ , then for every  $V_0 \in \mathcal{E}$ , there exists  $t_{\max} \leq +\infty$  such that the initial value problem

$$\begin{cases} V_t(t) = \mathcal{A}(V(t)) + \mathcal{F}(t, V(t), w(t)), \ t \ge 0\\ 0 = \mathcal{L}(w(t)) - \mathcal{G}(t, V(t))\\ V(0) = V_0. \end{cases}$$
(4)

has a unique solution  $U \in \mathcal{C}([0, tmax[, \mathcal{X}).$ 

*Proof.* By the invertibility of  $\mathcal{L}$ , we have  $w(t) = \mathcal{L}^{-1}\mathcal{G}(t, V(t))$ , where  $w(t) \in \mathcal{S}$  and  $V(t) \in \mathcal{E}$  for  $t \ge 0$ . Substituting w(t) into the first equation, we obtain

$$\begin{cases} V_t(t) = \mathcal{A}(V(t)) + \mathcal{F}(t, V(t), \mathcal{L}^{-1}\mathcal{G}(t, V(t))), \ t \ge 0\\ V(0) = V_0. \end{cases}$$
(5)

We define the operator  $\mathcal{K}$  as follows:

$$\mathcal{K}: \quad \mathbb{R}_+ \times \mathcal{E} \quad \longrightarrow \quad \mathcal{E}$$

$$(t, V(t)) \quad \longmapsto \quad \mathcal{K}(t, V(t)) = \mathcal{F}(t, V(t), \mathcal{L}^{-1}\mathcal{G}(t, V(t)))$$

So, the problem (5) take the form

$$\begin{cases} V_t(t) = \mathcal{A}(V(t)) + \mathcal{K}(t, V(t)), \ t \ge 0\\ V(0) = V_0. \end{cases}$$
(6)

Thus, the problem is defined on  $\mathcal{E}$ . To resolve problem (4), it suffices to solve problem (6). Since  $\mathcal{E}$  is a Banach space and  $\mathcal{A}$  generates a  $C_0$ -semigroup on  $\mathcal{E}$ , it suffices to prove that  $\mathcal{K}$  is continuous in t for  $t \geq 0$ , locally Lipschitz on  $\mathcal{E}$ , and uniformly Lipschitz in t on bounded intervals.

We have  $\mathcal{L}$  is continuous, linear, and invertible, and that both  $\mathcal{F}$  and  $\mathcal{G}$  are continuous in t for  $t \ge 0$ , therefore,  $\mathcal{G}$  is continuous in t for  $t \ge 0$ .

Let  $t' \ge 0$ ,  $C \ge 0$  and  $V_1(t), V_2(t) \in \mathcal{E}$  with  $||V_1(t)||_{\mathcal{E}} \le C$ ,  $||V_2(t)||_{\mathcal{E}} \le C$  and  $t \in [0, t']$ , so

$$\|\mathcal{K}(t,V_1(t)) - \mathcal{K}(t,V_2(t))\|_{\mathcal{E}} = \|F(t,V_1(t),L^{-1}\mathcal{G}(V_1(t))) - F(t,V_2,L^{-1}\mathcal{G}(t,V_2(t)))\|_{\mathcal{E}}$$

Since  $\mathcal{F}$  is locally Lipchitz in U and uniformly in t on bounded intervals, then there exists a positive constant  $L(t', C) \geq 0$  that depends on t' and C such that

$$\begin{aligned} \|\mathcal{K}(t,V_{1}(t)) - \mathcal{K}(t,V_{2}(t))\|_{\mathcal{E}} &\leq L(t',C) \|(V_{1}(t),\mathcal{L}^{-1}\mathcal{G}(t,V_{1}(t))) - (V_{2}(t),\mathcal{L}^{-1}\mathcal{G}(t,V_{2}(t)))\|_{\mathcal{X}}, \\ &= L(t',C)(\|V_{1}(t) - V_{2}(t)\|_{\mathcal{E}} + \|\mathcal{L}^{-1}\mathcal{G}(t,V_{1}(t)) - \mathcal{L}^{-1}\mathcal{G}(t,V_{2}(t))\|_{\mathcal{S}}) \end{aligned}$$

We have that  $\mathcal{L}$  is a continuous, linear, and invertible operator between two Banach spaces. Then,  $\mathcal{L}^{-1}$  is also linear and continuous. Therefore, there exists a positive constant  $L_1(C) \geq 0$ , which depends on C, such that

$$\|\mathcal{K}(t, V_1(t)) - \mathcal{K}(t, V_2)\|_{\mathcal{E}} \leq L(t', C)(\|V_1(t) - V_2(t)\|_{\mathcal{E}} + L_1(C)\|\mathcal{G}(t, (V_1(t)) - \mathcal{G}(t, V_2(t)))\|_{\mathcal{H}})$$

As  $\mathcal{G}$  is locally Lipchitz in V and uniformly in t on bounded intervals, then there exists a positive constant  $L_2(t', C) \geq 0$  that depends on t' and C such that

$$\begin{aligned} \|\mathcal{K}(t,V_{1}(t)) - \mathcal{K}(t,V_{2}(t))\|_{\mathcal{E}} &\leq L(t',C)(\|V_{1}(t) - V_{2}(t)\|_{\mathcal{E}} + L_{1}(C)L_{2}(t',C)\|V_{1}(t) - V_{2}(t)\|_{\mathcal{E}}), \\ &\leq L(t',C)(1 + L_{1}(t',C)L_{2}(t',C))\|V_{1}(t) - V_{2}(t)\|_{\mathcal{E}}. \end{aligned}$$

Then,

$$\|\mathcal{K}(t, V_1(t)) - \mathcal{K}(t, V_2(t))\|_{\mathcal{E}} \le L_3(t', C) \|V_1(t) - V_2(t)\|_{\mathcal{E}}.$$

where  $L_3(t', C) = L(t', C)(1 + L_1(C)L_2(t', C))$ . So,  $\mathcal{K}$  is continuous in t for  $t \ge 0$  and locally Lipchitz in  $\mathcal{E}$  and uniformly in t in bounded intervals. Therefore, by Theorem 2.1, there exists a positive  $t_{max} \le +\infty$  such that problem (6) admits a unique mild solution  $V \in \mathcal{C}([0, t_{max}], \mathcal{E})$ . Moreover, if  $t_{max} < +\infty$  then

$$\lim_{t \to t_{max}} \|V(t)\|_0 = +\infty.$$

By the continuity of  $\mathcal{L}^{-1}$  and  $\mathcal{G}$ , there exists a unique w that satisfies

$$w(t) = \mathcal{L}^{-1}\mathcal{G}(t, V(t)), \ w \in \mathcal{C}([0, t_{max}[, \mathcal{S}).$$

Therefore, problem (4) has a unique solution U = (V, w) that satisfies

$$U \in \mathcal{C}([0, t_{max}[, \mathcal{X})$$

## 4 Application

In this section, we apply the theorem to non-linear problem.

Let  $\mathcal{P}'$  be a nonlinear problem defined as follows.

$$\begin{cases} u_t = u_{xx} + wv + uw, \\ v_t = v_{xx} + wv + u, \\ 0 = w_{xxxx} + w + u + v, \end{cases} \quad t \ge 0 \text{ and } x \in \Omega, \qquad (\mathcal{P}')$$

where  $\Omega$  is a bounded open set in  $\mathbb{R}$  and U = (u, v, w) is an unknown function.

The initial conditions are defined as follows:

$$u(0,x) = u_0$$
,  $v(0,x) = v_0$ ,

and boundary conditions

$$u_x|_{\partial\Omega} = v_x|_{\partial\Omega} = 0.$$

We introduce the Hilbert spaces

$$\mathcal{X} = L^2(\Omega) \times L^2(\Omega) \times H^2_0(\Omega) \text{ and } \mathcal{E} = L^2(\Omega) \times L^2(\Omega),$$

with the inner products given by

$$\langle U_1, U_2 \rangle_{\mathcal{X}} = \int_{\Omega} u_1 u_2 + \int_{\Omega} v_1 v_2 + \int_{\Omega} w_{1xx} w_{2xx} \text{ and } \langle V_1, V_2 \rangle_{\mathcal{E}} = \int_{\Omega} u_1 u_2 + \int_{\Omega} v_1 v_2,$$

and the norms

$$||U||_{\mathcal{X}} = \left(\int_{\Omega} u^2 + \int_{\Omega} v^2 + \int_{\Omega} w_{xx}^2\right)^{\frac{1}{2}} and ||U||_{\mathcal{E}} = \left(\int_{\Omega} u^2 + \int_{\Omega} v^2\right)^{\frac{1}{2}}$$

where  $U_1, U_2 \in \mathcal{X}$ ,  $V_1, V_2 \in \mathcal{E}$  and  $U_1 = (V_1, w_1)$ ,  $U_2 = (V_2, w_2)$ ,  $V_1 = (u_1, v_1)$  and  $V_2 = (u_2, v_2)$ .

We reformulate the problem  $(\mathcal{P}')$  along with the initial condition of U as an abstract problem in the Hilbert space  $\mathcal{X}$  as follows:

$$\begin{cases} V_t(t) = \mathcal{A}(V(t)) + \mathcal{F}(U(t)), \\ 0 = \mathcal{L}(w(t)) - \mathcal{G}(V(t)) \\ V(0) = V_0, \end{cases} \quad t \ge 0,$$

$$(7)$$

where  $\mathcal{A} : \mathcal{D}(A) \subset \mathcal{E} \to \mathcal{E}, \ \mathcal{F} : \mathcal{X} \to \mathcal{E}, \ \mathcal{L} : H_0^2(\Omega) \to H^{-2}(\Omega) \text{ and } \mathcal{G} : \mathcal{E} \to H^{-2}(\Omega)$ where  $\mathcal{D}(\mathcal{A})$  is a domain of  $\mathcal{A}$ . These applications are given in the following form:

For each  $U = (V, w) \in \mathcal{X}$  such that  $V = (u, v) \in \mathcal{D}(A)$ ,

$$\mathcal{A}(V) = \begin{pmatrix} u_{xx} \\ v_{xx} \end{pmatrix}, \mathcal{F}(U) = \begin{pmatrix} wv + uw \\ wv + u \end{pmatrix}, \mathcal{L}(w) = w_{xxxx} + w$$

and

$$\mathcal{G}(V) = u + v.$$

The domain of  $\mathcal{A}$  is given by

$$\mathcal{D}(\mathcal{A}) = H^2(\Omega) \cap H^1_0(\Omega) \times H^2(\Omega) \cap H^1_0(\Omega).$$

**proposition 4.1.** The operator  $\mathcal{L} : H_0^2(\Omega) \to H^{-2}(\Omega)$  is continuous, linear, and invertible.

*Proof.* We decompose this proof into three steps. First, we show that  $\mathcal{L}$  is well-defined on  $H_0^2(\Omega)$ . Then, we prove that it is linear and continuous. Finally, we demonstrate that  $\mathcal{L}$  is bijective.

### Step 1: Show that the operator $\mathcal{L}$ is well-defined on the space $H_0^2(\Omega)$ .

Let  $w \in H^2_0(\Omega)$  and  $\psi \in \mathcal{D}(\Omega)$ :

$$\begin{aligned} |\langle \mathcal{L}(w), \psi \rangle| &= |\langle w_{xxxx} + w, \psi \rangle|, \\ &\leq |\langle w_{xxxx}, \psi \rangle| + |\langle w, \psi \rangle|, \\ &\leq |\langle w_{xx}, \psi_{xx} \rangle| + |\langle w, \psi \rangle|, \\ &\leq \|w\|_{H^{2}_{0}(\Omega)} \|\psi\|_{H^{2}_{0}(\Omega)} + \|w\|_{L^{2}(\Omega)} \|\psi\|_{L^{2}(\Omega)}, \end{aligned}$$

$$(8)$$

hence  $\mathcal{L}(w)$  is well-defined on the space  $\mathcal{D}(\Omega)$  for all  $w \in H_0^2(\Omega)$ .

The linearity is clear, now let us show the continuity. Let  $w \in H^2_0(\Omega)$  and  $\psi_n$  be a sequence convergent to  $\psi$  in  $\mathcal{D}(\Omega)$ , then

$$\begin{aligned} |\langle \mathcal{L}(w), \psi_n - \psi \rangle| &= |\langle w_{xxxx} + w, \psi_n - \psi \rangle|, \\ &\leq |\langle w_{xxxx}, \psi_n - \psi \rangle| + |\langle w, \psi_n - \psi \rangle|, \\ &\leq |\langle w_{xx}, (\psi_n)_{xx} - \psi_{xx} \rangle| + |\langle w, \psi_n - \psi \rangle|, \\ &\leq ||w_{xx}||_{L^2(\Omega)} ||(\psi_n)_{xx} - \psi_{xx}||_{L^2(\Omega)} + ||w||_{L^2(\Omega)} ||\psi_n - \psi||_{L^2(\Omega)}, \\ &\leq C(\Omega) \left( ||w_{xx}||_{L^2(\Omega)} ||(\psi_n)_{xx} - \psi_{xx}||_{\infty} + ||w||_{L^2(\Omega)} ||\psi_n - \psi||_{\infty} \right), \end{aligned}$$

where  $C(\Omega) > 0$ . As  $n \to \infty$ , we have

$$\lim_{n \to \infty} |\langle \mathcal{L}(w), \psi_n - \psi \rangle| = 0,$$

which implies that  $\mathcal{L}(w)$  is continuous on  $\mathcal{D}(\Omega)$ . Therefore,  $\mathcal{L}(w) \in \mathcal{D}'(\Omega)$  for all  $w \in H^2_0(\Omega)$ .

Let  $w \in H^2_0(\Omega)$ , by (8) and Poincarre inequality, there exists a positive constant C > 0 such that

$$|\langle \mathcal{L}(w), \psi \rangle| \le C \|w\|_{H^2_0(\Omega)} \|\psi\|_{H^2_0(\Omega)}, \text{ for all } \psi \in \mathcal{D}(\Omega).$$
(9)

Since  $\mathcal{L}(w) \in \mathcal{D}'(\Omega)$  and satisfies (9) for all  $w \in H^2_0(\Omega)$ , it follows that  $\mathcal{L}(w) \in H^{-2}(\Omega)$  for all  $w \in H^2_0(\Omega)$ . Therefore, the operator  $\mathcal{L}$  is well-defined.

#### Step 3: Show that the operator $\mathcal{L}$ is linear and continuous.

Let  $w, v \in H^2_0(\Omega)$ , then

$$\begin{aligned} |\langle \mathcal{L}(w), v \rangle| &= |\langle w_{xxxx} + w, v \rangle|, \\ &= |\int_{\Omega} w_{xxxx}v + \int_{\Omega} wv|, \\ &\leq |\int_{\Omega} w_{xx}v_{xx}| + |\int_{\Omega} wv|, \\ &\leq ||w_{xx}||_{L^{2}(\Omega)} ||v_{xx}||_{L^{2}(\Omega)} + ||w||_{L^{2}(\Omega)} ||v||_{L^{2}(\Omega)}, \end{aligned}$$

by Poincare's inequality, there exists C > 0 such that

 $|\langle \mathcal{L}(w), v \rangle| \le C ||w||_{H^2_0(\Omega)} ||v||_{H^2_0(\Omega)}.$ 

 $\operatorname{So}$ 

$$\|\mathcal{L}(w)\|_{H^{-2}(\Omega)} = \sup_{v \in H^{2}_{0}(\Omega), \, \|v\|_{H^{2}_{0}(\Omega)} \le 1} |\langle \mathcal{L}(w), v \rangle| \le C \|w\|_{H^{2}_{0}(\Omega)}.$$
 (10)

Therefore,  $\mathcal{L}$  is continuous on  $H_0^2(\Omega)$ .

### Step 2: Show that the operator $\mathcal{L}$ is bijective.

Let  $g \in H^{-2}(\Omega)$ , such that  $\mathcal{L}(w) = g$ , then the equation becomes:

$$w_{xxxx} + w = g$$

Let  $\phi \in H^2_0(\Omega)$ , Multiplying by  $\phi$  and integrating by parts, we obtain

$$\int_{\Omega} w_{xx} \phi_{xx} + \int_{\Omega} w \phi = \int_{\Omega} g \phi$$

We define

$$a: H^2_0(\Omega) \times H^2_0(\Omega) \longrightarrow \mathbb{R}$$

$$(w,v) \longmapsto a(w,\phi) = \int_{\Omega} w_{xx}\phi_{xx} + \int_{\Omega} w\phi$$

and

$$l: H^2_0(\Omega) \longrightarrow \mathbb{R}$$

$$\phi \qquad \longmapsto \ l(\phi) = \int_{\Omega} g\phi.$$

Since a is bilinear, it suffices to show that a is continuous and coercive. Let  $w, \phi \in H^2_0(\Omega)$ 

$$|a(w,\phi)| = \left| \int_{\Omega} w_{xx} \phi_{xx} + \int_{\Omega} w \phi \right|.$$

By Poincare's lemma, there exists C > 0 depending only on  $\Omega$  such that

$$|a(w,\phi)| \le C ||w||_{H^2_0(\Omega)} ||\phi||_{H^2_0(\Omega)},$$

then a is continuous.

So, for  $w \in H^2_0(\Omega)$ 

$$a(w,w) = \left(\int_{\Omega} w_{xx}^2 + \int_{\Omega} w^2\right) \ge ||w||_{H^2_0(\Omega)}^2,$$

then a is coercive.

Since  $g \in H^{-2}(\Omega)$ , there exists a constant C > 0 such that

$$|\langle g, \psi \rangle| \le C \|\psi\|_{H^2_0(\Omega)}, \text{ for all } \psi \in \mathcal{D}(\Omega).$$

Now, for any  $\psi \in H_0^2(\Omega)$ , there exists a sequence  $(\psi_n) \subset \mathcal{D}(\Omega)$  such that  $\psi_n \to \psi$  in  $H_0^2(\Omega)$ . Therefore,

$$|l(\psi)| = |\langle g, \psi \rangle| = \left| \langle g, \lim_{n \to \infty} \psi_n \rangle \right|.$$

By the continuity of g as a functional in  $H^{-2}(\Omega)$ ,

$$|l(\psi)| \le \lim_{n \to \infty} |\langle g, \psi_n \rangle| \le C \lim_{n \to \infty} \|\psi_n\|_{H^2_0(\Omega)} = C \|\psi\|_{H^2_0(\Omega)}.$$

Hence, l is a continuous linear on  $H_0^2(\Omega)$ .

Since l is linear continuous and we have a is bilinear, continuous, and coercive, by applying the Lax-Milgram lemma, there exists unique solution  $w \in H_0^2(\Omega)$  that satisfies

$$a(w,\phi) = l(\phi), \ \forall \phi \in \mathcal{S}.$$

Therefore,  $\mathcal{L}$  is invertible.

**proposition 4.2.** The operator  $\mathcal{G} : \mathcal{E} \longrightarrow \mathcal{S}$  given by

$$\mathcal{G}(V) = u + v$$

is a continuous linear operator.

Proof. Let  $V = (u, v) \in \mathcal{E}$ , then  $\mathcal{G}(V) = (u + v) \in L^2(\Omega)$ , hence  $\mathcal{G}(V) \in \mathcal{D}'(\Omega)$ . By the characterization of  $H^{-1}(\Omega)$ , it follows that  $\mathcal{G}(V) \in H^{-1}(\Omega)$ . Therefore, the operator  $\mathcal{G}(V) \in H^{-2}(\Omega)$ .

$$\|\mathcal{G}(V)\|_{H^{-2}(\Omega)} = \sup_{v \in H^{2}_{0}(\Omega), \|v\|_{H^{2}_{0}(\Omega)} \leq 1} |\langle \mathcal{G}(V), v \rangle| \leq \|u + v\|_{L^{2}(\Omega)} \leq \sqrt{2} \left( \|u\|_{L^{2}(\Omega)}^{2} + \|v\|_{L^{2}(\Omega)}^{2} \right)^{\frac{1}{2}}.$$
(11)

 ${\mathcal G}$  is a continuous linear operator.

Since the domain of  $\mathcal{A}$  is dense, it suffices to show that it is dissipative and maximal.

**proposition 4.3.** The operator  $\mathcal{A}$  is dissipative, that is  $Re\langle \mathcal{A}V, V \rangle_{\mathcal{E}} \leq 0$ .

*Proof.* Let  $V = (u, v)^T \in \mathcal{D}(\mathcal{A})$ , then

$$\langle \mathcal{A}V, V \rangle_{\mathcal{E}} = \int_{\Omega} u_{xx} u + \int_{\Omega} v_{xx} v,$$
  
$$= -\int_{\Omega} (u_x)^2 - \int_{\Omega} (v_x)^2,$$
  
$$\leq 0.$$

So  $\mathcal{A}$  is dissipative.

**proposition 4.4.** The operator  $\mathcal{A}$  is maximal, that is  $\mathcal{R}(I - \mathcal{A}) = \mathcal{E}$ .

*Proof.* Let  $g = (g_1, g_2)^T \in \mathcal{E}$ . We will prove that there exists  $V \in \mathcal{D}(\Omega)$  satisfying

$$(I - \mathcal{A})V = g \Leftrightarrow \begin{cases} -u_{xx} + u = g_1, \\ -v_{xx} + v = g_2. \end{cases}$$

By applying the Lax-Milgram lemma, along with the regularity of elliptic problems and the boundary conditions, there exists a unique  $V = (u, v) \in \mathcal{D}(\mathcal{A})$  such that

$$(I - \mathcal{A})V = g$$

Consequently,

$$\mathcal{R}(I-\mathcal{A})=\mathcal{E}.$$

Thus,  $\mathcal{A}$  is maximal.

**Theorem 4.5.** The operator  $\mathcal{A}$  generates a  $C_0$ -semi-group of contractions  $T(t) = e^{t\mathcal{A}}$  on  $\mathcal{E}$ .

*Proof.* Since  $\mathcal{D}(\mathcal{A})$  is dense in  $\mathcal{E}$ , from the propositions 4.3 and 4.4  $\mathcal{A}$  is dissipative, maximal, and by the Lumer-Phillips theorem  $\mathcal{A}$  generates a  $C_0$  semi-group of contractions in  $\mathcal{E}$ .

**Lemma 4.6.** The operator  $\mathcal{F}: \mathcal{X} \longrightarrow \mathcal{E}$  given by

$$\mathcal{F}(U) = \left(\begin{array}{c} wv + wu\\ wv + u \end{array}\right)$$

is locally Lipschitz.

*Proof.* Let C > 0 and  $U_1, U_2 \in \mathcal{X}$  such that  $||U_1||_{\mathcal{X}} \leq C$  and  $||U_2||_{\mathcal{X}} \leq C$ , then

$$\|\mathcal{F}(U_1) - \mathcal{F}(U_2)\|_{\mathcal{E}}^2 = \int_{\Omega} (w_1 v_1 - w_2 v_2 + u_1 w_1 - u_2 w_2)^2 + \int_{\Omega} (w_1 u_1 - w_2 u_2 + u_1 - u_2)^2.$$

Firstly, we analyze the term

$$\int_{\Omega} (w_1 v_1 - w_2 v_2 + u_1 w_1 - u_2 w_2)^2 \le 2 \left( \int_{\Omega} (w_1 v_1 - w_2 v_2)^2 + \int_{\Omega} (u_1 w_1 - u_2 w_2)^2 \right),$$

Next, we estimate

$$\int_{\Omega} (u_1 w_1 - u_2 w_2)^2 = \int_{\Omega} ((u_1 - u_2) w_1 + u_2 (w_1 - w_2))^2, \\
\leq 2 \left( \int_{\Omega} ((u_1 - u_2) w_1)^2 + \int_{\Omega} u_2 (w_1 - w_2)^2 \right), \\
\leq 2 \left( \int_{\Omega} ((u_1 - u_2) w_1)^2 + \int_{\Omega} u_2 (w_1 - w_2)^2 \right), \\
\leq 2 \left( \|w_1\|_{\infty}^2 \int_{\Omega} (u_1 - u_2)^2 + \|w_1 - w_2\|_{\infty}^2 \int_{\Omega} u_2^2 \right),$$
(12)

By Poincaré's inequality and the continuity of  $H^2(\Omega) \hookrightarrow L^{\infty}(\Omega)$ , there exists a positive constant  $C(\Omega)$  such that

$$\int_{\Omega} (u_1 w_1 - u_2 w_2)^2 \leq 2C(\Omega) \left( \|w_1\|_{H_0^2(\Omega)}^2 \|u_2 - u_1\|_{L^2(\Omega)}^2 + \|w_1 - w_2\|_{H_0^2(\Omega)}^2 \|u_2\|_{L^2(\Omega)}^2 \right),$$

$$\leq 2C^2 C(\Omega) \left( \|u_2 - u_1\|_{L^2(\Omega)}^2 + \|w_1 - w_2\|_{H_0^2(\Omega)}^2 \right).$$
(13)

Similarly, we have

$$\int_{\Omega} (w_1 v_1 - w_2 v_2)^2 \le 2C^2 C(\Omega) \left( \|v_2 - v_1\|_{L^2(\Omega)}^2 + \|w_1 - w_2\|_{H^2_0(\Omega)}^2 \right).$$
(14)

Combining (13) and (14), we obtain

$$\int_{\Omega} (w_1 v_1 - w_2 v_2 + u_1 w_1 - u_2 w_2)^2 \leq 8C^2 C(\Omega) \left( \|u_2 - u_1\|_{L^2(\Omega)}^2 + \|v_2 - v_1\|_{L^2(\Omega)}^2 + \|w_1 - w_2\|_{H^2_0(\Omega)}^2 \right) \\
\leq 8C^2 C(\Omega) \|U_1 - U_2\|_{\mathcal{X}}^2.$$
(15)

Secondly, we estimate

$$\begin{split} \int_{\Omega} (w_1 v_1 - w_2 v_2 + u_1 - u_2)^2 &\leq 2 \left( \int_{\Omega} (w_1 v_1 - w_2 v_2)^2 + \int_{\Omega} (u_1 - u_2)^2 \right), \\ &\leq 2 \left( 2C^2 C(\Omega) \left( \|v_2 - v_1\|_{L^2(\Omega)}^2 + \|w_1 - w_2\|_{H^2_0(\Omega)}^2 \right) + \|u_2 - u_1\|_{L^2(\Omega)}^2 \right), \\ &\leq 2M \|U_1 - U_2\|_{\mathcal{X}}^2, \end{split}$$
(16)

where  $M = max(1, 2C^2C(\Omega))$ . Combining the results (15) and (16), we find

$$\|\mathcal{F}(U_1) - \mathcal{F}(U_2)\|_{\mathcal{E}}^2 \leq L^2(C) \|U_1 - U_2\|_{\mathcal{X}}^2,$$

where  $L^2(C) = max(2M, 8C^2C(\Omega))$ . Therefor

$$\|\mathcal{F}(U_1) - \mathcal{F}(U_2)\|_{\mathcal{E}} \le L(C) \|U_1 - U_2\|_{\mathcal{X}}.$$

This proves that  $\mathcal{F}$  is locally Lipschitz continuous.

**Theorem 4.7.** If  $V_0 \in \mathcal{E}$ , there is a  $t_{max} \leq +\infty$  such that the initial value problem  $(\mathcal{P}')$  has a unique solution  $U \in \mathcal{C}([0, t_{max}[, \mathcal{X}).$ 

Proof. By Propositions 4.1, 4.2, Lemma 4.6 and Theorem 4.5, we have the following: the operator  $\mathcal{L}$  is continuous, linear, and invertible,  $\mathcal{G}$  is a continuous linear operator,  $\mathcal{F}$  is locally Lipschitz, and  $\mathcal{A}$  is the infinitesimal generator of a  $C_0$ -semi-group of contractions T(t). Then, by Theorem 3.1, there exists  $t_{max} \leq +\infty$  such that the initial value problem (7) has a unique solution  $U \in \mathcal{C}([0, t_{max}[, \mathcal{X}), \text{ and this solution satisfies the problem } (\mathcal{P}')$ .

## References

- A. Pazy, \*semi-groups of Linear Operators and Applications to Partial Differential Equations\*, vol. 44, Springer Science & Business Media, 2012.
- [2] K. E. Brenan, S. L. Campbell, and L. R. Petzold, \*Numerical Solution of Initial-Value Problems in Differential-Algebraic Equations\*, 2nd ed., Classics in Applied Mathematics 14, Society for Industrial and Applied Mathematics, 1987.
- [3] R. März, "On initial value problems in differential-algebraic equations and their numerical treatment," \*Computing\*, vol. 35, no. 1, pp. 13–37, 1985.
- [4] C. W. Gear and L. R. Petzold, "ODE methods for the solution of differential/algebraic systems," \*SIAM Journal on Numerical Analysis\*, vol. 21, no. 4, pp. 716–728, 1984.
- [5] Debrabant, Kristian, and Karl Strehmel. "Convergence of Runge-Kutta methods applied to linear partial differential-algebraic equations." Applied Numerical Mathematics 53.2-4 (2005): 213-229.
- [6] C. Schneider, "Rosenbrock-type methods adapted to differential-algebraic systems," \*Mathematics of Computation\*, vol. 56, no. 193, pp. 201–213, 1991.
- [7] A. Fortin and S. Garon, \*Les Éléments Finis: De la Théorie à la Pratique\*, Université Laval, 2011.

- [8] W. Lambert, A. Alvarez, I. Ledoino, D. Tadeu, D. Marchesin, and J. Bruining, "Mathematics and numerics for balance partial differential-algebraic equations (PDAEs)," \*Journal of Scientific Computing\*, vol. 84, no. 2, p. 29, 2020.
- [9] W. Lucht and K. Strehmel, "Discretization-based indices for semilinear partial differential-algebraic equations," \*Applied Numerical Mathematics\*, vol. 28, no. 2–4, pp. 371–386, 1998.
- [10] M. Jafari, M. M. Hosseini, S. T. Mohyud-Din, and M. Ghovatmand, "Modified homotopy perturbation method for solving nonlinear PDAEs and its applications in nanoelectronics," \*International Journal of Nonlinear Sciences and Numerical Simulation\*, vol. 11, no. 12, pp. 1047–1058, 2010.
- [11] L. Angermann and J. Rang, "Perturbation index of linear partial differentialalgebraic equations with a hyperbolic part," \*Central European Journal of Mathematics\*, vol. 5, pp. 19–49, 2007.
- [12] R. Riaza, \*Differential-Algebraic Systems: Analytical Aspects and Circuit Applications\*, World Scientific, 2008.
- [13] Schwarz, Diana Estévez. "Consistent initialization for index-2 differential algebraic equations and its application to circuit simulation." (2000).
- [14] Krzywicki, A., and T. Nadzieja. "A nonstationary problem in the theory of electrolytes." Quarterly of applied mathematics 50.1 (1992): 105-107.
- [15] Temam, R. "Qualitative Properties of Navier-Stokes Equations." North-Holland Mathematics Studies. Vol. 30. North-Holland, 1978. 485-494.
- [16] Lamour, René, Roswitha März, and Caren Tischendorf. PDAEs and further mixed systems as abstract differential algebraic systems. Humboldt-Universität zu Berlin, Mathematisch-Naturwissenschaftliche Fakultät II, Institut für Mathematik, 2005.
- [17] Marszalek, Wieslaw. Analysis of partial differential algebraic equations. North Carolina State University, 1997.
- [18] Brezis, Haïm, and Petru Mironescu. Sobolev maps to the circle. Vol. 96. Springer US, 2021.