Bob Krekelberg \square

Department of Information and Computing Sciences, Utrecht University, The Netherlands

Alison Hsiang-Hsuan Liu 🖂 🕩

Department of Information and Computing Sciences, Utrecht University, The Netherlands

— Abstract

Online interval coloring is a fundamental problem in graph algorithms and scheduling. Although the optimal online algorithm for coloring arbitrary-length intervals is known as 3-competitive (Kierstead and Trotter, 1981), the interest in coloring bounded-length intervals arose recently (Chybowska-Sokół et al., 2024). On the other hand, people are also interested in the FirstFit algorithm's performance because of its elegance. The competitive ratio of FirstFit on unit-length intervals is exact $2 - \frac{1}{\omega}$, where ω is the optimal number of colors needed (Epstein and Levy, 2005). However, for arbitrary-length intervals, the competitive ratio is only known to be between 5 (Kierstead et al., 2005) and 8 (Narayanaswamy and Subhash Babu, 2008). It has been open for a long time what the actual competitive ratio of FirstFit is for arbitrary-length intervals.

In this paper, we study the performance of the FirstFit algorithm for the online unit-length intervals coloring problem where the intervals can be either open or closed, which serves a further investigation towards the actual performance of FirstFit. We develop a sophisticated counting method by generalizing the classic neighborhood bound, which limits the color FirstFit can assign an interval by counting the potential intersections. In the generalization, we show that for any interval, there is a critical interval intersecting it that can help reduce the overestimation of the number of intersections, and it further helps bound the color an interval can be assigned. The technical challenge then falls on identifying these critical intervals that guarantee the effectiveness of counting. Using this new mechanism for bounding the color that FirstFit can assign an interval, we provide a tight analysis of 2ω colors when all intervals have integral endpoints and an upper bound of $\lceil \frac{7}{3}\omega \rceil - 2$ colors for the general case, where ω is the optimal number of colors needed for the input set of intervals.

2012 ACM Subject Classification Theory of computation \rightarrow Online algorithms

Keywords and phrases Online algorithms, First-Fit algorithm, Interval coloring, Unit intervals, Open and closed intervals

1 Introduction

This work studies the online open/closed unit-length interval coloring problem, which is a variant of online interval coloring problem. The input consists of a sequence of unit-length intervals, where each interval is either an open interval $(r, r + 1) = \{i \mid r < i < r + 1\}$ or a closed interval $[r, r + 1] = \{i \mid r \leq i \leq r + 1\}$. The intervals are released to the algorithm one by one. Once an interval is released, the online algorithm has to irrevocably assign the interval a color that is not assigned to any previously released interval that overlaps with the released interval. The aim is to minimize the colors used. In this work, we study the performance of the FIRSTFIT algorithm for coloring the open/closed unit-length intervals.

As a fundamental problem in graph theory and scheduling areas, intensive research is being conducted on the online interval coloring problem. The optimal online algorithm for the general case where intervals have arbitrary lengths uses at most $3\omega - 2$ colors where ω is the optimal number of colors [1, 10, 13]. Recently, interest in coloring bounded-length intervals arose, and better algorithms were found for special families of instances. For intervals with the length within some fixed range between 1 and $\sigma \geq 1$, a ($\sigma + 1$)-competitive algorithm

was proposed [2], which improved the performance of online algorithms for the cases where $\sigma < 2$.

On the other hand, the FIRSTFIT algorithm, a classical greedy algorithm, is interesting for its elegance. The FIRSTFIT algorithm is currently the best algorithm for (closed) unit-length intervals, which uses $2\omega - 1$ colors for any ω -colorable set of unit-length intervals [1, 5]. For arbitrary-length intervals, the competitive ratio is between 5 and 8 [9, 11], and it is open for a long time what the actual competitive ratio is.

The motivation to look into the performance of the FIRSTFIT algorithm on open/closed unit-length intervals coloring is two-fold. First, the performance of the FIRSTFIT algorithm is not fully understood. It is known that for unit-length closed intervals, the FIRSTFIT algorithm is exactly $(2 - \frac{1}{\omega})$ -competitive, where ω is the optimal number of colors needed. However, for general instances, the FIRSTFIT algorithm is at least 5-competitive. The performance of the FIRSTFIT algorithm is unclear for the instances between these cases. As the open/closed unit-length intervals case is the smallest possible instance of the coloring problem on bounded length intervals, this case is key to knowing the actual competitive ratio of the FIRSTFIT algorithm.

Moreover, this research helps us understand the impact of " ε -uncertainty". It can be considered that closed unit-length intervals are $\varepsilon \approx 0$ longer than the open unit-length intervals. This difference of ε increases the complexity of the problem as an open interval can be a proper subset of a closed interval, which was previously not the case when only closed unit-length intervals were considered. It was shown that the FIRSTFIT algorithm is at least 2-competitive [3], meaning that the open/closed unit-length intervals case is strictly "harder" for the FIRSTFIT algorithm than the closed unit-length intervals case. Finding the competitive ratio of the FIRSTFIT algorithm on open/closed unit-length intervals coloring quantifies the power of the uncertainty of ε -difference by understanding how much the adversary can use this uncertainty to trap the FIRSTFIT algorithm.

Related work

For the online interval coloring problem, Kierstead and Trotter designed an optimal online algorithm that uses at most $3\omega - 2$ colors on an ω -colorable interval graph [10]. Independently, Chrobak and Ślusarek found the same upper and lower bounds [1, 13].

Recently, interest in coloring bounded-length intervals arose. Chybowska-Sokół et al. [2] studied the online interval coloring problem where the size of the intervals is within some fixed range $[1, \sigma_{\geq 1}]$. When $\sigma = \infty$, it is the previously mentioned general case, and when $\sigma = 1$, it is the unit-length case. They proposed a $(\sigma + 1)$ -competitive algorithm, which improved the performance of the online algorithm for $1 < \sigma < 2$. Curbelo [4] then shows that, in the setting where the interval representation is unknown for the algorithm, for any $\varepsilon > 0$, there exists an $\sigma > 1$, such that any algorithm is at least $(3 - \varepsilon)$ -competitive. Matching the upper bound from Kierstead and Trotter for coloring intervals of general length.

FirstFit. On the other hand, despite the aforementioned results that included sophisticated algorithms, people are also interested in the performance of the naive but easy-toimplement FIRSTFIT algorithm. The exact competitive ratio of the FIRSTFIT algorithm is a long standing open problem. As early as 1976, Witsenhausen [14], and independently, in 1988, Chrobak and Ślusarek [1], proved that FIRSTFIT is at least 4 competitive. Later, Ślusarek [13] improved this bound to 4.45. The best lower bound known to date is from Kierstead et al. [9]. They prove that for every $\varepsilon > 0$, there exist an input such that the FIRSTFIT algorithm uses strictly more than $(5 - \varepsilon)\omega$ colors.

For the upper bounds of the competitive ratio of FIRSTFIT, Kierstead [7] first proved a

constant competitive of 40 in 1988. Later, this result was improved to 25.72 by Kierstead and Qin [8]. The next breakthrough was in 2003, when Pemmaraju et al. [12] showed that the FIRSTFIT algorithm is at most 10-competitive. The current best analysis of FIRSTFIT was by Narayanaswamy and Subhash Babu [11], who showed that FIRSTFIT uses at most $8\omega - 3$ colors on a set of ω -colorable intervals.

The performance of the FIRSTFIT algorithm has also been of interest in the special cases. Chybowska-Sokół et al. [2] studied the FIRSTFIT algorithm where the size of the intervals is within some fixed range $[1, \sigma_{\geq 1}]$. The authors first show a trivial bound of $\lceil \sigma + 1 \rceil \omega$ on the number of colors FIRSTFIT uses, where ω is the optimal coloring number.

In the case where only unit-length (closed) intervals are considered, Chrobak and Ślusarek [1], and independently also Epstein and Levy [5] proved that the FIRSTFIT algorithm uses exactly $2\omega - 1$ colors for a set of ω -colorable unit-length intervals.

In 2022, instead of only closed intervals as is typically considered, Biró and Curbelo proved that FIRSTFIT uses at least 2ω and at most $3\omega - 3$ colors when the optimal number of colors is ω [3, Theorem 6.0.1]. Furthermore, Curbelo claimed that FIRSTFIT uses at most $3\omega - f(\omega)$ colors for a non-specified function f, which increases with ω . In this paper, we find a better bound for this case.

FirstFit and counting. The analysis of the FIRSTFIT algorithm is generally done via careful counting intersections a critical interval can have. The elegance of the analysis lies in determining from which interval to count its number of intersecting intervals. The first analysis by Kierstead and Qin [7, 8] identify a large set of mutual-intersecting intervals and then carefully count structural properties of the graph, such as the number of overlapping intervals. The later results by Pemmaraju et al. [12] and by Narayanaswamy and Subhash Babu [11] first represent the overlapping structure of intervals using "columns". The columns are assigned different labels that encrypt the information of the intervals structure and colors assigned. Then, the analysis carefully counts the occurrences of each label to bound the total number of colors used by the FIRSTFIT algorithm. Also, for unit-length (closed) intervals, the analysis carefully counts for each interval at what positions an interval could intersect it and how many of such intervals there could be [1, 5].

Our contribution

Following the traditional counting approach, we analyze the FIRSTFIT algorithm on open/closed unit-length intervals by counting how many intersections an interval can have.

Our main technical contribution is a generalization of the traditional neighborhood bound of the FIRSTFIT algorithm (Lemma 4). Informally, this lemma shows that the color assigned to interval I by FIRSTFIT can be bounded via another interval intersecting I that is critical in the sense that it is a relatively high-colored interval but with relatively few intersections. Our results are built on this newly-observed property of FIRSTFIT. In a nutshell, for any interval I, by the fact that the input instance consists only of unit-length open or closed intervals, our analysis identifies the critical interval intersecting I. It shows that the number of intersections is bounded by sophisticated counting.

For a special case where all intervals have integral endpoints, we prove that FIRSTFIT is 2-competitive, which is tight as it matches the FIRSTFIT algorithm lower bound by Biró and Curbelo [3]. Formally, for any instance that can be colored by ω colors, we show that

▶ **Theorem 1.** For any unit interval graph with open and closed intervals with integral endpoints, the FIRSTFIT algorithm for online coloring uses at most 2ω colors.

Then, for unit-length intervals with arbitrary endpoints, we prove the following.

▶ **Theorem 2.** For any unit interval graph with open and closed intervals with arbitrary endpoints, the FIRSTFIT algorithm for online coloring uses at most $\lfloor \frac{7}{3}\omega \rfloor - 2$ colors.

Paper organization

Section 2 defines the problem formally. We also introduce the key lemmas, the framework of our analysis, and important concepts that are used heavily throughout the paper. As a starter, Section 3 shows the proof of 2-competitiveness (Theorem 1) of FIRSTFIT on open/closed unit-length intervals with integral endpoints. We also prove some interesting facts that hold in both the restricted and the more general setting where intervals have arbitrary endpoints. Section 4 then analyze the case where the unit-length intervals have arbitrary endpoints and prove Theorem 2. Finally, we end in Section 5 with concluding remarks. Due to the page limit, we leave most of the proofs in Section 4 to the full version in the appendix.

2 Preliminaries, key lemmas, and the framework

The input to our online coloring problem is an ordered set of open and closed unit-length intervals $\mathcal{I} = \{I_1, I_2, \cdots\}$, where the ordering of the set of intervals is the order in which they are revealed to the online algorithm. That is, I_j is revealed earlier than I_k if j < k. Each interval $I_j \in \mathcal{I}$ is either an *open* unit-length interval $(r_j, r_j + 1)$ or a *closed* unit-length interval $[r_j, r_j + 1]$ for some real number r. In either case, we say r_j and $r_j + 1$ are the *endpoints* of the interval. Note that the value of r_j is unrelated to the order of I_j in \mathcal{I} . Namely, r_j can be smaller than, equal to, or larger than r_k when j < k. We ignore the suffix of intervals when the reveal ordering is not necessarily to be emphasized.

Two intervals are referred to as *twins* when they are identical in terms of the exact location. That is, for interval I = [r, r+1], all intervals I' = [r, r+1] are its twins. Similarly, for interval I = (r, r+1), all intervals I' = (r, r+1) are its twins. However, an interval (r, r+1) is not a twin of [r, r+1], and vice versa. When intervals I and I' are twins, we write $I \equiv I'$. We further denote the set of twins of I by $\mathcal{T}(I)$. Note that we define $I \notin \mathcal{T}(I)$ for later usage.

A proper coloring of a set of intervals is a function $c : \mathcal{I} \to \mathbb{N}$, such that $c(I) \neq c(I')$ if $I \cap I' \neq \emptyset$. The algorithm aims to find a proper coloring of the intervals using the least number of colors.

We use terms from graph algorithms. The intervals in \mathcal{I} form an *interval graph*, where each $I \in \mathcal{I}$ is a vertex in the interval graph, and there is an edge between the vertices corresponding to I and I' if and only if $I \cap I' \neq \emptyset$. We denote the *neighborhood* of I by $\mathcal{N}(I)$, which is the set of all the intervals interesting with I.

As the coloring depends on intersecting intervals as well as the ordering of the input, we define the ℓ -neighborhood of an interval as its neighborhood at the moment when I_{ℓ} is revealed to the algorithm. Formally, $\mathcal{N}_{\ell}(I_j) = \{I_k \mid I_j \cap I_k \neq \emptyset \text{ and } k < \ell\}$.¹ Naturally, $\mathcal{N}(I_j) = \mathcal{N}_{|\mathcal{I}|}(I_j) \geq \mathcal{N}_{\ell}(I_j)$ for any $\ell \in [j, |\mathcal{I}|]$. Note that $\mathcal{N}_j(I_j)$ is the set of all intervals that intersect with I_j when I_j is revealed.

Interval graphs are *perfect graphs* [6], and the chromatic number of the graph is equal to the maximum clique size ω . This clique corresponds to the maximum set of mutually

¹ We only use $\mathcal{N}_{\ell}(I_i)$ for $\ell \geq j$.

intersecting intervals for the corresponding intervals. We denote ω as the number of colors an optimal solution uses for the given input \mathcal{I} .

FirstFit algorithm. The FIRSTFIT algorithm assigns each interval the least available color, that is, using $FF(I_j)$ to denote the color assigned to I_j by FIRSTFIT, $FF(I_j) \leftarrow \min\{\mathbb{N} \setminus \{FF(I) \mid I \in N_j(I_j)\}\}$. This greedy strategy provides an upper bound for the color of an interval. That is, the color of an interval can never be larger than 1 color above the number of intervals it intersects. More specifically, the color of the interval will be at most 1 greater than the size of the neighborhood when it is revealed. Formally,

▶ Lemma 3. (Neighborhood bound) For any interval $I_j \in \mathcal{I}$, its color assigned by FIRSTFIT is at most $1 + |N_j(I_j)|$.

Proof. Assume aiming towards contradiction that $FF(I_j) \ge 2 + |N_j(I_j)|$. Then, according to the pigeonhole principle, there must exist at least one color $1 \le c < FF(I_j)$ that is not assigned to any of the intervals in $N_j(I_j)$. Then $FF(I_j) = \min\{\mathbb{N} \setminus \{FF(I) \mid I \in N_j(I_j)\}\} \le c$, and it leads to a contradiction.

Pivot interval and ideas of further bonding the color. In this work, we further generalize Lemma 3 by considering a *pivot* interval in the neighborhood of the interval I and use the pivot interval to bound the color FF(I) more carefully. Informally, given any interval I and a pivot interval I^* that can be any interval intersecting with I, FF(I) is no more than 1 plus the number of intervals in I's neighborhood with color greater than the color of I^* . More formally,

▶ Lemma 4. (Pivot bound) For any interval $I_j \in \mathcal{I}$, let $I^* \in N(I_j)$ be any interval in $\mathcal{N}(I_j)$ and let $\mathcal{S}^* \subset \mathcal{N}(I_j)$ be the set of intervals in the neighborhood of I_j such that for all $I' \in \mathcal{S}^*$, $FF(I') > FF(I^*)$. Then, $FF(I_j) \leq FF(I^*) + |\mathcal{S}^*| + 1$.

Proof. Assume aiming toward contradiction that $FF(I_j) \ge 2 + FF(I^*) + |\mathcal{S}^*|$. Then, according to the pigeonhole principle, there must exist at least one color $FF(I^*) < c < FF(I_j)$ that is not assigned to any of the intervals in \mathcal{S}^* . Then $FF(I_j) = \min\{\mathbb{N} \setminus \{FF(I) \mid I \in N_j(I_j)\}\} \le c$, and it leads to a contradiction.

Note that the Neighborhood bound is a special case of the Pivot bound by selecting an empty set as its pivot. In this extreme case, the pivot is colored by 0 by FIRSTFIT, and $S^* \subseteq \mathcal{N}(I)$.

Intuitively, for a smaller upper bound of $FF(I_j)$, by the Pivot bound, we can choose a pivot I^* with a small color $FF(I^*)$ or choose a I^* with a small $|\mathcal{S}^*|$. However, we have a two-fold challenge. First, it is not trivial to find the best choice between minimizing $FF(I^*)$ or minimizing $|\mathcal{S}^*|$. Second, even when a good pivot I^* is given, it does not necessarily have non-trivial upper bounds of $FF(I^*)$ and $|\mathcal{S}^*|$. To cope with the difficulties, we construct a mechanism to balance the two choices of minimizing $FF(I^*)$ or $|\mathcal{S}^*|$ and different manners to bound $FF(I^*)$ and $|\mathcal{S}^*|$.

Optimal coloring and the image of walls of rocks and bricks. In the analysis of FIRSTFIT algorithm, we compare the solution of the FIRSTFIT algorithm to an *optimal offline algorithm* **OPT** that knows the complete input in advance.

We first imagine the intervals in \mathcal{I} as rocks (open intervals) and bricks (closed intervals) with fixed horizontal positions. That is, the rocks and bricks can be shifted vertically but not horizontally. We imagine that the optimal solution orients these rocks and bricks into a wall row by row using the fewest rows (without violating the fixed horizontal position constraints).



Figure 1 Rows denote the color assigned by OPT, where actual colors denote the color assigned by FIRSTFIT, where the order is based on the index.

More specifically, the intervals on the *i*-th row (from the bottom of the wall) are the intervals colored by *i* according to the optimal solution OPT. Since OPT uses precisely ω colors, the number of rows is exactly ω . Furthermore, it naturally follows that no two intervals drawn in the same row will intersect with each other, as otherwise OPT admits no proper coloring.

To avoid confusion between the color assigned by the optimal offline algorithm and the color assigned by the FIRSTFIT algorithm, from now on, we distinguish the color of an interval by FIRSTFIT or by OPT by its *color* or its *row*. More specifically, the color of an interval is the color assigned by FIRSTFIT algorithm, and the row refers to the color assigned by OPT. An example of the distinction between colors and rows is shown in Figure 1.

Relate the intervals and their structure in the wall. With the image of rock-andbrick wall in mind, given any interval (which may be a rock or a brick), we partition the rows according to their alignment with the interval *I*.

▶ **Definition 5.** For any interval $I \in \mathcal{I}$, let $\mathcal{R}_i(I)$ be the set of rows in the optimal solution $OPT(\mathcal{I})$ that contain *i* intervals intersecting *I*. And let $r_i(I)$ denote $|\mathcal{R}_i(I)|$.

Strictly speaking, the set $\mathcal{R}_{i}(I)$ contains rows instead of containing intervals. However, by slight abuse of notation, we say that interval $I' \in \mathcal{R}_{i}(I)$ if I' is in $\mathcal{N}(I)$ and colored by one of the colors in $R_{i}(I)$ by the optimal solution. In other words, $I' \in \mathcal{R}_{i}(I)$ if I' is an interval overlapping with I and colored by one of the colors in $\mathcal{R}_{i}(I)$ by OPT.

Since the intervals are either open unit intervals or closed-unit intervals, it is clear that not for all i the set $\mathcal{R}_i(I)$ is non-empty. Formally,

▶ Lemma 6. Consider any $I \in \mathcal{I}$.

- (a) $\mathcal{R}_{o}(I)$ contains at least the row where I is, and $r_{o}(I) \geq 1$.
- (b) If I is closed, $r_i(I) = 0$ for all $i \ge 4$.
- (c) If I is open, $r_i(I) = 0$ for all $i \ge 3$.
- (d) Since OPT uses ω rows, $r_0(I) + r_1(I) + r_2(I) + r_3(I) = \omega$.

Proof. (a) By definition, the rows $\mathcal{R}_i(I)$ correspond to the rows of an optimal solution OPT. Therefore, as OPT admits a proper coloring, there is no interval on the same row where I is contained and intersects I. Hence, on the row that contains I, no interval intersects I. Thus, this row is contained in $\mathcal{R}_0(I)$. It follows that $r_0(I) \geq 1$.

(b) Given a closed unit-length interval I = [a, a + 1], consider the following 3 points; $a, a + \frac{1}{2}$ and a + 1. Observe that the distance between these points is strictly smaller than the length of any (open or closed) unit-length interval, and that the endpoints of I are within these points. Then it follows that any interval that intersects interval I must intersect at least one of the points $a, a + \frac{1}{2}$, or a + 1. Thus since intervals which are drawn on the same row cannot intersect each other, at most 3 intervals per row can intersect interval I.

(c) Given an open unit-length interval I = (a, a + 1) and an infinitesimal number ε , consider the following 2 points; $a + \varepsilon$ and $a + 1 - \varepsilon$. Observe that the distance between these points is strictly smaller than the length of any (open or closed) unit-length interval, and that there exist no points between a and $a + \varepsilon$ and similarly there exist no points between a + 1 and $a + 1 - \varepsilon$. Then it follows that any interval that intersects interval I must intersect at least one of the points $a + \varepsilon$ or $a + 1 - \varepsilon$. Thus, since intervals that are drawn on the same row cannot intersect each other, at most 2 intervals per row can intersect interval I.

(d) By definition, the rows in $\mathcal{R}_i(I)$ correspond to the rows of an optimal solution OPT, of which there are ω such rows. We know from (b) that for all $i \geq 4$, $r_i(I) = 0$. Then, it follows that $r_0(I) + r_1(I) + r_2(I) + r_3(I) = \omega$. Note that this is for any interval I an exact equality because of the inclusion of $r_0(I)$.

By the definition of $\mathcal{R}_i(I)$, the number of intervals that intersect with I can be upper bounded in terms of $r_i(I)$. More specifically,

▶ Observation 7. For any interval $I \in \mathcal{I}$, $|\mathcal{N}_I(I)| \leq |\mathcal{N}(I)| \leq r_1(I) + 2 \cdot r_2(I) + 3 \cdot r_3(I)$.

Recall from Lemma 6 (b) and (c) that only closed intervals I can have non-empty $\mathcal{R}_3(I)$. Further, there is only a single way for a closed interval to intersect three intervals in the same row. Therefore, all rows in $\mathcal{R}_3(I)$ contain three sets of intervals, where the intervals in each set are twins. Formally, given any interval I = [r, r + 1] where r is some real number, we partition the intervals in $\mathcal{N}(I)$ into three sets:

- $= \mathcal{T}(I):$ Intervals $I' \equiv [r, r+1].$
- $\mathcal{A}(I)$: Intervals that are twins of intervals in $\mathcal{R}_3(I)$. More specifically, $\mathcal{A}(I)$ consists of the intervals $I' \equiv [r-1,r], I' \equiv (r,r+1)$, and $I' \equiv [r+1,r+2]$.
- $\blacksquare \mathcal{M}(I): \text{ All other intervals in } \mathcal{N}(I).$

Framework

Given that the optimal coloring uses ω colors on \mathcal{I} . For any interval $I \in \mathcal{I}$, we bound its color from FIRSTFIT by using the Neighborhood bound (Lemma 3) and the Pivot bound (Lemma 4). More specifically, we pick a pivot I^* with special structural property. Using the property, we bound the size of the corresponding \mathcal{S}^* . On the other hand, we bound the color $FF(I^*)$ using the Neighborhood bound. Finally, we use the Pivot bound to bound the color of I using the pivot I^* .

Practically, we first identify the "easy cases" where the FF(I) is at most 2ω as desired. Then, after peeling off these easy cases, we focus on the tough kernel of the analysis where there is no trivial choice for I^* such that the sum of $FF(I^*)$ and $|\mathcal{S}^*|$ is small. For this tough kernel of analysis, we identify two mediocre choices for I^* for which the sum of $FF(I^*)$ and $|\mathcal{S}^*|$ cannot both be large at the same time.

Intervals I with small $r_3(I)$. These intervals have relatively small neighborhoods. Formally, by the Neighborhood bound (Lemma 3) and Lemma 6 (d), we can show the following lemma.

▶ Lemma 8. For each interval $I \in \mathcal{I}$, if $r_3(I) < r_1(I) + 2$, then $FF(I) \leq 2\omega$.

Proof. By the Neighborhood bound (Lemma 3), $FF(I) \leq 1 + |\mathcal{N}_I(I)| \leq 1 + |\mathcal{N}(I)|$. By Lemma 6 (d), $FF(I) \leq 1 + r_1(I) + 2r_2(I) + 3r_3(I)$. The given condition $r_3(I) < r_1(I) + 2$ implies that $FF(I) \leq 2 + 2r_1(I) + 2r_2(I) + 2r_3(I)$. Since $r_0(I) \geq 1$, it follows that $r_1(I) + r_2(I) + r_3(I) \leq \omega - 1$. Thus, $FF(I) \leq 2\omega$.

Since any interval in $\mathcal{R}_3(I)$ must be in $\mathcal{A}(I)$, an empty $\mathcal{A}(I)$ implies $r_3(I) = 0$. Hence, we have a useful corollary:

▶ Corollary 9. If $\mathcal{A}(I) = \emptyset$, then $FF(I) \leq 2\omega$.

Open intervals.² By Lemma 6 (c), open intervals cannot have too many intersections per row. More specifically,

▶ Theorem 10. For each open interval $I \in \mathcal{I}$, $FF(I) \leq 2\omega - 1$.

Proof. By Lemma 6 (c), an open interval I intersects at most 2 intervals per row. Furthermore, since there are ω rows and $I \in \mathcal{R}_0(I)$, at most $\omega - 1$ rows remain to have any intervals intersecting I. It follows that $|\mathcal{N}_I(I)| \leq |\mathcal{N}(I)| \leq 2\omega - 2$. By the Neighborhood bound, $FF(I) \leq 2\omega - 1$.

Closed intervals I with large $r_3(I)$. Intuitively, because the intervals in $\mathcal{T}(I)$ have an identical neighborhood to interval I, these intervals are tricky to deal with. This leaves us to consider the intervals in the sets $\mathcal{A}(I)$ and $\mathcal{M}(I)$ as potential pivot intervals. In the ideal scenario, by picking as a pivot the interval with the largest color in the sets $\mathcal{A}(I)$ and $\mathcal{M}(I)$, $\mathcal{S}^* \subseteq \mathcal{T}(I)$. Using the Neighborhood bound, we bound the color of this largest interval. However, it is not always possible to pick such a pivot such that $\mathcal{S}^* \subseteq \mathcal{T}(I)$. In the case that $\mathcal{S}^* \not\subseteq \mathcal{T}(I)$, we look more closely at what additional intervals are part of this set. Finally, we apply the Pivot bound in order to prove an upper bound on the color of interval I.

In Sections 3 and 4, we will focus on the last case where I is a closed interval with large $r_3(I)$.

3 Warm-up: Regular case where endpoints are integral

As a starter, we consider a special case where all intervals have integral endpoints. More formally, for any interval $I \in \mathcal{I}$, I = (i, i + 1) for some integer *i* if *I* is open, and I = [i, i + 1]for some integer *i* if *I* is closed. In this section, we show that any $I \in \mathcal{I}$ has color $FF(I) \leq 2\omega$, where ω is the number of colors used by the optimal solution. Note that in the following, we focus on the closed intervals $I \in \mathcal{I}$ with $r_3(I) \geq r_1(I) + 2$ (by Lemma 10 and Lemma 8).

For such a set of integral-endpoints intervals, it is an important property that $\mathcal{M}(I) = \emptyset$ for any $I \in \mathcal{I}$. That is, the intervals in $\mathcal{N}(I)$ are either in $\mathcal{T}(I)$ (twins of I) or in $\mathcal{A}(I)$ (twins of the intervals in $\mathcal{R}_3(I)$). An example is shown in Figure 2.

Pick a pivot I^*

For any closed intervals $I \in \mathcal{I}$ with $r_3(I) \ge r_1(I) + 2$, we first identify the interval in $\mathcal{A}(I)$ with the highest color by FIRSTFIT.

▶ Definition 11. (Dominating interval $D^{\mathsf{A}}(I)$ in $\mathcal{A}(I)$) Let interval $D^{\mathsf{A}}(I) \in \mathcal{A}(I)$ be the highest colored interval. That is, there is no $I' \in \mathcal{A}(I)$ such that $FF(I') > FF(D^{\mathsf{A}}(I))$.

Note that $D^{\mathbb{A}}(I)$ must exist as $|\mathcal{A}(I)| \ge r_3(I) \ge r_1(I) + 2 \ge 2$.

In this analysis, we want to bound FF(I) by using $D^{\mathbb{A}}(I)$ as a pivot I^* and applying the Pivot bound (Lemma 4).

² This is a generalization from the results of Chrobak and Ślusarek [1] and Epstein and Levy [5], which showed for closed unit-length intervals, FIRSTFIT uses exactly $2\omega - 1$ colors.

Bound the size of \mathcal{S}^*

Given that we use $D^{\mathbb{A}}(I)$ as the pivot I^* , we bound the size of \mathcal{S}^* . Recall that \mathcal{S}^* is the set of intervals in $\mathcal{N}(I)$ which are assigned higher color than $FF(I^*)$. Since all intervals in \mathcal{I} have integral endpoints, any interval in $\mathcal{N}(I)$ is either in $\mathcal{A}(I)$ or in $\mathcal{T}(I)$. Therefore, by the definition of $D^{\mathbb{A}}(I)$, only intervals in $\mathcal{T}(I)$ are possible to contribute to \mathcal{S}^* . Moreover, since we focus on I that is closed, all intervals in $\mathcal{T}(I)$ must be in the rows in $\mathcal{R}_1(I)$. We use the following definition to represent the fraction of rows in $\mathcal{R}_1(I)$ that cannot contribute to \mathcal{S}^* .

▶ Definition 12. Given any $\hat{I} \in \mathcal{R}_1(I)$, $\alpha(\hat{I}) \in [0,1]$ denotes the fraction of rows $\mathcal{R} \in \mathcal{R}_1(I)$ where the interval I' in $\mathcal{R} \cap \mathcal{N}(I)$ satisfies 1) $I' \notin \mathcal{A}(I)$ and 2) $FF(I') \leq FF(\hat{I})$.

Note that we make Definition 12 more general for later usage. For the integral-endpoints case, it is sufficient to set \hat{I} as $D^{\mathbb{A}}(I)$, and $\alpha(D^{\mathbb{A}}(I)) \cdot r_1(I)$ is the number of intervals in $\mathcal{T}(I)$ that are not in \mathcal{S}^* .

► Lemma 13. If all intervals in \mathcal{I} have integral endpoints, by selecting $D^{\mathbb{A}}(I)$ as a pivot I^* , $|\mathcal{S}^*| = (1 - \alpha(D^{\mathbb{A}}(I))) \cdot r_1(I).$

Proof. First, by the definition of $\mathcal{R}_0(I)$, I does not intersect any interval in $\mathcal{R}_0(I)$. Thus, no interval in $\mathcal{R}_0(I)$ can be in \mathcal{S}^* . Next, as all intervals have integral endpoints, all intervals in $\mathcal{R}_2(I)$, $\mathcal{R}_3(I)$, or $\mathcal{R}_1(I) \setminus \mathcal{T}(I)$ are in the sets $\mathcal{A}(I)$. Thus, all these intervals are assigned a color below $FF(D^{\mathbb{A}}(I))$ and cannot contribute to \mathcal{S}^* . The remaining intervals are those in $\mathcal{T}(I)$. By definition, $\alpha(D^{\mathbb{A}}(I)) \cdot r_1(I)$ of these intervals have a color at most $FF(D^{\mathbb{A}}(I))$. Hence, the $(1 - \alpha(D^{\mathbb{A}}(I))) \cdot r_1(I)$ intervals are the only intervals intersecting I that have a color greater than $FF(D^{\mathbb{A}}(I))$. That is, $|\mathcal{S}^*| = (1 - \alpha(D^{\mathbb{A}}(I))) \cdot r_1(I)$.

Bound the color $FF(I^*)$

We bound the color of $D^{\mathbb{A}}(I)$ using the Neighborhood bound (Lemma 3). In general, apart from intervals in $\mathcal{T}(I)$, the number of intervals any interval in $\mathcal{A}(I)$ can intersect is shown in the following observation (also see Figure 3):

- ▶ Observation 14. Any interval in the set $\mathcal{A}(I)$ intersects
- (a) at most 2 intervals per row in $\mathcal{R}_{\mathcal{O}}(I)$,
- (b) at most 2 intervals which are not twins with I per row in $\mathcal{R}_1(I)$,
- (c) at most 2 intervals per row in $\mathcal{R}_2(I)$, and
- (d) exactly 1 interval per row in $\mathcal{R}_{\mathcal{J}}(I)$.

According to the definition of $D^{\mathbb{A}}(I)$, Observation 14 applies to $D^{\mathbb{A}}(I)$.

▶ Lemma 15. The color of interval
$$D^{A}(I)$$
, $FF(D^{A}(I)) \leq 2\omega + \alpha(D^{A}(I)) \cdot r_{1}(I) - r_{3}(I)$.

Proof. By the Neighborhood bound, $FF(D^{\mathbb{A}}(I)) \leq 1 + |N(D^{\mathbb{A}}(I))|$. By Observation 14, there are at most $2r_0(I) + 2r_1(I) + 2r_2(I) + r_3(I)$ intervals in $\mathcal{A}(I)$ that are in $\mathcal{N}(D^{\mathbb{A}}(I))$. Together with the $\alpha(D^{\mathbb{A}}(I)) \cdot r_1(I)$ intervals that "occupied" the smaller colors from the choice of $D^{\mathbb{A}}(I)$ at the moment when $D^{\mathbb{A}}(I)$ arrives, $|\mathcal{N}_{D^{\mathbb{A}}(I)}(D^{\mathbb{A}}(I))| \leq 2r_0(I) + 2r_1(I) + \alpha(D^{\mathbb{A}}(I)) \cdot r_1(I) + 2r_2(I) + r_3(I)$. Observe that by this bound, we count the row that contains interval $D^{\mathbb{A}}(I)$ itself. Since $D^{\mathbb{A}}(I)$ cannot intersect any interval on this row, we should subtract at least 1 from this bound. Thus,

$$\begin{aligned} \mathrm{FF}(D^{\mathbb{A}}(I)) &\leq 1 + 2r_{0}(I) + 2r_{1}(I) + \alpha(D^{\mathbb{A}}(I)) \cdot r_{1}(I) + 2r_{2}(I) + r_{3}(I) - 1 \\ &= 2\omega + \alpha(D^{\mathbb{A}}(I)) \cdot r_{1}(I) - r_{3}(I) \end{aligned}$$



Figure 2 All possible positions of intervals intersecting I per type of row. The intervals in $\mathcal{A}(I)$ are drawn in orange, green and blue, the intervals in $\mathcal{T}(I)$ in purple.

Figure 3 All possible intersections with interval $D^{\mathbb{A}}(I)$ per type of row. The intervals from $\mathcal{A}(I)$ that intersect with $D^{\mathbb{A}}(I)$ are drawn in orange, the intervals in $\mathcal{T}(I)$ in purple and the remaining intervals intersecting $D^{\mathbb{A}}(I)$ are drawn in yellow.

Note that due to the abstraction of $\alpha(D^{\mathbb{A}}(I))$, this proof does not rely on the property of integral endpoints. Therefore, Observation 14 also holds for the general case where the intervals in \mathcal{I} have arbitrary endpoints.

► Corollary 16. Given instance \mathcal{I} with arbitrary endpoints, the color of interval $D^{\mathbb{A}}(I)$ is at most $2\omega + \alpha(D^{\mathbb{A}}(I)) \cdot r_1(I) - r_3(I)$.

Proof of Theorem 1

Now, we are ready to prove Theorem 1 that for any \mathcal{I} with open or closed unit-length intervals that have integral endpoints, $FF(I) \leq 2\omega$ for all $I \in \mathcal{I}$, where ω is the optimal color needed for properly coloring \mathcal{I} .

Proof. By Lemma 8, we assume that $r_1(I) \leq r_3(I) - 2$. Moreover, by Lemma 10, we focus on closed I. According to Corollary 9, we assume that there exists a dominating interval $D^{\mathbb{A}}(I)$ in $\mathcal{A}(I)$ that has the highest color by FIRSTFIT. By Lemma 15, $FF(D^{\mathbb{A}}(I)) \leq 2\omega + \alpha(D^{\mathbb{A}}(I)) \cdot r_1(I) - r_3(I)$.

We now use the Pivot bound to bound the color of interval I from above, using $D^{\mathbb{A}}(I)$ as

the role of I^* . By Lemma 13, $|\mathcal{S}^*| = (1 - \alpha(D^{\mathbb{A}}(I))) \cdot r_1(I)$. It follows that

$$\begin{split} \mathrm{FF}(I) &\leq \mathrm{FF}(I^*) + |\mathcal{S}^*| + 1 \\ &= \mathrm{FF}(D^{\mathbb{A}}(I)) + (1 - \alpha(D^{\mathbb{A}}(I))) \cdot r_1(I) + 1 \\ &\leq 2\omega + \alpha(D^{\mathbb{A}}(I)) \cdot r_1(I) - r_3(I) + (1 - \alpha(D^{\mathbb{A}}(I))) \cdot r_1(I) + 1 \\ &\leq 2\omega + r_1(I) - r_3(I) + 1 \\ &\leq 2\omega - 1 \end{split}$$

◀

4 General case: Intervals with arbitrary endpoints

In this section, we consider the instance \mathcal{I} , where intervals have arbitrary endpoints. More specifically, by Lemma 8 and Lemma 10, we focus on $I \in \mathcal{I}$ that are closed and $r_3(I) \geq r_1(I)+2$. Recall from Section 3 that we partition $\mathcal{N}(I)$ into $\mathcal{T}(I)$, $\mathcal{A}(I)$, and $\mathcal{M}(I)$, that is, the set of twins, the set of intervals that are twins of intervals in $\mathcal{R}_3(I)$, and the rest of intervals with endpoints "misaligned" with the endpoints of I.

Applying the Pivot bound to bound the color FF(I) in this general case is more challenging than the integral-endpoints case. First, unlike in the integral-endpoints case where all intervals that intersect I must be either in $\mathcal{T}(I)$ or in $\mathcal{A}(I)$, the candidates in \mathcal{S}^* can also be in $\mathcal{M}(I)$. These intervals in $\mathcal{M}(I)$ can cross-interact with other intervals, making it difficult to bound the size of \mathcal{S}^* . Second, selecting a good pivot interval I^* in the general case is more challenging. We first show that using the same strategy as we used for the integral-endpoints case only guarantees a bound of $FF(I) \leq 3\omega$.

A naive attempt

As in the integral-endpoints case, let I^* be $D^{\mathbb{A}}(I)$, the interval with the highest color in $\mathcal{A}(I)$. In the general case, any interval in the sets $\mathcal{R}_1(I)$ and $\mathcal{R}_2(I)$ can be in $\mathcal{M}(I)$ and can therefore be a candidate of \mathcal{S}^* . If we now apply the same strategy as in the integral-endpoints setting, it is no longer the case that the set \mathcal{S}^* is a subset of $\mathcal{R}_1(I)$. Then, in each row $\mathcal{R} \in \mathcal{R}_2(I)$, it can be the case that two intervals (of which one intersects I and the other does not) are in the neighborhood of the pivot I^* . Furthermore, the remaining interval on row \mathcal{R} that also intersects with I can be considered for the set \mathcal{S}^* , resulting in us counting 3 intervals per row in $\mathcal{R}_2(I)$. Thus, the bound tends to 3ω when $r_1(I)$ and $r_3(I)$ are both small. To work around this, we need to closely look at the color of the intervals in $\mathcal{R}_2(I)$.

4.1 Roadmap

Primarily, we pick a pivot interval I^* and bound its color using the Neighborhood bound. The choice of pivot should provide a plausible way to bound the size of the corresponding S^* .

In the general case, we have another possible candidate of I^* . Symmetric to the dominating aligned interval $D^{\mathbb{A}}(I)$, we define interval $D^{\mathbb{M}}(I)$ as the dominating interval in rows in $\mathcal{R}_2(I)$ that is misaligned with I and assigned the highest color by FIRSTFIT. Formally,

▶ Definition 17. (Dominating interval $D^{\mathbb{M}}(I)$ in $\mathcal{R}_2(I) \cap \mathcal{M}(I)$) Let interval $D^{\mathbb{M}}(I)$ be the interval in $\mathcal{R}_2(I) \cap \mathcal{M}(I)$ with the highest color by FIRSTFIT. That is, for all $I' \in \mathcal{R}_2(I)$, $FF(I') \leq FF(D^{\mathbb{M}}(I))$.

A special property of the selection of $D^{\mathbb{M}}(I)$ is that since $D^{\mathbb{M}}(I) \in \mathcal{M}(I)$, any row in $\mathcal{R}_{0}(I), \mathcal{R}_{1}(I)$ or $\mathcal{R}_{3}(I)$, cannot be in $\mathcal{R}_{3}(D^{\mathbb{M}}(I))$ (Lemma ??). Therefore, all rows in $\mathcal{R}_{3}(D^{\mathbb{M}}(I))$ must also be in $\mathcal{R}_{2}(I)$.

In the general case, we pick one of $D^{\mathbb{A}}(I)$ and $D^{\mathbb{M}}(I)$ as the pivot and use the Pivot bound to bound the color FF(I). More specifically, we consider the following two cases.

(1) $\operatorname{FF}(D^{\mathbb{M}}(I)) \leq \operatorname{FF}(D^{\mathbb{A}}(I)).$

In this case, the misaligned intervals in $\mathcal{R}_2(I)$ are not in \mathcal{S}^* since they all have colors of at most $FF(D^{\mathbb{M}}(I)) \leq FF(D^{\mathbb{A}}(I))$. Using $D^{\mathbb{A}}(I)$ as the pivot and following a strategy similar to the case of integral-endpoints, we can show that $FF(I) \leq 2\omega$ (Lemma 21).

(2) $\operatorname{FF}(D^{\mathbb{M}}(I)) > \operatorname{FF}(D^{\mathbb{A}}(I)).$

In this case, applying the Pivot bound with $I^* = D^{\mathbb{A}}(I)$ leads to a bound of $FF(I) \leq 3\omega$ as shown in the naive attempt. To be precise, it is no longer true that $S^* \subseteq \mathcal{R}_1(I)$, and any interval in $\mathcal{R}_2(I)$ with a color greater than $FF(D^{\mathbb{A}}(I))$ is also a candidate of S^* .

An alternative strategy is to pick the interval $D^{\mathbb{M}}(I)$ as the pivot in this case. However, in the general case, the intervals in $\mathcal{R}_1(D^{\mathbb{M}}(I))$ are not particularly structured, More precisely, we do not know which intervals also intersect with I, and the neighborhood of $D^{\mathbb{M}}(I)$ can be huge. When $D^{\mathbb{M}}(I)$ has a large number of intersections, using the Neighborhood bound to bound the color $FF(D^{\mathbb{M}}(I))$ can lead to a high bound and further lead to a high bound of FF(I) by the Pivot bound. To deal with this situation, we first identify the rows containing intervals in $\mathcal{R}_1(D^{\mathbb{M}}(I))$ that we know their locations, namely, the intervals that are twins with $D^{\mathbb{M}}(I)$.

▶ Definition 18. Let the set of rows in $\mathcal{R}_2(I)$ with intervals identical to interval $D^{\mathbb{M}}(I)$ be

 $\mathcal{R}^{\text{TDM}}(I) = \{ row \ \mathcal{R} \in \mathcal{R}_2(I) \mid \mathcal{R} \text{ contains at least one interval that is in } \mathcal{T}(D^{\mathbb{M}}(I)) \}.$

Note that $\mathcal{R}^{\text{TDM}}(I) \subseteq \mathcal{R}_1(D^{\mathbb{M}}(I))$, and thus $r_3(D^{\mathbb{M}}(I)) \leq |\mathcal{R}^{\text{TDM}}(I)|$ implies that $r_3(D^{\mathbb{M}}(I)) \leq r_1(D^{\mathbb{M}}(I))$. On the contrary, $D^{\mathbb{M}}(I)$ potentially has a large neighborhood if $r_3(D^{\mathbb{M}}(I)) > |\mathcal{R}^{\text{TDM}}(I)|$. According to the size of $|\mathcal{R}^{\text{TDM}}(I)|$, we consider the following cases **2.a** and **2.b**.

(2.a) $r_3(D^{M}(I)) \leq |\mathcal{R}^{TDM}(I)|.$

In this case, we pick $D^{\mathbb{M}}(I)$ as the pivot I^* . By the selection of $D^{\mathbb{M}}(I)$ and $FF(D^{\mathbb{M}}(I)) > FF(D^{\mathbb{A}}(I))$, the only intervals that can contribute to \mathcal{S}^* are those in the set $\mathcal{R}_1(I)$. Furthermore, by $|\mathcal{R}^{\mathsf{TDM}}(I)| \geq r_3(D^{\mathbb{M}}(I))$, the neighborhood of $D^{\mathbb{M}}(I)$ is small, which enables us to use the Neighborhood bound to prove an upper bound the color of interval $D^{\mathbb{M}}(I)$ (Lemma 26). Then, by the Pivot bound (FF(I) is bounded by 2ω Lemma 28).

(2.b) $r_3(D^{M}(I)) > |\mathcal{R}^{\text{TDM}}(I)|.$

This is the most technical part of our analysis. In this case, $D^{\mathbb{M}}(I)$ may intersect more than 2 intervals per row on average and have a large neighborhood, and using the Neighborhood bound on $FF(D^{\mathbb{M}}(I))$ can lead to a large bound. We apply our framework recursively on $D^{\mathbb{M}}(I)$ by first showing that the intervals in $\mathcal{R}_3(D^{\mathbb{M}}(I))$ rows do not intersect many intervals when $r_3(D^{\mathbb{M}}(I))$ is large (Lemma 32). Specifically, we let $D^{\mathbb{A}\mathbb{M}}(I)$ be the highest colored interval in $\mathcal{R}_3(D^{\mathbb{M}}(I))$ that intersects with interval x.

▶ Definition 19. Let interval $D^{AM}(I) \in \mathcal{R}_{3}(D^{M}(I))$ be the dominating interval that has the highest color such that there exists no $I' \in \mathcal{R}_{3}(D^{M}(I))$ that intersects I such that $FF(I') > FF(D^{AM}(I))$.³

It is critical to use $D^{\mathbb{A}}(I)$, $D^{\mathbb{M}}(I)$, or $D^{\mathbb{A}\mathbb{M}}(I)$ as the pivot according to whether $FF(D^{\mathbb{A}\mathbb{M}}(I))$ is larger than $FF(D^{\mathbb{A}}(I))$.

(2.b.i) $FF(D^{AM}(I)) \ge FF(D^{A}(I)).$

In this case, we apply the Pivot bound by selecting $D^{\text{AM}}(I)$ as the pivot I^* . Recall that $D^{\text{A}}(I)$ is the dominating interval in $\mathcal{R}_3(I)$. The condition $\text{FF}(D^{\text{AM}}(I)) \geq \text{FF}(D^{\text{A}}(I))$ and the selection of $D^{\text{AM}}(I)$ ensure that no interval in $\mathcal{R}_3(I)$ can be in \mathcal{S}^* . Furthermore, by the definition of $D^{\text{AM}}(I)$, no interval in $\mathcal{R}_3(D^{\text{M}}(I))$ can contribute to \mathcal{S}^* . Therefore, \mathcal{S}^* is a subset of intervals in $\mathcal{R}_1(I)$ and $\mathcal{R}_2(I) \setminus \mathcal{R}_3(D^{\text{M}}(I))$.

The definition of $\mathcal{R}_3(D^{\mathbb{M}}(I))$ and the selection of $D^{\mathbb{A}\mathbb{M}}(I)$ guarantee that a row in the set $\mathcal{R}_3(D^{\mathbb{A}\mathbb{M}}(I))$ must contain an interval identical to the interval $D^{\mathbb{M}}(I)$ (see Figure 5). Thus, $\mathcal{R}_3(D^{\mathbb{A}\mathbb{M}}(I)) \subseteq \mathcal{R}^{\mathsf{TD}\mathbb{M}}(I)$, and $r_3(D^{\mathbb{A}\mathbb{M}}(I)) \leq |\mathcal{R}^{\mathsf{TD}\mathbb{M}}(I)|$. Moreover, since $\mathcal{T}(D^{\mathbb{A}\mathbb{M}}(I)) \subseteq \mathcal{R}_1(D^{\mathbb{A}\mathbb{M}}(I))$, and every $\mathcal{R}_3(D^{\mathbb{M}}(I))$ contains a twin of $D^{\mathbb{A}\mathbb{M}}(I)$, $r_1(D^{\mathbb{A}\mathbb{M}}(I)) \geq r_3(D^{\mathbb{M}}(I))$. Together with the condition $r_3(D^{\mathbb{M}}(I)) > |\mathcal{R}^{\mathsf{TD}\mathbb{M}}(I)|$, it guarantees that $r_1(D^{\mathbb{A}\mathbb{M}}(I)) \geq r_3(D^{\mathbb{M}}(I)) > |\mathcal{R}^{\mathsf{TD}\mathbb{M}}(I)| \geq r_3(D^{\mathbb{A}\mathbb{M}}(I))$. Therefore, $D^{\mathbb{A}\mathbb{M}}(I)$ intersects at most 2 intervals per row on average, and $\mathsf{FF}(D^{\mathbb{A}\mathbb{M}}(I))$ can be bounded by the Neighborhood bound effectively. With the bound $\mathcal{S}^* \subseteq \mathcal{R}_1(I) \cup (\mathcal{R}_2(I) \setminus \mathcal{R}_3(D^{\mathbb{M}}(I)))$, the upper bound $\mathsf{FF}(I) \leq 2\omega$ (Lemma 34).

(2.b.ii) $FF(D^{AM}(I)) < FF(D^{A}(I)).$

This case is the tough kernel of the analysis. In this case, considering using $D^{\mathbb{A}}(I)$, $D^{\mathbb{M}}(I)$, or $D^{\mathbb{A}\mathbb{M}}(I)$ as the pivot only is not sufficient to have an upper bound of FF(I) that is smaller than 3ω . Recall that this case is a subcase of case $FF(D^{\mathbb{A}}(I)) < FF(D^{\mathbb{M}}(I))$, picking $D^{\mathbb{A}}(I)$ as the pivot does not stop any interval in $\mathcal{R}_2(I)$ from contributing to \mathcal{S}^* . On the other hand, as $r_3(D^{\mathbb{M}}(I)) > |\mathcal{R}^{\mathbb{T}\mathbb{D}\mathbb{M}}(I)|$, $r_3(D^{\mathbb{M}}(I))$ may be much larger than $r_1(D^{\mathbb{M}}(I))$. Applying the Neighborhood bound yields a huge upper bound of $FF(I^*)$. Finally, condition $FF(D^{\mathbb{A}\mathbb{M}}(I)) < FF(D^{\mathbb{A}}(I))$ implies that when $D^{\mathbb{A}\mathbb{M}}(I)$ is selected as the pivot, \mathcal{S}^* could contain any interval in $\mathcal{R}_3(I)$ and become too large.

To deal with this tough case, we express the upper bounds obtained by using $D^{\mathbb{A}}(I)$ as the pivot and using $D^{\mathbb{M}}(I)$ as the pivot in terms of $|\mathcal{R}^{\mathsf{TDM}}(I)|$ and $r_2(D^{\mathbb{M}}(I))$ (Lemmas 26 and 36). These formulations show that the upper bounds from applying the Pivot bound using interval $D^{\mathbb{A}}(I)$ as I^* and using interval $D^{\mathbb{M}}(I)$ as I^* form a trade-off. More specifically, the upper bound using $D^{\mathbb{A}}(I)$ is maximized when $|\mathcal{R}^{\mathsf{TDM}}(I)| = \emptyset$, while upper bound using $D^{\mathbb{M}}(I)$ is maximized when $|\mathcal{R}^{\mathsf{TDM}}(I)| = \mathcal{R}_2(I)$. Then, we use that in any circumstance, the upper bound is at most $2\omega + \frac{|\mathcal{R}^{\mathsf{TDM}}(I)|}{2}$, which is strictly less than $2\omega + \frac{r_2(I)}{2} \cdot \omega$ (Definition 18). Since $r_2(I)$ can be upper bounded by $\frac{2}{3}\omega - 1$ (Corollary 39), the color assigned to interval Iis bounded by $\lceil \frac{7}{3}\omega \rceil - 2$ (Theorem 2).

³ Recall that we say an interval I' is in $\mathcal{R}_i(I)$ if it is in $\mathcal{N}(I)$ and is colored by one of the colors in $\mathcal{R}_i(I)$ by the optimal solution.

4.2 Analysis

Case (1): $FF(D^{A}(I)) \ge FF(D^{M}(I))$

We start by proving that in case (1), that is, when $FF(D^{\mathbb{A}}(I)) \geq FF(D^{\mathbb{M}}(I))$, the color of interval I can be bounded by 2ω .

Pick a pivot I^* . Given the condition where interval $D^{\mathbb{A}}(I)$ has a color larger or equal to interval $D^{\mathbb{M}}(I)$, picking $D^{\mathbb{A}}(I)$ as a pivot provides a smaller bound of $|\mathcal{S}^*|$. Therefore, we pick $D^{\mathbb{A}}(I)$ as the pivot I^* .

Bound the color of $FF(I^*)$. Next, we bound on the color of the pivot $FF(I^*)$. By Corollary 16, $FF(I^*) = FF(D^{\mathbb{A}}(I)) \leq 2\omega + \alpha(D^{\mathbb{A}}(I)) \cdot r_1(I) - r_3(I)$, where $\alpha(D^{\mathbb{A}}(I))$ is the fraction of rows in $\mathcal{R}_1(I)$ that contains intervals in $\mathcal{T}(I) \cup \mathcal{M}(I)$ and having color not higher than $D^{\mathbb{A}}(I)$.

Bound the size of S^* . Given that we use $D^{\mathbb{A}}(I)$ as pivot I^* , we bound the size of S^* . Recall that S^* is the set of intervals in $\mathcal{N}(I)$ which are assigned colors strictly higher than $FF(I^*)$. Since $FF(D^{\mathbb{A}}(I)) \geq FF(D^{\mathbb{M}}(I))$, only intervals in $\mathcal{T}(I)$ are possible to contribute to S^* .

▶ Lemma 20. When $FF(D^{\mathbb{A}}(I)) \ge FF(D^{\mathbb{M}}(I))$, by selecting $D^{\mathbb{A}}(I)$ as pivot I^* , $|\mathcal{S}^*| = (1 - \alpha(D^{\mathbb{A}}(I))) \cdot r_1(I)$.

Proof. First, by the definition of $\mathcal{R}_0(I)$, I does not intersect any interval in $\mathcal{R}_0(I)$. Thus, no interval in $\mathcal{R}_0(I)$ can be in \mathcal{S}^* . Next, the intervals in $\mathcal{R}_2(I)$ are either in $\mathcal{A}(I)$ or in $\mathcal{M}(I)$. Moreover, since $FF(D^{\mathbb{A}}(I)) \geq FF(D^{\mathbb{M}}(I))$, all intervals in $\mathcal{R}_2(I)$ have a color below $FF(D^{\mathbb{A}}(I))$ and cannot contribute to \mathcal{S}^* . Similarly, as all intervals in $\mathcal{R}_3(I)$ are in $\mathcal{A}(I)$, all intervals in $\mathcal{R}_3(I)$ have a color below $FF(D^{\mathbb{A}}(I))$ and cannot contribute to \mathcal{S}^* . The remaining intervals are the intervals in $\mathcal{R}_1(I)$. By definition, $\alpha(D^{\mathbb{A}}(I)) \cdot r_1(I)$ of these intervals have a color at most $FF(D^{\mathbb{A}}(I))$. Hence, the remaining $(1 - \alpha(D^{\mathbb{A}}(I))) \cdot r_1(I)$ are the only intervals intersecting I that have a color greater than $FF(D^{\mathbb{A}}(I))$. That is, $|\mathcal{S}^*| = (1 - \alpha(D^{\mathbb{A}}(I))) \cdot r_1(I)$.

Bound the color of FF(I). Now, we wrap up the analysis of the case where $FF(D^{\mathbb{A}}(I)) \geq FF(D^{\mathbb{M}}(I))$ using the bounds of $FF(I^*)$ and $|\mathcal{S}^*|$.

▶ Lemma 21. For any interval $I \in \mathcal{I}$, if $FF(D^{\mathbb{A}}(I)) \ge FF(D^{\mathbb{M}}(I))$, then $FF(I) \le 2\omega$.

Proof. By Lemma 8 and Lemma 10, $FF(I) \leq 2\omega$ if I is open or $r_3(I) < r_1(I) + 2$. Thus, in the following, we focus on closed intervals I with $r_1(I) \leq r_3(I) - 2$.

By taking $D^{\mathtt{A}}(I)$ as pivot I^* , according to Corollary 16, $\mathrm{FF}(D^{\mathtt{A}}(I)) \leq 2\omega + \alpha(D^{\mathtt{A}}(I)) \cdot r_1(I) - r_3(I)$. On the other hand, by Lemma 20, $|\mathcal{S}^*| = (1 - \alpha(D^{\mathtt{A}}(I))) \cdot r_1(I)$. Thus,

$$\begin{split} \operatorname{FF}(I) &\leq \operatorname{FF}(I^*) + |\mathcal{S}^*| + 1 \\ &\leq \operatorname{FF}(D^{\mathbb{A}}(I)) + (1 - \alpha(D^{\mathbb{A}}(I))) \cdot r_1(I) + 1 \\ &\leq 2\omega + r_1(I) - r_3(I) + 1 \\ &\leq 2\omega - 1 \end{split}$$

-



Figure 4 An exemplary non-exhaustive list of possible rows per type of row. Yellow intervals intersect only interval $D^{\mathbb{M}}(I)$, purple intervals intersect only I and orange intervals intersect both $D^{\mathbb{M}}(I)$ and I.

Case (2): $FF(D^{A}(I)) < FF(D^{M}(I))$

From now on, we focus on the case where $FF(D^{\mathbb{A}}(I)) < FF(D^{\mathbb{M}}(I))$.

Pick a pivot I^* . The condition where $FF(D^{\mathbb{A}}(I)) < FF(D^{\mathbb{M}}(I))$ guarantees that all intervals in the set $\mathcal{A}(I)$ have colors at most $FF(D^{\mathbb{M}}(I))$. Therefore, for the sake of the size of the \mathcal{S}^* , it is more efficient to pick the interval $D^{\mathbb{M}}(I)$ as a pivot than picking the interval $D^{\mathbb{A}}(I)$. Therefore, we pick interval $D^{\mathbb{M}}(I)$ as pivot I^* .

Recall from Definition 17 that $D^{\mathbb{M}}(I)$ is the interval with the highest color in the rows in $\mathcal{R}_2(I)$. The selection of $D^{\mathbb{M}}(I)$ attains a good structural property on $\mathcal{R}_3(D^{\mathbb{M}}(I))$ with regards to $\mathcal{R}_2(I)$:

▶ Lemma 22. $\mathcal{R}_{\mathcal{J}}(D^{M}(I)) \subseteq \mathcal{R}_{\mathcal{Z}}(I).$

Proof. By case distinction on $\mathcal{R}_i(I)$, we prove by contradiction that there is no intersection between $\mathcal{R}_3(D^{\mathbb{M}}(I)) \cap \mathcal{R}_i(I)$.

(1) Suppose on the contrary that $\mathcal{R}_3(D^{\mathbb{M}}(I)) \cap \mathcal{R}_0(I) \neq \emptyset$. If there exists a row with 3 intervals that intersect $D^{\mathbb{M}}(I)$, but none of them intersects I, then the intervals I and $D^{\mathbb{M}}(I)$ must not intersect. Thus, $D^{\mathbb{M}}(I) \notin \mathcal{M}(I)$, which is a contradiction by the definition of $D^{\mathbb{M}}(I)$.

(2) Suppose on the contrary that $\mathcal{R}_3(D^{\mathbb{M}}(I)) \cap \mathcal{R}_1(I) \neq \emptyset$. If there exists a row with 3 intervals that intersect $D^{\mathbb{M}}(I)$, but only one of those intervals intersects I. Then, either the interval $D^{\mathbb{M}}(I)$ is in the set $\mathcal{A}(I)$ or I and $D^{\mathbb{M}}(I)$ do not intersect. In either case, $D^{\mathbb{M}}(I) \notin \mathcal{M}(I)$, which is a contradiction by the definition of $D^{\mathbb{M}}(I)$.

(3) Suppose on the contrary that $\mathcal{R}_3(D^{\mathbb{M}}(I)) \cap \mathcal{R}_3(I) \neq \emptyset$. If there exists a row with 3 intervals that intersect both I and $D^{\mathbb{M}}(I)$, then interval $D^{\mathbb{M}}(I)$ must be in the set $\mathcal{T}(I)$. Then, $D^{\mathbb{M}}(I) \notin \mathcal{M}(I)$, which is a contradiction by the definition of $D^{\mathbb{M}}(I)$.

By (1) to (3), $\mathcal{R}_3(D^{\mathbb{M}}(I)) \cap (\mathcal{R}_0(I) \cup \mathcal{R}_1(I) \cup \mathcal{R}_3(I)) = \emptyset$. Therefore, it follows that $\mathcal{R}_3(D^{\mathbb{M}}(I)) \subseteq \mathcal{R}_2(I)$.

Recall from Definition 18 that the set $\mathcal{R}^{\text{TDM}}(I) \subset \mathcal{R}_2(I)$ contains all intervals in an $\mathcal{R}_2(I)$ row that are twins of interval $D^{\mathbb{M}}(I)$. Furthermore, recall from Definition 19 that interval $D^{\mathbb{A}\mathbb{M}}(I)$ is the highest colored interval in $\mathcal{R}_3(D^{\mathbb{M}}(I))$ that intersects with I. Observe that the intersection of the sets $\mathcal{R}^{\text{TDM}}(I)$ and $\mathcal{R}_3(D^{\mathbb{M}}(I))$ must be empty, as a row cannot have both 3 intervals intersecting $D^{\mathbb{M}}(I)$ and an interval identical to $D^{\mathbb{M}}(I)$, and therefore $|\mathcal{R}^{\text{TDM}}(I)| + r_3(D^{\mathbb{M}}(I)) \leq r_2(I)$.

Bound the color of $FF(I^*)$. We first provide observations on the maximum number of intersections intervals in $\mathcal{M}(I)$ and specifically $D^{\mathbb{M}}(I)$ have in each type of row (also see Figure 4).

▶ Observation 23. Any interval in the set $\mathcal{M}(I)$ intersects

- (a) at most 1 interval, other than I, per row in $\mathcal{R}_{\mathcal{O}}(I)$,
- (b) at most 1 interval, which does not intersect interval I, per row in $\mathcal{R}_1(I)$, and
- (c) exactly 2 intervals per row in $\mathcal{R}_{\mathcal{S}}(I)$.

▶ Observation 24. Interval $D^{M}(I)$ intersects

- (a) exactly 3 intervals per row in $\mathcal{R}_{\mathcal{J}}(D^{\mathbb{M}}(I))$,
- (b) exactly 1 interval per row in $\mathcal{R}^{\text{TDM}}(I)$, and
- (c) at most 2 intervals per row in $\mathcal{R}_2(I) \setminus (\mathcal{R}^{TDM}(I) \cup \mathcal{R}_3(D^M(I))).$

Furthermore, we define two variables with respect to the sets $\mathcal{R}^{\text{TDM}}(I)$ and $\mathcal{R}_3(D^{\mathbb{M}}(I))$. Similarly to $\alpha(i)$ being a fraction of the $\mathcal{R}_1(I)$ rows, we define β and γ as fractions of the set $\mathcal{R}_2(I)$ with respect to interval $D^{\mathbb{M}}(I)$.

▶ Definition 25. Given $D^{M}(I) \in \mathcal{R}_{2}(I)$,

- (a) $\beta \in [0,1]$ denotes the fraction of rows $\mathcal{R} \in \mathcal{R}_2(I)$ where the interval I' in $\mathcal{R} \cap \mathcal{N}(I)$ is in $D^{\mathbb{A}\mathbb{M}}(I)$, and
- (b) $\gamma \in [0,1]$ denotes the fraction of rows $\mathcal{R} \in \mathcal{R}_2(I)$ where the interval I' in $\mathcal{R} \cap \mathcal{N}(I)$ is in $\mathcal{R}_3(D^{M}(I))$.

Now we are equipped to compute an upper bound on the color of interval $D^{\mathbb{M}}(I)$.

▶ Lemma 26. The color of interval $D^{\mathbb{M}}(I)$ is at most $c(D^{\mathbb{M}}(I)) \leq \omega + \alpha(D^{\mathbb{M}}(I)) \cdot r_1(I) - \beta \cdot r_2(I) + \gamma \cdot r_2(I) + r_2(I) + r_3(I)$.

Proof. Using the Neighborhood bound and Observations 23 and 24, we can describe the maximum color of interval $D^{\mathbb{M}}(I)$ as 1 plus the number of intersections per type of row. Observe that by doing this, we count the row that contains interval $D^{\mathbb{M}}(I)$ itself, and since $D^{\mathbb{M}}(I)$ cannot intersect any interval in this row, we may subtract at least 1. Then it follows that

$$\begin{split} \operatorname{FF}(D^{\mathtt{M}}(I)) &\leq r_{\mathtt{0}}(I) + (1 + \alpha(D^{\mathtt{M}}(I))) \cdot r_{\mathtt{1}}(I) + \beta \cdot r_{\mathtt{2}}(I) + \gamma \cdot 3r_{\mathtt{2}}(I) \\ &+ (1 - \beta - \gamma) \cdot 2r_{\mathtt{2}}(I) + 2r_{\mathtt{3}}(I) + 1 - 1 \\ &= \omega + \alpha(D^{\mathtt{M}}(I)) \cdot r_{\mathtt{1}}(I) - \beta \cdot r_{\mathtt{2}}(I) + \gamma \cdot r_{\mathtt{2}}(I) + r_{\mathtt{2}}(I) + r_{\mathtt{3}}(I) \end{split}$$

Next, we make a case distinction based on the relative sizes of the sets $\mathcal{R}^{\text{TDM}}(I)$ and $\mathcal{R}_3(D^{\mathbb{M}}(I))$.

Case (2.a): $|\mathcal{R}^{\text{TDM}}(I)| \ge r_3(D^{M}(I))$

We start with the case where the number of intervals in the set $\mathcal{R}^{\text{TDM}}(I)$ is at least as large as the number of intervals in the set $\mathcal{R}_3(D^{\mathbb{M}}(I))$.

Bound the size of S^* .

▶ Lemma 27. When $FF(D^{M}(I)) > FF(D^{A}(I))$, by selecting $D^{M}(I)$ as pivot I^{*} , $|\mathcal{S}^{*}| = (1 - \alpha(D^{M}(I))) \cdot r_{1}(I)$.

Proof. First, by the definition of $\mathcal{R}_0(I)$, I does not intersect any interval in $\mathcal{R}_0(I)$. Thus, no interval in $\mathcal{R}_0(I)$ can be in \mathcal{S}^* . Next, all intervals in $\mathcal{R}_2(I)$ are either in $\mathcal{A}(I)$ or $\mathcal{M}(I)$. Since $FF(D^{\mathbb{M}}(I)) > FF(D^{\mathbb{A}}(I))$, it follows that all intervals in $\mathcal{R}_2(I)$ have a color below $FF(D^{\mathbb{M}}(I))$ and cannot contribute to \mathcal{S}^* . Similarly, as all intervals in $\mathcal{R}_3(I)$ are in $\mathcal{A}(I)$, all intervals in $\mathcal{R}_3(I)$ have a color below $FF(D^{\mathbb{M}}(I))$ and cannot contribute to \mathcal{S}^* . The remaining intervals are those in $\mathcal{R}_1(I)$. By definition, $\alpha(I) \cdot r_1(I)$ of these intervals have a color at most $FF(D^{\mathbb{M}}(I))$. Hence, the remaining $(1 - \alpha(D^{\mathbb{M}}(I))) \cdot r_1(I)$ are the only intervals intersecting I that have a color greater than $FF(D^{\mathbb{M}}(I))$. That is, $|\mathcal{S}^*| = (1 - \alpha(D^{\mathbb{M}}(I))) \cdot r_1(I)$.

Bound the color of FF(I).

▶ Lemma 28. If $|\mathcal{R}^{TDM}(I)| \ge r_{\mathcal{S}}(D^{M}(I))$, then $FF(I) \le 2\omega$.

Proof. Assume that $FF(D^{\mathbb{M}}(I)) > FF(D^{\mathbb{A}}(I))$ and interval I is a closed interval, as otherwise by Lemma 21 and Lemma 10 it follows that $FF(I) \leq 2\omega$. From the definition of β and γ it follows that since $|\mathcal{R}^{\mathsf{TDM}}(I)| \geq r_3(D^{\mathbb{M}}(I))$, also $\beta \geq \gamma$.

Now we can compute a bound on the color of interval I, using the Pivot bound where we take interval $D^{\mathbb{M}}(I)$ as I^* . By Lemma 26, the color of interval $D^{\mathbb{M}}(I)$ is bounded by $FF(D^{\mathbb{M}}(I)) \leq \omega + \alpha(D^{\mathbb{M}}(I)) \cdot r_1(I) - \beta \cdot r_2(I) + \gamma \cdot r_2(I) + r_2(I) + r_3(I)$. And, by Lemma 27, the size of \mathcal{S}^* equals $(1 - \alpha(D^{\mathbb{M}}(I))) \cdot r_1(I)$. Then it follows that,

$$\begin{split} \mathrm{FF}(I) &\leq \mathrm{FF}(I^*) + |\mathcal{S}^*| + 1 \\ &= \mathrm{FF}(D^{\mathrm{M}}(I)) + (1 - \alpha(D^{\mathrm{M}}(I))) \cdot r_1(I) + 1 \\ &\leq \omega + r_1(I) - \beta \cdot r_2(I) + \gamma \cdot r_2(I) + r_2(I) + r_3(I) + 1 \\ &\leq \omega + r_1(I) + r_2(I) + r_3(I) + 1 \\ &\leq 2\omega \end{split}$$

_	
_	

Case (2.b): $|\mathcal{R}^{\text{TDM}}(I)| < r_3(D^{M}(I))$

We continue with the case where the number of intervals in the set $\mathcal{R}^{\text{TDM}}(I)$ is strictly smaller than the number of intervals in the set $\mathcal{R}_3(D^{\mathbb{M}}(I))$. This relation between $\mathcal{R}^{\text{TDM}}(I)$ and $\mathcal{R}_3(D^{\mathbb{M}}(I))$ indicates that the average number of intersections interval $D^{\mathbb{M}}(I)$ has per row can be greater than 2. Then, it might be interesting to explore other options as a pivot. For this we make another case distinction on the relation between the colors of interval $D^{\mathbb{A}}(I)$ and interval $D^{\mathbb{A}}(I)$.

Case (2.b.i): $FF(D^{AM}(I)) \ge FF(D^{A}(I))$

First, let us explore the case where the color of interval $D^{\mathbb{A}M}(I)$ is at least as large as the color of interval $D^{\mathbb{A}}(I)$.

Pick a pivot I^* The condition that $FF(D^{AM}(I)) \ge FF(D^A(I))$ guarantees that all intervals in the set $\mathcal{A}(I)$ have colors at most $FF(D^{AM}(I))$. Therefore, for the sake of the size of the size of the set \mathcal{S}^* , it is more efficient to pick interval $D^{AM}(I)$ as a pivot compared to picking interval $D^A(I)$. Therefore, we pick interval $D^{AM}(I)$ as pivot I^* .

Bound the color $FF(I^*)$

As we now want to compute an upper bound on the color of interval $D^{\text{AM}}(I)$, it is of importance to know where interval $D^{\text{AM}}(I)$ might be located.



Figure 5 An exemplary non-exhaustive list of possible rows per type of $\mathcal{R}_2(I)$. Yellow intervals intersect only interval $D^{AM}(I)$ (resp. $D^{A}(I)$), purple intervals intersect only I and orange intervals intersect both $D^{AM}(I)$ (resp. $D^{A}(I)$) and I.

▶ Lemma 29. For any interval $I' \in \mathcal{R}_{3}(D^{M}(I)), I' \notin \mathcal{A}(I)$

Proof. Assume aiming towards a contradiction that $I' \in \mathcal{A}(I)$.

(1) I' is a closed interval. Assume without loss of generality that interval I' intersects interval I on the left-hand side of interval I. Since $I' \in \mathcal{R}_3(D^{\mathbb{M}}(I))$, and I' is a closed interval, it follows that there are precisely two positions where interval $D^{\mathbb{M}}(I)$ could lie. The first possible position is to the left of interval I'. In this case, interval $D^{\mathbb{M}}(I)$ and interval I cannot intersect. Then it follows that $D^{\mathbb{M}}(I) \notin \mathcal{M}(I)$, which is a contradiction by the definition of $D^{\mathbb{M}}(I)$. The other possible position is to the right of interval I'. In this case, interval $D^{\mathbb{M}}(I)$ must be identical to interval I, i.e., $D^{\mathbb{M}}(I) \in \mathcal{T}(I)$. Then it follows that $D^{\mathbb{M}}(I) \notin \mathcal{M}(I)$, which is a contradiction by the definition of $D^{\mathbb{M}}(I)$.

(2) I' is an open interval. Since $I' \in \mathcal{A}(I)$ and I' is an open interval, it follows that $I' \in \mathcal{T}(I)$. Similarly, since $I' \in \mathcal{R}_3(D^{\mathbb{M}}(I))$, and I' is an open interval, it follows that $I' \in \mathcal{T}(D^{\mathbb{M}}(I))$. An interval can only be in both $\mathcal{T}(I)$ and $\mathcal{T}(D^{\mathbb{M}}(I))$ if interval I and interval $D^{\mathbb{M}}(I)$ are identical, i.e., $D^{\mathbb{M}}(I) \in \mathcal{T}(I)$. Then it follows that $D^{\mathbb{M}}(I) \notin \mathcal{M}(I)$, which is a contradiction by the definition of $D^{\mathbb{M}}(I)$.

Note that Observation 23 remains true for interval $D^{\text{AM}}(I)$. Now, let us make an additional observation regarding the intersections interval $D^{\text{AM}}(I)$ has with intervals in $\mathcal{R}_2(I)$ rows (also see Figure 5):

▶ Observation 30. Interval $D^{\text{AM}}(I)$ intersects

- (a) exactly 1 interval per row contained in $\mathcal{R}_{\mathcal{J}}(D^{\mathbb{M}}(I))$,
- (b) exactly 3 intervals per row contained in $\mathcal{R}^{\text{TDM}}(I)$, and
- (c) at most 2 intervals per row contained in $\mathcal{R}_2(I) \setminus (\mathcal{R}^{TDM}(I) \cup \mathcal{R}_3(D^M(I)))$.

Furthermore, it is important to specify how many of the intervals in $\mathcal{R}_2(I) \setminus \mathcal{R}_3(D^{\mathbb{M}}(I))$ are assigned a color larger than $FF(D^{\mathbb{A}\mathbb{M}}(I))$ or smaller than $FF(D^{\mathbb{A}\mathbb{M}}(I))$. This was previously not necessary, when we considered interval $D^{\mathbb{M}}(I)$, which is, by definition, the interval in $\mathcal{R}_2(I)$ assigned the largest color. Analogously, we do not need to consider the intervals in $\mathcal{R}_3(D^{\mathbb{M}}(I))$ as per definition interval $D^{\mathbb{A}\mathbb{M}}(I)$ is the largest colored interval in this set.

▶ Definition 31. Given any interval \hat{I} , $\delta(\hat{I}) \in [0,1]$ denotes the fraction of rows $\mathcal{R} \in \mathcal{R}_2(I) \setminus \mathcal{R}_3(D^{\mathsf{M}}(I))$ where interval I' in $\mathcal{R} \cap \mathcal{N}(I) \cap \mathcal{N}(D^{\mathsf{M}}(I))$ satisfies $FF(I') \leq FF(\hat{I})$.

Now we are equipped to compute a bound on the color of interval $D^{\text{AM}}(I)$.

▶ Lemma 32. The color of interval $D^{AM}(I)$, $FF(D^{AM}(I)) \le r_0(I) + (1 + \alpha(D^{AM}(I))) \cdot r_1(I) + \delta(D^{AM}(I)) \cdot (1 - \gamma) \cdot 2r_2(I) + \beta \cdot r_2(I) + \gamma \cdot r_2(I) + 2r_3(I)$.

Proof. By the Neighborhood bound $\operatorname{FF}(D^{\operatorname{AM}}(I)) \leq 1 + |\mathcal{N}(D^{\operatorname{AM}}(I))|$. By Observation 23, there are at most $r_0(I) + 2r_1(I) + 2r_3(I)$ intervals in $\mathcal{R}_0(I), \mathcal{R}_1(I)$ and $\mathcal{R}_3(I)$ that are in $\mathcal{N}(D^{\operatorname{AM}}(I))$. However, by Definition 12, $(1 - \alpha(D^{\operatorname{AM}}(I)))$ of those intervals have a color larger than $\operatorname{FF}(D^{\operatorname{AM}}(I))$ and hence we do not consider them for this bound. By Observation 30, there are at most $r_3(D^{\operatorname{M}}(I)) + 3 \cdot |\mathcal{R}^{\operatorname{TDM}}(I)| + 2 \cdot |\mathcal{R}_2(I) \setminus (\mathcal{R}^{\operatorname{TDM}}(I) \cup \mathcal{R}_3(D^{\operatorname{M}}(I)))| = \gamma \cdot r_2(I) + \beta \cdot 3r_2(I) + (1 - \beta - \gamma) \cdot 2r_2(I)$ intervals in $\mathcal{R}_2(I)$ that are in $\mathcal{N}(D^{\operatorname{AM}}(I))$. However, by Definition 31, $(1 - \delta(D^{\operatorname{AM}}(I))) \cdot (\beta \cdot 2r_2(I) + (1 - \beta - \gamma) \cdot 2r_2(I))$ of those intervals have a color larger than $\operatorname{FF}(D^{\operatorname{AM}}(I))$ and hence we do not consider them for this bound. Observe that by this bound, we count the row that contains interval $D^{\operatorname{AM}}(I)$ itself. Since $D^{\operatorname{AM}}(I)$ cannot intersect any interval on this row, we should subtract at least 1 from this bound. Thus,

$$\begin{split} \mathrm{FF}(D^{\mathrm{AM}}(I)) &\leq r_0(I) + 2r_1(I) - (1 - \alpha(D^{\mathrm{AM}}(I))) \cdot r_1(I) + 2r_3(I) \\ &+ \beta \cdot 3r_2(I) + \gamma \cdot r_2(I) + (1 - \beta - \gamma) \cdot 2r_2(I) \\ &- (1 - \delta(D^{\mathrm{AM}}(I))) \cdot (\beta \cdot 2r_2(I) + (1 - \beta - \gamma) \cdot 2r_2(I)) - 1 + 1 \\ &= r_0(I) + (1 + \alpha(D^{\mathrm{AM}}(I))) \cdot r_1(I) + \delta(D^{\mathrm{AM}}(I)) \cdot (1 - \gamma) \cdot 2r_2(I) + \beta \cdot r_2(I) \\ &+ \gamma \cdot r_2(I) + 2r_3(I) \end{split}$$

-

Bound the size of S^* .

► Lemma 33. By selecting $D^{\text{AM}}(I)$ as pivot I^* , $|\mathcal{S}^*| = (1 - \alpha(D^{\text{AM}}(I))) \cdot r_1(I) + (1 - \delta(D^{\text{AM}}(I))) \cdot (1 - \gamma) \cdot 2r_2(I)$.

Proof. First, by the definition of $\mathcal{R}_0(I)$, I does not intersect any interval in $\mathcal{R}_0(I)$. Thus, no interval in $\mathcal{R}_0(I)$ can be in \mathcal{S}^* . Next, all intervals in $\mathcal{R}_3(I)$ are in the set $\mathcal{A}(I)$. Thus, all these intervals are assigned a color below $FF(D^{AM}(I))$ and cannot contribute to \mathcal{S}^* . Of the intervals in $\mathcal{R}_1(I)$, by definition of α , only $(1 - \alpha(D^{AM}(I))) \cdot r_1(I)$ intervals have a color larger than $FF(D^{AM}(I))$, and can contribute to set \mathcal{S}^* . Of the $\mathcal{R}_2(I)$ rows, out of the intervals intersecting I that are in the rows that are also in the set $\mathcal{R}_3(D^M(I))$, none have a color larger than $FF(D^{AM}(I))$ by the definition of $D^{AM}(I)$. Then, of the remaining $(1 - \gamma) \cdot r_2(I)$ $\mathcal{R}_2(I)$ rows, by the definition of δ , only $(1 - \delta(D^{AM}(I))) \cdot (1 - \beta - \gamma) \cdot 2r_2(I)$ intervals have a color larger than $FF(D^{AM}(I))$ and can contribute to the set \mathcal{S}^* . Then it follows that $|\mathcal{S}^*| = (1 - \alpha(D^{AM}(I))) \cdot r_1(I) + (1 - \delta(D^{AM}(I))) \cdot (1 - \gamma) \cdot 2r_2(I)$

Bound the color FF(I).

Now we can use this result to prove an upper bound on the color of interval I, when the color of interval $D^{\mathbb{A}\mathbb{M}}(I)$ is at least as large as the color of interval $D^{\mathbb{A}}(I)$.

▶ Lemma 34. If $FF(D^{AM}(I)) \ge FF(D^{A}(I))$, then $FF(I) \le 2\omega$

Proof. By Lemma 28, we assume that $|\mathcal{R}^{\text{TDM}}(I)| < r_3(D^{\text{M}}(I))$, and thus $\beta < \gamma$. By Lemma 32, $FF(D^{\text{AM}}(I)) \leq r_0(I) + (1 + \alpha(D^{\text{AM}}(I))) \cdot r_1(I) + \delta(D^{\text{AM}}(I)) \cdot (1 - \gamma) \cdot 2r_2(I) + \beta \cdot r_2(I) + \gamma \cdot r_2(I) + 2r_3(I)$. By Lemma 33, $|\mathcal{S}^*| = (1 - \alpha(D^{\text{AM}}(I))) \cdot r_1(I) + (1 - \delta(D^{\text{AM}}(I))) \cdot (1 - \gamma) \cdot 2r_2(I)$. It follows that,

$$\begin{split} \mathrm{FF}(I) &\leq \mathrm{FF}(I^*) + |\mathcal{S}^*| + 1 \\ &\leq \mathrm{FF}(D^{\mathrm{AM}}(I)) + (1 - \alpha(D^{\mathrm{AM}}(I))) \cdot r_1(I) + (1 - \delta(D^{\mathrm{AM}}(I))) \cdot (1 - \gamma) \cdot 2r_2(I) + 1 \\ &\leq r_0(I) + 2r_1(I) + \beta \cdot r_2(I) + \gamma \cdot r_2(I) + (1 - \gamma) \cdot 2r_2(I) + 2r_3(I) + 1 \\ &< r_0(I) + 2r_1(I) + \beta \cdot r_2(I) + (1 - \beta) \cdot r_2(I) + r_2(I) + 2r_3(I) + 1 \\ &\leq 2\omega \end{split}$$

◀

Case (2.b.ii): $FF(D^{A}(I)) > FF(D^{AM}(I))$

In order to prove Theorem 2 we still need to consider the tough kernel of the analysis. That is, the case where $FF(D^{AM}(I)) < FF(D^A(I)) < FF(D^M(I))$ and $\beta < \gamma$. For this case we abandon our usual mechanism slightly, where instead of focusing on a single pivot I^* and a single set S^* , we explore the option where two distinct pairs of a pivot and a set cannot both admit a large solution.

Pick the pivots I^* . Although we have reasoned earlier that picking interval $D^{\mathbb{A}}(I)$ and interval $D^{\mathbb{M}}(I)$ as the pivot is not sufficient to have an upper bound on FF(I) that is smaller than 3ω , we can deal with this tough case by expressing the bound obtained by using $D^{\mathbb{A}}(I)$ or $D^{\mathbb{M}}(I)$ as the pivot in terms of $|\mathcal{R}^{\text{TDM}}(I)|$.

Bound the color of I^* Let us first take a closer look at the color of interval $D^{\mathbb{A}}(I)$. Although Lemma 15 is technically still correct, knowing that $FF(D^{\mathbb{A}}(I)) < FF(D^{\mathbb{A}}(I))$, we can be slightly more precise about which intersections are assigned a color greater than $FF(D^{\mathbb{A}}(I))$. In order to be more precise, let us first make additional observations regarding the intersections of interval $D^{\mathbb{A}}(I)$ with interval in $\mathcal{R}_2(I)$ rows (also see Figure 5).

▶ Observation 35. Interval $D^{A}(I)$ intersects

- (a) at most 1 interval that is not in $\mathcal{N}(I)$ per row contained in $\mathcal{R}_2(I)$, and
- (b) at most 1 interval that is in $\mathcal{N}(I)$ per row contained in $\mathcal{R}_3(D^{\mathbb{M}}(I))$,
- (c) at most 1 intervals that is in $\mathcal{N}(I)$ per row contained in $\mathcal{R}^{\text{TDM}}(I)$, and
- (d) at most 1 intervals that is in $\mathcal{N}(I)$ per row contained in $\mathcal{R}_2(I) \setminus (\mathcal{R}^{TDM}(I) \cup \mathcal{R}_3(D^M(I)))$.

▶ Lemma 36. The color of interval $D^{A}(I)$, $FF(D^{A}(I)) \leq 2r_{0}(I) + 2r_{1}(I) + \alpha(D^{A}(I)) \cdot r_{1}(I) + \gamma \cdot r_{2}(I) + r_{2}(I) + \delta(D^{A}(I)) \cdot (1 - \gamma) \cdot r_{2}(I) + r_{3}(I)$.

Proof. By the Neighborhood bound, $FF(D^{\mathbb{A}}(I)) \leq 1 + |\mathcal{N}(D^{\mathbb{A}}(I))|$. By Observation 14, there are at most $2r_0(I) + 2r_1(I) + r_3(I)$ intervals in $\mathcal{R}_0(I), \mathcal{R}_1(I)$ and $\mathcal{R}_3(I)$ that are in $\mathcal{N}(D^{\mathbb{A}}(I))$. Together with the at most $\alpha(D^{\mathbb{A}}(I)) \cdot r_1(I)$ intervals that were assigned a color below $FF(D^{\mathbb{A}}(I))$ which are either in $\mathcal{T}(I)$ or do not intersect $D^{\mathbb{A}}(I)$ and are in $\mathcal{M}(I)$.

By Observation 35, there are at most $r_2(I) + \gamma \cdot r_2(I) + \beta \cdot r_2(I) + (1 - \gamma - \beta) \cdot r_2(I)$ intervals that are in $\mathcal{R}_2(I)$ and in $\mathcal{N}(D^{\mathbb{A}}(I))$. By our assumption that $FF(D^{\mathbb{A}\mathbb{M}}(I)) < FF(D^{\mathbb{A}}(I))$, the $\gamma \cdot r_2(I)$ intervals in the set $\mathcal{R}_3(D^{\mathbb{M}}(I)) \cap \mathcal{N}(I)$, must have a color smaller than $FF(D^{\mathbb{A}}(I))$. For the remaining $(1 - \gamma) \cdot r_2(I)$ intervals that are in $\mathcal{N}(I)$, by the definition of δ , $(1 - \delta(D^{\mathbb{A}}(I))) \cdot (1 - \gamma) \cdot r_2(I)$ are assigned a color larger than $FF(D^{\mathbb{M}}(I))$ and hence we do not consider them for this bound.

Observe that by this bound we count the row that contains interval $D^{\mathbb{A}}(I)$ itself. Since interval $D^{\mathbb{A}}(I)$ cannot intersect any interval on this row, we should subtract at least 1 from this bound. Thus,

$$\begin{split} \operatorname{FF}(D^{\mathtt{A}}(I)) &\leq |\mathcal{N}(D^{\mathtt{A}}(I))| + 1 \\ &\leq 2r_{\mathtt{0}}(I) + 2r_{\mathtt{1}}(I) + \alpha(D^{\mathtt{A}}(I)) \cdot r_{\mathtt{1}}(I) + \gamma \cdot r_{\mathtt{2}}(I) \\ &+ r_{\mathtt{2}}(I) + \delta(D^{\mathtt{A}}(I)) \cdot (1 - \gamma) \cdot r_{\mathtt{2}}(I) + r_{\mathtt{3}}(I) \end{split}$$

◀

As for the bound on the color of interval $D^{\mathbb{M}}(I)$, Lemma 26 still suffices.

Bound the sizes of \mathcal{S}^* . Given that we use $D^{\mathbb{A}}(I)$ as pivot I^* , we bound the size of \mathcal{S}^* .

▶ Lemma 37. When $FF(D^{A}(I)) \geq FF(D^{AM}(I))$, by selecting $D^{A}(I)$ as pivot I^{*} , $|\mathcal{S}^{*}| \leq (1 - \alpha(D^{A}(I))) \cdot r_{1}(I) + (1 - \delta(D^{M}(I))) \cdot (1 - \gamma) \cdot r_{2}(I) + (1 - \gamma) \cdot r_{2}(I)$

Proof. First, by the definition of $\mathcal{R}_0(I)$, I does not intersect any interval in $\mathcal{R}_0(I)$. Thus, no interval in $\mathcal{R}_0(I)$ can be in \mathcal{S}^* . By the definition of α , only $(1 - \alpha(D^{\mathsf{A}}(I))) \cdot r_1(I)$ intervals in $\mathcal{R}_1(I)$ are assigned a color strictly larger than $\operatorname{FF}(D^{\mathsf{A}}(I))$. Thus the only intervals in an $\mathcal{R}_1(I)$ that are eligible for \mathcal{S}^* are those $(1 - \alpha(D^{\mathsf{A}}(I))) \cdot r_1(I)$ intervals. Since $\operatorname{FF}(D^{\mathsf{A}}(I)) > \operatorname{FF}(D^{\mathsf{AM}}(I))$, no interval in an $\mathcal{R}_3(D^{\mathsf{M}}(I))$ row is assigned a color larger than $\operatorname{FF}(D^{\mathsf{A}}(I))$. Thus, none of the $\gamma \cdot 2r_2(I)$ intervals in $\mathcal{R}_3(D^{\mathsf{M}}(I)) \cap \mathcal{N}(I)$ can contribute to \mathcal{S}^* . By the definition of δ , only $(1 - \delta(D^{\mathsf{A}}(I))) \cdot (1 - \gamma) \cdot r_2(I)$ intervals both in $\mathcal{N}(D^{\mathsf{A}}(I))$ and $\mathcal{R}_2(I)$ are assigned a color strictly larger than $\operatorname{FF}(D^{\mathsf{A}}(I))$. Furthermore, all $(1 - \gamma) \cdot r_2(I)$ intervals which are in $\mathcal{R}_2(I)$ but not in $\mathcal{N}(D^{\mathsf{A}}(I))$ and not in $\mathcal{R}_3(D^{\mathsf{M}}(I))$ could potentially be assigned a color larger than $\operatorname{FF}(D^{\mathsf{A}}(I))$ and therefore contribute to \mathcal{S}^* . Finally, as all intervals in $\mathcal{R}_3(I)$ are in $\mathcal{A}(I)$, all intervals in $\mathcal{R}_3(I)$ have a color below $\operatorname{FF}(D^{\mathsf{A}}(I))$ and cannot contribute to \mathcal{S}^* . Thus, $|\mathcal{S}^*| \leq (1 - \alpha(D^{\mathsf{A}}(I))) \cdot r_1(I) + (1 - \delta(D^{\mathsf{M}}(I))) \cdot (1 - \gamma) \cdot r_2(I) + (1 - \gamma) \cdot r_2(I)$.

As for the set \mathcal{S}^* when we pick interval $D^{\mathbb{M}}(I)$ as pivot, Lemma 27 still suffices.

Bound the size of FF(I).

As the bound we are going to prove in the proof of Theorem 2 is based on the size of set $\mathcal{R}^{\text{TDM}}(I)$, which in turn is strictly smaller than the size of set $\mathcal{R}_2(I)$, an upper bound on $r_2(I)$ improves the bound on the color of I we would otherwise obtain. This upper bound on the number of rows in $\mathcal{R}_2(I)$ is obtained by a lower bound on the number of rows in $\mathcal{R}_3(I)$.

▶ Lemma 38. For any interval $I \in \mathcal{I}$, if $r_3(I) < \frac{1}{3}\omega$, then $FF(I) \leq \frac{7}{3}\omega - 2$.

Proof. We show this using the Neighborhood bound. Observe that $r_0(I) \ge 1$, since interval I does not intersect any intervals on its own row. Furthermore, note that the Neighborhood bound is maximized when we maximize the number of intervals that intersect interval I. That is, we maximize the bound when we maximize $r_3(I)$. Then, $r_3(I) = \frac{1}{3}\omega - 1$. For the remaining rows, we maximize the bound by maximizing $r_2(I)$, and hence $r_2(I) = \frac{2}{3}\omega$. Then, $r_1(I) = 0$, and we get the following upper bound on the color of interval I,

$$\begin{aligned} \mathrm{FF}(I) &\leq r_1(I) + 2r_2(I) + 3r_3(I) + 1 \\ &\leq 0 + 2 \cdot \frac{2}{3}\omega + 3 \cdot (\frac{1}{3}\omega - 1) + 1 \\ &= \frac{7}{3}\omega - 2 \end{aligned}$$

Now we have shown that $r_3(I) \ge \frac{1}{3}\omega$, it simply follows that $r_2(I) < \frac{2}{3}\omega$.

<

▶ Corollary 39. For any interval $I \in \mathcal{I}$, if $r_2(I) \geq \frac{2}{3}\omega$, then $FF(I) \leq \frac{7}{3}\omega - 2$.

We have now shown everything we need to show in order to compute a bound on the color of interval I.

Proof of Theorem 2.

For any interval $I \in \mathcal{I}$, the color of I is at most $FF(I) \leq \left\lceil \frac{7}{3}\omega \right\rceil - 2$.

Proof. By Lemma 10, 8, 21, 28 and 34, $FF(I) \leq 2\omega$ if I is open, $r_1(I) \leq r_3(I) - 2$, $FF(D^{\mathbb{A}}(I)) < FF(D^{\mathbb{M}}(I))$, $\beta < \gamma$ or $FF(D^{\mathbb{A}\mathbb{M}}(I)) < FF(D^{\mathbb{A}}(I))$. Thus in the following we focus on closed intervals with $r_1(I) \leq r_3(I) - 2$, $FF(D^{\mathbb{A}\mathbb{M}}(I)) < FF(D^{\mathbb{A}}(I)) < FF(D^{\mathbb{M}}(I))$ and $\beta < \gamma$.

Now we can compute the first bound on the color of interval I, using the Pivot bound with interval $D^{\mathsf{A}}(I)$ as pivot I^* . According to Lemma 36, $\operatorname{FF}(D^{\mathsf{A}}(I)) \leq 2r_0(I) + 2r_1(I) + \alpha(D^{\mathsf{A}}(I)) \cdot r_1(I) + \gamma \cdot r_2(I) + r_2(I) + \delta(D^{\mathsf{M}}(I)) \cdot (1 - \gamma) \cdot r_2(I) + r_3(I)$ and, by taking $D^{\mathsf{A}}(I)$ as pivot I^* , Lemma 37, $|\mathcal{S}^*| \leq (1 - \alpha(D^{\mathsf{A}}(I))) \cdot r_1(I) + (1 - \delta(D^{\mathsf{M}}(I))) \cdot (1 - \gamma) \cdot r_2(I) + (1 - \gamma) \cdot r_2(I)$. Thus,

$$\begin{split} \mathrm{FF}(I) &\leq \mathrm{FF}(I^*) + |\mathcal{S}^*| + 1 \\ &\leq \mathrm{FF}(D^{\mathsf{A}}(I)) + (1 - \alpha(D^{\mathsf{A}}(I))) \cdot r_1(I) + (1 - \delta(D^{\mathsf{M}}(I))) \cdot (1 - \gamma) \cdot r_2(I) \\ &\quad + (1 - \gamma) \cdot r_2(I) + 1 \\ &\leq 2r_0(I) + 3r_1(I) + 2r_2(I) + (1 - \gamma) \cdot r_2(I) + r_3(I) + 1 \\ &\leq 2r_0(I) + 2r_1(I) + (1 - \gamma) \cdot r_2(I) + 2r_2(I) + 2r_3(I) - 1 \\ &= 2\omega + (1 - \gamma) \cdot r_2(I) - 1 \end{split}$$

Next we move to the other bound. By taking $D^{\mathbb{M}}(I)$ as pivot I^* , according to Lemma 26, $FF(D^{\mathbb{M}}(I)) \leq \omega + \alpha(D^{\mathbb{M}}(I)) \cdot r_1(I) - \beta \cdot r_2(I) + \gamma \cdot r_2(I) + r_2(I) + r_3(I)$, and Lemma 27, $|\mathcal{S}^*| = (1 - \alpha(D^{\mathbb{M}}(I))) \cdot r_1(I)$. Then, it follows that,

$$\begin{split} \mathrm{FF}(I) &\leq \mathrm{FF}(I^*) + |\mathcal{S}^*| + 1 \\ &\leq \mathrm{FF}(D^{\mathsf{M}}(I)) + (1 - \alpha(D^{\mathsf{M}}(I))) \cdot r_1(I) + 1 \\ &\leq \omega + r_1(I) - \beta \cdot r_2(I) + \gamma \cdot r_2(I) + r_2(I) + r_3(I) + 1 \\ &\leq 2\omega + \gamma \cdot r_2(I) - 1 \end{split}$$

We have proven that simultaneously, the color of interval I is at most $FF(I) \leq 2\omega + \gamma \cdot r_2(I) - 1$ and at most $FF(I) \leq 2\omega + (1 - \gamma) \cdot r_2(I) - 1$. Then, it follows that for any value of γ the color of interval I is at most $FF(I) \leq 2\omega + \frac{1}{2} \cdot r_2(I) - 1$ Then, it follows from Corollary 39 that the number of $\mathcal{R}_2(I)$ rows must be less than $r_2(I) < \frac{2}{3}\omega$, as otherwise the color of interval I is bounded by $FF(I) \leq \frac{7}{3}\omega - 2$. Thus, the color of interval I is at most $FF(I) < 2\omega + \frac{1}{3} \cdot \omega - 1 = \frac{7}{3}\omega - 1$. Which, by the integrality of FF(I), is at most $FF(I) \leq [\frac{7}{3}\omega] - 2$

5 Conclusion

In this work, we develop a sophisticated counting method based on the Pivot bound and show that FIRSTFIT uses at most 2ω colors in the case that all open and closed unit-length intervals have integral endpoints, which matches the lower bound. We also show that FIRSTFIT uses

In an attempt to improve the current upper bound for the number of colors used by the FIRSTFIT algorithm for the general case, it may be of interest to investigate the number of colors used by the FIRSTFIT algorithm for the so-called bounded length intervals as introduced by Chybowska-Sokól et al. [2]. The technique and the results on open/closed unit-length intervals developed in this work serve as a first step towards investigating FIRSTFIT's performance via gradually relaxing the lengths bound.

— References

- Marek Chrobak and Maciej Ślusarek. On Some Packing Problem Related to Dynamic Storage Allocation. *RAIRO Theor. Informatics Appl.*, 22(4):487-499, 1988. URL: https: //doi.org/10.1051/ita/1988220404871, doi:10.1051/ITA/1988220404871.
- 2 Joanna Chybowska-Sokół, Grzegorz Gutowski, Konstanty Junosza-Szaniawski, Patryk Mikos, and Adam Polak. Online coloring of short intervals. *Eur. J. Comb.*, 118:103919, 2024. URL: https://doi.org/10.1016/j.ejc.2024.103919, doi:10.1016/J.EJC.2024.103919.
- 3 Israel R. Curbelo. The on-line Width of Various Classes of Posets. PhD thesis, University of Louisville, 2022. doi:10.18297/etd/3974.
- 4 Israel R. Curbelo. Online coloring of short interval graphs and two-count interval graphs. CoRR, abs/2412.17193, 2024. URL: https://doi.org/10.48550/arXiv.2412.17193, arXiv: 2412.17193, doi:10.48550/ARXIV.2412.17193.
- 5 Leah Epstein and Meital Levy. Online Interval Coloring and Variants. In Luís Caires, Giuseppe F. Italiano, Luís Monteiro, Catuscia Palamidessi, and Moti Yung, editors, Automata, Languages and Programming, 32nd International Colloquium, ICALP 2005, Lisbon, Portugal, July 11-15, 2005, Proceedings, volume 3580 of Lecture Notes in Computer Science, pages 602-613. Springer, 2005. doi:10.1007/11523468_49.
- 6 Martin Charles Golumbic. Algorithmic graph theory and perfect graphs. Elsevier, 2004.
- 7 Hal A. Kierstead. The linearity of first-fit coloring of interval graphs. SIAM J. Discret. Math., 1(4):526-530, 1988. doi:10.1137/0401048.
- 8 Hal A. Kierstead and Jun Qin. Coloring interval graphs with first-fit. Discret. Math., 144(1-3):47-57, 1995. doi:10.1016/0012-365X(94)00285-Q.
- 9 Hal A. Kierstead, David A. Smith, and William T. Trotter. First-fit coloring on interval graphs has performance ratio at least 5. Eur. J. Comb., 51:236-254, 2016. URL: https://doi.org/10.1016/j.ejc.2015.05.015, doi:10.1016/J.EJC.2015.05.015.
- 10 Hal A. Kierstead and William T. Trotter. An extremal problem in recursive combinatorics. Congressus Numerantium, 33:143–153, 1981.
- 11 N. S. Narayanaswamy and R. Subhash Babu. A Note on First-Fit Coloring of Interval Graphs. Order, 25(1):49-53, 2008. URL: https://doi.org/10.1007/s11083-008-9076-6, doi:10.1007/S11083-008-9076-6.
- 12 Sriram V. Pemmaraju, Rajiv Raman, and Kasturi R. Varadarajan. Max-coloring and online coloring with bandwidths on interval graphs. ACM Trans. Algorithms, 7(3):35:1–35:21, 2011. doi:10.1145/1978782.1978790.
- 13 Maciej Ślusarek. A Coloring Algorithm for Interval Graphs. In Antoni Kreczmar and Grazyna Mirkowska, editors, Mathematical Foundations of Computer Science 1989, MFCS'89, Porabka-Kozubnik, Poland, August 28 September 1, 1989, Proceedings, volume 379 of Lecture Notes in Computer Science, pages 471–480. Springer, 1989. doi:10.1007/3-540-51486-4_93.
- 14 Hans S. Witsenhausen. On woodall's interval problem. J. Comb. Theory A, 21(2):222–229, 1976. doi:10.1016/0097-3165(76)90066-2.