MEDIAL AXIS IN PSEUDO-EUCLIDEAN SPACES

ADAM BIAŁOŻYT

ABSTRACT. In the paper we will investigate the notion of medial axis for psuedo-Euclidean spaces. For most of the article we follow the path of Birbrair and Denkowski [1] checking its feasibility in the new context.

1. Preliminaries

Definition 1.1. We call the pair (\mathbb{R}^n, Q) , where Q is a nondegenerated quadratic form, *a pseudo-Euclidean space*.

For any pseudo-Euclidean space (\mathbb{R}^n, Q) there exists exactly one symmetric matrix $A \in M(n)$ such that

$$Q(x) = x^T A^T x.$$

For such a matrix and any two vectors $x, y \in \mathbb{R}^n$ we introduce their pseudo-scalar product $\langle \langle x, y \rangle \rangle := x^T A y$. It is easy to check that the pseudoscalar product is symmetrical, bilinear and nongenerated (meaning $\exists x \forall y \langle \langle x, y \rangle \rangle = 0 \Rightarrow x = 0$). From the general theory of quadratic forms it is known that we can fix a orthonormal basis of \mathbb{R}^n in such a way that $A = diag(\lambda_1, \ldots, \lambda_n), \lambda_i \neq 0, \lambda_i \geq \lambda_{i+1}$. Then $l := \max\{i \mid \lambda_i > 0\}$ is called the index of the pseudo-Eculidean space, and a pair (l, p) := (l, n - l) is called its signature. In place of \mathbb{R}^n it customed to write in such a setting $\mathbb{R}_{(l,p)}$. This notation makes sense as all $\mathbb{R}_{(l,p)}$ are lineary isometric. In this spirit it will not affect the generalicity of our study to assume that all λ_i have absolute value equal one. In such setting we denote

$$||x||_l = (\sum_{i=1}^l x_i^2)^{1/2}$$
 and $||x||_p = (\sum_{i=l+1}^p x_i^2)^{1/2}$.

For any $x \in \mathbb{R}_{(l,p)}$ the value Q(x) is a generalisation of the vector x norm squared. Since we do not assume Q to be positive definite, Q(x) can admit any real value. Henceforth we distinguish in the pseudo-Eulidean spaces the vectors that are

- space like Q(x) > 0
- light like Q(x) < 0

²⁰¹⁰ Mathematics Subject Classification. 32B20, 53A07, 54F45.

Key words and phrases. Medial Axis, Skeleton, Radius of Curvature, Dimension, Tangent Cone, O-minimal Geometry.

• isotropic Q(x) = 0

Analogously the two dimensional subspace V of $\mathbb{R}_{(l,p)}$ is called

- Euclidean if Q|V is definite (either positive or negative).
- pseudo-Euclidean if Q|V is non-singular indefininite.
- isotropic in the remaining cases.

Since the quadratic form Q is not convex the triangle inequality does not hold in general. However for any two $x, y \in \mathbb{R}_{(l,p)}$ there is

$$|Q(x+y)| \leq |Q(x)| + |Q(y)|$$
, if x, y span an Euclidean subspace of $\mathbb{R}_{(l,p)}$

$$|Q(x+y)| \geq |Q(x)| + |Q(y)|$$
, if x, y span a pseudo-Euclidean subspace of $\mathbb{R}_{(l,p)}$

For a real number r and $x \in \mathbb{R}_{(l,p)}$ we can define also the equivalents of open and closed balls

$$B(x,r) := \{ y \in \mathbb{R}_{(l,p)} \mid Q(x-y) < r \}, \ \overline{B}(x,r) := \{ y \in \mathbb{R}_{(l,p)} \mid Q(x-y) \le r \}.$$

Mind that we will use the sets B(x, r) and $\overline{B}(x, r)$ only as a tool in analysis of the mutual position of points in pseudo-Euclidean spaces. When it comes to the typically topological and metric properties like boundedness or compactness we will still mean the properties inherited from the natural topology of \mathbb{R}^n Mind also that for the simplicity of the definition of B(x, r) we take r to be equal the usual radius squared.

2. Medial Axis

For the whole article we will assume that $X \subset \mathbb{R}_{(l,p)}$ is closed in the Euclidean topology. Then for a point $a \in \mathbb{R}_{(l,p)}$ we define the squared distance function

$$\rho(a, X) := \inf\{Q(x-a) \mid x \in X\}$$

and a multifunction of the closest points

$$m(a) = \{ x \in X \mid Q(x - a) = \rho(a, X) \}.$$

Note here that m has also another equivalent and extremely useful definition

$$m(a) = \{ x \in X \mid \forall p \in X, Q(a-x) \le Q(a-p) \}.$$

Contrary to the Euclidean case, in general the function ρ is not bounded from below, and not all points *a* bring nonempty or compact m(a). Indeed for the set $X = \{x_2 = \sqrt{x_1^2 + 1}\} \subset \mathbb{R}_{(1,1)}$ we have m(0) = X and $m(a) = \emptyset$ for any other *a* on x_1 -axis.

Therefore besides the usual notion of the medial axis

$$M_X := \{ a \in \mathbb{R}_{(l,p)} \mid \#m(a) > 1 \}$$

a new potentially interesting set appears – the vacant axis of X

$$W_X := \{ a \in \mathbb{R}_{(l,p)} \mid m(a) = \emptyset \}.$$

Moreover, it is not all certain that the multifunction m seen as a relation is reflexive for a general set X (consider points of $X = 0 \times [0, 1]$)

in $\mathbb{R}_{(1,1)}$). To enforce reflexivity, we will assume X to be $acausal^1$ meaning for any two distinguished points $x, y \in X$ we assume to have Q(x-y) > 0. It is plain to see that in such a case $\pi_l : \mathbb{R}_{(l,p)} \to \mathbb{R}^l$, the natural projection on the first l coordinates, must be injective on X. If we consider now $\pi_p : \mathbb{R}_{(l,p)} \to \mathbb{R}^p$ to be the natural projection on the last p coordinates, and the composition $\pi_p \circ (\pi_l | X)^{-1} : \pi_l(X) \to \mathbb{R}^p$, it is easy to check that $\pi_p \circ (\pi_l | X)^{-1}$ is 1–Lipschitz for acausal X. Based on that we will call X to be L-pseudo-Lipschitz if $\pi_p \circ (\pi_l | X)^{-1}$ is L-Lipschitz. It translates to the condition

- $L||x y||_l \ge ||x y||_p, \text{ for all } x, y \in X.$ 3. MULTIFUNCTIONS *m* AND \mathcal{N}

Before delving any deeper into the analysis of medial axis in pseudo-Euclidean spaces, let us justify the choice of restriction we imposed on sets of our interest.

Theorem 3.1. For any point x of an acausal set X, $m(x) = \{x\}$ and the intersection $M_X \cap X$ is empty.

Proof. Taking $x \in X$ we obviously have Q(x - x) = 0, whereas $Q(x-y) = ||x-y||_l^2 - ||x-y||_p^2 = ||\pi_l(x) - \pi_l(y)||^2 - ||\pi_p(x) - \pi_p(y)||^2 > 0$ for any point $y \in X$ distinct from x. Hence $m(x) = \{x\}$ and x is not a point of M_X .

Thanks to the Theorem 3.1, we can define *the normal sets* of an acausal set just like in the Euclidean case.

$$\mathcal{N}(a) := \{ x \in \mathbb{R}_{(l,p)} \mid a \in m(x) \} = \{ x \in \mathbb{R}_{(l,p)} \mid Q(x-a) = \rho(x,X) \}$$

$$\mathcal{N}'(a) := \{ x \in \mathbb{R}_{(l,p)} \mid m(a) = \{a\} \}.$$

Proposition 3.2. For any *a* in an acausal definable subset *X* of $\mathbb{R}_{(l,p)}$ there is

(1) $a \in \mathcal{N}'(a) \subset \mathcal{N}(a)$, (2) $\mathcal{N}(a)$ is close convex and definable (3) $\mathcal{N}(a) - a \subset \mathcal{N}_a X := \{w \in \mathbb{R}_{(l,p)} \mid \forall v \in C_a X : \langle \langle w, v \rangle \rangle \leq 0 \}.$ (4) $x \in \mathcal{N}'(a) \Rightarrow [a, x] \subset \mathcal{N}'(a), x \in \mathcal{N}(a) \setminus \{a\} \Rightarrow [a, x) \subset \mathcal{N}'(a)$ (5) $\limsup_{X \in b \to a} \mathcal{N}(b) \subset \mathcal{N}(a)$ (6) $\mathcal{N}'(a)$ is convex and definable.

Proof. (1) $a \in \mathcal{N}'(a)$ is given by the previous theorem, the inclusion is trivial.

(2) The set is definable due to the description

$$\mathcal{N}(a) = \{ x \in \mathbb{R}_{(l,p)} \mid \forall b \in X : \ Q(x-a) \le Q(x-b) \}$$

¹The name comes from the Lorentzian geometry

since Q and X are definable. If we move the quantifier in front of the set braces we obtain

$$\mathcal{N}(a) = \bigcap_{b \in X} \{ x \in \mathbb{R}_{(l,p)} \mid Q(x-a) \le Q(x-b) \}$$

which gives closedness and convexity of $\mathcal{N}(a)$ as for any $b \in X$ the set $\{x \in \mathbb{R}_{(l,p)} \mid Q(x-a) \leq Q(x-b)\}$ is a closed halfspace.

(3) For any $b \in X$, the set $\mathcal{N}(a)$ is a subset of

$$\{x \in \mathbb{R}_{(l,p)} \mid Q(x-a) \leq Q(x-b)\} = \{x+a \mid Q(x+a-a) \leq Q(x+a-b)\} =$$
$$= \{x \mid Q(x) \leq Q(x+(a-b))\} + a = \{x \mid 0 \leq 2\langle\langle x, a-b \rangle\rangle + Q(a-b)\} + a.$$
Now, taking $v \in C_a X$ and sequences $b_{\nu} \to a, l_{\nu} \to 0^+$ such that $l_{\nu}(b_{\nu}-a) \to v$ we have

$$\mathcal{N}(a) \subset \{x \mid 2\langle\langle x, l_{\nu}(b_{\nu} - a)\rangle\rangle \leq \langle\langle a - b_{\nu}, l_{\nu}(b_{\nu} - a)\rangle\rangle\} + a$$

and by passing to the limit we obtain

$$\mathcal{N}(a) \subset \{x \mid 2\langle\langle x, v\rangle\rangle \leq 0\} + a, \,\forall v \in C_a X$$

(4) For any triple of points $a, b \in X, x \in \mathcal{N}(a)$ let us define a polynomial

$$R(t) := Q(tx + (1 - t)a - b) - Q(tx + (1 - t)a - a).$$

After simplification, R reduces to $R(t) = 2t\langle\langle x - a, a - b\rangle\rangle + Q(a - b)$, thus it is a degree 1 polynomial and R(0) > 0, $R(1) \ge 0$. Therefore, R|(0,1) is positive and consequently $m(tx+(1-t)a) = \{a\}$ thus $[a, x) \subset \mathcal{N}'(a)$.

(5) Take any sequence $X \ni b_{\nu} \to a$ and $x_{\nu} \in \mathcal{N}(b_{\nu}) \to x$. Then $b_{\nu} \in m(x_{\nu})$ thus $\forall b \in X : Q(b_{\nu} - x_{\nu}) \leq Q(b - x_{\nu})$. After passing to the limit we obtain $Q(a - x) \leq Q(b - x)$ for all $b \in X$ thus $a \in m(b)$. (6) Follows the same reasoning as (2).

Proposition 3.3. For any point $a \in X$, there is $\mathcal{N}(a) = \overline{\mathcal{N}'(a)}$

Proof. $\mathcal{N}(a)$ is closed and $\mathcal{N}'(a) \subset \mathcal{N}(a)$ thus $\overline{\mathcal{N}'(a)} \subset \mathcal{N}(a)$. On the other hand for any $x \in \mathcal{N}(a) \setminus \{a\}$ there is $[a, x) \subset \mathcal{N}'(a)$ thus $x \in \overline{\mathcal{N}'(a)}$.

Theorem 3.4. For any closed X

$$M_X = \bigcup \mathcal{N}(a) \setminus \mathcal{N}'(a) = \bigcup \mathcal{N}(a) \setminus \bigcup \mathcal{N}'(a) = \mathbb{R}_{(l,p)} \setminus \bigcup \mathcal{N}'(a) \cup W_X.$$

Proof. The proof is basically the same as in [1]

Recall that, we call X to be L-pseudo-Lipschitz if $\pi_p \circ (\pi_l | X)^{-1}$ (where π_l and π_p are natural projections on the first l and last p coordinates respectively) is L-Lipschitz. We have the following result concerning the multifunction m. **Theorem 3.5.** Let X be a closed L-pseudo-Lipschitz definable subset of $\mathbb{R}_{(l,p)}$ with L < 1, then $W_X = \emptyset$. Moreover, the multifunction m has compact values, is locally bounded and upper-semicontinuous, meaning $\limsup_{D \ni x \to a} m(x) = m(a)$ for any dense subset D.

Proof. Since X is L-pseudo-Lipschitz, for any $b \in X$ there is

$$X \subset B := \{ x \in \mathbb{R}_{(l,p)} \mid L \| b - x \|_l \ge \| b - x \|_p \}.$$

In such a case for $a \in \mathbb{R}_{(l,p)}$ and any sequence $X \ni x_{\nu} \to \infty$ the values of $Q(x_{\nu} - a)$ diverge to infinity, since the cones at infinity of $\{x \in \mathbb{R}_{(l,p)} \mid Q(x-a) \leq const.\}$ and B are disjoint. It means that $\{x \in X \mid Q(x-a) \leq \rho(a) + \varepsilon\}$ is bounded, and therefore compact and from any sequence realising the infimum $\rho(a)$ we can pick a subsequence convergent in X. Since Q is continuous the limiting point in X realises the value $\rho(a)$. Moreover,

$$m(a) = \bigcap_{\varepsilon > 0} \{ x \in X \mid Q(x - a) \le \rho(a) + \varepsilon \}$$

thus m(a) is compact, as it is closed and bounded.

Regarding local boundedness of m, let us star by remarking that ρ is upppersemi-continuous, as it is an infimum of a collection of continuous functions. It means that for any $\varepsilon > 0$ we can find U-a neighbourhood of a such that for all $\tilde{a} \in U$ there is $\rho(\tilde{a}) < \rho(a) + \varepsilon$. Taking a union of $\{x \in X \mid Q(x - \tilde{a}) \leq \rho(a) + \varepsilon\}$ over U we obtain a set with the cone at infinity equal to $Q^{-1}((-\infty, 0])$ which is disjoint with $C_{\infty}X$.

To prove one of the inclusions needed for m semicontinuity, take a sequence $D \ni x_{\nu} \to a$. Then for any sequence $y_{\nu} \in m(x_{\nu})$ convergent to y there is

$$Q(y_{\nu} - x_{\nu}) \le Q(y_{\nu} - p), \, \forall p \in X.$$

The inequality is preserved upon passing to the limit since Q is continuous, thus

$$\forall p \in X : Q(y-a) \le Q(y-p), \text{ and } y \in m(a).$$

The opposite inclusion is first proved for $a \notin M_X$. In such a case $m(a) = \{x_0\}$. Since *m* has nonempty values and is locally bounded, we can pick a convergent sequence from any $m(a_{\nu})$ where $a_{\nu} \to a$. Any such sequence must converge to x_0 due to the previous paragraph.

If $a \in M_X$ then from the first part we can approximate points of m(b) for any $b \in (a, x)$ where $x \in m(a)$. Consequently $m(a) \subset \limsup m(x)$.

Example 3.6. The condition concerning the Lipschitz constant in the previous theorem is optimal. Indeed, consider

$$X = \{ (t, t^2/(1+t^2), t) \in \mathbb{R}_{(2,1)} \mid t \in \mathbb{R} \}.$$

It is clearly a 1-pseudo-Lipschitz set, yet $\rho(a) = -\infty$ and m(a) is empty for any $a \notin V := \{(v, w, v) \mid w, v \in \mathbb{R}\}$. In the same time for

points $v = (v, w, v) \in V$ there is $\rho(v) = 0$ and $m(v) = (v, v^2/(1 + v^2), v)$.

Corollary 3.7. Let X be a L-pseudo-Lipschitz definable subset of $\mathbb{R}_{(l,p)}$ with L < 1, then the function ρ is continuous.

Proof. Choose $x \in \mathbb{R}_{(l,p)}$. Since the multifunction m is upper-semicontinuous, for any V - the neighbourhood of m(x) there exists U - a neighbourhood of x such that $m(U) \subset V$. Take V relatively compact, then for any point $y \in U$ there is $\rho(y) = \inf\{Q(y-a) \mid a \in X \cap V\}$. Now

$$\inf_{a \in X \cap V} \{Q(y-a)\} = Q(y-x) + \inf_{a \in X \cap V} \{2\langle \langle y-x, x-a \rangle \rangle + Q(x-a)\}.$$

The term Q(y-x) tends to zero when $y \to x$, the infimum on the other hand can be bounded from below by

$$\inf_{a \in X \cap V} \{2\langle \langle y - x, x - a \rangle \rangle + Q(x - a)\} \ge \inf_{a \in X \cap V} \{2\langle \langle y - x, x - a \rangle \rangle\} + \rho(x).$$

Since V was relatively compact the norm of x - a is bounded and $\langle \langle y - x, x - a \rangle \rangle$ tends to zero as $y \to x$, thus $\liminf \rho(y) \ge \rho(x)$. On the other hand $\rho(y) \le Q(y - b)$ for any $b \in m(x)$. Since Q is continuous we obtain $\limsup \rho(y) \le Q(x - b) = \rho(x)$.

Theorem 3.8. Let X be a L-pseudo-Lipschitz definable subset of $\mathbb{R}_{(l,p)}$ with L < 1, then ρ is locally lipschitz and differentiable for any point $x \in \mathbb{R}_{(l,p)} \setminus M_X$. Moreover for any such x there is $\nabla \rho(x) = 2A(x-m(x))$ thus ρ is \mathscr{C}^1 smooth in $\mathbb{R}_{(l,p)} \setminus \overline{M_X}$

Proof. For any $a \in \mathbb{R}_{(l,p)}$ the multifunction m is locally bounded. It means that we can find an open Euclidean ball B_a and R > 0 such that for any $b \in B_a$ there is $m(b) \subset B(0, R)$. Then ρ restricted to B_a is given as an infimum of functions $Q_b(x) := Q(x-b)$ with $b \in B(0, R) \cap X$. Every Q_b is Lipschitz in B_a and the Lipschitz constants are uniformly bounded by certain M > 0. Thus, ρ is M-Lipschitz in B_a as well.

Since ρ is locally lipschitz, the Rademacher theorem asserts that it is differentiable almost everywhere. In o-minimal structures it translates to the differentiability on an open and dense subset of $\mathbb{R}_{(l,p)}$.

For $x \notin M_X$ we investigate now the quotient

$$(\rho(x+h) - \rho(x) - 2(x - m(x))^T Ah) / \|h\| =$$

$$(Q(x+h-m(x+h)) - Q(x-m(x)) - 2\langle\langle x-m(x),h\rangle\rangle)/||h|| =$$

 $(Q(m(x+h)) - Q(m(x)) - 2\langle\langle x+h, m(x+h) - m(x)\rangle\rangle + o(||h||))/||h||.$ Where the second equality is due to $Q(x+h) - Q(x) - 2\langle\langle x,h\rangle\rangle =$

o(||h||). We can further simplify the last line, down to

(3.1)
$$\langle \langle m(x+h) + m(x) - 2x - 2h, \frac{m(x+h) - m(x)}{\|h\|} \rangle \rangle$$

6

Now, since m is definable it is differentiable in an open and dense subset of $\mathbb{R}_{(l,p)}$. Therefore, for almost all x we can write

(3.2)
$$\frac{m(x+h) - m(x)}{\|h\|} = \langle \nabla m(x), h/\|h\| \rangle + o(\|h\|).$$

Clearly, the quotient 3.2 is bounded and implies the boundedness of the expression 3.1.

Moreover $m(x+h) \in X$ and $m(x+h) \to m(x)$ when $h \to 0$. It means, that any accumulation point of $\frac{m(x+h)-m(x)}{\|h\|}$ ($h \to 0$) is an element of the tangent cone $C_{m(x)}X$. For any sequence $h \to 0$ yielding a convergent expression 3.1 we obtain therefore a limit

$$2\langle\langle m(x) - x, v\rangle\rangle, v \in C_{m(x)}X$$

which has to equal zero. Indeed, a lower estimate is given by Proposition 3.2. On the other hand, should $\langle \langle m(x)-x,v \rangle \rangle$ become positive, for h small enough we would obtain Q(x+h-m(x+h)) > Q(x+h-m(x)) which is a contradiction.

For an open and dense subset of $\mathbb{R}_{(l,p)}$ we have then

$$\nabla \rho(x) = 2A(x - m(x)).$$

The general theory of Clarke's subdifferential and theorem 3.5 assert now that ρ is differentiable whenever m(x) is a singleton. Moreover ρ is \mathscr{C}^1 smooth outside the closure of M_X .

Corollary 3.9. For any $x \notin M_X$ there is $m(x) = x - A\nabla \rho/2$.

Proof. It simply suffices to observe that $A^2 = I$. Then the previous theorem gives the assertion.

Theorem 3.10. Let X be a L-pseudo-Lipschitz definable subset of $\mathbb{R}_{(l,p)}$ with L < 1 then

- (1) $\partial \rho(x) = cnv\{2A(x-y) \mid y \in m(x)\} = \{2A(x-y) \mid y \in cnv m(x)\}$
- (2) $x \in M_X \iff \# \partial \rho(x) \neq 1 \iff x \notin D_{\rho}.$
- (3) $\nabla \rho = 0 \iff x \in X$
- (4) $\nabla \rho$ is continuous in $\mathbb{R}_{(l,p)} \setminus M_X$

Question 3.11. Is it true that $M_X \neq \emptyset$ if $\pi_l(X)$ do not contain any half-space?

Lemma 3.12. Let $X \subset \mathbb{R}^n$ be a closed, L-pseudo-Lipschitz set, $x_0 \in \mathbb{R}^n \setminus X$ a point and $B = B(x_0, r)$ a pseudo-ball such that $B \cap X = \bigcup_{j=1}^k X_j$ where the sets X_j are nonempty and pairwise disjoint, and for at least one $i, X_i \cap B \neq \emptyset$. Then there exists a neighbourhood U of x_0 such that

$$\rho(x,X) = \min_{j=1}^{k} \rho(x,X_j), \ x \in U$$

4. Central set

The definition of a central set in the Euclidean space \mathbb{E} is as follows. Firstly we define a collection of open balls included in the complement of X

$$\mathcal{B} := \{ B(x, r) \mid B(x, r) \subset \mathbb{E} \setminus X \}.$$

Then we call the ball $B \in \mathcal{B}$ maximal if

$$\forall B' \in \mathcal{B} : B \subset B' \Rightarrow B = B'.$$

The central set C_X of X collects the centres of all maximal balls in \mathcal{B} .

While adopting the notion in the pseudo-Euclidean setting one stumbles upon a problem. If Q is not positive definite the inclusion between any two balls B, B' implies their common centre. Therefore the central set C_X equals the whole complement of X. Indeed, every point x of Xcomplement is a centre of a maximal ball $B(x, \rho(x))$. The definition fails to yield any valuable study.

Hence a different, more local, approach is needed. As a source of our motivation we pick an observation that for any maximal ball B of centre c and $x \in \overline{B} \cap X \subset \mathbb{E}$ there is

$$\sup\{t \ge 0 \mid x \in m(x + t(c - x) / \|c - x\|)\} = \|c - x\|.$$

We construct the central set in the pseudo-Euclidean spaces by collecting all the points in the form $x + r_v v$ where $x \in X, v \in \mathcal{N}_x X$ and $r_v = \sup\{t \ge 0 \mid x \in m(x+tv)\}$. The main point in the definition is to establish a suitable object for the following Theorem which is crucial while investigating the limit passing of the medial axis families.

Theorem 4.1. Let X be a closed subset of pseudo-Euclidean space. Then $M_X \subset C_X \subset \overline{M_X}$.

Proof. The first inclusion is a consequence of Proposition 3.2(4). The proof of the second one is almost the same as in the standard Euclidean case (cf. [2]). Take a point $a \in C_X$ and assume that a is separated from the medial axis. Then there exists a radius r > 0 such that the multifunction m is univalued on $\mathbb{B}(a, r)$ and the image $m(\mathbb{B}(a, r))$ is a subset of $\mathbb{B}(m(a), ||m(a) - a||)$. In such a case, due to Proposition3.2, $a \in C_X$ implies $a \neq tx + (1 - t)m(x)$ for any $x \in \mathbb{B}(a, r)$. The function $H : (t, x) \to tx + (1 - t)m(x)$ is therefore a homothopy. Now, its composition with the spherical projection with the centre at a is a homothopy that transports the sphere $\mathbb{S}(a, r)$ onto its proper subset which is a contradiction.

Question 4.2. Is the definition of the Central set equivalent to the inclusion of a germ of the ball near the intersection points with X?

5. NASH LEMMA & KURATOWSKI LIMITS

The following theorem is a generalisation of a famous Nash lemma [3] to the pseudo-Euclidean spaces.

Theorem 5.1. Let $X \subset \mathbb{R}^n$ be a closed, k-dimensional L-pseudo-Lipschitz set with L < 1, then $\overline{M_X} \cap \operatorname{Reg}_2 X = \emptyset$

Proof. Take $x \in Reg_2 X$ and a local \mathscr{C}^2 parametrisation of X, a function $g: V \to X \cap U, g(0) = x$. Regard function

$$F: U \times V \ni (u, t) \to (\langle \langle u - g(t), \frac{\partial g}{\partial t_i}(t) \rangle \rangle)_{i=1}^k \in \mathbb{R}^k$$

Since the partial derivatives of g span $T_{g(t)}X$ we have F(u,t) = 0iff $u - g(t) \in \mathcal{N}_{g(t)}X$. Moreover it is possible to pick a neighbourhood U' of the point x that $m(x') \in U \cap X$ for all $x' \in U'$. In other words $\forall x' \in U' \exists t \in V : g(t) \in m(x')$.

Let us compute now

$$\det \frac{\partial F}{\partial t}(x,0) = -\det \left[\left\langle \left\langle \frac{\partial g}{\partial t_i}(0), \frac{\partial g}{\partial t_j}(0) \right\rangle \right]_{i,j=1}^k$$

and note that by a linear change of variables in V we can assure that

$$\langle \langle \frac{\partial g}{\partial t_i}(0), \frac{\partial g}{\partial t_j}(0) \rangle \rangle = 0 \text{ for } i \neq j.$$

In the end, the determinant will be equal to $-\sum Q(\frac{\partial g}{\partial t_i}(0))$. Since all the partial derivatives are vectors of the tangent cone T_xX , the values of $Q(\frac{\partial g}{\partial t_i}(0))$ are positive. Therefore the sum cannot vanish.

Now, the implicit function theorem allows us to uniquely solve the equation F(u,t) = 0 with respect to the variable t in a certain neighbourhood of (x,0). In details, we obtain a neighbourhood $W \times T \subset U \times V$ and a function $\tau : W \to V$ such that F(x,t) = 0 iff $t = \tau(x)$. Shrinking U' again to enforce $m(U') \subset g(T)$ we obtain therefore by setting $g(\tau(u))$ a unique vector satisfying $u - g(\tau(u)) \in \mathcal{N}_{g(\tau(u))}X$ for any given $u \in U'$. Thus the multifunction m is univalued in U'. Consequently M_X is disjoint with U' thus it is separated from $Reg_2 X$. \Box

An immediate corollary of the theorem forces us to search for the intersection points solely in the Sng_2X . The last set splits naturally into Sng_1X and $Reg_1X \cap Sng_2X$.

Question 5.2. Do the psuedo-Euclidean skeletons reach the set X under the same conditions as when treated as a subset of Euclidan space?

Lemma 5.3. Let $\{X_t\}_{t\in T}$ be a family of closed pseudo-L-lipschitz sets with $X_t \xrightarrow{K} X_0$, L < 1 and $0 \in \overline{T \setminus \{0\}}$. Posit $\rho_t(x) := \rho(x, X_t)$ and $m_t(x) = m(x, X_t)$. Then $\lim_{(t,x)\to(0,a)} \rho_t(x) = \rho_0(a)$ and $\limsup_{t\to 0} m_t(a) \subset m_0(a)$.

Proof. Take any $y \in m(a)$, then for t in a certain neighbourhood of 0 there is $X_t \cap \mathbb{B}(y, 1) \neq \emptyset$ due to the Kuratowski convergence of X_t .

Since sets X_t are pseudo-L-lipschitz, the inclusion occurs

$$X_t \subset \bigcup_{x \in \mathbb{B}(y,1)} S_x$$
, where $S_x = \{x' \mid ||x' - x||_l > L ||x' - x||_p\}.$

Thus, following the argument similar to the one of 3.8, all ρ_t are L-Lipschitz with an universal constant L_0 in a neighbourhood of a. Thus, for x_1, x_2 close to a there is

$$|\rho_t(x_1) - \rho_t(x_2)| \le L_0 ||x_1 - x_2||.$$

In the same time for small $\varepsilon > 0$ we have

$$B(a, \rho_0(a) + \varepsilon) \cap X_0 \neq \emptyset$$
 and $\overline{B(a, \rho_0(a) - \varepsilon)} \cap X_0 = \emptyset$.

Thus for t in suitably small neighbourhood of 0 there is $\rho_0(a) - \varepsilon < \rho_t(a) < \rho_0(a) + \varepsilon$. In the end we obtain thus

$$|\rho_t(x) - \rho_0(a)| \le |\rho_t(x) - \rho_t(a)| + |\rho_t(a) - \rho_0(a)| \le L_0 ||x - a|| + \varepsilon$$

which proves the convergence.

The inner continuity of m_t follows, as for any point $x \in X_0 \setminus m_0(a)$ by definition there is $Q(x-a) > \rho_0(a)$. However for a sequence of points $x_t \in X_t$ converging to $x \in \limsup m_t(a) \setminus m_0(a)$ that would mean

$$\rho_0(a) = \lim_{t \to 0} \rho_t(a) = \lim_{t \to 0} Q(x_t - a) = Q(x - a) > \rho_0(a)$$

which forms a clear contradiction.

Theorem 5.4. Let $\{X_t\}_{t\in T}$ be a family of closed pseudo-L-lipschitz sets with $X_t \xrightarrow{K} X_0$, L < 1 and $0 \in \overline{T \setminus \{0\}}$. Then

$$\liminf_{t \to 0} M_t \supset M_0,$$

where $M_t := M_{X_t}$.

Proof. Suppose that there exists a point

$$a \in M_0 \setminus \liminf_{t \to 0} M_t.$$

It implies existence of a sequence $T \ni t_{\nu} \to 0$ and an euclidean ball B centred at a such that

$$B \cap M_{t_{\nu}} = \emptyset$$
, for all $\nu \in \mathbb{N}$.

By shrinking the parameter space we can therefore assume that the intersection is empty for all $t \in T$. Mind that it does not change the convergence $X_t \to X_0$ in the induced topology.

For every $t \in T$ let m_t denote the closest points multifunction for X_t . Since a has a single closest point in X_t , it is possible to define

$$r(t) := \sup\{s \ge 0 \mid m_t(a) \in m_t(a(s,t))\}$$

for every $t \in T$, where $a(s,t) := sa + (1-s)m_t(a)$.

Note that since a(1,t) = a, the supremum is always greater or equal to one. Moreover, the point a(r(t),t) belongs to the central set of X_t whenever $r(t) < \infty$.

10

Our aim now is to prove that $\lim_{t\to 0} r(t) = 1$. Since $C_X \subset \overline{M_X}$ such limit would contradict $B \cap M_t = \emptyset$ for small t. Clearly $\liminf r(t) \ge 1$ since r(t) is at least equal to one. Suppose then that $\limsup r(t) > 1$. In such a case it is possible to pick c > 0 and a sequence $t_{\nu} \to 0$ for which the pseudoballs $B_{\nu} := B(a(1+c,t_{\nu}), Q(a(1+c,t_{\nu})-m_{t_{\nu}}(a)))$ are disjoint with $X_{t_{\nu}}$. This sequence of balls converges to the closure of a ball $B_0 := B(a(1+c,0), Q(a(1+c,0)-y))$, where y is a certain point in $m_0(a)$ (cf Lemma 5.3). Now, should $B_0 \cap X_0 = \emptyset$, the point y would be the closest to a(1+c,0) and #m(a) = 1 since $a \in (y, a(1+c,0))$ which contradicts $a \in M_0$. On the other hand any point $x \in B_0 \cap X_0$ would contradict the convergence of $X_t \to X_0$.

Therefore, the set $M_0 \setminus \liminf M_t$ ought to be empty.

6. Medial axis for hypersurfaces

Observe firstly that under the restrictions introduced so far, the only scenarios for X to be a hypersurface would be to consider $\mathbb{R}_{(l,p)}$ with p equal zero or one. The first instance is precisely the Euclidean case. For p = 1 we have what follows.

Theorem 6.1. Assume that X is k-codimensional 1-pseudo-Lipschitz closed subset of $\mathbb{R}_{(l,p)}$ and $a \in X$. Let x_1, \ldots, x_{k+1} be points in \mathbb{R}^n with $a \in m(x_i)$ for $i = 1 \ldots k + 1$ such that the vectors $x_i - a$ are linearly independent. Then $a \in Sng_1X$.

Proof. Define the hyperplanes H_i by

 $H_i := \{ v \in \mathbb{R}_{(l,p)} \mid \langle \langle x_i - a, v \rangle \rangle = 0 \}.$

If we had $a \in Reg_1X$ then by the choice of a we would have $x_i - a \in \mathcal{N}_aX$ for $i = 1, \ldots, k + 1$, and $T_aX \subset \bigcap_{i=1}^{k+1} H_i$. However, the intersection of H_i is transversal, thus its codimension equals k + 1 which is a contradiction.

Theorem 6.2. Take $a \in X$ and assume that for all x in some neighbourhood U of $x_0 \in \mathbb{R}^n$ we have $a \in m(x)$. Then $a \in Sng_1X$.

Proof. Simply apply the last theorem to any $\operatorname{codim} X + 1$ linearly independent vectors $x_i - a \in U - a$.

Theorem 6.3. Let X be a 1-pseudo-Lipschitz closed hypersurface of $\mathbb{R}_{(l,p)}$ and $a \in \mathbb{R}_{(l,p)} \setminus M_X$ be such that $m(a) \in \operatorname{Reg}_1 X$. Then for $S = B \cap \Gamma$ where Γ is any \mathscr{C}^1 submanifold with $T_a\Gamma$ transversal to $\mathbb{R}(a - m(a))$ and B is sufficiently small open ball centred at a, the mapping $m|S : S \to m(S)$ is a homeomorphism onto the open subset $m(S) \subset \operatorname{Reg}_1 X$.

Proof. By assumption, m is univaluead in a neighbourhood U of a, and thus continuous. Therefore for an open V such that $V \cap X \subset Reg_1 X$ we can find an open ball B centered at a with $m(B) \subset V$.

By the previous proposition for ant two points $x_1, x_2 \in S$ we have an inclusion between segments $[x_1, m(a)], [x_2, m(a)]$ which (provided B is small enough) means that $x_1 = x_2$. Thus, m|S is injective, and so by Brouwer Domain Invariance Theorem it is a homeomorphism onto an open subset $m(S) \subset Reg_1 X$

Remark 6.4. The previous theorem includes, in particular, the case when Γ equals the sphere $\mathbb{S}(m(a), ||a - m(a)||)$.

References

- L. Birbrair and M. Denkowski. "Medial axis and Singularieties". In: J. Geom. Anal. 27.3 (2017), pp. 2339–2380.
- [2] D. Fremlin. "Skeletons and central sets". In: P. Lon. Math. Soc. 74.3 (1997), pp. 701–720.
- [3] J. Nash. "Real Algebraic Manifolds". In: Ann. Math. 56.3 (1952), pp. 405–421.

JAGIELLONIAN UNIVERSITY, FACULTY OF MATHEMATICS AND COMPUTER SCI-ENCE, INSTITUTE OF MATHEMATICS, LOJASIEWICZA 6, 30-348 KRAKÓW, PO-LAND

Email address: adam.bialozyt@uj.edu.pl

12