

# Poincaré Inequality for Local Log-Polyak-Łojasiewicz Measures : Non-asymptotic Analysis in Low-temperature Regime

Yun Gong  
Peking University

STUDENT\_GONGYUN@PKU.EDU.CN

Niao He  
ETH Zürich

NIAO.HE@INF.ETHZ.CH

Zebang Shen  
ETH Zürich

ZEBANG.SHEN@INF.ETHZ.CH

## Abstract

Potential functions in highly pertinent applications, such as deep learning in over-parameterized regime, are empirically observed to admit non-isolated minima. To understand the convergence behavior of stochastic dynamics in such landscapes, we propose to study the class of  $\text{Log-PL}^\circ$  measures  $\mu_\epsilon \propto \exp(-V/\epsilon)$ , where the potential  $V$  satisfies a local Polyak-Łojasiewicz (PL) inequality, and its set of local minima is provably *connected*. Notably, potentials in this class can exhibit local maxima and we characterize its optimal set  $S$  to be a compact  $C^2$  embedding submanifold of  $\mathbb{R}^d$  without boundary. The *non-contractibility* of  $S$  distinguishes our function class from the classical convex setting topologically. Moreover, the embedding structure induces a naturally defined Laplacian-Beltrami operator on  $S$ , and we show that its first non-trivial eigenvalue provides an  $\epsilon$ -independent lower bound for the Poincaré constant in the Poincaré inequality of  $\mu_\epsilon$ . As a direct consequence, Langevin dynamics with such non-convex potential  $V$  and diffusion coefficient  $\epsilon$  converges to its equilibrium  $\mu_\epsilon$  at a rate of  $\tilde{O}(1/\epsilon)$ , provided  $\epsilon$  is sufficiently small. Here  $\tilde{O}$  hides logarithmic terms.

**Keywords:** Poincaré inequality, non-log-concave measure

## 1. Introduction

Consider the Langevin dynamics

$$dX(t) = -\nabla V(X(t))dt + \sqrt{2\epsilon}dW(t), \quad (1)$$

where  $V \in C^2(\mathbb{R}^d, \mathbb{R})$  and  $\epsilon > 0$  are the *potential* and *temperature* of the above system respectively, and  $W(t)$  denotes the  $d$ -dimensional Brownian motion. Under mild conditions, the above Stochastic Differential Equation (SDE) yields a unique equilibrium  $\mu_\epsilon$ , commonly known as the Gibbs measure:

$$\mu_\epsilon(x) = \frac{\exp(-V(x)/\epsilon)}{Z_\epsilon}, \text{ where } Z_\epsilon = \int_{\mathbb{R}^d} \exp(-V(x)/\epsilon)dx. \quad (2)$$

Langevin dynamics have various applications in domains like statistics, optimization, and machine learning, including sampling from target distribution  $\mu_\epsilon$  (Wibisono, 2018), minimizing non-convex objectives (Raginsky et al., 2017; Zhang et al., 2017), modelling Stochastic Gradient Descent (SGD) through SDE approximation (Li et al., 2017; Ben Arous et al., 2022; Paquette et al., 2022; Li et al., 2024). In practice, the low-temperature regime ( $\epsilon \ll 1$ ) is particularly relevant: In the context of sampling, low temperatures enable sharper concentration of samples around the modes, while in the context of SGD approximations,  $\epsilon$  corresponds to the step size, which is typically small.

An important aspect of Langevin dynamics is its convergence behavior toward equilibrium, also known as ergodicity (Cattiaux and Guillin, 2017). This is often studied through functional inequalities like the Poincaré inequality (PI) and the Log-Sobolev inequality (LSI), which quantify the rate of convergence. In this work, we focus on PI. Here, we recall that a measure  $\mu$  satisfies PI

\*Authors are listed in alphabetical order.

with Poincaré constant  $\rho_\mu$  (formally defined in Definition 1) if for any test function  $f$  in the Sobolev space weighted by  $\mu$ , its variance times  $\rho_\mu$  is bounded by its Dirichlet energy (both measured w.r.t.  $\mu$ ). Clearly, the PI constant of  $\mu_\epsilon$  is a function of the temperature  $\epsilon$ .

The convergence of Langevin dynamics under the  $\chi^2$ -divergence is closely tied to the constant  $\rho_{\mu_\epsilon}$ . Specifically, to achieve  $\chi^2(X(t), \mu_\epsilon) \leq \omega$ , the required time is  $t = \mathcal{O}\left(\frac{1}{\epsilon\rho_{\mu_\epsilon}} \log \frac{1}{\omega}\right)$ . Hence, in the low-temperature region, the dependence on  $\epsilon$  is the determining factor in the rate of convergence, which will be the *main focus of the paper*. Notably, this convergence behavior varies significantly between uni-modal and multi-modal distributions, and is reflected in the dependence of  $\rho_{\mu_\epsilon}$  on  $\epsilon$ :

- (constant-time convergence) When  $V$  is a strongly convex function or is close to one up to a perturbation of order  $\epsilon$ , a classical result is that  $\rho_{\mu_\epsilon}$  is of order  $\Omega(\frac{1}{\epsilon})$  (Bakry et al., 2014). Hence the mixing time  $t = \tilde{\mathcal{O}}(1)$  for all low temperatures.
- (sub-exponential-time convergence) For a general log-concave measure, existing research primarily focuses on how the PI constant depends on the problem dimension  $d$ , commonly known as the Kannan-Lovász-Simonovits (KLS) conjecture (Chen, 2021; Lee and Vempala, 2024), but gives little emphasis to its dependence on  $\epsilon$ . Nevertheless, existing technique can be combined to show that  $\rho_{\mu_\epsilon} = \Omega(1)$  under a mild exponential integrability assumption, i.e.  $\int_{\mathbb{R}^d} \exp(-V(x))dx < \infty$ . See a proof in Appendix B. Moreover, one can easily construct a convex function such that the corresponding  $\rho_{\mu_\epsilon}$  is constant. Consequently, for the general class of log-concave measures,  $\rho_{\mu_\epsilon} = \Theta(1)$ , and hence, the mixing time is  $t = \tilde{\mathcal{O}}(\frac{1}{\epsilon})$ , i.e. sub-exponential.
- (exponential-time convergence) When  $V$  has at least two *separated* local minima, convergence occurs in two distinct time-scales: a sub-exponential-time scale describes  $X(t)$  reaching a meta-stable equilibrium in one of  $V$ 's local regions of attraction; and an exponential scale describes the transition between meta-stable equilibria, which typically takes  $\Omega(\exp(\frac{1}{\epsilon}))$  time (Bovier et al., 2004; Gayraud et al., 2005; Menz and Schlichting, 2014). This exponential-time estimation is commonly known as the the Eyring-Kramers law (Eyring, 1935; Kramers, 1940).

A more detailed literature review is deferred to Appendix A.

In this paper, we focus on uni-modal measures, which allows convergence to occur within sub-exponential time<sup>1</sup>. Multi-modal measures, as discussed above, exhibit global convergence on an exponential time scale, a behavior which is less relevant in practical applications. We leave its investigation for future work.

For uni-modal measures, existing research often assumes that the optimal set have a simplistic structure, such as a singleton (when  $V$  is strongly convex) or a convex set (when  $V$  is convex). These assumptions, while analytically convenient, limit the scope of applications. For example, in key applications like deep learning, the function  $V$  is high-dimensional, highly *non-convex*, and many research suggest that, for over-parameterized models, the local minima are often *degenerate* and are *non-singletons* (Sagun et al., 2016; Safran and Shamir, 2016; Freeman and Bruna, 2017; Sagun et al., 2017; Venturi et al., 2018; Liang et al., 2018b; Draxler et al., 2018; Garipov et al., 2018; Liang et al., 2018a; Nguyen, 2019; Kawaguchi and Kaelbling, 2020; Kuditipudi et al., 2019; Lin et al., 2024). In emerging domains like Large Language Model, the number of parameters is of order billions and the scaling laws implies that the over-parameterization phenomenon will be increasingly more prominent. These important cases fall outside the scope of existing studies.

---

1. Our analysis can also address convergence toward a metastable equilibrium for multi-modal measures if the domain is properly partitioned, and appropriate boundary conditions are imposed. However, this is orthogonal to the primary focus of this project and is therefore not discussed here.

**Log-PL<sup>◦</sup> measures** We aim to address more complex structural possibilities within the uni-modal framework. To define the uni-modal measure class of interest, we state the necessary assumptions.

**Assumption 1** *Let  $S$  the collection of all local minima of the potential  $V \in \mathcal{C}^2$ . For every connected component  $S'$  in  $S$ , there exists an open neighborhood  $\mathcal{N}(S') \supset S'$  such that, in  $\mathcal{N}(S')$ ,  $V \in \mathcal{C}^3$  and moreover it is locally PL(Lojasiewicz, 1963; Polyak, 1963):*

$$\forall x \in \mathcal{N}(S'), |\nabla V(x)|^2 \geq \nu \left( V(x) - \min_{x \in \mathcal{N}(S')} V(x) \right). \quad (3)$$

The above assumption ensures “sharp boundaries” of the local optimal sets, i.e. the landscape within  $\mathcal{N}(S')$  is not overly flat: Rebjock and Boumal (2024) prove that the local PL condition implies local quadratic growth, thereby enforcing a curvature that prevents flatness in this region. Next, we need to exclude the possibility of saddle points so as to ensure the uni-modality. In words, the following assumption states that a critical point of  $V$  is either a local minimum or maximum<sup>2</sup>.

**Assumption 2** *Let  $\mathcal{N}(S)$  be the union of all the neighborhoods  $\mathcal{N}(S')$  defined in Assumption 1. For any  $x \in \mathbb{R}^d \setminus \mathcal{N}(S)$ , if  $\nabla V(x) = 0$ , one has  $\nabla^2 V(x) < 0$ .*

We further need all the local minima to be within a compact set. This is a technical assumption and we believe it can be relaxed to the coercivity of  $V$ . The latter is typically necessary for  $Z_\epsilon < \infty$ .

**Assumption 3'**  *$V$  is coercive and all local minima of  $V$  are contained in a compact set.*

We call  $V$  a PL<sup>◦</sup> function if it satisfies the above conditions, and refer to the Gibbs measure  $\mu_\epsilon$  as a Log-PL<sup>◦</sup> measure<sup>3</sup>. Note that if  $V$  is globally PL and coercive, the above assumptions, apart from the regularity ones, follow directly. However, our assumptions allow for the existence of local maxima, making them strictly weaker. This distribution class is of interest for the following reasons:

- (Relevance to crucial problems.) The local PL inequality is established for a class of over-parameterized neural networks (Oymak and Soltanolkotabi, 2020; Liu et al., 2022), and PL functions are an important class in the optimization literature (Karimi et al., 2016; Yang et al., 2020; Rebjock and Boumal, 2024).
- (Connectivity of optimal set.) We prove in Proposition 3 that, when the ambient dimension  $d \geq 2$ , Assumptions 1 to 3' together imply that the collection of all local minima has only one connected component. Hence, Log-PL<sup>◦</sup> measures are uni-modal. Our proof is built on a generalized version of the famous Mountain Passing Theorem (Katriel, 1994).
- (Optimal set with pertinent structures.) Built on the connectivity result above, we prove that the optimal set  $S$  is a  $\mathcal{C}^2$  embedding submanifold of the ambient space  $\mathbb{R}^d$  without boundary. This class of optimal set is highly pertinent to the machine learning community (Cooper, 2018, 2020; Fehrman et al., 2020; Li et al., 2022; Wojtowytsch, 2024; Levin et al., 2024)<sup>4</sup>. In particular, the optimal set  $S$  can be *non-contractible*, thus *topologically* different from convex sets. A simple example satisfying all assumptions is  $V(x) = \|x\|^3/3 - \|x\|^2/2$ , whose global optimal set is  $\|x\| = 1$ , forming a non-convex, non-contractible embedding submanifold.

Consequently, we believe understanding the convergence of the Langevin dynamics towards a Log-PL<sup>◦</sup> measure can provide a rich template for studying crucial problems like deep learning.

Further, to prove a fast sub-exponential convergence, w.r.t.  $1/\epsilon$ , of the Langevin dynamics, we make some technical assumptions, compiled in Section 2.5 for the ease of reference.

---

2. If we know a priori that  $\mu_\epsilon$  is uni-modal, Assumption 2 can be relaxed to “for all  $x \in \mathbb{R}^d \setminus \mathcal{N}(S)$ , if  $\nabla V(x) = 0$ ,  $\lambda_{\min}(\nabla^2 V(x)) < 0$ ”. However, this relaxed assumption is not sufficient to guarantee the uni-modality of  $\mu_\epsilon$ .  
 3. The superscript <sup>◦</sup> highlights that the mode of  $\mu_\epsilon$  can be a  $d$ -sphere, a representative embedding submanifold.  
 4. Existing papers assume this structure of  $S$  without proof, but we derive this result from our assumptions on  $V$ .

**Our result.** We prove that a  $\text{Log-PL}^\circ$  measure, while far from being log-concave, possesses a PI constant that is *non-asymptotically* lower bounded by a *temperature-independent* constant, for a sufficiently small  $\epsilon$ . The applicable temperature region depends on the geometric structure of the optimal set  $S$ . Our result is briefly summarized as follows, and formally stated in Theorem 4.

**Theorem 1 (informal)** *Suppose that the potential  $V$  satisfies Assumptions 1 to 3' and some additional regularity assumptions in Section 2.5. Consider the case where  $S$  is not a singleton. When  $\epsilon$  is sufficiently small, the Poincaré constant (4) of the measure  $\mu_\epsilon$  satisfies  $\rho_{\mu_\epsilon} = \Omega(\lambda_1(S))$ . Here  $\lambda_1(S) > 0$  denotes the first non-trivial eigenvalue of the Laplacian-Beltrami operator on  $S$ .*

A direct consequence of the result above is a quantitative characterization of the convergence behavior of Langevin dynamics and its discrete-time implementation in the low temperature region, such as the Langevin Monte Carlo (LMC) (Chewi et al., 2024, Theorem 7), for a  $\text{Log-PL}^\circ$  target measure. Our result also represents a significant step toward proving the stronger LSI, as the PI constant is often a crucial intermediary for estimating the LSI constant (Cattiaux et al., 2010, Theorem 1.2).

Furthermore, to the best of our knowledge, *in the low temperature region*, our work marks the first attempt to explore the behavior of Langevin dynamics in general non-convex landscapes with *non-isolated minimizers*. In addition, technique-wise, we are the first to connect the Poincaré constant of a measure on  $\mathbb{R}^d$  with the stability of the Laplacian-Beltrami eigenvalue on the optimal solution manifold, offering a novel perspective.

**Our proof strategy.** Our proof is split into two steps. Let  $U$  be a neighborhood of the optimal set  $S$ . First, we reduce the PI constant for a measure supported on  $\mathbb{R}^d$  to an eigenvalue problem of the Laplacian operator on  $U$ . Seeing  $U$  as an expansion of  $S$  (since  $S \subset U$ ), we then relate the eigenvalues on  $U$  and  $S$  through a stability analysis. A more technical summary is as follows.

1. First, we partition the domain  $\mathbb{R}^d$  into a collection of subdomains, allowing us to apply existing Lyapunov methods in (Menz and Schlichting, 2014) for establishing the PI constant, as stated in Proposition 1. This reduces the estimation of the PI constant  $\rho_{\mu_\epsilon}$  to the estimation of the Neumann eigenvalue of the Laplacian operator on a domain  $U = S^{\sqrt{C}\epsilon}$ . Here  $C$  is a constant independent of  $\epsilon$ , and  $S^\eta := \{x \in \mathbb{R}^d : \text{dist}(x, S) \leq \eta\}$ . See the precise statement in Theorem 2.
2. Second, we establish a temperature-independent lower bound for the Neumann eigenvalue on  $U$ . Since  $S$  is a  $C^2$  embedding submanifold,  $S^\eta$  matches the *tubular neighborhood* of  $S$ . By the tubular neighborhood theorem (Milnor and Stasheff, 1974), up to a diffeomorphism, we can decompose the uniform distribution on  $U$  as a pair of decoupled distributions along the tangent and the normal directions respectively. With the tensorization property of the Poincaré inequality (Bakry et al., 2014), we show that the Neumann eigenvalue on  $U$  is determined by the first non-trivial eigenvalue of the Laplacian-Beltrami operator on  $S$ , when  $\epsilon$  is sufficiently small. Here, the Laplacian-Beltrami operator is defined based on the aforementioned embedding structure of the submanifold  $S$ .

### Summary of contributions

- We identify the  $\text{PL}^\circ$  class as a rich and suitable template for studying the (sub-exponential time) convergence behavior of stochastic dynamics on potentials that admit non-isolated minima, a phenomenon empirically observed in crucial applications like deep learning in the over-parameterized regime. We prove that all local minima of a  $\text{PL}^\circ$  function are connected and its global optimal set  $S$  formulates a compact  $C^2$  embedding submanifold of  $\mathbb{R}^d$  without boundary.
- Built on the above characterization of the optimal set  $S$ , we show that the Poincaré constant of the Gibbs measure  $\mu_\epsilon$  (2) is lower bounded by the first non-trivial eigenvalue of the Laplacian-Beltrami operator on  $S$ , when  $S$  is non-singleton. This eigenvalue is temperature-independent. As a direct consequence, the Langevin dynamics (1) converges to  $\mu_\epsilon$  at a rate of  $\tilde{O}(1/\epsilon)$ .

**Relation to the sampling literature** To better position our work within the literature, we restate our primary focus: understanding the dependence of the PI constant  $\rho_{\mu_\epsilon}$  on  $\epsilon$ , which directly impacts the convergence rate of the Langevin dynamics. Many excellent works in the sampling literature focus on analyzing the convergence of the discretized algorithms like LMC in the *constant-temperature* ( $\epsilon = 1$ ) region, for (strongly) convex or nonconvex potentials under weak conditions, e.g. a joint of dissipativity and tail growth conditions (Erdogdu and Hosseinzadeh, 2021). We do not directly compare with them here since we focus on different temperature regions.

## 2. Preliminaries and Assumptions

### 2.1. Poincaré Inequality

**Definition 1 (Poincaré-Wirtinger Inequality)** A probability measure  $\mu$  with support  $\Omega \subseteq \mathbb{R}^d$  satisfies the Poincaré inequality with parameter  $\rho_\mu$ , or shortly  $PI(\rho_\mu)$ , if one has for any  $f \in H^1(\mu)$

$$\left\{ \text{Var}_\mu(f) := \int_\Omega \left( f - \int_\Omega f d\mu \right)^2 d\mu \right\} \leq \frac{1}{\rho_\mu} \left\{ \mathcal{D}_\mu(f) := \int_\Omega |\nabla f|^2 d\mu \right\}, \quad (4)$$

where  $\rho_\mu$  is called the PI constant and  $H^1(\mu)$  denotes the Sobolev space weighted by  $\mu$ .

Here  $\text{Var}_\mu(f)$  and  $\mathcal{D}_\mu(f)$  are the variance and the Dirichlet energy of the test function  $f \in H^1(\mu)$  w.r.t.  $\mu$ . For the Gibbs measure  $\mu_\epsilon$ , we have  $\Omega = \mathbb{R}^d$ .

### 2.2. The Lyapunov Function Approach and the Perturbation Principle

Define the truncated Gibbs measure on a given domain  $U \subset \mathbb{R}^d$  as

$$\mu_{\epsilon,U}(dx) = \frac{1_U}{Z_{\epsilon,U}} \exp\left(-\frac{V(x)}{\epsilon}\right) dx, \quad \text{with } Z_{\epsilon,U} = \int_U \exp\left(-\frac{V(x)}{\epsilon}\right) dx. \quad (5)$$

The next statement shows that a Lyapunov function and the PI for the truncated measure  $\mu_{\epsilon,U}$  can be combined to get the PI for the original Gibbs measure. Our work is built on this framework.

**Proposition 1 (Menz and Schlichting, 2014, Theorem 3.8)** Let  $\mathcal{L} := -\nabla V \cdot \nabla + \epsilon \Delta$  be the infinitesimal generator associated with the Langevin dynamics in eq. (1). A function  $\mathcal{W} : \mathbb{R}^d \rightarrow [1, \infty)$  is a Lyapunov function for  $\mathcal{L}$  if there exists  $U \subseteq \mathbb{R}^d$ ,  $b > 0$ ,  $\sigma > 0$ , such that

$$\forall x \in \mathbb{R}^d, \quad \epsilon^{-1} \mathcal{L} \mathcal{W}(x) \leq -\sigma \mathcal{W}(x) + b 1_U(x). \quad (6)$$

Given the existence of such a Lyapunov function  $\mathcal{W}$ , if one further has that the truncated Gibbs measure  $\mu_{\epsilon,U}$  satisfies PI with constant  $\rho_{\epsilon,U} > 0$ , the Gibbs measure  $\mu_\epsilon$  satisfies PI with constant

$$\rho_\epsilon \geq \frac{\sigma}{b + \rho_{\epsilon,U}} \rho_{\epsilon,U}. \quad (7)$$

Following Menz and Schlichting (2014), we adopt  $\mathcal{W}(x) = \exp\left(\frac{1}{2\epsilon} V\right)$  as the Lyapunov function throughout this work. This function satisfies  $\mathcal{W}(x) \geq 1$  since we assume  $\text{WLOG } V^* = 0$ . The only remaining argument is to establish the condition in eq. (6). To be more precise, we need to find two constants  $\sigma > 0$ ,  $b > 0$  and some set  $U \subset \mathbb{R}^d$  such that

$$\frac{\mathcal{L} \mathcal{W}}{\epsilon \mathcal{W}} = \frac{\Delta V}{2\epsilon} - \frac{|\nabla V|^2}{4\epsilon^2} \leq -\sigma + b 1_U. \quad (8)$$

We will find these two constants in Lemma 3. In addition to the above Lyapunov function framework, the following standard perturbation principle will also be helpful to us.

**Proposition 2 (Holley-Stroock perturbation principle)** Let  $V$  and  $\tilde{V}$  be two potential functions defined on a domain  $U$ . If the truncated Gibbs measures, defined in eq. (5), with energies  $V$  and  $\tilde{V}$  satisfy  $PI(\rho)$  and  $PI(\tilde{\rho})$  respectively, one has  $\rho \geq \exp\left\{-\frac{1}{\epsilon} (\sup_{x \in U} (V - \tilde{V}) - \inf_{x \in U} (V - \tilde{V}))\right\} \tilde{\rho}$ .

### 2.3. Properties of a $C^2$ Embedding Submanifold

The Laplacian-Beltrami operator on the optimal set  $S$  is crucial to our analysis. The definition of this operator is built on a pullback metric induced by the embedding structure of  $S$ , outlined as follows. Note that in the rest of the paper, we use  $k$  to denote the dimension of the manifold  $S$  and focus on the case  $k \geq 1$ . For  $k = 0$ ,  $S$  becomes a singleton, which is not the focus of our work.

**Definition 2 (Embedding submanifold in  $\mathbb{R}^d$ )** Consider a  $C^2$  manifold  $M$  such that  $M \subseteq \mathbb{R}^d$ . If the including map  $i_M : M \rightarrow \mathbb{R}^d$  is  $C^2$  and satisfies following two conditions: (1) The tangent map  $Di_M(x)$  has rank equal to  $\dim M$  for all  $x \in M$ ; (2)  $i_M$  is a homeomorphism of  $M$  onto its image  $i_M(M) \subset \mathbb{R}^d$ , where  $i_M(M)$  inherits the subspace topology from  $\mathbb{R}^d$ . We say that the including map  $i_M$  is an embedding and  $M$  is a  $C^2$  embedding submanifold of  $\mathbb{R}^d$ .

If  $M$  is a  $k$ -dimensional embedding submanifold of  $\mathbb{R}^d$ , the including map can be represented using the local coordinates of  $M$  as follows: Assume that  $\{\Gamma_i, \phi_i\}_{i \in \Lambda}$  is the maximal atlas of  $M$ . For  $u = (u^1, \dots, u^k) \in \Gamma_i \subset \mathbb{R}^k$ , the including map  $i_M : M \rightarrow \mathbb{R}^d$  can be written as

$$x^j = m_j^i(u^1, u^2, \dots, u^k), j \in \{1, \dots, d\}, \quad (9)$$

where  $m_j^i : \Gamma_i \subset \mathbb{R}^k \rightarrow \mathbb{R}$ ,  $j = 1, \dots, d$  are  $C^2$  coordinate functions. We also denote this embedding structure as  $\mathcal{M}^i(u) = (m_1^i(u), \dots, m_d^i(u))$  on a local chart  $(\Gamma_i, \phi_i)$ .

**Remark 1** *WLOG, we assume that there is only one local chart  $(\Gamma, \phi)$  in the rest of the paper, since we can always extend local results to a global one by the standard technique of partition of unity (Tu, 2010, Chapter 13). We write the corresponding embedding structure as  $\mathcal{M}$ , omitting the superscript. Any non-trivial differences encountered in related proofs will be explicitly highlighted.*

Given the above embedding structure, the embedding submanifold  $M$  naturally inherits Riemannian structures from the ambient space  $\mathbb{R}^d$ . In the following, we describe the first and second fundamental forms on  $M$ . The reader can find more details about these structures in Appendix D.

**The first fundamental form (or Riemannian metric).** We define the Riemannian metric  $g_M$  on  $M$  as the pullback metric induced by the including map  $i_M : M \hookrightarrow \mathbb{R}^d$ , i.e.  $g_M = i_M^*(g_E)$ , where  $g_E$  is the standard Riemannian metric on  $\mathbb{R}^d$  and  $i_M^*$  is the pullback map associated with  $i_M$ . Now we can say that  $(M, g_M)$  is a  $k$ -dimensional Riemannian submanifold on  $\mathbb{R}^d$  and the including map  $i_M : (M, g_M) \hookrightarrow (\mathbb{R}^d, g_E)$  is a Riemannian embedding. Based on this Riemannian metric, on the local chart  $(\Gamma, \phi)$ , we can define the *Laplacian-Beltrami operator*  $\Delta_{g_M}$  as

$$-\Delta_{g_M} = -\frac{1}{\sqrt{\det(g_M)}} \sum_{i,j=1}^k \frac{\partial}{\partial u^i} \left( \sqrt{\det(g_M)} g^{ij} \frac{\partial}{\partial u^j} \right) \quad u \in \Gamma, \quad (10)$$

and the standard volume form  $d\mathcal{M}$  as  $d\mathcal{M}(u) = \sqrt{\det(g_M)} |du^1 \wedge \dots \wedge du^k|$ ,  $u \in \Gamma$ , where  $\det(g_M)$  is the determinant of the matrix  $g_M = (g_{ij})$ , and  $(g^{ij})$  is the inverse matrix of  $g_M$ .

**The second fundamental form.** Recall that  $\mathcal{M}(u)$  is the embedding structure defined above. Let  $\mathcal{N}_{k+1}, \dots, \mathcal{N}_d : \Gamma \rightarrow \mathbb{R}^d$  be  $d - k$  normal vectors on  $M$  which are orthogonal to each other. Define the matrix  $G(l) = [G_{ij}(l)]$  with  $G_{ij}(l) = -\frac{\partial^2 \mathcal{M}(u)}{\partial u^i \partial u^j} \cdot \mathcal{N}_l(u)$ ,  $l = k + 1, \dots, d$ , for  $u \in \Gamma \subset \mathbb{R}^k$ . With this notation, we can define the second fundamental form of the manifold  $\mathcal{M}$  by  $\Pi = -\sum_{l=k+1}^d \left\{ r^l \sum_{i,j=1}^k G_{ij}(l) du^i du^j \right\}$ , for some small  $(r^l)_{l=k+1}^d$ . We further define a matrix  $\tilde{G}(l) = [G_j^i(l)]$  with  $G_j^i(l) = \sum_{s=1}^k g^{is} G_{sj}(l)$ ,  $l = k + 1, \dots, d$ , which will be useful to our presentation.

## 2.4. The Eigenvalue Problems of Differential Operators

In the next, we introduce the eigenvalue problem of the Laplacian operator with Neumann boundary condition on a compact set  $\Omega$ .

**Definition 3 (Neumann eigenvalue)** *Consider the eigenvalue problem for the Laplacian operator on a closed domain  $\Omega$ , subject to the Neumann boundary condition*

$$-\Delta u = \lambda u, x \in \text{int } \Omega \text{ and } \partial u / \partial \nu = 0, x \in \partial \Omega,$$

where  $\nu$  is the outward normal vector to  $\partial \Omega$  and  $u \in H^1(\Omega)$ . The Neumann eigenvalue  $\lambda_1^n(\Omega)$  is defined to be the minimum non-zero eigenvalue  $\lambda$  to the above problem.

Recall the Poincaré inequality in Definition 1. It is known that  $\lambda_1^n(\Omega)$  matches the best Poincaré constant for the Lebesgue measure on  $\Omega$ . We can use min-max formulation for Neumann eigenvalue of Laplacian operator to derive this fact, see (Davies, 1995, Theorem 4.5.1), i.e., it admits the following variational formulation:

$$\lambda_1^n(\Omega) = \inf_{\substack{L \subseteq H^1(\Omega) \\ \dim(L)=2}} \sup_{u \in L} \left\{ \frac{|\nabla u|_{L^2(\Omega)}^2}{|u|_{L^2(\Omega)}^2} \right\} = \min \left\{ \frac{|\nabla u|_{L^2(\Omega)}^2}{|u|_{L^2(\Omega)}^2} : u \in W^{1,2}(\Omega) \setminus \{0\}, \int_{\Omega} u(x) dx = 0 \right\}. \quad (11)$$

We will exploit the above formulation for  $\Omega = U = S^{\sqrt{C}\epsilon}$  in our analysis.

Next, we introduce the eigenvalue problem of the Laplacian-Beltrami operator on the compact Riemannian submanifold  $M$ , which strongly depends on the non-trivial metric  $g_M$ .

**Definition 4 (Eigenvalue of the Laplacian-Beltrami operator)** *Consider the eigenvalue problem for the Laplacian-Beltrami operator on the Riemannian submanifold  $(M, g_M)$  without boundary,*

$$-\Delta_{g_M} u = \lambda u, \quad x \in M, \quad (12)$$

where  $u \in H^1(M)$  and  $-\Delta_{g_M}$  is the Laplacian-Beltrami operator on  $M$  associated with metric  $g_M$ . The eigenvalue  $\lambda_1(M)$  is defined to be the minimum non-zero eigenvalue  $\lambda$  to the above problem.

Define the Dirichlet energy of Laplacian-Beltrami operator on  $S$ , for  $f \in W^{1,2}(M)$

$$\mathcal{Q}_M(f, f) = \int_M \langle f, -\Delta_{g_M} f \rangle_{g_E} d\mathcal{M} = \int_M \langle df, df \rangle_{g_M} d\mathcal{M}.$$

Here  $d$  is the exterior derivative on cotangent bundle  $T^*M$ , which can be written as  $df = \sum_{i,j=1}^k g^{ij} \frac{\partial f}{\partial u^i} \frac{\partial}{\partial u^j} = \sum_{i=1}^k g^{ij} \nabla_{u^j} f$ , on the local chart  $(\Gamma, \phi)$  by the duality between tangent bundle  $TM$  and cotangent bundle  $T^*M$ . Then min-max theory tells us that

$$\lambda_1(M) = \inf_{\substack{\tilde{L} \subseteq H^1(M) \\ \dim \tilde{L}=2}} \sup_{u \in \tilde{L}} \frac{\mathcal{Q}_M(u, u)}{\int_M |u|^2 d\mathcal{M}}.$$

The reader could find more materials about this part in (Bérard, 1986, Chapter3). In our analysis, we will take  $M = S$  in any manifold-related content.

## 2.5. Additional Regularity Assumptions

For the ease of reference, we summarize the required assumptions in this subsection.

**Assumption 3 (Behavior of  $V$  beyond a compact set)** *Beyond a compact set,  $V$  satisfies the error bound inequality, i.e.  $\exists R_0 > 0$  such that  $\forall |x| \geq R_0$ <sup>5</sup>,*

$$|\nabla V(x)| \geq \nu_{eb} \text{dist}(x, S). \quad (13)$$

*Moreover,  $\Delta V := \text{div} \nabla V$  grows at most polynomially beyond a compact set, i.e.  $\forall |x| \geq R_0$ ,  $|\Delta V(x)| \leq C_g |x|^2$ . WLOG, we assume that  $R_0$  is sufficiently large so that for all  $x \in S$ ,  $|x| \leq R_0$ .*

**Assumption 4** *Let  $k$  be the dimension of  $S$ . We assume  $k \geq 1$ , i.e.  $S$  is not a singleton. Moreover, we assume  $S$  to have a bounded second fundamental form: On the local chart  $(\Gamma, \phi)$  of  $S$ ,  $\sup_{k+1 \leq l \leq d} \|\tilde{G}(l)\|_\infty < \infty$ , where  $\tilde{G}(l), l = k+1, \dots, d$  are defined in the end of Section 2.3.*

**Remark 2** *The boundedness condition of the second fundamental form is necessary. There exist curves that do not have a bounded curvature: Consider the ‘‘Tractrix Curve’’ parameterized by  $t$ , defined as  $(x(t), y(t)) = (a \sin t, a \ln(\tan(t/2)) + a \cos t)$ . Its second fundamental form is  $\kappa(t) = \frac{|x''(t)y'(t) - x'(t)y''(t)|}{(x'^2(t) + y'^2(t))^{\frac{3}{2}}} = \left| \frac{\tan t}{a} \right|$ , which is not bounded at the point  $(a, 0)$  or equivalently  $t = \frac{\pi}{2}$ .*

**Remark 3 (Dimension of  $S$ )** *We prove in Corollary 1 that  $S$  has no boundary and hence its dimension  $k$  is strictly smaller than  $d$ : If  $k = d$ , then  $V$  is a constant function, which contradicts with Assumption 3. Consequently, we focus on  $1 \leq k \leq d - 1$ . For  $k = 0$ , i.e.  $S$  degenerates to a singleton, the PI constant under a global PL condition has been recently established in (Chewi and Stromme, 2024). In this case, the PI constant is of order  $\Omega(\frac{1}{\epsilon})$  since  $V \in C^2$  and the PL condition implies that  $V$  is locally strongly convex near the unique minimum.*

**Remark 4 ( $\lambda_1(S)$  is non-trivial)** *The Poincaré inequality on Riemannian manifold has been well-studied. We refer readers (Hebey, 1999, Theorem 2.10) to the case of compact Riemannian manifold, showing that  $0 < \lambda_1(S) < \infty$ .*

### 3. Step 1: Reduction to the Neumann Eigenvalue Problem

We show that, when the temperature  $\epsilon$  is sufficiently small, the Poincaré constant of  $\mu_\epsilon$  can be lower bounded by the Neumann eigenvalue of the Laplacian operator on a closed domain  $U$  in  $\mathbb{R}^d$ . We will first list a few useful properties of the Log-PL<sup>o</sup> measures and then present our proof.

#### 3.1. Properties of the Log-PL<sup>o</sup> measures

The most important property of the Log-PL<sup>o</sup> measures is its uni-modality, and the manifold characterization of its mode, which we state in the following. Detailed proof can be found in Appendix C.1.

**Proposition 3 (Uni-modality)** *Under Assumptions 1 to 3, the all local minima of the potential function  $V$  are connected. Hence,  $V$  has only one connected global minima set  $S$ .*

The proof the following corollary is built on (Rebjock and Boumal, 2024, theorem 2.16). The purpose of this restatement is to explicitly exclude the possibility of boundary, which is not discussed in the previous work, and strengthen their local manifold structure to a global one using the above connectivity result. The absence of boundary significantly simplifies the eigenvalue problem (12).

**Corollary 1 (Manifold structure)**  *$S$  is a  $C^2$ -embedding submanifold of  $\mathbb{R}^d$  without boundary.*

We then characterize the properties of  $V$  in three different regions: (1) when  $x$  is close to the global minima set; (2) when  $x$  is close to some local maximum; (3) otherwise.

5. Clearly, eq. (13) implies Assumption 3'. Henceforce, we refer to Assumption 3 when Assumption 3' is required.

**Lemma 1** *Under Assumptions 1 to 3, the function  $V$  satisfies the following properties:*

- For any  $x \in \partial\mathcal{N}(S)$ , there exists some constant  $\delta_0 > 0$  such that  $\text{dist}(x, S) \geq \delta_0$ .
- Let  $X$  denote the set of all local maxima of the potential  $V$ . If  $X \neq \emptyset$ , there exists constants  $R_1 > 0$  and  $\mu^- > 0$  such that for all  $x \in \mathcal{N}(X) := \{x : \text{dist}(x, X) \leq R_1\}$ ,  $\nabla^2 V(x) \leq -\mu^- I_d$ .
- For  $x \notin \mathcal{N}(X) \cup \mathcal{N}(S)$ , there exists some constant  $g_0 > 0$  such that  $\|\nabla V(x)\| \geq g_0$ .

The following results are direct implications of our assumptions. The constants therein will be used in the statement of Step 1.

**Lemma 2** *Under Assumptions 1 to 3, the function  $V$  satisfies the following properties:*

- Since  $V \in \mathcal{C}^2$ , there exists some constant  $L$  such that  $\|\nabla^2 V(x)\| \leq L$  for all  $x \in \mathcal{N}(S)$ .
- Recall  $R_0$  from Assumption 3. Since  $V \in \mathcal{C}^2$ , there exists some constant  $M_\Delta > 0$  such that  $|\Delta V(x)| \leq M_\Delta$  for  $\|x\| \leq R_0$ . WLOG, assume  $M_\Delta \geq d\mu^-$ . Otherwise, simply set  $\mu^- = M_\Delta/d$ .
- There exists a neighborhood  $\mathcal{N}'(S)$  of  $S$  on which the error bound inequality eq. (13) holds with a constant  $\nu'$  (Rebjock and Boumal, 2024). For simplicity, we assume  $\mathcal{N}'(S) = \mathcal{N}(S)$ , and  $\nu' = \nu$  since otherwise we can set  $\mathcal{N}(S) = \mathcal{N}(S) \cap \mathcal{N}'(S)$  and  $\nu = \min\{\nu, \nu'\}$  in Assumption 1. All derivations remain unchanged.

### 3.2. Proof Sketch of Step 1

Our proof is built on the Lyapunov approach described in Proposition 1. To meet the requirements in eq. (8), for any  $x$  outside of a closed domain  $U = S^{\sqrt{C}\epsilon}$ , we need (i) a lower bound for the gradient norm  $\|\nabla V(x)\|$  and (ii) an upper bound for the Laplacian  $|\Delta V(x)|$ .

**(i) Lower bound of gradient norm  $\|\nabla V(x)\|$ .** There are four situations: (a) When  $x$  is outside of a compact set, we utilize the error bound inequality in Assumption 3; (b) When  $x \in \mathcal{N}(S)$ , we utilize the error bound inequality in Lemma 2; (c) When  $x$  is close to a local maximum, it suffices to use the trivial bound  $\|\nabla V(x)\| \geq 0$ ; (d) Otherwise, we utilize the third property in Lemma 1.

**(ii) Upper bound of Laplacian  $|\Delta V(x)|$ .** To bound the Laplacian  $|\Delta V(x)|$ , we partition  $\mathbb{R}^d \setminus S^{\sqrt{C}\epsilon}$  into two of subdomains  $\Xi_1$  and  $\Xi_2$  and sketch the treatments: For  $x \in \Xi_1 = \{x : 2R_0 \leq \|x\|\}$ , we bound the Laplacian term  $|\Delta V(x)|$  by utilizing the growth of  $\Delta V$  from Assumption 3, in which  $R_0$  is defined. For  $x \in \Xi_2 = \{x : \sqrt{C}\epsilon \leq \text{dist}(x, S) \text{ and } \|x\| \leq 2R_0\}$ , there are three situations: (1)  $x$  is close to some local maximum; (2)  $x \in \mathcal{N}(S)$ ; (3) otherwise. We treat these three situations separately. The power index  $\frac{1}{2}$  in  $\Xi_2$  is the largest value such that the positivity of  $\sigma$  in eq. (8) still holds when  $x \in \mathcal{N}(S)$ . Here  $C$  is some constant independent of  $\epsilon$  defined in Lemma 3. Note that for each subdomain  $\Xi_i, i = 1, 2$ , there will be a corresponding value of  $\sigma_i$  and we set  $\sigma = \inf_{i \in \{1, 2\}} \sigma_i$ . Besides, we estimate  $b$  by restricting  $x$  on  $U = S^{\sqrt{C}\epsilon}$ .

### 3.3. Establishing the bound on $b$ and $\sigma$ .

The following lemma formally states the above results, whose proof is in Appendix E.2.

**Lemma 3** *Suppose that Assumptions 1 to 4 hold. Define a constant  $C = \frac{4M_\Delta}{\nu^2}$ . Suppose that  $\epsilon$  is sufficiently small such that  $\epsilon \leq \min\{\frac{\nu_{eb}^2 R_0^2}{64Cg}, \frac{\delta_0^2}{C}, \frac{g_0^2}{4M_\Delta}\}$ . Choose  $U = S^{\sqrt{C}\epsilon}$ . Equation (8) holds with*

$$\sigma = \min \left\{ \frac{\nu_{eb}^2 R_0^2}{128} \frac{1}{\epsilon^2}, \frac{d\mu^-}{2\epsilon} \right\}, \text{ and } b = \sigma + \frac{M_\Delta}{2\epsilon}. \quad (14)$$

### 3.4. Perturbation near the optimal set

Applying the estimates from Lemma 3 to Proposition 1, we have reduced the estimation for the Poincaré constant of the Gibbs measure  $\mu_\epsilon$  to the estimation for the one of the truncated Gibbs measure  $\mu_{\epsilon,U}$ . Unfortunately, the latter remains elusive. In this section, we further reduce the estimation of  $\rho_{\epsilon,U}$  to the estimation of the Neumann eigenvalue of the Laplacian operator on  $U$ .

Since all the points in the subdomain  $U$  are sufficiently close to the optimal set  $S$ , one can utilize the Taylor expansion of the potential  $V$  to show that the density function of the truncated Gibbs measure  $\mu_{\epsilon,U}$  is an  $\epsilon$ -perturbation of the uniform density function on  $U$ . We can utilize the perturbation principle in Proposition 2 and the smoothness of  $V$  (see the first point in Lemma 2) to derive the following result.

**Lemma 4** *Suppose that the assumptions and requirements in Lemma 3 hold. If the uniform measure  $\mu_U$  satisfies  $PI(\rho_U)$ , then the truncated Gibbs measure  $\mu_{\epsilon,U}$  also satisfies  $PI(\rho_{\epsilon,U})$  with*

$$\exp\{\bar{C}\}\rho_{\epsilon,U} \geq \rho_U = \lambda_1^n(U),$$

where  $\bar{C} = 4LC$  and  $\lambda_1^n(U)$  is the Neumann eigenvalue on  $U$ .

Combining Lemmas 3 and 4, we obtain the main conclusion of this section.

**Theorem 2** *Suppose that the requirements in Lemma 3 hold and further suppose that  $\epsilon$  is sufficiently small so that  $\sigma = \frac{d\mu^-}{2\epsilon}$  in eq. (14). By choosing  $U = S^{\sqrt{C}\epsilon}$ , we have  $\rho_{\mu_\epsilon} \geq \frac{1}{2} \frac{d\mu^-}{d\mu^- + M_\Delta} \exp(-\bar{C})\lambda_1^n(U)$ .*

## 4. Step 2: Stability Analysis of the Neumann Eigenvalue

In the previous section, we reduce the estimation of the Poincaré constant of  $\mu_\epsilon$  to  $\lambda_1^n(U)$ , the Neumann eigenvalue of the Laplacian operator on the subdomain  $U$ . In this section, we justify the choice of  $U = S^{\sqrt{C}\epsilon}$ . Recall that  $S$  is a  $C^2$  embedding submanifold. Hence,  $U$  matches the tubular neighborhood of  $S$ , a special kind of *neighborhood in the ambient space*, as described in Theorem 3. For the simplicity of notation, we denote  $\tilde{\epsilon} = \sqrt{C}\epsilon$  in this section.

### 4.1. Tubular Neighborhood of a $C^2$ Embedding Submanifold


As an important property of embedding submanifold in  $\mathbb{R}^d$ , let us introduce the tubular neighborhood theorem (see Guillemin (1974), Page 69).

**Theorem 3 (Tubular neighborhood theorem)** *Let  $M$  be a compact  $C^2$  embedding submanifold in  $\mathbb{R}^d$ . Let  $T(\tilde{\epsilon}) = M^{\tilde{\epsilon}} = \{y \in \mathbb{R}^d : |y - m| \leq \tilde{\epsilon}, m \in M\}$  be the tubular neighborhood of  $M$ . There exists a positive number  $\tilde{\epsilon}_{\text{TN}}$ , such that for all  $0 < \tilde{\epsilon} \leq \tilde{\epsilon}_{\text{TN}}$ , one has (1) each  $y \in T(\tilde{\epsilon})$  possesses a unique closest point  $\pi(y) \in M$ ; (2) the projection map  $\pi : T(\tilde{\epsilon}) \rightarrow M$  is a submersion. That is to say, the linear map  $D_y\pi : T_yT(\tilde{\epsilon}) \rightarrow T_{\pi(y)}M$  is surjective at each point  $y \in T(\tilde{\epsilon})$ .*

There is a more concrete representation for the tubular neighborhood, which we adopt in the rest of the paper: For any  $y \in T(\tilde{\epsilon})$ ,  $y$  can be written as  $y = m + v$ , where  $m$  is a point on  $M$  and  $v \perp M$  at  $m$  with  $|v| \leq \tilde{\epsilon}$ , and the map  $y \rightarrow (m, v)$  is a diffeomorphism. More precisely, under the local chart  $(\Gamma, \phi)$ , the diffeomorphism  $y \rightarrow (m, v)$  can be written as

$$y(u, r) = \mathcal{M}(u) + \sum_{l=k+1}^d r^l \mathcal{N}_l(u), \quad u \in \Gamma \subset \mathbb{R}^k, \quad (r^{k+1}, \dots, r^d) \in B(\tilde{\epsilon}) \subset \mathbb{R}^{d-k}. \quad (15)$$

Here  $\mathcal{M}(u)$  is local coordinate representation of the including map  $i_M : M \rightarrow \mathbb{R}^d$  in (9),  $\mathcal{N}_{k+1}, \dots, \mathcal{N}_d : \Gamma \rightarrow \mathbb{R}^d$  are  $d - k$  normal vector fields on  $M$  which are also orthogonal to each other, and  $B(\tilde{\epsilon})$



**Figure 1:** The circle in (a) can be represented using two local charts (blue and green). Using the tubular neighborhood theorem, in a local region of  $U$  (outlined with the red dashed line) we transform the uniform measure to a pair of decoupled measures on the tangent and normal directions. (a) Uniform measure  $\mu_U$  (over  $(x, y)$ ) under the Cartesian coordinate  $(x, y)$ ; (b) Uniform measure  $\mu_U$  (over  $(x, y)$ ) under the local coordinate  $(\theta, r)$ ; (c) Uniform measure (over  $(\theta, r)$ ) under the local coordinate  $(\theta, r)$ . Importantly, when the radius of the tubular neighborhood is small, the densities in (b) and (c) point-wisely control each other.

denotes ball with radius  $\tilde{\epsilon}$  in  $\mathbb{R}^{d-k}$ . We refer readers to Appendix D for more details about these vector fields on the local chart  $(\Gamma, \phi)$ . For brevity, we also denote  $r = (r^{k+1}, \dots, r^d)$  with  $|r| \leq \tilde{\epsilon}$ .

## 4.2. Stability of the Neumann Eigenvalue on the Tubular Neighborhood

We now focus on the stability of  $\lambda_1^n(T(\tilde{\epsilon}))$  w.r.t.  $\tilde{\epsilon}$ , where  $T(\tilde{\epsilon})$  is a tubular neighborhood of  $S$ . Given the special structure of  $T(\tilde{\epsilon})$ , our idea is to exploit the tensorization of the Poincaré inequality.

**Proposition 4 (Proposition 4.3.1 in (Bakry et al., 2014))** *Let  $(E_1, \mu_1)$  and  $(E_2, \mu_2)$  be two probability spaces with measure  $\mu_1$  and  $\mu_2$ , and they satisfy PI with constants  $C_1$  and  $C_2$  respectively. Then the product space  $(E_1 \times E_2, \mu_1 \times \mu_2)$  satisfies a PI with constant  $C = \max\{C_1, C_2\}$ .*

To utilize the above theorem, recall the min-max variational principle of PI in Section 2.4, which consists of the  $L^2$  norm and the  $H^1$  norm on  $T(\tilde{\epsilon})$ . To bound these integrals, we show that the uniform measure on  $T(\tilde{\epsilon})$  can be decomposed as the product of a pair of decoupled measures on the manifold  $S$  and the subspace of the normal coordinates  $r$ , up to an  $\mathcal{O}(\tilde{\epsilon})$  perturbation:

- We decompose the integral in  $T(\tilde{\epsilon})$  as the integral in the product space  $S \times B(\tilde{\epsilon})$  with an additional factor of order  $1 + \mathcal{O}(\tilde{\epsilon})$ . This is possible since for any  $x \in S$ , an  $\epsilon_1$ -neighborhood of  $x$  under the topology of  $\mathbb{R}^d$  can be viewed, up to a diffeomorphism, as the product space  $B^S(x, \epsilon_2) \times B(\epsilon_3)$ . Here  $B^S(x, \epsilon_2) \subset S$  is a ball in  $S$  with center  $x$  and radius  $\epsilon_2$  (defined according to the geodesic distance on  $S$ ) and  $B(\epsilon_3)$  is a ball in  $d-k$  normal directions of  $S$  at  $x$  with radius  $\epsilon_3$ . Equivalently, we turn the uniform measure on  $T(\tilde{\epsilon})$  to the product of the volume measure induced by the including map  $i_S$  on  $S$  and the uniform measure on  $B(\tilde{\epsilon})$ .
- With the above decomposition, we show that both  $L^2$  and  $H^1$  norm on  $T(\tilde{\epsilon})$  are bounded by their counterparts on the the product space  $S \times B(\tilde{\epsilon})$  with an  $\tilde{\epsilon}$  perturbation.

With the above derivation, we turn our focus to the PIs of the two decoupled measures. The PI constant of the volume measure on  $S$  is inherent to  $S$  and is *temperature-independent*. The PI constant of the measure on the normal coordinates has been explicitly calculated in the literature.

**Proposition 5** (PI for Lebesgue measure on a ball, (Evans, 2010, Page 293, Theorem 2)) Let  $B(\tilde{\epsilon}) \subseteq \mathbb{R}^{d-k}$  be a ball with radius  $\tilde{\epsilon}$ . Let  $\mu_{B(\tilde{\epsilon})}$  be the uniform measure over  $B(\tilde{\epsilon})$ . There exists a constant  $\tilde{C}$  depending only on the dimension  $(d-k)$  such that the PI constant  $\rho_{\mu_{B(\tilde{\epsilon})}} \geq \frac{1}{\tilde{C}\tilde{\epsilon}}$ .

Combining with Proposition 4, we know  $\lambda_1^n(B(\tilde{\epsilon}))$  is dominated by  $\lambda_1(S)$  when  $\tilde{\epsilon}$  is sufficiently small, and hence,  $\lambda_1^n(T(\tilde{\epsilon}))$  is determined by  $\lambda_1(S)$ . We now make the above reasoning rigorous.

**Lemma 5** (Weyl, 1939) Let  $\varphi : T(\tilde{\epsilon}) \rightarrow \mathbb{R}$  be an integrable function, then we have

$$\int_{T(\tilde{\epsilon})} \varphi(y) dy = \int_S \left\{ \int_{B(\tilde{\epsilon})} \varphi(y(u, r)) \left| \det \left( I_k + \sum_{l=k+1}^d r^l \tilde{G}(l) \right) \right| dr^{k+1} \dots dr^d \right\} dM,$$

where the variable  $y$  on the right hand side uses the expression of the local coordinate (15),  $I_k$  is  $k \times k$  identity matrix and  $\tilde{G}(l), l = k+1, \dots, d$  are defined in the end of Section 2.3.

To relate  $\lambda_1^n(U)$  to  $\lambda_1(S)$  through their min-max variational principles, we need the following expression of the gradient under change of variables.

**Lemma 6** Let  $\varphi(y) \in W^{1,2}(T(\tilde{\epsilon}))$ , then, on each local chart  $(\Gamma, \phi)$ , we have in the weak sense

$$\nabla_{(u,r)} \varphi(y(u, r)) = \nabla_y \varphi \cdot \left( \left[ \frac{\partial M}{\partial u^1}, \dots, \frac{\partial M}{\partial u^k}, \mathcal{N}_{k+1}, \dots, \mathcal{N}_d \right] \begin{bmatrix} I_k + \sum_{l=k+1}^d r^l \tilde{G}(l) & 0 \\ 0 & I_{d-k} \end{bmatrix} \right).$$

We are now ready to state the estimation of  $\lambda_1^n(U)$ . Note that the choice of  $U = T(\epsilon)$  can also be regarded as a domain expansion, so the following result is a stability analysis. However, one should keep in mind that the domain expansion is performed under the topology of the ambient space  $\mathbb{R}^d$ , so  $U$  and  $S$  do not have the same dimension.

**Proposition 6** Suppose that Assumptions 1 to 4 hold, then we have the non-asymptotic estimates of  $\lambda_1^n(T(\tilde{\epsilon}))$  based on  $\lambda_1(S)$

$$\lambda_1(S)(1 - B\tilde{\epsilon}) \leq \lambda_1^n(T(\tilde{\epsilon})) \leq \lambda_1(S)(1 + B\tilde{\epsilon}),$$

for some constant  $B = B(d, k, \tilde{G}(l)) > 0$  when  $\tilde{\epsilon}$  is small enough.

## 5. Poincaré Inequality for the Log-PL<sup>o</sup> Measure

We now combine the results in Sections 3 and 4 to conclude the Poincaré inequality for the Gibbs measure  $\mu_\epsilon$ . Please find the proof in Appendix G.1.

**Theorem 4** Suppose that the requirements in Lemma 3 hold. Suppose that the temperature  $\epsilon$  in addition satisfies  $\epsilon \leq \min \left\{ \tilde{\epsilon}_{\text{TN}}, \frac{1}{C} \left( \frac{1}{2B} \right)^2 \right\}$ , where  $\tilde{\epsilon}_{\text{TN}}$  and  $B$  appear in Theorem 3 and Proposition 6 respectively. Recall that  $\lambda_1(S)$  denotes the eigenvalue of the Laplacian-Beltrami operator (Definition 4). Define a constant  $C_P = \frac{1}{4} \frac{d\mu^-}{d\mu^- + M_\Delta} \exp(-4LC)$ . We have  $\rho_{\mu_\epsilon} \geq C_P \lambda_1(S)$ .

By noting that  $\lambda_1(S)$  is an inherent property of the optimal set  $S$  and hence the potential function  $V$ , we reach the target conclusion, i.e.  $\rho_{\mu_\epsilon} = \Omega(1)$ . With the connection between the convergence of Langevin dynamics and the Poincaré inequality, we have that for a PL<sup>o</sup> potential, the Langevin dynamics converges at the rate  $\tilde{O}(\frac{1}{\epsilon})$ . The convergence of discrete-time algorithms (in the sense of Rényi-divergence) like LMC can be readily derived by combining our result with (Chewi et al., 2024, Theorem 7). To the best of our knowledge, this is the first work to study the PI constant for potentials with non-isolated minima in the low temperature region. Interestingly, for the non-log-concave Log-PL<sup>o</sup> measures, we still yield an  $\Omega(1)$  lower bound for the PI constant.

**Conclusion and future work** We study the Poincaré constant of the Log-PL<sup>o</sup> measures as a template to understand the convergence behavior of stochastic dynamics on potentials with non-isolated minima. We relate the corresponding Poincaré constant to the spectral property of the Laplacian-Beltrami operator on the optimal set  $S$ , and establish a *temperature-independent* lower bound. Our next steps are (1) improving PI to LSI, (2) relaxing the  $C^2$  manifold to the  $C^1$  case, and (3) studying the non-isotropic noise case.

## References

- Svante. Arrhenius. On the reaction velocity of the inversion of cane sugar by acids. In *Selected readings in chemical kinetics*, pages 31–35. Elsevier, 1967.
- Dominique Bakry and Michel Émery. Diffusions hypercontractives. In *Séminaire de Probabilités XIX 1983/84: Proceedings*, pages 177–206. Springer, 2006.
- Dominique Bakry, Franck Barthe, Patrick Cattiaux, and Arnaud Guillin. A simple proof of the Poincaré inequality for a large class of probability measures. *Electronic Communications in Probability*, 13(none):60 – 66, 2008a. doi: 10.1214/ECP.v13-1352. URL <https://doi.org/10.1214/ECP.v13-1352>.
- Dominique Bakry, Patrick Cattiaux, and Arnaud Guillin. Rate of convergence for ergodic continuous markov processes: Lyapunov versus poincaré. *Journal of Functional Analysis*, 254(3): 727–759, 2008b.
- Dominique Bakry, Ivan Gentil, Michel Ledoux, et al. *Analysis and geometry of Markov diffusion operators*, volume 103. Springer, 2014.
- Gerard Ben Arous, Reza Gheissari, and Aukosh Jagannath. High-dimensional limit theorems for sgd: Effective dynamics and critical scaling. *Advances in Neural Information Processing Systems*, 35:25349–25362, 2022.
- Pierre H. Bérard. *Spectral Geometry: Direct and Inverse Problems*. 1986. URL <https://api.semanticscholar.org/CorpusID:117992829>.
- Nicolas Boumal. *An Introduction to Optimization on Smooth Manifolds*. Cambridge University Press, 2023.
- Anton Bovier, Michael Eckhoff, Véronique Gayrard, and Markus Klein. Metastability in reversible diffusion processes. i. sharp asymptotics for capacities and exit times. *J. Eur. Math. Soc.(JEMS)*, 6(4):399–424, 2004.
- Patrick Cattiaux and Arnaud Guillin. Hitting times, functional inequalities, lyapunov conditions and uniform ergodicity. *Journal of Functional Analysis*, 272(6):2361–2391, 2017. ISSN 0022-1236. doi: <https://doi.org/10.1016/j.jfa.2016.10.003>. URL <https://www.sciencedirect.com/science/article/pii/S0022123616303068>.
- Patrick Cattiaux, Arnaud Guillin, and Li-Ming Wu. A note on talagrand’s transportation inequality and logarithmic sobolev inequality. *Probability theory and related fields*, 148:285–304, 2010.
- Jeff Cheeger. A lower bound for the smallest eigenvalue of the laplacian. *Problems in analysis*, 625 (195-199):110, 1970.

- Yuansi Chen. An almost constant lower bound of the isoperimetric coefficient in the kls conjecture. *Geometric and Functional Analysis*, 31:34–61, 2020. URL <https://api.semanticscholar.org/CorpusID:227209486>.
- Yuansi Chen. An almost constant lower bound of the isoperimetric coefficient in the kls conjecture. *Geometric and Functional Analysis*, 31:34–61, 2021.
- Sinho Chewi and Austin J Stromme. The ballistic limit of the log-sobolev constant equals the Polyak-Łojasiewicz constant. November 2024.
- Sinho Chewi, Murat A Erdogdu, Mufan Li, Ruoqi Shen, and Matthew S Zhang. Analysis of langevin monte carlo from poincare to log-sobolev. *Foundations of Computational Mathematics*, pages 1–51, 2024.
- Y Cooper. The critical locus of overparameterized neural networks. *arXiv preprint arXiv:2005.04210*, 2020.
- Yaim Cooper. The loss landscape of overparameterized neural networks. *arXiv preprint arXiv:1804.10200*, 2018.
- E. Brian Davies. *Spectral Theory and Differential Operators*. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 1995.
- Felix Draxler, Kambis Veschgini, Manfred Salmhofer, and Fred Hamprecht. Essentially no barriers in neural network energy landscape. In *International conference on machine learning*, pages 1309–1318. PMLR, 2018.
- Ronen Eldan. Thin shell implies spectral gap up to polylog via a stochastic localization scheme. *Geometric and Functional Analysis*, 23:532 – 569, 2013. URL <https://api.semanticscholar.org/CorpusID:253637768>.
- Murat A Erdogdu and Rasa Hosseinzadeh. On the convergence of langevin monte carlo: The interplay between tail growth and smoothness. In *Conference on Learning Theory*, pages 1776–1822. PMLR, 2021.
- Lawrence C. Evans. *Partial differential equations*. American Mathematical Society, 2010.
- Henry Eyring. The activated complex in chemical reactions. *The Journal of Chemical Physics*, 3(2):107–115, 1935.
- Benjamin Fehrman, Benjamin Gess, and Arnulf Jentzen. Convergence rates for the stochastic gradient descent method for non-convex objective functions. *Journal of Machine Learning Research*, 21(136):1–48, 2020.
- C Daniel Freeman and Joan Bruna. Topology and geometry of half-rectified network optimization. In *5th International Conference on Learning Representations, ICLR 2017*, 2017.
- Mark I Freidlin and Alexander D Wentzell. *Random Perturbations of Dynamical Systems*, volume 260. Springer Science & Business Media, 2012.
- Timur Garipov, Pavel Izmailov, Dmitrii Podoprikhin, Dmitry P Vetrov, and Andrew G Wilson. Loss surfaces, mode connectivity, and fast ensembling of dnns. *Advances in neural information processing systems*, 31, 2018.

- Véronique Gayraud, Anton Bovier, and Markus Klein. Metastability in reversible diffusion processes ii: Precise asymptotics for small eigenvalues. *Journal of the European Mathematical Society*, 7(1):69–99, 2005.
- Alan Pollack Victor Guillemin. *Differential Topology*. 1974.
- Emmanuel Hebey. *Nonlinear analysis on manifolds: Sobolev spaces and inequalities*. 1999. URL <https://api.semanticscholar.org/CorpusID:118747316>.
- Ravi Kannan, László Miklós Lovász, and Miklós Simonovits. Isoperimetric problems for convex bodies and a localization lemma. *Discrete & Computational Geometry*, 13:541–559, 1995. URL <https://api.semanticscholar.org/CorpusID:14881695>.
- Hamed Karimi, Julie Nutini, and Mark Schmidt. Linear convergence of gradient and proximal-gradient methods under the polyak-łojasiewicz condition. In *Machine Learning and Knowledge Discovery in Databases: European Conference, ECML PKDD 2016, Riva del Garda, Italy, September 19-23, 2016, Proceedings, Part I 16*, pages 795–811. Springer, 2016.
- Guy Katriel. Mountain pass theorems and global homeomorphism theorems. In *Annales de l’Institut Henri Poincaré C, Analyse non linéaire*, volume 11, pages 189–209. Elsevier, 1994.
- Kenji Kawaguchi and Leslie Kaelbling. Elimination of all bad local minima in deep learning. In *International Conference on Artificial Intelligence and Statistics*, pages 853–863. PMLR, 2020.
- Hendrik Anthony Kramers. Brownian motion in a field of force and the diffusion model of chemical reactions. *physica*, 7(4):284–304, 1940.
- Rohith Kuditipudi, Xiang Wang, Holden Lee, Yi Zhang, Zhiyuan Li, Wei Hu, Rong Ge, and Sanjeev Arora. Explaining landscape connectivity of low-cost solutions for multilayer nets. *Advances in neural information processing systems*, 32, 2019.
- John M. Lee. *Introduction to Smooth Manifolds*. 2020.
- Yin Tat Lee and Santosh S. Vempala. The kannan-lovász-simonovits conjecture, 2018. URL <https://arxiv.org/abs/1807.03465>.
- Yin Tat Lee and Santosh S Vempala. Eldan’s stochastic localization and the kls conjecture: Isoperimetry, concentration and mixing. *Annals of Mathematics*, 199(3):1043–1092, 2024.
- Eitan Levin, Joe Kileel, and Nicolas Boumal. The effect of smooth parametrizations on nonconvex optimization landscapes. *Mathematical Programming*, pages 1–49, 2024.
- Qianxiao Li, Cheng Tai, and E Weinan. Stochastic modified equations and adaptive stochastic gradient algorithms. In *International Conference on Machine Learning*, pages 2101–2110. PMLR, 2017.
- Xiang Li, Zebang Shen, Liang Zhang, and Niao He. A hessian-aware stochastic differential equation for modelling sgd. *arXiv preprint arXiv:2405.18373*, 2024.
- Zhiyuan Li, Tianhao Wang, and Sanjeev Arora. What happens after sgd reaches zero loss?-a mathematical framework. In *10th International Conference on Learning Representations, ICLR 2022*, 2022.

- Shiyu Liang, Ruoyu Sun, and Jason D Lee. Adding one neuron can eliminate all bad local minima. *Advances in Neural Information Processing Systems*, 31, 2018a.
- Shiyu Liang, Ruoyu Sun, Yixuan Li, and Rayadurgam Srikant. Understanding the loss surface of neural networks for binary classification. In *International Conference on Machine Learning*, pages 2835–2843. PMLR, 2018b.
- Zhanran Lin, Puheng Li, and Lei Wu. Exploring neural network landscapes: Star-shaped and geodesic connectivity. *arXiv preprint arXiv:2404.06391*, 2024.
- Chaoyue Liu, Libin Zhu, and Mikhail Belkin. Loss landscapes and optimization in over-parameterized non-linear systems and neural networks. *Applied and Computational Harmonic Analysis*, 59:85–116, 2022.
- Stanislaw Lojasiewicz. A topological property of real analytic subsets. *Coll. du CNRS, Les équations aux dérivées partielles*, 117(87-89):2, 1963.
- Georg Menz and André Schlichting. Poincaré and logarithmic Sobolev inequalities by decomposition of the energy landscape. *The Annals of Probability*, 42(5):1809 – 1884, 2014. doi: 10.1214/14-AOP908. URL <https://doi.org/10.1214/14-AOP908>.
- John Milnor and James D. Stasheff. *Characteristic Classes. (AM-76), Volume 76*. Princeton University Press, Princeton, 1974. ISBN 9781400881826. doi: doi:10.1515/9781400881826. URL <https://doi.org/10.1515/9781400881826>.
- Quynh Nguyen. On connected sublevel sets in deep learning. In *International conference on machine learning*, pages 4790–4799. PMLR, 2019.
- Samet Oymak and Mahdi Soltanolkotabi. Toward moderate overparameterization: Global convergence guarantees for training shallow neural networks. *IEEE Journal on Selected Areas in Information Theory*, 1(1):84–105, 2020.
- Courtney Paquette, Elliot Paquette, Ben Adlam, and Jeffrey Pennington. Homogenization of sgd in high-dimensions: Exact dynamics and generalization properties. *arXiv preprint arXiv:2205.07069*, 2022.
- Boris Teodorovich Polyak. Gradient methods for minimizing functionals. *Zhurnal vychislitel'noi matematiki i matematicheskoi fiziki*, 3(4):643–653, 1963.
- Maxim Raginsky, Alexander Rakhlin, and Matus Telgarsky. Non-convex learning via stochastic gradient langevin dynamics: a nonasymptotic analysis. In *Conference on Learning Theory*, pages 1674–1703. PMLR, 2017.
- Quentin Rebjock and Nicolas Boumal. Fast convergence to non-isolated minima: four equivalent conditions for  $c^2$  functions. *Mathematical Programming*, pages 1–49, 2024.
- Itay Safran and Ohad Shamir. On the quality of the initial basin in overspecified neural networks. In *International Conference on Machine Learning*, pages 774–782. PMLR, 2016.
- Levent Sagun, Leon Bottou, and Yann LeCun. Eigenvalues of the hessian in deep learning: Singularity and beyond. *arXiv preprint arXiv:1611.07476*, 2016.
- Levent Sagun, Utku Evci, V Ugur Guney, Yann Dauphin, and Leon Bottou. Empirical analysis of the hessian of over-parametrized neural networks. *arXiv preprint arXiv:1706.04454*, 2017.

- L.W. Tu. *An Introduction to Manifolds*. Universitext. Springer New York, 2010. ISBN 9781441973993. URL <https://books.google.com.hk/books?id=br1KngEACAAJ>.
- Luca Venturi, Afonso S Bandeira, and Joan Bruna. Spurious valleys in two-layer neural network optimization landscapes. *arXiv preprint arXiv:1802.06384*, 2018.
- Hermann Weyl. On the volume of tubes. *American Journal of Mathematics*, 61:461, 1939. URL <https://api.semanticscholar.org/CorpusID:124362885>.
- Andre Wibisono. Sampling as optimization in the space of measures: The langevin dynamics as a composite optimization problem. In *Conference on Learning Theory*, pages 2093–3027. PMLR, 2018.
- Stephan Wojtowytsch. Stochastic gradient descent with noise of machine learning type part ii: Continuous time analysis. *Journal of Nonlinear Science*, 34(1):16, 2024.
- Junchi Yang, Negar Kiyavash, and Niao He. Global convergence and variance reduction for a class of nonconvex-nonconcave minimax problems. *Advances in Neural Information Processing Systems*, 33:1153–1165, 2020.
- Yuchen Zhang, Percy Liang, and Moses Charikar. A hitting time analysis of stochastic gradient langevin dynamics. In *Conference on Learning Theory*, pages 1980–2022. PMLR, 2017.

## Appendix A. Related Work

Poincaré inequality is a crucial topic in domains like probability, analysis, geometry and so on. We categorize the results according to the property of the energy.

- When the potential function  $V$  is strongly convex, the famous Bakry-Emery criterion ensures that the PI constant is of order  $\Theta(\frac{1}{\epsilon})$  (Bakry and Émery, 2006). See for more detailed discussion in the book (Bakry et al., 2014).

- When the potential function  $V$  is convex, there two prominent strategies to study the PI constant: the Lyapunov function approach (Bakry et al., 2008a) and the approach initiated by Cheeger (1970) which relates the PI constant to the isoperimetric constant of the target measure and the KLS conjecture (Kannan et al., 1995; Lee and Vempala, 2024).

Following (Bakry et al., 2008b), Bakry et al. (2008a) reduce the Poincaré inequality on  $\mathbb{R}^d$  into a small compact region  $U$  if  $V$  satisfies a Lyapunov condition. Convex functions that are exponentially integrable is proved to satisfy this condition and hence the corresponding log-concave measure satisfies Poincaré inequality. However, they do not consider the low temperature region and if we naively utilize the Holley-Stroock perturbation principle to derive the Poincaré inequality in the said compact region  $U$ , the resulting Poincaré constant on  $\mathbb{R}^d$  is  $\Omega(\exp(\frac{1}{\epsilon}))$ .

Another important research line is Cheeger’s inequality (Cheeger, 1970), which relates the Poincaré constant on a compact set with the Cheeger constant, describing the geometrical property of the set. Later on, the KLS conjecture extends the Cheeger constant in Euclidean space with log-concave measure in (Kannan et al., 1995). We recommend a good survey (Lee and Vempala, 2018) and reference therein for more precise introduction and recent progress. We mention a very important method to prove this conjecture — stochastic localization, which starts from Eldan’s work in (Eldan, 2013), and then generalized by Lee and Vempala (2024) and Chen (2020). We note that KLS conjecture pays more attention to the independence of dimension. However, it also helps to derive the relationship of temperature and Poincaré constant for log-concave measure. In fact, if we combine the Lyapunov function approach with the KLS conjecture, we can prove that for a log-concave measure, the Poincaré constant remains temperature-independent in the low temperature regime.

- The convergence behavior of the Langevin dynamics on a non-log-concave measure is also a crucial research problem in various domains. In particular, here we focus on the case where the potential function  $V$  has at least two separated (local) minima. As mentioned in the introduction, in this case, there is a two time-scale phenomenon in the convergence behavior in the low temperature region. The exponential dependence on the inverse temperature in the slow scale is classically known as the Arrhenius law (Arrhenius, 1967), which can be proved for example by the Freidlin-Wentzell theory on large deviation (Freidlin and Wentzell, 2012). With additional assumptions that  $V$  is a Morse function and its saddle points has exactly one negative eigenvalue, this subexponential factor in the convergence behavior is captured in (Eyring, 1935; Kramers, 1940) and rigorously proved by (Bovier et al., 2004; Gayraud et al., 2005) through potential theory. Menz and Schlichting (2014) study the same problem, but through the functional inequality perspective. Following a two-scale approach, Menz and Schlichting (2014) split the variance (the term  $\text{Var}_\mu(f)$  in eq. (4)) into local variances and coarse-grained variances. The estimations on both variances can be combined together to obtain the global Poincaré inequality.

## Appendix B. Poincaré constant for log-concave measure in the low temperature regime

Suppose the potential function  $V$  is convex and WLOG  $V^* = 0$ . Suppose that  $V$  is exponentially integrable for  $\epsilon = 1$ , i.e.  $\int_{\mathbb{R}^d} \exp(-V(x)) dx < \infty$ . We prove that the Poincaré constant  $\rho_{\mu_\epsilon}$  is  $\Omega(1)$ . Our proof is a combination of the Lyapunov function approach (Bakry et al., 2008a) and the KLS conjecture (Lee and Vempala, 2024).

Recall point (2) in (Bakry et al., 2008a, Lemma 2.2) which states that under the integrability assumption, one has

$$\exists \alpha > 0, R > 0, \text{ s.t. } \forall |x| \geq R, V(x) - V(0) \geq \alpha|x|.$$

Moreover, let us choose a Lyapunov function as  $W(x) = \exp(\tilde{W}(\gamma|x|))$  where  $\gamma = \frac{\alpha}{3}$

$$\tilde{W}(z) = \begin{cases} z & z \geq R \\ -\frac{12}{R^2}z^3 + \frac{28}{R}z^2 - 19z + 4R & R/2 \leq z \leq R \\ 0 & z \leq R/2 \end{cases}$$

We can compute for  $|x| \geq R$

$$\epsilon^{-1} \mathcal{L}W(x) = \gamma \left( \frac{n-1}{|x|} + \gamma - \frac{x \cdot \nabla V(x)}{\epsilon|x|} \right) W(x).$$

Note that  $W \in C^2$  and  $W(x) \geq 1$  for all  $x \in \mathbb{R}^d$ . Recall Proposition 1. Set  $\theta = \frac{\alpha}{\epsilon} - \gamma - \frac{(d-1)}{R}$  and set

$$b = \theta + \sup_{\|x\| \leq R} \epsilon^{-1} \mathcal{L}W(x).$$

With with parameters  $\theta, b, U = B(0, R)$ ,  $W$  is a valid Lyapunov function. Moreover, it can be easily checked that  $b \leq \theta + \frac{C_1}{\epsilon} + C_2$  for some temperature-independent constants  $C_1$  and  $C_2$  since both  $V$  and  $W$  are  $C^2$  and  $W$  is a constant for  $\|x\| \leq R/2$ .

Finally, the Poincaré constant for the truncated Gibbs measure can be bounded using the KLS conjecture: According to (Lee and Vempala, 2024, Theorem 13), the Cheeger constant of a log-concave measure can be lower bounded by (a polynomial of) the spectral norm of its covariance matrix. Since the truncated Gibbs measure is supported on a compact set, its Cheeger constant is lower bounded by a temperature-independent constant. Using the Cheeger's inequality, we have the conclusion.

## Appendix C. Properties of PL<sup>o</sup> functions and Log-PL<sup>o</sup> measures

### C.1. Log-PL<sup>o</sup> measures has a single modal

**Proof** [Proposition 3] Note that by Assumption 3',  $V$  is not a constant function. Recall Assumption 1. We have that for any  $S'$  and any  $x \in \mathcal{N}(S')$ , if  $\nabla V(x) = 0$ ,  $V(x) = \min_{x \in \mathcal{N}(S')} V(x)$ , i.e.  $x \in \mathcal{N}(S')$  is a local minimizer if it is critical point.

Let  $X$  denote the set of local maxima of  $V$ , which implies that  $\forall x' \in X, \nabla V(x') = 0$ .

Some useful facts are listed as follows

- $X \cap \mathcal{N}(S') = \emptyset$  for any  $S'$ : Let  $x' \in X \cap \mathcal{N}(S')$ . One has  $\nabla V(x') = 0$ , i.e.  $x'$  is a local minimizer in  $\mathcal{N}(S')$ . But this contradicts with the assumption that  $x'$  is a local maximizer.

- $X$  is either empty or a collection of singletons: Since for any  $x' \in X$  (if  $X \neq \emptyset$ ), one has  $\nabla V(x') = 0$ . Therefore, by Assumption 2,  $\nabla^2 V(x') < 0$ . Hence every  $x'$  is a strict local maximizer, i.e.  $X$  is a collection of singletons.
- The set of all local minima of  $V$  has at most a finite number of separated components: We prove via contradiction. Suppose that the set of all local minima of  $V$  has infinitely many separated components. From Assumption 3', all local minima are contained within a compact set. Denote the collection of all separated components of local minima of  $V$  as  $\{S'_i\}$ . For every  $i$ , pick a representative point  $x_i \in S'_i$ . Clearly all  $x_i$  are contained in a compact set. From Bolzano–Weierstrass theorem, we have that  $\{x_i\}$  (as an infinite sequence) has at least one accumulation point  $x$ . Since  $V$  is  $C^1$ ,  $\nabla V(x) = 0$ , i.e.  $x$  is also a critical point.
  - If  $x \notin \mathcal{N}(S)$  (recall the definition of  $\mathcal{N}(S)$  in Assumption 2),  $x$  is a local maximum by Assumption 2, which contradicts with the fact that in every neighborhood of  $x$  there is a local minimum (since  $x$  is a accumulation point of  $\{x_i\}$ ).
  - If  $x \in \mathcal{N}(S)$ , there exists some  $S_{i'}$  such that  $x \in \mathcal{N}(S_{i'})$ . If  $x \in S_{i'}$ , it cannot be an accumulation point of  $\{x_i\}$ , as in  $\mathcal{N}(S_{i'})$  there is only one representative point  $x_{i'}$ . If  $x \in \mathcal{N}(S_{i'}) \setminus S_{i'}$ ,  $\nabla V(x) \neq 0$ , which again contradicts with the fact that  $\nabla V(x) = 0$ .

We can now prove that Log-PL<sup>o</sup> is uni-modal via contradiction. Suppose that the local minima of  $V$  has at least two separated components. Pick any two separated components  $S'_1$  and  $S'_2$  and let  $x_1$  and  $x_2$  be two points in these two components respectively. WLOG, assume  $f(x_1) \geq f(x_2)$ .

**Theorem 5 (Katriel (1994), Theorem 2.1)** *Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be  $C^1$  and coercive. Let  $x_1, x_2 \in \mathbb{R}^n$  and let  $P \subset \mathbb{R}^n$  separate  $x_1$  and  $x_2$  (that is,  $x_1$  and  $x_2$  lie in different components of  $\mathbb{R}^n \setminus P$ ), and:*

$$\max \{f(x_1), f(x_2)\} < \inf_{x \in P} f(x) = p$$

*Then there exists a point  $x_3$  which is a critical point of  $f$ , with:  $f(x_3) > \max \{f(x_1), f(x_2)\}$ . Moreover,  $x_3$  is either a local minimum or a global mountain passing point<sup>6</sup>.*

To apply the above theorem, take  $P$  to be a subset of  $\mathcal{N}(S'_1) \setminus S'_1$  such that  $\forall y \in P, V(y) > V(x_1) \geq \max\{V(x_1), V(x_2)\}$ . This is always possible according to Assumption 1. Consequently, according to the above theorem, we can find a critical point  $x_3$  which is either a local minimum or a global mountain passing point. Consider two cases:

1.  $x_3$  is not a local minimum but a global mountain passing point. Since  $x_3$  is a critical point, Assumption 2 implies that  $x_3$  is a strict local maximum. However, for  $d \geq 2$ , this is not possible, as in this case a strict local minimum point is not a global mountain passing point.
2.  $x_3$  is a local minimum. If this is the case, we now pick  $x_1$  and  $x_3$  and apply Theorem 5. It gives us a new local minimum  $x_4$  (note that  $x_4$  cannot be a global mountain passing point as discussed in the first case). Important, note that every time we have  $V(x_{i+1}) > V(x_i)$  and that every  $x_i$  is a local minimum. We can only do this a finite number of times since the collection of local minima of  $V$  has at most a finite number of separated components (i.e. there can only be a finite number of different values of  $V$  on the collection of local minima of  $V$ ).

Consequently, none of the above two cases is possible and the local minima of  $V$  has only one connected component, i.e.  $\mu_\epsilon$  is uni-modal. ■

---

6. A point  $x$  is called a global mountain passing point of  $f$  if for every neighborhood  $\mathcal{N}(x)$ , the set  $\{y : f(y) < f(x)\} \cap \mathcal{N}(x)$  is disconnected.

### C.2. PL° functions admits an embedding submanifold as its global optimal set

**Lemma 6** (*Global embedding submanifold*) *Suppose that  $V$  satisfies Assumptions 1 to 3', then the optimal set  $S$  is an embedding submanifold of  $\mathbb{R}^d$ .*

**Proof** According to Assumption 1, we know that for each  $x \in S$ ,  $V(x)$  satisfies the 2-PL condition around  $x$  in  $\mathbb{R}^d$  with constant  $\nu$ . Then we immediately see that  $S$  is a  $C^2$  embedding submanifold locally around  $x$  by (Rebjoek and Boumal, 2024, Lemma 2.15). Moreover, we also know that  $\text{rank}(\nabla^2 V)$  is a constant by (Rebjoek and Boumal, 2024, Corollary 2.13) since we assume that  $S$  is compact in Assumption 3'. Using (Boumal, 2023, Theorem 8.75), we obtain that  $S$  is a global embedding submanifold with dimension  $d - \text{rank}(\nabla^2 V)$ . ■

**Lemma 7** (*No boundary*) *Suppose that  $V$  satisfies Assumptions 1 to 3', then the optimal set  $S$  is an embedding submanifold of  $\mathbb{R}^d$  without boundary.*

**Proof** Assume the optimal set  $S$  is a  $k$  dimensional embedding submanifold with boundary  $\partial S$ . We immediately know that  $\partial S$  is a  $k - 1$  dimensional submanifold around  $\bar{x}$  by Theorem 5.11 in (Lee, 2020). Then we take  $\bar{x} \in \partial S$  and  $\{x_n\}_{n=1}^{+\infty} \subset \text{int}S$  around  $\bar{x}$  such that  $x_n \rightarrow \bar{x}$  as  $n \rightarrow +\infty$ . Moreover, we can take  $\mathbf{n}$  as a normal direction of  $\partial S$  at  $\bar{x}$  but  $\mathbf{n} \notin N_{\bar{x}}S$  such that

$$x(r) = \exp(r\mathbf{n}), \quad x(0) = \bar{x}, \quad x(r_n) = x_n \quad r, r_n \in [0, \epsilon]$$

with  $r < \text{inj}(\bar{x})$  small enough. We denote that

$$\mathbf{n}(r) = \frac{\partial}{\partial r} \exp(r\mathbf{n}) = x_* (\mathbf{n}) \Big|_r \in T_{x(r)}S.$$

By equivalence of 2-PL condition in Assumption 1 and Morse-Bott property of interior point of  $S$  in (Rebjoek and Boumal, 2024) when  $V(x) \in C^2$ , we have

$$\nabla^2 V(x_n)[\mathbf{n}(r_n)] = 0,$$

since  $\mathbf{n}(r_n) \in \text{Ker} \nabla^2 V(x_n)$ . Again by continuity of  $\nabla^2 V$ , we have

$$\nabla^2 V(\bar{x})[\mathbf{n}] = \lim_{n \rightarrow +\infty} \nabla^2 V(x_n)[\mathbf{n}(r)] = 0. \quad (16)$$

On the other hand, we already know that  $\mathbf{n}$  is normal to  $\partial S$  at  $\bar{x}$  and  $x(r) \in S \setminus \partial S$  when we take  $r > 0$ , hence  $-\mathbf{n}$ <sup>7</sup> is the outward normal direction to  $S$  at  $\bar{x}$  according to local orientation. Again by Quadratic Growth property in (Rebjoek and Boumal, 2024, Proposition 2.2), we have

$$0 < C \text{dist}^2(x(r), S) \leq V(x(r)) - V(\bar{x}), \quad r \in [-\epsilon, 0). \quad (17)$$

7. Let us clarify the definition of “tangent space” and “normal space” of a point which is at the boundary of a manifold. Taking  $M$  to be a  $k$  dimensional manifold with boundary  $\partial M$ , and  $W$  to be an open set of  $H^k = \{(x_1, \dots, x_k) \in \mathbb{R}^k | x_k \geq 0\}$ , then we actually have  $\partial W = W \cap \partial H^k$ . Now we keep the notation  $\bar{x} \in \partial M \subset M$ , then there exists a local chart  $(W, \phi)$  around  $\bar{x}$ , i.e.  $\phi(a) = \bar{x}, a \in W$  and  $\phi : W \rightarrow M$  is an embedding.

When we consider  $\bar{x} \in M$ , we have  $T_{\bar{x}}M = \mathbb{R}^k$ . This conclusion can be verified according to the definition of tangent space by  $\mathbb{R}$ -algebra  $C_{\bar{x}}^\infty(M)$  in (Tu, 2010, Section 22.4 and Figure 22.5). If we also have  $M$  is a embedding submanifold of  $\mathbb{R}^d$ , then the normal space  $N_{\bar{x}}(M)$  is the complement space of  $T_{\bar{x}}M$  in  $\mathbb{R}^d$ .

However, when we consider  $\bar{x} \in \partial M$ , we need define  $\bar{\phi} = \phi|_{\partial W} : \partial W \rightarrow \partial M$ , which is a local chart on  $\mathbb{R}^{k-1}$ . Then we can say that  $T_{\bar{x}}(\partial M) = \bar{\phi}_*(T_a(\partial W)) = \bar{\phi}_*(\mathbb{R}^{k-1})$ , which is a  $k - 1$  dimensional subspace of  $T_{\bar{x}}(M)$ . The normal space  $N_{\bar{x}}(\partial M)$  is the complement space of  $T_{\bar{x}}(\partial M)$  in  $T_{\bar{x}}(M)$ . This definition is not contradicted to the conclusion that  $\partial M$  is a  $k - 1$  submanifold of  $M$  without boundary (See (Tu, 2010, Section 22.3)).

for some constant  $C(\nu) > 0$ . However, if we consider the Taylor expansion on the right hand side,

$$\begin{aligned} V(x(r)) - V(x(0)) &= \nabla^2 V(x(r))(x(r) - x(0))^2 + o(\|x(r) - x(0)\|^2) \\ &= r^2 \langle \nabla^2 V(\bar{x})[-\mathbf{n}], [-\mathbf{n}] \rangle + o(r^2) \\ &= o(r^2), \end{aligned}$$

which is contradictory to (17) since

$$r^2 \sim \text{dist}^2(x(r), S) \leq o(r^2), \quad r \in [-\epsilon, 0)$$

when we take  $\epsilon$  small enough. Now we finish the proof.  $\blacksquare$

## Appendix D. Local coordinate representation of geometric structures in Section 2.3

**The first fundamental form.** Using the local chart  $(\Gamma, \phi)$  of  $S$ , the Riemannian metric  $g_S$  can be written as

$$g_S = \sum_{i,j=1}^k g_{ij}(u) du^i du^j, \quad u \in \Gamma \subset \mathbb{R}^k, \quad (18)$$

where

$$g_{ij}(u) = \frac{\partial \mathcal{M}(u)}{\partial u^i} \cdot \frac{\partial \mathcal{M}(u)}{\partial u^j}, \quad u \in \Gamma \subset \mathbb{R}^k.$$

Here  $\{\partial \mathcal{M}/\partial u^i\}_{i=1}^k$  are actually  $k$  tangent vector fields of  $k$ -dimensional embedding submanifold  $S$ , they generate the tangent plane on each point of  $S$ , i.e.

$$T_m S = \text{Span} \left\langle \frac{\partial \mathcal{M}(u)}{\partial u^1}, \dots, \frac{\partial \mathcal{M}(u)}{\partial u^k} \right\rangle, \quad m = (m_1(u), \dots, m_d(u)) \in S,$$

and “ $\cdot$ ” is standard inner product on  $\mathbb{R}^d$ . Also, these  $k$  tangent vector fields decide  $d - k$  normal vector fields on  $S$  by following global equation group

$$\frac{\partial \mathcal{M}(u)}{\partial u^i} \cdot \mathcal{N}_l(u) \equiv 0, \quad i = 1, \dots, k, \quad l = k + 1, \dots, d, \quad u \in \Gamma \subset \mathbb{R}^k, \quad (19)$$

moreover, we can take  $\mathcal{N}_{k+1}, \dots, \mathcal{N}_d$  as standard normal vector fields by following global equation group,

$$\mathcal{N}_i(u) \cdot \mathcal{N}_j(u) \equiv \delta_{ij}, \quad i, j = k + 1, \dots, d, \quad u \in \Gamma \subset \mathbb{R}^k, \quad (20)$$

they generate the normal bundle  $N$  on  $S$ , i.e.

$$N(m) = \text{Span} \langle \mathcal{N}_{k+1}(u), \dots, \mathcal{N}_d(u) \rangle, \quad m = \mathcal{M}(u) \in M, \quad u \in \Gamma \subset \mathbb{R}^k,$$

**The second fundamental form.** We define the second fundamental form as a symmetric quadratic form on the local chart  $(\Gamma, \phi)$ ,

$$\Pi = - \sum_{l=k+1}^d \left\{ t^l \sum_{i,j=1}^k G_{ij}(l) du^i du^j \right\}, \quad u \in \Gamma \subset \mathbb{R}^k, \quad t \in B(\epsilon) \subset \mathbb{R}^{d-k}, \quad (21)$$

where the matrix  $G(l) = (G_{ij}(l))$  is symmetric for each  $l = k + 1, \dots, d$ . We also use the following notation

$$G_j^i(l) = \sum_{s=1}^k g^{is} G_{sj}(l), \quad l = k + 1, \dots, d. \quad (22)$$

It is easy to see that the matrix  $\tilde{G}(l) = (G_j^i(l))$  is also symmetric since the metric tensor  $(g_{ij})$  is symmetric. Then, matrices  $\{G(l)\}_{l=k+1}^d$  of the second fundamental form  $\Pi$  can be locally written as

$$G_{ij}(l)(u) = \frac{\partial \mathcal{M}(u)}{\partial u^i} \cdot \frac{\partial \mathcal{N}_l(u)}{\partial u^j}, \quad u \in \Gamma \subset \mathbb{R}^k.$$

Moreover, by constraint (20) of normal vector fields  $\mathcal{N}_l$ ,  $l = k + 1, \dots, d$ , we also have

$$G_{ij}(l)(u) = \frac{\partial \mathcal{M}(u)}{\partial u^i} \cdot \frac{\partial \mathcal{N}_l(u)}{\partial u^j} = -\frac{\partial^2 \mathcal{M}(u)}{\partial u^i \partial u^j} \cdot \mathcal{N}_l(u), \quad u \in \Gamma \subset \mathbb{R}^k,$$

which implies that the matrix  $G(l) = (G_{ij}(l))$  is naturally symmetric and  $\Pi$  is a symmetric quadratic form. The geometric meaning of the second fundamental form of Riemannian submanifold  $S$  by embedding structure (9) is the projection of the variation of normal vector fields along the tangent space of Riemannian submanifold  $(S, g_S)$  based on the ambient space  $(\mathbb{R}^d, g_E)$ .

## Appendix E. Proofs of results in Section 3

### E.1. Proof of Lemma 1

**Property 1.** We prove via contradiction. Suppose that there is a sequence  $x_i \in \partial \mathcal{N}(S)$  such that  $\lim_{i \rightarrow \infty} \text{dist}(x_i, S) = 0$ . Note that  $\partial \mathcal{N}(S)$  is bounded. Hence, WLOG, we assume  $x_i$  is convergent, since otherwise we can always take a convergent subsequence. Denote  $x = \lim_{i \rightarrow \infty} x_i$ . By construction,  $x \in \partial \mathcal{N}(S)$ , boundary of an open neighborhood of  $S$ , but since  $\text{dist}(x, S) = 0$ ,  $x \in S$ , which leads to a contradiction.

**Property 2.** Note that under Assumptions 1 to 3,  $X$  contains at most finitely many singletons. This can be proved via contradiction: Suppose that there is an infinite sequence  $x_i \in X$ . Note that  $X$  is bounded by Assumption 3. Hence, WLOG, we assume  $x_i$  is convergent, since otherwise we can always take a convergent subsequence. Denote  $x = \lim_{i \rightarrow \infty} x_i$ . From the construction of  $\{x_i\}$ , in any neighborhood of  $x$ , there is another local maximum point, which contracts with Assumption 2 and  $V \in \mathcal{C}^2$ , since  $x$  is a strict local maximum.

**Property 3.** We know from Assumption 3,  $\nabla V(x) \neq 0$  for any  $x$  beyond a compact set. Denote this compact set by  $\mathcal{Y}$ . We now focus our discussion in  $\mathcal{Y}$  and prove via contradiction. Suppose that within the said compact set there is a sequence  $x_i \in \mathcal{Y} \cap (\mathcal{N}(X) \cup \mathcal{N}(S))^c$  such that  $\lim_{i \rightarrow \infty} \nabla V(x_i) = 0$ . Note that  $(\mathcal{N}(X) \cup \mathcal{N}(S))^c$  is bound. WLOG, we assume  $x_i$  is convergent, since otherwise we can always take a convergent subsequence. Denote  $x = \lim_{i \rightarrow \infty} x_i$ . We know that  $x \in (\mathcal{N}(X) \cup \mathcal{N}(S))^c$  since this set is closed. Moreover, since  $\nabla V(x) = 0$ , we have  $x \in S$  or  $x \in X$ , which leads to a contradiction.

### E.2. Proof of Lemma 3

**Subdomain  $\Xi_1 = \{x : 2R_0 \leq \|x\|\}$ .** Recall  $R_0$  from Assumption 3. One has

$$\text{dist}(x, S) \geq |x| - R_0 \Rightarrow \text{dist}(x, S) \geq \frac{1}{2}|x|.$$

From Assumption 3, one has

$$\frac{\mathcal{L}W}{\epsilon W} \leq \frac{C_g |x|^2}{2\epsilon} - \frac{1}{4\epsilon^2} v_{eb}^2 \cdot \text{dist}^2(x, S) \leq \frac{C_g |x|^2}{2\epsilon} - \frac{1}{64\epsilon^2} v_{eb}^2 |x|^2.$$

Recall that  $\epsilon \leq \frac{v_{eb}^2}{64C_g}$  and  $|x| \geq 2R_0 \geq R_0$ . We have

$$\frac{\mathcal{L}W}{\epsilon W} \leq \frac{C_g |x|^2}{2\epsilon} - \frac{1}{64\epsilon^2} v_{eb}^2 |x|^2 \leq -\frac{v_{eb}^2 |x|^2}{128} \frac{1}{\epsilon^2} \leq -\frac{v_{eb}^2 R_0^2}{128} \frac{1}{\epsilon^2}.$$

Now we can select the parameter  $\sigma$  in (6) as  $\sigma_1 = \frac{v_{eb}^2 R_0^2}{128} \frac{1}{\epsilon^2}$  in this region. Consequently we can establish the inequality (6) for  $|x| \geq 2R_0$ .

**Subdomain**  $\Xi_2 = \left\{x : \sqrt{C}\epsilon \leq \text{dist}(x, S) \text{ and } \|x\| \leq 2R_0\right\}$ . There are three cases.

- $x$  is in the  $R_1$  neighborhood of  $X$ , where we recall that  $X$  is the collection of all local maxima of  $V$  in Lemma 1: Using the second property of Lemma 1, we have

$$\frac{\mathcal{L}W}{\epsilon W} \leq -\frac{d\mu^-}{2\epsilon}.$$

Note that the value  $\sigma_2^1 = \frac{d\mu^-}{2\epsilon}$  in this region.

- $x \in \mathcal{N}(S)$ : Recall  $M_\Delta$  from Lemma 2. Using the error bound in Lemma 2, one has

$$\frac{\mathcal{L}W}{\epsilon W} \leq \frac{M_\Delta}{2\epsilon} - \frac{1}{4\epsilon^2} v^2 \cdot \text{dist}^2(x, S).$$

Consequently, we can establish the inequality (6) for  $x \in \mathcal{N}(S)$  with  $C = \frac{4M_\Delta}{v^2}$  by

$$-\frac{1}{4\epsilon^2} v^2 \cdot \text{dist}^{\frac{2}{a-1}}(x, S) \leq -\frac{M_\Delta}{\epsilon}.$$

Note that the value  $\sigma_2^2 = \frac{M_\Delta}{2\epsilon}$  in this region.

- $x$  is in the compact set, but not in the above two cases, i.e.

$$x \in (\{x : \text{dist}(x, X) \leq R_1\} \cup \mathcal{N}(S))^c \cap \{x : \|x\| \leq 2R_0\}.$$

According to Lemma 1, there exists a constant lower bound of  $\|\nabla V(x)\| \geq g_0 > 0$  in this regime. One has

$$\frac{\mathcal{L}W}{\epsilon W} \leq \frac{M_\Delta}{2\epsilon} - \frac{g_0^2}{4\epsilon^2} \leq -\frac{M_\Delta}{4\epsilon}.$$

For a sufficiently small  $\epsilon$ , such that  $\frac{g_0^2}{4\epsilon^2} \geq \frac{dL}{\epsilon}$ . Note that the value  $\sigma_2^3 = \frac{M_\Delta}{4\epsilon}$  in this region.

WLOG, we assume  $\frac{M_\Delta}{4\epsilon} \geq \frac{d\mu^-}{2\epsilon}$  (otherwise simply set  $\mu^- = M_\Delta/2d$  in Lemma 1). We have  $\sigma_2 = \min\{\sigma_2^1, \sigma_2^2, \sigma_2^3\} = \frac{d\mu^-}{2\epsilon}$ .

**Global estimation of  $\sigma$ .** Since we need eq. (8) to hold on  $\mathbb{R}^d \setminus U$ , we take

$$\sigma = \inf_{i \in \{1,2\}} \sigma_i \tag{23}$$

**Estimation of  $b$ .** Since we will pick  $U \subseteq S^{\sqrt{C}\epsilon}$ , from Lemma 1, we can set

$$b := \sigma + \frac{M_\Delta}{2\epsilon}. \quad (24)$$

## Appendix F. Proof of results in Section 4

### F.1. Proof of Lemma 5

**Proof** Based on the representation of local coordinate (15), we use the formula of change of variables,

$$\int_{T(\epsilon)} \phi(y) dy = \int_{T(\epsilon)} \phi(y) |J(u, r)| du dr,$$

where  $J(u, r)$  is Jacobian determinant of change of variables of diffeomorphism  $y \rightarrow m + \nu$  defined by (15). We can easily compute the Jacobian determinant  $J(u, t)$  on each local chart  $(\Gamma, \phi)$  as

$$\begin{aligned} J(u, r) &= \left| \left[ \frac{\partial y}{\partial u^1}, \dots, \frac{\partial y}{\partial u^k}, \frac{\partial y}{\partial r^{k+1}}, \dots, \frac{\partial y}{\partial r^d} \right] \right| \\ &= \left| \left[ \frac{\partial y}{\partial u^1}, \dots, \frac{\partial y}{\partial u^k}, \mathcal{N}_{k+1}, \dots, \mathcal{N}_d \right] \right| \\ &= \left| \left[ \left( \frac{\partial \mathcal{M}}{\partial u^1} + \sum_{l=k+1}^d r^l \frac{\partial \mathcal{N}_l}{\partial u^1} \right), \dots, \left( \frac{\partial \mathcal{M}}{\partial u^k} + \sum_{l=k+1}^d r^l \frac{\partial \mathcal{N}_l}{\partial u^k} \right), \mathcal{N}_{k+1}, \dots, \mathcal{N}_d \right] \right| \\ &= \left| \left[ \left( \mathcal{M}_1 + \sum_{l=k+1}^d r^l \mathcal{N}_{l,1} \right), \dots, \left( \mathcal{M}_k + \sum_{l=k+1}^d r^l \mathcal{N}_{l,k} \right), \mathcal{N}_{k+1}, \dots, \mathcal{N}_d \right] \right|. \end{aligned}$$

We emphasize that these vector fields

$$\{\mathcal{M}_i\}_{i=1}^k, \quad \{\mathcal{N}_i\}_{i=1}^{d-k}, \quad \{\{\mathcal{N}_{l,j}\}_{l=k+1}^d\}_{j=1}^k,$$

only depend on variable  $u \in \Gamma \subset \mathbb{R}^k$ , i.e. they are only defined on Riemannian submanifold  $S$ . Moreover, each vector at the point  $m = \mathcal{M}(u) \in S$  is a linear combination of these  $d$  basic vectors  $\{\{\mathcal{M}_i(u)\}_{i=1}^k, \{\mathcal{N}_l(u)\}_{l=k+1}^d\}$ . Then we have

$$\mathcal{N}_{l,j} = \sum_{i=1}^k T_j^i(l) \mathcal{M}_i + \sum_{s=k+1}^d T_j^s(l) \mathcal{N}_s.$$

By constrain (19) and (20), we can easily compute that

$$\begin{cases} T_j^i(l) = \sum_{p=1}^k g^{ip} G_{pj}(l) = G_j^i(l) & i = 1, \dots, k, \\ T_j^s(l) \equiv 0, & s = k+1, \dots, d, \end{cases}$$

where  $(g^{ip}) = (g_{ip})^{-1}$  is the inverse matrix of  $k \times k$  matrix  $(g_{ij})$  in the Riemannian metric tensor (18). Hence we have

$$\mathcal{N}_{l,j} = \sum_{i=1}^k G_j^i(l) \mathcal{M}_i.$$

Now we back to the original computation,

$$\begin{aligned} \int_{T(\epsilon)} \phi(y) dy &= \int_{T(\epsilon)} \phi(y) |J(u, r)| du dr \\ &= \int_{T(\epsilon)} \phi(y) \left| \det \left( I_k + \sum_{l=k+1}^d r^l \tilde{G}(l) \right) \left[ \mathcal{M}_1, \dots, \mathcal{M}_k, \mathcal{N}_{k+1}, \dots, \mathcal{N}_d \right] \right| du dr, \end{aligned}$$

we observe that

$$\det \left( \left[ \mathcal{M}_1, \dots, \mathcal{M}_k, \mathcal{N}_{k+1}, \dots, \mathcal{N}_d \right]^T \left[ \mathcal{M}_1, \dots, \mathcal{M}_k, \mathcal{N}_{k+1}, \dots, \mathcal{N}_d \right] \right) = \det(g)$$

by (19), (20) and (18), and the determinant  $\det(g)$  does not depend on variable  $r \in B(\epsilon) \subset \mathbb{R}^{d-k}$ , we finally have

$$\begin{aligned} \int_{T(\epsilon)} \phi(y) dy &= \int_{T(\epsilon)} \phi(y) \left| \det \left( I_k + \sum_{l=k+1}^d r^l \tilde{G}(l) \right) \left[ \mathcal{M}_1, \dots, \mathcal{M}_k, \mathcal{N}_{k+1}, \dots, \mathcal{N}_d \right] \right| du dr \\ &= \int_{T(\epsilon)} \phi(y) \left| \det \left( I_k + \sum_{l=k+1}^d r^l \tilde{G}(l) \right) \right| \sqrt{\det(g)} du dr \\ &= \int_{\Gamma} \left\{ \int_{B(\epsilon)} \phi(y) \left| \det \left( I_k + \sum_{l=k+1}^d r^l \tilde{G}(l) \right) \right| dr^{k+1} \dots dr^d \right\} d\mathcal{M}(u), \end{aligned}$$

now we finish the proof. ■

## F.2. Proof of Lemma 6

**Proof** For the direction of the parameter  $u^i$ , by chain rule we have

$$\begin{aligned} \nabla_{u^i} \phi(y(u, r)) &= \nabla_y \phi \cdot \frac{\partial y}{\partial u^i} = \nabla_y \phi \cdot \left[ \frac{\partial \mathcal{M}}{\partial u^i} + \sum_{l=k+1}^d r^l \frac{\partial \mathcal{N}_l}{\partial u^i} \right] \\ &= \nabla_y \phi \cdot \left[ \mathcal{M}_i + \sum_{l=k+1}^d r^l \frac{\partial \mathcal{N}_l}{\partial u^i} \right] \\ &= \nabla_y \phi \cdot \left[ \mathcal{M}_i + \sum_{l=k+1}^d r_l \sum_{j=1}^k G_i^j(l) \mathcal{M}_j \right] \\ &= \nabla_y \phi \cdot [\mathcal{M}_1, \dots, \mathcal{M}_d] \cdot \left[ I_k + \sum_{l=k+1}^d r^l \tilde{G}(l) \right]. \end{aligned}$$

For the direction of the parameter  $r^i$ , by chain rule we have

$$\nabla_{r^i} \phi(y(u, r)) = \nabla_y \phi \cdot \frac{\partial y}{\partial r^i} = \nabla_y \phi \cdot \mathcal{N}_i.$$

Combining these two results, we have

$$\nabla_{(u,r)}\phi(y(u,r)) = \nabla_y\phi \cdot [\mathcal{M}_1, \dots, \mathcal{M}_k, \mathcal{N}_{k+1}, \dots, \mathcal{N}_d] \cdot \begin{bmatrix} I_k + \sum_{l=k+1}^d r^l \tilde{G}(l) & 0 \\ 0 & I_{d-k} \end{bmatrix}.$$

Now we finish the proof. ■

### F.3. Proof of Proposition 6

**Proof** Our main idea to describe the asymptotic behavior of Poincaré constant  $\lambda_1(T(\epsilon))$  is try to obtain the lower and upper bound of  $\lambda_1(T(\epsilon))$  based on the first eigenvalue of Laplacian-Beltrami operator on a trivial product Riemannian manifold  $(S \times B(\epsilon), g_S + g_{B(\epsilon)})$  and small  $\epsilon$  perturbation issues. Here  $g_{B(\epsilon)} = i_{B(\epsilon)}^*(g_E)$  is the standard Riemannian metric on  $B(\epsilon) \subset \mathbb{R}^{d-k}$ , i.e. the pullback of  $g_E$  by the including map  $i_{B(\epsilon)}$ . In the next we try to show

$$\lambda_1(T(\epsilon)) \sim \lambda_1(S \times B(\epsilon)) + O(\epsilon).$$

We start from the min-max formula and divide the estimates into two parts.

- The lower and upper bound of  $L^2$  norm in  $(T(\epsilon), g_E)$  based on the  $L^2$  norm in  $(S \times B(\epsilon), g_S + g_{B(\epsilon)})$  and  $\epsilon$  perturbation. By expansion formula of determinant,

$$\det(I_k + \sum_{l=k+1}^d r^l \tilde{G}(l)) = 1 + \sum_{i=1}^k \sum_{l=k+1}^d r^l G_i^i(l) + \dots,$$

and combine with bounded condition in Assumption 4 about the second fundamental form  $\Pi$  for Riemannian submanifold  $S$ , we have

$$1 - A_2\epsilon \leq \det(I_k + \sum_{l=k+1}^d r^l \tilde{G}(l)) \leq 1 + A_2\epsilon.$$

for some constant  $A_2 = A_2(d, k, \tilde{G}(l))$ . Using integral formula in Lemma 5, we have

$$\begin{aligned} \int_{T(\epsilon)} |\phi(y)|^2 dy &\leq (1 + A_2\epsilon) \int_{\Gamma} \left\{ \int_{B(\epsilon)} |\phi(y)|^2 dr^{k+1} \dots dr^d \right\} d\mathcal{M}, \\ \int_{T(\epsilon)} |\phi(y)|^2 dy &\geq (1 - A_2\epsilon) \int_{\Gamma} \left\{ \int_{B(\epsilon)} |\phi(y)|^2 dr^{k+1} \dots dr^d \right\} d\mathcal{M}. \end{aligned}$$

- The lower and upper bound of Dirichlet energy in  $(T(\epsilon), g_E)$  based on the Dirichlet energy in  $(S \times B(\epsilon), g_S + g_{B(\epsilon)})$  and  $\epsilon$  perturbation. By Lemma 6, we have

$$\nabla_y\phi(y) = \nabla_{(u,r)}\phi(y(u,r)) \cdot \begin{bmatrix} (I_k + \sum_{l=k+1}^d r^l \tilde{G}(l))^{-1} & 0 \\ 0 & I_{d-k} \end{bmatrix} \cdot [\mathcal{M}_1, \dots, \mathcal{M}_k, \mathcal{N}_1, \dots, \mathcal{N}_{d-k}]^{-1},$$

then we obtain

$$|\nabla_y\phi(y)|^2 = \nabla_{(u,r)}\phi(y(u,r)) \cdot \begin{bmatrix} (I_k + \sum_{l=k+1}^d r^l \tilde{G}(l))^{-1} (g)^{-1} (I_k + \sum_{l=k+1}^d r^l \tilde{G}(l))^{-1} & 0 \\ 0 & I_{d-k} \end{bmatrix} \cdot \nabla_{(u,r)}\phi(y(u,r))^T.$$

Selecting  $t$  small enough and using bounded condition in Assumption 4, we have

$$\begin{cases} |\nabla_y \phi(y)|^2 \leq (1 + A_1 \epsilon) (\nabla_u \phi(y(u, r))(g)^{-1} \nabla_u \phi(y(u, r))^T + |\nabla_r \phi(y(u, r))|^2), \\ |\nabla_y \phi(y)|^2 \geq (1 - A_1 \epsilon) (\nabla_u \phi(y(u, r))(g)^{-1} \nabla_u \phi(y(u, r))^T + |\nabla_r \phi(y(u, r))|^2), \end{cases}$$

for some constant  $A_1 = A_1(d, k, \tilde{G}(l)) > 0$ . Recall the duality between  $TS$  and  $T^*S$  induced by metric  $g$ ,

$$d^u \phi = \sum_{i,j=1}^k g^{ij} \frac{\partial \phi}{\partial u^j} \frac{\partial}{\partial u^i} = \sum_{i=1}^k g^{ij} \nabla_{u^j} \phi,$$

where  $d^u$  is external derivative associated with parameter  $u$ , then we have

$$\begin{aligned} \int_{T(\epsilon)} |\nabla_y \phi(y)|^2 dy &= \int_S \left\{ \int_{B(\epsilon)} |\nabla_y \phi(y)|^2 \left| \det \left( I_k + \sum_{l=k+1}^d r^l \tilde{G}(l) \right) \right| dr^{k+1} \dots dr^d \right\} d\mathcal{M} \\ &\leq (1 + A_2 \epsilon) \int_S \left\{ \int_{B(\epsilon)} |\nabla_y \phi(y)|^2 dr^{k+1} \dots dr^d \right\} d\mathcal{M} \\ &\leq (1 + A_2 \epsilon) (1 + A_1 \epsilon) \int_S \left\{ \int_{B(\epsilon)} (|d^u \phi|^2 + |\nabla_r \phi(y)|^2) dr^{k+1} \dots dr^d \right\} d\mathcal{M}. \end{aligned}$$

Similarly, we also have

$$\int_{T(\epsilon)} |\nabla_y \phi(y)|^2 dy \geq (1 - A_2 \epsilon) (1 - A_1 \epsilon) \int_S \left\{ \int_{B(\epsilon)} (|d^u \phi(y)|^2 + |\nabla_r \phi(y)|^2) dr^{k+1} \dots dr^d \right\} d\mathcal{M}.$$

Combining all these estimates together, we have

$$\frac{\int_{T(\epsilon)} |\nabla_y \phi(y)|^2 dy}{\int_{T(\epsilon)} |\phi(y)|^2 dy} \leq (1 + A_1 \epsilon) (1 + A_2 \epsilon)^2 \frac{\int_S \left\{ \int_{B(\epsilon)} (|d^u \phi(y)|^2 + |\nabla_r \phi(y)|^2) dr^{k+1} \dots dr^d \right\} d\mathcal{M}}{\int_S \left\{ \int_{B(\epsilon)} |\phi(y)|^2 dr^{k+1} \dots dr^d \right\} d\mathcal{M}}$$

and

$$\frac{\int_{T(\epsilon)} |\nabla_y \phi(y)|^2 dy}{\int_{T(\epsilon)} |\phi(y)|^2 dy} \geq (1 - A_1 \epsilon) (1 - A_2 \epsilon)^2 \frac{\int_S \left\{ \int_{B(\epsilon)} (|d^u \phi(y)|^2 + |\nabla_r \phi(y)|^2) dr^{k+1} \dots dr^d \right\} d\mathcal{M}}{\int_S \left\{ \int_{B(\epsilon)} |\phi(y)|^2 dr^{k+1} \dots dr^d \right\} d\mathcal{M}}.$$

These two kinds of estimates imply us that the eigenvalues of Laplacian-Beltrami operator on  $T(\epsilon)$  is equivalent to the eigenvalues of Laplacian-Beltrami operator of the following product Riemannian manifold

$$(S \times B(\epsilon), g_S + g_{B(\epsilon)}).$$

It is easy to know that

$$\lambda_1(S \times B(\epsilon)) = \min \{ \lambda_1(S, g_S), \lambda_1(B(\epsilon), g_{B(\epsilon)}) \}.$$

Now we complete our discussion of this problem and get conclusion

$$\lambda_1(S)(1 - B\epsilon) \leq \lambda(T(\epsilon)) \leq \lambda_1(S)(1 + B\epsilon),$$

for some constant  $B = B(A_1, A_2) > 0$  when  $\epsilon$  small enough. ■

## Appendix G. Proof of results in Section 5

### G.1. Proof of Theorem 4

The proof need to combine Lyapunov approach in Section 3 with spectral stability analysis in Section 4. Recall the final result in Theorem 2,

$$\rho_{\mu_\epsilon} \geq \frac{1}{2} \exp(-\bar{C}) \lambda_1^n(U)$$

with  $U = S^{\sqrt{C\epsilon}}$ . Let us focus on dealing with  $\lambda_1^n(U)$  by conclusions in Section 4,

We use Proposition 6 with  $\tilde{\epsilon} = \sqrt{C\epsilon}$ , then we have

$$\lambda_1(S)(1 - Bc(C\epsilon)^{\frac{1}{2}}) \leq \lambda_1^n(U) \leq \lambda_1(S)(1 + Bc(C\epsilon)^{\frac{1}{2}}).$$

We finally have

$$\rho_{\mu_\epsilon} \geq \frac{1}{2} \exp(-\bar{C}) \lambda_1^n(U) \geq \frac{1}{4} \exp(-\bar{C}) \lambda_1(S),$$

when  $\epsilon \leq \frac{1}{C} \left(\frac{1}{2B}\right)^2$  is small enough. Now we finish the proof.