# Critical exponent for the one-dimensional wave equation with a space-dependent scale invariant damping and time derivative nonlinearity

Ahmad Z. Fino

College of Engineering and Technology, American University of the Middle East, Kuwait Mohamed Ali Hamza

Imam Abdulrahman Bin Faisal University, Dammam, 34212, Saudi Arabia

### Abstract

We investigate in this paper the Cauchy problem of the one-dimensional wave equation with space-dependent damping of the form  $\mu_0(1 + x^2)^{-1/2}$ , where  $\mu_0 > 0$ , and time derivative nonlinearity. We establish global existence of mild solutions for small data compactly supported by employing energy estimates within suitable Sobolev spaces of the associated homogeneous problem. Furthermore, we derive a blow-up result under some positive initial data by employing the test function method. This shows that the critical exponent is given by  $p_G(1 + \mu_0) = 1 + 2/\mu_0$ , when  $\mu_0 \in (0, 1]$ , where  $p_G$  is the Glassey exponent. To the best of our knowledge, this constitutes the first identification of the critical exponent range for this class of equations. As by product, we extend the global existence result to a more general class of space/time nonlinearities of the form  $f(\partial_t u, \partial_x u) = |\partial_x u|^q$  or  $f(\partial_t u, \partial_x u) = |\partial_t u|^p |\partial_x u|^q$ , with p, q > 1.

MSC 2020 Classification: 35A01, 35B33, 35L15, 35D35, 35B44

**Keywords:** Nonlinear wave equations, global existence, blow-up, lifespan, critical exponent, scale-invariant damping, time-derivative nonlinearity.

### 1 Introduction

In this work, we consider the Cauchy problem for the wave equations with critical spacedependent damping and power-nonlinearity of derivative type

$$\begin{cases} \partial_t^2 u - \partial_x^2 u + V(x)\partial_t u = f(\partial_t u), & x \in \mathbb{R}, t > 0, \\ u(x,0) = u_0(x), & \partial_t u(x,0) = u_1(x) & x \in \mathbb{R}, \end{cases}$$
(1.1) {NLW}

where p > 1,  $V(x) = \mu_0(1 + x^2)^{-1/2}$ ,  $\mu_0 > 0$ ,  $f(\partial_t u) = |\partial_t u|^p$  or  $|\partial_t u|^{p-1}\partial_t u$ , and  $(u_0, u_1)$  are compactly supported initial data in the following Sobolev space

$$(u_0, u_1) \in H^2(\mathbb{R}) \times H^1(\mathbb{R}).$$
 (1.2) {initialdat

Such problems appear in models for wave propagation in a nonhomogeneous gas with damping, where the space-dependent coefficients represent either friction coefficients or potential (see [19]).

In this paper, we investigate both the global existence of small data solutions and the finite-time blow-up of solutions to the Cauchy problem for the nonlinear wave equation (1.1). These results allow us to determine the critical exponent for  $\mu_0 \in (0, 1]$ .

### **1.1** Prior results

<br/>
Before going to the main result, we shall review prior studies on the global existence/nonexistence<br/>
of solutions to some nonlinear damped wave equations with various type of potentials and non-<br/>
linearities.

For the Cauchy problem of the semilinear damped wave equation with power nonlinearity and when the variable coefficients are missing,

$$u_{tt} - \Delta u + u_t = |u|^p, \qquad t > 0, \ x \in \mathbb{R}^n, \tag{1.3} \quad \{\texttt{classicalcal}\}$$

The study of the analogous problem of (1.3) with space-dependent damping, namely

$$\partial_t^2 u - \Delta u + \frac{\mu_0}{(1+|x|^2)^{\alpha/2}} \partial_t u = |u|^p \text{ in } \mathbb{R}^n \times (0,\infty),$$
 (1.4) {D1}

has been the subject of several works in the literature. First, for  $\alpha \in (0, 1)$  the critical exponent is given by the several works in the literature is the literature. If is determined by the several works in the literature is the literature of the several work in the literature is given by the several works in the literature. First, if  $\alpha < (0, 1)$  the critical exponent is given by the several work in the literature. The literature is given by the several work is given by the several work of the several work is given by the several work of the several work

in a recent paper by Sobajima [37]. In the case of whole space, a blow-up result and lifespan estimate were also established by Ikeda and Sobajima [20].

Now, we turn back to the classical semilinear wave equation with time derivative nonlinearity without damping term, namely

$$\partial_t^2 u - \Delta u = |\partial_t u|^p, \quad \text{in } \mathbb{R}^n \times (0, \infty), \tag{1.5}$$

for which the critical exponent is determined by the Glassey exponent

$$p_G = p_G(n) := 1 + \frac{2}{n-1},$$
 (1.6) {Glassey}

see e.g. [24, 35, 36, 46].

$$\partial_t^2 u - \Delta u + \frac{\mu_0}{1+t} \partial_t u = |\partial_t u|^p, \quad \text{in } \mathbb{R}^n \times (0, \infty), \tag{1.7}$$

$$\partial_t^2 u - \Delta u + \frac{\mu_0}{(1+|x|^2)^{\alpha/2}} \partial_t u = |\partial_t u|^p \quad \text{in } \mathbb{R}^n \times (0,\infty), \tag{1.8}$$

which corresponds to the scattering case, it is reasonable to expect that the blow-up region of the solution of (1.8) is similar to the case of the pure wave equation and the scattering damping term has no influence in the dynamics. The predicted blow-up result was obtained

Recently, in [9], we proved that the blow-up region of the present problem (1.1) is given by  $p \in (1, p_G(n + \mu_0)]$  in higher dimensions  $n \ge 2$ . However, up to our knowledge, there are no results on the blow-up in finite time in the one dimensional case as well as the small data global existence of solutions is still an open problem in any dimension.

This paper aims to address and resolve the critical exponent problem associated with equation (1.1). As previously explained resolve the critical exponent a short in the one-dimensional case, where damping induces in the critical exponent as the problem (1.1). As previously explained resolve the problem (1.1), as previously explained resolve the problem (1.1). As previously explained resolve the problem (1.1), as previously explained resolve the problem (1.1),

• Part I: Establish global existence results for the problem (1.1), it is essential to analyze the associated linear problem and utilize decay rates in appropriate Sobolev spaces. That question was done by Ikehata-Todorova-Yordanov in [21]. Indeed, they showed that, in the case where the initial data are compactly supported, the solution satisfies the same kind of energy estimates, namely  $\|(\nabla u(t), \partial_t u(t))\|_{L^2} \leq C(1+t)^{-\gamma}$ , for some  $\gamma > 0$ , and where the constant C depends on the support of the initial data. We would like to draw the attention of the reader to the fact that a better understanding of the dependence of the constant related to the initial data is crucial for establishing a global existence result for the corresponding nonlinear problem. In this context, in recent work in [37], the author successfully established a global existence result for the nonlinear problem with space-dependent damping in an exterior domain. This achievement was made by employing weighted energy estimates for the associated linear problem. In the present paper, we start by studying the associated homogeneous problem, by using energy estimates for compactly supported initial data. Subsequently, we address the nonlinearity using the fixed-point theorem. Therefore, the following upper bound for the critical exponent of problem (1.1) is obtained:

$$p_c(1,\mu_0) \le 1 + \frac{2}{\alpha_0},$$
 (1.9) {critical1}

where

$$\alpha_0 := \min(1, \mu_0).$$
 (1.10) {alpha00}

• Part II: For initial data with compact support, we derive a blow-up result for the nonlinear wave equation (1.1), by employing the test function method. Specifically, we employ a test function constructed as the product of a cut-off function and an explicit solution to the adjoint equation associated with the linear part of (1.1). This approach establishes a lower bound for the blow-up region for the solution of equation (1.1). Hence, the following lower bound for the critical exponent of problem (1.1) is established:

$$p_c(1,\mu_0) \ge 1 + \frac{2}{\mu_0}.$$
 (1.11) {critical2}

By combining the results from Parts I and II, we easily deduce that

$$p_c(1,\mu_0) = 1 + \frac{2}{\mu_0}, \quad \text{if } \mu_0 \in (0,1].$$
 (1.12) {critical12

At the end of this subsection we prepare notation and several definitions used throughout this paper. We denote by C a positive constant, which may change from line to line.  $L^p =$  $L^p(\mathbb{R})$  stands for the usual Lebesgue space, and  $H^k = H^k(\mathbb{R})$  for  $k \in \mathbb{Z}_{>0}$  is the Sobolev space defined by:

$$H^{k}(\mathbb{R}) = \left\{ f \in L^{2}(\mathbb{R}); \|f\|_{H^{k}} = \sum_{\ell=0}^{k} \|f^{(\ell)}\|_{L^{2}} < \infty \right\}.$$

#### 1.2Main Result

The purpose of this subsection is to state our main results. Recall that f(d<sub>i</sub>u) = |d<sub>i</sub>u|<sup>p</sup> or  $|\partial_t u|^{p-1} \partial_t u$ . We start by giving the definition of mild solution of the system (1.1).

**Definition 1.1.** (Mild solution) <br/>*Assume that* (u<sub>0</sub>, u<sub>1</sub>) ∈ H<sup>2</sup>(ℝ) × H<sup>1</sup>(ℝ). We say that a function u is a mild solution of (1.1) if

$$u \in \mathcal{C}^1([0,T], H^1(\mathbb{R}))$$

and u has the initial data  $u(0) = u_0$ ,  $u_t(0) = u_1$  and satisfies the integral equation

$$u(t,x) = R(t)(u_0, u_1) + \int_0^t S(t,s)f(\partial_t u) \, ds =: u^{lin}(t) + u^{nl}(t) \tag{1.13}$$
 {mildsoluti

solution.

**Theorem 1.1.** (Small data global existence)  $R_0 > 0.$  If

$$p > 1 + \frac{2}{\alpha_0},$$

 
 where a\_0 := min(\mu\_0, 1), then there exists a positive constant 0 < \varepsilon\_0 \ll 1, small enough, such that for any initial data satisfying

$$||u_0||_{H^2} + ||u_1||_{H^1} \le \varepsilon_0,$$

problem (1.1) possesses a uniquely global mild solution

$$u \in \mathcal{C}([0,\infty), H^2(\mathbb{R})) \cap \mathcal{C}^1([0,\infty), H^1(\mathbb{R})) \cap \mathcal{C}^2([0,\infty), L^2(\mathbb{R})).$$

Moreover, the solution satisfies the following estimate

$$(1+t)^{-1} \|u(t)\|_{L^2} + \|\partial_x u(t)\|_{H^1} + \|\partial_t u(t)\|_{H^1} + \|\partial_t^2 u(t)\|_{L^2} \le C (1+t)^{-\frac{\alpha}{2}},$$

where  $\alpha$  is defined by (2.31).

{subsec3}

{globalexist

For the purpose of stating the blow-up result, we present the weak formulation of (1.1) in the associated energy space, as follows

ไbefinition 1. Let  $u_0 \in H^1(\mathbb{R}), u_1 \in L^2(\mathbb{R})$  and T > 0. Let u be such that  $u \in \mathcal{C}([0, T), H^1(\mathbb{R})) \cap$  $\mathcal{C}^1([0, T), L^2(\mathbb{R}))$  and  $\partial_t u \in L^p_{loc}((0, T) \times \mathbb{R}),$  verifies, for any  $\varphi \in \mathcal{C}^0_0([0, T) \times \mathbb{R}) \cap \mathcal{C}^\infty((0, T) \times \mathbb{R}),$  the following identity:

$$\begin{split} \varepsilon & \int_{\mathbb{R}} u_1(x)\varphi(x,0)dx + \int_0^T \int_{\mathbb{R}} f(\partial_t u)\varphi(x,t)\,dxdt \\ = & -\int_0^T \int_{\mathbb{R}} \partial_t u(t,x)\partial_t \varphi(x,t)\,dxdt + \int_0^T \int_{\mathbb{R}} \partial_x u(t,x)\partial_x \varphi(x,t)\,dxdt \qquad (1.14) \quad \{\text{weaksol}\} \\ & +\int_0^T \int_{\mathbb{R}} V(x)\partial_t u(t,x)\varphi(x,t)\,dxdt, \end{split}$$

{def1}

 $\{th2\}$ 

and the condition u(x, 0) = \varepsilon u(x), u, u, u, u) = \varepsilon u(x) are satisfied. Then, u is called an energy solution of (1.1) on [0, T).

We denote the lifespan for the energy solution by:

 $T_{\varepsilon}(u_0, u_1) := \sup\{T \in (0, \infty]; \text{ there exists a unique energy solution } u \text{ of } (1.1)\}.$ 

moreover, if T > 0 can be arbitrary chosen, i.e. T<sub>e</sub>(u<sub>0</sub>, u<sub>1</sub>) = ∞, then u is called a global energy solution of (1.1).

Theorem 1.2. (Blow-up)

$$\int_{\mathbb{R}} \left( u_0''(x) + u_1(x) \right) \phi(x) dx > 0, \tag{1.15} \quad \text{{CO}}$$

where  $\phi(x)$  is a solution of the elliptic problem (5.3). If

$$1$$

then the energy solution u of (1.1) with compact support

$$\operatorname{supp} u \in \{(x,t) \in \mathbb{R} \times [0,T) \colon |x| \le R_0 + t\}$$

$$(1.16) \quad \{\operatorname{suppcond}\}$$

blows-up in finite time. More precisely, there exists a constant  $\varepsilon_0 = \varepsilon_0(u_0, u_1, \mu_0, p, R_0) > 0$ such that the lifespan  $T_{\varepsilon}$  verifies

$$T_{\varepsilon} \leq \begin{cases} C \varepsilon^{-\frac{2(p-1)}{2-\mu_0(p-1)}} & \text{if } 1 (1.17) {T-epss}$$

for  $0 < \varepsilon \leq \varepsilon_0$  and some constant C independent of  $\varepsilon$ .

**Remark 1.1.** We emphasize, by Theorems 1.1 and 1.2, that when  $\mu_0 \in (0,1]$  the critical exponent in the one-dimensional case is given by

$$p_c(1,\mu_0) = 1 + \frac{2}{\mu_0}.$$

Let us mention that, When  $\mu_0 > 1$ , the critical exponent  $p_c(1, \mu_0)$  is known to lie within the interval  $[1 + 2/\mu_0, 3]$ . However, the precise value of  $p_c(1, \mu_0)$  in this case remains an open question.

Finally, we generalize our problem to the following Cauchy problem

$$\begin{cases} \partial_t^2 u - \partial_x^2 u + V(x)\partial_t u = f(\partial_t u, \partial_x u), & x \in \mathbb{R}, t > 0, \\ u(x, 0) = u_0(x), & \partial_t u(x, 0) = u_1(x) & x \in \mathbb{R}, \end{cases}$$
(1.18) {General}

where  $V(x) = \mu_0 (1 + x^2)^{-1/2}, \ \mu_0 > 0,$ 

$$f(\partial_t u, \partial_x u) = |\partial_t u|^{\beta} |\partial_x u|^{\gamma},$$

with  $(\beta, \gamma) = (0, q)$  or  $(\beta, \gamma) = (p, q)$  with p, q > 1.

$$q > 1 + \frac{2}{\alpha_0}$$

where a := min(µ0, 1), then there exists a positive constant 0 < a ... (1, small enough, such that for any initial data satisfying

 $||u_0||_{H^2} + ||u_1||_{H^1} \le \varepsilon_0,$ 

problem (1.18) possesses a uniquely global mild solution

$$u \in \mathcal{C}([0,\infty), H^2(\mathbb{R})) \cap \mathcal{C}^1([0,\infty), H^1(\mathbb{R})) \cap \mathcal{C}^2([0,\infty), L^2(\mathbb{R})).$$

Moreover, the solution satisfies the following estimate

$$(1+t)^{-1} \|u(t)\|_{L^2} + \|\partial_x u(t)\|_{H^1} + \|\partial_t u(t)\|_{H^1} + \|\partial_t^2 u(t)\|_{L^2} \le C (1+t)^{-\frac{\alpha}{2}},$$

where  $\alpha$  is defined by (2.31).

 Global existence. Case of (*β*, *γ*) = (*p*, *q*))
 Assume that u<sub>0</sub> ∈ H<sup>2</sup>(ℝ) and u<sub>1</sub> ∈ H<sup>1</sup>(ℝ) which are compactly supported on (−R<sub>0</sub>, R<sub>0</sub>), with
 R<sub>0</sub> > 0. If (*β*, *γ*) = (*p*, *q*), *p*, *q* > 1, and

$$p+q > 1 + \frac{2}{\alpha_0},$$

طi the constant of the constant

 $||u_0||_{H^2} + ||u_1||_{H^1} \le \varepsilon_0,$ 

problem (1.18) possesses a uniquely global mild solution

$$u \in \mathcal{C}([0,\infty), H^2(\mathbb{R})) \cap \mathcal{C}^1([0,\infty), H^1(\mathbb{R})) \cap \mathcal{C}^2([0,\infty), L^2(\mathbb{R})).$$

Moreover, the solution satisfies the following estimate

$$(1+t)^{-1} \|u(t)\|_{L^2} + \|\partial_x u(t)\|_{H^1} + \|\partial_t u(t)\|_{H^1} + \|\partial_t^2 u(t)\|_{L^2} \le C (1+t)^{-\frac{\alpha}{2}},$$

where  $\alpha$  is defined by (2.31).

This paper is organized as follows: Section 2 is devoted to the energy estimates of the solution of the corresponding linear equation of (1.1). Section 3 is dedicated to the proof of the solution of the solution of the corresponding linear equation of (1.1). While the proof of the blow-up result (Theorem 1.2) is given in Section 5. Finally, Section 4 generalizes the global existence results to a class of nonlinearities (Theorems 1.3 and 1.4)

### 2 Homogeneous equation

 nmind that the asymptotic profile of solutions to the linear part of the equation influences the critical exponent for the problem, we consider the following homogeneous problem

$$\begin{cases} \partial_t^2 u - \partial_x^2 u + V(x) \partial_t u = 0, & x \in \mathbb{R}, \ t > 0, \\ u(x,0) = u_0(x), \ \partial_t u(x,0) = u_1(x), & x \in \mathbb{R}, \end{cases}$$
(2.1) [1]

where V(x) =  $\mu_0(1+x^2)^{-1/2}, \, \mu_0 > 0$ . In addition, we assume that  $(u_0, u_1)$  are compactly where  $V(x) = \mu_0(1+x^2)^{-1/2}, \, \mu_0 > 0$ . In addition, we assume that  $(u_0, u_1) \in H^2(\mathbb{R}) \times H^1(\mathbb{R})$ . To begin with, we give the definition of a strong solution to (2.1).

**Definition 2.1** (Strong solution). Let  $(u_0, u_1) \in H^2(\mathbb{R}) \times H^1(\mathbb{R})$ . A function u is said to be a strong solution to (2.1) if

$$u \in \mathcal{C}\left([0,\infty), H^2(\mathbb{R})\right) \cap \mathcal{C}^1\left([0,\infty), H^1(\mathbb{R})\right) \cap \mathcal{C}^2\left([0,\infty), L^2(\mathbb{R})\right),$$

and u has initial data  $u(x,0) = u_0(x)$ ,  $\partial_t u(x,0) = u_1(x)$  and satisfies the equation in (2.1) in the sense of  $L^2(\mathbb{R})$ .

**Theorem 2.1** (Theorem 2.27 in [19]).

Let  $m \in \mathbb{N}$ . For each  $(u_0, u_1) \in H^{m+2}(\mathbb{R}) \times H^{m+1}(\mathbb{R})$ , there exists a unique strong solution u to (2.1) such that  $supp u(t) \subset \{x \in R; |x| \leq t + R_0\}$  and

$$u \in \bigcap_{j=0}^{m+2} \mathcal{C}^{m+2-j}\left([0,\infty), H^j(\mathbb{R})\right).$$

{sec2}

{existencet}

This section is devoted to the statement and the proof of Proposition 2.5. It is divided into three subsections:

- In the first one, we give the two classical energy estimates following from the multiplication of the equation (2.1) by  $\partial_t u$  and u and using the fact that the initial data  $(u_0, u_1)$ are compactly supported on  $(-R_0, R_0)$ . Combining these estimates yields Lemma 2.2.
- The second subsection is devoted to improving the previous energy estimates.
- Subsequently, we combine the above energy estimates to conclude Proposition 2.5.

# **2.1** A bound of the $H^1 \times L^2$ norm of $(u, \partial_t u)$

This subsection presents Lemma 2.2, which relies on the construction of a Lyapunov functional. We begin by introducing three crucial new functionals, defined as follows

$$E_0(u(t),\partial_t u(t)) := \frac{1}{2} \int_{\mathbb{R}} \left( (\partial_x u(t))^2 + (\partial_t u(t))^2 \right) \mathrm{d}x, \qquad (2.2) \quad \{\mathrm{En}\}$$

$$I(u(t), \partial_t u(t), t) := \int_{\mathbb{R}} \left( u(t) \partial_t u(t) + \frac{1}{2(t+1)} u^2(t) + \frac{V(x)}{2} u^2(t) \right) \mathrm{d}x,$$
(2.3) {F2}

$$F_A(u(t), \partial_t u(t), t) := AE_0(u(t), \partial_t u(t)) + \frac{1}{2(t+1)}I(u(t), \partial_t u(t), t),$$
(2.4) [F4]

 where A > 0 is a constant that will be defined later on. By evaluating the time derivative of E\_0(u(t), \partial\_t u(t), t), I(u(t), \partial\_t u(t), t) and F\_A(u(t), \partial\_t u(t), t), we prove the following

**Lemma 2.2.** Assume  $(u_0, u_1) \in H^2(\mathbb{R}) \times H^1(\mathbb{R})$  that are compactly supported on  $(-R_0, R_0)$ , with  $R_0 > 0$ , then there exists a constant  $C_0 = C_0(R_0, \mu_0) > 0$  such that the strong solution uof (2.1) satisfies, for all  $t \ge s \ge 0$ ,

$$\frac{\|u(t)\|_{L^{2}}^{2}}{(1+t)^{2}} + \frac{\|\sqrt{V}u(t)\|_{L^{2}}^{2}}{1+t} + \|\partial_{t}u(t)\|_{L^{2}}^{2} + \|\partial_{x}u(t)\|_{L^{2}}^{2} \\
\leq C_{0}\left(\frac{\|u(s)\|_{L^{2}}^{2}}{(1+s)^{2}} + \frac{\|\sqrt{V}u(s)\|_{L^{2}}^{2}}{1+s} + \|\partial_{t}u(s)\|_{L^{2}}^{2} + \|\partial_{x}u(s)\|_{L^{2}}^{2}\right).$$

Proof. Since  $(u_0, u_1) \in H^2(\mathbb{R}) \times H^1(\mathbb{R})$ , thanks to Theorem 2.1, then the solution u of (2.1) satisfies  $u \in \bigcap_{j=0}^2 \mathcal{C}^j([0,\infty), H^{2-j}(\mathbb{R}))$ . Therefore,  $E_0(u(t), \partial_t u(t))$  is a differentiable function in time. Moreover, by multiplying (2.1) by  $\partial_t u(t)$ , and integration by parts, we get for all t > 0,

$$\frac{d}{dt}E_0(u(t),\partial_t u(t)) = -\int_{\mathbb{R}} V(x)(\partial_t u(t))^2 \mathrm{d}x.$$
(2.5) {E00}

Similarly,  $I(u(t), \partial_t u(t), t)$  is a differentiable function. In a similar way, by multiplying (2.1) by u(t), and using integration by parts, we get for all t > 0,

$$\frac{d}{dt}I(u(t),\partial_t u(t),t) = \int_{\mathbb{R}} (\partial_t u(t))^2 dx - \int_{\mathbb{R}} (\partial_x u(t))^2 dx \qquad (2.6) \quad \{F1\}$$

$$-\frac{1}{2(1+t)^2} \int_{\mathbb{R}} (u(t))^2 dx + \frac{1}{1+t} \int_{\mathbb{R}} u(t)\partial_t u(t) dx.$$

{LEE}

On the other hand, exploiting (2.5), (2.6) and the definition of  $F_A(u(t), \partial_t u(t), t)$  given by (2.4), we infer

$$\frac{d}{dt}F_{A}(u(t),\partial_{t}u(t),t) = -A\int_{\mathbb{R}}V(x)(\partial_{t}u(t))^{2}dx + \frac{1}{2(1+t)}\int_{\mathbb{R}}(\partial_{t}u(t))^{2}dx - \frac{1}{2(1+t)}\int_{\mathbb{R}}(\partial_{x}u(t))^{2}dx 
- \frac{1}{2(1+t)^{3}}\int_{\mathbb{R}}(u(t))^{2}dx - \frac{1}{4(1+t)^{2}}\int_{\mathbb{R}}V(x)(u(t))^{2}dx 
\leq -A\int_{\mathbb{R}}V(x)(\partial_{t}u(t))^{2}dx + \frac{1}{2(1+t)}\int_{\mathbb{R}}(\partial_{t}u(t))^{2}dx.$$

By using the fact that supp(u) is included in  $|x| \leq R_0 + t$ , we have

$$V(x) \ge \frac{\mu_0}{\sqrt{1 + (R_0 + t)^2}} \ge \frac{\mu_0}{R_0 + 1 + t} \ge \frac{\mu_0}{(1 + R_0)(1 + t)},$$

for any  $t \ge 0$ , therefore for all  $A \ge (1 + R_0)/2\mu_0$ , we have

$$\frac{d}{dt}F_A(u(t),\partial_t u(t),t) \leq -\left(\frac{A\mu_0}{1+R_0}-\frac{1}{2}\right)\int_{\mathbb{R}}\frac{(\partial_t u(t))^2}{1+t}\mathrm{d}x \leq 0.$$

So, by integrating over [s, t] we write

$$F_A(u(t), \partial_t u(t), t) \le F_A(u(s), \partial_t u(s), s), \quad \text{for any } t \ge s \ge 0.$$
(2.7) **{F19B}**

Furthermore, by the following Young's inequality

$$|ab| \le \frac{1}{4}a^2 + b^2$$
 with  $a = \frac{u}{1+t}, b = \partial_t u,$ 

we have

$$\frac{1}{2} \int_{\mathbb{R}} \frac{u(t)\partial_t u(t)}{1+t} \mathrm{d}x \ge -\frac{1}{8} \int_{\mathbb{R}} \frac{(u(t))^2}{(1+t)^2} \mathrm{d}x - \frac{1}{2} \int_{\mathbb{R}} (\partial_t u(t))^2 \mathrm{d}x,$$

therefore,

$$F_{A}(u(t), \partial_{t}u(t), t) \geq \frac{A}{2} \int_{\mathbb{R}} (\partial_{x}u(t))^{2} dx + \frac{A-1}{2} \int_{\mathbb{R}} (\partial_{t}u(t))^{2} dx + \frac{1}{8} \int_{\mathbb{R}} \frac{(u(t))^{2}}{(1+t)^{2}} dx + \frac{1}{4} \int_{\mathbb{R}} \frac{V(x)(u(t))^{2}}{1+t} dx.$$

By choosing  $A = \max(5/4, (1+R_0)/2\mu_0)$ , we arrive at

$$F_A(u(t), \partial_t u(t), t) \ge \frac{1}{8} \int_{\mathbb{R}} \left( \frac{(u(t))^2}{(t+1)^2} + \frac{V(x)(u(t))^2}{t+1} + (\partial_x u(t))^2 + (\partial_t u(t))^2 \right) \mathrm{d}x, \quad \text{for all } t \ge 0.$$

$$(2.8) \quad \{\text{F17B}\}$$

On the other hand, using the following Young's inequality

$$|ab| \le \frac{a^2}{2} + \frac{b^2}{2}$$
 with  $a = \frac{u}{1+t}, b = \partial_t u,$ 

we conclude that

$$F_A(u(t), \partial_t u(t), t) \le \frac{2A+1}{4} \int_{\mathbb{R}} \left( \frac{(u(t))^2}{(t+1)^2} + \frac{V(x)(u(t))^2}{t+1} + (\partial_x u(t))^2 + (\partial_t u(t))^2 \right) dx, \quad \text{for all } t \ge 0.$$
(2.9) **[F18B]**

By (2.8) and (2.9), there exists  $C = C(R_0, \mu_0)$  such that

$$C^{-1}F_{A}(u(t),\partial_{t}u(t),t) \leq \int_{\mathbb{R}} \left( \frac{(u(t))^{2}}{(t+1)^{2}} + \frac{V(x)(u(t))^{2}}{t+1} + (\partial_{x}u(t))^{2} + (\partial_{t}u(t))^{2} \right) dx \leq CF_{A}(u(t),\partial_{t}u(t),t),$$
for all  $t > 0$ . Combining (2.7), and (2.10), this ends the proof of Lemma 2.2.

for all  $t \ge 0$ . Combining (2.7), and (2.10), this ends the proof of Lemma 2.2.

#### Decay estimates of the $H^1 \times L^2$ norm of $(u, \partial_t u)$ for $H^2 \times H^1$ data 2.2

In order to improve the decay estimate in Lemma 2.2, we need to introduce different set of energy functionals which depend on the value of the parameter  $\mu_0$ . Therefore, we have two cases to distinguish.

First case:  $\mu_0 \in (0, 1]$ .

Let  $\mu \in (0, \mu_0)$ . Now, we introduce the following functionals

$$E_1(u(t), \partial_t u(t), t) := \int_{\mathbb{R}} \left( u(t) \partial_t u(t) + \frac{1-\mu}{2(t+1)} u^2(t) + \frac{V(x)}{2} u^2(t) \right) \mathrm{d}x, \quad (2.11) \quad \{\mathrm{In}\}$$

$$E_2(u(t), \partial_t u(t), t) := E_0(u(t), \partial_t u(t)) + \frac{\mu}{2(t+1)} E_1(u(t), \partial_t u(t), t),$$
(2.12) {E3}

{LE}

i = 1, 2, we prove the following

**Lemma 2.3.** Let  $\mu_0 \in (0,1]$  and  $\mu \in (0,\mu_0)$ . Assume  $(u_0,u_1) \in H^2(\mathbb{R}) \times H^1(\mathbb{R})$  that are compactly supported on  $(-R_0, R_0)$ , with  $R_0 > 0$ , then there exists a constant  $C_1 = C_1(R_0, \mu, \mu_0) > 0$ such that the strong solution u of (2.1) satisfies, for all  $t \ge s \ge 0$ ,

$$\frac{\|u(t)\|_{L^{2}}^{2}}{(1+t)^{2}} + \frac{\|\sqrt{V}u(t)\|_{L^{2}}^{2}}{1+t} + \|\partial_{t}u(t)\|_{L^{2}}^{2} + \|\partial_{x}u(t)\|_{L^{2}}^{2} 
\leq C_{1}\left(\frac{1+s}{1+t}\right)^{\mu}\left(\frac{\|u(s)\|_{L^{2}}^{2}}{(1+s)^{2}} + \frac{\|\sqrt{V}u(s)\|_{L^{2}}^{2}}{1+s} + \|\partial_{t}u(s)\|_{L^{2}}^{2} + \|\partial_{x}u(s)\|_{L^{2}}^{2}\right). \quad (2.13) \quad \text{(A1)}$$

t > 0,

$$\frac{d}{dt}E_1(u(t),\partial_t u(t),t) = \int_{\mathbb{R}} (\partial_t u(t))^2 dx - \int_{\mathbb{R}} (\partial_x u(t))^2 dx \qquad (2.14) \quad \{\text{E1}\}$$

$$-\frac{1-\mu}{2(1+t)^2} \int_{\mathbb{R}} (u(t))^2 dx + \frac{1-\mu}{1+t} \int_{\mathbb{R}} u(t)\partial_t u(t) dx.$$

Therefore, exploiting (2.5), (2.14) and the definition of  $E_2(u(t), \partial_t u(t), t)$  given by (2.12), we infer

$$\begin{aligned} \frac{d}{dt} E_2(u(t), \partial_t u(t), t) &= -\int_{\mathbb{R}} V(x) (\partial_t u(t))^2 dx + \frac{\mu}{2(1+t)} \int_{\mathbb{R}} (\partial_t u(t))^2 dx - \frac{\mu}{2(1+t)} \int_{\mathbb{R}} (\partial_x u(t))^2 dx \\ &- \frac{\mu(1-\mu)}{2(1+t)^3} \int_{\mathbb{R}} (u(t))^2 dx - \frac{\mu^2}{2(1+t)^2} \int_{\mathbb{R}} u(t) \partial_t u(t) dx \\ &- \frac{\mu}{4(1+t)^2} \int_{\mathbb{R}} V(x) (u(t))^2 dx. \end{aligned}$$

Therefore,

$$\begin{aligned} \frac{d}{dt} E_2(u(t), \partial_t u(t), t) &+ \frac{\mu}{1+t} E_2(u(t), \partial_t u(t), t) &= -\int_{\mathbb{R}} V(x) (\partial_t u(t))^2 dx + \frac{\mu}{1+t} \int_{\mathbb{R}} (\partial_t u(t))^2 dx \\ &- \frac{\mu (1-\mu)(2-\mu)}{4(1+t)^3} \int_{\mathbb{R}} (u(t))^2 dx \\ &- \frac{\mu (1-\mu)}{4(1+t)^2} \int_{\mathbb{R}} V(x) (u(t))^2 dx \\ &\leq -\int_{\mathbb{R}} V(x) (\partial_t u(t))^2 dx + \frac{\mu}{1+t} \int_{\mathbb{R}} (\partial_t u(t))^2 dx. \end{aligned}$$

$$\frac{d}{dt}E_2(u(t),\partial_t u(t),t) + \frac{\mu}{1+t}E_2(u(t),\partial_t u(t),t) \le -\int_{\mathbb{R}} \left(V(x) - \frac{\mu}{1+t}\right)(\partial_t u(t))^2 \mathrm{d}x \le 0.$$
(2.15) **[E12]**

Multiplying (2.15) by  $(1+t)^{\mu}$  and integrating over [s, t], we deduce that, for all  $t \ge s \ge t_0$ ,

$$(1+t)^{\mu}E_2(u(t),\partial_t u(t),t) \le (1+s)^{\mu}E_2(u(s),\partial_t u(s),s).$$
(2.16) {E14}

In addition, using Young's inequality, we have

$$\frac{\mu}{2} \int_{\mathbb{R}} \frac{u(t)\partial_t u(t)}{1+t} \mathrm{d}x \ge -\frac{\mu}{8} \int_{\mathbb{R}} \frac{(u(t))^2}{(1+t)^2} \mathrm{d}x - \frac{\mu}{2} \int_{\mathbb{R}} (\partial_t u(t))^2 \mathrm{d}x.$$

thus,

$$E_{2}(u(t), \partial_{t}u(t), t) \geq \frac{1}{2} \int_{\mathbb{R}} (\partial_{x}u(t))^{2} dx + \frac{1-\mu}{2} \int_{\mathbb{R}} (\partial_{t}u(t))^{2} dx + \frac{\mu - 2\mu^{2}}{8} \int_{\mathbb{R}} \frac{(u(t))^{2}}{(1+t)^{2}} dx + \frac{\mu}{8} \int_{\mathbb{R}} \frac{V(x)(u(t))^{2}}{1+t} dx + \frac{\mu}{8} \int_{\mathbb{R}} \frac{V(x)(u(t))^{2}}{1+t} dx.$$

Plugging the above inequality into the fact that  $V(x) \ge \mu/(1+t)$  for all  $t \ge t_0$ , we obtain

$$E_2(u(t), \partial_t u(t), t) \ge \frac{1}{2} \int_{\mathbb{R}} (\partial_x u(t))^2 \mathrm{d}x + \frac{1-\mu}{2} \int_{\mathbb{R}} (\partial_t u(t))^2 \mathrm{d}x + \frac{\mu(1-\mu)}{8} \int_{\mathbb{R}} \frac{(u(t))^2}{(1+t)^2} \mathrm{d}x + \frac{\mu}{8} \int_{\mathbb{R}} \frac{V(x)(u(t))^2}{1+t} \mathrm{d}x$$

Consequently, we have

$$E_2(u(t), \partial_t u(t), t) \ge C \int_{\mathbb{R}} \left( \frac{(u(t))^2}{(t+1)^2} + \frac{V(x)(u(t))^2}{t+1} + (\partial_x u(t))^2 + (\partial_t u(t))^2 \right) \mathrm{d}x, \quad \text{for all } t \ge t_0.$$
(2.17) **[E17]**

On the other hand, by Young's inequality, we conclude that there exists  $C = C(\mu)$  such that

$$E_2(u(t), \partial_t u(t), t) \le C \int_{\mathbb{R}} \left( \frac{(u(t))^2}{(t+1)^2} + \frac{V(x)(u(t))^2}{t+1} + (\partial_x u(t))^2 + (\partial_t u(t))^2 \right) \mathrm{d}x, \quad \text{for all } t \ge 0.$$
(2.18) {E18}

By (2.17) and (2.18), there exists  $C = C(\mu)$  such that

$$C^{-1}E_{2}(u(t),\partial_{t}u(t),t) \leq \int_{\mathbb{R}} \left( \frac{(u(t))^{2}}{(t+1)^{2}} + \frac{V(x)(u(t))^{2}}{t+1} + (\partial_{x}u(t))^{2} + (\partial_{t}u(t))^{2} \right) \mathrm{d}x \leq CE_{2}(u(t),\partial_{t}u(t),t),$$
(2.19) **[E15]**

for all  $t \ge t_0$ . Finally, using (2.16), and (2.19), we deduce that

$$\int_{\mathbb{R}} \left( \frac{(u(t))^{2}}{(t+1)^{2}} + \frac{V(x)(u(t))^{2}}{t+1} + (\partial_{x}u(t))^{2} + (\partial_{t}u(t))^{2} \right) \mathrm{d}x$$

$$\leq \tilde{C} \left( \frac{1+s}{1+t} \right)^{\mu} \int_{\mathbb{R}} \left( \frac{(u(s))^{2}}{(s+1)^{2}} + \frac{V(x)(u(s))^{2}}{s+1} + (\partial_{x}u(s))^{2} + (\partial_{t}u(s))^{2} \right) \mathrm{d}x, \quad (2.20) \quad \{\text{E16}\}$$

for all  $t \ge s \ge t_0$ . Furthermore, by exploiting Lemma 2.2, we deduce

$$\frac{\|u(t)\|_{L^{2}}^{2}}{(1+t)^{2}} + \frac{\|\sqrt{V}u(t)\|_{L^{2}}^{2}}{1+t} + \|\partial_{t}u(t)\|_{L^{2}}^{2} + \|\partial_{x}u(t)\|_{L^{2}}^{2} \\
\leq \tilde{C}_{1}\left(\frac{1+s}{1+t}\right)^{\mu}\left(\frac{\|u(s)\|_{L^{2}}^{2}}{(1+s)^{2}} + \frac{\|\sqrt{V}u(s)\|_{L^{2}}^{2}}{1+s} + \|\partial_{t}u(s)\|_{L^{2}}^{2} + \|\partial_{x}u(s)\|_{L^{2}}^{2}\right), \quad (2.21) \quad \{\text{E16C}\}$$

Second case:  $\mu_0 \in (1, \infty)$ . We introduce the following functionals:

$$E_{3}(u(t), \partial_{t}u(t)) := \int_{\mathbb{R}} \left( u(t)\partial_{t}u(t) + \frac{V(x)}{2}u^{2}(t) \right) \mathrm{d}x,$$
  

$$E_{4}(u(t), \partial_{t}u(t), t) := E_{0}(u(t), \partial_{t}u(t)) + \frac{1}{2(t+1)}E_{3}(u(t), \partial_{t}u(t)),$$

where  $E_0(u(t), \partial_t u(t))$  is defined by (2.2).

$$\frac{\|u(t)\|_{L^{2}}^{2}}{(1+t)^{2}} + \frac{\|\sqrt{V}u(t)\|_{L^{2}}^{2}}{1+t} + \|\partial_{t}u(t)\|_{L^{2}}^{2} + \|\partial_{x}u(t)\|_{L^{2}}^{2} \\
\leq C\left(\frac{1+s}{1+t}\right)\left(\frac{\|u(s)\|_{L^{2}}^{2}}{(1+s)^{2}} + \frac{\|\sqrt{V}u(s)\|_{L^{2}}^{2}}{1+s} + \|\partial_{t}u(s)\|_{L^{2}}^{2} + \|\partial_{x}u(s)\|_{L^{2}}^{2}\right). \quad (2.22) \quad \{A1A\}$$

$$\frac{d}{dt}E_3(u(t),\partial_t u(t)) = \int_{\mathbb{R}} (\partial_t u(t))^2 \mathrm{d}x - \int_{\mathbb{R}} (\partial_x u(t))^2 \mathrm{d}x.$$
(2.23) {E3A}

Hence, by exploiting (2.5), (2.23) and the definition of  $E_4(u(t), \partial_t u(t), t)$ , we infer

$$\frac{d}{dt}E_4(u(t),\partial_t u(t),t) = -\int_{\mathbb{R}} V(x)(\partial_t u(t))^2 dx + \frac{1}{2(1+t)} \int_{\mathbb{R}} (\partial_t u(t))^2 dx - \frac{1}{2(1+t)} \int_{\mathbb{R}} (\partial_x u(t))^2 dx - \frac{1}{2(1+t)^2} \int_{\mathbb{R}} u(t)\partial_t u(t) dx - \frac{1}{4(1+t)^2} \int_{\mathbb{R}} V(x)(u(t))^2 dx.$$

Consequently,

$$\frac{d}{dt}E_4(u(t),\partial_t u(t),t) + \frac{1}{1+t}E_4(u(t),\partial_t u(t),t) = -\int_{\mathbb{R}} \left(V(x) - \frac{1}{1+t}\right) (\partial_t u(t))^2 \mathrm{d}x.$$

$$\frac{d}{dt}E_4(u(t), \partial_t u(t), t) + \frac{1}{1+t}E_4(u(t), \partial_t u(t), t) \le 0, \quad \text{for all } t \ge t_1.$$
(2.24) **[E12A]**

Multiplying (2.24) by (1 + t) and integrating over [s, t], we deduce that, for all  $t \ge s \ge t_1$ ,

$$(1+t)E_4(u(t),\partial_t u(t),t) \le (1+s)E_4(u(s),\partial_t u(s),s).$$
(2.25) {E14A}

In addition, by using the following Young's inequality

$$|ab| \le \frac{\varepsilon}{2}a^2 + \frac{1}{2\varepsilon}b^2$$
 with  $a = \frac{u}{1+t}, b = \partial_t u_t$ 

we have

$$\frac{1}{2} \int_{\mathbb{R}} \frac{u \,\partial_t u}{1+t} \mathrm{d}x \ge -\frac{\varepsilon}{4} \int_{\mathbb{R}} \frac{u^2}{(1+t)^2} \mathrm{d}x - \frac{1}{4\varepsilon} \int_{\mathbb{R}} (\partial_t u(t))^2 \mathrm{d}x,$$

{LEA}

thus, by using the fact that  $V(x) \ge 1/(1+t)$  for all  $t \ge t_1$ , we conclude that

$$E_{4}(u(t),\partial_{t}u(t),t) \geq \frac{1}{2} \int_{\mathbb{R}} (\partial_{x}u(t))^{2} dx + \frac{1}{2} \left(1 - \frac{1}{2\varepsilon}\right) \int_{\mathbb{R}} (\partial_{t}u(t))^{2} dx + \frac{1}{16} \int_{\mathbb{R}} \frac{V(x)u^{2}(t)}{1+t} dx + \frac{3}{16} \int_{\mathbb{R}} \frac{V(x)u^{2}(t)}{1+t} dx - \frac{\varepsilon}{4} \int_{\mathbb{R}} \frac{u^{2}(t)}{(1+t)^{2}} dx \geq \frac{1}{2} \int_{\mathbb{R}} (\partial_{x}u(t))^{2} dx + \frac{1}{2} \left(1 - \frac{1}{2\varepsilon}\right) \int_{\mathbb{R}} (\partial_{t}u(t))^{2} dx + \frac{1}{16} \int_{\mathbb{R}} \frac{V(x)u^{2}(t)}{1+t} dx + \frac{3-4\varepsilon}{16} \int_{\mathbb{R}} \frac{u^{2}(t)}{(1+t)^{2}} dx.$$

By choosing  $\varepsilon = 5/8$ , we obtain

$$E_4(u(t), \partial_t u(t), t) \ge \frac{1}{32} \int_{\mathbb{R}} \left( \frac{(u(t))^2}{(t+1)^2} + \frac{V(x)(u(t))^2}{t+1} + (\partial_x u(t))^2 + (\partial_t u(t))^2 \right) \mathrm{d}x, \quad \text{for all } t \ge t_1.$$
(2.26) **{E17A}**

On the other hand, using Young's inequality, we conclude that

$$E_4(u(t), \partial_t u(t), t) \le \frac{3}{4} \int_{\mathbb{R}} \left( \frac{(u(t))^2}{(t+1)^2} + \frac{V(x)(u(t))^2}{t+1} + (\partial_x u(t))^2 + (\partial_t u(t))^2 \right) \mathrm{d}x, \quad \text{for all } t \ge t_1.$$
(2.27) **{E18A}**

By (2.26) and (2.27),

$$C^{-1}E_4(u(t),\partial_t u(t),t) \le \int_{\mathbb{R}} \left( \frac{(u(t))^2}{(t+1)^2} + \frac{V(x)(u(t))^2}{t+1} + (\partial_x u(t))^2 + (\partial_t u(t))^2 \right) dx \le CE_4(u(t),\partial_t u(t),t),$$
(2.28) {E15A}

for all  $t \ge t_1$ . Finally, using (2.25), and (2.28), we deduce that

$$\int_{\mathbb{R}} \left( \frac{(u(t))^2}{(t+1)^2} + \frac{V(x)(u(t))^2}{t+1} + (\partial_x u(t))^2 + (\partial_t u(t))^2 \right) dx \\
\leq C \left( \frac{1+s}{1+t} \right) \int_{\mathbb{R}} \left( \frac{(u(s))^2}{(s+1)^2} + \frac{V(x)(u(s))^2}{s+1} + (\partial_x u(s))^2 + (\partial_t u(s))^2 \right) dx, \quad (2.29) \quad \{\text{E16A}\}$$

for some constant C and all  $t \ge s \ge t_1$ . Now, by exploiting Lemma 2.2, we arrive at

$$\frac{\|u(t)\|_{L^{2}}^{2}}{(1+t)^{2}} + \frac{\|\sqrt{V}u(t)\|_{L^{2}}^{2}}{1+t} + \|\partial_{t}u(t)\|_{L^{2}}^{2} + \|\partial_{x}u(t)\|_{L^{2}}^{2} \\
\leq \tilde{C}_{2}\left(\frac{1+s}{1+t}\right)\left(\frac{\|u(s)\|_{L^{2}}^{2}}{(1+s)^{2}} + \frac{\|\sqrt{V}u(s)\|_{L^{2}}^{2}}{1+s} + \|\partial_{t}u(s)\|_{L^{2}}^{2} + \|\partial_{x}u(s)\|_{L^{2}}^{2}\right), \quad (2.30) \quad \{\text{E16D}\}$$

### **2.3** Decay estimates of the $H^1 \times L^2$ norm of $(u, \partial_t u)$ for $H^1 \times L^2$ data

Based on the Lemmas proved in the previous subsections, we can now proceed to prove the main result of this section, namely Proposition 2.5.

ਂ In order to prove this proposition, let us denote by R(t,s) the operator which maps the initial data (u(s), u<sub>t</sub>(s))  $\in$  H<sup>2</sup>(model), let us denote us denote by R(t,s) the operator which maps the initial data (u(s), u<sub>t</sub>(s))  $\in$  H<sup>2</sup>(model), let us denote the us denote the proposition, let us denote the let us de

Let  $\mu \in (0, \mu_0)$  and let

$$\alpha := \alpha(\mu, \mu_0) = \begin{cases} \mu & \text{if } \mu_0 \in (0, 1], \\ 1 & \text{if } \mu_0 > 1. \end{cases}$$
(2.31) {alpha} (2.31) {prop2.1}

đ Proposition 2.5. Assume (u\_0, u\_1) \in H^1(\mathbb{R}) \times L^2(\mathbb{R}) that are compactly supported on (-R<sub>0</sub>, R<sub>0</sub>), with R<sub>0</sub> > 0, then there exist a constant C = C(R<sub>0</sub>, \alpha, \alpha\_0) > 0 and a unique mild solution

$$u \in \mathcal{C}\left([0,\infty), H^1(\mathbb{R})\right) \cap \mathcal{C}^1\left([0,\infty), L^2(\mathbb{R})\right),$$

of (2.1) satisfies

$$\frac{\|u(t)\|_{L^{2}}^{2}}{(1+t)^{2}} + \frac{\|\sqrt{V}u(t)\|_{L^{2}}^{2}}{1+t} + \|\partial_{t}u(t)\|_{L^{2}}^{2} + \|\partial_{x}u(t)\|_{L^{2}}^{2} \\
\leq C\left(\frac{1+s}{1+t}\right)^{\alpha}\left(\frac{\|u(s)\|_{L^{2}}^{2}}{(1+s)^{2}} + \frac{\|\sqrt{V}u(s)\|_{L^{2}}^{2}}{1+s} + \|\partial_{t}u(s)\|_{L^{2}}^{2} + \|\partial_{x}u(s)\|_{L^{2}}^{2}\right), \quad (2.32) \quad \text{(DA1)}$$

$$\begin{aligned} \frac{\|\sqrt{V}\partial_t u(t)\|_{L^2}^2}{1+t} + \|\partial_t^2 u(t)\|_{L^2}^2 + \|\partial_{xt}^2 u(t)\|_{L^2}^2 + \|\partial_x^2 u(t)\|_{L^2}^2 \\ &\leq C\left(\frac{1+s}{1+t}\right)^{\alpha} \left(\frac{\|\partial_t u(s)\|_{L^2}^2}{(1+s)^2} + \frac{\|\sqrt{V}\partial_t u(s)\|_{L^2}^2}{1+s} + \|\partial_t^2 u(s)\|_{L^2}^2 + \|\partial_x \partial_t u(s)\|_{L^2}^2\right), (2.33) \end{aligned}$$

for all  $t \geq s \geq 0$ .

*Proof.* Existence and Uniqueness. Let  $T_0 > 0$  be an arbitrary number, and let  $(u_0, u_1) \in H^1(\mathbb{R}) \times L^2(\mathbb{R})$ . By the density argument, there exist sequences

$$\left\{ (u_0^{(j)}, u_1^{(j)}) \right\}_{j=1}^{\infty} \subseteq H^2(\mathbb{R}) \times H^1(\mathbb{R}),$$

such that  $supp(u_0^{(j)})$ ,  $supp(u_1^{(j)}) \subseteq (-1 - R_0, 1 + R_0)$ ,  $1 \le j \le \infty$ , and  $\lim_{j \to \infty} (u_0^{(j)}, u_1^{(j)}) = (u_0, u_1) \text{ in } H^1(\mathbb{R}) \times L^2(\mathbb{R}).$ 

$$\begin{cases} \partial_t^2 u - \partial_x^2 u + \frac{\mu}{1+t} \partial_t u = 0, & x \in \mathbb{R}, \ t > 0 \\ u(x,0) = u_0^{(j)}(x) - u_0^{(k)}(x), \ \partial_t u(x,0) = u_1^{(j)}(x) - u_1^{(k)}(x), & x \in \mathbb{R}. \end{cases}$$

Apply Lemmas 2.3, and 2.4, to  $u^{(j)} - u^{(k)}$ , we have in particular

$$\begin{aligned} &\|u^{(j)} - u^{(k)}\|_{L^{2}}^{2} + \|\partial_{t}(u^{(j)} - u^{(k)})\|_{L^{2}}^{2} + \|\partial_{x}(u^{(j)} - u^{(k)})\|_{L^{2}}^{2} \\ &\leq C(1 + T_{0})^{2-\alpha} \left( \|u^{(j)}_{0} - u^{(k)}_{0}\|_{L^{2}}^{2} + \|u^{(j)}_{1} - u^{(k)}_{1}\|_{L^{2}}^{2} + \|\partial_{x}(u^{(j)}_{0} - u^{(k)}_{0})\|_{L^{2}}^{2} \right), \end{aligned}$$

ι in the section of the

$$\lim_{j \to \infty} u^{(j)} = u \in C([0,\infty); H^1(\mathbb{R})) \cap C^1([0,\infty); L^2(\mathbb{R})),$$
(2.34) {eq2.9}

since  $T_0 > 0$  is arbitrary. As  $u^{(j)}$  satisfies  $u^{(j)}(t, x) = R(t)(u_0^{(j)}, u_1^{(j)})$ , then

$$u(t) = \lim_{j \to \infty} R(t)(u_0^j, u_1^j),$$

this means that the operator R(t) has been extended uniquely to a new operator

$$\widetilde{R}(t): H^1(\mathbb{R}) \times L^2(\mathbb{R}) \longrightarrow X_0$$

$$(u_0, u_1) \longmapsto u(t)$$
(2.35) {Extend.Operative of the second second

also denoted by R(t), where

$$X_0 := \mathcal{C}\left([0,\infty), H^1(\mathbb{R})\right) \cap \mathcal{C}^1\left([0,\infty), L^2(\mathbb{R})\right),$$

which indicates that  $u(t) = R(t)(u_0, u_1)$  is a unique mild solution of (2.1).

**Energy estimate** (2.32). By Lemmas 2.3, and 2.4, each strong solution u<sup>(j)</sup> constructed above satisfies the estimates (2.13), (2.22), where R<sub>0</sub> has been replaced by R<sub>0</sub> + 1. By letting j → ∞ and using (2.34), the same estimates hold for the mild solution u, and we deduce that estimate (2.32) holds.

**Energy estimate** (2.33). Let  $1 \le i \le n$ . A straightforward computation implies that  $v = \partial_t u$  satisfies

$$\begin{cases} \partial_t^2 v - \partial_x^2 v + V(x) \partial_t v = 0, & x \in \mathbb{R}, \ t > 0, \\ v(x,0) = v_0(x), \ \partial_t v(x,0) = v_1(x), & x \in \mathbb{R}, \end{cases}$$
(2.36) {ut}

ろうろうろうろうろう Computer Co

$$\frac{\|v(t)\|_{L^{2}}^{2}}{(1+t)^{2}} + \frac{\|\sqrt{V}v(t)\|_{L^{2}}^{2}}{1+t} + \|\partial_{t}v(t)\|_{L^{2}}^{2} + \|\partial_{x}v(t)\|_{L^{2}}^{2}$$
  
$$\leq C\left(\frac{1+s}{1+t}\right)^{\alpha} \left(\frac{\|v(s)\|_{L^{2}}^{2}}{(1+s)^{2}} + \frac{\|\sqrt{V}v(s)\|_{L^{2}}^{2}}{1+s} + \|\partial_{t}v(s)\|_{L^{2}}^{2} + \|\partial_{x}v(s)\|_{L^{2}}^{2}\right),$$

which implies, together with  $\partial_x^2 u = \partial_t^2 u + V(x)\partial_t u$ , the estimate (2.33). This concludes the proof of Proposition 2.5.

**Remark 2.1.** Similarly to the existence part in the proof of Proposition 2.5 and using Lemmas 2.3, and 2.4, the operator R(t, s) and S(t, s) can be extended uniquely such that

$$R(t,s): H^1(\mathbb{R}) \times L^2(\mathbb{R}) \longrightarrow X_s$$
 (2.37) {Extend.Operation (2.37)

where

$$X_s := \mathcal{C}\left([s,\infty), H^1(\mathbb{R})\right) \cap \mathcal{C}^1\left([s,\infty), L^2(\mathbb{R})\right).$$

# 3 Proof of Theorems 1.1

This Section is devoted to the prove Theorem 1.1. To handle the nonlinear term in the proofs, we recall the following elementary inequalities and lemmas that are useful in our proofs.

**Lemma 3.1.** Let  $p \ge 1, a, b \in \mathbb{R}$ . Then

$$||a|^{p} - |b|^{p}| \leq C(|a|^{p-1} + |b|^{p-1})|a - b|, \qquad (3.1) \quad \text{{ab}}$$

$$||a|^{p-1}a - |b|^{p-1}b| \leq C(|a|^{p-1} + |b|^{p-1})|a - b|.$$
(3.2) {abb}

Then, we are in the position to state and prove the following nonlinear estimates.

$$|F(u(x,t)) - F(v(x,t))| \le C(|u(x,t)|^{p-1} + |v(x,t)|^{p-1})|u(x,t) - v(x,t)|, \quad \forall x \in \mathbb{R}, \ t > 0, \ (3.3)$$

$$|G(u(x,t)) - G(v(x,t))| \le C(|u(x,t)|^{p-1} + |v(x,t)|^{p-1})|u(x,t) - v(x,t)|, \quad \forall x \in \mathbb{R}, \ t > 0, \ (3.4)$$

$$\begin{aligned} & \left| \partial_x \left[ F(u(x,t)) \right] - \partial_x \left[ F(v(x,t)) \right] \right| & (3.5) \\ & \leq C \left| u(x,t) \right|^{p-1} \left| \partial_x (u(x,t) - v(x,t)) \right| + C \left| \partial_x v(x,t) \right| (|u(x,t)|^{p-2} + |v(x,t)|^{p-2}) |u(x,t) - v(x,t)|, \end{aligned}$$

for all  $x \in \mathbb{R}, t > 0$ ,

$$\begin{aligned} & \left| \partial_x \big[ G(u(x,t)) \big] - \partial_x \big[ G(v(x,t)) \big] \big| & (3.6) \\ & \leq C \ |u(x,t)|^{p-1} |\partial_x (u(x,t) - v(x,t))| + C |\partial_x v(x,t)| (|u(x,t)|^{p-2} + |v(x,t)|^{p-2}) |u(x,t) - v(x,t)|, \end{aligned} \end{aligned}$$

{handle}

{sec3}

{basic}

{rmk2.1}

a.e.  $x \in \mathbb{R}, t > 0$ ,

$$\begin{aligned} & \left|\partial_t \big[ F(u(x,t)) \big] - \partial_t \big[ F(v(x,t)) \big] \big| & (3.7) \\ & \leq C \ |u(x,t)|^{p-1} |\partial_t (u(x,t) - v(x,t))| + C |\partial_t v(x,t)| (|u(x,t)|^{p-2} + |v(x,t)|^{p-2}) |u(x,t) - v(x,t)|, \end{aligned}$$

for all  $x \in \mathbb{R}$ , t > 0, and

$$\begin{aligned} & \left| \partial_t \big[ G(u(x,t)) \big] - \partial_t \big[ G(v(x,t)) \big] \big| & (3.8) \\ & \leq C \ |u(x,t)|^{p-1} |\partial_t (u(x,t) - v(x,t))| + C |\partial_t v(x,t)| (|u(x,t)|^{p-2} + |v(x,t)|^{p-2}) |u(x,t) - v(x,t)|, \end{aligned} \end{aligned}$$

a.e.  $x \in \mathbb{R}, t > 0$ .

*Proof.* Note that (3.3) follows from (3.1). A straightforward calculation implies

$$\begin{aligned} \partial_x F(u(x,t)) &- \partial_x F(v(x,t)) &= p\Big(|u(x,t)|^{p-2}u(x,t)\partial_x u(x,t) - |v(x,t)|^{p-2}v(x,t)\partial_x v(x,t)\Big) \\ &= p|u(x,t)|^{p-2}u(x,t)(\partial_x u(x,t) - \partial_x v(x,t)) \\ &+ p\partial_x v(x,t)(|u(x,t)|^{p-2}u(x,t) - |v(x,t)|^{p-2}v(x,t)), \quad (3.9) \end{aligned}$$

for all  $x \in \mathbb{R}$ , t > 0. Consequently, we derive

$$\left|\partial_{x} \left[F(u)\right] - \partial_{x} \left[F(v)\right]\right| \leq C \left|u\right|^{p-1} \left|\partial_{x} u - \partial_{x} v\right| + C \left|\partial_{x} v\right| \left||u|^{p-2} u - |v|^{p-2} v\right|, \quad (3.10) \quad \text{{21d}}$$

寓, popp) popp)

Now, we are ready to give the

$$X(T) = \mathcal{C}([0,T], H^{2}(\mathbb{R})) \cap \mathcal{C}^{1}([0,T], H^{1}(\mathbb{R})) \cap \mathcal{C}^{2}([0,T], L^{2}(\mathbb{R})),$$

equipped with the norm

$$\|v\|_{X(T)} = \sup_{0 \le t \le T} (1+t)^{\frac{\alpha}{2}} \left\{ (1+t)^{-1} \|v(t)\|_{L^2} + \|\partial_x v(t)\|_{H^1} + \|\partial_t v(t)\|_{H^1} + \|\partial_t^2 v(t)\|_{L^2} \right\}$$

ਡ in indiceses ind

$$\Phi(u)(t) = R(t)(u_0, u_1) + \int_0^t S(t, s) f(\partial_t u) \, ds =: \Phi(u)^{lin}(t) + \Phi(u)^{nl}(t), \quad \text{for } u \in B_M(T).$$

Our goal is to prove that  $\Phi$  is a contraction mapping on  $B_M(T)$ .

Step 1.  $\Phi: B_M(T) \longrightarrow B_M(T)$ . Let  $u \in B_M(T)$ . Estimation of  $(1+t)^{-1} \| \Phi(u)(t) \|_{L^2} + \| \partial_t \Phi(u)(t) \|_{L^2} + \| \partial_x \Phi(u)(t) \|_{L^2}$ . Using (2.32) in the particular case where s = 0, we have

$$(1+t)^{-1} \|\Phi(u)^{lin}(t)\|_{L^2} + \|\partial_x \Phi(u)^{lin}(t)\|_{L^2} + \|\partial_t \Phi(u)^{lin}(t)\|_{L^2} \le C \varepsilon_0 (1+t)^{-\frac{\alpha}{2}}.$$
 (3.11) (3.11)

Moreover, by using (2.32) again in the case where the initial data is  $(0, f(\partial_t u(s)))$ , we obtain

$$(1+t)^{-1} \|\Phi(u)^{nl}(t)\|_{L^2} + \|\partial_t \Phi(u)^{nl}(t)\|_{L^2} + \|\partial_x \Phi(u)^{nl}(t)\|_{L^2} \le C \int_0^t \left(\frac{1+s}{1+t}\right)^{\frac{\alpha}{2}} \|f(\partial_t u(s))\|_{L^2} \, ds.$$

By the Sobolev embedding  $H^1(\mathbb{R}) \hookrightarrow L^{\infty}(\mathbb{R})$ , and the fact that  $u \in B_M(T)$ , we infer

$$\|\partial_t u(s)\|_{L^{\infty}} \leqslant C \|\partial_t u(s)\|_{H^1} \leqslant C(1+s)^{-\frac{\alpha}{2}} \|u\|_{X(T)} \le CM(1+s)^{-\frac{\alpha}{2}}.$$
 (3.12) {SO1}

Therefore, by (3.12) and exploiting again the fact that  $u \in B_M(T)$ , we conclude

$$\|f(\partial_t u(s))\|_{L^2} \le \|\partial_t u(s)\|_{\infty}^{p-1} \|\partial_t u(s)\|_{L^2} \le CM^p (1+s)^{-\frac{\alpha p}{2}}.$$
(3.13) {An1}

Then

$$(1+t)^{-1} \|\Phi(u)^{nl}(t)\|_{L^2} + \|\partial_t \Phi(u)^{nl}(t)\|_{L^2} + \|\partial_x \Phi(u)^{nl}(t)\|_{L^2} \le C M^p (1+t)^{-\frac{\alpha}{2}} \int_0^t (1+s)^{-\frac{(p-1)\alpha}{2}} ds.$$

$$\frac{(p-1)\alpha}{2} > 1. \tag{3.14} \quad \texttt{(alpha-alpha)}$$

Therefore

$$(1+t)^{-1} \|\Phi(u)^{nl}(t)\|_{L^2} + \|\partial_t \Phi(u)^{nl}(t)\|_{L^2} + \|\partial_x \Phi(u)^{nl}(t)\|_{L^2} \le C M^p (1+t)^{-\frac{\alpha}{2}}.$$
 (3.15) **(G200)**

So, by (3.11), and (3.15), we get

$$(1+t)^{-1} \|\Phi(u)(t)\|_{L^2} + \|\partial_t \Phi(u)(t)\|_{L^2} + \|\partial_x \Phi(u)(t)\|_{L^2} \le C \varepsilon_0 (1+t)^{-\frac{\alpha}{2}} + C M^p (1+t)^{-\frac{\alpha}{2}}.$$
 (3.16) **{G2T}**  
Estimation of  $\|\partial_{xt}^2 \Phi(u)(t)\|_{L^2} + \|\partial_t^2 \Phi(u)(t)\|_{L^2} + \|\partial_x^2 \Phi(u)(t)\|_{L^2}.$  By (2.33), we have

$$\|\partial_x^2 \Phi(u)^{lin}(t)\|_{L^2} + \|\partial_{xt}^2 \Phi(u)^{lin}(t)\|_{L^2} + \|\partial_t^2 \Phi(u)^{lin}(t)\|_{L^2} \le C \varepsilon_0 (1+t)^{-\frac{\alpha}{2}}.$$
(3.17) **{H6}**

Moreover, by using (2.33) again in the case where the initial data is  $(0, f(\partial_t u(s)))$ , we obtain

$$\begin{aligned} \|\partial_x^2 \Phi(u)^{nl}(t)\|_{L^2} &+ \|\partial_{xt}^2 \Phi(u)^{nl}(t)\|_{L^2} + \|\partial_t^2 \Phi(u)^{nl}(t)\|_{L^2} \\ &\leq C \, \int_0^t \left(\frac{1+s}{1+t}\right)^{\frac{\alpha}{2}} \left(\|f(\partial_t u(s))\|_{H^1} + \|\partial_t f(\partial_t u(s))\|_{L^2}\right) \, ds. \end{aligned} \tag{3.18}$$

Based on (3.12) and exploiting again the fact that  $u \in B_M(T)$ , we have

$$\|\partial_x f(\partial_t u(s))\|_{L^2} = \|p|\partial_t u\|_{L^2}^{p-1} \partial_{xt}^2 u\|_{L^2} \le p\|\partial_t u\|_{\infty}^{p-1} \|\partial_{xt}^2 u\|_{L^2} \le CM^p (1+s)^{-\frac{p\alpha}{2}}, \tag{3.19}$$

and

$$\|\partial_t f(\partial_t u(s))\|_{L^2} = \|p|\partial_t u|^{p-1}\partial_t^2 u\|_{L^2} \le p\|\partial_t u\|_{\infty}^{p-1}\|\partial_t^2 u\|_{L^2} \le CM^p(1+s)^{-\frac{p\alpha}{2}}.$$
 (3.20) **{Bn2**

Hence, by (3.13), (3.19), (3.20) and (3.18), we infer

$$\|\partial_x^2 \Phi(u)^{nl}(t)\|_{L^2} + \|\partial_{xt}^2 \Phi(u)^{nl}(t)\|_{L^2} + \|\partial_t^2 \Phi(u)^{nl}(t)\|_{L^2} \le C M^p (1+t)^{-\frac{\alpha}{2}} \int_0^t (1+s)^{-\frac{(p-1)\alpha}{2}} ds.$$
(3.21)

From (3.14), it follows that

$$\|\partial_x^2 \Phi(u)^{nl}(t)\|_{L^2} + \|\partial_{xt}^2 \Phi(u)^{nl}(t)\|_{L^2} + \|\partial_t^2 \Phi(u)^{nl}(t)\|_{L^2} \le C M^p (1+t)^{-\frac{\alpha}{2}}.$$
(3.22) [7mai3]

Therefore, by (3.17), and (3.22), we get

$$\|\partial_x^2 \Phi(u)(t)\|_{L^2} + \|\partial_{xt}^2 \Phi(u)(t)\|_{L^2} + \|\partial_t^2 \Phi(u)(t)\|_{L^2} \le C \varepsilon_0 (1+t)^{-\frac{\alpha}{2}} + C M^p (1+t)^{-\frac{\alpha}{2}}.$$
 (3.23) [G2000]

Summing up the estimates (3.16) and (3.23), we conclude that

$$\|\Phi(u)\|_{X(T)} \le C \varepsilon_0 + C M^p.$$

<br/> By choosing M > 0 such that C M^{p-1} < 1/2, and then 0 < < () < 1/2 such that C < () < 1/2 M, we arrive at

$$\|\Phi(u)\|_{X(T)} \le M$$

i.e.  $\Phi(u) \in B_M(T)$ .

$$(1+t)^{-1} \|\Phi(u)(t) - \Phi(v)(t)\|_{L^{2}} + \|\partial_{t}\Phi(u)(t) - \partial_{t}\Phi(v)(t)\|_{L^{2}} + \|\partial_{x}\Phi(u)(t) - \partial_{x}\Phi(v)(t)\|_{L^{2}} \\ \leq C \int_{0}^{t} \left(\frac{1+s}{1+t}\right)^{\frac{\alpha}{2}} \|f(\partial_{t}u(s)) - f(\partial_{t}v(s))\|_{L^{2}} \, ds. \quad (3.24)$$

By means of the basic inequalities (3.1)-(3.2), we obtain

$$\|f(\partial_t u(s)) - f(\partial_t v(s))\|_{L^2} \le \|\partial_t u(s) - \partial_t v(s)\|_{L^2} \left(\|\partial_t u(s)\|_{\infty}^{p-1} + \|\partial_t v(s)\|_{\infty}^{p-1}\right).$$

Note that

$$\|\partial_t u(s) - \partial_t v(s)\|_{L^2} \le (1+s)^{-\frac{\alpha}{2}} \|u-v\|_{X(T)}.$$

On the other hand, using the Sobolev embedding  $H^1(\mathbb{R}) \hookrightarrow L^{\infty}(\mathbb{R})$  we get

$$\|\partial_t u(s)\|_{\infty}^{p-1} \le C \|\partial_t u(s)\|_{H^1}^{p-1} \le C(1+s)^{-\frac{\alpha(p-1)}{2}} \|u\|_{X(T)}^{p-1} \le CM^{p-1}(1+s)^{-\frac{\alpha(p-1)}{2}},$$

and

$$\|\partial_t v(s)\|_{\infty}^{p-1} \le C \|\partial_t v(s)\|_{H^1}^{p-1} \le C(1+s)^{-\frac{\alpha(p-1)}{2}} \|v\|_{X(T)}^{p-1} \le CM^{p-1}(1+s)^{-\frac{\alpha(p-1)}{2}}.$$

Accordingly,

$$\|f(\partial_t u(s)) - f(\partial_t v(s))\|_{L^2} \le CM^{p-1}(1+s)^{-\frac{p\alpha}{2}} \|u - v\|_{X(T)},$$
(3.25) {Fn1}

and therefore

$$\begin{aligned} (1+t)^{-1} \|\Phi(u)(t) - \Phi(v)(t)\|_{L^2} + \|\partial_t \Phi(u)(t) - \partial_t \Phi(v)(t)\|_{L^2} + \|\partial_x \Phi(u)(t) - \partial_x \Phi(v)(t)\|_{L^2} \\ &\leq C M^{p-1} \|u - v\|_{X(T)} (1+t)^{-\frac{\alpha}{2}} \int_0^t (1+s)^{-\frac{(p-1)\alpha}{2}} ds \end{aligned}$$

Referring to (3.14), we deduce that

$$(1+t)^{-1} \|\Phi(u)(t) - \Phi(v)(t)\|_{L^2} + \|\partial_t \Phi(u)(t) - \partial_t \Phi(v)(t)\|_{L^2} + \|\partial_x \Phi(u)(t) - \partial_x \Phi(v)(t)\|_{L^2} \\ \leq C M^{p-1} \|u - v\|_{X(T)} (1+t)^{-\frac{\alpha}{2}}.$$

Estimation of  $\|\partial_t^2 (\Phi(u) - \Phi(v))(t)\|_{L^2} + \|\partial_{xt}^2 (\Phi(u) - \Phi(v))(t)\|_{L^2} + \|\partial_x^2 (\Phi(u) - \Phi(v))(t)\|_{L^2}$ . By applying (2.33), we obtain

$$\begin{aligned} \|\partial_t^2 \left( \Phi(u) - \Phi(v) \right)(t) \|_{L^2} + \|\partial_{xt}^2 \left( \Phi(u) - \Phi(v) \right)(t) \|_{L^2} + \|\partial_x^2 \left( \Phi(u) - \Phi(v) \right)(t) \|_{L^2} \\ &\leq C \int_0^t \left( \frac{1+s}{1+t} \right)^{\frac{\alpha}{2}} \left( \|f(\partial_t u) - f(\partial_t v)\|_{H^1} + \|\partial_t \left( f(\partial_t u) - f(\partial_t v) \right)\|_{L^2} \right) \, ds. \end{aligned}$$
(3.26) {Hn1}

Since  $p > 1 + \frac{2}{\alpha_0} \ge 3$ , it follows that (3.5)-(3.6) can be applied. Therefore

$$\begin{aligned} \|\partial_x \left( f(\partial_t u(s)) - f(\partial_t v(s)) \right)\|_{L^2} \\ \leqslant C \|\partial_{xt}^2 w(s)\|_{L^2} \|\partial_t u(s)\|_{L^{\infty}}^{p-1} + C \|\partial_t w(s)\|_{L^{\infty}} \left( \|\partial_t u(s)\|_{L^{\infty}}^{p-2} + \|\partial_t v(s)\|_{L^{\infty}}^{p-2} \right) \|\partial_{xt}^2 v(s)\|_{L^2}, \end{aligned}$$

where  $w(t,x) \doteq u(t,x) - v(t,x)$ . We observe from the Sobolev embedding  $H^1(\mathbb{R}) \hookrightarrow L^{\infty}(\mathbb{R})$  that

$$\begin{aligned} \|\partial_t u(s)\|_{L^{\infty}} &\leq C \|\partial_t u(s)\|_{H^1} \leq C(1+s)^{-\frac{\alpha}{2}} \|u\|_{X(T)} \leq CM(1+s)^{-\frac{\alpha}{2}}, \\ \|\partial_t v(s)\|_{L^{\infty}} &\leq C \|\partial_t v(s)\|_{H^1} \leq C(1+s)^{-\frac{\alpha}{2}} \|v\|_{X(T)} \leq CM(1+s)^{-\frac{\alpha}{2}}, \\ \|\partial_t w(s)\|_{L^{\infty}} &\leq C \|\partial_t w(s)\|_{H^1} \leq C(1+s)^{-\frac{\alpha}{2}} \|w\|_{X(T)} = C(1+s)^{-\frac{\alpha}{2}} \|u-v\|_{X(T)}, \\ \|\partial_{xt}^2 v(s)\|_{L^2} &\leq (1+s)^{-\frac{\alpha}{2}} \|v\|_{X(T)} \leq CM(1+s)^{-\frac{\alpha}{2}}, \\ \|\partial_{xt}^2 w(s)\|_{L^2} &\leq (1+s)^{-\frac{\alpha}{2}} \|w\|_{X(T)} = (1+s)^{-\frac{\alpha}{2}} \|u-v\|_{X(T)}, \end{aligned}$$

whence

$$\|\partial_x \left( f(\partial_t u(s)) - f(\partial_t v(s)) \right)\|_{L^2} \leqslant C M^{p-1} (1+s)^{-\frac{p\alpha}{2}} \|u-v\|_{X(T)}.$$
(3.27) (3.27)

Likewise, (3.7)-(3.8) enable us to conclude that

$$\begin{aligned} \|\partial_t \left( f(\partial_t u(s)) - f(\partial_t v(s)) \right)\|_{L^2} \\ \leqslant C \|\partial_t^2 w(s)\|_{L^2} \|\partial_t u(s)\|_{L^{\infty}}^{p-1} + C \|\partial_t w(s)\|_{L^{\infty}} \left( \|\partial_t u(s)\|_{L^{\infty}}^{p-2} + \|\partial_t v(s)\|_{L^{\infty}}^{p-2} \right) \|\partial_t^2 v(s)\|_{L^2}. \end{aligned}$$

By applying the estimates

$$\begin{aligned} \|\partial_t^2 v(s)\|_{L^2} &\leqslant (1+s)^{-\frac{\alpha}{2}} \|v\|_{X(T)} \le CM(1+s)^{-\frac{\alpha}{2}}, \\ \|\partial_t^2 w(s)\|_{L^2} &\leqslant (1+s)^{-\frac{\alpha}{2}} \|w\|_{X(T)} = (1+s)^{-\frac{\alpha}{2}} \|u-v\|_{X(T)}, \end{aligned}$$

we arrive at

$$\|\partial_t \left( f(\partial_t u(s)) - f(\partial_t v(s)) \right)\|_{L^2} \leqslant C M^{p-1} (1+s)^{-\frac{p\alpha}{2}} \|u-v\|_{X(T)}.$$
(3.28) (3.28)

By inserting (3.25), (3.27), and (3.28) into (3.26) and taking into account (3.14), we derive

$$\begin{aligned} \|\partial_t^2 \left( \Phi(u) - \Phi(v) \right)(t) \|_{L^2} + \|\partial_{xt}^2 \left( \Phi(u) - \Phi(v) \right)(t) \|_{L^2} + \|\partial_x^2 \left( \Phi(u) - \Phi(v) \right)(t) \|_{L^2} \\ &\leq C M^{p-1} \|u - v\|_{X(T)} (1+t)^{-\frac{\alpha}{2}}. \end{aligned}$$

By summing the preceding estimates, it follows that

$$|\Phi(u) - \Phi(v)||_{X(T)} \le C M^{p-1} ||u - v||_{X(T)}.$$

Choosing M > 0 so that  $C M^{p-1} \leq 1/2$ , we arrive at

$$\|\Phi(u) - \Phi(v)\|_{X(T)} \le \frac{1}{2} \|u - v\|_{X(T)}$$

Applying the Banach fixed-point theorem, our proof is complete.

### 4 Proof of Theorems 1.3 and 1.4

<br/>the proof of Theorem 1.3 follow a similar approach to that of Theorem 1.1, we will<br/>outline the key steps while highlighting any necessary modifications.

$$\|f(\partial_t u(s), \partial_x u(s))\|_{L^2} = \||\partial_x u(s)|^q\|_{L^2} \le \|\partial_x u(s)\|_{\infty}^{q-1} \|\partial_x u(s)\|_{L^2} \le C \|\partial_x u(s)\|_{H^1}^q \le CM^p (1+s)^{-\frac{q\alpha}{2}},$$
  
$$\|\partial_x f(\partial_t u(s), \partial_x u(s))\|_{L^2} = \|\partial_x |\partial_x u(s)|^q\|_{L^2} \le q \|\partial_x u\|_{\infty}^{q-1} \|\partial_x^2 u\|_{L^2} \le C \|\partial_x u\|_{H^1}^q \le CM^p (1+s)^{-\frac{q\alpha}{2}},$$

and

$$\|\partial_t f(\partial_t u(s), \partial_x u(s))\|_{L^2} = \|\partial_t |\partial_x u(s)|^q \|_{L^2} \le q \|\partial_x u\|_{\infty}^{q-1} \|\partial_{xt}^2 u\|_{L^2} \le C \|\partial_x u\|_{H^1}^{q-1} \|\partial_{xt}^2 u\|_{L^2} \le C M^p (1+s)^{-\frac{q\alpha}{2}}.$$

For the contraction mapping argument, we require the estimates

$$\begin{aligned} \|f(\partial_{t}u(s),\partial_{x}u(s)) - f(\partial_{t}v(s),\partial_{x}v(s))\|_{L^{2}} &= \||\partial_{x}u(s)|^{q} - |\partial_{x}v(s)|^{q}\|_{L^{2}} \\ &\leq \|\partial_{x}u(s) - \partial_{x}v(s)\|_{L^{2}} \left(\|\partial_{x}u(s)\|_{\infty}^{q-1} + \|\partial_{x}v(s)\|_{\infty}^{q-1}\right) \\ &\leq C\left(1+s\right)^{-\frac{\alpha}{2}}\|u-v\|_{X(T)} \left(\|\partial_{x}u(s)\|_{H^{1}}^{q-1} + \|\partial_{x}v(s)\|_{H^{1}}^{q-1}\right) \\ &\leq CM^{q-1}(1+s)^{-\frac{q\alpha}{2}}\|u-v\|_{X(T)},\end{aligned}$$

 $\{sec4\}$ 

$$\begin{aligned} &\|\partial_x \left( f(\partial_t u(s), \partial_x u(s)) - f(\partial_t v(s), \partial_x v(s)) \right) \|_{L^2} \\ &= \|\partial_x \left( |\partial_x u(s)|^q - |\partial_x v(s)|^q \right) \|_{L^2} \\ &\leq C \|\partial_x^2 w(s)\|_{L^2} \|\partial_x u(s)\|_{L^{\infty}}^{q-1} + C \|\partial_x w(s)\|_{L^{\infty}} \left( \|\partial_x u(s)\|_{L^{\infty}}^{q-2} + \|\partial_x v(s)\|_{L^{\infty}}^{q-2} \right) \|\partial_x^2 v(s)\|_{L^2} \\ &\leq C M^{q-1} (1+s)^{-\frac{q\alpha}{2}} \|u-v\|_{X(T)}, \end{aligned}$$

and

$$\begin{aligned} \|\partial_t \left( f(\partial_t u(s), \partial_x u(s)) - f(\partial_t v(s), \partial_x v(s)) \right) \|_{L^2} \\ &= \|\partial_t \left( |\partial_x u(s)|^q - |\partial_x v(s)|^q \right) \|_{L^2} \\ &\leq C \|\partial_{xt}^2 w(s)\|_{L^2} \|\partial_x u(s)\|_{L^{\infty}}^{q-1} + C \|\partial_t w(s)\|_{L^{\infty}} \left( \|\partial_x u(s)\|_{L^{\infty}}^{q-2} + \|\partial_x v(s)\|_{L^{\infty}}^{q-2} \right) \|\partial_{xt}^2 v(s)\|_{L^2} \\ &\leq C M^{q-1} (1+s)^{-\frac{q\alpha}{2}} \|u-v\|_{X(T)}, \end{aligned}$$

where  $w(t, x) \doteq u(t, x) - v(t, x)$ . These estimates ensure the existence and uniqueness of solutions via the Banach fixed point theorem.

**.**</sup>

$$\begin{aligned} \|f(\partial_{t}u(s),\partial_{x}u(s))\|_{L^{2}} &= \||\partial_{t}u(s)|^{p}|\partial_{x}u(s)|^{q}\|_{L^{2}} \\ &\leq \|\partial_{t}u(s)\|_{L^{\infty}}^{p}\||\partial_{x}u(s)|^{q}\|_{L^{2}} \\ &\leq C \|\partial_{t}u(s)\|_{H^{1}}^{p}\||\partial_{x}u(s)|^{q}\|_{L^{2}} \\ &\leq C M^{p}(1+s)^{-\frac{p\alpha}{2}}M^{q}(1+s)^{-\frac{q\alpha}{2}} \\ &= C M^{p+q}(1+s)^{-\frac{\alpha(p+q)}{2}}, \end{aligned}$$

which implies that

$$(1+t)^{-1} \|\Phi(u)^{nl}(t)\|_{L^2} + \|\partial_t \Phi(u)^{nl}(t)\|_{L^2} + \|\partial_x \Phi(u)^{nl}(t)\|_{L^2} \le C M^{p+q} (1+t)^{-\frac{\alpha}{2}} \int_0^t (1+s)^{-\frac{(p+q-1)\alpha}{2}} ds$$

Since  $p + q > 1 + \frac{2}{\alpha_0}$ , we conclude that

$$(1+t)^{-1} \|\Phi(u)^{nl}(t)\|_{L^2} + \|\partial_t \Phi(u)^{nl}(t)\|_{L^2} + \|\partial_x \Phi(u)^{nl}(t)\|_{L^2} \le C M^{p+q} (1+t)^{-\frac{\alpha}{2}}.$$

This ensures that  $\Phi$  maps into  $B_M(T)$ , and a similar argument establishes the contraction property, thereby concluding the proof.

### 5 Proof of Theorem 1.2

This section is dedicated to the proof of Theorem 1.2, which aims to determine the blow-up region and establish lifespan estimates for problem (1.1) in the one-dimensional space.

{sec5}

$$\partial_t^2 \psi(x,t) - \partial_x^2 \psi(x,t) - V(x) \partial_t \psi(x,t) = 0.$$
(5.1) {lambda-eq}

More precisely, we choose the function  $\psi$  given by

$$\psi(x,t) := \rho(t)\phi(x); \quad \rho(t) := e^{-t},$$
(5.2) {psi33}

where  $\phi(x) = \phi(-x)$  is a solution of the problem

$$\phi''(x) = (1 + V(x))\phi(x), \quad \text{for all } x \in \mathbb{R}, \tag{5.3} \quad \text{{\bf [phi]}}$$

$$\phi(r) = 1 + \int_0^r (r-s)(1+V(s))\phi(s) \, ds, \qquad r > 0.$$

Note that the existence of a positive solution for the associated problem of (5.3) is studied in [21, 26] in higher dimension. However, the specific case of one spatial dimension has not been addressed in these works. However, following the strategy used in [21] in the case of higher dimension, we easily prove the following properties of  $\phi$ .

Lemma 5.1. There exists an even function \$\phi\$ solution of (5.3) verifying \$\phi(0) = 1\$, \$\phi'(0) = 0\$, which satisfies

$$0 < \phi(x) \le C(1+|x|)^{\frac{\mu_0}{2}} e^{|x|}, \quad \text{for all } x \in \mathbb{R}.$$
(5.4) [elip1]

 $\{1em0\}$ 

$$q''(x) + 2q'(x) + (q'(x))^2 = V(x), \quad \text{for all } x \in \mathbb{R}, \quad (5.5) \quad \{\texttt{functionq}\}$$

$$e^{2x}q'(x) \le -1 + \int_0^x e^{2y}V(y)dy$$
, for all  $x > 0$ . (5.6) {functionq1

By using the expression of V, and integrating by parts we infer

$$\int_0^x e^{2y} V(y) dy = \frac{\mu_0}{2\sqrt{1+x^2}} e^{2x} - \frac{1}{2} \int_0^x e^{2y} V'(y) dy, \quad \text{for all } x > 0, \quad (5.7) \quad \{\text{function}(x) < 0\}$$

which is asymptotically

$$\frac{\mu_0}{2}x^{-1}e^{2x} + \mathcal{O}(x^{-2})e^{2x},$$

< i since V(x) ~ µ\_0x^{-1} and V'(x) ~ ~ µ\_0x^{-2} for large x. Hence, by (5.6) and (5.7), we write for x large enough,

$$q'(x) \le \frac{\mu_0}{2x} + \mathcal{O}(x^{-2}).$$
 (5.8) {functionq3}

Therefore, for x large enough,

$$q(x) \le \frac{\mu_0}{2} \ln(x) + \mathcal{O}(1). \tag{5.9} \quad \text{{functiong}}$$

Inequality (5.4) is a direct consequence of (5.9) owing to the positivity and the symmetry of

Furthermore, by employing the estimate (5.4) along with the expression of  $\psi$  given by (5.2), and proceeding along the same lines as in the derivation of inequality (2.5) in [44] or [9, Lemma 3.1], we obtain the following result.

**Lemma 5.2.** There exists a constant  $C = C(R_0, \mu_0) > 0$  such that

$$\int_{|x| \le R_0 + t} \psi(x, t) dx \le C(1 + t)^{\frac{\mu_0}{2}}, \quad \text{for all } t \ge 0.$$
(5.10) {psi}

**Proof of Theorem 1.2.** The proof relies on Lemma 5.2 and follows a similar strategy to the one used in [9, Theorem 2.3]. Specifically, we employ the test function method, originally introduced by Baras and Pierre [1], later refined by Zhang [45], and further developed in the works of Mitidieri and Pohozaev [30], which has since been widely applied in various contexts, including the works of Kirane et al. [25] and Fino et al. [4, 5, 6, 8].

### References

- P. Baras & M. Pierre, Critère d'existence de solutions positives pour des équations semilinéaires non monotones, Ann. Inst. H. Poincaré Anal. Non Linéaire 2 (1985), 185–212.
- [2] H. Bellout, A. Friedman, Blow-up estimates for a nonlinear hyperbolic heat equation, Math. Z. 145 (1975), 251 – 254.
- [3] M.F. Ben Hassen, M. Hamouda, M.A. Hamza and H.K. Teka, Nonexistence result for the generalized Tricomi equation with the scale-invariant damping, mass term and time derivative nonlinearity. Asymptotic Analysis, 128 (2022) 495–515, DOI 10.3233/ASY-211714.
- [4] W. Chen, A. Z. Fino, Blow-up of solutions to semilinear strongly damped wave equations with different nonlinear terms in an exterior domain. Math. Meth. Appl. Sci. 44(8), 6787– 6807 (2021)
- [5] T. A. Dao, A.Z. Fino, Critical exponent for semi-linear structurally damped wave equation of derivative type. Math. Meth. Appl. Sci. 43, 9766–9775 (2020)
- [6] A. Z. Fino, Finite time blow up for wave equations with strong damping in an exterior domain. Mediterranean Journal of Mathematics 17(6), Paper No. 174, 21 pp. (2020)

 $\{lem1\}$ 

4`

- [7] A. Z. Fino, M. Kirane, Qualitative properties of solutions to a nonlocal evolution system. Math. Meth. Appl. Sci. 34(9), 1125–1143 (2011)
- [8] A. Z. Fino & M. A. Hamza, Blow-up of solutions to semilinear wave equations with a time-dependent strong damping. Evolution Equations and Control Theory, 2022, 11(6): 1955-1966. doi: 10.3934/eect.2022006
- [9] A. Z. Fino & M. A. Hamza, Blow-up and lifespan estimate for wave equations with critical damping term of space-dependent type related to Glassey conjecture. Discrete and Continuous Dynamical Systems - S, 16 (6), 2023, 1383–1400. doi: 10.3934/dcdss.2022192.
- [10] A. Z. Fino & M. A. Hamza, Global existence for wave equations with scale-invariant timedependent damping and time derivative nonlinearity. arXiv:2409.13353v1
- [11] M. Hamouda and M.A. Hamza, Blow-up for wave equation with the scale-invariant damping and combined nonlinearities. Math Meth. Appl. Sci. Volume 44, Issue 1, 2021, Pages 1127–1136.
- [12] M. Hamouda and M.A. Hamza, Improvement on the blow-up of the wave equation with the scale-invariant damping and combined nonlinearities. Nonlinear Anal. Real World Appl. Volume 59, 2021, 103275, ISSN 1468-1218, 2021. Available from: https://doi.org/10.1016/j.nonrwa.2020.103275.
- [13] M. Hamouda and M.A. Hamza, Improvement on the blow-up for a weakly coupled wave equations with scale-invariant damping and mass and time derivative nonlinearity. Discrete and Continuous Dynamical Systems - S. 2023, Vol. 16, Issue 6, 1468–1483. Doi: 10.3934/dcdss.2022173.
- [14] M. Hamouda and M.A. Hamza, A blow-up result for the wave equation with localized initial data: the scale-invariant damping and mass term with combined nonlinearities. Journal of Applied Analysis and Computation, 2022, 12(5): 1816-1841. doi: 10.11948/20210361.
- [15] M. Hamouda and M.A. Hamza, Blow-up and lifespan estimate for the generalized Tricomi equation with mixed nonlinearities. Adv. Pure Appl. Math.12 (2021), Special issue, 54–70.
- [16] M. Hamouda, M.A. Hamza and A. Palmieri, A note on the nonexistence of global solutions to the semilinear wave equation with nonlinearity of derivative-type in the generalized Einstein-de Sitter spacetime. Communications on Pure & Applied Analysis20 (2021), no. 11, 3703–3721.Doi: 10.3934/cpaa.2021127.
- [17] M. Hamouda, M.A. Hamza and A. Palmieri, Blow-up and lifespan estimates for a damped wave equation in the Einstein-de Sitter spacetime with nonlinearity of derivative type, NoDEA Nonlinear Differential Equations Appl. 29 (2022), no. 2, Paper No. 19, 15 pp.
- [18] H. Fujita, On the blowing up of solutions of the problem for  $u_t = \Delta u + u^{1+\alpha}$ , J. Fac. Sci. Univ. Tokyo **13** (1966), 109–124.

- [19] M. Ikawa, Hyperbolic partial differential equations and wave phenomena, American Mathematical Society, 2000.
- [20] M. Ikeda & M. Sobajima, Life-span of Blowup Solutions to Semilinear Wave Equation with Space-dependent Critical Damping, Funkcialaj Ekvacioj 64, (2021), 137–162.
- [21] R. Ikehata, G. Todorova & B. Yordanov, Optimal decay rate of the energy for wave equations with critical potential, J. Math. Soc. Japan 65 no. 1 (2013), 183–236.
- [22] R. Ikehata, K. Tanizawa, Global existence for solutions for semilinear damped wave equation in ℝ<sup>N</sup> with noncompactly supported initial data, Nonlinear Analysis 61 (2005), 1189– 1208.
- [23] R. Ikehata, G. Todorova, B.Yordanov.: Critical exponent for semilinear wave equations with space-dependent potential. Funkcialaj Ekvacioj (2009), 52, 411–435.
- [24] F. John, Blow-up for quasilinear wave equations in three space dimensions. Comm. Pure Appl. Math., 34 (1981), 29–51.
- [25] M. Kirane, Y. Laskri, N.E. Tatar, Critical exponents of Fujita type for certain evolution equations and systems with Spatio-Temporal Fractional derivatives. J. Math. Anal. Appl. 312, (2005), 488–501.
- [27] N.A. Lai and Z. Tu, Strauss exponent for semilinear wave equations with scattering space dependent damping. J. Math. Anal. Appl. (2020), 489, 124–189.
- [28] X. Li, Critical exponent for semilinear wave equation with critical potential. Nonlinear Differ. Equ. Appl. (2013), 20, 1379–1391.
- [29] T.-T. Li,Y. Zhou, Breakdown of solutions to  $\Box u + u_t = |u|^{1+\alpha}$ , Discrete Contin. Dyn. Syst. 1 (1995), 503–520.
- [30] E. Mitidieri, S. I. Pokhozhaev, Apriori estimates and the absence of solutions of nonlinear partial differential equations and inequalities. Tr. Mat. Inst. Steklova 234, (2001), 1–384.
- [31] K. Nishihara, M. Sobajima, Y. Wakasugi, Critical exponent for the semilinear wave equations with a damping increasing in the far field. Nonlinear Differ. Equ. Appl. 25, 55 (2018). https://doi.org/10.1007/s00030-018-0546-2.
- [32] N.A. Lai and H. Takamura, Nonexistence of global solutions of nonlinear wave equations with weak time-dependent damping related to Glassey's conjecture, Differential Integral Equations, **32**, 2019, no. 1-2, 37–48.

- [34] A. Palmieri and Z. Tu, A blow-up result for a semilinear wave equation with scaleinvariant damping and mass and nonlinearity of derivative type, Calc. Var. 60, 72, 2021. Available from: https://doi.org/10.1007/s00526-021-01948-0.
- [35] M.A. Rammaha, Finite-time blow-up for nonlinear wave equations in high dimensions. Comm. Partial Differential Equations, 12 (1987), (6), 677–700.
- [36] T.C. Sideris, Global behavior of solutions to nonlinear wave equations in three space dimensions. Comm. Partial Differential Equations, 8 (1983), no. 12, 1291–1323.
- [37] M. Sobajima, On global existence for semilinear wave equations with spacedependent critical damping. J. Math. Soc. Japan 75(2), (2023), 603–627. https://doi.org/10.2969/jmsj/87388738
- [38] M. Sobajima, Y. Wakasugi, Diffusion phenomena for the wave equation with spacedependent damping in an exterior domain. J. Differ. Equ. 261, (2016), 5690–5718.
- [39] M. Sobajima, Y. Wakasugi, Diffusion phenomena for the wave equation with spacedependent damping term growing at infinity. Adv. Differ. Equ. 23, (2018), 581–614.
- [40] M. Sobajima, Y. Wakasugi, Weighted energy estimates for wave equation with spacedependent damping term for slowly decaying initial data, Communications in Contemporary Mathematics 21(5), (2019), 1850035.
- [41] G. Todorova, B. Yordanov, Critical exponent for a nonlinear wave equation with damping, J. Differential Equations 174 (2001), 464 - 489.
- [42] Y. Wakasugi, Small data global existence for the semilinear wave equation with space–time dependent damping, J. Math. Anal. Appl, 2012, 393, 66–79.
- men sig i no sig in sig i no si
- n [45] Q.S. Zhang, A blow-up result for a nonlinear wave equation with damping: the critical case. C R Acad Sci Paris Sér I Math., **333** (2) (2001), 109–114.
- [46] Y. Zhou, Blow-up of solutions to the Cauchy problem for nonlinear wave equations. Chin. Ann. Math., 22B (3) (2001), 275–280.