

# Exact Schwinger functions for a class of bounded interactions in $d \geq 2$

Wojciech Dybalski

Faculty of Mathematics and Computer Science,  
Adam Mickiewicz University in Poznań,  
ul. Uniwersytetu Poznańskiego 4, 61-614 Poznań, Poland  
E-mail: wojciech.dybalski@amu.edu.pl

## Abstract

We consider a scalar Euclidean QFT with interaction given by a bounded, measurable function  $V$  such that  $V^\pm := \lim_{w \rightarrow \pm\infty} V(w)$  exist. We find a field renormalization such that all the  $n$ -point connected Schwinger functions for  $n \neq 2$  exist non-perturbatively in the UV limit. They coincide with the tree-level one-particle irreducible Schwinger functions of the  $\text{erf}(\phi/\sqrt{2})$  interaction with a coupling constant  $\frac{1}{2}(V^+ - V^-)$ . By a slight modification of our construction we can change this coupling constant to  $\frac{1}{2}(V_+ - V_-)$ , where  $V_\pm := \lim_{w \rightarrow 0^\pm} V(w)$ . Thereby non-Gaussianity of these theories is governed by a discontinuity of  $V$  at zero. The open problem of controlling also the two-point function of these QFTs is discussed.

## 1 Introduction

Rigorous construction of interacting scalar quantum field theories in spacetime dimension  $d = 2, 3$  is among the most important achievements of the constructive programme [GJ]. These efforts focussed on polynomial interactions and were limited by the requirement of super-renormalizability: For  $d = 2$  all (bounded from below) polynomials are covered, for  $d = 3$  only polynomials up to the fourth order, whereas for  $d \geq 4$  all the polynomial theories are expected to be trivial [Ai82, AD21, Fr82]. These limitations suggest some room for non-polynomial scalar theories in  $d = 2$ . In fact, theories based on exponential interaction functions, such as the Sine-Gordon and Sinh-Gordon model, are among the classical topics of constructive QFT [FP77] and still constitute a very active field of research [Le08, CT16, HS16, BFR21, BD21, BH22, Ko23, BDR24, CFM24, GM24]. On the other hand, for  $d > 2$ , there seems to be no room for non-polynomial interactions, at least if we think about them in terms of their Taylor expansions. But this leaves aside bounded interaction functions like  $\arctan(\phi)$ , whose Taylor expansion has finite radius of convergence and thus does not capture the large-field behaviour. Not to speak of functions like  $\text{sgn}(\phi)$  which lack a Taylor expansion around zero altogether. Clearly, polynomial interactions are not a reliable guide to study such bounded interactions and we are not aware of any systematic treatment. In this short note we demonstrate that these models are exactly solvable for any  $d \geq 2$  in a sense which we now explain.

We consider interaction functions  $V : \mathbb{R} \rightarrow \mathbb{R}$  which are bounded and measurable. Depending on a situation we will need the existence of the following limits:

$$V_\pm := \lim_{w \rightarrow 0^\pm} V(w), \tag{A_1}$$

$$V^\pm := \lim_{w \rightarrow \pm\infty} V(w). \tag{A_2}$$

We will refer to these requirements as assumptions (A<sub>1</sub>) and (A<sub>2</sub>), respectively. The corresponding interaction term in the finite volume  $B := [-\frac{L}{2}, \frac{L}{2}]^d$ ,  $L > 0$ , is  $\mathcal{V}(\phi) := \int_B dx V(\phi(x))$ . We do not impose any Wick ordering on the interaction.

Next, we introduce the free covariance with mass  $m > 0$  and a UV cut-off  $\Lambda$  s.t.  $\log(\Lambda) \geq 1$ :

$$C_\Lambda(x) := \int_{\mathbb{R}^d} \frac{dp}{(2\pi)^d} e^{ipx} \frac{e^{-(p^2+m^2)/\Lambda^2}}{p^2 + m^2}, \quad C(x) := \lim_{\Lambda \rightarrow \infty} C_\Lambda(x). \quad (1.1)$$

We allow for a field renormalization  $Z_\Lambda > 0$  and denote the renormalized covariance by  $\tilde{C}_\Lambda := Z_\Lambda C_\Lambda$ . Let  $\nu_{\tilde{C}_\Lambda}$  be the corresponding Gaussian measure with mean zero and denote the expectation w.r.t. this measure by  $\langle \dots \rangle_{\tilde{C}_\Lambda} = \int \dots d\nu_{\tilde{C}_\Lambda}$ . In particular,

$$S_{0,\Lambda}(J) := \langle e^{\phi(J)} \rangle_{\tilde{C}_\Lambda} = e^{\frac{1}{2}\langle J, \tilde{C}_\Lambda J \rangle}, \quad J \in S(\mathbb{R}^d; \mathbb{R}) \quad (1.2)$$

is the generating functional of the Schwinger functions of the free theory, where  $\langle \cdot, \cdot \rangle$  on the r.h.s. is the scalar product in  $L^2(\mathbb{R}^d; \mathbb{R})$ . The generating functional of the interacting theory has the form

$$S_\Lambda(J) := \langle e^{\phi(J)} e^{-\lambda V(\phi)} \rangle_{\tilde{C}_\Lambda}, \quad \lambda \in \mathbb{R}, \quad (1.3)$$

disregarding a  $J$ -independent normalization constant, which has no effect on connected Schwinger functions. We refer to Appendix A for routine arguments on the well-definiteness of this functional. A standard tool to study the effect of the interaction is the following modified generating functional, which we state together with its connected counterpart, cf. e.g. [Sa, Eq. (2.104)],

$$\Sigma_\Lambda(J) := \frac{S_\Lambda(J)}{S_{0,\Lambda}(J)}, \quad \Sigma_\Lambda^c(J) := \log \Sigma_\Lambda(J). \quad (1.4)$$

It is our main result, stated in Theorem 1.1, that  $\Sigma_\Lambda^c(J)$  has a finite and exactly computable UV limit. The proof is given in Section 2.

**Theorem 1.1.** *Set  $Z_\Lambda = C_\Lambda(0)^\eta$ ,  $\eta \in \mathbb{R}$ . Then the limit  $\Sigma^c(J) := \lim_{\Lambda \rightarrow \infty} \Sigma_\Lambda^c(J)$  exists for any  $J \in S(\mathbb{R}^d; \mathbb{R})$  under the assumptions specified below. It has the form:*

(a) *Let  $-\infty < \eta < -1$ . Then, assuming (A<sub>1</sub>),*

$$\Sigma^c(J) = -\lambda \frac{V_+ + V_-}{2} |B|. \quad (1.5)$$

(b) *Let  $\eta = -1$ . Then, without additional assumptions,*

$$\Sigma^c(J) = -\lambda |B| \frac{1}{(2\pi)^{1/2}} \int_{\mathbb{R}} dw V(w) e^{-\frac{1}{2}w^2}. \quad (1.6)$$

(c) *Let  $-1 < \eta < 1$ . Then, assuming (A<sub>2</sub>),*

$$\Sigma^c(J) = -\lambda \frac{V^+ + V^-}{2} |B|. \quad (1.7)$$

(d) *Let  $\eta = 1$ . Then, assuming (A<sub>2</sub>),*

$$\Sigma^c(J) = -\lambda \frac{V^+ + V^-}{2} |B| - \lambda \frac{V^+ - V^-}{2} \frac{1}{(2\pi)^{1/2}} \int_B dx \int dw \operatorname{sgn}(w) e^{-\frac{1}{2}(w - \langle \delta_x, CJ \rangle)^2}. \quad (1.8)$$

(e) *Let  $1 < \eta < \infty$ . Then, assuming (A<sub>2</sub>), and choosing  $J$  compactly supported in Fourier space,*

$$\Sigma^c(J) = -\lambda \frac{V^+ + V^-}{2} |B| - \lambda \frac{V^+ - V^-}{2} \mathbb{1}(J \neq 0) \int_B dx \operatorname{sgn}(\langle \delta_x, CJ \rangle). \quad (1.9)$$

We note that  $\Sigma_\Lambda^c(J)$  is the generating functional of connected  $n$ -point Schwinger functions  $S_{\Lambda,n}^c$  of the theory with cut-offs for  $n \neq 2$ , cf. [PT11, Chapter 3]. Therefore, we interpret  $\Sigma^c(J)$  as the generating functional of the connected Schwinger functions  $S_{n \neq 2}^c$  of the limiting theory. Part (d) of Theorem 1.1 is the most interesting one as the connected Schwinger functions  $S_{n \neq 2}^c$  of the limiting theory are non-trivial and readily computable, cf. Appendix B,

$$\begin{aligned} S_n^c(x_1, \dots, x_n) &:= \frac{\delta}{\delta J(x_1)} \cdots \frac{\delta}{\delta J(x_n)} \Sigma^c(J)|_{J=0} \\ &= -\lambda \frac{(V^+ - V^-)}{2} [(\partial_w)^n \operatorname{erf}(w/\sqrt{2})]_{w=0} \int_B dx C(x - x_1) \cdots C(x - x_n). \end{aligned} \quad (1.10)$$

We observe that these  $n$ -point functions coincide with the tree-level one-particle irreducible Schwinger functions of a perturbative theory with the interaction function  $\operatorname{erf}(\phi/\sqrt{2})$  and coupling constant  $\lambda \frac{(V^+ - V^-)}{2}$ . We find it quite surprising that for all the bounded interaction functions  $V$  as defined above we obtain the same Schwinger functions for  $n \neq 2$  up to a finite coupling constant renormalization. We also note that the infinite volume limit  $B \nearrow \mathbb{R}^d$  can easily be taken in (1.10) after smearing with test functions.

The remaining parts of Theorem 1.1 also carry physically relevant information, although the Schwinger functions  $S_{n \neq 2}^c$  are zero or undefined. From the point of view of statistical physics, the quantity  $\Sigma^c(J)$  is the change of the free energy due to interaction, which is an interesting quantity. Depending on the parameter  $\eta$ , it probes various properties of the interaction function  $V$ , such as the behaviour at zero or infinity. For example, in the cases (c)-(e), we have

$$\Sigma^c(0) = -\lambda \frac{(V^+ + V^-)}{2} |B|. \quad (1.11)$$

We set  $J = 0$  here to make this quantity insensitive to the denominator  $S_{0,\Lambda}(J = 0) = 1$  in (1.4) and thus consistent with the standard definition of the free energy in constructive QFT. We recall that upper and lower bounds on free energy, called UV stability, are difficult to prove for polynomial interactions, see e.g. [Di13]. For bounded interactions treated in this note the free energy is readily computable.

Our paper is organized as follows: Section 2 starts with a non-technical explanation of a factorization mechanism which is behind Theorem 1.1 and then moves on to the proof of our main result. In Section 3 we generalize our construction to a certain class of  $\Lambda$ -dependent interaction functions. In Section 4 we discuss the problem of controlling also the two-point Schwinger function for our class of bounded interactions.

**Acknowledgements:** I would like to thank Paweł Duch and Yoh Tanimoto for useful comments. Financial support of the National Science Centre, Poland, via the grant ‘Sonata Bis’ 2019/34/E/ST1/00053, is gratefully acknowledged.

## 2 Analysis

All constants  $c, c_1, c_2, \dots$  in the following depend only on  $m$  and  $d$  and may change from line to line. Our proof of Theorem 1.1 is based on the following proposition:

**Proposition 2.1.** *The following bound holds true for  $\ell \in \mathbb{N}$*

$$\left| \frac{\langle e^{\phi(J)} \mathcal{V}(\phi)^\ell \rangle_{\tilde{C}_\Lambda}}{\langle e^{\phi(J)} \rangle_{\tilde{C}_\Lambda}} - \left( \frac{\langle e^{\phi(J)} \mathcal{V}(\phi) \rangle_{\tilde{C}_\Lambda}}{\langle e^{\phi(J)} \rangle_{\tilde{C}_\Lambda}} \right)^\ell \right| \leq c \ell^6 \|V\|_\infty^\ell |B|^\ell \frac{1}{\sqrt{\log(\Lambda)}}. \quad (2.1)$$

Let us first explain in a simple example the mechanism, which is behind the factorization property (2.1). We choose  $V(w) = \operatorname{sgn}(w)$ , which is invariant under scaling  $w \mapsto \alpha w$ ,  $\alpha > 0$ . Hence the Fourier transform satisfies

$\hat{V}(\alpha u) = \alpha^{-1} \hat{V}(u)$ . Using this property, representation  $V(\phi(x)) = \frac{1}{\sqrt{2\pi}} \int du \hat{V}(u) e^{i\phi(x)u}$  and relation (1.2), we obtain

$$\frac{\langle e^{\phi(J)} \mathcal{V}(\phi)^2 \rangle_{\tilde{C}_\Lambda}}{\langle e^{\phi(J)} \rangle_{\tilde{C}_\Lambda}} = \frac{1}{2\pi} \int_{B^2} dx_1 dx_2 \int_{\mathbb{R}^2} du_1 du_2 \hat{V}(u_1) \hat{V}(u_2) e^{-\frac{1}{2} \sum_{i=1}^2 u_i^2 - u_1 u_2 \frac{C_\Lambda(x_1-x_2)}{C_\Lambda(0)} + i \frac{(\sum_{j=1}^2 u_j \delta_{x_j}, \tilde{C}_\Lambda J)}{\tilde{C}_\Lambda(0)^{1/2}}}. \quad (2.2)$$

Clearly, the only obstacle to factorization of the  $u$ -integral is the term involving  $\frac{C_\Lambda(x_1-x_2)}{C_\Lambda(0)}$ . But for  $x_1 \neq x_2$  this expression tends to zero with  $\Lambda \rightarrow \infty$  by Lemma 2.2 below. On the other hand, the region close to the diagonal  $x_1 = x_2$  gives a small contribution for bounded interactions, cf. estimates (2.19), (2.20) below. We stress that this factorization mechanism is independent of the field renormalization and relevant to all parts of Theorem 1.1. It is also distinct from the classical limit, which effects a similar factorization in perturbation theory, cf. (4.2) below. The field renormalization is needed to ensure the non-triviality of (2.2), that is, to keep the  $J$ -dependent term in the exponent in (2.2) non-zero and finite. Since  $\frac{\tilde{C}_\Lambda}{\tilde{C}_\Lambda(0)^{1/2}} = Z_\Lambda^{1/2} \frac{C_\Lambda}{C_\Lambda(0)^{1/2}}$ , this can be achieved e.g. by setting  $Z_\Lambda = C_\Lambda(0)$ .

Moving on to the proof of Proposition 2.1, we list the relevant properties of the covariance  $C$ . The non-optimal bounds from Lemma 2.2 below are convenient for us, as they facilitate uniform treatment of all dimensions  $d \geq 2$ . We postpone the proof to Appendix C.

**Lemma 2.2.** *The propagator  $C_\Lambda$  has the following properties for  $d \geq 2$ :*

$$0 < C_\Lambda(x) < C_\Lambda(0) \quad \text{for } x \neq 0, \quad (2.3)$$

$$C_\Lambda(x) \leq \frac{c}{|x|^{d-3/2}}, \quad (2.4)$$

$$C_\Lambda(0) \geq c_1 \log(\Lambda), \quad \text{for } c_1 > 0. \quad (2.5)$$

Next, we introduce the following  $\ell \times \ell$  matrices for  $\mathbf{x} := (x_1, \dots, x_\ell) \in \mathbb{R}^{\ell d}$ ,  $0 \leq \alpha \leq 1$ ,

$$\mathbf{M}_\alpha(\mathbf{x}) := I + \alpha \mathbf{m}(\mathbf{x}), \quad (2.6)$$

where  $\mathbf{m}_{i,j}(\mathbf{x}) := \frac{C_\Lambda(x_i-x_j)}{C_\Lambda(0)}$  for  $i \neq j$  and  $\mathbf{m}_{i,i}(\mathbf{x}) = 0$ . We set  $\mathbf{M}(\mathbf{x}) := \mathbf{M}_{\alpha=1}(\mathbf{x})$  and for  $\ell = 1$  it is understood that  $\mathbf{M}_\alpha(\mathbf{x}) = 1$ . Also, we will often abbreviate  $\mathbf{M}_\alpha := \mathbf{M}_\alpha(\mathbf{x})$ ,  $\mathbf{m} := \mathbf{m}(\mathbf{x})$ . Next, we define the following neighbourhood of the diagonal in  $B^\ell$

$$D_\delta := \{ \mathbf{x} \in B^\ell \mid \exists_{i \neq j} \text{ s.t. } |x_i - x_j| \leq \delta \} \quad (2.7)$$

and denote its complement in  $B^\ell$  by  $D'_\delta$ . (For  $\ell = 1$  we set  $D'_\delta := B$ ). We choose  $\delta$ , depending on  $\Lambda$ ,  $\ell$ ,  $d$ , as follows

$$\delta := \left( \frac{\ell}{\sqrt{\log(\Lambda)}} \right)^{\frac{1}{d-3/2}}. \quad (2.8)$$

**Lemma 2.3.** *For  $\mathbf{x} \in D'_\delta$ , and  $\Lambda \geq \Lambda_0$  (for some  $\Lambda_0$  depending only on  $d, m$ ) the matrices  $\mathbf{M}_\alpha(\mathbf{x})$  are positive definite. Furthermore,*

$$\|\mathbf{M}_\alpha(\mathbf{x})^{-1}\| \leq c, \quad \|\mathbf{M}_\alpha(\mathbf{x})^{-1} \partial_\alpha \mathbf{M}_\alpha(\mathbf{x})\| \leq \frac{c}{\sqrt{\log(\Lambda)}}, \quad (2.9)$$

for  $c$  possibly depending on  $d, m$  but independent of  $\Lambda, \mathbf{x}$ .

**Proof.** For  $\mathbf{x} \in D'_\delta$  we have  $|x_i - x_j| > \delta$  for all  $1 \leq i < j \leq \ell$ . By Lemma 2.2,

$$\mathbf{m}_{i,j}(\mathbf{x}) = \frac{C_\Lambda(x_i - x_j)}{C_\Lambda(0)} < c \delta^{-(d-3/2)} \frac{1}{\log(\Lambda)} = \frac{c}{\ell} \frac{1}{\sqrt{\log(\Lambda)}}, \quad (2.10)$$

where we used definition (2.8). Thus, recalling that  $\mathbf{m}_{i,i} = 0$  and using Schur's test

$$\|\mathbf{m}\| \leq \sup_i \sum_j |\mathbf{m}_{i,j}| \leq c \frac{1}{\sqrt{\log(\Lambda)}} \leq \frac{1}{2} \quad (2.11)$$

for  $\Lambda \geq \Lambda_0$ , where  $\Lambda_0$  depends only on  $d, m$ . Consequently,  $\mathbf{M}_\alpha = I + \alpha \mathbf{m}$ ,  $0 \leq \alpha \leq 1$ , is positive definite. Furthermore,

$$\|\mathbf{M}_\alpha^{-1}\| \leq \sum_{\ell'=0}^{\infty} \|\mathbf{m}\|^{\ell'}, \quad (2.12)$$

$$\|\mathbf{M}_\alpha^{-1} \partial_\alpha \mathbf{M}\| \leq \|\mathbf{M}_\alpha^{-1}\| \|\mathbf{m}\|. \quad (2.13)$$

Substituting (2.11) to (2.12), (2.13) we complete the proof.  $\square$

**Lemma 2.4.** For  $\mathbf{x} \in D'_\delta$  and  $\Lambda \geq \Lambda_0$  as in Lemma 2.3, the following equality holds true

$$\begin{aligned} & \frac{\langle e^{\phi(J)} V(\phi(x_1)) \dots V(\phi(x_\ell)) \rangle_{\tilde{C}_\Lambda}}{\langle e^{\phi(J)} \rangle_{\tilde{C}_\Lambda}} \\ &= \frac{1}{(2\pi)^{\ell/2}} \det(\mathbf{M}(\mathbf{x}))^{-1/2} \int_{\mathbb{R}^\ell} d\mathbf{w} V(\tilde{C}_\Lambda(0)^{1/2} \mathbf{w}) e^{-\frac{1}{2}(\mathbf{w}-\mathbf{q})^T \mathbf{M}(\mathbf{x})^{-1}(\mathbf{w}-\mathbf{q})}, \end{aligned} \quad (2.14)$$

where we set  $V(\mathbf{w}) := V(w_1) \dots V(w_\ell)$  and defined the vector

$$\mathbf{q} := \left( \frac{\langle \delta_{x_1}, \tilde{C}_\Lambda J \rangle}{\tilde{C}_\Lambda(0)^{1/2}}, \dots, \frac{\langle \delta_{x_\ell}, \tilde{C}_\Lambda J \rangle}{\tilde{C}_\Lambda(0)^{1/2}} \right). \quad (2.15)$$

**Proof.** Let us choose  $V_\varepsilon \in S(\mathbb{R})$  s.t.  $V_\varepsilon \nearrow V$  pointwise. Then, using that (1.2) holds for complex valued  $J$  if we treat  $\langle \cdot, \cdot \rangle$  as complex-linear in both arguments,

$$\begin{aligned} & e^{-\frac{1}{2}\langle J, \tilde{C}_\Lambda J \rangle} \langle e^{\phi(J)} V_\varepsilon(\phi(x_1)) \dots V_\varepsilon(\phi(x_\ell)) \rangle_{\tilde{C}_\Lambda} \\ &= e^{-\frac{1}{2}\langle J, \tilde{C}_\Lambda J \rangle} \frac{1}{(2\pi)^{\ell/2}} \int_{\mathbb{R}^\ell} d\mathbf{u} \widehat{V}_\varepsilon(\mathbf{u}) \langle e^{i\phi(\sum_{j=1}^\ell u_j \delta_{x_j} - iJ)} \rangle_{\tilde{C}_\Lambda} \\ &= e^{-\frac{1}{2}\langle J, \tilde{C}_\Lambda J \rangle} \frac{1}{(2\pi)^{\ell/2}} \int_{\mathbb{R}^\ell} d\mathbf{u} \widehat{V}_\varepsilon(\mathbf{u}) e^{-\frac{1}{2}\langle \sum_{j=1}^\ell u_j \delta_{x_j} - iJ, \tilde{C}_\Lambda(\sum_{j=1}^\ell u_j \delta_{x_j} - iJ) \rangle} \\ &= \frac{1}{(2\pi)^{\ell/2}} \tilde{C}_\Lambda(0)^{-\ell/2} \int_{\mathbb{R}^\ell} d\mathbf{u} \widehat{V}_\varepsilon(\tilde{C}_\Lambda(0)^{-1/2} \mathbf{u}) e^{-\frac{1}{2} \sum_{j=1}^\ell u_j^2 - \sum_{i < j} u_i u_j \frac{\tilde{C}_\Lambda(x_i - x_j)}{\tilde{C}_\Lambda(0)} + i \frac{\langle \sum_{j=1}^\ell u_j \delta_{x_j}, \tilde{C}_\Lambda J \rangle}{\tilde{C}_\Lambda(0)^{1/2}}}, \end{aligned} \quad (2.16)$$

and we made a change of variables  $\mathbf{u} \mapsto \tilde{C}_\Lambda(0)^{-1/2} \mathbf{u}$  in the last step. Thus we can rearrange this expression as follows, relying on the fact that  $\mathbf{M}(\mathbf{x})$  is positive definite for  $\mathbf{x} \in D'_\delta$  by Lemma 2.3:

$$\begin{aligned} & \frac{1}{(2\pi)^{\frac{1}{2}\ell}} \tilde{C}_\Lambda(0)^{-\ell/2} \int_{\mathbb{R}^\ell} d\mathbf{u} \widehat{V}_\varepsilon(\tilde{C}_\Lambda(0)^{-1/2} \mathbf{u}) e^{-\frac{1}{2} \mathbf{u}^T \mathbf{M}(\mathbf{x}) \mathbf{u} + i \frac{\langle \sum_{j=1}^\ell u_j \delta_{x_j}, \tilde{C}_\Lambda J \rangle}{\tilde{C}_\Lambda(0)^{1/2}}} \\ &= \frac{1}{(2\pi)^\ell} \tilde{C}_\Lambda(0)^{-\ell/2} \int_{\mathbb{R}^\ell} d\mathbf{w} V_\varepsilon(\mathbf{w}) \int_{\mathbb{R}^\ell} d\mathbf{u} e^{-i\tilde{C}_\Lambda(0)^{-1/2} \mathbf{w}^T \mathbf{u}} e^{-\frac{1}{2} \mathbf{u}^T \mathbf{M}(\mathbf{x}) \mathbf{u}} e^{i\mathbf{q}^T \mathbf{u}} \\ &= \frac{1}{(2\pi)^\ell} \tilde{C}_\Lambda(0)^{-\ell/2} \int_{\mathbb{R}^\ell} d\mathbf{w} V_\varepsilon(\mathbf{w}) \int_{\mathbb{R}^\ell} d\mathbf{u} e^{-i(\tilde{C}_\Lambda(0)^{-1/2} \mathbf{w} - \mathbf{q})^T \mathbf{u}} e^{-\frac{1}{2} \mathbf{u}^T \mathbf{M}(\mathbf{x}) \mathbf{u}} \\ &= \frac{1}{(2\pi)^\ell} \int_{\mathbb{R}^\ell} d\mathbf{w} V_\varepsilon(\tilde{C}_\Lambda(0)^{1/2} \mathbf{w}) \int_{\mathbb{R}^\ell} d\mathbf{u} e^{-i(\mathbf{w} - \mathbf{q})^T \mathbf{u}} e^{-\frac{1}{2} \mathbf{u}^T \mathbf{M}(\mathbf{x}) \mathbf{u}} \\ &= \frac{1}{(2\pi)^{\ell/2}} \det(\mathbf{M}(\mathbf{x}))^{-1/2} \int_{\mathbb{R}^\ell} d\mathbf{w} V_\varepsilon(\tilde{C}_\Lambda(0)^{1/2} \mathbf{w}) e^{-\frac{1}{2}(\mathbf{w} - \mathbf{q})^T \mathbf{M}(\mathbf{x})^{-1}(\mathbf{w} - \mathbf{q})}. \end{aligned} \quad (2.17)$$

By taking the limit  $\varepsilon \rightarrow 0$  in the first line of (2.16) and in the last line of (2.17) and applying dominated convergence we conclude the proof.  $\square$

**Proof of Proposition 2.1.** First we estimate the volume of  $D_\delta$  for  $\delta$  given by (2.8). We define  $D_{\delta;i,j} := \{\mathbf{x} \in B^\ell \mid |x_i - x_j| \leq \delta\}$ . Clearly,

$$D_\delta \subset \bigcup_{i < j} D_{\delta;i,j} \quad \Rightarrow \quad |D_\delta| \leq \sum_{i < j} |D_{\delta;i,j}|. \quad (2.18)$$

We note that  $|D_{\delta;i,j}| \leq cL^{d(\ell-1)}\delta^d$  hence  $|D_\delta| \leq c\ell^2 L^{d(\ell-1)}\delta^d$ . Now we observe that

$$\begin{aligned} \left| \int_{B^\ell} d\mathbf{x} \mathbb{1}_{D_\delta}(\mathbf{x}) \frac{\langle e^{\phi(J)} V(\phi(x_1)) \dots V(\phi(x_\ell)) \rangle_{\tilde{C}_\Lambda}}{\langle e^{\phi(J)} \rangle_{\tilde{C}_\Lambda}} \right| \\ \leq c \|V\|_\infty^\ell \ell^2 L^{d(\ell-1)} \delta^d \leq c \|V\|_\infty^\ell \ell^6 L^{d\ell} \frac{1}{\sqrt{\log(\Lambda)}}, \end{aligned} \quad (2.19)$$

$$\begin{aligned} \left| \int_{B^\ell} d\mathbf{x} \mathbb{1}_{D_\delta}(\mathbf{x}) \frac{\langle e^{\phi(J)} V(\phi(x_1)) \rangle_{\tilde{C}_\Lambda}}{\langle e^{\phi(J)} \rangle_{\tilde{C}_\Lambda}} \dots \frac{\langle e^{\phi(J)} V(\phi(x_\ell)) \rangle_{\tilde{C}_\Lambda}}{\langle e^{\phi(J)} \rangle_{\tilde{C}_\Lambda}} \right| \\ \leq c \|V\|_\infty^\ell \ell^2 L^{d(\ell-1)} \delta^d \leq c \|V\|_\infty^\ell \ell^6 L^{d\ell} \frac{1}{\sqrt{\log(\Lambda)}}, \end{aligned} \quad (2.20)$$

where we entered with the modulus under the Gaussian integrals and used positivity of  $e^{\phi(J)}$ . We also exploited that  $\log(\Lambda) \geq 1$ , hence

$$\delta^d = \left( \frac{\ell}{\sqrt{\log(\Lambda)}} \right)^{\frac{d}{d-3/2}} \leq \ell^4 \frac{1}{\sqrt{\log(\Lambda)}}. \quad (2.21)$$

We also note that the bounds (2.19), (2.20) remain true if we replace  $\mathbb{1}_{D_\delta}(\cdot)$  under the integral by one and  $\delta^d$  on the r.h.s by  $L^d$ . Thereby we obtain the factorization property (2.1) for  $\Lambda \leq \Lambda_0$ , where  $\Lambda_0$  appeared in Lemma 2.3. Thus it suffices to consider  $\Lambda \geq \Lambda_0$  in the following.

Let us look at the  $D'_\delta$  part. By Lemma 2.4, the relevant quantity has the form:

$$\begin{aligned} \int_{B^\ell} d\mathbf{x} \mathbb{1}_{D'_\delta}(\mathbf{x}) \left[ \frac{\langle e^{\phi(J)} V(\phi(x_1)) \dots V(\phi(x_\ell)) \rangle_{\tilde{C}_\Lambda}}{\langle e^{\phi(J)} \rangle_{\tilde{C}_\Lambda}} - \frac{\langle e^{\phi(J)} V(\phi(x_1)) \rangle_{\tilde{C}_\Lambda}}{\langle e^{\phi(J)} \rangle_{\tilde{C}_\Lambda}} \dots \frac{\langle e^{\phi(J)} V(\phi(x_\ell)) \rangle_{\tilde{C}_\Lambda}}{\langle e^{\phi(J)} \rangle_{\tilde{C}_\Lambda}} \right] \\ = \int_{B^\ell} d\mathbf{x} \mathbb{1}_{D'_\delta}(\mathbf{x}) \left\{ (2\pi)^{-\ell/2} \det(\mathbf{M})^{-1/2} \int_{\mathbb{R}^\ell} d\mathbf{w} V(\tilde{C}_\Lambda(0)^{1/2} \mathbf{w}) e^{-\frac{1}{2}(\mathbf{w}-\mathbf{q})^T \mathbf{M}^{-1}(\mathbf{w}-\mathbf{q})} \right. \\ \left. - (2\pi)^{-\ell/2} \int d\mathbf{w} V(\tilde{C}_\Lambda(0)^{1/2} \mathbf{w}) e^{-\frac{1}{2}(\mathbf{w}-\mathbf{q})^T (\mathbf{w}-\mathbf{q})} \right\}. \end{aligned} \quad (2.22)$$

We rewrite the expression in curly brackets as follows:

$$\begin{aligned} (2\pi)^{-\ell/2} \int_0^1 d\alpha \partial_\alpha \left[ \det(\mathbf{M}_\alpha)^{-1/2} \int_{\mathbb{R}^\ell} d\mathbf{w} V(\tilde{C}_\Lambda(0)^{1/2} \mathbf{w}) e^{-\frac{1}{2}(\mathbf{w}-\mathbf{q})^T \mathbf{M}_\alpha^{-1}(\mathbf{w}-\mathbf{q})} \right] \\ = -\frac{1}{2} (2\pi)^{-\ell/2} \int_0^1 d\alpha \left[ \det(\mathbf{M}_\alpha)^{-1/2} \text{Tr}(\mathbf{M}_\alpha^{-1} \partial_\alpha \mathbf{M}_\alpha) \int_{\mathbb{R}^\ell} d\mathbf{w} V(\tilde{C}_\Lambda(0)^{1/2} \mathbf{w}) e^{-\frac{1}{2}(\mathbf{w}-\mathbf{q})^T \mathbf{M}_\alpha^{-1}(\mathbf{w}-\mathbf{q})} \right] \end{aligned} \quad (2.23)$$

$$\begin{aligned} + \frac{1}{2} (2\pi)^{-\ell/2} \int_0^1 d\alpha \left[ \det(\mathbf{M}_\alpha)^{-1/2} \int_{\mathbb{R}^\ell} d\mathbf{w} V(\tilde{C}_\Lambda(0)^{1/2} \mathbf{w}) \times \right. \\ \left. \times (\mathbf{w}-\mathbf{q})^T \mathbf{M}_\alpha^{-2} \partial_\alpha \mathbf{M}_\alpha (\mathbf{w}-\mathbf{q}) e^{-\frac{1}{2}(\mathbf{w}-\mathbf{q})^T \mathbf{M}_\alpha^{-1}(\mathbf{w}-\mathbf{q})} \right], \end{aligned} \quad (2.24)$$

where we used that  $\mathbf{M}_\alpha(\mathbf{x})$  is positive definite for  $\mathbf{x} \in D'_\delta$ , thus  $\det(\mathbf{M}_\alpha)^{-1/2} = e^{-\frac{1}{2}\text{Tr}(\log(\mathbf{M}_\alpha))}$ . To estimate the respective terms above we bound  $V$  by its supremum and change variables  $\mathbf{w} \mapsto \mathbf{w} + \mathbf{q}$

$$\begin{aligned}
|(2.23)| &\leq (2\pi)^{-\ell/2} \|V\|_\infty^\ell \frac{1}{2} \int_0^1 d\alpha |\text{Tr}(\mathbf{M}_\alpha^{-1} \partial_\alpha \mathbf{M}_\alpha)| \int_{\mathbb{R}^\ell} d\mathbf{w} e^{-\frac{1}{2}\mathbf{w}^T \mathbf{w}} \\
&= \frac{1}{2} \|V\|_\infty^\ell \int_0^1 d\alpha |\text{Tr}(\mathbf{M}_\alpha^{-1} \partial_\alpha \mathbf{M}_\alpha)| \leq \frac{1}{2} \|V\|_\infty^\ell \int_0^1 d\alpha \ell \|\mathbf{M}_\alpha^{-1} \partial_\alpha \mathbf{M}_\alpha\| \\
&\leq c\ell \|V\|_\infty^\ell \frac{1}{\sqrt{\log(\Lambda)}}, \tag{2.25}
\end{aligned}$$

where we used the bound (2.9). Similarly,

$$\begin{aligned}
|(2.24)| &\leq (2\pi)^{-\ell/2} \|V\|_\infty^\ell \frac{1}{2} \int_0^1 d\alpha \|\mathbf{M}_\alpha^{-1} \partial_\alpha \mathbf{M}_\alpha\| \int_{\mathbb{R}^\ell} d\mathbf{w} \mathbf{w}^T \mathbf{w} e^{-\frac{1}{2}\mathbf{w}^T \mathbf{w}} \\
&= \ell \|V\|_\infty^\ell \frac{1}{2} \int_0^1 d\alpha \|\mathbf{M}_\alpha^{-1} \partial_\alpha \mathbf{M}_\alpha\| \leq c\ell \frac{1}{\sqrt{\log(\Lambda)}}. \tag{2.26}
\end{aligned}$$

Thus the claim follows from (2.19)–(2.22).  $\square$

**Proof of Theorem 1.1.** We consider the functional  $\Sigma_\Lambda(J)$  and expand  $e^{-\lambda\mathcal{V}(\phi)}$  into a power series in  $\lambda$ . Since the function  $V$  is bounded, the series is absolutely convergent for any  $\lambda \in \mathbb{R}$ . We write

$$\begin{aligned}
\Sigma_\Lambda(J) &= \frac{\langle e^{\phi(J)} e^{-\lambda\mathcal{V}(\phi)} \rangle_{\tilde{C}_\Lambda}}{\langle e^{\phi(J)} \rangle_{\tilde{C}_\Lambda}} = \exp\left(-\lambda \frac{\langle e^{\phi(J)} \mathcal{V}(\phi) \rangle_{\tilde{C}_\Lambda}}{\langle e^{\phi(J)} \rangle_{\tilde{C}_\Lambda}}\right) \\
&\quad + \underbrace{\sum_{\ell=0}^{\infty} \frac{(-\lambda)^\ell}{\ell!} \left( \frac{\langle e^{\phi(J)} \mathcal{V}(\phi)^\ell \rangle_{\tilde{C}_\Lambda}}{\langle e^{\phi(J)} \rangle_{\tilde{C}_\Lambda}} - \left( \frac{\langle e^{\phi(J)} \mathcal{V}(\phi) \rangle_{\tilde{C}_\Lambda}}{\langle e^{\phi(J)} \rangle_{\tilde{C}_\Lambda}} \right)^\ell \right)}_{R_\Lambda}. \tag{2.27}
\end{aligned}$$

By Proposition 2.1, we obtain that the rest term  $R_\Lambda$ , defined by (2.27), satisfies

$$|R_\Lambda| \leq c \frac{e^{c_1 \lambda |B| \|V\|_\infty}}{\sqrt{\log(\Lambda)}}. \tag{2.28}$$

Consequently,

$$\Sigma_\Lambda^c(J) = \log(\Sigma_\Lambda(J)) = -\lambda \frac{\langle e^{\phi(J)} \mathcal{V}(\phi) \rangle_{\tilde{C}_\Lambda}}{\langle e^{\phi(J)} \rangle_{\tilde{C}_\Lambda}} + \log\left(1 + \exp\left(\lambda \frac{\langle e^{\phi(J)} \mathcal{V}(\phi) \rangle_{\tilde{C}_\Lambda}}{\langle e^{\phi(J)} \rangle_{\tilde{C}_\Lambda}}\right) R_\Lambda\right). \tag{2.29}$$

Now, making use of Lemma 2.4 for  $\ell = 1$ , we can write

$$\frac{\langle e^{\phi(J)} \mathcal{V}(\phi) \rangle_{\tilde{C}_\Lambda}}{\langle e^{\phi(J)} \rangle_{\tilde{C}_\Lambda}} = \frac{1}{(2\pi)^{1/2}} \int_B dx \int_{\mathbb{R}} dw V((Z_\Lambda C_\Lambda(0))^{1/2} w) e^{-\frac{1}{2}(w - (\frac{Z_\Lambda}{C_\Lambda(0)})^{1/2} \langle \delta_x, C_\Lambda J \rangle)^2}. \tag{2.30}$$

This expression is bounded in  $\Lambda$ , since we can estimate  $V$  by its supremum and then shift the  $w$ -variable. Thus the logarithm on the r.h.s. of (2.29) tends to zero by estimate (2.28). Thus it suffices to compute the limit  $\Lambda \rightarrow \infty$  of (2.30) for the respective items of Theorem 1.1:

- (a) For  $-\infty < \eta < -1$ ,  $Z_\Lambda = C_\Lambda(0)^\eta$ , we have  $Z_\Lambda/C_\Lambda(0) \rightarrow 0$  and  $Z_\Lambda C_\Lambda(0) \rightarrow 0$ . Then, by the dominated convergence and assumption (A<sub>1</sub>),

$$(2.30) \rightarrow \frac{1}{(2\pi)^{1/2}} \int_B dx \int_{\mathbb{R}} dw [V_+ \theta(w) + V_- \theta(-w)] e^{-\frac{1}{2}w^2}. \tag{2.31}$$

As  $\theta(w) = \frac{1}{2}(\text{sgn}(w) + 1)$ , and  $\text{sgn}$  is antisymmetric, we obtain the claim.

(b) For  $\eta = -1$  we have  $Z_\Lambda/C_\Lambda(0) \rightarrow 0$  but  $Z_\Lambda C_\Lambda(0) \rightarrow 1$ . Then, recalling that  $V$  may not be continuous, we change variables before computing the limit:

$$\begin{aligned}
(2.30) &= \frac{1}{(2\pi)^{1/2}} \int_B dx \int_{\mathbb{R}} dw V((Z_\Lambda C_\Lambda(0))^{1/2} w) e^{-\frac{1}{2} \left( w - \left( \frac{Z_\Lambda}{C_\Lambda(0)} \right)^{1/2} \langle \delta_x, C_\Lambda J \rangle \right)^2} \\
&= \frac{1}{(2\pi)^{1/2}} \int_B dx \int_{\mathbb{R}} \frac{dw}{(Z_\Lambda C_\Lambda(0))^{1/2}} V(w) e^{-\frac{1}{2} \left( \frac{w}{(Z_\Lambda C_\Lambda(0))^{1/2}} - \left( \frac{Z_\Lambda}{C_\Lambda(0)} \right)^{1/2} \langle \delta_x, C_\Lambda J \rangle \right)^2} \\
&\rightarrow \frac{1}{(2\pi)^{1/2}} \int_B dx \int_{\mathbb{R}} dw V(w) e^{-\frac{1}{2} w^2}.
\end{aligned} \tag{2.32}$$

(c) For  $-1 < \eta < 1$  we have  $Z_\Lambda/C_\Lambda(0) \rightarrow 0$  and  $Z_\Lambda C_\Lambda(0) \rightarrow \infty$ . Then, by assumption (A<sub>2</sub>),

$$(2.30) \rightarrow \frac{1}{(2\pi)^{1/2}} \int_B dx \int_{\mathbb{R}} dw [V^+ \theta(w) + V^- \theta(-w)] e^{-\frac{1}{2} w^2}. \tag{2.33}$$

(d) For  $\eta = 1$  we have  $Z_\Lambda/C_\Lambda(0) \rightarrow 1$  and  $Z_\Lambda C_\Lambda(0) \rightarrow \infty$ . Hence, again by (A<sub>2</sub>),

$$(2.30) \rightarrow \frac{1}{(2\pi)^{1/2}} \int_B dx \int_{\mathbb{R}} dw [V^+ \theta(w) + V^- \theta(-w)] e^{-\frac{1}{2} (w - \langle \delta_x, C_\Lambda J \rangle)^2}. \tag{2.34}$$

(e) For  $1 < \eta < \infty$  we have  $Z_\Lambda/C_\Lambda(0) \rightarrow \infty$  and  $Z_\Lambda C_\Lambda(0) \rightarrow \infty$ . Then, assuming that  $\hat{J}$  is compactly supported, vanishing of  $x \mapsto \langle \delta_x, C J \rangle$  on a set of non-zero Lebesgue measure implies  $J = 0$ . Thus inserting  $1 = \mathbb{1}(\langle \delta_x, C J \rangle = 0) + \mathbb{1}(\langle \delta_x, C J \rangle \neq 0)$  and using (A<sub>2</sub>), we obtain

$$\begin{aligned}
(2.30) &= \frac{1}{(2\pi)^{1/2}} \int_B dx \int_{\mathbb{R}} dw V((Z_\Lambda C_\Lambda(0))^{1/2} (w + \left( \frac{Z_\Lambda}{C_\Lambda(0)} \right)^{1/2} \langle \delta_x, C_\Lambda J \rangle)) e^{-\frac{1}{2} w^2} \\
&\rightarrow \mathbb{1}(J = 0) \frac{1}{(2\pi)^{1/2}} \int_B dx \int_{\mathbb{R}} dw (V^+ \theta(w) + V^- \theta(-w)) e^{-\frac{1}{2} w^2} \\
&\quad + \mathbb{1}(J \neq 0) \int_B dx (V^+ \theta(\langle \delta_x, C J \rangle) + V^- \theta(-\langle \delta_x, C J \rangle)).
\end{aligned} \tag{2.35}$$

After obvious rearrangements this gives the claim. (Without the assumption that  $\hat{J}$  is compactly supported, the limit in the  $\langle \delta_x, C J \rangle = 0$  case would depend on the convergence properties of the sequences  $\Lambda \mapsto \left( \frac{Z_\Lambda}{C_\Lambda(0)} \right)^{1/2} \langle \delta_x, (C_\Lambda - C) J \rangle$ . These may a priori vary in a complicated manner with  $x$  and  $J$ ).  $\square$

One unfortunate feature of Theorem 1.1 is the absence of any  $J$  dependence of  $\hat{S}^c(J)$  in parts (a)–(c), i.e., for  $\eta < 1$ . With the Sine-Gordon model in mind, one could hope that such dependence emerges if we relax the assumptions (A<sub>1</sub>), (A<sub>2</sub>), e.g., by admitting some oscillations of  $V$  near zero or infinity. It turns out that this would not help. In fact, let us set  $t_\Lambda := C_\Lambda(0)^{1/2}$ ,  $Z_\Lambda = C_\Lambda(0)^\eta = t_\Lambda^{2\eta}$  and  $a(x) := \langle \delta_x, C_\Lambda J \rangle$ . Then, by (2.30), we can write

$$\lambda \frac{\langle e^{\phi(J)} \mathcal{V}(\phi) \rangle_{\tilde{C}_\Lambda}}{\langle e^{\phi(J)} \rangle_{\tilde{C}_\Lambda}} = \frac{\lambda}{(2\pi)^{1/2}} \int_B dx \int_{\mathbb{R}} dw V(t_\Lambda^{\eta+1} w) e^{-\frac{1}{2} (w - t_\Lambda^{\eta-1} a(x))^2}. \tag{2.36}$$

Now we introduce a parameter  $s \in [0, 1]$ , replace  $a(x)$  with  $sa(x)$  on the r.h.s. of (2.36) and call the resulting function  $f_\Lambda(s)$ . By estimating  $|f_\Lambda(1) - f_\Lambda(0)| \leq \int_0^1 ds |\partial_s f_\Lambda(s)|$  and taking the limit  $\Lambda \rightarrow \infty$  we obtain that the  $J$  dependence is trivial for  $\eta < 1$  relying only on the boundedness of  $V$ . This follows from

$$\partial_s \int_{\mathbb{R}} dw V(t_\Lambda^{\eta+1} w) e^{-\frac{1}{2} (w - t_\Lambda^{\eta-1} sa(x))^2} = t_\Lambda^{\eta-1} a(x) \int_{\mathbb{R}} dw V(t_\Lambda^{\eta+1} w) \partial_w e^{-\frac{1}{2} (w - t_\Lambda^{\eta-1} sa(x))^2} \tag{2.37}$$



due to the pre-factor  $t_\Lambda^{\eta-1}$ . One could try to compensate this pre-factor by choosing  $\lambda$  growing with  $\Lambda$ , but such a growth would immediately spoil the bound (2.28) and thus undermine the factorization mechanism from Proposition 2.1. Incidentally, this is how the Sine-Gordon model  $V(\phi) = \cos(\beta\phi)$  evades our analysis: The Wick ordering gives effective behaviour of the coupling constant  $\lambda \sim e^{\frac{1}{2}\beta C_\Lambda(0)}$  which is incompatible with the factorization.

### 3 Generalization

It is clear from estimate (2.28) that our methods apply to interaction functions  $V$  depending on  $\Lambda$ , as long as their supremum norms remain uniformly bounded in  $\Lambda$ . We consider an  $\Lambda$  dependent interaction function of the form  $V_\Lambda(w) := V(z_\Lambda w)$  and the corresponding interaction term  $\mathcal{V}_\Lambda(\phi) := \int_B dx V_\Lambda(\phi(x))$ . We can compute the UV limit of the resulting modified generating functional, which we still denote by  $\Sigma_\Lambda^c(J)$ , using obvious modifications of formulas (2.29), (2.30). Thus the problem reduces to computing the limit  $\Lambda \rightarrow \infty$  of

$$\frac{\langle e^{\phi(J)} \mathcal{V}_\Lambda(\phi) \rangle_{\tilde{C}_\Lambda}}{\langle e^{\phi(J)} \rangle_{\tilde{C}_\Lambda}} = \frac{1}{(2\pi)^{1/2}} \int_B dx \int_{\mathbb{R}} dw V(z_\Lambda (Z_\Lambda C_\Lambda(0))^{1/2} w) e^{-\frac{1}{2} \left( w - \left( \frac{Z_\Lambda}{C_\Lambda(0)} \right)^{1/2} \langle \delta_x, C_\Lambda J \rangle \right)^2}. \quad (3.1)$$

Setting for concreteness  $z_\Lambda := C_\Lambda(0)^\kappa$ ,  $\kappa \in \mathbb{R}$ , Theorem 1.1 can be generalized<sup>1</sup>. We focus here on part (d), as it brings interesting new insights. It splits into three cases depending on  $\kappa$ :

**Corollary 3.1.** *Set  $Z_\Lambda = C_\Lambda(0)$ ,  $z_\Lambda = C_\Lambda(0)^\kappa$ ,  $\kappa \in \mathbb{R}$ . Then the limit  $\Sigma^c(J) := \lim_{\Lambda \rightarrow \infty} \Sigma_\Lambda^c(J)$  exists for any  $J \in S(\mathbb{R}^d; \mathbb{R})$  under the assumptions specified below. It has the form:*

(d.1) *Let  $\kappa < -1$ . Then, assuming (A<sub>1</sub>),*

$$\Sigma^c(J) = -\lambda \frac{V_+ + V_-}{2} |B| - \lambda \frac{V_+ - V_-}{2} \frac{1}{(2\pi)^{1/2}} \int_B dx \int dw \operatorname{sgn}(w) e^{-\frac{1}{2} (w - \langle \delta_x, CJ \rangle)^2}. \quad (3.2)$$

(d.2) *Let  $\kappa = -1$ . Then, without additional assumptions,*

$$\Sigma^c(J) = -\lambda \frac{1}{(2\pi)^{1/2}} \int_B dx \int dw V(w) e^{-\frac{1}{2} (w - \langle \delta_x, CJ \rangle)^2}. \quad (3.3)$$

(d.3) *Let  $\kappa > -1$ . Then, assuming (A<sub>2</sub>),*

$$\Sigma^c(J) = -\lambda \frac{V^+ + V^-}{2} |B| - \lambda \frac{V^+ - V^-}{2} \frac{1}{(2\pi)^{1/2}} \int_B dx \int dw \operatorname{sgn}(w) e^{-\frac{1}{2} (w - \langle \delta_x, CJ \rangle)^2}. \quad (3.4)$$

We note that in the case (d.1) we obtain Schwinger functions of the form (1.10) up to the coupling constant renormalization, which is now  $\frac{V_+ - V_-}{2}$ . Hence non-Gaussianity holds if and only if  $V$  is discontinuous at zero. Thus our theories turn out to be sensitive to local regularity properties of the interaction functions.

In the case (d.2) the Schwinger functions differ more substantially from (1.10). By repeating the steps from Appendix B, we obtain, for  $n \neq 2$ ,

$$S_n^c(x_1, \dots, x_n) = -\lambda \left( \frac{1}{(2\pi)^{1/2}} \int dw (\partial^n V)(w) e^{-\frac{1}{2} w^2} \right) \int_B dx C(x - x_1) \dots C(x - x_n). \quad (3.5)$$

These  $n$ -point functions coincide with the tree level one-particle irreducible Schwinger functions of a perturbative theory with the interaction function given by the convolution  $V * \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}(\cdot)^2}$ . Thus the class of theories tractable by our methods is richer than one could expect from (1.10).

---

<sup>1</sup>Strictly speaking, in part (e) additional regularity assumptions on  $V$  may be needed to control the entire range of  $\kappa$ .

## 4 Conclusion and outlook

In this paper we succeeded in computing exactly the UV limits  $\Sigma^c(J)$  of the modified generating functionals (1.4) for a large class of bounded measurable interaction functions  $V$ . We computed the connected Schwinger functions  $S_{n \neq 2}^c$  as derivatives of  $\Sigma^c(J)$  w.r.t.  $J$  and observed that they coincide with the tree-level one-particle irreducible Schwinger functions of the  $\text{erf}(\phi/\sqrt{2})$  theory with the coupling constant renormalized by the factor  $\frac{V^+ - V^-}{2}$ . In Section 3 we noted that  $V$  may depend on  $\Lambda$  as long as it is uniformly bounded in the supremum norm. In the class of interactions  $\phi \mapsto V(z_\Lambda \phi)$ , we could change the coupling constant renormalization to  $\frac{V_+ - V_-}{2}$  so that the Schwinger functions probe the continuity of  $V$  at zero.

A more intriguing direction is to look at functions  $V$  which become increasingly rough as  $\Lambda$  increases, but remain uniformly bounded in  $\Lambda$ . For example, the number of discontinuities could increase with  $\Lambda$ . The hope is that thinking in this direction one could invalidate (2.37) by disintegration of the  $w$ -integrals and obtain non-trivial Schwinger functions  $S_{n \neq 2}^c$  also for  $\eta < 1$ . This would be an important step forward, as only for  $\eta \leq 0$  a finite two-point connected Schwinger function is immediately available, as we now explain.

Strictly speaking, the limiting modified functionals  $\Sigma^c(J)$  do not carry any information about the two-point function of the limiting theory. But it appears consistent with (1.4), (1.2) to define the two-point connected Schwinger function by

$$S_2^c(x_1, x_2) := \frac{\delta}{\delta J(x_1)} \frac{\delta}{\delta J(x_2)} \Sigma^c(J)|_{J=0} + \lim_{\Lambda \rightarrow \infty} Z_\Lambda C_\Lambda(x_1, x_2). \quad (4.1)$$

We note that it is finite and well defined only for  $\eta \leq 0$ , in which case we have  $S_{n \neq 2}^c = 0$ . For  $\eta < 0$  we have  $S_2^c(x_1, x_2) = 0$ , i.e., all the Schwinger functions are zero, whereas for  $\eta = 0$  we obtain  $S_2^c(x_1, x_2) = C(x_1, x_2)$  and reproduce the free field theory. In the case of  $\eta = 1$ , in which we obtained the Schwinger functions (1.10), we have  $S_2^c(x_1, x_2) = \infty$ . We observe that this divergence coexists with relation (1.11), which demonstrates the limitations of the conventional UV stability criterion.

From the point of view of Euclidean constructive QFT the blow-up of the two-point function is an obstacle, which undermines the Osterwalder-Schrader reconstruction of the model as a Minkowskian QFT. From the Minkowskian perspective this divergence may be a manifestation of the fact that the vacuum vector is not in the domain of the quantum field or that pointlike-localized fields do not exist at all in these models. Actually, there are general arguments pointing in a similar direction: if  $Z_\Lambda > 1$  or  $d > 4$  then interacting scalar fields must violate the canonical commutation relations [We, Eq (10.7.22)], [Ba87]. As a singular behaviour of pointlike localized fields is not an obstacle to the existence of a QFT in the Haag-Kastler sense, it may be fruitful to pursue a direct Minkowskian construction of models with bounded interactions in the spirit of [BF20, BFR21].

Nevertheless, it would be preferable to tame the blow-up of the two-point function. As the problem occurs at the level of free theory, it may be solvable by adding some spectator field, not interacting with the original system. To illustrate the idea, we note that the denominator in (1.4) satisfies  $\frac{1}{S_{0,\Lambda}(J)} = \langle e^{i\varphi(J)} \rangle_{\tilde{C}_\Lambda}$ , thus differs from  $S_{0,\Lambda}(J)$  only by the imaginary unit in the exponent. Thus we can express  $\Sigma_\Lambda(J)$  as a (non-modified) generating functional of the complex field  $\psi := \phi + i\varphi$  and interpret the  $n$ -point functions (1.10), including  $n = 2$ , as connected Schwinger functions of  $\psi$ . By an analogous construction we obtain the Schwinger functions of  $\psi^*$ , but  $n$ -point functions involving both  $\psi$  and  $\psi^*$  are not immediately available. Their construction amounts to the problem of coupling of two probability measures, whose solution is, in general, not unique. While naive choices lead to the blow-up of such Schwinger functions, one can hope for a solution consistent with Osterwalder-Schrader axioms.

Another direction is to take simultaneously the UV limit and the classical limit, i.e., to analyse

$$\check{S}_\Lambda^c(J) := \hbar_\Lambda \log \langle e^{\hbar_\Lambda^{-1} \phi(J)} e^{-\lambda \hbar_\Lambda^{-1} \mathcal{V}(\phi)} \rangle_{\hbar_\Lambda C_\Lambda}. \quad (4.2)$$

It is easy to show by our methods that  $\check{S}_\Lambda^c(J)$  is bounded in  $\Lambda$  for  $\hbar_\Lambda = C_\Lambda(0)^{-1}$ . Based on preliminary computations, we conjecture that it has a limit and  $\lim_{\Lambda \rightarrow \infty} \check{S}_\Lambda^c(J) = \Sigma^c(J) + \frac{1}{2} \langle J, C J \rangle$ . In this case all the

Schwinger functions are finite and the family (1.10) is naturally complemented by the free covariance  $C$ . This is reminiscent of formal perturbation theory, where the classical limit extracts the tree-level connected Schwinger functions. Unfortunately, the Osterwalder-Schrader reconstruction would fail to give a non-trivial QFT, due to the Jost-Schroer theorem. But the construction would give an interesting, non-Gaussian, statistical physics model.

## A Existence of the functional measure with cut-offs

We define the Sobolev spaces  $L^{2,s}(\mathbb{R}^d) := \{ \phi \in S'(\mathbb{R}^d) \mid (1 - \Delta)^{s/2} \phi \in L^2(\mathbb{R}^d) \}$  for  $s \in \mathbb{N}$ .

**Lemma A.1.** *Let  $\chi \in C_0^\infty(\mathbb{R}^d)$  be an approximate characteristic function of the set  $B$ . Then the Gaussian measure  $\nu_{\tilde{C}_\Lambda}$  is concentrated on  $\phi$  s.t.  $\chi\phi \in L^{2,s}(\mathbb{R}^d)$  for any  $s \in \mathbb{N}$ .*

**Proof.** For any  $c > 0$

$$\begin{aligned} \int d\nu_{\tilde{C}_\Lambda}(\phi) \mathbb{1}(\|(1 - \Delta)^{s/2} \chi\phi\|_2^2 = \infty) &\leq \frac{1}{c} \int d\nu_{\tilde{C}_\Lambda}(\phi) \|(1 - \Delta)^{s/2} \chi\phi\|_2^2 \\ &= \frac{1}{c} \int dx \chi(x) \int d\nu_{\tilde{C}_\Lambda}(\phi) (1 - \Delta_{x'})^s \chi(x') \phi(x) \phi(x')|_{x'=x} \\ &= \frac{1}{c} \int dx \chi(x) (1 - \Delta_{x'})^s \chi(x') \tilde{C}_\Lambda(x - x')|_{x'=x}. \end{aligned} \quad (\text{A.1})$$

Recalling formula (1.1) for  $C_\Lambda$ , we obtain that (A.1) is finite for any fixed  $\Lambda$ . As  $c$  can be chosen arbitrarily large, this completes the proof.  $\square$

We recall the Sobolev embedding theorem [BCD, Theorem 1.66]:

**Theorem A.2.** *The space  $L^{2,s}(\mathbb{R}^d)$  embeds continuously in the Hölder space  $C^{k,\rho}(\mathbb{R}^d)$  if  $s \geq d/2 + k + \rho$  for some  $k \in \mathbb{N}$  and  $\rho \in ]0, 1[$ .*

For the Hölder space  $C^{k,\rho}(\mathbb{R}^d)$  we refer to [BCD, Definition 1.49]. We only need here that it consists of continuous functions.

**Corollary A.3.** *For any bounded measurable function  $V$  the measure*

$$d\mu_{\tilde{C}_\Lambda}(\phi) = \frac{1}{\mathcal{Z}_\Lambda} \exp\left(-\lambda \int_B dx V(\phi(x))\right) d\nu_{\tilde{C}_\Lambda}(\phi), \quad (\text{A.2})$$

where  $\mathcal{Z}_\Lambda$  is the normalization constant, is well defined.

**Proof.** We note that  $\int_B dx V(\phi(x)) = \int_B dx V((\chi\phi)(x))$ . By Lemma A.1, we can assume that  $\chi\phi \in L^{2,s}(\mathbb{R}^d)$ . Thus, by Theorem A.2, it has a continuous representative which we can compose with  $V$ . By the Jensen inequality, boundedness of  $V$  and compactness of  $B$  we have  $\mathcal{Z}_\Lambda > 0$ .  $\square$

## B Computation of Schwinger functions

Recall the generating functional (1.8)

$$\Sigma^c(J) = -\lambda(V^+ + V^-)|_B - \lambda \frac{(V^+ - V^-)}{2} \frac{1}{(2\pi)^{1/2}} \int_B dx \int dw \operatorname{sgn}(w) e^{-\frac{1}{2}(w - \langle \delta_x, CJ \rangle)^2}. \quad (\text{B.1})$$

We can write, by Plancherel,

$$\int dw \operatorname{sgn}(w) e^{-\frac{1}{2}(w - \langle \delta_x, CJ \rangle)^2} = \int du \widehat{\operatorname{sgn}}(u) e^{-\frac{1}{2}u^2} e^{iu \langle \delta_x, CJ \rangle}, \quad (\text{B.2})$$

which is the action of the distribution  $\widehat{\text{sgn}}$  on a test function. As we are only interested in Schwinger functions smeared with test functions  $f_1 \dots f_n \in S(\mathbb{R}^d; \mathbb{R})$ , it suffices to compute functional derivatives w.r.t.  $J$  at zero in the direction of such functions. This amounts to:

$$\partial_{\varepsilon_1} \dots \partial_{\varepsilon_n} \int du \widehat{\text{sgn}}(u) e^{-\frac{1}{2}u^2} e^{iu \sum_{i=1}^n \varepsilon_i \langle \delta_x, C f_i \rangle} \Big|_{\varepsilon_1, \dots, \varepsilon_n = 0}. \quad (\text{B.3})$$

It is easily seen that these derivatives commute with the action of  $\widehat{\text{sgn}}$  as the approximating sequences converge in  $S(\mathbb{R}^d; \mathbb{R})$ . This expression equals

$$\int du (iu)^n \widehat{\text{sgn}}(u) e^{-\frac{1}{2}u^2} (\langle \delta_x, C f_1 \rangle \dots \langle \delta_x, C f_n \rangle). \quad (\text{B.4})$$

We note the simple identities

$$\begin{aligned} \int du (iu)^n \widehat{\text{sgn}}(u) e^{-\frac{1}{2}u^2} &= 2 \int du (\widehat{\partial^{n-1} \delta})(u) e^{-\frac{1}{2}u^2} \\ &= 2 \int dw (\partial^{n-1} \delta)(w) e^{-\frac{1}{2}w^2} \\ &= 2(-1)^{n-1} (\partial^{n-1} e^{-\frac{1}{2}w^2})_{w=0} = \sqrt{2\pi} (\partial_w^n \text{erf}(w/\sqrt{2})) \Big|_{w=0}, \end{aligned} \quad (\text{B.5})$$

where we used the definition of the error function  $\text{erf}(w) := \frac{2}{\sqrt{\pi}} \int_0^w dw' e^{-(w')^2}$  and the fact that it is antisymmetric.

## C Proof of Lemma 2.2

By performing the  $p_0$ -integral in (1.1) we obtain

$$C_\Lambda(x) = \int \frac{d\vec{p}}{(2\pi)^d} \int_{1/\Lambda^2}^\infty d\alpha \sqrt{\frac{\pi}{\alpha}} e^{-\alpha((\vec{p})^2 + m^2)} e^{-\frac{1}{2}(2\alpha)^{-1}|x|^2}, \quad (\text{C.1})$$

hence  $0 < C_\Lambda(x)$  and  $C_\Lambda(x) < C_\Lambda(0)$  for  $x \neq 0$ . Property (2.4) follows directly from [GJ, Prop. 7.2.1]. To obtain (2.5), we compute also the  $\vec{p}$ -integral in (C.1):

$$C_\Lambda(x) = \frac{1}{2^d \pi^{d/2}} \int_{1/\Lambda^2}^\infty d\alpha \alpha^{-d/2} e^{-\alpha m^2} e^{-\frac{1}{2}(2\alpha)^{-1}|x|^2}. \quad (\text{C.2})$$

Consequently  $C_\Lambda(0) \geq c \int_{1/\Lambda^2}^2 d\alpha \alpha^{-d/2}$  for some  $c > 0$ .

## References

- [Ai82] M. Aizenman. *Geometric analysis of  $\phi^4$  fields and Ising models. I, II*. Commun. Math. Phys. **86**, 1–48 (1982).
- [AD21] M. Aizenman and H. Duminil-Copin. *Marginal triviality of the scaling limits of critical 4D Ising and  $\phi^4$  models*. Ann. of Math. (2) **194** no. 1 163–235 (2021).
- [BFR21] D. Bahns, K. Fredenhagen and K. Rejzner. *Local nets of von Neumann algebras in the Sine–Gordon model*. Commun. Math. Phys. **383**, (2021) 1–33.
- [BCD] H. Bahouri, J.-Y. Chemin and R. Danchin. *Fourier analysis and nonlinear partial differential equations*. Springer, 2011.

- [BD21] N. Barashkov and F. C. De Vecchi. *Elliptic stochastic quantization of Sinh-Gordon QFT*. arXiv:2108.12664.
- [BH22] R. Bauerschmidt and M. Hofstetter. *Maximum and coupling of the sine-Gordon field*. Ann. Probab. **50**, (2022) 455–508.
- [Ba87] K. Baumann. *On relativistic irreducible quantum fields fulfilling CCR*. J. Math. Phys. **28**, (1987) 697–704.
- [BDR24] A. Bonicelli, C. Dappiaggi and P. Rinaldi. *On the stochastic Sine-Gordon model: An interacting field theory approach*. Commun. Math. Phys. **405**, (2024) 288.
- [BF20] D. Buchholz and K. Fredenhagen. *A  $C^*$ -algebraic approach to interacting quantum field theories*. Commun. Math. Phys. **377**, (2020) 947–969.
- [CFM24] D. Cadamuro, M.B. Fröb and C. Moreira Ferrera. *The Sine–Gordon QFT in de Sitter spacetime*. Lett Math Phys **114**, (2024) 138.
- [CT16] D. Cadamuro and Y. Tanimoto. *Wedge-local observables in the deformed sine-Gordon model*. arXiv:1612.02073
- [Di13] J. Dimock. *The Renormalization Group According to Balaban - I. Small fields*. Rev. Math. Phys. **25**, (2013) 1330010.
- [FP77] J. Fröhlich and Y.M. Park. *Remarks on exponential interactions and the quantum sine-Gordon equation in two space-time dimensions*. Helv. Phys. Acta **50**, (1977) 315–329.
- [Fr82] J. Fröhlich. *On the triviality of  $\lambda\phi_d^4$  theories and the approach to the critical point in  $d > 4$  dimensions*. Nuclear Physics B **200**, (1982) 281–296.
- [GJ] J. Glimm and A. Jaffe. *Quantum Physics. A Functional Integral Point of View*. Second Edition. Springer, 1987.
- [GM24] M. Gubinelli and S.-J. Meyer. *The FBSDE approach to sine–Gordon up to  $6\pi$* . arXiv:2401.13648.
- [HS16] M. Hairer and H. Shen. *The dynamical Sine-Gordon model*. Commun. Math. Phys. **341**, (2016) 933–989.
- [Ko23] K.K. Kozłowski. *On convergence of form factor expansions in the infinite volume quantum Sinh-Gordon model in 1+1 dimensions*. Invent. math. **233**, (2023) 725–827.
- [Le08] G. Lechner. *Construction of quantum field theories with factorizing  $S$ -matrices*. Commun. Math. Phys. **277**, (2008) 821–860.
- [PT11] G. Peccati and M.S. Taqqu. *Wiener Chaos: Moments, Cumulants and Diagrams*. Springer, 2011.
- [Sa] M. Salmhofer. *Renormalization. An Introduction*. Springer, 1999.
- [We] S. Weinberg. *The quantum theory of fields. Volume 1: Foundations*. Cambridge University Press, 2005.