# CHERN-SCHWARTZ-MACPHERSON CLASSES IN THE POINT OF VIEW OF OBSTRUCTION THEORY AND LIPSCHITZ FRAMEWORK.

JEAN-PAUL BRASSELET, TADEUSZ MOSTOWSKI, AND THỦY NGUYỄN THỊ BÍCH

ABSTRACT. Since Chern and Grothendieck, Chern's characteristic class theory has made significant progress. In particular with regard to the classes of singular varieties. Conjectured by Grothendieck and Deligne and demonstrated by MacPherson, Chern classes of singular varieties have been defined in several ways, such as using polar varieties, Lagrangian theory... However, the initial definition using obstruction theory, due to Marie-Hélène Schwartz, has been forgotten. Despite the simple ideas that enabled the obstruction definition, their implementation using Whitney stratifications requires delicate and technical constructions. In the present article, we show that in the Lipschitz framework, the ideas of Marie-Hélène Schwartz lead to a simplified definition and construction of Chern classes of complex analytic varieties.

# 0. INTRODUCTION.

In his creative article [5], Chern gives various definitions of classes of hermitian manifolds. One of them uses obstruction theory for the construction of frames tangent to the manifold. In 1965, M.-H. Schwartz provides a definition of Chern classes for singular analytic varieties, using obstruction theory in the framework of Whitney stratifications. Her construction involves rather technical and delicate systems of tubular neighbourhoods of the strata.

The aim of this article is to show that the M.-H. Schwartz' ideas for defining Chern classes in the singular setting can be implemented in the framework of Lipschitz stratifications, instead of Whitney ones and that, in this framework, the M.-H. Schwartz' construction is simplified.

In this article, manifolds are smooth analytic complex varieties. A complex r-field tangent to a subset A of a complex manifold M is a set of r continuous complex vectors fields tangent to the manifold. When the vectors are  $\mathbb{C}$ -linearly independent at all points of A, we will speak of an r-frame. A singular point of an r-field is a point at which the r vectors fail to be  $\mathbb{C}$ -linearly independent. The classical obstruction theory (see [24]) says that, on an m-dimensional complex manifold M, it is possible to construct such a complex r-field without singularities on the 2m - 2r + 1-skeleton of a triangulation of M and with isolated singular points situated at the barycenters of the 2p = 2(m - r + 1) simplices of the triangulation. Let  $I(v^{(r)}, a)$  be the index of the r-field  $v^{(r)} = (v_1, \ldots, v_r)$  at the barycenter a of the 2p-simplex  $\sigma$ . The Chern

The first named author was supported by CNRS (UMR 7373) and Aix-Marseille University (I2M). The third named author was supported by the Brazilian São Paulo Research Foundation (FAPESP) grant #2023/07802-7.

class (in  $H^{2p}(M,\mathbb{Z})$ ) is the class of the cocycle whose value on the simplex  $\sigma$  is the index  $I(v^{(r)}, a)$ .

In this article, we consider a singular complex analytic variety X embedded in the complex Euclidean space  $\mathbb{C}^m$ . Note that our result applies to situations which admit Lipschitz stratifications (T. Mostowski [14]).

When using a triangulation compatible with a stratification of X, the obstruction dimension for the construction of a complex *r*-frame tangent to a stratum depends on the considered stratum. It is then clearly impossible to define a cocycle in the same way than in the smooth case because the dimension of the cocycle depends on the dimension of each stratum.

Given a stratification of X, a stratification of  $\mathbb{C}^m$  is obtained by adding the stratum  $\mathbb{C}^m \setminus X$ . Let (K) be a locally finite triangulation of  $\mathbb{C}^m$  compatible with the stratification of X, *i.e.* every open simplex lies in one and only one stratum. The first idea of Marie-Hélène Schwartz is to consider in  $\mathbb{C}^m$  a barycentric cellular decomposition (D) dual of the triangulation (K). A dual cell d which is an element of (D) is then topologically transverse to all strata S, that means (see Lemma 1.3) that

# $\operatorname{codim}(d) + \operatorname{codim}(S) = \operatorname{codim}(d \cap S).$

The intersection of a 2(m-r+1)-dimensional dual cell with a stratum is a cell whose dimension is equal to the obstruction dimension for constructing an *r*-frame tangent to that stratum. It is then possible to define a 2(m-r+1)-cocycle by attaching to each 2(m-r+1) cell of (D) meeting X the index of a stratified complex *r*-field with isolated singularities at the barycenters of the cells. Here, a stratified *r*-field means an *r*-field such that at each point x of the stratification of  $\mathbb{C}^m$ , the *r*-field is tangent to the stratum containing x.

The second idea of M.-H. Schwartz concerns the frames. When proving the Poincaré-Hopf theorem for singular varieties, M.-H. Schwartz has showed that considering stratified vector fields is not sufficient. She provides counter-examples for the Poincaré-Hopf theorem using vector fields which are only stratified. She has showed that one has to consider radial extension of stratified vector fields. The same situation appears in the definition of characteristic classes (Chern classes). M.-H. Schwartz considers stratified *r*-fields, obtained by suitable radial extensions. Such an *r*-field defines an element of the relative cohomology  $H^{2(m-r+1)}(\mathbb{C}^m, \mathbb{C}^m \setminus X)$ , called the Schwartz (cohomology) class of X. It is independent on the choices of stratification, triangulation and of the radial extension.

The M.-H. Schwartz' ideas are very clear and adapted. However, their implementation is rather technical. Under Whitney stratifications it uses delicate constructions of systems of adapted tubular neighbourhoods. Moreover, it needs to consider technical homotopies between frames issued from different strata.

Let us recall that, in his original article [14], T. Mostowski proved that analytic complex varieties admit Lipschitz stratifications.

In this article, we use the M.-H. Schwartz' ideas in the framework of Lipschitz stratifications instead of Whitney stratifications. That allows to simplify her construction. For example, the continuity of *r*-frames comes directly from Lipschitz properties without using homotopy between frames.

By Alexander duality isomorphism

$$H^{2(m-r+1)}(\mathbb{C}^m,\mathbb{C}^m\backslash X) \xrightarrow{\cong} H_{2(r-1)}(X),$$

the obtained homology classes are the MacPherson classes of the constructible function  $\mathbf{1}_X$  [4]. These homology classes have been defined by MacPherson [13], answering a conjecture by Deligne and Grothendieck [8, Note 87<sub>1</sub>].

Nowadays, these classes are called (Chern-)Schwartz-MacPherson classes.

The paper is organized as follows: In the first section, we recall the classical obstruction theory for manifolds and we explain the M.-H. Schwartz' idea for considering obstruction theory in the case of stratified singular varieties. The second section deals with Lipschitz stratifications: their definition and properties. In the third section, we show how Lipschitz properties allow to construct radial extensions of Lipschitz vector fields and frames. The last section is devoted to the definition of obstruction cocycles and classes in our setting.

### 1. A Few on obstruction theory

1.1. The smooth case. Let M be a locally compact complex analytic manifold with complex dimension m. The complex tangent bundle, denoted by TM is locally trivial.

For a fixed r, with  $1 \leq r \leq m$ , an *r*-field on a subset A of M is a set  $v^{(r)} = \{v_1, \ldots, v_r\}$  of r continuous complex vector fields defined on A and tangent to M. A singular point of  $v^{(r)}$  is a point where the vectors  $\{v_1, \ldots, v_r\}$  fail to be  $\mathbb{C}$ -linearly independent. An *r*-frame is a non-singular *r*-field.

The objective of the obstruction theory is to evaluate the obstruction to the construction of r sections of TM linearly independent at each point, *i.e.* an r-frame.

Let us denote by  $V_r(TM)$  the fibre bundle over M associated to TM and whose fibre at the point x of M is the set of r-frames of the fiber  $T_xM$  of TM at x. An r-frame on a subset A of M is then a section of  $V_r(TM)$  over A. The typical fibre of  $V_r(TM)$  is the Stiefel manifold of complex r-frames in  $\mathbb{C}^m$  denoted by  $V_r(\mathbb{C}^m)$ . In the same way that TM is locally trivial, the associated bundle  $V_r(TM)$  is locally trivial with the same triviality open subsets.

In this section, we use a triangulation of the manifold M. In the singular case (section 1.2.2), we will see the usefulness of considering a cellular decomposition, with a similar construction.

We assume that the triangulation (K) of M is sufficiently small so that every simplex  $\sigma$  is included in an open subset U over which the bundle  $V_r(TM)$  is trivial.

Let us assume that one has an r-frame  $v^{(r)}$  defined on the boundary  $\partial \sigma$  of a k-simplex  $\sigma$ . It provides a map

(1) 
$$\partial \sigma \xrightarrow{v^{(r)}} V_r(TM)|_U \cong U \times V_r(\mathbb{C}^m) \xrightarrow{pr_2} V_r(\mathbb{C}^m),$$

where  $pr_2$  is the second projection. One obtains a map

(2) 
$$\xi: \mathbb{S}^{k-1} \cong \partial \sigma \xrightarrow{pr_2 \circ v^{(r)}} V_r(\mathbb{C}^m)$$

that defines an element of the homotopy group  $\pi_{k-1}(V_r(\mathbb{C}^m))$ , denoted by  $[\xi]$ .

By classical homotopy theory, if  $[\xi] = 0$ , then the map  $\partial \sigma \to V_r(\mathbb{C}^m)$  defined on the boundary of the simplex  $\sigma$  can be extended inside  $\sigma$ . In other words, if  $[\xi] = 0$ , then there is no obstruction for the extension of the *r*-frame  $v^{(r)}$ , section of  $V_r(TM)$ , inside  $\sigma$ . This happens in particular when  $\pi_{k-1}(V_r(\mathbb{C}^m)) = 0$ .

The homotopy groups  $\pi_*(V_r(\mathbb{C}^m))$  are computed by Stiefel and by Whitney (see [24]). One has:

(3) 
$$\pi_{k-1}(V_r(\mathbb{C}^m)) = \begin{cases} 0 & \text{for } k < 2m - 2r + 2\\ \mathbb{Z} & \text{for } k = 2m - 2r + 2. \end{cases}$$

Moreover, one has the following results:

**Proposition 1.1.** Let us consider a complex r-frame  $v^{(r)}$  defined on the boundary  $\partial \sigma$  of the k-simplex  $\sigma$ .

- If k < 2(m-r+1), then  $[\xi] = 0$  and one can extend the r-frame  $v^{(r)}$  inside  $\sigma$  without singularity.
- If k = 2(m r + 1), then one can extend the r-frame  $v^{(r)}$  inside  $\sigma$  with an isolated singularity at the barycenter  $\hat{\sigma}$  of the simplex. In that case, one obtains an index  $I(v^{(r)}, \hat{\sigma})$  defined by the class  $[\xi] \in \pi_{k-1}(V_r(\mathbb{C}^m)) \cong \mathbb{Z}$ . The index measures the obstruction to the extension of  $v^{(r)}$ , defined on the boundary  $\partial \sigma$ , inside  $\sigma$ .

The dimension 2p = 2(m - r + 1) is called the *obstruction dimension* for the construction of a complex r-frame tangent to M.

For q = 0, ..., 2m, we denote by  $(K)^{(q)}$  the (q)-skeleton of (K), union of simplices of dimension less than or equal to q.

By Proposition 1.1, it is possible to construct an *r*-frame  $v^{(r)}$  without obstruction on the (2p-1)-skeleton  $(K)^{(2p-1)}$  and with isolated singularities at the barycenters  $\hat{\sigma}$  of the 2*p*-simplices  $\sigma$  with indices  $I(v^{(r)}, \hat{\sigma})$ . The cochain  $c^p : (K)^{(2p)} \to \mathbb{Z}$ defined by  $c^p(\sigma) = I(v^{(r)}, \hat{\sigma})$  is a 2*p*-cocycle of (K) (for instance see [6]). Its class in  $H^{2p}(K;\mathbb{Z}) = H^{2p}(M;\mathbb{Z})$ , denoted by  $c^p(M)$ , is called the *p*-th cohomology Chern class of M.

The Chern class does not depend on the choices of the triangulation and of the r-frame.

1.2. The singular case. In the following,  $X \subset \mathbb{C}^m$  will denote an analytic complex variety of complex dimension n, subset of a compact region.

1.2.1. Stratifications. A stratification (S) of X is a family of closed analytic subsets of X

$$(\mathcal{S}): \qquad X = X^n \supset X^{n-1} \supset X^{n-2} \supset \dots \supset X^1 \supset X^0 \supset \emptyset = X^{-1},$$

where each stratum  $S^{j} = X^{j} - X^{j-1}$  is either empty or a (non necessarily connected) smooth complex manifold of pure (complex) dimension j.

A stratification of X gives rise to a stratification of  $\mathbb{C}^m$  by adding the stratum  $\mathbb{C}^m \setminus X$ .

There are many ways to consider and define a stratification of an analytic variety. In her original work on characteristic classes for singular varieties, Marie-Hélène Schwartz uses Whitney stratifications. In this article, we consider Lipschitz stratifications (*cf.* Section 2). For a complete presentation of stratifications, see for instance [25].

5

1.2.2. Triangulations and cellular decompositions. Let (K) be a triangulation of  $\mathbb{C}^m$  compatible with the stratification of  $\mathbb{C}^m$ . That means that every open simplex of (K) lies in one and only one stratum. The existence of such a triangulation has been proved by Lefschetz [10] in the analytic case.

We consider a barycentric subdivision  $(\hat{K})$  of (K). The barycenter of a simplex  $(\sigma \in K)$  will be denoted by  $\hat{\sigma}$ . Every simplex in  $\hat{K}$  can be written as

$$(\hat{\sigma}_{i_1}, \hat{\sigma}_{i_2}, \dots, \hat{\sigma}_{i_p})$$

where  $\sigma_{i_1} < \sigma_{i_2} < \cdots < \sigma_{i_p}$ . Here the symbol  $\sigma < \sigma'$  means that the simplex  $\sigma$  is a face of  $\sigma'$ .

The dual cell of a simplex  $\sigma$ , denoted by  $d(\sigma)$ , is the set of all (closed) simplices  $\tau$  in  $(\hat{K})$  such that  $\tau \cap \sigma = \{\hat{\sigma}\}$ . That is the set of all simplices in  $(\hat{K})$  on the form  $(\hat{\sigma}, \hat{\sigma}_{i_1}, \ldots, \hat{\sigma}_{i_k})$  with  $\sigma < \sigma_{i_1} < \cdots < \sigma_{i_k}$ .



FIGURE 1. Dual cells: The barycenter of  $\sigma_0 = \{A\}$  is A itself. The dual cell of  $\sigma_0$  is the dark gray cell. The dual cell of  $\sigma_1 = AB$  is composed of the two segments (double lines)  $A'\widehat{\sigma_1}$  and  $\widehat{\sigma_1}B'$ . The dual cell of the triangle  $\sigma_2 = ABC$  is the barycenter of  $\sigma_2 = A'$ . The stratum  $S^j$  is the thick dashed line

The dual cells satisfy the nice properties (see for instance [16]):

**Lemma 1.2.** (i) The dual cell of a k-simplex is a (2m-k)-cell, homeomorphic to the unit ball  $\mathbb{B}^{2m-k} \subset \mathbb{R}^{2m-k}$  and its boundary is homeomorphic to the corresponding sphere  $\mathbb{S}^{2m-k-1}$ .

(ii) Given a triangulation (K) of  $\mathbb{C}^m$  compatible with the stratification of  $\mathbb{C}^m$ , the set of dual cells provides a cellular decomposition of  $\mathbb{C}^m$ , denoted by (D) and called dual cellular decomposition associated to the barycentric subdivision  $(\hat{K})$  of (K).

If the cell  $d(\sigma)$  meets X, the simplex  $\sigma$  of which it is dual lies in X. The unique intersection point  $\hat{\sigma} = d(\sigma) \cap \sigma$  is the barycenter of  $\sigma$  and is also the barycenter of  $d(\sigma)$ . It will be denoted by  $\hat{d}$ .

The following straightforward Lemma plays an important role in the construction of M.-H. Schwartz [22].

**Lemma 1.3.** Let (S) be a stratification of  $\mathbb{C}^m$  and (K) be a triangulation of  $\mathbb{C}^m$  compatible with the stratification. Let (D) be a cellular decomposition of  $\mathbb{C}^m$  dual of (K). Then the dual cells are (topologically) transverse to the strata.

Here, the topological transversality means that if  $S^j$  is a stratum and d is a dual cell, then the (real) codimensions in  $\mathbb{C}^m$  satisfy

 $\operatorname{codim}(d \cap S^j) = \operatorname{codim}(d) + \operatorname{codim}(S^j).$ 

**Remark 1.4.** The dual cells of all simplices  $\sigma$  in the stratum  $S^j$  form a cellular neighbourhood of  $S^j$  (see the grey neighbourhood of  $S^j$  in Figure ??). This fact will be useful in the following.

1.2.3. Obstruction in a stratified space. According to Proposition 1.1, the obstruction for constructing a stratified r-frame tangent to the stratum  $\mathbb{C}^m \setminus X$  lies in dimension 2(m-r+1). The obstruction for constructing an r-frame tangent to a stratum  $S^j$  with complex dimension j lies in (real) dimension 2(j-r+1).

When using a triangulation (K) of  $\mathbb{C}^m$  compatible with a stratification of  $\mathbb{C}^m$ , the obstruction dimension for the construction of a complex *r*-frame tangent to a stratum depends on the considered stratum. It is then clearly impossible to define a cocycle in the same way than in the smooth case, the dimension of the obstruction cocycle depends on the dimension of each stratum.

The first idea of Marie-Hélène Schwartz is to consider in  $\mathbb{C}^m$  a cellular decomposition (D) dual of (K). In that case, as (K) is compatible with the stratification, the cells of (D) are topologically transverse to the strata (Lemma 1.3). When not empty, the intersection of an 2(m - r + 1) cell of (D) with a stratum  $S^j$  of (real) dimension 2j is a cell of (real) dimension 2(j - r + 1). That is exactly the obstruction dimension for the construction of an *r*-frame tangent to  $S^j$ . It is then possible to define a 2p = 2(m - r + 1)-cocycle  $\gamma^p$  (see formula (12)) by attaching to each 2(m - r + 1) cell of (D) meeting X the index of a stratified complex *r*-field at isolated singularities. We will use this first idea in the same way.

In the previous paper ([3, Section 2]), we explicited the second idea of Marie-Hélène Schwartz. To prove the Poincaré-Hopf theorem in the case of real semianalytic varieties we need to use radial vector fields, which are obtained as a sum of parallel and transversal extensions. The same idea is valid for constructing the Schwartz classes. In particular this guarantees that the index of the considered *r*-field at an isolated singular point is the same, whether it is calculated on the stratum or in the ambient space. Then, the cocycle  $\gamma^p$  is well defined. In section 3 we will apply the M.-H. Schwartz' idea in the context of Lipschitz stratifications. The Schwartz class  $c^p(X)$  will be defined as the class of the cocycle  $\gamma^p$ .

# 2. Lipschitz stratifications

Motivated by a question of Sullivan ([23], 1979) (for a complete discussion see [19]), Lipschitz stratifications are defined in 1985 by Tadeusz Mostowski [14]. Mostowski proved existence of stratifications satisfying the so-called Lipschitz *L*-conditions for complex analytic varieties. Some basic properties of Lipschitz stratifications are improved and clarified by Adam Parusiński [18].

The first ingredient for the definition of Lipschitz stratifications is the notion of chains (see [14, 15, 19]).

**Definition 2.1** (Definition of chains). Let us fix a (real) constant K > 1. A chain starting at a point  $q_{j_1}$  in an  $j_1$ -dimensional stratum  $S^{j_1}$  is a sequence of points  $\{q_{j_1}, \ldots, q_{j_t}, \ldots, q_{j_\ell}\}$  where  $q_{j_k} \in S^{j_k}$ , such that

$$X^{j_1} \supset X^{j_2} \supset \cdots \supset X^{j_t} \supset \cdots \supset X^{j_k} \supset \cdots \supset X^{j_\ell}$$

(with complex dimensions  $j_1 > j_2 > \cdots > j_t > \cdots > j_\ell$ ) and where  $j_t$  (depending on the constant K) is the greatest integer for which

(4) 
$$d(q_{j_1}, X^{j_k}) \ge 2K^2 d(q_{j_1}, X^{j_t})$$
 for all  $k$  such that  $j_t > j_k \ge j_\ell$   
and

(5) 
$$|q_{j_1} - q_{j_t}| \leq K d(q_{j_1}, X^{j_t})$$

**Remark 2.2.** The condition (4) can be written

$$d(q_{j_1}, X^{j_t}) \leq \frac{1}{2K^2} d(q_{j_1}, X^{j_k})$$

and means that when  $q_{j_1}$  is going far away from  $X^{j_t}$  to a stratum  $S^{j_k}$  of lower dimension  $j_k < j_t$ , the distances

$$d(q_{i_1}, X^{j_t})$$
 and  $d(q_{i_1}, X^{j_k})$ 

are controlled by the inequality (4).

**Remark 2.3.** The condition (5) says that in the same stratum  $S^{j_t}$ , the distances from the starting point  $q_{j_1}$  to the point  $q_{j_t}$  and to the stratum  $S^{j_t}$  are controlled by the inequality (5).

The following example (here in  $\mathbb{R}^3$ , due to limitation of dimension in figures) illustrates the definition.



FIGURE 2. Example of chain:  $j_1 = 2, j_t = j_2 = 1, j_3 = 0$ , the chain is:  $(q_{j_1}, q_{j_2}, q_{j_3})$ .

Before providing the definition of Lipschitz stratifications, we fix some more notations: For  $q \in S^{j}$ , let

$$P_q: \mathbb{C}^m \to T_q S^j \subset T_q \mathbb{C}^m$$

be the orthogonal projection onto the tangent space to the stratum  $S^j$  at the point q and let  $P_q^{\perp} = Id - P_q$  be the orthogonal projection onto the normal space  $T_q^{\perp}S^j$ .

Identifying  $T_q \mathbb{C}^m$  by  $\mathbb{C}^m$ , then  $P_q$  and  $P_q^{\perp}$  are linear maps from  $\mathbb{C}^m$  to  $\mathbb{C}^m$ . The norm of the linear map  $P_q$  can be defined as any norm of endomorphims of  $\mathbb{C}^m$ :

$$||P_q|| := \sup_{||x||=1} |P_q(x)|.$$

The same stands for  $P_q^{\perp}$ .

**Remark 2.4.** An important remark (philosophy) concerning exponents and multiplicative constants is that exponents are important and interesting while multiplicative constants appearing in plenty of inequalities are not important. We shall use occasionally the following notations:

$$f \lesssim g \iff$$
 for some constant  $C > 0, \ f \leqslant Cg,$   
 $f \sim g \iff |f| \lesssim |g| \lesssim |f|.$ 

**Definition 2.5** (Lipschitz stratification). The stratification (S) is said a Lipschitz stratification if there exists a constant C > 0 (called Lipschitz constant) such that a) for every chain  $\{q_{j_1}, q_{j_2}, \ldots, q_{j_\ell}\}$  (Definition 2.1), we have

(6) 
$$\|P_{q_{j_1}}^{\perp}P_{q_{j_2}}\cdots P_{q_{j_k}}\| \leqslant C \frac{|q_{j_1}-q_{j_2}|}{d(q_{j_1}, X^{j_k-1})},$$

for all  $1 < k \leq \ell$ .

b) If, further,  $q_{j_1}$  and  $q'_{j_1}$  are points in  $S^{j_1}$  such that

$$q'_{j_1} - q_{j_1} \le \frac{1}{2K} d(q_{j_1}, X^{j_1 - 1})$$

(here K is the constant in Definition 2.1), then

(7) 
$$\|(P_{q'_{j_1}} - P_{q_{j_1}})P_{q_{j_2}} \cdots P_{q_{j_k}}\| \leq C \frac{|q'_{j_1} - q_{j_1}|}{d(q_{j_1}, X^{j_k-1})},$$

for all  $1 \leq k \leq \ell$ . In particular,

(8) 
$$\|P_{q'_{j_1}} - P_{q_{j_1}}\| \le C \frac{|q'_{j_1} - q_{j_1}|}{d(q_{j_1}, X^{j_1 - 1})}$$

**Remark 2.6.** The inequality (8) says that if  $q_{j_1}$  and  $q'_{j_1}$  are sufficiently close points in the stratum  $S^{j_1}$ , then the norm  $||P_{q'_{j_1}} - P_{q_{j_1}}||$  is roughly given geometrically by the angle between the tangent spaces  $T_{q_{j_1}}S^{j_1}$  and  $T_{q'_{j_1}}S^{j_1}$ . More precisely, if

$$\theta = \text{angle}(T_{q_{j_1}}S^{j_1}, T_{q'_{j_1}}S^{j_1})$$

then one has  $||P_{q'_{j_1}} - P_{q_{j_1}}|| \approx \sin(\theta)$ . This angle is controlled by the inequality (8). Similarly one has  $||P_{q'_{j_1}}^{\perp}P_{q_{j_1}}|| = \sin(\theta)$  (see also [3, Figure 5]).

The inequalities (6) and (7) are more general in the sense that the angles of all the tangent spaces at points appearing in the chain are taken into account.

In the following, one provides the relation between the different stratifications: Lipschitz, Kuo-Verdier and Whitney. In fact one explains (or recalls the proof of) the relation

Lipschitz (L) 
$$\Rightarrow$$
 Kuo-Verdier (w)  $\Rightarrow$  Whitney (b)  $\Rightarrow$  Whitney (a).

9

**Remark 2.7.** a) The Lipschitz conditions are stronger than Whitney conditions [27]. First, let us remark that for a chain of two points  $\{q_j, q_k\}$ , then the norm  $\|P_{q_j}^{\perp}P_{q_k}\|$  in the inequality (6) is geometrically given by  $\sin(\theta)$  where  $\theta$  is the angle between the two tangent spaces  $T_{q_j}S^j$  and  $T_{q_k}S^k$ .

Now, recall that the Whitney condition (a) for a pair of adjacent strata  $S^j$  and  $S^k$  with  $S^k \subset \overline{S^j}$ , says that the angle between the tangent spaces  $T_y S^k$  and  $T_x S^j$  tends to zero when  $x \in S^j$  tends to  $y \in S^k$  (see [25, 27]).

The Lipschitz conditions, showed in the inequality (6), are stronger than Whitney condition (a) in the sense that this condition is satisfied for all pairs of strata involved in the considered chain.

b) The Lipschitz conditions are stronger than Kuo-Verdier conditions (w) which are themselves stronger than Whitney condition (b) [9, 26, 25].

The Kuo-Verdier conditions (w) says that, in a neighbourhood sufficiently small of the stratum  $S^k$  (in  $\mathbb{C}^m$ ), the angle  $\theta$  between the tangent spaces  $T_x S^j$  and  $T_y S^k$ is estimated by an inequality

$$\sin(\theta) \leqslant G|x-y|,$$

for some constant G.

(9)

The Whitney condition (b) is verified for a pair of adjacent strata  $S^j$  and  $S^k$ with  $S^k \subset \overline{S^j}$ , if for each sequence  $\{x_i\}$  of points in  $S^j$  and each sequence  $\{y_i\}$  of points in  $S^k$ , both converging to the same point y in  $S^k$ , such that the sequence of secant lines  $(x_i, y_i)$  converges to a line  $\ell$  and the sequence of tangent spaces  $T_x S^j$ converges to a space T (in a suitable Grassmannian space) as i tends to infinity, then  $\ell$  is contained in T.

Now apply the inequality (6) for two adjacent strata  $(S^j, S^k)$ . We have

$$\sin(\theta) \leqslant C \frac{|x-y|}{d(x, X^{j-1})}.$$

In formula (9), put

$$G = \frac{C}{\inf_{x \in \mathcal{U}} \{ d(x, X^{j-1}) \}}$$

where  $\mathcal{U}$  is a sufficiently small neighbourhood of  $S^k$  (in  $\mathbb{C}^m$ ). Then the Kuo-Verdier conditions (w) are satisfied.

We emphasize again that the Lipschitz conditions are stronger than the Whitney condition (b) in the sense that this condition is satisfied for all pairs of strata appearing in the chain.

#### 3. Lipschitz vector fields

In [21] Marie-Hélène Schwartz shows that considering stratified vector fields is not sufficient to prove the Poincaré-Hopf theorem for singular stratified varieties. She gives counterexamples for which the index of the considered vector field at an isolated singular point calculated on the stratum is not the same when calculated in the ambient space (see also [3, §2.1]). She shows that the theorem can be proved when considering radial extensions, obtained as the sum of two extensions: parallel and transversal. She constructs these extensions in the framework of Whitney stratifications, that is very technical and delicate. In [22], Marie-Hélène Schwartz uses the same construction to define Chern classes of stratified varieties.

In [3], we show how the Lipschitz stratifications framework allows us to obtain a simpler proof of the Poincaré-Hopf theorem, in particular concerning the construction of parallel and transversal extensions. This later is named "basic normal" in [3, §4.2], we will keep the terminology. The aim of this section is to show that the construction of these extensions in the Lipschitz framework allows us to simplify the definition of characteristic classes of complex singular analytic varieties.

The main advantage of using Lipschitz stratifications is that, instead of constructing the parallel and transverse extensions by pairs of strata, the construction is done all strata at once, taking into account the distance to the strata to determine the angle of the extension. Moreover, the continuity of constructed vector fields (resp. *r*-frames) comes directly from Lipschitz properties without using any homotopy.

3.1. The parallel Lipschitz extension. Lipschitz stratifications have (among others) the following property (see Mostowski (1985) [14, Prop. 2.1]):

**Proposition 3.1.** Let  $\{X^j\}_{j=1}^n$  be a Lipschitz stratification of X and let v be a Lipschitz stratified vector field on  $X^j$ . Then v can be extended to a Lipschitz stratified vector field on  $S^{j+1}$ .

A Lipschitz vector field v defined on  $X^j$  and tangent to all strata of dimension less than (or equal to) j can be extended to a Lipschitz vector field  $\hat{v}$  defined on  $\mathbb{C}^m$  with the same Lipschitz constant (see Kirszbraun Theorem 1934, for instance in [7]). Mostowski defines the parallel extension of v by

(10) 
$$\tilde{v}(x) = \begin{cases} \hat{v}(x) & \text{if } x \in X^j \\ P_x \hat{v}(x) & \text{if } x \in S^{j+1}. \end{cases}$$

Then  $\tilde{v}$  is a stratified Lipschitz vector field defined on  $S^{j+1}$  (no matter which extension  $\hat{v}$  is chosen).



FIGURE 3. The parallel Lipschitz extension.

The following results are direct consequences of the construction and of the definition of Lipschitz contidions (angles between tangent spaces to the strata are small).

**Corollary 3.2.** If  $v_1$  and  $v_2$  are  $\mathbb{C}$ -linearly independent on a stratum  $S^j$ , then their Lipschitz parallel extensions are  $\mathbb{C}$ -linearly independent in a neighbourhood of  $S^j$ .

In a more general way, one has:

**Corollary 3.3.** If  $v^{(r)}$  is an r-frame in a stratum  $S^j$ , then its Lipschitz parallel extension is an r-frame in a neighbourhood of  $S^j$ .

3.2. The basic normal Lipschitz vector field. The M.-H. Schwartz' idea is given a field of vectors tangent to a stratum and having an isolated singular point, how to extend it in a neighbourhood of the singular point into a stratified vector field, with the same isolated singular point and the same index (calculated in the ambient space) at that point as the initial vector field.

The previous paragraph and the corollaries 3.2 and 3.3 show how to extend an *r*-field tangent to a stratum into a stratified *r*-field in a neighbourhood of this stratum. This technique provides an *r*-field comprising a set of non-isolated singular points, namely a disk, slice of the tubular neighbourhood of the stratum.

In order to meet the conditions of the extension, the idea is to add to the preceding parallel *r*-field a "transversal" vector field "tangent" to the slice and having an isolated singular point of index +1 in the center of the disk. This vector field is in a sense orthogonal to the stratum, it is constructed in the real setting and named by "basic normal" vector field in [3, §4.2].

The following Lemmas have been proved in the real case of semi-analytic sets in [3]. The proof of the complex analytic case is similar.

Let  $X \subset \mathbb{C}^m$  be a smooth analytic subset of some compact region, of dimension n and let  $X' = \overline{X} \setminus X$ . For  $q \in \mathbb{C}^m$ , let us denote by  $d_X(q) = d(q, X)$  the distance from q to X.

**Lemma 3.4.** [3, Lemma 4.1] There exists an integer  $K \in \mathbb{N}$  and an  $\varepsilon > 0$  such that for

$$U = \{ x \in \mathbb{C}^m : d_X(x) < \varepsilon d_{X'}^K(x) \},\$$

then for very  $x \in U$  there exists exactly one point  $\pi(x) \in X$  for which  $d_X(x) = |x - \pi(x)|$ . The mapping  $\pi$  is smooth in U.

Define the basic normal vector field  $r_X$  by  $r_X(x) = x - \pi(x)$  after a translation to x.

**Lemma 3.5.** [3, Lemma 4.3] The basic normal vector field  $r_X$  is smooth, Lipschitz in U and satisfies  $||r_X(x)|| = d_X(x)$ . It can be extended to a Lipschitz vector field on  $\mathbb{C}^m$ , again denoted by  $r_X$ , so that  $|r_X| \leq d_X$  and  $r_X = 0$  outside of  $\widetilde{U} = \{x : d_X(x) < 2\varepsilon d_{X'}^K(x)\}.$ 

In the following, we will apply these two Lemmas 3.4 and 3.5 to the situation of the stratum  $S^{j}$  and will denote by  $U_{j}$  and  $r_{S^{j}} = r_{j}$  respectively the neighbourhood and the basic normal vector field corresponding.

3.3. "Champs sortants" in Lipschitz stratifications. Let  $S = \{X^j\}$  be a Lipschitz stratification of the analytic variety X, subset of some compact region in  $\mathbb{C}^m$ .

The notion of "champ sortant" (literally "outgoing field") has been introduced by Marie-Hélène Schwartz under the terminology "transversal vector field" in her proof



FIGURE 4. The neighbourhood U.

of the Poincaré-Hopf theorem, using Whitney stratification. In the previous paper [3, §4.2] we introduce the notion of "champ sortant" in the framework of Lipschitz stratifications of real varieties. This construction applies also in the complex setting. Here we give a summary of this construction with the important elements.

Let  $q \in S^j$ . Consider the space  $T_q^{\perp}S^j$  as a subspace of  $U_j$  (it is  $\pi^{-1}(q)$  in the previous terminology). If  $x \in T_q^{\perp}S^j$  and  $z \in T_x \mathbb{C}^m$ , let  $\omega_j$  be the orthogonal projection of z onto  $T_x(T_q^{\perp}S^j)$ .

**Definition 3.6.** (see [3, Definition 5.1] in the real setting). Let  $\lambda$  be a constant,  $0 < \lambda < 1$ . A  $\lambda$ -"champ sortant" is a Lipschitz stratified vector field w on  $\mathbb{C}^m$  such that, for every j, in some neighbourhood  $U'_j \subset U_j$  of  $S^j$ ,

$$\|\omega_j(w - \psi_j r_j)\| \leq \lambda \|\psi_j r_j\| = \lambda \|\omega_j(\psi_j r_j)\|$$

for some (scalar valued) functions  $\psi_j$  taking values in some interval  $[a_1, a_2]$  with  $0 < a_1 < a_2 < \infty$ ; and so

$$\not\prec(w,r_j) < \lambda \quad in \ U'_j.$$

Actually, the function  $\psi_j$  will be constant.

For each point  $q \in X$  and each stratum  $X^j$ , let us denote by  $d_j(q) = d_{X^j}(q)$  the distance from q to  $X^j$ , and, for the points q, q'

$$d_j(q,q') = \min(d_j(q), d_j(q')).$$

Consider a pair of strata  $(S^k, S^j)$  such that  $S^j \subset \overline{S^k} = X^k$ . By [3, Lemma 1.1] which applies also in the complex setting, for  $q \in S^j$ ,  $q' \in S^k$ , there exists numbers C > 0,  $\beta > 0$  such that

$$\measuredangle(q'-q, T_q S^j) \lesssim C \frac{|q'-q|^\beta}{d_{j-1}(q, q')}$$

It follows that if  $|q'-q| \leq \min(d_{j-1}(q))^K, \frac{1}{2}d_{j-1}(q))$ , then

$$\measuredangle(q'-q, T_q S^j) \leqslant C \frac{|q'-q|^{\beta}}{d_{j-1}(q)} \lesssim d_{j-1}(q)^{jK}$$

It follows that if K appearing in the definition of  $U_j$  satisfies the estimate  $K \ge 2/\beta$ , then in our stratification, we have the above estimate in  $U_j$ .

Let us fix a stratum  $S^j$  and take K so big that in  $U_j^* = \{d_j < 2d_{j-1}^K\}$  we have the basic normal vector field  $r_{S^j} = r_j$  with all properties of Lemma 3.5 and the above form of (b) holds in  $U_j^*$ . Let  $U_j = \{d_j < d_{j-1}^K\}$ , and  $\varepsilon = jK$ .

The proof of the following Lemma ([3, Lemma 5.2] in the real setting) applies in the complex setting:

**Lemma 3.7.** There exists  $\varepsilon > 0$  and a Lipschitz S-stratified vector field  $w_j$  such that

$$\|w_j - r_j\| \leq d_j^{\varepsilon+1} \text{ in } U_j,$$
  
$$\|w_j\| \leq \|r_j\| = d_j \text{ in } U_j^*,$$
  
$$w_j = 0 \text{ outside of } U_j^*.$$

**Remark 3.8.** Of course, we have  $||w_j|| \leq \min(d_j, d_{j-1}^K)$ .

**Proposition 3.9.** ([3, Proposition 5.4] in the real setting). Let  $w_i$ 's be as in Lemma 3.7 and  $\lambda$  any number,  $0 < \lambda < 1$ . Then, if  $0 < c_0 < c_1 < \cdots < c_n$  in a sufficiently rapidly increasing sequence of numbers (i.e.  $\frac{c_{i+1}}{c_i}$  is sufficiently big, for all i), then the vector field  $w = \sum c_i w_i$  is a S-stratified Lipschitz vector field  $\lambda$ -"champ sortant".

*Proof.* Let us fix a stratum  $S^j$  and consider  $w_j$  in the neighbourhood  $U_j$  of  $S^j$ . Let i < j. Since  $w_i$  is Lipschitz and stratified, we have, in  $U_j$ ,

$$\|\omega_j(w_i)\| \leq d_j$$

because  $\omega_j(w_i) = 0$  on  $S^j$ . Since

$$\|\omega_j(w_j)\| \sim \|w_j\| \sim d_j,$$

we have, for a sufficiently big constant A,

$$\|\omega_j(w_i)\| < A\|\omega_j(w_j)\|.$$

Let i > j. Then  $||w_i|| \leq d_{i-1}^K \leq d_j^K$  (here K is as in Lemma 3.7; we may assume that  $K \geq 2$ ). It follows that, no matter how big is a constant A, in a suitable neighbourhood of  $S^j$ , we have

$$\|\omega_j(w_j)\| > A \|\omega_j(w_i)\|.$$

A bound below for  $\omega_j(w_j)$  is given by

$$\|\omega_j(w_j)\| \sim \|w_j\| \sim d_j.$$

Thus, if  $c_i$  increases sufficiently rapidly,

$$||w - c_j w_j|| = ||\sum_{i \neq j} c_i w_i|| < \frac{\lambda}{2} ||c_j w_j||,$$

and, by Lemma 3.7

$$\|w - c_j r_j\| < \lambda \|c_j r_j\|$$

in some neighbourhood of  $S^j$ .

The "champ sortant" is the stratified Lipschitz vector field w.

3.4. **Applications.** According to section 1.2.3, the aim of the construction is to attach to each 2p = 2(m - r + 1)-cell of a cellular dual decomposition, the index of a complex *r*-field with isolated singularities. The *r*-field will be constructed by using the parallel Lipschitz extension and the "champ sortant".

The parallel Lipschitz extension and the Lipschitz "champ sortant" are well defined within a neighbourhood  $\mathcal{U}_j$  for each stratum  $S^j$ . That means that we have to refine the triangulation (K) and consequently also the dual cellular decomposition so that each dual cell meeting the stratum  $S^j$  is included in  $\mathcal{U}_j$ , this for all strata.

Now, let d be a 2p-cell dual of a (K)-simplex  $\sigma^{2(r-1)} \subset S^j$ , where  $S^j$  is a stratum of (complex) dimension  $j \ge (r-1)$ .

The obstruction dimension to construct an r-frame tangent in  $S^j$  is  $2p_j = 2(j - r + 1)$ , that is the dimension of  $d \cap S^j$  (note that d is topologically transverse to  $S^j$  - see Lemma 1.3). One can construct a Lipschitz r-frame  $v^{(r)} = (v_1, \ldots, v_{r-1}, v_r)$  on  $d \cap S^j$  with an isolated singularity due to cancellation of the last vector  $v_r$  at the barycenter  $\hat{d}$  of the cell d. Let us recall that  $\hat{d}$  is also barycenter of the simplex  $\sigma^{2(r-1)}$ .

Let us denote by  $\overline{v}^{(r)}$  the parallel Lipschitz extension of  $v^{(r)}$  (defined in section 3.1) to the last vector of which we add the "champ sortant" w (defined in section 3.3), that means

$$\overline{v}^{(r)} = (\widetilde{v}_1, \dots, \widetilde{v}_{r-1}, \widetilde{v}_r + w).$$

That is a Lipschitz stratified r-field well defined on the 2p-cell d with an isolated singularity at the barycenter  $\hat{d}$  of d.

**Proposition 3.10.** (a) Let  $v^{(r)} = (v_1, \ldots, v_{r-1}, v_r)$  be a Lipschitz r-field tangent to  $S^j$ , with j > r-1 and with an isolated singularity at  $\hat{d} \in S^j$  due to cancellation of the last vector  $v_r$ . Let us denote by  $\overline{v}^{(r)}$  the parallel extension of  $v^{(r)}$  to the last vector of which we add the Lipschitz champ sortant:

$$\overline{v}^{(r)} = (\widetilde{v}_1, \dots, \widetilde{v}_{r-1}, \widetilde{v}_r + w).$$

The point  $\hat{d}$  is also an isolated singularity of  $\overline{v}^{(r)}$  and one has equality of indices

$$I(v^{(r)}, \hat{d}; S^j) = I(\overline{v}^{(r)}, \hat{d}; \mathbb{C}^m)$$

where  $I(v^{(r)}, \hat{d}; S^j)$  is the index of the r-field  $v^{(r)}$  at its singular point  $\hat{d}$ , calculated on the stratum  $S^j$ , and  $I(\overline{v}^{(r)}, \hat{d}; \mathbb{C}^m)$  is the index of its extension, the r-field  $\overline{v}^{(r)}$ , in the ambient space, at its same singular point  $\hat{d}$ ,

(b) If j = r - 1, then one can consider only the (r - 1)-field  $(v_1, \ldots, v_{r-1})$  on  $S^j$ . Then the r-field  $(\tilde{v}_1, \ldots, \tilde{v}_{r-1}, w)$  is a Lipschitz stratified r-field well defined on d and its index  $I(\overline{v}^{(r)}, \hat{d}; \mathbb{C}^m)$  is equal to +1.

3.5. Radial extension of r-frames. In the same way as in ([4], [20], [22]), we prove the following theorem, existence of radial extension of frames, in the Lipschitz framework:

**Theorem 3.11.** Let X be an analytic variety, subset of some compact region in the complex space  $\mathbb{C}^m$  and (S) a Lipschitz stratification of  $\mathbb{C}^m$  which is composed of a stratification of X to which we add the stratum ( $\mathbb{C}^m \setminus X$ ). Let (K) be a triangulation of  $\mathbb{C}^m$  compatible with the stratification and denote by (D) a cellular decomposition of  $\mathbb{C}^m$  dual of (K). Let  $\mathcal{D}$  be the cellular neighbourhood of X in  $\mathbb{C}^m$ , union of cells

which meet X. We can construct, on the 2*p*-skeleton  $(D)^{2p}$ , a Lipschitz stratified r-field  $v^{(r)}$  which is called radial field, whose singularities satisfy the following properties:

(i)  $v^{(r)}$  has only isolated singular points, which are zeroes of the last vector field  $v_r$  and which are barycenters  $\hat{d}$  of 2p-cells. On  $(D)^{2p-1}$ , the r-field  $v^{(r)}$  has no singular point and on  $(D)^{2p}$  the (r-1)-field  $v^{(r-1)}$  has no singular point.

(ii) Let  $\hat{d} \in S^j \cap (D)^{2p}$  be a singular point of  $v^{(r)}$  in the 2*j*-dimensional stratum  $S^j$ . If j > r - 1, the index of  $v^{(r)}$  at  $\hat{d}$ , computed in the ambient space is the same as the index of the restriction of  $v^{(r)}$  to  $S^j \cap (D)^{2p}$  considered as an *r*-field tangent to  $S^j$ . If j = r - 1, then  $I(v^{(r)}, \hat{d}) = +1$ .

(iii) Inside a 2p-cell d which meets several strata, the only singularities of  $v^{(r)}$  are inside the lowest dimensional one (in fact located at the barycenter of d). (iv) The r-field  $v^{(r)}$  has no singularity on  $\partial \mathcal{D} \cap (D)^{2p}$ .

Basically, the proof follows the same process as the proof of the Poincaré-Hopf theorem [3, §6]. The proof of the Poincaré-Hopf Theorem is a proof by induction starting with a radial vector field pointing outwards balls centered at points corresponding to the 0-dimensional strata. Here the role of the strata of dimension 0 is played by the strata  $S^j$  such that the intersection of a cell d of dimension 2p = 2(m-r+1) with  $S^j$  is the barycenter  $\hat{d}$  of d. We have therefore 2(j-r+1) = 0, i.e. j = (r-1).

*Proof.* Let  $S^j$  be a stratum of (real) dimension 2j = 2(r-1). The intersection of a cell d of dimension 2p = 2(m-r+1) with  $S^j$  is the barycenter  $\hat{d}$  of d. At the points  $\hat{d}$ , we give ourselves a complex Lipschitz (r-1)-field tangent to  $S^j$ . It can be extended along d by Lipschitz extension, with the condition that d is sufficiently small to be included in  $\mathcal{U}_j$ .



FIGURE 5. The starting point of induction (real picture of a complex situation). The cell d is dual of the simplex  $\sigma = ABC$ .

More precisely, the starting point of the induction is as follows:

If  $S^{r-1}$  is a stratum whose real dimension is 2r - 2 = 2(m - p), the intersection of a 2*p*-cell with  $S^{r-1}$  is a point: the barycenter  $\hat{d}$  of d, which is also the barycenter of the simplex  $\sigma$  (in  $S^{r-1}$ ) of which d is the dual cell (see Figure 5).

The obstruction dimension for the construction of an *r*-frame tangent to  $S^{r-1}$  is zero. One takes any Lipschitz (r-1)-frame  $v^{(r-1)}$  tangent to  $S^{r-1}$  at the barycenter  $\hat{d}$  and the last vector  $v_r$  is taken as zero at these points.

We construct the Lipschitz parallel extension (see section 3.1) of the (r-1)-frame  $v^{(r-1)}$  in the cellular tubular neighbourhood  $\mathcal{U}_{r-1}$  as an (r-1)-frame denoted by  $\tilde{v}^{(r-1)}$ . The extension, together with the Lipschitz champ sortant w (see section 3.2) provide an r-field  $\tilde{v}^{(r)} = (\tilde{v}^{(r-1)}, w)$  which has an isolated singularity at  $\hat{d}$ . According to Proposition 3.10 (b), one has  $I(\tilde{v}^{(r)}, \hat{d}) = +1$ .

Now, the induction process goes as follows: Let us suppose k > r - 1 and the construction already performed on all strata  $S^j$  whose (real) dimension is less than 2k. That means that the construction provides an *r*-field on the 2*p*-skeleton of a cellular neighbourhood  $\mathcal{D}^j$  of  $S^j$  for all strata  $S^j$  with dimension j < k. As in Proposition 3.10, the *r*-field on  $S^j$  is denoted by  $v^{(r)} = (v^{(r-1)}, v_r)$  and its Lipschitz extension by  $\overline{v}^{(r)}$ . If a 2*p* cell is transverse to a stratum  $S^j$ , then the *r*-field has an isolated singularity at the barycenter  $\hat{d}$ , due to cancellation of the last vector  $v_r$ . The indices satisfy then

(11) 
$$I(v^{(r)}, \hat{d}; S^j) = I(\overline{v}^{(r)}, \hat{d}; \mathbb{C}^m)$$

If the 2*p*-cell *d* does not meet a stratum and lies in the boundary of the cellular neighbourhood  $\mathcal{D}^{j}$  of a stratum  $S^{j}$ , then the basic normal vector field *w* does not vanish on *d* and the *r*-field  $\tilde{v}^{(r)}$  has no singularity on *d*. One remarks that such cells are dual of simplices in (K) which have a face in  $S^{j}$  but are not included in  $S^{j}$ .

The boundary of  $S^k$  is union of strata  $S^j$ , each of them equipped with a cellular neighbourhood  $\mathcal{D}^j$  and an *r*-field satisfying (11) at each singular point.

The *r*-frame is constructed on the 2p-skeleton of a cellular neighbourhood of the boundary of  $S^k$ .

We extend the (r-1)-frame  $\tilde{v}^{(r-1)}$  inside  $S^k$ , more precisely on the  $2p_k$ -skeleton of the intersection  $(D)^{(2p)} \cap S^k$ , with  $2p_k = 2(k-r+1)$ . This is performed without singularity.

Using the Lipschitz extension (3.1), one extends the last vector  $v_r$  with singularities at the barycenters  $\hat{d}$  of cells d in  $(D)^{(2p)} \cap S^k$ . One obtains, on the stratum  $S^k$  an *r*-frame with isolated singularities at the barycenters  $\hat{d}$ , which are zeroes of the last vector  $v_r$ . One has (Proposition 3.10 (a))

$$I(\widetilde{v}^{(r)}, \widehat{d}; \mathbb{C}^m) = I(v^{(r)}, \widehat{d}; S^k).$$

In summary, an r-frame already known on a neighbourhood of the boundary of a stratum is extended with isolated singularities inside (a suitable skeleton of) the stratum and then extended with property (a) of the Proposition 3.10 to a cellular neighbourhood of this stratum.

We denote by  $\mathcal{D}$  the tubular neighbourhood of X in  $\mathbb{C}^m$  consisting of the (D)cells which meet X. By subdividing if necessary, we can assume that  $\mathcal{D}$  is contained
in the union of the Lipschitz neighbourhood  $\mathcal{U}^j$ .

## 4. Obstruction cocycles and classes

We define a 2*p*-dimensional (*D*)-cochain in  $C^{2p}(\mathcal{D}, \partial \mathcal{D})$  by:

(12) 
$$\gamma^p: d \to I(\widetilde{v}^{(r)}, d).$$

The cochain  $\gamma^p$  actually is a cocycle whose class  $c^p(X)$  lies in

$$H^{2p}(\mathcal{D},\partial\mathcal{D}) \cong H^{2p}(\mathcal{D},\mathcal{D}\backslash X) \cong H^{2p}(\mathbb{C}^m,\mathbb{C}^m\backslash X),$$

where the first isomorphism is given by retraction along the rays of  $\mathcal{D}$  and the second by excision (by  $\mathbb{C}^m \setminus \mathcal{D}$ ). The class  $c^p(X)$  coincides with the Schwartz class, because on the one hand, the Lipschitz frame is a particular case of the frames defined by M.-H. Schwartz and we know that the Schwartz class does not depend on the choice of frame in her framework. On the other hand, a Lipschitz stratification is particularly a Whitney stratification.

**Definition 4.1.** The p-th Schwartz class of X is the class

$$c^p(X) \in H^{2p}(\mathbb{C}^m, \mathbb{C}^m \setminus X).$$

Marie-Hélène Schwartz proved "by hand" that the resulting class is independent of the stratification, of the triangulation (compatible with the stratification) and of the choice of the radial r-frame. Here, within the Lipschitz framework, the resulting class is independent of the Lipschitz stratification (including the Lipschitz constant C), of the triangulation and of the choice of the Lipschitz extensions of frames. The proof of independence is now facilitated as we know [4] that by Alexander duality isomorphism

$$H^{2(m-r+1)}(\mathbb{C}^m,\mathbb{C}^m\backslash X)\to H_{2(r-1)}(X), \qquad d\to\sigma$$

image of the Schwartz class is the MacPherson class [13] of the constructible function  $\mathbf{1}_X$  and that the MacPherson class does not use neither triangulation, nor *r*-frames. It is proved independent of the used stratification. Nowadays, the classes are called Schwartz-MacPherson classes.

#### References

- [1] J.-P. Brasselet, An introduction to Characteristic Classes, Publications IMPA, 33° Colóquio Brasileiro de Matemática.
- [2] J.-P. Brasselet and Thủy Nguyễn T.B, O Teorema de Poincaré-Hopf, C.Q.D.- Revista Electrônica Paulista de Matemática 16 (2019), p. 134-162 (in Portuguese).
- [3] J.-P. Brasselet, T. Mostowski and Thủy Nguyễn T.B, Poincaré-Hopf Theorem for singular analytic varieties. The Marie-Hélène Schwartz' ideas in the Lipschitz framework. Banach Center Publications, Volume 128, Warszawa, 2024, 23–43.
- [4] J.P. Brasselet and M.-H. Schwartz: Sur les classes de Chern d'un ensemble analytique complexe, Astérisque 82-83 (1981), 93-147.
- [5] S.S. Chern, Characteristic classes of hermitian manifolds, Ann. Math. 47 (1946), 85-121.
- [6] J. F. Davis and P. Kirk, Lecture notes in algebraic topology, Vol. 35. Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, (2001).
- [7] H. Federer, Geometric measure theory. Classics in Mathematics, Springer 1996.
- [8] A. Grothendieck, Récoltes et Semailles, Réflexions et témoignage sur un passé de mathématicien, 1986, Gallimard, 2023, available in english translation at https://web.ma.utexas.edu/users/slaoui/notes/recoltes\_et\_semailles.pdf
- [9] D. Juniati, L. Noirel and D. Trotman, Whitney, Kuo-Verdier and Lipschitz stratifications for the surfaces  $y^a = z^b x^c + x^d$ . Topology and Applications, 234 (2018), 335-347.
- [10] S. Lefschetz, Topology, Amer. Math. Soc, Coll. Publications, New York, 1930.
- [11] S. Lojasiewicz, Triangulation of semi-analytic sets, Ann. Scuola Norm. Sup. di Pisa, Ser. 3, 18-4 (1964), 449-474.
- [12] S. Lojasiewicz, On semi-analytic and subanalytic geometry, Panoramas of Mathematics, Banach Center Publications, Volume 34, Institute of Mathematics, Polish Academy of Sciences, Warszawa 1995, pp 89 – 104.

- 18 JEAN-PAUL BRASSELET, TADEUSZ MOSTOWSKI, AND THỦY NGUYỄN THỊ BÍCH
- [13] R. D. MacPherson, Chern classes for singular algebraic varieties. Ann. of Math. (2) 100, (1974), 423–432.
- [14] T. Mostowski, *Lipschitz Equisingularity*, Dissertationes Math. (Rozprawy Mat.) 243, PWN, Warszawa, 1985.
- [15] T. Mostowski, Tangent Cones and Lipschitz Stratifications, Singularities, Banach Center Publications (S. Lojasiewicz, ed.), Vol. XX, PWN, Warszawa, 1988, pp. 303-322.
- [16] J. R. Munkres, *Elements of algebraic topology*. Addison-Wesley Publishing Company, Menlo Park, CA, 1984.
- [17] A. Parusiński, Lipschitz Properties of Semi-analytic Sets, Ann. Inst. Fourier, Grenoble, Vol. 38, 1988, no. 4, pp. 189-213.
- [18] A. Parusiński, *Lipschitz stratifications*, in: Global Analysis in Modern Mathematics (Orono, 1991; Waltham, 1992), K. Uhlenbeck (ed.), Publish or Perish, Houston, 1993, 73–89.
- [19] A. Parusiński, Lipschitz stratification of subanalytic sets, Ann. Sci. École Norm. Sup. (4) 27 (1994), no. 6, 661–696.
- [20] M.H.Schwartz, Classes caractéristiques définies par une stratification d'une variété analytique complexe, C. R. Acad. Sci. Paris Sér. I Math. 260, (1965), 3262-3264 et 3535–3537.
- [21] M.-H. Schwartz: Champs radiaux sur une stratification analytique, Travaux en cours, 39 (1997), Hermann, Paris.
- [22] M.-H. Schwartz, Classes de Chern des ensembles analytiques. Actualités mathématiques, Hermann, Paris, 2000.
- [23] L. Siebenmann and D. Sullivan, On Complexes that are Lipschitz Manifolds, Geometric Topology (J. Cantrell, ed.), Academic Press, New York, 1979, pp. 503-552.
- [24] N. Steenrod, The Topology of Fibre Bundles (1951). Vol. 14. Princeton Mathematical Series. Princeton University Press, Princeton, N. J.
- [25] D. Trotman, Stratification theory, Chapter 4 In: Cisneros-Molina, J.L., Lê, D.T., Seade, J. (eds.) Handbook of Geometry and Topology of Singularities, Volume I. Springer, Cham, 231 – 260.
- [26] J. L. Verdier, Stratifications de Whitney et théorème de Bertini-Sard, Invent. Math., Vol. 36, 1976, pp. 295-312.
- [27] H. Whitney, Local properties of analytic varieties, in Differential and Combinatorial Topology (ed. S. S. Cairns), Princeton Univ. Press (1965), 205-244.

I2M CNRS, AIX-MARSEILLE UNIVERSITY, MARSEILLE, FRANCE. JEAN-PAUL.BRASSELET@UNIV-AMU.FR

UNIWERSYTET WARSZAWSKI, POLSKA, TMOSTOWS@MIMUW.EDU.PL

São Paulo State University, UNESP/Ibilce, São José do Rio do Preto, Brazil, bich.thuy@unesp.br