

FREE PRODUCTS AND RESCALINGS INVOLVING NON-SEPARABLE ABELIAN VON NEUMANN ALGEBRAS

KEN DYKEMA AND JUNCHEN ZHAO

ABSTRACT. For a self-symmetric tracial von Neumann algebra A , we study rescalings of $A^{*n} * L\mathbb{F}_r$ for $n \in \mathbb{N}$ and $r \in (1, \infty]$ and use them to obtain an interpolation $\mathcal{F}_{s,r}(A)$ for all real numbers $s > 0$ and $1 - s < r \leq \infty$. We get formulas for their free products, and free products with finite-dimensional or hyperfinite von Neumann algebras. In particular, for any such A , we can compute compressions $(A^{*n})^t$ for $0 < t < 1$, and the Murray-von Neumann fundamental group of $A^{*\infty}$. When A is also non-separable and abelian, this answers two questions in Section 4.3 of [3].

1. INTRODUCTION

The free group factors $L\mathbb{F}_n$ for natural numbers $n \geq 2$ are among the first examples of II_1 factors considered by Murray and von Neumann in the 1940s in [9]. They showed that $L\mathbb{F}_2 \not\cong \mathcal{R}$, where \mathcal{R} is the hyperfinite II_1 factor. However, the free group factor isomorphism problem, first asked in 1967 by Kadison, namely, whether $L\mathbb{F}_n \cong L\mathbb{F}_m$ or not for $n \neq m$, remains open until this day. In the 1990s, using Voiculescu's theory of free probability, the first author in [7] and Rădulescu in [10] independently found a continuous family of II_1 factors $L\mathbb{F}_r$ for $r \in (1, \infty]$ as an interpolation of the free group factors. The family $L\mathbb{F}_r$ satisfies the following rescaling and free product addition formulas:

$$(L\mathbb{F}_r)^t = L\mathbb{F}_{1+\frac{r-1}{t^2}}$$

and

$$L\mathbb{F}_r * L\mathbb{F}_{r'} = L\mathbb{F}_{r+r'}$$

for all $1 < r, r' \leq \infty$ and $0 < t < \infty$. These formulas led to the proof of the dichotomy that the interpolated free group factors, and thus the free group factors, are either all non-isomorphic or all isomorphic to each other.

Recently in Theorem 1.1 of [3], Boutonnet, Drimbe, Ioana, and Popa settled the non-separable version of the free group factor isomorphism problem: for any non-separable diffuse abelian tracial von Neumann algebra A , the II_1 factors A^{*n} are all non-isomorphic and have trivial fundamental group, for all natural numbers $n \geq 2$. Examples of such abelian algebras A include $L\mathbb{R}$, namely, the group von Neumann algebra of \mathbb{R} taken with discrete topology, and the ultrapower $(L\mathbb{Z})^\omega$, where ω is a free ultrafilter on \mathbb{N} . In Proposition

4.1 and Corollary 4.4 in [3], using tools in free probability, compressions $(A^{*n})^t$ are computed to be

$$(A^{*n})^t = A^{*(n/t)} * L\mathbb{F}_{\frac{n-1}{t^2} - \frac{n}{t} + 1} \quad (1)$$

for any $n = 2, 3, \dots, \infty$, $t = 1/m$ for some $m \in \mathbb{N}$, and any *homogeneous* abelian tracial von Neumann algebra A , namely, such that for every $k \in \mathbb{N}$, there exists a partition of unity into k projections $p_1, \dots, p_k \in A$ such that for every $1 \leq j \leq k$ we have that $\tau(p_j) = 1/k$ and $(Ap_j, k\tau|_{Ap_j}) \cong (A, \tau)$ (see Section 4.3 of [3]). Our goal is to study rescalings of $(A^{*n})^t$ for more general $t \in (0, \infty)$.

In order to avoid tracking different pieces of A in the rescalings of the free product, we impose a condition that roughly says A can be continuously decomposed into pieces that are all the same as A itself in a trace-scaling way:

Definition 1.1. A tracial von Neumann algebra (A, τ) is called *self-symmetric* if, for all $0 < t < 1$, there exists a central projection $p \in A$ with $\tau(p) = t$ and tracial isomorphisms $(Ap, t^{-1}\tau|_{Ap}) \cong (A, \tau) \cong (A(1-p), (1-t)^{-1}\tau|_{A(1-p)})$.

Remark 1.2. Note here we are not requiring A to be abelian itself. Rather, we just require A to have central projections that do the job. Although most, if not all, literature related to our results in this paper concerns the cases where A is abelian, we decide to use this definition that witnesses the interpolation of free product phenomenon for a more general class of tracial von Neumann algebras. Note that direct sums of self-symmetric tracial von Neumann algebras are still self-symmetric. Abelian examples include $L\mathbb{Z}$, $(L\mathbb{Z})^\omega$, and $(L\mathbb{Z})^\omega \oplus (L\mathbb{Z})$. Non-abelian examples include any self-symmetric abelian tracial von Neumann algebra tensor product with any non-abelian tracial von Neumann algebra.

Remark 1.3. Note that for an abelian tracial von Neumann algebra (A, τ) , it follows trivially from definitions that being self-symmetric implies being homogeneous, and being homogeneous implies being diffuse. We actually show in Theorem 2.8 that any diffuse abelian tracial von Neumann algebra is self-symmetric, regardless of separability.

For a self-symmetric tracial von Neumann algebra A , $s \in (0, \infty)$, and $r \in (1-s, \infty]$, we define II_1 -factors $\mathcal{F}_{s,r}$ (Definition 3.3) so that $\mathcal{F}_{s,r}$ is isomorphic to $A^{*s} * L\mathbb{F}_r$ when $s \in \mathbb{N}$ and $r \in (1, \infty]$. We prove the rescaling formula

$$(\mathcal{F}_{s,r})^t \cong \mathcal{F}_{\frac{s}{t}, \frac{s+r-1}{t^2} - \frac{s}{t} + 1} \quad (2)$$

for all $s \in (0, \infty)$, $r \in (1-s, \infty]$, $t \in (0, \infty)$. We also prove the addition formula for free products

$$\mathcal{F}_{s,r} * \mathcal{F}_{s',r'} \cong \mathcal{F}_{s+s', r+r'} \quad (3)$$

for all $s, s' \in (0, \infty)$, $r \in (1-s, \infty]$, $r' \in (1-s', \infty]$.

We comment that, by results in [3], if A is also non-separable and abelian, then $\mathcal{F}_{s,r} \not\cong \mathcal{F}_{s',r'}$ whenever $s \neq s'$. We will also develop theorems about countably infinite free products of members of the family $\mathcal{F}_{s,r}$. Combining with our formulas (2) and (3), those allow us to show the fundamental group of $A^{*\infty}$ is \mathbb{R}_+^* , for all self-symmetric tracial von Neumann algebras A .

2. BACKGROUND AND NOTATION

2.1. Tracial von Neumann algebras and their isomorphisms. A tracial von Neumann algebra is a pair (M, τ) of a von Neumann algebra M endowed with a normal, faithful, tracial state τ . By an isomorphism between tracial von Neumann algebras (M, τ_M) and (N, τ_N) , we mean there exists a trace-preserving $*$ -isomorphism, namely, $\phi : M \rightarrow N$ such that $\tau_N \circ \phi = \tau_M$ on M . The isomorphisms throughout this paper will all be tracial. We, however, simply write as $M \cong N$ when there is no ambiguity about the traces involved.

2.2. Rescaling of a II_1 factor. For a II_1 factor M and $t \in (0, \infty)$, we denote the rescaling of M by t as M^t . These were first considered by Murray and von Neumann in [9] and they used them to define the fundamental group of a II_1 factor.

2.3. Tracial ultrapowers and non-separability. See Section 2.2 of [4] for an excellent exposition of ultrafilters and ultrapowers. We observe the following corollary to the proofs of Lemma 2.2 and 2.3(1) in [4]: for any free ultrafilter ω on \mathbb{N} , $(L\mathbb{Z})^\omega$ is not only non-separable but also *purely non-separable*, which in the abelian case means having no separable direct summand.

2.4. Free dimension. Free dimension, first defined in [5], is a tool for computing various free products involving finite-dimensional or hyperfinite von Neumann algebras, interpolated free group factors, and direct sums of those. See [5] for details.

2.5. Weighted direct sum. For a finite or countably infinite index set I , and tracial von Neumann algebras $\{(A_i, \tau_i)\}_{i \in I}$, we denote as

$$A = \bigoplus_{i \in I} \begin{matrix} p_i \\ A_i \\ t_i \end{matrix}$$

the von Neumann algebra direct sum of $\{A_i\}_{i \in I}$ equipped with the trace $\tau((a_i)_{i \in I}) = \sum_{i \in I} t_i \tau_i(a_i)$, where $0 < t_i < 1$ for all $i \in I$, $\sum_{i \in I} t_i = 1$, and p_i is the projection in A that is the identity in A_i and zero elsewhere.

Here we prove a lemma on weighted direct sums of self-symmetric tracial von Neumann algebras that will be of use later.

Lemma 2.1. *Let I be a finite or countably infinite index set, and $\{t_i\}_{i \in I}$ be numbers in $(0, 1)$ such that $\sum_{i \in I} t_i = 1$. Let (A, τ) be a self-symmetric tracial von Neumann algebra. Then*

$$A \cong \bigoplus_{i \in I}^{t_i} A_i$$

where each A_i is a copy of A .

Proof. Using the pair of isomorphisms in Definition 1.1, we can recursively choose mutually orthogonal central projections $\{p_i\}$ in A with $\tau(p_i) = t_i$ and $(Ap_i, t_i^{-1}\tau|_{Ap_i}) \cong (A, \tau)$. Then we have $\sum_{i \in I} p_i = 1_A$ since

$$\tau\left(\sum_{i \in I} p_i\right) = \sum_{i \in I} \tau(p_i) = \sum_{i \in I} t_i = 1,$$

and thus

$$A \cong \bigoplus_{i \in I} Ap_i \cong \bigoplus_{i \in I}^{t_i} A_i.$$

□

2.6. Diffuse abelian tracial von Neumann algebras. See [2] for a comprehensive account of abelian von Neumann algebras. It contains classification results of *von Neumann measure algebras* (von Neumann algebras with a faithful normal semi-finite weight) that enable us to prove that every diffuse abelian tracial von Neumann algebra is self-symmetric (see Theorem 2.8 below).

Definition 2.2 (Section 9.1 in [2]). Let κ be a cardinal. Let $(\mathcal{A}_\kappa, \tau_\kappa)$ denote the tensor product of the family of von Neumann probability algebras $\{(A_i, \tau_i)\}_{i \in \kappa}$, where each (A_i, τ_i) is the tracial von Neumann algebra

$$\mathbb{C} \oplus_{1/2} \mathbb{C}.$$

In particular, τ_κ is a normal faithful tracial state on \mathcal{A}_κ .

Definition 2.3 (Definition 9.7 in [2]). Let A be an abelian von Neumann algebra. We denote by $\mathfrak{k}(A)$ the smallest cardinality of a set P of projections in A not containing 1_A such that $P \cup \{1_A\}$ generates A as a von Neumann algebra.

Definition 2.4 (Definition 9.9 in [2]). Let κ be a cardinal. We say that an abelian von Neumann algebra A is κ -homogeneous if for any non-zero projection $p \in A$, we have $\kappa = \mathfrak{k}(A) = \mathfrak{k}(Ap)$.

The following is a special case of Theorem 9.16 in [2].

Theorem 2.5. *Any two κ -homogeneous abelian von Neumann measure algebras equipped with normal, faithful, tracial states are isomorphic, via a trace-preserving von Neumann algebra isomorphism.*

Theorem 2.6 (Proposition 9.18 in [2]). *For every infinite cardinal κ , the von Neumann algebra \mathcal{A}_κ (Definition 2.2) is abelian, σ -finite, diffuse, and κ -homogeneous.*

Theorem 2.7 (Theorem 9.20 in [2]). *Let (A, τ) be a diffuse abelian von Neumann measure algebra. Then there are families $(\kappa_i)_{i \in I}$ and $(\lambda_i)_{i \in I}$, where each κ_i is an infinite cardinal and each λ_i is a positive real number, such that*

$$(A, \tau) \cong \bigoplus_{i \in I} (\mathcal{A}_{\kappa_i}, \lambda_i \tau_{\kappa_i}).$$

Theorem 2.8. *Let (A, τ) be a diffuse abelian tracial von Neumann algebra. Then (A, τ) is self-symmetric.*

Proof. Write (A, τ) in the sense of Theorem 2.7 above. Since τ is a faithful normal tracial state, I must be countable, and $\sum_{i \in I} \lambda_i = 1$. Fix $0 < t < 1$, we need to show there exists a projection $p \in A$ with $\tau(p) = t$ and

$$(Ap, t^{-1}\tau|_{Ap}) \cong (A, \tau) \cong (A(1-p), (1-t)^{-1}\tau|_{A(1-p)}). \quad (4)$$

For each $i \in I$, we take a projection $p_i \in \mathcal{A}_{\kappa_i}$ with $\tau_{\kappa_i}(p_i) = t$. Now note for all non-zero projections $q \in \mathcal{A}_{\kappa_i} p_i$,

$$\mathfrak{k}((\mathcal{A}_{\kappa_i} p_i)q) = \mathfrak{k}(\mathcal{A}_{\kappa_i} q) = \mathfrak{k}(\mathcal{A}_{\kappa_i}) = \mathfrak{k}(\mathcal{A}_{\kappa_i} p_i) = \kappa_i,$$

because \mathcal{A}_{κ_i} is κ_i -homogeneous, and q and p_i are both non-zero projections in \mathcal{A}_{κ_i} . This implies $\mathcal{A}_{\kappa_i} p_i$ is κ_i -homogeneous, and $(\mathcal{A}_{\kappa_i} p_i, t^{-1}\lambda_i \tau_{\kappa_i}|_{\mathcal{A}_{\kappa_i} p_i})$ is a κ_i -homogeneous von Neumann measure algebra with

$$t^{-1}\lambda_i \tau_{\kappa_i}(p_i) = \lambda_i = \lambda_i \tau_{\kappa_i}(1_{\kappa_i}),$$

where p_i is the identity in $\mathcal{A}_{\kappa_i} p_i$, and 1_{κ_i} is the identity in \mathcal{A}_{κ_i} . Then by Theorem 2.5 and 2.6, we have a tracial isomorphism

$$(\mathcal{A}_{\kappa_i} p_i, t^{-1}\lambda_i \tau_{\kappa_i}|_{\mathcal{A}_{\kappa_i} p_i}) \cong (\mathcal{A}_{\kappa_i}, \lambda_i \tau_{\kappa_i}).$$

Doing so for all $i \in I$, and letting $p = \sum_{i \in I} p_i$, we obtain in the direct sum

$$(Ap, t^{-1}\tau|_{Ap}) \cong \bigoplus_{i \in I} (\mathcal{A}_{\kappa_i} p_i, \lambda_i t^{-1}\tau_{\kappa_i}) \cong \bigoplus_{i \in I} (\mathcal{A}_{\kappa_i}, \lambda_i \tau_{\kappa_i}) \cong (A, \tau),$$

which proves the first isomorphism in (4). Similarly, we can obtain the second isomorphism in (4) and thus (A, τ) is self-symmetric. \square

Remark 2.9. This theorem provides us with many examples of self-symmetric tracial von Neumann algebras, including the group von Neumann algebras of all infinite discrete abelian groups, regardless of the cardinality. (See, for example, Proposition 5.1 of [5] for a proof of the well-known fact that all such algebras are diffuse.)

3. WELL-DEFINEDNESS AND PROPERTIES OF $\mathcal{F}_{s,r}$

Throughout this section, unless otherwise noted, we fix a self-symmetric tracial von Neumann algebra A . We denote $\mathcal{F}_{n,r} = A^{*n} * L\mathbb{F}_r$ for $n \in \mathbb{N}$ and $r \in (1, \infty]$. Since the free product of any diffuse tracial von Neumann algebra and any non-trivial tracial von Neumann algebra is a II_1 factor (see Corollary 5.3.8 in [1] for example). Our plan is to define $\mathcal{F}_{s,r}$ for more general s and r by rescalings of these $\mathcal{F}_{n,r}$. We first compute $(\mathcal{F}_{n,r})^{1/k}$.

Lemma 3.1. *Let $n, k \in \mathbb{N}$ and $r \in (1, \infty]$, then*

$$(A^{*n} * L\mathbb{F}_r)^{1/k} \cong A^{*(nk)} * L\mathbb{F}_{(n+r-1)k^2 - nk + 1},$$

in other words

$$(\mathcal{F}_{n,r})^{1/k} \cong \mathcal{F}_{nk, (n+r-1)k^2 - nk + 1}.$$

Proof. We first prove the case $n = 1$ and proceed by induction on n . From the self-symmetry of A , we take projections $p_1, \dots, p_k \in A$, where $\sum_{j=1}^k p_j = 1_A$, $\tau(p_j) = 1/k$, and $(Ap_j, k\tau|_{Ap_j}) \cong (A, \tau)$ for all $1 \leq j \leq k$. Then

$$\begin{aligned} (A * L\mathbb{F}_r)^{1/k} &\cong p_1((Ap_1 \oplus \dots \oplus Ap_k) * L\mathbb{F}_r)p_1 \\ &\cong Ap_1 * p_1((\mathbb{C}p_1 \oplus Ap_2 \oplus \dots \oplus Ap_k) * L\mathbb{F}_r)p_1 \\ &\cong Ap_1 * p_2((\mathbb{C}p_1 \oplus Ap_2 \oplus \dots \oplus Ap_k) * L\mathbb{F}_r)p_2 \\ &\cong \dots \\ &\cong Ap_1 * \dots * Ap_k * ((\mathbb{C}p_1 \oplus \dots \oplus \mathbb{C}p_k) * L\mathbb{F}_r)^{1/k} \\ &\cong A^{*k} * (L\mathbb{F}_{\frac{k(k-1)}{k^2} + r})^{1/k} \cong A^{*k} * L\mathbb{F}_{rk^2 - k + 1}. \end{aligned}$$

follows from Claim 4.3 in [3] and Proposition 2.4 in [5]. Next suppose the lemma holds for some $n \in \mathbb{N}$. Then

$$\begin{aligned} (A^{*(n+1)} * L\mathbb{F}_r)^{1/k} &= (A * A^{*n} * L\mathbb{F}_r)^{1/k} \\ &\cong p_1((Ap_1 \oplus \dots \oplus Ap_k) * A^{*n} * L\mathbb{F}_r)p_1 \\ &\cong Ap_1 * \dots * Ap_k * p_1((\mathbb{C}p_1 \oplus \dots \oplus \mathbb{C}p_k) * A^{*n} * L\mathbb{F}_r)p_1 \\ &\cong A^{*k} * (A^{*n} * L\mathbb{F}_{\frac{k(k-1)}{k^2} + r})^{1/k} \\ &\cong A^{*k} * (A^{*(nk)} * L\mathbb{F}_{(n+r)k^2 - (n+1)k + 1}) \\ &\cong A^{*(n+1)k} * L\mathbb{F}_{(n+r)k^2 - (n+1)k + 1} \end{aligned}$$

follows similarly by associativity and commutativity of the free product, and the inductive hypothesis. \square

The next lemma is key to well-definedness:

Lemma 3.2. *Let $n, m \in \mathbb{N}$, $r, u \in (1, \infty]$, and $t, s \in (0, \infty)$ satisfy*

$$\frac{n}{t} = \frac{m}{s} \quad \text{and} \quad \frac{n+r-1}{t^2} = \frac{m+u-1}{s^2}$$

then

$$(\mathcal{F}_{n,r})^t \cong (\mathcal{F}_{m,u})^s.$$

Proof. Using the previous lemma, we can compute

$$((\mathcal{F}_{n,r})^t)^{1/m} \cong ((\mathcal{F}_{n,r})^{1/m})^t \cong (\mathcal{F}_{nm,(n+r-1)m^2-nm+1})^t$$

and

$$((\mathcal{F}_{m,u})^s)^{1/m} \cong ((\mathcal{F}_{m,u})^{1/n})^t \cong (\mathcal{F}_{nm,(m+u-1)n^2-nm+1})^t.$$

The right-hand sides are precisely the same by our assumption, so

$$(\mathcal{F}_{n,r})^t \cong M_m(\mathbb{C}) \otimes (\mathcal{F}_{nm,(m+u-1)n^2-nm+1})^t \cong (\mathcal{F}_{m,u})^s.$$

□

Definition 3.3. For a self-symmetric tracial von Neumann algebra A , for $s \in (0, \infty)$, and $r \in (1 - s, \infty]$, we choose $n \in \mathbb{N}$ large enough so that

$$\frac{s+r-1}{s^2}n^2 - n + 1 > 1$$

and define

$$\mathcal{F}_{s,r}(A) := (A^{*n} * L\mathbb{F}_{\frac{s+r-1}{s^2}n^2-n+1})^{n/s}. \quad (5)$$

When the algebra A is clear from context, we abbreviate the above as $\mathcal{F}_{s,r}$. When $r = \infty$, we use its usual arithmetic in the extended real number system.

Remark 3.4. This definition gives isomorphic copies of II_1 factors regardless of n by Lemma 3.2. It is also elementary to verify that this definition recovers $\mathcal{F}_{s,r} \cong A^{*s} * L\mathbb{F}_r$ when $s \in \mathbb{N}$ and $r \in (1, \infty]$, and $\mathcal{F}_{s,0} \cong A^{*s}$ when $s \in \mathbb{N}$ and $s \geq 2$. The first isomorphism is obtained by setting $n = s$ in (5). The second isomorphism follows from setting $n = 2s$ in (5), and then taking the $1/2$ compression:

$$(\mathcal{F}_{s,0})^{1/2} \cong A^{*(2s)} * L\mathbb{F}_{2s-3} \cong (A^{*s})^{1/2},$$

which follows from (1).

The rescaling formula (2) now follows directly from well-definedness.

Theorem 3.5. Let $s \in (0, \infty)$, $r \in (1 - s, \infty]$, and $t \in (0, \infty)$. Then

$$(\mathcal{F}_{s,r})^t \cong \mathcal{F}_{\frac{s}{t}, \frac{s+r-1}{t^2} - \frac{s}{t} + 1}.$$

Proof. It suffices to choose $n \in \mathbb{N}$ with

$$\frac{s+r-1}{s^2}n^2 - n + 1 > 1$$

to realize both sides of the isomorphism above as in (5). Then it follows

$$\begin{aligned} \mathcal{F}_{\frac{s}{t}, \frac{s+r-1}{t^2} - \frac{s}{t} + 1} &\cong (A^{*n} * L\mathbb{F}_{\frac{s+r-1}{s^2}n^2-n+1})^{nt/s} \\ &\cong (\mathcal{F}_{s,r})^t. \end{aligned}$$

□

Remark 3.6. The rescaling formula in particular extends the classical rescaling formula of the interpolated free group factors when A is separable, and the result in Corollary 4.4(1) of [3] when $s \geq 2$ is a natural number, $r = 0$, and $t = 1/m$ for some $m \in \mathbb{N}$.

Theorem 3.7. *Let A be a non-separable self-symmetric abelian tracial von Neumann algebra. Let $0 < s \neq s' < \infty$, $r \in (1 - s, \infty]$, and $r' \in (1 - s', \infty]$. Then*

$$\mathcal{F}_{s,r}(A) \not\cong \mathcal{F}_{s',r'}(A).$$

Proof. We use the sans-rank of [3] to distinguish. From Remark 2.11, Proposition 2.12, Theorem 3.8, and proof of Theorem 1.1 of [3],

$$\mathrm{r}_{\mathrm{ns}}(\mathcal{F}_{s,r}(A)) = s \cdot \mathrm{r}_{\mathrm{ns}}(A) = s \cdot \tau(p) \neq s' \cdot \tau(p) = \mathrm{r}_{\mathrm{ns}}(\mathcal{F}_{s',r'}(A)),$$

where $p \in A$ is the maximal, necessarily non-zero projection such that Ap is purely non-separable. Therefore, the non-isomorphism holds regardless of r and r' . \square

Now we work on proving the addition formula (3).

Lemma 3.8. *Let $n \in \mathbb{N}$ and $s \in (0, \infty)$ satisfy*

- (i) $n \geq 3$,
 - (ii) $\max\{t, 1 - t\} < \frac{n}{n+1}$,
 - (iii) $r = \frac{n-1}{t^2} - \frac{n}{t} + 1$,
- where $t = \frac{n}{s+n}$. Then we have

$$A^{*n} * \mathcal{F}_{s,r} \cong \mathcal{F}_{s+n,r}.$$

Proof. Denote the n copies of A in A^{*n} by A_1, \dots, A_n respectively. For all $1 \leq j \leq n$, by self-symmetry of A_j , we take a projection $e_j \in A_j$ of trace t with the trace-preserving isomorphisms as in Definition 1.1. We know $e_1 \sim \dots \sim e_n$ in

$$\left(\begin{smallmatrix} \mathbb{C} & 1-e_1 \\ t & 1-t \end{smallmatrix} \right) * \dots * \left(\begin{smallmatrix} \mathbb{C} & 1-e_n \\ t & 1-t \end{smallmatrix} \right),$$

since, using conditions (i) and (ii), the above is known to be an interpolated free group factor by Theorem 1.1 and 4.6 in [5]. From Fact 4.2 of [3],

$$\begin{aligned} & (A_1 * \dots * A_n)^t \\ & \cong e_1((A_1 e_1 \oplus A_1(1 - e_1)) * A_2 * \dots * A_n) e_1 \\ & \cong A_1 e_1 * e_1((\mathbb{C} e_1 \oplus A_1(1 - e_1)) * A_2 * \dots * A_n) e_1 \\ & \cong A_1 e_1 * e_2 \left(\begin{smallmatrix} \mathbb{C} & 1-e_1 \\ t & 1-t \end{smallmatrix} \oplus A_1 \right) * (A_2 e_2 \oplus A_2(1 - e_2)) * \dots * A_n) e_2 \\ & \cong \dots \\ & \cong A_1 e_1 * \dots * A_n e_n * e_n \left(\begin{smallmatrix} \mathbb{C} & 1-e_1 \\ t & 1-t \end{smallmatrix} \oplus A_1 \right) * \dots * \left(\begin{smallmatrix} \mathbb{C} & 1-e_n \\ t & 1-t \end{smallmatrix} \oplus A_n \right) e_n \\ & \cong A^{*n} * \left(\begin{smallmatrix} \mathbb{C} & A \\ t & 1-t \end{smallmatrix} \right)^{*n}{}^t. \end{aligned}$$

In the same way, using Theorem 1.1 and 4.6 of [5],

$$\begin{aligned} ((\mathbb{C} \oplus A)_{t \ 1-t}^{*n})^{1-t} &\cong A^{*n} * ((\mathbb{C} \oplus \mathbb{C})_{t \ 1-t}^{*n})^{1-t} \\ &\cong A^{*n} * (L\mathbb{F}_{2nt(1-t)})^{1-t}. \end{aligned}$$

Employing Definition 3.3 and formula (2), we obtain:

$$\begin{aligned} \mathcal{F}_{\frac{n}{t}, \frac{n-1}{t^2} - \frac{n}{t} + 1} &\cong (\mathcal{F}_{n,0})^t \cong (A^{*n})^t \\ &\cong A^{*n} * (A^{*n} * (L\mathbb{F}_{2nt(1-t)})^{1-t})^{\frac{t}{1-t}} \\ &\cong A^{*n} * (\mathcal{F}_{n,1+\frac{2nt(1-t)-1}{(1-t)^2}})^{\frac{t}{1-t}} \\ &\cong A^{*n} * \mathcal{F}_{\frac{n}{t} - n, \frac{n-1}{t^2} - \frac{n}{t} + 1}. \end{aligned}$$

which is exactly

$$A^{*n} * \mathcal{F}_{s,r} \cong \mathcal{F}_{s+n,r}.$$

□

Lemma 3.9. *Let $s \in (0, \infty)$, $r \in (1 - s, \infty]$, and $u \in (1, \infty]$. Then*

$$\mathcal{F}_{s,r} * L\mathbb{F}_u \cong \mathcal{F}_{s,r+u}.$$

Proof. We choose $n \in \mathbb{N}$ large enough so that $\frac{n^2}{s^2} - 1 > 1$ and $\mathcal{F}_{s,r} \cong (\mathcal{F}_{n, \frac{r+s-1}{s^2}n^2 - n + 1})^{n/s}$ as in Definition 3.3. It then follows from Theorem 1.5 in [8] that

$$\begin{aligned} (\mathcal{F}_{s,r} * L\mathbb{F}_u)^{s/n} &\cong (\mathcal{F}_{s,r})^{s/n} * (L\mathbb{F}_u)^{s/n} * L\mathbb{F}_{\frac{n^2}{s^2} - 1} \\ &\cong A^{*n} * L\mathbb{F}_{\frac{r+s-1}{s^2}n^2 - n + 1} * L\mathbb{F}_{1+(u-1)\frac{n^2}{s^2}} * L\mathbb{F}_{\frac{n^2}{s^2} - 1} \\ &\cong A^{*n} * L\mathbb{F}_{\frac{r+s-1}{s^2}n^2 - n + 1 + u\frac{n^2}{s^2}}. \end{aligned}$$

Hence, by the rescaling formula (2),

$$\mathcal{F}_{s,r} * L\mathbb{F}_u \cong (\mathcal{F}_{n, \frac{r+s-1}{s^2}n^2 - n + 1 + u\frac{n^2}{s^2}})^{n/s} \cong \mathcal{F}_{s,r+u}.$$

□

Theorem 3.10. *Let $s, v \in (0, \infty)$, $r \in (1 - s, \infty]$, $u \in (1 - v, \infty]$, then*

$$\mathcal{F}_{s,r} * \mathcal{F}_{v,u} \cong \mathcal{F}_{s+v,r+u}.$$

Proof. Without loss of generality we assume $v \geq s$. We realize $\mathcal{F}_{s,r}$ and $\mathcal{F}_{v,u}$ by choosing $n, m \in \mathbb{N}$ large enough so that

$$(i) \quad n^2/s^2 - 1 > 1,$$

$$(ii) \quad n \geq 3,$$

$$(iii) \quad \frac{v}{s+v} < \frac{n}{n+1},$$

(iv)

$$\frac{r+s+u+v-1}{s^2}n^2 - \frac{(s+v)^2}{s^2}n + \frac{(s+v)^2}{s^2} > 1,$$

as well as

$$\mathcal{F}_{s,r} \cong (\mathcal{F}_{n, \frac{r+s-1}{s^2}n^2-n+1})^{n/s}$$

and

$$\mathcal{F}_{v,u} \cong (\mathcal{F}_{m, \frac{u+v-1}{v^2}m^2-m+1})^{m/v}$$

as in Definition 3.3. We let $t = s/n$. It follows from Theorem 1.5 in [8] using condition (i), rescaling formula (2), and Lemma 3.9 that

$$\begin{aligned} (\mathcal{F}_{s,r} * \mathcal{F}_{v,u})^t &\cong (\mathcal{F}_{n, \frac{r+s-1}{s^2}n^2-n+1})^{nt/s} * (\mathcal{F}_{m, \frac{u+v-1}{v^2}m^2-m+1})^{mt/v} * L\mathbb{F}_{t^{-2}-1} \\ &\cong \mathcal{F}_{n, \frac{r+s-1}{s^2}n^2-n+1} * \mathcal{F}_{\frac{v}{s}n, \frac{u+v-1}{s^2}n^2-\frac{v}{s}n+1} * L\mathbb{F}_{t^{-2}-1} \\ &\cong \mathcal{F}_{n,0} * L\mathbb{F}_{\frac{r+s-1}{s^2}n^2-n+1} * \mathcal{F}_{\frac{v}{s}n, \frac{u+v-1}{s^2}n^2-\frac{v}{s}n+1} * L\mathbb{F}_{t^{-2}-1} \\ &\cong A^{*n} * \mathcal{F}_{\frac{v}{s}n, \frac{r+s+u+v-1}{s^2}n^2-\frac{s+v}{s}n+1}. \end{aligned}$$

By Lemma 3.8 using conditions (ii) and (iii), A^{*n} can be absorbed like so:

$$A^{*n} * \mathcal{F}_{\frac{v}{s}n, (\frac{(s+v)^2}{s^2} - \frac{s+v}{s})n - \frac{(s+v)^2}{s^2} + 1} \cong \mathcal{F}_{\frac{v}{s}n+n, (\frac{(s+v)^2}{s^2} - \frac{s+v}{s})n - \frac{(s+v)^2}{s^2} + 1}.$$

Now by Lemma 3.9 using condition (iv), we add back an interpolated free group factor to obtain:

$$\begin{aligned} (\mathcal{F}_{s,r} * \mathcal{F}_{v,u})^t &\cong A^{*n} * \mathcal{F}_{\frac{v}{s}n, \frac{r+s+u+v-1}{s^2}n^2-\frac{s+v}{s}n+1} \\ &\cong A^{*n} * \mathcal{F}_{\frac{v}{s}n, (\frac{(s+v)^2}{s^2} - \frac{s+v}{s})n - \frac{(s+v)^2}{s^2} + 1} * L\mathbb{F}_{\frac{r+s+u+v-1}{s^2}n^2 - \frac{(s+v)^2}{s^2}n + \frac{(s+v)^2}{s^2}} \\ &\cong \mathcal{F}_{\frac{v}{s}n+n, (\frac{(s+v)^2}{s^2} - \frac{s+v}{s})n - \frac{(s+v)^2}{s^2} + 1} * L\mathbb{F}_{\frac{r+s+u+v-1}{s^2}n^2 - \frac{(s+v)^2}{s^2}n + \frac{(s+v)^2}{s^2}} \\ &\cong \mathcal{F}_{\frac{v}{s}n+n, \frac{r+s+u+v-1}{s^2}n^2-\frac{s+v}{s}n+1} \\ &\cong (\mathcal{F}_{s+v, r+u})^t, \end{aligned}$$

where the last isomorphism is just the rescaling formula (2). The theorem is then proved by an amplification by $1/t$ on both sides. \square

4. MORE RESULTS AND APPLICATIONS

For a non-separable self-symmetric tracial von Neumann algebra A , heuristically, we can think of the parameter r in $\mathcal{F}_{s,r}$ as the free dimension of the separable component, and the parameter s as the free dimension of the non-separable component. The next proposition shows different von Neumann algebras with the same free dimension would give isomorphic free products with our family $\mathcal{F}_{s,r}$. It is therefore an extension of Lemma 3.9 above.

Proposition 4.1. *Fix some $\mathcal{F}_{s,r}$ and $u \in (0, \infty]$. Let B_u be any finite-dimensional (with dimension ≥ 2) or hyperfinite von Neumann algebra, or an interpolated free group factor, or direct sums of those, with free dimension u . Then*

$$\mathcal{F}_{s,r} * B_u \cong \mathcal{F}_{s,r+u}.$$

Proof. We choose $n \in \mathbb{N}$ large enough so that $B_u * M_n(\mathbb{C})$ is a factor. By Theorem 4.6 of [5], it suffices for the minimal projections in B_u to have trace less than $1 - n^{-2}$, and then $B_u * M_n(\mathbb{C}) \cong L\mathbb{F}_{u+1-n^{-2}}$. We have by Theorem 1.2 of [5]

$$\begin{aligned} (\mathcal{F}_{s,r} * B_u)^{1/n} &\cong ((M_n(\mathbb{C}) \otimes (\mathcal{F}_{s,r})^{1/n}) * B_u)^{1/n} \\ &\cong (\mathcal{F}_{s,r})^{1/n} * (B_u * M_n(\mathbb{C}))^{1/n}. \end{aligned}$$

Using the rescaling formula (2) and Lemma 3.9:

$$\begin{aligned} (\mathcal{F}_{s,r} * B_u)^{1/n} &\cong \mathcal{F}_{ns, n^2(s+r-1)-ns+1} * (L\mathbb{F}_{u+1-n^{-2}})^{1/n} \\ &\cong \mathcal{F}_{ns, n^2(s+r-1)-ns+1} * L\mathbb{F}_{1+n^2(u-n^{-2})} \\ &\cong \mathcal{F}_{ns, n^2(u+s+r-1)-ns+1} \\ &\cong (\mathcal{F}_{s,r+u})^{1/n}. \end{aligned}$$

Now our proposition follows by tensoring with $M_n(\mathbb{C})$. \square

The next proposition collects some more free product results involving the family $\mathcal{F}_{s,r}$.

Proposition 4.2. *For a self-symmetric tracial von Neumann algebra A , $n, k \in \mathbb{N}$, $r \in (1, \infty]$ and $t \in (0, 1)$, we have*

(i)

$$A * LZ \cong (\mathcal{F}_{k, k^2-k+1})^k \cong A * \mathcal{R},$$

(ii)

$$A^{*n} * \left(A \oplus_t \mathbb{C} \right) \cong \mathcal{F}_{n+t, t-t^2},$$

(iii)

$$(A \otimes M_k(\mathbb{C})) * L\mathbb{F}_r \cong \mathcal{F}_{\frac{1}{k}, r-\frac{1}{k}+1},$$

(iv)

$$\left(A \oplus_t \mathbb{C} \right) * L(\mathbb{F}_r) \cong \mathcal{F}_{t, r+t-t^2},$$

(v)

$$\left(A \oplus_t LZ \right)^{*n} \cong \mathcal{F}_{nt, n(1-t)}.$$

Proof. We first comment that (i) is analogous to Theorem 4.1 of [6], and (ii) is analogous to Remark 1.5 of [5].

For (i), it suffices to compute the $1/k$ compression of $A * LZ$ for $k \in \mathbb{N}$. As in the beginning of the proof of Lemma 3.1, we choose projections p_1, \dots, p_k in A to obtain

$$\begin{aligned}
(A * LZ)^{1/k} &\cong ((Ap_1 \oplus \dots \oplus Ap_k) * LZ)^{1/k} \\
&\cong Ap_1 * \dots * Ap_k * \left(\left(\underset{1/k}{\mathbb{C}} \oplus \dots \oplus \underset{1/k}{\mathbb{C}} \right) * LZ \right)^{1/k} \\
&\cong A^{*k} * \left(L\mathbb{F}_{\frac{k(k-1)}{k^2}+1} \right)^{1/k} \\
&\cong A^{*k} * L\mathbb{F}_{k^2-k+1} \\
&\cong \mathcal{F}_{k, k^2-k+1}.
\end{aligned}$$

The computation with \mathcal{R} in place of LZ is exactly the same because both have free dimension 1.

For (ii), it follows from the same argument as in the proof of Lemma 3.8.

For (iii), it is a direct application of Theorem 1.2 of [5].

For (iv), let

$$\mathcal{M} = \left(\underset{t}{A} \oplus \underset{1-t}{\mathbb{C}} \right) * L(\mathbb{F}_r).$$

By Theorem 1.2 of [5],

$$\mathcal{M}^t \cong p\mathcal{M}p \cong A * \left(\left(\underset{t}{\mathbb{C}} \oplus \underset{1-t}{\mathbb{C}} \right) * L(\mathbb{F}_r) \right)^t \cong A * L(\mathbb{F}_{r+2t(1-t)})^t \cong \mathcal{F}_{1, \frac{r-1}{t^2} + \frac{2}{t} - 1}.$$

Thus,

$$\mathcal{M} \cong \mathcal{F}_{1, \frac{r-1}{t^2} + \frac{2}{t} - 1}^{1/t} \cong \mathcal{F}_{t, r+t-t^2}.$$

For (v), we choose $k \in \mathbb{N}$ large enough so that

$$\left(\underset{t/k}{\mathbb{C}} \oplus \dots \oplus \underset{t/k}{\mathbb{C}} \oplus \underset{1-t}{LZ} \right)^{*n}$$

is a factor. Then the same argument as in Lemma 3.8 shows

$$\begin{aligned}
\left(\left(\underset{t}{A} \oplus \underset{1-t}{LZ} \right)^{*n} \right)^{t/k} &\cong A^{*(nk)} * \left(\left(\underset{t/k}{\mathbb{C}} \oplus \dots \oplus \underset{t/k}{\mathbb{C}} \oplus \underset{1-t}{LZ} \right)^{*n} \right)^{t/k} \\
&\cong \mathcal{F}_{nk, (n-1)\frac{k^2}{t^2} - nk + 1}.
\end{aligned}$$

Rescaling by k/t gives (v). \square

Remark 4.3. Note that for all $t \in (0, 1)$,

$$A_t := \underset{t}{A} \oplus \underset{1-t}{LZ}$$

is still a non-separable self-symmetric abelian tracial von Neumann algebra if A is so, and $A_t \not\cong A$ since $r_{\text{ns}}(A_t) = t \cdot r_{\text{ns}}(A)$. For all $s \in (0, \infty)$ and $r \in (1-s, \infty]$, we have

$$\mathcal{F}_{s,r}(A_t) \cong \mathcal{F}_{st, s+r-st}(A)$$

by (v) above.

We now consider countably infinite free products of $\mathcal{F}_{s_i, r_i}(A)$ for a self-symmetric tracial von Neumann algebra A . (Note: the result below is generalized in Theorem 4.10.)

Theorem 4.4. *Fix a self-symmetric tracial von Neumann algebra A . Assume for each $i \in \mathbb{N}$, $s_i \in (0, \infty)$, $r_i \in (1 - s_i, \infty]$, and*

$$\sum_{i \in \mathbb{N}} s_i = \infty.$$

Then

$$\ast_{i \in \mathbb{N}} \mathcal{F}_{s_i, r_i} \cong A^{\ast\infty}.$$

Proof. From Theorem 1.5 of [8] and Proposition 4.1, we have

$$\ast_{i \in \mathbb{N}} \mathcal{F}_{s_i, r_i} \cong \left(\ast_{i \in \mathbb{N}} \mathcal{F}_{s_i, r_i} \right) \ast L\mathbb{F}_\infty \cong \ast_{i \in \mathbb{N}} (\mathcal{F}_{s_i, r_i} \ast L\mathbb{F}_\infty) \cong \ast_{i \in \mathbb{N}} \mathcal{F}_{s_i, \infty},$$

so we may without loss of generality assume $r_i = \infty$ for all $i \in \mathbb{N}$. From the free product addition formula (3), we have for $l \in \mathbb{N}$,

$$\ast_{i=k}^{k+l} \mathcal{F}_{s_i, \infty} \cong \mathcal{F}_{s, \infty},$$

where $s = s_k + \dots + s_{k+l}$, so we may without loss of generality assume $s_i \geq 3$ for all $i \in \mathbb{N}$ by regrouping the infinite free product, if necessary. Moreover, from Proposition 4.2, we have

$$\mathcal{F}_{s, \infty} \ast \left(A \oplus_t \mathbb{C} \right) \cong \mathcal{F}_{s+t, \infty} \quad (6)$$

for all $s > 1$ and $0 \leq t < 1$, where we abuse notation to mean

$$A \oplus_0 \mathbb{C} \cong \mathbb{C}.$$

Now let

$$n_1 = \lfloor s_1 \rfloor, \quad t_1 = s_1 - n_1$$

and recursively choose, for all $i \geq 2$,

$$n_i = \lfloor s_i - (1 - t_{i-1}) \rfloor, \quad t_i = s_i - (1 - t_{i-1}) - n_i.$$

In particular, we have $n_1 \geq 1$, $0 \leq t_1 < 1$, and

$$\mathcal{F}_{s_1, \infty} \cong A^{\ast n_1} \ast L\mathbb{F}_\infty \ast \left(A \oplus_{t_1} \mathbb{C} \right),$$

which follows from Remark 3.4 and (6). For all $i \geq 2$, we have $n_i \geq 1$, $t_i \in [0, 1)$ and $s_i = (1 - t_{i-1}) + n_i + t_i$, so

$$\mathcal{F}_{s_i, \infty} \cong \left(A \oplus_{1-t_{i-1}} \mathbb{C} \right) \ast A^{\ast n_i} \ast L\mathbb{F}_\infty \ast \left(A \oplus_{t_i} \mathbb{C} \right).$$

Substituting these into the infinite free product and using associativity and commutativity of the free product, we get

$$\begin{aligned}
& \bigstar_{i=1}^{\infty} \mathcal{F}_{s_i, \infty} \\
& \cong A^{*n_1} * L\mathbb{F}_{\infty} * \left(A \oplus_{t_1} \mathbb{C} \oplus_{1-t_1} \right) * \left(\bigstar_{i=2}^{\infty} \left(A \oplus_{1-t_{i-1}} \mathbb{C} \oplus_{t_{i-1}} \right) * A^{*n_i} * L\mathbb{F}_{\infty} * \left(A \oplus_{t_i} \mathbb{C} \oplus_{1-t_i} \right) \right) \\
& \cong A^{*n_1} * L\mathbb{F}_{\infty} * \left(\bigstar_{i=1}^{\infty} \left(\left(A \oplus_{t_i} \mathbb{C} \oplus_{1-t_i} \right) * \left(A \oplus_{1-t_i} \mathbb{C} \oplus_{t_i} \right) * A^{*n_{i+1}} * L\mathbb{F}_{\infty} \right) \right) \\
& \cong A^{*n_1} * L\mathbb{F}_{\infty} * \left(\bigstar_{i=1}^{\infty} \left(A^{*(1+n_{i+1})} * L\mathbb{F}_{\infty} \right) \right) \\
& \cong A^{*\infty} * L\mathbb{F}_{\infty} \\
& \cong A^{*\infty},
\end{aligned}$$

where we have used (6) twice to obtain the third isomorphism above. \square

Theorem 4.5. *The Murray-von Neumann fundamental group of $A^{*\infty}$ is \mathbb{R}_+^* for all self-symmetric tracial von Neumann algebras A .*

Proof. Let $t \in (0, 1)$ be arbitrary. Then by Theorem 1.5 of [8], the rescaling formula (2), and the above Theorem 4.4,

$$(A^{*\infty})^t \cong \bigstar_{n \in \mathbb{N}} (A * A)^t \cong \bigstar_{n \in \mathbb{N}} \mathcal{F}_{\frac{2}{t}, \frac{1}{t^2} - \frac{2}{t} + 1} \cong A^{*\infty}.$$

This shows the fundamental group of $A^{*\infty}$ contains $(0, 1)$ so is all of \mathbb{R}_+^* . \square

The remainder of this section is dedicated to proving Theorem 4.10, which is a generalization of Theorem 4.4. We nonetheless keep the proof of Theorem 4.4 presented above in our exposition, because it is considerably easier than the proof of the more general result, and because Theorem 4.10 is not needed for the above result about the fundamental group.

The following is well known, but for convenience we provide a proof.

Lemma 4.6. *Let (\mathcal{M}, τ) be a tracial von Neumann algebra. Suppose p and q are projections in \mathcal{M} that are free and satisfy $\tau(p) = \tau(q)$. Then there exists a unitary element, u , in the von Neumann algebra generated by $\{p, q, 1\}$, such that $u^*qu = p$. Moreover, the central support of p in \mathcal{M} equals the central support of q in \mathcal{M} .*

Proof. Let $\alpha = \tau(p)$. If $\alpha = 0$ or $\alpha = 1$, then we may take $u = 1$. Suppose $0 < \alpha < 1$. Let μ be the distribution of pqp . Then μ is equal to the multiplicative free convolution of the measure $(1 - \alpha)\delta_0 + \alpha\delta_1$ with itself. By Example 2.8 in [11], μ has an atom at 0 with mass equal to $1 - \alpha$. Consequently, the support projection of pqp is equal to p . Let $qp = v(pqp)^{1/2}$ be the polar decomposition of qp . Then v is a partial isometry satisfying $v^*v = p$ and $vv^* = q$. Similarly, from the polar decomposition of $(1 - q)(1 - p)$, we obtain a partial isometry w satisfying $w^*w = 1 - p$ and $w w^* = 1 - q$. Let $u = v + w$. Then $u^*qu = p$.

The existence of the unitary u implies the statement about the central supports. \square

Lemma 4.7. *Let B be a tracial von Neumann algebra and let $\alpha \in (0, 1)$. Let*

$$\mathcal{M} = \left(\overset{p}{\underset{\alpha}{\mathbb{C}}} \oplus B\right) * \left(\overset{q}{\underset{\alpha}{\mathbb{C}}} \oplus \tilde{B}\right),$$

where \tilde{B} is a copy of B . Consider the von Neumann subalgebras

$$\mathcal{N} = \left(\overset{p}{\underset{\alpha}{\mathbb{C}}} \oplus B\right) * \left(\overset{q}{\underset{\alpha}{\mathbb{C}}} \oplus \mathbb{C}\right),$$

$$\tilde{\mathcal{N}} = \left(\overset{p}{\underset{\alpha}{\mathbb{C}}} \oplus \mathbb{C}\right) * \left(\overset{q}{\underset{\alpha}{\mathbb{C}}} \oplus \tilde{B}\right)$$

of \mathcal{M} . Then there is a trace-preserving automorphism β of \mathcal{M} satisfying

$$\beta(p) = p, \quad \beta(\mathcal{N}) = \tilde{\mathcal{N}} \quad \text{and} \quad \beta(\tilde{\mathcal{N}}) = \mathcal{N}.$$

Proof. Let σ be the free flip automorphism of \mathcal{M} , namely, that interchanges the two copies $\mathbb{C} \oplus B$ and $\mathbb{C} \oplus \tilde{B}$, mapping each identically onto the other. Note that σ is trace-preserving, $\sigma^2 = \text{id}$, $\sigma(p) = q$ and $\sigma(\mathcal{N}) = \tilde{\mathcal{N}}$. Let $u \in W^*(\{p, q, 1\})$ be the unitary from Lemma 4.6 satisfying $u^*qu = p$. Note that we have $u \in \mathcal{N} \cap \tilde{\mathcal{N}}$. Let $\beta(\cdot) = u^*\sigma(\cdot)u$. Then β is a trace-preserving automorphism of \mathcal{M} and

$$\begin{aligned} \beta(p) &= u^*qu = p, \\ \beta(\mathcal{N}) &= u^*\tilde{\mathcal{N}}u = \tilde{\mathcal{N}} \\ \beta(\tilde{\mathcal{N}}) &= u^*\mathcal{N}u = \mathcal{N}. \end{aligned}$$

\square

Lemma 4.8. *Let A and B be tracial von Neumann algebras and let $\alpha \in (0, 1)$. Let*

$$(\mathcal{M}, \tau_{\mathcal{M}}) = \left(\overset{p}{\underset{\alpha}{A}} \oplus B\right) * \left(\overset{q}{\underset{\alpha}{\mathbb{C}}} \oplus \tilde{B}\right),$$

where \tilde{B} is a copy of B . Consider the von Neumann subalgebras

$$\mathcal{N} = \left(\overset{p}{\underset{\alpha}{\mathbb{C}}} \oplus B\right) * \left(\overset{q}{\underset{\alpha}{\mathbb{C}}} \oplus \mathbb{C}\right),$$

$$\tilde{\mathcal{N}} = \left(\overset{p}{\underset{\alpha}{\mathbb{C}}} \oplus \mathbb{C}\right) * \left(\overset{q}{\underset{\alpha}{\mathbb{C}}} \oplus \tilde{B}\right).$$

Then there is a trace-preserving automorphism γ of \mathcal{M} such that

$$\gamma(p) = p, \quad \gamma(\mathcal{N}) = \tilde{\mathcal{N}}, \quad \gamma(\tilde{\mathcal{N}}) = \mathcal{N}$$

and γ maps the subalgebra $A \oplus \mathbb{C}$ of \mathcal{M} identically to itself (by which we mean the restriction of γ to this subalgebra is the identity map of the subalgebra to itself).

Proof. Consider the von Neumann subalgebra

$$\mathcal{Q} = \left(\overset{p}{\mathbb{C}} \oplus B \right) * \left(\overset{q}{\mathbb{C}} \oplus \tilde{B} \right)$$

of \mathcal{M} . Clearly, $A = A \oplus 0$ and \mathcal{Q} together generate \mathcal{M} . By Theorem 1.2 of [5], A and $p\mathcal{Q}p$ are free in $p\mathcal{M}p$ with respect to the renormalized trace $\alpha^{-1}\tau_{\mathcal{M}}|_{p\mathcal{M}p}$. Let β be the trace-preserving automorphism of \mathcal{Q} obtained from Lemma 4.7 and satisfying $\beta(p) = p$, $\beta(\mathcal{N}) = \tilde{\mathcal{N}}$ and $\beta(\tilde{\mathcal{N}}) = \mathcal{N}$. Since A and $p\mathcal{Q}p$ are free, there is a trace-preserving automorphism $\tilde{\gamma}$ of $p\mathcal{M}p$ mapping A identically to itself and whose restriction to $p\mathcal{Q}p$ agrees with β .

We claim that $\tilde{\gamma}$ extends to an automorphism of \mathcal{M} whose restriction to \mathcal{Q} is β . Indeed, we may choose a maximal family $(v_i)_{i \in I}$ of partial isometries in \mathcal{Q} subject to the conditions

$$\begin{aligned} \forall i \quad v_i^* v_i \leq p \text{ and } v_i v_i^* \leq 1 - p, \\ i, j \in I, i \neq j \implies v_i v_i^* \perp v_j v_j^*. \end{aligned}$$

Then

$$\text{span} \left((1-p)\mathcal{Q}(1-p) \cup p\mathcal{M}p \cup \bigcup_{i \in I} (v_i p\mathcal{M}p \cup p\mathcal{M}p v_i^*) \right)$$

is dense in \mathcal{M} . Indeed, given $x \in \mathcal{M}$, we have

$$x = p x p + (1-p)x(1-p) + \sum_{i \in I} (v_i (v_i^* x p) + (p x v_i) v_i^*),$$

where the sum converges in s.o.t. We may now define

$$\gamma(x) = \tilde{\gamma}(p x p) + \beta((1-p)x(1-p)) + \sum_{i \in I} (\beta(v_i) \tilde{\gamma}(v_i^* x p) + \tilde{\gamma}(p x v_i) \beta(v_i^*)).$$

Using that $\tilde{\gamma}$ and β are trace-preserving automorphisms and that $\tilde{\gamma}|_{p\mathcal{Q}p} = \beta|_{p\mathcal{Q}p}$, we can show that γ is a trace-preserving automorphism of \mathcal{M} . It satisfies the desired properties by construction. \square

Proposition 4.9. *Let I be a finite or countable set and let $\alpha_i > 0$ satisfy $\sum_{i \in I} \alpha_i = 1$. For each $i \in I$, let B_i be a tracial von Neumann algebra. Let*

$$\mathcal{M} = \left(\bigoplus_{i \in I} \overset{p_i}{B_i} \right) * \left(*_{i \in I} \left(\overset{q_i}{\mathbb{C}} \oplus \mathbb{C} \right) \right)$$

and

$$\mathcal{P} = \left(\bigoplus_{i \in I} \overset{r_i}{\mathbb{C}} \right) * \left(*_{i \in I} \left(\overset{s_i}{B_i} \oplus \mathbb{C} \right) \right).$$

Then there is a normal, trace-preserving $$ -isomorphism $\pi : \mathcal{M} \rightarrow \mathcal{P}$ such that for all $i \in I$, $\pi(p_i) = r_i$.*

Proof. If $|I| = 1$, there is nothing to show, so assume $|I| \geq 2$. Let

$$\mathcal{Q} = \left(\bigoplus_{i \in I} \begin{matrix} r_i & \\ & B_i \\ \alpha_i & \end{matrix} \right) * \left(\bigoplus_{i \in I} \begin{matrix} s_i & \\ & \tilde{B}_i \oplus \mathbb{C} \\ \alpha_i & \end{matrix} \right),$$

where \tilde{B}_i is a copy of B_i . Let $\pi_0 : \mathcal{M} \rightarrow \mathcal{Q}$ be the trace-preserving, injective, normal $*$ -homomorphism sending $\bigoplus_{i \in I} B_i$ identically to itself and sending each projection q_i to s_i . Fixing $j \in I$, we may write

$$\bigoplus_{i \in I} \begin{matrix} r_i & & 1-r_j & & r_j \\ & B_i & & & \oplus B_j \\ \alpha_i & & 1-\alpha_j & & \alpha_j \end{matrix},$$

where

$$A_j = \bigoplus_{i \in I \setminus \{j\}} \begin{matrix} r_i \\ B_i \\ \alpha_i / (1-\alpha_j) \end{matrix}.$$

Consider the subalgebra

$$\mathcal{Q}_j = \left(\begin{matrix} 1-r_j & & r_j \\ A_j & \oplus & B_j \\ 1-\alpha_j & & \alpha_j \end{matrix} \right) * \left(\begin{matrix} s_j & \\ \tilde{B}_j \oplus \mathbb{C} \\ \alpha_j & 1-\alpha_j \end{matrix} \right)$$

of \mathcal{Q} . By Lemma 4.8, there is a trace-preserving automorphism $\tilde{\gamma}_j$ of \mathcal{Q}_j that maps $A_j \oplus \mathbb{C}$ identically to itself and such that, letting

$$\begin{aligned} \mathcal{N}_j &= \left(\mathbb{C} \oplus \begin{matrix} r_j \\ B_j \\ \alpha_j \end{matrix} \right) * \left(\begin{matrix} s_j \\ \mathbb{C} \oplus \mathbb{C} \\ \alpha_j \end{matrix} \right) \subseteq \mathcal{Q}_j \\ \tilde{\mathcal{N}}_j &= \left(\mathbb{C} \oplus \begin{matrix} r_j \\ \mathbb{C} \\ \alpha_j \end{matrix} \right) * \left(\begin{matrix} s_j \\ \tilde{B}_j \oplus \mathbb{C} \\ \alpha_j \end{matrix} \right) \subseteq \mathcal{Q}_j, \end{aligned}$$

we have $\tilde{\gamma}_j(\mathcal{N}_j) = \tilde{\mathcal{N}}_j$ and $\tilde{\gamma}_j(\tilde{\mathcal{N}}_j) = \mathcal{N}_j$. We extend $\tilde{\gamma}_j$ to an automorphism γ_j of \mathcal{Q} by taking the free product with the identity maps on all the $\tilde{B}_i \oplus \mathbb{C}$ with $i \neq j$. Note that, if $i \neq j$, then γ_j sends each of

$$\begin{matrix} 1-r_i & r_i \\ \mathbb{C} \oplus B_i & \\ 1-\alpha_i & \alpha_i \end{matrix} \quad \text{and} \quad \begin{matrix} s_i & 1-s_i \\ \tilde{B}_i \oplus \mathbb{C} & \\ \alpha_i & 1-\alpha_i \end{matrix}$$

identically to itself, so sends each of \mathcal{N}_i and $\tilde{\mathcal{N}}_i$ identically to itself, while $\gamma_j(r_j) = r_j$ and, of course, $\gamma_j(\mathcal{N}_j) = \tilde{\mathcal{N}}_j$ and $\gamma_j(\tilde{\mathcal{N}}_j) = \mathcal{N}_j$.

Let

$$\mathcal{M}_i = \left(\begin{matrix} p_i & 1-p_i \\ \tilde{B}_i \oplus \mathbb{C} & \\ \alpha_i & 1-\alpha_i \end{matrix} \right) * \left(\begin{matrix} q_i & 1-q_i \\ \mathbb{C} \oplus \mathbb{C} & \\ \alpha_i & 1-\alpha_i \end{matrix} \right) \subseteq \mathcal{M}.$$

Then

$$\mathcal{M} = W^* \left(\bigcup_{i \in I} \mathcal{M}_i \right).$$

Moreover, $\pi_0(\mathcal{M}_i) = \mathcal{N}_i$. Suppose that I is finite, say, $I = \{1, 2, \dots, n\}$. Let

$$\pi = \gamma_n \circ \gamma_{n-1} \circ \dots \circ \gamma_2 \circ \gamma_1 \circ \pi_0 : \mathcal{M} \rightarrow \mathcal{Q}.$$

Note that π is a trace-preserving, injective, normal $*$ -homomorphism. Furthermore, for each $j \in I$,

$$\begin{aligned}\pi(\mathcal{M}_j) &= \gamma_n \circ \gamma_{n-1} \circ \cdots \circ \gamma_2 \circ \gamma_1(\mathcal{N}_j) \\ &= \gamma_n \circ \gamma_{n-1} \circ \cdots \circ \gamma_j(\mathcal{N}_j) \\ &= \gamma_n \circ \gamma_{n-1} \circ \cdots \circ \gamma_{j+1}(\tilde{\mathcal{N}}_j) = \tilde{\mathcal{N}}_j.\end{aligned}\tag{7}$$

Thus,

$$\pi(\mathcal{M}) = W^*\left(\bigcup_{j=1}^n \pi(\mathcal{M}_j)\right) = W^*\left(\bigcup_{j=1}^n \tilde{\mathcal{N}}_j\right) = \mathcal{P},$$

and we are done in the case that I is finite.

Now suppose that I is infinite, say, $I = \{1, 2, \dots\}$. For each $n \geq 1$, let

$$\pi_n = \gamma_n \circ \gamma_{n-1} \circ \cdots \circ \gamma_2 \circ \gamma_1 \circ \pi_0 : \mathcal{M} \rightarrow \mathcal{Q}.$$

Of course, each π_n is an injective, trace-preserving, normal $*$ -homomorphism. Modifying the argument seen at equation (7), we get

$$\pi(\mathcal{M}_j) = \begin{cases} \mathcal{N}_j, & \text{if } n < j \\ \tilde{\mathcal{N}}_j, & \text{if } n \geq j \end{cases}$$

and, furthermore, since the restriction of γ_n to $\tilde{\mathcal{N}}_j$ is the identity map when $n > j$, we see that for all $x \in \bigcup_{j=1}^{\infty} \mathcal{M}_j$, the sequence $(\pi_n(x))_{n=1}^{\infty}$ is eventually constant. Since $\bigcup_{j=1}^{\infty} \mathcal{M}_j$ is dense in \mathcal{M} , we may define an injective, trace perservation, normal $*$ -homomorphism $\pi : \mathcal{M} \rightarrow \mathcal{Q}$ by

$$\pi(x) = \lim_{n \rightarrow \infty} \pi_n(x),$$

for all $x \in \mathcal{M}$, where the limit is in s.o.t. Now we have

$$\pi(\mathcal{M}) = W^*\left(\bigcup_{j=1}^{\infty} \pi(\mathcal{M}_j)\right) = W^*\left(\bigcup_{j=1}^{\infty} \tilde{\mathcal{N}}_j\right) = \mathcal{P}.$$

□

We are finally ready to prove the generalization of Theorem 4.4.

Theorem 4.10. *Fix a self-symmetric tracial von Neumann algebra A . Assume for each $i \in \mathbb{N}$, $s_i \in (0, \infty)$, $r_i \in (1 - s_i, \infty]$, and let $s = \sum_{i \in \mathbb{N}} s_i$. Then*

$$* \mathcal{F}_{s_i, r_i} \cong \mathcal{F}_{s, \infty}.$$

Proof. If $s = \infty$, then this was proved in Theorem 4.4, so let us assume $s < \infty$. Arguing as at the start of the proof of Theorem 4.4, we may without loss of generality assume $r_i = \infty$ for all i . Now, employing the rescaling formula (2), proved in Theorem 3.5, and Theorem 1.5 of [8] (directly for compression by s if $0 < s < 1$, or indirectly for compression by $1/s$ if $s > 1$), we have

$$\left(* \mathcal{F}_{s_i, \infty} \right)^s \cong * \mathcal{F}_{\frac{s_i}{s}, \infty},$$

while $(\mathcal{F}_{s,\infty})^s \cong \mathcal{F}_{1,\infty}$. Hence, we may without loss of generality assume $s = 1$. Using Proposition 4.2,

$$\mathcal{F}_{s_i,\infty} = (A \oplus_{s_i} \mathbb{C}) * L(\mathbb{F}_\infty).$$

Employing Proposition 4.9, we get

$$\begin{aligned} *_{i \in \mathbb{N}} \mathcal{F}_{s_i,r_i} &\cong *_{i \in \mathbb{N}} \left((A \oplus_{s_i} \mathbb{C}) * L(\mathbb{F}_\infty) \right) \\ &\cong \left(*_{i \in \mathbb{N}} (A \oplus_{s_i} \mathbb{C}) \right) * \left(\bigoplus_{i \in \mathbb{N}} \mathbb{C} \right) * L(\mathbb{F}_\infty) \\ &\cong \left(\bigoplus_{i \in \mathbb{N}} A \right) * \left(*_{i \in \mathbb{N}} (\mathbb{C} \oplus_{s_i} \mathbb{C}) \right) * L(\mathbb{F}_\infty) \\ &\cong A * L(\mathbb{F}_\infty) \cong \mathcal{F}_{1,\infty} \end{aligned} \tag{8}$$

where in the second last isomorphism above we used the self-symmetry property of A and Lemma 2.1. \square

We conclude our paper with the following theorem that uses the full power of Proposition 4.9.

Theorem 4.11. *Let I be a finite or countably infinite index set. Let $\{A_i\}_{i \in I}$ be self-symmetric tracial von Neumann algebras. For each $i \in I$, let $s_i \in (0, \infty)$, $r_i \in (1 - s_i, \infty]$. Assume $s = \sum_{i \in I} s_i < \infty$, and denote $r = \sum_{i \in I} r_i$,*

$$A = \bigoplus_{i \in I} A_i \oplus_{s_i/s}$$

Then

$$*_{i \in I} \mathcal{F}_{s_i,r_i}(A_i) \cong \mathcal{F}_{s,r}(A).$$

Proof. When I is infinite, $r = \infty$ necessarily. Then the result follows from the same proof as of Theorem 4.10, except in the series of isomorphisms (8), we replace each

$$A \oplus_{s_i} \text{ with } A_i \oplus_{s_i}$$

When I is finite, it suffices to show the case $|I| = 2$, and then the theorem follows by induction on $|I|$. In this case, A is

$$A_1 \oplus_t A_2 \oplus_{1-t}$$

where $t = s_1/s$. We realize $\mathcal{F}_{s,r}(A)$ by choosing $n \in \mathbb{N}$ large enough so that $(st/n)^{-2} - 1 > 1$, and that

$$(\mathbb{C} \oplus_t \mathbb{C})^{*n} \oplus_{1-t}$$

is a factor, in addition to

$$\mathcal{F}_{s,r}(A) \cong (A^{*n} * L\mathbb{F}_{\frac{s+r-1}{s^2}n^2-n+1})^{n/s}.$$

We use Theorem 1.5 of [8] and the rescaling formula (2):

$$\begin{aligned} & (\mathcal{F}_{st,r_1}(A_1) * \mathcal{F}_{s(1-t),r_2}(A_2))^{st/n} \\ & \cong (\mathcal{F}_{st,r_1}(A_1))^{st/n} * (\mathcal{F}_{s(1-t),r_2}(A_2))^{st/n} * L\mathbb{F}_{(st/n)^{-2}-1} \\ & \cong \mathcal{F}_{n,0}(A_1) * (\mathcal{F}_{n,1+\frac{2nt(1-t)-1}{(1-t)^2}}(A_2))^{\frac{t}{1-t}} * L\mathbb{F}_{t^{-2}(\frac{s+r-1}{s^2}n^2-n+1)}. \end{aligned}$$

Then we use the same technique as the proof of Lemma 3.8 to compute

$$\left(\begin{array}{c} (A_1 \oplus A_2)^{*n} \\ t \quad 1-t \end{array} \right)^t \cong A_1^{*n} * \left(\begin{array}{c} \mathbb{C} \oplus A_2 \\ t \quad 1-t \end{array} \right)^{*n}{}^t$$

and

$$\left(\begin{array}{c} \mathbb{C} \oplus A_2 \\ t \quad 1-t \end{array} \right)^{*n}{}^{1-t} \cong A_2^{*n} * \left(\begin{array}{c} \mathbb{C} \oplus \mathbb{C} \\ t \quad 1-t \end{array} \right)^{*n}{}^{1-t} \cong \mathcal{F}_{n,1+\frac{2nt(1-t)-1}{(1-t)^2}}(A_2).$$

The above two lines imply the following:

$$\left(\begin{array}{c} (A_1 \oplus A_2)^{*n} \\ t \quad 1-t \end{array} \right)^t \cong A_1^{*n} * \left(\mathcal{F}_{n,1+\frac{2nt(1-t)-1}{(1-t)^2}}(A_2) \right)^{\frac{t}{1-t}}.$$

Finally, using Theorem 1.5 of [8], we compute

$$\begin{aligned} (\mathcal{F}_{s,r}(A))^{st/n} & \cong (A^{*n} * L\mathbb{F}_{\frac{s+r-1}{s^2}n^2-n+1})^t \\ & \cong (A^{*n})^t * (L\mathbb{F}_{\frac{s+r-1}{s^2}n^2-n+1})^t * B_{t^{-2}-1} \\ & \cong (A^{*n})^t * L\mathbb{F}_{t^{-2}(\frac{s+r-1}{s^2}n^2-n+1)}, \end{aligned}$$

where $B_{t^{-2}-1}$ is a tracial von Neumann algebra having free dimension $t^{-2}-1$. Now putting them all together, we obtain

$$(\mathcal{F}_{s,r}(A))^{st/n} \cong (\mathcal{F}_{st,r_1}(A_1) * \mathcal{F}_{s(1-t),r_2}(A_2))^{st/n}.$$

Rescaling will then prove the case $|I| = 2$. \square

REFERENCES

- [1] C. Anantharaman and S. Popa, *An introduction to II_1 factors*, 2020. <https://www.math.ucla.edu/~popa/Books/IIunV15.pdf>.
- [2] D. P. Blecher, S. Goldstein, and L. E. Labuschagne, *Abelian von Neumann algebras, measure algebras and L^∞ -spaces*, *Expositiones Mathematicae* **40** (2022), no. 3, 758–818.
- [3] R. Boutonnet, D. Drimbe, A. Ioana, and S. Popa, *Non-isomorphism of A^{*n} , $2 \leq n \leq \infty$, for a non-separable abelian von Neumann algebra A* , *Geometric and Functional Analysis* **34** (2024), no. 2, 393–408.
- [4] R. Boutonnet, I. Chifan, and A. Ioana, *II_1 factors with non-isomorphic ultrapowers*, *Duke Mathematical Journal* **166** (2017), no. 11, 2023–2051.
- [5] K. Dykema, *Free products of hyperfinite von Neumann algebras and free dimension*, *Duke Mathematical Journal* **69** (1993), no. 1, 97–119.
- [6] ———, *On certain free product factors via an extended matrix model*, *Journal of Functional Analysis* **112** (1993), 31–60.
- [7] ———, *Interpolated free group factors*, *Pacific Journal of Mathematics* **163** (1994), no. 1, 123–135.
- [8] K. Dykema and F. Rădulescu, *Compressions of free products of von Neumann algebras*, *Math. Ann.* **316** (2000), no. 1, 61–82.

- [9] F. J. Murray and J. von Neumann, *On rings of operators. IV*, Ann. of Math. **44** (1943), 716–808.
- [10] F. Rădulescu, *Random matrices, amalgamated free products and subfactors of the von Neumann algebra of a free group, of noninteger index*, Invent. Math. **115** (1994), no. 2, 347–389.
- [11] D. Voiculescu, *Multiplication of certain noncommuting random variables*, J. Operator Theory **18** (1987), no. 2, 223–235.

KEN DYKEMA, DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TX 77843-3368, USA.

Email address: `kdykema@math.tamu.edu`

JUNCHEN ZHAO, DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TX 77843-3368, USA.

Email address: `junchenzhao@tamu.edu`