

# A POSITIVE HOMOGENEOUS FRACTIONAL HARMONIC FUNCTION IN A LIPSCHITZ CONE

CHILIN ZHANG

ABSTRACT. We construct a positive function  $u$  supported and solving  $(-\Delta)^s u = 0$  in a Lipschitz cone. Such a function is unique up to a constant multiplication. Moreover, we show that it is homogeneous of some degree  $0 < \alpha < 2s$ .

## 1. INTRODUCTION

Let  $\Sigma \subseteq \partial B_1$  be an open spherical domain with respect to the north pole  $e_n^+$ . Let  $Cone_\Sigma$  be the cone of  $\Sigma$  defined as

$$Cone_\Sigma = \{X \in \mathbb{R}^n \setminus \{0\} : \frac{X}{|X|} \in \Sigma\}.$$

If  $\partial Cone_\Sigma = \{x_n = g(x')\}$  for some  $g \in C^{0,1}$ , then we say  $Cone_\Sigma$  is a Lipschitz cone.

The purpose of this paper is to prove the following result.

**Theorem 1.1.** *Let  $0 < s < 1$  and fix a Lipschitz cone  $Cone_\Sigma$ . Then there exists a function  $u : \mathbb{R}^n \rightarrow \mathbb{R}$  such that:*

- (1)  $u > 0$  in  $Cone_\Sigma$ , and it can be continuously extended to zero elsewhere;
- (2)  $u \in \mathcal{L}_{2s}(\mathbb{R}^n)$ , and it is a classical solution to  $(-\Delta)^s u = 0$  inside  $Cone_\Sigma$ ;
- (3)  $u$  is a homogeneous function with degree  $0 < \alpha < 2s$ , and  $\max_{|X| \leq 1} u(X) = 1$ ;
- (4) Any other function satisfying (1) and (2) must be a constant multiple of  $u$ .

Here the non-local operator  $(-\Delta)^s$  is defined using the singular integration

$$(1.1) \quad (-\Delta)^s u(X) := C_{n,s} PV \int_{\mathbb{R}^n} \frac{u(X) - u(Y)}{|X - Y|^{n+2s}} dY.$$

If  $u$  is compactly supported (or has a fast decay), then the fractional Laplacian can be equivalently defined using the Caffarelli-Silvestre extension [2].

The problem is a natural extension of the spherical harmonic of the local operator. Suppose that  $\lambda_\Sigma$  is the first eigenvalue of  $\Sigma$  with respect to the Laplace-Beltrami operator

on the sphere, which can also be defined in the following variational way

$$(1.2) \quad \lambda_\Sigma = \inf_{F(\theta) \in C_0^\infty(\Sigma)} \frac{\int_\Sigma |\nabla F(\theta)|^2 d\theta}{\int_\Sigma |F(\theta)|^2 d\theta},$$

we then let  $\alpha_\Sigma > 0$  satisfying  $\alpha_\Sigma(\alpha_\Sigma + n - 2) = \lambda_\Sigma$ . If  $E(\theta) \in C^\infty(\Sigma) \cap C^0(\overline{\Sigma})$  is a minimizer of (1.2), then we can obtain a homogeneous positive harmonic function

$$H_\Sigma(X) = r^{\alpha_\Sigma} E(\theta), \quad (r, \theta) = (|X|, \frac{X}{|X|}).$$

It is very surprising that the homogeneous degree  $\alpha$  is strictly less than  $2s$  in Theorem 1.1. Recall that for the local operator  $-\Delta$  (so  $s = 1$ ), the homogeneous degree  $\alpha$  is not guaranteed to be less than  $2s = 2$ , since the positive homogeneous harmonic function in a  $\theta$  angle has degree  $\alpha = \pi/\theta$ . If  $\theta \leq \pi/2$ , we will have  $\alpha \geq 2$ . Therefore, property (3) in Theorem 1.1 is a special feature of the fractional Laplace equation.

The author thinks that it is also an interesting question to find other homogeneous solutions to  $(-\Delta)^s u = 0$  in  $Cone_\Sigma$  with a changing sign. Such solutions might not belong to the class  $\mathcal{L}_{2s}$ , but  $(-\Delta)^s u$  might still be defined in some weaker sense (for instance in the PV sense) because there will be significant cancellation in the expression (1.1).

Besides, we will later see in Example 2.1 that when  $Cone_\Sigma = \mathbb{R}_+^n$ ,  $u(X)$  obtained in Theorem 1.1 is exactly the well-known  $s$ -harmonic function  $u(X) = \max\{x_n, 0\}^s$ .

The strategy in proving Theorem 1.1 is to "reverse the logic" of the Caffarelli-Silvestre extension. We intend to find a homogeneous solution to the extension problem, which is a degenerate or singular equation with  $A_2$ -Muckenhoupt weight. Equations with a generic  $A_2$ -Muckenhoupt weight were initially studied in [5, 6]. Recently, higher order estimates were obtained in [3, 4, 8, 9] for weights with asymptotic behavior  $y^a$ . Such a degenerate or singular equation could also be treated as an equation with a  $d$ -regular weight, see [7].

## 2. PROOF OF THEOREM 1.1

The easier part in proving Theorem 1.1 is the uniqueness part (4) and the key is to use the boundary Harnack principle for fractional Laplacian obtained by Bogdan in [1].

*Proof of Theorem 1.1 (4).* Let  $v$  be another function satisfying (1)(2) in Theorem 1.1. We further assume that  $u(\vec{e}_n) = v(\vec{e}_n)$  by multiplying a constant to  $v(X)$ . Let  $R > 2$  be a

sufficiently large radius and we apply [1] to  $u$  and  $v$  in  $B_R$ . It then follows that in  $B_{R/2}$ ,

$$\left| \frac{u(X)/u(\frac{R}{2}\vec{e}_n)}{v(X)/v(\frac{R}{2}\vec{e}_n)} - L_R \right| \leq C \left| \frac{X}{R} \right|^\epsilon, \quad C^{-1} \leq L_R \leq C.$$

Here,  $C$  and  $\epsilon$  depend only on the shape of  $Cone_\Sigma$ . By plugging in  $X = \vec{e}_n$ , we have

$$\left| L_R - \frac{v(\frac{R}{2}\vec{e}_n)}{u(\frac{R}{2}\vec{e}_n)} \right| \leq CR^{-\epsilon}, \quad \text{and thus } \frac{u(\frac{R}{2}\vec{e}_n)}{v(\frac{R}{2}\vec{e}_n)} \leq \frac{1}{C^{-1} - CR^{-\epsilon}}.$$

Then for any  $X \in Cone_\Sigma$ , without loss of generality we assume  $|X| \leq 1$ , then as  $R \rightarrow \infty$ ,

$$\left| \frac{u(X)}{v(X)} - 1 \right| \leq \left| \frac{u(X)}{v(X)} - \frac{u(\frac{R}{2}\vec{e}_n)}{v(\frac{R}{2}\vec{e}_n)} L_R \right| + \left| \frac{u(\frac{R}{2}\vec{e}_n)}{v(\frac{R}{2}\vec{e}_n)} L_R - 1 \right| \leq \frac{2CR^{-\epsilon}}{C^{-1} - CR^{-\epsilon}} \rightarrow 0.$$

This proves the uniqueness part of Theorem 1.1.  $\square$

The harder part in proving Theorem 1.1 is the construction part. Our strategy is to reverse the logic in the Caffarelli-Silvestre extension. We first consider a higher dimensional half-space  $\mathbb{R}_+^{n+1}$ . A point in  $\mathbb{R}_+^{n+1}$  will be written as

$$Z = (X, y) \in \mathbb{R}^n \times \mathbb{R}_+.$$

The polar coordinate for a non-zero point is  $(r, \Theta)$  such that  $r = |Z|$ ,  $\Theta = \frac{Z}{|Z|} \in \mathbb{S}^n$ .

We intend to find a positive function  $U(Z) = r^\alpha \Phi(\Theta) : \overline{\mathbb{R}_+^{n+1}} \rightarrow \mathbb{R}$  such that

$$(2.1) \quad \begin{cases} \operatorname{div}(y^{1-2s} \nabla U(Z)) = 0 & \text{in } \mathbb{R}_+^{n+1} \\ U(Z) = 0 & \text{on } \mathbb{R}^n \setminus Cone_\Sigma \\ - \lim_{y \rightarrow 0^+} y^{1-2s} \partial_y U(X, y) = 0 & \text{for } X \in Cone_\Sigma \end{cases}$$

We will then let  $u(X)$  be the restriction of  $U(Z)$  on  $\mathbb{R}^n = \{y = 0\}$ . Since we have reversed the logic of [2] and  $u(X)$  has a faster growth rate, we need to eventually verify that it is a fractional harmonic function in  $Cone_\Sigma$ .

We state the following Proposition, and it will be proven in the next section.

**Proposition 2.1.** *The problem (2.1) has a positive  $\alpha$ -homogeneous solution  $U(Z) = r^\alpha \Phi(\Theta) : \overline{\mathbb{R}_+^{n+1}} \rightarrow \mathbb{R}$ , which is locally continuous. Moreover, we have  $0 < \alpha < 2s$ .*

In the present section we just assume its correctness and continue on the proof of Theorem 1.1. Without loss of generality, we can assume that the solution  $U(Z)$  given by Proposition 2.1 is normalized such that  $\max_{B_1 \cap Cone_\Sigma} U(Z) = 1$ .

*Proof of Theorem 1.1 (1)(3).* We just let  $u(X)$  be the restriction of  $U(Z)$  to  $\mathbb{R}^n$ , that

$$u(X) = U(X, 0), \quad X \in \mathbb{R}^n.$$

One then easily verifies  $u(X)$  satisfies (1)(3).  $\square$

We then show that  $u(X)$  satisfies (2). We first notice that as  $u(X)$  is homogeneous of degree  $0 < \alpha < 2s$ , we have  $u(X) \in \mathcal{L}_{2s}$ . We now let  $\eta$  be a smooth positive function supported in  $B_2 \subseteq \mathbb{R}^n$  so that it equals 1 in  $B_1$ . Consider the following cut-off of  $u(X)$ , which equals  $u(X)$  in  $B_R$  and vanishes outside  $B_{2R}$ :

$$u_R(X) = u(X) \cdot \eta\left(\frac{X}{R}\right).$$

It then follows that  $\lim_{R \rightarrow \infty} \|u_R - u\|_{\mathcal{L}_{2s}(\mathbb{R}^n)} = 0$ . For a fixed  $X_0 \in Cone_\Sigma$ , when  $R \geq |X_0|$  we have  $u(X_0) = u_R(X_0)$ , so

$$(-\Delta)^s u(X_0) = (-\Delta)^s u_R(X_0) + C_{n,s} \int_{B_R^c} \frac{-u(Y) + u_R(Y)}{|X_0 - Y|^{n+2s}} dY = (-\Delta)^s u_R(X_0) + o(1).$$

We let  $U_R(Z) : \overline{\mathbb{R}_+^{n+1}} \rightarrow \mathbb{R}$  be the Caffarelli-Silvestre extension of  $u_R(X)$ , such that

$$\begin{cases} \operatorname{div}(y^{1-2s} \nabla U_R(Z)) = 0 & \text{in } \mathbb{R}_+^{n+1} \\ U_R(X, 0) = u_R(X) & \text{on } \mathbb{R}^n \setminus Cone_\Sigma, \\ U_R(Z) \rightarrow 0 & \text{as } |Z| \rightarrow \infty \end{cases}$$

existence of which lies on  $u_R(X)$ 's being compactly supported. Moreover, for  $X_0 \in Cone_\Sigma$ ,

$$(-\Delta)^s u_R(X_0) = - \lim_{y \rightarrow 0^+} y^{1-2s} \partial_y U_R(X_0, y).$$

The proof of Theorem 1.1 (2) is then reduced to showing

$$(2.2) \quad \lim_{R \rightarrow \infty} \left\{ \lim_{y \rightarrow 0^+} y^{1-2s} \partial_y U_R(X_0, y) \right\} = 0.$$

*Proof of Theorem 1.1 (2).* As is discussed above, it suffices to show (2.2). Recall that for  $X_0 \in Cone_\Sigma$  we already have from Proposition 2.1 that  $\lim_{y \rightarrow 0^+} y^{1-2s} \partial_y U(X_0, y) = 0$ , so we consider  $V_R(Z) = U_R(Z) - U(Z)$  and thus

$$\lim_{y \rightarrow 0^+} y^{1-2s} \partial_y U_R(X_0, y) = \lim_{y \rightarrow 0^+} y^{1-2s} \partial_y V_R(X_0, y).$$

We list some properties of  $V_R(Z)$ :

- (a)  $V_R(X, 0) = 0$  for  $|X| \leq R$ ;
- (b)  $\operatorname{div}(y^{1-2s} V_R(Z)) = 0$  in  $\mathbb{R}_+^{n+1}$ ;
- (c)  $|V_R(Z)| \leq C_1 R^\alpha$  if  $|Z| \leq 2R$  for some universal  $C_1$ .

One can easily verify (a)(b), while (c) needs some explanation. In fact, on the one hand, we know  $|U(Z)| \leq CR^\alpha$  by the homogeneity. On the other hand, since  $u_R(X) \leq u(X) \cdot \chi_{\{|X| \leq 2R\}}$ , we know by the maximal principle that

$$\|U_R(Z)\|_{L^\infty(\mathbb{R}^{n+1})} \leq \|u_R(X)\|_{L^\infty(\mathbb{R}^n)} \leq CR^\alpha.$$

For a fixed  $X_0 \in \text{Cone}_\Sigma$  and a sufficiently large  $R$ , we consider the barrier

$$\beta(Z) = 4^s C_1 \left(1 + \frac{n}{2-2s}\right) R^{\alpha-2s} y^{2s} + 4C_1 R^{\alpha-2} \left(|X - X_0|^2 - \frac{n}{2-2s} y^2\right)$$

in the cylinder  $\mathcal{C}_{R/2}(X_0) = \{Z = (X, y) : |X - X_0| \leq R/2, 0 \leq y \leq R/2\}$ . One can verify that  $\text{div}(y^{1-2s} \nabla \beta(Z)) = 0$ , and  $\beta(Z) \geq |V_R(Z)|$  on  $\partial(\mathcal{C}_{R/2}(X_0))$ . Therefore,  $|V_R(Z)| \leq \beta(Z)$  in  $\mathcal{C}_{R/2}(X_0)$ . In particular, there exists a uniform  $C_2$  such that

$$|V_R(Z)| \leq C_2 R^{\alpha-2s} |Z - (X_0, 0)|^{2s} \quad \text{in } \mathcal{C}_{R/2}(X_0).$$

We then apply the interior Schauder estimate to  $\text{div}(y^{1-2s} V_R(Z)) = 0$  in each ball  $B_{y/2}(X_0, y) \subseteq \mathbb{R}_+^{n+1}$  (so that the equation is uniformly elliptic) and conclude that  $|\nabla V_R(X_0, y)| \leq C_3 R^{\alpha-2s} y^{2s-1}$ . We then have

$$\left| \lim_{y \rightarrow 0^+} y^{1-2s} \partial_y U_R(X_0, y) \right| = \left| \lim_{y \rightarrow 0^+} y^{1-2s} \partial_y V_R(X_0, y) \right| \leq C_3 R^{\alpha-2s}.$$

Recall that  $\alpha < 2s$ , so we send  $R \rightarrow \infty$  and get (2.2). □

**Example 2.1.** In fact, for the simplest cone  $\text{Cone}_\Sigma = \mathbb{R}_+^n$ , we would consider

$$U(Z) = \left( \frac{x_n + \sqrt{x_n^2 + y^2}}{2} \right)^s.$$

It can be easily verified that  $U(Z)$  satisfies (2.1). By restricting  $U(Z)$  to  $\mathbb{R}^n$ , we get  $u(X) = \max\{x_n, 0\}^s$ . This is the well-known  $s$ -harmonic function in a half space.

Now we have completed the proof of Theorem 1.1. However, we still owe the readers the proof of Proposition 2.1, and it will be proven in the next section.

### 3. PROOF OF PROPOSITION 2.1

**3.1. Setting up.** Let's now denote  $\Sigma^c \subseteq \mathbb{S}^{n-1}$  as the complement of  $\Sigma$  in  $\mathbb{S}^{n-1}$ , then  $\Sigma^c \cup \Sigma = \mathbb{S}^{n-1}$  forms the boundary of  $\mathbb{S}_+^n$ .

As we aim at finding a homogeneous  $U(Z) = r^\alpha \Phi(\Theta)$  solving  $\text{div}(y^{1-2s} \nabla U(Z)) = 0$ , we need to turn that into an equation satisfied by  $\Phi(\Theta)$  in  $\overline{\mathbb{S}_+^n}$ . We let

$$L\Phi(\Theta) = y^{\frac{1}{2}-s} \Delta_\Theta(y^{\frac{1}{2}-s} \Phi) - y^{\frac{1}{2}-s} \Phi \Delta_\Theta(y^{\frac{1}{2}-s}) = \text{div}_\Theta(y^{1-2s} \nabla_\Theta \Phi(\Theta)),$$

where  $\Delta_\Theta$  is the Laplace-Beltrami operator in  $\mathbb{S}^n$ . As  $y^{\frac{1}{2}-s}U$  is a homogeneous function of degree  $(\frac{1}{2} - s + \alpha)$ , we have

$$\begin{aligned} 0 &= \operatorname{div}(y^{1-2s}\nabla U(Z)) = y^{\frac{1}{2}-s}\Delta(y^{\frac{1}{2}-s}U) - y^{\frac{1}{2}-s}U\Delta(y^{\frac{1}{2}-s}) \\ &= (\frac{1}{2} - s + \alpha)(n - \frac{1}{2} - s + \alpha)\frac{y^{1-2s}U}{r^2} + \frac{y^{\frac{1}{2}-s}}{r^2}\Delta_\Theta(y^{\frac{1}{2}-s}U) - y^{\frac{1}{2}-s}U\Delta(y^{\frac{1}{2}-s}) \\ &= \Lambda\frac{y^{1-2s}U}{r^2} + \frac{y^{\frac{1}{2}-s}}{r^2}\Delta_\Theta(y^{\frac{1}{2}-s}U) + (\frac{1}{2} - s)(n - \frac{1}{2} - s)\frac{y^{1-2s}U}{r^2} - y^{\frac{1}{2}-s}U\Delta(y^{\frac{1}{2}-s}) \\ &= \Lambda\frac{y^{1-2s}U}{r^2} + \frac{y^{\frac{1}{2}-s}}{r^2}\Delta_\Theta(y^{\frac{1}{2}-s}U) - \frac{y^{\frac{1}{2}-s}U}{r^2}\Delta_\Theta(y^{\frac{1}{2}-s}), \quad \text{where } \Lambda = \alpha(n - 2s + \alpha). \end{aligned}$$

By setting  $r = 1$ , we require

$$(3.1) \quad L\Phi(\Theta) + \Lambda y^{1-2s}\Phi(\Theta) = 0 \quad \text{in } \mathbb{S}_+^n.$$

Moreover, to get (2.1), we also require

$$(3.2) \quad \lim_{y(\Theta) \rightarrow 0^+} y^{1-2s}\nabla_\Theta y \cdot \nabla_\Theta \Phi(\Theta) = 0 \quad \text{on } \Sigma,$$

$$(3.3) \quad \Phi(\Theta) = 0 \quad \text{on } \Sigma^c.$$

It then suffices to find such a pair  $(\Phi, \Lambda)$  satisfying (3.1)(3.2)(3.3). The strategy is to consider a variational problem in a weighted Sobolev space.

**Definition 3.1.** We denote  $\overline{\mathcal{S}} = H^1(\mathbb{S}_+^n, y^{1-2s}d\Theta)$  as the linear subspace of  $H_{loc}^1(\mathbb{S}_+^n)$  functions satisfying

$$\|\Psi\|_{H^1(\mathbb{S}_+^n, y^{1-2s}d\Theta)} = \left\{ \int_{\mathbb{S}_+^n} y^{1-2s}|\nabla_\Theta \Psi|^2 d\Theta + \int_{\mathbb{S}_+^n} y^{1-2s}\Psi^2 d\Theta \right\}^{1/2} < \infty.$$

We also denote  $\mathcal{S} \subseteq \overline{\mathcal{S}}$  as a subspace of  $C_{loc}^\infty(\mathbb{S}_+^n) \cap C(\overline{\mathbb{S}_+^n})$  functions inside  $\overline{\mathcal{S}}$ .

In fact, as the weighted measure  $y^{1-2s}d\Theta$  is a finite positive measure in  $\mathbb{S}_+^n$ , we can then apply Lusin's theorem. Precisely we mean that if a function  $f$  satisfies

$$\|f\|_{L^2(\mathbb{S}_+^n, y^{1-2s}d\Theta)} = \left\{ \int_{\mathbb{S}_+^n} y^{1-2s}f^2 d\Theta \right\}^{1/2} < \infty,$$

then  $f$  can be approximated by a sequence of  $C(\overline{\mathbb{S}_+^n})$  functions in the  $L^2(\mathbb{S}_+^n, y^{1-2s}d\Theta)$  sense, see [7]. We conclude the following density result.

**Lemma 3.1.**  $\mathcal{S}$  is dense in  $\overline{\mathcal{S}}$  with respect to the  $H^1(\mathbb{S}_+^n, y^{1-2s}d\Theta)$  norm.

*Proof.* Let's parametrize  $\mathbb{S}_+^n$  using the parameters  $(\vec{\theta}, \gamma) \in \mathbb{S}^{n-1} \times (0, \frac{\pi}{2})$ , that

$$\Theta(\vec{\theta}, \gamma) = \sin \gamma \cdot \vec{\theta} + \cos \gamma \cdot e_{n+1}^{\vec{}}.$$

We would then like to map  $\mathbb{S}_+^n$  to itself via the map

$$F_\epsilon(\vec{\theta}, \gamma) = \sin[(1-2\epsilon)\gamma] \cdot \vec{\theta} + \cos[(1-2\epsilon)\gamma] \cdot e_{n+1} = \Theta(\vec{\theta}, (1-2\epsilon)\gamma).$$

For any  $X, Y \in \mathbb{S}^n$ , we let  $\eta_{\epsilon, Y}(X) \geq 0$  be a smooth mollifier on  $\mathbb{S}^n$  such that

- (1)  $\eta_{\epsilon, Y}(X)$  depends only on  $|X - Y|$ ,  $\eta_{\epsilon, Y} > 0$  if and only if  $|X - Y| \leq \sin(\frac{\epsilon\pi}{2})$ ;
- (2) For each fixed  $Y \in \mathbb{S}^n$ ,  $\int_{X \in \mathbb{S}^n} \eta_{\epsilon, Y}(X) dX = 1$ ;
- (3) There is a constant  $C_1$  independent of  $\epsilon$ , such that  $|\nabla \eta_{\epsilon, Y}| \leq \frac{C_1}{\epsilon^{n+1}}$ .

Let  $\Psi \in \overline{\mathcal{S}}$ , we then consider the following approximation

$$(3.4) \quad \Psi_\epsilon(\Theta) = (\Psi \circ F_\epsilon) * \eta_\epsilon := \int_{X \in \mathbb{S}^n} \Psi(X) \eta_{\epsilon, F_\epsilon(\Theta)}(X) dX.$$

Notice that  $X \in \mathbb{S}_+^n$  when  $\eta_{\epsilon, F_\epsilon(\Theta)}(X) > 0$ , so (3.4) is a well-defined expression for any  $\Theta \in \mathbb{S}_+^n$ . Next we show the convergence of  $\Psi_\epsilon$  to  $\Psi$ . We have

$$\|\Psi_\epsilon - \Psi\|_{H^1(\mathbb{S}_+^n, y^{1-2s} d\Theta)} \leq \|\Psi_\epsilon - \Psi \circ F_\epsilon\|_{H^1(\mathbb{S}_+^n, y^{1-2s} d\Theta)} + \|\Psi \circ F_\epsilon - \Psi\|_{H^1(\mathbb{S}_+^n, y^{1-2s} d\Theta)}.$$

By Lusin's theorem, both  $\Psi$  and  $\nabla \Psi$  can be approximated by  $C(\overline{\mathbb{S}_+^n})$  functions in the  $L^2(\mathbb{S}_+^n, y^{1-2s} d\Theta)$  sense, and thus so do  $\Psi \circ F_\epsilon$  and  $\nabla(\Psi \circ F_\epsilon)$ . These guarantee the smallness of  $\|\Psi_\epsilon - \Psi\|_{H^1(\mathbb{S}_+^n, y^{1-2s} d\Theta)}$ .  $\square$

In the weak sense, we would rephrase (3.1)(3.2)(3.3) as follows.

**Definition 3.2.** Let  $\Psi \in \overline{\mathcal{S}}$ . We say  $\Psi$  satisfies (3.3) in the trace sense, denoted by  $\Psi \in \overline{\mathcal{S}}_\Sigma$ , if it can be approximated by  $\Psi_k \in \mathcal{S}$  vanishing on  $\Sigma^c$ , under the distance of the  $H^1(\mathbb{S}_+^n, y^{1-2s} d\Theta)$  norm.

**Remark 3.1.** One can verify that  $\overline{\mathcal{S}}_\Sigma$  with the  $H^1(\mathbb{S}_+^n, y^{1-2s} d\Theta)$  norm is a Hilbert space.

**Definition 3.3.** Assume that  $\Phi \in \overline{\mathcal{S}}_\Sigma$ . Then we say  $\Phi$  satisfies (3.1)(3.2) in the weak sense, if for any  $\eta \in \overline{\mathcal{S}}$  (or just  $\eta \in \mathcal{S}$ ) satisfying (3.3), we have

$$\int_{\mathbb{S}_+^n} y^{1-2s} \nabla \Phi \cdot \nabla \eta d\Theta = \Lambda \int_{\mathbb{S}_+^n} y^{1-2s} \Phi \cdot \eta d\Theta.$$

One can easily verify that a weak solution of (3.1)(3.2)(3.3) can generate a homogeneous solution of (2.1) in the weak sense.

**Lemma 3.2.** Let  $U(Z) = r^\alpha \Phi(\Theta)$  and  $\Lambda = \alpha(n-2s+\alpha)$ . Then (3.1)(3.2)(3.3) can imply (2.1) in the weak sense. Moreover, if  $0 < \Lambda < 2ns$ , then  $0 < \alpha < 2s$ .

Now we define  $J(\Psi)$  and  $I(\Psi)$  which take value in  $[0, \infty]$ , such that

$$J(\Psi) = \int_{\mathbb{S}_+^n} y^{1-2s} |\nabla_{\Theta} \Psi|^2 d\Theta, \quad I(\Psi) = \int_{\mathbb{S}_+^n} y^{1-2s} \Psi^2 d\Theta.$$

**Lemma 3.3.** *Assume that some  $\Phi \in \overline{\mathcal{S}}_{\Sigma}$  minimizes  $J(\Psi)/I(\Psi)$  among all  $\Psi \in \overline{\mathcal{S}}_{\Sigma}$ , and  $J(\Phi)/I(\Phi) = \Lambda$ . Then the pair  $(\Phi, \Lambda)$  satisfies (3.1) and (3.2) in the weak sense.*

*Proof.* Let  $\eta \in \mathcal{S}$  be a test function satisfying (3.3). We let  $\Psi_t = \Phi + t\eta$ , then

$$f(t) = J(\Psi_t) - \Lambda \cdot I(\Psi_t) \geq 0, \quad f(0) = 0.$$

We then have  $f'(0) = 0$ , or in other words,

$$\int_{\mathbb{S}_+^n} y^{1-2s} \nabla \Phi \cdot \nabla \eta d\Theta - \Lambda \int_{\mathbb{S}_+^n} y^{1-2s} \Phi \cdot \eta d\Theta = 0.$$

By arbitrarily choosing  $\eta$ , we see  $(\Phi, \Lambda)$  satisfies (3.1) and (3.2).  $\square$

**3.2. Solving the variational problem.** Now let's prove that  $J(\Psi)/I(\Psi)$  has a minimizer in  $\overline{\mathcal{S}}_{\Sigma}$ . We first show that such a ratio could be less than  $2ns$ .

**Lemma 3.4.** *We can guarantee that  $\inf_{\Psi \in \overline{\mathcal{S}}_{\Sigma}} \frac{J(\Psi)}{I(\Psi)} < 2ns$ .*

*Proof.* We consider  $\Psi_0(\Theta) = y^{2s} \in \mathcal{S}$  vanishing on  $\partial\mathbb{S}_+^n$ , then  $J(\Psi_0)/I(\Psi_0) = 2ns$ . Moreover, as  $\Psi_0$  does not satisfy (3.2) (even in the weak sense), we see from Lemma 3.3 that  $\Psi_0$  is not a minimizer. Therefore, the infimum of  $J(\Psi)/I(\Psi)$  can be even smaller.  $\square$

Next, such a ratio must be bounded away from zero.

**Lemma 3.5.** *For every  $\Psi \in \overline{\mathcal{S}}_{\Sigma}$ ,  $J(\Psi)/I(\Psi) \geq c$  for some  $c > 0$ .*

*Proof.* Let  $\mathcal{D}$  be a small spherical disc inside  $\Sigma^c$ , then we see  $\Psi = 0$  on  $\mathcal{D}$ . For simplicity, let's assume that after a rotation, there exists some  $t > 1$  such that

$$\mathcal{D} = \mathbb{S}^{n-1} \cap \{|x'|^2 + |x_n + t|^2 \leq t^2 - 1\}.$$

Then we would like to parametrize  $\mathbb{S}_+^n$  by two parameters  $(\vec{\beta}, \gamma)$  with

$$\vec{\beta} \in \mathbb{S}^{n-1}, \quad \angle(\vec{\beta}, \vec{e}_n) < \sin^{-1}\left(\frac{1}{t}\right), \quad \gamma \in (0, \pi),$$



such that  $X \in \mathbb{S}_+^n$  can be expressed as

$$\begin{aligned} X(\vec{\beta}, \gamma) &= -t\vec{e}_n + \left\{ \sqrt{1 - (t \sin \angle(\vec{\beta}, \vec{e}_n))^2} \sin \gamma \right\} \vec{e}_{n+1} \\ &\quad + \left\{ t \cos \angle(\vec{\beta}, \vec{e}_n) - \sqrt{1 - (t \sin \angle(\vec{\beta}, \vec{e}_n))^2} \cos \gamma \right\} \vec{\beta}. \end{aligned}$$

Let  $\Psi(\vec{\beta}, \gamma) = \Psi(X(\vec{\beta}, \gamma))$ , by integrating

$$\begin{aligned} \Psi(\vec{\beta}, \gamma)^2 &\leq \left\{ \int_0^\gamma \left| \frac{\partial \Psi(\vec{\beta}, t)}{\partial \gamma} \right| dt \right\}^2 \leq \left\{ \int_0^\gamma (\sin t)^{2s-1} dt \right\} \left\{ \int_0^\gamma (\sin t)^{1-2s} \left| \frac{\partial \Psi(\vec{\beta}, t)}{\partial \gamma} \right|^2 dt \right\} \\ &\leq C \left\{ \int_0^\pi (\sin \gamma)^{1-2s} \left| \frac{\partial \Psi(\vec{\beta}, \gamma)}{\partial \gamma} \right|^2 d\gamma \right\}, \end{aligned}$$

we see that there exists some positive constant  $c$  independent of  $\beta$ , such that

$$(3.5) \quad \int_0^\pi (\sin \gamma)^{1-2s} \left| \frac{\partial \Psi(\vec{\beta}, \gamma)}{\partial \gamma} \right|^2 d\gamma \geq c \int_0^\pi (\sin \gamma)^{1-2s} \Psi(\vec{\beta}, \gamma)^2 d\gamma.$$

Besides, there exists a density function  $\mu(\vec{\beta}, \gamma)$ , such that the area measure  $d\Theta$  on  $\mathbb{S}_+^n$  can be expressed as  $d\Theta = \mu(\vec{\beta}, \gamma) d\vec{\beta} d\gamma$ . Moreover, there exists a constant  $C = C(t)$  depending only on  $t$ , so that for any  $\vec{\beta}$  we have

$$(3.6) \quad \sup_{\gamma \in (0, \pi)} \mu(\vec{\beta}, \gamma) \leq C \inf_{\gamma \in (0, \pi)} \mu(\vec{\beta}, \gamma).$$

Notice that  $\left| \nabla \Psi(\vec{\beta}, \gamma) \right| \geq \left| \frac{\partial \Psi(\vec{\beta}, \gamma)}{\partial \gamma} \right|$  for any  $\vec{\beta}$ , then we have

$$\begin{aligned} J(\Psi) &= \int_{\angle(\vec{\beta}, \vec{e}_n) < \sin^{-1}(\frac{1}{t})} \int_{\gamma \in (0, \pi)} (\sin \gamma)^{1-2s} \left| \nabla \Psi(\vec{\beta}, \gamma) \right|^2 \mu(\vec{\beta}, \gamma) d\vec{\beta} d\gamma \\ &\geq \int_{\angle(\vec{\beta}, \vec{e}_n) < \sin^{-1}(\frac{1}{t})} \left\{ \int_{\gamma \in (0, \pi)} (\sin \gamma)^{1-2s} \left| \frac{\partial \Psi(\vec{\beta}, \gamma)}{\partial \gamma} \right|^2 \mu(\vec{\beta}, \gamma) d\vec{\beta} \right\} d\gamma \\ &\gtrsim \int_{\angle(\vec{\beta}, \vec{e}_n) < \sin^{-1}(\frac{1}{t})} \left\{ \int_{\gamma \in (0, \pi)} (\sin \gamma)^{1-2s} \Psi(\vec{\beta}, \gamma)^2 \mu(\vec{\beta}, \gamma) d\vec{\beta} \right\} d\gamma = I(\Psi). \end{aligned}$$

Here, the third line is obtained from the second line by using (3.5) and (3.6).  $\square$

We have a Sobolev inequality for the weighted measure  $y^{1-2s} d\Theta$  on  $\mathbb{S}_+^n$ .

**Lemma 3.6.** *There exist constants  $q = q(n, s) > 2$  and  $C = C(n, s, \Sigma)$ , such that*

$$\left\{ \int_{\mathbb{S}_+^n} y^{1-2s} \Psi^q d\Theta \right\}^{1/q} \leq C \left\{ \int_{\mathbb{S}_+^n} y^{1-2s} |\nabla \Psi|^2 d\Theta \right\}^{1/2} \quad \text{for any } \Psi \in \overline{\mathcal{S}}_\Sigma.$$

*Proof.* It was introduced in [7] the terminology of  $d$ -regularity for a bounded metric space equipped with a measure. We treat  $\mathbb{S}_+^n$  as a metric space with the distance function generated by geodesics, and equip it with the measure  $\mu = y^{1-2s}d\Theta$ . Then  $\mathbb{S}_+^n$  has finite diameter (less than  $\pi$ ) and finite total measure, meaning  $\mu(\mathbb{S}_+^n) < \infty$ .

For any  $\Theta \in \mathbb{S}_+^n$  and any  $r \leq \pi$ , we see the geodesic ball of radius  $r$  centered at  $\Theta$ , denoted as  $\mathcal{B}_r(\Theta)$ , has measure  $\mu(\mathcal{B}_r(\Theta))$  greater than  $c_1 \cdot r^d$  with  $d = \max\{n+1-2s, n\} \geq 2$ . Moreover, the measure  $\mu$  has a doubling property, that  $\mu(\mathcal{B}_{2r}(\Theta)) \leq C \cdot \mu(\mathcal{B}_r(\Theta))$ .

In [7], the gradient of  $\Psi(x)$  is defined as  $g(x) \geq 0$ , such that

$$|\Psi(x) - \Psi(y)| \leq (g(x) + g(y)) \cdot \text{dist}(x, y)$$

for two points  $x, y$  in the metric space. In our situation,  $g(x)$  can be defined as the Hardy-Littlewood maximal function of  $|\nabla\Psi|$ .

By [7, Theorem 6], there exists some  $q > 2$ , such that

$$\|\Psi - \Psi_{avg}\|_{L^q(\mathbb{S}_+^n, y^{1-2s}d\Theta)} \leq C_2 \|\Psi\|_{H^1(\mathbb{S}_+^n, y^{1-2s}d\Theta)},$$

where

$$|\Psi_{avg}| = \frac{|\int_{\mathbb{S}_+^n} y^{1-2s}\Psi d\Theta|}{\int_{\mathbb{S}_+^n} y^{1-2s}d\Theta} \leq C_3 \|\Psi\|_{L^2(\mathbb{S}_+^n, y^{1-2s}d\Theta)} \leq C_3 \|\Psi\|_{H^1(\mathbb{S}_+^n, y^{1-2s}d\Theta)}.$$

Moreover, by Lemma 3.5, we see for  $\Psi \in \overline{\mathcal{S}}_\Sigma$ ,

$$\|\Psi\|_{H^1(\mathbb{S}_+^n, y^{1-2s}d\Theta)} \leq C_4 \left\{ \int_{\mathbb{S}_+^n} y^{1-2s} |\nabla\Psi|^2 d\Theta \right\}^{1/2}.$$

Therefore, we can bound  $\|\Psi\|_{L^q(\mathbb{S}_+^n, y^{1-2s}d\Theta)}$  using  $\int_{\mathbb{S}_+^n} y^{1-2s} |\nabla\Psi|^2 d\Theta$ . □

We immediately obtain the following corollary using the Hölder inequality.

**Corollary 3.1.** *Assume that  $\Psi \in \overline{\mathcal{S}}_\Sigma$  and  $J(\Psi) \leq 1$ . Then for any  $\epsilon \leq 1/2$  we have*

$$\int_{\mathbb{S}_+^n \cap \{y \leq \epsilon\}} y^{1-2s} \Psi^2 d\Theta \leq C \epsilon^{2(1-s)(1-\frac{2}{q})} = o(1).$$

Therefore, we can prove the existence of a minimizer of  $J(\Psi)/I(\Psi)$  in  $\overline{\mathcal{S}}_\Sigma$ .

**Lemma 3.7.** *There exists some  $\Phi \in \overline{\mathcal{S}}_\Sigma$  minimizing the ratio  $J(\Psi)/I(\Psi)$ .*

*Proof.* Let's just write  $\Lambda = \inf_{\Psi \in \overline{\mathcal{S}}_\Sigma} \frac{J(\Psi)}{I(\Psi)}$ . Let  $\Psi_k \in \overline{\mathcal{S}}_\Sigma$  be a minimizing sequence so that

$$I(\Psi_k) = 1, \quad \Lambda \leq J(\Psi_k) \leq \Lambda + \frac{1}{k}.$$

In the argument below, each time we find a subsequence of  $\Psi_k$ , we still write it as  $\Psi_k$ .

- Step 1: We can first find a subsequence  $\Psi_k$  converging weakly to  $\Phi \in \overline{\mathcal{S}}_\Sigma$  in the  $H^1(\mathbb{S}_+^n, y^{1-2s}d\Theta)$  sense. This implies that

$$I(\Phi) \leq \liminf_{k \rightarrow \infty} I(\Psi_k) \leq 1, \quad J(\Phi) \leq \liminf_{k \rightarrow \infty} J(\Psi_k) \leq \Lambda.$$

- Step 2: For each  $\epsilon$ , we can apply the classical Rellich compactness theorem, and find a subsequence of  $\Psi_k$  converging in  $L^2(\mathbb{S}_+^n \cap \{y \geq \epsilon\}, y^{1-2s}d\Theta)$ . Its limit could be nothing but  $\Phi$  because it is the weak limit of  $\Psi_k$ . By a diagonal argument, there is a subsequence  $\Psi_k$ , such that for any fixed  $\epsilon > 0$ ,

$$\lim_{k \rightarrow \infty} \|\Psi_k - \Phi\|_{L^2(\mathbb{S}_+^n \cap \{y \geq \epsilon\}, y^{1-2s}d\Theta)} = 0.$$

For any given  $\delta$ , we can find  $\epsilon$  sufficiently small so that by Corollary 3.1,

$$\|\Psi_k\|_{L^2(\mathbb{S}_+^n \cap \{y \leq \epsilon\}, y^{1-2s}d\Theta)}, \|\Phi\|_{L^2(\mathbb{S}_+^n \cap \{y \leq \epsilon\}, y^{1-2s}d\Theta)} \leq \delta/3.$$

We let  $k$  sufficiently large so that  $\|\Psi_k - \Phi\|_{L^2(\mathbb{S}_+^n \cap \{y \geq \epsilon\}, y^{1-2s}d\Theta)} \leq \delta/3$ , then

$$\|\Psi_k - \Phi\|_{L^2(\mathbb{S}_+^n, y^{1-2s}d\Theta)} \leq \|\Psi_k - \Phi\|_{L^2(\mathbb{S}_+^n \cap \{y \geq \epsilon\}, y^{1-2s}d\Theta)} + \|\Psi_k - \Phi\|_{L^2(\mathbb{S}_+^n \cap \{y \leq \epsilon\}, y^{1-2s}d\Theta)} \leq \delta.$$

This means  $\|\Psi_k - \Phi\|_{L^2(\mathbb{S}_+^n, y^{1-2s}d\Theta)} \rightarrow 0$ , so  $I(\Phi) = 1$ . We then have  $J(\Phi) = \Lambda$  since  $J(\Psi)/I(\Psi) \geq \Lambda$  for all  $\Psi \in \overline{\mathcal{S}}_\Sigma$ . Therefore,  $\Phi \in \overline{\mathcal{S}}_\Sigma$  is the desired minimizer and we have completed the proof of Lemma 3.7.  $\square$

Moreover, by Lemma 3.2 and Lemma 3.3,  $U(Z) = r^\alpha \Phi(\Theta)$  satisfies (2.1) in the weak sense. One can also easily verify that  $U(Z)$  does not change sign in  $\mathbb{R}_+^{n+1}$ , and

$$(3.7) \quad \int_{\mathbb{R}_+^{n+1} \cap B_4} y^{1-2s} \left( |\nabla U(Z)|^2 + U(Z)^2 \right) dZ \leq C.$$

**3.3. Continuity estimates.** In the previous subsection, we have found a minimizer  $\Phi$ , and  $U(Z) = r^\alpha \Phi(\Theta)$  is a weak solution to (2.1). In this part, we prove that  $U(Z)$  is in fact a continuous classical solution to (2.1), thus finishing the proof of Proposition 2.1.

By the homogeneity of  $U(Z)$ , it suffices to consider  $Z^* = (X^*, y^*) \in \overline{\mathbb{R}_+^{n+1}} \cap B_1$  and show that  $U(Z)$  satisfies (2.1) in the classical sense near  $Z^*$ .

Let's first consider the simplest case where  $y^* > 0$ , then (2.1) is uniformly elliptic in  $B_{y^*/2}(Z^*) \subseteq \mathbb{R}_+^{n+1} \cap B_4$ . By the standard elliptic theory, we see  $U$  is  $C^{2,\alpha}$  in  $B_{y^*/4}(Z^*)$ . Here we have used (3.7) to bound the  $L^2$  norm of  $U$  in  $B_{y^*/2}(Z^*)$ . In other words, we see  $U(Z)$  satisfies the first line of (2.1) in  $\mathbb{R}_+^{n+1}$  in the classical sense.

Next, we consider the case where  $y^* = 0$  and  $X^* \in Cone_\Sigma$ . We consider a half ball  $B_\epsilon(Z^*)^+ = \mathbb{R}_+^{n+1} \cap B_\epsilon(Z^*)$ , where  $\epsilon$  is so small that  $B_\epsilon(X^*)' \subseteq Cone_\Sigma$ . Then the second line of (2.1) is in fact useless in  $B_\epsilon(Z^*)^+$ . Besides,  $\int_{B_\epsilon(Z^*)^+} y^{1-2s} U(Z)^2 dZ$  is bounded because of (3.7). We then apply [8, Theorem 1.6] and conclude the  $C^{1,\alpha}$  regularity of  $U$  near  $Z^*$ . This implies that the third line of (2.1) holds at  $Z^*$  in the classical sense.

We can argue similarly for the case where  $y^* = 0$  and  $X^* \in int(\mathbb{R}^n \setminus Cone_\Sigma)$ . This time we can apply [6, Theorem 2.4.6] or [9, Theorem 1.6] because the weight  $y^{1-2s} dZ$  is  $A_2$ -Muckenhoupt, and  $U$  vanishes in  $\partial\mathbb{R}_+^{n+1}$  near  $X^*$  in the weak sense. We see  $U(Z)$  is locally Hölder continuous near  $Z^*$  and  $U(Z^*) = 0$  in the classical sense.

By discussion above, the remaining job in proving Proposition 2.1 is to show

$$(3.8) \quad \lim_{r \rightarrow 0} \sup_{B_r(Z^*)^+} |U| = 0$$

for a fixed  $Z^* = (X^*, y^*)$  satisfying  $y^* = 0$  and  $X^* \in \partial(Cone_\Sigma)$ . To achieve this, we first prove the following  $L^\infty$  estimate using the De Giorgi iteration.

**Lemma 3.8.** *Assume that  $U(Z)$  is a weak solution to (2.1) in  $B_4^+ = \mathbb{R}_+^{n+1} \cap B_4$  and that  $\int_{B_4^+} y^{1-2s} U^2 dZ$  is sufficiently small, then  $\|U\|_{L^\infty(B_2^+)} \leq 1$ .*

*Proof.* Let's assume  $U \geq 0$  for simplicity because it does not change sign. We denote  $(U - t)_+ = \max\{U - t, 0\}$  for any  $t \geq 0$ , then it satisfies the second and the third line of (2.1) in the weak sense, and satisfies  $\operatorname{div}(y^{1-2s} \nabla(U - t)_+) \geq 0$  in the weak sense.

We first prove a Caccioppoli inequality. For any smooth  $\eta \geq 0$  supported in  $B_4$ ,

$$(3.9) \quad \int_{B_4^+} y^{1-2s} |\nabla(\eta(U - t)_+)|^2 dZ \leq \int_{B_4^+} y^{1-2s} (U - t)_+^2 |\nabla\eta|^2 dZ.$$

In fact, we multiply  $\eta^2(U - t)_+$  on both sides of  $\operatorname{div}(y^{1-2s} \nabla(U - t)_+) \geq 0$ . Notice that by the second and the third line of (2.1), no boundary term is produced during the integration

by parts. Precisely we have

$$\int_{B_4^+} y^{1-2s} (|\nabla(\eta(U-t)_+)|^2 - (U-t)_+^2 |\nabla\eta|^2) dZ = \int_{B_4^+} y^{1-2s} \nabla(U-t)_+ \cdot \nabla(\eta^2(U-t)_+) dZ \leq 0.$$

We next apply [6, Theorem 1.2] to the weight  $y^{1-2s} dZ$ , which is  $A_2$ -Muckenhoupt, and obtain a Sobolev inequality. If  $V(Z) = 0$  for all  $Z \geq 4$ , then

$$\left( \int_{B_4^+} y^{1-2s} V(Z)^p dZ \right)^{1/p} \leq C \left( \int_{B_4^+} y^{1-2s} |\nabla V(Z)|^2 dZ \right)^{1/2} \quad \text{for some } p > 2.$$

In the last step we prove  $U \leq 1$  in  $B_2^+$ . Denote  $t_k = 1 - 2^{-k}$  and  $r_k = 2 + 2^{-k}$  and let  $U_k = (U - t_k)_+$ . We consider a smooth cut-off function  $\eta_k \geq 0$  so that

$$\text{supp}(\eta_k) = B_{r_k}^+, \quad \eta_k \Big|_{B_{r_{k+1}}^+} = 1, \quad |\nabla\eta_k| \leq C2^k.$$

Let's denote  $A_k = \int_{B_{r_k}^+} y^{1-2s} U_k^2 dZ$ , then it follows from (3.9) that

$$\int_{B_{r_k}^+} y^{1-2s} |\nabla(\eta_k U_k)|^2 dZ \leq C4^k \int_{B_{r_k}^+} y^{1-2s} U_k^2 dZ = C4^k A_k.$$

We apply the Sobolev inequality to  $\eta_k U_k$ , then apply the Hölder inequality and obtain

$$\int_{B_{r_{k+1}}^+} y^{1-2s} U_k^2 dZ \leq \int_{B_{r_k}^+} y^{1-2s} \eta_k^2 U_k^2 dZ \leq C4^k A_k \left( \int_{B_{r_k}^+} y^{1-2s} \chi_{\{U_k > 0\}} dZ \right)^{\frac{p-2}{p}}.$$

As  $U_{k+1} \leq U_k$ , we have  $\int_{B_{r_{k+1}}^+} y^{1-2s} U_k^2 \geq A_{k+1}$ . On the other hand, we first notice that as  $\{U_k > 0\} = \{U_{k-1} > t_k - t_{k-1} = 2^{-k}\}$ , we have

$$A_{k-1} \geq \int_{B_{r_k}^+} y^{1-2s} U_{k-1}^2 dZ \geq 4^{-k} \int_{B_{r_k}^+} y^{1-2s} \chi_{\{U_k > 0\}} dZ.$$

Now we have an inductive inequality  $A_{k+1} \leq C16^k A_k A_{k-1}^{\frac{p-2}{p}}$ . Taking the logarithm on both sides, we know  $A_k \rightarrow 0$  as the initial data  $A_1, A_2$  is small. This means  $U \leq 1$  in  $B_2^+$ .  $\square$

We can now prove Equation (3.8) using the weak Harnack principle.

*Proof of Equation (3.8).* Once more we write  $Z^* = (X^*, 0)$  for  $X^* \in \partial\text{Cone}_\Sigma$ . Without loss of generality, we assume  $\sup_{B_1(Z^*)^+} U \leq 1$  using Lemma 3.8. The key is to show that

$$(3.10) \quad \sup_{B_{1/2}(Z^*)^+} U \leq \sigma \quad \text{for some } \sigma < 1.$$

Then after an iterative argument, we have  $\sup_{B_{2^{-i}}(Z^*)^+} U \leq \sigma^i \rightarrow 0$ .

We let  $t_k = 1 - 2^{-k}$  and  $U_k = 2^k(U - t_k)_+$ . We need to prove

$$(3.11) \quad \int_{Z \in B_1(Z^*)^+} y^{1-2s} \chi_{\{U_K(Z) > 0\}} dZ \leq c_1 \quad \text{for some large } K \text{ and small } c_1.$$

Then we can apply Lemma 3.8 to  $U_K$  and conclude  $U_K \leq 1/2$  in  $B_{1/2}(Z^*)^+$ . This means we can choose  $\sigma = 1 - 2^{-K-1}$  in (3.10). We prove (3.11) using contradiction.

Recall that  $\partial \text{Cone}_\Sigma = \{x_n = g(x')\}$  is a Lipschitz graph, we then have some constant  $c_2$  depending only on  $[g]_{C^{0,1}}$  such that  $|\Omega'| = |B_{1/2}(X^*)' \cap \{U = 0\}| \geq c_2$ . Using the Caccioppoli inequality (3.9), we see for  $h \leq 1/4$ ,

$$\begin{aligned} \int_{X \in \Omega'} U(X, h) dX &\leq \int_{\Omega' \times [0, h]} |\nabla U(X, t)| dt dX \\ &\leq \left\{ \int_{\Omega' \times [0, h]} t^{2s-1} dt dX \right\}^{1/2} \left\{ \int_{\Omega' \times [0, h]} t^{1-2s} |\nabla U(X, t)|^2 dt dX \right\}^{1/2} \\ &\leq Ch^s \left\{ \int_{Z \in B_{3/4}(Z^*)^2} y^{1-2s} |\nabla U(Z)|^2 dZ \right\}^{1/2} \leq Ch^s. \end{aligned}$$

In particular, it implies that  $|B_{3/4}(Z^*)^+ \cap \{y \geq \epsilon\} \cap \{U \leq 1/2\}| \geq c_3$  for any small  $\epsilon$ . Suppose (3.11) is false for large  $K$ , and thus even for any  $1 \leq k \leq K - 1$ . We then see

$$\int_{B_{1-\epsilon}(Z^*)^+ \cap \{y \geq \epsilon\}} \chi_{\{U_k \geq 1/2\}} dZ \geq c_4, \quad \int_{B_{1-\epsilon}(Z^*)^+ \cap \{y \geq \epsilon\}} \chi_{\{U_k = 0\}} dZ \geq c_4 \quad \text{for some small } \epsilon.$$

We apply the Poincaré inequality to  $\bar{U}_k(Z) = \min\{U_k, \frac{1}{2}\}$ , and have

$$\int_{B_{1-\epsilon}(Z^*)^+} y^{1-2s} |\nabla \bar{U}_k(Z)| dZ \geq C \int_{B_{1-\epsilon}(Z^*)^+ \cap \{y \geq \epsilon\}} |\nabla \bar{U}_k(Z)| dZ \geq c_5.$$

Notice that by applying (3.9) to  $U_k$  we have  $\int_{B_{1-\epsilon}(Z^*)^+} y^{1-2s} |\nabla U_k(Z)|^2 dZ \leq C_6$ , so

$$\int_{B_{1-\epsilon}(Z^*)^+} y^{1-2s} \chi_{\{t_k \leq U \leq t_{k+1}\}} dZ \geq \int_{B_{1-\epsilon}(Z^*)^+} y^{1-2s} \chi_{\{|\nabla \bar{U}_k \neq 0\}} dZ \geq c_7$$

for all  $1 \leq k \leq K - 1$ . This prevents  $K$  from being too large (recall that  $y^{1-2s} dZ$  is a finite measure), and then (3.11) is proven.  $\square$

In conclusion, now we have completed the proof of Proposition 2.1.

## ACKNOWLEDGMENT

The author would like to thank Yahong Guo, Congming Li, and Hui Yu for their useful suggestions.

## REFERENCES

- [1] Krzysztof Bogdan. The boundary Harnack principle for the fractional Laplacian. *Studia Math.*, 123(1):43–80, 1997.
- [2] Luis Caffarelli and Luis Silvestre. An extension problem related to the fractional Laplacian. *Comm. Partial Differential Equations*, 32(7-9):1245–1260, 2007.
- [3] Hongjie Dong and Tuoc Phan. On parabolic and elliptic equations with singular or degenerate coefficients. *Indiana Univ. Math. J.*, 72(4):1461–1502, 2023.
- [4] Hongjie Dong and Tuoc Phan. Weighted mixed-norm  $L_p$  estimates for equations in non-divergence form with singular coefficients: the Dirichlet problem. *J. Funct. Anal.*, 285(2):Paper No. 109964, 43, 2023.
- [5] Eugene B. Fabes. Properties of nonnegative solutions of degenerate elliptic equations. In *Proceedings of the international conference on partial differential equations dedicated to Luigi Amerio on his 70th birthday (Milan/Como, 1982)*, volume 52, pages 11–21, 1982.
- [6] Eugene B. Fabes, Carlos E. Kenig, and Raul P. Serapioni. The local regularity of solutions of degenerate elliptic equations. *Comm. Partial Differential Equations*, 7(1):77–116, 1982.
- [7] Piotr Hajłasz. Sobolev spaces on an arbitrary metric space. *Potential Anal.*, 5(4):403–415, 1996.
- [8] Yannick Sire, Susanna Terracini, and Stefano Vita. Liouville type theorems and regularity of solutions to degenerate or singular problems part I: even solutions. *Comm. Partial Differential Equations*, 46(2):310–361, 2021.
- [9] Yannick Sire, Susanna Terracini, and Stefano Vita. Liouville type theorems and regularity of solutions to degenerate or singular problems part II: odd solutions. *Math. Eng.*, 3(1):Paper No. 5, 50, 2021.

SCHOOL OF MATHEMATICAL SCIENCES, FUDAN UNIVERSITY, SHANGHAI 200433, CHINA

*Email address:* zhangchilin@fudan.edu.cn