

# OBSERVABILITY AND CONTROLLABILITY FOR SCHRÖDINGER EQUATIONS IN THE SEMI-PERIODIC SETTING

JINGRUI NIU AND ZEHUA ZHAO

ABSTRACT. Strichartz estimates, well-posedness theory and long time behavior for (nonlinear) Schrödinger equations on waveguide manifolds  $\mathbb{R}^m \times \mathbb{T}^n$  are intensively studied in recent decades while the corresponding control theory and observability estimates remain incomplete. The purpose of this short paper is to investigate the observability and controllability for Schrödinger equations in the waveguide (semi-periodic) setting.

Our main result establishes local exact controllability for the cubic nonlinear Schrödinger equations (NLS) on  $\mathbb{R}^2 \times \mathbb{T}$ , under certain geometric conditions on the control region. To address the nonlinear control problem, we begin by analyzing the observability properties of the linear Schrödinger operator on a general waveguide manifold  $\mathbb{R}^m \times \mathbb{T}^n$ . Utilizing  $H^s$  estimates of the Hilbert Uniqueness Method (HUM) operator and Bourgain spaces, we then prove local exact controllability through a fixed-point method.

## CONTENTS

1. Introduction	1
2. Linear observability and Stationary estimates	5
3. Well-posedness of NLS	10
4. Local null controllability of NLS: Proof of Theorem 1.2	11
5. Exact controllability: Proof of Theorem 1.1	14
Appendix A. Some properties of $X^{s,b}$ spaces	14
References	16

## 1. INTRODUCTION

In this paper, we study the internal exact controllability for the semi-periodic nonlinear Schrödinger equation on the waveguide manifold  $\mathbb{R}^m \times \mathbb{T}^n$  (with dimension  $d = m + n$ ):

$$(1.1) \quad \begin{cases} i\partial_t u + \Delta_{\mathbb{R}^m \times \mathbb{T}^n} u + \epsilon |u|^{p-1} u = f & \text{on } [0, +\infty) \times \mathbb{R}^m \times \mathbb{T}^n \\ u|_{t=0} = u_0 & \text{on } \mathbb{R}^m \times \mathbb{T}^n \end{cases}$$

where  $m$  and  $n$  are two positive integers, and  $\epsilon \in \mathbb{R}$ .  $\Delta_{\mathbb{R}^m \times \mathbb{T}^n}$  is the canonical Laplacian defined in  $\mathbb{R}^m \times \mathbb{T}^n$ , and sometimes we may write  $\Delta_{x,y}$  for short.

**1.1. Background and motivations.** In the literature, there has been extensive research on the controllability and observability of (nonlinear) Schrödinger equations on compact manifolds and bounded domains. We begin with the linear setting, particularly addressing observability problems. When the control region is an open set satisfying the geometric control condition (GCC), Lebeau [35] proved that observability is true for an arbitrarily short time  $T > 0$ . However, GCC is not always a

---

2020 *Mathematics Subject Classification.* Primary: 93B05; Secondary: 35R01, 93C20, 35Q55.

*Key words and phrases.* nonlinear Schrödinger equations, waveguide manifolds, well-posedness, controllability, observability.

necessary condition. In various contexts, including the torus [2, 7, 9, 27], compact hyperbolic surfaces [28] (also see [3] for negatively curved manifolds), and the disk [1], observability has been established for any  $T > 0$  and for any non-empty open control region, provided the control region encompasses a neighborhood of some portion of the boundary. It is also worth mentioning that for certain subelliptic Schrödinger equations, the minimal observability time can be strictly positive (see [8]).

Regarding the controllability of nonlinear Schrödinger equations (NLS), significant results have been obtained for different dimensions on compact manifolds or bounded domains (though the list here is not exhaustive). In one dimension, Laurent [32] proved global internal controllability in large time for NLS on a bounded interval, while local results were also achieved in [40]. For two dimensions, Dehman, Gérard, and Lebeau [15] established exact controllability in  $H^1$  for the defocusing NLS on compact surfaces. In the three-dimensional case, Laurent [33] demonstrated global internal controllability over large time intervals for NLS on certain compact manifolds. Concerning the boundary controls, we refer to [19, 39] and the references therein. For a broader overview of these results, we refer to the surveys [52, 34].

More recently, attention has shifted towards understanding observability for Schrödinger equations in non-compact settings [24, 45, 38]. The specific case of periodic domains has been explored for the observability of the free Schrödinger equation [42] and for settings with a periodic potential [4]. Notably, in [4], the control region, although  $2\pi\mathbb{Z}^2$ -periodic, might not satisfy GCC. Despite these advances, the study of controllability and observability in semi-periodic settings remains largely unexplored.

We now provide a concise overview of our research focus: “**(nonlinear) Schrödinger equations on waveguides**”. This area has garnered significant attention over the past few decades, emerging as a prominent subject within nonlinear dispersive equations. Our approach blends traditional dispersive techniques with innovative analysis tools to delve into this intriguing domain.

Waveguide manifolds, denoted as  $\mathbb{R}^m \times \mathbb{T}^n$ , represent the product of Euclidean space with tori and are of particular relevance in nonlinear optics, especially within telecommunications. Currently, data signals in backbone networks predominantly travel via optical carriers through fibers, which function as specialized waveguides. As applications like the internet demand greater bandwidth and cost-effective data transmission, there is increasing emphasis on optimizing these network infrastructures. The nonlinear Schrödinger equation plays a pivotal role in modeling nonlinear effects in optical fibers, essential for enhancing performance and efficiency. In physics, an optical waveguide directs light along a defined path, and the study of solutions on waveguide manifolds is particularly intriguing due to the unique combination of properties inherited from both Euclidean spaces and tori, offering deeper insights into the underlying physics.

Given the nature of these combined spaces, the nonlinear Schrödinger Equation (NLS) posed on the waveguide manifold inherits characteristics from both Euclidean spaces and tori. The Euclidean case is studied and the theory, at least in the defocusing setting, is well established. (See [13, 14, 17, 41] and the references therein.) Moreover, we refer to [22, 25, 30, 48] for a few works on tori. Due to the nature of such product spaces, we see NLS posed on the waveguide manifold mixed inheriting properties from those on classical Euclidean spaces and tori. The techniques used in Euclidean and tori settings are frequently combined and applied to waveguide problems. We refer to [10, 11, 12, 20, 22, 43, 44, 23, 25, 26, 47, 49, 50, 51] for some NLS results in the waveguide setting. At last, we note that, though scattering behavior is not expected for the periodic case because of the lack of dispersive, for some specific models of waveguides, modified scattering, and scattering results can be obtained as in the Euclidean. (See [10, 20, 44, 21] for example.)

**Geometric setting.** We work in a  $d$ -dimensional waveguide manifold as  $\mathbb{R}^m \times \mathbb{T}^n$ , where  $m$  and  $n$  are two positive integers with (whole dimension)  $d = m + n$  ( $m, n \geq 1$ ). We consider a control region  $\Omega$  of the following type :

(**G**)  $\Omega = (\Omega_1, \Omega_2) \subset \mathbb{R}^m \times \mathbb{T}^n$ . Let  $\Omega_1 \subset \mathbb{R}^m$  be a nonempty, open,  $2\pi\mathbb{Z}^m$ -invariant set. Let  $\Omega_2 \subset \mathbb{T}^{d-m}$  be open and nonempty.

This kind of region may not satisfy the Geometric Control Condition, which is raised in e.g. [35]

**1.2. The statement of the main results.** In this subsection, we present the main results of this paper respectively. We note that all of the results can be generalized to all dimensions in a natural way. Since 3D cubic NLS model is typical and popular<sup>2</sup>, in Theorem 1.1 and Theorem 1.2, we adopt 3D cubic NLS model<sup>3</sup> to illustrate the nonlinear results.

**Theorem 1.1** (Exact controllability). *Let  $T > 0$ , and  $\epsilon = \pm 1$ . Let  $\Omega = (\Omega_1, \Omega_2)$  satisfy the condition (**G**). For any  $s \geq 1$ , there exists  $\delta > 0$ , such that for all  $u_0, u_f \in H^s(\mathbb{R}^2 \times \mathbb{T})$  satisfying that  $\|u_0\|_{H^s(\mathbb{R}^2 \times \mathbb{T})} + \|u_f\|_{H^s(\mathbb{R}^2 \times \mathbb{T})} < \delta$ , there exists a control function  $f \in C([0, T]; H^s(\mathbb{R}^2 \times \mathbb{T}))$  supported in  $[0, T] \times \Omega$  such that the unique solution  $u \in X_T^{s,b}$  to*

$$(1.2) \quad \begin{cases} i\partial_t u + \Delta_{x,y} u + \epsilon|u|^2 u = f & \text{on } [0, T] \times \mathbb{R}^2 \times \mathbb{T}, \\ u|_{t=0} = u_0 & \text{on } \mathbb{R}^2 \times \mathbb{T}, \end{cases}$$

fulfills  $u|_{t=T} = u_f$ .

This is a local exact controllability result for NLS in the semi-periodic setting. The general strategy to establish the exact controllability is to reduce it to the *null controllability*, i.e. the exact controllability with null final data ( $u_f = 0$ ). To be more specific, we first prove the following result.

**Theorem 1.2** (Null controllability). *Let  $T > 0$ . Let  $\epsilon$  and  $\Omega$  be the same as in Theorem 1.1. For any  $s \geq 1$ , there exists  $\delta > 0$ , such that for all  $u_0 \in H^s(\mathbb{R}^2 \times \mathbb{T})$  satisfying that  $\|u_0\|_{H^s(\mathbb{R}^2 \times \mathbb{T})} < \delta$ , there exists a control function  $g$  such that the unique solution  $u \in X_T^{s,b}$  to*

$$(1.3) \quad \begin{cases} i\partial_t u + \Delta_{x,y} u + \epsilon|u|^2 u = \varphi_T \chi_\Omega (1 - \Delta_{x,y})^{-s} \varphi_T \chi_\Omega g & \text{on } [0, T] \times \mathbb{R}^2 \times \mathbb{T}, \\ u|_{t=0} = u_0 & \text{on } \mathbb{R}^2 \times \mathbb{T}, \end{cases}$$

fulfills  $u|_{t=T} = 0$ . Furthermore, the control function  $g$  verifies the following conditions

- (1)  $g \in C([0, T]; H^{-s}(\mathbb{R}^2 \times \mathbb{T}))$
- (2)  $\varphi_T(\cdot) = \varphi_1(\cdot/T) \in C^\infty(\mathbb{R})$  where  $\varphi_1(t) = 1$  for  $t \leq \frac{1}{2}$  and  $\varphi_1(t) = 0$  for  $t \geq \frac{3}{4}$ ;
- (3)  $0 \leq \chi_\Omega \in C^\infty(\mathbb{R}^2 \times \mathbb{T})$  satisfies  $\mathbf{1}_{\Omega' \times \Omega_2} \leq \chi_\Omega \leq \mathbf{1}_{\Omega_1 \times \Omega_2}$ , where  $\overline{\Omega'} \subset \Omega_1$ .

To investigate the nonlinear case, a classic approach is to first study the linearized Schrödinger equation. The following theorem concerns solutions of the stationary Schrödinger equation and is applicable to high-energy eigenfunctions.

**Theorem 1.3** (Stationary estimate). *Let  $\Omega = (\Omega_1, \Omega_2)$  satisfy the condition (**G**). For all  $\lambda \in \mathbb{R}$  we have the following estimate: for*

$$(1.4) \quad (\Delta_{x,y} - \lambda)u = f,$$

<sup>1</sup>This means  $\Omega = \{(x, y) | x \in \Omega_1, y \in \Omega_2\}$ .

<sup>2</sup>We refer to Kenig-Merle [29] for the scattering result in the Euclidean setting. See also the references therein.

<sup>3</sup>We consider cubic NLS on  $\mathbb{R}^2 \times \mathbb{T}$  or  $\mathbb{R} \times \mathbb{T}^2$ . Since the proofs work in the same way, we only consider the  $\mathbb{R}^2 \times \mathbb{T}$  case.

we have,

$$(1.5) \quad \|u(x, y)\|_{L^2(\mathbb{R}^n \times \mathbb{T}^m)} \leq C(\|f(x, y)\|_{L^2(\mathbb{R}^n \times \mathbb{T}^m)} + \|u(x, y)\|_{L^2(\Omega)}),$$

with  $C$  independent of  $\lambda$ .

*Remark 1.4.* In particular, if  $f = 0$ , this estimate implies a unique continuation property for the eigenfunctions of  $-\Delta_{x,y}$ . Indeed, if  $f = 0$ , the total  $L^2$ -norm of  $u$  can be bounded by its local  $L^2$ -norm in  $\Omega$ . As a consequence, if an eigenfunction  $u$  vanishes in a subdomain  $\Omega$ , which verifies the condition **(G)**, then, it vanishes everywhere.

Related to this stationary estimate, we have the following *observability estimate* for linear Schrödinger propagator. Once we have it, using *Hilbert uniqueness method* (HUM for short, see [36] and Section 4.1), we obtain the controllability result automatically for the linearized equation.

**Theorem 1.5** (Observability). *Let  $\Omega = (\Omega_1, \Omega_2)$  satisfy the condition **(G)**. For every  $T > 0$ , there exists a constant  $C = C(T, \Omega)$  such that for  $\forall u_0 \in L^2(\mathbb{R}^n \times \mathbb{T}^m)$ ,*

$$\|u_0\|_{L^2(\mathbb{R}^n \times \mathbb{T}^m)}^2 \leq C \int_0^T \|e^{it\Delta_{x,y}} u_0\|_{L^2(\Omega)}^2 dt.$$

**1.3. Strategy of the proofs.** Now we discuss the strategy of the proofs for the main theorems stated in the previous subsection.

Our primary objective is to prove Theorem 1.1, i.e., the local exact controllability for nonlinear Schrödinger equations on  $\mathbb{R}^2 \times \mathbb{T}$ . We first reduce it to the null controllability with certain control in a specific form (Theorem 1.2). Following the classic approach, we need to analyze the control problem for the corresponding linear equation:

$$\begin{cases} i\partial_t u + \Delta_{x,y} u = f & \text{on } [0, T] \times \mathbb{R}^2 \times \mathbb{T}, \\ u|_{t=0} = u_0, u|_{t=T} = 0. & \text{on } \mathbb{R}^2 \times \mathbb{T}, \end{cases}$$

As a consequence of *Hilbert Uniqueness Method* (see Section 4.1 for details), constructing the control operator  $\mathcal{L} : u_0 \mapsto f$  is equivalent to proving observability for any  $s \in \mathbb{R}$ ,

$$(1.6) \quad \|u_0\|_{H^s(\mathbb{R}^2 \times \mathbb{T})}^2 \leq C \int_0^T \|\chi_\Omega e^{it\Delta_{x,y}} u_0\|_{H^s(\mathbb{R}^2 \times \mathbb{T})}^2 dt.$$

In order to prove (1.6) for any  $s \in \mathbb{R}$ , we combine the  $L^2$ -observability (Theorem 1.5) with a unique continuation property, which is ensured by the resolvent estimate (Theorem 1.3). Inspired by [46], we prove these two results by reducing them into a tori setting by the application of *Floquet-Bloch transform*. After a precise analysis of the linear problem, we follow the ideas in [16, 33] and decompose the solution  $u$  to (1.3) into  $u = v + \Psi$ , where  $v$  solves the following equation

$$\begin{cases} i\partial_t v + \Delta_{x,y} v = -\epsilon|u|^2 u, \\ v|_{t=T} = 0, \end{cases}$$

and  $\Psi$  is the solution to

$$\begin{cases} i\partial_t \Psi + \Delta_{x,y} \Psi = \varphi_T \chi_\Omega (1 - \Delta_{x,y})^{-s} \varphi_T \chi_\Omega g, \\ \Psi|_{t=0} = \Psi_0, \Psi|_{t=T} = 0. \end{cases}$$

The purpose is then to choose the adequate  $\Psi_0$ , and the system is completely determined. Let us define  $\mathcal{J} : \Psi_0 \mapsto u_0 - v|_{t=0}$ . Thus, seeking the control function  $g$  is reduced to finding a fixed point for  $\mathcal{J}$ , provided that  $\|u_0\|_{H^s}$  is small enough (see Section 4 for details). The general result will follow by reversing time (presented in Section 5).

**1.4. Organization of the rest of this paper.** In Section 2, we prove linear observability and stationary estimates, which includes the proof for Theorem 1.3; in Section 3, we discuss well-posedness theory for NLS which will be used for the nonlinear case; in Section 4, we show the local null controllability of NLS, which completes the proof of Theorem 1.2; in Section 5, we obtain the exact controllability, which finishes the proof of Theorem 1.1.

**1.5. Notations.** Throughout this paper, we use  $C$  to denote the universal constant and  $C$  may change line by line. We say  $A \lesssim B$ , if  $A \leq CB$ . We say  $A \sim B$  if  $A \lesssim B$  and  $B \lesssim A$ . We also use the notation  $C_B$  to denote a constant depends on  $B$ . We use the standard notation for  $L^p$  spaces and  $L^2$ -based Sobolev spaces  $H^s$ .

We define the Fourier transform on  $\mathbb{R}^n \times \mathbb{T}^m$  as follows:

$$(1.7) \quad (\mathcal{F}f)(\xi) = \int_{\mathbb{R}^n \times \mathbb{T}^m} f(z) e^{-iz \cdot \xi} dz,$$

where  $\xi = (\xi_1, \xi_2, \dots, \xi_d) \in \mathbb{R}^n \times \mathbb{Z}^m$  and  $d = m + n$ . We also note the Fourier inversion formula

$$(1.8) \quad f(z) = c \sum_{(\xi_{n+1}, \dots, \xi_d) \in \mathbb{Z}^m} \int_{(\xi_1, \dots, \xi_n) \in \mathbb{R}^n} (\mathcal{F}f)(\xi) e^{iz \cdot \xi} d\xi_1 \dots d\xi_n.$$

For convenience, we may consider the discrete sum to be the integral with discrete measure so we can combine the above integrals together and treat them to be one integral. Moreover, we define the Schrödinger propagator  $e^{it\Delta}$  by

$$(1.9) \quad (\mathcal{F}e^{it\Delta} f)(\xi) = e^{-it|\xi|^2} (\mathcal{F}f)(\xi).$$

At last, we refer to [18, 51] for the definitions and properties of Fourier restriction spaces (also known as Bourgain spaces) in the waveguide setting. They share very similar properties as the tori case. For the sake of completeness, we include some properties of these function spaces in the Appendix A. These spaces are very useful and frequently used for the study of NLS on tori or waveguide manifolds.<sup>4</sup>

**Acknowledgment.** We highly appreciate Prof. N. Burq for helpful suggestions and insightful discussions. J. Niu is supported by Defi Inria EQIP. Z. Zhao is supported by the NSF grant of China (No. 12101046, 12271032, 2426205) and the Beijing Institute of Technology Research Fund Program for Young Scholars.

## 2. LINEAR OBSERVABILITY AND STATIONARY ESTIMATES

**2.1. The Floquet-Bloch transform.** In this part, we introduce the partial Floquet-Bloch transform. This tool is instrumental in the proof of Theorem 1.3. For more details, we refer to [46], [31, Section 4] and [4, Section 2.1]. We first introduce the partial Floquet transform.

**Definition 2.1.** We define the partial Floquet-Bloch transform  $\mathcal{T}^p : L^2(\mathbb{R}^n \times \mathbb{T}^m) \rightarrow L^2(\mathbb{T}^{m+n} \times [0, 1)^n)$  by

$$\mathcal{T}^p u(x, y, \alpha) := e^{ix \cdot \alpha} \sum_{k \in \mathbb{Z}^n} e^{2\pi i \alpha \cdot k} u(x + 2k\pi, y), \quad \forall u \in L^2(\mathbb{R}^n \times \mathbb{T}^m), \forall (x, y, \alpha) \in \mathbb{T}^n \times \mathbb{T}^m \times [0, 1)^n.$$

*Remark 2.2.* In particular, if  $m = 0$ , we obtain the usual Floquet-Bloch transform. Furthermore, removing the factor  $e^{i\alpha \cdot x}$ , we obtain a partial Floquet transform

$$\Pi^p u(x, y, \alpha) := \sum_{k \in \mathbb{Z}^n} e^{2\pi i \alpha \cdot k} u(x + 2k\pi, y), \quad \forall u \in L^2(\mathbb{R}^n \times \mathbb{T}^m), \forall (x, y, \alpha) \in \mathbb{T}^n \times \mathbb{T}^m \times [0, 1)^n.$$

<sup>4</sup>We also refer to [22, 23] and the references therein for another type of function spaces, which are also very useful for the study of NLS on tori or waveguide manifolds (especially for the critical case).

For convenience, for any  $\alpha \in [0, 1)^n$  and  $u \in L^2(\mathbb{R}^n \times \mathbb{T}^m)$ , we define  $\Pi_\alpha : L^2(\mathbb{R}^n \times \mathbb{T}^m) \rightarrow L^2(\mathbb{T}^{m+n})$  by

$$(2.1) \quad \Pi_\alpha u(x, y) := e^{ix \cdot \alpha} \sum_{k \in \mathbb{Z}^n} e^{2\pi i \alpha \cdot k} u(x + 2k\pi, y), \quad \forall (x, y) \in \mathbb{T}^n \times \mathbb{T}^m.$$

The next proposition ensures that the partial Floquet-Bloch transform is an isometry from  $L^2(\mathbb{R}^n \times \mathbb{T}^m)$  to  $L^2(\mathbb{T}^{m+n} \times [0, 1)^n)$ .

**Proposition 2.3.** *The map  $\mathcal{T}^p$  is an isometric isomorphism from  $L^2(\mathbb{R}^n \times \mathbb{T}^m)$  to  $L^2(\mathbb{T}^{m+n} \times [0, 1)^n)$ .*

This proposition is based on the following lemma.

**Lemma 2.4.** *We have the equality of  $L^2$  norms*

$$(2.2) \quad \|g\|_{L^2(\mathbb{R}^n \times \mathbb{T}^m)}^2 = \int_{[0, 1)^n} \|\Pi_\alpha g\|_{L^2(\mathbb{T}^n \times \mathbb{T}^m)}^2 d\alpha.$$

More generally, if  $\Omega_1 \in \mathbb{R}^n$  is  $2\pi\mathbb{Z}^n$ -invariant and  $\Omega_0$  denotes its projection to  $\mathbb{T}^n$

$$(2.3) \quad \|g\|_{L^2(\Omega_1 \times \mathbb{T}^m)}^2 = \int_{[0, 1)^n} \|\Pi_\alpha g\|_{L^2(\Omega_0 \times \mathbb{T}^m)}^2 d\alpha.$$

*Proof.* We use Fubini to compute the Fourier coefficients of the periodic functions  $\Pi_\alpha g$  on  $\mathbb{T}^{m+n}$ .

$$\begin{aligned} \widehat{\Pi_\alpha g}(k, l) &= \frac{1}{(2\pi)^{\frac{n+m}{2}}} \int_{\mathbb{T}^m} \int_{\mathbb{T}^n} e^{-ix \cdot k} e^{-iy \cdot l} \sum_{j \in \mathbb{Z}^n} e^{ix \cdot \alpha} e^{2\pi i \alpha \cdot j} g(x + 2j\pi, y) dx dy \\ &= \frac{1}{(2\pi)^{\frac{n+m}{2}}} \int_{\mathbb{T}^m} \int_{\mathbb{R}^n} e^{ix \cdot (\alpha - k)} e^{-iy \cdot l} g(x, y) dx dy \\ &= \mathcal{F}_x(\widehat{g}(\cdot, l))(k - \alpha). \end{aligned}$$

Integrating the sum of squares of the right hand side over the unit cube  $[0, 1)^n$  gives  $\|g\|_{L^2(\mathbb{R}^n \times \mathbb{T}^m)}^2$  by Fubini and Plancherel theorems on  $\mathbb{R}^n \times \mathbb{T}^m$ , while on the left hand side, we get

$$\int_{[0, 1)^n} \sum_{k \in \mathbb{Z}^n, l \in \mathbb{Z}^m} |\widehat{\Pi_\alpha g}(k, l)|^2 d\alpha = \int_{[0, 1)^n} \|\Pi_\alpha g\|_{L^2(\mathbb{T}^{n+m})}^2 d\alpha$$

by Plancherel on the torus. The generalization to taking the norm over  $\Omega_1 \times \Omega_2$  is proved simply by applying (2.2) to the function  $\mathbb{1}_{\Omega_1 \times \Omega_2} g$ .  $\square$

**2.2. Observability for partial-twisted Laplacian.** Let  $\alpha \in \mathbb{R}^n$ , we set

$$H_\alpha := (\partial_x - i\alpha)^2 + \Delta_y.$$

Note that these operators are all self-adjoint with the same domain independent of  $\alpha$ . We first recall the Schrödinger observability on the torus

**Theorem 2.5.** [2, Theorem 4] *For every non-empty open set  $\Omega \subset \mathbb{T}^d$  and every  $T > 0$  there exists a constant  $C = C(T, \Omega)$  such that*

$$\|u_0\|_{L^2(\mathbb{T}^d)}^2 \leq C \int_0^T \|e^{it\Delta} u_0\|_{L^2(\Omega)}^2 dt$$

for every initial datum  $u_0 \in L^2(\mathbb{T}^d)$ .

We use this classical theorem to prove the following lemma

**Lemma 2.6.** *For every non-empty open set  $\Omega = (\Omega_0, \Omega_2) \subset \mathbb{T}^n \times \mathbb{T}^m$ , and every  $T > 0$  there exists a constant  $C = C(T, \Omega)$  such that*

$$\|u_0\|_{L^2(\mathbb{T}^{n+m})}^2 \leq C \int_0^T \|e^{itH_\alpha} u_0\|_{L^2(\Omega)}^2 dt$$

for every initial datum  $u_0 \in L^2(\mathbb{T}^{n+m})$ .

*Proof.* We point out that this lemma is almost a restatement of [46, Proposition 4]. When  $m = 0$ , the twisted Laplacian is defined in [46, Eq. (3)] satisfies a similar observability. The proof of [46, Proposition 4] can also be applied here without changes. For the completeness of our paper, we present the proof adapted to our setting. For  $\Omega_0 \subset \mathbb{T}^n$ , we can find a nonempty open set  $\omega_0 \subset \mathbb{T}^n$  such that  $\overline{\omega_0} \subset \Omega_0$ . We set  $\omega = (\omega_0, \Omega_2) \subset \Omega$ . Hence there exists  $T > 0$  such that  $x_1 \in \omega_0$  and  $d(x_1, x_2) < 4\sqrt{n}T$  imply  $x_2 \in \Omega_0$ . Using the operator identity

$$e^{itH_\alpha} = e^{it(\partial_x - i\alpha)^2} e^{i\Delta_y} = e^{-it|\alpha|^2} \tau_{(2t\alpha, 0)} e^{it\Delta_{x,y}},$$

where for  $\theta = (\theta_1, \theta_2) \in \mathbb{R}^{n+m}$ ,  $\tau_\theta$  denotes the translation operator  $\tau_\theta f(x, y) := f(x + \theta_1, y + \theta_2)$ . Thus, by the  $H_0$ -observability and the choice of  $T \ll 1$  so that  $\tau_{(-2t\alpha, 0)}(\omega) \subset \Omega$  for  $\forall t \in [0, T]$ , we obtain

$$\begin{aligned} \|u_0\|_{L^2(\mathbb{T}^{n+m})}^2 &\leq C \int_0^T \|e^{it\Delta_{x,y}} u_0\|_{L^2(\omega)}^2 dt \\ &\leq C \int_0^T \|e^{-it|\alpha|^2} \tau_{(2t\alpha, 0)} e^{it\Delta_{x,y}} u_0\|_{L^2(\tau_{(-2t\alpha, 0)}(\omega))}^2 dt \\ &\leq C \int_0^T \|e^{-it|\alpha|^2} \tau_{(2t\alpha, 0)} e^{it\Delta_{x,y}} u_0\|_{L^2(\Omega)}^2 dt \\ &\leq C \int_0^T \|e^{itH_\alpha} u_0\|_{L^2(\Omega)}^2 dt. \end{aligned}$$

□

Thanks to [37, Theorem 5.1], we have the following proposition.

**Proposition 2.7.** *Let  $\Omega = (\Omega_0, \Omega_2) \subset \mathbb{T}^n \times \mathbb{T}^m$  be open and nonempty. For all  $\alpha \in [0, 1]^n$  and*

$$(2.4) \quad (H_\alpha - \lambda)u = f,$$

posed on  $\mathbb{T}^{n+m}$ , we have,

$$(2.5) \quad \|u\|_{L^2(\mathbb{T}^{n+m})} \leq C(\|f\|_{L^2(\mathbb{T}^{n+m})} + \|u\|_{L^2(\Omega)}),$$

with constants independent of  $\alpha$  and  $\lambda \in \mathbb{R}$ .

### 2.3. Stationary estimate: proof of Theorem 1.3.

*Proof of Theorem 1.3.* First, we notice that

$$(2.6) \quad H_\alpha - \lambda = (\partial_x - i\alpha)^2 + \Delta_y - \lambda = e^{i\alpha \cdot x} (\Delta_{x,y} - \lambda) e^{-i\alpha \cdot x},$$

where  $\alpha \in [0, 1]^n$ . Thus  $(\Delta_{x,y} - \lambda)u = f$  yields

$$(2.7) \quad (H_\alpha - \lambda)e^{i\alpha \cdot x} u = e^{i\alpha \cdot x} f.$$

Applying the partial Floquet transform  $\Pi^p$  on both sides and using translation-invariance of  $H_\alpha$ , we get an equation on the torus:

$$(2.8) \quad (H_\alpha - \lambda)\Pi_\alpha u = \Pi_\alpha f \text{ on } \mathbb{T}^{n+m}.$$

Applying Proposition 2.7, we obtain for every  $\alpha$  in a fundamental domain (and with constants independent of  $\alpha$ )

$$\|\Pi_\alpha u\|_{L^2(\mathbb{T}^{n+m})} \leq C(\|\Pi_\alpha f\|_{L^2(\mathbb{T}^{n+m})} + \|\Pi_\alpha u\|_{L^2(\Omega_0)}).$$

Now by Lemma 2.4 we may integrate both sides in  $\alpha_1$  to obtain Theorem 1.3.  $\square$

*Remark 2.8.* The idea of establishing the estimate (1.5) in a periodic setting through a twisted Laplacian was first raised by Wunsch in [46]. The correspondence between  $H_\alpha$  on tori and  $\Delta$  in periodic setting introduced in [46] may look simple but turns out to be a powerful tool.

#### 2.4. Observability in $\mathbb{R}^m \times \mathbb{T}^n$ : Proof of Theorem 1.5.

*Proof of Theorem 1.5.* Let  $u(t) := e^{it\Delta_{x,y}}u_0$  and  $v(t) := \Pi_\alpha u(t)$ , for  $\alpha \in [0, 1]^n$ . Then  $u$  is the solution to

$$\begin{cases} (i\partial_t + \Delta_{x,y})u = 0 & \text{in } (0, T) \times \mathbb{R}^n \times \mathbb{T}^m, \\ u|_{t=0} = u_0 & \text{in } \mathbb{R}^n \times \mathbb{T}^m, \end{cases}$$

and  $v$  is the solution to

$$\begin{cases} (i\partial_t + H_\alpha)v = 0 & \text{in } (0, T) \times \mathbb{T}^n \times \mathbb{T}^m, \\ v|_{t=0} = \Pi_\alpha u_0 & \text{in } \mathbb{T}^n \times \mathbb{T}^m. \end{cases}$$

By Lemma 2.6, we obtain

$$\|\Pi_\alpha u_0\|_{L^2(\mathbb{T}^n \times \mathbb{T}^m)}^2 \leq C \int_0^T \|v(t)\|_{L^2(\Omega)}^2 dt = C \int_0^T \|\Pi_\alpha u(t)\|_{L^2(\Omega)}^2 dt.$$

Integrating the LHS over  $[0, 1]^n$  gives  $\|u_0\|_{L^2(\mathbb{R}^n \times \mathbb{T}^m)}^2$  by Lemma 2.4, while on the RHS, we get

$$\int_{[0,1]^n} \int_0^T \|\Pi_\alpha u(t)\|_{L^2(\Omega)}^2 dt d\alpha = \int_0^T \|u(t)\|_{L^2(\Omega)}^2 dt,$$

by Fubini and Lemma 2.4. We conclude that

$$\|u_0\|_{L^2(\mathbb{R}^n \times \mathbb{T}^m)}^2 \leq C \int_0^T \|e^{it\Delta_{x,y}}u_0\|_{L^2(\Omega)}^2 dt$$

$\square$

**Proposition 2.9.** *Let  $\Omega = (\Omega_1, \Omega_2)$  satisfy the condition  $(\mathbf{G})$ . For every  $T > 0$  and  $s \geq 1$ , there exists a constant  $C = C(T, \Omega, s, \chi_\Omega)$  such that for  $\forall u_0 \in H^s(\mathbb{R}^n \times \mathbb{T}^m)$*

$$(2.9) \quad \|u_0\|_{H^s(\mathbb{R}^n \times \mathbb{T}^m)}^2 \leq C \int_0^T \|\chi_\Omega e^{it\Delta_{x,y}}u_0\|_{H^s(\mathbb{R}^n \times \mathbb{T}^m)}^2 dt,$$

where  $0 \leq \chi_\Omega \in C^\infty(\mathbb{R}^n \times \mathbb{T}^m)$  satisfies  $\mathbf{1}_\Omega \leq \chi_\Omega$ .

*Proof.* By the definition of  $\chi_\Omega$  and Theorem 1.5, we can easily derive that  $\exists C_0 = C_0(T, \Omega) > 0$  such that for  $\forall u_0 \in L^2(\mathbb{R}^n \times \mathbb{T}^m)$

$$(2.10) \quad \|u_0\|_{L^2(\mathbb{R}^n \times \mathbb{T}^m)}^2 \leq C_0 \int_0^T \|\chi_\Omega e^{it\Delta_{x,y}}u_0\|_{L^2(\mathbb{R}^n \times \mathbb{T}^m)}^2 dt.$$

For the initial datum  $u_0 \in H^s(\mathbb{R}^n \times \mathbb{T}^m)$ ,

$$\|u_0\|_{H^s(\mathbb{R}^n \times \mathbb{T}^m)}^2 = \|(1 - \Delta_{x,y})^{\frac{s}{2}}u_0\|_{L^2(\mathbb{R}^n \times \mathbb{T}^m)}^2 \leq C_0 \int_0^T \|\chi_\Omega e^{it\Delta_{x,y}}(1 - \Delta_{x,y})^{\frac{s}{2}}u_0\|_{L^2(\mathbb{R}^n \times \mathbb{T}^m)}^2 dt.$$

Thanks to Cauchy-Schwarz inequality and the fact that

$$\chi_\Omega e^{it\Delta_{x,y}}(1 - \Delta_{x,y})^{\frac{s}{2}}u_0 = (1 - \Delta_{x,y})^{\frac{s}{2}}\chi_\Omega e^{it\Delta_{x,y}}u_0 + [\chi_\Omega, (1 - \Delta_{x,y})^{\frac{s}{2}}]e^{it\Delta_{x,y}}u_0,$$



we obtain

$$\begin{aligned} \|u_0\|_{H^s(\mathbb{R}^n \times \mathbb{T}^m)}^2 &\leq 2C_0 \int_0^T \|(1 - \Delta_{x,y})^{\frac{s}{2}} \chi_\Omega e^{it\Delta_{x,y}} u_0\|_{L^2(\mathbb{R}^n \times \mathbb{T}^m)}^2 dt \\ &\quad + 2C_0 \int_0^T \|[\chi_\Omega, (1 - \Delta_{x,y})^{\frac{s}{2}}] e^{it\Delta_{x,y}} u_0\|_{L^2(\mathbb{R}^n \times \mathbb{T}^m)}^2 dt. \end{aligned}$$

Since the commutator  $[\chi_\Omega, (1 - \Delta_{x,y})^{\frac{s}{2}}]$  is a pseudo-differential operator of order  $s-1$ , we know that  $\|[\chi_\Omega, (1 - \Delta_{x,y})^{\frac{s}{2}}] e^{it\Delta_{x,y}} u_0\|_{L^2(\mathbb{R}^n \times \mathbb{T}^m)}^2 \leq C \|e^{it\Delta_{x,y}} u_0\|_{H^{s-1}(\mathbb{R}^n \times \mathbb{T}^m)}^2 \leq C \|u_0\|_{H^{s-1}(\mathbb{R}^n \times \mathbb{T}^m)}^2$ . Thus,

$$(2.11) \quad \|u_0\|_{H^s(\mathbb{R}^n \times \mathbb{T}^m)}^2 \leq 2C_0 \int_0^T \|\chi_\Omega e^{it\Delta_{x,y}} u_0\|_{H^s(\mathbb{R}^n \times \mathbb{T}^m)}^2 dt + 2C_0 C T \|u_0\|_{H^{s-1}(\mathbb{R}^n \times \mathbb{T}^m)}^2.$$

This is the so-called *weak observability* in  $H^s(\mathbb{R}^n \times \mathbb{T}^m)$ . We apply a standard procedure based on compactness-uniqueness arguments to derive the observability (2.9) from the weak observability (2.11). Suppose that (2.9) is false. Then we find a sequence  $\{u_0^k\}_{k \in \mathbb{N}^*} \subset H^s(\mathbb{R}^n \times \mathbb{T}^m)$  such that

$$(2.12) \quad \|u_0^k\|_{H^s(\mathbb{R}^n \times \mathbb{T}^m)}^2 = 1,$$

$$(2.13) \quad \int_0^T \|\chi_\Omega e^{it\Delta_{x,y}} u_0^k\|_{H^s(\mathbb{R}^n \times \mathbb{T}^m)}^2 dt \rightarrow 0.$$

The second statement (2.13) leads to

$$(2.14) \quad \chi_\Omega e^{it\Delta_{x,y}} u_0^k \rightarrow 0, \text{ strongly in } L^2((0, T); H^s(\mathbb{R}^n \times \mathbb{T}^m)), \text{ as } k \rightarrow \infty.$$

Thanks to (2.12), we can extract a subsequence (still denoted by  $u_0^k$  for simplicity) such that  $u_0^k \rightharpoonup u$  weakly in  $H^s(\mathbb{R}^n \times \mathbb{T}^m)$ . By the compact inclusion  $i : H^s(\mathbb{R}^n \times \mathbb{T}^m) \rightarrow H^{s-1}(\mathbb{R}^n \times \mathbb{T}^m)$ , we deduce that  $u_0^k \rightarrow u$  strongly in  $H^{s-1}(\mathbb{R}^n \times \mathbb{T}^m)$ . Then we define a subspace  $\mathcal{N}_{T_0}$  in  $H^s(\mathbb{R}^n \times \mathbb{T}^m)$ , collecting initial data that generate invisible solutions in  $(0, T_0)$ , by

$$(2.15) \quad \mathcal{N}_{T_0} := \{f \in H^s(\mathbb{R}^n \times \mathbb{T}^m) : \chi_\Omega e^{it\Delta_{x,y}} f = 0 \text{ for } t \in (0, T_0)\}.$$

**Lemma 2.10.**  $\mathcal{N}_T = \{0\}$ .

Using Lemma 2.10, we finish the proof of the observability. Using the weak observability (2.11) and the condition (2.13), we obtain

$$\begin{aligned} 1 &= \lim_{k \rightarrow \infty} \|u_0^k\|_{H^s(\mathbb{R}^n \times \mathbb{T}^m)}^2 \leq \lim_{k \rightarrow \infty} \left( 2C_0 \int_0^T \|\chi_\Omega e^{it\Delta_{x,y}} u_0^k\|_{H^s(\mathbb{R}^n \times \mathbb{T}^m)}^2 dt + 2C_0 C T \|u_0^k\|_{H^{s-1}(\mathbb{R}^n \times \mathbb{T}^m)}^2 \right) \\ &\leq \lim_{k \rightarrow \infty} 2C_0 C T \|u_0^k\|_{H^{s-1}(\mathbb{R}^n \times \mathbb{T}^m)}^2 \\ &\leq 2C_0 C T \lim_{k \rightarrow \infty} \|u_0^k\|_{H^{s-1}(\mathbb{R}^n \times \mathbb{T}^m)}^2. \end{aligned}$$

By the strong convergence of  $u_0^k$  in  $H^{s-1}(\mathbb{R}^n \times \mathbb{T}^m)$  and the condition (2.14), we know that  $u \in \mathcal{N}_T = \{0\}$  and  $1 \leq 2C_0 C T \|u\|_{H^{s-1}(\mathbb{R}^n \times \mathbb{T}^m)}^2$ , which is a contradiction. Hence, the observability (2.9) holds.  $\square$

Now we are in a position to prove Lemma 2.10.

*Proof of Lemma 2.10.* By the definition of  $\mathcal{N}_T$ , for any  $\delta > 0$ , we deduce that  $\mathcal{N}_{T+\delta} \subset \mathcal{N}_T$ . Then, we claim that  $\mathcal{N}_T$  is of finite dimension  $\forall T > 0$ . Indeed, thanks to the

weak observability and the definition of  $\mathcal{N}_T$ ,  $\forall f \in \mathcal{N}_T$ ,

$$\begin{aligned} \|f\|_{H^s(\mathbb{R}^n \times \mathbb{T}^m)}^2 &\leq 2C_0 \int_0^T \|\chi_\Omega e^{it\Delta_{x,y}} f\|_{H^s(\mathbb{R}^n \times \mathbb{T}^m)}^2 dt + 2C_0 CT \|f\|_{H^{s-1}(\mathbb{R}^n \times \mathbb{T}^m)}^2 \\ &= 2C_0 CT \|f\|_{H^{s-1}(\mathbb{R}^n \times \mathbb{T}^m)}^2. \end{aligned}$$

This implies that  $\mathcal{N}_T$  is of finite dimension. Moreover, combining with the fact that  $\mathcal{N}_{T_0+\delta} \subset \mathcal{N}_{T_0}$ ,  $\forall \delta > 0$ , there exists  $\delta_0 > 0$  such that  $\forall \delta \in (0, \delta_0)$ ,  $\mathcal{N}_{T_0+\delta} = \mathcal{N}_{T_0}$ . For any  $\delta \in (0, \delta_0)$  and  $\forall f \in \mathcal{N}_{T_0+\delta} = \mathcal{N}_{T_0}$ ,  $\chi_\Omega e^{it\Delta_{x,y}} e^{i\delta\Delta_{x,y}} f = 0$ , for  $\forall t \in (0, T_0)$ . Consequently, we know that  $e^{i\delta\Delta_{x,y}} f \in \mathcal{N}_{T_0}$ . Thus,  $\Delta_{x,y} f = \lim_{\delta \rightarrow 0^+} \frac{e^{i\delta\Delta_{x,y}} f - f}{i\delta} \in \mathcal{N}_{T_0}$ , which implies that  $\mathcal{N}_{T_0}$  is stable under  $\Delta_{x,y}$ . Suppose  $\mathcal{N}_{T_0} \neq \{0\}$ . Since  $\mathcal{N}_{T_0}$  is of finite dimension, it must contain an eigenfunction of  $\Delta_{x,y}$ . Let us denote this eigenfunction by  $\phi \neq 0$  and its associated eigenvalue  $\lambda_0 \in \mathbb{R}$ . Thus,  $\phi$  satisfies

$$\Delta_{x,y}\phi = \lambda_0\phi, \chi_\Omega e^{i\lambda_0 t}\phi = 0, \forall t \in (0, T_0).$$

Applying Theorem 1.5, we know that  $\phi \equiv 0$ , which is a contradiction. This leads to  $\mathcal{N}_{T_0} = \{0\}$ .  $\square$

### 3. WELL-POSEDNESS OF NLS

In this section, we include some well-posedness results of NLS. As in Theorem 1.1 and Theorem 1.2, we consider a specific and popular model: 3D cubic NLS in energy space and higher.

**Theorem 3.1.** *Let  $T > 0$ ,  $s \geq 1$ , and  $\epsilon = \pm 1$ . For every  $f \in L^2([-T, T], H^s)$  and  $u_0 \in H^s$ , there is a unique solution  $u \in X_T^{s,b}$  to*

$$(3.1) \quad \begin{cases} i\partial_t u + \Delta u + \epsilon|u|^2 u = f & \text{on } [-T, T] \times \mathbb{R}^2 \times \mathbb{T} \\ u|_{t=0} = u_0 & \text{on } \mathbb{R}^2 \times \mathbb{T} \end{cases}$$

Moreover the flow map is Lipschitz on every bounded subset.

*Remark 3.2.* We can replace  $\mathbb{R}^2 \times \mathbb{T}$  by  $\mathbb{R} \times \mathbb{T}^2$  (still 3-dimensional manifold) and the proof follows in the same way since the corresponding estimates still work.

*Proof.* The proof is similar to the proof of Theorem 2.1 in [32]. In fact, our case is easier since we do not have a damping term. We only need to handle the source term  $f$ .

First, we notice that if  $f \in L^2([-T, T], H^s)$ , it also belongs to  $X_T^{s,-b'}$  as  $b' \geq 0$  due to the property of function spaces. Moreover, it suffices to consider positive times. The solution on  $[-T, 0]$  can be obtained similarly.

We consider the functional

$$\Phi(u)(t) = e^{it\Delta_x} u_0 - i \int_0^t e^{i(t-\tau)\Delta_x} [\lambda|u|^2 u + f](\tau) d\tau.$$

One can apply a standard fixed point argument on the Banach space  $X_T^{s,b}$ .

We recall that, for  $T \leq 1$  we have

$$\|e^{it\Delta_x} u_0\|_{X_T^{s,b}} \leq C \|u_0\|_{H^s}.$$

This handles the linear term. Next, regarding the source term, one can use  $\|\psi(t/T) \int e^{i(t-\tau)\Delta_x} F(\tau)\|_{X^{s,b}} \leq CT^{1-b-b'} \|F\|_{X^{s,-b'}}$  to deal with the source term  $f$ .

Then the proof can be reduced to the case without the source. We refer to Bourgain [5].  $\square$

For the nonlinear equation (3.1), we can decompose it near a linear solution  $\Psi$  given by

$$(3.2) \quad \begin{cases} i\partial_t \Psi + \Delta_{x,y} \Psi = f, \\ \Psi|_{t=0} = \Psi_0 \in H^s(\mathbb{R}^2 \times \mathbb{T}) \end{cases}$$

We write  $u = \Psi + v$ . Then  $v$  satisfies

$$(3.3) \quad \begin{cases} i\partial_t v + \Delta_{x,y} v + \epsilon \left( |\Psi|^2 \Psi + 2|\Psi|^2 v + \Psi^2 \bar{v} \right) = -\epsilon F(\Psi, v), \\ v|_{t=T} = 0. \end{cases}$$

where  $F(\Psi, v) := |v|^2 v + 2|v|^2 \Psi + v^2 \bar{\Psi}$ . In the following proposition, we present the well-posedness of  $v$ .

**Proposition 3.3.** *Let  $T > 0$ ,  $s \geq 1$ , and  $\epsilon = \pm 1$ . For every  $\Psi_0 \in H^s$ , there is a unique solution  $v \in X_T^{s,b}$  to the equation (3.3).*

*Proof.* Since the proof is very similar to Theorem 3.1, we only give a sketch for it.

First, we note that, for any  $\Psi_0 \in H^s$ , the solution  $\Psi$  to the linear equation (4.4) satisfies  $\Psi \in X_T^{s,b}$ . This observation allows us to control for  $\Psi$ .

Then, compared to Theorem 3.1, there is no source term (that is good for us) in (4.6) and there are more nonlinear terms. We note that all nonlinear terms are cubic and we can estimate them using Strichartz estimate in a standard way. The proof can be done as in Bourgain [5] since we already have control for  $\Psi$ .  $\square$

#### 4. LOCAL NULL CONTROLLABILITY OF NLS: PROOF OF THEOREM 1.2

**4.1. Control for linear problem: Hilbert uniqueness method.** Consider a linear Schrödinger equation

$$(4.1) \quad \begin{cases} i\partial_t u + \Delta u = \varphi_T \chi_\Omega (1 - \Delta_{x,y})^{-s} \varphi_T \chi_\Omega f & \text{on } \mathbb{R} \times \mathbb{R}^2 \times \mathbb{T}, \\ u|_{t=0} = u_0 & \text{on } \mathbb{R}^2 \times \mathbb{T}, \end{cases}$$

such that  $u(T) \equiv 0$ . Define the range operator  $R$  by

$$R : H^{-s}(\mathbb{R} \times \mathbb{R}^2 \times \mathbb{T}) \rightarrow H^s(\mathbb{R}^2 \times \mathbb{T}) \\ f \mapsto u_0.$$

Consider the adjoint system

$$(4.2) \quad \begin{cases} i\partial_t w + \Delta w = 0 & \text{on } \mathbb{R} \times \mathbb{R}^2 \times \mathbb{T}, \\ w|_{t=0} = w_0 & \text{on } \mathbb{R}^2 \times \mathbb{T}, \end{cases}$$

Define the solution operator  $S$  by

$$S : H^s(\mathbb{R}^2 \times \mathbb{T}) \rightarrow C^0(\mathbb{R}; L^2(\mathbb{R}^2 \times \mathbb{T})) \subset H^s(\mathbb{R} \times \mathbb{R}^2 \times \mathbb{T}) \\ w_0 \mapsto e^{it\Delta} w_0.$$

**Proposition 4.1.** *Let  $s \in \mathbb{R}$ . Let  $f \in H^{-s}(\mathbb{R} \times \mathbb{R}^2 \times \mathbb{T})$  and  $w_0 \in H^{-s}(\mathbb{R}^2 \times \mathbb{T})$ . Then the duality holds as follows:*

$$(4.3) \quad \langle \varphi_T \chi_\Omega (1 - \Delta_{x,y})^{-s} \varphi_T \chi_\Omega f, S w_0 \rangle_{H^s(\mathbb{R}^2 \times \mathbb{T}), H^{-s}(\mathbb{R}^2 \times \mathbb{T})} = \langle iR(f), w_0 \rangle_{H^s(\mathbb{R}^2 \times \mathbb{T}), H^{-s}(\mathbb{R}^2 \times \mathbb{T})}.$$

*Proof.* We only consider  $f$  and  $w_0$  are smooth in the space variables. Then the general case is proved by standard approximation arguments.

$$\begin{aligned} \langle \varphi_T \chi_\Omega (1 - \Delta_{x,y})^{-s} \varphi_T \chi_\Omega f, S w_0 \rangle_{H^s, H^{-s}} &= \langle i\partial_t u + \Delta u, e^{it\Delta} w_0 \rangle_{H^s, H^{-s}} \\ &= \langle u, (i\partial_t + \Delta) e^{it\Delta} w_0 \rangle_{H^s, H^{-s}} + \langle iu_0, w_0 \rangle_{H^s, H^{-s}} \\ &= \langle iR(f), w_0 \rangle_{H^s, H^{-s}}. \end{aligned}$$

$\square$

Then we can define the HUM operator  $\mathcal{K} = iR \circ S$ , which satisfies the following proposition.

**Proposition 4.2.**  *$\mathcal{K}$  is an isomorphism from  $H^{-s}$  to  $H^s$ .*

*Proof.* We define a continuous form by  $\alpha(u, v) := \langle \mathcal{K}u, v \rangle_{H^s(\mathbb{R}^2 \times \mathbb{T}), H^{-s}(\mathbb{R}^2 \times \mathbb{T})}$ , for  $u, v \in H^{-s}(\mathbb{R}^2 \times \mathbb{T})$ . By the definition of  $\mathcal{K}$ , it is easy to verify that  $\mathcal{K}$  is self-adjoint operator. Then we check the coercivity of  $\alpha$ . Thanks to the observability estimate in Theorem 1.5, for  $\forall v \in L^2(\mathbb{R}^2 \times \mathbb{T})$ ,

$$\begin{aligned} \alpha(v, v) &= \langle \mathcal{K}v, v \rangle_{H^s, H^{-s}} = \left( (1 - \Delta_{x,y})^{-\frac{s}{2}} \varphi_T \chi_\Omega S v, (1 - \Delta_{x,y})^{-\frac{s}{2}} \varphi_T \chi_\Omega S v \right)_{L^2(\mathbb{R}^2 \times \mathbb{T})} \\ &= \|\varphi_T \chi_\Omega S v\|_{H^{-s}(\mathbb{R}^2 \times \mathbb{T})}^2. \end{aligned}$$

By Proposition 2.9, there exists a constant  $C = C(T, \Omega, s, \chi_\Omega) > 0$  such that

$$\|\varphi_T \chi_\Omega S v\|_{H^{-s}(\mathbb{R}^2 \times \mathbb{T})}^2 \geq C \|v\|_{H^{-s}(\mathbb{R}^2 \times \mathbb{T})}^2.$$

By Lax-Milgram's theorem, we prove that the HUM operator  $\mathcal{K}$  is an isomorphism  $H^{-s}$  to  $H^s$ .  $\square$

Define the control operator  $\mathcal{L}$  by  $\mathcal{L} = iS \circ \mathcal{K}^{-1} : H^s \rightarrow H^{-s}$ . Since  $R \circ \mathcal{L}\Psi_0 = \Psi_0$ , the solution  $\Psi$  to the linear Schrödinger equation with an initial datum  $\Psi_0$

$$(4.4) \quad \begin{cases} i\partial_t \Psi + \Delta_{x,y} \Psi = \varphi_T \chi_\Omega (1 - \Delta_{x,y})^{-s} \varphi_T \chi_\Omega \mathcal{L}\Psi_0, \\ \Psi|_{t=0} = \Psi_0 \in H^s(\mathbb{R}^2 \times \mathbb{T}) \end{cases}$$

satisfies that  $\Psi(T) = 0$ .

**4.2. Proof of Theorem 1.2.** Now we consider the null control problem for NLS. Recall that we seek a control function  $g$  such that the solution  $u$  to

$$(4.5) \quad \begin{cases} i\partial_t u + \Delta_{x,y} u + \epsilon |u|^2 u = \varphi_T \chi_\Omega (1 - \Delta_{x,y})^{-s} \varphi_T \chi_\Omega g & \text{on } [0, T] \times \mathbb{R}^2 \times \mathbb{T}, \\ u|_{t=0} = u_0 & \text{on } \mathbb{R}^2 \times \mathbb{T}, \end{cases}$$

such that  $u|_{t=T} = 0$ .

*Proof of Theorem 1.2.* First, we decompose  $u = v + \Psi$ , where  $\Psi$  is defined by (4.4). Then,  $v$  is the solution to

$$(4.6) \quad \begin{cases} i\partial_t v + \Delta_{x,y} v + \epsilon \left( |\Psi|^2 \Psi + 2|\Psi|^2 v + \Psi^2 \bar{v} \right) = -\epsilon F(\Psi, v), \\ v|_{t=T} = 0. \end{cases}$$

where  $F(\Psi, v) := |v|^2 v + 2|v|^2 \Psi + v^2 \bar{\Psi}$ . Thus,  $i\partial_t v + \Delta_{x,y} v + \epsilon |\Psi + v|^2 (\Psi + v) = 0$ . Thanks to Proposition 3.3, we know that  $v \in X_T^{s,b}$ . Let us define a nonlinear solution operator  $S_N^v : H^s \rightarrow H^s$  associated with the equation (4.6) by  $S_N^v \Psi_0 = v|_{t=0}$ , where  $v$  solves the equation (4.6) and  $\Psi$  solves the equation (4.4) with an initial datum  $\Psi_0$ . Due to  $u = v + \Psi$ ,  $u$  satisfies

$$(4.7) \quad \begin{cases} i\partial_t u + \Delta_{x,y} u + \epsilon |u|^2 u = \varphi_T \chi_\Omega (1 - \Delta_{x,y})^{-s} \varphi_T \chi_\Omega \mathcal{L}\Psi_0, \\ u|_{t=T} = v|_{t=T} + \Psi|_{t=T} = 0. \end{cases}$$

As a consequence, for any  $\Psi_0 \in H^s$ , the control function  $g = \mathcal{L}\Psi_0$  steers the initial state  $\Psi|_{t=0} + v|_{t=0}$  to 0.

Define another nonlinear operator  $\mathcal{J} : H^s \rightarrow H^s$  by  $\mathcal{J}\vartheta = u_0 - S_N^v \vartheta$ . If  $\mathcal{J}$  has a fixed point  $\Psi_0$ , i.e.,  $\mathcal{J}\Psi_0 = \Psi_0$ , then, this  $\Psi_0$  produces a function  $g = \mathcal{L}\Psi_0$ , which achieves the null controllability, due to  $u_0 = \Psi|_{t=0} + v|_{t=0} = \Psi_0 + S_N^v \Psi_0$ . Hence, the null controllability of nonlinear Schrödinger equation (1.3) is reduced to the following lemma.

**Lemma 4.3.** *There exists a constant  $\eta > 0$  such that  $\mathcal{J}$  has a fixed point in a ball  $B_{H^s}(0, \eta)$  in  $H^s$ .*

□

We complete the proof of Theorem 1.2 by demonstrating Lemma 4.3.

*Proof of Lemma 4.3.* We only need to prove that  $\mathcal{J}$  is contracting on a small ball  $B_{H^s}(0, \eta)$  in  $H^s$ , provided with  $\|u_0\|_{H^s}$  small enough. In this proof, for simplicity, we can assume that  $T < 1$ . Note that our constant  $C$  which may depend on  $T$  and  $s$  could vary from line to line.

For any  $\Psi_0 \in B_{H^s}(0, \eta)$ , we first show that  $\mathcal{J}\Psi_0 \in B_{H^s}(0, \eta)$  for  $\eta$  sufficiently small. By the definition of the map  $\mathcal{J}$ ,  $\|\mathcal{J}\Psi_0\|_{H^s} \leq \|u|_{t=0}\|_{H^s} + \|v|_{t=0}\|_{H^s}$ . Thanks to Proposition 3.3,

$$\|v\|_{X_T^{s,b}} \leq \| |u|^2 u \|_{X_T^{s,-b'}} \leq \|u\|_{X_T^{s,b'}}^3.$$

Applying Theorem 3.1 and Proposition 4.2, we obtain

$$\|u\|_{X_T^{s,b'}} \leq C\|\mathcal{L}\Psi_0\|_{H^{-s}} \leq C\|\Psi_0\|_{H^s} < C\eta.$$

As a consequence, we obtain

$$\|v|_{t=0}\|_{H^s} \leq \|v\|_{X_T^{s,b}} \leq \|u\|_{X_T^{s,b'}}^3 < C\eta^3.$$

Thus,  $\|\mathcal{J}\Psi_0\|_{H^s} \leq \|u_0\|_{H^s} + \|v|_{t=0}\|_{H^s} \leq \|u_0\|_{H^s} + C\eta^3$ . Choosing  $\eta$  such that  $C\eta^2 < \frac{1}{2}$  and  $\|u_0\|_{H^s} < \frac{\eta}{2}$ , we obtain  $\|\mathcal{J}\Psi_0\|_{H^s} < \eta$  and  $\mathcal{J}$  reproduces the ball  $B_{H^s}(0, \eta)$ .

Let us prove that  $\mathcal{J}$  is contracting for  $H^s$ -norm. For any  $\vartheta, \tilde{\vartheta} \in B_{H^s}(0, \eta)$ , let  $u$  and  $\tilde{u}$  be the solutions to (4.5) with  $g = \mathcal{L}\vartheta$  and  $g = \mathcal{L}\tilde{\vartheta}$ , respectively. Similarly, we denote by  $v$  and  $\tilde{v}$ . Then,  $u - \tilde{u}$  solves the equation:

$$\begin{cases} i\partial_t(u - \tilde{u}) + \Delta_{x,y}(u - \tilde{u}) + \epsilon|u|^2u - \epsilon|\tilde{u}|^2\tilde{u} = \varphi_T\chi_\Omega(1 - \Delta_{x,y})^{-s}\varphi_T\chi_\Omega\mathcal{L}(\vartheta - \tilde{\vartheta}), \\ (u - \tilde{u})|_{t=0} = (u - \tilde{u})|_{t=T} = 0. \end{cases}$$

And  $v - \tilde{v}$  solves the equation:

$$\begin{cases} i\partial_t(v - \tilde{v}) + \Delta_{x,y}(v - \tilde{v}) + \epsilon|u|^2u - \epsilon|\tilde{u}|^2\tilde{u} = 0, \\ (v - \tilde{v})|_{t=T} = 0. \end{cases}$$

As we presented in the proof of Theorem 3.1, we have

$$\begin{aligned} \|u - \tilde{u}\|_{X_T^{s,b}} &\leq C\| |u|^2u - |\tilde{u}|^2\tilde{u} \|_{X_T^{s,-b'}} + C\|\mathcal{L}(\vartheta - \tilde{\vartheta})\|_{H^{-s}} \\ &\leq C(\|u\|_{X_T^{s,b'}}^2 + \|\tilde{u}\|_{X_T^{s,b'}}^2)\|u - \tilde{u}\|_{X_T^{s,b'}} + C\|\vartheta - \tilde{\vartheta}\|_{H^s} \\ &\leq C\eta^2\|u - \tilde{u}\|_{X_T^{s,b'}} + C\|\vartheta - \tilde{\vartheta}\|_{H^s}. \end{aligned}$$

For  $\eta$  sufficiently small, it yields  $C\eta^2 < \frac{1}{2}$ , and we obtain  $\|u - \tilde{u}\|_{X_T^{s,b}} \leq C\|\vartheta - \tilde{\vartheta}\|_{H^s}$ . Similarly, for  $v - \tilde{v}$ , we have

$$\begin{aligned} \|v - \tilde{v}\|_{X_T^{s,b}} &\leq C\| |u|^2u - |\tilde{u}|^2\tilde{u} \|_{X_T^{s,-b'}} \\ &\leq C\eta^2\|u - \tilde{u}\|_{X_T^{s,b'}} \\ &\leq C\eta^2\|\vartheta - \tilde{\vartheta}\|_{H^s}. \end{aligned}$$

We deduce that  $\|v|_{t=0} - \tilde{v}|_{t=0}\|_{H^s} \leq \|v - \tilde{v}\|_{X_T^{s,b}} < C\eta^2\|\vartheta - \tilde{\vartheta}\|_{H^s}$ . Therefore, we obtain

$$\begin{aligned} \|\mathcal{J}\vartheta - \mathcal{J}\tilde{\vartheta}\|_{H^s} &= \|(u_0 - S_N^v\vartheta) - (u_0 - S_N^v\tilde{\vartheta})\|_{H^s} \\ &= \|v|_{t=0} - \tilde{v}|_{t=0}\|_{H^s} \\ &\leq C\eta^2\|\vartheta - \tilde{\vartheta}\|_{H^s} \end{aligned}$$

This yields that  $\mathcal{J}$  is a contraction on a small ball  $B_{H^s}(0, \eta)$  in  $H^s$ , which completes the proof. □

## 5. EXACT CONTROLLABILITY: PROOF OF THEOREM 1.1

In general, in order to Theorem 1.1, we only need to reverse time and glue the forward and backward solutions together. For the completeness of our paper, we choose to present the proof as follows.

*Proof of Theorem 1.1.* We first consider the time-reversed equation (that is, the equation obtained by the change of variable  $t \mapsto T - t$ ) of (4.5):

$$-i\partial_t w + \Delta_{x,y} w + \epsilon|w|^2 w = \varphi_T \chi_\Omega (1 - \Delta_{x,y})^{-s} \varphi_T \chi_\Omega h,$$

where  $w(t, x) = u(T - t, x)$ . For this equation, we repeated the procedure as the proof for Theorem 1.2. There exists a control function  $h$  such that for  $\|w_0\|_{H^s}$  small enough,  $h$  achieves null controllability for  $w$ . Now we choose  $u_0, u_f \in H^s$ , satisfying that

$$\|u_0\|_{H^s} + \|u_f\|_{H^s} < \delta,$$

with  $\delta$  sufficiently small. There exists  $g$  such that

$$\begin{cases} i\partial_t u + \Delta_{x,y} u + \epsilon|u|^2 u = \varphi_T \chi_\Omega (1 - \Delta_{x,y})^{-s} \varphi_T \chi_\Omega g \\ u|_{t=0} = u_0 \quad u|_{t=\frac{T}{2}} = 0. \end{cases}$$

And there exists  $h$  such that

$$\begin{cases} -i\partial_t w + \Delta_{x,y} w + \epsilon|w|^2 w = \varphi_T \chi_\Omega (1 - \Delta_{x,y})^{-s} \varphi_T \chi_\Omega h \\ w|_{t=0} = u_f \quad w|_{t=\frac{T}{2}} = 0. \end{cases}$$

Moreover, we know that the solutions  $u, w \in C([0, \frac{T}{2}], H^s)$ . Now we define  $U \in C([0, T], H^s)$  and  $f \in C([0, T], H^s)$  by

$$U(t) = \begin{cases} u(t) & t \in [0, \frac{T}{2}], \\ w(T-t) & t \in (\frac{T}{2}, T]. \end{cases} \quad \text{and}$$

$$f(t) = \begin{cases} \varphi_T \chi_\Omega (1 - \Delta_{x,y})^{-s} \varphi_T \chi_\Omega g, & t \in [0, \frac{T}{2}], \\ \varphi_T (T-t) \chi_\Omega (1 - \Delta_{x,y})^{-s} \varphi_T (T-t) \chi_\Omega h(T-t), & t \in (\frac{T}{2}, T]. \end{cases}$$

Indeed,  $f$  continues in time, since the cut-off function vanishes near  $t = \frac{T}{2}$ .  $U$  solves the equation:

$$(5.1) \quad \begin{cases} i\partial_t U + \Delta_{x,y} U + \epsilon|U|^2 U = f, \\ U|_{t=0} = u|_{t=0} = u_0, \\ U|_{t=T} = w(T-t)|_{t=T} = w|_{t=0} = u_f. \end{cases}$$

This completes the proof of Theorem 1.1.  $\square$

APPENDIX A. SOME PROPERTIES OF  $X^{s,b}$  SPACES

For the sake of completeness, in Appendix, we include the definitions for  $X^{s,b}$  spaces in the waveguide setting, together with some useful properties. We refer to Tao [41] for more details. See also [32]. Since the results are well-known, we only present them and omit the proofs.

Following [5, 6], one may define  $X^{s,b}$  norm as

$$(A.1) \quad \|u\|_{X^{s,b}} := \|\langle \tau - |\xi|^2 \rangle^b \langle \xi \rangle^s \tilde{u}\|_{L^2_{\xi,\tau}},$$

where  $u = u(z, t)$  is a function defined on  $\mathbb{R}^m \times \mathbb{T}^n \times \mathbb{R}$ , and  $z \in \mathbb{R}^m \times \mathbb{T}^n, t \in \mathbb{R}$ . And  $\tilde{u}(\xi, \tau)$  is the space-time Fourier transform of  $u$ , where  $\xi \in \mathbb{R}^m \times \mathbb{Z}^n, \tau \in \mathbb{R}$ .

And  $X^{s,b}$  spaces are just all those functions with finite  $X^{s,b}$  norm.

In practice, one mainly works on  $s \geq 0, b > \frac{1}{2}$ .

One key property for  $X^{s,b}$  space is that inherits the estimates of linear solutions, which is known to be ‘‘transference principle’’.

$X_T^{s,b}$  is the associated restriction space with the norm

$$\|u\|_{X_T^{s,b}} = \inf \{ \|\tilde{u}\|_{X^{s,b}} \mid \tilde{u} = u \text{ on } (0, T) \times M \}.$$

We also write  $\|u\|_{X_T^{s,b}}$  if the infimum is taken on functions  $\tilde{u}$  equalling  $u$  on an interval

$I$ . The following properties of  $X_T^{s,b}$  spaces are easily verified.

- (1)  $X^{s,b}$  and  $X_T^{s,b}$  are Hilbert spaces.
- (2) If  $s_1 \leq s_2$ ,  $b_1 \leq b_2$  we have  $X^{s_2, b_2} \subset X^{s_1, b_1}$  with continuous embedding.
- (3) For every  $s_1 < s_2$ ,  $b_1 < b_2$  and  $T > 0$ , we have  $X_T^{s_2, b_2} \subset X_T^{s_1, b_1}$  with compact embedding.
- (4) For  $0 < \theta < 1$ , the complex interpolation space  $(X^{s_1, b_1}, X^{s_2, b_2})_{[\theta]}$  is  $X^{(1-\theta)s_1 + \theta s_2, (1-\theta)b_1 + \theta b_2}$ .

We note that (4) can be proved with the interpolation theorem of Stein-Weiss for weighted  $L^p$  spaces.

Then, we list some additional trilinear estimates that will be used all along the paper.

**Lemma A.1.** *For every  $r \geq s > s_0$ , there exist  $0 < b' < 1/2$  and  $C > 0$  such that for any  $u$  and  $\tilde{u} \in X^{r, b'}$*

$$(A.2) \quad \| |u|^2 u \|_{X^{r, -b'}} \leq C \|u\|_{X^{s, b'}}^2 \|u\|_{X^{r, b'}}$$

$$(A.3) \quad \| |u|^2 \tilde{u} \|_{X^{r, -b'}} \leq C \|u\|_{X^{s, b'}} \|u\|_{X^{r, b'}} \|\tilde{u}\|_{X^{r, b'}}$$

$$(A.4) \quad \| |u|^2 u - |\tilde{u}|^2 \tilde{u} \|_{X^{s, -b'}} \leq C \left( \|u\|_{X^{s, b'}}^2 + \|\tilde{u}\|_{X^{s, b'}}^2 \right) \|u - \tilde{u}\|_{X^{s, b'}}.$$

Moreover, the same estimates hold with  $z_1 \bar{z}_2 z_3$  replaced by any  $\mathbb{R}$ -trilinear form on  $\mathbb{C}$ .

**Lemma A.2.** *For every  $-1 \leq s \leq 1$  and any  $s_0 < r \leq 1$ , there exist  $0 < b' < 1/2$  and  $C > 0$  such that for any  $u \in X^{s, b'}$  and  $a_1, a_2 \in X^{1, b'}$*

$$(A.5) \quad \| a_1 \bar{a}_2 u \|_{X^{s, -b'}} \leq C \|a_1\|_{X^{1, b'}} \|a_2\|_{X^{1, b'}} \|u\|_{X^{s, b'}}$$

$$(A.6) \quad \| |a_1|^2 u \|_{X^{s, -b'}} \leq C \|a_1\|_{X^{1, b'}} \|a_1\|_{X^{r, b'}} \|u\|_{X^{s, b'}}.$$

Moreover, the same estimates hold with  $z_1 \bar{z}_2 z_3$  replaced by any  $\mathbb{R}$ -trilinear form on  $\mathbb{C}$ .

Let us study the stability of the  $X^{s,b}$  spaces with respect to some particular operations.

**Lemma A.3.** *Let  $\varphi \in C_0^\infty(\mathbb{R})$  and  $u \in X^{s,b}$  then  $\varphi(t)u \in X^{s,b}$ .*

*If  $u \in X_T^{s,b}$  then we have  $\varphi(t)u \in X_T^{s,b}$ .*

In the case of pseudo-differential operators in the space variable, we have to deal with a loss in  $X^{s,b}$  regularity compared to what we could expect. Some regularity in the index  $b$  is lost, due to the fact that a pseudo-differential operator does not keep the structure in time of the harmonics.

This loss is unavoidable as we can see, for simplicity on the torus  $\mathbb{T}^1$ : we take  $u_n = \psi(t) e^{inx} e^{i|n^2|t}$  (where  $\psi \in C_0^\infty$  equal to 1 on  $[-1, 1]$ ) which is uniformly bounded in  $X^{0,b}$  for every  $b \geq 0$ . Yet, if we consider the operator  $B$  of order 0 of multiplication by  $e^{ix}$ , we get  $\|e^{ix} u_n\|_{X^{0,b}} \approx n^b$ . Yet, we do not have such loss for operator of the form  $(-\Delta)^r$  which acts from any  $X^{s,b}$  to  $X^{s-2r,b}$ . But if we do not make any further assumption on the pseudo-differential operator, we can show that our example is the worst one:

**Lemma A.4.** *Let  $-1 \leq b \leq 1$  and  $B$  be a pseudo-differential operator in the space variable of order  $\rho$ . For any  $u \in X^{s,b}$  we have  $Bu \in X^{s-\rho-|b|, b}$ .*

*Similarly,  $B$  maps  $X^{s,b}t$  into  $X_T^{s-\rho-|b|, b}$ .*

We will also use the following elementary estimate.

**Lemma A.5.** *Let  $(b, b')$  satisfying*

$$(A.7) \quad 0 < b' < \frac{1}{2} < b, \quad b + b' \leq 1.$$

If we note  $F(t) = \Psi\left(\frac{t}{T}\right) \int_0^t f(t') dt'$ , we have for  $T \leq 1$

$$\|F\|_{H^b} \leq CT^{1-b-b'} \|f\|_{H^{-b'}}.$$

In the future aim of using a boot-strap argument, we will need some continuity in  $T$  of the  $X_T^{s,b}$  norm of a fixed function :

**Lemma A.6.** *Let  $0 < b < 1$  and  $u$  in  $X^{s,b}$  then the function*

$$\begin{cases} f & : (0, T] \longrightarrow \mathbb{R} \\ t & \longmapsto \|u\|_{X_t^{s,b}} \end{cases}$$

is continuous. Moreover, if  $b > 1/2$ , there exists  $C_b$  such that

$$\lim_{t \rightarrow 0} f(t) \leq C_b \|u(0)\|_{H^s}.$$

The following lemma will be useful to control solutions on large intervals that will be obtained by piecing together solutions on smaller ones.

**Lemma A.7.** *Let  $0 < b < 1$ . If  $\bigcup(a_k, b_k)$  is a finite covering of  $[0, 1]$ , then there exists a constant  $C$  depending only of the covering such that for every  $u \in X^{s,b}$ ,*

$$\|u\|_{X_{[0,1]}^{s,b}} \leq C \sum_k \|u\|_{X_{[a_k, b_k]}^{s,b}}.$$

#### REFERENCES

- [1] N. Anantharaman, M. Léautaud, and F. Macià. Wigner measures and observability for the Schrödinger equation on the disk. *Invent. Math.*, 206(2):485–599, 2016.
- [2] N. Anantharaman and F. Macià. Semiclassical measures for the Schrödinger equation on the torus. *J. Eur. Math. Soc. (JEMS)*, 16(6):1253–1288, 2014.
- [3] N. Anantharaman and G. Rivière. Dispersion and controllability for the Schrödinger equation on negatively curved manifolds. *Anal. PDE*, 5(2):313–338, 2012.
- [4] K. Le Balc'h and J. Martin. Observability estimates for the Schrödinger equation in the plane with periodic bounded potentials from measurable sets, 2023.
- [5] J. Bourgain. Fourier transform restriction phenomena for certain lattice subsets and applications to nonlinear evolution equations. II. The KdV-equation. *Geom. Funct. Anal.*, 3(3):209–262, 1993.
- [6] J. Bourgain. Refinements of Strichartz inequality and applications to 2D-NLS with critical nonlinearity. *International Mathematics Research Notices*, 1998(5):253–283, 1998.
- [7] J. Bourgain, N. Burq, and M. Zworski. Control for Schrödinger operators on 2-tori: rough potentials. *J. Eur. Math. Soc. (JEMS)*, 15(5):1597–1628, 2013.
- [8] N. Burq and C. Sun. Time optimal observability for Grushin Schrödinger equation. *Anal. PDE*, 15(6):1487–1530, 2022.
- [9] N. Burq and M. Zworski. Control for Schrödinger operators on tori. *Math. Res. Lett.*, 19(2):309–324, 2012.
- [10] X. Cheng, Z. Guo, K. Yang, and L. Zhao. On scattering for the cubic defocusing nonlinear Schrödinger equation on the waveguide  $\mathbb{R}^2 \times \mathbb{T}$ . *Rev. Mat. Iberoam.*, 36(4):985–1011, 2020.
- [11] X. Cheng, Z. Guo, and Z. Zhao. On scattering for the defocusing quintic nonlinear Schrödinger equation on the two-dimensional cylinder. *SIAM J. Math. Anal.*, 52(5):4185–4237, 2020.
- [12] X. Cheng, Z. Zhao, and J. Zheng. Well-posedness for energy-critical nonlinear Schrödinger equation on waveguide manifold. *J. Math. Anal. Appl.*, 494(2):Paper No. 124654, 14, 2021.
- [13] F. M. Christ and M. I. Weinstein. Dispersion of small amplitude solutions of the generalized Korteweg-de Vries equation. *J. Funct. Anal.*, 100(1):87–109, 1991.
- [14] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao. Global well-posedness for Schrödinger equations with derivative. *SIAM J. Math. Anal.*, 33(3):649–669, 2001.
- [15] B. Dehman, P. Gérard, and G. Lebeau. Stabilization and control for the nonlinear Schrödinger equation on a compact surface. *Math. Z.*, 254(4):729–749, 2006.



- [16] B. Dehman, P. Gérard, and G. Lebeau. Stabilization and control for the nonlinear Schrödinger equation on a compact surface. *Math. Z.*, 254(4):729–749, 2006.
- [17] B. Dodson. Global well-posedness and scattering for the defocusing,  $L^2$ -critical nonlinear Schrödinger equation when  $d \geq 3$ . *J. Amer. Math. Soc.*, 25(2):429–463, 2012.
- [18] C. Fan and Z. Zhao. On long time behavior for stochastic nonlinear schrödinger equations with a multiplicative noise. *International Mathematics Research Notices*, page rnae035, 2024.
- [19] L. Gagnon. Ground state solitary waves local controllability for the nonlinear focusing Schrödinger equation in the mass critical and mass slightly subcritical case. *J. Differential Equations*, 376:235–282, 2023.
- [20] Z. Hani and B. Pausader. On scattering for the quintic defocusing nonlinear Schrödinger equation on  $\mathbb{R} \times \mathbb{T}^2$ . *Comm. Pure Appl. Math.*, 67(9):1466–1542, 2014.
- [21] Z. Hani, B. Pausader, N. Tzvetkov, and N. Visciglia. Modified scattering for the cubic Schrödinger equation on product spaces and applications. *Forum Math. Pi*, 3:e4, 63, 2015.
- [22] S. Herr, D. Tataru, and N. Tzvetkov. Global well-posedness of the energy-critical nonlinear Schrödinger equation with small initial data in  $H^1(\mathbb{T}^3)$ . *Duke Math. J.*, 159(2):329–349, 2011.
- [23] S. Herr, D. Tataru, and N. Tzvetkov. Strichartz estimates for partially periodic solutions to Schrödinger equations in  $4d$  and applications. *J. Reine Angew. Math.*, 690:65–78, 2014.
- [24] S. Huang, G. Wang, and M. Wang. Observable sets, potentials and Schrödinger equations. *Comm. Math. Phys.*, 395(3):1297–1343, 2022.
- [25] A. D. Ionescu and B. Pausader. The energy-critical defocusing NLS on  $\mathbb{T}^3$ . *Duke Math. J.*, 161(8):1581–1612, 2012.
- [26] A. D. Ionescu and B. Pausader. Global well-posedness of the energy-critical defocusing NLS on  $\mathbb{R} \times \mathbb{T}^3$ . *Comm. Math. Phys.*, 312(3):781–831, 2012.
- [27] S. Jaffard. Contrôle interne exact des vibrations d’une plaque rectangulaire. *Portugal. Math.*, 47(4):423–429, 1990.
- [28] L. Jin. Control for Schrödinger equation on hyperbolic surfaces. *Math. Res. Lett.*, 25(6):1865–1877, 2018.
- [29] C. Kenig and F. Merle. Scattering for  $H^{1/2}$  bounded solutions to the cubic, defocusing NLS in 3 dimensions. *Transactions of the American Mathematical Society*, 362(4):1937–1962, 2010.
- [30] R. Killip and M. Viřan. Scale invariant Strichartz estimates on tori and applications. *Math. Res. Lett.*, 23(2):445–472, 2016.
- [31] P. Kuchment. *Floquet theory for partial differential equations*, volume 60 of *Operator Theory: Advances and Applications*. Birkhäuser Verlag, Basel, 1993.
- [32] C. Laurent. Global controllability and stabilization for the nonlinear Schrödinger equation on an interval. *ESAIM: Control, Optimisation and Calculus of Variations*, 16(2):356–379, 2010.
- [33] C. Laurent. Global controllability and stabilization for the nonlinear Schrödinger equation on some compact manifolds of dimension 3. *SIAM J. Math. Anal.*, 42(2):785–832, 2010.
- [34] C. Laurent. Internal control of the Schrödinger equation. *Math. Control Relat. Fields*, 4(2):161–186, 2014.
- [35] G. Lebeau. Contrôle de l’équation de Schrödinger. *J. Math. Pures Appl. (9)*, 71(3):267–291, 1992.
- [36] J-L. Lions. *Contrôlabilité exacte, perturbations et stabilisation de systèmes distribués. Tome 1*, volume 8 of *Recherches en Mathématiques Appliquées [Research in Applied Mathematics]*. Masson, Paris, 1988. Contrôlabilité exacte. [Exact controllability], With appendices by E. Zuazua, C. Bardos, G. Lebeau and J. Rauch.
- [37] L. Miller. Controllability cost of conservative systems: resolvent condition and transmutation. *J. Funct. Anal.*, 218(2):425–444, 2005.
- [38] A. Prouff. Observability of the Schrödinger equation with subquadratic confining potential in the euclidean space, 2023.
- [39] L. Rosier and B. Zhang. Exact boundary controllability of the nonlinear Schrödinger equation. *J. Differential Equations*, 246(10):4129–4153, 2009.
- [40] L. Rosier and B. Zhang. Local exact controllability and stabilizability of the nonlinear Schrödinger equation on a bounded interval. *SIAM J. Control Optim.*, 48(2):972–992, 2009.
- [41] T. Tao. *Nonlinear dispersive equations*, volume 106 of *CBMS Regional Conference Series in Mathematics*. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 2006. Local and global analysis.
- [42] M. Täufer. Controllability of the Schrödinger equation on unbounded domains without geometric control condition. *ESAIM Control Optim. Calc. Var.*, 29:Paper No. 59, 11, 2023.
- [43] N. Tzvetkov and N. Visciglia. Small data scattering for the nonlinear Schrödinger equation on product spaces. *Comm. Partial Differential Equations*, 37(1):125–135, 2012.
- [44] N. Tzvetkov and N. Visciglia. Well-posedness and scattering for nonlinear Schrödinger equations on  $\mathbb{R}^d \times \mathbb{T}$  in the energy space. *Rev. Mat. Iberoam.*, 32(4):1163–1188, 2016.

- [45] G. Wang, M. Wang, and Y. Zhang. Observability and unique continuation inequalities for the Schrödinger equation. *J. Eur. Math. Soc. (JEMS)*, 21(11):3513–3572, 2019.
- [46] J. Wunsch. Periodic damping gives polynomial energy decay. *Math. Res. Lett.*, 24(2):571–580, 2017.
- [47] K. Yang and Z. Zhao. On scattering asymptotics for the 2D cubic resonant system. *Journal of Differential Equations*, 345:447–484, 2023.
- [48] H. Yue. Global well-posedness for the energy-critical focusing nonlinear Schrödinger equation on  $\mathbb{T}^4$ . *J. Differential Equations*, 280:754–804, 2021.
- [49] Z. Zhao. Global well-posedness and scattering for the defocusing cubic Schrödinger equation on waveguide  $\mathbb{R}^2 \times \mathbb{T}^2$ . *J. Hyperbolic Differ. Equ.*, 16(1):73–129, 2019.
- [50] Z. Zhao. On scattering for the defocusing nonlinear Schrödinger equation on waveguide  $\mathbb{R}^m \times \mathbb{T}$  (when  $m = 2, 3$ ). *J. Differential Equations*, 275:598–637, 2021.
- [51] Z. Zhao and J. Zheng. Long time dynamics for defocusing cubic nonlinear Schrödinger equations on three dimensional product space. *SIAM J. Math. Anal.*, 53(3):3644–3660, 2021.
- [52] E. Zuazua. Remarks on the controllability of the Schrödinger equation. In *Quantum control: mathematical and numerical challenges*, volume 33 of *CRM Proc. Lecture Notes*, pages 193–211. Amer. Math. Soc., Providence, RI, 2003.

JINGRUI NIU

SORBONNE UNIVERSITÉ, CNRS, UNIVERSITÉ PARIS CITÉ, INRIA TEAM CAGE, LABORATOIRE JACQUES-LOUIS LIONS (LJLL), F-75005 PARIS, FRANCE

*Email address:* `jingrui.niu@sorbonne-universite.fr`

ZEHUA ZHAO

DEPARTMENT OF MATHEMATICS AND STATISTICS, BEIJING INSTITUTE OF TECHNOLOGY, BEIJING, CHINA.

KEY LABORATORY OF ALGEBRAIC LIE THEORY AND ANALYSIS OF MINISTRY OF EDUCATION, BEIJING, CHINA.

*Email address:* `zzh@bit.edu.cn`