STABILITY OF COMPOSITE WAVE OF PLANAR VISCOUS SHOCK AND RAREFACTION FOR 3D BAROTROPIC NAVIER-STOKES EQUATIONS

JIAJIN SHI AND YI WANG

We prove the nonlinear time-asymptotic stability of the composite wave consisting of a planar rarefaction wave and a planar viscous shock for the three-dimensional (3D) compressible barotropic Navier-Stokes equations under generic perturbations, in particular, without zero-mass conditions. It is shown that if the composite wave strength and the initial perturbations are suitably small, then 3D Navier-Stokes system admits a unique global-in-time strong solution which time-asymptotically converges to the corresponding composite wave up to a time-dependent shift for planar viscous shock. Our proof is based on the *a*-contraction method with time-dependent shift and suitable weight function.

1. INTRODUCTION

We are concerned with the large-time behavior of the solution to 3D compressible barotropic Navier-Stokes equations

(1.1)
$$\begin{cases} \partial_t \rho + \operatorname{div}(\rho u) = 0, & x \in \Omega, \ t \ge 0, \\ \partial_t(\rho u) + \operatorname{div}(\rho u \otimes u) + \nabla p(\rho) = \mu \Delta u + (\mu + \lambda) \nabla \operatorname{div} u, \end{cases}$$

where the unknowns $\rho = \rho(t, x) > 0$ and u = u(t, x) represent respectively the mass density and the velocity of the fluid with $x = (x_1, x_2, x_3) \in \Omega := \mathbb{R} \times \mathbb{T}^2$ and $\mathbb{T} := \mathbb{R}/\mathbb{Z}$. The pressure $p = p(\rho)$ is given by the well-known γ -law

$$p(\rho) = b\rho^{\gamma}$$

where b > 0, $\gamma > 1$ are both constants depending on the fluid. Without loss of generality, we will take b = 1 hereafter. The constants μ and λ are viscosity coefficients satisfying the following physical constraints

$$\mu > 0, \quad 2\mu + 3\lambda \ge 0.$$

The system (1.1) is endowed with initial data with prescribed far-field states in the x_1 -direction:

(1.2)
$$(\rho, u)(t, x)|_{t=0} = (\rho_0(x), u_0(x)) \to (\rho_{\pm}, u_{\pm}), \text{ as } x_1 \to \pm \infty$$

where $\rho_{\pm} > 0$, $u_{\pm} = (u_{1\pm}, 0, 0)^t$ are given planar constant states, and the periodic boundary conditions are imposed on $(x_2, x_3) \in \mathbb{T}^2$ for the solution (ρ, u) .

When the density ρ is away from vacuum ($\rho > 0$), it is convenient to introduce the specific volume $v := 1/\rho$, and then (1.1) can be transformed into the equivalent system with respect to (v, u):

(1.3)
$$\begin{cases} \rho(\partial_t v + u \cdot \nabla v) = \operatorname{div} u, \\ \rho(\partial_t u + u \cdot \nabla u) + \nabla p = (2\mu + \lambda) \nabla \operatorname{div} u - \mu \nabla \times \nabla \times u, \end{cases}$$

where we invoke the identity

$$\nabla \times \nabla \times u = \nabla \operatorname{div} u - \Delta u \,.$$

The large-time behavior of solutions to (1.1)-(1.2) is expected to be determined by the planar Riemann problem of the corresponding 3D compressible Euler system

(1.4)
$$\begin{cases} \partial_t \rho + \operatorname{div}(\rho u) = 0, \\ \partial_t(\rho u) + \operatorname{div}(\rho u \otimes u) + \nabla p(\rho) = 0, \\ (\rho, u)(0, x) = \begin{cases} (\rho_-, u_-), & x_1 < 0, \\ (\rho_+, u_+), & x_1 > 0, \end{cases} \end{cases}$$

which contains in general two nonlinear planar waves, i.e. planar shock and planar rarefaction wave.

Such expectation has been well substantiated in one-dimensional case. In 1960, Il'in-Oleĭnik [12] first validate the stability of single viscous shock and rarefaction wave for scalar quasi-linear equation with convex flux. Later, under zero-mass conditions on initial perturbations, Goodman [5] and Matsumura-Nishihara [27] independently proved the stability of single viscous shock to uniformly viscous conservation

Acknowledgment. The work of Yi Wang is partially supported by the NSFC grants (Grant No.s 12171459, 12090014, 12288201) and CAS Project for Young Scientists in Basic Research, Grant No. YSBR-031.

laws and compressible Navier-Stokes system, respectively. Then, Liu [23] and Szepessy-Xin [35] removed the zero-mass conditions in [5] by introducing a constant shift to the shock profile together with extra diffusion waves in the transverse characteristic fields, and the physical viscosity case was tackled by Liu-Zeng [25]. For the stability of rarefaction wave, we refer to Matsumura-Nishihara [28, 29] for the isentropic Navier-Stokes system, and Liu-Xin [24], Nishihara-Yang-Zhao [32] for the non-isentropic case. As to the stability of composite waves, we refer to Huang-Matsumura [10] for two viscous shocks, and Huang-Li-Matsumura [9] for the combination of a viscous contact wave and a rarefaction wave. The case of the superposition of a viscous shock and a rarefaction wave is more challenging [26], for the incompatibility between the standard anti-derivative method, used to study the stability of viscous shocks, and the direct energy method for rarefactions. Recently, Kang-Vasseur-Wang [19] conquered such difficulty by the method of *a*-contraction with shifts and then [14] showed the stability of generic Riemann solutions consisting of a viscous shock, a rarefaction and a viscous contact wave for Navier-Stokes-Fourier system.

Comparatively speaking, the stability of elementary wave patterns for multi-dimensional viscous conservation laws remains less explored. For scalar viscous conservation laws, we refer to Goodman [6]. Hoff-Zumbrun [7, 8] and Kang-Vasseur-Wang [18] for the stability of planar viscous shocks, as well as Xin [37], Ito [13] and Nishikawa-Nishihara [33] for the stability of planar rarefaction waves. As to the multi-dimensional Navier-Stokes system, Humpherys-Lyng-Zumbrun [11] proved the spectral stability of a planar viscous shock under some spectral conditions in \mathbb{R}^3 , while Wang-Wang [36] and Kang-Lee [15] proved the nonlinear stability for single and two planar viscous shocks under generic perturbations in $\mathbb{R} \times \mathbb{T}^2$ respectively. In addition, Li-Wang-Wang [22] proved the stability of planar rarefaction waves for the 3D Navier-Stokes-Fourier system in $\mathbb{R} \times \mathbb{T}^2$. Very recently, Meng [30] proved the stability of composite waves for the general $n \times n$ multi-dimensional conservation laws system in $\mathbb{R} \times \mathbb{T}^{d-1}$ with uniformly positive viscosities, in particular, including compressible barotropic Brenner-Navier-Stokes equations. However, this general result can not be directly applicable to the compressible barotropic Navier-Stokes system (1.1) with physical viscosities. Meanwhile, it should be remarked that the nonlinear stability of a planar viscous shock and its composite waves for the multi-dimensional compressible Navier-Stokes-Fourier system remains largely open even with the aid of a-contraction method, due to the temperature effects in the energy equation and their couplings.

The purpose of this paper is to prove the nonlinear time-asymptotic stability of the composite wave consisting of a planar rarefaction wave and a planar viscous shock for 3D barotropic Navier-Stokes system (1.1) under generic perturbations without zero-mass conditions, which can be viewed as a 3D version of [19]. Compared with the tractable Lagrangian structure in one-dimensional case, 3D Navier-Stokes system is expressed in Eulerian coordinates, whose diffusive terms are more difficult to be fully utilized. Motivated by [36], effective velocity is introduced to make full use of the dissipative effect. Also, some underlying physical structures are used to overcome the difficulties arising from the wave propagation along transverse directions and the interactions between the planar rarefaction wave and the planar viscous shock.

The rest of the paper is organized as follows. In Section 2, we first outline the properties of planar rarefaction and viscous shock, then we state our main result. In Section 3, we begin by introducing some useful inequalities, followed by the construction of the weight function and the shift function. The proof of our main result is then provided, based on the existence of local solution in Proposition 3.1 and uniform-in-time a priori estimates in Proposition 3.2. In Section 4, we first reformulate the problem in terms of (v, h), and then validate the uniform-in-time a priori estimates as stated in Proposition 3.2.

2. Preliminaries and main result

In this section, we first exhibit the construction and properties of planar rarefaction and viscous shock. Then we state our main result on the nonlinear time-asymptotic stability of composite waves to 3D compressible Navier-Stokes equations (1.1) under generic H^2 -perturbations without zero-mass conditions.

2.1. Planar rarefaction wave. Throughout this paper, we consider the planar wave that depends only on x_1 and has null component in the (x_2, x_3) direction. Thus, in order to construct planar rarefaction wave, we only need to consider the 1D Euler system

(2.1)
$$\begin{cases} \rho_t + (\rho u_1)_{x_1} = 0, \\ (\rho u_1)_t + (\rho u_1^2 + p(\rho))_{x_1} = 0, \end{cases}$$

$$\begin{pmatrix} \rho \\ u_1 \end{pmatrix}_t + \begin{pmatrix} u_1 & \rho \\ p'(\rho)/\rho & u_1 \end{pmatrix} \begin{pmatrix} \rho \\ u_1 \end{pmatrix}_{x_1} = 0.$$

The Jacobi matrix

$$\left(\begin{array}{cc} u_1 & \rho \\ p'(\rho)/\rho & u_1 \end{array}\right)$$

has two distinct eigenvalues

$$\lambda_1(\rho, u_1) = u_1 - \sqrt{p'(\rho)}, \qquad \lambda_2(\rho, u_1) = u_1 + \sqrt{p'(\rho)}$$

with corresponding right eigenvectors

$$r_i(\rho, u_1) = \left(1, \, (-1)^i \frac{\sqrt{p'(\rho)}}{\rho}\right)^t, \qquad i = 1, 2,$$

such that

$$r_i(\rho, u) \cdot \nabla_{(\rho, u)} \lambda_i(\rho, u) = (-1)^i \frac{\rho p''(\rho) + 2p'(\rho)}{2\rho \sqrt{p'(\rho)}} \neq 0, \ i = 1, 2, \text{ if } \rho > 0.$$

Define the *i*-Riemann invariant (i = 1, 2) as

$$\Sigma_i(\rho, u_1) := u_1 + (-1)^{i+1} \int^{\rho} \frac{\sqrt{p'(s)}}{s} ds$$

which satisfies

$$\nabla_{(\rho,u_1)} \Sigma_i(\rho, u_1) \cdot r_i(\rho, u) \equiv 0, \qquad \forall \, \rho > 0, u$$

For a given right state $(\rho_0, u_{10}) \in \mathbb{R}_+ \times \mathbb{R}$, the 1-rarefaction wave curve $R_1(\rho_0, u_{10})$ can be defined as the integral curve of the 1-characteristic field, that is

$$R_1(\rho_0, u_{10}): \frac{du_1}{d\rho} = -\frac{\sqrt{p'(\rho)}}{\rho}$$

When $p(\rho) = \rho^{\gamma}, \gamma > 1$, it has the following explicit expression:

$$R_1(\rho, u_1) = \left\{ (\rho, u_1) : u_1 - u_{10} = \frac{2\sqrt{\gamma}}{\gamma - 1} \left(\rho^{-\frac{\gamma - 1}{2}} - \rho_0^{-\frac{\gamma - 1}{2}} \right) \right\}.$$

Now, we utilize the *i*-Riemann invariants to construct rarefaction waves for (2.1). For the sake of brevity, we only consider 1-rarefaction wave and 2-rarefaction wave can be obtained similarly. To do this, we first study the Riemann problem for the inviscid Burgers equation

(2.2)
$$\begin{cases} w_t + ww_{x_1} = 0, \quad t > 0, \ x_1 \in \mathbb{R}, \\ w(0, x_1) = w_0^r(x_1) = \begin{cases} w_{-}, \ x_1 < 0, \\ w_{m}, \ x_1 > 0. \end{cases} \end{cases}$$

If $w_{-} < w_{m}$, the Riemann problem (2.2) admits a rarefaction wave solution $w^{r}(t, x_{1}) = w^{r}(x_{1}/t)$ which reads as

(2.3)
$$w^{r}\left(\frac{x_{1}}{t}\right) = \begin{cases} w_{-}, & x_{1} \leq w_{-}t, \\ \frac{x_{1}}{t}, & w_{-}t \leq x_{1} \leq w_{m}t, \\ w_{m}, & x_{1} \geq w_{m}t. \end{cases}$$

For the given states $(\rho_m, u_{1m}) \in \mathbb{R}_+ \times \mathbb{R}$ and $(\rho_-, u_{1-}) \in R_1(\rho_m, u_{1m})$, let

$$w_{-} = \lambda_1(\rho_{-}, u_{1-}), \quad w_m = \lambda_1(\rho_m, u_{1m}),$$

and $w^r(t, x_1) := w^r(x_1/t)$ be the solution defined by (2.3).

Then we can construct the 1-rarefaction wave $(\rho^r, u_1^r)(x_1/t)$ as follows

(2.4)
$$\begin{cases} \lambda_1(\rho^r, u_1^r)(x_1/t) = w^r(x_1/t), \\ \Sigma_1(\rho^r, u_1^r)(x_1/t) = \Sigma_1(\rho_-, u_{1-}) = \Sigma_1(\rho_m, u_{1m}), \end{cases}$$

which is the self-similar solution of the 1D Euler system (2.1) with initial data

$$(\rho, u_1)|_{t=0} = \begin{cases} (\rho_-, u_{1-}), & x_1 < 0, \\ (\rho_m, u_{1m}), & x_1 > 0. \end{cases}$$

2.2. Approximate rarefaction wave. Motivated by [28], we construct the smooth approximate 1rarefaction wave with the aid of the smooth solution to the Burgers equation

(2.5)
$$\begin{cases} w_t + ww_{x_1} = 0, \quad t > 0, \ x_1 \in \mathbb{R}, \\ w(0, x_1) = w_0^R(x_1) = \frac{w_m + w_-}{2} + \frac{w_m - w_-}{2} \tanh x_1. \end{cases}$$

By the method of characteristics, the solution $w^{R}(t, x_{1})$ to (2.5) can be defined as

(2.6)
$$w^{R}(t,x_{1}) = w_{0}^{R}(x_{0}(t,x_{1})), \qquad x_{1} = x_{0}(t,x_{1}) + w_{0}^{R}(x_{0}(t,x_{1}))t.$$

Also, $w^{R}(t, x_{1})$ has the following properties whose proofs can be found in [28] or [29].

Lemma 2.1. Let $w_m > w_-$ and $\tilde{w} := w_m - w_-$, then the Cauchy problem (2.5) admits a unique global smooth solution $w^{R}(t, x_{1})$ satisfying

- (1) $w_{-} < w^{R}(t, x_{1}) < w_{m}$ and $w^{R}_{x_{1}} > 0$, for $x_{1} \in \mathbb{R}$ and $t \geq 0$. (2) For any t > 0 and $p \in [1, +\infty]$, there exists a positive constant $C = C_{p}$ such that

$$\begin{split} \|w_{x_1}^R(t,\cdot)\|_{L^p} &\leq C \min\left\{ |\tilde{w}|, |\tilde{w}|^{1/p} t^{-1+1/p} \right\}, \\ \|w_{x_1x_1}^R(t,\cdot)\|_{L^p} &\leq C \min\left\{ |\tilde{w}|, t^{-1} \right\}, \\ \|w_{x_1x_1x_1}^R(t,\cdot)\|_{L^p} &\leq C \min\left\{ |\tilde{w}|, t^{-1} \right\}. \end{split}$$

(3) For $x_1 \ge w_m t$, $t \ge 0$, it holds that

$$\begin{aligned} |w^{R}(t,x_{1}) - w_{m}| &\leq \tilde{w}e^{-2|x_{1} - w_{m}t|}, \\ |(w^{R}_{x_{1}},w^{R}_{x_{1}x_{1}})(t,x_{1})| &\leq C\tilde{w}e^{-2|x_{1} - w_{m}t|}. \end{aligned}$$

(4) For $x_1 \leq w_t$, $t \geq 0$, it holds that

$$|w^{R}(t, x_{1}) - w_{-}| \leq \tilde{w}e^{-2|x_{1} - w_{-}t|},$$

$$|(w^{R}_{x_{1}}, w^{R}_{x_{1}x_{1}})(t, x_{1})| \leq C\tilde{w}e^{-2|x_{1} - w_{-}t|}.$$

(5) $\lim_{t \to +\infty} \sup_{x, \in \mathbb{R}} \left| w^R(t, x_1) - w^r(\frac{x_1}{t}) \right| = 0.$

Correspondingly, the approximate 1-rarefaction wave $(\rho^R, u_1^R)(t, x_1)$ of the 1-rarefaction wave $(\rho^r, u_1^r)(x_1/t)$ can be defined by

(2.7)
$$\begin{cases} w_{-} = \lambda_{1}(\rho_{-}, u_{1-}), \ w_{m} = \lambda_{1}(\rho_{m}, u_{1m}), \\ \lambda_{1}(\rho^{R}, u_{1}^{R})(t, x_{1}) = w^{R}(1+t, x_{1}), \\ \Sigma_{1}(\rho^{R}, u_{1}^{R})(t, x_{1}) = \Sigma_{1}(\rho_{-}, u_{1-}) = \Sigma_{1}(\rho_{m}, u_{1m}) \end{cases}$$

where $w^{R}(t, x_{1})$ is the solution of the Burgers equation (2.5). By direct computations, it can be verified that (ρ^R, u_1^R) satisfies the 1D Euler system

$$\begin{cases} \rho_t + (\rho u_1)_{x_1} = 0, \\ (\rho u_1)_t + (\rho u_1^2 + p(\rho))_{x_1} = 0. \end{cases}$$

Also, the following lemma can be derived from Lemma 2.1 (cf. [28]).

Lemma 2.2. For any $(\rho_m, u_{1m}) \in \mathbb{R}_+ \times \mathbb{R}$ and $(\rho_-, u_{1-}) \in R_1(\rho_m, u_{1m})$, let $\lambda_{1-} := \lambda_1(\rho_-, u_{1-})$, $\lambda_{1m} := \lambda_1(\rho_m, u_{1m})$ and $\delta_R := |v_m - v_-| \sim |u_{1m} - u_-|$ denoting the rarefaction wave strength. Then the approximate 1-rarefaction wave $(\rho^R, u_1^R)(t, x)$ defined by (2.7) satisfies the following properties, where $v^R := 1/\rho^R,$

- (1) For any $x_1 \in \mathbb{R}$ and any $t \ge 0$, it holds $u_{1,x_1}^R > 0$, $v_{x_1}^R > 0$ and $u_{1,x_1}^R \sim v_{x_1}^R$.
- (2) For any $t \ge 0$ and any $p \in [1, +\infty]$, it holds

$$\begin{aligned} \|(v_{x_{1}}^{R}, u_{1,x_{1}}^{R})\|_{L^{p}} &\leq C \min\{\delta_{R}, \delta_{R}^{1/p}(1+t)^{-1+1/p}\},\\ \|(v_{x_{1}x_{1}}^{R}, u_{1,x_{1}x_{1}}^{R})\|_{L^{p}} &\leq C \min\{\delta_{R}, (1+t)^{-1}\},\\ \|u_{1,x_{1}x_{1}}^{R}\| &\leq C \|u_{1,x_{1}}^{R}\|, \|u_{1,x_{1}x_{1}x_{1}}^{R}\| &\leq C \|u_{1,x_{1}x_{1}}^{R}\|, \forall x \in \mathbb{R}. \end{aligned}$$

$$(3) \ For \ x_{1} &\geq \lambda_{1m}(1+t), \ t \geq 0, \ it \ holds \\ \|(v^{R}, u_{1}^{R})(t, x_{1}) - (v_{m}, u_{m})\| &\leq C\delta_{R} \ e^{-2|x_{1}-\lambda_{1m}(1+t)|},\\ \|(v_{x_{1}}^{R}, u_{1,x_{1}}^{R})(t, x_{1})\| &\leq C\delta_{R} \ e^{-2|x_{1}-\lambda_{1m}(1+t)|}. \end{aligned}$$

(4) For
$$x_1 \leq \lambda_{1-}(1+t), t \geq 0$$
, it holds

$$|(v^R, u_1^R)(t, x_1) - (\rho_-, u_-)| \leq C\delta_R e^{-2|x_1 - \lambda_{1-}(1+t)|}$$

$$|(v_{x_1}^R, u_{1,x_1}^R)(t, x_1)| \leq C\delta_R e^{-2|x_1 - \lambda_{1-}(1+t)|}.$$
(5) $\lim_{t \to +\infty} \sup_{x \in \mathbb{R}} \left| (v^R, u_1^R)(t, x_1) - (v^r, u_1^r) \left(\frac{x_1}{t}\right) \right| = 0.$

Remark 2.1. We wish to emphasize the distinction between the superscript \cdot^{r} and the superscript \cdot^{R} . Throughout this paper, the former denotes the standard rarefaction waves, while the latter signifies the approximated rarefaction waves.

2.3. Planar viscous shock. Like before, we consider the 1D barotropic Navier-Stokes system

(2.8)
$$\begin{cases} \partial_t \rho + \partial_1(\rho u_1) = 0, \\ \partial_t(\rho u_1) + \partial_1(\rho u_1^2 + p(\rho)) = (2\mu + \lambda)\partial_{11}u_1, \end{cases}$$

with initial data

(2.9)
$$(\rho, u_1)(0, x_1) = (\rho_0, u_{10})(x_1) = \begin{cases} (\rho_m, u_{1m}), & x_1 < 0, \\ (\rho_+, u_{1+}), & x_1 > 0. \end{cases}$$

We hope to seek for travelling wave solutions of the form $(\rho, u_1)(t, x_1) = (\rho^s, u_1^s)(\xi)$, where $\xi = x_1 - \sigma t$ and σ is a constant to be determined. Inserting the Ansatz $(\rho, u_1)(t, x_1) = (\rho^s, u_1^s)(\xi)$ into (2.8) leads to

(2.10)
$$\begin{cases} -\sigma(\rho^s)' + (\rho^s u_1^s)' = 0, & ' := \frac{d}{d\xi}, \\ -\sigma(\rho^s u_1^s)' + (\rho^s(u_1^s)^2)' + p(\rho^s)' = (2\mu + \lambda)(u_1^s)'', \end{cases}$$

which is an ODE for $(\rho^s, u_1^s)(\xi)$. The initial data (2.9) are then converted to

(2.11)
$$(\rho^s, u_1^s)(-\infty) = (\rho_m, u_{1m}), \quad (\rho^s, u_1^s)(+\infty) = (\rho_+, u_{1+}).$$

Integrating (2.10) with respect to ξ over \mathbb{R} results in the following Rankine-Hugoniot condition

(2.12)
$$\begin{cases} -\sigma(\rho_{+} - \rho_{m}) + (\rho_{+}u_{1+} - \rho_{m}u_{1m}) = 0, \\ -\sigma(\rho_{+}u_{1+} - \rho_{m}u_{1m}) + (\rho_{+}u_{1+}^{2} - \rho_{m}u_{1m}^{2}) + (p(\rho_{+}) - p(\rho_{m})) = 0, \end{cases}$$

which is necessary for (2.10)-(2.11) to be solvable. Furthermore, we say $(\rho_m, u_{1m}) \in S_2(\rho_+, u_{1+})$ if it also satisfies the Lax entropy conditions

(2.13)
$$\begin{cases} \sigma > \lambda_1(\rho_m, u_{1m}), \\ \lambda_2(\rho_+, u_{1+}) < \sigma < \lambda_2(\rho_m, u_{1m}), \end{cases}$$

where $\lambda_1(\rho, u_1) = u_1 - \sqrt{p'(\rho)}, \ \lambda_2(\rho, u_1) = u_1 + \sqrt{p'(\rho)}.$

When away from vacuum, let $v^s := 1/\rho^s$, then the ODE (2.10) can be converted to an ODE for $(v^s, u_1^s)(\xi)$:

(2.14)
$$\begin{cases} \rho_s \big(-\sigma(v^s)' + u_1^s(v^s)' \big) = (u_1^s)', \\ \rho_s \big(-\sigma(u_1^s)' + u_1^s(u_1^s)' \big) + p(v^s)' = (2\mu + \lambda)(u_1^s)'', \end{cases}$$

where $p(v^s) = (v^s)^{-\gamma}$. Integrating $(2.10)_1$ over $(-\infty, \xi)$ leads to

(2.15)
$$-\sigma\rho^{s} + \rho^{s}u_{1}^{s} = -\sigma\rho_{-} + \rho_{-}u_{1-} =: \sigma_{*}.$$

By virtue of the σ_* defined above, the system (2.14) and the far-field conditions (2.11) can be rewritten as

(2.16)
$$\begin{cases} -\sigma_*(v^s)' = (u_1^s)', \\ -\sigma_*(u_1^s)' + p(v^s)' = (2\mu + \lambda)(u_1^s)'', \end{cases}$$

and

(2.17)
$$(v^s, u_1^s)(-\infty) = (v_m, u_{1m}), \quad (v^s, u_1^s)(+\infty) = (v_+, u_{1+}).$$

where $v_m = 1/\rho_m$, $v_+ = 1/\rho_+$. In addition, integrating (2.16) over \mathbb{R} and combining (2.15), we have

$$\begin{cases} -\sigma_*(v_+ - v_m) = u_{1+} - u_{1m}, \\ -\sigma_*(u_{1+} - u_{1m}) + p(v_+) - p(v_m) = 0, \end{cases}$$

then we can obtain another useful expression for σ_* :

(2.18)
$$\sigma_* = \sqrt{-\frac{p(v_+) - p(v_m)}{v_+ - v_m}} > 0.$$

Then, we summarize the properties of the 2-viscous shock in the following lemma, whose proof can be found in [27].

Lemma 2.3. For any given state (v_+, u_{1+}) and $(v_m, u_{1m}) \in S_2(v_+, u_{1+})$, let

$$\delta_S := |p(v_m) - p(v_+)| \sim |v_m - v_+| \sim |u_{1m} - u_{1+}|,$$

which denotes the shock wave strength. Then, up to a constant shift, the ODE (2.16)-(2.17) admits a unique solution $(v^s, u_1^s)(\xi)$ satisfying

$$(v^s)' > 0, \quad (u_1^s)' = -\sigma_*(v^s)' < 0.$$

In addition, there exists a constant C > 0 such that the following holds

$$\begin{aligned} |v^{s}(\xi) - v_{m}| &\leq C\delta_{S}e^{-C\delta_{S}|\xi|}, \quad \forall \xi < 0, \\ |v^{s}(\xi) - v_{+}| &\leq C\delta_{S}e^{-C\delta_{S}|\xi|}, \quad \forall \xi > 0, \\ |((v^{s})', (u_{1}^{s})')| &\leq C\delta_{S}^{2}e^{-C\delta_{S}|\xi|}, \quad \forall \xi \in \mathbb{R}, \\ |((v^{s})'', (u_{1}^{s})'')| &\leq C\delta_{S} |((v^{s})', (u_{1}^{s})')|, \quad \forall \xi \in \mathbb{R}, \\ |((v^{s})''', (u_{1}^{s})''')| &\leq C\delta_{S}^{2} |((v^{s})', (u_{1}^{s})')|, \quad \forall \xi \in \mathbb{R}. \end{aligned}$$

2.4. Notations. For the convenience of the readers, we list some notational conventions that taken throughout this paper.

(1) Let Ω denote the domain $\mathbb{R} \times \mathbb{T}^2$. For $x = (x_1, x_2, x_3) \in \Omega$, we define

$$x' := (x_2, x_3), \quad dx' := dx_2 dx_3.$$

(2) Let $1 \leq r \leq +\infty$, we denote $L^r := L^r(\Omega)$ and

$$||f|| := ||f||_{L^2} = \left(\int |f|^2 \, dx\right)^{\frac{1}{2}}$$

For $m \in \mathbb{N}$, $H^m := H^m(\Omega)$ denotes the usual Sobolev space based on L^2 with the norm

$$\|f\|_{H^m} := \left(\sum_{k=0}^m \|\nabla^k f\|^2\right)^{\frac{1}{2}}, \quad \|(f,g)\|_{H^m} := \|f\|_{H^m} + \|g\|_{H^m}$$

(3) Let (v^s, u_1^s) be the 2-viscous shock defined in Lemma 2.3 and $\mathbf{X}(t)$ be any time-dependent shift, we define the shifted shock with uppercase superscript to distinguish

(2.19)
$$(v^S, u_1^S)(t, x) := (v^s, u_1^s) (x_1 - \sigma t - \mathbf{X}(t))$$
where σ is the constant defined by (2.12).

(4) (v^r, u_1^r) and (v^R, u_1^R) and given by (2.4) and (2.7), respectively. Let $u^R := (u_1^R, 0, 0)^t$, $u^S := (u_1^S, 0, 0)^t$ and

(2.20)
$$\begin{cases} \bar{v}(t,x) := (v^S + v^R - v_m)(t,x), \\ \bar{u}_1(t,x) := (u_1^S + u_1^R - u_{1m})(t,x), \\ \bar{u}(t,x) := (\bar{u}_1(t,x), 0, 0)^t, \end{cases}$$

all of which depend only on (t, x_1) . In addition, we define the error terms as follows

(2.21)
$$\begin{cases} \phi := v - \bar{v}, \\ \psi := u - \bar{u} = (u_1 - \bar{u}_1, 0, 0)^t = (\psi_1, 0, 0)^t \end{cases}$$

(5) For the pressure function $p(v) = v^{-\gamma}$, we often take the following abbreviations

$$p^S:=p(v^S), \quad p^R:=p(v^R), \quad \bar{p}:=p(\bar{v})$$

(6) δ_R , δ_S and δ denote the strength of the 1-rarefaction, 2-viscous shock and the composite wave, respectively, which are defined as

(2.22)
$$\begin{cases} \delta_R := |v_m - v_-| \sim |u_{1m} - u_-|, \\ \delta_S := |p(v_m) - p(v_+)| \sim |v_m - v_+| \sim |u_{1m} - u_{1+}| \\ \delta := \delta_R + \delta_S. \end{cases}$$

2.5. Main result. Now we can state our main result as follows. Roughly speaking, as long as the strength of the composite wave is suitably small, then it is nonlinearly stable under generic H^2 -perturbations without zero-mass conditions.

Theorem 2.1. For any given state $(v_+, u_{1+}) \in \mathbb{R}_+ \times \mathbb{R}$ there exist constants $\delta_0, \varepsilon_0 > 0$ such that for each $(v_m, u_{1m}) \in S_2(v_+, u_{1+}), (v_-, u_{1-}) \in R_1(v_m, u_{1m})$ with

$$\delta := |v_{+} - v_{m}| + |v_{m} - v_{-}| \le \delta_{0}$$

and each initial data with

(2.23)
$$\| (v_0 - \bar{v}(0, \cdot), u_0 - \bar{u}(0, \cdot)) \|_{H^2(\mathbb{R} \times \mathbb{T}^2)} \le \varepsilon_0,$$

the 3D barotropic Navier-Stokes system (1.3) admits a unique global-in-time solution (v, u)(t, x) with $v := \frac{1}{a}$. In addition, there exists an absolutely continuous shift $\mathbf{X}(t)$ such that

(2.24)
$$v(t,x) - \bar{v}(t,x) \in C([0,+\infty); H^{2}(\mathbb{R} \times \mathbb{T}^{2})),$$
$$u(t,x) - \bar{u}(t,x) \in C([0,+\infty); H^{2}(\mathbb{R} \times \mathbb{T}^{2})),$$
$$\nabla(v(t,x) - \bar{v}(t,x)) \in L^{2}(0,+\infty; H^{1}(\mathbb{R} \times \mathbb{T}^{2})),$$
$$\nabla(u(t,x) - \bar{u}(t,x)) \in L^{2}(0,+\infty; H^{2}(\mathbb{R} \times \mathbb{T}^{2})).$$

Here,

(2.25)
$$\bar{v}(t,x) := v^R(t,x_1) + v^s(x_1 - \sigma t - \mathbf{X}(t)) - v_m \cdot \\ \bar{u}(t,x) := u^R(t,x_1) + u^s(x_1 - \sigma t - \mathbf{X}(t)) - u_m$$

where $(v^R, u^R)(t, x_1)$ and $(v^s, u^s)(x_1 - \sigma t)$ represent the planar approximate 1-rarefaction and the planar 2-viscous shock, i.e.

$$(v^{R}, u^{R})(t, x_{1}) := (v^{R}, (u_{1}^{R}, 0, 0)^{t})(t, x_{1}),$$

$$(v^{s}, u^{s})(x_{1} - \sigma t) := (v^{s}, (u_{1}^{s}, 0, 0)^{t})(x_{1} - \sigma t),$$

$$u_{m} := (u_{1m}, 0, 0)^{t}.$$

Furthermore, we have the following time-asymptotically stability. As $t \to +\infty$,

(2.26)
$$\sup_{x \in \mathbb{R}} \left| (v, u)(t, x) - \left(v^r(\frac{x_1}{t}) + v^s(x_1 - \sigma t - \mathbf{X}(t)) - v_m, \right. \\ \left. u^r(\frac{x_1}{t}) + u^s(x_1 - \sigma t - \mathbf{X}(t)) - u_m \right) \right| \to 0,$$

and

(2.27)
$$\lim_{t \to +\infty} \left| \dot{\mathbf{X}}(t) \right| = 0$$

where

$$(v^r, u^r)\left(\frac{x_1}{t}\right) := \left(v^r, (u_1^r, 0, 0)^t\right)\left(\frac{x_1}{t}\right)$$

and $(v^r, u_1^r)\left(\frac{x_1}{t}\right)$ is the 1-rarefaction defined by (2.4) with end states (v_-, u_{1-}) and (v_m, u_{1m}) .

Remark 2.2. Theorem 2.1 states that if the two far-field states (ρ_{\pm}, u_{\pm}) in (1.2) are connected by the superposition of a rarefaction wave and a shock, then the solution to the three-dimensional compressible Navier-Stokes system (1.1) tends to the superposition wave of the planar inviscid rarefaction wave and the planar viscous shock with shift $\mathbf{X}(t)$ in long time.

Remark 2.3. Compared with the 1D result in [19], the 3D Navier-Stokes system is expressed in Eulerian coordinates, whose diffusive terms are more difficult to be fully utilized. Therefore, motivated by [36], we need to introduce an effective velocity $h := u - (2\mu + \lambda)\nabla v$ to make full use of the dissipative effect. In addition, some underlying physical structures are used to overcome the difficulties arising from the wave propagation along transverse directions and the interactions between the planar rarefaction wave and the planar viscous shock.

Remark 2.4. The shift function is proved to satisfy the time-asymptotic behavior (2.27), which implies that

$$\lim_{t \to +\infty} \frac{\mathbf{X}(t)}{t} = 0.$$

Hence, the shift function grows at most sub-linearly with respect to the time t, which time-asymptotically preserves the traveling wave profile of the shifted viscous shock.

3. Proof of the main result

The main strategy of our proof is the use of the method of *a*-contraction with shifts [16, 20], which is based on the relative entropy introduced by DiPerna [4] and Dafermos [3]. To achieve this, we first give some technical lemmas and then construct the weight function and the time-dependent shift. Finally, we present the proof of Theorem 2.1 based on a local existence result and uniform-in-time a priori estimates.

3.1. Some inequalities. Firstly, we introduce a 3D weighted sharp Poincaré type inequality that given in [36], which can be viewed as a 3D version of the 1D case in [17].

Lemma 3.1. For every $f:[0,1] \times \mathbb{T}^2 \to \mathbb{R}$ satisfying

$$\int_{\mathbb{T}^2} \int_0^1 y_1(1-y_1) |\partial_{y_1} f|^2 + \frac{|\nabla_{y'} f|^2}{y_1(1-y_1)} \, dy_1 dy' < \infty,$$

it holds

(3.1)

$$\begin{aligned} \int_{\mathbb{T}^2} \int_0^1 |f - \bar{f}|^2 \, dy_1 dy' \\
&\leq \frac{1}{2} \int_{\mathbb{T}^2} \int_0^1 y_1 (1 - y_1) |\partial_{y_1} f|^2 dy_1 dy' + \frac{1}{16\pi^2} \int_{\mathbb{T}^2} \int_0^1 \frac{|\nabla_{y'} f|^2}{y_1 (1 - y_1)} dy_1 dy', \\
&\text{where } \bar{f} := \int_{\mathbb{T}^2} \int_0^1 f dy_1 dy', \, y' := (y_2, y_3).
\end{aligned}$$

The following 3D Gagliardo-Nirenberg type inequality in the domain $\Omega = \mathbb{R} \times \mathbb{T}^2$ is also useful, whose proof can be found in [21].

Lemma 3.2. Let $\Omega := \mathbb{R} \times \mathbb{T}^2$. There exists constant C > 0, such that for each $g \in H^2(\Omega)$, it holds

(3.2)
$$\|g\|_{L^{\infty}(\Omega)} \leq \sqrt{2} \|g\|_{L^{2}(\Omega)}^{\frac{1}{2}} \|\partial_{1}g\|_{L^{2}(\Omega)}^{\frac{1}{2}} + C \|\nabla g\|_{L^{2}(\Omega)}^{\frac{1}{2}} \|\nabla^{2}g\|_{L^{2}(\Omega)}^{\frac{1}{2}}$$

3.2. Relative quantities. Then we present some estimates concerning the relative quantities. For any function F defined on some interval I of \mathbb{R} , we define the associated relative quantity with $v, u \in I$,

$$F(v|w) := F(v) - F(w) - F'(w)(v - w).$$

We will state some useful inequalities about the relative quantities associated to the pressure $p(v) = v^{-\gamma}$ and the internal energy $Q(v) := \frac{v^{1-\gamma}}{\gamma-1}$. The proofs, which are based on Taylor expansions, can be accessed in [17].

Lemma 3.3. Let $p(v) = v^{-\gamma}$ and $Q(v) := \frac{v^{1-\gamma}}{\gamma-1}$ with $\gamma > 1$. For given constants $v_{-} > 0$, there exist constants $C, \delta_* > 0$ such that the following holds true.

(1) For any $v, w \in \mathbb{R}_+$ such that $0 < w < 2v_-$ and $0 < v \le 3v_-$, it holds

(3.3)
$$|v - w|^2 \le CQ(v|w), \quad |v - w|^2 \le Cp(v|w).$$

(2) For any
$$v, w > \frac{v_-}{2}$$
, it holds

(3.4)
$$|p(v) - p(w)| \le C|v - w|.$$

(3) For any $0 < \delta < \delta_*$ and $(v, w) \in \mathbb{R}^2_+$ satisfying $|p(v) - p(w)| < \delta$ and $|p(w) - p(v_-)| < \delta$, the following holds true:

(3.5)
$$p(v|w) \le \left(\frac{\gamma+1}{2\gamma} \frac{1}{p(w)} + C\delta\right) |p(v) - p(w)|^2,$$

(3.6)
$$Q(v|w) \ge \frac{|p(v) - p(w)|^2}{2\gamma p^{1+\frac{1}{\gamma}}(w)} - \frac{1+\gamma}{3\gamma^2} \frac{(p(v) - p(w))^3}{p^{2+\frac{1}{\gamma}}(w)}$$

(3.7)
$$Q(v|w) \le \left(\frac{1}{2\gamma p^{1+\frac{1}{\gamma}}(w)} + C\delta\right)|p(v) - p(w)|^2.$$

At last, we list an estimate about the inverse function of $p(v) = v^{-\gamma}$, whose proof can be found in [17].

Lemma 3.4. For given constant $v_- > 0$, there exist $\delta_0 > 0, C > 0$ such that for any $v_+ > 0$ with $0 < \delta := p(v_-) - p(v_+) \le \delta_0$ and $v_- \le v \le v_+$, it holds

(3.8)
$$\left| \frac{v - v_{-}}{p(v) - p(v_{-})} + \frac{v - v_{+}}{p(v_{+}) - p(v)} + \frac{1}{2} \frac{p''(v_{-})}{p'(v_{-})^2} (v_{-} - v_{+}) \right| \le C\delta^2.$$

3.3. Construction of the weight function and the time-dependent shift. We define the weight function a(t, x) by

 $\sqrt{\delta_{S}}$

(3.9)
$$a(t,x) := 1 + \frac{\nu}{\delta_S} (p(v_m) - p(v^S)) \\ = 1 + \frac{\nu}{\delta_S} (p(v_-) - p(v^s(x_1 - \sigma t - \mathbf{X}(t)))),$$

where ν is a constant to be determined with

$$\delta \ll \nu \lesssim \sqrt{\delta_S}.$$

For the sake of clarity, we just choose

(3.11)
$$\nu =$$

Notice that a(t, x) depends only on (t, x_1) and satisfies

(3.12)
$$1 < a(t, x) < 1 + \nu$$
.

Also, we have

(3.13)
$$\partial_1 a = -\frac{\nu}{\delta_S} p'(v^S) \partial_1 v^S > 0$$

and then

(3.14)
$$\partial_1 a \sim \frac{\nu}{\delta_S} \partial_1 v^S = \frac{1}{\nu} \partial_1 v^S.$$

With the above weight function a(t, x) given, we can define the shift $\mathbf{X}(t)$ as the solution of the following Cauchy problem:

(3.15)
$$\begin{cases} \dot{\mathbf{X}}(t) = -\frac{M}{\delta_S} \left[\int_{\Omega} \frac{a\rho}{\sigma_*} \cdot \partial_1 h_1^S \cdot \left(p(v) - p(\bar{v}) \right) - a\rho \cdot \partial_1 [p(v^S)] \cdot (v - \bar{v}) \, dx \right], \\ \mathbf{X}(0) = 0, \end{cases}$$

where $h_1^S := u_1^S - (2\mu + \lambda)\partial_1 v^S$, and

(3.16)
$$M := \frac{5(\gamma+1)}{8\gamma} \cdot \frac{\sigma_m^3 v_m^2}{p(v_m)}, \quad \sigma_m := \sqrt{-p'(v_m)}.$$

The shift $\mathbf{X}(t)$ is well-defined for the well-posedness of the ODE (3.15), which is guaranteed by the following lemma (cf. [2]).

Lemma 3.5. Let p > 1 and T > 0. Suppose that a function $F : [0, T] \times \mathbb{R} \to \mathbb{R}$ satisfies

$$\sup_{x \in \mathbb{R}} |F(t,x)| \le f(t) \quad and \quad \sup_{x,y \in \mathbb{R}, x \ne y} \left| \frac{F(t,x) - F(t,y)}{x - y} \right| \le g(t) \text{ for } t \in [0,T]$$

for some functions $f \in L^1(0,T)$ and $g \in L^p(0,T)$. Then for any $x_0 \in \mathbb{R}$, there exists a unique absolutely continuous function $\mathbf{X} : [0,T] \to \mathbb{R}$ satisfying

(3.17)
$$\begin{cases} \dot{\mathbf{X}}(t) = F(t, \mathbf{X}(t)) & \text{for a.e. } t \in [0, T] \\ X(0) = x_0 \,. \end{cases}$$

Let $F(t, \mathbf{X})$ be the right-hand side of the ODE (3.15). Due to the facts that

$$||a||_{C^1} \le 2, \quad ||v^S||_{C^2} \le \max\{1, v_+\}, \quad ||\partial_1 v^S||_{L^1} \le \delta_S,$$

we find that for some constant C > 0,

(3.18)
$$\sup_{\mathbf{X}\in\mathbb{R}}|F(t,\mathbf{X})| \leq \frac{C}{\delta_S} \left(\|v\|_{L^{\infty}} + \|v^S\|_{L^{\infty}} \right) \int_{\Omega} \partial_1 v^S \, dx \leq C,$$

and

$$\sup_{\mathbf{X}\in\mathbb{R}} \left|\partial_{\mathbf{X}}F(t,\mathbf{X})\right| \le \frac{C}{\delta_S} \left\|a\right\|_{C^1} \left(\left\|v\right\|_{L^{\infty}} + \left\|v^S\right\|_{L^{\infty}}\right) \int_{\Omega} \partial_1 v^S \, dx \le C$$

Thus by Lemma 3.5, the shift $\mathbf{X}(t)$ is well-defined and absolutely continuous. In addition, we have $|\dot{\mathbf{X}}(t)| \leq C$ from (3.18), and consequently

$$(3.19) |\mathbf{X}(t)| \le Ct, \forall t \in [0, T].$$

3.4. **Proof of Theorem 2.1.** Roughly speaking, we prove Theorem 2.1 by continuation arguments in virtue of a local existence result together with a priori estimates. For that, we first state the following local existence result, which can be proved by standard methods (see [31] or [34]).

Proposition 3.1 (Local existence). Let $(\tilde{v}, \tilde{u})(t, x)$ be the composite wave without shift, i.e.

$$\tilde{v}(t,x) := v^{R}(t,x_{1}) + v^{s}(x_{1} - \sigma t) - v_{m},\\ \tilde{u}(t,x) := u^{R}(t,x_{1}) + u^{s}(x_{1} - \sigma t) - u_{m}$$

For any $\Xi > 0$, there exists $T_0 > 0$ such that for each initial data (v_0, u_0) with

$$\left\| \left(v_0 - \tilde{v}(0, \cdot) , u_0 - \tilde{u}(0, \cdot) \right) \right\|_{H^2(\mathbb{R} \times \mathbb{T}^2)} \le \Xi ,$$

the 3D Navier-Stokes system (1.3) admits a unique solution (v, u) on $[0, T_0]$ satisfying

$$v - \tilde{v} \in C([0, T_0]; H^2(\mathbb{R} \times \mathbb{T}^2)), \quad \nabla(v - \tilde{v}) \in L^2(0, T_0; H^1(\mathbb{R} \times \mathbb{T}^2)), \\ u - \tilde{u} \in C([0, T_0]; H^2(\mathbb{R} \times \mathbb{T}^2)), \quad \nabla(u - \tilde{u}) \in L^2(0, T_0; H^2(\mathbb{R} \times \mathbb{T}^2)),$$

and for $t \in [0, T_0]$, it holds that

The pivotal part in the proof of Theorem 2.1 is the following a priori estimates.

Proposition 3.2 (A priori estimates). Let (v, u) be the solution of (1.3) on [0, T] for some T > 0, (\bar{v}, \bar{u}) be the composite wave given by (2.25). Then there exist positive constants $\delta_0 \leq 1$, $\chi_0 \leq 1$, C_0 independent of T, such that if the composite wave strength $\delta := \delta_R + \delta_S < \delta_0$ and

$$\begin{split} &v-\bar{v}\in C([0,T];H^2(\mathbb{R}\times\mathbb{T}^2)),\quad \nabla(v-\bar{v})\in L^2(0,T;H^1(\mathbb{R}\times\mathbb{T}^2)),\\ &u-\bar{u}\in C([0,T];H^2(\mathbb{R}\times\mathbb{T}^2)),\quad \nabla(u-\bar{u})\in L^2(0,T;H^2(\mathbb{R}\times\mathbb{T}^2)), \end{split}$$

with

(3.21)
$$\chi := \sup_{0 \le t \le T} \| (v - \bar{v}, u - \bar{u})(t, \cdot) \|_{H^2} \le \chi_0,$$

then it holds that

(3.22)

$$\sup_{0 \le t \le T} \|(v - \bar{v}, u - \bar{u})(t, \cdot)\|_{H^{2}}^{2} + \delta_{S} \int_{0}^{T} |\dot{\mathbf{X}}(t)|^{2} dt \\
+ \int_{0}^{T} (\partial_{1}v^{S} + \partial_{1}v^{R})|v - \bar{v}|^{2} + \|\nabla(v - \bar{v})\|_{H^{1}}^{2} + \|\nabla(u - \bar{u})\|_{H^{2}}^{2} dt \\
\le C_{0} \|v(0, \cdot) - \bar{v}(0, \cdot)\|_{H^{2}}^{2} + C_{0} \|u(0, \cdot) - \bar{u}(0, \cdot)\|_{H^{2}}^{2} + C_{0}\delta_{R}^{\frac{1}{3}}.$$

In addition, by (3.15), we have

$$(3.23) \qquad |\dot{\mathbf{X}}(t)| \le C_0 \left\| (v - \bar{v})(t, \cdot) \right\|_{L^{\infty}}, \quad \forall t \le T$$

Based on Propositions 3.1 and 3.2, we can employ continuation arguments to establish the global-intime existence of perturbations as stated in (2.24). Furthermore, Proposition 3.2 can be used to validate (2.26) for the large-time behavior. These proofs are typical and utilize the same arguments presented in the preceding paper [19, 36]. Consequently, we will omit these details and complete the proof of Theorem 2.1.

Thus, the proof of Theorem 2.1 is reduced to that of Proposition 3.2, which is the heart of the whole proof. The next section is then devoted to the proof of Proposition 3.2.

4. PROOF OF PROPOSITION 3.2

The proof of Proposition 3.2 will be given in the following subsections. Throughout this section, C denotes a positive generic constant independent of δ and ν (see (2.22) and (3.11) for their definitions). Also, we use $f \leq g$ to denote $f \leq Cg$ for some generic constant C > 0.

4.1. Reformulation of the problem. Motivated by [1], we introduce the multi-dimensional effective velocity h as follows:

(4.1)
$$h := u - (2\mu + \lambda)\nabla v \,.$$

By (4.1), the system (1.3) can be deformed into

(4.2)
$$\begin{cases} \rho(\partial_t v + u \cdot \nabla v) - \operatorname{div} h = (2\mu + \lambda)\Delta v, \\ \rho(\partial_t h + u \cdot \nabla h) + \nabla p = R, \end{cases}$$

where

(4.3)
$$R := \frac{2\mu + \lambda}{v} (\nabla u^t \cdot \nabla v - \operatorname{div} u \nabla v) - \mu \nabla \times \nabla \times u$$
$$= (2\mu + \lambda) (\nabla u^t - \operatorname{div} u \mathbb{I}) \nabla \ln v - \mu \nabla \times \nabla \times u$$

and \mathbb{I} denote the 3×3 identity matrix. Similarly, we define

(4.4)
$$h_1^S := u_1^S - (2\mu + \lambda)\partial_1 v^S, \quad h^S := (h_1^S, 0, 0).$$

For (v^S, h_1^S) , it is easy to see that

(4.5)
$$\begin{cases} \rho^{S}(-\sigma\partial_{1}v^{S}+u_{1}^{S}\cdot\partial_{1}v^{S})-\partial_{1}h_{1}^{S}=(2\mu+\lambda)\partial_{11}v^{S}\\ \rho^{S}(-\sigma\partial_{1}h_{1}^{s}+u_{1}^{S}\cdot\partial_{1}h_{1}^{S})+\partial_{1}[p(v^{S})]=0. \end{cases}$$

With the constant σ_* defined in (2.15), (4.5) can be rewritten as

(4.6)
$$\begin{cases} -\sigma_*\partial_1 v^S - \partial_1 h_1^S = (2\mu + \lambda)\partial_{11} v^S \\ -\sigma_*\partial_1 h_1^S + \partial_1 [p(v^S)] = 0. \end{cases}$$

Let

(4.7)
$$\bar{h}_1 := u_1^R + h_1^S - u_{1m}, \quad \bar{h} := (\bar{h}_1, 0, 0).$$

Thus,

$$\bar{h} = \bar{u} - (2\mu + \lambda)\nabla v^{S} = \bar{u} - (2\mu + \lambda)\nabla \bar{v} + (2\mu + \lambda)\nabla v^{R}$$

It can be directly verified that

(4.8)
$$\begin{cases} \rho(\partial_t \bar{v} + u \cdot \nabla \bar{v}) - \operatorname{div}\bar{h} + \rho \dot{\mathbf{X}}(t) \cdot \partial_1 v^S = (2\mu + \lambda)\Delta(\bar{v} - v^R) + Q_1, \\ \rho(\partial_t \bar{h} + u \cdot \nabla \bar{h}) + \nabla p(\bar{v}) + \rho \dot{\mathbf{X}}(t) \cdot \partial_1 h^S = Q_2 \cdot e_1, \end{cases}$$

where $e_1 := (1, 0, 0)^t$,

(4.9)
$$Q_1 := \rho(u_1 - u_1^R) \cdot \partial_1 v^R - \rho(v - v^R) \cdot \partial_1 u_1^R + F \cdot \partial_1 v^S,$$

(4.10)
$$Q_2 := \rho(u_1 - u_1^R) \cdot \partial_1 u_1^R + \rho(v - v^R) \cdot \partial_1 p^R + F \cdot \partial_1 h_1^S + \left[\partial_1 \bar{p} - \partial_1 p^R - \partial_1 p^S\right],$$

(4.11)
$$F := -\sigma(\rho - \rho^S) + (\rho u_1 - \rho^S u_1^S) = \sigma_* \rho(v - v^S) + \rho(u_1 - u_1^S)$$

Subtracting (4.8) from (4.2), we obtain the perturbed system for $(\phi, h - \bar{h}) = (v - \bar{v}, h - \bar{h})$

(4.12)
$$\begin{cases} \rho \partial_t \phi + \rho u \cdot \nabla \phi - \operatorname{div}(h - \bar{h}) - \rho \dot{\mathbf{X}}(t) \cdot \partial_1 v^S = (2\mu + \lambda) \Delta(\phi + v^R) - Q_1, \\ \rho \partial_t (h - \bar{h}) + \rho u \cdot \nabla (h - \bar{h}) + \nabla (p - \bar{p}) - \rho \dot{\mathbf{X}}(t) \cdot \partial_1 h^S = R - Q_2 \cdot e_1. \end{cases}$$

4.2. Estimates on $||(v - \bar{v}, h - \bar{h})||$. In this subsection, we focus on the following lemma.

Lemma 4.1. Under the assumptions of Proposition 3.2, there exists constant C > 0 (independent of ν, δ, χ and T) such that for all $t \in [0, T]$, it holds

at

(4.13)
$$\sup_{t\in[0,T]} \left\| (v-\bar{v},h-\bar{h})(t,\cdot) \right\|^2 + \delta_S \int_0^t |\dot{\mathbf{X}}(\tau)|^2 d\tau + \int_0^t G_2(\tau) + G_3(\tau) + G^R(\tau) + G^S(\tau) + D(\tau) d\tau \leq C \left\| (v-\bar{v},h-\bar{h})(0,\cdot) \right\|^2 + C(\delta+\chi) \int_0^t \left\| \nabla (u-\bar{u}) \right\|_{H^1}^2 d\tau + C \delta_R^{\frac{1}{3}}$$

where

$$G_{2}(t) := \frac{\nu}{\delta_{S}} \int_{\Omega} \partial_{1} v^{S} \left| h_{1} - \bar{h}_{1} - \frac{p - \bar{p}}{\sigma_{*}} \right|^{2} dx,$$

$$G_{3}(t) := \frac{\nu}{\delta_{S}} \int_{\Omega} \partial_{1} v^{S} (h_{2}^{2} + h_{3}^{2}) dx,$$

$$G^{R}(t) := \int_{\Omega} \partial_{1} u_{1}^{R} \cdot p(v|\bar{v}) dx,$$

$$G^{S}(t) := \int_{\Omega} \partial_{1} v^{S} \cdot |p - \bar{p}|^{2} dx,$$

$$D(t) := \int_{\Omega} |\nabla(p - \bar{p})|^{2} dx.$$

First of all, by direct calculations, we can obtain the following lemma.

Lemma 4.2. Let a(t, x) be the weighted function defined in (3.9), then we have

(4.15)
$$\frac{d}{dt} \int_{\Omega} a\rho \left(Q(v|\bar{v}) + \frac{1}{2}|h - \bar{h}|^2 \right) dx = \dot{\mathbf{X}}(t)\mathbf{Y}(t) + \mathbf{B}(t) + \mathbf{S}(t) - \mathbf{G}(t) - \mathbf{D}(t),$$

where

$$\begin{split} \mathbf{Y}(t) &:= -\int_{\Omega} \partial_1 a \cdot \rho \left(Q(v|\bar{v}) + \frac{1}{2} |h - \bar{h}|^2 \right) \, dx - \int_{\Omega} a \rho \partial_1 v^S \cdot p'(\bar{v}) (v - \bar{v}) \, dx \\ &+ \int_{\Omega} a \rho \partial_1 h_1^S \cdot (h_1 - \bar{h}_1) \, dx, \\ \mathbf{B}(t) &:= \sum_{i=1}^8 \mathbf{B}_i(t), \end{split}$$

with

$$\begin{split} \mathbf{B}_{1}(t) &:= \frac{1}{2\sigma_{*}} \int_{\Omega} \partial_{1} a \cdot |p - \bar{p}|^{2} dx, \\ \mathbf{B}_{2}(t) &:= \sigma_{*} \int_{\Omega} a \partial_{1} v^{S} \cdot p(v|\bar{v}) dx, \\ \mathbf{B}_{3}(t) &:= \int_{\Omega} F \cdot \partial_{1} a \left(Q(v|\bar{v}) + \frac{1}{2} |h - \bar{h}|^{2} \right) dx, \\ \mathbf{B}_{4}(t) &:= (2\mu + \lambda) \int_{\Omega} \frac{\partial_{1} a}{p'(v)} (p - \bar{p}) \cdot \partial_{1} (p - \bar{p}) dx, \\ \mathbf{B}_{5}(t) &:= (2\mu + \lambda) \int_{\Omega} \partial_{1} a \cdot (p - \bar{p}) \left(\frac{1}{p'(v)} - \frac{1}{p'(\bar{v})} \right) \cdot \partial_{1} \bar{p} dx, \\ \mathbf{B}_{6}(t) &:= (2\mu + \lambda) \int_{\Omega} a \partial_{1} (p - \bar{p}) \cdot \partial_{1} \bar{p} \left(\frac{1}{p'(v)} - \frac{1}{p'(\bar{v})} \right) dx, \\ \mathbf{B}_{7}(t) &:= -\int_{\Omega} a (p - \bar{p}) \cdot \Delta v^{R} dx, \\ \mathbf{B}_{8}(t) &:= \int_{\Omega} a (h - \bar{h}) \cdot R dx, \end{split}$$

and

$$\begin{split} \mathbf{S}(t) &:= \int_{\Omega} a \left[p'(\bar{v})(v - \bar{v}) \cdot Q_1 - (h_1 - \bar{h}_1) \cdot Q_2 \right] \, dx, \\ \mathbf{D}(t) &:= (2\mu + \lambda) \int_{\Omega} \frac{a}{|p'(v)|} |\nabla(p - \bar{p})|^2 \, dx, \\ \mathbf{G}(t) &:= \sum_{i=1}^3 \mathbf{G}_i(t) + \mathbf{G}^R(t), \\ \mathbf{G}_1(t) &:= \sigma_* \int_{\Omega} \partial_1 a \cdot Q(v|\bar{v}) \, dx, \\ \mathbf{G}_2(t) &:= \frac{\sigma_*}{2} \int_{\Omega} \partial_1 a \cdot \left| h_1 - \bar{h}_1 - \frac{p - \bar{p}}{\sigma_*} \right|^2 \, dx, \end{split}$$

$$\mathbf{G}_{3}(t) := \sigma_{*} \int_{\Omega} \partial_{1} a \cdot \frac{h_{2}^{2} + h_{3}^{2}}{2} \, dx,$$
$$\mathbf{G}^{R}(t) := \int_{\Omega} a \partial_{1} u_{1}^{R} \cdot p(v|\bar{v}) \, dx.$$

Proof. Multiplying $(4.12)_1$ by $-a(p-\bar{p})$, we have

$$\begin{aligned} \partial_t \left[a\rho Q(v|\bar{v}) \right] + \operatorname{div} \left[a\rho Q(v|\bar{v})u \right] + a(p-\bar{p})\operatorname{div}(h-\bar{h}) \\ &= \partial_1 a \left[u_1 - \sigma - \dot{\mathbf{X}}(t) \right] \rho Q(v|\bar{v}) - a\rho \dot{\mathbf{X}}(t) \partial_1 v^S \cdot (p-\bar{p}) \\ &- ap(v|\bar{v}) \left[\rho \partial_t \bar{v} + \rho u \cdot \nabla \bar{v} \right] - a(2\mu + \lambda)(p-\bar{p}) \Delta(\phi + v^R) + aQ_1(p-\bar{p}) \\ &= (\rho u_1 - \rho \sigma) \partial_1 a \cdot Q(v|\bar{v}) - \dot{\mathbf{X}}(t) \left[\rho \partial_1 a \cdot Q(v|\bar{v}) + a\rho \partial_1 v^S \cdot p'(\bar{v})(v-\bar{v}) \right] \\ &- a\operatorname{div} \bar{u} \cdot p(v|\bar{v}) - a(2\mu + \lambda)(p-\bar{p}) \Delta(\phi + v^R) + aQ_1 p'(\bar{v})(v-\bar{v}), \end{aligned}$$

Here, we use the fact that

$$\rho \partial_t \bar{v} + \rho u \cdot \nabla \bar{v} = -\rho \dot{\mathbf{X}}(t) \partial_1 v^S + \operatorname{div} \bar{u} + Q_1.$$

Recall the definition of F and σ_* in (4.11) and (2.15), it holds

$$\rho u_1 - \rho \sigma = -\sigma \rho^S + \rho^S u_1^S - \sigma (\rho - \rho^S) + (\rho u_1 - \rho^S u_1^S) = -\sigma_* + F.$$

Also, with the fact $-\sigma_*\partial_1 v^S = \partial_1 u_1^S$ form (2.16), we have

(4.17)

$$\frac{\partial_t \left[a\rho Q(v|\bar{v}) \right] + \operatorname{div} \left[a\rho Q(v|\bar{v})u \right] + a(p-\bar{p})\operatorname{div}(h-h) \\
= (F - \sigma_*)\partial_1 a \cdot Q(v|\bar{v}) - \dot{\mathbf{X}}(t) \left[\rho \partial_1 a \cdot Q(v|\bar{v}) + a\rho \partial_1 v^S \cdot p'(\bar{v})(v-\bar{v}) \right] \\
+ a(\sigma_* \partial_1 v^S - \partial_1 u_1^R) \cdot p(v|\bar{v}) - a(2\mu + \lambda)(p-\bar{p})\Delta v^R \\
- (2\mu + \lambda)\operatorname{div} \left[a(p-\bar{p})\nabla \phi \right] + (2\mu + \lambda)\nabla \left[a(p-\bar{p}) \right] \cdot \nabla \phi + aQ_1 p'(\bar{v})(v-\bar{v}).$$

Since

$$\nabla v = \frac{\nabla p(v)}{p'(v)} = \frac{\nabla p(v)}{-\gamma p^{1+\frac{1}{\gamma}}(v)},$$

we can make the following transformations

$$(2\mu + \lambda)\nabla[a(p - \bar{p})] \cdot \nabla\phi$$

$$= (2\mu + \lambda)a\nabla(p - \bar{p}) \cdot \left(\frac{\nabla p(v)}{p'(v)} - \frac{\nabla p(\bar{v})}{p'(\bar{v})}\right) + (2\mu + \lambda)\partial_1 a \cdot (p - \bar{p}) \left(\frac{\partial_1 p(v)}{p'(v)} - \frac{\partial_1 p(\bar{v})}{p'(\bar{v})}\right)$$

$$(4.18)$$

$$= -(2\mu + \lambda)a\frac{|\nabla(p - \bar{p})|^2}{|p'(v)|} - (2\mu + \lambda)a\partial_1(p - \bar{p})\partial_1\bar{p} \cdot \left(\frac{1}{p'(v)} - \frac{1}{p'(\bar{v})}\right)$$

$$+ (2\mu + \lambda)\partial_1 a \cdot (p - \bar{p})\frac{\partial_1(p - \bar{p})}{p'(v)} + (2\mu + \lambda)\partial_1 a \cdot (p - \bar{p})\partial_1\bar{p} \cdot \left(\frac{1}{p'(v)} - \frac{1}{p'(\bar{v})}\right).$$

Then, multiplying $(4.12)_2$ by $a(h - \bar{h})$, we have

$$\partial_t \left[a\rho \frac{|h-\bar{h}|^2}{2} \right] + \operatorname{div} \left[a\rho \frac{|h-\bar{h}|^2}{2} u \right] + \operatorname{div} \left[a(p-\bar{p})(h-\bar{h}) \right] - a(p-\bar{p})\operatorname{div}(h-\bar{h})$$
$$= \partial_1 a \left[u_1 - \sigma - \dot{\mathbf{X}}(t) \right] \rho \frac{|h-\bar{h}|^2}{2} + a\rho \dot{\mathbf{X}}(t) \partial_1 h^S \cdot (h-\bar{h}) + \partial_1 a(p-\bar{p})(h_1 - \bar{h}_1)$$
$$= \partial_1 a \left[u_1 - \sigma - \dot{\mathbf{X}}(t) \right] \rho \frac{|h-\bar{h}|^2}{2} + a\rho \dot{\mathbf{X}}(t) \partial_1 h^S \cdot (h-\bar{h}) + \partial_1 a(p-\bar{p})(h_1 - \bar{h}_1)$$

(4.19)

$$= \partial_{1}a \cdot F \frac{|h-\bar{h}|^{2}}{2} + \dot{\mathbf{X}}(t) \left[a\rho\partial_{1}h^{S} \cdot (h-\bar{h}) - \rho\partial_{1}a \cdot \frac{|h-\bar{h}|^{2}}{2} \right] - \sigma_{*}\partial_{1}a \cdot F \frac{|h-\bar{h}|^{2}}{2} + \partial_{1}a(p-\bar{p})(h_{1}-\bar{h}_{1}) + aR \cdot (h-\bar{h}) - Q_{2}(h_{1}-\bar{h}_{1}).$$

By direct computation, we have

(4.20)
$$= -\frac{\sigma_* \partial_1 a \cdot F \frac{|h - \bar{h}|^2}{2} + \partial_1 a (p - \bar{p}) (h_1 - \bar{h}_1) \\ = -\frac{\sigma_*}{2} \partial_1 a \left| h_1 - \bar{h}_1 - \frac{p - \bar{p}}{\sigma_*} \right|^2 + \partial_1 a \cdot \frac{|p - \bar{p}|^2}{2\sigma_*} - \sigma_* \partial_1 a \cdot \frac{h_2^2 + h_3^2}{2},$$

Adding (4.17) and (4.19) together, and integrating the resultant equation on Ω , we can achieve (4.15). \Box

For notational simplicity, we decompose the $\mathbf{Y}(t)$ in Lemma 4.2 as follows:

$$\mathbf{Y}(t) = -\int_{\Omega} \partial_1 a \cdot \rho \left(Q(v|\bar{v}) + \frac{1}{2}|h - \bar{h}|^2 \right) dx - \int_{\Omega} a\rho \partial_1 v^S \cdot p'(\bar{v})(v - \bar{v}) dx + \int_{\Omega} a\rho \partial_1 h_1^S \cdot (h_1 - \bar{h}_1) dx =: \sum_{i=1}^5 \mathbf{Y}_i(t),$$

where

$$\begin{split} \mathbf{Y}_{1}(t) &:= -\int_{\Omega} a\rho \partial_{1} v^{S} \cdot p'(v^{S})(v-\bar{v}) \, dx, \\ \mathbf{Y}_{2}(t) &:= \int_{\Omega} \frac{a\rho}{\sigma_{*}} \cdot \partial_{1} h_{1}^{S} \cdot (p-\bar{p}) \, dx, \\ \mathbf{Y}_{3}(t) &:= \int_{\Omega} a\rho \partial_{1} v^{S} \cdot [p'(v^{S}) - p'(\bar{v})](v-\bar{v}) \, dx, \\ \mathbf{Y}_{4}(t) &:= \int_{\Omega} a\rho \partial_{1} h_{1}^{S} \cdot \left(h_{1} - \bar{h}_{1} - \frac{p-\bar{p}}{\sigma_{*}}\right) \, dx, \\ \mathbf{Y}_{5}(t) &:= -\frac{1}{2} \int_{\Omega} \partial_{1} a \cdot \rho \left(h_{1} - \bar{h}_{1} - \frac{p-\bar{p}}{\sigma_{*}}\right) \left(h_{1} - \bar{h}_{1} + \frac{p-\bar{p}}{\sigma_{*}}\right) \, dx, \\ \mathbf{Y}_{6}(t) &:= -\int_{\Omega} \partial_{1} a \cdot \rho \left(Q(v|\bar{v}) + \frac{h_{2}^{2} + h_{3}^{2}}{2}\right) \, dx - \int_{\Omega} \partial_{1} a \cdot \rho \frac{|p-\bar{p}|^{2}}{2\sigma_{*}^{2}} \, dx \end{split}$$

Then, according to (3.15), we have

(4.21)
$$\dot{\mathbf{X}}(t) = -\frac{M}{\delta_S} \big[\mathbf{Y}_1(t) + \mathbf{Y}_2(t) \big],$$

and

(4.22)
$$\dot{\mathbf{X}}(t)\mathbf{Y}(t) = -\frac{M}{\delta_S}|\dot{\mathbf{X}}(t)|^2 + \dot{\mathbf{X}}(t)\sum_{i=3}^{5}\mathbf{Y}_i(t).$$

Secondly, we give the following leading order estimates, which is crucial for the proof of Lemma 4.1.

Lemma 4.3. There exists uniform-in-time $C_0 > 0$ such that for all $t \in [0, T]$, it holds

(4.23)
$$-\frac{\delta_S}{2M}|\dot{\mathbf{X}}(t)|^2 + \mathbf{B}_1(t) + \mathbf{B}_2(t) - \mathbf{G}_1(t) - \frac{3}{4}\mathbf{D}(t) \le -C_0 G^S(t) + C \int_{\Omega} \partial_1 a \cdot |p - \bar{p}|^3 \, dx,$$

where $G^S(t) := \int_{\Omega} \partial_1 v^S \cdot |p - \bar{p}|^2 \, dx.$

Proof. First, we introduce the following time-dependent change of variables:

(4.24)
$$\begin{cases} y_1 = \frac{p(v_m) - p(v^S)}{\delta_S} = \frac{p(v_m) - p(v^s(x_1 - \sigma t - \mathbf{X}(t)))}{\delta_S}, \\ y_2 = x_2, \\ y_3 = x_3, \end{cases}$$

and set

(4.25)
$$w(t,y) := (p - \bar{p})(t, x(y)), \qquad y' := (y_2, y_3)$$

By direct computation, we know that $y = (y_1, y') \in (0, 1) \times \mathbb{T}^2$. Also, it follows from (3.9) that $a = 1 + \nu y_1$ and

(4.26)
$$\frac{dy_1}{dx_1} = -\frac{1}{\delta_S} \partial_{x_1} p^S, \quad \partial_{x_1} a = \nu \frac{dy_1}{dx_1}, \quad |a-1| \le \nu = \sqrt{\delta_S}.$$

In the above new variables, we have

(4.27)
$$\int_{\mathbb{T}^2} \int_{\mathbb{R}} \partial_1 a |p - \bar{p}|^2 \, dx_1 dx' = \nu \int_{\mathbb{T}^2} \int_0^1 w^2 \, dy_1 dy',$$

and so

(4.28)
$$\int_{\mathbb{T}^2} \int_{\mathbb{R}} \partial_1 v^S |p - \bar{p}|^2 dx_1 dx' \sim \delta_S \int_{\mathbb{T}^2} \int_0^1 w^2 dy_1 dy'.$$

For a finer estimate, we introduce two constants

$$\sigma_m := \sqrt{-p'(v_m)}, \quad \alpha_m := \frac{\gamma + 1}{2\gamma \sigma_m p(v_m)}.$$

Then, from (2.18) and the fact that $\sigma_m^2 = -p'(v_m) = \gamma p^{1+\frac{1}{\gamma}}(v_m)$, we obtain (4.29) $|\sigma_* - \sigma_m| \le C\delta_S$,

and

(4.30)
$$|\sigma_m^2 + p'(\bar{v})| \le C\delta, \quad \left|\frac{1}{\sigma_m^2} - \frac{1}{\gamma p^{1+\frac{1}{\gamma}}(\bar{v})}\right| \le C\delta.$$

• Estimates on $-\frac{\delta_S}{2M} |\dot{\mathbf{X}}(t)|^2$: According to (4.21), we only have to estimate $\mathbf{Y}_1(t)$ and $\mathbf{Y}_2(t)$. For one thing,

$$\mathbf{Y}_1(t) = -\int_{\Omega} a\rho \partial_1 p^S(v-\bar{v}) \, dx = \delta_S \int_{\mathbb{T}^2} \int_0^1 \frac{a}{v} (v-\bar{v}) \, dy_1 dy'.$$

Thus, by Taylor expansion and that $|v - \bar{v}| \sim |p - \bar{p}|$, it holds

$$|p(v) - p(\bar{v}) - p'(\bar{v})(v - \bar{v})| \le C|p - \bar{p}|^2$$

and so

(4.31)
$$\left| v - \bar{v} + \frac{p - \bar{p}}{\sigma_m^2} \right| \le C(\delta + \chi)|p - \bar{p}|, \quad \left| v - \bar{v} + \frac{p - \bar{p}}{\sigma_*^2} \right| \le C(\delta + \chi)|p - \bar{p}|.$$

Together with $|a - 1| \leq \nu$ form (3.9), we get

(4.32)
$$\left| \mathbf{Y}_1(t) + \frac{\delta_S}{\sigma_m^2 v_m} \int_{\mathbb{T}^2} \int_0^1 w \, dy_1 dy' \right| \le C \delta_S(\nu + \delta + \chi) \int_{\mathbb{T}^2} \int_0^1 |w| \, dy_1 dy'.$$

For another thing, by (4.6), we yield

$$\mathbf{Y}_2(t) = \int_{\Omega} \frac{a}{v\sigma_*^2} \cdot \partial_1 p^S \cdot (p - \bar{p}) \, dx = -\frac{\delta_S}{\sigma_*^2} \int_{\mathbb{T}^2} \int_0^1 \frac{a}{v} \cdot w \, dy_1 dy',$$

and thus

(4.33)
$$\left| \mathbf{Y}_{2}(t) + \frac{\delta_{S}}{\sigma_{m}^{2} v_{m}} \int_{\mathbb{T}^{2}} \int_{0}^{1} w \, dy_{1} dy' \right| \leq C \delta_{S}(\nu + \delta + \chi) \int_{\mathbb{T}^{2}} \int_{0}^{1} |w| \, dy_{1} dy'.$$

By (4.21), (4.32) and (4.33), we have

$$\begin{aligned} \left| \dot{\mathbf{X}}(t) - \frac{2M}{\sigma_m^2 v_m} \int_{\mathbb{T}^2} \int_0^1 w \, dy_1 dy' \right| &= \left| \sum_{i=1}^2 \frac{M}{\delta_S} \left(\mathbf{Y}_i(t) + \frac{\delta_S}{\sigma_m^2 v_m} \int_{\mathbb{T}^2} \int_0^1 w \, dy_1 dy' \right) \right| \\ &\leq C(\nu + \delta + \chi) \int_{\mathbb{T}^2} \int_0^1 |w| \, dy_1 dy', \end{aligned}$$

which leads to

$$\left(\left| \frac{2M}{\sigma_m^2 v_m} \int_{\mathbb{T}^2} \int_0^1 w \, dy_1 dy' \right| - |\dot{\mathbf{X}}(t)| \right)^2 \le C(\nu + \delta + \chi)^2 \left(\int_{\mathbb{T}^2} \int_0^1 |w| \, dy_1 dy' \right)^2 \\ \le C(\nu + \delta + \chi)^2 \int_{\mathbb{T}^2} \int_0^1 |w|^2 \, dy_1 dy'.$$

Finally, with the algebraic inequality $\frac{p^2}{2} - q^2 \le (p-q)^2, \forall p, q \ge 0$, we arrive at

$$\frac{2M^2}{\sigma_m^4 v_m^2} \left(\int_{\mathbb{T}^2} \int_0^1 w \, dy_1 dy' \right)^2 - |\dot{\mathbf{X}}(t)|^2 \le C(\nu + \delta + \chi)^2 \int_{\mathbb{T}^2} \int_0^1 |w|^2 \, dy_1 dy',$$

and consequently

(4.34)
$$-\frac{\delta_S}{2M}|\dot{\mathbf{X}}(t)|^2 \le -\frac{M\delta_S}{\sigma_m^4 v_m^2} \left(\int_{\mathbb{T}^2} \int_0^1 w \, dy_1 dy'\right)^2 + C\delta_S(\nu + \delta + \chi)^2 \int_{\mathbb{T}^2} \int_0^1 |w|^2 \, dy_1 dy'.$$

• Estimates on $\mathbf{B}_1(t) - \mathbf{G}_1(t)$ and $\mathbf{B}_2(t)$: First, by (4.29),

(4.35)
$$\mathbf{B}_{1}(t) = \frac{1}{2\sigma_{*}} \int_{\Omega} \partial_{1} a \cdot |p - \bar{p}|^{2} dx = \frac{\nu}{2\sigma_{*}} \int_{\mathbb{T}^{2}} \int_{0}^{1} w^{2} dy_{1} dy' \\ \leq \frac{\nu}{2\sigma_{m}} \int_{\mathbb{T}^{2}} \int_{0}^{1} w^{2} dy_{1} dy' + C\nu \delta_{S} \int_{\mathbb{T}^{2}} \int_{0}^{1} w^{2} dy_{1} dy'.$$

For $\mathbf{B}_2(t)$, by (3.5), we have

$$(4.36) \qquad \mathbf{B}_{2}(t) = \sigma_{*} \int_{\Omega} a\partial_{1}v^{S} \cdot p(v|\bar{v}) \, dx = \sigma_{*}\delta_{S} \int_{\mathbb{T}^{2}} \int_{0}^{1} \frac{a}{|p'(v^{S})|} \cdot p(v|\bar{v}) \, dy_{1} dy' \\ \leq \sigma_{*}\delta_{S}(1+\nu) \int_{\mathbb{T}^{2}} \int_{0}^{1} \left(\frac{\gamma+1}{2\gamma} \frac{1}{p(v^{S})} + C\chi\right) \frac{|p-\bar{p}|^{2}}{|p'(v^{S})|} \, dy_{1} dy' \\ \leq \sigma_{*}\delta_{S}(1+\nu) \int_{\mathbb{T}^{2}} \int_{0}^{1} \left(\alpha_{m} \frac{\sigma_{m}p(v_{m})}{p(v^{S})} + C\chi\right) \frac{w^{2}}{|p'(v^{S})|} \, dy_{1} dy' \\ \leq \delta_{S}\alpha_{m} \int_{\mathbb{T}^{2}} \int_{0}^{1} w^{2} \, dy_{1} dy' + C(\nu+\delta+\chi) \int_{\mathbb{T}^{2}} \int_{0}^{1} w^{2} \, dy_{1} dy'.$$

For $G_1(t)$, by (3.6) and (4.30), we get

$$(4.37) \qquad \mathbf{G}_{1}(t) = \sigma_{*} \int_{\Omega} \partial_{1} a \cdot Q(v|\bar{v}) \, dx,$$

$$\geq \sigma_{*} \int_{\Omega} \partial_{1} a \cdot \frac{|p - \bar{p}|^{2}}{2\gamma p^{1 + \frac{1}{\gamma}}(\bar{v})} \, dx - \sigma_{*} \int_{\Omega} \partial_{1} a \cdot \frac{1 + \gamma}{3\gamma^{2}} \frac{(p - \bar{p})^{3}}{p^{2 + \frac{1}{\gamma}}(\bar{v})} \, dx$$

$$\geq \frac{\nu \sigma_{*}}{2\sigma_{m}^{2}} (1 - C\delta) \int_{\mathbb{T}^{2}} \int_{0}^{1} w^{2} \, dy_{1} dy' - C\nu \int_{\mathbb{T}^{2}} \int_{0}^{1} w^{3} \, dy_{1} dy'$$

$$\geq \frac{\nu}{2\sigma_{m}} \int_{\mathbb{T}^{2}} \int_{0}^{1} w^{2} \, dy_{1} dy' - C\nu \delta_{S} \int_{\mathbb{T}^{2}} \int_{0}^{1} w^{2} \, dy_{1} dy' - C\nu \int_{\mathbb{T}^{2}} \int_{0}^{1} w^{3} \, dy_{1} dy',$$

Hence, together with (4.35), we acquire

(4.38)
$$\mathbf{B}_{new}(t) := \mathbf{B}_1(t) - \mathbf{G}_1(t) \le C\nu\delta_S \int_{\mathbb{T}^2} \int_0^1 w^2 \, dy_1 dy' + C\nu \int_{\mathbb{T}^2} \int_0^1 w^3 \, dy_1 dy'.$$

• Estimate on D(t): First, by (3.9) and the change of variables (4.24), we get

$$\begin{split} \mathbf{D}(t) &\geq (2\mu + \lambda) \int_{\mathbb{T}^2} \int_{\mathbb{R}} \frac{|\partial_{x_1} \left(p(v) - p(\bar{v}) \right)|^2}{\gamma p^{1 + \frac{1}{\gamma}}(v)} + \frac{|\nabla_{x'} \left(p(v) - p(\bar{v}) \right)|^2}{\gamma p^{1 + \frac{1}{\gamma}}(v)} dx_1 dx' \\ &= \underbrace{(2\mu + \lambda) \!\int_{\mathbb{T}^2} \!\int_0^1 \frac{|\partial_{y_1} w|^2}{\gamma p^{1 + \frac{1}{\gamma}}(v)} \Big(\frac{dy_1}{dx_1} \Big) dy_1 dy'}_{\mathbf{D}_{\mathrm{I}}(t)} + \underbrace{(2\mu + \lambda) \!\int_{\mathbb{T}^2} \!\int_0^1 \frac{|\nabla_{y'} w|^2}{\gamma p^{1 + \frac{1}{\gamma}}(v)} \Big(\frac{dx_1}{dy_1} \Big) dy_1 dy'}_{\mathbf{D}_{\mathrm{II}}(t)} + \underbrace{(2\mu + \lambda) \!\int_{\mathbb{T}^2} \int_0^1 \frac{|\nabla_{y'} w|^2}{\gamma p^{1 + \frac{1}{\gamma}}(v)} \Big(\frac{dx_1}{dy_1} \Big) dy_1 dy'}_{\mathbf{D}_{\mathrm{II}}(t)} + \underbrace{(2\mu + \lambda) \!\int_{\mathbb{T}^2} \int_0^1 \frac{|\nabla_{y'} w|^2}{\gamma p^{1 + \frac{1}{\gamma}}(v)} \Big(\frac{dx_1}{dy_1} \Big) dy_1 dy'}_{\mathbf{D}_{\mathrm{II}}(t)} + \underbrace{(2\mu + \lambda) \!\int_{\mathbb{T}^2} \int_0^1 \frac{|\nabla_{y'} w|^2}{\gamma p^{1 + \frac{1}{\gamma}}(v)} \Big(\frac{dx_1}{dy_1} \Big) dy_1 dy'}_{\mathbf{D}_{\mathrm{II}}(t)} + \underbrace{(2\mu + \lambda) \!\int_{\mathbb{T}^2} \int_0^1 \frac{|\nabla_{y'} w|^2}{\gamma p^{1 + \frac{1}{\gamma}}(v)} \Big(\frac{dx_1}{dy_1} \Big) dy_1 dy'}_{\mathbf{D}_{\mathrm{II}}(t)} + \underbrace{(2\mu + \lambda) \!\int_{\mathbb{T}^2} \int_0^1 \frac{|\nabla_{y'} w|^2}{\gamma p^{1 + \frac{1}{\gamma}}(v)} \Big(\frac{dx_1}{dy_1} \Big) dy_1 dy'}_{\mathbf{D}_{\mathrm{II}}(t)} + \underbrace{(2\mu + \lambda) \!\int_{\mathbb{T}^2} \int_0^1 \frac{|\nabla_{y'} w|^2}{\gamma p^{1 + \frac{1}{\gamma}}(v)} \Big(\frac{dx_1}{dy_1} \Big) dy_1 dy'}_{\mathbf{D}_{\mathrm{II}}(t)} + \underbrace{(2\mu + \lambda) \!\int_{\mathbb{T}^2} \int_0^1 \frac{|\nabla_{y'} w|^2}{\gamma p^{1 + \frac{1}{\gamma}}(v)} \Big(\frac{dx_1}{dy_1} \Big) dy_1 dy'}_{\mathbf{D}_{\mathrm{II}}(t)} + \underbrace{(2\mu + \lambda) \!\int_{\mathbb{T}^2} \int_0^1 \frac{|\nabla_{y'} w|^2}{\gamma p^{1 + \frac{1}{\gamma}}(v)} \Big(\frac{dx_1}{dy_1} \Big) dy_1 dy'}_{\mathbf{D}_{\mathrm{II}}(t)} + \underbrace{(2\mu + \lambda) \!\int_{\mathbb{T}^2} \int_0^1 \frac{|\nabla_{y'} w|^2}{\gamma p^{1 + \frac{1}{\gamma}}(v)} \Big(\frac{dx_1}{dy_1} \Big) dy_1 dy'}_{\mathbf{D}_{\mathrm{II}}(t)} + \underbrace{(2\mu + \lambda) \!\int_{\mathbb{T}^2} \int_0^1 \frac{|\nabla_{y'} w|^2}{\gamma p^{1 + \frac{1}{\gamma}}(v)} \Big(\frac{dx_1}{dy_1} \Big) dy_1 dy'}_{\mathbf{D}_{\mathrm{II}}(t)} + \underbrace{(2\mu + \lambda) \!\int_{\mathbb{T}^2} \int_0^1 \frac{dx_1}{\gamma p^{1 + \frac{1}{\gamma}}(v)} \Big(\frac{dx_1}{dy_1} \Big) dy_1 dy'}_{\mathbf{D}_{\mathrm{II}}(t)} + \underbrace{(2\mu + \lambda) \!\int_{\mathbb{T}^2} \int_0^1 \frac{dx_1}{\gamma p^{1 + \frac{1}{\gamma}}(v)} \Big(\frac{dx_1}{dy_1} \Big) dy_1 dy'}_{\mathbf{D}_{\mathrm{II}}(t)} + \underbrace{(2\mu + \lambda) \!\int_{\mathbb{T}^2} \int_0^1 \frac{dx_1}{\gamma p^{1 + \frac{1}{\gamma}}(v)} \Big(\frac{dx_1}{dy_1} \Big) dy_1 dy'}_{\mathbf{D}_{\mathrm{II}}(t)} + \underbrace{(2\mu + \lambda) \!\int_{\mathbb{T}^2} \int_0^1 \frac{dx_1}{\gamma p^{1 + \frac{1}{\gamma}}(v)} \Big(\frac{dx_1}{\eta p^{1 + \frac{1}{\gamma}}(v)} \Big(\frac{dx_1}{\eta p^{1 + \frac{1}{\gamma}}(v)} \Big) dy_1 dy'}_$$

For one thing, integrating (4.6) over $(-\infty, x_1]$ leads to

$$(2\mu + \lambda)\partial_{x_1}v^S = -\sigma_*(v^S - v_m) - \frac{1}{\sigma_*}(p(v^S) - p(v_m)).$$

For another thing,

$$\partial_{x_1} v^S = \frac{\partial_{x_1} p(v^S)}{p'(v^S)} = \frac{\delta_S}{\gamma p^{1+\frac{1}{\gamma}}(v^S)} \frac{dy_1}{dx_1} \ .$$

Therefore, we have

$$(2\mu + \lambda) \frac{\delta_S}{\gamma p^{1 + \frac{1}{\gamma}} (v^S)} \frac{dy_1}{dx_1} = -\sigma_* (v^S - v_m) - \frac{1}{\sigma_*} (p(v^S) - p(v_m))$$
$$= -\frac{1}{\sigma_*} \Big(\sigma_*^2 (v^S - v_m) + (p(v^S) - p(v_m)) \Big),$$

which together with $\sigma_*^2 = -\frac{p(v_m) - p(v_+)}{v_m - v_+}$ implies

$$(2\mu + \lambda) \frac{\delta_S}{\gamma p^{1+\frac{1}{\gamma}} (v^S)} \frac{dy_1}{dx_1}$$

= $\frac{-1}{\sigma_* (v_+ - v_m)} \Big((p(v_m) - p(v_+))(v^S - v_m) + (p(v^S) - p(v_m))(v_+ - v_m) \Big)$
= $\frac{-1}{\sigma_* (v_+ - v_m)} \Big((p(v^S) - p(v_+))(v^S - v_m) - (v^S - v_+)(p(v^S) - p(v_m)) \Big).$

By (4.24) and (2.22), it holds $y_1 = \frac{p(v_m) - p(v^S)}{\delta_S}$ and $1 - y_1 = \frac{p(v^S) - p(v_+)}{\delta_S}$, then

$$\frac{1}{y_1(1-y_1)}\frac{2\mu+\lambda}{\gamma p^{1+\frac{1}{\gamma}}(v^S)}\frac{dy_1}{dx_1} = \frac{\delta_S}{\sigma_*(v_+-v_m)}\left(\frac{v^S-v_m}{p(v^S)-p(v_m)} - \frac{v^S-v_+}{p(v^S)-p(v_+)}\right).$$

Thus

$$\begin{aligned} & \left| \frac{1}{y_1(1-y_1)} \frac{2\mu + \lambda}{\gamma p^{1+\frac{1}{\gamma}}(v^S)} \frac{dy_1}{dx_1} - \frac{\delta_S p''(v_m)}{2\sigma_m (p'(v_m))^2} \right| \\ & \leq \left| \frac{1}{y_1(1-y_1)} \frac{2\mu + \lambda}{\gamma p^{1+\frac{1}{\gamma}}(v^S)} \frac{dy_1}{dx_1} - \frac{\delta_S p''(v_m)}{2\sigma_* (p'(v_m))^2} \right| + \frac{\delta_S p''(v_m)}{2(p'(v_m))^2} \left| \frac{1}{\sigma_*} - \frac{1}{\sigma_m} \right| \\ & \leq \left| \frac{\delta_S}{\sigma_* (v_+ - v_m)} \left(\frac{v^S - v_m}{p(v^S) - p(v_m)} + \frac{v^S - v_+}{p(v_+) - p(v^S)} \right) \right| + \frac{\delta_S p''(v_m)}{2(p'(v_m))^2} \left| \frac{1}{\sigma_*} - \frac{1}{\sigma_m} \right| \end{aligned}$$

Using Lemma 3.4, we have

$$\frac{\delta_S}{\sigma_*(v_+ - v_m)} \left(\frac{v^S - v_m}{p(v^S) - p(v_m)} + \frac{v^S - v_+}{p(v_+) - p(v^S)} \right) - \frac{\delta_S p''(v_m)}{2\sigma_*(p'(v_m))^2} \bigg| \le C\delta_S^2.$$

Also, it follows from (4.29) that

$$\frac{\delta_S p''(v_m)}{2(p'(v_m))^2} \left| \frac{1}{\sigma_*} - \frac{1}{\sigma_m} \right| \le C \delta_S^2 \,,$$

Hence, we get

(4.39)
$$\left| \frac{1}{y_1(1-y_1)} \frac{2\mu + \lambda}{\gamma p^{1+\frac{1}{\gamma}} (v^S)} \frac{dy_1}{dx_1} - \frac{\delta_S p''(v_m)}{2\sigma_m (p'(v_m))^2} \right| \le C \delta_S^2.$$

In addition,

$$\left(\frac{p(v^S)}{p(v)}\right)^{1+\frac{1}{\gamma}} - 1 \right| \le C|v - v^S| \le C(\chi + \delta_R).$$

. 1

Thus, we obtain

$$\begin{aligned} \mathbf{D}_{\mathrm{I}}(t) &= \int_{\mathbb{T}^2} \int_0^1 y_1(1-y_1) |\partial_{y_1}w|^2 \left(\frac{p(v^S)}{p(v)}\right)^{1+\frac{1}{\gamma}} \frac{1}{y_1(1-y_1)} \frac{2\mu+\lambda}{\gamma p^{1+\frac{1}{\gamma}}(v^S)} \left(\frac{dy_1}{dx_1}\right) dy_1 dy' \\ &\geq \left(1 - C(\chi+\delta_R)\right) \left(\frac{\delta_S p''(v_m)}{2\sigma_m (p'(v_m))^2} - C\delta_S^2\right) \int_{\mathbb{T}^2} \int_0^1 y_1(1-y_1) |\partial_{y_1}w|^2 dy_1 dy'. \end{aligned}$$

A direct computation gives rise to

$$\frac{p''(v_m)}{2\sigma_m(p'(v_m))^2} = \frac{\gamma+1}{2\gamma\sigma_m p(v_m)} = \alpha_m,$$

which leads to

$$\mathbf{D}_{\mathrm{I}}(t) \ge \alpha_m \delta_S (1 - C(\delta + \chi)) \int_{\mathbb{T}^2} \int_0^1 y_1 (1 - y_1) |\partial_{y_1} w|^2 dy_1 dy'.$$

It follows from (4.39) that

$$y_1(1-y_1)\frac{dx_1}{dy_1} \ge \frac{2\mu + \lambda}{\gamma p^{1+\frac{1}{\gamma}}(v^S)} \frac{1}{\alpha_m \delta_S + C\delta_S^2} \ge \frac{2\mu + \lambda}{2\alpha_m \delta_S |p'(v_m)|}$$

Hence, we get

$$\begin{aligned} \mathbf{D}_{\mathrm{II}}(t) &= (2\mu + \lambda) \int_{\mathbb{T}^2} \int_0^1 \frac{|\nabla_{y'}w|^2}{y_1(1-y_1)} \left(\frac{p(v_m)}{p(v)}\right)^{1+\frac{1}{\gamma}} \frac{y_1(1-y_1)}{\gamma p^{1+\frac{1}{\gamma}}(v_m)} \left(\frac{dx_1}{dy_1}\right) dy_1 dy' \\ &\geq (1 - C(\delta + \chi))(2\mu + \lambda) \int_{\mathbb{T}^2} \int_0^1 \frac{|\nabla_{y'}w|^2}{y_1(1-y_1)} \frac{2\mu + \lambda}{|p'(v_m)|^2} \frac{1}{2\alpha_m \delta_S} dy_1 dy' \\ &\geq (1 - C(\delta + \chi)) \frac{\sigma_m}{\delta_S} \frac{(2\mu + \lambda)^2}{p''(v_m)} \int_{\mathbb{T}^2} \int_0^1 \frac{|\nabla_{y'}w|^2}{y_1(1-y_1)} dy_1 dy'. \end{aligned}$$

Combining the estimates on $\mathbf{D}_{\mathrm{I}}(t)$ and $\mathbf{D}_{\mathrm{II}}(t)$, we arrive at

(4.40)
$$\mathbf{D}(t) \ge \alpha_m \delta_S (1 - C(\delta + \chi)) \int_{\mathbb{T}^2} \int_0^1 y_1 (1 - y_1) |\partial_{y_1} w|^2 dy_1 dy' + (1 - C(\delta + \chi)) \frac{\sigma_m}{\delta_S} \frac{(2\mu + \lambda)^2}{p''(v_m)} \int_{\mathbb{T}^2} \int_0^1 \frac{|\nabla_{y'} w|^2}{y_1 (1 - y_1)} dy_1 dy'.$$

• Proof of Lemma 4.3 : First of all, in view of (4.36), (4.38) and (4.40), it follows that

$$\begin{split} \mathbf{B}_{1}(t) + \mathbf{B}_{2}(t) - \mathbf{G}_{1}(t) &- \frac{3}{4}\mathbf{D}(t) \\ \leq \alpha_{m}\delta_{S}\int_{\mathbb{T}^{2}}\int_{0}^{1}w^{2}\,dy_{1}dy' - \frac{3}{4}\alpha_{m}\delta_{S}\int_{\mathbb{T}^{2}}\int_{0}^{1}y_{1}(1-y_{1})|\partial_{y_{1}}w|^{2}dy_{1}dy' \\ &- \frac{C}{\delta_{S}}\int_{\mathbb{T}^{2}}\int_{0}^{1}\frac{|\nabla_{y'}w|^{2}}{y_{1}(1-y_{1})}dy_{1}dy' + C(\delta+\chi)\int_{\mathbb{T}^{2}}\int_{0}^{1}y_{1}(1-y_{1})|\partial_{y_{1}}w|^{2}dy_{1}dy' \\ &+ C\nu\delta_{S}\int_{\mathbb{T}^{2}}\int_{0}^{1}w^{2}\,dy_{1}dy' + C\nu\int_{\mathbb{T}^{2}}\int_{0}^{1}w^{3}\,dy_{1}dy'. \end{split}$$

We denote the average of w by \bar{w} , i.e.

$$\bar{w} := \int_{\mathbb{T}^2} \int_0^1 w \, dy_1 dy',$$

It is east to show that

$$\int_{\mathbb{T}^2} \int_0^1 |w - \bar{w}|^2 \, dy_1 dy' = \int_{\mathbb{T}^2} \int_0^1 w^2 \, dy_1 dy' - \bar{w}^2.$$

2

Taking ν, δ, χ suitably small and using the weighted Poincaré inequality (3.1), we acquire

(4.41)
$$\mathbf{B}_{1}(t) + \mathbf{B}_{2}(t) - \mathbf{G}_{1}(t) - \frac{5}{4}\mathbf{D}(t)$$
$$\leq \frac{6}{5}\alpha_{m}\delta_{S}\int_{\mathbb{T}^{2}}\int_{0}^{1}w^{2}\,dy_{1}dy' - \frac{5}{4}\alpha_{m}\delta_{S}\int_{\mathbb{T}^{2}}\int_{0}^{1}|w - \bar{w}|^{2}dy_{1}dy'$$
$$- \frac{C}{\delta_{S}}\int_{\mathbb{T}^{2}}\int_{0}^{1}\frac{|\nabla_{y'}w|^{2}}{y_{1}(1 - y_{1})}dy_{1}dy' + C\nu\int_{\mathbb{T}^{2}}\int_{0}^{1}w^{3}\,dy_{1}dy'.$$

Finally, choosing $M = \frac{5}{4} \alpha_m \sigma_m^4 v_m^2$ and combining (4.34) and (4.41), we arrive at

$$\begin{aligned} &-\frac{\delta_S}{2M} |\dot{\mathbf{X}}(t)|^2 + \mathbf{B}_1(t) + \mathbf{B}_2(t) - \mathbf{G}_1(t) - \frac{3}{4} \mathbf{D}(t) \\ &\leq \left(-\frac{\alpha_m \delta_S}{20} + C \delta_S (\nu + \delta + \chi)^2 \right) \int_{\mathbb{T}^2} \int_0^1 w^2 \, dy_1 dy' \\ &- \frac{C}{\delta_S} \int_{\mathbb{T}^2} \int_0^1 \frac{|\nabla_{y'} w|^2}{y_1(1 - y_1)} dy_1 dy' + C \nu \int_{\mathbb{T}^2} \int_0^1 w^3 \, dy_1 dy' \\ &\leq - C_0 \delta_S \int_{\mathbb{T}^2} \int_0^1 w^2 \, dy_1 dy' + C \nu \int_{\mathbb{T}^2} \int_0^1 w^3 \, dy_1 dy'. \end{aligned}$$

Notice that

$$G^{S}(t) \sim \delta_{S} \int_{\mathbb{T}^{2}} \int_{0}^{1} w^{2} \, dy_{1} dy',$$

so we obtain the desired inequality (4.23) and complete the proof of Lemma 4.3.

At last, we give the proof of Lemma 4.1. To do so, we only need to estimate the remaining terms in (4.13) individually.

Proof of Lemma 4.1: First, by (4.15) and (4.21), it holds

$$\begin{aligned} \frac{d}{dt} &\int_{\Omega} a\rho \left(Q(v|\bar{v}) + \frac{1}{2}|h - \bar{h}|^2 \right) dx = \dot{\mathbf{X}}(t)\mathbf{Y}(t) + \mathbf{B}(t) + \mathbf{S}(t) - \mathbf{G}(t) - \mathbf{D}(t) \\ &= -\frac{\delta_S}{2M} |\dot{\mathbf{X}}(t)|^2 + \mathbf{B}_1(t) + \mathbf{B}_2(t) - \mathbf{G}_1(t) - \frac{3}{4}\mathbf{D}(t) \\ &- \frac{\delta_S}{2M} |\dot{\mathbf{X}}(t)|^2 + \dot{\mathbf{X}}(t) \sum_{i=3}^{6} \mathbf{Y}_i(t) + \sum_{i=3}^{8} \mathbf{B}_i(t) + \mathbf{S}(t) - \sum_{i=2}^{3} \mathbf{G}_i(t) - \mathbf{G}^R(t) - \frac{1}{4}\mathbf{D}(t). \end{aligned}$$

Using Lemma 4.3 and Cauchy's inequality, we get

$$\frac{d}{dt} \int_{\Omega} a\rho \left(Q(v|\bar{v}) + \frac{1}{2}|h - \bar{h}|^2 \right) dx$$

$$\leq -C_0 G^S(t) + C \int_{\Omega} \partial_1 a \cdot |p - \bar{p}|^3 dx - \frac{\delta_S}{4M} |\dot{\mathbf{X}}(t)|^2 + \frac{M}{\delta_S} \sum_{i=3}^6 |\mathbf{Y}_i(t)|^2$$

+
$$\sum_{i=3}^{8} \mathbf{B}_{i}(t) + \mathbf{S}(t) - \sum_{i=2}^{3} \mathbf{G}_{i}(t) - \frac{1}{4}\mathbf{D}(t),$$

Before estimating the above bad terms individually, we notice that, by (3.21) and the Gagliardo-Nirenberg inequality (3.2), we have

$$C \int_{\Omega} \partial_{1} a \cdot |p - \bar{p}|^{3} dx \leq \frac{C}{\nu} \|p - \bar{p}\|_{L^{\infty}}^{2} \int_{\Omega} \partial_{1} v^{S} \cdot |p - \bar{p}|^{3} dx$$

$$(4.42) \qquad \leq \frac{C}{\nu} \|\nabla(p - \bar{p})\| \left(\|p - \bar{p}\| + \|\nabla^{2}(p - \bar{p})\| \right) \left(\int_{\Omega} \partial_{1} v^{S} \cdot |p - \bar{p}|^{2} dx \right)^{\frac{1}{2}} \left(\int_{\Omega} \partial_{1} v^{S} dx \right)^{\frac{1}{2}}$$

$$\leq C\chi \|\nabla(p - \bar{p})\| \sqrt{G^{S}(t)} \leq C\chi \mathbf{D}(t) + C\chi G^{S}(t).$$

• Estimates on $\mathbf{Y}_i(t)$ (i = 3, 4, 5, 6): First, by (3.21) and the wave interaction estimates in Lemma 4.4, we have

$$\begin{aligned} |\mathbf{Y}_{3}(t)| &= \left| \int_{\Omega} a\rho \partial_{1} v^{S} \cdot [p'(v^{S}) - p'(\bar{v})](v - \bar{v}) \, dx \right| \\ &\leq C \int_{\Omega} \partial_{1} v^{S} \cdot |\bar{v} - v^{S}| |\phi| \, dx \leq C \left\| \partial_{1} v^{S} (\bar{v} - v^{S}) \right\| \|\phi\| \leq C \chi \delta_{R} \delta_{S}^{3/2} e^{-C \delta_{S} t}, \end{aligned}$$

and so

$$\frac{M}{\delta_S} |\mathbf{Y}_3(t)|^2 \le C \chi^2 \delta_R^2 \delta_S^2 \cdot e^{-C \delta_S t}.$$

For $\mathbf{Y}_4(t)$, by Holder's inequality, we have

$$|\mathbf{Y}_4(t)| \le C\nu \int_{\Omega} \partial_1 a \cdot \left| h_1 - \bar{h}_1 - \frac{p - \bar{p}}{\sigma_*} \right| \, dx \le C\nu^{3/2} \sqrt{\mathbf{G}_2(t)},$$

and thus

$$\frac{M}{\delta_S} |\mathbf{Y}_4(t)|^2 \le C \nu \mathbf{G}_2(t).$$

Recall from (4.1) and (4.7) that

$$h - \bar{h} = \psi - (2\mu + \lambda)\nabla\phi - (2\mu + \lambda)\nabla v^R,$$

thus, by (3.21) and Lemma 2.2, it follows

$$\left\|h_1 - \bar{h}_1 - \frac{p - \bar{p}}{\sigma_*}\right\| \lesssim \|\phi\| + \|\nabla\phi\| + \|\psi\| + \|\partial_1 v^R\| \lesssim \chi + \delta_R.$$

Hence, we yield

$$|\mathbf{Y}_{5}(t)| \leq C \left\| \sqrt{\partial_{1} a} \right\|_{L^{\infty}} \sqrt{\mathbf{G}_{2}(t)} (\chi + \delta_{R}) \leq C \sqrt{\nu \delta_{S}} (\chi + \delta_{R}) \sqrt{\mathbf{G}_{2}(t)},$$

and so

$$\frac{M}{\delta_S} |\mathbf{Y}_5(t)|^2 \le C\nu(\chi + \delta_R)^2 \mathbf{G}_2(t).$$

For $\mathbf{Y}_6(t)$, by (3.7) and (3.21), we get

$$\begin{split} \frac{M}{\delta_S} |\mathbf{Y}_6(t)|^2 &\leq \frac{C}{\delta_S} \left(\int_{\Omega} \partial_1 a \cdot |p - \bar{p}|^2 \, dx \right)^2 + \frac{C}{\delta_S} \left(\int_{\Omega} \partial_1 a \cdot \frac{|h_2^2 + h_3^2|}{2} \, dx \right)^2 \\ &\leq \frac{C}{\delta_S^2} \left(\int_{\Omega} \partial_1 v^S \cdot |p - \bar{p}|^2 \, dx \right)^2 + C\nu \left\| h - \bar{h} \right\|^2 \mathbf{G}_3(t) \\ &\leq C \left\| p - \bar{p} \right\|^2 G^S(t) + C\nu \left\| h - \bar{h} \right\|^2 \mathbf{G}_3(t) \\ &\leq C\chi^2 G^S(t) + C\nu\chi^2 \mathbf{G}_3(t). \end{split}$$

• Estimates on $\mathbf{B}_i(t)$ (i = 3, ..., 8) : First, it follows from (4.11) that

$$F = \rho \sigma_* (v - v^S) + \rho (u_1 - u_1^S) = \rho (\sigma_* \phi + \psi_1) + [\rho \sigma_* (\bar{v} - v^S) + \rho (\bar{u}_1 - u_1^S)]$$

Thus, by assumption (3.21), we get

(4.43)
$$|F| \lesssim |\phi| + |\psi_1| + |\bar{v} - v^S| + |\bar{u}_1 - u_1^S| \lesssim \chi + \delta_R.$$

For $\mathbf{B}_3(t)$, using (3.7) and (4.43), we yield

$$\mathbf{B}_{3}(t) = \int_{\Omega} F \cdot \partial_{1} a \left(Q(v|\bar{v}) + \frac{|h_{1} - \bar{h}_{1}|^{2}}{2} + \frac{h_{2}^{2} + h_{3}^{2}}{2} \right) dx$$

$$\lesssim \int_{\Omega} |F| \cdot \partial_1 a \left(|p - \bar{p}|^2 + \left| h_1 - \bar{h}_1 - \frac{p - \bar{p}}{\sigma_*} \right|^2 + \frac{h_2^2 + h_3^2}{2} \right) dx$$

$$\lesssim \int_{\Omega} |F| \cdot \partial_1 a \cdot |p - \bar{p}|^2 dx + (\chi + \delta_R) (\mathbf{G}_2(t) + \mathbf{G}_3(t)).$$

Recall that

$$\psi_1 = h_1 - \bar{h}_1 + (2\mu + \lambda)\partial_1\phi + (2\mu + \lambda)\partial_1v^R,$$

which together with (4.43) leads to

$$\int_{\Omega} |F| \cdot \partial_1 a \cdot |p - \bar{p}|^2 dx \lesssim \sum_{i=1}^{4} \mathbf{B}_{3,i}(t),$$

where

$$\begin{split} \mathbf{B}_{3,1}(t) &:= \int_{\Omega} \partial_1 a \cdot |p - \bar{p}|^3 \, dx, \\ \mathbf{B}_{3,2}(t) &:= \int_{\Omega} \partial_1 a \cdot |h_1 - \bar{h}_1| \cdot |p - \bar{p}|^2 \, dx, \\ \mathbf{B}_{3,3}(t) &:= \int_{\Omega} \partial_1 a \cdot |\partial_1 \phi| \cdot |p - \bar{p}|^2 \, dx, \\ \mathbf{B}_{3,4}(t) &:= \int_{\Omega} \partial_1 a \cdot \left(|\bar{v} - v^S| + |\bar{u}_1 - u_1^S| + \partial_1 v^R\right) \cdot |p - \bar{p}|^2 \, dx. \end{split}$$

Essentially the same with (4.42), it holds

$$\mathbf{B}_{3,1}(t) \le C\chi \mathbf{D}(t) + C\chi G^S(t).$$

Using Lemma 2.3 and the Gagliardo-Nirenberg inequality (3.2), we yield

$$\begin{aligned} \mathbf{B}_{3,2}(t) &\leq C \int_{\Omega} \partial_{1} a \cdot \left| h_{1} - \bar{h}_{1} - \frac{p - \bar{p}}{\sigma_{*}} \right| \cdot |p - \bar{p}|^{2} \, dx + C \int_{\Omega} \partial_{1} a \cdot |p - \bar{p}|^{3} \, dx \\ &\leq C \left\| p - \bar{p} \right\|_{L^{\infty}}^{2} \sqrt{\mathbf{G}_{2}(t)} \left(\int_{\Omega} \partial_{1} a \, dx \right)^{\frac{1}{2}} + C \chi \mathbf{D}(t) + C \chi G^{S}(t) \\ &\leq C \sqrt{\nu} \sqrt{\mathbf{G}_{2}(t)} \left(\left\| p - \bar{p} \right\| + \left\| \nabla^{2}(p - \bar{p}) \right\| \right) \left\| \nabla(p - \bar{p}) \right\| + C \chi \mathbf{D}(t) + C \chi G^{S}(t) \\ &\leq C \chi \sqrt{\nu} \sqrt{\mathbf{G}_{2}(t)} \sqrt{\mathbf{D}(t)} + C \chi \mathbf{D}(t) + C \chi G^{S}(t) \\ &\leq C \chi^{2} \nu \mathbf{G}_{2}(t) + C \chi \mathbf{D}(t) + C \chi G^{S}(t). \end{aligned}$$

Noticing the fact $|\phi| \sim |p-\bar{p}|$ and that

$$\nabla(p-\bar{p}) = p'(v)\nabla(v-\bar{v}) + (p'(v) - p'(\bar{v}))\nabla\bar{v},$$

we have

(4.44)
$$|\nabla\phi| \lesssim |\nabla(p-\bar{p})| + |\partial_1 \bar{v}| \cdot |p-\bar{p}| \lesssim |\nabla(p-\bar{p})| + (\partial_1 v^R + \partial_1 v^S) \cdot |p-\bar{p}|.$$
 Using (4.44), we have

1g. (4 .44),

$$\mathbf{B}_{3,3}(t) \leq C \int_{\Omega} \partial_1 a \cdot |\nabla(p-\bar{p})| \cdot |p-\bar{p}|^2 \, dx + C \int_{\Omega} \partial_1 a \cdot (\partial_1 v^R + \partial_1 v^S) \cdot |p-\bar{p}|^3 \, dx$$
$$\leq C \chi^2 \delta_S^2 \cdot \mathbf{D}(t) + C \delta \int_{\Omega} \partial_1 a \cdot |p-\bar{p}|^3 \, dx \leq C \delta \chi \mathbf{D}(t) + C \delta \chi G^S(t).$$

Using Lemma 4.4 and assumption (3.21), we get

$$\mathbf{B}_{3,4}(t) \leq \frac{C}{\nu} \chi \left\| p - \bar{p} \right\| \left(\left\| \partial_1 v^S (\bar{v} - v^S) \right\| + \left\| \partial_1 v^S (\bar{u}_1 - u_1^S) \right\| + \left\| \partial_1 v^R \cdot \partial_1 v^S \right\| \right)$$

$$\leq C \chi^2 \delta_R \delta_S e^{-C\delta_S t}.$$

Combining the above estimates, we obtain

(4.45)
$$\mathbf{B}_{3}(t) \leq C\chi \mathbf{D}(t) + C\chi G^{S}(t) + C\chi \mathbf{G}_{2}(t) + C\chi^{2} \delta_{R} \delta_{S} e^{-C\delta_{S} t}$$

For $\mathbf{B}_4(t)$, we have

(4.46)
$$\mathbf{B}_{4}(t) \leq \frac{C}{\nu} \int_{\Omega} \partial_{1} v^{S} \cdot |p - \bar{p}| \cdot |\nabla(p - \bar{p})| \, dx$$
$$\leq \frac{C}{\nu} \sqrt{\|\partial_{1} v^{S}\|_{L^{\infty}}} \sqrt{G^{S}(t)} \sqrt{\mathbf{D}(t)} \leq C \nu G^{S}(t) + C \nu \mathbf{D}(t)$$

For $\mathbf{B}_5(t)$, by Lemma 4.4 and assumption (3.21), we yield

(4.47)

$$\mathbf{B}_{5}(t) \leq C \int_{\Omega} \partial_{1} a \cdot |p - \bar{p}|^{2} \cdot (\partial_{1} v^{R} + \partial_{1} v^{S}) dx$$

$$\leq C \nu \delta_{S} G^{S}(t) + \frac{C \chi}{\nu} \|p - \bar{p}\| \|\partial_{1} v^{R} \cdot \partial_{1} v^{S}\|$$

$$\leq C \nu \delta_{S} G^{S}(t) + C \chi^{2} \delta_{R} \delta_{S} e^{-C \delta_{S} t}.$$

For $\mathbf{B}_6(t)$, using Lemma 2.2, assumption (3.21) and the fact $p(v|\bar{v}) \sim |p-\bar{p}|^2$, we obtain

(4.48)

$$\mathbf{B}_{6}(t) \leq C \int_{\Omega} |\nabla(p-\bar{p})| \cdot |p-\bar{p}| \cdot (\partial_{1}v^{R} + \partial_{1}v^{S}) dx$$

$$\leq C \delta_{S} \sqrt{G^{S}(t)} \sqrt{\mathbf{D}(t)} + C \|\nabla(p-\bar{p})\| \left(\int_{\Omega} |\partial_{1}v^{R}|^{2} |p-\bar{p}|^{2} dx \right)^{\frac{1}{2}}$$

$$\leq C \delta_{S} \sqrt{G^{S}(t)} \sqrt{\mathbf{D}(t)} + C \sqrt{\delta_{R}} \sqrt{\mathbf{G}^{R}(t)} \sqrt{\mathbf{D}(t)}$$

$$\leq C \delta_{S} G^{S}(t) + C \left(\delta_{S} + \sqrt{\delta_{R}} \right) \mathbf{D}(t) + C \sqrt{\delta_{R}} \mathbf{G}^{R}(t).$$

For $\mathbf{B}_7(t)$, by assumption (3.21), the Gagliardo-Nirenberg inequality (3.2), (4.44) and Young's inequality, we have

(4.49)
$$\mathbf{B}_{7}(t) \leq C \|\partial_{11}v^{R}\|_{L^{1}} \|\phi\|_{L^{\infty}} \leq C \|\partial_{11}v^{R}\|_{L^{1}} \cdot \left(\|\phi\|^{\frac{1}{2}} + \|\nabla^{2}\phi\|^{\frac{1}{2}}\right) \|\nabla\phi\|^{\frac{1}{2}} \leq C\chi \|\nabla\phi\|^{2} + C\chi^{\frac{1}{3}} \|\partial_{11}v^{R}\|_{L^{1}}^{\frac{4}{3}} \leq C\chi \mathbf{D}(t) + C\chi\delta_{R}\mathbf{G}^{R}(t) + C\chi\delta_{S}^{2}G^{S}(t) + C\chi^{\frac{1}{3}} \|\partial_{11}v^{R}\|_{L^{1}}^{\frac{4}{3}}$$

To estimate $\mathbf{B}_8(t)$, we first notice that, by the definition of R in (4.3), it holds

$$R = \rho(2\mu + \lambda)(\nabla u^{t} - \operatorname{div} u \mathbb{I})\nabla v - \mu \nabla \times \nabla \times u$$
$$= \rho(2\mu + \lambda)(\nabla \psi^{t} - \operatorname{div} \psi \mathbb{I})\nabla v - \rho(2\mu + \lambda)\partial_{1}\bar{u}_{1} \begin{bmatrix} 0 & & \\ & 1 & \\ & & 1 \end{bmatrix} \nabla v - \mu \nabla \times \nabla \times u,$$

where $h' := (h_2, h_3)$ and \mathbb{I} denotes the 3×3 identity matrix. Meanwhile, it follows from (3.21) and Sobolev inequality that

(4.50)
$$\begin{aligned} \|\phi\|_{L^{3}} + \|\psi\|_{L^{3}} \lesssim \|\phi\|^{\frac{1}{2}} \|\phi\|_{L^{6}}^{\frac{1}{2}} + \|\psi\|^{\frac{1}{2}} \|\psi\|_{L^{6}}^{\frac{1}{2}} \lesssim \|\phi\|_{H^{1}} + \|\psi\|_{H^{1}} \lesssim \chi \,, \\ \|\nabla\phi\|_{L^{3}} + \|\nabla\psi\|_{L^{3}} \lesssim \|\nabla\phi\|^{\frac{1}{2}} \|\nabla\phi\|_{L^{6}}^{\frac{1}{2}} + \|\nabla\psi\|^{\frac{1}{2}} \|\nabla\psi\|_{L^{6}}^{\frac{1}{2}} \lesssim \|\nabla(\phi,\psi)\|_{H^{1}} \lesssim \chi \,. \end{aligned}$$

which together with Lemma 2.2 and the fact that $|h - \bar{h}| \lesssim |\psi| + |\nabla \phi| + |\partial_1 v^R|$ leads to (4.51) $||h - \bar{h}||_{L^3} \lesssim \chi + \delta_R.$

(4.51) $\|h - \bar{h}\|_{L^3} \lesssim \chi + \delta_R.$

For simplicity of presentation, we divide
$$\mathbf{B}_8(t)$$
 as follows:

$$\mathbf{B}_8(t) =: \sum_{i=1}^3 \mathbf{B}_{8,i}(t),$$

where

$$\begin{split} \mathbf{B}_{8,1}(t) &:= \int_{\Omega} a\rho(2\mu + \lambda)(h - \bar{h})^t (\nabla \psi^t - \operatorname{div} \psi \,\mathbb{I}) \nabla v \, dx, \\ \mathbf{B}_{8,2}(t) &:= -\int_{\Omega} a\rho(2\mu + \lambda) \partial_1 \bar{u}_1 \cdot h' \cdot \nabla_{x'} v \, dx = -\int_{\Omega} a\rho(2\mu + \lambda) \partial_1 \bar{u}_1 \cdot h' \cdot \frac{\nabla_{x'} p(v)}{p'(v)} \, dx \\ &= -\int_{\Omega} a\rho(2\mu + \lambda) \partial_1 \bar{u}_1 \cdot h' \cdot \frac{\nabla_{x'} (p - \bar{p})}{p'(v)} \, dx, \\ \mathbf{B}_{8,3}(t) &:= -\int_{\Omega} a\mu(h - \bar{h}) \cdot \nabla \times \nabla \times u \, dx. \end{split}$$

For $\mathbf{B}_{8,1}(t)$, by (3.21), (4.44), (4.51) and Sobolev inequality, we have

$$\mathbf{B}_{8,1}(t) \le C \int_{\Omega} |h - \bar{h}| \cdot |\nabla \psi| \cdot (|\nabla \phi| + |\nabla \bar{v}|) \, dx$$

$$\begin{split} &\leq C \int_{\Omega} |h - \bar{h}| \cdot |\nabla \psi| \cdot \left(|\nabla (p - \bar{p})| + 2|\partial_1 \bar{v}| \right) dx \\ &\leq C \int_{\Omega} |h - \bar{h}| \cdot |\nabla \psi| \cdot |\nabla (p - \bar{p})| \, dx + C \int_{\Omega} \partial_1 v^R \cdot |h - \bar{h}| \cdot |\nabla \psi| \, dx \\ &\quad + C \int_{\Omega} \partial_1 v^S \cdot |h - \bar{h}| \cdot |\nabla \psi| \, dx \\ &\leq C \left\| h - \bar{h} \right\|_{L^3} \|\nabla \psi\|_{L^6} \|\nabla (p - \bar{p})\| + C \left\| h - \bar{h} \right\|_{L^3} \left\| \partial_1 v^R \right\|_{L^6} \|\nabla \psi\| \\ &\quad + C \delta_S \int_{\Omega} |\nabla \psi|^2 \, dx + C \delta_S \int_{\Omega} \partial_1 v^S \cdot |h - \bar{h}|^2 \, dx \\ &\leq C (\chi + \delta) \left(\mathbf{D}(t) + \|\nabla \psi\|_{H^1}^2 + \left\| \partial_1 v^R \right\|_{L^6}^2 \right) + C \delta_S \int_{\Omega} \partial_1 v^S \cdot (h_2^2 + h_3^2) \, dx \\ &\quad + C \delta_S \int_{\Omega} \partial_1 v^S \cdot \left| h_1 - \bar{h}_1 - \frac{p - \bar{p}}{\sigma_*} \right|^2 \, dx + C \delta_S \int_{\Omega} \partial_1 v^S \cdot |p - \bar{p}|^2 \, dx \\ &\leq C (\chi + \delta) \left(\mathbf{D}(t) + \|\nabla \psi\|_{H^1}^2 + \left\| \partial_1 v^R \right\|_{L^6}^2 \right) + C \delta_S \left[\mathbf{G}_2(t) + \mathbf{G}_3(t) + G^S(t) \right]. \end{split}$$

Also, it follows from Lemma 2.2 that

$$\left\|\partial_1 v^R\right\|_{L^6}^2 \lesssim \delta_R^{\frac{1}{3}} \cdot (1+t)^{-2+\frac{1}{3}} = \delta_R^{\frac{1}{3}} \cdot (1+t)^{-\frac{5}{3}}.$$

Hence, we yield

(4.52)

$$\mathbf{B}_{8,1}(t) \le C(\chi + \delta) \left(\mathbf{D}(t) + \|\nabla\psi\|_{H^1}^2 \right) + C\delta_S \big[\mathbf{G}_2(t) + \mathbf{G}_3(t) + G^S(t) \big] \\ + C(\chi + \delta) \delta_B^{\frac{1}{3}} \cdot (1+t)^{-\frac{5}{3}}.$$

By (4.51), it holds

$$\begin{aligned} \mathbf{B}_{8,2}(t) &\leq C \int_{\Omega} \partial_1 v^S \cdot |h'| \cdot |\nabla_{x'}(p-\bar{p})| \, dx + C \int_{\Omega} \partial_1 v^R \cdot |h'| \cdot |\nabla_{x'}(p-\bar{p})| \, dx \\ &\leq C \delta_S \int_{\Omega} \partial_1 v^S \cdot (h_2^2 + h_3^2) \, dx + C \delta_S \mathbf{D}(t) + C \left\| h - \bar{h} \right\|_{L^3} \left\| \partial_1 v^R \right\|_{L^6} \left\| \nabla(p-\bar{p}) \right\| \\ &\leq C \delta_S \mathbf{G}_3(t) + C(\chi + \delta_S) \mathbf{D}(t) + C(\chi + \delta_R) \delta_R^{\frac{1}{3}} \cdot (1+t)^{-\frac{5}{3}}. \end{aligned}$$

In view of the property of curl operator, we have

$$\nabla \times u = \nabla \times \left[u - (2\mu + \lambda) \nabla v \right] = \nabla \times h,$$

and so

(4.53)

$$\nabla \times \nabla \times u = \nabla \times \nabla \times h = \nabla \times \nabla \times (h - \bar{h}).$$

Thus, it holds

$$\begin{split} \mathbf{B}_{8,3}(t) &= -\mu \int_{\Omega} a(h-\bar{h}) \cdot \nabla \times \nabla \times (h-\bar{h}) \, dx \\ &= -\mu \int_{\Omega} \nabla \times \left[a(h-\bar{h}) \right] \cdot \nabla \times (h-\bar{h}) \, dx \\ &= -\mu \int_{\Omega} a |\nabla \times (h-\bar{h})|^2 \, dx - \mu \int_{\Omega} \left[(h-\bar{h}) \times \nabla a \right] \cdot \nabla \times (h-\bar{h}) \, dx \\ &= -\mu \int_{\Omega} a |\nabla \times (h-\bar{h})|^2 \, dx - \mu \int_{\Omega} \partial_1 a(0, -h_3, h_2) \cdot \nabla \times (h-\bar{h}) \, dx \\ &\leq -\frac{\mu}{2} \int_{\Omega} a |\nabla \times (h-\bar{h})|^2 \, dx + C \int_{\Omega} |\partial_1 a|^2 \cdot (h_2^2 + h_3^2) \, dx \\ &\leq -\frac{\mu}{2} \int_{\Omega} a |\nabla \times (h-\bar{h})|^2 \, dx + C \nu \delta_S \mathbf{G}_3(t). \end{split}$$

To sum up, we obtain

$$\mathbf{B}_{8}(t) \leq C(\chi + \delta) \left(\mathbf{D}(t) + \|\nabla\psi\|_{H^{1}}^{2} \right) + C\delta_{S} \left[\mathbf{G}_{2}(t) + \mathbf{G}_{3}(t) + G^{S}(t) \right]$$
$$+ C(\chi + \delta) \delta_{R}^{\frac{1}{3}} \cdot (1+t)^{-\frac{5}{3}}.$$

• Estimates on $\mathbf{S}(t)$: By definition,

$$\mathbf{S}(t) = \int_{\Omega} a\rho \left[p'(\bar{v})(v-\bar{v}) \cdot vQ_1 - (h_1 - \bar{h}_1) \cdot vQ_2 \right].$$

Recall that

$$\begin{cases} Q_1 = \rho(u_1 - u_1^R) \cdot \partial_1 v^R - \rho(v - v^R) \cdot \partial_1 u_1^R + F \cdot \partial_1 v^S ,\\ Q_2 = \rho(u_1 - u_1^R) \cdot \partial_1 u_1^R + \rho(v - v^R) \cdot \partial_1 p^R + F \cdot \partial_1 h_1^S + \left[\partial_1 \bar{p} - \partial_1 p^R - \partial_1 p^S\right] ,\\ F = \rho \sigma_*(v - v^S) + \rho(u_1 - u_1^S) . \end{cases}$$

So we can devide vQ_1 as follows:

(4.54)
$$vQ_1 = (u_1 - u_1^R) \cdot \partial_1 v^R - (v - v^R) \cdot \partial_1 u_1^R + \left[\sigma_*(v - v^S) + (u_1 - u_1^S)\right] \cdot \partial_1 v^S$$
$$=: Q_1^R + Q_1^S + Q_1^I,$$

where

$$\begin{aligned} Q_1^R &:= \psi_1 \partial_1 v^R - \phi \partial_1 u_1^R \,, \qquad Q_1^S := (\sigma_* \phi + \psi_1) \cdot \partial_1 v^S \,, \\ Q_1^I &:= (\bar{u}_1 - u_1^R) \cdot \partial_1 v^R - (\bar{v} - v^R) \cdot \partial_1 u_1^R + \left[\sigma_* (\bar{v} - v^S) + (\bar{u}_1 - u_1^S) \right] \cdot \partial_1 v^S \,. \end{aligned}$$

Similarly, we let

(4.55)
$$vQ_{2} = (u_{1} - u_{1}^{R}) \cdot \partial_{1} u_{1}^{R} + (v - v^{R}) \cdot \partial_{1} p^{R} + \left[\sigma_{*}(v - v^{S}) + (u_{1} - u_{1}^{S})\right] \cdot \partial_{1} h_{1}^{S} + v \left[\partial_{1} \bar{p} - \partial_{1} p^{R} - \partial_{1} p^{S}\right] =: Q_{2}^{R} + Q_{2}^{S} + Q_{2}^{I},$$

where

$$Q_{2}^{R} := \psi_{1}\partial_{1}u_{1}^{R} + \phi\partial_{1}p^{R}, \qquad Q_{2}^{S} := (\sigma_{*}\phi + \psi_{1}) \cdot \partial_{1}h_{1}^{S},$$
$$Q_{2}^{I} := (\bar{u}_{1} - u_{1}^{R}) \cdot \partial_{1}u_{1}^{R} - (\bar{v} - v^{R})\partial_{1}p^{R} + [\sigma_{*}(\bar{v} - v^{S}) + (\bar{u}_{1} - u_{1}^{S})]\partial_{1}h_{1}^{S}$$
$$+ v \left[\partial_{1}\bar{p} - \partial_{1}p^{R} - \partial_{1}p^{S}\right].$$

With the aid of the above notations, we divide S(t) as follows:

$$\mathbf{S}(t) = \int_{\Omega} a\rho \left[p'(\bar{v})(v-\bar{v}) \cdot vQ_1 - (h_1 - \bar{h}_1) \cdot vQ_2 \right] dx =: \mathbf{S}_1(t) + \mathbf{S}_2(t) + \mathbf{S}_3(t),$$

where

$$\begin{aligned} \mathbf{S}_{1}(t) &:= \int_{\Omega} a\rho \left[p'(\bar{v})(v-\bar{v}) \cdot Q_{1}^{S} - (h_{1}-\bar{h}_{1}) \cdot Q_{2}^{S} \right] dx, \\ \mathbf{S}_{2}(t) &:= \int_{\Omega} a\rho \left[p'(\bar{v})(v-\bar{v}) \cdot Q_{1}^{R} - (h_{1}-\bar{h}_{1}) \cdot Q_{2}^{R} \right] dx, \\ \mathbf{S}_{3}(t) &:= \int_{\Omega} a\rho \left[p'(\bar{v})(v-\bar{v}) \cdot Q_{1}^{I} - (h_{1}-\bar{h}_{1}) \cdot Q_{2}^{I} \right] dx. \end{aligned}$$

For $\mathbf{S}_1(t)$, we have

$$\begin{aligned} \mathbf{S}_{1}(t) &= \int_{\Omega} a\rho(\sigma_{*}\phi + \psi_{1}) \left[p'(\bar{v})\phi \cdot \partial_{1}v^{S} - (h_{1} - \bar{h}_{1}) \cdot \partial_{1}h_{1}^{S} \right] dx \\ &= \int_{\Omega} a\rho \left[\sigma_{*}p'(\bar{v})\partial_{1}v^{S} \cdot \phi^{2} - \partial_{1}h_{1}^{S} \cdot (h_{1} - \bar{h}_{1})\psi_{1} \right] dx \\ &+ \int_{\Omega} a\rho\phi \left[p'(\bar{v})\partial_{1}v^{S} \cdot \psi_{1} - \sigma_{*}\partial_{1}h_{1}^{S} \cdot (h_{1} - \bar{h}_{1}) \right] dx =: \mathbf{S}_{1,1}(t) + \mathbf{S}_{1,2}(t). \end{aligned}$$

Using the fact that $\sigma_*\partial_1 h_1^S = \partial_1 p^S$, we get

$$\begin{aligned} \mathbf{S}_{1,1}(t) &= \int_{\Omega} a\rho \left[\sigma_* p'(\bar{v}) \partial_1 v^S \cdot \phi^2 - \partial_1 h_1^S \cdot (h_1 - \bar{h}_1) \psi_1 \right] dx \\ &= \int_{\Omega} a\rho \left[\sigma_* \partial_1 p^S \phi^2 - \partial_1 h_1^S \cdot |h_1 - \bar{h}_1|^2 \right] dx + \int_{\Omega} a\rho \sigma_* [p'(\bar{v}) - p'(v^S)] \partial_1 v^S \phi^2 dx \\ &+ \int_{\Omega} a\rho \partial_1 h_1^S \cdot (h_1 - \bar{h}_1) (h_1 - \bar{h}_1 - \psi_1) dx \\ &\leq \int_{\Omega} a\rho \frac{\partial_1 p^S}{\sigma_*} \left[\sigma_*^2 \phi^2 - |h_1 - \bar{h}_1|^2 \right] dx + C \delta_R G^S(t) \\ &+ C \int_{\Omega} \partial_1 v^S \cdot |h_1 - \bar{h}_1| \cdot |\partial_1 \phi| \, dx + C \int_{\Omega} \partial_1 v^S \cdot |h_1 - \bar{h}_1| \cdot \partial_1 v^R \, dx. \end{aligned}$$

In addition, by (4.31), it holds

$$\left| (\sigma_* \phi)^2 - \frac{|p - \bar{p}|^2}{\sigma_*^2} \right| \lesssim |p - \bar{p}| \cdot \left| \sigma_* \phi + \frac{p - \bar{p}}{\sigma_*} \right| \lesssim |p - \bar{p}|^3,$$

Thus, using (3.21), (4.44) and Lemma 4.4, we obtain

$$\begin{split} \mathbf{S}_{1,1}(t) &\leq C \int_{\Omega} \partial_{1} v^{S} \cdot \left| |h_{1} - \bar{h}_{1}|^{2} - \frac{|p - \bar{p}|^{2}}{\sigma_{*}^{2}} \right| dx + C \int_{\Omega} \partial_{1} v^{S} \cdot |p - \bar{p}|^{3} dx + C \delta_{R} G^{S}(t) \\ &+ C \int_{\Omega} \partial_{1} v^{S} |h_{1} - \bar{h}_{1}| \cdot (|\nabla(p - \bar{p})| + \delta|p - \bar{p}|) dx + C \int_{\Omega} \partial_{1} v^{S} |h_{1} - \bar{h}_{1}| \cdot \partial_{1} v^{R} dx \\ &\leq C \int_{\Omega} \partial_{1} v^{S} \left| h_{1} - \bar{h}_{1} - \frac{p - \bar{p}}{\sigma_{*}} \right|^{2} dx + C \int_{\Omega} \partial_{1} v^{S} |p - \bar{p}| \cdot \left| h_{1} - \bar{h}_{1} - \frac{p - \bar{p}}{\sigma_{*}} \right| dx \\ &+ C(\chi + \delta_{R}) \left(G^{S}(t) + \left\| \partial_{1} v^{S} \cdot \partial_{1} v^{R} \right\| \right) + C \int_{\Omega} \partial_{1} v^{S} |p - \bar{p}| \cdot |\nabla(p - \bar{p})| dx \\ &+ C \int_{\Omega} \partial_{1} v^{S} \left| h_{1} - \bar{h}_{1} - \frac{p - \bar{p}}{\sigma_{*}} \right| |\nabla(p - \bar{p})| dx + C \delta \int_{\Omega} \partial_{1} v^{S} \cdot |p - \bar{p}|^{2} dx \\ &\leq C \nu \mathbf{G}_{2}(t) + C \sqrt{\nu} (\mathbf{G}_{2}(t) + G^{S}(t)) + C(\chi + \delta_{R}) \left(G^{S}(t) + \left\| \partial_{1} v^{S} \cdot \partial_{1} v^{R} \right\| \right) \\ &+ C \delta_{S} \mathbf{G}_{2}(t) + C \delta_{S} \mathbf{D}(t) + C \delta G^{S}(t) \\ &\leq C(\chi + \delta) G^{S}(t) + C(\sqrt{\nu} + \delta_{S}) \mathbf{G}_{2}(t) + C \delta_{S} \mathbf{D}(t) + C(\chi + \delta_{R}) \delta_{R} \delta_{S}^{3/2} e^{-C \delta_{S} t}. \end{split}$$

Similarly, by (3.21), (4.44) and Lemma 4.4, we yield

$$\begin{aligned} \mathbf{S}_{1,2}(t) &\leq C \int_{\Omega} \partial_1 v^S \cdot |\bar{v} - v^S| |\phi| |\psi_1| \, dx + C \int_{\Omega} |\partial_1 p^S| \cdot |\phi| |h_1 - \bar{h}_1 - \psi_1| \, dx \\ &\leq C \chi \, \|\phi\| \left\| \partial_1 v^S (\bar{v} - v^S) \right\| + C \int_{\Omega} \partial_1 v^S \cdot |p - \bar{p}| (|\nabla \phi| + \partial_1 v^R) \, dx \\ &\leq C \chi \delta_R \delta_S^{3/2} e^{-C\delta_S t} + C \delta G^S(t) + C \delta_S \mathbf{D}(t). \end{aligned}$$

Hence, it holds

(4.56) $\mathbf{S}_{1}(t) \leq C(\chi + \delta)G^{S}(t) + C(\sqrt{\nu} + \delta_{S})\mathbf{G}_{2}(t) + C\delta_{S}\mathbf{D}(t) + C(\chi + \delta_{R})\delta_{R}\delta_{S}^{3/2}e^{-C\delta_{S}t}.$ For $\mathbf{S}_{2}(t)$, using (4.44), (4.50), (4.52) and assumption (3.21), we have

$$\begin{split} \mathbf{S}_{2}(t) &= \int_{\Omega} a\rho \left[p'(\bar{v})\phi \cdot Q_{1}^{R} - (h_{1} - \bar{h}_{1}) \cdot Q_{2}^{R} \right] dx \\ &= -\int_{\Omega} a\partial_{1}u_{1}^{R} \cdot \rho p'(\bar{v})\phi^{2} \, dx + \int_{\Omega} a\rho\partial_{1}v^{R} \cdot p'(\bar{v})\phi\psi_{1} \, dx - \int_{\Omega} a\rho\partial_{1}p^{R}\phi\psi_{1} \, dx \\ &- \int_{\Omega} a\rho\partial_{1}u_{1}^{R} \cdot \psi_{1}^{2} \, dx + \int_{\Omega} a\rho \left[(2\mu + \lambda)\partial_{1}\phi + \partial_{1}v^{R} \right] (\psi_{1}\partial_{1}u_{1}^{R} + \phi\partial_{1}p^{R}) \, dx \\ &\leq -\int_{\Omega} a\partial_{1}u_{1}^{R} \cdot \bar{\rho}p'(\bar{v})\phi^{2} \, dx - \int_{\Omega} a\rho\partial_{1}u_{1}^{R} \cdot \psi^{2} \, dx + C\chi \int_{\Omega} \partial_{1}u_{1}^{R} \cdot |\phi\psi_{1}| \, dx \\ &+ C\int_{\Omega} \left| \partial_{1}v^{R} \right|^{2} (|\psi| + |\phi|) \, dx + C\int_{\Omega} \partial_{1}v^{R} |\partial_{1}\bar{v}| \cdot |p - \bar{p}|(|\psi| + |\phi|) \, dx \\ &+ C\chi \mathbf{G}^{R}(t) + C\int_{\Omega} \partial_{1}v^{R} \cdot |\nabla(p - \bar{p})|(|\psi| + |\phi|) \, dx \\ &\leq -\int_{\Omega} a\partial_{1}u_{1}^{R} \cdot \bar{\rho}p'(\bar{v})\phi^{2} \, dx - \int_{\Omega} a\rho\partial_{1}u_{1}^{R} \cdot \psi^{2} \, dx + C\chi \int_{\Omega} \partial_{1}u_{1}^{R} \cdot \psi_{1}^{2} \, dx \\ &+ C\chi \left\| \partial_{1}v^{R} \right\|_{L^{4}}^{2} + C\chi \left\| \partial_{1}v^{R} \cdot \partial_{1}v^{S} \right\| + C\chi \mathbf{G}^{R}(t) + C\chi \left\| \partial_{1}v^{R} \right\|_{L^{6}} \left\| \nabla(p - \bar{p}) \right\| \\ &\leq -\int_{\Omega} a\partial_{1}u_{1}^{R} \cdot \bar{\rho}p'(\bar{v})\phi^{2} \, dx - \int_{\Omega} a\rho\partial_{1}u_{1}^{R} \cdot \psi^{2} \, dx + C\chi \int_{\Omega} \partial_{1}u_{1}^{R} \cdot \psi_{1}^{2} \, dx \\ &+ C\chi \left\| \partial_{1}v^{R} \right\|_{L^{4}}^{2} + C\chi \mathbf{G}^{R}(t) + C\chi \mathbf{D}(t) + C\chi \delta_{R} \delta_{S}^{2/3} e^{-C\delta_{S}t} + C\chi \delta_{R}^{\frac{1}{3}} \cdot (1 + t)^{-\frac{5}{3}}. \end{split}$$

Notice that

$$\mathbf{G}^{R}(t) = \int_{\Omega} a\partial_{1}u_{1}^{R} \cdot p(v|\bar{v}) \, dx \ge \int_{\Omega} a\partial_{1}u_{1}^{R} \cdot \frac{p''(\bar{v})}{2} \phi^{2} \, dx + C\chi \mathbf{G}^{R}(t),$$

and that

$$-\bar{\rho}p'(\bar{v}) - \frac{p''(\bar{v})}{2} = \gamma \bar{v}^{\gamma-2} - \frac{\gamma(\gamma+1)}{2}\bar{v}^{\gamma-2} = -\frac{\gamma-1}{\gamma+1} \cdot \frac{p''(\bar{v})}{2},$$

so it follows

$$-\int_{\Omega} a\partial_1 u_1^R \cdot \bar{\rho} p'(\bar{v}) \phi^2 \, dx - \mathbf{G}^R(t) \leq -\frac{\gamma - 1}{\gamma + 1} \mathbf{G}^R(t) + C\chi \mathbf{G}^R(t).$$

Therefore, choosing χ sufficiently small, we acquire

$$\begin{aligned} \mathbf{S}_{2}(t) - \mathbf{G}^{R}(t) &\leq -\frac{\gamma - 1}{\gamma + 1} \mathbf{G}^{R}(t) + C\chi \mathbf{G}^{R}(t) + C\chi \mathbf{D}(t) + C\chi \delta_{R} \delta_{S}^{2/3} e^{-C\delta_{S}t} \\ &+ C\chi \delta_{R}^{\frac{1}{3}} \cdot (1 + t)^{-\frac{5}{3}} + C\chi \left\| \partial_{1} v^{R} \right\|_{L^{4}}^{2}. \end{aligned}$$

For $\mathbf{S}_3(t)$, $\partial_1 \bar{p} - \partial_1 p^R - \partial_1 p^S$ is the only nontrivial term in Q_2^I . Notice that

$$\begin{aligned} |\partial_1 \bar{p} - \partial_1 p^R - \partial_1 p^S| &= |p'(\bar{v})(\partial_1 v^R + \partial_1 v^S) - p'(v^R)\partial_1 v^R - p'(v^S)\partial_1 v^S \\ &\lesssim \partial_1 v^R \cdot |\bar{v} - v^R| + \partial_1 v^S \cdot |\bar{v} - v^S|. \end{aligned}$$

According to Lemma 4.4, it holds

$$\left\|Q_{i}^{I}\right\| \lesssim \delta_{R}\delta_{S} \cdot e^{-C\delta_{S}t}, \quad i=1,2.$$

Also, by assumption (3.21), we have

$$\left\|h - \bar{h}\right\| \lesssim \|\psi\| + \|\nabla\phi\| + \left\|\partial_1 v^R\right\| \lesssim \chi + \delta_R.$$

Thus, using Lemma 4.4, we obtain

(4.57)
$$\mathbf{S}_{3}(t) \leq C \|\phi\| \|Q_{1}^{I}\| + C \|h - \bar{h}\| \|Q_{2}^{I}\| \\ \leq C(\chi + \delta_{R})\delta_{R}\delta_{S} \cdot e^{-C\delta_{S}t}.$$

Combining the above estimates, we arrive at

(4.58)
$$\mathbf{S}(t) - \mathbf{G}^{R}(t)$$
$$\leq -\frac{\gamma - 1}{\gamma + 1} \mathbf{G}^{R}(t) + C(\sqrt{\nu} + \delta_{S}) \mathbf{G}_{2}(t) + C\chi \mathbf{G}^{R}(t) + C(\chi + \delta_{S}) \mathbf{D}(t) + C(\chi + \delta) G^{S}(t)$$
$$+ C(\chi + \delta_{R}) \delta_{R} \delta_{S} \cdot e^{-C\delta_{S}t} + C(\chi + \delta) \delta_{R}^{\frac{1}{3}} \cdot (1 + t)^{-\frac{5}{3}} + C\chi \left\| \partial_{1} v^{R} \right\|_{L^{4}}^{2}.$$

Finally, using Lemma 4.2 and Lemma 4.3 and choosing $\nu, \delta_R, \delta_S, \chi$ sufficiently small, we yield

(4.59)
$$\begin{aligned} \frac{d}{dt} \int_{\Omega} a\rho \left[Q(v|\bar{v}) + \frac{|h - \bar{h}|^2}{2} \right] dx + \delta_S |\dot{\mathbf{X}}(t)|^2 \\ + \mathbf{G}_2(t) + \mathbf{G}_3(t) + \mathbf{G}^R(t) + \mathbf{D}(t) + G^S(t) \\ \lesssim (\chi + \delta) \left\| \nabla \psi \right\|_{H^1}^2 + \delta_R \delta_S \cdot e^{-C\delta_S t} + \delta_R^{\frac{1}{3}} \cdot (1 + t)^{-\frac{5}{3}} + \left\| \partial_1 v^R \right\|_{L^4}^2 + \left\| \partial_{11} v^R \right\|_{L^1}^{\frac{4}{3}}. \end{aligned}$$

According to Lemma 2.2, it holds

$$\left\|\partial_{11}v^R\right\|_{L^1} \lesssim \begin{cases} \delta_R, & 1+t \le \frac{1}{\delta_R}, \\ \frac{1}{1+t}, & 1+t \ge \frac{1}{\delta_R}, \end{cases}$$

and

$$\left\|\partial_1 v^R\right\|_{L^4} \lesssim \begin{cases} \delta_R, & 1+t \le \frac{1}{\delta_R}, \\ \frac{\delta_R^{\frac{1}{4}}}{1+t}, & 1+t \ge \frac{1}{\delta_R}. \end{cases}$$

So direct computations give

(4.60)
$$\int_{0}^{\infty} \|\partial_{11}v^{R}\|_{L^{1}}^{\frac{4}{3}} d\tau \lesssim \delta_{R}^{\frac{1}{3}}, \qquad \int_{0}^{\infty} \|\partial_{1}v^{R}\|_{L^{4}}^{2} d\tau \lesssim \delta_{R}.$$

Notice that

$$Q(v|\bar{v}) \sim |v - \bar{v}|^2,$$

and that

$$\mathbf{G}_2(t) \sim G_2(t), \quad \mathbf{G}_3(t) \sim G_3(t), \quad \mathbf{G}^R(t) \sim G^R(t), \quad \mathbf{D}(T) \sim D(t).$$

Hence, integrating (4.59) over [0, t], we acquire the desired inequality (4.13), which completes the proof of Lemma 4.1.

4.3. Wave interaction estimates. According to (4.9) and (4.11), it holds

$$vQ_1 = (u_1 - u_1^R)\partial_1 v^R - (v - v^R)\partial_1 u_1^R + \left[\sigma_*(v - v^S) + (u_1 - u_1^S)\right]\partial_1 v^S$$

=: $Q_1^R + Q_1^S + Q_1^I$,

where

$$Q_1^R := \psi_1 \partial_1 v^R - \phi \partial_1 u_1^R, \qquad Q_1^S := (\sigma_* \phi + \psi_1) \partial_1 v^S, Q_1^I := (\bar{u}_1 - u_1^R) \partial_1 v^R - (\bar{v} - v^R) \partial_1 u_1^R + \left[\sigma_* (\bar{v} - v^S) + (\bar{u}_1 - u_1^S) \right] \partial_1 v^S,$$

Note that the above Q_1^I is a nonlinear term reflecting the interaction of the approximate rarefaction and viscous shock, which can be estimated as follows.

Lemma 4.4. Under the assumptions of Proposition 3.2, there exists constant C > 0 (independent of ν, δ, χ and T) such that for all $t \in [0, T]$, it holds

(4.61)
$$\left\|\partial_1 v^R (\bar{v} - v^R)\right\| + \left\|\partial_1 v^R (\bar{u}_1 - u_1^R)\right\| \le C \delta_R \delta_S e^{-C \delta_S t}$$

(4.62)
$$\|\partial_1 v^S (\bar{v} - v^S)\| + \|\partial_1 v^S (\bar{u}_1 - u_1^S)\| + \|\partial_1 v^R \cdot \partial_1 v^S\| \le C \delta_R \delta_S^{3/2} e^{-C \delta_S t},$$

Proof. Notice that the Navier-Stokes system and Euler system are of Galilean invariance, so we can assume $u_{1m} = 0$ without loss of generality. It follows from (2.13) that $\lambda_{1m} < \sigma < u_{1m} = 0$. Since $\bar{v} = v^R + v^S - v_m$, it holds

$$\left\|\partial_{1}v^{R}(\bar{v}-v^{R})\right\|^{2} = \int_{\Omega} |\partial_{1}v^{R}|^{2} |v^{S}-v_{m}|^{2} dx = \int_{\mathbb{R}} |\partial_{1}v^{R}|^{2} |v^{S}-v_{m}|^{2} dx_{1}.$$

To estimate the above integral, we introduce two time-dependent sets:

$$E_1(t) := \left\{ x_1 \in \mathbb{R} : x_1 > \frac{\lambda_{1m} + \sigma}{2} (1+t) \right\},$$
$$E_2(t) := \left\{ x_1 \in \mathbb{R} : x_1 \le \frac{\lambda_{1m} + \sigma}{2} (1+t) \right\}.$$

First, by assumption (3.21) and Sobolev inequality, it follows

(4.63)
$$\|p - \bar{p}\|_{L^{\infty}} \le C \|v - \bar{v}\|_{L^{\infty}} \le C\chi,$$

In addition, by the definition of $\mathbf{X}(t)$ (3.15), we have

(4.64)
$$|\dot{\mathbf{X}}(t)| \leq \frac{C}{\delta_S} \|v - \bar{v}\|_{L^{\infty}} \int_{\Omega} \partial_1 v^S \, dx \leq C \|v - \bar{v}\|_{L^{\infty}} \leq C\chi, \quad t \in [0, T],$$

which together with $\mathbf{X}(0) = 0$ leads to

$$|\mathbf{X}(t)| \le C\chi t, \quad t \in [0, T].$$

According to the Lax entropy conditions (2.13), it holds $\sigma - \lambda_{1m} > 0$. Thus choosing χ sufficiently small, we have

$$|\mathbf{X}(t)| \le C\chi t < \frac{\sigma - \lambda_{1m}}{4}t, \quad t \in [0, T]$$

Hence, for each fixed $t \in [0, T]$, if $x_1 \in E_2$, then

$$x_1 - \sigma t - \mathbf{X}(t) \le \frac{\lambda_{1m} + \sigma}{2} (1+t) - \sigma t - \mathbf{X}(t)$$
$$\le \frac{\lambda_{1m} + \sigma}{2} + \frac{1}{2} (\lambda_{1m} - \sigma)t - \mathbf{X}(t) < -\frac{1}{4} (\sigma - \lambda_{1m})t < 0,$$

Also, according to Lemma 2.3 and the definition of v^{S} (2.19), it holds

(4.65)
$$|v^{S} - v_{m}| \leq C\delta_{S}e^{-C\delta_{S}|x_{1} - \sigma t - \mathbf{X}(t)|} \leq C\delta_{S}e^{\frac{-C\delta_{S}|x_{1} - \sigma t - \mathbf{X}(t)|}{2}} \cdot e^{-C\delta_{S}t}, \quad x_{1} \in E_{2},$$

Therefore, using Lemma 2.2, we obtain

(4.66)
$$\int_{E_2} |\partial_1 v^R|^2 |v^S - v_m|^2 \, dx_1 \le C \delta_S^2 e^{-2C\delta_S t} \int_{\mathbb{R}} |\partial_1 v^R|^2 \, dx_1 \le C \delta_R^2 \delta_S^2 e^{-2C\delta_S t}.$$

Secondly, for each fixed $t \in [0, T]$, if $x_1 \in E_1$, then

$$x_1 - \lambda_{1m}(1+t) = x_1 - \frac{\lambda_{1m} + \sigma}{2}(1+t) + \frac{\sigma - \lambda_{1m}}{2}(1+t) > 0.$$

Notice that

$$e^{-2|x_1-\lambda_{1m}(1+t)|} = e^{-2[x_1-\frac{\lambda_{1m}+\sigma}{2}(1+t)+\frac{\sigma-\lambda_{1m}}{2}(1+t)]} = e^{[x_1-\frac{\lambda_{1m}+\sigma}{2}(1+t)]} \cdot e^{[\frac{\sigma-\lambda_{1m}}{2}(1+t)]}$$

Thus, by Lemma 2.2, it holds

(4.67)
$$\int_{E_1} |\partial_1 v^R|^2 |v^S - v_m|^2 \, dx_1 \le C \delta_R^2 \delta_S^2 \cdot e^{\left[\frac{\sigma - \lambda_{1m}}{2}(1+t)\right]} \int_{E_1} e^{\left[x_1 - \frac{\lambda_{1m} + \sigma}{2}(1+t)\right]} \, dx_1,$$
$$\le C \delta_R^2 \delta_S^2 \cdot e^{-Ct}.$$

To sum up, we have

$$\left\|\partial_1 v^R (\bar{v} - v^R)\right\| \le C \delta_R \delta_S \cdot e^{-C \delta_S t}$$

Similarly, we can acquire the same estimate for $\|\partial_1 v^R(\bar{u}_1 - u_1^R)\|$ and then verify (4.61). Similar to the proof of (4.61), in order to prove (4.62), we only need to verify

$$\|\partial_1 v^S(\bar{v} - v^S)\| < C \delta_R \delta_S^{3/2} e^{-C \delta_S t}.$$

$$\left\|\partial_1 v^S (\bar{v} - v^S)\right\| \le C \delta_R \delta_S^{3/2} e^{-C \delta_S}$$

Notice that

$$\begin{split} \left\| \partial_1 v^S (\bar{v} - v^S) \right\|^2 &= \int_{\Omega} |\partial_1 v^R|^2 |v^R - v_m|^2 \, dx = \int_{\mathbb{R}} |\partial_1 v^S|^2 |v^R - v_m|^2 \, dx_1 \\ &= \int_{E_1} |\partial_1 v^S|^2 |v^R - v_m|^2 \, dx_1 + \int_{E_2} |\partial_1 v^S|^2 |v^R - v_m|^2 \, dx_1. \end{split}$$

Thus for each fixed $t \in [0, T]$, if $x_1 \in E_2$, then by Lemma 2.3 and the definition of v^S (2.19), it follows

(4.68)
$$\begin{aligned} |\partial_1 v^S| &\leq C \delta_S^2 \cdot e^{-C\delta_S |x_1 - \sigma t - \mathbf{X}(t)|} \\ &\leq C \delta_\sigma^2 \cdot e^{\frac{-C\delta_S |x_1 - \sigma t - \mathbf{X}(t)|}{2}} \cdot e^{-C\delta_S t} \end{aligned}$$

$$\leq C\delta_S^2 \cdot e^{\frac{-C\delta_S|x_1 - \sigma t - \mathbf{X}(t)|}{2}} \cdot e^{-C\delta_S t}, \quad x_1 \in E_2.$$

Hence,

$$\begin{split} \int_{E_2} |\partial_1 v^S|^2 |v^R - v_m|^2 \, dx_1 &\leq C \delta_R^2 \delta_S^2 \cdot e^{-C\delta_S t} \int_{E_2} \partial_1 v^S \, dx_1 \\ &\leq C \delta_R^2 \delta_S^3 \cdot e^{-C\delta_S t}. \end{split}$$

Meanwhile,

$$\int_{E_1} |\partial_1 v^S|^2 |v^R - v_m|^2 \, dx_1 \le C \delta_R^2 \delta_S^4 \cdot e^{\left[\frac{\sigma - \lambda_{1m}}{2} (1+t)\right]} \int_{E_1} e^{\left[x_1 - \frac{\lambda_{1m} + \sigma}{2} (1+t)\right]} \, dx_1,$$
$$\le C \delta_R^2 \delta_S^4 \cdot e^{-Ct}.$$

Therefore, we have

$$\left\|\partial_1 v^S (\bar{v} - v^S)\right\| \le C \delta_R \delta_S^{3/2} e^{-C \delta_S t}$$

This completes the proof of Lemma 4.4.

4.4. Estimates on $||u - \bar{u}||$ and $||v - \bar{v}||_{H^1}$. From now on, we will consider the energy estimates with respect to $(v - \bar{v}, u - \bar{u})$. In this subsection, we first give the zero-th order estimates.

Lemma 4.5. Under the assumptions of Proposition 3.2, there exists constant C > 0 (independent of ν, δ, χ and T) such that for all $t \in [0, T]$, it holds

$$(4.69) \qquad \|(v-\bar{v})(t)\|_{H^{1}}^{2} + \|(u-\bar{u})(t)\|^{2} + \delta_{S} \int_{0}^{t} |\dot{\mathbf{X}}(\tau)|^{2} d\tau + \int_{0}^{t} G_{2}(\tau) + G_{3}(\tau) + \mathcal{G}^{R}(\tau) + G^{S}(\tau) + D(\tau) + \|\nabla(u-\bar{u})\|^{2} d\tau \leq C \left(\|v_{0}-\bar{v}(0,\cdot)\|_{H^{1}}^{2} + \|u_{0}-\bar{u}(0,\cdot)\|^{2}\right) + C(\delta+\chi) \int_{0}^{t} \|\nabla^{2}(u-\bar{u})\|^{2} d\tau + C\delta_{R}^{\frac{1}{3}},$$

where $G^{S}(t), G^{R}(t), D(t)$ are as in (4.14) and

$$\mathcal{G}^R(t) := \frac{\gamma - 1}{\gamma + 1} G^R(t) + \int_{\Omega} \rho \partial_1 u_1^R \cdot \psi_1^2 \, dx = \int_{\Omega} \frac{\gamma - 1}{\gamma + 1} \partial_1 u_1^R \cdot p(v|\bar{v}) + \rho \partial_1 u_1^R \cdot \psi_1^2 \, dx.$$

Proof. Since $\bar{v} = v^S + v^R - v_m$, $\bar{u} = u^S - u^R - u_m$, direct computations give

(4.70)
$$\begin{cases} \rho \partial_t \bar{v} + \rho u \cdot \nabla \bar{v} - \operatorname{div} \bar{u} + \rho \dot{\mathbf{X}}(t) \partial_1 v^S = Q_1, \\ \rho \partial_t \bar{u} + \rho u \cdot \nabla \bar{u} + \nabla \bar{p} + \rho \dot{\mathbf{X}}(t) \partial_1 u^S = (2\mu + \lambda) \Delta (\bar{u} - u^R) + Q_3 \cdot e_1. \end{cases}$$

where Q_1 is as in (4.9) and $e_1 = (1, 0, 0)^t$. Q_3 is defined as follows:

(4.71)
$$Q_3 := \rho(v - v^R)\partial_1 p^R + \rho(u_1 - u_1^R)\partial_1 u_1^R + F \cdot \partial_1 u_1^S + \partial_1 (\bar{p} - p^S - p^R),$$

where F is as defined in (4.11) . Subtracting (4.70) from (1.3), we acquire the perturbed system for $(\phi, \psi) = (v - \bar{v}, u - \bar{u})$ which read as

(4.72)
$$\begin{cases} \rho \partial_t \phi + \rho u \cdot \nabla \phi - \operatorname{div} \psi - \rho \dot{\mathbf{X}}(t) \partial_1 v^S = -Q_1, \\ \rho \partial_t \psi + \rho u \cdot \nabla \psi + \nabla (p - \bar{p}) - \rho \dot{\mathbf{X}}(t) \partial_1 u^S = \mu \Delta \psi + (\mu + \lambda) \nabla \operatorname{div} \psi \\ + (2\mu + \lambda) \Delta u^R - Q_3 \cdot e_1. \end{cases}$$

Multiplying $(4.72)_1$ by $-(p-\bar{p})$, we have

(4.73)

$$\begin{aligned}
\partial_t \left[\rho Q(v|\bar{v}) \right] + \operatorname{div} \left[\rho Q(v|\bar{v})u \right] + (p - \bar{p}) \operatorname{div}\psi \\
&= -\rho \dot{\mathbf{X}}(t) \partial_1 v^S \cdot (p - \bar{p}) - p(v|\bar{v}) \left[\rho \partial_t \bar{v} + \rho u \cdot \nabla \bar{v} \right] + Q_1 (p - \bar{p}) \\
&= -\rho \dot{\mathbf{X}}(t) \partial_1 v^S \cdot p'(\bar{v}) (v - \bar{v}) - p(v|\bar{v}) \cdot \operatorname{div}\bar{u} + p'(\bar{v}) (v - \bar{v}) Q_1.
\end{aligned}$$

Multiplying $(4.72)_2$ by $\psi = u - \bar{u}$, we have

(4.74)
$$\partial_t \left[\rho \frac{|\psi|^2}{2} \right] + \operatorname{div} \left[\rho \frac{|\psi|^2}{2} u \right] + \operatorname{div} \left[(p - \bar{p}) \psi \right] - (p - \bar{p}) \operatorname{div} \psi$$
$$= \rho \dot{\mathbf{X}}(t) \partial_1 u_1^S \cdot \psi_1 + (2\mu + \lambda) \partial_{11} u_1^R \cdot \psi_1 - Q_3 \cdot \psi_1 + \mu \psi \Delta \psi + (\mu + \lambda) \nabla \operatorname{div} \psi.$$

Adding (4.73) and (4.74) together and integrating the resultant equation over Ω , we yield

(4.75)
$$\frac{d}{dt} \int_{\Omega} \rho Q(v|\bar{v}) + \rho \frac{|\psi|^2}{2} dx + G^R(t) + D_1(t) = \dot{\mathbf{X}}(t) \mathcal{Y}(t) + \sum_{i=1}^3 I_i(t),$$

where

$$\begin{split} D_1(t) &:= \int_{\Omega} \mu |\nabla \psi|^2 + (\mu + \lambda) |\operatorname{div} \psi|^2 \, dx, \\ \mathcal{Y}(t) &:= -\int_{\Omega} \rho \partial_1 v^S \cdot p'(\bar{v}) \phi \, dx + \int_{\Omega} \rho \partial_1 u_1^S \cdot \psi_1 \, dx =: \mathcal{Y}_1(t) + \mathcal{Y}_2(t), \\ I_1(t) &:= \int_{\Omega} -\partial_1 u_1^S \cdot p(v|\bar{v}) \, dx, \\ I_2(t) &:= \int_{\Omega} p'(\bar{v}) \phi \cdot Q_1 - Q_3 \cdot \psi_1 \, dx, \\ I_3(t) &:= \int_{\Omega} (2\mu + \lambda) \partial_{11} u_1^R \cdot \psi_1 \, dx. \end{split}$$

For $I_1(t)$, it is easy to see that

$$I_1(t) \le CG^S(t).$$

The treatment for $I_2(t)$ is similar to that for $\mathbf{S}(t)$. We first divide vQ_3 as follows, where Q_3 is defined in (4.71).

(4.76)
$$vQ_3 = (u_1 - u_1^R) \cdot \partial_1 u_1^R + (v - v^R) \cdot \partial_1 p^R + \left[\sigma_*(v - v^S) + (u_1 - u_1^S)\right] \cdot \partial_1 u_1^S \\ + v \left[\partial_1 \bar{p} - \partial_1 p^R - \partial_1 p^S\right] =: Q_3^R + Q_3^S + Q_3^I,$$

where

$$\begin{aligned} Q_3^R &:= \psi_1 \partial_1 u_1^R + \phi \partial_1 p^R \,, \qquad Q_3^S := (\sigma_* \phi + \psi_1) \cdot \partial_1 u_1^S \,, \\ Q_3^I &:= (\bar{u}_1 - u_1^R) \cdot \partial_1 u_1^R - (\bar{v} - v^R) \partial_1 p^R + \left[\sigma_* (\bar{v} - v^S) + (\bar{u}_1 - u_1^S) \right] \partial_1 u_1^S \\ &+ v \left[\partial_1 \bar{p} - \partial_1 p^R - \partial_1 p^S \right] . \end{aligned}$$

Thus,

$$I_2(t) =: \sum_{i=1}^3 I_{2,i}(t) = \int_{\Omega} \rho \left[p'(\bar{v})\phi \cdot Q_1^S - \psi_1 \cdot Q_3^S \right] dx$$
$$+ \int_{\Omega} \rho \left[p'(\bar{v})\phi \cdot Q_1^R - \psi_1 \cdot Q_3^R \right] dx$$
$$+ \int_{\Omega} \rho \left[p'(\bar{v})\phi \cdot Q_1^I - \psi_1 \cdot Q_3^I \right] dx.$$

For $I_{2,1}(t)$, by Cauchy's inequality, it holds

$$I_{2,1}(t) \lesssim \int_{\Omega} \partial_1 v^S(\phi^2 + \psi_1^2) \, dx \lesssim G^S(t) + \int_{\Omega} \partial_1 v^S \cdot \psi_1^2 \, dx.$$

Notice that

$$\begin{aligned} |\psi_1| \lesssim \left| h_1 - \bar{h}_1 - \frac{p - \bar{p}}{\sigma_*} \right| + |p - \bar{p}| + |\nabla \phi| + \partial_1 v^R \\ \lesssim \left| h_1 - \bar{h}_1 - \frac{p - \bar{p}}{\sigma_*} \right| + |p - \bar{p}| + |\nabla (p - \bar{p})| + \partial_1 v^R \end{aligned}$$

which together with Lemma 4.4 leads to

$$I_{2,1}(t) \le CG^{S}(t) + C\nu G_{2}(t) + C\delta_{S}D(t) + C\chi^{2}\delta_{R}\delta_{S}^{\frac{3}{2}}e^{-C\delta_{S}t}.$$

Similar to the estimate for $\mathbf{S}_2(t)$, we have

$$\begin{split} I_{2,2}(t) - G^R(t) &= \int_{\Omega} \rho \left[p'(\bar{v})\phi \cdot Q_1^R - \psi_1 \cdot Q_3^R \right] - \partial_1 u_1^R \cdot p(v|\bar{v}) \, dx \\ &= \int_{\Omega} \rho \left[-p'(\bar{v})\partial_1 u_1^R \cdot \phi^2 + p'(\bar{v})\partial_1 v^R \cdot \phi \psi_1 - \partial_1 p^R \cdot \phi \psi_1 - \partial_1 u_1^R \cdot \psi_1^2 \right] - \partial_1 u_1^R \cdot p(v|\bar{v}) \, dx \\ &\leq -\frac{\gamma - 1}{\gamma + 1} G^R(t) - \int_{\Omega} \rho \partial_1 u_1^R \cdot \psi_1^2 \, dx + C(\chi + \delta) G^R(t) + C\chi^2 \left\| \partial_1 v^R(\bar{v} - v^R) \right\| \\ &\leq -\mathcal{G}^R(t) + C(\chi + \delta) G^R(t) + C\chi^2 \delta_R \delta_R \cdot e^{-C\delta_S t}. \end{split}$$

The estimate for $I_{2,3}(t)$ is exactly the same with that for $\mathbf{S}_3(t)$, which reads as

$$I_{2,3}(t) \le C(\chi + \delta_R)\delta_R\delta_S \cdot e^{-C\delta_S t}.$$

Hence, we obtain

(4.77)
$$I_2(t) - G^R(t) \le -\mathcal{G}^R(t) + C(\chi + \delta)G^R(t) + C\delta_R\delta_R \cdot e^{-C\delta_S t} + CG^S(t) + C\nu G_2(t) + C\delta_S D(t).$$

For $I_3(t)$, by assumption (3.21), the Gagliardo-Nirenberg inequality (3.2) and (4.44), we have

(4.78)
$$I_{3}(t) \leq C \left\| \partial_{11} u_{1}^{R} \right\|_{L^{1}} \|\psi_{1}\|_{L^{\infty}} \leq C \left\| \partial_{11} u_{1}^{R} \right\|_{L^{1}} \cdot \left(\|\psi_{1}\|^{\frac{1}{2}} + \left\| \nabla^{2} \psi_{1} \right\|^{\frac{1}{2}} \right) \left\| \nabla \psi_{1} \right\|^{\frac{1}{2}} \leq C \chi D_{1}(t) + C \chi^{\frac{1}{3}} \left\| \partial_{11} v^{R} \right\|_{L^{1}}^{\frac{4}{3}}.$$

Since $|p - \bar{p}| \sim |\phi|$, it follows from Holder inequality that

$$\mathcal{Y}_1(t)| \le \left(\int_{\Omega} \partial_1 v^S \, dx\right)^{\frac{1}{2}} \left(\int_{\Omega} \partial_1 v^S \cdot \phi^2 \, dx\right) \le C\nu \sqrt{G^S(t)}.$$

Recall that

$$\phi_1 = h_1 - \bar{h}_1 + (2\mu + \lambda)\partial_1\phi + (2\mu + \lambda)\partial_1v^R,$$

and that

$$|\nabla \phi| \lesssim |\nabla (p - \bar{p})| + |\partial_1 \bar{v}| |p - \bar{p}|.$$

Thus, using Lemma 4.4, we have

$$\begin{aligned} |\mathcal{Y}_2(t)| &\leq C \int_{\Omega} \partial_1 v^S \left(\left| h_1 - \bar{h}_1 - \frac{p - \bar{p}}{\sigma_*} \right| + |p - \bar{p}| + |\nabla(p - \bar{p})| + \partial_1 v^R \right) dx \\ &\leq C \nu^{\frac{3}{4}} \sqrt{G_2(t)} + C \nu \sqrt{G^S(t)} + \delta_S \sqrt{D(t)} + C \delta_R \delta_S \cdot e^{-C \delta_S t}. \end{aligned}$$

Therefore, we obtain

$$\begin{aligned} \dot{\mathbf{X}}(t)\mathcal{Y}(t) &\leq |\dot{\mathbf{X}}(t)| \left(|\mathcal{Y}_{1}(t)| + |\mathcal{Y}_{1}(t)|\right) \\ &\leq \frac{\delta_{S}}{4} |\dot{\mathbf{X}}(t)|^{2} + C\nu G_{2}(t) + C_{1}G^{S}(t) + C\delta_{S}D(t) + C\delta_{R}^{2}\delta_{S} \cdot e^{-C\delta_{S}t}. \end{aligned}$$

Integrating (4.75) over [0, t] for any $t \in [0, T]$, using the above estimates and (4.60), we can find some constant $C_2 > 0$ independent of t such that

$$\int_{\Omega} \rho Q(v|\bar{v}) + \frac{\rho|\psi|^2}{2} dx + \int_0^t \frac{\gamma - 1}{\gamma + 1} G^R(\tau) + D_1(\tau) d\tau$$

$$\leq C \int_{\Omega} |v_0 - \bar{v}(0, x)|^2 + |u_0 - \bar{u}(0, x)|^2 dx + \int_0^t \frac{\delta_S}{4} |\dot{\mathbf{X}}(\tau)|^2 + C_1 G^S(\tau) d\tau + \int_0^t C \nu G_2(\tau) + C \delta_S D(\tau) + C(\chi + \delta) G^R(\tau) + C \chi D_1(\tau) d\tau + C \delta_R^{\frac{1}{3}}.$$

Multiplying the above inequality by $\frac{1}{2 \max\{1, C_2\}}$, adding the resultant expression to (4.13), and choosing ν, χ, δ sufficiently small, we yield

$$\begin{aligned} \|(v-\bar{v})(t)\|^{2} + \|(h-\bar{h})(t)\|^{2} + \|(u-\bar{u})(t)\|^{2} + \delta_{S} \int_{0}^{t} \dot{\mathbf{X}}(\tau) d\tau \\ + \int_{0}^{t} G_{2}(\tau) + G_{3}(\tau) + G^{R}(\tau) + G^{S}(\tau) + D(\tau) + D_{1}(t) d\tau \\ \leq C \left(\|v_{0} - \bar{v}(0, \cdot)\|^{2} + \|h_{0} - \bar{h}(0, \cdot)\|^{2} + \|u_{0} - \bar{u}(0, \cdot)\|^{2} \right) \\ + C(\chi + \delta) \int_{0}^{t} \|\nabla^{2}(u-\bar{u})\| d\tau + C\delta_{R}^{\frac{1}{3}}. \end{aligned}$$

Finally, since

$$\begin{aligned} |\nabla(v-\bar{v})| &\lesssim |h-\bar{h}| + |u-\bar{u}| + \partial_1 v^R, \\ |h_0 - \bar{h}_0(0,\cdot)| &\lesssim |u_0 - \bar{u}_0(0,\cdot)| + |v_0 - \bar{v}_0(0,\cdot)| + \partial_1 v^R, \end{aligned}$$

and

$$D_1(t) \sim \left\|\nabla(u - \bar{u})\right\|^2$$

we can acquire the desired inequality (4.69), which completes the proof of Lemma 4.5.

4.5. Estimates on $\|\nabla(u-\bar{u})\|$.

Lemma 4.6. Under the assumptions of Proposition 3.2, there exists constant C > 0 (independent of ν, δ, χ and T) such that for all $t \in [0, T]$, it holds

(4.79)
$$\begin{aligned} \|\nabla(u-\bar{u})(t)\|^{2} + \int_{0}^{t} \|\nabla^{2}(u-\bar{u})\|^{2} d\tau \\ \leq C\left(\|v_{0}-\bar{v}(0,\cdot)\|^{2}_{H^{1}} + \|u_{0}-\bar{u}(0,\cdot)\|^{2}_{H^{1}}\right) + C\delta_{R}^{\frac{1}{3}}. \end{aligned}$$

Proof. Multiplying $(4.72)_2$ by $-v\Delta\psi$, and integrating the resultant equation over Ω , it follows

$$\frac{d}{dt} \int_{\Omega} \frac{|\nabla \psi|^2}{2} \, dx + D_2(t) = \sum_{i=1}^5 J_i(t),$$

where

$$D_{2}(t) := \int_{\Omega} \mu v |\Delta \psi|^{2} + (\mu + \lambda) v |\nabla \operatorname{div} \psi|^{2} dx,$$

$$J_{1}(t) := \int_{\Omega} \operatorname{div} u \frac{|\nabla \psi|^{2}}{2} - (\nabla \psi \cdot \nabla u) : \nabla \psi dx,$$

$$J_{2}(t) := -\dot{\mathbf{X}}(t) \int_{\Omega} \partial_{1} u_{1}^{S} \cdot \Delta \psi dx,$$

$$J_{3}(t) := (\mu + \lambda) \int_{\Omega} \operatorname{div} \psi (\Delta \psi - \nabla \operatorname{div} \psi) \cdot \nabla v dx,$$

$$J_{4}(t) := \int_{\Omega} v \Delta \psi \cdot \nabla (p - \bar{p}) dx,$$

$$J_{5}(t) := \int_{\Omega} v Q_{3} \cdot \Delta \psi_{1} dx.$$

Using assumption (3.21), Sobolev inequality and the L^2 theory for elliptic equations, we have

$$J_{1}(t) \leq C \|\nabla\psi\|_{L^{3}} \|\nabla\psi\|_{L^{6}} \|\nabla\psi\| + C\delta \|\nabla\psi\|^{2}$$

$$\leq C\chi \|\nabla\psi\|_{H^{1}} \|\nabla\psi\| + C\delta \|\nabla\psi\|^{2}$$

$$\leq C\chi D_{2}(t) + C(\chi + \delta) \|\nabla\psi\|^{2}.$$

By Holder inequality and Lemma 2.3, we yield

$$J_2(t) \le C |\dot{\mathbf{X}}(t)| \cdot \left\| \partial_1 u_1^S \right\| \sqrt{D_2(t)} \le C \delta_S^{\frac{3}{2}} |\dot{\mathbf{X}}(t)| \sqrt{D_2(t)} \le C \delta_S^2 |\dot{\mathbf{X}}(t)|^2 + C \delta_S D_2(t).$$

Using assumption (3.21) and the fact that $\|\nabla \phi\|_{L^3} \lesssim \|\nabla \phi\|_{H^1} \lesssim \chi$, we obtain

$$J_{3}(t) \leq C\delta \|\Delta\psi\| \|\operatorname{div}\psi\| + C \|\Delta\psi\| \|\nabla\phi\|_{L^{3}} \|\operatorname{div}\psi\|_{L^{6}}$$
$$\leq C\delta\sqrt{D_{2}(t)} \|\nabla\psi\| + C\chi\sqrt{D_{2}(t)} \|\operatorname{div}\psi\|_{H^{1}}$$
$$\leq C(\chi + \delta)D_{2}(t) + C(\chi + \delta) \|\nabla\psi\|^{2}.$$

It follows from Cauchy's inequality that

$$J_4(t) \le \frac{1}{4}D_2(t) + CD(t).$$

Using (4.76), we decompose $J_5(t)$ as follows:

$$J_{5}(t) = \int_{\Omega} Q_{3}^{S} \cdot \Delta \psi_{1} \, dx + \int_{\Omega} Q_{3}^{R} \cdot \Delta \psi_{1} \, dx + \int_{\Omega} Q_{3}^{I} \cdot \Delta \psi_{1} \, dx =: \sum_{i=1}^{3} J_{5,i}(t).$$

According to Lemma 2.3, it holds

$$J_{5,1}(t) \leq C \int_{\Omega} \partial_1 v^S \cdot \Delta \psi_1 \left(\left| h_1 - \bar{h}_1 - \frac{p - \bar{p}}{\sigma_*} \right| + |p - \bar{p}| + |\nabla(p - \bar{p})| \right) dx$$

$$\leq C \left(\delta_S \sqrt{G_3(t)} + \delta_S \sqrt{G^S(t)} + \delta_S^2 \sqrt{D(t)} \right) \sqrt{D_2(t)}$$

$$\leq C \delta_S G_3(t) + C \delta_S G^S(t) + C \delta_S D(t) + C \delta_S D_2(t),$$

Using Lemma 2.2 and Cauchy's inequality, we have

$$J_{5,2}(t) \le C\sqrt{\delta_R}\mathcal{G}^R(t) + C\sqrt{\delta_R}D_2(t).$$

By assumption (3.21) and Lemma 4.4, we get

$$J_{5,3}(t) \le C \left\| Q_3^I \right\| \left\| \Delta \psi_1 \right\| \le C \chi \delta_R \delta_S \cdot e^{-C \delta_S t},$$

Combining the above estimates, we have

$$\frac{d}{dt} \left\| \nabla \psi \right\|^2 + D_2(t) \le C\delta_S^2 |\dot{\mathbf{X}}(t)|^2 + C\delta_S G_3(t) + C\delta_S G^S(t) + CD(t) + C(\chi + \delta) \left\| \nabla \psi \right\|^2 + C\sqrt{\delta_R} \mathcal{G}^R(t) + C\chi \delta_R \delta_S \cdot e^{-C\delta_S t}.$$

Finally, integrating the above inequality over [0, t], using Lemma 4.5 and taking $\chi, \delta, \delta_R, \delta_S$ sufficiently small, we can obtain the desired inequality (4.79), where we use the fact

$$D_2(t) \sim \left\|\nabla^2 \psi\right\|^2.$$

The proof of Lemma 4.6 is completed.

4.6. Estimates on $\|\nabla^2(v-\bar{v})\|$.

Lemma 4.7. Under the assumptions of Proposition 3.2, there exists constant C > 0 (independent of ν, δ, χ and T) such that for all $t \in [0, T]$, it holds

(4.80)
$$\begin{aligned} \left\| \nabla^2 (v - \bar{v})(t) \right\|^2 + \int_0^t \left\| \nabla^2 (v - \bar{v}) \right\|^2 d\tau \\ &\leq C \left(\left\| v_0 - \bar{v}(0, \cdot) \right\|_{H^2}^2 + \left\| u_0 - \bar{u}(0, \cdot) \right\|_{H^1}^2 \right) + C(\chi + \delta) \int_0^t \left\| \nabla^3 (u - \bar{u}) \right\|^2 d\tau + C \delta_R^{\frac{1}{3}}. \end{aligned}$$

Proof. We first deform the perturbed system (4.72) as follows:

(4.81)
$$\begin{cases} \partial_t \phi + u \cdot \nabla \phi - v \operatorname{div} \psi - \dot{\mathbf{X}}(t) \partial_1 v^S = -vQ_1, \\ \partial_t \psi + u \cdot \nabla \psi + v p'(v) \nabla \phi + v (p'(v) - p'(\bar{v})) \nabla \bar{v} - \dot{\mathbf{X}}(t) \partial_1 u^S \\ = \mu v \Delta \psi + (\mu + \lambda) v \nabla \operatorname{div} \psi + (2\mu + \lambda) v \Delta u^R - vQ_3 \cdot e_1. \end{cases}$$

Applying ∂_{ij} (i, j = 1, 2, 3) to $(4.81)_1$ leads to

(4.82)
$$\frac{\partial_t \partial_{ij} \phi + u \cdot \nabla \partial_{ij} \phi + \partial_i u \cdot \nabla \partial_j u + \partial_j u \cdot \nabla \partial_i \phi + \partial_{ij} u \cdot \nabla \phi - \dot{\mathbf{X}}(t) \partial_{1ij} v^S}{\partial_{ij} v \cdot \operatorname{div} \psi + \partial_i v \cdot \partial_j \operatorname{div} \psi + \partial_j v \cdot \partial_i \operatorname{div} \psi + v \partial_{ij} \operatorname{div} \psi - \partial_{ij} (vQ_1). }$$

Multiplying (4.82) by $\rho(2\mu + \lambda)\partial_{ij}\phi$ and integrating the resultant equation over Ω , we have

$$(4.83) \qquad (2\mu+\lambda)\frac{d}{dt}\int_{\Omega}\rho\frac{|\nabla^{2}\phi|^{2}}{2}\,dx - (2\mu+\lambda)\int_{\Omega}\nabla\partial_{i}\phi\cdot\nabla\partial_{i}\mathrm{div}\psi\,dx \\ = -(2\mu+\lambda)\int_{\Omega}2\rho\partial_{ij}\phi\cdot\partial_{i}u\cdot\nabla\partial_{j}\phi\,dx - (2\mu+\lambda)\int_{\Omega}\rho\partial_{ij}\phi\cdot\partial_{ij}u\cdot\nabla\phi\,dx \\ + (2\mu+\lambda)\int_{\Omega}\rho\partial_{ij}\phi\cdot\partial_{ij}v\cdot\mathrm{div}\psi\,dx + (2\mu+\lambda)\int_{\Omega}2\rho\partial_{ij}\phi\cdot\partial_{i}v\cdot\partial_{j}\mathrm{div}\psi\,dx \\ + (2\mu+\lambda)\dot{\mathbf{X}}(t)\int_{\Omega}\rho\partial_{111}v^{S}\cdot\partial_{11}\phi\,dx - (2\mu+\lambda)\int_{\Omega}\rho\partial_{ij}\phi\cdot\partial_{ij}(vQ_{1})\,dx. \end{cases}$$

Note that throughout this paper we obey the Einstein summation convention, i.e. repeated indices are implicitly summed from 1 to 3.

Applying ∂_i (i = 1, 2, 3) to $(4.81)_2$, it follows

$$(4.84) \qquad \begin{aligned} \partial_{t}\partial_{i}\psi + u\cdot\nabla\partial_{i}\psi + \partial_{i}u\cdot\nabla\psi + vp'(v)\nabla\partial_{i}\phi + \partial_{i}(vp'(v))\nabla\phi - \dot{\mathbf{X}}(t)\partial_{1i}u_{1}^{S} \\ &= -v(p'(v) - p'(\bar{v}))\nabla\partial_{i}\bar{v} - \partial_{i}\left[v(p'(v) - p'(\bar{v}))\right]\nabla\bar{v} + \mu v\Delta\partial_{i}\psi + (\mu + \lambda)v\nabla\partial_{i}\operatorname{div}\psi \\ &+ (2\mu + \lambda)v\Delta\partial_{i}u^{R} - \partial_{i}v\left(\mu\Delta\psi + (\mu + \lambda)\nabla\operatorname{div}\psi + (2\mu + \lambda)\Delta u^{R}\right) - \partial_{i}(vQ_{3})\cdot e_{1}. \end{aligned}$$

Multiplying (4.84) by $-\rho \nabla \partial_i \phi$ and integrating the resultant equation over Ω , we have

$$(4.85) \qquad \begin{aligned} & \int_{\Omega} -p'(v) |\nabla^{2}\phi|^{2} dx + (2\mu + \lambda) \int_{\Omega} \nabla \partial_{i}\phi \cdot \nabla \partial_{i} \operatorname{div}\psi dx \\ &= \frac{d}{dt} \int_{\Omega} \rho \partial_{i}\psi \cdot \nabla \partial_{i}\phi dx - \int_{\Omega} \rho \partial_{i}\psi_{j} [\partial_{t}\partial_{ij}\phi + u \cdot \nabla \partial_{ij}\phi] dx + \int_{\Omega} \rho (\partial_{i}u \cdot \nabla \psi) \cdot \nabla \partial_{i}\phi dx \\ &+ \int_{\Omega} \rho \partial_{i}(vp'(v)) \nabla \phi \cdot \nabla \partial_{i}\phi dx + \int_{\Omega} \rho \partial_{i} [v(p'(v) - p'(\bar{v}))] \partial_{1}\bar{v} \cdot \partial_{1i}\phi dx \\ &+ \int_{\Omega} (p'(v) - p'(\bar{v})) \partial_{11}\bar{v} \cdot \partial_{11}\phi dx - \dot{\mathbf{X}}(t) \int_{\Omega} \rho \partial_{11}u_{1}^{S} \cdot \partial_{11}\phi dx + \int_{\Omega} \rho \partial_{i}(vQ_{3}) \partial_{1i}\phi dx \\ &- (2\mu + \lambda) \int_{\Omega} \partial_{111}u_{1}^{R} \cdot \partial_{11}\phi dx - (2\mu + \lambda) \int_{\Omega} \rho \partial_{i}v \cdot \partial_{11}u_{1}^{R} \cdot \partial_{11}\phi dx \\ &- (2\mu + \lambda) \int_{\Omega} \rho \partial_{i}v(\mu\Delta\psi + (\mu + \lambda)\nabla \operatorname{div}\psi) \cdot \nabla \partial_{i}\phi dx \,. \end{aligned}$$

Adding (4.83) and (4.85) together, we arrive at

(4.86)
$$(2\mu + \lambda)\frac{d}{dt} \int_{\Omega} \rho \frac{|\nabla^2 \phi|^2}{2} \, dx + D_{\phi}(t) = \sum_{i=1}^9 K_i(t),$$

where

$$\begin{split} D_{\phi}(t) &= \int_{\Omega} |p'(v)| \cdot |\nabla^{2} \phi|^{2} dx \,, \\ K_{1}(t) &= \frac{d}{dt} \int_{\Omega} \rho \partial_{i} \psi \cdot \nabla \partial_{i} \phi \, dx \,, \\ K_{2}(t) &= \dot{\mathbf{X}}(t) \int_{\Omega} \rho \partial_{11} \phi \big[(2\mu + \lambda) \partial_{111} v^{S} - \partial_{11} u_{1}^{S} \big] \, dx \,, \\ K_{3}(t) &= -(2\mu + \lambda) \int_{\Omega} \rho \partial_{ij} \phi \, \partial_{ij} (vQ_{1}) \, dx \,, \\ K_{4}(t) &= \int_{\Omega} \rho \partial_{1i} \phi \, \partial_{i} (vQ_{3}) \, dx \,, \\ K_{5}(t) &= - \int_{\Omega} \rho \partial_{i} \psi_{j} \big[\partial_{t} \partial_{ij} \phi + u \cdot \nabla \partial_{ij} \phi \big] \, dx \,, \\ K_{6}(t) &= -(2\mu + \lambda) \int_{\Omega} \rho \partial_{ij} \phi \, (2\partial_{i} u \cdot \nabla \partial_{j} \phi + \partial_{ij} u \cdot \nabla \phi) \, dx \\ &+ \int_{\Omega} \rho \nabla \partial_{i} \phi \cdot \big[\partial_{i} u \cdot \nabla \psi + \partial_{i} (vp'(v)) \nabla \phi \big] \, dx \,, \\ K_{7}(t) &= \int_{\Omega} \partial_{11} \bar{v} \big[p'(v) - p'(\bar{v}) \big] \partial_{11} \phi \, dx \,+ \int_{\Omega} \rho \partial_{1} \bar{v} \, \partial_{i} \big[v \big(p'(v) - p'(\bar{v}) \big) \big] \partial_{1i} \phi \, dx \,, \end{split}$$

$$K_{8}(t) = (2\mu + \lambda) \int_{\Omega} \rho \partial_{ij} \phi \left[\partial_{ij} v \operatorname{div} \psi + 2\partial_{i} v \partial_{j} \operatorname{div} \psi \right] dx$$
$$- \int_{\Omega} \rho \partial_{i} v \left[\mu \Delta \psi + (\mu + \lambda) \nabla \operatorname{div} \psi \right] \nabla \partial_{i} \phi \, dx \,,$$
$$K_{9}(t) = -(2\mu + \lambda) \int_{\Omega} \partial_{111} u_{1}^{R} \, \partial_{11} \phi + \rho \partial_{11} u_{1}^{R} \, \partial_{i} v \partial_{1i} \phi \, dx \,.$$

Using Cauchy's inequality, we have

$$\int_{0}^{t} K_{1}(\tau) d\tau = \int_{\Omega} \rho \partial_{i} \psi \cdot \nabla \partial_{i} \phi \, dx \Big|_{\tau=0}^{\tau=t}$$

$$\leq \frac{2\mu + \lambda}{8} \left\| \sqrt{\rho} \nabla \phi(t) \right\|^{2} + C \left\| \nabla \psi(t) \right\|^{2} + C \left(\left\| \nabla \psi(0, \cdot) \right\|^{2} + \left\| \nabla^{2} \phi(0, \cdot) \right\|^{2} \right).$$

By Lemma 2.3 and Holder inequality, it holds

$$K_2(t) \lesssim \delta_S |\dot{\mathbf{X}}(t)| \int_{\Omega} \partial_1 v^S \sqrt{|p'(v)|} |\nabla^2 \phi| \, dx \lesssim \delta_S^2 |\dot{\mathbf{X}}(t)|^2 + \delta_S^2 D_{\phi}(t)$$

Using the decomposition of vQ_1 in (4.54), it follows from Cauchy's inequality that

$$K_{3}(t) \leq C \int_{\Omega} |\nabla^{2} \phi| \cdot |\nabla^{2}(vQ_{1})| \, dx \leq C \sum_{i=1}^{3} K_{3,i}(t),$$

where

$$K_{3,1}(t) = \int_{\Omega} |\nabla^2 \phi| \cdot |\nabla^2 (Q_1^R)| \, dx \,,$$

$$K_{3,2}(t) = \int_{\Omega} |\nabla^2 \phi| \cdot |\nabla^2 (Q_1^S)| \, dx \,,$$

$$K_{3,3}(t) = \int_{\Omega} |\nabla^2 \phi| \cdot |\nabla^2 (Q_1^I)| \, dx \,.$$

Using (4.50), Lemma 2.2 and Holder inequality, we have

$$\begin{split} K_{3,1}(t) &\leq C \int_{\Omega} |\nabla^{2}\phi| \cdot \partial_{1} v^{R} \big(|\psi_{1}| + |\phi| + |\nabla\psi_{1}| + |\nabla\phi| + |\nabla^{2}\psi_{1}| + |\nabla^{2}\phi| \big) \, dx \\ &\leq C \left\| \partial_{1} u_{1}^{R} \right\|_{L^{6}} \|(\phi, \psi)\|_{L^{3}} \left\| \nabla^{2}\phi \right\| + C\delta \big(\left\| \nabla(\phi, \psi) \right\|^{2} + \left\| \nabla^{2}(\phi, \psi) \right\|^{2} \big) \\ &\leq C\chi \left\| \partial_{1} u_{1}^{R} \right\|_{L^{6}}^{2} + C(\chi + \delta) \big(\left\| \nabla(\phi, \psi) \right\|^{2} + \left\| \nabla^{2}(\phi, \psi) \right\|^{2} \big) \\ &\leq C\chi \, \delta_{R}^{\frac{1}{3}} \cdot (1 + t)^{-\frac{5}{3}} + C(\chi + \delta) \big(\left\| \nabla(\phi, \psi) \right\|^{2} + \left\| \nabla^{2}(\phi, \psi) \right\|^{2} \big). \end{split}$$

Notice that

$$\|\nabla\phi\|^2 \lesssim D(t) + \delta_S^2 G^S(t) + \delta_R G^R(t),$$

 \mathbf{SO}

(4.87)

$$K_{3,1}(t) \leq C\chi \,\delta_R^{\frac{1}{3}} \cdot (1+t)^{-\frac{5}{3}} + C(\chi+\delta) \left(\left\| \nabla^2 \phi \right\|^2 + \left\| \nabla \psi \right\|_{H^1}^2 \right) \\ + C(\chi+\delta) \left(D(t) + G^S(t) + G^R(t) \right).$$

By the definition of h, it holds

$$\psi_1 = (h_1 - \bar{h}_1) + (2\mu + \lambda)\partial_1\phi - (2\mu + \lambda)\partial_1v^R,$$

and so

$$\begin{aligned} |\psi_1| \lesssim |h_1 - \bar{h}_1| + |\nabla(p - \bar{p})| + \partial_1 \bar{v} \cdot |p - \bar{p}| + \partial_1 v^R \\ \lesssim \left| h_1 - \bar{h}_1 - \frac{p - \bar{p}}{\sigma_*} \right| + |\nabla(p - \bar{p})| + |p - \bar{p}| + \partial_1 u_1^R, \end{aligned}$$

which together with $|\phi| \sim |p - \bar{p}|$ lead to

$$\begin{split} K_{3,2}(t) &\lesssim \int_{\Omega} |\nabla^2 \phi| \cdot \partial_1 v^S \left(|\psi_1| + |\phi| + |\nabla \psi_1| + |\nabla \phi| + |\nabla^2 \psi_1| + |\nabla^2 \phi| \right) dx \\ &\lesssim \int_{\Omega} |\nabla^2 \phi| \cdot \partial_1 v^S \left(\left| h_1 - \bar{h}_1 - \frac{p - \bar{p}}{\sigma_*} \right| + |\nabla (p - \bar{p})| + |p - \bar{p}| + \partial_1 u_1^R \right) dx \\ &\quad + \delta \left(\left\| \nabla (\phi, \psi) \right\|^2 + \left\| \nabla^2 (\phi, \psi) \right\|^2 \right) \\ &\lesssim \chi \left\| \partial_1 u_1^R \cdot \partial_1 v^S \right\| + \delta \left(G_2(t) + D(t) + G^S(t) \right) + \delta \left(\left\| \nabla (\phi, \psi) \right\|^2 + \left\| \nabla^2 (\phi, \psi) \right\|^2 \right) \end{split}$$

.

According to 4.4, it holds

In addition,

$$\left\|\partial_1 u_1^R \cdot \partial_1 v^S\right\| \le C \delta_R \delta_S e^{-C \delta_S t}.$$

$$\|\nabla\phi\|^2 \lesssim D(t) + \delta_S^2 G^S(t) + \delta_R^2 G^R(t).$$

Hence, we have

$$K_{3,2}(t) \leq C\delta_R \delta_S e^{-C\delta_S t} + C\delta \big(G_2(t) + D(t) + G^R(t) + G^S(t)\big) + C\delta \left\|\nabla^2 \phi\right\|^2 + C\delta \left\|\nabla \psi\right\|_{H^1}^2.$$

It follows from Lemma 4.4 that

$$\left\|\nabla^2 Q_1^I\right\| \le C\delta_R \delta_S e^{-C\delta_S t},$$

 \mathbf{SO}

$$K_{3,3}(t) \le \left\| \nabla^2 \phi \right\| \cdot \left\| \nabla^2 Q_1^I \right\| \le C \chi \delta_R \delta_S e^{-C \delta_S t}.$$

Combining the above estimates, we yield

$$K_{3}(t) \leq C\chi \,\delta_{R}^{\frac{1}{3}} \cdot (1+t)^{-\frac{5}{3}} + C\delta_{R}\delta_{S}e^{-C\delta_{S}t} + C\delta D_{\phi}(t) + C\delta \,\|\nabla\psi\|_{H^{1}}^{2} + C(\chi+\delta) \big(G_{2}(t) + D(t) + G^{R}(t) + G^{S}(t)\big).$$

Using the decomposition of vQ_3 (4.76), we have

$$K_4(t) \le C \int_{\Omega} |\nabla^2 \phi| \cdot |\nabla(vQ_3)| \, dx \le C \sum_{i=1}^3 K_{4,i}(t),$$

where

$$K_{4,1}(t) = \int_{\Omega} |\nabla^2 \phi| \cdot |\nabla(Q_3^R)| \, dx \,,$$

$$K_{4,2}(t) = \int_{\Omega} |\nabla^2 \phi| \cdot |\nabla(Q_3^S)| \, dx \,,$$

$$K_{4,3}(t) = \int_{\Omega} |\nabla^2 \phi| \cdot |\nabla(Q_3^I)| \, dx \,.$$

Similarly, we can obtain

$$\begin{split} K_{4,1}(t) &\leq C \int_{\Omega} |\nabla^{2} \phi| \cdot \partial_{1} u_{1}^{R} (|\psi_{1}| + |\phi| + |\nabla \psi_{1}| + |\nabla \phi|) \, dx \\ &\leq C \left\| \partial_{1} u_{1}^{R} \right\|_{L^{6}} \|(\phi, \psi)\|_{L^{3}} \left\| \nabla^{2} \phi \right\| + C \delta_{R} \left\| \nabla(\phi, \psi) \right\|^{2} + C \delta_{R} \left\| \nabla^{2} \phi \right\|^{2} \\ &\leq C \chi \left\| \partial_{1} u_{1}^{R} \right\|_{L^{6}}^{2} + C(\chi + \delta) \left\| \nabla(\phi, \psi) \right\|^{2} + C(\chi + \delta) \left\| \nabla^{2} \phi \right\|^{2} \\ &\leq C \chi \, \delta_{R}^{\frac{1}{3}} \cdot (1 + t)^{-\frac{5}{3}} + C(\chi + \delta) \big(\left\| \nabla(\phi, \psi) \right\|^{2} + \left\| \nabla^{2} \phi \right\|^{2} \big) \\ &\leq C \chi \, \delta_{R}^{\frac{1}{3}} \cdot (1 + t)^{-\frac{5}{3}} + C(\chi + \delta) \big(\left\| \nabla^{2} \phi \right\|^{2} + \left\| \nabla \psi \right\|^{2} \big) \\ &+ C(\chi + \delta) \big(D(t) + G^{S}(t) + G^{R}(t) \big), \end{split}$$

and

$$\begin{split} K_{4,2}(t) &\leq C \int_{\Omega} |\nabla^2 \phi| \cdot \partial_1 v^S \left(|\psi_1| + |\phi| + |\nabla \psi_1| + |\nabla \phi| \right) dx \\ &\leq C \int_{\Omega} |\nabla^2 \phi| \cdot \partial_1 v^S \left(\left| h_1 - \bar{h}_1 - \frac{p - \bar{p}}{\sigma_*} \right| + |\nabla (p - \bar{p})| + |p - \bar{p}| + \partial_1 u_1^R \right) dx \\ &\quad + C\delta \left\| \nabla (\phi, \psi) \right\|^2 + C\delta \left\| \nabla^2 \phi \right\|^2 \\ &\leq C \delta_R \delta_S e^{-C\delta_S t} + C\delta \big(G_2(t) + D(t) + G^R(t) + G^S(t) + \left\| \nabla^2 \phi \right\|^2 + \left\| \nabla \psi \right\|^2 \big). \end{split}$$

For $K_{4,3}(t)$, the only nontrivial thing is to estimate

$$\int_{\Omega} |\nabla^2 \phi| \cdot \left| \nabla \left(v \partial_1 (\bar{p} - p^S - p^R) \right) \right| dx.$$

It follows from direct computations that

$$\begin{aligned} |\nabla \partial_1 (\bar{p} - p^S - p^R)| \\ \lesssim & \left| \nabla \left[(p'(\bar{v}) - p'(v^R)) \partial_1 v^R \right] \right| + \left| \nabla \left[(p'(\bar{v}) - p'(v^S)) \partial_1 v^S \right] \right| \\ \lesssim & \left| \bar{v} - v^R \right| \partial_1 v^R + \left| \bar{v} - v^S \right| \partial_1 v^S + \partial_1 v^R \cdot \partial_1 v^S \,, \end{aligned}$$

thus by Lemma 4.4, we have

$$\left\|\nabla \partial_1 (\bar{p} - p^S - p^R)\right\| \le C \delta_R \delta_S e^{-C \delta_S t}.$$

Notice that

$$\left\|\partial_1(\bar{p}-p^S-p^R)\right\|_{L^{\infty}} \lesssim \left\|\partial_1 v^R\right\|_{L^{\infty}} + \left\|\partial_1 v^S\right\|_{L^{\infty}} \lesssim \delta_R + \delta_S \lesssim \delta.$$

Hence, it holds

$$\begin{split} &\int_{\Omega} |\nabla^2 \phi| \cdot \left| \nabla \left(v \partial_1 (\bar{p} - p^S - p^R) \right) \right| dx \\ &\leq C \int_{\Omega} |\nabla^2 \phi| \cdot |\nabla \partial_1 (\bar{p} - p^S - p^R)| dx + C \int_{\Omega} |\nabla^2 \phi| \cdot |\nabla v| \cdot |\partial_1 (\bar{p} - p^S - p^R)| dx \\ &\leq C \chi \, \delta_R \delta_S e^{-C \delta_S t} + C \delta \int_{\Omega} |\nabla^2 \phi| \cdot |\nabla \phi| \, dx \\ &\leq C \chi \, \delta_R \delta_S e^{-C \delta_S t} + C \delta \left(D(t) + G^S(t) + G^R(t) \right) + C \delta \left\| \nabla^2 \phi \right\|^2 \, . \end{split}$$

Therefore, we have

$$K_4(t) \le C\chi \,\delta_R^{\frac{1}{3}} \cdot (1+t)^{-\frac{5}{3}} + C\delta_R \delta_S e^{-C\delta_S t} + C\delta D_\phi(t) + C\delta \, \|\nabla\psi\|^2 + C(\chi+\delta) \big(G_2(t) + D(t) + G^R(t) + G^S(t)\big).$$

For $K_5(t)$, according to (4.82), it holds

$$K_5(t) = -\int_{\Omega} \rho \partial_i \psi_j \left[\partial_t \partial_{ij} \phi + u \cdot \nabla \partial_{ij} \phi \right] dx =: \sum_{i=1}^5 K_{5,i}(t),$$

where

$$\begin{split} K_{5,1}(t) &= -\dot{\mathbf{X}}(t) \int_{\Omega} \rho \partial_1 \psi_1 \partial_{111} v^S \, dx \,, \\ K_{5,2}(t) &= \int_{\Omega} \rho \partial_i \psi_j \left[\partial_i u \cdot \nabla \partial_j \phi + \partial_j u \cdot \nabla \partial_i \phi + \partial_{ij} u \cdot \nabla \phi \right] \, dx \,, \\ K_{5,3}(t) &= -\int_{\Omega} \rho \partial_i \psi_j \left[\partial_j \psi \cdot \partial_i \mathrm{div} \psi + \partial_i \psi \cdot \partial_j \mathrm{div} \psi + \partial_{ij} v \cdot \mathrm{div} \psi \right] \, dx \,, \\ K_{5,4}(t) &= -\int_{\Omega} \partial_i \psi_j \cdot \partial_{ij} \mathrm{div} \psi \, dx \,, \\ K_{5,5}(t) &= \int_{\Omega} \rho \partial_i \psi_j \cdot \partial_{ij} (vQ_1) \, dx. \end{split}$$

Using Lemma 2.3 and Cauchy's inequality, we have

$$K_{5,1}(t) \le C\delta_S^2 |\dot{\mathbf{X}}(t)|^2 + C\delta_S^2 \|\nabla \psi\|^2.$$

By (4.50) and (4.87), it holds

$$\begin{aligned} K_{5,2}(t) &\leq C \int_{\Omega} |\nabla \psi| \left(|\nabla^2 \psi| |\nabla \phi| + |\nabla \psi| |\nabla^2 \phi| \right) + |\nabla \psi| \left(|\partial_{11} \bar{u}_1| |\nabla \phi| + |\partial_1 \bar{u}_1| |\nabla^2 \phi| \right) dx \\ &\leq C \left\| \nabla \psi \right\|_{L^6} \left\| \nabla (\phi, \psi) \right\|_{L^3} \left\| \nabla^2 (\phi, \psi) \right\| + C\delta \left\| \nabla \psi \right\| \left(\left\| \nabla \phi \right\| + \left\| \nabla^2 \phi \right\| \right) \\ &\leq C(\chi + \delta) \left(\left\| \nabla^2 \phi \right\|^2 + \left\| \nabla \psi \right\|_{H^1}^2 \right) + C\delta \left(G^R(t) + G^S(t) + D(t) \right). \end{aligned}$$

Similar to $K_{5,2}(t)$, we have

$$K_{5,3}(t) \leq C \int_{\Omega} |\nabla \psi| \left(|\nabla \psi| |\nabla^2 \phi| + |\nabla^2 \psi| |\nabla \phi| \right) + |\nabla \psi| \left(|\partial_{11} \bar{u}_1| |\nabla \psi| + |\partial_1 \bar{u}_1| |\nabla^2 \psi| \right) dx$$

$$\leq C \|\nabla \psi\|_{L^6} \|\nabla (\phi, \psi)\|_{L^3} \|\nabla^2 (\phi, \psi)\| + C\delta \|\nabla \psi\| \left(\|\nabla \psi\| + \|\nabla^2 \psi\| \right)$$

$$\leq C(\chi + \delta) \left(\|\nabla^2 \phi\|^2 + \|\nabla \psi\|_{H^1}^2 \right).$$

Integration by parts over Ω leads to

$$K_{5,4}(t) = -\int_{\Omega} \partial_i \psi_j \cdot \partial_{ij} \operatorname{div} \psi \, dx = \int_{\Omega} \partial_i \operatorname{div} \psi \cdot \partial_i \operatorname{div} \psi \, dx = \|\nabla \operatorname{div} \psi\|^2 \lesssim \|\nabla^2 \psi\|^2.$$

Essentially the same with $K_3(t)$, we have

$$K_{5,5}(t) \le C\chi \,\delta_R^{\frac{1}{3}} \cdot (1+t)^{-\frac{5}{3}} + C\delta_R \delta_S e^{-C\delta_S t} + C\delta D_\phi(t) + C\delta \left\|\nabla\psi\right\|_{H^1}^2 + C(\chi+\delta) \big(G_2(t) + D(t) + G^R(t) + G^S(t)\big).$$

Combining the above estimates, we yield

$$K_{5}(t) \leq C\chi \,\delta_{R}^{\frac{1}{3}} \cdot (1+t)^{-\frac{5}{3}} + C\delta_{R}\delta_{S}e^{-C\delta_{S}t} + C\delta D_{\phi}(t) + C\delta_{S}^{2}|\dot{\mathbf{X}}(t)|^{2} + C \|\nabla\psi\|_{H^{1}}^{2} + C(\chi+\delta) \big(G_{2}(t) + D(t) + G^{R}(t) + G^{S}(t)\big).$$

Using (4.50), Cauchy's inequality and the fact that

$$\left\|\nabla^{2}\phi\right\|^{2}\left\|\nabla\psi\right\|_{L^{\infty}} \lesssim \left\|\nabla^{2}\phi\right\|^{2}\left\|\nabla\psi\right\|_{H^{2}} \lesssim \chi\left\|\nabla^{2}\phi\right\|\left\|\nabla\psi\right\|_{H^{2}} \lesssim \chi D_{\phi}(t) + \chi\left\|\nabla\psi\right\|_{H^{2}}^{2},$$

we have

$$\begin{split} K_{6}(t) &\leq C \int_{\Omega} |\nabla^{2}\phi| \left(|\nabla^{2}\psi| |\nabla\phi| + |\nabla\psi| |\nabla^{2}\phi| + |\nabla\psi|^{2} + |\nabla\phi|^{2} \right) \\ &+ |\nabla^{2}\phi| \left(|\partial_{11}\bar{u}_{1}| |\nabla\phi| + |\partial_{1}\bar{u}_{1}| |\nabla\psi| + |\partial_{1}\bar{u}_{1}| |\nabla^{2}\phi| + |\partial_{1}\bar{v}| \nabla\phi| \right) dx \\ &\leq C \left\| \nabla^{2}\phi \right\| \left(\left\| \nabla^{2}\psi \right\|_{L^{6}} \left\| \nabla\phi \right\|_{L^{3}} + \left\| \nabla\psi \right\|_{L^{6}} \left\| \nabla\psi \right\|_{L^{3}} + \left\| \nabla^{2}\phi \right\| \left\| \nabla\psi \right\|_{L^{\infty}} \right) \\ &+ C \left\| \nabla\phi \right\|_{L^{6}} \left\| \nabla\phi \right\|_{L^{3}} \left\| \nabla^{2}\phi \right\| + C\delta \left\| \nabla^{2}\phi \right\| \left(\left\| \nabla(\phi,\psi) \right\| + \left\| \nabla^{2}\phi \right\| \right) \\ &\leq C(\chi + \delta) \left(D_{\phi}(t) + D(t) + G^{R}(t) + G^{S}(t) + \left\| \nabla\psi \right\|_{H^{2}}^{2} \right). \end{split}$$

Notice that $vp'(v) = -\gamma p(v)$, so it holds

$$\nabla \left[v \left(p'(v) - p'(\bar{v}) \right) \right] = \nabla \left[v p'(v) - \bar{v} p'(\bar{v}) - \phi p'(\bar{v}) \right] \lesssim |\nabla (p - \bar{p})| + |\nabla \phi| + |\phi| |\partial_1 \bar{v}|.$$

Hence, we have

$$\begin{aligned} K_7(t) &\leq C \int_{\Omega} |\partial_{11}\bar{v}| |\phi| |\nabla^2 \phi | dx + C \int_{\Omega} |\partial_1 \bar{v}| |\nabla^2 \phi| \left(|\nabla(p - \bar{p})| + |\nabla \phi| + |\phi| |\partial_1 \bar{v}| \right) dx \\ &\leq C \int_{\Omega} \left(\partial_{11} v^R + |\partial_1 v^R|^2 \right) |\phi| |\nabla^2 \phi| \, dx + C \int_{\Omega} \partial_1 v^S |p - \bar{p}| |\nabla^2 \phi| \, dx \\ &\quad + C\delta \int_{\Omega} |\nabla^2 \phi|^2 + |\nabla(p - \bar{p})|^2 + |\nabla \phi|^2 \, dx \\ &\leq C \sqrt{\delta} \left(G^R(t) + D_{\phi}(t) \right) + C\delta \left(D(t) + G^S(t) \right). \end{aligned}$$

Similar to $K_6(t)$, it holds

$$K_{8}(t) \leq C \int_{\Omega} |\nabla^{2}\phi| (|\nabla^{2}\psi||\nabla\phi| + |\nabla\psi||\nabla^{2}\phi| + |\partial_{1}\bar{v}||\nabla^{2}\psi| + |\partial_{11}\bar{v}||\nabla\psi|) dx$$

$$\leq C \|\nabla^{2}\phi\| (\|\nabla^{2}\psi\|_{L^{6}} \|\nabla\phi\|_{L^{3}} + \|\nabla\psi\|_{L^{\infty}} \|\nabla^{2}\phi\| + \delta \|\nabla\psi\|_{H^{1}})$$

$$\leq C\chi D_{\phi}(t) + C(\chi + \delta) \|\nabla\psi\|_{H^{2}}^{2}.$$

Using Lemma 2.2, we get

$$K_{9}(t) \leq C \int_{\Omega} |\partial_{11}v^{R}| \cdot |\nabla^{2}\phi| \, dx + C \int_{\Omega} |\partial_{1}v^{R}| \cdot |\nabla\phi| \cdot |\nabla^{2}\phi| \, dx$$

$$\leq C \left\| \partial_{11}v^{R} \right\| \left\| \nabla^{2}\phi \right\| + \left\| \partial_{1}v^{R} \right\|_{L^{6}} \left\| \nabla\phi \right\|_{L^{3}} \left\| \nabla^{2}\phi \right\|$$

$$\leq C(\delta_{R})^{-\frac{1}{2}} \left\| \partial_{11}v^{R} \right\|^{2} + C\chi \left\| \partial_{1}v^{R} \right\|_{L^{6}}^{2} + C(\chi + \sqrt{\delta_{R}}) \left\| \nabla^{2}\phi \right\|^{2}$$

$$\leq C(\delta_{R})^{-\frac{1}{2}} \left\| \partial_{11}v^{R} \right\|^{2} + C\chi \, \delta_{R}^{\frac{1}{3}} \cdot (1+t)^{-\frac{5}{3}} + C(\chi + \sqrt{\delta_{R}}) D_{\phi}(t).$$

In addition, it follows from Lemma 2.2 that

$$\left\|\partial_{11}v^{R}\right\|^{2} \lesssim \begin{cases} \delta_{R}, & 1+t \leq \delta_{R}^{-1} \\ \frac{1}{(1+t)^{2}}, & 1+t \geq \delta_{R}^{-1} \end{cases},$$

which yields

$$\int_0^t (\delta_R)^{-\frac{1}{2}} \left\| \partial_{11} v^R \right\|^2 d\tau \le \int_0^\infty (\delta_R)^{-\frac{1}{2}} \left\| \partial_{11} v^R \right\|^2 d\tau \lesssim \sqrt{\delta_R} \,.$$

Finally, notice the fact that

$$D_{\phi}(t) \sim \left\| \nabla^2 \phi \right\|^2.$$

Thus, integrating (4.86) over [0, t] and using the above estimates, we can deduce the desired inequality (4.80) from Lemma 4.5 and Lemma 4.6. The proof of Lemma 4.7 is completed.

4.7. Estimates on $\|\nabla^2(u-\bar{u})\|$.

Lemma 4.8. Under the assumptions of Proposition 3.2, there exists constant C > 0 (independent of ν, δ, χ and T) such that for all $t \in [0, T]$, it holds

(4.88)
$$\begin{aligned} \left\| \nabla^2 (u - \bar{u})(t) \right\|^2 + \int_0^t \left\| \nabla^3 (u - \bar{u}) \right\|^2 d\tau \\ \leq C \left(\left\| v_0 - \bar{v}(0, \cdot) \right\|_{H^2}^2 + \left\| u_0 - \bar{u}(0, \cdot) \right\|_{H^2}^2 \right) + C \delta_R^{\frac{1}{3}}. \end{aligned}$$

Proof. Multiplying (4.84) by $-\Delta \partial_i \psi$ and summing *i* from 1 to 3, then integrating the resultant equation over Ω , we have

$$\frac{d}{dt} \int_{\Omega} \frac{|\nabla^2 \psi|^2}{2} \, dx + D_3(t) =: \sum_{i=1}^8 L_i(t),$$

where

(4.89)

$$\begin{split} D_{3}(t) &:= \mu \int_{\Omega} v |\nabla \Delta \psi|^{2} \, dx + (\mu + \lambda) \int_{\Omega} v |\nabla^{2} \mathrm{div}\psi|^{2} \, dx, \\ L_{1}(t) &:= \int_{\Omega} \frac{|\nabla^{2}\psi|^{2}}{2} \cdot \mathrm{div}u \, dx - \int_{\Omega} (\partial_{j}u \cdot \nabla) \partial_{i}\psi \cdot \partial_{ij}\psi \, dx, \\ L_{2}(t) &:= \int_{\Omega} (\partial_{i}u \cdot \nabla)\psi \cdot \Delta \partial_{i}\psi + \partial_{i}(vp'(v))\nabla \phi \cdot \Delta \partial_{i}\psi \, dx, \\ L_{3}(t) &:= \int_{\Omega} vp'(v)\nabla \partial_{i}\phi \cdot \Delta \partial_{i}\psi \, dx, \\ L_{4}(t) &:= \int_{\Omega} v(p'(v) - p'(\bar{v}))\partial_{11}\bar{v} \cdot \Delta \partial_{1}\psi_{1} + \partial_{1}\bar{v}\nabla \left[v(p'(v) - p'(\bar{v}))\right] \cdot \nabla \Delta \psi_{1} \, dx, \\ L_{5}(t) &:= -\dot{\mathbf{X}}(t) \int_{\Omega} \partial_{11}u_{1}^{S} \cdot \Delta \partial_{1}\psi_{1} \, dx, \\ L_{6}(t) &:= \int_{\Omega} \nabla (vQ_{3}) \cdot \nabla \Delta \psi_{1} \, dx, \\ L_{7}(t) &:= -(2\mu + \lambda) \int_{\Omega} \nabla (v \cdot \partial_{11}u_{1}^{R}) \cdot \nabla \Delta \psi_{1} \, dx, \\ L_{8}(t) &:= (\mu + \lambda) \int_{\Omega} \partial_{i} \mathrm{div}\psi (\Delta \partial_{i}\psi - \nabla \partial_{i}\mathrm{div}\psi) \cdot \nabla v \, dx \\ &- \int_{\Omega} \partial_{i}v \left(\mu \Delta \psi + (\mu + \lambda)\nabla \mathrm{div}\psi\right) \cdot \Delta \partial_{i}\psi \, dx. \end{split}$$

It follows from Holder inequality and (4.50) that

$$L_{1}(t) \leq C\delta \left\| \nabla^{2}\psi \right\|^{2} + C \left\| \nabla\psi \right\|_{L^{3}} \left\| \nabla^{2}\psi \right\|_{L^{6}} \left\| \nabla^{2}\psi \right\|$$
$$\leq C(\delta + \chi) \left\| \nabla^{2}\psi \right\|^{2} + C\chi \left\| \nabla^{3}\psi \right\|^{2}$$
$$\leq C(\delta + \chi) \left\| \nabla^{2}\psi \right\|^{2} + C\chi D_{3}(t).$$

Notice that $vp'(v) = -\gamma p(v)$ and that

$$|\nabla (p - \bar{p})| \lesssim |\nabla \phi| + |\partial_1 \bar{v}| |\phi| \lesssim |\nabla \phi| + \delta |\phi|.$$

Thus, using Holder inequality, Sobolev inequality and (4.50), we have

$$L_{2}(t) \leq C\delta \|\nabla\psi\| \|\nabla\Delta\psi\| + C \|\nabla\psi\|_{L^{3}} \|\nabla\psi\|_{L^{6}} \|\nabla\Delta\psi\| + C\delta \|\nabla\phi\| \|\nabla\Delta\psi\| + \int_{\Omega} |\nabla(p-\bar{p})| \cdot |\nabla\phi| \cdot |\nabla\Delta\psi| \, dx$$
$$\leq C(\delta+\chi) \Big(D_{3}(t) + \|\nabla\psi\|_{H^{1}}^{2} + \|\nabla\phi\|^{2} + \|\nabla^{2}\phi\|^{2} \Big)$$
$$\leq C(\delta+\chi) \Big(D_{3}(t) + G^{R}(t) + G^{S}(t) + D(t) + \|\nabla\psi\|_{H^{1}}^{2} + \|\nabla^{2}\phi\|^{2} \Big).$$

By Cauchy's inequality, it holds

$$L_3(t) \le \frac{1}{8}D_3(t) + C \|\nabla^2 \phi\|^2.$$

Notice that

$$\nabla \left[v(p'(v) - p'(\bar{v})) \right] = -\gamma \nabla (p - \bar{p}) - \nabla [p'(\bar{v})\phi]$$

$$\lesssim |\nabla(p-\bar{p})| + (\partial_1 v^S + \partial_1 v^R)|\phi| + |\nabla\phi| \lesssim |\nabla(p-\bar{p})| + (\partial_1 v^S + \partial_1 v^R)|\phi|$$

Thus, using Lemma 2.2, Lemma 2.3 and (4.50), we have

$$L_4(t) \le C \int_{\Omega} |\partial_{11}\bar{v}| |\phi| |\nabla \Delta \psi| \, dx + C\delta \Big(D_3(t) + G^R(t) + G^S(t) + D(t) \Big)$$

$$\le C(\delta + \sqrt{\delta_R}) \Big(D_3(t) + G^R(t) \Big) + C\delta \Big(G^S(t) + D(t) \Big).$$

By Lemma 2.3, we have

$$L_{5}(t) \leq C\delta_{S} |\dot{\mathbf{X}}(t)| \sqrt{D_{3}(t)} \left\| \partial_{1} v^{S} \right\| \leq C\delta_{S}^{\frac{5}{2}} |\dot{\mathbf{X}}(t)| \sqrt{D_{3}(t)} \leq C\delta_{S}^{\frac{5}{2}} |\dot{\mathbf{X}}(t)|^{2} + C\delta_{S}^{\frac{5}{2}} D_{3}(t).$$

Using the decomposition of vQ_3 in (4.76), we divide $L_6(t)$ as follows:

$$L_{6}(t) = \int_{\Omega} \nabla(Q_{3}^{R}) \cdot \nabla \Delta \psi_{1} \, dx + \int_{\Omega} \nabla(Q_{3}^{S}) \cdot \nabla \Delta \psi_{1} \, dx$$
$$+ \int_{\Omega} \nabla(Q_{3}^{I}) \cdot \nabla \Delta \psi_{1} \, dx =: \sum_{i=1}^{3} L_{6,i}(t).$$

Using Lemma 2.2, assumption (3.21) and (4.50), we yield

$$\begin{split} L_{6,1}(t) &\leq C \int_{\Omega} |\nabla \Delta \psi| \cdot \partial_{1} u_{1}^{R} (|\psi_{1}| + |\phi| + |\nabla \psi_{1}| + |\nabla \phi|) \, dx \\ &\leq C \left\| \partial_{1} u_{1}^{R} \right\|_{L^{6}} \|(\phi, \psi)\|_{L^{3}} \left\| \nabla \Delta \psi \right\| + C \delta_{R} \left\| \nabla (\phi, \psi) \right\|^{2} + C \delta_{R} \left\| \nabla \Delta \psi \right\|^{2} \\ &\leq C \chi \left\| \partial_{1} u_{1}^{R} \right\|_{L^{6}}^{2} + C(\chi + \delta) \left\| \nabla (\phi, \psi) \right\|^{2} + C(\chi + \delta) D_{3}(t) \\ &\leq C \chi \, \delta_{R}^{\frac{1}{3}} \cdot (1 + t)^{-\frac{5}{3}} + C(\chi + \delta) \left(\left\| \nabla (\phi, \psi) \right\|^{2} + D_{3}(t) \right) \\ &\leq C \chi \, \delta_{R}^{\frac{1}{3}} \cdot (1 + t)^{-\frac{5}{3}} + C(\chi + \delta) \left(\left\| \nabla \psi \right\|^{2} + D_{3}(t) + D(t) + G^{S}(t) + G^{R}(t) \right) \end{split}$$

It follows from Lemma 2.3 and Lemma 4.4 that

$$\begin{split} L_{6,2}(t) &\leq C \int_{\Omega} |\nabla \Delta \psi| \cdot \partial_1 v^S \left(|\psi_1| + |\phi| + |\nabla \psi_1| + |\nabla \phi| \right) dx \\ &\leq C \int_{\Omega} |\nabla \Delta \psi| \cdot \partial_1 v^S \left(\left| h_1 - \bar{h}_1 - \frac{p - \bar{p}}{\sigma_*} \right| + |\nabla (p - \bar{p})| + |p - \bar{p}| + \partial_1 u_1^R \right) dx \\ &+ C\delta \left\| \nabla (\phi, \psi) \right\|^2 + C\delta \left\| \nabla \Delta \psi \right\|^2 \\ &\leq C\delta_R \delta_S e^{-C\delta_S t} + C\delta \left(G_2(t) + D(t) + G^R(t) + G^S(t) + D_3(t) + \left\| \nabla \psi \right\|^2 \right). \end{split}$$

Similar to $K_{4,3}(t)$, we have

$$L_{6,3}(t) \le C\delta_R\delta_S e^{-C\delta_S t} + C\delta \left(D(t) + G^S(t) + G^R(t) + D_3(t) \right)$$

Using Lemma 2.2, assumption (3.21) and (4.50), we get

$$L_{7}(t) \leq C \int_{\Omega} |\partial_{11}u_{1}^{R}| \cdot |\nabla\Delta\psi| \, dx + C \int_{\Omega} |\partial_{1}u_{1}^{R}| \cdot |\partial_{1}\bar{v} + \nabla\phi| \cdot |\nabla\Delta\psi| \, dx$$

$$\leq C \left\| \partial_{11}u_{1}^{R} \right\| \left\| \nabla\Delta\psi \right\| + C \left\| \partial_{1}u_{1}^{R} \right\|_{L^{6}} \left\| \nabla\phi \right\|_{L^{3}} \left\| \nabla\Delta\psi \right\|$$

$$\leq C(\delta_{R})^{-\frac{1}{2}} \left\| \partial_{11}u_{1}^{R} \right\|^{2} + C(\sqrt{\delta_{R}} + \chi)D_{3}(t) + C\chi \left\| \partial_{1}u_{1}^{R} \right\|_{L^{6}}^{2}.$$

It follows Sobolev inequality and (4.50) that

$$\begin{split} L_8(t) &\leq C \int_{\Omega} |\partial_1 \bar{v} + \nabla \phi| \cdot |\nabla \mathrm{div}\psi| \cdot (|\nabla \Delta \psi| + |\nabla^2 \mathrm{div}\psi|) \, dx \\ &+ C \int_{\Omega} |\partial_1 \bar{v} + \nabla \phi| \cdot |\mu \Delta \psi + (\mu + \lambda) \nabla \mathrm{div}\psi| \cdot |\nabla \Delta \psi| \, dx \\ &\leq C\delta \left(\left\| \nabla^2 \psi \right\|^2 + D_3(t) \right) + C \left\| \nabla^2 \mathrm{div}\psi \right\| \left\| \nabla \phi \right\|_{L^3} \left\| \nabla \mathrm{div}\psi \right\|_{L^6} \\ &+ C \left\| \nabla \Delta \psi \right\| \left\| \nabla \phi \right\|_{L^3} \left(\left\| \Delta \psi \right\|_{L^6} + \left\| \nabla \mathrm{div}\psi \right\|_{L^6} \right) \\ &\leq C\delta \left(\left\| \nabla^2 \psi \right\|^2 + D_3(t) \right) + C\chi \left(\left\| \nabla^3 \psi \right\|^2 + D_3(t) \right). \end{split}$$

Finally, notice that

$$D_3(t) \sim \left\|\nabla^3 \psi\right\|^2$$
.

4.8. Proof of Proposition 3.2. Notice that

$$\|\nabla(v-\bar{v})\|^2 \lesssim D(t) + \delta_S^2 G^S(t) + \delta_R G^R(t).$$

Thus, using Lemma 4.5-4.8, we immediately obtain (3.22). In addition, by the definition of $\mathbf{X}(t)$ (3.15), it follows from Lemma 2.3 and assumption (3.21) that

$$|\dot{\mathbf{X}}(t)| \le \frac{C}{\delta_S} \|v - \bar{v}\|_{L^{\infty}} \int_{\Omega} \partial_1 v^S \, dx \le C \, \|v - \bar{v}\|_{L^{\infty}} \, ,$$

which implies (3.23). The proof of Proposition 3.2 is completed.

References

- Didier Bresch and Benoît Desjardins. On the construction of approximate solutions for the 2D viscous shallow water model and for compressible Navier-Stokes models. J. Math. Pures Appl. (9), 86(4):362–368, 2006.
- [2] Kyudong Choi, Moon-Jin Kang, Young-Sam Kwon, and Alexis F. Vasseur. Contraction for large perturbations of traveling waves in a hyperbolic-parabolic system arising from a chemotaxis model. *Math. Models Methods Appl. Sci.*, 30(2):387–437, 2020.
- [3] C. M. Dafermos. Entropy and the stability of classical solutions of hyperbolic systems of conservation laws. In Recent mathematical methods in nonlinear wave propagation (Montecatini Terme, 1994), volume 1640 of Lecture Notes in Math., pages 48–69. Springer, Berlin, 1996.
- [4] Ronald J. DiPerna. Uniqueness of solutions to hyperbolic conservation laws. Indiana Univ. Math. J., 28(1):137–188, 1979.
- [5] Jonathan Goodman. Nonlinear asymptotic stability of viscous shock profiles for conservation laws. Arch. Ration. Mech. Anal., 95(4):325–344, 1986.
- [6] Jonathan Goodman. Stability of viscous scalar shock fronts in several dimensions. Trans. Amer. Math. Soc., 311(2):683–695, 1989.
- [7] David Hoff and Kevin Zumbrun. Asymptotic behavior of multidimensional scalar viscous shock fronts. Indiana Univ. Math. J., 49(2):427-474, 2000.
- [8] David Hoff and Kevin Zumbrun. Pointwise Green's function bounds for multidimensional scalar viscous shock fronts. J. Differential Equations, 183(2):368-408, 2002.
- [9] Feimin Huang, Jing Li, and Akitaka Matsumura. Asymptotic stability of combination of viscous contact wave with rarefaction waves for one-dimensional compressible Navier-Stokes system. Arch. Ration. Mech. Anal., 197(1):89–116, 2010.
- [10] Feimin Huang and Akitaka Matsumura. Stability of a composite wave of two viscous shock waves for the full compressible Navier-Stokes equation. Comm. Math. Phys., 289(3):841–861, 2009.
- [11] Jeffrey Humpherys, Gregory Lyng, and Kevin Zumbrun. Multidimensional stability of large-amplitude Navier-Stokes shocks. Arch. Ration. Mech. Anal., 226(3):923–973, 2017.
- [12] A. M. Il'in and O. A. Oleňnik. Asymptotic behavior of solutions of the Cauchy problem for some quasi-linear equations for large values of the time. *Mat. Sb.* (N.S.), 51(93):191–216, 1960.
- [13] Kazuo Ito. Asymptotic decay toward the planar rarefaction waves of solutions for viscous conservation laws in several space dimensions. Number 966, pages 116–135. 1996. Nonlinear evolution equations and their applications (Japanese) (Kyoto, 1995).
- [14] Moon-Jin Kang, Alexis F., and Yi Wang. Time-asymptotic stability of generic Riemann solutions for compressible Navier-Stokes-Fourier equations, 2023. arXiv: 2306.05604.
- [15] Moon-Jin Kang and Hobin Lee. Long-time behavior toward composite wave of shocks for 3D barotropic Navier-Stokes system, 2024. arXiv: 2406.11215.
- [16] Moon-Jin Kang and Alexis F. Vasseur. L²-contraction for shock waves of scalar viscous conservation laws. Ann. Inst. H. Poincaré C Anal. Non Linéaire, 34(1):139–156, 2017.
- [17] Moon-Jin Kang and Alexis F. Vasseur. Contraction property for large perturbations of shocks of the barotropic Navier-Stokes system. J. Eur. Math. Soc. (JEMS), 23(2):585–638, 2021.
- [18] Moon-Jin Kang, Alexis F. Vasseur, and Yi Wang. L²-contraction of large planar shock waves for multi-dimensional scalar viscous conservation laws. J. Differential Equations, 267(5):2737–2791, 2019.
- [19] Moon-Jin Kang, Alexis F. Vasseur, and Yi Wang. Time-asymptotic stability of composite waves of viscous shock and rarefaction for barotropic Navier-Stokes equations. Adv. Math., 419:Paper No. 108963, 66, 2023.
- [20] Nicholas Leger and Alexis F. Vasseur. Relative entropy and the stability of shocks and contact discontinuities for systems of conservation laws with non-BV perturbations. Arch. Ration. Mech. Anal., 201(1):271–302, 2011.
- [21] Hai-Liang Li, Teng Wang, and Yi Wang. Wave phenomena to the three-dimensional fluid-particle model. Arch. Ration. Mech. Anal., 243(2):1019–1089, 2022.
- [22] Lin-an Li, Teng Wang, and Yi Wang. Stability of planar rarefaction wave to 3D full compressible Navier-Stokes equations. Arch. Ration. Mech. Anal., 230(3):911–937, 2018.
- [23] Tai-Ping Liu. Nonlinear stability of shock waves for viscous conservation laws. Mem. Amer. Math. Soc., 56(328):v+108, 1985.
- [24] Tai-Ping Liu and Zhouping Xin. Pointwise decay to contact discontinuities for systems of viscous conservation laws. Asian J. Math., 1(1):34–84, 1997.
- [25] Tai-Ping Liu and Yanni Zeng. Shock waves in conservation laws with physical viscosity. Mem. Amer. Math. Soc., 234(1105):vi+168, 2015.

JIAJIN SHI AND YI WANG

- [26] Akitaka Matsumura. Waves in compressible fluids: viscous shock, rarefaction, and contact waves. In Handbook of mathematical analysis in mechanics of viscous fluids, pages 2495–2548. Springer, Cham, 2018.
- [27] Akitaka Matsumura and Kenji Nishihara. On the stability of travelling wave solutions of a one-dimensional model system for compressible viscous gas. Japan J. Appl. Math., 2(1):17–25, 1985.
- [28] Akitaka Matsumura and Kenji Nishihara. Asymptotics toward the rarefaction waves of the solutions of a onedimensional model system for compressible viscous gas. Japan J. Appl. Math., 3(1):1–13, 1986.
- [29] Akitaka Matsumura and Kenji Nishihara. Global stability of the rarefaction wave of a one-dimensional model system for compressible viscous gas. Comm. Math. Phys., 144(2):325–335, 1992.
- [30] Jiayun Meng. Time-asymptotic stability of composite weak planar waves for a general $n \times n$ multi-D viscous system, 2025. arXiv: 2501.14188.
- [31] John Nash. Le problème de Cauchy pour les équations différentielles d'un fluide général. Bull. Soc. Math. France, 90:487–497, 1962.
- [32] Kenji Nishihara, Tong Yang, and Huijiang Zhao. Nonlinear stability of strong rarefaction waves for compressible Navier-Stokes equations. SIAM J. Math. Anal., 35(6):1561–1597, 2004.
- [33] Masataka Nishikawa and Kenji Nishihara. Asymptotics toward the planar rarefaction wave for viscous conservation law in two space dimensions. Trans. Amer. Math. Soc., 352(3):1203–1215, 2000.
- [34] V. A. Solonnikov. The solvability of the initial-boundary value problem for the equations of motion of a viscous compressible fluid. Zap. Naučn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI), 56:128–142, 197, 1976. Investigations on linear operators and theory of functions, VI.
- [35] Anders Szepessy and Zhouping Xin. Nonlinear stability of viscous shock waves. Arch. Ration. Mech. Anal., 122(1):53– 103, 1993.
- [36] Teng Wang and Yi Wang. Nonlinear stability of planar viscous shock wave to three-dimensional compressible Navier-Stokes equations. J. Eur. Math. Soc. (JEMS), 2023.
- [37] Zhouping Xin. Asymptotic stability of planar rarefaction waves for viscous conservation laws in several dimensions. Trans. Amer. Math. Soc., 319(2):805–820, 1990.

SCHOOL OF MATHEMATICAL SCIENCES, NANJING NORMAL UNIVERSITY, NANJING 210023, P. R. CHINA

AND INSTITUTE OF APPLIED MATHEMATICS, ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, CHINESE ACADEMY OF SCIENCES, BEIJING 100190, P. R. CHINA

Email address: shijiajin@amss.ac.cn

INSTITUTE OF APPLIED MATHEMATICS, ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, CHINESE ACADEMY OF SCIENCES, BEIJING 100190, P. R. CHINA

AND SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF CHINESE ACADEMY OF SCIENCES, BEIJING 100049, P. R. CHINA *Email address*: wangyi@amss.ac.cn