

ANALYTICAL STUDY OF A GENERALIZED DIRICHLET–NEUMANN OPERATOR FOR THREE-DIMENSIONAL WATER WAVES WITH VORTICITY

S. PASQUALI^(*)

ABSTRACT. In this paper we consider three-dimensional water waves with vorticity, under the action of gravity. We discuss a generalized Zakharov–Craig–Sulem formulation of the problem introduced by Castro and Lannes, which involves a generalized Dirichlet–Neumann operator. We study this operator in detail, extending some well-known results about the classical Dirichlet–Neumann operator for irrotational water waves, such as the Taylor expansion in homogeneous powers of the wave profile, the computation of its differential and a parilinearization result. We stress the fact that no geometric condition on either the velocity field or the vorticity is assumed.

Keywords: Vorticity, Water Waves, Dirichlet–Neumann
MSC2020: 37K45, 76B03, 76B15

CONTENTS

1. Introduction	1
2. Main results	10
3. The straightened boundary value problem	15
4. Corollaries of Proposition 1.4	16
5. Regularity of the velocity field	17
6. Analyticity of the generalized Dirichlet–Neumann operator	23
7. Parilinearization of the generalized Dirichlet–Neumann operator	32
Acknowledgements	40
Appendix A. Proof of Proposition 1.4	40
Appendix B. Paradifferential calculus	49
References	52

1. INTRODUCTION

In this paper we study three-dimensional pure gravity waves on the free surface of a perfect fluid. We assume that the fluid has flat bottom with finite depth. We denote by \mathbf{U} the velocity field of the fluid, and by $\boldsymbol{\omega} := \operatorname{curl} \mathbf{U}$ its vorticity.

The problem of existence of small-amplitude water waves has a long history. The first results for two-dimensional travelling water waves have been proved in the irrotational setting by Stokes [Sto80], and later for periodic waves by Levi-Civita [LC25], and in the case of nonzero vorticity by Gerstner [Ger09]. The problem of existence of two-dimensional water waves has been extensively studied, both in the irrotational setting (see [IPT05] [Wu09] [AD15] [IP15] [BBHM18] [Bam21] [AIT22] [FG24] and references therein; see [AB15] [BD18] for gravity-capillary water waves), and in the case of nonzero vorticity (we mention [Wah07] [IT18] [BFM21] [BFM24] for the case of constant vorticity; see [CS04] [Wah06] [EWZ22] [WW23] for the case of an arbitrary vorticity distribution).

On the other hand, there are fewer results about the existence of three-dimensional small-amplitude water waves. Regarding travelling waves in the irrotational case, we mention the results by Reeder and Shinbrot [RS81] and by Craig and Nicholls [CN00, CN02] for *gravity-capillary* water waves, and the results by Iooss and Plotnikov for *pure gravity* water waves [IP09, IP11]; we also mention the survey [HHS⁺22] regarding travelling water waves. We also mention [ABZ14] [ABZ16] regarding the local Cauchy theory for gravity water waves, [Wu11] [GMS12] [DIPP17] regarding global existence for small initial data, [IP19] about long-time existence of gravity-capillary water waves in the periodic setting, [BGW18] [BGW22] regarding the existence of fully localised solitary waves for the three-dimensional gravity-capillary water waves system.

Recently there has been some interest in Beltrami fields, namely those fields for which $\text{curl } \mathbf{U} = \alpha \mathbf{U}$, for some $\alpha \in \mathbb{R}$. We mention a variational formulation by Lokharu and Wahlén [LW19] for doubly periodic waves with relative velocities given by Beltrami fields, which is valid under general assumptions on the wave profile (which admits for example the case of overhanging wave profiles). More recently, Groves and Horn proved another variational formulation for Beltrami fields [GH20], under the more classical assumption that the free surface is given by the graph of an unknown function η depending only on the horizontal directions: such formulation, which can be considered as a generalization of an alternative variational framework for three-dimensional irrotational water waves by Benjamin [Ben84], is not only more explicit, but it allows to recover the classical Zakharov–Craig–Sulem formulation in the irrotational case $\alpha = 0$. Moreover, this formulation leads naturally to the definition of a generalized Dirichlet–Neumann operator, which reduces to the classical Dirichlet–Neumann operator in the irrotational case; for a more detailed study of the properties of the generalized Dirichlet–Neumann operator for Beltrami flows see [GNPW24]. These formulations lead to existence results for small amplitude doubly periodic waves [LSW20] [GNPW24].

Regarding three-dimensional water waves with general vorticity, we mention the paper by Castro and Lannes [CL15], in which the authors introduced a generalized Zakharov–Craig–Sulem formulation (widely used in the present work) and proved a local well-posedness result; moreover, the authors proved that their formulation is formally Hamiltonian, and that it reduces to the Zakharov–Craig–Sulem formulation in the irrotational case. We mention [WZZZ21], in which the authors establish local well-posedness for solutions with low regularity, using a different formulation compared to Castro–Lannes, see (6.1)–(6.8) in [WZZZ21]. More recently, in [SVW24] the authors constructed symmetric doubly periodic gravity-capillary three-dimensional water waves with small vorticity, by adapting an ansatz introduced by Lortz in the context of existence of symmetric magnetohydrostatic equilibria in toroidal domains. We mention a recent result by Ifrim, Pineau, Tataru and Taylor [IPTT23], in which the authors provide a complete local well-posedness theory in Sobolev spaces for the free boundary incompressible Euler equations with zero surface tension on a connected fluid domain, working in an Eulerian framework which allow the authors to prove sharp results without using Nash–Moser theory.

We also mention the papers [CL00] [CS07] [Lin05] [SZ11], which deal with the local well-posedness theory of the free boundary Euler equations (and its variants) and take into account the effect of vorticity.

Finally, we mention a recent and important result by Ginsberg and Pusateri [GP24], in which the authors proved long-time existence for small amplitude three-dimensional water waves with vorticity. From a technical point of view, the result by Ginsberg and Pusateri requires the following geometric assumption:

- i. the initial vorticity ω_0 vanishes on the boundary of the fluid domain at time $t = 0$, see statement of Theorem 2.2 of [GP24].

As mentioned in pag.7 of [GP24], condition i. is preserved in time. We point out that in [GP24] the authors decompose the velocity field such that its rotational part is tangential to the free surface, see (1.1d) and (2.1) of [GP24]; a consequence of such decomposition is the fact that the system can be reduced to a system of two equations evaluated on the free boundary, like the Zakharov–Craig–Sulem formulation for the irrotational case (see Appendix B of [GP24]; see also Sec.3 of [Gin18]).

In the present paper we perform a detailed study of the water waves system with general vorticity, proving analytic properties of the generalized Dirichlet–Neumann operator $\mathcal{G}_{gen}[\eta]$, where η is a function whose graph determines the free surface of the fluid domain. Regarding the velocity field and the generalized Dirichlet–Neumann operator, we prove:

- an estimate for the velocity field of the fluid;
- an analyticity result for the generalized Dirichlet–Neumann operator;
- a formula for the differential of the rotational part of the generalized Dirichlet–Neumann operator;
- an explicit (up to recursive formulae) Taylor expansion of the generalized Dirichlet–Neumann operator around $\eta = 0$;
- a parilinearization result for the rotational part of the generalized Dirichlet–Neumann operator.

The assumptions used in the present paper involve only the regularity and the smallness of $(\eta, \Phi, \tilde{\omega})$ in suitable Sobolev and Hölder spaces, where Φ denotes the trace of the velocity potential at the surface, and $\tilde{\omega}$ denotes the straightened vorticity. We emphasise the fact that no geometric condition on either the velocity field or the vorticity is required; the lack of geometric assumptions leads to a highly non-trivial analysis of the rotational part of both the velocity field and of the generalized Dirichlet–Neumann operator.

As pointed out by Castro and Lannes (see pag. 1172 in [CL15]), the advantage of using their formulation is to highlight the role of vorticity as explicitly as possible. We believe that the present work complements the formulation by Castro and Lannes, providing a very flexible framework which generalizes in a natural way well-known results for the Zakharov–Craig–Sulem formulation in the irrotational case, and that it is suited to deal with many different problems, such as long-time existence of solutions, analysis of periodic wave patterns or solitary waves, and derivation of asymptotic models in various physical regimes.

1.1. The Model. Consider an incompressible inviscid fluid occupying a three-dimensional domain with flat bottom, under the action of gravity. The fluid domain can be parametrized by

$$(1.1) \quad D_\eta(t) := \{(\mathbf{x}', z) \in \mathbb{R}^2 \times \mathbb{R} : -h < z < \eta(t, \mathbf{x}')\},$$

so that the free surface is given by the graph of an unknown function $\eta : \mathbb{R} \times \mathbb{R}^2 \rightarrow \mathbb{R}$, and $h > 0$ is the depth of the fluid. We also assume that the fluid has constant density equal to 1.

The equations describing the flow are given by

$$(1.2) \quad \operatorname{div} \mathbf{U} = 0, \quad \text{in } D_\eta(t),$$

$$(1.3) \quad \mathbf{U}_t + (\mathbf{U} \cdot \nabla_{\mathbf{x}', z}) \mathbf{U} = -\nabla_{\mathbf{x}', z} \mathcal{P} - g \mathbf{e}_3, \quad \text{in } D_\eta(t),$$

$$(1.4) \quad \mathbf{U} \cdot \mathbf{e}_3 = 0, \quad \text{at } z = -h,$$

$$(1.5) \quad \eta_t - \mathbf{U} \cdot \mathbf{N} = 0, \quad \text{at } z = \eta,$$

$$(1.6) \quad \mathcal{P} = \mathcal{P}_0, \quad \text{at } z = \eta,$$

where $\mathbf{U} : \overline{D_\eta(t)} \rightarrow \mathbb{R}^3$ is the velocity field, $\mathbf{e}_3 = (0, 0, 1)^T$, \mathbf{N} denotes the outward normal vector, g is the gravity constant, \mathcal{P}_0 is the atmospheric pressure (assumed to be

constant). The curl and divergence operator used above are defined as $\text{curl } \mathbf{U} := \nabla_{\mathbf{x}',z} \times \mathbf{U}$ and $\text{div } \mathbf{U} := \nabla_{\mathbf{x}',z} \cdot \mathbf{U}$.

Whenever it is not important, we omit the time-dependence of $D_\eta(t)$ and we will denote it by D_η . We also denote by D_0 the fixed domain $\mathbb{R}^2 \times (-h, 0)$.

Remark 1.1. *In addition to (1.2)-(1.6), in the literature it is usually assumed that the vorticity of the fluid satisfies additional properties. Two examples are irrotational velocity fields, which correspond to*

$$(1.7) \quad \text{curl } \mathbf{U} = \mathbf{0}, \quad \text{in } D_\eta,$$

and (strong) Beltrami fields, which correspond to

$$(1.8) \quad \text{curl } \mathbf{U} = \alpha \mathbf{U}, \quad \text{in } D_\eta, \quad \text{for some } \alpha \in \mathbb{R}.$$

1.2. Notation. Let $\mathbf{F} = (F_1, F_2, F_3)^T$ be a three-dimensional vector field, and denote by $\mathbf{F}_h = (F_1, F_2)^T$ and by $\mathbf{F}_\parallel = \mathbf{F}_h + F_3 \nabla \eta|_{z=\eta}$ the horizontal component of the tangential part of \mathbf{F} . In order to simplify the notation, we will sometimes indicate the evaluation at the free surface with an underscore.

For the two-dimensional vector field $\mathbf{f} = (f_1, f_2)^T$ we denote $\mathbf{f}^\perp = (f_2, -f_1)^T$; we also introduce the following operators

$$(1.9) \quad \nabla = (\partial_x, \partial_y)^T, \quad \nabla^\perp = (\partial_y, -\partial_x)^T,$$

(this notation is consistent with [GNPW24], and differs from [CL15]). According to the Hodge-Weyl decomposition for vector fields in \mathbb{R}^2 , for any three-dimensional vector field \mathbf{F} we have

$$(1.10) \quad \mathbf{F}_\parallel = \nabla \Phi + \nabla^\perp \Psi,$$

$$(1.11) \quad \Phi = \Delta^{-1}(\nabla \cdot \mathbf{F}_\parallel), \quad \Psi = \Delta^{-1}(\nabla^\perp \cdot \mathbf{F}_\parallel),$$

where Δ^{-1} is the two-dimensional Newtonian potential.

We denote by $\mathbf{N} := (-\eta_x, -\eta_y, 1)^T$ the normal vector at the free surface. Therefore for any vector field $\mathbf{A} : D_\eta \rightarrow \mathbb{R}^3$ we have

$$\underline{\mathbf{A}} \times \mathbf{N} = \begin{pmatrix} \mathbf{A}_\parallel^\perp \\ \mathbf{A}_\parallel^\perp \cdot \nabla \eta \end{pmatrix}.$$

We denote by \mathbf{v} the horizontal component of the velocity field \mathbf{U} , and by w its vertical component, so that

$$\mathbf{U} = \begin{pmatrix} \mathbf{v} \\ w \end{pmatrix}.$$

We denote by $\mathcal{S}(\mathbb{R}^2)$ the space of Schwartz functions on \mathbb{R}^2 .

We set $\Lambda := (1 - \Delta)^{1/2}$, and we define for all $s \in \mathbb{R}$ the Sobolev space

$$H^s(\mathbb{R}^2) := \{u \in \mathcal{S}'(\mathbb{R}^2) : \Lambda^s f \in L^2(\mathbb{R}^2)\},$$

$$\|u\|_{H^s(\mathbb{R}^2)} := \|\Lambda^s u\|_{L^2(\mathbb{R}^2)}, \quad \forall u \in H^s(\mathbb{R}^2),$$

and for all $k \in \mathbb{N}$ the Sobolev space

$$H^k(D_\eta) := \{u \in \mathcal{S}'(D_\eta) : \partial^\alpha f \in L^2(D_\eta) \quad \forall \alpha \in \mathbb{N}^3 : |\alpha| \leq k\},$$

$$\|u\|_{H^k(D_\eta)} := \sum_{\alpha \in \mathbb{N}^3 : |\alpha| \leq k} \|\partial^\alpha f\|_{L^2(D_\eta)}, \quad \forall u \in H^k(D_\eta).$$

Notice that by an extension argument standard Sobolev embedding theorems holds true also if we replace \mathbb{R}^3 by D_0 (see also proof of Theorem 4.10 in [GH20]).

We also introduce for all $s \in \mathbb{R}$ the space

$$\begin{aligned} H_0^s(\mathbb{R}^2) &:= \{u \in H^s(\mathbb{R}^2) : \exists v \in H^{s+1}(\mathbb{R}^2) \text{ s.t. } u = |\mathbf{D}|v\}, \\ \mathbf{D} &:= -i\nabla, \\ \|u\|_{H_0^s(\mathbb{R}^2)} &:= \|\mathbf{D}|^{-1}u\|_{H^{s+1}(\mathbb{R}^2)}, \quad \forall u \in H_0^s(\mathbb{R}^2). \end{aligned}$$

Moreover, we define for all $s \in \mathbb{R}$ and $k \in \mathbb{N}$ the space $H^{s,k}(D_0)$ by

$$\begin{aligned} H^{s,k}(D_0) &:= \bigcap_{j=0}^k H^j((-h, 0); H^{s-j}(\mathbb{R}^2)), \\ \|u\|_{H^{s,k}(D_0)} &:= \sum_{j=0}^k \|\Lambda^{s-j} \partial_z^j u\|_{L^2(D_0)}, \quad \forall u \in H^{s,k}(D_0), \end{aligned}$$

and we recall the following embedding (see Proposition 2.12 in [Lan13])

$$H^{s+1/2,1}(D_0) \subset L^\infty([-h, 0]; H^s(\mathbb{R}^2)), \quad \forall s \in \mathbb{R},$$

where the space $L^\infty([-h, 0]; H^s(\mathbb{R}^2))$ is endowed with the canonical norm

$$\|f\|_{L^\infty([-h, 0]; H^s(\mathbb{R}^2))} := \sup_{w \in [-h, 0]} \|f(\cdot, w)\|_{H^s(\mathbb{R}^2)}, \quad \forall f \in L^\infty([-h, 0]; H^s(\mathbb{R}^2)).$$

We also recall the definition of Hölder and Zygmund spaces. Following Appendix A of [AD15], we denote by $C^0(\mathbb{R}^2)$ the space of *bounded* continuous functions; for any $s \in \mathbb{N}$, we denote by $C^s(\mathbb{R}^2)$ the space of $C^0(\mathbb{R}^2)$ -functions whose derivative of order less or equal to s are in $C^0(\mathbb{R}^2)$. For any $s \in (0, +\infty) \setminus \mathbb{N}$, we denote by $C^s(\mathbb{R}^2)$ the space of *bounded* functions on \mathbb{R}^2 whose derivatives of order $[s]$ are uniformly Hölder continuous with exponent $s - [s]$.

Next, choose a function $\Gamma \in C_0^\infty(B(0, 1])$ which is equal to 1 on $B(0, 1/2]$; set $\gamma(\xi) := \Gamma(\xi/2) - \Gamma(\xi)$ (γ is supported on the annulus $\{1/2 \leq |\xi| \leq 2\}$), then we have that for any $\xi \in \mathbb{R}^2$

$$1 = \Gamma(\xi) + \sum_{j \in \mathbb{N}} \gamma(2^{-j}\xi),$$

and, recalling that $\mathbf{D} = -i\nabla$, we set $\Delta_j := \gamma(2^{-j}\mathbf{D})$ for any $j \in \mathbb{Z}$, and $S_0 := \Gamma(\mathbf{D})$. Then for any $s \in \mathbb{R}$ we define the Zygmund space $C_*^s(\mathbb{R}^2)$ as the space of temperate distributions $f \in \mathcal{S}'(\mathbb{R}^2)$ such that

$$\|f\|_{C_*^s(\mathbb{R}^2)} := \|S_0 f\|_{L^\infty(\mathbb{R}^2)} + \sup_{j \in \mathbb{N}} 2^{js} \|\Delta_j f\|_{L^\infty(\mathbb{R}^2)} < +\infty.$$

We mention that for $s \in (0, +\infty) \setminus \mathbb{N}$ we have that $C_*^s(\mathbb{R}^2) = C^s(\mathbb{R}^2)$ and that the norms $\|\cdot\|_{C_*^s(\mathbb{R}^2)}$ and $\|\cdot\|_{C^s(\mathbb{R}^2)}$ are equivalent (see also Proposition 41.16 of [Mét08]). We also use the notations $C^0(D_0)$ and $C^s(D_0)$ to denote bounded continuous functions on D_0 and bounded functions on D_0 whose derivatives of order $[s]$ are uniformly Hölder continuous with exponent $s - [s]$, respectively. We also mention the following consequences of the Sobolev embedding theorem in Zygmund spaces (see Proposition 8.4 in [Tay11]),

$$H^s(\mathbb{R}^2) \subset C^{s-1}(\mathbb{R}^2), \quad H^s(D_0) \subset C^{s-3/2}(D_0) \quad \forall s \in \mathbb{R} \setminus \frac{1}{2}\mathbb{N}.$$

Next, we define for all $k \in \mathbb{N}$ the homogeneous Sobolev space

$$\dot{H}^{k+1}(D_\eta) := \{f \in L_{\text{loc}}^2(D_\eta) : \nabla_{\mathbf{x}', z} f \in (H^k(D_\eta))^3\},$$

endowed with the norm $\|f\|_{\dot{H}^{k+1}(D_\eta)} := \|\nabla_{\mathbf{x}', z} f\|_{H^k(D_\eta)}$, and for all $s \in \mathbb{R}$

$$\dot{H}^{s+1}(\mathbb{R}^2) := \{f \in L_{\text{loc}}^2(\mathbb{R}^2) : \nabla f \in (H^s(\mathbb{R}^2))^2\},$$

endowed with the norm $\|f\|_{\dot{H}^{s+1}(\mathbb{R}^2)} := \|\nabla f\|_{H^s(\mathbb{R}^2)}$.

By using the definitions of the above norms one can check the following embedding of homogeneous Sobolev space,

$$\dot{H}^s(\mathbb{R}^2) \cap C^0(\mathbb{R}^2) \subset C_*^{s-1}(\mathbb{R}^2), \quad \forall s \in \mathbb{R},$$

see also Theorem 1.50 of [BCD11] for a closely related result.

1.3. The generalized Zakharov–Craig–Sulem formulation. We now introduce a recent formulation for water waves with vorticity by Castro and Lannes, which generalizes the well-known Zakharov–Craig–Sulem formulation for irrotational velocity fields. We defer to Sec. 2 of [CL15] for more details.

Taking the trace of (1.3) at the free surface and then taking the cross product of that equation with \mathbf{N} , we obtain

$$(1.12) \quad -\partial_t \mathbf{U}_{\parallel} - g \nabla \eta - \frac{1}{2} \nabla |\mathbf{U}_{\parallel}|^2 + \frac{1}{2} \nabla ((1 + |\nabla \eta|^2) \underline{w}^2) - \underline{\omega} \cdot \mathbf{N} \underline{v}^{\perp} = \mathbf{0}.$$

Denoting by Π and Π_{\perp} respectively the projection onto gradient vector fields and the projection onto orthogonal gradient vector fields, we obtain

$$\Pi = \nabla \nabla^T \Delta^{-1}, \quad \Pi_{\perp} = \nabla^{\perp} (\nabla^{\perp})^T \Delta^{-1}.$$

Now we can use the Hodge–Weyl decomposition (1.10) in order to decompose \mathbf{U}_{\parallel} in the following way,

$$\mathbf{U}_{\parallel} = \Pi \mathbf{U}_{\parallel} + \Pi_{\perp} \mathbf{U}_{\parallel} = \nabla \Phi + \nabla^{\perp} \Psi,$$

for some scalar functions Φ and Ψ . Hence, by applying Π to the above Eq. (1.12), we obtain the following equation for Φ ,

$$(1.13) \quad \Phi_t + g \eta + \frac{1}{2} |\mathbf{U}_{\parallel}|^2 - \frac{1}{2} ((1 + |\nabla \eta|^2) \underline{w}^2) - \nabla \cdot \Delta^{-1} (\underline{\omega} \cdot \mathbf{N} \underline{v}^{\perp}) = 0.$$

We also observe that by taking the curl of (1.3), we obtain the vorticity equation

$$(1.14) \quad \omega_t + (\mathbf{U} \cdot \nabla_{\mathbf{x}',z}) \omega = (\omega \cdot \nabla_{\mathbf{x}',z}) \mathbf{U}, \quad \text{in } D_{\eta}.$$

Remark 1.2. *We just point out that the function Ψ in the decomposition of \mathbf{U}_{\parallel} is completely determined by ω and η . Indeed, since for any three-dimensional vector field \mathbf{F} the following identity holds true,*

$$\underline{\text{curl}} \mathbf{F} \cdot \mathbf{N} = \nabla \cdot \mathbf{F}_{\parallel}^{\perp},$$

we get

$$\underline{\omega} \cdot \mathbf{N} = \nabla \cdot \mathbf{U}_{\parallel}^{\perp},$$

so that, if $\omega \in (L^2(D_{\eta}))^3$ is divergence-free, we can obtain Ψ as the unique solution in $\dot{H}^{3/2}(\mathbb{R}^2)$ of

$$-\Delta \Psi = \underline{\omega} \cdot \mathbf{N}.$$

Remark 1.3. *At the bottom $z = -h$ the vorticity satisfies the following relation,*

$$\omega \cdot \mathbf{e}_3 = \nabla \cdot \mathbf{v}^{\perp}, \quad \text{at } z = -h,$$

so that if $\mathbf{v} \in (H^1(D_{\eta}))^2$, then $\omega \cdot \mathbf{e}_3|_{z=-h} \in H_0^{-1/2}(\mathbb{R}^2)$.

Now let us assume that $\eta \in C^2(\mathbb{R}^2)$, and let us denote by D_{η} the domain

$$(1.15) \quad D_{\eta} := \{(\mathbf{x}', z) \in \mathbb{R}^2 \times \mathbb{R} : -h < z < \eta(\mathbf{x}')\}.$$

Assume moreover that the above domain is *strictly connected*, namely that there exists $h_0 > 0$ such that

$$(1.16) \quad h + \eta(\mathbf{x}') \geq h_0, \quad \forall \mathbf{x}' \in \mathbb{R}^2.$$

Then Castro and Lannes proved that it is possible to reconstruct the velocity field \mathbf{U} in terms of $\boldsymbol{\omega}$, $\nabla\Phi$ and η (see Theorem 2.4 in [CL15], which holds also for $\eta \in W^{2,\infty}(\mathbb{R}^2)$; see also Corollary 2.44 in [Lan13] for the corresponding result for the irrotational case). More precisely, they proved that there exists a unique $\mathbf{U} \in (H^1(D_\eta))^3$ which solves the following boundary value problem,

$$(1.17) \quad \begin{cases} \operatorname{curl} \mathbf{U} = \boldsymbol{\omega}, & \text{in } D_\eta, \\ \operatorname{div} \mathbf{U} = 0, & \text{in } D_\eta, \\ \mathbf{U} \cdot \mathbf{e}_3 = 0, & \text{at } z = -h, \\ \mathbf{U}_\parallel = \nabla\Phi - \nabla^\perp \Delta^{-1}(\boldsymbol{\omega} \cdot \mathbf{N}), & \text{at } z = \eta. \end{cases}$$

In order to state the result, we have to introduce some auxiliary definitions. Indeed, let $\eta \in H^k(\mathbb{R}^2)$ with $k > \frac{5}{2}$ be such that (1.16) holds true, and let us denote by D_η the domain as in (1.15). We define the subspace of $(L^2(D_\eta))^3$ of divergence-free vector fields as

$$(1.18) \quad H(\operatorname{div}_0, D_\eta) := \{\mathbf{F} \in (L^2(D_\eta))^3 : \operatorname{div} \mathbf{F} = 0\}.$$

Moreover, we introduce the set

$$(1.19) \quad H_b(\operatorname{div}_0, D_\eta) := \{\mathbf{F} \in (L^2(D_\eta))^3 : \operatorname{div} \mathbf{F} = 0, \mathbf{F} \cdot \mathbf{N}|_{z=-h} \in H_0^{-1/2}(\mathbb{R}^2)\}$$

which we endow with the norm

$$\|\mathbf{F}\|_{2,b} := \|\mathbf{F}\|_{(L^2(D_\eta))^3} + \|\mathbf{F} \cdot \mathbf{N}|_{z=-h}\|_{H_0^{-1/2}(\mathbb{R}^2)}, \quad \forall \mathbf{F} \in H_b(\operatorname{div}_0, D_\eta).$$

Proposition 1.4. *Let $k > \frac{5}{2}$, let $\eta \in H^k(\mathbb{R}^2)$ be such that (1.16) holds true, and let us denote by D_η the domain as in (1.15). Let also $\boldsymbol{\omega} \in H_b(\operatorname{div}_0, D_\eta)$ and $\Phi \in \dot{H}^{3/2}(\mathbb{R}^2)$.*

Then there exists a unique $\mathbf{U} \in (H^1(D_\eta))^3$ which solves the boundary value problem (1.17). Moreover, one has

$$(1.20) \quad \mathbf{U} = \operatorname{curl} \mathbf{A} + \nabla_{\mathbf{x}',z} \varphi,$$

where $\mathbf{A} \in (H^2(D_\eta))^3$ solves

$$(1.21) \quad \begin{cases} \operatorname{curl} \operatorname{curl} \mathbf{A} = \boldsymbol{\omega}, & \text{in } D_\eta, \\ \operatorname{div} \mathbf{A} = 0, & \text{in } D_\eta, \\ \mathbf{A} \times \mathbf{e}_3 = \mathbf{0}, & \text{at } z = -h, \\ \operatorname{curl} \mathbf{A} \cdot \mathbf{e}_3 = 0, & \text{at } z = -h, \\ \mathbf{A} \cdot \mathbf{N} = 0, & \text{at } z = \eta, \\ (\operatorname{curl} \mathbf{A})_\parallel = -\nabla^\perp \Delta^{-1}(\boldsymbol{\omega} \cdot \mathbf{N}), & \text{at } z = \eta, \end{cases}$$

while $\varphi \in H^2(D_\eta)$ solves

$$(1.22) \quad \begin{cases} \Delta_{\mathbf{x}',z} \varphi = 0, & \text{in } D_\eta, \\ \varphi = \Phi, & \text{at } z = \eta, \\ \partial_n \varphi = 0, & \text{at } z = -h. \end{cases}$$

Then

$$(1.23) \quad \begin{aligned} & \|\mathbf{U}\|_{(L^2(D_\eta))^3} + \|\nabla_{\mathbf{x}',z} \mathbf{U}\|_{(L^2(D_\eta))^3 \times 3} \\ & \leq C(h_0^{-1}, h, \|\eta\|_{H^k(\mathbb{R}^2)}) \left(\|\boldsymbol{\omega}\|_{2,b} + \|\nabla\Phi\|_{(H^{1/2}(\mathbb{R}^2))^2} \right). \end{aligned}$$

For completeness, we prove Proposition 1.4 in Appendix A. We just mention that the statement of Proposition 1.4 improves the corresponding result by Castro and Lannes (see Theorem 2.4 of [CL15]); indeed, by Sobolev embedding we have that $\eta \in C^{3/2+\varepsilon}(\mathbb{R}^2)$, for some $\varepsilon > 0$. This is due to more refined estimates in Lemma A.9 (see Sec. 3.1.1 of [ABZ14] for a similar approach; see also Proposition 4.1 and Lemma 4.3 of [WZZZ21]).

Moreover, the proof allows us to reformulate the boundary value problem (1.21) (see Corollary 4.3; see also Sec. 4 of [GH20] for a similar formulation for Beltrami flows); such formulation allows us to obtain a solution of (1.21) by solving an equivalent boundary value problem, whose solution can be written explicitly in terms of a Green matrix (see Sec. 6).

Remark 1.5. *We can check that the first two equations in (1.21) imply that*

$$(1.24) \quad \begin{aligned} \boldsymbol{\omega} &= \operatorname{curl} \operatorname{curl} \mathbf{A} = \nabla_{\mathbf{x}',z}(\nabla_{\mathbf{x}',z} \cdot \mathbf{A}) - (\nabla_{\mathbf{x}',z} \cdot \nabla_{\mathbf{x}',z})\mathbf{A} \\ &= -\Delta_{\mathbf{x}',z}\mathbf{A}, \text{ in } D_\eta. \end{aligned}$$

Let $k > \frac{5}{2}$, $\eta \in H^k(\mathbb{R}^2)$ be such that (1.16) holds true, and let us denote by D_η the domain as in (1.15). We now define the following linear maps:

$$\begin{aligned} \mathbb{A}[\eta] &: H_b(\operatorname{div}_0, D_\eta) \rightarrow (H^2(D_\eta))^3, \quad \mathbb{A}[\eta]\boldsymbol{\omega} := \mathbf{A}, \\ \mathbb{U}_I[\eta] &: \dot{H}^{3/2}(\mathbb{R}^2) \rightarrow (H^1(D_\eta))^3, \quad \mathbb{U}_I[\eta]\Phi := \nabla_{\mathbf{x}',z}\varphi, \\ \mathbb{U}_{II}[\eta] &: H_b(\operatorname{div}_0, D_\eta) \rightarrow (H^1(D_\eta))^3, \quad \mathbb{U}_{II}[\eta]\boldsymbol{\omega} := \operatorname{curl} \mathbf{A}, \end{aligned}$$

where \mathbf{A} and φ are given by the decomposition (1.20). Moreover, we introduce the linear map that reconstructs the velocity field

$$\begin{aligned} \mathbb{U}[\eta] &: \dot{H}^{1/2}(\mathbb{R}^2) \times H_b(\operatorname{div}_0, D_\eta) \rightarrow (H^1(D_\eta))^3, \\ \mathbb{U}[\eta](\Phi, \boldsymbol{\omega}) &:= \mathbb{U}_I[\eta]\Phi + \mathbb{U}_{II}[\eta]\boldsymbol{\omega}. \end{aligned}$$

We also define the map which gives the tangential velocity field at the free surface,

$$\begin{aligned} \mathbb{U}_\parallel[\eta] &: \dot{H}^{3/2}(\mathbb{R}^2) \times H_b(\operatorname{div}_0, D_\eta) \rightarrow (H^1(D_\eta))^2, \\ \mathbb{U}_\parallel[\eta](\Phi, \boldsymbol{\omega}) &:= \nabla\Phi - \nabla^\perp \Delta^{-1}(\boldsymbol{\omega} \cdot \mathbf{N}). \end{aligned}$$

Similarly, if we denote by $\mathbb{V}[\eta]$ and $\mathbb{W}[\eta]$ the operators that reconstruct respectively the horizontal and the vertical part of the velocity field, we can define

$$\underline{\mathbb{V}}[\eta](\Phi, \boldsymbol{\omega}) := \mathbb{V}[\eta](\Phi, \boldsymbol{\omega})|_{z=\eta}, \quad \underline{\mathbb{W}}[\eta](\Phi, \boldsymbol{\omega}) := \mathbb{W}[\eta](\Phi, \boldsymbol{\omega})|_{z=\eta},$$

so that

$$\mathbb{U}_\parallel[\eta](\Phi, \boldsymbol{\omega}) = \underline{\mathbb{V}}[\eta](\Phi, \boldsymbol{\omega}) + \underline{\mathbb{W}}[\eta](\Phi, \boldsymbol{\omega}) \nabla\eta.$$

Moreover, under the same assumptions on η we can also define a *generalized Dirichlet-Neumann operator*

$$(1.25) \quad \begin{aligned} \mathcal{G}_{gen}[\eta] &: \dot{H}^{3/2}(\mathbb{R}^2) \times H_b(\operatorname{div}_0, D_\eta) \rightarrow H^{1/2}(\mathbb{R}^2), \\ \mathcal{G}_{gen}[\eta](\Phi, \boldsymbol{\omega}) &:= \mathbb{U}[\eta](\Phi, \boldsymbol{\omega}) \cdot \mathbf{N}|_{z=\eta}. \end{aligned}$$

With this definition we have

$$\underline{\mathbb{W}}[\eta](\Phi, \boldsymbol{\omega}) = \frac{\mathcal{G}_{gen}[\eta](\Phi, \boldsymbol{\omega}) + \nabla\eta \cdot \mathbb{U}_\parallel[\eta](\Phi, \boldsymbol{\omega})}{1 + |\nabla\eta|^2}.$$

Hence the system (1.2)-(1.6) is equivalent to

$$(1.26) \quad \operatorname{div} \boldsymbol{\omega} = 0, \text{ in } D_\eta.$$

$$(1.27) \quad \eta_t - \mathcal{G}_{gen}[\eta](\Phi, \boldsymbol{\omega}) = 0, \text{ at } z = \eta,$$

$$(1.28) \quad \begin{aligned} \Phi_t + g\eta + \frac{1}{2}|\mathbb{U}_\parallel[\eta](\Phi, \boldsymbol{\omega})|^2 - \frac{(\mathcal{G}_{gen}[\eta](\Phi, \boldsymbol{\omega}) + \nabla\eta \cdot \mathbb{U}_\parallel[\eta](\Phi, \boldsymbol{\omega}))^2}{2(1 + |\nabla\eta|^2)} \\ + \nabla^\perp \cdot \Delta^{-1}(\boldsymbol{\omega} \cdot \mathbf{N} \underline{\mathbb{V}}[\eta](\Phi, \boldsymbol{\omega})) = 0, \text{ at } z = \eta, \end{aligned}$$

$$(1.29) \quad \boldsymbol{\omega}_t + (\mathbb{U}[\eta](\Phi, \boldsymbol{\omega}) \cdot \nabla_{\mathbf{x}',z}) \boldsymbol{\omega} = (\boldsymbol{\omega} \cdot \nabla_{\mathbf{x}',z}) \mathbb{U}[\eta](\Phi, \boldsymbol{\omega}), \text{ in } D_\eta.$$

Remark 1.6. *Eq. (1.26) is automatically satisfied for all times if it is satisfied at time $t = 0$.*

Remark 1.7. *In the irrotational case $\boldsymbol{\omega} \equiv \mathbf{0}$ we have that*

$$\begin{aligned} \mathbb{U}_{\parallel}[\eta](\Phi, \mathbf{0}) &= \nabla\Phi, \\ \mathcal{G}_{gen}[\eta](\Phi, \mathbf{0}) &= \mathcal{G}[\eta]\Phi, \end{aligned}$$

where $\mathcal{G}[\eta]$ denotes the classical Dirichlet–Neumann operator. Moreover, the system (1.26)–(1.29) reduces to

$$(1.30) \quad \begin{aligned} \eta_t - \mathcal{G}[\eta]\Phi &= 0, \\ \Phi_t + g\eta + \frac{1}{2}|\nabla\Phi|^2 - \frac{1}{2(1+|\nabla\eta|^2)} (\mathcal{G}[\eta]\Phi + \nabla\eta \cdot \nabla\Phi)^2 &= 0, \end{aligned}$$

which is the usual Zakharov–Craig–Sulem formulation.

Observe that in the irrotational case the system (1.30) involve functions depending only on the horizontal variables, while in the system (1.26)–(1.29) the vorticity $\boldsymbol{\omega}$ is a function defined on the time-dependent fluid domain $D_{\eta}(t)$. In order to deal with this difficulty we introduce the following straightening diffeomorphism (also called admissible diffeomorphism in the literature, see Definition 2.13 in [Lan13]),

$$(1.31) \quad \begin{aligned} \Sigma(t) : D_0 &\rightarrow D_{\eta}(t), \\ \Sigma(t)(\mathbf{x}', w) &:= (\mathbf{x}', w + \sigma(t, \mathbf{x}', w)), \\ \sigma(t, \mathbf{x}', -h) &= 0, \quad \sigma(t, \mathbf{x}', 0) = \eta(t, \mathbf{x}') \in H^s(\mathbb{R}^2), \quad s > 2, \end{aligned}$$

where η satisfies condition (1.16), the coefficients of the Jacobian matrix $J_{\Sigma} = \nabla_{\mathbf{x}', w}\Sigma$ belong to $L^{\infty}(D_0)$, and there exists a constant

$$M_0 = M_0(h_0^{-1}, \|\eta\|_{H^s(\mathbb{R}^2)}, \|\eta\|_{C^1(\mathbb{R}^2)}) > 0$$

such that

$$\|J_{\Sigma, i, j}\|_{L^{\infty}(D_0)} \leq M_0, \quad \forall i, j = 1, 2, 3,$$

and such that the determinant of the Jacobian matrix J_{Σ} is uniformly bounded from below on D_0 by $c_0 > 0$ satisfying $M_0 \geq 1/c_0$.

Moreover, we say that a straightening diffeomorphism is *regularizing* if

$$\|\nabla_{\mathbf{x}', w}\sigma\|_{(H^{s-1/2, 0}(D_0))^3} \leq M_0 \|\eta\|_{H^s(\mathbb{R}^2)}.$$

The introduction of regularizing diffeomorphisms is motivated by a classical issue in elliptic PDEs, where it is important to keep track of the regularity of the boundary of the domain. We now make some remarks about straightening diffeomorphisms, and we discuss two common examples of them.

Remark 1.8. *From the above definition of the straightening diffeomorphism, and in particular from the fact that the determinant of the Jacobian matrix J_{Σ} is bounded from below, we deduce that the function σ satisfies*

$$1 + \partial_w\sigma(t) \neq 0, \quad \text{in } D_0.$$

Remark 1.9. *The so-called trivial diffeomorphism between D_0 and $D_{\eta}(t)$ is given by choosing*

$$(1.32) \quad \sigma(t, \mathbf{x}', w) := \left(1 + \frac{w}{h}\right) \eta(t, \mathbf{x}')$$

in the formula (1.31).

We mention the following result for a common choice of regularizing diffeomorphism (see Proposition 2.16 and Proposition 2.18 of [Lan13]).

Proposition 1.10. *Let $s > 2$ and assume that $\eta \in H^s(\mathbb{R}^2)$ satisfies condition (1.16). Let $\chi \in C_c^\infty(\mathbb{R})$ be a positive function equal to 1 in a neighborhood of the origin, and consider the diffeomorphism (1.31), where*

$$(1.33) \quad \sigma(t, \mathbf{x}', w) := \left(1 + \frac{w}{h}\right) \eta^{(\delta)}(t, \mathbf{x}', w), \quad \eta^{(\delta)}(t, \mathbf{x}', w) := \chi(\delta w |\mathbf{D}|) \eta(t, \mathbf{x}').$$

Then

- i. if $\delta > 0$ in (1.33) satisfies

$$\delta < \frac{h_0}{C(\chi) \|\eta\|_{H^s(\mathbb{R}^2)}},$$

$$C(\chi) := \|\chi'\|_{L^\infty(\mathbb{R})} \left[\int_{\mathbb{R}^2} (1 + |\xi|^2)^{-(s-1)} d\xi \right]^{1/2} < +\infty,$$

then the diffeomorphism defined in (1.33) is such that $J_\Sigma = \nabla_{\mathbf{x}', z} \Sigma$ belong to $(L^\infty(D_0))^0$, and there exists $M_0 > 0$ such that

$$\|J_{\Sigma, i, j}\|_{L^\infty(D_0)} \leq M_0, \quad \forall i, j = 1, 2, 3.$$

Moreover, the determinant of the Jacobian matrix J_Σ is uniformly bounded from below on D_0 such that $M_0 \geq 1/c_0$, where $c_0 = h_0 - \delta C(\chi) \|\eta\|_{H^s(\mathbb{R}^2)}$.

- ii. if the assumption of i. holds true, then the diffeomorphism defined in (1.33) is regularizing.

We also mention the following smoothing property (see Lemma 2.20 in [Lan13]).

Lemma 1.11. *Let $s \in \mathbb{R}$ and $\lambda_1 > 0$, then for all $f \in C_c(\mathbb{R})$ there exists a constant $C = C(f) > 0$ such that*

$$(1.34) \quad \left\| f(\lambda_1^{1/2} w |\mathbf{D}|) u \right\|_{H^s(D_0)}^2 \leq C \left\| (1 + \lambda_1^{1/2} |\mathbf{D}|)^{-1/2} u \right\|_{H^s(\mathbb{R}^2)}^2.$$

Observe that by Lemma 1.11 (with $\lambda_1 = \delta^2$) we have that

$$\|\eta^{(\delta)}\|_{H^{s+1/2, 2}(D_0)} \leq C(1/\delta) \|\eta\|_{H^s(\mathbb{R}^2)}.$$

As a comparison, recall that for the trivial diffeomorphism of Remark 1.9 one need to control $H^s(\mathbb{R}^2)$ -norm of the surface parameterization η in order to control the $H^s(D_0)$ -norm of σ in (1.32).

By using straightening diffeomorphisms the role of the vorticity ω is replaced by the straightened vorticity $\tilde{\omega} := \omega \circ \Sigma(t)$, and role of the velocity field \mathbf{U} is replaced by the straightened velocity field $\tilde{\mathbf{U}} := \mathbf{U} \circ \Sigma(t)$. The consequence is that with this approach we can define a straightened version of the generalized Zakharov–Craig–Sulem formulation (see (6.20)–(6.23)), where the unknowns $(\eta, \Phi, \tilde{\omega})$ are defined on the time-independent domain D_0 .

Finally, we mention that Castro and Lannes, starting from the generalized Zakharov–Craig–Sulem formulation (1.27)–(1.29), exhibited a Hamiltonian formulation for water waves with vorticity (see Sec. 6 of [CL15] for more details), which reduces to the classical Zakharov–Craig–Sulem formulation in the irrotational case.

2. MAIN RESULTS

Here we present the main results of the paper.

2.1. Analytical results for the generalized Dirichlet–Neumann operator. Exploiting the generalized Zakharov–Craig–Sulem formulation of Sec. 1.3, and requiring more regularity with respect to the assumptions of Proposition 1.4, we first prove a regularity result on the velocity field of the fluid. Recall that we denote $\Lambda = (1 - \Delta)^{1/2}$ and that the Sobolev norm $\|\cdot\|_{H^{k,\ell}(D_0)}$ controls at most ℓ vertical derivatives (when $\ell \leq k$). Introducing the operator $\partial_w^\Sigma = \frac{\partial_w}{1 + \partial_w \sigma}$, we define the *good unknowns*

$$(2.1) \quad \tilde{\varphi}_{(j)} := \partial^j \tilde{\varphi} - (\partial^j \sigma) \partial_w^\Sigma \tilde{\varphi}, \quad j \in \mathbb{N}^2 \setminus \{\mathbf{0}\},$$

$$(2.2) \quad \tilde{\Phi}_{(j)} := \partial^j \tilde{\Phi} - \tilde{\omega}_I \partial^j \eta, \quad j \in \mathbb{N}^2 \setminus \{\mathbf{0}\}, \quad \tilde{\omega}_I := \tilde{\mathcal{U}}_I[\eta]\Phi|_{w=0},$$

$$(2.3) \quad \tilde{\mathbf{A}}_{(j)} := \partial^j \tilde{\mathbf{A}} - (\partial^j \sigma) \partial_w^\Sigma \tilde{\mathbf{A}}, \quad j \in \mathbb{N}^2 \setminus \{\mathbf{0}\}.$$

Moreover, we define

$$\mathfrak{P} := \frac{|\mathbf{D}|}{(1 + |\mathbf{D}|)^{1/2}},$$

which appears already in the irrotational case, see for example Lemma 2.34 in [Lan13].

Theorem 1. *Let $N > 3/2$, let $\eta \in C^N(\mathbb{R}^2)$ be such that (1.16) holds true. Assume that the assumptions of Proposition 1.4 hold true, and that σ is a regularizing diffeomorphism such that*

$$(2.4) \quad \|\sigma\|_{C^N(D_0)} \leq C \|\eta\|_{C^N(\mathbb{R}^2)}$$

holds true. Then there exists a unique solution $\tilde{\mathcal{U}} \in (H^1(D_0))^3$ which solves the boundary value problem (3.3). If $0 \leq k < N - 3/2$, $k \in \mathbb{N}$, $\Phi \in \dot{H}^{k+3/2}(\mathbb{R}^2)$, $\Lambda^k \tilde{\omega} \in (L^2(D_0))^3$, then

$$\begin{aligned} \|\tilde{\mathcal{U}}\|_{(H^{k+1,1}(D_0))^3} &\leq C(h_0^{-1}, h, \|\eta\|_{C^N(\mathbb{R}^2)}) \times \\ &\times \left[\|\mathfrak{P}\Phi\|_{H^1(\mathbb{R}^2)} + \sum_{1 < |j| \leq k+1} \|\mathfrak{P}\Phi_{(j)}\|_{L^2(\mathbb{R}^2)} + \|\Lambda^k \tilde{\omega}\|_{2,b} \right]. \end{aligned}$$

Moreover, if $N > 3$, $0 \leq \ell \leq k < N - 3/2$ and $\tilde{\omega} \in (H^{k,\ell}(D_0))^3$, then the following higher-order estimate hold

$$\begin{aligned} \|\tilde{\mathcal{U}}\|_{(H^{k+1,\ell+1}(D_0))^3} &\leq C(h_0^{-1}, h, \|\eta\|_{C^N(\mathbb{R}^2)}) \times \\ &\times \left[\|\mathfrak{P}\Phi\|_{H^1(\mathbb{R}^2)} + \sum_{1 < |j| \leq k+1} \|\mathfrak{P}\Phi_{(j)}\|_{L^2(\mathbb{R}^2)} \right. \\ &\left. + \|\tilde{\omega}\|_{(H^{k,\ell}(D_0))^3} + \|\Lambda^k(\tilde{\omega} \cdot \mathbf{e}_3)|_{w=-h}\|_{H_0^{-1/2}} \right]. \end{aligned}$$

The above result is proved in Sec. 5. From Theorem 1 we can immediately derive the following higher order estimates on the generalized Dirichlet–Neumann operator defined in (1.25). More precisely, we consider the straightened version of the generalized Dirichlet–Neumann operator, given by

$$\tilde{\mathcal{G}}_{gen}[\eta](\Phi, \tilde{\omega}) = \tilde{\mathcal{U}}[\eta](\Phi, \tilde{\omega}) \cdot \mathbf{N}|_{w=0}.$$

Corollary 2.1. *Let $N > 3/2$, let $\eta \in C^N(\mathbb{R}^2)$ be such that (1.16) holds true. Assume that $0 \leq k < N - 3/2$, $k \in \mathbb{N}$, $\Phi \in \dot{H}^{k+3/2}(\mathbb{R}^2)$, $\tilde{\omega} \in (H^{k,0}(D_0))^3$ and that σ is a regularizing diffeomorphism such that (2.4) holds true. Then*

$$\tilde{\mathcal{G}}_{gen}[\eta](\Phi, \tilde{\omega}) \in H^{k+1/2}(\mathbb{R}^2),$$

and the following higher-order estimate holds true,

$$\begin{aligned} & \left\| \tilde{\mathcal{G}}_{gen}[\eta](\Phi, \tilde{\omega}) \right\|_{H^{k+1/2}(\mathbb{R}^2)} \leq C(h_0^{-1}, h, \|\eta\|_{C^N(\mathbb{R}^2)}) \times \\ & \times \left[\|\mathfrak{P}\Phi\|_{H^1(\mathbb{R}^2)} + \sum_{1 < |j| \leq k+1} \|\mathfrak{P}\Phi_{(j)}\|_{L^2(\mathbb{R}^2)} \right. \\ & \left. + \|\tilde{\omega}\|_{(H^{k,0}(D_0))^3} + \|\Lambda^k(\tilde{\omega} \cdot \mathbf{e}_3)|_{w=-h}\|_{H_0^{-1/2}} \right]. \end{aligned}$$

Next, we give an analyticity result for the generalized Dirichlet–Neumann operator (see also Theorem 6 and Corollary 6.2).

Theorem 2. *Let $s \geq 2$, and let $\tilde{\omega} \in (H^{s-2}(D_0))^3$, $\Phi \in \dot{H}^{s-1/2}(\mathbb{R}^2)$. Assume that the condition*

$$(2.5) \quad \|\sigma\|_{H^{s+1/2}(D_0)} \leq C \|\eta\|_{H^{s+1/2}(\mathbb{R}^2)}$$

holds true. Then there exists an open neighbourhood V of the origin in $H^{s+1/2}(\mathbb{R}^2)$ such that formulae

$$\begin{aligned} \tilde{\mathcal{G}}_{gen,I}[\eta]\Phi &= \tilde{\mathbb{U}}_I[\eta]\Phi \cdot \mathbf{N}|_{w=0}, \\ \tilde{\mathcal{G}}_{gen,II}[\eta]\tilde{\omega} &= \tilde{\mathbb{U}}_{II}[\eta]\tilde{\omega} \cdot \mathbf{N}|_{w=0} \end{aligned}$$

define analytic maps

$$\begin{aligned} \tilde{\mathcal{G}}_{gen,I}[\cdot] &: V \rightarrow L(\dot{H}^{s-1/2}(\mathbb{R}^2), H^{s-3/2}(\mathbb{R}^2)), \\ \tilde{\mathcal{G}}_{gen,II}[\cdot] &: V \rightarrow L((H^{s-2}(D_0))^3, H^{s-3/2}(\mathbb{R}^2)). \end{aligned}$$

We point out that the technical condition (2.5) is satisfied both by the trivial diffeomorphism (1.32) and by the regularizing diffeomorphism (1.33). We mention that in Sec. 6 we also compute the differential of the rotational part of the generalized Dirichlet–Neumann operator, which is a result that could be interesting in itself (see Theorem 3.21 in [Lan13] and Theorem 1.7 of [GNPW24] for corresponding results for the classical Dirichlet–Neumann operator and for the generalized Dirichlet–Neumann operator for Beltrami flows, respectively).

Proposition 2.2. *Let $s \geq 3$, and let $\tilde{\omega} \in (H^{s-2}(D_0))^3$, $\Phi \in \dot{H}^{s-1/2}(\mathbb{R}^2)$. Assume that the condition (2.5) holds true. Then there exists an open neighbourhood V of the origin in $H^{s+1/2}(\mathbb{R}^2)$ such that the differential of the operator*

$$\tilde{\mathcal{G}}_{gen,II}[\cdot] : V \rightarrow L((H^{s-2}(D_0))^3, H^{s-3/2}(\mathbb{R}^2))$$

is given by

$$(2.6) \quad d\tilde{\mathcal{G}}_{gen,II}[\eta](\delta\eta)\tilde{\omega} = -\tilde{\mathcal{G}}_{gen,II}[\eta]\tilde{\gamma} - \nabla \cdot [(\tilde{\mathbb{K}}_{II}[\eta]\tilde{\omega} - \tilde{\mathbb{W}}_{II}[\eta]\tilde{\omega} \cdot \nabla\eta) \delta\eta],$$

$$(2.7) \quad \tilde{\gamma} := \tilde{\gamma}[\eta](\delta\eta)\tilde{\omega} = \partial_w^\Sigma \tilde{\omega} \, d\sigma(\delta\eta),$$

$$\tilde{\mathbb{K}}_{II}[\eta]\tilde{\omega} := -\nabla^\perp \Delta^{-1}(\tilde{\omega} \cdot \mathbf{N}|_{w=0}), \quad \tilde{\mathbb{W}}_{II}[\eta]\tilde{\omega} := \frac{1}{1+|\eta|^2} [\tilde{\mathbb{K}}_{II}[\eta]\tilde{\omega} \cdot \nabla\eta + \tilde{\mathcal{G}}_{gen,II}[\eta]\tilde{\omega}].$$

Next, we study the Taylor expansion of the generalized Dirichlet–Neumann operator (1.25) around $\eta = 0$. Recall that from (1.25) and from Remark 1.2 we have

$$\mathcal{G}_{gen}[\eta](\Phi, \omega) = \nabla \cdot \mathbf{A}_\parallel^\perp + \underline{\nabla_{\mathbf{x}', z} \varphi} \cdot \mathbf{N}.$$

From (1.25) we recall the decomposition

$$\begin{aligned} \mathcal{G}_{gen}[\eta](\Phi, \omega) &= \mathcal{G}_{gen,I}[\eta]\Phi + \mathcal{G}_{gen,II}[\eta]\omega, \\ \mathcal{G}_{gen,I}[\eta]\Phi &:= \mathbb{U}_I[\eta]\Phi \cdot \mathbf{N}|_{z=\eta} = \underline{\nabla_{\mathbf{x}', z} \varphi} \cdot \mathbf{N}, \\ \mathcal{G}_{gen,II}[\eta]\omega &:= \mathbb{U}_{II}[\eta]\omega \cdot \mathbf{N}|_{z=\eta} = \nabla \cdot \mathbf{A}_\parallel^\perp, \end{aligned}$$

together with its straightened version

$$\begin{aligned}\tilde{\mathcal{G}}_{gen}[\eta](\Phi, \tilde{\omega}) &= \tilde{\mathcal{G}}_{gen,I}[\eta]\Phi + \tilde{\mathcal{G}}_{gen,II}[\eta]\tilde{\omega}, \\ \tilde{\mathcal{G}}_{gen,I}[\eta]\Phi &:= \tilde{\mathcal{U}}_I[\eta]\Phi \cdot \mathbf{N}|_{w=0} = \nabla^\Sigma \tilde{\varphi} \cdot \mathbf{N}|_{w=0}, \\ \tilde{\mathcal{G}}_{gen,II}[\eta]\tilde{\omega} &:= \tilde{\mathcal{U}}_{II}[\eta]\tilde{\omega} \cdot \mathbf{N}|_{w=0} = \text{curl}^\Sigma \tilde{\mathbf{A}} \cdot \mathbf{N}|_{w=0},\end{aligned}$$

and its associated expansion

$$(2.8) \quad \tilde{\mathcal{G}}_{gen}[\eta] = \sum_{j=0}^{\infty} \tilde{\mathcal{G}}_j[\eta], \quad \tilde{\mathcal{G}}_{gen,I}[\eta] = \sum_{j=0}^{\infty} \tilde{\mathcal{G}}_{j,I}[\eta], \quad \tilde{\mathcal{G}}_{gen,II}[\eta] = \sum_{j=0}^{\infty} \tilde{\mathcal{G}}_{j,II}[\eta].$$

where $\mathcal{G}_{j,I}[\eta]$ and $\mathcal{G}_{j,II}[\eta]$ are homogeneous of degree j in η . It is also useful to consider the corresponding expansion for the quantity $\tilde{\gamma}$ defined in (2.7) evaluated at $\delta\eta = \eta$, namely

$$\tilde{\gamma}|_{\delta\eta=\eta} = \sum_{j \geq 0} \tilde{\gamma}_j,$$

where $\tilde{\gamma}_j$ is homogeneous of degree j in η .

Theorem 3. *Under the assumptions of Proposition 2.2, the term $\tilde{\mathcal{G}}_0$ in the homogeneous expansion (2.8) of the operator $\tilde{\mathcal{G}}_{gen}[\eta]$ is given by*

$$\tilde{\mathcal{G}}_0 = \tilde{\mathcal{G}}_{0,I} + \tilde{\mathcal{G}}_{0,II},$$

where $\tilde{\mathcal{G}}_{0,I}$ and $\tilde{\mathcal{G}}_{0,II}$ are given by

$$\begin{aligned}\tilde{\mathcal{G}}_{0,I} \Phi &= \mathcal{F}^{-1} \left[|\boldsymbol{\xi}| \tanh(h |\boldsymbol{\xi}|) (\mathcal{F}\Phi) \right], \\ \tilde{\mathcal{G}}_{0,II} \tilde{\omega} &= \nabla \cdot \mathcal{F}^{-1} \left[\int_{-h}^0 \frac{\sinh(|\boldsymbol{\xi}|(\zeta + h))}{|\boldsymbol{\xi}| \cosh(h |\boldsymbol{\xi}|)} (\mathcal{F}\tilde{\omega})_{\mathbb{R}}^\perp(\cdot, \zeta) d\zeta \right. \\ &\quad \left. - \frac{\sinh(h |\boldsymbol{\xi}|)}{|\boldsymbol{\xi}| \cosh(h |\boldsymbol{\xi}|)} \mathcal{F} [\nabla^\perp \Delta^{-1}(\tilde{\omega}_3|_{w=0})] \right],\end{aligned}$$

respectively. Moreover, if $\tilde{\gamma}_0 = \mathbf{0}$, then for any $j \geq 1$ the term $\tilde{\mathcal{G}}_j[\eta]$ are given by

$$\tilde{\mathcal{G}}_j[\eta] = \tilde{\mathcal{G}}_{j,I}[\eta] + \tilde{\mathcal{G}}_{j,II}[\eta],$$

where $\tilde{\mathcal{G}}_{j,I}$ and $\tilde{\mathcal{G}}_{j,II}$ are given by (6.14) and (6.19), respectively.

The above result is proved in Sec. 6. As mentioned in Sec. 1.3 of [GNPW24], individual terms appearing in recursion formulae for $\mathcal{G}_j[\eta]$ involve more and more derivatives of η as j increases; the overall validity of the formulae arises from cancellations between the terms (for the irrotational case see Sec. 2.2 of [NR01]).

We just mention that the condition $\tilde{\gamma}_0 = \mathbf{0}$ depends both on the vorticity and on the choice of the straightening diffeomorphism in (1.31): it is trivially true in the irrotational case, but it is true also in more general cases (for example if we choose σ as the trivial diffeomorphism (1.32), and the vorticity $\tilde{\omega}$ in a open neighbourhood of the origin in $(H^{s-2}(D_0))^3$; see also Sec. 2.4 of [GNPW24] for a similar approach for Beltrami flows).

2.2. Paralinearization of the generalized Dirichlet–Neumann operator. In this paper we also prove a paralinearization result for the rotational part of the generalized Dirichlet–Neumann operator. We refer to Appendix B, and in particular to (B.2), for the definition of paradifferential operators.

By following the approach of Sec.3 of [ABZ11], we introduce the following localizing transform: we choose $\delta > 0$ such that the fluid domain D_η contains the strip

$$(2.9) \quad \Omega_\delta := \{(\mathbf{x}', z) \in \mathbb{R}^2 \times \mathbb{R} \mid \eta(\mathbf{x}') - \delta h \leq z < \eta(\mathbf{x}')\}.$$

Remark 2.3. We notice that if (1.16) holds true, then we can take any $\delta \in (0, h_0/h)$ in (2.9): indeed, $z \geq \eta(\mathbf{x}') - \delta h > -h$ implies $\eta(\mathbf{x}') > -h(1 - \delta)$. One can also consider more general fluid domains, see [ABZ11].

We define $\hat{\Sigma} : D_0 \rightarrow \Omega_\delta$,

$$\hat{\Sigma} : (\mathbf{x}', w) \mapsto (\mathbf{x}', \varrho(\mathbf{x}', w)), \quad \varrho(\mathbf{x}', w) := \delta w + \eta(\mathbf{x}'),$$

and write $\hat{\mathbf{A}}(\mathbf{x}', w) := \mathbf{A}(\mathbf{x}', \varrho(\mathbf{x}', w))$, where \mathbf{A} is the solution of the boundary value problem (1.21) restricted to Ω_δ .

We denote by $\nabla_{\mathbf{x}', z} = (\partial_x, \partial_y, \partial_z)^T$, and

$$\nabla_{\mathbf{x}', w}^e := (J_{\hat{\Sigma}}^{-1})^T \nabla_{\mathbf{x}', z}, \quad (J_{\hat{\Sigma}}^{-1})^T := \begin{pmatrix} \mathbb{I}_2 & -\frac{\nabla \eta}{\delta} \\ 0 & \frac{1}{\delta} \end{pmatrix},$$

$$\partial_i^e := \partial_i - \frac{\partial_i \eta}{\delta} \partial_w, \quad i = x, y, \quad \partial_w^e := \frac{\partial_w}{\delta},$$

$$\text{curl}^e \hat{f}(\mathbf{x}', w) := (\text{curl } f) \circ \hat{\Sigma}(\mathbf{x}', w), \quad \text{div}^e \hat{f}(\mathbf{x}', w) := (\text{div } f) \circ \hat{\Sigma}(\mathbf{x}', w).$$

The flattened version of the operators curl and Δ applied respectively to $\hat{\mathbf{F}}(x, y, w) = \mathbf{F}(x, y, z)$ and to $\hat{f}(x, y, w) = f(x, y, z)$ are given by

$$\begin{aligned} \text{curl}^e \hat{\mathbf{F}} &= \text{curl } \hat{\mathbf{F}} - \frac{1}{\delta} (\partial_y \eta \partial_w \hat{F}_3, -\partial_x \eta \partial_w \hat{F}_3, \partial_x \eta \partial_w \hat{F}_2 - \partial_y \eta \partial_w \hat{F}_1)^T \\ &\quad - \left(\frac{1}{\delta} - 1 \right) (\partial_w \hat{F}_2, -\partial_w \hat{F}_1, 0)^T, \\ -\Delta^e \hat{f} &= -\Delta \hat{f} + \frac{2}{\delta} (\partial_x \eta \partial_{xw}^2 \hat{f} + \partial_y \eta \partial_{yw}^2 \hat{f}) - \left(\frac{1}{\delta^2} - 1 \right) \partial_w^2 \hat{f} \\ &\quad + \frac{1}{\delta} (\Delta \eta) \partial_w \hat{f} - \frac{1}{\delta^2} |\nabla \eta|^2 \partial_w^2 \hat{f}. \end{aligned}$$

Theorem 4. Let $n \geq 4$ be such that $n \notin \mathbb{N}$, $\varepsilon > 0$ and assume that $\eta \in H^n(\mathbb{R}^2)$ is such that (1.16) holds true. If

$$(2.10) \quad \begin{aligned} \partial_w^k \hat{\mathbf{A}} &\in (C^0([-h, 0]; H^{n-k}(\mathbb{R}^2)))^3, \quad k = 0, 1, \\ \hat{\omega} &\in (C^0([-h, 0]; H^{n-2}(\mathbb{R}^2)))^3, \end{aligned}$$

then the operator

$$\hat{\mathcal{G}}_{gen, II}[\eta] \hat{\omega} := \text{curl}^e \hat{\mathbf{A}} \cdot \mathbf{N}|_{w=0},$$

is given by

$$\begin{aligned} \hat{\mathcal{G}}_{gen, II}[\eta] \hat{\omega} &= T_{\lambda_{II}} (\hat{\mathbf{A}} - T_{\partial_w^e \hat{\mathbf{A}}} \eta) + \frac{1}{\delta} T_{1+|\nabla \eta|^2} T_{\nabla^\perp \eta} \cdot \hat{\mathbf{f}}(\eta, \hat{\omega}) \\ &\quad - T_{\nabla \eta} \cdot (\hat{\mathbb{V}}_{II}[\eta] \hat{\omega}) - T_{\hat{\mathbb{V}}_{II}[\eta] \hat{\omega}} \cdot \nabla \eta - T_{|\nabla \eta|^2} (\hat{\mathbb{W}}_{II}[\eta] \hat{\omega}) \\ &\quad + T_{1+|\nabla \eta|^2} \left(T_{\partial_w^e \hat{A}_{2x}} \eta - T_{\partial_w^e \hat{A}_{1y}} \eta - T_{\eta_x} T_{(\partial_w^e)^2 \hat{A}_2} \eta + T_{\eta_y} T_{(\partial_w^e)^2 \hat{A}_1} \eta \right) \\ &\quad + \mathcal{R}_{II}, \end{aligned}$$

where the right-hand side is evaluated at $w = 0$; moreover, $\lambda_{II} \in M_{1,3}(\Sigma_{n-2}^1(\mathbb{R}^2))$ is given by (7.27), $\hat{\mathbf{f}}(\eta, \hat{\omega}) \in (H^{n-1}(\mathbb{R}^2))^2$,

$$\begin{aligned} \hat{\mathbb{V}}_{II}[\eta] \hat{\omega} &= (\text{curl}^e \hat{\mathbf{A}})_h|_{w=0} \in (H^{n-1}(\mathbb{R}^2))^2, \\ \hat{\mathbb{W}}_{II}[\eta] \hat{\omega} &= (\text{curl}^e \hat{\mathbf{A}})_3|_{w=0} \in H^{n-1}(\mathbb{R}^2), \end{aligned}$$

and $\mathcal{R}_{II}(\eta, \hat{\omega}) \in H^{2n-3-\varepsilon}(\mathbb{R}^2)$.

The above result is proved in Sec. 7.

3. THE STRAIGHTENED BOUNDARY VALUE PROBLEM

Recalling the straightening diffeomorphism (1.31), we want to study the straightened version of the boundary value problem (1.17).

For simplicity, in the following discussion we assume that η , the function σ in (1.31), the function $f : D_\eta \rightarrow \mathbb{R}$ and the vector field $\mathbf{F} : D_\eta \rightarrow \mathbb{R}^3$ are sufficiently smooth (either in Sobolev spaces or in Hölder spaces), so that the quantities we introduce below are well-defined.

We define $\tilde{\mathbf{F}}(x, y, w) := \mathbf{F} \circ \Sigma(x, y, w)$ and $\tilde{f}(x, y, w) = f \circ \Sigma(x, y, w)$; moreover, we write

$$\partial_i^\Sigma \tilde{\mathbf{F}} := (\partial_i \mathbf{F}) \circ \Sigma, \quad i = t, x, y, w,$$

and similarly for $\partial_i^\Sigma \tilde{f}$.

Notice that from the definition of the straightening diffeomorphism (1.31) we have that

$$\partial_j^\Sigma = \partial_j - \frac{\partial_j \sigma}{1 + \partial_w \sigma} \partial_w, \quad j = x, y, t, \quad \partial_w^\Sigma = \frac{\partial_w}{1 + \partial_w \sigma}.$$

Remark 3.1. *Similarly, we can define*

$$\operatorname{curl}^\Sigma \tilde{\mathbf{F}} = \nabla^\Sigma \times \tilde{\mathbf{F}} := (\operatorname{curl} \mathbf{F}) \circ \Sigma, \quad \operatorname{div}^\Sigma \tilde{\mathbf{F}} = \nabla^\Sigma \cdot \tilde{\mathbf{F}} := (\operatorname{div} \mathbf{F}) \circ \Sigma, \quad \Delta^\Sigma \tilde{f} := (\Delta f) \circ \Sigma.$$

One can check that the flattened version of the operators curl and Δ applied to $\tilde{\mathbf{F}}$ and to \tilde{f} are given by

$$\begin{aligned} \operatorname{div}^\Sigma \tilde{\mathbf{F}} &= \partial_x^\Sigma \tilde{F}_1 + \partial_y^\Sigma \tilde{F}_2 + \partial_w^\Sigma \tilde{F}_3 \\ &= \operatorname{div} \tilde{\mathbf{F}} - \frac{1}{1 + \partial_w \sigma} (\partial_x \sigma \partial_w \tilde{F}_1 + \partial_y \sigma \partial_w \tilde{F}_2) - \frac{\partial_w \sigma}{1 + \partial_w \sigma} \partial_w \tilde{F}_3, \\ \operatorname{curl}^\Sigma \tilde{\mathbf{F}} &= (\partial_y^\Sigma \tilde{F}_3 - \partial_w^\Sigma \tilde{F}_2, -\partial_x^\Sigma \tilde{F}_3 + \partial_w^\Sigma \tilde{F}_1, \partial_x^\Sigma \tilde{F}_2 - \partial_y^\Sigma \tilde{F}_1)^T \\ &= \operatorname{curl} \tilde{\mathbf{F}} - \frac{\partial_w \sigma}{1 + \partial_w \sigma} (-\partial_w \tilde{F}_2, \partial_w \tilde{F}_1, 0)^T \\ &\quad - \frac{1}{1 + \partial_w \sigma} (\partial_y \sigma \partial_w \tilde{F}_3, -\partial_x \sigma \partial_w \tilde{F}_3, \partial_x \sigma \partial_w \tilde{F}_2 - \partial_y \sigma \partial_w \tilde{F}_1)^T, \\ \Delta^\Sigma \tilde{f} &= (\partial_x^\Sigma)^2 \tilde{f} + (\partial_y^\Sigma)^2 \tilde{f} + (\partial_w^\Sigma)^2 \tilde{f} = [\mathbf{a} \partial_w^2 + \Delta + \mathbf{b} \cdot \nabla \partial_w - \mathbf{c} \partial_w] \tilde{f}, \end{aligned}$$

where

$$(3.1) \quad \mathbf{a} := \frac{1 + |\nabla_{\mathbf{x}'} \sigma|^2}{(1 + \partial_w \sigma)^2}, \quad \mathbf{b} := -2 \frac{\nabla_{\mathbf{x}'} \sigma}{1 + \partial_w \sigma}, \quad \mathbf{c} := \frac{1}{1 + \partial_w \sigma} [\Delta_{\mathbf{x}'} \sigma + \mathbf{b} \cdot \nabla_{\mathbf{x}'} \partial_w \sigma + \mathbf{a} \partial_w^2 \sigma].$$

Moreover, if we denote by J_Σ the Jacobian matrix of Σ , we denote by $P(\Sigma)$ the symmetric matrix given by

$$(3.2) \quad P(\Sigma) := |\det(J_\Sigma)| J_\Sigma^{-1} J_\Sigma^{-T} = (1 + \partial_w \sigma) \begin{pmatrix} \mathbb{I}_2 & \frac{-\nabla_{\mathbf{x}'} \sigma}{1 + \partial_w \sigma} \\ \frac{-(\nabla_{\mathbf{x}'} \sigma)^T}{1 + \partial_w \sigma} & \frac{1 + |\nabla_{\mathbf{x}'} \sigma|^2}{(1 + \partial_w \sigma)^2} \end{pmatrix},$$

so that

$$\Delta^\Sigma \tilde{f} = (1 + \partial_w \sigma)^{-1} \nabla_{\mathbf{x}', w} \cdot P(\Sigma) \nabla_{\mathbf{x}', w} \tilde{f},$$

see Lemma 2.5 in [Lan05].

From the above definitions and from Proposition 1.4 it follows that

Proposition 3.2. *Let $k > \frac{5}{2}$, let $\eta \in H^k(\mathbb{R}^2)$ and let σ be a regularizing diffeomorphism be such that (1.16) holds true, and let us denote by D_η the domain as in (1.15). Let also $\omega \in H_b(\operatorname{div}_0, D_\eta)$ and $\Phi \in \dot{H}^{3/2}(\mathbb{R}^2)$.*

Then there exists a unique $U \in (H^1(D_\eta))^3$ which solves the boundary value problem (1.17). Moreover, $\tilde{U} = U \circ \Sigma \in (H^1(D_0))^3$ solves

$$(3.3) \quad \begin{cases} \operatorname{curl}^\Sigma \tilde{U} = \tilde{\omega}, & \text{in } D_0, \\ \operatorname{div}^\Sigma \tilde{U} = 0 & \text{in } D_0, \\ \tilde{U} \cdot e_3 = 0, & \text{at } w = -h, \\ \tilde{U}_\parallel = \nabla \Phi - \nabla^\perp \Delta^{-1}(\tilde{\omega} \cdot N), & \text{at } w = 0. \end{cases}$$

Conversely, if $\tilde{U} \in (H^1(D_0))^3$ is the unique solution to the boundary value problem (3.3), then $U = \tilde{U} \circ \Sigma^{-1} \in (H^1(D_\eta))^3$ is the unique solution to the boundary value problem (1.17).

From Proposition 1.4 one obtains the following estimate on \tilde{U} ,

$$(3.4) \quad \begin{aligned} & \|\tilde{U}\|_{(L^2(D_0))^3} + \|\nabla^\Sigma \tilde{U}\|_{(L^2(D_0))^{3 \times 3}} \\ & \leq C(h_0^{-1}, h, \|\eta\|_{H^k(\mathbb{R}^2)}) \left(\|\tilde{\omega}\|_{2,b} + \|\nabla \Phi\|_{(H^{1/2}(\mathbb{R}^2))^2} \right), \end{aligned}$$

where we point out that Σ is a regularizing diffeomorphism (see Appendix A; see also Sec. 3.1.1 of [ABZ14] and Sec.4.2 of [WZZZ21]). We also remind that if $\eta \in C^2(\mathbb{R}^2)$, then Theorem 2.4 of [CL15] implies that the estimate (3.4) holds true with $\|\eta\|_{H^k(\mathbb{R}^2)}$ replaced by $\|\eta\|_{C^2(\mathbb{R}^2)}$.

4. COROLLARIES OF PROPOSITION 1.4

We now discuss some consequences of Proposition 1.4. Observe that Eq. (1.20) provides a decomposition of the velocity field into a rotational part and an irrotational part. We also mention an immediate corollary of the above Proposition 1.4 (see Corollary 3.9 in [CL15]), which provides an inverse of the curl operator on the space of divergence-free vector fields.

Corollary 4.1. *Let $k > \frac{5}{2}$, let $\eta \in H^k(\mathbb{R}^2)$ be such that (1.16) holds true, and let us denote by D_η the domain as in (1.15). Let also $C \in H(\operatorname{div}_0, D_\eta)$ be such that $C \cdot N|_{z=-h} = 0$.*

Then there exists a unique $B \in (H^1(D_\eta))^3$ which solves the boundary value problem

$$\begin{aligned} \operatorname{curl} B &= C, & \text{in } D_\eta, \\ \operatorname{div} B &= 0, & \text{in } D_\eta, \\ B &= \mathbf{0}, & \text{at } z = -h, \\ B_\parallel &= -\nabla^\perp \Delta^{-1}(C \cdot N), & \text{at } z = \eta, \end{aligned}$$

and we denote this solution by $B = \operatorname{curl}^{-1} C$.

From Proposition 1.4 and Remark B.11 we can immediately derive the following result on the generalized Dirichlet–Neumann operator defined in (1.25).

Corollary 4.2. *Let $k > \frac{5}{2}$, let $\eta \in H^k(\mathbb{R}^2)$ be such that (1.16) holds true. Then for any $\Phi \in \dot{H}^{3/2}(\mathbb{R}^2)$ and for any $\omega \in H_b(\operatorname{div}_0, D_\eta)$ we have*

$$\mathcal{G}_{gen}[\eta](\Phi, \omega) \in H^{1/2}(\mathbb{R}^2),$$

and the following estimate holds true,

$$(4.1) \quad \begin{aligned} & \|\mathcal{G}_{gen}[\eta](\Phi, \omega)\|_{H^{1/2}(\mathbb{R}^2)} \\ & \leq C(h_0^{-1}, h, \|\eta\|_{H^k(\mathbb{R}^2)}) \left(\|\omega\|_{2,b} + \|\nabla \Phi\|_{H^{1/2}(\mathbb{R}^2)} \right). \end{aligned}$$

We also reformulate the boundary value problem (1.21) in an equivalent way, along the lines of the approach outlined in Sec. 4 of [GH20] for Beltrami flows. This allows us to obtain a solution of (1.21) by solving (the flattened version of) an equivalent problem, whose solution can be explicitly written in terms of a Green matrix (see Sec. 6).

Corollary 4.3. *Let $k > \frac{5}{2}$, let $\eta \in H^k(\mathbb{R}^2)$ be such that (1.16) holds true. Let $\omega \in H_b(\text{div}_0, D_\eta)$ and $\Phi \in \dot{H}^{3/2}(\mathbb{R}^2)$, then the strong solutions of the boundary value problem (1.21) and of the boundary value problem*

$$(4.2) \quad \begin{cases} -\Delta \mathbf{A} = \omega, & \text{in } D_\eta, \\ A_{3z} = 0, & \text{at } z = -h, \\ \mathbf{A} \times \mathbf{e}_3 = \mathbf{0}, & \text{at } z = -h, \\ \mathbf{A} \cdot \mathbf{n} = 0, & \text{at } z = \eta, \\ (\text{curl } \mathbf{A})_{\parallel} = -\nabla^\perp \Delta^{-1}(\omega \cdot \mathbf{N}), & \text{at } z = \eta, \end{cases}$$

coincide.

Proof. Suppose that $\mathbf{A} \in (H^2(D_\eta))^3$ is a strong solution of (1.21), so that \mathbf{A} satisfies

$$-\Delta \mathbf{A} = \omega, \quad \text{in } D_\eta,$$

(see Remark 1.5) and $\text{div } \mathbf{A}|_{z=-h} = 0$.

Since $\mathbf{A}_h|_{z=-h} = 0$ and $\nabla \cdot \mathbf{A}_h|_{z=-h} = 0$, we can deduce that $A_{3z}|_{z=-h} = \text{div } \mathbf{A} - \nabla \cdot \mathbf{A}_h|_{z=-h} = 0$.

Conversely, any strong solution $\mathbf{A} \in (H^2(D_\eta))^3$ of (4.2) satisfies $\text{div } \mathbf{A}|_{z=-h} = 0$; we have to show that $\text{div } \mathbf{A} = 0$ in D_η . Since

$$-\Delta \mathbf{A} = \text{curl curl } \mathbf{A} - \nabla_{\mathbf{x}',z} \text{div } \mathbf{A},$$

taking the scalar product of the interior equation in (4.2) with a function $\mathbf{B} \in \mathfrak{X}_\eta$ and integrating by parts using the integral identity

$$\int_{D_\eta} \nabla_{\mathbf{x}',z} \text{div } \mathbf{A} \cdot \mathbf{B} = - \int_{D_\eta} \text{div } \mathbf{A} \text{div } \mathbf{B},$$

$$\forall \mathbf{A}, \mathbf{B} \text{ s.t. } \text{div } \mathbf{A}|_{z=-h} = 0, \quad \mathbf{B} \times \mathbf{e}_3|_{z=-h} = \mathbf{0}, \quad \underline{\mathbf{B}} \cdot \mathbf{n} = 0,$$

we can see that \mathbf{A} satisfies (A.10), hence \mathbf{A} is a weak solution of (1.21), so that

$$\text{div } \mathbf{A} = 0, \quad \text{in } D_\eta.$$

□

5. REGULARITY OF THE VELOCITY FIELD

All results for the velocity field \mathbf{U} have an analogue for the straightened field

$$\tilde{\mathbf{U}}[\eta](\Phi, \tilde{\omega}) := \mathbb{U}[\eta](\Phi, \omega) \circ \Sigma,$$

where we introduce the decomposition

$$\begin{aligned} \tilde{\mathbf{U}}[\eta](\Phi, \tilde{\omega}) &:= \tilde{\mathbf{U}}_I[\eta]\Phi + \tilde{\mathbf{U}}_{II}[\eta]\tilde{\omega}, \\ \tilde{\mathbf{U}}_I[\eta]\Phi &:= \mathbb{U}_I[\eta]\Phi \circ \Sigma, \quad \tilde{\mathbf{U}}_{II}[\eta]\tilde{\omega} := \mathbb{U}_{II}[\eta]\omega \circ \Sigma. \end{aligned}$$

Similarly, we define the straightened version of the generalized Dirichlet–Neumann operator defined in (1.25),

$$\begin{aligned} \tilde{\mathcal{G}}_{gen}[\eta](\Phi, \tilde{\omega}) &:= \tilde{\mathcal{G}}_{gen,I}[\eta]\Phi + \tilde{\mathcal{G}}_{gen,II}[\eta]\tilde{\omega}, \\ \tilde{\mathcal{G}}_{gen,I}[\eta]\Phi &:= \tilde{\mathbf{U}}_I[\eta]\Phi \cdot \mathbf{N}|_{w=0}, \quad \tilde{\mathcal{G}}_{gen,II}[\eta]\tilde{\omega} := \tilde{\mathbf{U}}_{II}[\eta]\tilde{\omega} \cdot \mathbf{N}|_{w=0}. \end{aligned}$$

For example, we have that

$$\int_{D_\eta} \nabla_{\mathbf{x}',z} \mathbf{U} \cdot \nabla_{\mathbf{x}',z} \mathbf{C} = \int_{D_\eta} \omega \cdot \text{curl } \mathbf{C} + \int_{\mathbb{R}^2} \mathbf{F} \cdot \underline{\mathbf{C}}, \quad \forall \mathbf{C} \in (H^1(D_\eta))^3,$$

$$\mathbf{F} := \mathbf{N} \cdot \underline{\nabla_{\mathbf{x}',z} \mathbf{U}} + \mathbf{N} \times \underline{\text{curl } \mathbf{U}}.$$

This is equivalent to

$$(5.1) \quad \int_{D_0} \nabla^\Sigma \tilde{U} \cdot P(\Sigma) \nabla^\Sigma \tilde{C} = \int_{D_0} (1 + \partial_w \sigma) \tilde{\omega} \cdot \text{curl}^\Sigma \tilde{C} + \int_{\mathbb{R}^2} \mathbf{F} \cdot \underline{\mathbf{C}},$$

for all $\tilde{C} \in (H^1(D_0))^3$, where we recall that $P(\Sigma) = (1 + \partial_w \sigma) J_\Sigma^{-1} (J_\Sigma^{-1})^T$.

Recalling that $\Lambda = (1 - \Delta)^{1/2}$ and that the Sobolev norm $\|\cdot\|_{H^{k,\ell}(D_0)}$ controls at most ℓ vertical derivatives ($\ell \leq k$), we recall the definition of the *good unknowns* (2.1)-(2.3),

$$\begin{aligned} \tilde{\varphi}_{(j)} &= \partial^j \tilde{\varphi} - (\partial^j \sigma) \partial_w^\Sigma \tilde{\varphi}, \quad j \in \mathbb{N}^2 \setminus \{\mathbf{0}\}, \\ \Phi_{(j)} &= \partial^j \Phi - \tilde{\mathbf{w}}_I \partial^j \eta, \quad j \in \mathbb{N}^2 \setminus \{\mathbf{0}\}, \quad \tilde{\mathbf{w}}_I := \tilde{\mathbb{W}}_I[\eta] \Phi|_{w=0}, \\ \tilde{A}_{(j)} &= \partial^j \tilde{A} - (\partial^j \sigma) \partial_w^\Sigma \tilde{A}, \quad j \in \mathbb{N}^2 \setminus \{\mathbf{0}\}. \end{aligned}$$

Remark 5.1. We mention that in (3.22) of [CL15] the authors define for all $j \in \mathbb{N}^2 \setminus \{\mathbf{0}\}$ the good unknowns $\Phi_{(j)} := \partial^j \Phi - \tilde{\mathbf{w}} \partial^j \eta$. On the other hand, we introduce the good unknowns (2.1)-(2.3) in order to better highlight the roles of the rotational and irrotational parts of the velocity field.

Moreover, we define

$$\mathfrak{P} := \frac{|\mathbf{D}|}{(1 + |\mathbf{D}|)^{1/2}};$$

the operator \mathfrak{P} appears already in the irrotational case, see for example Lemma 2.34 in [Lan13]. We state the following result (see Proposition 3.12 in [CL15]), which is a generalization of Proposition 1.4. In the rest of the section, we use the convention that a summation over an empty set is equal to zero.

Theorem 5. Let $N > 3/2$, let $\eta \in C^N(\mathbb{R}^2)$ be such that (1.16) holds true. Assume that the assumptions of Proposition 1.4 hold true, and that σ is a regularizing diffeomorphism such that (2.4) holds true. Then there exists a unique solution $\tilde{U} \in (H^1(D_0))^3$ which solves the boundary value problem (3.3). If $0 \leq k < N - 3/2$, $k \in \mathbb{N}$, $\Phi \in \dot{H}^{k+3/2}(\mathbb{R}^2)$, $\Lambda^k \tilde{\omega} \in (L^2(D_0))^3$, then

$$(5.2) \quad \begin{aligned} \|\tilde{U}\|_{(H^{k+1,1}(D_0))^3} &\leq C(h_0^{-1}, h, \|\eta\|_{C^N(\mathbb{R}^2)}) \times \\ &\times \left[\|\mathfrak{P}\Phi\|_{H^1(\mathbb{R}^2)} + \sum_{1 < |j| \leq k+1} \|\mathfrak{P}\Phi_{(j)}\|_{L^2(\mathbb{R}^2)} + \|\Lambda^k \tilde{\omega}\|_{2,b} \right]. \end{aligned}$$

If, moreover, $N > 3$, $0 \leq \ell \leq k < N - 3/2$, and $\tilde{\omega} \in (H^{k,\ell}(D_0))^3$, then the following higher-order estimate hold

$$(5.3) \quad \begin{aligned} \|\tilde{U}\|_{(H^{k+1,\ell+1}(D_0))^3} &\leq C(h_0^{-1}, h, \|\eta\|_{C^N(\mathbb{R}^2)}) \times \\ &\times \left[\|\mathfrak{P}\Phi\|_{H^1(\mathbb{R}^2)} + \sum_{1 < |j| \leq k+1} \|\mathfrak{P}\Phi_{(j)}\|_{L^2(\mathbb{R}^2)} \right. \\ &\left. + \|\tilde{\omega}\|_{(H^{k,\ell}(D_0))^3} + \|\Lambda^k(\tilde{\omega} \cdot \mathbf{e}_3)|_{w=-h}\|_{H_0^{-1/2}(\mathbb{R}^2)} \right]. \end{aligned}$$

Proof. We start by taking $\tilde{C} = \partial^{2j} \tilde{U}$ in (5.1), with $|j| = k < N - 3/2$; this is a slight abuse of notation, as mentioned in the proof of Proposition 3.12 in [CL15], because one should actually take $\tilde{C} = \chi(\delta w |\mathbf{D}|) \partial^{2j} \tilde{U}$ where $\delta > 0$ and $\chi \in C_c^\infty(\mathbb{R}^2)$, $\chi \equiv 1$ in a neighbourhood of the origin, and then deduce that the estimates are uniform as $\delta \rightarrow 0$, see for example Lemma 2.34 in [Lan13].

We obtain

$$\begin{aligned} & \int_{D_0} \nabla^\Sigma \tilde{U} \cdot P(\Sigma) \nabla^\Sigma \partial^{2j} \tilde{U} \\ &= \int_{D_0} (1 + \partial_w \sigma) \tilde{\omega} \cdot \operatorname{curl}^\Sigma \partial^{2j} \tilde{U} + \int_{\mathbb{R}^2} \tilde{F} \cdot \partial^{2j} \tilde{U}|_{w=0}, \end{aligned}$$

which, by Plancherel theorem, is equivalent to

$$\begin{aligned} \int_{D_0} \nabla^\Sigma \partial^j \tilde{U} \cdot P(\Sigma) \nabla^\Sigma \partial^j \tilde{U} &= I_1 + I_2 + I_3, \\ I_1 &:= \int_{D_0} \nabla^\Sigma \partial^j \tilde{U} \cdot [\partial^j, P(\Sigma)] \nabla^\Sigma \tilde{U}, \\ I_2 &:= \int_{D_0} \Lambda^k \tilde{\omega} \cdot \Lambda^{-k} \left((1 + \partial_w \sigma) \operatorname{curl}^\Sigma \partial^{2j} \tilde{U} \right), \\ I_3 &:= \int_{\mathbb{R}^2} \partial^j \tilde{F} \cdot \partial^j \tilde{U}|_{w=0}. \end{aligned}$$

By Lemma B.10 and by condition (2.4) we have that for any $0 < \varepsilon \ll 1$ the following estimate holds true,

$$\left\| \partial^j \nabla_{\mathbf{x}', w} \tilde{U} \right\|_{(L^2(D_0))^3}^2 \leq C \left(h_0^{-1}, \|\eta\|_{C^{1+\varepsilon}(\mathbb{R}^2)} \right) (|I_1| + |I_2| + |I_3|).$$

Regarding I_1 , if we denote by $Q(\Sigma) := P(\Sigma) - \mathbb{I}_3$, we have again by Lemma B.10 and by (2.4) that for any $\varepsilon > 0$ such that $N \geq k + 1 + \varepsilon$

$$\begin{aligned} |I_1| &\leq \|\nabla^\Sigma \partial^j \tilde{U}\|_{(L^2(D_0))^3} \|\partial^j, Q(\Sigma)\| \nabla^\Sigma \tilde{U} \|_{(L^2(D_0))^3} \\ &\leq C \|\nabla^\Sigma \partial^j \tilde{U}\|_{(L^2(D_0))^3} \|Q(\Sigma)\|_{M_3(C^{k+\varepsilon}(D_0))} \|\Lambda^{k-1} \nabla^\Sigma \tilde{U}\|_{(L^2(D_0))^3} \\ (5.4) \quad &\leq C(h_0^{-1}, h, \|\eta\|_{C^N(\mathbb{R}^2)}) \|\nabla^\Sigma \partial^j \tilde{U}\|_{(L^2(D_0))^3} \|\Lambda^{k-1} \nabla^\Sigma \tilde{U}\|_{(L^2(D_0))^3}. \end{aligned}$$

Next, we estimate I_2 . We have by Lemma B.10 that for any $\varepsilon > 0$ such that $N \geq k + 1 + \varepsilon$

$$\begin{aligned} & \|\Lambda^{-k} \left((1 + \partial_w \sigma) \operatorname{curl}^\Sigma \partial^{2j} \tilde{U} \right)\|_{(L^2(D_0))^3} \\ & \leq C \left(h_0^{-1}, \|\nabla_{\mathbf{x}', w} \sigma\|_{(C^{k+\varepsilon}(D_0))^3} \right) \|\partial^j \nabla^\Sigma \tilde{U}\|_{(L^2(D_0))^3}, \end{aligned}$$

so that by (2.4)

$$(5.5) \quad |I_2| \leq C \left(h_0^{-1}, \|\eta\|_{C^N(\mathbb{R}^2)} \right) \|\Lambda^k \tilde{\omega}\|_{(L^2(D_0))^3} \|\partial^j \nabla^\Sigma \tilde{U}\|_{(L^2(D_0))^3}.$$

Finally, we estimate I_3 . We have

$$I_3 = \int_{\mathbb{R}^2} \partial^j \left[\mathbf{N} \cdot \nabla^\Sigma \tilde{\mathbf{v}}|_{w=0} + \mathbf{N} \times \operatorname{curl}^\Sigma \tilde{\mathbf{v}}|_{w=0} \right] \cdot \partial^j \tilde{U}|_{w=0}.$$

We decompose I_3 in the following way: we denote $\tilde{U}_h|_{w=0} = \underline{\tilde{v}}$, $\tilde{u}_3|_{w=0} = \underline{\tilde{w}}$, and we write

$$\begin{aligned} I_3 &= I_{31} + I_{32} + I_{33}, \\ I_{31} &:= \int_{\mathbb{R}^2} 2\partial^j \underline{\tilde{v}} \cdot \partial^j \nabla \underline{\tilde{w}}, \quad I_{32} := - \int_{\mathbb{R}^2} (\nabla^\perp \eta \cdot \nabla) \partial^j \underline{\tilde{v}}^\perp \cdot \partial^j \underline{\tilde{v}}, \\ I_{33} &:= - \int_{\mathbb{R}^2} [\partial^j, \nabla^\perp \eta] \cdot \nabla \underline{\tilde{v}}^\perp \cdot \partial^j \underline{\tilde{v}}. \end{aligned}$$

Next, we estimate I_{31} : by writing $\partial^j \tilde{\mathbf{v}} = \partial^j (\nabla \Phi - \tilde{\mathbf{w}}_I \nabla \eta) + \partial^j (\nabla^\perp \Psi - \tilde{\mathbf{w}}_{II} \nabla \eta)$, we obtain

$$\begin{aligned} I_{31} &= \int_{\mathbb{R}^2} 2 \left[\partial^j (\nabla \Phi - \tilde{\mathbf{w}}_I \nabla \eta) + \partial^j (\nabla^\perp \Psi - \tilde{\mathbf{w}}_{II} \nabla \eta) \right] \cdot \partial^j \nabla \tilde{\mathbf{w}} \\ &= 2 \int_{\mathbb{R}^2} (\nabla \partial^j \Phi - \tilde{\mathbf{w}}_I \nabla \partial^j \eta) \cdot \partial^j \nabla \tilde{\mathbf{w}} \\ &\quad + 2 \int_{\mathbb{R}^2} (-\partial^j \tilde{\mathbf{w}}_{II} \nabla \eta - [\partial^j, \tilde{\mathbf{w}}_{II}, \nabla \eta]) \cdot \nabla \partial^j \tilde{\mathbf{w}} \\ &\quad + 2 \int_{\mathbb{R}^2} \partial^j ((\text{curl}^\Sigma \tilde{\mathbf{A}})_\parallel - \tilde{\mathbf{w}}_{II} \nabla \eta) \cdot \nabla \partial^j \tilde{\mathbf{w}}, \end{aligned}$$

where $[\partial^j, f, g] := \partial^j (fg) - (\partial^j f)g - f \partial^j g$, and where the term involving $\nabla^\perp \Psi$ has been rewritten using (1.21).

Recalling the definition of the operator \mathfrak{P} , we now exploit the following product estimate

$$(5.6) \quad \int_{\mathbb{R}^2} f \partial_j g \leq C \|\mathfrak{P}f\|_{L^2(\mathbb{R}^2)} \|g\|_{H^{1/2}(\mathbb{R}^2)}, \quad j = x, y, \quad \forall f, g \in H^{1/2}(\mathbb{R}^2),$$

see (3.38) in [CL15].

Indeed, by using Lemma B.10 and by condition (2.4) we have that for any $\varepsilon > 0$ such that $N \geq k + 3/2 + \varepsilon$

$$\begin{aligned} |I_{31}| &\leq 2 \|\Lambda^k \tilde{\mathbf{w}}\|_{H^{1/2}(\mathbb{R}^2)} \times \\ &\quad \times \left[\|\mathfrak{P}(\nabla \partial^j \Phi - \tilde{\mathbf{w}}_I \nabla \partial^j \eta)\|_{(L^2(\mathbb{R}^2))^2} + \|\mathfrak{P}(\partial^j \tilde{\mathbf{w}}_{II} \nabla \eta)\|_{(L^2(\mathbb{R}^2))^2} \right] \\ &\quad + \|\nabla [\partial^j, \tilde{\mathbf{w}}_{II}, \nabla \eta]\|_{(L^2(\mathbb{R}^2))^2} \|\Lambda^k \tilde{\mathbf{w}}\|_{L^2(\mathbb{R}^2)} \\ &\quad + \|\mathbf{D}^{k-1} (\text{curl}^\Sigma \tilde{\mathbf{A}})_\parallel\|_{(H^{1/2}(\mathbb{R}^2))^2} \|\mathfrak{P} \Lambda^k \tilde{\mathbf{w}}\|_{L^2(\mathbb{R}^2)} \\ &\leq C (h_0^{-1}, h, \|\eta\|_{C^N(\mathbb{R}^2)}) \|\Lambda^k \nabla^\Sigma \tilde{\mathbf{U}}\|_{(L^2(D_0))^3} \times \\ &\quad \times \left[\|\mathfrak{P}(\nabla \partial^j \Phi - \tilde{\mathbf{w}}_I \nabla \partial^j \eta)\|_{(L^2(\mathbb{R}^2))^2} + \|\Lambda^k \tilde{\omega}\|_{2,b} \right. \\ &\quad \left. + \|\Lambda^{k-1} \nabla^\Sigma \tilde{\mathbf{U}}\|_{(L^2(D_0))^3} \right]. \end{aligned}$$

Similarly, under the same assumptions one can prove that

$$\begin{aligned} |I_{32}| &\leq C (h_0^{-1}, h, \|\eta\|_{C^N(\mathbb{R}^2)}) \|\Lambda^k \nabla^\Sigma \tilde{\mathbf{U}}\|_{(L^2(D_0))^3} \times \\ &\quad \times \left[\|\mathfrak{P}(\nabla \partial^j \Phi - \tilde{\mathbf{w}}_I \nabla \partial^j \eta)\|_{L^2(\mathbb{R}^2)} + \|\mathfrak{P} \partial^j \nabla^\perp \Psi\|_{L^2(\mathbb{R}^2)} \right. \\ &\quad \left. + \|\Lambda^{k-1} \nabla^\Sigma \tilde{\mathbf{U}}\|_{(L^2(D_0))^3} \right], \\ |I_{33}| &\leq C (h_0^{-1}, h, \|\eta\|_{C^N(\mathbb{R}^2)}) \|\Lambda^{k-1} \nabla^\Sigma \tilde{\mathbf{U}}\|_{(L^2(D_0))^3} \|\Lambda^k \nabla^\Sigma \tilde{\mathbf{U}}\|_{(L^2(D_0))^3}, \end{aligned}$$

(see proof of Proposition 3.12 in [CL15]), so that by combining the estimates for I_{31} , for I_{32} and for I_{33} together with the following estimate (see Lemma 3.13 of [CL15])

$$\|\Lambda^k \nabla \tilde{\Psi}\|_{H^{1/2}(\mathbb{R}^2)} \leq C(h, h_0^{-1}, \|\eta\|_{C^N(\mathbb{R}^2)}) \|\Lambda^k \tilde{\omega}\|_{2,b},$$

we derive that

$$(5.7) \quad \begin{aligned} |I_3| &\leq C(h_0^{-1}, \|\eta\|_{C^N(\mathbb{R}^2)}) \|\Lambda^k \nabla^\Sigma \tilde{U}\|_{(L^2(D_0))^3} \times \\ &\quad \times \left[\|\mathfrak{P}(\nabla \partial^j \Phi - \tilde{\omega} \nabla \partial^j \eta)\|_{L^2(\mathbb{R}^2)} + \|\Lambda^k \tilde{\omega}\|_{2,b} \right. \\ &\quad \left. + \|\Lambda^{k-1} \nabla^\Sigma \tilde{U}\|_{(L^2(D_0))^3} \right]. \end{aligned}$$

Combining (5.4), (5.5) and (5.7), and using (5.1) and (3.4), we have that for any $\varepsilon > 0$ such that $N \geq k + 3/2 + \varepsilon$ the bound (5.2) holds true.

Finally, observe that for all $N > 3$ and $1 \leq m < N - 1$ there exists $C > 0$ such that for any $\varepsilon > 0$ such that $m + \varepsilon \leq N - 1$

$$(5.8) \quad \begin{aligned} \|\Lambda^{m-j} \partial_z^j (fg)\|_{L^2(D_0)} &\leq C \|f\|_{C^{m+\varepsilon}(D_0)} \|g\|_{H^{m,j}(D_0)}, \quad \forall j \in \{0, \dots, m\}, \\ \forall f &\in C^{N-1}(D_0), \quad \forall g \in H^{m,j}(D_0), \end{aligned}$$

(this is a consequence of Lemma B.10; see Lemma 3.15 of [CL15]). Let $1 \leq \ell \leq k$: combining the identity

$$\partial_w^\Sigma \tilde{U} = \frac{1}{1 + |\nabla \eta|^2} \left[(\mathbf{N} \cdot \partial_w^\Sigma \tilde{U}) \mathbf{N} + (\mathbf{N} \times \partial_w^\Sigma \tilde{U}) \times \mathbf{N} \right]$$

with the above product rule (5.8) and with condition (2.4) we obtain

$$\|\partial_w \tilde{U}\|_{(H^{k,\ell}(D_0))^3} \leq C(\|\eta\|_{C^N(\mathbb{R}^2)}) \left[\|\tilde{U}\|_{(H^{k+1,\ell}(D_0))^3} + \|\tilde{\omega}\|_{(H^{k,\ell}(D_0))^3} \right],$$

hence

$$\|\tilde{U}\|_{(H^{k+1,\ell+1}(D_0))^3} \leq C(\|\eta\|_{C^N(\mathbb{R}^2)}) \left[\|\tilde{U}\|_{(H^{k+1,\ell}(D_0))^3} + \|\tilde{\omega}\|_{(H^{k,\ell}(D_0))^3} \right],$$

and by finite induction on ℓ we get

$$\|\tilde{U}\|_{(H^{k+1,\ell+1}(D_0))^3} \leq C(\|\eta\|_{C^N(\mathbb{R}^2)}) \left[\|\tilde{U}\|_{(H^{k+1,1}(D_0))^3} + \|\tilde{\omega}\|_{(H^{k,\ell}(D_0))^3} \right],$$

so that we can deduce (5.3). \square

Remark 5.2. We point out that the assumption $\Phi \in \dot{H}^{k+3/2}(\mathbb{R}^2)$ can be weakened; like in Chap. 2 of [Lan13] and Theorem 2.4 of [CL15], it suffices to assume that

$$\|\mathfrak{P}\Phi_{(j)}\|_{L^2(\mathbb{R}^2)} < +\infty, \quad 1 < \alpha \leq k + 1.$$

Remark 5.3. In the corresponding result by Castro and Lannes (see Proposition 3.12 in [CL15]) the control of the $H^{k,\ell}$ -norm of \tilde{U} is given in terms of the norms of the good unknowns $\Phi_{(j)}$, see Remark 5.1 regarding the different definition of the good unknowns.

From Theorem 5 and Remark B.11 we can immediately derive the following higher order estimates on the generalized Dirichlet–Neumann operator defined in (1.25).

Corollary 5.4. Let $N > 3/2$, let $\eta \in C^N(\mathbb{R}^2)$ be such that (1.16) holds true. Assume that $0 \leq k < N - 3/2$, $k \in \mathbb{N}$, $\Phi \in \dot{H}^{k+3/2}(\mathbb{R}^2)$, $\tilde{\omega} \in (H^{k,0}(D_0))^3$ and that σ is a regularizing diffeomorphism such that (2.4) holds true. Then

$$\tilde{\mathcal{G}}_{gen}[\eta](\Phi, \tilde{\omega}) \in H^{k+1/2}(\mathbb{R}^2),$$

and the following higher-order estimate holds true,

$$(5.9) \quad \begin{aligned} & \left\| \tilde{\mathcal{G}}_{gen}[\eta](\Phi, \tilde{\omega}) \right\|_{H^{k+1/2}(\mathbb{R}^2)} \leq C(h_0^{-1}, h, \|\eta\|_{C^N(\mathbb{R}^2)}) \times \\ & \times \left[\|\mathfrak{P}\Phi\|_{H^1(\mathbb{R}^2)} + \sum_{1 < |j| \leq k+1} \|\mathfrak{P}\Phi_{(j)}\|_{L^2(\mathbb{R}^2)} \right. \\ & \left. + \|\tilde{\omega}\|_{(H^{k,0}(D_0))^3} + \|\Lambda^k(\tilde{\omega} \cdot \mathbf{e}_3)|_{w=-h}\|_{H_0^{-1/2}(\mathbb{R}^2)} \right]. \end{aligned}$$

Finally, we mention the following higher order estimate; it can be regarded as a slight modification of Corollary 3.14 in [CL15], and it can be proved by following the same argument.

Corollary 5.5. *Let $s \in \mathbb{N}$, $s \geq 5$, and let $\eta \in H^s(\mathbb{R}^2)$ be such that (1.16) holds true, $\omega \in H_b(\text{div}_0, D_\eta)$, $\Phi \in \dot{H}^{3/2}(\mathbb{R}^2)$. Moreover, assume that $0 \leq k \leq s-1$,*

$$\Phi_{(j)} \in \dot{H}^{1/2}(\mathbb{R}^2), \quad \forall 1 < |j| \leq k+1,$$

$\tilde{\omega} \in (H^{k,0}(D_0))^3$ and that

$$\|\nabla_{\mathbf{x}', w} \sigma\|_{L^\infty([-h, 0]; (H^k(\mathbb{R}^2))^3)} \leq C \|\eta\|_{H^s(\mathbb{R}^2)}$$

holds true. Then

$$\tilde{\mathcal{U}}[\eta](\Phi, \tilde{\omega}) \in (H^{k+1,1}(D_0))^3,$$

and the following higher-order estimate holds true,

$$(5.10) \quad \begin{aligned} & \left\| \tilde{\mathcal{U}}[\eta](\Phi, \tilde{\omega}) \right\|_{(H^{k+1,1}(D_0))^3} \\ & \leq C(h_0^{-1}, h, \|\eta\|_{H^s(\mathbb{R}^2)}) \times \\ & \times \left[\|\mathfrak{P}\Phi\|_{H^1(\mathbb{R}^2)} + \sum_{1 < |j| \leq k+1} \|\mathfrak{P}\Phi_{(j)}\|_{L^2(\mathbb{R}^2)} \right. \\ & \left. + \|\tilde{\omega}\|_{(H^{k,0}(D_0))^3} + \|\Lambda^k(\tilde{\omega} \cdot \mathbf{e}_3)|_{w=-h}\|_{H_0^{-1/2}(\mathbb{R}^2)} \right]. \end{aligned}$$

If in addition we have $0 \leq k \leq s-3/2$, then

$$\tilde{\mathcal{G}}_{gen}[\eta](\Phi, \tilde{\omega}) \in H^{k+1/2}(\mathbb{R}^2),$$

and the following estimate holds true,

$$(5.11) \quad \begin{aligned} & \left\| \tilde{\mathcal{G}}_{gen}[\eta](\Phi, \tilde{\omega}) \right\|_{H^{k+1/2}(\mathbb{R}^2)} \\ & \leq C(h_0^{-1}, h, \|\eta\|_{H^s(\mathbb{R}^2)}) \times \\ & \times \left[\|\mathfrak{P}\Phi\|_{H^1(\mathbb{R}^2)} + \sum_{1 < |j| \leq k+1} \|\mathfrak{P}\Phi_{(j)}\|_{L^2(\mathbb{R}^2)} \right. \\ & \left. + \|\tilde{\omega}\|_{(H^{k,0}(D_0))^3} + \|\Lambda^k(\tilde{\omega} \cdot \mathbf{e}_3)|_{w=-h}\|_{H_0^{-1/2}(\mathbb{R}^2)} \right]. \end{aligned}$$

Remark 5.6. *In the statement of Corollary (5.5) we can replace the assumption on the good unknowns $\Phi_{(j)}$ and on $\tilde{\omega}$ with the assumptions $\Phi \in \dot{H}^{k+3/2}(\mathbb{R}^2)$ and $\tilde{\omega} \in (H^k(D_0))^3$, see also Remark 5.2 and Sec. 2.4 of [Lan13].*

6. ANALITICITY OF THE GENERALIZED DIRICHLET–NEUMANN OPERATOR

In this section we prove an analyticity result for the generalized Dirichlet–Neumann operator; we also show a formula for the differential of its rotational part. Moreover, as for the classical Dirichlet–Neumann operator for irrotational flows (see [CS93] [CG94]) and for Beltrami flows (see [GNPW24]), one can write the Taylor expansion of $\tilde{\mathcal{G}}_{gen}[\eta]$ around $\eta = 0$,

$$(6.1) \quad \tilde{\mathcal{G}}_{gen}[\eta] = \sum_{j=0}^{\infty} \tilde{\mathcal{G}}_j[\eta],$$

where $\tilde{\mathcal{G}}_j[\eta]$ is homogeneous of degree j in η , and for $j \geq 1$ the terms $\tilde{\mathcal{G}}_j[\eta]$ may be computed systematically by using a recursion formula.

Indeed, from (1.25) and from Remark 1.2 we have

$$\mathcal{G}_{gen}[\eta](\Phi, \boldsymbol{\omega}) = \nabla \cdot \mathbf{A}_{\parallel}^{\perp} + \underline{\nabla_{\mathbf{x}', z} \varphi} \cdot \mathbf{N}.$$

We look for a scalar function φ and for a vector potential \mathbf{A} which solve the systems (1.21) and (1.22) respectively. While the uniqueness of the solution of the boundary value problem (1.22) is well-known (see Proposition 2.9 and Corollary 2.44 in [Lan13]), we now exploit Corollary 4.3 in order to obtain a solution of (1.21) by solving (the flattened version of) an equivalent problem, whose solution can be explicitly written in terms of a Green matrix (see Sec.4(c) of [GH20] for a similar approach).

From (1.25) we recall the decomposition

$$\begin{aligned} \tilde{\mathcal{G}}_{gen}[\eta](\Phi, \tilde{\boldsymbol{\omega}}) &= \tilde{\mathcal{G}}_{gen, I}[\eta]\Phi + \tilde{\mathcal{G}}_{gen, II}[\eta]\tilde{\boldsymbol{\omega}}, \\ \tilde{\mathcal{G}}_{gen, I}[\eta]\Phi &:= \tilde{\mathcal{U}}_I[\eta]\Phi \cdot \mathbf{N}|_{w=0} = \nabla^{\Sigma} \tilde{\varphi} \cdot \mathbf{N}|_{w=0}, \\ \tilde{\mathcal{G}}_{gen, II}[\eta]\tilde{\boldsymbol{\omega}} &:= \tilde{\mathcal{U}}_{II}[\eta]\tilde{\boldsymbol{\omega}} \cdot \mathbf{N}|_{w=0} = \text{curl}^{\Sigma} \tilde{\mathbf{A}} \cdot \mathbf{N}|_{w=0}, \end{aligned}$$

and the associated expansions

$$\tilde{\mathcal{G}}_{gen, I}[\eta] = \sum_{j=0}^{\infty} \tilde{\mathcal{G}}_{j, I}[\eta], \quad \tilde{\mathcal{G}}_{gen, II}[\eta] = \sum_{j=0}^{\infty} \tilde{\mathcal{G}}_{j, II}[\eta].$$

We consider the flattened version of (1.21) and (1.22); by exploiting Corollary 4.3, we obtain that $\tilde{\mathbf{A}}$ and $\tilde{\varphi}$ satisfy

$$(6.2) \quad (|\mathbf{D}|^2 - \partial_w^2) \tilde{\mathbf{A}}(\mathbf{x}', w) = \tilde{\boldsymbol{\omega}}(\mathbf{x}', w) + \mathbf{R}_{11}[\sigma, \tilde{\mathbf{A}}](\mathbf{x}', w), \quad \text{in } D_0,$$

$$(6.3) \quad \partial_w \tilde{\mathbf{A}}_3(\cdot, -h) = 0,$$

$$(6.4) \quad \tilde{\mathbf{A}}_1(\cdot, -h) = \tilde{\mathbf{A}}_2(\cdot, -h) = 0,$$

$$(6.5) \quad \tilde{\mathbf{A}}_3(\cdot, 0) = r_{21}[\sigma, \tilde{\mathbf{A}}](\mathbf{x}'),$$

$$(6.6) \quad \mathbf{i} \mathbf{D}^{\perp} \tilde{\mathbf{A}}_3(\cdot, 0) - \partial_w \tilde{\mathbf{A}}_h(\cdot, 0)^{\perp} = -\nabla^{\perp} \Delta^{-1} (\tilde{\boldsymbol{\omega}}_3|_{w=0}) + r_{31}[\sigma, \tilde{\mathbf{A}}, \tilde{\boldsymbol{\omega}}],$$

$$(6.7) \quad (|\mathbf{D}|^2 - \partial_w^2) \tilde{\varphi}(\mathbf{x}', w) = r_{12}[\sigma, \tilde{\varphi}](\mathbf{x}', w), \quad \text{in } D_0,$$

$$(6.8) \quad \partial_w \tilde{\varphi}(\cdot, -h) = r_{22}[\sigma, \tilde{\varphi}],$$

$$(6.9) \quad \tilde{\varphi}(\cdot, 0) = \Phi,$$

where

$$\begin{aligned} \mathbf{R}_{11}[\sigma, \tilde{\mathbf{A}}](\mathbf{x}', w) &:= \Delta^\Sigma \tilde{\mathbf{A}}(\mathbf{x}', w) - \Delta_{\mathbf{x}', w} \tilde{\mathbf{A}}(\mathbf{x}', w), \\ r_{21}[\sigma, \tilde{\mathbf{A}}](\mathbf{x}') &:= \tilde{\mathbf{A}}_{\text{h}}(\mathbf{x}', 0) \cdot \nabla \eta(\mathbf{x}'), \\ r_{31}[\sigma, \tilde{\mathbf{A}}, \tilde{\omega}](\mathbf{x}') &:= - \left((\text{curl}^\Sigma \tilde{\mathbf{A}})|_{w=0} \right)_{\text{h}} + \left(\text{curl} \tilde{\mathbf{A}} \right)_{\text{h}}(\mathbf{x}', 0) \\ &\quad - \left((\text{curl}^\Sigma \tilde{\mathbf{A}})|_{w=0} \right)_3 \nabla \eta \\ &\quad + \left(\nabla^\perp \Delta^{-1}(\tilde{\omega}(\mathbf{x}', 0) \cdot \nabla \eta) \right), \end{aligned}$$

and

$$\begin{aligned} r_{12}[\sigma, \tilde{\varphi}](\mathbf{x}', w) &:= \Delta^\Sigma \tilde{\varphi}(\mathbf{x}', w) - \Delta_{\mathbf{x}', w} \tilde{\varphi}(\mathbf{x}', w), \\ r_{22}[\sigma, \tilde{\varphi}](\mathbf{x}') &:= -(\partial_w^\Sigma \tilde{\varphi})(\mathbf{x}', -h) + \partial_w \tilde{\varphi}(\mathbf{x}', -h). \end{aligned}$$

In the rest of the section we denote by \mathcal{F} the Fourier transform with respect to \mathbf{x}' , and we denote by \tilde{f}' the partial derivative $\partial_w \tilde{f}$.

Lemma 6.1. *Assume that $s \geq 2$, and let $\tilde{\omega} \in (H^{s-2}(D_0))^3$, $\Phi \in \dot{H}^{s-1/2}(\mathbb{R}^2)$. Then the boundary value problems*

$$\begin{cases} (|\mathbf{D}|^2 - \partial_w^2) \tilde{\mathbf{A}} &= \tilde{\omega} + \mathbf{R}_1, \text{ in } D_0, \\ \partial_w \tilde{A}_3(\cdot, -h) &= 0, \\ \tilde{A}_1(\cdot, -h) = \tilde{A}_2(\cdot, -h) &= 0, \\ \tilde{A}_3(\cdot, 0) &= r_2, \\ i\mathbf{D}^\perp \tilde{A}_3(\cdot, 0) - \partial_w \tilde{\mathbf{A}}_{\text{h}}(\cdot, 0)^\perp &= -\nabla^\perp \Delta^{-1}(\tilde{\omega}_3|_{w=0}) + \mathbf{r}_3, \end{cases}$$

$$\begin{cases} (|\mathbf{D}|^2 - \partial_w^2) \tilde{\varphi} &= r_4, \text{ in } D_0, \\ \partial_w \tilde{\varphi}(\cdot, -h) &= r_5, \\ \tilde{\varphi}(\cdot, 0) &= \Phi, \end{cases}$$

admit unique solutions $\tilde{\mathbf{A}} \in (H^s(D_0))^3$ and $\tilde{\varphi} \in \dot{H}^s(D_0)$ for any $\mathbf{R}_1 \in (H^{s-2}(D_0))^3$, $r_2 \in H^{s-1/2}(\mathbb{R}^2)$, $\mathbf{r}_3 \in (H^{s-3/2}(\mathbb{R}^2))^2$, $r_4 \in \dot{H}^{s-2}(D_0)$ and $r_5 \in \dot{H}^{s-3/2}(\mathbb{R}^2)$.

Proof. We rewrite the first boundary value problem for $\tilde{\mathbf{F}} := \mathcal{F} \tilde{\mathbf{A}}$ in the following way,

$$\begin{aligned} \mathbf{M} \tilde{\mathbf{F}} &= \mathcal{F} \tilde{\omega} + \mathcal{F} \mathbf{R}_1 \text{ in } D_0, \\ \mathbf{L}_{-h} \tilde{\mathbf{F}} &= \mathbf{0}, \text{ at } w = -h, \\ \mathbf{L}_0 \tilde{\mathbf{F}} &= \left(\mathcal{F} \left[\nabla \Delta^{-1}(\tilde{\omega}_3|_{w=0}) \right] + \mathcal{F} \mathbf{r}_3^\perp + i \boldsymbol{\xi} \mathcal{F} r_2 \right), \text{ at } w = 0, \end{aligned}$$

where

$$\mathbf{M} = (|\boldsymbol{\xi}|^2 - \partial_w^2) \mathbb{I}_3, \quad \mathbf{L}_{-h} = \text{diag}(1, 1, \partial_w), \quad \mathbf{L}_0 = \text{diag}(\partial_w, \partial_w, 1).$$

Let us define the Green's matrix

$$\mathbf{G}_{\tilde{\mathbf{A}}}(w, \zeta) = \begin{cases} \mathbf{U}(w+h) \overline{\mathbf{C}^{-1} \mathbf{W}(\zeta)}^T, & \text{for } -h \leq w \leq \zeta \leq 0, \\ \mathbf{W}(w) \mathbf{C}^{-1} \overline{\mathbf{U}(\zeta+h)}^T, & \text{for } -h \leq \zeta \leq w \leq 0, \end{cases}$$

where $\mathbf{U}(\cdot+h)$ (resp. \mathbf{W}) is a 3×3 matrix whose columns $\mathbf{U}_j(\cdot+h)$ (resp. \mathbf{W}_j), $j = 1, 2, 3$, are linearly independent solutions of

$$\begin{aligned} \mathbf{M} \tilde{\mathbf{F}} &= \mathbf{0}, \text{ in } D_0, \quad \mathbf{L}_{-h} \tilde{\mathbf{F}} = \mathbf{0}, \text{ at } w = -h, \\ (\text{resp. } \mathbf{M} \tilde{\mathbf{F}} &= \mathbf{0}, \text{ in } D_0, \quad \mathbf{L}_0 \tilde{\mathbf{F}} = \mathbf{0}, \text{ at } w = 0,) \end{aligned}$$

and

$$\mathbf{C} = -\overline{\mathbf{U}}^T(w+h) \mathbf{W}'(w) + \overline{\mathbf{U}}^T(w+h) \mathbf{W}(w).$$

More explicitly, we choose

$$\begin{aligned} \mathbf{U}_j(w) &= \sinh(|\xi|w) \mathbf{e}_j, \quad j = 1, 2, \quad \mathbf{U}_3(w) = \cosh(|\xi|w) \mathbf{e}_3, \\ \mathbf{W}_j(w) &= \cosh(|\xi|w) \mathbf{e}_j, \quad j = 1, 2, \quad \mathbf{W}_3(w) = \sinh(|\xi|w) \mathbf{e}_3, \end{aligned}$$

so that

$$\mathbf{C} = |\xi| \cosh(h|\xi|) \operatorname{diag}(1, 1, -1)$$

and

$$\mathbf{G}_{\tilde{\mathbf{A}}}(w, \zeta) = \begin{cases} \frac{1}{|\xi| \cosh(h|\xi|)} \operatorname{diag}(\sinh(|\xi|(w+h)) \cosh(|\xi|\zeta), \\ \quad \sinh(|\xi|(w+h)) \cosh(|\xi|\zeta), \\ \quad -\cosh(|\xi|(w+h)) \sinh(|\xi|\zeta)), \\ \quad \text{for } -h \leq w \leq \zeta \leq 0, \\ \\ \frac{1}{|\xi| \cosh(h|\xi|)} \operatorname{diag}(\sinh(|\xi|(\zeta+h)) \cosh(|\xi|w), \\ \quad \sinh(|\xi|(\zeta+h)) \cosh(|\xi|w), \\ \quad -\cosh(|\xi|(\zeta+h)) \sinh(|\xi|w)), \\ \quad \text{for } -h \leq \zeta \leq w \leq 0. \end{cases}$$

Hence, the solution $\tilde{\mathbf{A}}$ of the boundary value problem (6.2)-(6.6) is given by

$$\begin{aligned} \tilde{\mathbf{A}}(\cdot, w) &= \mathcal{F}^{-1} \left[\int_{-h}^0 \mathbf{G}_{\tilde{\mathbf{A}}}(w, \zeta) [(\mathcal{F}\tilde{\omega})(\cdot, \zeta) + \mathcal{F}\mathbf{R}_1(\cdot, \zeta)] d\zeta \right. \\ &\quad - \mathbf{G}_{\tilde{\mathbf{A}}}(w, 0) \left(\begin{array}{c} \mathcal{F} [\nabla \Delta^{-1}(\tilde{\omega}_3|_{w=0})] + \mathcal{F}\mathbf{r}_3^\perp + \mathbf{i}\xi \mathcal{F}\mathbf{r}_2 \\ 0 \end{array} \right) \\ &\quad \left. - \mathbf{G}_{\tilde{\mathbf{A}}, \zeta}(w, 0) \begin{pmatrix} 0 \\ 0 \\ \mathcal{F}\mathbf{r}_2 \end{pmatrix} \right]. \end{aligned} \tag{6.10}$$

Similarly, we write the Green's matrix for the boundary value problem for $\tilde{g} := \mathcal{F}\tilde{\varphi}$,

$$\mathbf{G}_{\tilde{\varphi}}(w, \zeta) = \begin{cases} \mathbf{u}(w+h) \mathbf{c}^{-1} \overline{\mathbf{w}(\zeta)}, & \text{for } -h \leq w \leq \zeta \leq 0, \\ \overline{\mathbf{w}(w)} \mathbf{c}^{-1} \mathbf{u}(\zeta+h), & \text{for } -h \leq \zeta \leq w \leq 0, \end{cases}$$

where $\mathbf{u}(\cdot+h)$ (resp. $\overline{\mathbf{w}}$) is a solution of

$$\begin{aligned} (|\xi|^2 - \partial_w^2) \tilde{g} &= 0, \quad \text{in } D_0, \quad \tilde{g}'(-h) = 0, \\ (\text{resp. } (|\xi|^2 - \partial_w^2) \tilde{g} &= 0, \quad \text{in } D_0, \quad \tilde{g}(0) = 0) \end{aligned}$$

and

$$\mathbf{c} = -\overline{\mathbf{u}}(w+h) \mathbf{w}'(w) + \overline{\mathbf{u}}'(w+h) \mathbf{w}(w).$$

More explicitly, we choose

$$\mathbf{u}(w) = \cosh(|\xi|w) \quad \mathbf{w}(w) = \sinh(|\xi|w),$$

so that

$$\mathbf{c} = -|\xi| \cosh(h|\xi|),$$

and

$$\mathbf{G}_{\tilde{\varphi}}(w, \zeta) = \begin{cases} -\frac{1}{|\xi| \cosh(h|\xi|)} \cosh(|\xi|(w+h)) \sinh(|\xi|\zeta), \\ \quad \text{for } -h \leq w \leq \zeta \leq 0, \\ \\ -\frac{1}{|\xi| \cosh(h|\xi|)} \cosh(|\xi|(\zeta+h)) \sinh(|\xi|w), \\ \quad \text{for } -h \leq \zeta \leq w \leq 0. \end{cases}$$

Hence the solution $\tilde{\varphi}$ of the boundary value problem (6.7)-(6.9) is given by

$$\begin{aligned} & \tilde{\varphi}(\cdot, w) \\ &= \mathcal{F}^{-1} \left[\int_{-h}^0 \mathbf{G}_{\tilde{\varphi}}(w, \zeta) \mathcal{F}r_4(\cdot, \zeta) d\zeta - \mathbf{G}_{\tilde{\varphi}}(w, -h) \mathcal{F}r_5 - \mathbf{G}_{\tilde{\varphi}, \zeta}(w, 0) (\mathcal{F}\Phi) \right]. \end{aligned} \quad (6.11)$$

□

Following an argument analogous to the proof of Theorem 4.10 in [GH20], one can deduce from Lemma 6.1 the following result.

Theorem 6. *Suppose that $s \geq 2$, and let $\tilde{\omega} \in (H^{s-2}(D_0))^3$, $\Phi \in \dot{H}^{s-1/2}(\mathbb{R}^2)$. There exists an open neighbourhood W of the origin in $H^{s+1/2}(D_0)$ such that the boundary value problems (6.2)-(6.6) and (6.7)-(6.9) admit unique solutions $\tilde{\mathbf{A}} = \tilde{\mathbf{A}}(\sigma, \tilde{\omega}) \in (H^s(D_0))^3$ and $\tilde{\varphi} = \tilde{\varphi}(\sigma, \Phi) \in H^s(D_0)$ for each $\sigma \in W$. Furthermore, $\tilde{\mathbf{A}}(\sigma, \tilde{\omega})$ depends analytically upon σ and $\tilde{\omega}$ and linearly upon $\tilde{\omega}$, while $\tilde{\varphi}(\sigma, \Phi)$ depends analytically upon σ and Φ and linearly upon Φ .*

Proof. Let $s \geq 2$. By (B.5) (which holds true also for Sobolev spaces on D_0 , as mentioned at pag.16 of [GH20]), one can see that the maps

$$(\sigma, \tilde{\mathbf{A}}) \mapsto \mathbf{R}_{11}[\sigma, \tilde{\mathbf{A}}], \quad (\sigma, \tilde{\mathbf{A}}) \mapsto r_{21}[\sigma, \tilde{\mathbf{A}}],$$

are analytic in $W \times (H^s(D_0))^3$, where W is an open neighbourhood of $H^{s+1/2}(\mathbb{R}^2)$. Similarly, the map

$$(\sigma, \tilde{\mathbf{A}}, \tilde{\omega}) \mapsto \mathbf{r}_{31}[\sigma, \tilde{\mathbf{A}}, \tilde{\omega}]$$

is analytic in $W \times (H^s(D_0))^3 \times (H^{s-2}(D_0))^3$, and the maps

$$(\sigma, \tilde{\varphi}) \mapsto r_{12}[\sigma, \tilde{\varphi}], \quad (\sigma, \tilde{\varphi}) \mapsto r_{22}[\sigma, \tilde{\varphi}]$$

are analytic in $W \times (H^s(D_0))^3$. Hence the maps

$$\begin{aligned} & \mathcal{F}_1(\tilde{\mathbf{A}}, \sigma, \tilde{\omega}) \\ &:= \begin{pmatrix} (|\mathbf{D}|^2 - \partial_w^2) \tilde{\mathbf{A}} - \tilde{\omega} - \mathbf{R}_{11}[\sigma, \tilde{\mathbf{A}}] \\ \tilde{\mathbf{A}}_3(\cdot, 0) - r_{21}[\sigma, \tilde{\mathbf{A}}] \\ i\mathbf{D}^\perp \tilde{\mathbf{A}}_3(\cdot, 0) - \partial_w \tilde{\mathbf{A}}_h(\cdot, 0)^\perp - \nabla^\perp \Delta^{-1} (\tilde{\omega}_3|_{w=0}) - \mathbf{r}_{31}[\sigma, \tilde{\mathbf{A}}] \end{pmatrix} \end{aligned}$$

and

$$\mathcal{F}_2(\tilde{\varphi}, \sigma, \Phi) := \begin{pmatrix} (|\mathbf{D}|^2 - \partial_w^2) \tilde{\varphi} - r_{12}[\sigma, \tilde{\varphi}] \\ \partial_w \tilde{\varphi}(\cdot, -h) - r_{22}[\sigma, \tilde{\varphi}] \\ \tilde{\varphi}(\cdot, 0) - \Phi \end{pmatrix}$$

define analytic maps $\mathcal{F}_1 : S \times W \times (H^{s-2}(D_0))^3 \rightarrow (H^{s-2}(D_0))^3 \times H^{s-1/2}(\mathbb{R}^2) \times (H^{s-3/2}(\mathbb{R}^2))^2$, where

$$S := \{ \tilde{\mathbf{A}} \in (H^s(D_0))^3 : \tilde{\mathbf{A}}_1(\cdot, -h) = \tilde{\mathbf{A}}_2(\cdot, -h) = 0, \partial_w \tilde{\mathbf{A}}_3(\cdot, -h) = 0 \},$$

and $\mathcal{F}_2 : \dot{H}^s(D_0) \times W \times \dot{H}^{s-1/2}(\mathbb{R}^2) \rightarrow \dot{H}^{s-2}(D_0) \times H^{s-3/2}(\mathbb{R}^2) \times \dot{H}^{s-1/2}(\mathbb{R}^2)$, where $\mathcal{F}_1(\mathbf{0}, \mathbf{0}, \mathbf{0}) = (\mathbf{0}, \mathbf{0}, \mathbf{0})$ and $\mathcal{F}_2(0, 0, 0) = (0, 0, 0)$. Moreover, Lemma 6.1 implies that the maps $d_1 \mathcal{F}_1[\mathbf{0}, \mathbf{0}, \mathbf{0}] : S \times H^{s+1/2}(D_0) \times (H^{s-2}(D_0))^3 \rightarrow (H^{s-2}(D_0))^3 \times H^{s-1/2}(\mathbb{R}^2) \times (H^{s-3/2}(\mathbb{R}^2))^2$ and $d_1 \mathcal{F}_2[0, 0, 0] : \dot{H}^s(D_0) \times H^{s+1/2}(D_0) \times \dot{H}^{s-1/2}(\mathbb{R}^2) \rightarrow \dot{H}^{s-2}(D_0) \times H^{s-3/2}(\mathbb{R}^2) \times \dot{H}^{s-1/2}(\mathbb{R}^2)$ are isomorphisms. The thesis follows by the analytic implicit function theorem (see Theorem 4.5.3 of [BT03]).

□

From Theorem 6 we can deduce the following analyticity result (see also Corollary 4.12 of [GH20]).

Corollary 6.2. *Let $s \geq 2$, and let $\tilde{\omega} \in (H^{s-2}(D_0))^3$, $\Phi \in \dot{H}^{s-1/2}(\mathbb{R}^2)$. Assume that the condition (2.5) holds true. Then there exists an open neighbourhood V of the origin in $H^{s+1/2}(\mathbb{R}^2)$ such that formulae*

$$\begin{aligned}\tilde{\mathcal{G}}_{gen,I}[\eta](\Phi, \tilde{\omega}) &= \tilde{U}_I[\eta]\Phi \cdot \mathbf{N}|_{w=0}, \\ \tilde{\mathcal{G}}_{gen,II}[\eta](\Phi, \tilde{\omega}) &= \tilde{U}_{II}[\eta]\tilde{\omega} \cdot \mathbf{N}|_{w=0}\end{aligned}$$

define analytic maps

$$\begin{aligned}\tilde{\mathcal{G}}_{gen,I}[\cdot] : V &\rightarrow L(\dot{H}^{s-1/2}(\mathbb{R}^2), H^{s-3/2}(\mathbb{R}^2)), \\ \tilde{\mathcal{G}}_{gen,II}[\cdot] : V &\rightarrow L((H^{s-2}(D_0))^3, H^{s-3/2}(\mathbb{R}^2)).\end{aligned}$$

We notice that the formula for $\tilde{\mathcal{G}}_{gen,II}[\eta]\tilde{\omega}$ can be rewritten as

$$\begin{aligned}\tilde{\mathcal{G}}_{gen,II}[\eta]\tilde{\omega} &= \nabla \cdot (\tilde{A}_h^\perp + \tilde{A}_3 \nabla^\perp \eta)|_{w=0} \\ &= \tilde{A}_{2x} + \eta_y \tilde{A}_{3x} - \tilde{A}_{1y} - \eta_x \tilde{A}_{3y}|_{w=0}.\end{aligned}$$

We compute the term $\tilde{\mathcal{G}}_0$ in the Taylor expansion (6.1), where we write

$$\tilde{\mathcal{G}}_0(\Phi, \tilde{\omega}) = \tilde{\mathcal{G}}_{0,I}\Phi + \tilde{\mathcal{G}}_{0,II}\tilde{\omega}.$$

From formulae (6.10) and (6.11) we obtain that for $\eta = 0$

$$\begin{aligned}\tilde{\mathbf{A}}_0(\cdot, w) &= \tilde{\mathbf{A}}(\cdot, w)|_{\mathbf{R}_{11}=0, r_{21}=0, r_{31}=0} \\ &= \mathcal{F}^{-1} \left[\int_{-h}^0 \mathbf{G}_{\tilde{\mathbf{A}}}(w, \zeta) (\mathcal{F}\tilde{\omega})(\cdot, \zeta) d\zeta - \mathbf{G}_{\tilde{\mathbf{A}}}(w, 0) \left(\mathcal{F} \left[\begin{array}{c} \nabla \Delta^{-1}(\tilde{\omega}_3|_{w=0}) \\ 0 \end{array} \right] \right) \right], \\ \tilde{\varphi}_0(\cdot, w) &= \tilde{\varphi}(\cdot, w)|_{r_{12}=r_{22}=0} \\ &= \mathcal{F}^{-1} \left[-\mathbf{G}_{\tilde{\varphi}, \zeta}(w, 0) (\mathcal{F}\Phi) \right] = \mathcal{F}^{-1} \left[\frac{\cosh(|\xi|(w+h))}{\cosh(h|\xi|)} (\mathcal{F}\Phi) \right],\end{aligned}$$

so that

$$\begin{aligned}\tilde{\mathcal{G}}_{0,I}\Phi &= \partial_w \tilde{\varphi}_0|_{w=0} = -\mathcal{F}^{-1} \left[\mathbf{G}_{\tilde{\varphi}, w\zeta}(0, 0) (\mathcal{F}\Phi) \right] \\ (6.12) \quad &= \mathcal{F}^{-1} \left[|\xi| \tanh(h|\xi|) (\mathcal{F}\Phi) \right],\end{aligned}$$

$$\begin{aligned}\tilde{\mathcal{G}}_{0,II}\tilde{\omega} &= \tilde{\mathbf{A}}_{0,2x} - \tilde{\mathbf{A}}_{0,1y}|_{w=0} \\ &= \nabla \cdot \mathcal{F}^{-1} \left[\int_{-h}^0 \mathbf{G}_{\tilde{\mathbf{A}}}(0, \zeta) (\mathcal{F}\tilde{\omega})(\cdot, \zeta) d\zeta \right. \\ &\quad \left. - \mathbf{G}_{\tilde{\mathbf{A}}}(0, 0) \left(\mathcal{F} \left[\begin{array}{c} \nabla \Delta^{-1}(\tilde{\omega}_3|_{w=0}) \\ 0 \end{array} \right] \right) \right]_{\mathbf{h}}^\perp, \\ &= \nabla \cdot \mathcal{F}^{-1} \left[\int_{-h}^0 \frac{\sinh(|\xi|(\zeta+h))}{|\xi| \cosh(h|\xi|)} (\mathcal{F}\tilde{\omega})_{\mathbf{h}}^\perp(\cdot, \zeta) d\zeta \right. \\ (6.13) \quad &\quad \left. - \frac{\sinh(h|\xi|)}{|\xi| \cosh(h|\xi|)} \mathcal{F} \left[\nabla^\perp \Delta^{-1}(\tilde{\omega}_3|_{w=0}) \right] \right],\end{aligned}$$

since

$$\mathbf{G}_{\tilde{\mathbf{A}}}(0, \zeta) = \frac{\sinh(|\xi|(\zeta+h))}{|\xi| \cosh(h|\xi|)} \text{diag}(1, 1, 0), \quad -h \leq \zeta \leq 0.$$

From (6.11) we can deduce a formula for the expansion of the operator $\tilde{\mathcal{G}}_I[\eta]$ around $\eta = 0$ (see Sec. 2.4 of [GNPW24], where such expansion is proved in the more general case of Beltrami flows; see also the argument by Craig-Sulem in [CS93])

$$\begin{aligned}
\tilde{\mathcal{G}}_{1,I}[\eta]\Phi &= -\tilde{\mathcal{G}}_{0,I}\left(\eta\tilde{\mathcal{G}}_{0,I}\Phi\right) - \nabla \cdot (\eta \nabla \Phi), \\
\tilde{\mathcal{G}}_{j,I}[\eta]\Phi &= \frac{1}{j} \left\{ -\tilde{\mathcal{G}}_{j-1,I}[\eta](\eta\tilde{\mathcal{G}}_{0,I}\Phi) \right. \\
&\quad \left. - \sum_{k=0}^{j-2} \tilde{\mathcal{G}}_{k,I}[\eta]\left(u_{j-1-k}(\eta)(\Phi)\eta\right) \right. \\
(6.14) \quad &\quad \left. + \nabla \cdot \left((u_{j-2}(\eta)(\Phi) \nabla \eta) \eta \right) \right\}, \quad j \geq 2,
\end{aligned}$$

where

$$\begin{aligned}
u_0(\Phi) &= \tilde{\mathcal{G}}_{0,I}\Phi, \\
u_k(\eta)(\Phi) &= \begin{cases} (-1)^{k/2} |\nabla \eta|^k \tilde{\mathcal{G}}_{0,I}\Phi \\ \quad + \sum_{i+2j=k} (\mathbf{K}_{i-1}\Phi \cdot \nabla \eta + \tilde{\mathcal{G}}_{i,I}[\eta]\Phi) (-1)^j |\nabla \eta|^{2j}, & \text{if } k \in 2\mathbb{N}, \\ \sum_{i+2j=k} (\mathbf{K}_{i-1}\Phi \cdot \nabla \eta + \tilde{\mathcal{G}}_{i,I}[\eta]\Phi) (-1)^j |\nabla \eta|^{2j}, & \text{if } k \notin 2\mathbb{N}, \end{cases}
\end{aligned}$$

for $k \geq 1$, with $\mathbf{K}_0\Phi = \nabla\Phi$ and $\mathbf{K}_i\Phi = \mathbf{0} \forall i \geq 1$.

Next, we obtain the terms $\tilde{\mathcal{G}}_{j,II}[\eta]$ in the Taylor expansion of the operator $\tilde{\mathcal{G}}_{II}[\eta]$ by deducing a formula for the differential of $d\tilde{\mathcal{G}}_{gen,II}[\eta](\delta\eta)$, and then by evaluating it at $\delta\eta = \eta$.

Proposition 6.3. *Let $s \geq 3$, and let $\tilde{\omega} \in (H^{s-2}(D_0))^3$, $\Phi \in \dot{H}^{s-1/2}(\mathbb{R}^2)$. Assume that the condition (2.5) holds true. Then there exists an open neighbourhood V of the origin in $H^{s+1/2}(\mathbb{R}^2)$ such that the differential of the operator*

$$\tilde{\mathcal{G}}_{gen,II}[\cdot] : V \rightarrow L\left((H^{s-2}(D_0))^3, H^{s-3/2}(\mathbb{R}^2)\right)$$

is given by

$$\begin{aligned}
(6.15) \quad d\tilde{\mathcal{G}}_{gen,II}[\eta](\delta\eta)\tilde{\omega} &= -\tilde{\mathcal{G}}_{gen,II}[\eta]\tilde{\gamma} - \nabla \cdot \left[(\tilde{\mathbb{K}}_{II}[\eta]\tilde{\omega} - \tilde{\mathbb{W}}_{II}[\eta]\tilde{\omega} \nabla \eta) \delta\eta \right], \\
\tilde{\gamma} &= \partial_w^\Sigma \tilde{\omega} \, d\sigma,
\end{aligned}$$

$$\tilde{\mathbb{K}}_{II}[\eta]\tilde{\omega} = -\nabla^\perp \Delta^{-1}(\tilde{\omega} \cdot \mathbf{N}|_{w=0}), \quad \tilde{\mathbb{W}}_{II}[\eta]\tilde{\omega} = \frac{1}{1+|\eta|^2} \left[\tilde{\mathbb{K}}_{II}[\eta]\tilde{\omega} \cdot \nabla \eta + \tilde{\mathcal{G}}_{gen,II}[\eta]\tilde{\omega} \right].$$

Remark 6.4. *Due to the decomposition (1.20) of the velocity field, we have that*

$$\tilde{\mathbb{K}}_{II}[\eta]\tilde{\omega} = (\text{curl}^\Sigma \tilde{\mathbf{A}})_\parallel, \quad \tilde{\mathbb{W}}_{II}[\eta]\tilde{\omega} = (\text{curl}^\Sigma \tilde{\mathbf{A}})_3|_{w=0}.$$

Proof of Proposition 6.3. Recall that

$$\tilde{\mathcal{G}}_{gen,II}[\eta]\tilde{\omega} = (\text{curl}^\Sigma \tilde{\mathbf{A}}) \cdot \mathbf{N}|_{w=0},$$

where $\tilde{\mathbf{A}}$ solves

$$(6.16) \quad \begin{cases} -\Delta^\Sigma \tilde{\mathbf{A}} = \tilde{\omega}, & \text{in } D_0, \\ \partial_w^\Sigma \tilde{\mathbf{A}}_3 = 0, & \text{at } w = -h, \\ \tilde{\mathbf{A}} \times \mathbf{e}_3 = \mathbf{0}, & \text{at } w = -h, \\ \tilde{\mathbf{A}} \cdot \mathbf{N} = 0, & \text{at } w = 0, \\ (\text{curl}^\Sigma \tilde{\mathbf{A}})_\parallel = -\nabla^\perp \Delta^{-1}(\tilde{\omega} \cdot \mathbf{N}), & \text{at } w = 0. \end{cases}$$

Let us recall the following identity (see (3.41) in [CL15])

$$(6.17) \quad \delta(\partial_j^\Sigma f) = \partial_j^\Sigma(\delta f - \delta\sigma \partial_w^\Sigma f) + \delta\sigma \partial_w^\Sigma \partial_j^\Sigma f, \quad j = x, y, w, t,$$

where δ is any linearization operator. If we introduce

$$\tilde{C} := d\tilde{A} - d\sigma \partial_w^\Sigma \tilde{A},$$

where $df = df[\eta](\delta\eta)$, and we apply (6.17), we have that

$$\begin{aligned} & d\tilde{\mathcal{G}}_{gen,II}[\eta](\delta\eta)\tilde{\omega} \\ &= (\operatorname{curl}^\Sigma \tilde{C}) \cdot \mathbf{N}|_{w=0} + \partial_w^\Sigma(\operatorname{curl}^\Sigma \tilde{A}) \cdot \mathbf{N}|_{w=0} \delta\eta - (\operatorname{curl}^\Sigma \tilde{A})_h|_{w=0} \cdot \nabla \delta\eta \\ &= (\operatorname{curl}^\Sigma \tilde{C}) \cdot \mathbf{N}|_{w=0} - \nabla \cdot ((\operatorname{curl}^\Sigma \tilde{A})_h|_{w=0} \delta\eta). \end{aligned}$$

Since $\partial_w^\Sigma \tilde{A}|_{w=0} \cdot \mathbf{N} = -\nabla \cdot \tilde{A}_h|_{w=0}$, we have

$$\left\{ \begin{array}{ll} -\Delta^\Sigma \tilde{C} = -\partial_w^\Sigma \tilde{\omega} d\sigma, & \text{in } D_0, \\ \partial_w^\Sigma \tilde{C}_3 = 0, & \text{at } w = -h, \\ \tilde{C} \times \mathbf{e}_3 = \mathbf{0}, & \text{at } w = -h, \\ \tilde{C} \cdot \mathbf{N} = \nabla \cdot (\tilde{A}_h|_{w=0} \delta\eta), & \text{at } w = 0, \\ (\operatorname{curl}^\Sigma \tilde{C})_\parallel = \nabla^\perp \Delta^{-1} (\tilde{\omega}_h|_{w=0} \cdot \nabla \delta\eta) \\ \quad - \delta\eta \partial_w^\Sigma (\operatorname{curl}^\Sigma \tilde{A})_h|_{w=0} \\ \quad - \delta\eta \partial_w^\Sigma (\operatorname{curl}^\Sigma \tilde{A})_3|_{w=0} \nabla \eta \\ \quad - (\operatorname{curl}^\Sigma \tilde{A})_3|_{w=0} \nabla \delta\eta, & \text{at } w = 0, \end{array} \right.$$

where we used that $d\sigma|_{w=0} = \delta\eta$, $d\sigma|_{w=-h} = 0$, that

$$-\delta\eta \partial_w^\Sigma \tilde{A} \cdot \mathbf{N} + \nabla(\delta\eta) \cdot \tilde{A}_h|_{w=0} = \nabla \cdot (\tilde{A}_h|_{w=0} \delta\eta),$$

and that

$$-d \nabla^\perp \Delta^{-1} (\tilde{\omega} \cdot \mathbf{N}) = -\nabla^\perp \Delta^{-1} [-\tilde{\omega}_h \cdot \nabla \delta\eta].$$

Now we rewrite the last boundary condition using that

$$\partial_w^\Sigma (\operatorname{curl}^\Sigma \tilde{A})_h|_{w=0} + \partial_w^\Sigma (\operatorname{curl}^\Sigma \tilde{A})_3|_{w=0} \nabla \eta = \nabla (\operatorname{curl}^\Sigma \tilde{A})_3|_{w=0} + \tilde{\omega}_h^\perp|_{w=0},$$

hence

$$\begin{aligned} & (\operatorname{curl}^\Sigma \tilde{C})_\parallel \\ &= \nabla^\perp \Delta^{-1} (\tilde{\omega}_h|_{w=0} \cdot \nabla \delta\eta) - \tilde{\omega}_h^\perp|_{w=0} \delta\eta - \nabla ((\operatorname{curl}^\Sigma \tilde{A})_3|_{w=0} \delta\eta) \\ &= \nabla^\perp \Delta^{-1} (\tilde{\omega}_h|_{w=0} \cdot \nabla \delta\eta) - \nabla^\perp \Delta^{-1} (\nabla^\perp \cdot (\tilde{\omega}_h^\perp|_{w=0} \delta\eta)) \\ &\quad - \nabla \Delta^{-1} (\nabla \cdot (\tilde{\omega}_h^\perp|_{w=0} \delta\eta)) - \nabla ((\operatorname{curl}^\Sigma \tilde{A})_3|_{w=0} \delta\eta) \\ &= -\nabla^\perp \Delta^{-1} [(\nabla \cdot \tilde{\omega}_h|_{w=0}) \delta\eta] \\ &\quad + \nabla [-\Delta^{-1} \nabla \cdot (\tilde{\omega}_h^\perp|_{w=0} \delta\eta) - (\operatorname{curl}^\Sigma \tilde{A})_3|_{w=0} \delta\eta] \\ &= -\nabla^\perp \Delta^{-1} [-\partial_w^\Sigma \tilde{\omega}|_{w=0} \cdot \mathbf{N} \delta\eta] \\ &\quad + \nabla [-\Delta^{-1} \nabla \cdot (\tilde{\omega}_h^\perp|_{w=0} \delta\eta) - (\operatorname{curl}^\Sigma \tilde{A})_3|_{w=0} \delta\eta] \end{aligned}$$

and we replace the interior equation and the last two boundary conditions with

$$\begin{aligned} & -\Delta^\Sigma \tilde{C} = -\tilde{\gamma}, \quad \text{in } D_0, \quad \tilde{\gamma} := \partial_w^\Sigma \tilde{\omega} d\sigma, \\ & \tilde{C} \cdot \mathbf{N} = 0, \quad \text{at } w = 0, \\ & (\operatorname{curl}^\Sigma \tilde{C})_\parallel = -\nabla^\perp \Delta^{-1} (-\tilde{\gamma} \cdot \mathbf{N}), \quad \text{at } w = 0, \end{aligned}$$

by replacing \tilde{C} with $\tilde{C} + \nabla^\Sigma \tilde{\phi}_1 + \nabla^\Sigma \tilde{\phi}_2$, where

$$\begin{cases} \Delta^\Sigma \tilde{\phi}_1 = 0, & \text{in } D_0, \\ \nabla^\Sigma \tilde{\phi}_1 \cdot \mathbf{N} = \nabla \cdot (\tilde{\mathbf{A}}_h|_{w=0} \delta\eta), & \text{at } w = 0, \\ \tilde{\phi}_1 = 0, & \text{at } w = -h. \end{cases}$$

and

$$\begin{cases} \Delta^\Sigma \tilde{\phi}_2 = 0, & \text{in } D_0, \\ \tilde{\phi}_2 = -\Delta^{-1} \nabla \cdot (\tilde{\omega}_h^\perp|_{w=0} \delta\eta) - (\text{curl}^\Sigma \tilde{\mathbf{A}})_3|_{w=0} \delta\eta, & \text{at } w = 0, \\ \partial_n \tilde{\phi}_2 = 0, & \text{at } w = -h. \end{cases}$$

Therefore

$$d\tilde{\mathcal{G}}_{gen,II}[\eta](\delta\eta)\tilde{\omega} = -\tilde{\mathcal{G}}_{gen,II}[\eta]\tilde{\gamma} - \nabla \cdot [(\tilde{\mathbb{K}}_{II}[\eta]\tilde{\omega} - \tilde{\mathbb{W}}_{II}[\eta]\tilde{\omega} \nabla\eta) \delta\eta],$$

where $\tilde{\gamma}$, $\tilde{\mathbb{K}}_{II}[\eta]\tilde{\omega}$ and $\tilde{\mathbb{W}}_{II}[\eta]\tilde{\omega}$ are as in the statement. \square

In order to compute the terms in the expansion

$$\tilde{\mathcal{G}}_{gen,II}[\eta] = \sum_{j=0}^{\infty} \tilde{\mathcal{G}}_{j,II}[\eta], \quad \tilde{\mathcal{G}}_{j,II}[\eta] = \frac{1}{j!} d^j \tilde{\mathcal{G}}_{gen,II}[0][\eta, \dots, \eta],$$

where $\tilde{\mathcal{G}}_{j,II}(\eta)$ is homogeneous of degree j in η , we first evaluate formula (6.15) at $\delta\eta = \eta$. Then we expand the different terms appearing in (6.15): if $\tilde{\gamma}|_{\delta\eta=\eta}$ has no homogeneous term of degree 0 in η we have $\tilde{\gamma}_0 = \mathbf{0}$, therefore

$$\tilde{\gamma}|_{\delta\eta=\eta} = \sum_{j \geq 1} \tilde{\gamma}_j, \quad \tilde{\gamma}_j := \frac{1}{j!} d^j (\tilde{\gamma}[0]\tilde{\omega})[\eta, \dots, \eta],$$

while

$$\tilde{\nu} := \tilde{\nu}[\eta]\tilde{\omega} := \tilde{\omega} \cdot \mathbf{N}|_{w=0} = \sum_{j \geq 0} \tilde{\nu}_j, \quad \tilde{\nu}_j := \frac{1}{j!} d^j (\tilde{\nu}[0]\tilde{\omega})[\eta, \dots, \eta],$$

$$\tilde{\mathbb{K}}_{II}[\eta]\tilde{\omega} = \sum_{k=0}^{\infty} \tilde{\mathbb{K}}_{k,II}[\eta]\tilde{\omega}, \quad \tilde{\mathbb{W}}_{II}[\eta]\tilde{\omega} = \sum_{k=0}^{\infty} \tilde{\mathbb{W}}_{k,II}[\eta]\tilde{\omega},$$

where

$$\begin{aligned} \tilde{\mathbb{K}}_{0,II}\tilde{\omega} &= -\nabla^\perp \Delta^{-1} \tilde{\nu}_0, \quad \tilde{\mathbb{K}}_{k,II}[\eta]\tilde{\omega} = -\nabla^\perp \Delta^{-1} \tilde{\nu}_k, \quad \forall k \geq 1, \\ \tilde{\mathbb{W}}_{0,II}\tilde{\omega} &= \tilde{\mathcal{G}}_{0,II}\tilde{\omega}, \\ \tilde{\mathbb{W}}_{k,II}[\eta]\tilde{\omega} &= \tilde{\mathbb{Z}}_k[\eta]\tilde{\omega} + \sum_{i+2j=k} (\tilde{\mathbb{K}}_{i-1,II}[\eta]\tilde{\omega} \cdot \nabla\eta + \tilde{\mathcal{G}}_{i,II}[\eta]\tilde{\omega}) (-1)^j |\nabla\eta|^{2j}, \\ &\quad \forall k \geq 1, \\ \tilde{\mathbb{Z}}_k[\eta]\tilde{\omega} &= \begin{cases} (-1)^{k/2} |\nabla\eta|^k \tilde{\mathcal{G}}_{0,II}\tilde{\omega}, & \text{if } k \in 2\mathbb{N}, \\ 0, & \text{if } k \notin 2\mathbb{N}. \end{cases} \end{aligned}$$

Therefore, we obtain

$$\begin{aligned} \sum_{j \geq 0} j \tilde{\mathcal{G}}_{j,II}[\eta]\tilde{\omega} &= - \left[\sum_{h \geq 0} \tilde{\mathcal{G}}_{h,II}[\eta] \right] \left[\sum_{k \geq 1} \tilde{\gamma}_k \right] - \nabla \cdot [\tilde{\mathbb{K}}_{0,II}\tilde{\omega} \eta] \\ &\quad - \sum_{k \geq 1} \nabla \cdot [(\tilde{\mathbb{K}}_{k,II}[\eta]\tilde{\omega} - \tilde{\mathbb{W}}_{k-1,II}[\eta]\tilde{\omega} \nabla\eta) \eta], \end{aligned}$$

hence

$$(6.18) \quad \tilde{\mathcal{G}}_{1,II}[\eta]\tilde{\omega} = -\tilde{\mathcal{G}}_{0,II}\tilde{\gamma}_1 - \nabla \cdot [\tilde{\mathbb{K}}_{0,II}\tilde{\omega} \eta],$$

and for any $j \geq 2$

$$(6.19) \quad \tilde{\mathcal{G}}_{j,II}[\eta]\tilde{\omega} = \frac{1}{j} \left\{ - \sum_{h=0}^{j-1} \tilde{\mathcal{G}}_{h,II}[\eta]\tilde{\gamma}_{j-h} - \nabla \cdot [(\tilde{\mathbb{K}}_{j-1,II}[\eta]\tilde{\omega} - \tilde{\mathbb{V}}_{j-2,II}[\eta]\tilde{\omega} \nabla \eta) \eta] \right\}.$$

By summarising, we have shown the following result.

Theorem 7. *Under the assumptions of Proposition 6.3, the term $\tilde{\mathcal{G}}_0$ in the homogeneous expansion (6.1) of the operator $\tilde{\mathcal{G}}_{gen}[\eta]$ is given by*

$$\tilde{\mathcal{G}}_0 = \tilde{\mathcal{G}}_{0,I} + \tilde{\mathcal{G}}_{0,II},$$

where $\tilde{\mathcal{G}}_{0,I}$ and $\tilde{\mathcal{G}}_{0,II}$ are given by (6.12) and (6.13), respectively. Moreover, if $\tilde{\gamma}_0 = \mathbf{0}$, then for any $j \geq 1$ the term $\tilde{\mathcal{G}}_j[\eta]$ are given by

$$\tilde{\mathcal{G}}_j[\eta] = \tilde{\mathcal{G}}_{j,I}[\eta] + \tilde{\mathcal{G}}_{j,II}[\eta],$$

where $\tilde{\mathcal{G}}_{j,I}$ and $\tilde{\mathcal{G}}_{j,II}$ are given by (6.14) and (6.19), respectively.

Remark 6.5. *As an application of Theorem 7, we consider the water waves system in the generalized Zakharov–Craig–Sulem formulation, see (1.26)-(1.29), and we compute its linearization at $(\eta, \Phi, \tilde{\omega}) = (0, 0, \mathbf{0})$. We consider the straightened version of (1.26)-(1.29),*

$$(6.20) \quad \operatorname{div}^\Sigma \tilde{\omega} = 0, \quad \text{in } D_0,$$

$$(6.21) \quad \eta_t - \tilde{\mathcal{G}}_{gen}[\eta](\Phi, \tilde{\omega}) = 0, \quad \text{at } w = 0,$$

$$(6.22) \quad \begin{aligned} \Phi_t + g\eta + \frac{1}{2}|\tilde{\mathbb{U}}_\parallel[\eta](\Phi, \tilde{\omega})|^2 - \frac{(\tilde{\mathcal{G}}_{gen}[\eta](\Phi, \tilde{\omega}) + \nabla\eta \cdot \tilde{\mathbb{U}}_\parallel[\eta](\Phi, \tilde{\omega}))^2}{2(1 + |\nabla\eta|^2)} \\ + \nabla^\perp \cdot \Delta^{-1}(\tilde{\omega} \cdot \mathbf{N} \tilde{\mathbb{V}}[\eta](\Phi, \tilde{\omega})) = 0, \quad \text{at } w = 0, \end{aligned}$$

$$(6.23) \quad \tilde{\omega}_t + (\tilde{\mathbb{U}}[\eta](\Phi, \tilde{\omega}) \cdot \nabla^\Sigma) \tilde{\omega} = (\tilde{\omega} \cdot \nabla^\Sigma) \tilde{\mathbb{U}}[\eta](\Phi, \tilde{\omega}), \quad \text{in } D_0,$$

and we study its linearization at $(\eta, \Phi, \tilde{\omega}) = (0, 0, \mathbf{0})$. We obtain

$$(6.24) \quad \begin{cases} \operatorname{div}^\Sigma \tilde{\omega} = 0, & \text{in } D_0, \\ \eta_t = \tilde{\mathcal{G}}_0(\Phi, \tilde{\omega}), & \text{at } w = 0, \\ \tilde{\omega}_t = \mathbf{0}, & \text{in } D_0, \\ \Phi_t + g\eta = 0, & \text{at } w = 0, \end{cases}$$

where by Theorem 7 we have

$$\begin{aligned} \tilde{\mathcal{G}}_0(\Phi, \tilde{\omega}) &= \tilde{\mathcal{G}}_{0,I}\Phi + \tilde{\mathcal{G}}_{0,II}\tilde{\omega}, \\ \tilde{\mathcal{G}}_{0,I}\Phi &= \mathcal{F}^{-1} \left[|\boldsymbol{\xi}| \tanh(h|\boldsymbol{\xi}|) (\mathcal{F}\Phi) \right], \\ \tilde{\mathcal{G}}_{0,II}\tilde{\omega} &= \nabla \cdot \mathcal{F}^{-1} \left[\int_{-h}^0 \frac{\sinh(|\boldsymbol{\xi}|(\zeta + h))}{|\boldsymbol{\xi}| \cosh(h|\boldsymbol{\xi}|)} (\mathcal{F}\tilde{\omega})_\mathbb{R}^\perp(\cdot, \zeta) d\zeta \right. \\ &\quad \left. - \frac{\sinh(h|\boldsymbol{\xi}|)}{|\boldsymbol{\xi}| \cosh(h|\boldsymbol{\xi}|)} \mathcal{F} [\nabla^\perp \Delta^{-1}(\tilde{\omega}_3|_{w=0})] \right]. \end{aligned}$$

7. PARALINEARIZATION OF THE GENERALIZED DIRICHLET–NEUMANN OPERATOR

Now we study the parlinearization of the generalized Dirichlet–Neumann operator $\mathcal{G}_{gen}[\eta]$ defined in (1.25). Recalling the decomposition

$$\begin{aligned}\mathcal{G}_{gen}[\eta](\Phi, \omega) &= \mathcal{G}_{gen,I}[\eta]\Phi + \mathcal{G}_{gen,II}[\eta]\omega, \\ \mathcal{G}_{gen,I}[\eta]\Phi &= \mathcal{G}[\eta]\Phi = \mathbb{U}_I[\eta]\Phi \cdot \mathbf{N} = \underline{\nabla_{\mathbf{x}',z}\varphi} \cdot \mathbf{N}, \\ \mathcal{G}_{gen,II}[\eta]\omega &= \mathbb{U}_{II}[\eta]\omega \cdot \mathbf{N} = \nabla \cdot \mathbf{A}_{\parallel}^{\dagger},\end{aligned}$$

used in Section 6, we parlinearize $\mathcal{G}_{gen,I}[\eta]$ and $\mathcal{G}_{gen,II}[\eta]$ separately. Recall that given a symbol $a \in \Sigma_{n-2}^m(\mathbb{R}^2)$, the corresponding paradifferential operator T_a is given by

$$\mathcal{F}(T_a u)(\xi) = (2\pi)^{-2} \int_{\mathbb{R}^2} \chi(\xi - \xi') \mathcal{F}a(\xi - \xi') \mathcal{F}u(\xi') d\xi',$$

see (B.2).

Regarding the irrotational Dirichlet–Neumann operator $\mathcal{G}_{gen,I}[\eta]$, we recall the following result by Alazard and Métivier (see Theorem 2.12 of [AM09] for a result for the periodic case; as mentioned in Remark 2.14 of the same paper, the result is valid also in the non periodic case).

Proposition 7.1. *Let $n \geq 4$ be such that $n \notin \mathbb{N}$, $\varepsilon > 0$. If $\eta \in H^{n+1/2}(\mathbb{R}^2)$, $\Phi \in \dot{H}^n(\mathbb{R}^2)$, then*

$$(7.1) \quad \mathcal{G}_{gen,I}[\eta]\Phi = T_{\lambda_I}(\Phi - T_{\underline{\mathbb{W}}_I[\eta]\Phi}\eta) - T_{\underline{\mathbb{V}}_I[\eta]\Phi} \cdot \nabla\eta - T_{\text{div}\underline{\mathbb{V}}_I[\eta]\Phi}\eta + \mathcal{R}_I(\eta, \Phi),$$

where the symbol $\lambda_I \in \Sigma_{n-2}^1(\mathbb{R}^2)$. Moreover, we have that

$$\underline{\mathbb{W}}_I[\eta]\Phi \in H^{n-1}(\mathbb{R}^2), \quad \underline{\mathbb{V}}_I[\eta]\Phi \in (H^{n-1}(\mathbb{R}^2))^2,$$

and are given by

$$\underline{\mathbb{W}}_I[\eta]\Phi = \frac{\mathcal{G}[\eta]\Phi + \nabla\eta \cdot \nabla\Phi}{1 + |\nabla\eta|^2}, \quad \underline{\mathbb{V}}_I[\eta]\Phi = \nabla\Phi - (\underline{\mathbb{W}}_I[\eta]\Phi)\nabla\eta,$$

where $\mathcal{G}[\eta]$ denotes the classical Dirichlet–Neumann operator, and $\mathcal{R}_I(\eta, \Phi) \in H^{2n-3-\varepsilon}(\mathbb{R}^2)$ for any $\varepsilon > 0$.

Next, we proceed to study the rotational part $\mathcal{G}_{gen,II}[\eta]$ of the generalized Dirichlet–Neumann operator.

Theorem 8. *Let $n \geq 4$ be such that $n \notin \mathbb{N}$, $\varepsilon > 0$ and assume that $\eta \in H^n(\mathbb{R}^2)$ is such that (1.16) holds true. If*

$$(7.2) \quad \begin{aligned}\partial_w^k \hat{\mathbf{A}} &\in (C^0([-h, 0]; H^{n-k}(\mathbb{R}^2)))^3, \quad k = 0, 1, \\ \hat{\omega} &\in (C^0([-h, 0]; H^{n-2}(\mathbb{R}^2)))^3,\end{aligned}$$

then

$$(7.3) \quad \begin{aligned}\hat{\mathcal{G}}_{gen,II}[\eta]\hat{\omega} &= T_{\lambda_{II}}(\hat{\mathbf{A}} - T_{\partial_w^0 \hat{\mathbf{A}}}\eta) + \frac{1}{\delta} T_{1+|\nabla\eta|^2} T_{\nabla^\perp \eta} \cdot \hat{\mathbf{f}}(\eta, \hat{\omega}) \\ &+ T_{1+|\nabla\eta|^2} \left(\frac{1}{\delta} T_{\hat{\mathbf{A}}_{2xw}} \eta - \frac{1}{\delta} T_{\hat{\mathbf{A}}_{1yw}} \eta - \frac{1}{\delta^2} T_{\eta_x} T_{\hat{\mathbf{A}}_{2ww}} \eta + \frac{1}{\delta^2} T_{\eta_y} T_{\hat{\mathbf{A}}_{1ww}} \eta \right) \\ &- T_{\nabla\eta} \cdot (\hat{\mathbb{V}}_{II}[\eta]\hat{\omega}) - T_{\hat{\mathbb{V}}_{II}[\eta]\hat{\omega}} \cdot \nabla\eta - T_{|\nabla\eta|^2} (\hat{\mathbb{W}}_{II}[\eta]\hat{\omega}) + \mathcal{R}_{II},\end{aligned}$$

where the right-hand side is evaluated at $w = 0$; moreover, $\lambda_{II} \in M_{1,3}(\Sigma_{n-2}^1(\mathbb{R}^2))$ is given by (7.27), $\hat{\mathbf{f}}(\eta, \hat{\omega}) \in (H^{n-1}(\mathbb{R}^2))^2$,

$$\begin{aligned}\hat{\mathbb{V}}_{II}[\eta]\hat{\omega} &= (\text{curl}^e \hat{\mathbf{A}})_h|_{w=0} \in (H^{n-1}(\mathbb{R}^2))^2, \\ \hat{\mathbb{W}}_{II}[\eta]\hat{\omega} &= (\text{curl}^e \hat{\mathbf{A}})_3|_{w=0} \in H^{n-1}(\mathbb{R}^2),\end{aligned}$$

and $\mathcal{R}_{II}(\eta, \hat{\omega}) \in H^{2n-3-\varepsilon}(\mathbb{R}^2)$ for any $\varepsilon > 0$.

Remark 7.2. By Sobolev embedding, Corollary 5.5 and Remark 5.6, formula (7.3) is still valid if we can replace the regularity assumptions (7.2) on η , $\hat{\mathbf{A}}$ and $\hat{\boldsymbol{\omega}}$ with

$$n \geq 5, \quad \eta \in H^{n+\varepsilon}(\mathbb{R}^2), \quad \hat{\boldsymbol{\omega}} \in (H^{n-3/2+\varepsilon}(D_0))^3, \quad \varepsilon > 0.$$

In the rest of the section we prove the above Theorem 8, by following the approach by Alazard and Métivier (see Sec. 4 in [AM09]).

By flattening (4.2) we find that $\hat{\mathbf{A}}$ satisfies

$$(7.4) \quad -\Delta^e \hat{\mathbf{A}} - \hat{\boldsymbol{\omega}} = 0 \quad \text{in } D_0,$$

with boundary conditions

$$(7.5) \quad \hat{\mathbf{A}} \cdot \mathbf{N} = 0, \quad w = 0,$$

$$(7.6) \quad (\text{curl}^e \hat{\mathbf{A}})_{||} = -\nabla^\perp \Delta^{-1}(\hat{\boldsymbol{\omega}} \cdot \mathbf{N}), \quad w = 0.$$

Moreover, in the new coordinates the rotational part of the straightened generalized Dirichlet–Neumann operator can be written as

$$(7.7) \quad \hat{\mathcal{G}}_{gen,II}[\eta]\hat{\boldsymbol{\omega}} = \hat{A}_{2x} + \eta_y \hat{A}_{3x} - \hat{A}_{1y} - \eta_x \hat{A}_{3y}|_{w=0}.$$

Equation (7.4) is equivalent to the system

$$(7.8) \quad E \hat{\mathbf{A}} = -\check{\mathbf{a}} \hat{\boldsymbol{\omega}},$$

where

$$\begin{aligned} E &:= \partial_w^2 + \check{\mathbf{a}} \Delta + \check{\mathbf{b}} \cdot \nabla \partial_w - \check{\mathbf{c}} \partial_w, \\ \check{\mathbf{a}} &= \mathbf{a}_0^{-1}, \quad \check{\mathbf{b}} = \mathbf{b}_0 / \mathbf{a}_0, \quad \check{\mathbf{c}} = \mathbf{c}_0 / \mathbf{a}_0, \\ \mathbf{a}_0 &= \frac{1 + |\nabla \eta|^2}{\delta^2}, \quad \mathbf{b}_0 = -\frac{2\nabla \eta}{\delta}, \quad \mathbf{c}_0 = \frac{\nabla \eta}{\delta}. \end{aligned}$$

Proposition 7.3. *Set*

$$(7.9) \quad \hat{\mathbf{B}} := \hat{\mathbf{A}} - T_{\partial_w^e \hat{\mathbf{A}}} \eta,$$

and let $T_E := \partial_w^2 + (Id + T_{\check{\mathbf{a}}-1})\Delta + T_{\check{\mathbf{b}}} \cdot \nabla \partial_w - T_{\check{\mathbf{c}}} \partial_w$, then $\hat{\mathbf{B}}$ satisfies

$$(7.10) \quad T_E \hat{\mathbf{B}} = \hat{\mathbf{F}}_0 + \hat{\mathbf{F}}_1,$$

where

$$(7.11) \quad \begin{aligned} \hat{\mathbf{F}}_0 &:= \hat{\mathbf{F}}_0(\hat{\boldsymbol{\omega}}, \eta) = -T_{\check{\mathbf{a}}-1} \hat{\boldsymbol{\omega}} - T_{\check{\boldsymbol{\omega}}}(\check{\mathbf{a}} - 1) + T_{\partial_w^e(\check{\mathbf{a}} \hat{\boldsymbol{\omega}})} \eta + T_{[\partial_w^e, E] \hat{\mathbf{A}}} \eta \\ &\quad - (Id + T_{\check{\mathbf{a}}-1}) [T_{\Delta \partial_w^e \hat{\mathbf{A}}} \eta + 2T_{\nabla \partial_w^e \hat{\mathbf{A}}} \cdot \nabla \eta] - T_{\check{\mathbf{b}}} \cdot T_{\partial_w \partial_w^e \hat{\mathbf{A}}} \nabla \eta \\ &\quad - T_{\Delta \hat{\mathbf{A}}}(\check{\mathbf{a}} - 1) - T_{\nabla \partial_w \hat{\mathbf{A}}} \cdot \check{\mathbf{b}} + T_{\partial_w \hat{\mathbf{A}}} \check{\mathbf{c}}, \end{aligned}$$

and $\hat{\mathbf{F}}_1 \in (C^0([-h, 0]; H^{2n-4}(\mathbb{R}^2)))^3$.

Proof. We follow an argument similar to the proof of Proposition 4.12 in [AM09]. First, recall that by assumption (7.2) we have that

$$\begin{aligned} \partial_w^k \hat{\mathbf{A}} &\in (C^0([-h, 0]; H^{n-k}(\mathbb{R}^2)))^3, \quad k = 0, 1, \\ \hat{\boldsymbol{\omega}} &\in (C^0([-h, 0]; H^{n-2}(\mathbb{R}^2)))^3, \end{aligned}$$

and that by the regularity assumption of η we have

$$\check{\mathbf{a}} - 1 \in H^{n-1}(\mathbb{R}^2), \quad \check{\mathbf{b}} \in (H^{n-1}(\mathbb{R}^2))^2, \quad \check{\mathbf{c}} \in H^{n-2}(\mathbb{R}^2).$$

By Lemma B.8 we have that if we set

$$\mathcal{T}(\hat{\mathbf{A}}) := T_{\Delta \hat{\mathbf{A}}}(\check{\mathbf{a}} - 1) + T_{\nabla \partial_w \hat{\mathbf{A}}} \cdot \check{\mathbf{b}} - T_{\partial_w \hat{\mathbf{A}}} \check{\mathbf{c}},$$

then

$$E \hat{\mathbf{A}} + \hat{\boldsymbol{\omega}} + (\check{\mathbf{a}} - 1) \hat{\boldsymbol{\omega}} - T_E \hat{\mathbf{A}} - \mathcal{T}(\hat{\mathbf{A}}) - T_{\check{\mathbf{a}}-1} \hat{\boldsymbol{\omega}} - T_{\check{\boldsymbol{\omega}}}(\check{\mathbf{a}} - 1)$$

belongs to $(C^0([-h, 0]; H^{2n-4}(\mathbb{R}^2)))^3$.

Next, we substitute $\hat{\mathbf{A}} = \hat{\mathbf{B}} + T_{\partial_w^e} \hat{\mathbf{A}} \eta$ in the above interior equation. Recall that $E\hat{\mathbf{A}} = -\check{\mathbf{a}}\hat{\omega}$, and that $\partial_w^k \hat{\mathbf{A}} \in (C^0([-h, 0]; H^{n-k}(\mathbb{R}^2)))^3$ for $k \leq 3$.

From the definition of $\hat{\mathbf{B}}$, we get (up to a smooth remainder)

$$\begin{aligned} \partial_w^2 \hat{\mathbf{B}} &= \partial_w^2 \hat{\mathbf{A}} - \partial_w^2 (T_{\partial_w^e} \hat{\mathbf{A}} \eta) = \partial_w^2 \hat{\mathbf{A}} - T_{\partial_w^2 \partial_w^e} \hat{\mathbf{A}} \eta, \\ \Delta \hat{\mathbf{B}} &= \Delta \hat{\mathbf{A}} - T_{\Delta \partial_w^e} \hat{\mathbf{A}} \eta - 2T_{\nabla \partial_w^e} \hat{\mathbf{A}} \cdot \nabla \eta - T_{\partial_w^e} \hat{\mathbf{A}} \Delta \eta, \\ \nabla \partial_w \hat{\mathbf{B}} &= \nabla \partial_w \hat{\mathbf{A}} - T_{\nabla \partial_w \partial_w^e} \hat{\mathbf{A}} \eta - T_{\partial_w \partial_w^e} \hat{\mathbf{A}} \nabla \eta, \\ \partial_w \hat{\mathbf{B}} &= \partial_w \hat{\mathbf{A}} - T_{\partial_w \partial_w^e} \hat{\mathbf{A}} \eta. \end{aligned}$$

By Lemma B.8 we can deduce from the interior equation $E\hat{\mathbf{A}} + \check{\mathbf{a}}\hat{\omega} = \mathbf{0}$ that

$$\begin{aligned} \eta [\partial_w^e E\hat{\mathbf{A}} + \partial_w^e (\check{\mathbf{a}}\hat{\omega})] &= \mathbf{0}; \\ T_E \partial_w^e \hat{\mathbf{A}} \eta + T_{\partial_w^e (\check{\mathbf{a}}\hat{\omega})} \eta + T_{[\partial_w^e, E] \hat{\mathbf{A}}} \eta &\in (C^0([-h, 0]; H^{2n-4}(\mathbb{R}^2)))^3. \end{aligned}$$

Therefore, we have (up to a smooth remainder)

$$\begin{aligned} T_E \hat{\mathbf{A}} - T_E \hat{\mathbf{B}} &= T_{\partial_w^2 \partial_w^e} \hat{\mathbf{A}} \eta + (Id + T_{\check{\mathbf{a}}-1}) [T_{\Delta \partial_w^e} \hat{\mathbf{A}} \eta + 2T_{\nabla \partial_w^e} \hat{\mathbf{A}} \cdot \nabla \eta + T_{\partial_w^e} \hat{\mathbf{A}} \Delta \eta] \\ &\quad + T_{\check{\mathbf{b}}} \cdot [T_{\nabla \partial_w \partial_w^e} \hat{\mathbf{A}} \eta + T_{\partial_w \partial_w^e} \hat{\mathbf{A}} \nabla \eta] - T_{\check{\mathbf{c}}} T_{\partial_w \partial_w^e} \hat{\mathbf{A}} \eta, \end{aligned}$$

which, up to a remainder in $(C^0([-h, 0]; H^{2n-4}(\mathbb{R}^2)))^3$, it implies that

$$\begin{aligned} T_E \hat{\mathbf{A}} - T_E \hat{\mathbf{B}} &= -T_{\partial_w^e (\check{\mathbf{a}}\hat{\omega})} \eta - T_{[\partial_w^e, E] \hat{\mathbf{A}}} \eta + (Id + T_{\check{\mathbf{a}}-1}) [T_{\Delta \partial_w^e} \hat{\mathbf{A}} \eta + 2T_{\nabla \partial_w^e} \hat{\mathbf{A}} \cdot \nabla \eta] \\ &\quad + T_{\check{\mathbf{b}}} \cdot T_{\partial_w \partial_w^e} \hat{\mathbf{A}} \nabla \eta. \end{aligned}$$

If we plug in the above formula in the parilinearized interior equation we obtain that, up to a remainder in $(C^0([-h, 0]; H^{2n-4}(\mathbb{R}^2)))^3$,

$$\begin{aligned} T_E \hat{\mathbf{B}} &= T_{\partial_w^e (\check{\mathbf{a}}\hat{\omega})} \eta + T_{[\partial_w^e, E] \hat{\mathbf{A}}} \eta - (Id + T_{\check{\mathbf{a}}-1}) [T_{\Delta \partial_w^e} \hat{\mathbf{A}} \eta + 2T_{\nabla \partial_w^e} \hat{\mathbf{A}} \cdot \nabla \eta] \\ &\quad - T_{\check{\mathbf{b}}} \cdot T_{\partial_w \partial_w^e} \hat{\mathbf{A}} \nabla \eta - \mathcal{T}(\hat{\mathbf{A}}) - T_{\check{\mathbf{a}}-1} \hat{\omega} - T_{\check{\mathbf{a}}} (\check{\mathbf{a}} - 1). \end{aligned}$$

□

Next we consider the reduction to the boundary. We want to find two matrix-valued symbols \mathbb{M}, \mathbb{N} such that (up to a smoothing remainder)

$$T_E \mathbb{I}_3 = (\partial_w \mathbb{I}_3 - T_{\mathbb{N}}) (\partial_w \mathbb{I}_3 - T_{\mathbb{M}}),$$

see Sec. 3.4.2 of [ABZ11] and Sec. 4.3 of [AM09] for a corresponding argument for the irrotational case, and Sec. 3.1 of [GNPW24] for the case of Beltrami flows.

Lemma 7.4. *There exist operators M, N with associated symbols*

$$\mathbb{M}, \mathbb{N} \in M_3(\Sigma_{n-2}^1(\mathbb{R}^2))$$

such that

- (i) $T_E \mathbb{I}_3 - (\partial_w \mathbb{I}_3 - T_{\mathbb{N}}) (\partial_w \mathbb{I}_3 - T_{\mathbb{M}}) \in M_3(\Sigma_{n-2}^{1-(n-4)}(\mathbb{R}^2))$,
- (ii) the principal symbols $\mathbb{M}^{(1)}, \mathbb{N}^{(1)}$ of M, N take the form $\mathbb{M}^{(1)} = \mathbf{m}^{(1)} \mathbb{I}_3$, $\mathbb{N}^{(1)} = \mathbf{n}^{(1)} \mathbb{I}_3$, where the scalar-valued symbols $\mathbf{m}^{(1)}, -\mathbf{n}^{(1)} \in \Sigma_{n-2}^1(\mathbb{R}^2)$ are strongly elliptic.

Proof. We first set $L_1 := [\check{\mathbf{b}} \cdot \nabla - \check{\mathbf{c}}] \mathbb{I}_3$, $L_0 := \check{\mathbf{a}} \Delta$, so that $E = \partial_w^2 \mathbb{I}_3 + L_1 \partial_w + L_0$. Recall that $\eta \in H^n(\mathbb{R}^2)$. Since

$$E - (\partial_w \mathbb{I}_3 - N) (\partial_w \mathbb{I}_3 - M) = (L_1 + (M + N)) \partial_w + (L_0 - NM)$$

we set

$$(7.12) \quad N = -L_1 - M$$

and, by following Proposition B.6, we seek M such that

$$L_0 + L_1 M + M^2 = \mathbb{O}_3$$

by constructing a symbol $\mathbf{M} \in M_3(\Sigma_{n-2}^1(\mathbb{R}^2))$ such that

$$\begin{aligned} & -\check{\mathbf{a}}|\boldsymbol{\xi}|^2 \mathbb{I}_3 - \sum_{\alpha=(\alpha_1, \alpha_2) \in \mathbb{N}_0^2} [\partial_{\xi_1}^{\alpha_1} \partial_{\xi_2}^{\alpha_2} |\boldsymbol{\xi}|^2 D_x^{\alpha_1} D_y^{\alpha_2} \check{\mathbf{a}}] \mathbb{I}_3 + [\mathbf{i} \check{\mathbf{b}} \cdot \boldsymbol{\xi} - \check{c}] \mathbf{M} \\ & + (\check{\mathbf{b}} \cdot \nabla) \mathbf{M} + \sum_{\alpha=(\alpha_1, \alpha_2) \in \mathbb{N}^2} \partial_{\xi_1}^{\alpha_1} \partial_{\xi_2}^{\alpha_2} \mathbf{M} D_x^{\alpha_1} D_y^{\alpha_2} \mathbf{M} = \mathbb{O}_3, \end{aligned}$$

up to a remainder in $M_3(\Sigma_{n-2}^{1-(n-4)}(\mathbb{R}^2))$, and such that

$$\mathbf{M} = \sum_{1-(n-4) < j \leq 1} \mathbf{M}^{(j)}, \quad \mathbf{M}^{(j)} \in M_3(\Gamma_{n-3+j}^j(\mathbb{R}^2)).$$

We proceed by computing the terms in the asymptotic expansion of \mathbf{M} inductively.

- At leading order we have

$$-\check{\mathbf{a}}|\boldsymbol{\xi}|^2 \mathbb{I}_3 + \mathbf{i} \check{\mathbf{b}} \cdot \boldsymbol{\xi} \mathbf{M}^{(1)} + (\mathbf{M}^{(1)})^2 = \mathbb{O}_3,$$

so that

$$\mathbf{M}^{(1)} = \mathbf{m}^{(1)} \mathbb{I}_3,$$

where

$$\mathbf{m}^{(1)} := \frac{1}{2} \left[-\mathbf{i} \check{\mathbf{b}} \cdot \boldsymbol{\xi} + \sqrt{-(\check{\mathbf{b}} \cdot \boldsymbol{\xi})^2 + 4\check{\mathbf{a}}|\boldsymbol{\xi}|^2} \right].$$

Note that

$$\mathbf{m}^{(1)} = \delta \frac{\mathbf{i} \boldsymbol{\xi} \cdot \nabla \eta + \lambda^{(1)}}{1 + |\nabla \eta|^2}, \quad \lambda^{(1)} := \sqrt{(1 + |\nabla \eta|^2)|\boldsymbol{\xi}|^2 - (\boldsymbol{\xi} \cdot \nabla \eta)^2},$$

where $\lambda^{(1)}$ is the leading order symbol of the classical Dirichlet–Neumann operator.

- The subprincipal symbol is found from the equation

$$\begin{aligned} & \mathbf{i} (\nabla \check{\mathbf{a}}) \cdot 2\boldsymbol{\xi} \mathbb{I}_3 - \check{c} \mathbf{M}^{(1)} + \mathbf{i} \check{\mathbf{b}} \cdot \boldsymbol{\xi} \mathbf{M}^{(0)} + \mathbf{M}^{(0)} \mathbf{M}^{(1)} + \mathbf{M}^{(1)} \mathbf{M}^{(0)} \\ & + (\check{\mathbf{b}} \cdot \nabla) \mathbf{M}^{(1)} - \mathbf{i} \partial_{\xi_1} \mathbf{M}^{(1)} \partial_x \mathbf{M}^{(1)} - \mathbf{i} \partial_{\xi_2} \mathbf{M}^{(1)} \partial_y \mathbf{M}^{(1)} = \mathbb{O}_3, \end{aligned}$$

which yields

$$\mathbf{M}^{(0)} = \mathbf{m}^{(0)} \mathbb{I}_3,$$

where

$$\begin{aligned} \mathbf{m}^{(0)} &= \frac{1}{\mathbf{i} \check{\mathbf{b}} \cdot \boldsymbol{\xi} + 2\mathbf{m}^{(1)}} \times \\ & \times [\mathbf{i} \nabla_{\boldsymbol{\xi}} \mathbf{m}^{(1)} \cdot \nabla \mathbf{m}^{(1)} - (\check{\mathbf{b}} \cdot \nabla) \mathbf{m}^{(1)} + \check{c} \mathbf{m}^{(1)} - \mathbf{i} (\nabla \check{\mathbf{a}}) \cdot 2\boldsymbol{\xi}]. \end{aligned}$$

- Suppose that $\mathbf{M}^{(j)}$ has been calculated for $j = 1, 0, \dots, -J$ for some $0 \leq J < 1 - (n - 4)$. The term $\mathbf{M}^{(-J-1)}$ can be found from the equation

$$(2\mathbf{m}^{(1)} + \mathbf{i} \check{\mathbf{b}} \cdot \boldsymbol{\xi}) \mathbf{M}^{(-J-1)} = \mathbf{G}^{(-J)},$$

where $\mathbf{G}^{(-J)} \in M_3(\Sigma_{n-2}^{(-J)}(\mathbb{R}^2))$ is given by

$$\begin{aligned} & \mathbf{G}^{(-J)} \\ &= \sum_{|\alpha|=2=J} \partial_{\xi_1}^{\alpha_1} \partial_{\xi_2}^{\alpha_2} (|\xi|^2) D_x^{\alpha_1} D_y^{\alpha_2} \check{\mathbf{a}} + \check{\mathbf{c}} \mathbf{M}^{(-J)} - \check{\mathbf{b}} \cdot \nabla M^{(-J)} \\ & \quad - \sum_{\substack{j_1, j_2 \leq 0 \\ |\alpha|+j_1+j_2=J}} \partial_{\xi_1}^{\alpha_1} \partial_{\xi_2}^{\alpha_2} \mathbf{M}^{(-j_1)} D_x^{\alpha_1} D_y^{\alpha_2} \mathbf{M}^{(-j_2)} - \sum_{|\alpha|=J+2} \partial_{\xi_1}^{\alpha_1} \partial_{\xi_2}^{\alpha_2} \mathbf{M}^{(1)} D_x^{\alpha_1} D_y^{\alpha_2} \mathbf{M}^{(1)} \\ & \quad - \sum_{|\alpha|=J+1} \partial_{\xi_1}^{\alpha_1} \partial_{\xi_2}^{\alpha_2} \mathbf{M}^{(1)} D_x^{\alpha_1} D_y^{\alpha_2} \mathbf{M}^{(0)} - \sum_{|\alpha|=J+1} \partial_{\xi_1}^{\alpha_1} \partial_{\xi_2}^{\alpha_2} \mathbf{M}^{(0)} D_x^{\alpha_1} D_y^{\alpha_2} \mathbf{M}^{(1)}. \end{aligned}$$

The construction is completed by noting that there exists a symbol

$$\mathbf{M} \in M_3(\Sigma_{n-2}^1(\mathbb{R}^2))$$

such that

$$\mathbf{M} = \sum_{1-(n-4) < j \leq 1} \mathbf{M}^{(j)}$$

(see Shubin [Shu01, §3.3]).

Defining N by equation (7.12), we can compute the terms in the asymptotic expansion

$$\mathbf{N} = \sum_{1-(n-4) < j \leq 1} \mathbf{N}^{(j)}$$

of the operator N ; in particular, we find that

$$\mathbf{n}^{(1)} = \mathbf{n}^{(1)} \mathbb{I}_3, \quad \mathbf{n}^{(1)} := \frac{1}{2} \left[-i\check{\mathbf{b}} \cdot \xi - \sqrt{-(\check{\mathbf{b}} \cdot \xi)^2 + 4\check{\mathbf{a}}|\xi|^2} \right],$$

so that

$$\mathbf{n}^{(1)} = \delta \frac{i\xi \cdot \nabla \eta - \lambda^{(1)}}{1 + |\nabla \eta|^2}.$$

Finally, note that

$$\operatorname{Re} \mathbf{n}^{(1)} = -\operatorname{Re} \mathbf{n}^{(1)} = \delta \frac{\lambda^{(1)}}{1 + |\nabla \eta|^2} \geq \delta \frac{|\xi|}{1 + \max |\nabla \eta|^2} \gtrsim \delta \langle \xi \rangle$$

for sufficiently large $|\xi|$, so that $\mathbf{m}^{(1)}$ and $-\mathbf{n}^{(1)}$ are strongly elliptic. \square

Lemma 7.5. *There exist two matrix-valued symbols*

$$\mathbf{M}, \mathbf{N} \in M_3(\Sigma_{n-2}^1(\mathbb{R}^2))$$

such that

$$(7.13) \quad (\partial_w \mathbb{I}_3 - T_{\mathbf{N}}) (\partial_w \mathbb{I}_3 - T_{\mathbf{M}}) \hat{\mathbf{B}} = \hat{\mathbf{F}}_0 + \hat{\mathbf{F}}_1,$$

where $\hat{\mathbf{F}}_0$ is given by (7.11), and where $\hat{\mathbf{F}}_1 \in (C^0([-h, 0]; H^{2n-4}(\mathbb{R}^2)))^3$.

Proof. We argue by following the proof of Lemma 4.13 in [AM09]. We look for \mathbf{M}, \mathbf{N} of the form

$$\begin{aligned} \mathbf{M} &= \mathbf{M}^{(1)}(\mathbf{x}', \xi) + \dots + \mathbf{M}^{(1-(n-3))}(\mathbf{x}', \xi), \\ \mathbf{N} &= \mathbf{N}^{(1)}(\mathbf{x}', \xi) + \dots + \mathbf{N}^{(1-(n-3))}(\mathbf{x}', \xi), \end{aligned} \quad (7.14)$$

$$\mathbf{M}^{(j)}, \mathbf{N}^{(j)} \in M_3(\Gamma_{n-3+j}^j(\mathbb{R}^2)),$$

where the symbol class Γ_{ℓ}^m is defined in B.1. By repeating the same argument of Lemma 7.4 one can check that

$$\mathbf{M}^{(j)} = \mathbf{m}^{(j)} \mathbb{I}_3, \quad \mathbf{N}^{(j)} = \mathbf{n}^{(j)} \mathbb{I}_3, \quad n = 1, 0,$$

and define the lower-order symbols by induction. By construction we also have that

$$\mathbf{M}^{(\ell)}, \mathbf{N}^{(\ell)} \in M_3(\Gamma_{n-3+\ell}^\ell(\mathbb{R}^2)),$$

so that

$$\mathbf{M}, \mathbf{N} \in M_3(\Sigma_{n-2}^1(\mathbb{R}^2)).$$

□

Recalling the definition (7.9) of the good unknown $\hat{\mathbf{B}}$ and the regularity assumption on $\hat{\mathbf{A}}$, we have that

$$\partial_w \hat{\mathbf{B}} - T_{\mathbb{N}} \hat{\mathbf{B}}|_{w=0} \in (H^{n-1}(\mathbb{R}^2))^3.$$

For the irrotational case the elliptic regularity result given by Proposition 4.13 in [AM09] can improve the regularity in the right-hand side of the (analogue of the) above equation up to $H^{2n-3-\varepsilon}(\mathbb{R}^2)$ for any fixed $\varepsilon > 0$ (see Corollary 4.14 of [AM09]). However, this is not true if $\hat{\omega} \neq \mathbf{0}$.

Corollary 7.6. *Let $\varepsilon > 0$. Introduce $\hat{\mathbf{W}} := (\partial_w - T_{\mathbb{N}}) \hat{\mathbf{B}}$ and we write $N = N^{(1)} + N^{(\leq 0)}$, where $N^{(1)}$ is the principal part of N obtained in Lemma 7.5, and $N^{(\leq 0)} \in M_3(\Gamma_0^0(\mathbb{R}^2))$. Then*

$$\partial_w \hat{\mathbf{W}} - T_{N^{(1)}} \hat{\mathbf{W}} = T_{N^{(\leq 0)}} \hat{\mathbf{W}} + \hat{\mathbf{F}}_0 + \hat{\mathbf{F}}_1,$$

where $\hat{\mathbf{F}}_0$ is given by (7.11), and $\hat{\mathbf{F}}_1 \in (C^0([-h, 0]; H^{2n-4}(\mathbb{R}^2)))^3$.

Moreover, up to terms in $(C^0([-h, 0]; H^{2n-3-\varepsilon}(\mathbb{R}^2)))^3$, the following equality holds true,

$$(7.15) \quad \begin{aligned} \partial_w \hat{\mathbf{B}} - T_{\mathbb{N}} \hat{\mathbf{B}}|_{w=0} &= \int_{-h}^0 (T_{\mathcal{E}} \hat{\mathbf{F}}_0)(z) dz, \\ \mathcal{E}(z; \mathbf{x}', \boldsymbol{\xi}) &= \exp(z N^{(1)}(\mathbf{x}', \boldsymbol{\xi})) \\ &\quad - i \exp(z N^{(1)}(\mathbf{x}', \boldsymbol{\xi})) \frac{z^2}{2} \nabla_{\boldsymbol{\xi}} N^{(1)}(\mathbf{x}', \boldsymbol{\xi}) \cdot \nabla_{\mathbf{x}'} N^{(1)}(\mathbf{x}', \boldsymbol{\xi}). \end{aligned}$$

The above corollary can be proved by following the proof of Proposition B.14 (see also Proposition 4.13 in [AM09]).

Now, in order to simplify the notation we omit until the end of the section the fact that all functions depending on w are evaluated on $w = 0$.

First, from (7.5) we have

$$(7.16) \quad \hat{A}_3 = \eta_x \hat{A}_1 + \eta_y \hat{A}_2.$$

Next, we rewrite the boundary condition (7.6) in the following way,

$$(7.17) \quad \begin{aligned} \hat{A}_{3y} - \frac{1}{\delta} \eta_y \hat{A}_{3w} - \frac{1}{\delta} \hat{A}_{2w} + \eta_x \left(\hat{A}_{2x} - \hat{A}_{1y} - \frac{1}{\delta} \eta_x \hat{A}_{2w} + \frac{1}{\delta} \eta_y \hat{A}_{1w} \right) \\ = -\partial_y \Delta^{-1}(\hat{\omega} \cdot \mathbf{N}), \end{aligned}$$

$$(7.18) \quad \begin{aligned} -\hat{A}_{3x} + \frac{1}{\delta} \eta_x \hat{A}_{3w} + \frac{1}{\delta} \hat{A}_{1w} + \eta_y \left(\hat{A}_{2x} - \hat{A}_{1y} - \frac{1}{\delta} \eta_x \hat{A}_{2w} + \frac{1}{\delta} \eta_y \hat{A}_{1w} \right) \\ = \partial_x \Delta^{-1}(\hat{\omega} \cdot \mathbf{N}). \end{aligned}$$

Then if we multiply (7.17) by η_x and (7.18) by η_y and we sum them up, we obtain

$$\begin{aligned} -\frac{1}{\delta} \eta_x (1 + |\nabla \eta|^2) \hat{A}_{2w} + \frac{1}{\delta} \eta_y (1 + |\nabla \eta|^2) \hat{A}_{1w} + (\eta_x \eta_{xy} - \eta_y \eta_{xx}) \hat{A}_1 \\ + (\eta_x \eta_{yy} - \eta_y \eta_{xy}) \hat{A}_2 + \eta_x \eta_y \hat{A}_{2y} + \eta_x^2 \hat{A}_{2x} - \eta_x \eta_y \hat{A}_{1x} - \eta_y^2 \hat{A}_{1y} \\ = -\nabla \eta \cdot \nabla^\perp \Delta^{-1}(\hat{\omega} \cdot \mathbf{N}), \end{aligned}$$

and if we add $\operatorname{curl}^e \hat{\mathbf{A}} \cdot \mathbf{N}$ to both sides of the above equation we obtain

$$(7.19) \quad \begin{aligned} (\operatorname{curl}^e \hat{\mathbf{A}})_3 (1 + |\nabla \eta|^2) &= \operatorname{curl}^e \hat{\mathbf{A}} \cdot \mathbf{N} - \nabla \eta \cdot \nabla^\perp \Delta^{-1}(\hat{\omega} \cdot \mathbf{N}); \\ (\operatorname{curl}^e \hat{\mathbf{A}})_3 &= \partial_x^e \hat{A}_2 - \partial_y^e \hat{A}_1. \end{aligned}$$

Moreover, from the incompressibility we have

$$(7.20) \quad \frac{1}{\delta} \hat{A}_{3w} = -\partial_x^e \hat{A}_1 - \partial_y^e \hat{A}_2.$$

Moreover, plugging (7.16) into (7.17)-(7.18) we obtain

$$(7.21) \quad \begin{aligned} \frac{1}{\delta} (1 + |\nabla \eta|^2) \hat{A}_{1w} &= \eta_{xx} \hat{A}_1 + \eta_{xy} \hat{A}_2 + 2\eta_x \hat{A}_{1x} + \eta_y \hat{A}_{1y} + (\eta_x + \eta_y) \hat{A}_{2y} \\ &\quad - \eta_y \hat{A}_{2x} + \partial_x \Delta^{-1}(\hat{\omega} \cdot \mathbf{N}), \end{aligned}$$

$$(7.22) \quad \begin{aligned} \frac{1}{\delta} (1 + |\nabla \eta|^2) \hat{A}_{2w} &= \eta_{xy} \hat{A}_1 + \eta_{yy} \hat{A}_2 + (\eta_x + \eta_y) \hat{A}_{1x} - \eta_x \hat{A}_{1y} + 2\eta_y \hat{A}_{2y} \\ &\quad + \eta_x \hat{A}_{2x} + \partial_y \Delta^{-1}(\hat{\omega} \cdot \mathbf{N}). \end{aligned}$$

Consider the formula (7.7) for the rotational part $\hat{\mathcal{G}}_{gen,II}[\eta] \hat{\omega}$ of the straightened version of the generalized Dirichlet–Neumann operator. If we plug the boundary condition (7.16) into (7.7) we get

$$(7.23) \quad \begin{aligned} \hat{\mathcal{G}}_{gen,II}[\eta] \hat{\omega} &= (1 + \eta_y^2) \hat{A}_{2x} - (1 + \eta_x^2) \hat{A}_{1y} + (\eta_y \eta_{xx} - \eta_x \eta_{xy}) \hat{A}_1 + \eta_x \eta_y \hat{A}_{1x} \\ &\quad + (\eta_y \eta_{xy} - \eta_x \eta_{yy}) \hat{A}_2 - \eta_x \eta_y \hat{A}_{2y}, \quad \text{at } w = 0. \end{aligned}$$

Finally, we conclude with the parilinearization of the generalized Dirichlet–Neumann operator. By (7.19), we have

$$\hat{\mathcal{G}}_{gen,II}[\eta] \hat{\omega} = (1 + |\nabla \eta|^2) (\partial_x^e \hat{A}_2 - \partial_y^e \hat{A}_1) - \nabla \eta \cdot (\operatorname{curl}^e \hat{\mathbf{A}})_h - (\operatorname{curl}^e \hat{\mathbf{A}})_3 |\nabla \eta|^2 \quad \text{at } w = 0,$$

and if we parilinearize the above expression we get

$$(7.24) \quad \begin{aligned} T_{1+|\nabla \eta|^2} (\partial_x^e \hat{A}_2 - \partial_y^e \hat{A}_1) + T_{\partial_x^e \hat{A}_2 - \partial_y^e \hat{A}_1} |\nabla \eta|^2 - T_{\nabla \eta} \cdot (\hat{\mathbb{V}}_{II}[\eta] \hat{\omega}) \\ - T_{\hat{\mathbb{V}}_{II}[\eta] \hat{\omega}} \cdot \nabla \eta - T_{\hat{\mathbb{W}}_{II}[\eta] \hat{\omega}} |\nabla \eta|^2 - T_{|\nabla \eta|^2} (\hat{\mathbb{W}}_{II}[\eta] \hat{\omega}) - \hat{\mathcal{G}}_{gen,II}[\eta] \hat{\omega} \in H^{2n-3}(\mathbb{R}^2), \end{aligned}$$

where we recall that $\hat{\mathbb{V}}_{II}[\eta] \hat{\omega} = (\operatorname{curl} \hat{\mathbf{A}})_h|_{w=0}$ and $\hat{\mathbb{W}}_{II}[\eta] \hat{\omega} = (\operatorname{curl} \hat{\mathbf{A}})_3|_{w=0}$.

Now, observe that

$$\partial_y^e \hat{A}_1 = \hat{A}_{1y} - \frac{1}{\delta} \eta_y \hat{A}_{1w},$$

hence

$$\partial_y^e \hat{A}_1 + \frac{1}{\delta} T_{\eta_y} \hat{A}_{1w} + \frac{1}{\delta} T_{\hat{A}_{1w}} \eta_y - \hat{A}_{1y} \in H^{2n-3}(\mathbb{R}^2),$$

so that the good unknown $\hat{\mathbf{B}}$ given by (7.9) satisfies

$$\hat{B}_{1y} + \frac{1}{\delta} T_{\hat{A}_{1yw}} \eta - \frac{1}{\delta} T_{\eta_y} \hat{B}_{1w} - \frac{1}{\delta^2} T_{\eta_y} T_{\hat{A}_{1ww}} \eta - \partial_y^e \hat{A}_1 \in H^{2n-3}(\mathbb{R}^2),$$

and similarly

$$\hat{B}_{2x} + \frac{1}{\delta} T_{\hat{A}_{2xw}} \eta - \frac{1}{\delta} T_{\eta_x} \hat{B}_{2w} - \frac{1}{\delta^2} T_{\eta_x} T_{\hat{A}_{2ww}} \eta - \partial_x^e \hat{A}_2 \in H^{2n-3}(\mathbb{R}^2).$$

If we insert the above formulae in (7.24) we get

$$\begin{aligned} \hat{\mathcal{G}}_{gen,II}[\eta]\hat{\omega} &= T_{1+|\nabla\eta|^2} \left(\hat{B}_{2x} - \hat{B}_{1y} + \frac{1}{\delta} T_{\hat{A}_{2xw}} \eta - \frac{1}{\delta} T_{\hat{A}_{1yw}} \eta - \frac{1}{\delta} T_{\eta_x} \hat{B}_{2w} + \frac{1}{\delta} T_{\eta_y} \hat{B}_{1w} \right. \\ &\quad \left. - \frac{1}{\delta^2} T_{\eta_x} T_{\hat{A}_{2ww}} \eta + \frac{1}{\delta^2} T_{\eta_y} T_{\hat{A}_{1ww}} \eta \right) - T_{\nabla\eta} \cdot (\hat{\mathbb{V}}_{II}[\eta]\hat{\omega}) \\ &\quad - T_{\hat{\mathbb{V}}_{II}[\eta]\hat{\omega}} \cdot \nabla\eta - T_{|\nabla\eta|^2} (\hat{\mathbb{W}}_{II}[\eta]\hat{\omega}) + \mathcal{R}_{II}, \end{aligned} \quad (7.25)$$

$$\mathcal{R}_{II} \in H^{2n-3}(\mathbb{R}^2),$$

where by evaluating the interior equation (7.8) at the boundary we have

$$\begin{aligned} \hat{A}_{1ww} &= \frac{\delta^2}{1+|\nabla\eta|^2} \left[-\Delta\hat{A}_1 + \frac{2}{\delta} \nabla\eta \cdot \nabla\hat{A}_{1w} + \frac{1}{\delta} \Delta\eta \nabla\hat{A}_{1w} - \hat{\omega}_1 \right], \\ \hat{A}_{2ww} &= \frac{\delta^2}{1+|\nabla\eta|^2} \left[-\Delta\hat{A}_2 + \frac{2}{\delta} \nabla\eta \cdot \nabla\hat{A}_{2w} + \frac{1}{\delta} \Delta\eta \nabla\hat{A}_{2w} - \hat{\omega}_2 \right]. \end{aligned}$$

Now, recall that Corollary 7.6 implies that for $\varepsilon > 0$

$$\begin{cases} \hat{B}_{1w} - (T_{M_{11}}\hat{B}_1 + T_{M_{12}}\hat{B}_2 + T_{M_{13}}\hat{B}_3) - \int_{-h}^0 (T_{\mathcal{E}}\hat{F}_0)_1(z) dz \\ \in H^{2n-3-\varepsilon}(\mathbb{R}^2) \\ \hat{B}_{2w} - (T_{M_{21}}\hat{B}_1 + T_{M_{22}}\hat{B}_2 + T_{M_{23}}\hat{B}_3) - \int_{-h}^0 (T_{\mathcal{E}}\hat{F}_0)_2(z) dz \\ \in H^{2n-3-\varepsilon}(\mathbb{R}^2) \end{cases}.$$

so that the first term on the right-hand side of (7.25) can be rewritten as

$$\begin{aligned} &T_{\lambda_{II}}\hat{B} + \frac{1}{\delta} T_{1+|\nabla\eta|^2} T_{\nabla^\perp\eta} \cdot \left[\int_{-h}^0 (T_{\mathcal{E}}\hat{F}_0)_h(z) dz \right] \\ (7.26) \quad &+ T_{1+|\nabla\eta|^2} \left(\frac{1}{\delta} T_{\hat{A}_{2xw}} \eta - \frac{1}{\delta} T_{\hat{A}_{1yw}} \eta - \frac{1}{\delta^2} T_{\eta_x} T_{\hat{A}_{2ww}} \eta + \frac{1}{\delta^2} T_{\eta_y} T_{\hat{A}_{1ww}} \eta \right) + \mathcal{R}_{II}, \end{aligned}$$

where $\lambda_{II} \in M_{1,3}(\Sigma_{n-2}^1(\mathbb{R}^2))$ is given by

$$\begin{aligned} \lambda_{II} &= - \begin{pmatrix} 1 & 1 & 0 \end{pmatrix} (1+|\nabla\eta|^2) \times \\ (7.27) \quad &\times \left[\begin{pmatrix} 0 & i\xi_1 & 0 \\ -i\xi_2 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} + \frac{1}{\delta} \begin{pmatrix} T_{\eta_y} T_{M_{11}} & T_{\eta_y} T_{M_{12}} & T_{\eta_y} T_{M_{13}} \\ -T_{\eta_x} T_{M_{21}} & -T_{\eta_x} T_{M_{22}} & -T_{\eta_x} T_{M_{23}} \\ 0 & 0 & 0 \end{pmatrix} \right], \end{aligned}$$

and where $\mathcal{R}_{II} \in H^{2n-3-\varepsilon}(\mathbb{R}^2)$. From (7.25) we finally obtain

$$\begin{aligned} &\hat{\mathcal{G}}_{gen,II}[\eta]\hat{\omega} \\ &= T_{\lambda_{II}}\hat{B} + \frac{1}{\delta} T_{1+|\nabla\eta|^2} T_{\nabla^\perp\eta} \cdot \left[\int_{-h}^0 (T_{\mathcal{E}}\hat{F}_0)_h(z) dz \right] \\ &\quad + T_{1+|\nabla\eta|^2} \left(\frac{1}{\delta} T_{\hat{A}_{2xw}} \eta - \frac{1}{\delta} T_{\hat{A}_{1yw}} \eta - \frac{1}{\delta^2} T_{\eta_x} T_{\hat{A}_{2ww}} \eta + \frac{1}{\delta^2} T_{\eta_y} T_{\hat{A}_{1ww}} \eta \right) \\ &\quad - T_{\nabla\eta} \cdot (\hat{\mathbb{V}}_{II}[\eta]\hat{\omega}) - T_{\hat{\mathbb{V}}_{II}[\eta]\hat{\omega}} \cdot \nabla\eta - T_{|\nabla\eta|^2} (\hat{\mathbb{W}}_{II}[\eta]\hat{\omega}) + \mathcal{R}_{II}, \end{aligned} \quad (7.28)$$

where $\mathcal{R}_{II} \in H^{2n-3-\varepsilon}(\mathbb{R}^2)$.

ACKNOWLEDGEMENTS

The author would like to warmly thank Thomas Alazard and Erik Wahlén for useful comments and suggestions.

This research was supported by the Swedish Research Council under grant no. 2021-06594 while the author was in residence at Institut Mittag-Leffler in Djursholm, Sweden in Autumn 2023.

S. Pasquali is supported by PRIN 2022 ‘‘Turbulent effects vs Stability in Equations from Oceanography’’ (TESEO), project number: 2022HSSYPN. S. Pasquali is supported by INdAM-GNAMPA.

APPENDIX A. PROOF OF PROPOSITION 1.4

We now prove Proposition 1.4.

In order to solve the div-curl problem (1.17), and in particular to deal with the boundary conditions in a more convenient way, we recall the definitions

$$(A.1) \quad H(\operatorname{div}, D_\eta) := \{\mathbf{F} \in (L^2(D_\eta))^3 : \operatorname{div} \mathbf{F} \in L^2(D_\eta)\},$$

$$(A.2) \quad H(\operatorname{curl}, D_\eta) := \{\mathbf{F} \in (L^2(D_\eta))^3 : \operatorname{curl} \mathbf{F} \in (L^2(D_\eta))^3\}.$$

It follows that if $\boldsymbol{\omega} \in (L^2(D_\eta))^3$ and \mathbf{U} solves the first two equations of (1.17), one can take the normal and the tangential traces at the boundary, from which we derive the boundary conditions of (1.17).

First, it is easy to reduce (1.17) to the case $\Phi = 0$. Indeed, let $\varphi \in \dot{H}^2(D_\eta)$ be the solution of the boundary value problem

$$(A.3) \quad \begin{cases} \Delta_{\mathbf{x}', z} \varphi = 0, & \text{in } D_\eta, \\ \varphi = \Phi, & \text{at } z = \eta, \\ \partial_n \varphi = 0, & \text{at } z = -h, \end{cases}$$

(its existence is guaranteed by the results in Sec. 2.1.3 and Sec. 2.5.2 of [Lan13]), then, if we define $\mathbf{v} = \mathbf{U} - \nabla_{\mathbf{x}', z} \varphi$, we have

$$(A.4) \quad \begin{cases} \operatorname{curl} \mathbf{v} = \boldsymbol{\omega}, & \text{in } D_\eta, \\ \operatorname{div} \mathbf{v} = 0 & \text{in } D_\eta, \\ \mathbf{v}_\parallel = \nabla^\perp \Psi & \text{at } z = \eta, \\ \mathbf{v} \cdot \mathbf{e}_3 = 0, & \text{at } z = -h, \end{cases}$$

which is equivalent to (1.17) with $\Phi = 0$. We look for a solution of (A.4) of the form $\mathbf{v} = \operatorname{curl} \mathbf{A}$, where the vector potential \mathbf{A} satisfies

$$(A.5) \quad \begin{cases} \operatorname{curl} \operatorname{curl} \mathbf{A} = \boldsymbol{\omega}, & \text{in } D_\eta, \\ \mathbf{A} \times \mathbf{e}_3 = \mathbf{0}, & \text{at } z = -h, \\ \mathbf{A} \cdot \mathbf{N} = 0, & \text{at } z = \eta, \\ (\operatorname{curl} \mathbf{A})_\parallel = -\nabla^\perp \Delta^{-1}(\boldsymbol{\omega} \cdot \mathbf{N}), & \text{at } z = \eta. \end{cases}$$

Before proving the existence of a vector potential \mathbf{A} of (A.5), we prove some technical lemmas.

Lemma A.1. *Assume that $\eta \in C^1(\mathbb{R}^2)$ satisfies condition (1.16). Let also $\mathbf{A} \in H(\operatorname{div}, D_\eta) \cap H(\operatorname{curl}, D_\eta)$ be such that*

$$\begin{cases} \mathbf{A} \times \mathbf{e}_3 = \mathbf{0}, & \text{at } z = -h, \\ \mathbf{A} \cdot \mathbf{N} = 0, & \text{at } z = \eta. \end{cases}$$

Then

$$\|\mathbf{A}\|_{(L^2(D_\eta))^3} \leq C(h, \|\eta\|_{C^1(\mathbb{R}^2)}) \|\partial_z \mathbf{A}\|_{(L^2(D_\eta))^3}.$$

Proof. Let $\Pi_I(\mathbf{x}') : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be the projection onto $\text{span}(\mathbf{N}(\mathbf{x}'))$ parallel to $\text{span}(\mathbf{e}_1, \mathbf{e}_2)$, and $\Pi_{II}(\mathbf{x}') : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be the projection onto $\text{span}(\mathbf{e}_1, \mathbf{e}_2)$ parallel to $\text{span}(\mathbf{N}(\mathbf{x}'))$. One can write the following decomposition

$$\mathbf{A} = \Pi_I \mathbf{A} + \Pi_{II} \mathbf{A},$$

when $\Pi_I \mathbf{A}$ vanishes at $z = \eta$, and $\Pi_{II} \mathbf{A}$ vanishes at $z = -h$. Therefore

$$(A.6) \quad \begin{aligned} \|\mathbf{A}\|_{(L^2(D_\eta))^3} &\leq \|\Pi_I \mathbf{A}\|_{(L^2(D_\eta))^3} + \|\Pi_{II} \mathbf{A}\|_{(L^2(D_\eta))^3} \\ &\leq \|h + \eta\|_{L^\infty(\mathbb{R}^2)} \left[\|\partial_z \Pi_I \mathbf{A}\|_{(L^2(D_\eta))^3} + \|\partial_z \Pi_{II} \mathbf{A}\|_{(L^2(D_\eta))^3} \right]. \end{aligned}$$

Since the projectors do not depend on the vertical direction, and since the norms of the projectors is bounded by $C(\|\eta\|_{C^1(\mathbb{R}^2)})$, by exploiting Lemma B.10 we can deduce the thesis. \square

Lemma A.2. *Let $k > \frac{5}{2}$, let $\eta \in H^k(\mathbb{R}^2)$ satisfies condition (1.16). Let also $\mathbf{A} \in H(\text{div}, D_\eta) \cap H(\text{curl}, D_\eta)$ be such that*

$$\begin{cases} \mathbf{A} \times \mathbf{e}_3 = \mathbf{0}, & \text{at } z = -h, \\ \mathbf{A} \cdot \mathbf{N} = 0, & \text{at } z = \eta. \end{cases}$$

Then there exists $C > 0$ such that

$$\begin{aligned} &\|\nabla_{\mathbf{x}', z} \mathbf{A}\|_{(L^2(D_\eta))^{3 \times 3}}^2 \\ &\leq \|\text{curl } \mathbf{A}\|_{(L^2(D_\eta))^3}^2 + \|\text{div } \mathbf{A}\|_{L^2(D_\eta)}^2 + C \|\eta\|_{H^k(\mathbb{R}^2)} \|\underline{\mathbf{A}}\|_{(L^2(\mathbb{R}^2))^3}^2. \end{aligned}$$

Proof. Here we adopt the convention of summing over repeated indices and we write $(\partial_i)_{i=1,2,3} =: \nabla_{\mathbf{x}', z}$. By arguing as in the proof of Lemma 3.2 of [CL15] we obtain

$$(A.7) \quad \begin{aligned} \|\nabla_{\mathbf{x}', z} \mathbf{A}\|_{(L^2(D_\eta))^{3 \times 3}}^2 &= \sum_{i,j=1}^3 \int_{D_\eta} |\partial_j A_i|^2 \\ &= \|\text{curl } \mathbf{A}\|_{(L^2(D_\eta))^3}^2 + \int_{D_\eta} \partial_j A_i \partial_i A_j. \end{aligned}$$

Following Sec. 3.1.1 of [ABZ14] (see also Proposition 4.1 and Lemma 4.3 of [WZZZ21]), we rewrite the last integral exploiting a regularizing diffeomorphism in (1.31),

$$\begin{aligned} &\int_{D_0} \partial_j^\Sigma \tilde{A}_i \partial_i^\Sigma \tilde{A}_j (1 + \sigma_w) \\ &= - \int_{D_0} \tilde{A}_i \partial_j^\Sigma \partial_i^\Sigma \tilde{A}_j (1 + \sigma_w) - \sigma_{wj} \tilde{A}_i \partial_i^\Sigma \tilde{A}_j \\ &\quad + \int_{\partial D_0} \left[n_j \tilde{A}_i \partial_i^\Sigma \tilde{A}_j - \frac{\sigma_j}{1 + \sigma_w} \tilde{A}_i \partial_i^\Sigma \tilde{A}_j \right] (1 + \sigma_w), \end{aligned}$$

where we are using the notation of Sec. 3. Notice that

$$\begin{aligned}
& \left| \int_{D_0} \tilde{A}_i \partial_j^\Sigma \partial_i^\Sigma \tilde{A}_j (1 + \sigma_w) \right| \\
& \leq \left| - \int_{D_0} \partial_i^\Sigma \tilde{A}_i \partial_j^\Sigma \tilde{A}_j (1 + \sigma_w) + \sigma_{wi} \tilde{A}_i \partial_j^\Sigma \tilde{A}_j \right| \\
& \quad + \left| \int_{\partial D_0} \left[n_i \tilde{A}_i \partial_j^\Sigma \tilde{A}_j - \frac{\sigma_i}{1 + \sigma_w} \tilde{A}_i \partial_j^\Sigma \tilde{A}_j \right] (1 + \sigma_w) \right| \\
& \quad + C(\|\sigma\|_{C^1(D_0)}) \|\tilde{\mathbf{A}}\|_{(L^2(D_0))^3}^3 \\
& \stackrel{(1.34)}{\leq} \|\operatorname{div} \mathbf{A}\|_{L^2(D_\eta)}^2 + \|\eta\|_{H^k(\mathbb{R}^2)} \|\partial_j^\Sigma \tilde{A}_j\|_{L^2(D_0)} \|\tilde{A}_i\|_{L^2(D_0)} \\
& \quad + C(\|\sigma\|_{C^1(D_0)}) \left[\left| \int_{\partial D_0} n_i \tilde{A}_i \partial_j^\Sigma \tilde{A}_j \right| + \|\tilde{\mathbf{A}}\|_{(L^2(D_0))^3}^2 \right].
\end{aligned}$$

Hence,

$$\begin{aligned}
& \|\nabla_{\mathbf{x}',z} \mathbf{A}\|_{(L^2(D_\eta))^{3 \times 3}}^2 \\
& \leq \|\operatorname{curl} \mathbf{A}\|_{(L^2(D_\eta))^3}^2 + \|\operatorname{div} \mathbf{A}\|_{L^2(D_\eta)}^2 \\
& \quad + 2\|\eta\|_{H^k(\mathbb{R}^2)} \|\partial_j^\Sigma \tilde{A}_j\|_{L^2(D_0)} \|\tilde{A}_i\|_{L^2(D_0)} \\
& \quad + C(\|\sigma\|_{C^1(D_0)}) \left[\left| \int_{\partial D_0} n_i \tilde{A}_i \partial_j^\Sigma \tilde{A}_j \right| + \left| \int_{\partial D_0} n_j \tilde{A}_i \partial_i^\Sigma \tilde{A}_j \right| + \|\tilde{\mathbf{A}}\|_{(L^2(D_0))^3}^2 \right]
\end{aligned}$$

Therefore, by exploiting that $\mathbf{A} \in H(\operatorname{div}, D_\eta)$ and Lemma A.1

$$\begin{aligned}
& \|\nabla_{\mathbf{x}',z} \mathbf{A}\|_{(L^2(D_\eta))^{3 \times 3}}^2 \\
& \leq (1 + C(h, \|\sigma\|_{C^1(D_0)})) \left\{ \|\operatorname{curl} \mathbf{A}\|_{(L^2(D_\eta))^3}^2 + \|\operatorname{div} \mathbf{A}\|_{L^2(D_\eta)}^2 \right. \\
& \quad \left. + C(\|\sigma\|_{C^1(D_0)}) \left[\left| \int_{\partial D_0} n_i \tilde{A}_i \partial_j^\Sigma \tilde{A}_j \right| + \left| \int_{\partial D_0} n_j \tilde{A}_i \partial_i^\Sigma \tilde{A}_j \right| \right] \right\}.
\end{aligned}$$

Now we evaluate the boundary integral on the right-hand side of (A.7). When we consider the bottom contribution, we observe that

$$(\mathbf{n} \cdot \tilde{\mathbf{A}}) \operatorname{div}^\Sigma \tilde{\mathbf{A}}|_{w=-h} = \tilde{\mathbf{A}} \cdot \partial_n^\Sigma \tilde{\mathbf{A}},$$

because the bottom is flat. Since

$$\int_{\{w=-h\}} n_i \tilde{A}_j (\partial_j^\Sigma \tilde{A}_i) = \int_{\{w=-h\}} \tilde{\mathbf{A}} \cdot \partial_n^\Sigma \tilde{\mathbf{A}}$$

because $\tilde{\mathbf{A}}$ is normal to the bottom, we obtain that

$$\int_{\{w=-h\}} n_i \tilde{A}_j (\partial_j^\Sigma \tilde{A}_i) - (\mathbf{n} \cdot \tilde{\mathbf{A}}) \operatorname{div}^\Sigma \tilde{\mathbf{A}} = 0.$$

If we consider the contribution at the free surface, by using (1.34) we have that there exists $C > 0$ such that

$$\int_{\{w=0\}} n_j \tilde{A}_i (\partial_i^\Sigma \tilde{A}_j) \leq C \|\eta\|_{H^k(\mathbb{R}^2)} \|\tilde{\mathbf{A}}|_{w=0}\|_{(L^2(\mathbb{R}^2))^3}^2.$$

Hence, we can deduce that there exists $C > 0$ such that

$$\int_{\{w=0\}} n_i \tilde{A}_j (\partial_j^\Sigma \tilde{A}_i) - (\mathbf{n} \cdot \tilde{\mathbf{A}}) \operatorname{div}^\Sigma \tilde{\mathbf{A}} \leq C \|\eta\|_{H^k(\mathbb{R}^2)} \|\tilde{\mathbf{A}}|_{w=0}\|_{(L^2(\mathbb{R}^2))^3}^2.$$

□

Lemma A.3. Let $k > \frac{5}{2}$, and assume that $\eta \in H^k(\mathbb{R}^2)$ satisfies condition (1.16). Let also $\mathbf{A} \in H(\operatorname{div}, D_\eta) \cap H(\operatorname{curl}, D_\eta)$ be such that

$$\begin{cases} \mathbf{A} \times \mathbf{e}_3 = \mathbf{0}, & \text{at } z = -h, \\ \mathbf{A} \cdot \mathbf{N} = 0, & \text{at } z = \eta. \end{cases}$$

Then there exists $C > 0$ such that

$$\begin{aligned} & \|\mathbf{A}\|_{(L^2(D_\eta))^3} + \|\nabla_{\mathbf{x}', z} \mathbf{A}\|_{(L^2(D_\eta))^{3 \times 3}}^2 \\ & \leq C(h, \|\eta\|_{H^k(\mathbb{R}^2)}) \left(\|\operatorname{curl} \mathbf{A}\|_{(L^2(D_\eta))^3}^2 + \|\operatorname{div} \mathbf{A}\|_{L^2(D_\eta)}^2 \right). \end{aligned}$$

Proof. From the assumption that \mathbf{A} is tangential at the free surface we can deduce that $\underline{\mathbf{A}}_3 = \underline{\mathbf{A}}_h \cdot \nabla \eta$, hence there exists $C > 0$ such that

$$\|\underline{\mathbf{A}}\|_{(L^2(\mathbb{R}^2))^3} \leq C \|\eta\|_{C^1(\mathbb{R}^2)} \|\underline{\mathbf{A}}_h\|_{(L^2(\mathbb{R}^2))^3}.$$

Now we estimate the L^2 -norm of $\underline{\mathbf{A}}_h$: using the fact that $\mathbf{A}_h|_{z=-h} = \mathbf{0}$, we have

$$\int_{\mathbb{R}^2} |\underline{\mathbf{A}}_h|^2 = \int_{D_\eta} \partial_z |\mathbf{A}_h|^2 = 2 \int_{D_\eta} \mathbf{A}_h \cdot \partial_z \mathbf{A}_h,$$

and by observing that $\partial_z \mathbf{A}_h = (\operatorname{curl} \mathbf{A})_h^\perp + \nabla A_3$, we get

$$\begin{aligned} \int_{\mathbb{R}^2} |\underline{\mathbf{A}}_h|^2 &= 2 \int_{D_\eta} \mathbf{A}_h \cdot (\operatorname{curl} \mathbf{A}_h)^\perp + 2 \int_{D_\eta} \mathbf{A}_h \cdot \nabla A_3 \\ &= 2 \int_{D_\eta} \mathbf{A}_h \cdot (\operatorname{curl} \mathbf{A}_h)^\perp - 2 \int_{D_\eta} (\nabla \cdot \mathbf{A}_h) A_3 - 2 \int_{\mathbb{R}^2} (\nabla \eta \cdot \underline{\mathbf{A}}_h) \underline{\mathbf{A}}_3, \end{aligned}$$

where

$$-2 \int_{\mathbb{R}^2} (\nabla \cdot \mathbf{A}_h) A_3 = -2 \int_{D_\eta} (\operatorname{div} \mathbf{A}) A_3 + 2 \int_{D_\eta} (\partial_z A_3) A_3.$$

Recalling that $\underline{\mathbf{A}}_3 = \nabla \eta \cdot \underline{\mathbf{A}}_h$, we obtain

$$\int_{\mathbb{R}^2} |\underline{\mathbf{A}}_h|^2 = 2 \int_{D_\eta} \mathbf{A}_h \cdot (\operatorname{curl} \mathbf{A}_h)^\perp - 2 \int_{D_\eta} (\operatorname{div} \mathbf{A}) A_3 - \int_{\mathbb{R}^2} (\mathbf{A}_3|_{z=-h})^2 - \int_{\mathbb{R}^2} (\underline{\mathbf{A}}_3)^2,$$

from which we deduce that there exists $C > 0$ such that

$$\begin{aligned} & \|\underline{\mathbf{A}}\|_{(L^2(\mathbb{R}^2))^3} \\ & \leq C \left[\|\mathbf{A}\|_{(L^2(D_\eta))^3} \|\operatorname{curl} \mathbf{A}\|_{(L^2(D_\eta))^3} + \|\operatorname{div} \mathbf{A}\|_{L^2(D_\eta)} \|\mathbf{A}\|_{(L^2(D_\eta))^3} \|\eta\|_{C^{1+\varepsilon}(\mathbb{R}^2)} \right] \\ & \leq C(h, \|\eta\|_{C^1(\mathbb{R}^2)}) \|\nabla_{\mathbf{x}', z} \mathbf{A}\|_{(L^2(D_\eta))^{3 \times 3}} \left[\|\operatorname{curl} \mathbf{A}\|_{(L^2(D_\eta))^3} + \|\operatorname{div} \mathbf{A}\|_{L^2(D_\eta)} \right], \end{aligned}$$

and by combining Lemma A.2 with Young's inequality for products we get the thesis. \square

We now construct a solution to (A.5), imposing in addition that \mathbf{A} is divergence free. More explicitly, we study the boundary value problem

$$(A.8) \quad \begin{cases} \operatorname{curl} \operatorname{curl} \mathbf{A} = \boldsymbol{\omega}, & \text{in } D_\eta, \\ \operatorname{div} \mathbf{A} = 0, & \text{in } D_\eta, \\ \mathbf{A} \times \mathbf{e}_3 = \mathbf{0}, & \text{at } z = -h, \\ \mathbf{A} \cdot \mathbf{N} = 0, & \text{at } z = \eta, \\ (\operatorname{curl} \mathbf{A})_\parallel = \nabla^\perp \Psi, & \text{at } z = \eta, \end{cases}$$

with $\Psi \in \dot{H}^{1/2}(\mathbb{R}^2)$. Instead of adopting an argument analogous to Definition 3.4 - Lemma 3.6 in [CL15], we restate the boundary value problem (1.21) in an equivalent way, along the lines of the approach outlined in Sec. 4 of [GH20] for Beltrami flows; this will allow us

to obtain a solution of (1.21) by solving (the flattened version of) an equivalent problem, whose solution can be explicitly written in terms of a Green matrix.

Let $k > \frac{5}{2}$, and let $\eta \in H^k(\mathbb{R}^2)$ be such that (1.16) holds true. Let us also assume that $\omega \in H_b(\text{div}_0, D_\eta)$ and that $\Phi \in \dot{H}^{3/2}(\mathbb{R}^2)$.

Let us denote by \mathfrak{X}_η the following subspace of $(H^1(D_\eta))^3$,

$$(A.9) \quad \mathfrak{X}_\eta := \{\mathbf{F} \in (H^1(D_\eta))^3 : \mathbf{F} \times \mathbf{e}_3|_{z=-h} = \mathbf{0}, \mathbf{F} \cdot \mathbf{n} = 0\}.$$

By Corollary A.4 we introduce the following notion: we call a *weak solution* of (1.21) a function $\mathbf{A} \in \mathfrak{X}_\eta$ such that

$$(A.10) \quad \int_{D_\eta} \text{curl } \mathbf{A} \cdot \text{curl } \mathbf{B} + \text{div } \mathbf{A} \text{ div } \mathbf{B} = \int_{D_\eta} \omega \cdot \mathbf{B} + \int_{\mathbb{R}^2} \nabla \Delta^{-1}(\underline{\omega} \cdot \mathbf{N}) \cdot \mathbf{B}_\parallel, \\ \forall \mathbf{B} \in \mathfrak{X}_\eta,$$

while we call a *strong solution* of (1.21) a weak solution such that $\mathbf{A} \in (H^2(D_\eta))^3$ is solenoidal and satisfies

$$\text{curl curl } \mathbf{A} = \omega, \quad \text{in } D_\eta,$$

in $(L^2(D_\eta))^3$, and

$$(\text{curl } \mathbf{A})_\parallel = -\nabla^\perp \Delta^{-1}(\underline{\omega} \cdot \mathbf{N})$$

in $(H^{1/2}(\mathbb{R}^2))^2$.

We mention an immediate consequence of Lemma A.2 regarding a characterization of a closed subspace of $(H^1(D_\eta))^3$; such result will be used in Sec. 6 for the homogeneous expansion of the generalized Dirichlet–Neumann operator (see also Proposition 4.2 in [GH20] for an analogous result applied to Beltrami flows).

Corollary A.4. *Let $k > \frac{5}{2}$, and let $\eta \in H^k(\mathbb{R}^2)$ be such that (1.16) holds true. Then*

$$\mathfrak{X}_\eta = \left\{ \mathbf{F} \in (L^2(D_\eta))^3 : \text{curl } \mathbf{F} \in (L^2(D_\eta))^3, \text{div } \mathbf{F} \in L^2(D_\eta), \right. \\ \left. \mathbf{F} \times \mathbf{e}_3|_{z=-h} = \mathbf{0}, \mathbf{F} \cdot \mathbf{n} = 0 \right\},$$

and the function $\mathbf{F} \mapsto (\|\text{curl } \mathbf{F}\|_{(L^2(D_\eta))^3} + \|\text{div } \mathbf{F}\|_{L^2(D_\eta)})^{1/2}$ is equivalent to its usual norm.

We now discuss the existence of weak and strong solutions for (1.21). First we state the following auxiliary results (for the proof see Proposition 4.3 and Proposition 4.6 in [GH20]).

Lemma A.5. (i) *The function $\mathbf{F} \mapsto \underline{\mathbf{F}}$ is a continuous linear mapping $(H^1(D_\eta))^3 \rightarrow (H^{1/2}(\mathbb{R}^2))^3$ with continuous right inverse $(H^{1/2}(\mathbb{R}^2))^3 \rightarrow \{\mathbf{F} \in (H^1(D_\eta))^3 : \mathbf{F}|_{z=-h} = \mathbf{0}\}$.*

(ii) *The mapping $\mathbf{F} \rightarrow \mathbf{F}_\parallel^\perp$ defined on $\mathcal{D}(\overline{D_\eta})^3$ extends to a continuous linear mapping*

$$\{\mathbf{F} \in (L^2(D_\eta))^3 : \text{curl } \mathbf{F} \in (L^2(D_\eta))^3\} \rightarrow (H^{-1/2}(\mathbb{R}^2))^2,$$

where the former space is equipped with the norm

$$\mathbf{F} \mapsto (\|\mathbf{F}\|_{(L^2(D_\eta))^3}^2 + \|\text{curl } \mathbf{F}\|_{(L^2(D_\eta))^3}^2)^{1/2}.$$

Lemma A.6. (i) *The spaces*

$$\left\{ \mathbf{F} \in (L^2(D_\eta))^3 : \text{curl } \mathbf{F} \in (L^2(D_\eta))^3, \text{div } \mathbf{F} \in L^2(D_\eta), \right. \\ \left. \mathbf{F} \times \mathbf{e}_3|_{z=-h} \in (H^{1/2}(\mathbb{R}^2))^3, \underline{\mathbf{F}} \cdot \mathbf{n} \in H^{1/2}(\mathbb{R}^2) \right\}$$

and

$$\left\{ \mathbf{F} \in (L^2(D_\eta))^3 : \operatorname{curl} \mathbf{F} \in (L^2(D_\eta))^3, \operatorname{div} \mathbf{F} \in L^2(D_\eta), \right. \\ \left. \mathbf{F} \times \mathbf{e}_3|_{z=-h} \in (H^{1/2}(\mathbb{R}^2))^3, \mathbf{F}_\parallel^\perp \in H^{1/2}(\mathbb{R}^2) \right\}$$

coincide with $(H^1(D_\eta))^3$.

(ii) The space

$$\left\{ \mathbf{F} \in (L^2(D_\eta))^3 : \operatorname{curl} \mathbf{F} \in (H^1(D_\eta))^3, \operatorname{div} \mathbf{F} \in H^1(D_\eta), \right. \\ \left. \mathbf{F} \times \mathbf{e}_3|_{z=-h} = \mathbf{0}, \underline{\mathbf{F}} \cdot \mathbf{n} = 0 \right\}$$

coincides with

$$\{ \mathbf{F} \in (H^2(D_\eta))^3 : \mathbf{F} \times \mathbf{e}_3|_{z=-h} = \mathbf{0}, \underline{\mathbf{F}} \cdot \mathbf{n} = 0 \}.$$

We also recall the vector calculus identity

$$\int_{D_\eta} \mathbf{F} \cdot \operatorname{curl} \mathbf{G} - \operatorname{curl} \mathbf{F} \cdot \mathbf{G} = \int_{\mathbb{R}^2} \mathbf{F}_\parallel^\perp \cdot \mathbf{G}_\parallel, \quad \forall \mathbf{F} \in (\mathcal{D}(\overline{D_\eta}))^3, \\ \forall \mathbf{G} \in (H^1(D_\eta))^3, \text{ s.t. } \mathbf{G}|_{z=-h} = \mathbf{0},$$

see pag.11 of [GH20].

Now we prove the existence of weak and strong solutions for (1.21), together with an alternative characterization of strong solutions.

Proposition A.7. *Let $\boldsymbol{\omega} \in H_b(\operatorname{div}_0, D_\eta)$ and $\Phi \in \dot{H}^{1/2}(\mathbb{R}^2)$, then the boundary value problem (1.21) admits a weak solution \mathbf{A} . Moreover, the solution \mathbf{A} is solenoidal, it satisfies*

$$\operatorname{curl} \operatorname{curl} \mathbf{A} = \boldsymbol{\omega}, \quad \text{in } D_\eta,$$

in the sense of distributions, and it satisfies

$$(\operatorname{curl} \mathbf{A})_\parallel = -\nabla^\perp \Delta^{-1}(\underline{\boldsymbol{\omega}} \cdot \mathbf{N})$$

in $(H^{-1/2}(\mathbb{R}^2))^2$.

Proof. Since there exists $C > 0$ such that

$$\left| \int_{D_\eta} \operatorname{curl} \mathbf{A} \cdot \mathbf{B} \right| + \left| \int_{D_\eta} \operatorname{div} \mathbf{A} \operatorname{div} \mathbf{B} \right| \\ \leq C \left(\|\operatorname{curl} \mathbf{A}\|_{(L^2(D_\eta))^3} \|\operatorname{curl} \mathbf{B}\|_{(L^2(D_\eta))^3} + \|\operatorname{div} \mathbf{A}\|_{L^2(D_\eta)} \|\operatorname{div} \mathbf{B}\|_{L^2(D_\eta)} \right)$$

then the left-hand side of (A.10) is a continuous, coercive, bilinear form from $\mathfrak{X}_\eta \times \mathfrak{X}_\eta$ to \mathbb{R} . On the other hand, there exist $C_1, C_2 > 0$ (all depending on h and $\|\eta\|_{C^{3/2+\varepsilon}(\mathbb{R}^2)}$, where $0 < \varepsilon < \min(\frac{1}{2}, k - \frac{5}{2})$) such that

$$\left| \int_{D_\eta} \boldsymbol{\omega} \cdot \mathbf{B} \right| \leq C_1 \|\boldsymbol{\omega}\|_{(L^2(D_\eta))^3} \|\mathbf{B}\|_{(L^2(D_\eta))^3}, \\ \left| \int_{\mathbb{R}^2} \nabla \Delta^{-1}(\underline{\boldsymbol{\omega}} \cdot \mathbf{N}) \cdot \mathbf{B}_\parallel \right| \leq \|\nabla \Delta^{-1}(\underline{\boldsymbol{\omega}} \cdot \mathbf{N})\|_{(H^{-1/2}(\mathbb{R}^2))^2} \|\mathbf{B}_\parallel\|_{(H^{1/2}(\mathbb{R}^2))^2} \\ \leq C_2 \|\boldsymbol{\omega}\|_{(L^2(D_\eta))^3} \|\mathbf{B}\|_{(H^1(D_\eta))^3},$$

hence the right-hand side of (A.10) is a continuous, bilinear form from $(L^2(D_\eta))^3 \times (\dot{H}^{1/2}(\mathbb{R}^2))^2 \times \mathfrak{X}_\eta$ to \mathbb{R} (note that we used Lemma A.6 in the above estimates). The existence of a unique solution $\mathbf{A} \in \mathfrak{X}_\eta$ follows from the Lax-Milgram lemma.

Let $\varphi_{\mathbf{A}} \in H^2(D_\eta)$ be the unique function satisfying

$$\begin{aligned}\Delta\varphi_{\mathbf{A}} &= \operatorname{div} \mathbf{A}, \quad \text{in } D_\eta, \\ \partial_n \varphi_{\mathbf{A}} &= 0, \quad \text{at } z = \eta, \\ \varphi_{\mathbf{A}} &= 0, \quad \text{at } z = -h,\end{aligned}$$

(its existence is ensured by Corollary 2.44 in [Lan13]); then we have that $\mathbf{B} = \nabla_{x',z}\varphi_{\mathbf{A}} \in \mathfrak{X}_\eta$, hence $\mathbf{B}_\parallel = \nabla(\underline{\varphi}_{\mathbf{A}})$, and from (A.10) we get

$$\int_{D_\eta} (\operatorname{div} \mathbf{A})^2 = \int_{D_\eta} \boldsymbol{\omega} \cdot \nabla_{x',z}\varphi_{\mathbf{A}} + \int_{\mathbb{R}^2} \nabla \Delta^{-1}(\underline{\boldsymbol{\omega}} \cdot \mathbf{N}) \cdot \nabla(\underline{\varphi}_{\mathbf{A}}),$$

and since

$$\int_{D_\eta} \boldsymbol{\omega} \cdot \nabla_{x',z}\varphi_{\mathbf{A}} = - \int_{\mathbb{R}^2} \nabla \Delta^{-1}(\underline{\boldsymbol{\omega}} \cdot \mathbf{N}) \cdot \nabla(\underline{\varphi}_{\mathbf{A}}),$$

we can conclude that $\operatorname{div} \mathbf{A} = 0$ in D_η . Next, if we take $\mathbf{B} \in (\mathcal{D}(D_\eta))^3$, we can check that \mathbf{A} solves

$$\operatorname{curl} \operatorname{curl} \mathbf{A} = \boldsymbol{\omega}, \quad \text{in } D_\eta,$$

in the sense of distributions, and hence that $\operatorname{curl} \operatorname{curl} \mathbf{A} \in (L^2(D_\eta))^3$. Therefore by Lemma A.6 we have that $(\operatorname{curl} \mathbf{A})_\parallel^\perp \in (H^{1/2}(\mathbb{R}^2))^2$, and since we have

$$\int_{D_\eta} \operatorname{curl} \mathbf{A} \cdot \mathbf{B} = \int_{D_\eta} \boldsymbol{\omega} \cdot \mathbf{B} + \int_{\mathbb{R}^2} (\operatorname{curl} \mathbf{A})_\parallel^\perp \cdot \mathbf{B}_\parallel, \quad \forall \mathbf{B} \in \mathfrak{X}_\eta,$$

we can conclude by (A.10) that

$$(\operatorname{curl} \mathbf{A})_\parallel = -\nabla^\perp \Delta^{-1}(\underline{\boldsymbol{\omega}} \cdot \mathbf{N})$$

in $(H^{-1/2}(\mathbb{R}^2))^2$. □

Proposition A.8. *Let $\boldsymbol{\omega} \in H_b(\operatorname{div}_0, D_\eta)$ and $\Phi \in \dot{H}^{3/2}(\mathbb{R}^2)$, then any weak solution \mathbf{A} of the boundary value problem (1.21) is a strong solution.*

Proof. Recall that $\operatorname{curl} \operatorname{curl} \mathbf{A} \in (L^2(D_\eta))^3$ and

$$(\operatorname{curl} \mathbf{A})_\parallel = -\nabla^\perp \Delta^{-1}(\underline{\boldsymbol{\omega}} \cdot \mathbf{N})$$

in $(H^{-1/2}(\mathbb{R}^2))^2$, hence $(\operatorname{curl} \mathbf{A})_\parallel^\perp \in (H^{1/2}(\mathbb{R}^2))^2$. Since $0 = \operatorname{div} \operatorname{curl} \mathbf{A} \in L^2(D_\eta)$ and $\operatorname{curl} \mathbf{A} \cdot \mathbf{e}_3|_{z=-h} = 0$, it follows by Lemma A.6 that $\operatorname{curl} \mathbf{A} \in (H^1(D_\eta))^3$; moreover, since $\operatorname{curl} \mathbf{A} \in (H^1(D_\eta))^3$, $0 = \operatorname{div} \mathbf{A} \in H^1(D_\eta)$ with $\mathbf{A} \times \mathbf{e}_3|_{z=-h} = \mathbf{0}$, $\underline{\mathbf{A}} \cdot \mathbf{n} = 0$, we have by Lemma A.6 that $\mathbf{A} \in (H^2(D_\eta))^3$. Finally, note that

$$\operatorname{curl} \operatorname{curl} \mathbf{A} = \boldsymbol{\omega}, \quad \text{in } D_\eta,$$

holds in $(L^2(D_\eta))^3$ because it holds in the sense of distributions, and because $\mathbf{A} \in (H^2(D_\eta))^3$. □

A more quantitative reformulation of Proposition A.7 is the following one (analogous to Lemma 3.5 in [CL15]).

Lemma A.9. *Let $k > \frac{5}{2}$, and assume that $\eta \in H^k(\mathbb{R}^2)$ satisfies condition (1.16), and let $\boldsymbol{\omega} \in H_b(\operatorname{div}_0, D_\eta)$ and $\Psi \in \dot{H}^{1/2}(\mathbb{R}^2)$. Then there exists a unique $\mathbf{A} \in \mathfrak{X}_\eta$ to (A.8). Moreover, there exists $C > 0$ be such that*

$$\|\operatorname{curl} \mathbf{A}\|_{(L^2(D_\eta))^3} \leq C(h, \|\eta\|_{H^k(\mathbb{R}^2)}) \left(\|\boldsymbol{\omega}\|_{(L^2(D_\eta))^3} + \|\nabla \Psi\|_{(H^{-1/2}(\mathbb{R}^2))^2} \right).$$

Proof. Notice that the bilinear form

$$(\mathbf{A}, \mathbf{B}) \mapsto \int_{D_\eta} \operatorname{curl} \mathbf{A} \cdot \operatorname{curl} \mathbf{B} + \operatorname{div} \mathbf{A} \operatorname{div} \mathbf{B}$$

is continuous on $\mathfrak{X}_\eta \times \mathfrak{X}_\eta$; under the assumption that $\eta \in H^k(\mathbb{R}^2)$, it is also coercive by Lemma A.3. Moreover, if we consider the linear form

$$L(\mathbf{A}) := \int_{D_\eta} \underline{\omega} \cdot \mathbf{A} + \int_{\mathbb{R}^2} \nabla \Delta^{-1}(\underline{\omega} \cdot \mathbf{N}) \cdot \mathbf{A},$$

then by using (B.3) in order to estimate the second integral we obtain that for any $0 < \varepsilon < \min(\frac{1}{2}, k - \frac{5}{2})$

$$\begin{aligned} |L(\mathbf{A})| &\leq \|\underline{\omega}\|_{(L^2(D_\eta))^3} \|\mathbf{A}\|_{(L^2(D_\eta))^3} \\ &\quad + C(\|\eta\|_{C^{3/2+\varepsilon}(\mathbb{R}^2)}) \|\nabla \Psi\|_{(H^{-1/2}(\mathbb{R}^2))^2} \|\mathbf{A}\|_{(H^1(D_\eta))^3}, \end{aligned}$$

hence it is continuous on \mathfrak{X}_η , and we can deduce the existence and uniqueness of a variational solution by Lax-Milgram's theorem.

Taking $\mathbf{B} = \mathbf{A}$ in (A.10) and using the above estimate for $L(\mathbf{A})$

$$\|\operatorname{curl} \mathbf{A}\|_{(L^2(D_\eta))^3} \leq C(\|\eta\|_{C^{3/2+\varepsilon}(\mathbb{R}^2)}) \|\nabla \Psi\|_{(H^{-1/2}(\mathbb{R}^2))^2} \|\mathbf{A}\|_{(H^1(D_\eta))^3},$$

so that the thesis follows from Lemma A.3 and by the fact that $\operatorname{div} \mathbf{A} = 0$. \square

Now consider $\mathbf{A} \in \mathfrak{X}_\eta$ as the solution given by Proposition A.7 and Proposition A.8, and set $\mathbf{v} := \operatorname{curl} \mathbf{A}$. We have that $\mathbf{v} \in H(\operatorname{div}, D_\eta) \cap H(\operatorname{curl}, D_\eta)$, and from Lemma A.9 we get

$$(A.11) \quad \|\mathbf{v}\|_{(L^2(D_\eta))^3} \leq C(h, \|\eta\|_{H^k(\mathbb{R}^2)}) [\|\underline{\omega}\|_{(L^2(D_\eta))^3} + \|\nabla \Psi\|_{(H^{-1/2}(\mathbb{R}^2))^2}].$$

Lemma A.10. *Assume that $\eta \in C^1(\mathbb{R}^2)$ satisfies condition (1.16), and let $\underline{\omega} \in H_b(\operatorname{div}_0, D_\eta)$. Then there exists a unique $\Psi \in \dot{H}^{3/2}(\mathbb{R}^2)$ which satisfies $\Delta \Psi = \underline{\omega} \cdot \mathbf{N}$. Moreover, there exists $C > 0$ such that*

$$\|\nabla \Psi\|_{H^{1/2}(\mathbb{R}^2)} \leq C(h_0^{-1}, \|\eta\|_{C^1(\mathbb{R}^2)}) \|\underline{\omega}\|_{2,b}.$$

Proof. Let us define the following linear form on $\dot{H}^1(\mathbb{R}^2)$.

$$\ell(v) := \int_{\mathbb{R}^2} (\underline{\omega} \cdot \mathbf{N})v, \quad \forall v \in \dot{H}^1(\mathbb{R}^2).$$

Provided that ℓ is continuous, then the thesis follows from Lax-Milgram's theorem. Therefore we now prove that ℓ is continuous.

In the following we use the straightening diffeomorphism (1.31). By integration by parts we have

$$\ell(v) = \int_{\mathbb{R}^2} (\underline{\omega}|_{w=-h} \cdot \mathbf{e}_3) v^{ext}|_{w=-h} + \int_{D_0} (1 + \partial_w \sigma) \nabla^\Sigma v^{ext} \cdot \tilde{\omega},$$

where for all $v \in \dot{H}^1(\mathbb{R}^2)$, v^{ext} is the solution of the boundary value problem

$$(A.12) \quad \begin{cases} \Delta v^{ext} = 0, & \text{in } D_0, \\ \partial_w v^{ext} = 0, & \text{at } w = -h, \\ v^{ext} = 0, & \text{at } w = 0. \end{cases}$$

We have that there exists $c_1, c_2, C > 0$ such that

$$\begin{aligned} \ell(v) &\leq c_1 \|\underline{\omega}|_{z=-h} \cdot \mathbf{e}_3\|_{H_0^{-1/2}(\mathbb{R}^2)} \|v^{ext}|_{w=-h}\|_{H^{1/2}(\mathbb{R}^2)} \\ &\quad + c_2 \|(1 + \partial_w \sigma) \tilde{\omega}\|_{(L^2(D_0))^3} \|\nabla^\Sigma v^{ext}\|_{(L^2(D_0))^3} \end{aligned}$$

where by Lemma B.10 we have

$$\begin{aligned} \|(1 + \partial_w \sigma) \tilde{\omega}\|_{(L^2(D_0))^3} &\leq (1 + \|\partial_w \sigma\|_{C(D_0)}) \|\tilde{\omega}\|_{(L^2(D_0))^3} \\ &\leq (1 + M_0 \|\eta\|_{C^1(D_0)}) \|\tilde{\omega}\|_{(L^2(D_0))^3}, \end{aligned}$$

so that

$$\ell(v) \leq C (h_0^{-1}, \|\eta\|_{C^1(\mathbb{R}^2)}) \|\omega\|_{2,b} \|\nabla v\|_{L^2(\mathbb{R}^2)}.$$

Hence there exists a unique variational solution $\Psi \in \dot{H}^1(\mathbb{R}^2)$ which satisfies $-\Delta \Psi = \underline{\omega} \cdot \mathbf{N}$; we also have that

$$\|\nabla \Psi\|_{L^2(\mathbb{R}^2)} \leq C (h_0^{-1}, \|\eta\|_{C^{1/2+\varepsilon}(\mathbb{R}^2)}) \|\omega\|_{2,b}.$$

Moreover, if we set $\Lambda := (1 - \Delta)^{1/2}$, we have

$$\begin{aligned} \|\nabla \Psi\|_{H^{1/2}(\mathbb{R}^2)}^2 &\leq \left| \int_{\mathbb{R}^2} (\Lambda \Psi) (\underline{\omega} \cdot \mathbf{N}) \right| \\ &\leq C (h_0^{-1}, \|\eta\|_{C^1(\mathbb{R}^2)}) \|\omega\|_{2,b} \|\Lambda \nabla \Psi\|_{(L^2(\mathbb{R}^2))^2} \\ &\leq C (h_0^{-1}, \|\eta\|_{C^1(\mathbb{R}^2)}) \|\omega\|_{2,b} \|\nabla \Psi\|_{(H^{1/2}(\mathbb{R}^2))^2}, \end{aligned}$$

where we applied part ii. of Proposition 1.10 for the second-to-last estimate, and Lemma 1.11 for the last estimate. This allows us to deduce the thesis. \square

Lemma A.11. *Let $k > \frac{5}{2}$, and assume that $\eta \in H^k(\mathbb{R}^2)$ satisfies condition (1.16), let $\omega \in H_b(\text{div}_0, D_\eta)$, and let $\Psi \in \dot{H}^{3/2}(\mathbb{R}^2)$ be the solution of the equation $-\Delta \Psi = \underline{\omega} \cdot \mathbf{N}$ given by Lemma A.10. If $\mathbf{v} \in (L^2(D_\eta))^3$ solves (A.4), then $\mathbf{v} \in (H^1(D_\eta))^3$, and there exists $C > 0$ such that*

$$\|\mathbf{v}\|_{(L^2(D_\eta))^3} + \|\nabla_{\mathbf{x}', z} \mathbf{v}\|_{(L^2(D_\eta))^{3 \times 3}} \leq C(h, h_0^{-1}, \|\eta\|_{H^k(\mathbb{R}^2)}) \|\omega\|_{2,b}.$$

Proof. First notice that by combining the first two equations of (A.4) we get $-\Delta_{\mathbf{x}', z} \mathbf{v} = \omega$, hence

$$\int_{D_\eta} (-\Delta_{\mathbf{x}', z} \mathbf{v}) \cdot \mathbf{v} = \int_{D_\eta} \text{curl } \omega \cdot \mathbf{v}.$$

Now we consider a regularizing straightening diffeomorphism in (1.31). Hence, recalling that $\mathbf{n} = \mathbf{N}/|\mathbf{N}|$, we have

$$\begin{aligned} \int_{D_\eta} |\nabla_{\mathbf{x}', z} \mathbf{v}|^2 &= \int_{D_0} \text{curl}^\Sigma \tilde{\omega} \cdot \tilde{\mathbf{v}} (1 + \sigma_w) \\ &\leq C(\|\sigma\|_{C^1(D_0)}) \left[\int_{\partial D_0} (\partial_n^\Sigma \tilde{\mathbf{v}} + \mathbf{n} \times \tilde{\omega}) \cdot \tilde{\mathbf{v}} + \int_{D_0} |\tilde{\omega}|^2 \right. \\ &\quad \left. + \int_{D_0} (\text{curl}^\Sigma \tilde{\omega} - \text{curl } \tilde{\omega}) \cdot \tilde{\mathbf{v}} \right]. \end{aligned}$$

Observe that at the bottom $\mathbf{N} = -\mathbf{e}_3$ and $(\text{curl}^\Sigma \tilde{\mathbf{v}})_3 = 0$. Therefore, we obtain

$$\partial_n^\Sigma \tilde{\mathbf{v}} + \tilde{\mathbf{n}} \times \tilde{\omega}|_{w=-h} = \mathbf{0},$$

so that we get

$$\begin{aligned} &\int_{\{w=0\}} (\partial_n^\Sigma \tilde{\mathbf{v}} + \mathbf{n} \times \tilde{\omega}) \cdot \tilde{\mathbf{v}} + \int_{D_0} |\tilde{\omega}|^2 \\ &= \int_{\mathbb{R}^2} \left(\frac{\mathbf{N}}{|\mathbf{N}|} \cdot \nabla^\Sigma \tilde{\mathbf{v}}|_{w=0} + \frac{\mathbf{N}}{|\mathbf{N}|} \times \text{curl}^\Sigma \tilde{\mathbf{v}}|_{w=0} \right) \cdot \tilde{\mathbf{v}}|_{w=0} + \int_{D_0} |\tilde{\omega}|^2. \end{aligned}$$

Observe that in the original coordinates

$$\mathbf{N} \cdot \underline{\nabla}_{\mathbf{x}', z} \mathbf{v} + \mathbf{N} \times \underline{\text{curl}} \mathbf{v} = \begin{pmatrix} \nabla \underline{\mathbf{w}} - (\nabla^\perp \eta \cdot \nabla) \underline{\mathbf{v}}^\perp \\ -\nabla \cdot \underline{\mathbf{v}} \end{pmatrix};$$

moreover, we also have

$$\tilde{\mathbf{v}}|_{w=0} = -\nabla^\perp \Psi - \tilde{\mathbf{w}}|_{w=0} \nabla \eta.$$

By using integration by parts on the terms involving derivatives of either $\tilde{\mathbf{v}}$ or $\tilde{\mathbf{w}}$, by using (A.11) in order to estimate $\|\mathbf{v}\|_{(L^2(D_\eta))^3}$ and by using the smoothing property (1.34) of the regularizing diffeomorphism we can conclude that

$$\begin{aligned} & \|\mathbf{v}\|_{(L^2(D_\eta))^3} + \|\nabla_{\mathbf{x}', z} \mathbf{v}\|_{(L^2(D_\eta))^{3 \times 3}} \\ & \leq C(h, h_0^{-1}, \|\eta\|_{H^k(\mathbb{R}^2)}) \left[\|\boldsymbol{\omega}\|_{2,b} + \|\nabla \Psi\|_{(H^{1/2}(\mathbb{R}^2))^2} \right], \end{aligned}$$

and by Lemma A.10 we deduce the thesis. \square

We conclude the proof of Proposition 1.4 by setting $\mathbf{U} = \mathbf{v} + \nabla_{\mathbf{x}', z} \varphi$, where φ solves (A.3). The uniqueness of the solution \mathbf{U} follows from Lemma A.11, which also leads to the estimate

$$\|\nabla_{\mathbf{x}', z} \varphi\|_{(H^1(D_\eta))^3} \leq C(h_0^{-1}, \|\eta\|_{H^k(\mathbb{R}^2)}) \|\nabla \Phi\|_{(H^{1/2}(\mathbb{R}^2))^2}.$$

APPENDIX B. PARADIFFERENTIAL CALCULUS

Here we remind some standard rules of paradifferential calculus (see Sec. 5 of [Mét08] and Sec. 4 of [AM09]). The paradifferential calculus was introduced by Bony in order to deal with the quantization of symbols $a(\mathbf{x}', \boldsymbol{\xi})$ of degree m with respect to $\boldsymbol{\xi}$ and limited regularity in the \mathbf{x}' -variable, to which are associated operators of order $\leq m$ denoted by T_a .

For any $u \in \mathcal{S}'(\mathbb{R}^d)$ we denote its Fourier transform by $\mathcal{F}u$. If $p : \mathcal{S}(\mathbb{R}^d) \rightarrow \mathcal{S}'(\mathbb{R}^d)$, then

$$p(\mathbf{D})u := \mathcal{F}^{-1}(p \mathcal{F}u).$$

In the following we also write $\mathbb{R}_{\neq}^d := \mathbb{R}^d \setminus \{\mathbf{0}\}$.

Definition B.1. Let $\rho \geq 0$ and $m \in \mathbb{R}$, we denote by $\Gamma_\rho^m(\mathbb{R}^d)$ the space of locally bounded functions $a(\mathbf{x}', \boldsymbol{\xi}) \in \mathbb{R}^d \times \mathbb{R}_{\neq}^d$ which are of class C^∞ with respect to $\boldsymbol{\xi}$ for $\boldsymbol{\xi} \neq \mathbf{0}$ and such that for all $\alpha \in \mathbb{N}^2$ and all $\boldsymbol{\xi} \neq \mathbf{0}$ the map $\mathbf{x}' \mapsto \partial_{\boldsymbol{\xi}}^\alpha a(\mathbf{x}', \boldsymbol{\xi})$ belongs to $C^e(\mathbb{R}^d)$ and there exists $C_\alpha > 0$ such that

$$(B.1) \quad \forall |\boldsymbol{\xi}| \geq 1/2, \quad \|\partial_{\boldsymbol{\xi}}^\alpha a(\cdot, \boldsymbol{\xi})\|_{C^e(\mathbb{R}^d)} \leq C_\alpha (1 + |\boldsymbol{\xi}|)^{m-|\alpha|}.$$

Note that in the above definition we considered symbols which are not necessarily smooth for $\boldsymbol{\xi} = \mathbf{0}$. The motivation comes from the principal symbol $\lambda^{(1)}(\mathbf{x}', \boldsymbol{\xi})$ of the classical Dirichlet–Neumann operator. One can notice that this symbol is not of class C^∞ with respect to the variable $\boldsymbol{\xi}$; moreover, if $\eta \in C^s(\mathbb{R}^d)$, then $\lambda^{(1)} \in \Gamma_{s-1}^1(\mathbb{R}^d)$.

We now introduce the space of pluri-homogeneous symbols.

Definition B.2. Let $\rho \geq 1$ and $m \in \mathbb{R}$, we denote by $\Sigma_\rho^m(\mathbb{R}^d)$ the space of symbols of the form

$$a(\mathbf{x}', \boldsymbol{\xi}) = \sum_{0 \leq j < \rho} a_{m-j}(\mathbf{x}', \boldsymbol{\xi}),$$

where $a_{m-j} \in \Gamma_{\rho-j}^{m-j}(\mathbb{R}^d)$ is homogeneous of degree $m-j$ in the variable $\boldsymbol{\xi}$ and is of class C^∞ in the variable $\boldsymbol{\xi}$ for $\boldsymbol{\xi} \neq \mathbf{0}$, and with regularity C^{e-j} in the variable \mathbf{x}' . We say that a_m is the principal symbol of a .

In order to define rigorously paradifferential operators we first introduce a fixed cutoff functions χ . Fix $\varepsilon_1, \varepsilon_2$ such that $0 < 2\varepsilon_1 < \varepsilon_2 < 1/2$ and a function $f \in C_0^\infty(\mathbb{R}^d)$ satisfying $f(t) = f(-t)$, $f(t) = 1$ for $|t| \leq 2\varepsilon_1$ and $f(t) = 0$ for $|t| \geq \varepsilon_2$; then set

$$\chi(\xi_1, \xi_2) = (1 - f(\xi_2)) f(|\xi_1|/|\xi_2|).$$

Observe that the cutoff function satisfies the following properties:

- i. there exists $\varepsilon_1, \varepsilon_2$ satisfying $0 < 2\varepsilon_1 < \varepsilon_2 < 1/2$ such that

$$\begin{aligned} \chi(\xi_1, \xi_2) &= 1, \quad \text{if } |\xi_1| \leq \varepsilon_1 |\xi_2| \text{ and } |\xi_2| \geq 2, \\ \chi(\xi_1, \xi_2) &= 0, \quad \text{if } |\xi_1| \geq \varepsilon_2 |\xi_2| \text{ or } |\xi_2| \geq 1; \end{aligned}$$

- ii. for all $(\alpha, \beta) \in \mathbb{N}^d$, there exists $C_{\alpha, \beta} > 0$ such that

$$|\partial_{\xi_1}^\alpha \partial_{\xi_2}^\beta \chi(\xi_1, \xi_2)| \leq C_{\alpha, \beta} (1 + |\xi_2|)^{-|\alpha| - |\beta|};$$

- iii. χ satisfies the following symmetry condition,

$$\chi(\xi_1, \xi_2) = \chi(-\xi_1, -\xi_2) = \chi(-\xi_1, \xi_2).$$

Given a symbol $a \in \Sigma_\varrho^m(\mathbb{R}^d)$, we define the paradifferential operator T_a by

$$(B.2) \quad \mathcal{F}(T_a u)(\xi) := (2\pi)^{-d} \int_{\mathbb{R}^d} \chi(\xi - \xi', \xi') \mathcal{F}a(\xi - \xi', \xi') \mathcal{F}u(\xi') d\xi',$$

where $\mathcal{F}a$ is the Fourier transform of a with respect to the first variable.

We notice that if $Q(D)$ is a Fourier multiplier with symbol $q(\xi)$, we do not have $Q(D) = T_q$, because of the cut-off function used in the definition (B.2); however, we have $Q(D) = T_q + R$, where $R : H^t(\mathbb{R}^d) \rightarrow H^\infty(\mathbb{R}^d)$ for all $t \in \mathbb{R}$.

Definition B.3. Let $m \in \mathbb{R}$, then an operator T is of order m if, for all $s \in \mathbb{R}$, it is bounded from $H^{s+m}(\mathbb{R}^d)$ to $H^s(\mathbb{R}^d)$.

Proposition B.4. Let $m \in \mathbb{R}$ and let $a \in \Gamma_0^m(\mathbb{R}^d)$, then T_a is of order m .

Recall the following properties

Proposition B.5. Let $m_1, m_2 \in \mathbb{R}$ and let $a \in \Gamma_1^{m_1}(\mathbb{R}^d)$, $b \in \Gamma_1^{m_2}(\mathbb{R}^d)$, then $T_a T_b - T_{ab}$ is of order $m_1 + m_2 - 1$.

Proposition B.6. Let $\varrho > 0$, $m_1, m_2 \in \mathbb{R}$ and let $a \in \Gamma_\varrho^{m_1}(\mathbb{R}^d)$, $b \in \Gamma_\varrho^{m_2}(\mathbb{R}^d)$. Set

$$a \sharp b(x', \xi) := \sum_{|\alpha| < \varrho} \frac{(-i)^\alpha}{\alpha!} \partial_\xi^\alpha a(x', \xi) \partial_{x'}^\alpha b(x', \xi) \in \sum_{j < \varrho} \Gamma_{\varrho-j}^{m_1+m_2-j}(\mathbb{R}^d),$$

then $T_a T_b - T_{a \sharp b}$ is of order $\leq m_1 + m_2 - \varrho$.

If $a = a(x')$ does not depend on the variable ξ , then the paradifferential operator T_a is called *paraproduct*. From Proposition B.4 we have that if $b \in H^\beta(\mathbb{R}^d)$ with $\beta > \frac{d}{2}$, then T_b is of order 0.

Moreover, we recall the following properties

Lemma B.7. Let $m_1 \in \mathbb{R}$, $m_2 < d/2$, $a \in H^{m_1}(\mathbb{R}^d)$, $b \in H^{m_2}(\mathbb{R}^d)$. Then $T_b a \in H^{m_1+m_2-d/2}(\mathbb{R}^d)$.

Moreover, if $a \in L^\infty(\mathbb{R}^d)$, then T_a is an operator of order ≤ 0 , and there exists $C > 0$ such that

$$\|T_a u\|_{H^s(\mathbb{R}^d)} \leq C \|a\|_{L^\infty(\mathbb{R}^d)} \|u\|_{H^s(\mathbb{R}^d)}, \quad \forall u \in H^s(\mathbb{R}^d), \quad \forall s \in \mathbb{R}.$$

Similarly, if $a \in L^\infty(\mathbb{R}^d)$, then for all $s \in (0, +\infty) \setminus \mathbb{N}$ there exists $C > 0$ such that

$$\|T_a u\|_{C^s(\mathbb{R}^d)} \leq C \|a\|_{L^\infty(\mathbb{R}^d)} \|u\|_{C^s(\mathbb{R}^d)}, \quad \forall u \in C^s(\mathbb{R}^d).$$

Lemma B.8. *Let $m > d/2$, $a \in H^m(\mathbb{R}^d)$, $F \in C^\infty(\mathbb{R})$. Then $F(a) - T_{F'(a)}a \in H^{2m-d/2}(\mathbb{R}^d)$.*

Moreover, let $m_1, m_2 \in \mathbb{R}$ be such that $m_1 + m_2 > 0$, and let $a \in H^{m_1}(\mathbb{R}^d)$, $b \in H^{m_2}(\mathbb{R}^d)$. Then

$$R(a, b) := ab - T_a b - T_b a \in H^{m_1+m_2-d/2}(\mathbb{R}^d),$$

and there exists $K > 0$ such that

$$\|R(a, b)\|_{H^{m_1+m_2-d/2}(\mathbb{R}^d)} \leq K \|a\|_{H^{m_1}(\mathbb{R}^d)} \|b\|_{H^{m_2}(\mathbb{R}^d)}.$$

Similarly, let $m_1 \in \mathbb{R}_+$, $m_2 \in \mathbb{R}$ be such that $m_1 + m_2 > 0$, and let $a \in C^{m_1}(\mathbb{R}^d)$, $b \in H^{m_2}(\mathbb{R}^d)$. Then

$$R(a, b) = ab - T_a b - T_b a \in H^{m_1+m_2}(\mathbb{R}^d),$$

and there exists $K > 0$ such that

$$\|R(a, b)\|_{H^{m_1+m_2}(\mathbb{R}^d)} \leq K \|a\|_{C^{m_1}(\mathbb{R}^d)} \|b\|_{H^{m_2}(\mathbb{R}^d)}.$$

Lemma B.9. *Let $m > \frac{d}{2}$ be such that $m - \frac{d}{2} \notin \mathbb{N}$. If $a \in H^m(\mathbb{R}^d)$, $b \in H^m(\mathbb{R}^d)$, then $T_a T_b - T_{ab}$ is of order $-(m - \frac{d}{2})$.*

Next, we mention some product estimates (see (A.120)-(A.121) of [AD15] for the case $d = 2$; see also Proposition 9.10 of [Tay11] for (B.3) and Proposition 8.6.8 of [Hör97] for (B.4)).

Lemma B.10. *For any $(r, s, s') \in \mathbb{R}_+^3$ such that $s' > s \geq r$, then there exists a constant $C > 0$ such that*

$$\|T_a u\|_{H^{s-r}(\mathbb{R}^2)} \leq C \|a\|_{H^{-r}(\mathbb{R}^2)} \|u\|_{C^{s'}(\mathbb{R}^2)}, \quad \forall a \in H^{-r}(\mathbb{R}^2), \quad \forall u \in C^{s'}(\mathbb{R}^2).$$

Moreover, let $d \geq 1$, $s \in \mathbb{R}$ and $s' \in \mathbb{R}_+$ be such that $s' > |s|$, then there exists a constant $C > 0$ such that

$$(B.3) \quad \|f g\|_{H^s(\mathbb{R}^d)} \leq C \|f\|_{H^s(\mathbb{R}^d)} \|g\|_{C^{s'}(\mathbb{R}^d)}, \quad \forall f \in H^s(\mathbb{R}^d), \quad \forall g \in C^{s'}(\mathbb{R}^d).$$

Finally, let $d \geq 1$, $s_1, s_2 \in \mathbb{R}$ be such that $s_1 + s_2 > 0$, and set $s := \min(s_1, s_2)$; then there exists a constant $C = C(s_1, s_2) > 0$ such that

$$(B.4) \quad \|f g\|_{C^s(\mathbb{R}^d)} \leq C \|f\|_{C^{s_1}(\mathbb{R}^d)} \|g\|_{C^{s_2}(\mathbb{R}^d)}, \quad \forall f \in C^{s_1}(\mathbb{R}^d), \quad \forall g \in C^{s_2}(\mathbb{R}^d).$$

Remark B.11. *In addition to (B.3), we recall the following standard product estimates (see Theorem 8.3.1 in [Hör97]).*

Let $d \geq 1$, and let $r, r_1, r_2 \in \mathbb{R}$ be such that $r_1 + r_2 \geq 0$, $r \leq \min(r_1, r_2)$, $r \leq r_1 + r_2 - d/2$ ($r < r_1 + r_2 - d/2$ if either r_1, r_2 or $-r$ is equal to $d/2$), then

$$(B.5) \quad \|f g\|_{H^r(\mathbb{R}^d)} \leq C \|f\|_{H^{r_1}(\mathbb{R}^d)} \|g\|_{H^{r_2}(\mathbb{R}^d)}, \quad \forall f \in H^{r_1}(\mathbb{R}^d), \quad \forall g \in H^{r_2}(\mathbb{R}^d).$$

Moreover, let $d \geq 1$, $r > 0$ and $f, g \in H^r(\mathbb{R}^d) \cap L^\infty(\mathbb{R}^d)$, then

$$(B.6) \quad \|f g\|_{H^r(\mathbb{R}^d)} \leq C (\|f\|_{H^r(\mathbb{R}^d)} \|g\|_{L^\infty(\mathbb{R}^d)} + \|f\|_{L^\infty(\mathbb{R}^d)} \|g\|_{H^r(\mathbb{R}^d)}).$$

We also mention a result concerning the action of pseudodifferential operators on Hölder spaces, see (1.1) and Proposition 1.1 in [Tay15].

Lemma B.12. *Let $d \geq 1$, $m, r \in \mathbb{R}$, $P \in OPS_{1,0}^m(\mathbb{R}^d)$, then $P : C_*^r(\mathbb{R}^d) \rightarrow C_*^{r-m}(\mathbb{R}^d)$.*

Next, we mention the following result (see Lemma 2.19 in [ABZ14] for a slightly more general statement; see also Proposition 2.12 in [Lan13]).

Lemma B.13. *Let $h > 0$, $s \in \mathbb{R}$, and let $\tilde{f} : D_0 \rightarrow \mathbb{R}$ be such that*

$$\tilde{f} \in L^2([-h, 0]; H^{s+1/2}(\mathbb{R}^2)), \quad \partial_w \tilde{f} \in L^2([-h, 0]; H^{s-1/2}(\mathbb{R}^2)).$$

Then $\tilde{f} \in C^0([-h, 0]; H^s(\mathbb{R}^2))$, and there exists a constant $C > 0$ such that

$$\|\tilde{f}\|_{C^0([-h, 0]; H^s(\mathbb{R}^2))} \leq C [\|\tilde{f}\|_{L^2([-h, 0]; H^{s+1/2}(\mathbb{R}^2))} + \|\partial_w \tilde{f}\|_{L^2([-h, 0]; H^{s-1/2}(\mathbb{R}^2))}].$$

Finally, we state and we prove the following elliptic regularity result (see also Proposition 4.10 of [AM09]).

Proposition B.14. *Let $\tau > 0$, $0 \leq r < 1$, $A \in M_3(\Gamma_{1+r}^1(\mathbb{R}^2))$, and $B \in M_3(\Gamma_0^0(\mathbb{R}^2))$. Assume that there exists $c > 0$ such that*

$$\forall(\mathbf{x}', \boldsymbol{\xi}) \in \mathbb{R}^2 \times \mathbb{R}^2, |\operatorname{Re} A(\mathbf{x}', \boldsymbol{\xi})| \geq c|\boldsymbol{\xi}|.$$

If $\mathbf{U} \in C^1([-\tau, 0]; (H^{-\infty}(\mathbb{R}^2))^3)$ solves

$$\partial_w \mathbf{U} + T_A \mathbf{U} = T_B \mathbf{U} + \mathbf{F},$$

where $\mathbf{F} \in C^0([-\tau, 0]; (H^s(\mathbb{R}^2))^3)$ for some $s \in \mathbb{R}$, then

$$\mathbf{U}(0) \in (H^{s+r}(\mathbb{R}^2))^3.$$

Moreover, if $\mathbf{U} \in C^1([-\tau, 0]; (H^s(\mathbb{R}^2))^3) \cap C^0([-\tau, 0]; (H^{s-1}(\mathbb{R}^2))^3)$, then

$$\mathbf{U} \in C^0([-\tau, 0]; (H^{s+r}(\mathbb{R}^2))^3),$$

and there exists $K > 0$ such that

$$\begin{aligned} \|\mathbf{U}\|_{L^\infty([-\tau, 0]; H^{s+r}(\mathbb{R}^2, \mathbb{R}^3))} &\leq K \times \\ &\times \left[\|\mathbf{U}(0)\|_{(H^{s+r}(\mathbb{R}^2))^3} + \|\mathbf{F}\|_{L^\infty([-\tau, 0]; (H^s(\mathbb{R}^2))^3)} + \|\mathbf{U}\|_{L^\infty([-\tau, 0]; (H^s(\mathbb{R}^2))^3)} \right]. \end{aligned}$$

Proof. Without loss of generality we can assume that $B = \mathbb{O}_3$ and that

$$\mathbf{U} \in C^1([-\tau, 0]; (H^{-\infty}(\mathbb{R}^2))^3).$$

Moreover, we can assume that $\mathbf{U}(t, \mathbf{x}', w) = \mathbf{0}$ for $w \leq -\tau/2$.

We set

$$\begin{aligned} \mathcal{E}(w; \mathbf{x}', \boldsymbol{\xi}) &:= \mathcal{E}_0(w; \mathbf{x}', \boldsymbol{\xi}) + \mathcal{E}_{-1}(w; \mathbf{x}', \boldsymbol{\xi}) \\ &= \exp(w A(\mathbf{x}', \boldsymbol{\xi})) - i \exp(w A(\mathbf{x}', \boldsymbol{\xi})) \frac{w^2}{2} \nabla_{\boldsymbol{\xi}} A(\mathbf{x}', \boldsymbol{\xi}) \cdot \nabla_{\mathbf{x}'} A(\mathbf{x}', \boldsymbol{\xi}), \end{aligned}$$

so that $\mathcal{E}(0; \mathbf{x}', \boldsymbol{\xi}) = \mathbb{I}_3$, and

$$\partial_w \mathcal{E} = \mathcal{E}_0 A + \mathcal{E}_{-1} A - i \nabla_{\boldsymbol{\xi}} \mathcal{E}_0 \cdot \nabla_{\mathbf{x}'} A(\mathbf{x}', \boldsymbol{\xi}).$$

By assumption we have

$$\mathcal{E}_j \in C^0([-\tau, 0]; M_3(\Gamma_{1+r-j}^j(\mathbb{R}^2))), \quad j = 0, -1.$$

By Proposition B.6 we have that $T_{\partial_w \mathcal{E}} - T_{\mathcal{E}} T_A$ is of order $-r$. Write

$$\partial_w (T_{\mathcal{E}} \mathbf{U}) = T_{\mathcal{E}} \mathbf{f} + \mathbf{F}, \quad \mathbf{F} \in C^0([-\tau, 0]; (H^{s+r}(\mathbb{R}^2))^3).$$

By arguing as in Proposition 4.10 of [AM09] and as in Lemma 2.2.7 of [AD15] we can conclude. \square

REFERENCES

- [AB15] Thomas Alazard and Pietro Baldi. Gravity capillary standing water waves. *Archive for Rational Mechanics and Analysis*, 217(3):741–830, 2015.
- [ABZ11] Thomas Alazard, Nicolas Burq, and Claude Zuily. On the water-wave equations with surface tension. *Duke Mathematical Journal*, 158(3):413–499, 2011.
- [ABZ14] Thomas Alazard, Nicolas Burq, and Claude Zuily. On the Cauchy problem for gravity water waves. *Inventiones mathematicae*, 198(1):71–163, 2014.
- [ABZ16] Thomas Alazard, Nicolas Burq, and Claude Zuily. Cauchy theory for the gravity water waves system with non-localized initial data. *Annales de l'Institut Henri Poincaré C, Analyse non linéaire*, 33(2):337–395, 2016.
- [AD15] Thomas Alazard and Jean-Marc Delort. Sobolev estimates for two dimensional gravity water waves. *Astérisque*, 374:viii–241, 2015.
- [AIT22] Albert Ai, Mihaela Ifrim, and Daniel Tataru. Two-dimensional gravity waves at low regularity II: Global solutions. *Annales de l'Institut Henri Poincaré C*, 39(4):819–884, 2022.
- [AM09] Thomas Alazard and Guy Métivier. Paralinearization of the Dirichlet to Neumann operator, and regularity of three-dimensional water waves. *Communications in Partial Differential Equations*, 34(12):1632–1704, 2009.

- [Bam21] Dario Bambusi. Hamiltonian studies on counter-propagating water waves. *Water Waves*, 3(1):49–83, 2021.
- [BBHM18] Pietro Baldi, Massimiliano Berti, Emanuele Haus, and Riccardo Montalto. Time quasi-periodic gravity water waves in finite depth. *Inventiones mathematicae*, 214(2):739–911, 2018.
- [BCD11] Hajer Bahouri, Jean-Yves Chemin, and Raphaël Danchin. *Fourier analysis and nonlinear partial differential equations*. Springer, 2011.
- [BD18] Massimiliano Berti and Jean-Marc Delort. *Almost Global Solutions of Capillary-Gravity Water Waves Equations on the Circle*. Springer, 2018.
- [Ben84] T Brooke Benjamin. Impulse, flow force and variational principles. *IMA Journal of applied Mathematics*, 32(1-3):3–68, 1984.
- [BFM21] Massimiliano Berti, Luca Franzoi, and Alberto Maspero. Traveling quasi-periodic water waves with constant vorticity. *Archive for Rational Mechanics and Analysis*, 240(1):99–202, 2021.
- [BFM24] Massimiliano Berti, Luca Franzoi, and Alberto Maspero. Pure gravity traveling quasi-periodic water waves with constant vorticity. *Communications on Pure and Applied Mathematics*, 77(2):990–1064, 2024.
- [BGW18] Boris Buffoni, Mark D Groves, and Erik Wahlén. A variational reduction and the existence of a fully localised solitary wave for the three-dimensional water-wave problem with weak surface tension. *Archive for Rational Mechanics and Analysis*, 228:773–820, 2018.
- [BGW22] Boris Buffoni, Mark D Groves, and Erik Wahlén. Fully localised three-dimensional gravity-capillary solitary waves on water of infinite depth. *Journal of Mathematical Fluid Mechanics*, 24(2):55, 2022.
- [BT03] Boris Buffoni and John Toland. *Analytic theory of global bifurcation: an introduction*, volume 9. Princeton University Press, 2003.
- [CG94] Walter Craig and Mark D Groves. Hamiltonian long-wave approximations to the water-wave problem. *Wave motion*, 19(4):367–389, 1994.
- [CL00] Demetrios Christodoulou and Hans Lindblad. On the motion of the free surface of a liquid. *Communications on Pure and Applied Mathematics: A Journal Issued by the Courant Institute of Mathematical Sciences*, 53(12):1536–1602, 2000.
- [CL15] Angel Castro and David Lannes. Well-posedness and shallow-water stability for a new Hamiltonian formulation of the water waves equations with vorticity. *Indiana University Mathematics Journal*, pages 1169–1270, 2015.
- [CN00] Walter Craig and David P Nicholls. Traveling two and three dimensional capillary gravity water waves. *SIAM Journal on Mathematical Analysis*, 32(2):323–359, 2000.
- [CN02] Walter Craig and David P Nicholls. Traveling gravity water waves in two and three dimensions. *European Journal of Mechanics-B/Fluids*, 21(6):615–641, 2002.
- [CS93] Walter Craig and Catherine Sulem. Numerical simulation of gravity waves. *Journal of computational physics*, 108(1):73–83, 1993.
- [CS04] Adrian Constantin and Walter Strauss. Exact steady periodic water waves with vorticity. *Communications on Pure and Applied Mathematics*, 57(4):481–527, 2004.
- [CS07] Daniel Coutand and Steve Shkoller. Well-posedness of the free-surface incompressible Euler equations with or without surface tension. *Journal of the American Mathematical Society*, 20(3):829–930, 2007.
- [DIPP17] Yu Deng, Alexandru D Ionescu, Benoît Pausader, and Fabio Pusateri. Global solutions of the gravity-capillary water-wave system in three dimensions. *Acta Mathematica*, 219:213–402, 2017.
- [EWZ22] Mats Ehrnström, Samuel Walsh, and Chongchun Zeng. Smooth stationary water waves with exponentially localized vorticity. *Journal of the European Mathematical Society*, 25(3):1045–1090, 2022.
- [FG24] Roberto Feola and Filippo Giuliani. *Quasi-periodic traveling waves on an infinitely deep perfect fluid under gravity*, volume 295. American Mathematical Society, 2024.
- [Ger09] Franz Gerstner. Theorie der wellen. *Annalen der Physik*, 32(8):412–445, 1809.
- [GH20] Mark D Groves and J Horn. A variational formulation for steady surface water waves on a Beltrami flow. *Proceedings of the Royal Society A*, 476(2234):20190495, 2020.
- [Gin18] Daniel Ginsberg. On the lifespan of three-dimensional gravity water waves with vorticity. *arXiv preprint arXiv:1812.01583*, 2018.
- [GMS12] Pierre Germain, Nader Masmoudi, and Jalal Shatah. Global solutions for the gravity water waves equation in dimension 3. *Annals of Mathematics*, pages 691–754, 2012.
- [GNPW24] Mark D Groves, Dag Nilsson, Stefano Pasquali, and Erik Wahlén. Analytical study of a generalised Dirichlet–Neumann operator and application to three-dimensional water waves on Beltrami flows. *Journal of Differential Equations*, 413:129–189, 2024.
- [GP24] Daniel Ginsberg and Fabio Pusateri. Long time regularity for 3d gravity waves with vorticity. *arXiv preprint arXiv:2401.10096*, 2024.
- [HHS⁺22] Susanna Haziot, Vera Hur, Walter Strauss, John Toland, Erik Wahlén, Samuel Walsh, and Miles Wheeler. Traveling water waves—the ebb and flow of two centuries. *Quarterly of applied mathematics*, 80(2):317–401, 2022.
- [Hör97] Lars Hörmander. *Lectures on nonlinear hyperbolic differential equations*, volume 26. Springer Science & Business Media, 1997.
- [IP09] Gérard Iooss and Pavel I Plotnikov. *Small divisor problem in the theory of three-dimensional water gravity waves*. American Mathematical Soc., 2009.

- [IP11] Gérard Iooss and Pavel Plotnikov. Asymmetrical three-dimensional travelling gravity waves. *Archive for rational mechanics and analysis*, 200(3):789–880, 2011.
- [IP15] Alexandru D Ionescu and Fabio Pusateri. Global solutions for the gravity water waves system in 2d. *Inventiones mathematicae*, 199:653–804, 2015.
- [IP19] Alexandru D Ionescu and Fabio Pusateri. Long-time existence for multi-dimensional periodic water waves. *Geometric and Functional Analysis*, 29(3):811–870, 2019.
- [IPT05] Gérard Iooss, Pavel I Plotnikov, and John F Toland. Standing waves on an infinitely deep perfect fluid under gravity. *Archive for rational mechanics and analysis*, 177(3):367–478, 2005.
- [IPTT23] Mihaela Ifrim, Ben Pineau, Daniel Tataru, and Mitchell A Taylor. Sharp Hadamard local well-posedness, enhanced uniqueness and pointwise continuation criterion for the incompressible free boundary Euler equations. *arXiv preprint arXiv:2309.05625*, 2023.
- [IT18] Mihaela Ifrim and Daniel Tataru. Two-dimensional gravity water waves with constant vorticity, I: Cubic lifespan. *Analysis & PDE*, 12(4):903–967, 2018.
- [Lan05] David Lannes. Well-posedness of the water-waves equations. *Journal of the American Mathematical Society*, 18(3):605–654, 2005.
- [Lan13] David Lannes. *The water waves problem: mathematical analysis and asymptotics*, volume 188. American Mathematical Soc., 2013.
- [LC25] Tullio Levi-Civita. Détermination rigoureuse des ondes permanentes d’amplitude finie. *Mathematische Annalen*, 93(1):264–314, 1925.
- [Lin05] Hans Lindblad. Well-posedness for the motion of an incompressible liquid with free surface boundary. *Annals of mathematics*, pages 109–194, 2005.
- [LSW20] Evgeniy Lokharu, DS Seth, and E Wahlén. An Existence Theory for Small-Amplitude Doubly Periodic Water Waves with Vorticity. *Archive for Rational Mechanics and Analysis*, 238(2):607–637, 2020.
- [LW19] Evgeniy Lokharu and Erik Wahlén. A variational principle for three-dimensional water waves over Beltrami flows. *Nonlinear Analysis*, 184:193–209, 2019.
- [Mét08] Guy Métivier. Paradifferential Calculus and Application to the Cauchy Problem for Non-linear Systems. *CRM Series, Edizioni della Scuola Normale Superiore*, 2008.
- [NR01] David P Nicholls and Fernando Reitich. A new approach to analyticity of Dirichlet-Neumann operators. *Proceedings of the Royal Society of Edinburgh Section A: Mathematics*, 131(6):1411–1433, 2001.
- [RS81] John Reeder and Marvin Shinbrot. Three-dimensional, nonlinear wave interaction in water of constant depth. *Nonlinear Analysis: Theory, Methods & Applications*, 5(3):303–323, 1981.
- [Shu01] Mikhail Aleksandrovich Shubin. *Pseudodifferential operators and spectral theory*, volume 57. Springer, 2001.
- [Sto80] George Gabriel Stokes. On the theory of oscillatory waves. *Transactions of the Cambridge philosophical society*, 1880.
- [SVW24] Douglas S Seth, Kristoffer Varholm, and Erik Wahlén. Symmetric doubly periodic gravity-capillary waves with small vorticity. *Advances in Mathematics*, 447:109683, 2024.
- [SZ11] Jalal Shatah and Chongchun Zeng. Local well-posedness for fluid interface problems. *Archive for rational mechanics and analysis*, 199(2):653–705, 2011.
- [Tay11] Michael Taylor. *Partial differential equations III: Nonlinear Equations*, volume 117. Springer Science & Business Media, 2011.
- [Tay15] Michael Taylor. Commutator estimates for Hölder continuous and bmo-Sobolev multipliers. *Proceedings of the American Mathematical Society*, 143(12):5265–5274, 2015.
- [Wah06] Erik Wahlén. Steady periodic capillary-gravity waves with vorticity. *SIAM journal on mathematical analysis*, 38(3):921–943, 2006.
- [Wah07] Erik Wahlén. A Hamiltonian formulation of water waves with constant vorticity. *Letters in Mathematical Physics*, 79(3):303–315, 2007.
- [Wu09] Sijue Wu. Almost global wellposedness of the 2-D full water wave problem. *Inventiones mathematicae*, 177(1):45–135, 2009.
- [Wu11] Sijue Wu. Global wellposedness of the 3-D full water wave problem. *Inventiones mathematicae*, 184(1):125–220, 2011.
- [WW23] Erik Wahlén and Jörg Weber. Global bifurcation of capillary-gravity water waves with overhanging profiles and arbitrary vorticity. *International Mathematics Research Notices*, 2023(20):17377–17410, 2023.
- [WZZZ21] Chao Wang, Zhifei Zhang, Weiren Zhao, and Yunrui Zheng. Local well-posedness and breakdown criterion of the incompressible Euler equations with free boundary. *Memoirs of the American Mathematical Society*, 270(1318), 2021.

(*) INTERNATIONAL SCHOOL FOR ADVANCED STUDIED (SISSA), VIA BONOMEA 265, 34136, TRIESTE, ITALY
 Email address, *: stefano.pasquali@sissa.it