3-MANIFOLDS WITH POSITIVE SCALAR CURVATURE AND BOUNDED GEOMETRY

OTIS CHODOSH, YI LAI, AND KAI XU

ABSTRACT. We show that a complete contractible 3-manifold with positive scalar curvature and bounded geometry must be \mathbb{R}^3 . We also show that an open handlebody of genus larger than 1 does not admit complete metrics with positive scalar curvature and bounded geometry. Our results rely on the maximal weak solution to inverse mean curvature flow due to the third-named author.

1. INTRODUCTION

We are concerned here with the classification problem for 3-manifolds M admitting complete Riemannian metrics with positive scalar curvature; see Question 27 in Yau's list [41]. A fundamental observation of Schoen–Yau relates scalar curvature to the stability of minimal surfaces [32], which ultimately leads to several topological obstructions to the existence of complete nonnegative scalar curvature metrics on a 3-manifold M. First, if $\pi_1(M)$ contains a surface subgroup, then M is flat by Schoen-Yau [33, Theorem 4] and Gromov–Lawson [19, Theorem 8.4]. Second, J. Wang [37, 38] proved that if M is contractible and admits an exhaustion by solid tori, then $M \cong \mathbb{R}^3$ (cf. Lemma 2.4). In particular, the Whitehead manifold does not admit such a metric. See also relevant results [6, 8, 9, 14, 26, 42]. In spite of these results, the general classification for noncompact M admitting such a metric is widely open. In particular, we note the following special cases:

- If M is a contractible 3-manifold and admits a complete metric of nonnegative scalar curvature, do we have $M \cong \mathbb{R}^3$? (Asked by J. Wang [37, 38], cf. [7].)
- If M_{γ} is an open handlebody of genus γ , and admits a complete metric of nonnegative scalar curvature, do we have $\gamma \leq 1$? (Asked by Gromov [18, §3.10.2].)

In this paper, we resolve these two questions under the *additional* assumption that the metric has bounded geometry:

$$|\operatorname{Rm}| \le \Lambda, \quad \operatorname{inj} \ge \Lambda^{-1},$$
 (BG)

by combining minimal surface obstructions with new topological constraints obtained using inverse mean curvature flow. In particular, our main results are as follows. Hereafter, we use R to denote the scalar curvature.

Theorem 1.1. Let (M,g) be a complete, connected, contractible Riemannian 3-manifold satisfying $R \ge 0$ and (BG). Then M is diffeomorphic to \mathbb{R}^3 .

Theorem 1.2. Let M_{γ} denote the interior of the handlebody of genus γ . If (M_{γ}, g) is a complete Riemannian 3-manifold satisfying $R \geq 0$ and (BG), then $\gamma \leq 1$.

Note that \mathbb{R}^3 and $\mathbb{R}^2 \times \mathbb{S}^1$ (corresponding to $\gamma = 0, 1$ in Theorem 1.2) both admit complete metrics with $R \ge 0$ and bounded geometry. Concrete examples are a capped-off half-cylinder (which actually has $R \ge 1$) and the product of Cigar soliton and \mathbb{S}^1 , respectively. For the stronger uniformly positive scalar curvature condition $R \geq 1$, J. Wang has obtained a complete classification [36]: these 3-manifolds are infinite connect sums of spherical space forms and $S^2 \times S^1$. In particular, the only contractible manifold or handlebody admitting such a metric is \mathbb{R}^3 . We note that earlier work of Bessières–Besson–Maillot [2] used Ricci flow to prove such a classification with an additional bounded geometry assumption.

A crucial tool in J. Wang's classification [36] is the μ -bubbles introduced by Gromov [18, §3.7.2]. From this one obtains that a 3-manifold M with $R \ge 1$ admits an exhaustion by regions whose boundary consists of \mathbb{S}^2 's. This places very strong topological constraints on M (see also [15]). When the scalar curvature is only non-negative, it's not known how to use the μ -bubble method to obtain such an exhaustion (in particular, the genus one handlebody shows that one must allow for torus boundaries in this exhaustion). As we will discuss below, one of our main contributions here is to find an exhaustion with only sphere or tori boundary in certain 3-manifolds with non-negative scalar curvature.

1.1. Topological obstructions via inverse mean curvature flow. The key novelty introduced in this paper is the use of inverse mean curvature flow as a replacement for μ -bubbles in topological applications. A family of hypersurfaces is a smooth inverse mean curvature flow (IMCF) if it evolves in the outwards pointing direction with speed $\frac{1}{H}$, where H denotes the mean curvature.

The relevance of this flow to scalar curvature is the following: if (M, g) is a Riemannian 3-manifold with nonnegative scalar curvature, and $\Sigma_t \subset M$ is a compact family evolving by the smooth IMCF, then $|\Sigma_t| = e^t |\Sigma_0|$ and

$$\frac{d}{dt} \int_{\Sigma_t} H^2 \le -\frac{1}{2} \int_{\Sigma_t} H^2 + 4\pi \chi(\Sigma_t).$$
(1.1)

This is known as the Geroch monotonicity formula [17]. In particular, if the flow exists for all time $t \in [0, \infty)$ then Σ_t cannot have genus ≥ 2 for all large t, since otherwise (1.1) would force $\int_{\Sigma_t} H^2$ to be negative for $t \gg 1$, which is impossible. In particular, this implies that M admits an exhaustion by regions with sphere or torus boundaries, strongly constraining its topology.

However, there are major issues with the assumption of long-time existence in practice. First, singularities are likely to develop along the flow. Secondly, it's possible that the flow rushes to infinity in finite time if the infinity is not "large" enough.

To allow for singularities, we can use the notion of weak IMCF introduced by Huisken– Ilmanen [20] en route to their proof of the Riemannian Penrose inequality. Intuitively, this solution can be described as running the smooth flow except at each time replacing Σ_t by its least area enclosure. As proven by Huisken–Ilmanen, the Geroch monotonicity (1.1) remains true for weak solutions as long as they exist.

A weak IMCF that does not rush to infinity in finite time is called proper. In [20] and [3, 4, 5, 35, 10, 13, 23, 12, 11, 1, 21], proper IMCF are used as central tools in several scalar curvature problems. We also refer to [24, 28, 40] about proper IMCF.

In our current setting, we inevitably encounter non-proper weak IMCFs (i.e. weak IMCFs that rush to infinity within finite time). We make essential use of the third-named author's recent work [39], which shows that (M, g) always admits a "maximal" (or "innermost", "slowest") weak IMCF. Assuming bounded geometry and one-endedness of M, we show that the maximal weak solution satisfies exactly one of the following three possibilities, see Lemma 3.12:

- (i) Proper: The solution exists and remains bounded for all time.
- (ii) Sweeping: The solution entirely moves to infinity at some time $T \in (0, \infty)$.
- (iii) Escaping: The solution exists until a time $T \in (0, \infty)$, then "jumps" to infinity.

In the proper case (i), we can obtain a topological obstruction using the monotonicity formula (1.1) as above. Now we consider the remaining cases (ii) (iii); see Figure 1 for examples of each of these cases.



FIGURE 1. An IMCF sweeping out the manifold at t = T (left) and one that escapes at t = T (right).

First, we consider the case of sweeping flow. We can show that for a sequence of times $t_i \nearrow T$, the surfaces Σ_{t_i} have uniformly bounded diameters, are uniformly $C^{1,\alpha}$ -smooth, and are "almost" area-minimizing. Taking a subsequential limit, we obtain an area-minimizing hypersurface in some limit of M at infinity. By the scalar curvature lower bound, this limiting hypersurface must be \mathbb{S}^2 or \mathbb{T}^2 , which in turn implies that all but finitely many Σ_{t_i} are \mathbb{S}^2 or \mathbb{T}^2 . This again puts strong constraints on the topology of M.

Finally, we consider the case of escaping flow. In order to find a nice exhausting sequence and perform a limiting argument, we make a small perturbation of the metric so that it becomes "larger at infinity". This will delay the escape time of the maximal IMCF, thus some new level set will form in the edited region. Letting the edited region diverge to infinity and making the perturbation smaller and smaller, we obtain another diverging sequence of hypersurfaces which are "almost" area-minimizing as well. Then the limiting argument in case (ii) is employed to prove the main theorems. We refer to Section 4 for more details.



FIGURE 2. Iterated trefoils. The figure shows the embedding $\Omega_1 \subset \Omega_2 \subset \Omega_3$.

1.2. Manifolds that are nested by solid tori. Our work suggests that it might be possible to give a complete classification of 3-manifolds admitting a metric of non-negative scalar curvature with bounded geometry. The techniques developed here suggest that a key step in this classification (which we have not been able to resolve) would be the classification

of 3-manifolds that admit such metrics and also admit an exhaustion $\Omega_1 \subset \Omega_2 \subset \ldots$, $M = \bigcup_{i=1}^{\infty} \Omega_i$, with each Ω_i a solid torus. As a concrete example, suppose that Ω_i embeds in Ω_{i+1} as a trefoil knot (see Figure 2); we do not know if M admits a complete metric of non-negative scalar curvature and bounded geometry.

1.3. Acknowledgements. O.C. was supported by a Terman Fellowship and an NSF grant (DMS-2304432). Y.L. is supported by NSF grant DMS-2203310. The authors wish to thank Richard Bamler, Hubert Bray, Marcus Khuri, Chao Li, John Lott for their interest. Finally, we thank the hospitality of Simons Laufer Mathematical Institute, where a major part of this work was completed during the authors' visits.

2. TOPOLOGICAL PRELIMINARIES

In the following, we assume that all manifolds are noncompact without boundary.

Lemma 2.1. Assume M is contractible and dim $M \geq 2$. Then M has only one end.

Proof. If M has more than one end, then we can find a smooth domain $K \subseteq M$ such that $M \setminus K$ has more than one unbounded components. Let E be an unbounded component of $M \setminus K$, then $M \setminus \overline{E}$ is also unbounded, hence $[\partial E] \neq 0 \in H_{n-1}(M; \mathbb{Z})$. This is impossible since M is contractible.

Lemma 2.2. Assume M is a contractible 3-manifold. Then M is irreducible.

Proof. A smoothly embedded 2-sphere in M must bound a precompact region Ω . By Van Kampen's theorem we have $\pi_1(M) = \pi_1(\Omega) * \pi_1(M \setminus \Omega) = 1$, and thus $\pi_1(\Omega) = 1$. So the Poincaré conjecture [29, 30] implies that Ω is a ball.

In Lemma 2.3, 2.4 below, we suppose that M is a contractible 3-manifold, and that $\Omega_1 \Subset \Omega_2 \Subset \cdots \Subset M$ with $\bigcup_i \Omega_i = M$ is an exhaustion of M, with each $\partial \Omega_i$ being smooth and connected.

Lemma 2.3. If $\partial \Omega_i \cong \mathbb{S}^2$ for all $i = 1, 2, \ldots$, then $M \cong \mathbb{R}^3$.

Proof. As in Lemma 2.2, each Ω_i is a 3-ball. Thus M is an increasing union of smoothly embedded 3-balls. Thus, M is diffeomorphic to \mathbb{R}^3 . This final step follows from [22, Theorem 1] but since the present situation is much simpler, we sketch the proof here. Capping off Ω_{i+1} with a 3-ball we can use Alexander's theorem to see that $\overline{\Omega_{i+1}} \setminus \overline{\Omega_i} \cong S^2 \times [0, 1]$. We can assemble these diffeomorphisms together to show that $M \setminus \Omega_1 \cong S^2 \times [1, \infty)$. This completes the proof.

The next result essentially follows from the work of J. Wang [37] (see also [38]), but we need a slightly stronger result than the one stated there.

Lemma 2.4. If $\partial \Omega_i \cong \mathbb{T}^2$ for all i = 1, 2, ..., and M admits a complete metric of positive scalar curvature, then $M \cong \mathbb{R}^3$.

Proof. Since $\pi_1(M) = 1$, the map $\pi_1(\partial \Omega_i) \to \pi_1(M)$ is not injective, and thus the loop theorem implies that $\partial \Omega_i$ is compressible.

We claim we can find a compressing disk D of $\partial\Omega_i$ such that $D \subset \Omega_i$ or $D \subset M \setminus \Omega_i$. Let D be a compressing disk. We may assume D is transverse to $\partial\Omega_i$, thus $D \cap \partial\Omega_i$ consists of finitely many disjoint circles. Let γ be an innermost circle in the intersection. We may assume that γ is homotopically nontrivial in $\partial\Omega_i$: otherwise, it bounds a disk $D_1 \subset \partial\Omega_i$

and also a disk $D_2 \subset \Omega_i$ or $D_2 \subset M \setminus \Omega_i$, thus we can replace D_2 by D_1 and push it slightly outwards or inwards to remove γ from the intersection. Through this reduction process, we obtain a homotopically non-trivial circle $\gamma \subset \partial \Omega_i$, with a compressing disk $D' \subset \Omega_i$ or $D' \subset M \setminus \Omega_i$. The disk D' proves our claim.

If $D \subset M \setminus \Omega_i$ for some *i*, then by doing surgery along *D*, we can find another $\Omega'_i \supset \Omega_i$ with $\partial \Omega'_i \cong \mathbb{S}^2$. If this occurs for infinitely many *i*, we can then pass to a subsequence so that Lemma 2.3 applies. As such, it suffices to assume that $D \subset \Omega_i$ for all *i*. Then the surgery of $\partial \Omega_i$ along *D* is a sphere and thus (by Lemma 2.2) bounds a ball *B* in *M*. This implies that Ω_i is a solid torus.

Suppose that for infinitely many i, the torus $\partial \Omega_{i+1}$ is compressible in $\Omega_{i+1} \setminus \Omega_i$. Then, by surgery along the compressing disk, we can again find an exhaustion by regions with spherical boundary, and thus Lemma 2.3 applies. As such, we can assume that $\partial \Omega_{i+1}$ is not compressible in $\Omega_{i+1} \setminus \Omega_i$ for all i. This implies that M is a contractible genus-one manifold in the sense of [37, Definition 2.14]. Such an M cannot admit a complete metric of positive scalar curvature by [37, Theorem 1.2]. This completes the proof.

The following two lemmas concern the connectedness of boundaries.

Lemma 2.5. Suppose M is one-ended, and for all $K \Subset M$ the map $H_1(M \setminus K, \mathbb{Z}) \to H_1(M, \mathbb{Z})$ is surjective. If $E \Subset M$ is a connected C^1 domain such that $M \setminus E$ does not have bounded connected components, then ∂E is connected.

Proof. Since M is one-ended, $M \setminus E$ must be connected. If ∂E is disconnected, then there is a component of ∂E that has nonzero algebraic intersection with a closed loop γ . However, by the lemma's condition, γ is homologous to a loop that is disjoint from ∂E . This leads to a contradiction.

Lemma 2.6. Suppose M is one-ended with $b_1(M) < \infty$. Then there is a compact set S with the following property: if $E \subseteq M$ is a connected C^1 domain such that $M \setminus E$ does not have bounded connected components, then

- (i) either ∂E is connected,
- (ii) or every connected component of ∂E intersects S.

Proof. Let $\{\gamma_i\}$ be a finite collection of loops that generates $H_1(M,\mathbb{Z})$ modulo torsion. Choose any compact set $S \supseteq \bigcup \gamma_i$. We claim that S satisfies the lemma's condition. Let E be as stated in the lemma. Since M is one-ended, $M \setminus E$ must be connected and noncompact. Also, recall that E is connected. So if ∂E is disconnected, each connected component of ∂E must be non-separating. Let Σ be any component of ∂E ; thus Σ has nonzero algebraic intersection with some loop σ . Since σ is homologous to a linear combination of γ_i modulo torsion, and $\gamma_i \subset S$ have zero intersection numbers with Σ , it follows that the σ has zero intersection numbers with Σ , which is a contradiction.

Lemma 2.7. Let M^3 be an open handlebody. Let $\Omega_1 \Subset \Omega_2 \Subset \cdots \Subset M$ be a C^1 exhaustion of M, such that each Ω_i and $M \setminus \Omega_i$ is connected. Then genus $(\partial \Omega_i) \ge \text{genus}(M)$ for all large i.

Proof. We choose $U \cong \Sigma \times [0, 1]$ to be a collar neighborhood of $\partial M \cong \Sigma \times \{1\}$. For *i* sufficiently large, it holds $\Sigma \times \{0\} \subset \Omega_i \cap U$. For all large *i* we have

- (1) $\partial \Omega_i$ is connected (by Lemma 2.5),
- (2) $\partial \Omega_i$ separates $\Sigma \times \{0\}$ and $\Sigma \times \{1\}$.

Hence, the natural projection map $\partial \Omega_i \to \partial M$ has nonzero degree, and by Kneser's theorem (cf. [31]), we have genus($\partial \Omega_i$) \geq genus(∂M).

Lemma 2.8. Assume M has only one end, $E \subset M$ is connected, with $E \neq M$, ∂E compact, and $M \setminus E$ unbounded. Then $E \subseteq M$.

Proof. Since M is one-ended, there is a connected compact set $K \supseteq \partial E$ such that $M \setminus K$ is connected and unbounded. Since E is connected and $\partial E \cap (M \setminus K) = \emptyset$, we have either $E \cap (M \setminus K) = \emptyset$ or $E \supset M \setminus K$. The first case implies $E \subset K$ hence proves the lemma. The second case implies $M \setminus E \subset K$, which is excluded by the hypotheses of the lemma. \Box

3. IMCF Preliminaries

We provide a brief introduction to the weak IMCF. Here we only state results needed in the subsequent sections; we refer the reader to [20, 39] for much detailed introduction to this subject. We fix the following notations:

- (1) M will always denote a complete, connected, noncompact Riemannian manifold, with dimension $n \leq 7$ (so that minimal surface singularities do not appear).
- (2) For a function u, we denote $E_t(u) = \{u < t\}$ and $E_t^+(u) = \{u \le t\}$. When there is no ambiguity, we will write E_t, E_t^+ for brevity.
- (3) When u is defined in some domain Ω , we view E_t as a subset of Ω , hence a subset of M. The perimeter $|\partial E_t|$ is viewed as the perimeter in M, so it contains $|\partial E_t \cap \Omega|$ as well as $|\partial E_t \cap \partial \Omega|$.
- (4) We will use $C(\dots)$ to denote generic constants which depend on the items in the parentheses. The constant may change from line to line.

A smooth IMCF is defined as a smooth function u with non-vanishing gradient, satisfying the equation

$$\operatorname{div}\left(\frac{\nabla u}{|\nabla u|}\right) = |\nabla u|. \tag{3.1}$$

Note that this is a level set flow: setting $\Sigma_t = \{u = t\}$, one finds that $\{\Sigma_t\}$ is a family of hypersurfaces evolving at the speed of $1/H_t$, where H_t is the mean curvature of Σ_t . To see this, it can be calculated that $H_t = \text{div}\left(\frac{\nabla u}{|\nabla u|}\right)$ while the speed of evolution equals $\frac{1}{|\nabla u|}$.

The weak IMCF is defined as a variational form of (3.1).

Definition 3.1 (weak IMCF and its energy functional).

Given a domain $\Omega \subset M$. For a function $u \in \text{Lip}_{\text{loc}}(\Omega)$, a set E with locally finite perimeter, and a domain $K \subseteq \Omega$, we define the energy functional as

$$J_u^K(E) = |\partial^* E \cap K| - \int_{E \cap K} |\nabla u|$$

where $\partial^* E$ is the reduced boundary of E (see [27]).

We say that u is a weak solution of IMCF in Ω , if $u \in \text{Lip}_{\text{loc}}(\Omega)$, and each level set $E_t = \{u < t\}$ locally minimizes J_u in the following sense: for any $K \Subset \Omega$ and any domain F with $F\Delta E_t \Subset K$, it holds $J_u^K(E_t) \leq J_u^K(F)$.

Note that that a smooth IMCF is also a weak IMCF [20, Lemma 2.3]. Let us include a brief proof here: suppose u solves (3.1) in Ω , $t \in \mathbb{R}^2$, $K \subseteq \Omega$ and F is a competitor as in

Definition 3.1. Then

$$J_{u}^{K}(F) - J_{u}^{K}(E_{t}) = \left|\partial^{*}F \cap K\right| - \left|\partial^{*}E_{t} \cap K\right| - \int (\chi_{F} - \chi_{E_{t}})|\nabla u|$$

$$\geq \int_{\partial^{*}F \cap K} \nu_{F} \cdot \frac{\nabla u}{|\nabla u|} - \int_{\partial^{*}E_{t} \cap K} \nu_{E_{t}} \cdot \frac{\nabla u}{|\nabla u|} - \int (\chi_{F} - \chi_{E_{t}})|\nabla u|, \quad (3.2)$$

where we used $\nu_F \cdot \frac{\nabla u}{|\nabla u|} \leq 1$ and $\nu_{E_t} \cdot \frac{\nabla u}{|\nabla u|} = 1$. Then applying the divergence theorem and using the IMCF equation (3.1), the right side of (3.2) equals zero.

Next, we define the initial value problem for the weak IMCF.

Definition 3.2 (initial value problem). Given a domain $\Omega \subset M$ and a $C^{1,1}$ domain $E_0 \Subset \Omega$. We say that u is a weak solution of IMCF in Ω with initial value E_0 , if

(i) $u \in \operatorname{Lip}_{\operatorname{loc}}(\Omega)$ and $E_0 = \{u < 0\};$

(ii) $u|_{\Omega\setminus\overline{E_0}}$ is a weak solution of IMCF in $\Omega\setminus\overline{E_0}$.

Note that the specific value of u inside E_0 is not important, and a weak IMCF with initial value E_0 is not necessarily a weak IMCF in Ω .

Lemma 3.3 (immediate properties). For Ω , E_0 , u as in Definition 3.2, we have:

(i) each E_t (t > 0) is a $C^{1,\alpha}$ hypersurface;

(ii) each E_t (t > 0) is locally outward perimeter-minimizing¹ in Ω (in particular, $\Omega \setminus E$ has no precompact connected components);

(iii) $|\partial E_t| = e^{t-s} |\partial E_s| \le e^t |\partial E_0|$ for all 0 < s < t, as long as $E_t \subseteq \Omega$. If further E_0 is locally outward minimizing in Ω , then $|\partial E_t| = e^t |\partial E_0|$.

Proof. (i) follows from the standard regularity results in geometric measure theory, see [39, p.13]. For (ii) see [39, Lemma 2.6]. (iii) follows from [20, Lemma 1.4, 1.6]. \Box

The following maximum principle is useful in later arguments.

Lemma 3.4 (semilocal uniqueness, [20, Theorem 2.2]).

Let $E_0 \subseteq \Omega$, and u, v be weak solutions of IMCF in Ω with initial value E_0 . If for some t > 0 we have $E_t(u) \subseteq \Omega$ and $E_t(v) \subseteq \Omega$, then $E_t(u) = E_t(v)$, and u = v in $E_t(u) \setminus E_0$.

The following types of solutions will occur in the main proof:

Definition 3.5 (solution types).

Let u be a weak IMCF in M with an initial value E_0 , as in Definition 3.2. We say that: (i) u is proper, if $\lim_{x\to\infty} u(x) = +\infty$, or equivalently, if $E_t \in M$ for all $t \in [0, \infty)$;

(i) u is proper, if $\operatorname{Im}_{x\to\infty} u(x) = +\infty$, of equivalently, if $L_t \in \mathcal{M}$ for all $t \in [0,\infty)$, (ii) u is sweeping, if $T := \sup(u) \in (0,\infty)$, and $E_t \in \mathcal{M}$ for all t < T, and $E_T = \mathcal{M}$.

(ii) u is succepting, if $T := \sup(u) \in (0, \infty)$, and $E_T \subseteq M$ for all $v \in T$, and $E_T = M$. (iii) u is instantly escaping, if $T := \sup(u) \in (0, \infty)$, and $E_T \subseteq M$, and $u \equiv T$ in $M \setminus E_T$. In case (iii), we call T the escape time of u.

See Figure 1 for examples of sweeping and escaping flows. Also, see Figure 3 below for a complicated case of sweeping maximal IMCF, where there are infinitely many bubble-shapes hence infinitely many jumps in the flow.

Regarding proper IMCF, we have the well-known Geroch monotonicity formula:

¹We say that a set E is locally outward perimeter-minimizing in a domain Ω , if for all $K \Subset \Omega$ and $F \supset E$ with $F \setminus E \Subset K$, it holds $|\partial^* E \cap K| \le |\partial^* F \cap K|$.



FIGURE 3. A complicated case of sweeping IMCF (grey regions represent jumps)

Lemma 3.6 (Monotonicity).

Let M be 3-dimensional and has nonnegative scalar curvature. Let u be a proper weak IMCF in M with a $C^{1,1}$ initial value $E_0 \Subset M$. Then it holds (in the Gronwall sense)

$$\frac{d}{dt} \int_{\partial E_t} H^2 \le 4\pi \chi(\partial E_t) - \frac{1}{2} \int_{\partial E_t} H^2.$$

Proof. This follows from [20, Formula (5.22)]; see also [20, p. 395–396] for computations in the smooth case. \Box

In the case that no proper solutions exist, we will make use of maximal weak solutions. A weak solution u satisfying Definition 3.2 is called *maximal*, if $u \ge v$ for all weak solutions v having the same initial value as u. We refer to [39] for a systematic study of this object. In the following lemma, we summarize its properties that will be used later. Note that a proper IMCF is also a maximal IMCF, hence the results below are applicable to proper IMCF as well.

Lemma 3.7 (properties and approximation of maximal solutions).

(i) For any C^{1,1} initial value E₀ ∈ M, there exists a unique maximal solution u of IMCF in M with initial value E₀. Such solution has the following properties:
(a) if M satisfies (BG), then we have the gradient estimate

$$|\nabla u| \leq C(\Lambda, H^+_{\partial E_0})$$
 outside E_0 ,

where $H^+_{\partial E_0}$ is the maximum of the mean curvature of ∂E_0 ;

- (b) if E_0 is connected, then E_t is connected for each t > 0;
- (c) for each t > 0, $M \setminus E_t$ does not have bounded connected components;
- (d) $|\partial E_t| \leq e^t |\partial E_0|$ for all t > 0.
- (ii) For any sequence of smooth domains $E_0 \in \Omega_1 \in \Omega_2 \in \cdots \in M$ with $\bigcup_{l=1}^{\infty} \Omega_l = M$, there are functions $u_l \in \operatorname{Lip}_{\operatorname{loc}}(\Omega_l)$ such that:
 - (a) each u_l is the maximal weak solution of IMCF on Ω_l with initial value E_0 ;
 - (b) each $\partial E_t(u_l)$ is a $C^{1,\alpha}$ surface in M;
 - (c) $u_1 \ge u_2 \ge u_3 \ge \cdots$ outside E_0 , and $\lim_{l\to\infty} u_l = u$ in $C^0_{\text{loc}}(M \setminus E_0)$, where u is the solution given in (i);
 - (d) for each l and $t > s \ge 0$, we have $|\partial E_t(u_l)| \le e^{t-s} |\partial E_s(u_l)|$.

(e) if $E_t(u) \Subset \Omega_l$ for some t > 0 and $l \ge 1$, then $E_t(u_l) = E_t(u)$, and we have $u_l = u$ in $E_t(u_l)$.

Proof. For item (ii), we set each u_l to be the weak IMCF with initial value E_0 and outer obstacle $\partial \Omega_l$, given by [39, Theorem 6.1]. Items (iia, iib) follow from items (iii, iv) of [39, Theorem 6.1]. Item (iic) and the existence claim in item (i) follows from [39, Theorem 7.1] and its proof. Item (iid) follows from [39, Corollary 3.14]. Item (iie) is proved as follows: by item (iic) we have $E_t(u_l) \subset E_t(u) \Subset \Omega_l$, then the conclusion is implied by Lemma 3.4 (interior maximum principle). Items (ia, ib, id) follow directly from [39, Theorem 7.2]. Item (ic) follows from Lemma 3.3(ii).

Lemma 3.8 (regularity and density bounds).

Let M satisfy (BG), and u be a weak solution of IMCF in some domain $\Omega \subset M$. Moreover, assume the gradient bound $|\nabla u| \leq L$ on Ω . Then for all $t \in \mathbb{R}$ the following hold: (i) for all $x \in \partial E_t(u)$ and $r \leq \Lambda^{-1}$ with $B(x, 2r) \subset \Omega$, we have the density bound

$$\left|\partial E_t \cap B(x,r)\right| \ge C(L,\Lambda)r^{n-1}$$

(ii) if M further satisfies $|\nabla^k \operatorname{Rm}| \leq \Lambda', \forall 1 \leq k \leq 5$, then $\partial E_t(u)$ is a $C^{1,\alpha}$ surface, where the $C^{1,\alpha}$ norm² near $x \in \partial E_t(u)$ depends only on $\alpha, L, \Lambda, \Lambda'$, and $d(x, \partial \Omega)$.

Proof. Note that the gradient bound and Definition 3.1 implies

$$|\partial E_t \cap K| \le |\partial F \cap K| + L|E_t \Delta F|$$
, for all F with $E_t \Delta F \Subset K \Subset \Omega$.

Then item (i) follows from the classical area monotonicity formula, while (ii) follows from the standard regularity theory, see [27, Chapter 21]. \Box

Corollary 3.9 (diameter bounds for level sets).

Let $E_0 \subseteq M$ be a $C^{1,1}$ domain, and u be the maximal solution of IMCF in M with initial value E_0 , as given by Lemma 3.7(i). Denote $\mathcal{C}(\partial E_0) := \max \{ \operatorname{diam}(\partial E_0), |\partial E_0|, H_{\partial E_0}^+ \}$.

(i) Suppose M satisfies (BG), and $0 < t \leq T$ is such that $E_t \neq M$. Then ∂E_t is compact, and each connected component of ∂E_t has diameter bounded by $C(T, \Lambda, C(\partial E_0))$.

(ii) If further M has only one end, then $E_t \subseteq M$ for t as in (i).

Proof. (i) First, Lemma 3.7(ia) provides a uniform gradient estimate in $M \setminus E_0$. Let Σ be a connected component of ∂E_t . Note that one can find at least $\lfloor \operatorname{diam}(\Sigma)/\Lambda \rfloor$ disjoint balls of the form $B(x_i, 1/2\Lambda)$, with $x_i \in \Sigma$. If Σ is noncompact, then we can find infinitely many such disjoint balls. Now note that:

- (1) If $d(x_i, E_0) \ge 1/\Lambda$, then by Lemma 3.8(i) we have $|\Sigma \cap B(x_i, 1/2\Lambda)| \ge C(\Lambda, H^+_{\partial E_0})$.
- (2) By volume comparison, among all the x_i , there are at most $C(\Lambda, \operatorname{diam}(\partial E_0))$ many points such that $d(x_i, E_0) < 1/\Lambda$.

Furthermore, by Lemma 3.7(id) we have $|\Sigma| \leq |\partial E_t| \leq e^t |\partial E_0|$. The diameter bound then follows from area comparison. Since we have proved that each component of ∂E_t has a uniform area lower bound, the compactness of ∂E_t again follows from area comparison.

(ii) If M has only one end, then Lemma 3.7(ib)(ic) and the fact $E_t \neq M$ implies $E_t \Subset M$, due to Lemma 2.8.

²We say that the $C^{1,\alpha}$ norm of a hypersurface Σ is bounded by C near x, if there exists a geodesic normal coordinates in $B(x, C^{-1})$ in which Σ is the graph of a function f with $||f||_{C^{1,\alpha}} \leq C$.

Finally, the following lemma shows that for E_0 a sufficiently small geodesic ball, there is a weak solution that remains proper for some definite amount of time. See the last picture in Figure 4 for a weak solution that jumps to infinity at t = 0.

Lemma 3.10 (no instant escape).

Assume M satisfies (BG), and fix $p \in M$. There is a sufficiently small $r_0 > 0$ such that: setting $E_0 = B(p, r_0)$, the maximal solution u given by Lemma 3.7(i) satisfies $E_t(u) \Subset M$ for some t > 0.

Proof. Note that (BG) implies a uniform lower bound on the volume of balls: there exists V > 0, such that

$$|B(x,1)| \ge V, \qquad \forall x \in M.$$

This further implies $|M| = \infty$ since M is connected and noncompact. Let us show that there exists constant $C(\Lambda) > 0$ such that

$$|\partial \Omega| \ge C(\Lambda)^{-1} \min\left\{1, |\Omega|^{(n-1)/n}\right\}, \qquad \forall \, \Omega \Subset M \text{ with smooth boundary.}$$
(3.3)

Once this is proved, then we recall the following result:

Theorem 3.11 ([40, Theorem 4.1]). Let M be of infinite volume, and whose isoperimetric profile function $ip(v) := inf \{ |\partial E| : E \in M, |E| = v \}$ satisfies

$$\liminf_{v \to \infty} \operatorname{ip}(v) \ge A > 0, \qquad \int_0^{v_0} \frac{dv}{\operatorname{ip}(v)} < \infty \quad \text{for some } v_0 > 0.$$

Then for all E_0 with $|\partial E_0| < A$, there exists a weak IMCF u' in M with initial value E_0 , such that $E_t(u') \subseteq M$ for some t > 0.

Joining this result with (3.3), it follows that we may pick $r_0 \ll 1$ and obtain an IMCF u' starting from E_0 , such that $E_t(u') \Subset M$ for some t > 0. Since u is the maximal solution, it follows that $E_t(u) \Subset E_t(u') \Subset M$ for the same t. This proves the lemma.

It remains to prove (3.3). We divide into two cases.

Case 1: $|\Omega \cap B(x,1)| \ge V/2$ for some $x \in M$. By continuity, we can find another point $x' \in M$ such that $|\Omega \cap B(x',1)| = V/2$. Recall a well-known isoperimetric inequality:

$$\min\left\{|B(x',1)\cap\Omega|,|B(x',1)\setminus\Omega|\right\} \le C(\Lambda)\left|\partial\Omega\cap B(x',1)\right|^{\frac{n}{n-1}},\tag{3.4}$$

and note that the left hand side is at least V/2. Thus $|\partial \Omega| \ge C(\Lambda)^{-1}$ for this case.

Case 2: $|\Omega \cap B(x,1)| < V/2$ for all $x \in M$. Then (3.4) implies $|B(x,1) \cap \Omega| \leq C(\Lambda)|B(x,1) \cap \partial \Omega|^{n/(n-1)}$ for all $x \in M$. By volume doubling, we may find finitely many balls $\{B(x_i,1)\}_{i=1}^m$ that covers $\overline{\Omega}$, whose covering multiplicity is bounded by $C(\Lambda)$.

$$\begin{aligned} |\Omega| &\leq \sum_{i=1}^{m} \left| \Omega \cap B(x_{i},1) \right| \leq C(\Lambda) \sum_{i=1}^{m} \left| \partial \Omega \cap B(x_{i},1) \right|^{\frac{n}{n-1}} \\ &\leq C(\Lambda) \Big(\sum_{i=1}^{m} \left| \partial \Omega \cap B(x_{i},1) \right| \Big)^{\frac{n}{n-1}} \leq C(\Lambda) |\partial \Omega|^{\frac{n}{n-1}}. \end{aligned}$$

Then (3.3) follows by combining the two cases.

Lemma 3.12. Suppose (M,g) satisfies (BG) and has only one end. Let $E_0 = B(p,r_0)$ be as in Lemma 3.10, and u be the maximal IMCF with initial condition E_0 , as given by Lemma 3.7(i). Then u is either proper, sweeping, or instantly escaping.

Proof. Let $T' := \sup \{t \ge 0 : E_t \in M\}$. By Lemma 3.10 we have T' > 0. If $T' = \infty$ then u is proper. Now assume $T' < \infty$. By Corollary 3.9(ii), we must have $E_t = M$ for all t > T'. Therefore $T' = \sup(u)$. Finally, consider $E_{T'}$: if $E_{T'} \in M$ then u is instantly escaping by Definition 3.5. Otherwise, we must have $E_{T'} = M$ by Corollary 3.9(ii) again. Hence $M = \bigcup_{t < T'} E_t$ with $E_t \in M$ for each t < T'. This implies that u is sweeping by Definition 3.5.

Remark 3.13. Both the bounded geometry and the one-endedness assumption cannot be removed. Indeed, Figure 4 shows three examples where:

(i) M has bounded geometry, but the maximal IMCF partially rushes to infinity at time T due to the presence of a cylindrical end;

(ii) for a manifold with unbounded geometry, the maximal IMCF could partially rush to infinity in finite time;

(iii) M has an end with finite volume (thus has unbounded geometry), and the maximal IMCF rushes to infinity at time zero.



FIGURE 4. Examples of maximal IMCF that do not fall in Definition 3.5.

4. INSTANTLY ESCAPING IMCF

Consider an instantly escaping maximal solution u of the IMCF with a bounded initial value (see Definition 3.5). This means that there exists $T \in (0, \infty)$ such that

$$E_T(u) \Subset M, \qquad u \equiv T \text{ in } M \setminus E_T(u).$$

The proof of the main theorems 1.1, 1.2 are based on finding exhaustions of M by level sets of IMCFs. However, for an instantly escaping solution u, there is no level set outside $E_T(u)$. We will modify the underlying metric and slightly enlarge it at infinity, so that a level set of the new maximal IMCF will appear in the edited region. Letting the edited region tend to infinity and the perturbation tend to zero, we finally obtain the desired sequence of exhausting surfaces. The main result of this section is Theorem 4.3, which is based on Lemma 4.2 (the main perturbation lemma).



FIGURE 5. Left: the instantly escaping maixmal IMCF. Right: the metric perturbation and the new level set (see Lemma 4.2 below).

First, we prove the following lemma which says that $|\partial E_T|$ equals the "circumference at infinity" of M.

Lemma 4.1. Suppose u is an escaping maximal IMCF starting from some $C^{1,1}$ initial value $E_0 \Subset M$. Let T be the escaping time of u. Then there exists an exhaustion of M by $C^{1,\alpha}$ domains $\{F_k\}_{k=1}^{\infty}$ such that $E_T(u) \Subset F_1 \Subset F_2 \Subset \cdots \Subset M$ and $\lim_{k\to\infty} |\partial F_k| = |\partial E_T(u)|$.

Proof. First, we fix a sequence of precompact smooth domains $\{\Omega_l\}_{l=1}^{\infty}$ with $E_T(u) \in \Omega_1 \in \Omega_2 \in \cdots \in M$ and $\bigcup_{l \ge 1} \Omega_l = M$. By Lemma 3.7(ii), we know that there exists a sequence of weak solutions $u_l \in \text{Lip}_{\text{loc}}(\Omega_l)$ that converge to u in C_{loc}^0 , and by Lemma 3.7(ii), we have $E_T(u_l) = E_T(u)$ for all l. For each $\varepsilon > 0$, note the following facts:

- (1) $\bigcup_{l\geq 1} E_{T+\varepsilon}(u_l) = M$. This follows from $u_l \xrightarrow{C_{loc}^0} u$ and $u \leq T$.
- (2) By Lemma 3.7(iid) and the above discussions, we have

$$\left|\partial E_{T+\varepsilon}(u_l)\right| \le e^{\varepsilon} \left|\partial E_T(u_l)\right| = e^{\varepsilon} \left|\partial E_T(u)\right|$$
 for all l .

So we can inductively choose a sequence $l_k \to \infty$ so that

$$E_{T+1/k}(u_{l_k}) \supseteq B_g(p,k) \cup E_{T+1/(k-1)}(u_{l_{k-1}}).$$

Since $\partial E_T(u)$ is outward minimizing, this implies

$$|\partial E_T(u)| \le |\partial E_{T+1/k}(u_{l_k})| \le e^{1/k} |\partial E_T(u)|.$$

This proves the lemma by setting $F_k = E_{T+1/k}(u_{l_k})$.

The following lemma shows that for any compact subset K, we can perturb the metric near infinity, so that the maximal solution "jumps" out of K but stays compact until a slightly larger time t > T.

Lemma 4.2. Suppose (M^n, g) has only one end, and there are $\Lambda_k > 0$ for each k such that

$$\operatorname{inj} \ge \Lambda_0^{-1}, \qquad |\nabla^k \operatorname{Rm}| \le \Lambda_k.$$
 (4.1)

Suppose u is an instantly escaping maximal IMCF on M with an initial value $E_0 \subseteq M$. Let $T \in (0, \infty)$ be the escape time of u.

Then for any $K \subseteq M$ and $\varepsilon \in (0,1)$, there is a smooth metric \tilde{g} with $\|\tilde{g} - g\|_{C_g^{10}} \leq \varepsilon$ and with the following properties. If \tilde{u} is the maximal IMCF in (M, \tilde{g}) starting from E_0 , then:

- (i) $E_T(\widetilde{u}) = E_T(u),$
- (ii) there exists $t \in (T, T + \varepsilon)$ such that $E_t(\widetilde{u}) \Subset M$ and $\partial E_t(\widetilde{u}) \cap (M \setminus K) \neq \emptyset$.

Proof. Without loss of generality, we may assume $K \supseteq E_T(u)$. By Lemma 4.1, we can replace K by a larger domain and assume that it satisfies

$$\left|\partial K\right|_{g} < (1+\varepsilon)^{n-2} \left|\partial E_{T}(u)\right|. \tag{4.2}$$

Fix a point $p \in E_0$. Assume $K \subseteq B_g(p, R_1)$ for some $R_1 > 0$. We use C > 0 to denote a generic constant that only depends on finitely many Λ_k .

Claim 1. There exist $R_2 > R_1$ and a smooth Riemannian metric \tilde{g} , such that

- (i) $\widetilde{g} \ge g$ and $\|\widetilde{g} g\|_{C^{10}_a} \le C\varepsilon$;
- (ii) $\widetilde{g} = g$ in $B_q(p, R_1)$, and $\widetilde{g} = (1 + \varepsilon)g$ in $M \setminus B_q(p, R_2)$.

Proof. First, we smooth the distance function $d_g(p, \cdot)$ by evolving it by a heat equation: Let w_t be the solution to $\partial_t w_t = \Delta_g w_t$ with $w_0 = d_g(p, \cdot)$. Then w_t is smooth for any t > 0. For any fixed $x \in M$, we define a time-dependent function $v_t(y) := w_t(y) - w_0(x)$ for all $y \in M$, then by the triangle inequality we have $|v_0(y)| \leq d_g(x, y)$. By the assumption (4.1), the heat

kernel satisfies the Gaussian bound $H(x,t;y,s) \leq \frac{C}{(t-s)^{n/2}} e^{-\frac{d_g^2(x,y)}{C(t-s)}}$ for all $s < t \in [0, C^{-1}]$, see e.g. [16, Corollary 26.26]. So using the representation formula

$$w_t(x) = \int_M H(x,t;y,0)w_0(y) \, d\mathrm{vol}_g y,$$

it is easy to see $||v_t||_{C^0(B_g(x,1))} \leq C$ for all $t \in [0, C^{-1}]$. So by the standard parabolic estimate [25], it follows for each $t \in [\frac{1}{2}C^{-1}, C^{-1}]$ that

$$\|w_t - w_0(x)\|_{C_g^{10}(B_g(x,1))} = \|v_t\|_{C_g^{10}(B_g(x,1))} \le C \|v_t\|_{C_g^{0}(B_g(x,1)\times[0,C^{-1}])} \le C.$$

Therefore, the function $\widetilde{d} := w_{C^{-1}}$ satisfies

$$\|\nabla \widetilde{d}\|_{C_g^9} \le C$$
, and $|\widetilde{d}(x) - d_g(p, x)| \le C$ for all $x \in M$.

Let $R_2 > R_1 + 3C$ and let $\eta : \mathbb{R} \to [0,1]$ be a smooth non-decreasing function, such that $\eta \equiv 0$ on $(-\infty, R_1 + C], \eta \equiv 1$ on $[R_2 - C, \infty)$, and $\|\eta\|_{C_g^{10}} \leq 100$. Then the metric $\tilde{g} = (1 + \varepsilon \eta \circ \tilde{d}) \cdot g$ satisfies all assertions.

In the following, we fix the choice of $R_2 > R_1$ and \tilde{g} as in Claim 1. Then we see that \tilde{g} satisfies (4.1) for all $k \leq 8$ with possibly larger Λ_k depending only on C. Thus Lemma 3.8 and Corollary 3.9 hold in (M, \tilde{g}) with a weaker constant. It is helpful to recall the chain of inclusions

$$E_T(u) \Subset K \Subset B_g(p, R_1) \Subset \{ \widetilde{g} = g \} \Subset \{ \widetilde{g} \neq (1 + \varepsilon)g \} \Subset B_g(p, R_2).$$

Claim 2. u is also a weak solution of IMCF in (M, \tilde{g}) with initial value E_0 .

Proof. By Definition 3.2, it suffices to show that u is a weak solution of IMCF in $(M \setminus \overline{E_0}, \tilde{g})$. By Definition 3.1, this is to show that for any $t \in \mathbb{R}$, $K' \in M \setminus \overline{E_0}$ and set F with $F \Delta E_t \in K'$, we have $J_{u,\tilde{g}}^{K'}(E_t) \leq J_{u,\tilde{g}}^{K'}(F)$. If t > T then $E_t = M$, hence $F = M \setminus K''$ for some $K'' \in M \setminus \overline{E_0}$. In this case $J_{u,\tilde{g}}^{K'}(E_t) \leq J_{u,\tilde{g}}^{K'}(F)$ is equivalent to $|\partial K''| + \int_{K''} |\nabla_{\tilde{g}}u| \geq 0$, which is obviously true. Now suppose $t \leq T$. Note the following facts: $\tilde{g} \geq g$, and $\partial E_t(u) \subset \{\tilde{g} = g\}$, and $\int_A |\nabla_{\tilde{g}}u| dV_{\tilde{g}} = \int_A |\nabla_g u| dV_g$ for all set A (since $u \equiv T$ wherever $\tilde{g} \neq g$). As a result, we have

$$\begin{aligned} J_{u,\widetilde{g}}^{K'}(F) &= |\partial F \cap K'|_{\widetilde{g}} - \int_{F \cap K'} |\nabla_{\widetilde{g}} u| \, dV_{\widetilde{g}} \ge |\partial F \cap K'|_g - \int_{F \cap K'} |\nabla_g u| \, dV_g \\ &= J_{u,g}^{K'}(F) \ge J_{u,g}^{K'}(E_t) = J_{u,\widetilde{g}}^{K'}(E_t), \end{aligned}$$
he claim.

proving the claim.

Let \tilde{u} be the maximal weak IMCF in (M, \tilde{g}) with initial value E_0 . Hence we have $\tilde{u} \ge u$. This implies $E_T(\tilde{u}) \subset E_T(u) \Subset M$. Since u is also a weak solution in (M, \tilde{g}) with initial value E_0 , by Lemma 3.4 we have $E_T(\tilde{u}) = E_T(u)$.

Claim 3. For all $t < T + (n-1)\log(1+\varepsilon)$, we have $E_t(\widetilde{u}) \in M$.

Proof. Fix such a t. Let $\{\Omega_l\}_{l=1}^{\infty}$ be a smooth precompact exhaustion with

$$B_q(p, R_2) \Subset \Omega_1 \Subset \Omega_2 \Subset \cdots \Subset M.$$

Apply Lemma 3.7(ii) to this exhaustion: we obtain a sequence \tilde{u}_l of maximal weak IMCFs in (Ω_l, \tilde{g}) with the same initial value E_0 . Then $\{\tilde{u}_l\}$ is a descending sequence, and $\tilde{u}_l \searrow \tilde{u}$ in $C^0_{\text{loc}}(M \setminus E_0)$. In particular, note that $E_T(\tilde{u}_l) \subset E_T(\tilde{u}) \Subset \Omega_l$. Applying Lemma 3.4 in Ω_l , it follows that $E_T(\tilde{u}_l) = E_T(\tilde{u}) = E_T(u)$ for each l.

By Corollary 3.9(ii) and (4.1) and the fact that M is one-ended, to prove the claim it suffices to prove that $E_t(\tilde{u}) \neq M$. Suppose this is not the case. As $E_t(\tilde{u}) = \bigcup_{l \geq 1} E_t(\tilde{u}_l)$, we would have $E_t(\tilde{u}_l) \supseteq B_g(p, R_2)$ for some sufficiently large l. This implies

$$\partial E_t(\widetilde{u}_l) \subset M \setminus B_g(p, R_2) \subset \left\{ \widetilde{g} = (1 + \varepsilon)g \right\}$$

So by our definition of t and Lemma 3.7(iid), we have

$$|\partial E_t(\widetilde{u}_l)|_g = \frac{|\partial E_t(\widetilde{u}_l)|_{\widetilde{g}}}{(1+\varepsilon)^{n-1}} \le e^{t-T} \frac{|\partial E_T(\widetilde{u}_l)|_{\widetilde{g}}}{(1+\varepsilon)^{n-1}} = e^{t-T} \frac{|\partial E_T(u)|_g}{(1+\varepsilon)^{n-1}} < |\partial E_T(u)|_g.$$

So we find a surface outside of $E_T(u)$ that has a strictly smaller *g*-perimeter, but this contradicts the outward minimizing of $E_T(u)$.

Claim 4. If
$$T + (n-2)\log(1+\varepsilon) < t < T + (n-1)\log(1+\varepsilon)$$
, then $E_t(\widetilde{u}) \cap (M \setminus K) \neq \emptyset$.

Proof. Suppose the claim is false at such a time t. Note that $E_t(\widetilde{u}) \subset K \subseteq M$. By Lemma 3.3(iii) and noting that $\widetilde{g} = g$ in K and $E_T(\widetilde{u}) = E_T(u) \subseteq K$, we have

$$\left|\partial E_t(\widetilde{u})\right|_{\widetilde{g}} = e^{t-T} \left|\partial E_T(\widetilde{u})\right|_{\widetilde{g}} > (1+\varepsilon)^{n-2} |\partial E_T(u)|_g.$$

But recalling (4.2) and $\tilde{g} = g$ on K this implies

$$|\partial K|_{\widetilde{g}} = |\partial K|_g < (1+\varepsilon)^{n-2} |\partial E_T(u)|_g < |\partial E_t(\widetilde{u})|_{\widetilde{g}},$$

contradicting the outward minimizing property of $\partial E_t(\tilde{u})$ since we assumed $E_t(\tilde{u}) \subset K$. \Box

Combining Claim 2 and 3, the lemma is proven.

Theorem 4.3. Suppose that M^n is one-ended and satisfies $b_1(M) < \infty$, and that there are $\Lambda_k > 0$ for each k such that

$$\operatorname{inj} \ge \Lambda_0^{-1}, \quad |\nabla^k \operatorname{Rm}| \le \Lambda_k.$$
 (4.3)

Suppose u is an escaping maximal IMCF on M with a $C^{1,1}$ initial value $E_0 \Subset M$. Let $T \in (0, \infty)$ be the escape time of u.

Then there exists C > 0 such that for any compact subset $K \supset E_T(u)$ and $\delta > 0$, there exists a domain $\Omega \supseteq K$ such that

(i) $\partial \Omega$ is a connected $C^{1,\alpha}$ surface with diam $(\Omega) \leq C$ and $C^{1,\alpha}$ norm controlled by C;

(ii) For any F with $E_T(u) \in F \in M$, we have $|\partial \Omega|_g \leq (1+\delta)|\partial F|_g$.

In particular, there exists a sequence of exhaustion by precompact subsets $\{\Omega_k\}_{k=1}^{\infty}$ satisfying (i) and (ii) for a sequence $\delta_k \to 0$ as $k \to \infty$.

Proof. Note that M is one-ended. Thus for any $K' \in M$ and $\varepsilon < 1/100$, by Lemma 4.2 we find a metric \tilde{g} with $\|\tilde{g} - g\|_{C_g^{10}} \leq \varepsilon$ and a maximal IMCF \tilde{u} in (M^n, \tilde{g}) , such that $E_T(u) = E_T(\tilde{u})$, and for some $t \in (T, T + \varepsilon)$ we have $E_t(\tilde{u}) \in M$, $\partial E_t(\tilde{u}) \cap (M \setminus K') \neq \emptyset$.

By Lemma 3.7(ib,ic), $E_t(\tilde{u})$ is connected, and $M \setminus E_t(\tilde{u})$ has only unbounded connected components. Combining Corollary 3.9(i) and $\|\tilde{g} - g\|_{C_g^{10}} \leq 1/100$, we have that each connected component Σ of $\partial E_t(\tilde{u})$ satisfies

$$\operatorname{diam}_{q}(\Sigma) \leq C\left(T+1, \Lambda, |\partial E_{0}|, H^{+}_{\partial E_{0}}\right) =: C_{1}.$$
(4.4)

In particular, this bound is independent of K'. Now let S be the compact set obtained in Lemma 2.6, and choose K' such that $N_g(S \cup K, C_1) \Subset K' \Subset M$, where $N_g(X, r)$ denotes the *r*-neighborhood of a subset X. Then it follows from $\partial E_t(\widetilde{u}) \cap (M \setminus K') \neq \emptyset$ that some connected component of $\partial E_t(\widetilde{u})$ is disjoint from $S \cup K$. Inserting this into Lemma 2.6, we see that $\partial E_t(\widetilde{u})$ is connected. Hence by (4.4) again we have $\partial E_t(\widetilde{u}) \cap K = \emptyset$, which implies $E_t(\widetilde{u}) \supseteq K$.

Moreover, by Lemma 3.8(ii) and $\|\tilde{g} - g\|_{C_g^{10}} \leq 1/100$, $\partial E_t(\tilde{u})$ has bounded $C^{1,\alpha}$ norm with respect to \tilde{g} , hence with respect to g as well. This proves (i) by setting $\Omega = E_t(\tilde{u})$.

Since $E_T(\widetilde{u})$ is outward minimizing in (M, \widetilde{g}) , for any $F \supseteq E_T(\widetilde{u})$ we have

$$(1+\varepsilon)^{n-1}|\partial F|_g \ge |\partial F|_{\widetilde{g}} \ge |\partial E_T(\widetilde{u})|_{\widetilde{g}} \ge e^{-(t-T)}|\partial\Omega|_{\widetilde{g}} \ge e^{-\varepsilon}|\partial\Omega|_g.$$

By taking ε sufficiently small, this implies assertion (ii) since $E_T(\widetilde{u}) = E_T(u)$.

5. Proof of the main theorems

Now we can prove Theorem 1.1 and 1.2. Recall that in these two theorems, the manifolds have non-negative scalar curvature and bounded geometry.

Remark 5.1. Let (M, g) be a manifold as in Theorem 1.1 or 1.2. Without loss of generality, we may assume that the scalar curvature is strictly positive, and there are constants $\Lambda_k > 0$ for each $k \ge 0$ such that

$$\operatorname{inj} \ge \Lambda_0^{-1}, \qquad |\nabla^k \operatorname{Rm}| \le \Lambda_k.$$
 (5.1)

Indeed, thanks to the bounded geometry assumption, we can run a smooth complete Ricci flow (M, g(t)), $t \in [0, t_0]$ with g(0) = g for some $t_0 > 0$, and $(M, g(t_0))$ satisfies (5.1) by Shi's derivative estimates [34]. Moreover, if R > 0 does not hold everywhere on $(M, g(t_0))$, then by the strong maximum principle, (M, g(t)) is flat for all $t \in [0, t_0]$. In the context of Theorem 1.1, this implies $M \cong \mathbb{R}^3$. In the context of Theorem 1.2, this leads to a contradiction. Therefore, replacing g with $g(t_0)$, we may assume that (5.1) holds and R > 0 strictly.

Proof of Theorem 1.1. By Remark 5.1, we may assume R > 0 and (5.1) holds.

Fix $E_0 = B(p, r_0)$ as in Lemma 3.10, and let u be the maximal weak solution of IMCF starting from E_0 . Since (M, g) is contractible, by Lemma 2.1 it has one end, and then Lemma 3.12 implies that u is either proper, sweeping, or instantly escaping. We prove the main theorem in three cases.

Case 1: Suppose u is proper. Since ∂E_0 is connected and M is contractible, by Lemma 3.7(ib, ic) and Lemma 2.5 we see that all the ∂E_t (t > 0) are connected. By Lemma 3.6 we have

$$\int_0^t 4\pi \chi(\partial E_t) \, dt \ge \int_{\partial E_t} H^2 \, d\mu_t - \int_{\partial E_0} H^2 \, d\mu_0 \ge - \int_{\partial E_0} H^2 \, d\mu_0, \quad \forall t > 0.$$

Thus there exists a sequence $t_i \to \infty$ such that ∂E_{t_i} is either an \mathbb{S}^2 or \mathbb{T}^2 , for all *i*. Since $u \in \operatorname{Lip}_{\operatorname{loc}}(M)$, the sets E_{t_i} must form an exhaustion of M. Thus we obtain an exhaustion of M by precompact domains with boundaries either \mathbb{S}^2 or \mathbb{T}^2 . If there are infinitely many

 \mathbb{S}^2 , then Lemma 2.3 implies that $M \cong \mathbb{R}^3$. If there are infinitely many \mathbb{T}^2 , Lemma 2.4 implies that $M \cong \mathbb{R}^3$.

Case 2: Suppose u is sweeping. Then recall from Definition 3.5 that there exists $T \in (0,\infty)$ such that $E_t \in M$ for all t < T, and $E_T = \bigcup_{t \in [0,T)} E_t = M$. By Lemma 3.7(ib, ic) and Lemma 2.5, each ∂E_t is connected. Then by Corollary 3.9, for all $t \in [0,T)$ we have that ∂E_t is uniformly $C^{1,\alpha}$ -bounded and has uniformly bounded diameters. Thus, we can select a sequence $t_i \nearrow T$ such that

$$\lim_{i \to \infty} d_g \left(\partial E_{t_{i-1}}, \partial E_{t_i} \right) = \infty.$$
(5.2)

Next, recall that each E_{t_i} is outward area-minimizing. Hence for each i and set F with $E_{t_{i-1}} \subset F \Subset M$, we have

$$\left|\partial F\right| \ge \left|\partial E_{t_{i-1}}(u)\right| = e^{t_{i-1}-t_i} \left|\partial E_{t_i}(u)\right|.$$

$$(5.3)$$

Now fix basepoints $p_i \in \partial E_{t_i}$, so $p_i \to \infty$ as $i \to \infty$. By (5.1), we may pass to a subsequence and assume (M, g, p_i) converge smoothly to a manifold $(M_{\infty}, g_{\infty}, p_{\infty})$ with $R \ge 0$, and ∂E_{t_i} converge in the $C^{1,\beta}$ -sense for some $\beta \in (0, \alpha)$ to a connected closed surface $\Sigma_{\infty} \subset M_{\infty}$. Combining (5.3) and (5.2) and the standard set replacing argument (see for example [27, Theorem 21.14]), it follows that Σ_{∞} is locally area-minimizing. Since g_{∞} has nonnegative scalar curvature, it follows by Schoen-Yau [32] that Σ_{∞} must be \mathbb{S}^2 or \mathbb{T}^2 . Hence there is an infinite subsequence of $\{\partial E_{t_i}\}$ which either consists of \mathbb{S}^2 or consists of \mathbb{T}^2 . This proves the theorem as in the end of Case 1.

Case 3: Suppose u is escaping. Then let $\{\Omega_k\}_{k=1}^{\infty}$ be the exhaustion of $C^{1,\alpha}$ domains as in Theorem 4.3. Fixing basepoints $p_k \in \partial \Omega_k$, we have that up to a subsequence, (M, g, p_k) smoothly converges to $(M_{\infty}, g_{\infty}, p_{\infty})$ with $R \geq 0$, and $\partial \Omega_k$ converge in $C^{1,\beta}$ -sense to a connected closed surface $\Sigma_{\infty} \subset M_{\infty}$. By Theorem 4.3(ii) and set replacing, Σ_{∞} is in fact locally area-minimizing. Arguing as in Case 2, this proves the theorem.

Proof of Theorem 1.2. Suppose (M, g) is a solid handlebody, where g is complete with $R \ge 0$ and bounded geometry. Due to Remark 5.1, we may assume that R > 0 and (5.1) holds. Same as the proof of Theorem 1.1, consider a maximal weak IMCF u starting from a sufficiently small geodesic ball. In either case, the same argument in Theorem 1.1 implies that M can be exhausted by bounded domains with \mathbb{S}^2 or \mathbb{T}^2 boundaries. By Lemma 2.7, the genus of M is at most 1.

References

- V. Agostiniani, M. Fogagnolo, and L. Mazzieri. Sharp geometric inequalities for closed hypersurfaces in manifolds with nonnegative Ricci curvature. *Invent. Math.*, 222(3):1033–1101, 2020.
- [2] L. Bessières, G. Besson, and S. Maillot. Ricci flow on open 3-manifolds and positive scalar curvature. Geom. Topol., 15(2):927-975, 2011.
- [3] H. L. Bray and A. Neves. Classification of prime 3-manifolds with Yamabe invariant greater than ℝP³. Ann. of Math. (2), 159(1):407-424, 2004.
- [4] S. Brendle and O. Chodosh. A volume comparison theorem for asymptotically hyperbolic manifolds. Comm. Math. Phys., 332(2):839–846, 2014.
- [5] S. Brendle, P.-K. Hung, and M.-T. Wang. A Minkowski inequality for hypersurfaces in the anti-de Sitter–Schwarzschild manifold. Comm. Pure Appl. Math., 69(1):124–144, 2016.
- [6] S. Cecchini, D. Räde, and R. Zeidler. Nonnegative scalar curvature on manifolds with at least two ends. J. Topol., 16(3):855–876, 2023.
- [7] S. Chang, S. Weinberger, and G. Yu. Taming 3-manifolds using scalar curvature. Geom. Dedicata, 148:3–14, 2010.

- [8] J. Chen, P. Liu, Y. Shi, and J. Zhu. Incompressible hypersurface, positive scalar curvature and positive mass theorem. available at https://arxiv.org/abs/2112.14442, 2021.
- [9] S. Chen. A generalization of the Geroch conjecture with arbitrary ends. Math. Ann., 389(1):489–513, 2024.
- [10] O. Chodosh. Large isoperimetric regions in asymptotically hyperbolic manifolds. Comm. Math. Phys., 343(2):393–443, 2016.
- [11] O. Chodosh and M. Eichmair. Global uniqueness of large stable CMC spheres in asymptotically flat Riemannian 3-manifolds. Duke Math. J., 171(1):1–31, 2022.
- [12] O. Chodosh, M. Eichmair, Y. Shi, and H. Yu. Isoperimetry, scalar curvature, and mass in asymptotically flat Riemannian 3-manifolds. *Comm. Pure Appl. Math.*, 74(4):865–905, 2021.
- [13] O. Chodosh, M. Eichmair, Y. Shi, and J. Zhu. Characterization of large isoperimetric regions in asymptotically hyperbolic initial data. Comm. Math. Phys., 368(2):777–798, 2019.
- [14] O. Chodosh and C. Li. Generalized soap bubbles and the topology of manifolds with positive scalar curvature. Ann. of Math. (2), 199(2):707–740, 2024.
- [15] O. Chodosh, D. Maximo, and A. Mukherjee. Complete Riemannian 4-manifolds with uniformly positive scalar curvature. https://arxiv.org/abs/2407.05574, 2024.
- [16] B. Chow, S.-C. Chu, D. Glickenstein, C. Guenther, J. Isenberg, T. Ivey, D. Knopf, P. Lu, F. Luo, and L. Ni. The Ricci flow: techniques and applications. Part III. Geometric-analytic aspects, volume 163 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 2010.
- [17] R. Geroch. Energy extraction. Annals of the New York Academy of Sciences, 224(1):108–117, 1973.
- [18] M. Gromov. Four lectures on scalar curvature. In *Perspectives in scalar curvature*. Vol. 1, pages 1–514. World Sci. Publ., Hackensack, NJ, [2023] ©2023.
- [19] M. Gromov and H. B. Lawson, Jr. Positive scalar curvature and the Dirac operator on complete Riemannian manifolds. Inst. Hautes Études Sci. Publ. Math., (58):83–196, 1983.
- [20] G. Huisken and T. Ilmanen. The inverse mean curvature flow and the Riemannian Penrose inequality. J. Differential Geom., 59(3):353–437, 2001.
- [21] G. Huisken and T. Koerber. Inverse mean curvature flow and ricci-pinched three-manifolds. Journal für die reine und angewandte Mathematik (Crelles Journal), 2024(814):1–8, 2024.
- [22] L. S. Husch and T. M. Price. Finding a boundary for a 3-manifold. Ann. of Math. (2), 91:223–235, 1970.
- [23] J. L. Jauregui and D. A. Lee. Lower semicontinuity of mass under C⁰ convergence and Huisken's isoperimetric mass. J. Reine Angew. Math., 756:227–257, 2019.
- [24] B. Kotschwar and L. Ni. Local gradient estimates of p-harmonic functions, 1/H-flow, and an entropy formula. Ann. Sci. Éc. Norm. Supér. (4), 42(1):1–36, 2009.
- [25] N. V. Krylov. Lectures on Elliptic and Parabolic Equations in Hölder Spaces, volume 12 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 1996.
- [26] J. Lott. Some obstructions to positive scalar curvature on a noncompact manifold. available at https: //arxiv.org/abs/2402.13239, 2024.
- [27] F. Maggi. Sets of finite perimeter and geometric variational problems, volume 135 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 2012. An introduction to geometric measure theory.
- [28] L. Mari, M. Rigoli, and A. G. Setti. On the 1/H-flow by p-Laplace approximation: new estimates via fake distances under Ricci lower bounds. Amer. J. Math., 144(3):779–849, 2022.
- [29] G. Perelman. The entropy formula for the Ricci flow and its geometric applications. http://arxiv.org/abs/math/0211159, 2002.
- [30] G. Perelman. Finite extinction time for the solutions to the Ricci flow on certain three-manifolds. http://arxiv.org/abs/math/0307245, 2003.
- [31] A. Ryabichev. Short proof of the Kneser-Edmonds theorem on the degree of a map between closed surfaces. https://arxiv.org/abs/2401.01041, 2023.
- [32] R. Schoen and S.-T. Yau. Existence of incompressible minimal surfaces and the topology of three dimensional manifolds with non-negative scalar curvature. Annals of Mathematics, 110(1):127–142, 1979.
- [33] R. Schoen and S. T. Yau. Complete three-dimensional manifolds with positive ricci curvature and scalar curvature. In *Seminar on differential geometry*, volume 102, pages 209–228. Princeton University Press Princeton, 1982.

OTIS CHODOSH, YI LAI, AND KAI XU

- [34] W.-X. Shi. Deforming the metric on complete Riemannian manifolds. J. Differential Geom., 30(1):223– 301, 1989.
- [35] Y. Shi. The isoperimetric inequality on asymptotically flat manifolds with nonnegative scalar curvature. Int. Math. Res. Not. IMRN, (22):7038–7050, 2016.
- [36] J. Wang. Topology of 3-manifolds with uniformly positive scalar curvature, 2023.
- [37] J. Wang. Contractible 3-manifold and Positive scalar curvature (I). J. Differential Geom., 127(3):1267– 1304, 2024.
- [38] J. Wang. Contractible 3-manifolds and positive scalar curvature (II). J. Eur. Math. Soc. (JEMS), 26(2):537–572, 2024.
- [39] K. Xu. Inverse mean curvature flow with outer obstacle, 2024.
- [40] K. Xu. Isoperimetry and the properness of weak inverse mean curvature flow. Calculus of Variations and Partial Differential Equations, 63(8):216, 2024.
- [41] S. T. Yau. Problem section. In Seminar on differential geometry, volume 102, pages 669–706, 1982.
- [42] J. Zhu. Calabi-Yau type theorem for complete manifolds with nonnegative scalar curvature. available at https://arxiv.org/abs/2402.15118, 2024.

Otis Chodosh

DEPARTMENT OF MATHEMATICS, STANFORD UNIVERSITY, CA 94305, USA Email address: ochodosh@stanford.edu

Yi Lai

DEPARTMENT OF MATHEMATICS, UC IRVINE, CA 92697, USA Email address: ylai25@uci.edu

Kai Xu

DEPARTMENT OF MATHEMATICS, DUKE UNIVERSITY, DURHAM, NC 27705, USA Email address: kai.xu631@duke.edu

18