Algorithmic contiguity from low-degree conjecture and applications in correlated random graphs

Zhangsong Li Peking University

February 17, 2025

Abstract

In this paper, assuming a natural strengthening of the low-degree conjecture, we provide evidence of computational hardness for two problems: (1) the (partial) matching recovery problem in the sparse correlated Erdős-Rényi graphs $\mathcal{G}(n,q;\rho)$ when the edge-density $q = n^{-1+o(1)}$ and the correlation $\rho < \sqrt{\alpha}$ lies below the Otter's threshold, solving a remaining problem in [15]; (2) the detection problem between the correlated sparse stochastic block model $\mathcal{S}(n,\frac{\lambda}{n};k,\epsilon;s)$ and a pair of independent stochastic block models $\mathcal{S}(n,\frac{\lambda s}{n};k,\epsilon)$ when $\epsilon^2\lambda s < 1$ lies below the Kesten-Stigum (KS) threshold and $s < \sqrt{\alpha}$ lies below the Otter's threshold, solving a remaining problem in [9].

One of the main ingredient in our proof is to derive certain forms of *algorithmic* contiguity between two probability measures based on bounds on their low-degree advantage. To be more precise, consider the high-dimensional hypothesis testing problem between two probability measures \mathbb{P} and \mathbb{Q} based on the sample Y. We show that if the low-degree advantage $\operatorname{Adv}_{\leq D}\left(\frac{d\mathbb{P}}{d\mathbb{Q}}\right) = O(1)$, then (assuming the low-degree conjecture) there is no efficient algorithm \mathcal{A} such that $\mathbb{Q}(\mathcal{A}(Y) = 0) = 1 - o(1)$ and $\mathbb{P}(\mathcal{A}(Y) = 1) = \Omega(1)$. This framework provides a useful tool for performing reductions between different inference tasks.

Contents

T.	Introduction	2
	1.1 Key challenges and innovations	5
	1.2 Notation	7
	1.3 Organization of this paper	8
2	Low-degree framework and algorithmic contiguity	8
3	Partial recovery in correlated random graphs	14
3	Partial recovery in correlated random graphs 3.1 Proof of Item (1) in Proposition 3.3	14 15
3	Partial recovery in correlated random graphs3.1 Proof of Item (1) in Proposition 3.33.2 Proof of Item (2) in Proposition 3.3	14 15 22

0

4 Detection in correlated SBMs

A Preliminary results in graphs and probability

1 Introduction

Graph matching, also referred to as network alignment, is the problem of identifying a bijection between the vertex sets of two graphs that maximizes the number of common edges. When the two graphs are exactly isomorphic to each other, this problem reduces to the classical graph isomorphism problem, for which the best known algorithm runs in quasi-polynomial time [1]. In general, graph matching is an instance of the *quadratic assignment problem* [7], which is known to be NP-hard to solve or even approximate [38].

Motivated by real-world applications (such as social network de-anonymization [45] and computational biology [51]) as well as the need to understand the average-case computational complexity, recent research has focused on developing theoretical foundations and efficient algorithms for graph matching under statistical models. These models assume that the two graphs are randomly generated with correlated edges under a hidden vertex correspondence, and a canonical model among them is the following correlated random graph model. For any integer n, denote by $U = U_n$ the set of unordered pairs (i, j) with $1 \le i \ne j \le n$.

Definition 1.1 (Correlated random graph model). Given an integer $n \ge 1$, for $(i, j) \in U_n$ let $J_{i,j}$ and $K_{i,j}$ be independent Bernoulli variables with parameter s. In addition, let π_* be an independent uniform permutation on $[n] = \{1, \ldots, n\}$. Then, we define a triple of correlated random graphs (G, A, B) as follows: first we generate G independently with $\{I_{i,j}, K_{i,j}\}$ and π_* from a specific probability distribution over all graphs on [n], and then (conditioned on G) we define for each $(i, j) \in U_n$ that (note that we identify a graph with its adjacency matrix)

$$A_{i,j} = G_{i,j}J_{i,j}, \quad B_{i,j} = G_{i,j}K_{\pi_*^{-1}(i),\pi_*^{-1}(j)}.$$

In short, we will subsample A, B from G with subsampling probability s and then permute the vertices of B by a uniform permutation.

Of particular interest in our paper are the following two special cases, namely the correlated Erdős-Rényi models and the correlated stochastic block models (SBMs).

Definition 1.2 (Correlated Erdős-Rényi graph model). Given an integer $n \ge 1$ and two parameters $p, s \in (0, 1)$, we generate a triple of correlated random graphs (G, A, B) such that we first generate G according to an Erdős-Rényi graph distribution $\mathcal{G}(n, p)$ (i.e., for each $(i, j) \in U_n$ we connect (i, j) in G independently with probability p), and then generate (A, B) from G according to Definition 1.1. For ease of presentation, we shall reparameterize such that q = ps and $\rho = \frac{s(1-p)}{1-ps}$ respectively. We will denote the marginal law of (A, B) as $\mathcal{G}(n,q;\rho)$.

Definition 1.3 (Stochastic block model). Given an integer $n \ge 1$ and three parameters $k \in \mathbb{N}, \lambda > 0, \epsilon \in (0, 1)$, we define a random graph G as follows: (1) sample a labeling $\sigma_* \in [k]^n = \{1, \ldots, k\}^n$ uniformly at random; (2) for every distinct pair $(i, j) \in U_n$, we let $G_{i,j}$ be an independent Bernoulli variable such that $G_{i,j} = 1$ (which represents that there is an undirected edge between i and j) with probability $\frac{(1+(k-1)\epsilon)\lambda}{n}$ if $\sigma_*(i) = \sigma_*(j)$ and with probability $\frac{(1-\epsilon)\lambda}{n}$ if $\sigma_*(i) \neq \sigma_*(j)$. In this case, we say that G is sampled from a stochastic block model $S(n, \frac{\lambda}{n}; k, \epsilon)$.

Definition 1.4 (Correlated stochastic block models). Given an integer $n \ge 1$ and four parameters $k \in \mathbb{N}, \lambda > 0, \epsilon, s \in (0, 1)$, for $(i, j) \in U_n$ let $J_{i,j}$ and $K_{i,j}$ be independent Bernoulli variables with parameter s. We define a triple of correlated random graphs (G, A, B) as follows: we first sample G according to the law of a stochastic block model $S(n, \frac{\lambda}{n}; k, \epsilon)$ and then generate (A, B) from G according to Definition 1.1. We will denote the marginal law of (A, B) as $S(n, \frac{\lambda}{n}; k, \epsilon; s)$.

Two fundamental problems in the study of correlated random graph models are as follows: (1) the *detection* problem, which involves determining whether a given pair of graphs (A, B) is sampled from a pair of correlated random graphs or from a pair of independent random graphs; (2) the *matching* problem, which focuses on recovering the latent matching π_* from a sample (A, B) from the distribution of correlated random graphs. In recent years, significant progress has been made in understanding these problems for both the correlated Erdős-Rényi model and correlated stochastic block models (SBMs). Through the collective efforts of the community, the information-theoretic thresholds for detection and matching have been fully characterized for correlated Erdős-Rényi model and partially characterized for correlated SBMs [10, 29, 54, 55, 25, 13, 14, 47, 28]. Additionally, various efficient detecting and matching algorithms have been developed with performance guarantees [11, 4, 18, 20, 21, 24, 26, 27, 39, 40, 42, 41, 16, 17, 8, 9]. Notably, like many other inference tasks in high-dimensional statistics [57, 48, 37, 22], these problems appear to exhibit information-computation gaps. Specifically, for certain ranges of the correlation strength, detection/matching is information theoretically possible but no efficient algorithm is known to achieve these tasks. We now focus on the algorithmic side of these problems as they are more relevant to our work. Indeed, it has been shown that many inference tasks in the correlated random graph models exhibits sharp algorithmic phase transitions, as we elaborate below:

• For the detection problem between a pair of correlated Erdős-Rényi models $\mathcal{G}(n,q;\rho)$ and two independent Erdős-Rényi models $\mathcal{G}(n,q)$, we focus on the sparse regime where $q = n^{-1+o(1)}$. In this regime, on the one hand, it was shown in [42] that when $\rho > \sqrt{\alpha}$ where $\alpha \approx 0.338$ is the Otter's constant [46], there is an efficient algorithm that strongly distinguish these two models; on the other hand, it was shown in [15] that when $s < \sqrt{\alpha}$ there are evidences suggesting that all algorithms based on *low-degree polynomials* fail to strongly distinguish these two models.

• For the detection problem between a pair of correlated SBMs $S(n, \frac{\lambda}{n}; k, \epsilon; s)$ and two independent Erdős-Rényi graphs $\mathcal{G}(n, \frac{\lambda s}{n})$, we focus on the constant degree regime where $\lambda = O(1)$. In this regime, on the one hand, it was shown in [9] that when $s > \min\{\sqrt{\alpha}, \frac{1}{\epsilon^2 \lambda}\}$ where $\alpha \approx 0.338$ is the Otter's constant and $\frac{1}{\epsilon^2 \lambda}$ is the Kesten-Stigum threshold [33], there is an efficient algorithm that strongly distinguish these two models; on the other hand, it was also shown in [9] that when $s < \min\{\sqrt{\alpha}, \frac{1}{\epsilon^2 \lambda}\}$ there are evidences suggesting that all algorithms based on *low-degree polynomials* fail to strongly distinguish these two models.

The lower bound in the aforementioned results explored inherent computational barriers from the perspective of the *low-degree polynomial framework*. Indeed, it has been proved that the class of low-degree polynomial algorithms is a useful proxy for computationally efficient algorithms, in the sense that the best-known polynomial-time algorithms for a wide variety of high-dimensional inference problems are captured by the low-degree class; see e.g. [31, 30, 50, 37]. However, these aforementioned results suffer from two significant limitations, which we now discuss. Firstly, the aforementioned result cannot provide the evidence that *partial matching recovery* (that is, recover a positive fraction of the coordinates of the latent matching π_*) is impossible by efficient algorithms although efficient detection is already ruled out. This limitation arises because there is no *evident* reduction from partial matching to detection. Secondly, (in the case of correlated SBMs) they are only able to establish the computation threshold on the detection problem between the correlated model and a pair of independent Erdős-Rényi graphs. These limitations motivate two natural questions:

- **Question 1.5.** (1) Can we provide the evidences that partial matching recovery is impossible (by efficient algorithms) in the same parameter regime where detection is impossible (by efficient algorithms)?
 - (2) What can we say about the (arguably more natural) detection problem between a pair of correlated SBMs $S(n, \frac{\lambda}{n}; k, \epsilon; s)$ and a pair of independent SBMs $S(n, \frac{\lambda s}{n}; k, \epsilon)$?

The aim of this paper is to find (partial) answers for these two problems. Our main result can be informally summarized as follows:

Theorem 1.6 (Informal). Assuming a natural strengthening of the low-degree conjecture (see Conjecture 2.7 for its precise meaning), we have

(1) For the correlated Erdős-Rényi model $\mathcal{G}(n,q,\rho)$, when $q = n^{-1+o(1)}$ and $\rho < \sqrt{\alpha}$ it is impossible to recover a positive fraction of the coordinates of π_* with positive probability by efficient algorithms.

- (2) For the correlated stochastic block models $S(n, \frac{\lambda}{n}; k, \epsilon; s)$, when $\lambda = O(1)$ and $s < \min\{\sqrt{\alpha}, \frac{1}{\epsilon^2 \lambda}\}$ it is impossible to recover a positive fraction of the coordinates of π_* with probability tending to 1 as $n \to \infty$ by efficient algorithms.
- (3) For the correlated stochastic block models $S(n, \frac{\lambda}{n}; k, \epsilon; s)$, when $s < \min\{\sqrt{\alpha}, \frac{1}{\epsilon^2 \lambda}\}$ it is impossible to strongly distinguish this model and a pair of independent SBMs $S(n, \frac{\lambda s}{n}; k, \epsilon)$, provided that the average degree λs is sufficiently large.

Remark 1.7. Note that when $\rho > \sqrt{\alpha}$ (respectively, $s > \sqrt{\alpha}$), the results in [41, 24, 26, 27] shows (or naturally extends to show) that there exists an efficient algorithm that achieves partial recovery of π_* in a pair of correlated Erdős-Rényi models (respectively, correlated SBMs) with probability tending to 1 as $n \to \infty$. Thus, Items (1) and (2) in Theorem 1.6 is tight in some sense and the algorithmic partial recovery threshold is indeed given by $\sqrt{\alpha}$.

Remark 1.8. From Item (3) in Theorem 1.6 we see that for the correlated SBMs (A, B), when marginally both A and B are below the KS-threshold, there is no efficient algorithm that strongly distinguish (A, B) from a pair of independent stochastic block models when the correlation $s < \sqrt{\alpha}$. Since the result in [42] extends naturally to the case of stochastic block models which provides an efficient algorithm that strongly distinguish these two models when $s > \sqrt{\alpha}$, we see that in the subcritical regime (i.e., when marginally both A and B are below the KS-threshold) the algorithmic correlation detection threshold is given by $\sqrt{\alpha}$. On the contrary, in the supercritical regime where A and B are above the KS-threshold, we believe that the correlation detection threshold should be strict lower than $\sqrt{\alpha}$.

1.1 Key challenges and innovations

In this subsection, we first briefly discuss our approach of showing Theorem 1.6 and some conceptual innovations behind it. Our idea can be summarized as follows:

- (1) As for Items (1) and (2), for simplicity we take correlated Erdős-Rényi model for example. Denote \mathbb{P} to be the law of $\mathcal{G}(n, q, \rho)$ and \mathbb{Q} to be the law of two independent $\mathcal{G}(n, q)$. We will argue by contradiction and assume that there is a partial recovery algorithm. Then we show that we can use this algorithm to efficiently construct a family of statistics $\{g_i : 1 \leq i \leq n\}$ such that g_i approximates $\mathbf{1}_{\{\pi_*(1)=i\}}$ under \mathbb{P} in a certain sense.
- (2) We will show that the low-degree advantage between $\mathbb{P}(\cdot \mid \pi_*(1) = i)$ and \mathbb{Q} is bounded by an absolute constant. Then, from the standard low-degree conjecture (see Conjecture 2.2 for details) these two measures cannot be strongly distinguished by efficient algorithms. Thus, since g_i is "not small" under the measure $\mathbb{P}(\cdot \mid \pi_*(i) = 1)$ (as it should approximate $\mathbf{1}_{\{\pi_*(1)=i\}}$ in some sense) we expect that g_i should also be "not small" under \mathbb{Q} .

- (3) Define $g = g_1 + \ldots + g_n$. We then have a statistics g that can be efficiently calculated, and (I) under \mathbb{Q} we expect that g is "large" as it is the sum of n "not small" terms; (II) under \mathbb{P} we expect that g is "small" as each g_i should approximate $\mathbf{1}_{\{\pi_*(1)=i\}}$ in some sense. Thus, the statistics g accumulates more signals than all low-degree polynomials, which violates the low-degree conjecture.
- (4) As for Item (3), denote P to be the law of correlated SBMs and Q of independent SBMs. Also denote Q to be the law of independent Erdős-Rényi graphs. As it was already shown in [9] that P and Q cannot be strongly distinguished, (non-rigorously speaking) we only need to show that Q and Q are also "indistinguishable" in some sense.

However, there are certain obstacles when implementing the above ideas, as we shall discuss below. In Step (2) we need to "transfer" the behavior of a statistics g_i under $\mathbb{P}(\cdot \mid \pi_*(1) = i)$ to its behavior under \mathbb{Q} . If the low-degree advantage between $\mathbb{P}(\cdot \mid \pi_*(1) = i)$ and \mathbb{Q} is given by 1 + o(1), then the standard low-degree conjecture implies that we cannot distinguish between $\mathbb{P}(\cdot \mid \pi_*(1) = i)$ and \mathbb{Q} better than random by efficient algorithms. Consequently, we expect the behavior of g_i to be "almost identical" under both $\mathbb{P}(\cdot \mid \pi_*(1) = i)$ and \mathbb{Q} . However, in our case the low-degree advantage is just bounded by a large (but fixed) constant. This weaker condition means that the standard low-degree conjecture only rules out the possibility to distinguish these two measures with vanishing errors, leaving room for non-negligible distinctions. Similar issues also arise in Step (4), in which simply showing the low-degree advantage between \mathbb{P} and \mathbb{Q} and the low-degree advantage between \mathbb{Q} and $\widetilde{\mathbb{Q}}$ is bounded is not strong enough for our goal.

To address these issues, one of the main conceptual contributions in our work is to give a more refined characterization of the limitations of efficient algorithms when the low-degree advantage is O(1). Specifically, we show that (assuming low-degree conjecture) bounded low-degree advantage does not only excludes all algorithms that strongly distinguish \mathbb{P} and \mathbb{Q} , but also suggest a certain kind of *algorithmic contiguity*. To be more precise, if the low-degree advantage $\operatorname{Adv}_{\leq D}\left(\frac{\mathrm{dP}}{\mathrm{dQ}}\right)$ is bounded, then there is no efficient algorithm \mathcal{A} such that $\mathbb{Q}(\mathcal{A}=0) = 1 - o(1)$ and $\mathbb{P}(\mathcal{A}=1) = \Omega(1)$. This framework allows us to transfer the behavior of the behavior of a efficiently computable statistics g under different probability measures more easily. For example, if we know that $\mathbb{P}(g \geq c) \geq \Omega(1)$ it immediately holds that $\mathbb{Q}(g \geq c) \geq \Omega(1)$.

Another difficulty is that reducing between different inference tasks requires a framework that better constrains the behavior of efficiently computable estimators. To address this, we introduce a natural refinement of the low-degree conjecture: we posit that lowdegree polynomials achieve signal-to-noise ratios that are at least as good as those of any efficient algorithm (see Conjecture 2.7 for a more precise description). While similar modifications to the low-degree conjecture have been explored in [44], our motivations differ significantly. Their work focused on characterizing the limits of precise error for all computationally feasible algorithms, whereas our goal is to enable reductions between distinct inference tasks.

1.2 Notation

In this subsection, we record a list of notations that we shall use throughout the paper. Denote \mathfrak{S}_n the set of permutations over [n] and denote μ the uniform distribution on \mathfrak{S}_n . In addition, denote ν to be the uniform distribution on $[k]^n$. We will use the following notation conventions on graphs.

- Labeled graphs. Denote by \mathcal{K}_n the complete graph with vertex set [n] and edge set U_n . For any graph H, let V(H) denote the vertex set of H and let E(H) denote the edge set of H. We say H is a subgraph of G, denoted by $H \subset G$, if $V(H) \subset V(G)$ and $E(H) \subset E(G)$. Define the excess of the graph $\tau(H) = |E(H)| |V(H)|$.
- Isolated vertices. For $u \in V(H)$, we say u is an isolated vertex of H, if there is no edge in E(H) incident to u. Denote $\mathcal{I}(H)$ the set of isolated vertices of H. For two graphs H, S, we denote $H \ltimes S$ if $H \subset S$ and $\mathcal{I}(S) \subset \mathcal{I}(H)$, and we denote $H \Subset S$ if $H \subset S$ and $\mathcal{I}(H) = \emptyset$. For any graph $H \subset \mathcal{K}_n$, let \widetilde{H} be the subgraph of H induced by $V(H) \setminus \mathcal{I}(H)$.
- Graph intersections and unions. For $H, S \subset \mathcal{K}_n$, denote by $H \cap S$ the graph with vertex set given by $V(H) \cap V(S)$ and edge set given by $E(H) \cap E(S)$. Denote by $S \cup H$ the graph with vertex set given by $V(H) \cup V(S)$ and edge set $E(H) \cup E(S)$. In addition, denote by $S \cap H, S \setminus H$ and $S \triangle H$ the graph induced by the edge set $E(S) \cap E(H), E(S) \setminus E(H)$ and $E(S) \triangle E(H)$, respectively (in particular, these induced graphs have no isolated points).
- Paths. We say a subgraph $H \subset \mathcal{K}_n$ is a path with endpoints u, v (possibly with u = v), if there exist distinct $w_1, \ldots, w_m \neq u, v$ such that $V(H) = \{u, v, w_1, \ldots, w_m\}$ and $E(H) = \{(u, w_1), (w_1, w_2) \ldots, (w_m, v)\}$. We say H is a simple path if its endpoints $u \neq v$. Denote EndP(P) as the set of endpoints of a path P.
- Cycles and independent cycles. We say a subgraph H is an m-cycle if $V(H) = \{v_1, \ldots, v_m\}$ and $E(H) = \{(v_1, v_2), \ldots, (v_{m-1}, v_m), (v_m, v_1)\}$. For a subgraph $K \subset H$, we say K is an independent m-cycle of H, if K is an m-cycle and no edge in $E(H) \setminus E(K)$ is incident to V(K). Denote by $C_m(H)$ the set of m-cycles of H and denote by $C_m(H)$ the set of independent m-cycles of H. For $H \subset S$, we define $\mathfrak{C}_m(S, H)$ to be the set of independent m-cycles in S whose vertex set is disjoint from V(H). Define $\mathfrak{C}(S, H) = \bigcup_{m \geq 3} \mathfrak{C}_m(S, H)$.
- Leaves. A vertex $u \in V(H)$ is called a leaf of H, if the degree of u in H is 1; denote $\mathcal{L}(H)$ as the set of leaves of H.

• Graph isomorphisms and unlabeled graphs. Two graphs H and H' are isomorphic, denoted by $H \cong H'$, if there exists a bijection $\pi : V(H) \to V(H')$ such that $(\pi(u), \pi(v)) \in E(H')$ if and only if $(u, v) \in E(H)$. Denote by \mathcal{H} the isomorphism class of graphs; it is customary to refer to these isomorphic classes as unlabeled graphs. Let Aut(H) be the number of automorphisms of H (graph isomorphisms to itself).

For two real numbers a and b, we let $a \lor b = \max\{a, b\}$ and $a \land b = \min\{a, b\}$. We use standard asymptotic notations: for two sequences a_n and b_n of positive numbers, we write $a_n = O(b_n)$, if $a_n < Cb_n$ for an absolute constant C and for all n (similarly we use the notation O_h is the constant C is not absolute but depends only on h); we write $a_n = \Omega(b_n)$, if $b_n = O(a_n)$; we write $a_n = \Theta(b_n)$, if $a_n = O(b_n)$ and $a_n = \Omega(b_n)$; we write $a_n = o(b_n)$ or $b_n = \omega(a_n)$, if $a_n/b_n \to 0$ as $n \to \infty$. In addition, we write $a_n \triangleq b_n$ if $a_n = [1 + o(1)]b_n$. For a set A, we will use both #A and |A| to denote its cardinality. For two probability measures \mathbb{P} and \mathbb{Q} , we denote the total variation distance between them by $\mathrm{TV}(\mathbb{P}, \mathbb{Q})$.

1.3 Organization of this paper

The rest part of this paper is organized as follows: in Section 2 we state the precise framework of low-degree polynomials and how we relate it to the notion of algorithmic contiguity (see Theorem 2.4). In Section 3 we use this framework to deduce the hardness of partial matching in correlated Erdős-Rényi models and correlated SBMs, thus verifying Items (1) and (2) in Theorem 1.6 (see Theorem 3.2). In Section 4 we again use this framework to deduce the hardness of testing correlated SBMs against independent SBMs, thus verifying Item (3) in Theorem 1.6 (see Corollary 4.2). Several auxiliary results are moved to the appendix to ensure a smooth flow of presentation.

2 Low-degree framework and algorithmic contiguity

Initially emerged from the works of [5, 32, 31, 30], the low-degree polynomial framework was later refined and extended in various directions. It has found applications in a wide range of problems, including detection problems such as planted clique, planted dense subgraph, community detection, sparse-PCA, correlated random graphs (see [32, 31, 30, 37, 50, 12, 2, 19, 43, 15, 36]), optimization problems such as maximal independent sets in sparse random graphs [23, 53], and constraint satisfaction problems such as random k-SAT [6] (see also the survey [37]). Furthermore, it is conjectured in [30] that the failure of degree-D polynomials implies the failure of all "robust" algorithms with running time $n^{\tilde{O}(D)}$ (here \tilde{O} means having at most this order up to a poly log n factor). In the remaining of this paper, we will focus on applying this framework in the context of high-dimensional hypothesis testing problems.

To be more precise, consider the hypothesis testing problem between two probability measures \mathbb{P} and \mathbb{Q} based on the sample $\mathsf{Y} \in \mathbb{R}^N$. We will be especially interested in

asymptotic settings where $N = N_n$, $\mathbb{Q} = \mathbb{Q}_n$, $\mathbb{P} = \mathbb{P}_n$, $Y = Y_n$ scale with n as $n \to \infty$ in some prescribed way. The standard low-degree polynomial framework primarily focus on the following notions on strong and weak detection.

Definition 2.1 (Strong/weak detection). We say an algorithm \mathcal{A} that takes Y as input and outputs either 0 or 1 achieves

- strong detection, if the sum of type-I and type-II errors Q(A(Y) = 1)+P(A(Y) = 0) tends to 0 as n → ∞.
- weak detection, if the sum of type-I and type-II errors is uniformly bounded above by 1 − ε for some fixed ε > 0.

Roughly speaking, the approach of low-degree polynomials amounts to studying the capabilities and limitations of algorithms that can be represented as low-degree polynomial functions of the input variables (in our case, the entries of Y). More precisely, denote $\mathcal{P}_D = \mathcal{P}_{n,D}$ for the set of polynomials from \mathbb{R}^N to \mathbb{R} with degree no more than D. With a slight abuse of notation, we will often say "a polynomial" to mean a sequence of polynomials $f = f_n \in \mathcal{P}_{n,D}$, one for each problem size n; the degree $D = D_n$ of such a polynomial may scale with n. The polynomial degree is thought of as a measure of the algorithm's complexity and a proxy for runtime, with degree D corresponding to runtime roughly n^D (up to log n factors in the exponent), which is the number of terms in such a polynomial. As suggested by [30], the key quantity is the low-degree advantage

$$\mathsf{Adv}_{\leq D}\left(\frac{\mathrm{d}\mathbb{P}}{\mathrm{d}\mathbb{Q}}\right) := \sup_{f\in\mathcal{P}_D} \frac{\mathbb{E}_{\mathbb{P}}[f]}{\sqrt{\mathbb{E}_{\mathbb{Q}}[f^2]}}.$$
(2.1)

The low degree conjecture, proposed in [30], can be summarized as follows.

Conjecture 2.2 (Low-degree conjecture). For "natural" high-dimensional hypothesis testing problems between \mathbb{P} and \mathbb{Q} , the following statements are equivalent:

- (1) $\operatorname{Adv}_{\leq D}\left(\frac{\mathrm{d}\mathbb{P}}{\mathrm{d}\mathbb{Q}}\right) = O(1)$ (respectively, $\operatorname{Adv}_{\leq D}\left(\frac{\mathrm{d}\mathbb{P}}{\mathrm{d}\mathbb{Q}}\right) = 1 + o(1)$) as $n \to \infty$.
- (2) There is no algorithm with running time $n^{\widetilde{O}(D)}$ that achieves strong detection (respectively, weak detection) between \mathbb{P} and \mathbb{Q} .

This informal conjecture is inspired by [30, Hypothesis 2.1.5 and Conjecture 2.2.4], as well as the fact that low-degree polynomials capture the best known algorithms for a wide variety of statistical inference tasks. The conjecture appears to hold up for distributions \mathbb{P}, \mathbb{Q} of a particular style that often arises in high-dimensional statistics, and we refer the readers to [30, 37, 35, 56] for further discussion on which distributions \mathbb{P}, \mathbb{Q} are appropriate for this framework. In addition, we point out that although in most applications (and in the statement of [30, Hypothesis 2.1.5]) we will usually take \mathbb{Q} to be a "null" measure and \mathbb{P} to be a "planted" measure (which makes (2.1) more tractable), several recent works [49, 34] showed that this framework might also be applicable for many "planted-versus-planted" problems. Nevertheless, we will be more conservative and will explicitly indicate in this paper whenever \mathbb{Q} is treated as a planted measure.

The framework in Conjecture 2.2 provides a useful tool for probing the computational feasibility of strong or weak detection. However, as discussed in Subsection 1.1, it turns out that the failure of strong detection is not enough in our cases, especially when we hope to perform some reductions between statistical models in a regime where weak (but not strong) detection is possible. Thus, in this regime, we aim to characterize a stronger framework that rules out all *one-sided test*. This motivates thus to propose the following notion of *algorithmic contiguity*, which is defined as follows.

Definition 2.3. We say an algorithm \mathcal{A} that takes Y as input and outputs either 0 or 1 is a \mathbb{Q} -based one-sided test, if

$$\mathbb{Q}(\mathcal{A}(\mathsf{Y}) = 0) = 1 - o(1) \text{ and } \mathbb{P}(\mathcal{A}(\mathsf{Y}) = 1) = \Omega(1).$$
(2.2)

We say that \mathbb{Q} is algorithmic contiguous with respect to \mathbb{P} , denoted as $\mathbb{Q} \triangleleft_{\leq D} \mathbb{P}$, if no \mathbb{Q} based one-sided testing algorithm runs in time $n^{\tilde{O}(D)}$. We say that \mathbb{Q} and \mathbb{P} are algorithmic mutually contiguous, denoted as $\mathbb{Q} \bowtie_{\leq D} \mathbb{P}$, if both $\mathbb{Q} \triangleleft_{\leq D} \mathbb{P}$ and $\mathbb{P} \triangleleft_{\leq D} \mathbb{Q}$ hold.

Recall that in probability theory we say a probability measure $\mathbb{Q} = \mathbb{Q}_n$ is contiguous with respect to $\mathbb{P} = \mathbb{P}_n$, if for all sequence of events $\{A_n\}$ we have $\mathbb{P}(A_n) \to 0$ implies that $\mathbb{Q}(A_n) \to 0$. Thus, our definition can be regarded as the generalization of contiguity in algorithmic view. Our main result in this section can be stated as follows.

Theorem 2.4. Assuming the Conjecture 2.2, for the high-dimensional hypothesis testing problem between \mathbb{P} and \mathbb{Q} , if $\mathsf{Adv}_{\leq D}\left(\frac{\mathrm{d}\mathbb{P}'}{\mathrm{d}\mathbb{Q}'}\right) = O(1)$ for some \mathbb{P}', \mathbb{Q}' such that $\mathrm{TV}(\mathbb{P}, \mathbb{P}') = o(1)$ and $\mathrm{TV}(\mathbb{Q}, \mathbb{Q}') = o(1)$, then we have $\mathbb{Q} \triangleleft_{\leq D} \mathbb{P}$.

The main part of this section is devoted to the proof of Theorem 2.4. We first briefly explain our proof ideas. Suppose on the contrary that there is an efficient algorithm \mathcal{A} that takes Y as input and outputs either 0 or 1 with

$$\mathbb{P}(\mathcal{A}(\mathsf{Y}) = 1) = \Omega(1) \text{ and } \mathbb{Q}(\mathcal{A}(\mathsf{Y}) = 0) = 1 - o(1).$$
(2.3)

Since $TV(\mathbb{Q}, \mathbb{Q}'), TV(\mathbb{P}, \mathbb{P}') = o(1)$, we also have

$$\mathbb{P}'(\mathcal{A}(\mathsf{Y})=1) = \Omega(1) \text{ and } \mathbb{Q}'(\mathcal{A}(\mathsf{Y})=0) = 1 - \epsilon \text{ for some } \epsilon = \epsilon_n \to 0.$$
 (2.4)

Define

$$M = M_n = \epsilon_n^{-\frac{1}{2}} \,. \tag{2.5}$$

The crux of our argument is to consider the following *hidden informative sample* problem.

Definition 2.5. Consider the following hypothesis testing problem: we need to determine whether a sample $(Y_1, Y_2, ..., Y_M)$ where each $Y_i \in \mathbb{R}^N$ is generated by

- $\overline{\mathcal{H}}_0$: we let Y_1, \ldots, Y_M to be independently sampled from \mathbb{Q}' .
- $\overline{\mathcal{H}}_1$: we first sample $\kappa \in \{1, \ldots, M\}$ uniformly at random, and (conditioned on the value of κ) we let Y_1, \ldots, Y_M are independent samples with Y_{κ} generated from \mathbb{P}' and $\{Y_j : j \neq \kappa\}$ generated from \mathbb{Q}' .

In addition, denote $\overline{\mathbb{P}}$ and $\overline{\mathbb{Q}}$ to be the law of $(\mathsf{Y}_1, \ldots, \mathsf{Y}_M)$ under $\overline{\mathcal{H}}_1$ and $\overline{\mathcal{H}}_0$, respectively.

Now assuming that (2.4) holds, we see that

$$\overline{\mathbb{Q}}\Big(\big(\mathcal{A}(\mathsf{Y}_1),\ldots,\mathcal{A}(\mathsf{Y}_M)\big) = (0,\ldots,0)\Big) \ge 1 - M\epsilon \stackrel{(2.5)}{=} 1 - o(1); \qquad (2.6)$$

$$\overline{\mathbb{P}}\Big(\big(\mathcal{A}(\mathsf{Y}_1),\ldots,\mathcal{A}(\mathsf{Y}_M)\big)\neq(0,\ldots,0)\Big)\geq\Omega(1)\,.$$
(2.7)

Thus, there is an efficient algorithm that achieves weak detection between $\overline{\mathbb{P}}$ and $\overline{\mathbb{Q}}$. Our next result, however, shows the low-degree advantage $\operatorname{Adv}_{\leq D}\left(\frac{\mathrm{d}\overline{\mathbb{P}}}{\mathrm{d}\overline{\mathbb{Q}}}\right)$ is bounde by 1 + o(1).

Lemma 2.6. If $\operatorname{Adv}_{\leq D}\left(\frac{\mathrm{d}\mathbb{P}'}{\mathrm{d}\mathbb{Q}'}\right) = O(1)$, then $\operatorname{Adv}_{\leq D}\left(\frac{\mathrm{d}\mathbb{P}}{\mathrm{d}\mathbb{Q}}\right) = 1 + o(1)$.

Proof. Note that

$$\frac{\mathrm{d}\overline{\mathbb{P}}}{\mathrm{d}\overline{\mathbb{Q}}}(\mathsf{Y}_1,\ldots,\mathsf{Y}_M) = \frac{1}{M}\sum_{i=1}^M \frac{\mathrm{d}\overline{\mathbb{P}}(\cdot \mid \kappa = i)}{\mathrm{d}\overline{\mathbb{Q}}}(\mathsf{Y}_1,\ldots,\mathsf{Y}_M) = \frac{1}{M}\sum_{i=1}^M \frac{\mathrm{d}\mathbb{P}'}{\mathrm{d}\mathbb{Q}'}(\mathsf{Y}_i).$$
(2.8)

In addition, recall (2.1). If we identify \mathcal{P}_D as a (finite-dimensional) Hilbert space with the inner product $\langle f,g \rangle = \mathbb{E}_{\mathbb{Q}'}[fg]$, then using Schmidt orthogonalization procedure we can find a standard orthogonal basis $\{f_\alpha : \alpha \in \Lambda\}$ of this Hilbert space. In particular, we can choose Λ such that $0 \in \Lambda$ and $f_0 = 1$. Note that in some cases (e.g., when \mathbb{Q}' is a product measure), this standard orthogonal basis have a closed form; however, for general \mathbb{Q}' the explicit form of $\{f_\alpha\}$ is often intractable. We first show that

$$\mathsf{Adv}_{\leq D}\left(\frac{\mathrm{d}\mathbb{P}'}{\mathrm{d}\mathbb{Q}'}\right) = \left(\sum_{\alpha \in \Lambda} \mathbb{E}_{\mathbb{P}'}[f_{\alpha}(\mathsf{Y})]^2\right)^{\frac{1}{2}}.$$
(2.9)

Indeed, for any $f \in \mathcal{P}_D$, it can be uniquely expressed as

$$f = \sum_{\alpha \in \Lambda} C_{\alpha} f_{\alpha} \,,$$

where C_{α} 's are real constants. Applying Cauchy-Schwartz inequality one gets

$$\frac{\mathbb{E}_{\mathbb{P}'}[f]}{\sqrt{\mathbb{E}_{\mathbb{Q}'}[f^2]}} = \frac{\sum_{\alpha \in \Lambda} C_{\alpha} \mathbb{E}_{\mathbb{P}'}[f_{\alpha}(\mathsf{Y})]}{\sqrt{\sum_{\alpha \in \Lambda} C_{\alpha}^2}} \le \left(\sum_{\alpha \in \Lambda} \mathbb{E}_{\mathbb{P}'}[f_{\alpha}(\mathsf{Y})]^2\right)^{1/2},$$

with equality holds if and only if $C_{\alpha} \propto \mathbb{E}_{\mathbb{P}'}[f_{\alpha}]$. This yields (2.9). Now, note that $\overline{\mathbb{Q}} = (\mathbb{Q}')^{\otimes M}$ is a product measure of \mathbb{Q}' , there is a natural standard orthogonal basis under $\overline{\mathbb{Q}}$, given by

$$\left\{\prod_{i=1}^{M} f_{\alpha_i}(\mathsf{Y}_i) : \alpha_i \in \Lambda, \sum_{i=1}^{M} \deg(f_{\alpha_i}) \le D\right\}.$$

Thus, similarly as in (2.9), we see that

$$\left(\mathsf{Adv}_{\leq D}\left(\frac{\mathrm{d}\overline{\mathbb{P}}}{\mathrm{d}\overline{\mathbb{Q}}}\right)\right)^{2} = \sum_{\substack{(\alpha_{1},\dots,\alpha_{M}):\alpha_{i}\in\Lambda\\\sum_{i=1}^{M}\deg(f_{\alpha_{i}})\leq D}} \mathbb{E}_{\overline{\mathbb{P}}}\left[\prod_{i=1}^{M}f_{\alpha_{i}}(\mathsf{Y}_{i})\right]^{2}.$$
(2.10)

In addition, using (2.8), we see from direct calculation that

$$\mathbb{E}_{\overline{\mathbb{P}}} \Big[\prod_{i=1}^{M} f_{\alpha_{i}}(\mathsf{Y}_{i}) \Big] = \mathbb{E}_{\overline{\mathbb{Q}}} \Big[\prod_{i=1}^{M} f_{\alpha_{i}}(\mathsf{Y}_{i}) \cdot \frac{\mathrm{d}\overline{\mathbb{P}}}{\mathrm{d}\overline{\mathbb{Q}}}(\mathsf{Y}_{1}, \dots, \mathsf{Y}_{M}) \Big]$$

$$\stackrel{(2.8)}{=} \frac{1}{M} \sum_{i=1}^{M} \mathbb{E}_{\overline{\mathbb{Q}}} \Big[\prod_{i=1}^{M} f_{\alpha_{i}}(\mathsf{Y}_{i}) \cdot \frac{\mathrm{d}\mathbb{P}'}{\mathrm{d}\mathbb{Q}'}(\mathsf{Y}_{i}) \Big]$$

$$= \begin{cases} 1, & (\alpha_{1}, \dots, \alpha_{M}) = (0, \dots, 0); \\ \frac{1}{M} \mathbb{E}_{\mathbb{P}'} \big[f_{\alpha_{j}}(\mathsf{Y}_{j}) \big], & (\alpha_{1}, \dots, \alpha_{M}) = (0, \dots, 0, \alpha_{j}, 0, \dots, 0); \\ 0, & \text{otherwise}. \end{cases}$$

$$(2.11)$$

Plugging (2.11) into (2.10), we get that

$$(2.10) = 1 + \sum_{i=1}^{M} \sum_{\alpha_i \in \Lambda \setminus \{0\}} \left(\frac{1}{M} \mathbb{E}_{\mathbb{P}'} \left[f_{\alpha_i}(\mathsf{Y}_i) \right] \right)^2$$

$$\leq 1 + \frac{1}{M} \sum_{\alpha \in \Lambda} \mathbb{E}_{\mathbb{P}'} \left[f_{\alpha}(\mathsf{Y}_j) \right]^2 = 1 + \frac{1}{M} \cdot O(1) \stackrel{(2.5)}{=} 1 + o(1) ,$$

where in the second equality we use (2.9) and the assumption that $\operatorname{Adv}_{\leq D}\left(\frac{d\mathbb{P}'}{d\mathbb{Q}'}\right) = O(1)$. This completes our proof.

We can now finish the proof of Theorem 2.4.

Proof of Theorem 2.4. Suppose on the contrary that there is an efficient algorithm \mathcal{A} satisfying (2.3). Consider the hypothesis testing problem stated in Definition 2.5. Using (2.6) and (2.7), we see that there is an efficient algorithm that achieves weak detection between $\overline{\mathbb{P}}$ and $\overline{\mathbb{Q}}$, which contradicts with Lemma 2.6 and the equivalence between Items (1) and (2) in Conjecture 2.2.

In the remaining part of this work we will also need a strengthening of Conjecture 2.2, as introduced in Subsection 1.1. As discussed in the beginning of this section, the low-degree conjecture asserts that (for certain testing problems) low-degree polynomials are at least as powerful as all algorithms of the corresponding runtime (where the correspondence is described in Conjecture 2.2). For our purposes, we introduce a natural refinement of the low-degree conjecture: we posit that low-degree polynomials perform at least as well as all algorithms of the corresponding runtime in terms of the value of the ratio on the right-hand side of (2.1).

Conjecture 2.7 (Revised low-degree conjecture). For "natural" high-dimensional hypothesis testing problems between \mathbb{P} and \mathbb{Q} , the following statements are equivalent:

- (1) $\operatorname{Adv}_{\leq D}\left(\frac{\mathrm{d}\mathbb{P}'}{\mathrm{d}\mathbb{Q}'}\right) = O(1)$ (respectively, $\operatorname{Adv}_{\leq D}\left(\frac{\mathrm{d}\mathbb{P}'}{\mathrm{d}\mathbb{Q}'}\right) = 1 + o(1)$) as $n \to \infty$ for some \mathbb{P}', \mathbb{Q}' such that $\operatorname{TV}(\mathbb{P}, \mathbb{P}'), \operatorname{TV}(\mathbb{Q}, \mathbb{Q}') = o(1)$.
- (2) There is no algorithm with running time $n^{\widetilde{\Theta}(D)}$ that achieves strong detection (respectively, weak detection) between \mathbb{P} and \mathbb{Q} .
- (3) For all statistics f = f(Y) that can be computed in running time $n^{\widetilde{\Theta}(d)}$, there exists some \mathbb{P}', \mathbb{Q}' such that $\mathrm{TV}(\mathbb{P}, \mathbb{P}'), \mathrm{TV}(\mathbb{Q}, \mathbb{Q}') = o(1)$ such that $\mathsf{Adv}(f) = O(1)$ (respectively, $\mathsf{Adv}(f) = 1 + o(1)$), where

$$\mathsf{Adv}(f) = \frac{\mathbb{E}_{\mathbb{P}'}[f]}{\sqrt{\mathbb{E}_{\mathbb{Q}'}[f^2]}} = O(1)$$

Note that in Conjecture 2.7 we are allowed to replace \mathbb{P}, \mathbb{Q} with some \mathbb{P}', \mathbb{Q}' that are statistically indistinguishable with \mathbb{P}, \mathbb{Q} . This modification enables us to avoid some situations where the low-degree advantage explodes due to some "rare events" (see [3, 15, 12, 9] for example). This revised low-degree conjecture was first proposed in [44] (in a slightly different manner) for a different purpose where they aimed to study the limits of precise error of all computationally feasible algorithms. However, we will show in the next two sections that this conjecture is also useful in performing reductions between different inference tasks.

3 Partial recovery in correlated random graphs

In this section, we will use the framework we established in Section 2 to show the hardness of partial matching in correlated random graphs, thus justifying Items (1) and (2) in Theorem 1.6. To this end, we first state the precise meaning of an algorithm achieves partial matching.

Definition 3.1 (Partial recovery algorithm in correlated Erdős-Rényi model). Given a sample (A, B) from the law a pair of correlated random graphs in Definition 1.1 (we denote this law of \mathbb{P}_*). We say an algorithm \mathcal{A} achieves strong partial matching, if it takes (A, B) as input and outputs a family of estimators $\{h_{i,j} : 1 \leq i, j \leq n\}$ such that

- (1) $h_{i,j} \in \{0,1\}$ for all $1 \le i, j \le n$ a.s. under \mathbb{P}_* ;
- (2) $h_{i,1} + \ldots + h_{i,n} = 1$ for all $1 \le i \le n$ a.s. under \mathbb{P}_* ;
- (3) There exists a fixed constant $\iota > 0$ such that

$$\mathbb{P}_*(h_{1,\pi_*(1)} + \ldots + h_{n,\pi_*(n)} \ge \iota n) = 1 - o(1).$$

We say an algorithm \mathcal{A} achieves weak partial matching, if it takes (A, B) as input and outputs a family of estimators f_1, \ldots, f_n satisfying Items (1), (2) above and

$$\mathbb{P}_*(h_{1,\pi_*(1)} + \ldots + h_{n,\pi_*(n)} \ge \iota n) = \Omega(1).$$

We point out that in the above definition $h_{i,j}$ can be thought as the estimator of $\mathbf{1}_{\{\pi_*(i)=j\}}$, where Item (3) implies this algorithm correctly matches a positive fraction of $\pi_*(i)$ with probability 1-(1) (or with positive probability) and Items (1) and (2) are some regularity requirements. Our result in this section can be stated as follows:

Theorem 3.2. Assuming Conjecture 2.7, we have the following:

(1) For the correlated Erdős-Rényi graphs $\mathcal{G}(n,q,\rho)$ where $q = n^{-1+o(1)}$ and $\rho < \sqrt{\alpha} - \delta$ for a fixed constant $\delta > 0$. There is no algorithm with running time $n^{\tilde{O}(D)}$ that achieves weak partial matching, provided that

$$D = \exp\left(o\left(\frac{\log n}{\log(nq)} \land \sqrt{\log n}\right)\right). \tag{3.1}$$

(2) For the correlated SBMs $S(n, \frac{\lambda}{n}; k, \epsilon; s)$ where $\lambda = O(1)$ and $\epsilon^2 \lambda < 1 - \delta, s < \sqrt{\alpha} - \delta$ for a fixed constant $\delta > 0$. There is no algorithm with running time $n^{\tilde{O}(D)}$ that achieves strong partial matching, provided that

$$D = O(\log n). \tag{3.2}$$

The main step of proving Theorem 3.2 is to show the following proposition:

Proposition 3.3. Assuming Conjecture 2.7, we have the following:

- (1) If $(G, A, B) \sim \mathcal{G}(n, q, \rho)$ and let \mathbb{P}_* to be the joint low of (π_*, G, A, B) where π_* is the latent matching. Suppose that q, ρ, D satisfy the assumptions in Item (1) of Theorem 3.2. Then for all $\{f_{i,j} : 1 \leq i, j \leq n\}$ that can be computed in time $n^{\tilde{O}(D)}$, we have $\mathbb{E}_{\mathbb{P}_*}[f_{i,\pi_*(i)}] = o(1)$ for all $1 \leq i \leq n$.
- (1) If $(G, A, B) \sim S(n, \frac{\lambda}{n}; k, \epsilon, s)$ and let \mathbb{P}_* to be the joint low of (π_*, G, A, B) where π_* is the latent matching. Suppose that λ, k, ϵ, s satisfy the assumptions in Item (2) of Theorem 3.2. Then there exists an event \mathcal{E} such that $\mathbb{P}_*(\mathcal{E}) = \omega(1)$ such that for all $\{f_{i,j} : 1 \leq i, j \leq n\}$ that can be computed in time $n^{\widetilde{\Theta}(D)}$, we have $\mathbb{E}_{\mathbb{P}_*}[f_{i,\pi_*(i)} \mid \mathcal{E}] = o(1)$ for all $1 \leq i \leq n$.

Clearly, based on Proposition 3.3, we can deduce Theorem 3.2 via a simple Markov inequality. The rest of this section is devoted to the proof of Proposition 3.3. In the following subsections, our main focus is on proving Item (1) of Proposition 3.3. Given the similarity between the proofs of Item (1) and Item (2), for Item (2) we will provide an outline with the main differences while adapting arguments from proving Item (1) without presenting full details.

3.1 Proof of Item (1) in Proposition 3.3

This subsection is devoted to the proof of Item (1) in Proposition 3.3. Throughout this subsection, we will denote \mathbb{P}_* to be the law of (π_*, G, A, B) where $(G, A, B) \sim \mathcal{G}(n, q; \rho)$. We will also denote \mathbb{P} to be the marginal law of (A, B). In addition, we assume throughout this subsection that there exists a small constant $0 < \delta < 0.01$ such that

$$\rho^2 < \alpha - \delta$$
, $q = n^{-1+o(1)}$, $\log D = o\left(\frac{\log n}{\log(nq)} \land \sqrt{\log n}\right)$. (3.3)

We first introduce some notations used in [15].

Definition 3.4. Given a graph H = H(V, E), define

$$\Phi(H) = \left(n^{1+4/D} D^{20}\right)^{|V(H)|} \left(q D^6\right)^{|E(H)|}, \qquad (3.4)$$

and the graph H is said to be bad if $\Phi(H) < (\log n)^{-1}$. Furthermore, we say a graph is admissible if it contains no bad subgraph, and we say it is inadmissible otherwise.

Denote \mathcal{E} for the event that G does not contain any bad subgraph with no more than d^2 vertices. In addition, let $\overline{\mathbb{P}}_*$ be the conditional version of \mathbb{P}_* given \mathcal{E} , and let $\overline{\mathbb{P}}$ be the corresponding marginal distribution of $\overline{\mathbb{P}}_*$ on (A, B).

We remark here that our definition of "bad" amounts to an atypically large edge density, with a carefully chosen quantitative threshold on "large". Roughly speaking, we expect that any subgraph with size no more than $D^2 = n^{o(1)}$ of a sparse Erdős-Rényi graph has edge-to-vertex ratio 1 + o(1). In the definition of Φ , the term $n^{1+4/D}D^{20}$ should be thought as $n^{1+o(1)}$, and qD^6 as $n^{-1+o(1)}$. The o(1) terms are tuned carefully so that for a typical subgraph H of a sparse Erdős-Rényi graph, $\Phi(H)$ is much bigger than 1. The choice of $(\log n)^{-1}$ as the Φ -threshold for bad graph is somewhat arbitrary, which we will only need to be vanishing as $n \to \infty$. In [15], the authors showed that one the one hand, we have $\mathbb{P}_*(\mathcal{E}) = 1 - o(1)$ and thus $\mathrm{TV}(\mathbb{P}_*, \mathbb{P}_*), \mathrm{TV}(\mathbb{P}, \mathbb{P}) = o(1)$; on the other hand, we have $\mathrm{Adv}_{\leq D}\left(\frac{\mathrm{d}\mathbb{P}}{\mathrm{d}\mathbb{Q}}\right) = O(1)$, thus verifying the low-degree hardness for the detection problem. The first step of our proof is to slightly generalize the result in [15], as incorporated in the following lemma.

Lemma 3.5. For all
$$1 \le i, j \le n$$
, we have $\operatorname{Adv}_{\le D}\left(\frac{\mathrm{d}\overline{\mathbb{P}}(\cdot|\pi_*(i)=j)}{\mathrm{d}\mathbb{Q}}\right) = O_{\delta}(1)$.

The proof of Lemma 3.5 is quite technical and thus is postponed to the end of this subsection. Now, based on Lemma 3.5, we establish the following result, which basically suggests that it is impossible to obtain a "good approximation" of $\mathbf{1}_{\{\pi_*(i)=j\}}$ in some sense.

Lemma 3.6. Assuming Conjecture 2.7, we have for all $1 \leq i \leq n$ and all statistics $\{g_{i,j} = g_{i,j}(A,B) : 1 \leq j \leq n\}$ that can be computed in time $n^{\tilde{O}(D)}$, it holds that

$$\sum_{j=1}^{n} \mathbb{E}_{\mathbb{P}_{*}} \left[\left(\mathbf{1}_{\{\pi_{*}(i)=j\}} - g_{i,j} \right)^{2} \right] \ge 1 - o(1) \,. \tag{3.5}$$

Proof. Without losing of generality, we may assume that i = 1 in the following proof. Suppose on the contrary that there are statistics $\{g_j = g_j(A, B) : 1 \le j \le n\}$ that can be computed in time $n^{\tilde{O}(D)}$ such that

$$\sum_{j=1}^{n} \mathbb{E}_{\mathbb{P}_{*}} \left[\left(\mathbf{1}_{\{\pi_{*}(1)=j\}} - g_{j} \right)^{2} \right] \leq 1 - c \text{ for some constant } c > 0.$$
(3.6)

Without losing of generality, we may assume that $0 \le g_i \le 1$, since otherwise we may replace g_i with min $\{\max\{g_i, 0\}, 1\}$, which will only make the left hand side of (3.6) smaller. Denote

$$\Lambda := \left\{ 1 \le j \le n : \mathbb{E}_{\mathbb{P}_*} \left[(\mathbf{1}_{\{\pi_*(1)=j\}} - g_j)^2 \right] \le \frac{1 - \frac{c}{2}}{n} \right\}.$$
(3.7)

Using Markov inequality, we see that

$$|\Lambda| \ge n - \frac{1-c}{1-\frac{c}{2}}n \ge \frac{cn}{2}.$$
 (3.8)

In addition, it is straightforward to check that for all $i \in \Lambda$

$$\mathbb{E}_{\overline{\mathbb{P}}}\left[(1-g_i)^2 \mid \pi_*(1)=i\right] \le n \cdot \mathbb{E}_{\mathbb{P}_*}\left[(\mathbf{1}_{\{\pi_*(1)=i\}}-g_i)^2\right] \le 1-\frac{c}{2}.$$

Thus, using Markov inequality we see that (note that $0 \le g_i \le 1$)

$$\overline{\mathbb{P}}(g_i > \frac{c}{2}) \ge \Omega(1)$$

Since $\mathrm{TV}(\mathbb{P},\overline{\mathbb{P}}) = o(1)$, we have

$$\mathbb{P}'(g_i > \frac{c}{2}) \ge \Omega(1) \text{ for all } \operatorname{TV}(\mathbb{P}', \mathbb{P}) = o(1).$$
(3.9)

However, using Lemma 3.5 and Theorem 2.4, we see that (assuming Conjecture 2.7) we have $\mathbb{Q} \triangleleft_{\leq D} \mathbb{P}$. Thus we must have

$$\mathbb{Q}'(g_i > \frac{c}{2}) \ge \Omega(1) \text{ for all } \operatorname{TV}(\mathbb{Q}', \mathbb{Q}) = o(1).$$
 (3.10)

This yields that

$$\mathbb{E}_{\mathbb{Q}'}[g_i] = \Omega(1) \text{ for all } \operatorname{TV}(\mathbb{Q}, \mathbb{Q}') = o(1) \text{ and } i \in \Lambda.$$
(3.11)

To this end, define $g = g_1 + \ldots + g_n$. Using (3.8) and (3.11), we see that (recall again that $0 \le g_i \le 1$)

$$\mathbb{E}_{\mathbb{Q}'}[g] = \Omega(n) \text{ for all } \operatorname{TV}(\mathbb{Q}, \mathbb{Q}') = o(1).$$
(3.12)

However, using $0 \leq g \leq n$, we see that for all $\mathrm{TV}(\mathbb{P}, \mathbb{P}') = o(1)$ we have

$$\mathbb{E}_{\mathbb{P}'}[g^2] \leq \mathrm{TV}(\overline{\mathbb{P}}, \mathbb{P}') \cdot \|g\|_{\infty}^2 + \mathbb{E}_{\overline{\mathbb{P}}}[g^2] \leq o(n^2) + 2\left(1 + \mathbb{E}_{\overline{\mathbb{P}}}[(g-1)^2]\right) \\
\leq o(n^2) + 2\mathbb{E}_{\overline{\mathbb{P}}}\left[\left(\sum_{i=1}^n (\mathbf{1}_{\{\pi_*(1)=i\}} - g_i)\right)^2\right] \\
\leq o(n^2) + 2n \cdot \sum_{i=1}^n \mathbb{E}_{\overline{\mathbb{P}}}\left[(\mathbf{1}_{\{\pi_*(1)=i\}} - g_i)^2\right] \stackrel{(3.6)}{=} o(n^2) + O(n) = o(n^2). \quad (3.13)$$

Combining (3.12) and (3.13), we see that for all $TV(\mathbb{Q}, \mathbb{Q}'), TV(\mathbb{P}, \mathbb{P}') = o(1)$ we have

$$\frac{\mathbb{E}_{\mathbb{Q}'}[g]}{\sqrt{\mathbb{E}_{\mathbb{P}'}[g^2]}} = \omega(1).$$
(3.14)

Thus, using the equivalence of Items (2) and (3) in Conjecture 2.7 on the testing problem between \mathbb{P} and \mathbb{Q} , we see that there exists an algorithm with running time $n^{\widetilde{O}(D)}$ that strongly distinguish \mathbb{P} and \mathbb{Q} . However, it is impossible assuming Conjecture 2.7 as [15] showed that $\operatorname{Adv}_{\leq D}\left(\frac{d\mathbb{P}}{d\mathbb{Q}}\right) = O(1)$ where $\operatorname{TV}(\mathbb{P},\mathbb{P}) = o(1)$. This implies that (3.6) is impossible and thus completes our proof. Now we can finish the proof of Theorem 3.2.

Proof of Theorem 3.2 assuming Lemma 3.6. Suppose on the contrary there are statistics $\{f_j : 1 \leq j \leq n\}$ that can be computed in time $n^{\widetilde{O}(D)}$ with $\mathbb{E}_{\mathbb{P}_*}[f_{\pi_*(i)}] \geq 1-c$ for some fixed constant 0 < c < 0.01. Using Items (1) and (2) in Definition 3.1, we see that $f_j f_k = 0$ for all $k \neq j$, and thus

$$1 = \mathbb{E}_{\mathbb{P}}\left[(f_1 + \ldots + f_n)^2\right] = \sum_{j=1}^n \mathbb{E}_{\mathbb{P}}\left[f_j^2\right].$$
 (3.15)

In addition, we have

$$\mathbb{E}_{\mathbb{P}_{*}}[f_{\pi_{*}(i)}] = \frac{1}{n} \sum_{j=1}^{n} \mathbb{E}_{\mathbb{P}_{*}}[f_{j} \mid \pi_{*}(i) = j] \ge c.$$
(3.16)

Thus, for all $\lambda \in [0, 1]$ we have

$$\sum_{j=1}^{n} \mathbb{E}_{\mathbb{P}_{*}} \left[\left(\mathbf{1}_{\{\pi_{*}(i)=j\}} - \frac{1-\lambda}{n} - \lambda f_{j} \right)^{2} \right] \stackrel{(3.15),(3.16)}{\leq} 1 + \lambda^{2} - 2c\lambda + O\left(\frac{1}{n}\right).$$

Thus, by choosing $\lambda = \lambda(c)$ to be a sufficiently small positive constant we get that

$$\sum_{j=1}^{n} \mathbb{E}_{\mathbb{P}_{*}} \left[\left(\mathbf{1}_{\{\pi_{*}(i)=j\}} - g_{j} \right)^{2} \right] = 1 - \Omega(1) \text{ where } g_{j} = \frac{1-\lambda}{n} + \lambda f_{j},$$

contradicting to Lemma 3.6. This leads to the desired result.

Now we provide the postponed proof of Lemma 3.5. For two graphs $S_1, S_2 \in \mathcal{K}_n$, define the polynomial ϕ_{S_1,S_2} associated with S_1, S_2 by

$$\phi_{S_1,S_2}(\{A_{i,j}\},\{B_{i,j}\}) = (q(1-q))^{-\frac{|E(S_1)|+|E(S_2)|}{2}} \prod_{(i,j)\in E(S_1)} \overline{A}_{i,j} \prod_{(i,j)\in E(S_2)} \overline{B}_{i,j}, \quad (3.17)$$

where $\overline{A}_{i,j} = A_{i,j} - q$, $\overline{B}_{i,j} = B_{i,j} - q$ for all $(i, j) \in U$. In particular, $\phi_{\emptyset,\emptyset} \equiv 1$. It can be easily checked that $\mathcal{O}_D = \{\phi_{S_1,S_2} : |E(S_1)| + |E(S_2)| \leq D\}$ constitutes a standard orthogonal basis of \mathcal{P}_D under \mathbb{Q} . In addition, we say a polynomial $\phi_{S_1,S_2} \in \mathcal{O}_D$ is admissible if both S_1 and S_2 are admissible graphs. Furthermore, we define $\mathcal{O}'_D \subset \mathcal{O}_D$ as the set of admissible polynomials in \mathcal{O}_D , and define $\mathcal{P}'_D \subset \mathcal{P}_D$ as the linear subspace spanned by polynomials in \mathcal{O}'_D . It has been shown in [15, Proposition 3.4] that for any $f \in \mathcal{P}_D$, there exists $f' \in \mathcal{P}'_D$ such that $\mathbb{E}_{\mathbb{Q}}[(f')^2] \leq 8\mathbb{E}_{\mathbb{Q}}[f^2]$ and f' = f a.s. under both $\overline{\mathbb{P}}$ and $\overline{\mathbb{P}}_*$. Thus, we get that

$$\mathsf{Adv}_{\leq D}\left(\frac{\mathrm{d}\overline{\mathbb{P}}(\cdot|\pi_*(1)=i)}{\mathrm{d}\mathbb{Q}}\right) = \sup_{f\in\mathcal{P}_D}\frac{\mathbb{E}_{\overline{\mathbb{P}}}[f\mid\pi_*(1)=1]}{\sqrt{\mathbb{E}_{\mathbb{Q}}[f^2]}} \leq 2\sqrt{2}\sup_{f\in\mathcal{P}'_D}\frac{\mathbb{E}_{\overline{\mathbb{P}}}[f\mid\pi_*(1)=1]}{\sqrt{\mathbb{E}_{\mathbb{Q}}[f^2]}}\,.$$
 (3.18)

Thus, it suffices to show the right hand side of (3.18) is bounded by O(1). Similar as (2.9), we have

$$\sup_{f \in \mathcal{P}'_D} \frac{\mathbb{E}_{\overline{\mathbb{P}}}[f \mid \pi_*(1) = 1]^2}{\mathbb{E}_{\mathbb{Q}}[f^2]} = \sum_{\phi_{S_1, S_2} \in \mathcal{O}'_D} \mathbb{E}_{\overline{\mathbb{P}}}[\phi_{S_1, S_2} \mid \pi_*(1) = i]^2.$$

Without losing of generality, in the following we will only show that

$$\sum_{\phi_{S_1,S_2} \in \mathcal{O}'_D} \mathbb{E}_{\overline{\mathbb{P}}} [\phi_{S_1,S_2} \mid \pi_*(1) = 1]^2 = O(1) \,. \tag{3.19}$$

For a deterministic permutation $\pi \in \mathfrak{S}_n$, we use \mathbb{P}_{π} and $\overline{\mathbb{P}}_{\pi}$ to represent $\mathbb{P}_*(\cdot \mid \pi_* = \pi)$ and $\overline{\mathbb{P}}_*(\cdot \mid \pi_* = \pi)$, respectively. For $S_1, S_2 \in \mathcal{K}_n$ with $|E(S_1)|, |E(S_2)| \leq D$, define

$$\mathbf{F}(S_1, S_2) = \sum_{\substack{\mathbf{H}_0 \in \mathcal{H}\\\mathbf{H}_0 \hookrightarrow S_i, i=1,2}} n^{-\frac{|V(S_1)| + |V(S_2)|}{2}} \rho^{|E(\mathbf{H}_0)|} d^{-6(|E(S_1)| + |E(S_2)| - 2|E(\mathbf{H}_0)|)} \operatorname{Aut}(\mathbf{H}_0),$$
(3.20)

where the notation $\mathbf{H} \hookrightarrow S$ means that \mathbf{H} can be embedded into S as a subgraph. The key of our proof is the following estimation.

Proposition 3.7. We have

$$\left| \mathbb{E}_{\overline{\mathbb{P}}}[\phi_{S_1,S_2} \mid \pi_*(1) = 1] \right| \le [1 + o(1)] \left(\mathbf{1}_{1 \notin V(S_1) \cap V(S_2)} + n \mathbf{1}_{1 \in V(S_1) \cap V(S_2)} \right) \cdot \mathbb{F}(S_1, S_2) . \quad (3.21)$$
Proof. For $S_2 \hookrightarrow S_1$. S_2 define

Proof. For $S_0 \hookrightarrow S_1, S_2$, define

$$M(S_0, S_1, S_2) = \rho^{|E(S_0)|} n^{-\frac{|V(S_1)| + |V(S_2)|}{2} + |V(S_0)|} D^{-7(|E(S_1)| + |E(S_2)| - 2|E(S_0)|)}.$$
(3.22)

Using [15, Equation (3.29)], for any $S_1, S_2 \in \mathcal{K}_n$ with at most D edges and any permutation $\pi \in \mathfrak{S}_n$, we have that (denote $S_0 = S_1 \cap S_2$ below)

$$\left|\mathbb{E}_{\overline{\mathbb{P}}_{\pi}}[\phi_{S_1,S_2}]\right| \le M(S_0,S_1,S_2).$$
(3.23)

Thus, note that

$$\left| \mathbb{E}_{\overline{\mathbb{P}}}[\phi_{S_1,S_2} \mid \pi_*(1) = 1] \right| = \frac{1}{(n-1)!} \left| \sum_{\pi \in \mathfrak{S}_n, \pi(1) = 1} \mathbb{E}_{\overline{\mathbb{P}}_\pi}[\phi_{S_1,S_2}] \right| \\ \leq \frac{1}{(n-1)!} \sum_{\pi \in \mathfrak{S}_n, \pi(1) = 1} \left| \mathbb{E}_{\overline{\mathbb{P}}_\pi}[\phi_{S_1,S_2}] \right|,$$
(3.24)

we can group the permutations $\pi \in \mathfrak{S}_n$ according to the realization of $S_0 = S_1 \cap \pi^{-1}(S_2)$ and obtain that $|\mathbb{E}_{\mathbb{F}}[\phi_{S_1,S_2} \mid \pi_*(1) = 1]|$ is bounded by

$$\frac{1}{(n-1)!} \sum_{\substack{S_0, S_0': S_0 \cong S_0'\\S_0 \subset S_1, S_0' \subset S_2}} \#\{\pi \in \mathfrak{S}_n : \pi(1) = 1, \pi(S_0) = S_0', S_0 = S_1 \cap \pi^{-1}(S_2)\} M(S_0, S_1, S_2) .$$
(3.25)

For the case when $1 \notin V(S_1) \cap V(S_2)$, we must have $1 \notin V(S_0)$ and thus

Enum :=
$$\#\{\pi \in \mathfrak{S}_n : \pi(1) = 1, \pi(S_0) = S'_0\}$$
 (3.26)

is upper-bounded by

$$\operatorname{Aut}(S_0)(n-1-|V(S_0)|)! = [1+o(1)]\operatorname{Aut}(S_0)(n-1)! \cdot n^{-|V(S_0)|}.$$
(3.27)

Thus, we have that (3.25) is further upper-bounded by (up to a factor of (1 + o(1)))

$$\sum_{\substack{S_0, S'_0: S_0 \cong S'_0 \\ S_0 \subset S_1, S'_0 \subset S_2}} n^{-|V(S_0)|} \operatorname{Aut}(S_0) M(S_0, S_1, S_2)$$

$$= \sum_{\substack{\mathbf{H}_0 \in \mathcal{H} \\ \mathbf{H}_0 \hookrightarrow S_i, i=1,2}} \frac{\operatorname{Aut}(\mathbf{H}_0)}{n^{|V(\mathbf{H}_0)|}} M(\mathbf{H}_0, S_1, S_2) \cdot \#\{(S_0, S'_0) : S_0 \subset S_1, S'_0 \subset S_2, S_0 \cong S'_0 \cong \mathbf{H}_0\}$$

$$\leq \sum_{\substack{\mathbf{H}_0 \in \mathcal{H} \\ \mathbf{H}_0 \hookrightarrow S_i, i=1,2}} \frac{\operatorname{Aut}(\mathbf{H}_0)}{n^{|V(\mathbf{H}_0)|}} M(\mathbf{H}_0, S_1, S_2) \cdot D^{|E(S_1)| + E(S_2) - 2|E(\mathbf{H}_0)|},$$

where the last inequality follows from Lemma A.1 (v) and the assumption that $|E(S_1)|$ and $|E(S_2)|$ are bounded by D. For the case $1 \in V(S_1) \cap V(S_2)$, we have $\mathsf{Enum} \leq [1 + o(1)] \operatorname{Aut}(S_0)n! \cdot n^{-|V(S_0)|}$, and thus we get an additional factor of n in the final estimation. This completes the proof of Proposition 3.7.

We are now ready to establish (3.19), thus completing the proof of Lemma 3.5.

Proof of (3.19). We divide the left hand side of (3.19) into two parts: the first part consists of those S1, S2 such that $1 \notin V(S_1) \cap V(S_2)$, and the second part consists of those with $1 \in V(S_1) \cap V(S_2)$. We bound the first part

$$\sum_{\substack{\phi_{S_1,S_2} \in \mathcal{O}'_D\\ \notin V(S_1) \cap V(S_2)}} \mathbb{E}_{\mathbb{P}} [\phi_{S_1,S_2} \mid \pi_*(1) = 1]^2$$
(3.28)

via Proposition 3.7 by

1

$$[1+o(1)] \sum_{\substack{S_1,S_2 \in \mathcal{K}_n \text{ admissible} \\ |E(S_1)|+|E(S_2)| \le D}} \left(\sum_{\substack{\mathbf{H}_0 \in \mathcal{H} \\ \mathbf{H}_0 \hookrightarrow S_i, i=1,2}} \frac{\rho^{|E(\mathbf{H}_0)|} \operatorname{Aut}(\mathbf{H}_0)}{D^{6(|E(S_1)|+|E(S_2)|-2|E(\mathbf{H}_0)|)} n^{\frac{1}{2}(|V(S_1)|+|V(S_2)|)}} \right)^2$$

$$\le [1+o(1)] \sum_{\substack{S_1,S_2 \in \mathcal{K}_n \text{ admissible} \\ |E(S_1)|+|E(S_2)| \le D}} \left(\sum_{\substack{\mathbf{H}_0 \in \mathcal{H} \\ \mathbf{H}_0 \hookrightarrow S_i, i=1,2}} \frac{\rho^{2|E(\mathbf{H}_0)|} \operatorname{Aut}(\mathbf{H}_0)^2}{D^{6(|E(S_1)|+|E(S_2)|-2|E(\mathbf{H}_0)|)} n^{|V(S_1)|+|V(S_2)|}} \right)$$

$$\times \left(\sum_{\substack{\mathbf{H}_0 \in \mathcal{H} \\ \mathbf{H}_0 \hookrightarrow S_i, i=1,2}} D^{-6(|E(S_1)|+|E(S_2)|-2|E(\mathbf{H}_0)|)} \right), \qquad (3.29)$$

where the last inequality comes form Cauchy-Schwartz inequality. Using [15, Equations (3.28)], we have that the second bracket (3.29) is bounded by $O_{\delta}(1)$. Thus, (3.28) is bounded by a constant factor times

$$\sum_{\substack{\mathbf{H}_{0} \in \mathcal{H} \\ \mathbf{H}_{0} \hookrightarrow S_{i}, i=1,2}} \rho^{2|E(\mathbf{H}_{0})|} \operatorname{Aut}(\mathbf{H}_{0})^{2} \sum_{\substack{S_{1}, S_{2} \in \mathcal{K}_{n} \text{ admissible} \\ |E(S_{1})| + |E(S_{2})| \leq D}} \frac{n^{-|V(S_{1})| - |V(S_{2})|}}{D^{6(|E(S_{1})| + |E(S_{2})| - 2|E(\mathbf{H}_{0})|)}} .$$
(3.30)

By denoting $S_1 \cong \mathbf{S}_1$ and $S_2 \cong \mathbf{S}_2$ with $\mathbf{S}_1, \mathbf{S}_2 \in \mathcal{H}$, the right hand side of (3.30) reduces to

$$\sum_{\mathbf{H}_{0}\in\mathcal{H}} \rho^{2|E(\mathbf{H}_{0})|} \operatorname{Aut}(\mathbf{H}_{0})^{2} \sum_{\substack{\mathbf{S}_{1}, \mathbf{S}_{2}\in\mathcal{H} \text{ admissible}\\ \mathbf{H}_{0}\hookrightarrow\mathbf{S}_{i}, i=1,2\\ |E(\mathbf{S}_{1})|+|E(\mathbf{S}_{2})|\leq d}} \frac{\#\{S_{1}, S_{2}: S_{1}\cong\mathbf{S}_{1}, S_{2}\cong\mathbf{S}_{2}\}}{D^{6(|E(\mathbf{S}_{1})|+|E(\mathbf{S}_{2})|-2|E(\mathbf{H}_{0})|)}n^{|V(\mathbf{S}_{1})|+|V(\mathbf{S}_{2})|}}$$

$$\leq \sum_{\mathbf{H}_{0}\in\mathcal{H} \text{ admissible}} \rho^{2|E(\mathbf{H}_{0})|} \sum_{\substack{\mathbf{S}_{1}, \mathbf{S}_{2}\in\mathcal{H}, \mathbf{H}_{0}\hookrightarrow\mathbf{S}_{i}, i=1,2\\ |E(\mathbf{S}_{1})|+|E(\mathbf{S}_{2})|\leq d}} \frac{\operatorname{Aut}(\mathbf{H}_{0})^{2}}{\operatorname{Aut}(\mathbf{S}_{1})\operatorname{Aut}(\mathbf{S}_{2})D^{6(|E(\mathbf{S}_{1})|+|E(\mathbf{S}_{2})|-2|E(\mathbf{H}_{0})|)}}$$

$$\leq O_{\delta}(1) \cdot \sum_{\mathbf{H}_{0}\in\mathcal{H} \text{ admissible}} \rho^{2|E(\mathbf{H}_{0})|} \rho^{2|E(\mathbf{H}_{0})|} = O_{\delta}(1), \qquad (3.31)$$

where the first inequality follows from

$$\#\{S_1, S_2: S_1 \cong \mathbf{S}_1, S_2 \cong \mathbf{S}_2\} \le [1 + o(1)] \cdot \frac{n^{|V(\mathbf{S}_1)|}}{\operatorname{Aut}(\mathbf{S}_1)} \cdot \frac{n^{|V(\mathbf{S}_2)|}}{\operatorname{Aut}(\mathbf{S}_2)},$$

the second inequality follows from [15, Equations (3.31)-(3.32)], and the last equality follows from [15, Lemma A.3]. Thus, we have

$$(3.28) = O_{\delta}(1) \,. \tag{3.32}$$

We now deal with the second part

$$\sum_{\substack{\phi_{S_1,S_2} \in \mathcal{O}'_D\\1 \in V(S_1) \cap V(S_2)}} \mathbb{E}_{\overline{\mathbb{P}}} \left[\phi_{S_1,S_2} \mid \pi_*(1) = 1 \right]^2.$$
(3.33)

Following a similar derivation of (3.30), we see that (3.33) is bounded by a constant factor times

$$\sum_{\substack{\mathbf{H}_0 \in \mathcal{H} \\ \mathbf{H}_0 \hookrightarrow S_i, i=1,2}} \rho^{2|E(\mathbf{H}_0)|} \operatorname{Aut}(\mathbf{H}_0)^2 \sum_{\substack{S_1, S_2 \in \mathcal{K}_n \text{admissible} \\ 1 \in V(S_1) \cap V(S_2) \\ |E(S_1)| + |E(S_2)| \le D}} \frac{n^{-|V(S_1)| - |V(S_2)|}}{D^{6(|E(S_1)| + |E(S_2)| - 2|E(\mathbf{H}_0)|)}} \times n^2$$

(Note that in the above there is an additional factor of n^2 compared with (3.30).) However, due to the restriction $1 \in V(S_1) \cap V(S_2)$, for any fixed $\mathbf{S}_1, \mathbf{S}_2 \in \mathcal{H}$ the enumeration of (S_1, S_2) such that $S_1 \cong \mathbf{S}_1, S_2 \cong \mathbf{S}_2$ is upper-bounded by

$$\frac{n^{|V(\mathbf{S}_1)|-1}}{\operatorname{Aut}(\mathbf{S}_1)} \cdot \frac{n^{|V(\mathbf{S}_2)|-1}}{\operatorname{Aut}(\mathbf{S}_2)} = \frac{n^{|V(\mathbf{S}_1)|+|V(\mathbf{S}_2)|}}{\operatorname{Aut}(\mathbf{S}_1)\operatorname{Aut}(\mathbf{S}_2)} \times n^{-2}.$$

Thus, after canceling the factor of n^{-2} with the factor of n^2 above we arrive at the same upper bound as we obtain on (3.28). This shows that

$$(3.33) = O_{\delta}(1). \tag{3.34}$$

Combining (3.32) and (3.34) leads to the desired result.

3.2 Proof of Item (2) in Proposition 3.3

This subsection is devoted to the proof of Item (2) in Proposition 3.3. Throughout this subsection, we will denote \mathbb{P}_* to be the law of (π_*, G, A, B) where $(G, A, B) \sim S(n, \frac{\lambda}{n}; k, \epsilon; s)$ and \mathbb{P} the marginal law of (A, B). In addition, we denote \mathbb{Q} to be the law of a pair of independent Erdős-Rényi models $\mathcal{G}(n, \frac{\lambda s}{n})$. In addition, we assume throughout this subsection that there exists a small constant $0 < \delta < 0.01$ such that

$$s < \sqrt{\alpha} - \delta, \quad \epsilon^2 \lambda s < 1 - \delta.$$
 (3.35)

We also choose a sufficiently large constant $N = N(k, \lambda, \delta, \epsilon, s) \ge 2/\delta$ such that

$$(\sqrt{\alpha} - \delta)(1 + \epsilon^N k) \le \sqrt{\alpha} - \delta/2; \quad 10k(1 - \delta)^N \le (1 - \delta/2)^N; (\sqrt{\alpha} - \delta/2)(1 + (1 - \delta/2)^N)^2 \le \sqrt{\alpha} - \delta/4; \quad (1 - \delta/2)^N(N + 1) \le 1.$$
 (3.36)

Again, we begin by recalling some notations in [9].

Definition 3.8. Denote $\tilde{\lambda} = \lambda \vee 1$. Given a graph H = H(V, E), define

$$\Upsilon(H) = \left(\frac{2\tilde{\lambda}^2 k^2 n}{D^{50}}\right)^{|V(H)|} \left(\frac{1000\tilde{\lambda}^{20} k^{20} D^{50}}{n}\right)^{|E(H)|}.$$
(3.37)

Then we say the graph H is bad if $\Upsilon(H) < (\log n)^{-1}$, and we say a graph H is selfbad if H is bad and $\Upsilon(H) < \Upsilon(K)$ for all $K \subset H$. Furthermore, we say that a graph H is admissible if it contains no bad subgraph and $C_j(H) = \emptyset$ for $j \leq N$; we say H is inadmissible otherwise. Denote $\mathcal{E} = \mathcal{E}^{(1)} \cap \mathcal{E}^{(2)}$, where $\mathcal{E}^{(1)}$ is the event that G does not contain any bad subgraph with no more than D^3 vertices, and $\mathcal{E}^{(2)}$ is the event that G does not contain any cycles with length at most N. **Definition 3.9.** List all self-bad subgraphs of \mathcal{K}_n with at most D^3 vertices and all cycles of \mathcal{K}_n with lengths at most N in an arbitrary but prefixed order (B_1, \ldots, B_M) . Define a stochastic block model with "bad graphs" removed as follows: (1) sample $G \sim \mathcal{S}(n, \frac{\lambda}{n}; k, \epsilon)$; (2) for each $1 \leq i \leq M$ such that $B_i \subset G$, we independently uniformly remove one edge in B_i . The unremoved edges in G constitute a graph G', which is the output of our modified stochastic block model. Clearly, from this definition G' does not contain any cycle of length at most N nor any bad subgraph with at most D^3 vertices. Conditioned on G' and π_* , we define

$$A'_{i,j} = G'_{i,j}J'_{i,j}, B'_{i,j} = G'_{\pi^{-1}_*(i),\pi^{-1}_*(j)}K'_{i,j},$$

where J' and K' are independent Bernoulli variables with parameter s. Let $\widetilde{\mathbb{P}}_* = \widetilde{\mathbb{P}}_{*,n}$ be the law of $(\sigma_*, \pi_*, G, G', A', B')$ and denote $\widetilde{\mathbb{P}} = \widetilde{\mathbb{P}}_n$ the marginal law of (A', B').

It was shown in [9, Lemmas 4.2 and 4.4] that

$$\mathbb{P}_*(\mathcal{E}) = \Omega(1) \text{ and } \operatorname{TV}(\mathbb{P}, \mathbb{P}(\cdot \mid \mathcal{E})) = o(1).$$

We now show the following lemma.

Lemma 3.10. We have $\operatorname{Adv}_{\leq D}\left(\frac{\operatorname{d}\widetilde{\mathbb{P}}(\cdot|\pi_*(i)=j)}{\operatorname{d}\mathbb{Q}}\right) = O_{\delta}(1).$

Based on Lemma 3.10, we can deduce our main result just as how we deduce Theorem 3.2 from Lemma 3.5. The only difference is that we will replace all $\overline{\mathbb{P}}$ with $\widetilde{\mathbb{P}}$ and replace all \mathbb{P} with $\mathbb{P}(\cdot | \mathcal{E})$ so we omit further details here. Now it remains to prove Lemma 3.10. Recall the definition of ϕ_{S_1,S_2} in (3.17) (here we will take $q = \frac{\lambda s}{n}$). We first show the following estimation. Again, we say a polynomial $\phi_{S_1,S_2} \in \mathcal{O}_D$ is admissible if both S_1 and S_2 are admissible graphs. Furthermore, we define $\mathcal{O}'_D \subset \mathcal{O}_D$ as the set of admissible polynomials in \mathcal{O}_D , and define $\mathcal{P}'_D \subset \mathcal{P}_D$ as the linear subspace spanned by polynomials in \mathcal{O}'_D . Using [9, Proposition 4.5], we see that for any $f \in \mathcal{P}_D$, there exists some $f' \in \mathcal{P}'_D$ such that $\mathbb{E}_{\mathbb{Q}}[(f')^2] \leq O(1) \cdot \mathbb{E}_{\mathbb{Q}}[f^2]$ and f' = f a.s. under both $\widetilde{\mathbb{P}}_*$ and $\widetilde{\mathbb{P}}$. Thus, we get that

$$\mathsf{Adv}_{\leq D}\Big(\frac{\mathrm{d}\widetilde{\mathbb{P}}(\cdot|\pi_*(i)=j)}{\mathrm{d}\mathbb{Q}}\Big) = \sup_{f\in\mathcal{P}_D}\frac{\mathbb{E}_{\widetilde{\mathbb{P}}}[f\mid\pi_*(i)=j]}{\sqrt{\mathbb{E}_{\mathbb{Q}}[f^2]}} \leq 2\sqrt{2}\sup_{f\in\mathcal{P}'_D}\frac{\mathbb{E}_{\widetilde{\mathbb{P}}}[f\mid\pi_*(i)=j]}{\sqrt{\mathbb{E}_{\mathbb{Q}}[f^2]}} \,.$$

Thus, it suffices to show that

$$\sup_{f \in \mathcal{P}'_D} \frac{\mathbb{E}_{\widetilde{\mathbb{P}}}[f \mid \pi_*(1) = 1]}{\sqrt{\mathbb{E}_{\mathbb{Q}}[f^2]}} = \sum_{\phi_{S_1, S_2} \in \mathcal{O}'_D} \mathbb{E}_{\widetilde{\mathbb{P}}} \left[\phi_{S_1, S_2} \mid \pi_*(i) = j\right]^2 = O_{\delta}(1) \,.$$

For notational simplicity, in the following we will only show

$$\sum_{\phi_{S_1,S_2} \in \mathcal{O}'_D} \mathbb{E}_{\widetilde{\mathbb{P}}} \left[\phi_{S_1,S_2} \mid \pi_*(1) = 1 \right]^2 = O_\delta(1) \,, \tag{3.38}$$

and the general cases for $\pi_*(i) = j$ can be derived in a similar manner. For $H \subset S$, we define $\mathbb{N}(S, H)$ to be

$$\mathbb{N}(S,H) = \left(\frac{D^{28}}{n^{0.1}}\right)^{\frac{1}{2}(|\mathcal{L}(S)\setminus V(H)| + \tau(S) - \tau(H))} \left(1 - \frac{\delta}{2}\right)^{|E(S)| - |E(H)|}.$$
(3.39)

We first show the following estimation.

Proposition 3.11. For all admissible $S_1, S_2 \in \mathcal{K}_n$ with $|E(S_1)|, |E(S_2)| \leq D$, we have that $|\mathbb{E}_{\tilde{\mathbb{P}}}[\phi_{S_1,S_2} | \pi_*(1) = 1]|$ is bounded by $O_{\delta}(1)$ times (note that in the summation below H_1, H_2 may have isolated vertices)

$$\sum_{\substack{H_1 \subset S_1, H_2 \subset S_2 \\ H_1 \cong H_2}} \frac{(\sqrt{\alpha} - \frac{\delta}{2})^{|E(H_1)|} \operatorname{Aut}(H_1)}{n^{|V(H_1)| - \mathbf{1}_{\{1 \in V(H_1 \cap V(H_2))\}}}} * \frac{2^{|\mathfrak{C}(S_1, H_1)| + |\mathfrak{C}(S_2, H_2)|} \mathbb{N}(S_1, H_1) \mathbb{N}(S_2, H_2)}{n^{\frac{1}{2}(|V(S_1)| + |V(S_2)| - |V(H_1)| - |V(H_2)|)}}.$$
 (3.40)

Proof. For each deterministic permutation $\pi \in \mathfrak{S}_n$ and each labeling $\sigma \in [k]^n$, we denote $\widetilde{\mathbb{P}}_{\sigma,\pi} = \widetilde{\mathbb{P}}(\cdot \mid \sigma_* = \sigma, \pi_* = \pi)$, $\widetilde{\mathbb{P}}_{\pi} = \widetilde{\mathbb{P}}(\cdot \mid \pi_* = \pi)$ and $\widetilde{\mathbb{P}}_{\sigma} = \widetilde{\mathbb{P}}(\cdot \mid \sigma_* = \sigma)$ respectively. It is clear that

$$\left| \mathbb{E}_{\widetilde{\mathbb{P}}} \left[\phi_{S_1, S_2} \mid \pi_*(1) = 1 \right] \right| = \left| \frac{1}{(n-1)!} \sum_{\substack{\pi \in \mathfrak{S}_n \\ \pi(1) = 1}} \mathbb{E}_{\widetilde{\mathbb{P}}_{\pi}} [\phi_{S_1, S_2}] \right| \\ \leq \frac{1}{(n-1)!} \sum_{\substack{\pi \in \mathfrak{S}_n \\ \pi(1) = 1}} \left| \mathbb{E}_{\widetilde{\mathbb{P}}_{\pi}} [\phi_{S_1, S_2}] \right|.$$
(3.41)

For $H \subset S$, we define

$$\mathbb{M}(S,H) = \left(\frac{D^8}{n^{0.1}}\right)^{\frac{1}{2}(|\mathcal{L}(S)\setminus V(H)| + \tau(S) - \tau(H))} \left(1 - \frac{\delta}{2}\right)^{|E(S)| - |E(H)|}.$$
(3.42)

Applying [9, Lemma 4.10], we have that for all admissible $S_1, S_2 \in \mathcal{K}_n$ with at most D edges and for all permutation π on [n], denote $H_1 = S_1 \cap \pi^{-1}(S_2)$ and $H_2 = \pi(S_1) \cap S_2$. We have that $|\mathbb{E}_{\mathbb{F}_{\pi}}[\phi_{S_1,S_2}]|$ is bounded by O(1) times

$$\left(\sqrt{\alpha} - \frac{\delta}{2}\right)^{|E(H_1)|} \sum_{H_1 \ltimes K_1 \subset S_1} \sum_{H_2 \ltimes K_2 \subset S_2} \frac{\mathsf{M}(S_1, K_1) \mathsf{M}(S_2, K_2) \mathsf{M}(K_1, H_1) \mathsf{M}(K_2, H_2)}{n^{\frac{1}{2}(|V(S_1)| + |V(S_2)| - |V(H_1)| - |V(H_2)|)}} \,. \tag{3.43}$$

Note that we have

$$\mu(\{\pi:\pi(1)=1,\pi(H_1)=H_2\}) \le [1+o(1)] \cdot \operatorname{Aut}(H_1)n^{-|V(H_1)|-1+\mathbf{1}_{1\in V(H_1)\cap V(H_2)}}.$$
 (3.44)

Thus, it yields that the right-hand side of (3.41) is bounded by (up to a O(1) factor)

$$\sum_{\substack{H_1,H_2:H_1\cong H_2\\H_1\subset S_1,H_2\subset S_2}} \frac{\operatorname{Aut}(H_1)(\sqrt{\alpha}-\frac{\delta}{2})^{|E(H_1)|}}{n^{|V(H_1)|-\mathbf{1}_{1\in V(H_1)\cap V(H_2)}}} * \frac{\operatorname{P}(S_1,H_1)\operatorname{P}(S_2,H_2)}{n^{\frac{1}{2}}(|V(S_1)|+|V(S_2)|-|V(H_1)|-|V(H_2)|)}, \quad (3.45)$$

where (recall (3.42))

$$\mathbb{P}(S,H) = \sum_{H \ltimes K \subset S} \mathbb{M}(S,K) \mathbb{M}(K,H) \,. \tag{3.46}$$

Using [9, Claim 4.11], we have that $\mathbb{P}(S, H) \leq [1 + o(1)] \cdot 2^{|\mathfrak{C}(S,H)|} \mathbb{N}(S, H)$, thus leading to the desired result.

Now we prove (3.38) formally, thus finishing the proof of Lemma 3.10.

Proof of (3.38). By Proposition 3.11 and Cauchy-Schwartz inequality, (3.38) is upperbounded by $O_{\delta}(1)$ times

$$\sum_{\substack{\phi_{S_1,S_2} \in \mathcal{O}'_D \\ H_1 \subset S_1, H_2 \subset S_2 \\ H_1 \cong H_2}} \left(\sum_{\substack{H_1 \subset S_1, H_2 \subset S_2 \\ H_1 \cong H_2}} n^{0.02|\mathcal{I}(H_1)|} \mathbb{N}(S_1, H_1) \mathbb{N}(S_2, H_2) \right) \times$$
(3.47)
$$\left(\sum_{\substack{H_1 \subset S_1, H_2 \subset S_2 \\ H_1 \cong H_2 \\ H_1 \cong H_2}} \frac{(\sqrt{\alpha} - \frac{\delta}{2})^{2|E(H_1)|} \operatorname{Aut}(H_1)^2}{n^{2|V(H_1)| - 2 \cdot \mathbf{1}_{1 \in V(H_1) \cap V(H_2)} + 0.02|\mathcal{I}(H_1)|}} \cdot \frac{4^{|\mathfrak{C}(S_1, H_1)| + |\mathfrak{C}(S_2, H_2)|} \mathbb{N}(S_1, H_1) \mathbb{N}(S_2, H_2)}{n^{(|V(S_1)| + |V(S_2)| - |V(H_1)| - |V(H_2)|)}} \right)$$

It was shown in [9, Equation (4.35)] that (3.47) is bounded by O(1). Thus, we see that the left hand side of (3.38) is bounded by $O_{\delta}(1)$ times

$$\sum_{\substack{H_1 \cong H_2, H_1, H_2 \text{ admissible} \\ |E(H_1)| + |E(H_2)| \le D}} \frac{(\sqrt{\alpha} - \frac{\delta}{2})^{2|E(H_1)|} \operatorname{Aut}(H_1)^2}{n^{2|V(H_1)| - 2 \cdot \mathbf{1}_{1 \in V(H_1) \cap V(H_2)} + 0.02|\mathcal{I}(H_1)|} \cdot \operatorname{L}(H_1, H_2)},$$

where

$$\mathbf{L}(H_1, H_2) = \sum_{H_1 \subset S_1, H_2 \subset S_2} \frac{4^{|\mathfrak{C}(S_1, H_1)| + |\mathfrak{C}(S_2, H_2)|} \mathbf{N}(S_1, H_1) \mathbf{N}(S_2, H_2)}{n^{(|V(S_1)| + |V(S_2)| - |V(H_1)| - |V(H_2)|)}}$$

Using [9, Equation (4.37)], we see that $L(H_1, H_2) = O_{\delta}(1)$. Thus, the left hand side of (3.38) is bounded by $O_{\delta}(1)$ times

$$\sum_{\substack{H_{1}\cong H_{2},H_{1},H_{2} \text{ admissible}\\|E(H_{1})|+|E(H_{2})|\leq D}} \frac{(\sqrt{\alpha}-\frac{\delta}{2})^{2|E(H_{1})|} \operatorname{Aut}(H_{1})^{2}}{n^{2|V(H_{1})|-2\cdot\mathbf{1}_{1\in V(H_{1})\cap V(H_{2})}+0.02|\mathcal{I}(H_{1})|}} \\ = \sum_{\substack{H_{1}\cong H_{2},H_{1},H_{2} \text{ admissible}\\|E(H_{1})|+|E(H_{2})|\leq D\\1\notin V(H_{1})\cap V(H_{2})}} \frac{(\sqrt{\alpha}-\frac{\delta}{2})^{2|E(H_{1})|} \operatorname{Aut}(H_{1})^{2}}{n^{2|V(H_{1})|+0.02|\mathcal{I}(H_{1})|}} \\ + \sum_{\substack{H_{1}\cong H_{2},H_{1},H_{2} \text{ admissible}\\|E(H_{1})|+|E(H_{2})|\leq D\\1\in V(H_{1})\cap V(H_{2})}} \frac{(\sqrt{\alpha}-\frac{\delta}{2})^{2|E(H_{1})|} \operatorname{Aut}(H_{1})^{2}}{n^{2|V(H_{1})|+0.02|\mathcal{I}(H_{1})|}} \times n^{2}. \quad (3.49)$$

Recall that we use \widetilde{H}_1 to denote the subgraph of H_1 obtained by removing all the vertices in $\mathcal{I}(H_1)$. In addition, for $|V(H_1)| \leq |V(S_1)| \leq 2D$, we have $\operatorname{Aut}(H_1) = \operatorname{Aut}(\widetilde{H}_1) \cdot |\mathcal{I}(H_1)|! \leq (2D)^{|\mathcal{I}(H_1)|} \operatorname{Aut}(\widetilde{H}_1)$. Thus, we see that Thus, we have that

$$(3.48) \leq \sum_{\substack{|E(\mathbf{H})| \leq D, \mathcal{I}(\mathbf{H}) = \emptyset \\ \mathbf{H} \text{ is admissible}}} \sum_{\substack{j \geq 0 \\ |\mathcal{I}(H_1)| = |\mathcal{I}(H_2) : \tilde{H}_1 \cong \tilde{H}_2 \cong \mathbf{H} \\ |\mathcal{I}(H_1)| = |\mathcal{I}(H_2)| = j}} n^{-0.01j} \cdot \frac{\operatorname{Aut}(\mathbf{H})^2 (2D)^{2j} (\sqrt{\alpha} - \frac{\delta}{4})^{2|E(\mathbf{H})|}}{n^{2(|V(\mathbf{H})|+j)}}$$

$$\stackrel{\text{e}}{=} \sum_{\substack{|E(\mathbf{H})| \leq D, \mathcal{I}(\mathbf{H}) = \emptyset \\ \mathbf{H} \text{ is admissible}}} (\sqrt{\alpha} - \frac{\delta}{4})^{2|E(\mathbf{H})|} \leq O_{\delta}(1) ,$$

As for (3.49), due to the restriction $1 \in V(H_1) \cap V(H_2)$, for any fixed $\mathbf{H}_1, \mathbf{H}_2 \in \mathcal{H}$ the enumeration of (H_1, H_2) such that $H_1 \cong \mathbf{H}_1, H_2 \cong \mathbf{H}_2$ is upper-bounded by

$$\frac{n^{|V(\mathbf{H}_1)|-1}}{\operatorname{Aut}(\mathbf{H}_1)} \cdot \frac{n^{|V(\mathbf{H}_2)|-1}}{\operatorname{Aut}(\mathbf{H}_2)} = \frac{n^{|V(\mathbf{H}_1)|+|V(\mathbf{H}_2)|}}{\operatorname{Aut}(\mathbf{H}_1)\operatorname{Aut}(\mathbf{H}_2)} \times n^{-2}.$$

Thus, after canceling the factor of n^{-2} with the factor of n^2 above we arrive at the same upper bound as we obtain on (3.49). This yields that $(3.49) = O_{\delta}(1)$ and thus completes our proof.

4 Detection in correlated SBMs

In this section, we will use the framework we established in Section 2 to show the hardness of detection between correlated SBMs and independent SBMs. The main results of this section is incorporated as follows.

Theorem 4.1. For any constant $K \in \mathbb{N}$, denote \mathbb{P} to be the law of K independent stochastic block models $S(n, \frac{\lambda}{n}; k, \epsilon)$ and denote \mathbb{Q} to be the law of K independent Erdős-Rényi graphs $\mathcal{G}(n, \frac{\lambda}{n})$. Then, assuming Conjecture 2.2, for any $\delta > 0$ there exists $\lambda_0 = \lambda_0(\delta, k)$ to be a sufficiently large constant such that when $\epsilon^2 \lambda < 1 - \delta$ and $\lambda > \lambda_0$, we have we have $\mathbb{P} \bowtie_{\leq D} \mathbb{Q}$ for any $D = n^{o(1)}$.

Our result has an immediate corollary in the detection problem between a pair of correlated SBMs and a pair of independent SBMs, as incorporated in the following corollary.

Corollary 4.2. Assuming Conjecture 2.2, when $\epsilon^2 \lambda s < 1-\delta$, $s < \sqrt{\alpha}-\delta$ and $\lambda > \lambda_0$, there is no algorithms with polynomial running time that can strongly distinguish $S(n, \frac{\lambda}{n}; k, \epsilon; s)$ and two independent $S(n, \frac{\lambda s}{n}; k, \epsilon)$.

Proof. Denote $\mu_{\text{CorSBM}}, \mu_{\text{IndSBM}}$ and μ_{IndER} to be the law of a pair of correlated SBMs $S(n, \frac{\lambda}{n}; k, \epsilon; s)$, a pair of independent SBMs $S(n, \frac{\lambda s}{n}; k, \epsilon)$ and a pair of independent Erdős-Rényi graphs $\mathcal{G}(n, \frac{\lambda s}{n})$, respectively. Suppose on the contrary that there exists an efficient algorithm \mathcal{A} such that

$$\mu_{\text{CorSBM}}(\mathcal{A}(A,B)=1) = 1 - o(1), \quad \mu_{\text{IndSBM}}(\mathcal{A}(A,B)=0) = 1 - o(1).$$
(4.1)

Using Theorem 4.1 with K = 2, we see that $\mu_{\text{IndSBM}} \bowtie_{\leq D} \mu_{\text{IndER}}$ for $D = n^{o(1)}$. Thus

$$\mu_{\mathsf{IndER}}(\mathcal{A}(A,B)=0) = 1 - o(1)$$

Thus, this algorithm \mathcal{A} strongly distinguish μ_{CorSBM} and μ_{IndER} . This contradicts Conjecture 2.2 and the low-degree hardness established in [9, Theorem 1.3].

The rest part of this section is devoted to the proof of Theorem 4.1. For notational simplicity, in the following we will only prove the case where K = 1 and the proof for general K is similar. Note that $\epsilon^2 \lambda < 1$ and $\lambda > \lambda_0$ implies that $\epsilon < \epsilon_0 = \lambda_0^{-1/2}$. We choose λ_0 to be a sufficient large constant such that

$$(\epsilon^2 \lambda) \cdot \frac{(k-1)\sqrt{1-\epsilon} + \sqrt{1+\epsilon(k-1)}}{k} \ge 1 - \delta/2 \text{ for all } \epsilon < \epsilon_0 = \lambda_0^{-1/2}.$$
(4.2)

In the rest part of this section we will always assume that

$$D = n^{o(1)}, \lambda > \lambda_0 \text{ and } \epsilon^2 \lambda < 1 - \delta \text{ for some constant } 0 < \delta < 0.01.$$
(4.3)

Clearly, using Theorem 2.4, it suffices to show that under (4.3) and λ_0 we have

$$\operatorname{\mathsf{Adv}}_{\leq D}\left(\frac{\mathrm{d}\mathbb{P}}{\mathrm{d}\mathbb{Q}}\right) = O(1) \text{ and } \operatorname{\mathsf{Adv}}_{\leq D}\left(\frac{\mathrm{d}\mathbb{Q}}{\mathrm{d}\mathbb{P}}\right) = O(1).$$

Indeed, it has been shown in [32] that $\operatorname{Adv}_{\leq D}\left(\frac{\mathrm{d}\mathbb{P}}{\mathrm{d}\mathbb{Q}}\right) = O(1)$ provided with (4.3). It remains to show that under (4.3) we have (note that now \mathbb{P} is the planted measure)

$$\mathsf{Adv}_{\leq D}\left(\frac{\mathrm{d}\mathbb{Q}}{\mathrm{d}\mathbb{P}}\right) = \sup_{f \in \mathcal{P}_D} \frac{\mathbb{E}_{\mathbb{Q}}[f]}{\sqrt{\mathbb{E}_{\mathbb{P}}[f^2]}} = O_{\delta}(1) \,. \tag{4.4}$$

We point out that our approach to proving (4.4) is based on the the work [52]. While the authors of [52] mainly focus on showing the low-degree hardness for community recovery problems, extending their approach to bound the left hand side of (4.4) is relatively standard. To this end, define

$$\omega(\sigma_i, \sigma_j) = \begin{cases} k - 1, & \sigma_i = \sigma_j; \\ -1, & \sigma_i \neq \sigma_j \end{cases}$$
(4.5)

In addition, for all $S \in \mathcal{K}_n$ define

$$\phi_S(\{G_{i,j}\}) = \prod_{(i,j)\in E(S)} \frac{G_{i,j} - \frac{\lambda}{n}}{\sqrt{\frac{\lambda}{n}(1 - \frac{\lambda}{n})}}.$$
(4.6)

It is well known in [32] that $\{\phi_S : S \in \mathcal{K}_n, |E(S)| \leq D\}$ constitutes a standard orthogonal basis of \mathcal{P}_D under \mathbb{Q} . Thus, each $f \in \mathcal{P}_D$ can be written as

$$f(G) = \sum_{S \in \mathcal{K}_n, |E(S)| \le D} \widehat{f}_S \cdot \phi_S(G) , \qquad (4.7)$$

which means that f is uniquely characterized by a vector \hat{f} indexed by $\{S \in \mathcal{K}_n : |E(S)| \leq D\}$. In addition, direct calculation yields that

$$\mathbb{E}_{\mathbb{Q}}\left[\phi_{S}(G)\right] = \mathbf{1}_{\{S=\emptyset\}}.$$
(4.8)

Thus, we have

$$\mathbb{E}_{\mathbb{Q}}[f] \stackrel{(4.7)}{=} \widehat{f}_{\emptyset} = \langle \widehat{f}, c \rangle, \qquad (4.9)$$

where c is a vector indexed by $\{S \in \mathcal{K}_n : |E(S)| \le D\}$ with

$$c_S = \mathbf{1}_{\{S = \emptyset\}} \,. \tag{4.10}$$

We now turn to $\mathbb{E}_{\mathbb{P}}[f^2]$. For any $\sigma \in [k]^n$ and $S \Subset \mathcal{K}_n$, define

$$\psi_{\sigma,S}(\{G_{i,j}\}) = k^{\frac{n}{2}} \mathbf{1}_{\sigma_* = \sigma} \cdot \prod_{(i,j) \in E(S)} \frac{G_{i,j} - \frac{(1 + \epsilon \omega(\sigma_i, \sigma_j))\lambda}{n}}{\sqrt{\frac{(1 + \epsilon \omega(\sigma_i, \sigma_j))\lambda}{n} (1 - \frac{(1 + \epsilon \omega(\sigma_i, \sigma_j))\lambda}{n})}}$$
(4.11)

We can check that $\{\psi_{\sigma,S} : \sigma \in [k]^n, S \in \mathcal{K}_n\}$ is standard orthogonal under \mathbb{P}_* , i.e., we have

$$\mathbb{E}_{\mathbb{P}_*}\left[\psi_{\sigma,S}\psi_{\sigma',S'}\right] = \mathbf{1}_{\{\sigma=\sigma',S=S'\}}.$$
(4.12)

Lemma 4.3. We have

$$\mathbb{E}_{\mathbb{P}}\left[\phi_{S}(G)\psi_{\sigma,H}(G)\right] = \frac{\mathbf{1}_{H\subset S}}{k^{\frac{n}{2}}} \prod_{(i,j)\in E(H)} h(\sigma_{i},\sigma_{j}) \prod_{(i,j)\in E(S)\setminus E(H)} \omega(\sigma_{i},\sigma_{j})\sqrt{\frac{\epsilon^{2}\lambda}{n}}, \quad (4.13)$$

where

$$\mathbf{h}(\sigma_i, \sigma_j) = \sqrt{\frac{\left(1 - \frac{(1 + \epsilon\omega(\sigma_i, \sigma_j))\lambda}{n}\right)\left(1 + \epsilon\omega(\sigma_i, \sigma_j)\right)}{1 - \frac{\lambda}{n}}}.$$
(4.14)

Proof. Note that using (4.6) and (4.11), we have that $\mathbb{E}_{\mathbb{P}}[\phi_S(G)\psi_{\sigma,H}(G)]$ equals

$$\mathbb{E}_{\mathbb{P}}\left[k^{\frac{n}{2}}\mathbf{1}_{\sigma_{*}=\sigma}\cdot\prod_{(i,j)\in E(S)}\frac{G_{i,j}-\frac{\lambda}{n}}{\sqrt{\frac{\lambda}{n}(1-\frac{\lambda}{n})}}\cdot\prod_{(i,j)\in E(H)}\frac{G_{i,j}-\frac{(1+\epsilon\omega(\sigma_{i},\sigma_{j}))\lambda}{n}}{\sqrt{\frac{(1+\epsilon\omega(\sigma_{i},\sigma_{j}))\lambda}{n}(1-\frac{(1+\epsilon\omega(\sigma_{i},\sigma_{j}))\lambda}{n})}}\right]$$
$$=k^{-\frac{n}{2}}\mathbb{E}_{\mathbb{P}_{\sigma}}\left[\prod_{(i,j)\in E(S)}\frac{G_{i,j}-\frac{\lambda}{n}}{\sqrt{\frac{\lambda}{n}(1-\frac{\lambda}{n})}}\cdot\prod_{(i,j)\in E(H)}\frac{G_{i,j}-\frac{(1+\epsilon\omega(\sigma_{i},\sigma_{j}))\lambda}{n}}{\sqrt{\frac{(1+\epsilon\omega(\sigma_{i},\sigma_{j}))\lambda}{n}(1-\frac{(1+\epsilon\omega(\sigma_{i},\sigma_{j}))\lambda}{n})}}\right].$$
(4.15)

Clearly, when $H \not\subset S$ we have (4.15) cancels to 0, since (denote (i, j) to be an element in $E(H) \setminus E(S)$) under \mathbb{P}_{σ} we have $\mathbb{E}_{\mathbb{P}_{\sigma}}[G_{i,j} - \frac{(1+\epsilon\omega(\sigma_i,\sigma_j))\lambda}{n}] = 0$ and all the edges $\{G_{i,j} : (i, j) \in U_n\}$ are independent. In addition, from direct calculation we have that

$$\mathbb{E}_{\mathbb{P}_{\sigma}}\left[G_{i,j}-\frac{\lambda}{n}\right] = \frac{\epsilon\lambda\omega(\sigma_{i},\sigma_{j})}{n},\\ \mathbb{E}_{\mathbb{P}_{\sigma}}\left[(G_{i,j}-\frac{\lambda}{n})(G_{i,j}-\frac{(1+\epsilon\omega(\sigma_{i},\sigma_{j}))\lambda}{n})\right] = \frac{(1+\epsilon\omega(\sigma_{i},\sigma_{j}))\lambda}{n}\left(1-\frac{(1+\epsilon\omega(\sigma_{i},\sigma_{j}))\lambda}{n}\right).$$

Plugging this result into (4.15), we get (4.13) and thus completes the proof.

Based on Lemma 4.3, we define a matrix M with rows indexed by $\{S : S \in \mathcal{K}_n, |E(S)| \leq D\}$ and columns indexed by $\{(\sigma, S) : \sigma \in [k]^n, S \in \mathcal{K}_n, |E(S)| \leq D\}$, and entries given by

$$M_{S;(\sigma,H)} = \frac{\mathbf{1}_{H \subset S}}{k^{\frac{n}{2}}} \prod_{(i,j) \in E(H)} \mathbf{h}(\sigma_i, \sigma_j) \prod_{(i,j) \in E(S) \setminus E(H)} \omega(\sigma_i, \sigma_j) \sqrt{\frac{\epsilon^2 \lambda}{n}}.$$
 (4.16)

From Parserval's inequality, we see that

$$\mathbb{E}_{\mathbb{P}}[f^{2}] \geq \sum_{\substack{\sigma \in [k]^{n}, H \in \mathcal{K}_{n} \\ |E(H)| \leq D}} \mathbb{E}_{\mathbb{P}}[f \cdot \psi_{\sigma,H}]^{2} \stackrel{(4.7)}{=} \sum_{\substack{\sigma \in [k]^{n}, H \in \mathcal{K}_{n} \\ |E(H)| \leq D}} \left(\sum_{\substack{S \in \mathcal{K}_{n} \\ |E(S)| \leq D}} \widehat{f}_{S} \mathbb{E}_{\mathbb{P}}[\phi_{S} \cdot \psi_{\sigma,H}]\right)^{2}$$
$$= \sum_{\substack{\sigma \in [k]^{n}, H \in \mathcal{K}_{n} \\ |E(H)| \leq D}} \left(\sum_{\substack{S \in \mathcal{K}_{n} \\ |E(S)| \leq D}} \widehat{f}_{S} M_{S;(\sigma,H)}\right)^{2} = \|\widehat{f}M\|^{2}.$$
(4.17)

Thus, we have

$$\mathsf{Adv}_{\leq D}\left(\frac{\mathrm{d}\mathbb{Q}}{\mathrm{d}\mathbb{P}}\right) = \sup_{f \in \mathcal{P}_D} \left\{ \frac{\mathbb{E}_{\mathbb{Q}}[f]}{\sqrt{\mathbb{E}_{\mathbb{P}}[f^2]}} \right\} \leq \sup_{\widehat{f}} \left\{ \frac{\langle \widehat{f}, c \rangle}{\|\widehat{f}M\|} \right\} \leq \inf_{Mu^{\top} = c} \left\{ \|u\| \right\},\tag{4.18}$$

where the last inequality follows from the fact that for $Mu_0^{\top} = c$ we have

$$\langle \hat{f}, c \rangle = \langle \hat{f}, M u_0^\top \rangle = \langle \hat{f} M, u_0 \rangle \le \| u_0 \| \cdot \| \hat{f} M \|$$

Regarding (4.18), it suffices to show that there exists $Mu^{\top} = c$ and ||u|| = O(1). Note that $Mu^{\top} = c$ is equivalent to

$$\sum_{\sigma \in [k]^n} \sum_{H \subset S} \frac{u_{\sigma,H}}{k^{\frac{n}{2}}} \prod_{(i,j) \in E(H)} h(\sigma_i, \sigma_j) \prod_{(i,j) \in E(S) \setminus E(H)} \omega(\sigma_i, \sigma_j) \sqrt{\frac{\epsilon^2 \lambda}{n}} = \mathbf{1}_{S = \emptyset} .$$
(4.19)

The first step of our analysis is to simplify (4.19). Based on Lemma A.8, we see that for all $H \subset S$ with $\mathcal{L}(S) \setminus V(H) \neq \emptyset$, we have (denote $\mathbf{V} = V(H)$)

$$\begin{split} &\sum_{\sigma \in [k]^n} u_{\sigma,H} \prod_{(i,j) \in E(H)} \mathbf{h}(\sigma_i,\sigma_j) \prod_{(i,j) \in E(S) \setminus E(H)} \omega(\sigma_i,\sigma_j) \sqrt{\frac{\epsilon^2 \lambda}{n}} \\ &= \left(\frac{\epsilon^2 \lambda}{n}\right)^{\frac{|E(S)| - |E(H)|}{2}} \sum_{\sigma_{\mathbf{V}} \in [n]^{\mathbf{V}}} \prod_{(i,j) \in E(H)} \mathbf{h}(\sigma_i,\sigma_j) \times \sum_{\sigma_{[n] \setminus \mathbf{V}} \in [k]^{[n] \setminus \mathbf{V}}} \prod_{(i,j) \in E(S) \setminus E(H)} \omega(\sigma_i,\sigma_j) = 0 \,. \end{split}$$

Thus, (4.19) can be further simplified to

$$\sum_{\sigma \in [k]^n} \sum_{\substack{H \subset S \\ \mathcal{L}(S) \subset V(H)}} \frac{u_{\sigma,H}}{k^{\frac{n}{2}}} \prod_{(i,j) \in E(H)} h(\sigma_i, \sigma_j) \prod_{(i,j) \in E(S) \setminus E(H)} \omega(\sigma_i, \sigma_j) \sqrt{\frac{\epsilon^2 \lambda}{n}} = \mathbf{1}_{S = \emptyset} .$$
(4.20)

We now construct the solution $\{u_{\sigma,H} : \sigma \in [k]^n, H \in \mathcal{K}_n, |E(H)| \leq D\}$ of (4.20) as follows: let $u_{\sigma,H} = \frac{1}{k^{\frac{n}{2}}} \cdot \Xi(H)$, where

$$\Xi(\emptyset) = 1 \text{ and } \Xi(S) = 0 \text{ for } \mathcal{L}(S) \neq \emptyset$$
 (4.21)

and then iteratively define for all $\mathcal{L}(S) = \emptyset$ by

$$\Xi(S) = -\left(\mathbb{E}_{\sigma \sim \nu} \left[\prod_{(i,j) \in E(S)} \mathbf{h}(\sigma_i, \sigma_j)\right]\right)^{-1} \sum_{\substack{H \subset S \\ \mathcal{L}(H) = \emptyset}} \left(\frac{\epsilon^2 \lambda}{n}\right)^{\frac{|E(S)| - |E(H)|}{2}} \Xi(H) \\ \times \mathbb{E}_{\sigma \sim \nu} \left[\prod_{(i,j) \in E(H)} \mathbf{h}(\sigma_i, \sigma_j) \prod_{(i,j) \in E(S) \setminus E(H)} \omega(\sigma_i, \sigma_j)\right],$$

$$(4.22)$$

To prove (4.4), it suffices to show the following lemma

Lemma 4.4. The vector $\{u_{\sigma,H} : \sigma \in [k]^n, H \in \mathcal{K}_n, |E(H)| \leq D\}$ satisfies (4.20) and $||u|| = O_{\delta}(1)$.

The crucial input in our proof of Lemma 4.4 is the following estimation on $\Xi(H)$.

Lemma 4.5. Assuming (4.3) with $|E(S)| \leq D$ and $\mathcal{L}(S) = \emptyset$. We have the following estimation:

- (1) If $S = S_1 \cup S_2$ such that $V(S_1) \cap V(S_2) = \emptyset$, then $\Xi(S) = \Xi(S_1)\Xi(S_2)$.
- (2) If S can be decomposed into different cycles C_1, \ldots, C_m , then

$$|\Xi(S)| \le k^m (1 - \delta/2)^{\frac{|E(H)|}{2}} n^{-\frac{|E(H)|}{2}}.$$

(3) If $\tau(S) > 0$ and $\mathfrak{C}_k(S) = m_k$ for $3 \le k \le D$, then

$$|\Xi(S)| \le (10\tau(S))! \cdot (2kD)^{10\tau(H)} (1-\delta/2)^{\frac{|E(H)|}{2}} n^{-\frac{|E(H)|}{2}}.$$

Proof. We first prove Item (1). We will prove by induction on |E(S)|. The case $S = \emptyset$ is trivial. Now Suppose $S = S_1 \cup S_2$, for all $H \subset S$ with $\mathcal{L}(H) = \emptyset$, we have $H = H_1 \cup H_2$ with $H_1 \subset S_1, H_2 \subset S_2$ and $\mathcal{L}(H_1), \mathcal{L}(H_2) = \emptyset$. In addition, denote

$$\begin{split} \mathbf{P}(S) &= \mathbb{E}_{\sigma \sim \nu} \left[\prod_{(i,j) \in E(S)} \mathbf{h}(\sigma_i, \sigma_j) \right], \\ \mathbf{Q}(S, H) &= \mathbb{E}_{\sigma \sim \nu} \left[\prod_{(i,j) \in E(H)} \mathbf{h}(\sigma_i, \sigma_j) \prod_{(i,j) \in E(S) \setminus E(H)} \omega(\sigma_i, \sigma_j) \right]. \end{split}$$

We have $P(S) = P(S_1)P(S_2)$ and $Q(S, H) = Q(S_1, H_1)Q(S_2, H_2)$. Also, it is straightforward to verify that Q(S, S) = P(S). Thus, we have

$$\begin{split} -\Xi(S) &= \sum_{\substack{H_1 \subsetneq S_1, H_2 \subsetneq S_2\\ \mathcal{L}(H_1), \mathcal{L}(H_2) = \emptyset}} \frac{\mathbb{Q}(S_1, H_1) \mathbb{Q}(S_2, H_2) \Xi(H_1 \sqcup H_2)}{\mathbb{P}(S_1) \mathbb{P}(S_2)} + \sum_{\substack{H_2 \subsetneq S_2\\ \mathcal{L}(H_2) = \emptyset}} \frac{\mathbb{Q}(S_2, H_2) \Xi(S_1 \sqcup H_2)}{\mathbb{P}(S_2)} \\ &+ \sum_{\substack{H_1 \subsetneq S_1\\ \mathcal{L}(H_1) = \emptyset}} \frac{\mathbb{Q}(S_1, H_1) \Xi(H_1 \sqcup S_2)}{\mathbb{P}(S_1)} \\ &= \sum_{\substack{H_1 \subsetneq S_1, H_2 \subsetneq S_2\\ \mathcal{L}(H_1), \mathcal{L}(H_2) = \emptyset}} \frac{\mathbb{Q}(S_1, H_1) \mathbb{Q}(S_2, H_2) \Xi(H_1) \Xi(H_2)}{\mathbb{P}(S_1) \mathbb{P}(S_2)} + \sum_{\substack{H_2 \subseteq S_2\\ \mathcal{L}(H_2) = \emptyset}} \frac{\mathbb{Q}(S_2, H_2) \Xi(S_1) \Xi(H_2)}{\mathbb{P}(S_2)} \\ &+ \sum_{\substack{H_1 \subsetneq S_1\\ \mathcal{L}(H_1) = \emptyset}} \frac{\mathbb{Q}(S_1, H_1) \Xi(H_1) \Xi(S_2)}{\mathbb{P}(S_1)} \\ &= (-\Xi(S_1)) \cdot (-\Xi(S_2)) + \Xi(S_1) \cdot (-\Xi(S_2)) + (-\Xi(S_1)) \cdot \Xi(S_2) = -\Xi(S_1) \Xi(S_2) \,, \end{split}$$

where the second equality follows from induction hypothesis and the third equality follows from (4.22). This yields Item (1).

Now we focus on Item (2). Based on Item (1). it suffices to show that for any cycle C with $|E(C)| \leq D$ we have

$$|\Xi(C)| \le k(1-\delta)^{\frac{|E(C)|}{2}} n^{-\frac{|E(C)|}{2}}.$$
(4.23)

Note that using (4.21) and (4.22), we have that

$$\Xi(C) = \tfrac{\mathbb{Q}(S,\emptyset)}{\mathbb{P}(S)} \cdot \left(\tfrac{\epsilon^2\lambda}{n}\right)^{|E(C)|/2} = \tfrac{(k-1)}{\mathbb{P}(S)} \cdot (\epsilon^2\lambda)^{|E(C)|/2} n^{-|E(C)|/2} \,,$$

where in the second equality we use Lemma A.7. Note that since $\langle \sigma_i, \sigma_j \rangle \in \{-1, k-1\}$, we have

$$\mathbf{h}(\sigma_i,\sigma_j) = a\omega(\sigma_i,\sigma_j) + b\,,$$

where

$$a = [1 + o(1)] \cdot \frac{(k-1)\sqrt{1-\epsilon} + \sqrt{1+\epsilon(k-1)}}{k}$$
 and $b = [1 + o(1)] \cdot \frac{\sqrt{1+\epsilon(k-1)} - \sqrt{1-\epsilon}}{k}$

Thus, using Lemma A.7 we get that

$$\mathsf{P}(C) = a^{|E(C)|} + (k-1)b^{|E(C)|} \ge a^{|E(C)|} \,.$$

Plugging this result into the bound on $\Xi(C)$ yields Item (2) (recall (4.2)).

Finally we prove Item (3). Again based on Item (1), we may assume that S is connected. We prove Item (3) by induction on $\tau(S)$. The case where $\tau(S) = 0$ is proved in Item (2). Now assume Item (3) holds for $\tau(S) \leq m-1$, we now focus on $\tau(S) = m$. Using Lemma A.4, we can decompose S into t disjoint self-avoiding paths P_1, \ldots, P_t with $t \leq 3m$ satisfying Lemma A.4. In addition, applying Lemma A.4 for $H \subset S$ and $\mathcal{L}(H) = \mathcal{L}(S) = \emptyset$, we see that assuming $\tau(H) = m - r$ for some r > 0, we then have $S \setminus H = P_{i_1} \cup \ldots \cup P_{i_s}$ where $s \leq 3r$. Denote $\mathbb{V} = \bigcup_{1 \leq i \leq m} \operatorname{EndP}(P_i)$, by conditioning on $\{\sigma_u : u \in \mathbb{V}\}$ we see that

$$\Big\{\prod_{(i,j)\in E(P_{i})}\mathtt{h}(\sigma_{i},\sigma_{j}):1\leq\mathtt{i}\leq m\Big\}$$

are conditional independent, with (denote $EndP(P_i) = \{u_i, v_i\}$)

$$\mathbb{E}\Big[\prod_{(i,j)\in E(P_{\mathtt{i}})}\mathtt{h}(\sigma_i,\sigma_j)\mid \{\sigma_u: u\in \mathtt{V}\}\Big]=a^{|E(P_{\mathtt{i}})|}+b^{|E(P_{\mathtt{i}})|}\omega(u_{\mathtt{i}},v_{\mathtt{i}})\,.$$

Thus, we have

$$\begin{split} \mathbf{P}(S) &= \mathbb{E}_{\sigma_{\mathbf{V}} \sim \nu_{\mathbf{V}}} \mathbb{E}_{\sigma_{\mathbf{V}} \sim \nu_{\mathbf{V}}} \left[\prod_{1 \leq \mathbf{i} \leq \mathbf{t}} \prod_{(i,j) \in E(P_{\mathbf{i}})} \mathbf{h}(\sigma_{i},\sigma_{j}) \mid \{\sigma_{u} : u \in \mathbf{V}\} \right] \\ &= \mathbb{E}_{\sigma_{\mathbf{V}} \sim \nu_{\mathbf{V}}} \left[\prod_{1 \leq \mathbf{i} \leq \mathbf{t}} \prod_{\{u_{\mathbf{i}}, v_{\mathbf{i}}\} = \mathrm{EndP}(P_{\mathbf{i}})} \left(a^{|E(P_{\mathbf{i}})|} + b^{|E(P_{\mathbf{i}})|} \omega(u_{\mathbf{i}}, v_{\mathbf{i}}) \right) \right]. \end{split}$$

Similarly we can show that (denote $\Lambda = \{i_1, \ldots, i_s\}$)

$$\begin{aligned} \mathsf{Q}(S,H) &= \mathbb{E}_{\sigma_{\mathbf{V}} \sim \nu_{\mathbf{V}}} \left[\prod_{\mathbf{i} \in [\mathbf{t}] \setminus \Lambda} \prod_{\{u_{\mathbf{i}}, v_{\mathbf{i}}\} = \mathrm{EndP}(P_{\mathbf{i}})} \left(a^{|E(P_{\mathbf{i}})|} + b^{|E(P_{\mathbf{i}})|} \omega(u_{\mathbf{i}}, v_{\mathbf{i}}) \right) \prod_{\mathbf{i} \in \Lambda} \prod_{\{u_{\mathbf{i}}, v_{\mathbf{i}}\} = \mathrm{EndP}(P_{\mathbf{i}})} \prod_{\mathbf{i} \in \Lambda} \sum_{\{u_{\mathbf{i}}, v_{\mathbf{i}}\} = \mathrm{EndP}(P_{\mathbf{i}})} \left[\prod_{1 \leq \mathbf{i} \leq \mathbf{t}} \prod_{\{u_{\mathbf{i}}, v_{\mathbf{i}}\} = \mathrm{EndP}(P_{\mathbf{i}})} \left(a^{|E(P_{\mathbf{i}})|} + b^{|E(P_{\mathbf{i}})|} \omega(u_{\mathbf{i}}, v_{\mathbf{i}}) \right) \right] \\ &\leq k^{s} a^{-(|E(S)| - |E(H)|)} \mathsf{P}(S) \,. \end{aligned}$$

Thus, we have that

$$\frac{(\epsilon^2 \lambda)^{|E(S)| - |E(H)|} \mathbb{Q}(S, H)}{\mathbb{P}(S)} \leq (2k)^s (\epsilon^2 \lambda)^{|E(S)| - |E(H)|} a^{|E(S)| - |E(H)|} \leq (2k)^{3r} (1 - \delta/2)^{|E(S)| - |E(H)|}.$$

Thus, we have (combining the induction hypothesis on H)

$$\begin{split} |\Xi(S)| &\leq (1-\delta/2)^{|E(S)|} n^{-|E(S)|/2} \sum_{1 \leq r \leq m} (2k)^{3r} (10(m-r))! \# \{ H \subset S : \mathcal{L}(H) = \emptyset \} \\ &\leq (1-\delta/2)^{|E(S)|} n^{-|E(S)|/2} \sum_{1 \leq r \leq m} (2k)^{3r} (10(m-r))! D^{3r} \\ &\leq (10m)! (2kD)^{10m} (1-\delta/2)^{\frac{|E(S)|}{2}} n^{-\frac{|E(S)|}{2}}, \end{split}$$

where in the second inequality we use Lemma A.6 with $m_j = 0$, leading to Item (3).

Now we prove Lemma 4.4 based on Lemma 4.5.

Proof of Lemma 4.4. It is easy to check that u satisfies (4.20). We need to show that

$$\sum_{\sigma \in [k]^n} \sum_{H \in \mathcal{K}_n, |E(H)| \le D} u_{\sigma,H}^2 = \sum_{\substack{H \in \mathcal{K}_n, \mathcal{L}(H) = \emptyset\\|E(H)| \le D}} \Xi(H)^2$$
(4.24)

is bounded by $O_{\delta}(1)$. Using Lemma 4.5, we see that (4.24) is bounded by

$$\sum_{\substack{m_3,\dots,m_D \ge 0,t \ge 0\\v \ge 3m_3+\dots+Dm_D}} 2^{10k+m_3+\dots+m_D} k^{10t} n^{-(v+t)} (1-\delta/2)^{v+t} \cdot \mathsf{ENUM}'(m_3,\dots,m_D;t) , \quad (4.25)$$

where

$$\mathsf{ENUM}'(m_3,\ldots,m_D;t) = \# \left\{ S \Subset \mathcal{K}_n : \mathcal{L}(S) = \emptyset, \mathfrak{C}_k(S) = m_k \text{ for } k \le D, \tau(S) = t \right\}.$$
(4.26)

Using Lemma A.5 with $H = \emptyset$, we see that

$$\mathsf{ENUM}'(m_3, \dots, m_D; t) \le n^v D^{10t} \prod_{3 \le i \le D} \frac{1}{m_i!} \,. \tag{4.27}$$

Plugging (4.27) into (4.25), we get that

$$\begin{aligned} (4.25) &\leq \sum_{\substack{m_3, \dots, m_D \geq 0, t \geq 0 \\ v \geq 3m_3 + \dots + Dm_D}} 2^{10t + m_3 + \dots + m_D} k^{10t} n^{-(v+t)} (1 - \delta/2)^{v+t} \cdot n^v D^{10t} \prod_{3 \leq i \leq D} \frac{1}{i^{m_i} m_i!} \\ &\leq \sum_{\substack{m_3, \dots, m_D \geq 0, t \geq 0 \\ v \geq 3m_3 + \dots + Dm_D}} 2^{m_3 + \dots + m_D} (2^{10} D^{10} k/n)^{10t} (1 - \delta/2)^v \prod_{3 \leq i \leq D} \frac{1}{m_i!} \\ &\leq [1 + o(1)] \sum_{\substack{m_3, \dots, m_D \geq 0 \\ v \geq 3m_3 + \dots + Dm_D}} 2^{m_3 + \dots + m_D} (1 - \delta/2)^v \prod_{3 \leq i \leq D} \frac{1}{i^{m_i} m_i!} \\ &\leq O_{\delta}(1) \cdot \sum_{m_3, \dots, m_D \geq 0} 2^{m_3 + \dots + m_D} \prod_{3 \leq i \leq D} \frac{1}{i^{m_i} m_i!} = O_{\delta}(1) \,. \end{aligned}$$

This concludes the desired result.

Acknowledgment. The author thanks Hang Du for pointing him to the reference [52]; the author thanks Jingqiu Ding for stimulating discussions on the strengthening of low-degree conjecture. This work is partially supported by National Key R&D program of China (No. 2023YFA1010103) and NSFC Key Program (Project No. 12231002).

A Preliminary results in graphs and probability

In this section we include some preliminary results we will use in the main part of the paper. We first state some results on the structural properties of graphs established in [15, 9].

Lemma A.1 ([15], Lemma A.1). Let $S, T \in \mathcal{K}_n$ satisfy $S \cong \mathbf{S}$ and $T \cong \mathbf{T}$ for some $\mathbf{S}, \mathbf{T} \in \mathcal{H}$. Recall $S \cup T, S \cap T \in \mathcal{K}_n$ defined as edge-induced graphs of \mathcal{K}_n . We have the following hold:

- (i) $|V(S \cup T)| + |V(S \cap T)| \le |V(S)| + |V(T)|$, and $|E(S \cup T)| + |E(S \cap T)| = |E(S)| + |E(T)|$.
- (*ii*) $\Phi(S \cup T)\Phi(S \cap T) \le \Phi(S)\Phi(T)$.
- (iii) If $\mathbf{S} \subset \mathbf{T}$, then $|\operatorname{Aut}(\mathbf{S})| \leq |\operatorname{Aut}(\mathbf{T})| \cdot |V(\mathbf{T})|^{2(|E(\mathbf{T})| |E(\mathbf{S})|)}$.

$$(iv) # \{T' \in \mathcal{K}_n : S \subset T', |V(T')| - |V(S)| = k, |E(T')| - |E(S)| = l\} \le n^k (|V(S)| + k)^{2l}.$$

 $(v) \ \#\{T' \subset S : |E(S)| - |E(T')| = k\} \le |E(S)|^k.$

Lemma A.2. Let $S, T \subset \mathcal{K}_n$. Recall that $S \cap T \subseteq \mathcal{K}_n$ is defined as edge-induced subgraphs of \mathcal{K}_n . We have the following properties:

- (i) $\tau(S \cup T) + \tau(S \cap T) \ge \tau(S) + \tau(T)$ and $\Upsilon(S \cup T)\Upsilon(S \cap T) \le \Phi(S)\Phi(T)$.
- (ii) $|\mathsf{C}_j(S \cup T)| + |\mathsf{C}_j(S \cap T)| \ge |\mathsf{C}_j(S)| + |\mathsf{C}_j(T)|.$
- (iii) If $S \subset T$, S is self-bad and V(S) = V(T), then T is self-bad.
- (iv) If S and T are both self-bad, then $S \cup T$ is self-bad.

Lemma A.3 ([9], Lemma A.3). For $H \subset S$, we can decompose $E(S) \setminus E(H)$ into m cycles C_1, \ldots, C_m and t paths P_1, \ldots, P_t for some $m, t \ge 0$ such that the following holds:

- (1) C_1, \ldots, C_m are vertex-disjoint (i.e., $V(C_i) \cap V(C_j) = \emptyset$ for all $i \neq j$) and $V(C_i) \cap V(H) = \emptyset$ for all $1 \leq i \leq m$.
- (2) EndP(P_j) $\subset V(H) \cup (\cup_{i=1}^{m} V(C_i)) \cup (\cup_{k=1}^{j-1} V(P_k)) \cup \mathcal{L}(S)$ for $1 \leq j \leq t$.

(3) $(V(P_{j})\setminus \operatorname{EndP}(P_{j})) \cap (V(H)\cup (\cup_{i=1}^{m}V(C_{i}))\cup (\cup_{k=1}^{j-1}V(P_{k}))\cup \mathcal{L}(S)) = \emptyset \text{ for } 1 \leq j \leq t.$ (4) $t = |\mathcal{L}(S)\setminus V(H)| + \tau(S) - \tau(H).$

Lemma A.4 ([9], Corollary A.4). For $H \subset S$, we can decompose $E(S) \setminus E(H)$ into m cycles C_1, \ldots, C_m and t paths P_1, \ldots, P_t for some $m, t \ge 0$ such that the following hold:

- (1) C_1, \ldots, C_m are independent cycles in S.
- (2) $V(P_{j}) \cap \left(V(H) \cup \left(\bigcup_{i=1}^{m} V(C_{i})\right) \cup \left(\bigcup_{k \neq j} V(P_{k})\right) \cup \mathcal{L}(S)\right) = \operatorname{EndP}(P_{j}) \text{ for } 1 \leq j \leq t.$ (3) $t \leq 5(|\mathcal{L}(S) \setminus V(H)| + \tau(S) - \tau(H)).$

Lemma A.5 ([9], Lemma A.9). For $H \subset \mathcal{K}_n$, we have (below we write $\mathfrak{P} = \{(p_{N+1}, \ldots, p_D) : \sum_{i=N+1}^{D} p_i \leq p, \text{ for all } N+1 \leq l \leq D\}$ and $\sum_{\mathfrak{P}}$ for the summation over $(p_{N+1}, \ldots, p_D) \in \mathfrak{P}$)

$$\# \Big\{ S \ admissible: H \ltimes S; |\mathfrak{C}_{l}(S,H)| = 0 \ for \ l > N; |\mathcal{L}(S) \setminus V(H)| + \tau(S) - \tau(H) = m; \\ |E(S)| - |E(H)| = p, |V(S)| - |V(H)| = q, |E(S)| \le D \Big\} \le (2D)^{3m} n^{q} \sum_{\mathfrak{P}} \prod_{j=N+1}^{D} \frac{1}{p_{j}!}.$$
(A.1)

Lemma A.6 ([9], Lemma A.10). For $S \subset \mathcal{K}_n$ with $|E(S)| \leq D$, we have

$$#\left\{H: H \ltimes S, |\mathcal{L}(S) \setminus V(H)| + \tau(S) - \tau(H) = m, \\ \mathfrak{C}_{j}(S; H) = m_{j}, N + 1 \le j \le D\right\} \le D^{15m} \prod_{j=N+1}^{D} \binom{|\mathcal{C}_{j}(S)|}{m_{j}}.$$
(A.2)

We now show some preliminary results on calculating expectations under $\sigma \sim \nu$.

Lemma A.7. For a path \mathcal{P} with $V(\mathcal{P}) = \{v_0, \ldots, v_l\}$ and $\operatorname{EndP}(\mathcal{P}) = \{v_0, v_l\}$, we have

$$\mathbb{E}_{\sigma \sim \nu} \Big[\prod_{i=1}^{l} \Big(a + b\omega(\sigma_{i-1}, \sigma_i) \Big) \mid \sigma_0, \sigma_l \Big] = a^l + b^l \cdot \omega(\sigma_0, \sigma_l) \,. \tag{A.3}$$

Proof. By independence, we see that $\mathbb{E}_{\sigma \sim \nu} \left[\prod_{i \in I} \omega(\sigma_{i-1}, \sigma_i) \mid \sigma_0, \sigma_l \right] = 0$ if $I \subsetneq [l]$. Thus,

$$\mathbb{E}_{\sigma \sim \nu} \Big[\prod_{i=1}^{l} \Big(a + b\omega(\sigma_{i-1}, \sigma_i) \Big) \mid \sigma_0, \sigma_l \Big] = a^l + b^l \mathbb{E}_{\sigma \sim \nu} \Big[\prod_{i=1}^{l} \omega(\sigma_{i-1}, \sigma_i) \mid \sigma_0, \sigma_l \Big].$$

It remains to prove that

$$\mathbb{E}_{\sigma \sim \nu} \left[\prod_{i=1}^{l} \omega(\sigma_{i-1}, \sigma_i) \mid \sigma_0, \sigma_l \right] = \omega(\sigma_0, \sigma_l).$$
(A.4)

We shall show (A.4) by induction. The case l = 1 follows immediately. Now we assume that (A.4) holds for l. Then we have

$$\mathbb{E}_{\sigma \sim \nu} \Big[\prod_{i=1}^{l+1} \omega(\sigma_{i-1}, \sigma_i) \mid \sigma_0, \sigma_{l+1} \Big]$$

= $\mathbb{E}_{\sigma \sim \nu} \Big[\omega(\sigma_l, \sigma_{l+1}) \mathbb{E}_{\sigma \sim \nu} \Big[\prod_{i=1}^l \omega(\sigma_{i-1}, \sigma_i) \mid \sigma_0, \sigma_l, \sigma_{l+1} \Big] \mid \sigma_0, \sigma_{l+1} \Big]$
= $\mathbb{E}_{\sigma \sim \nu} \Big[\omega(\sigma_l, \sigma_{l+1}) \omega(\sigma_0, \sigma_l) \mid \sigma_0, \sigma_{l+1} \Big] = \omega(\sigma_0, \sigma_{l+1}),$

which completes the induction procedure.

Lemma A.8. For $H \subset S$ with $\mathcal{L}(S) \not\subset V(H)$, we have

$$\mathbb{E}_{\sigma \sim \nu} \left[\prod_{(i,j) \in E(S) \setminus E(H)} \omega(\sigma_i, \sigma_j) \mid \{ \sigma_u : u \in V(H) \} \right] = 0.$$
(A.5)

Proof. Denote $v \in \mathcal{L}(S) \setminus V(H)$ and $(v, w) \in E(S)$. Define σ_U and $\sigma_{\setminus U}$ to be the restriction of σ on U and on $[n] \setminus U$, respectively. Also define ν_U and $\nu_{\setminus U}$ to be the restriction of ν on U and on $[n] \setminus U$, respectively. Then we have (let $\mathbb{V} = V(H)$)

$$\mathbb{E}_{\sigma\sim\nu} \left[\prod_{(i,j)\in E(S)\setminus E(H)} \omega(\sigma_i,\sigma_j) \mid \{\sigma_u : u \in V(H)\} \right]$$

= $\mathbb{E}_{\sigma_{[n]\setminus \mathbb{V}}\sim\nu_{\mathbb{V}}} \left[\prod_{(i,j)\in E(S)\setminus E(H)} \omega(\sigma_i,\sigma_j) \right]$
= $\mathbb{E}_{\sigma_{[n]\setminus (\mathbb{V}\cup \{v\})}\sim\nu_{\mathbb{V}}} \mathbb{E}_{\sigma_{\{v\}}\sim\nu_{\{v\}}} \left[\omega(\sigma_v,\sigma_w) \prod_{(i,j)\in E(S)\setminus (E(H)\cup \{v,w\})} \omega(\sigma_i,\sigma_j) \right] = 0.$

References

- L. Babai. Graph isomorphism in quasipolynomial time. In Proceedings of the 48th Annual ACM Symposium on Theory of Computing (STOC), pages 684–697. ACM, 2016.
- [2] A. S. Bandeira, D. Kunisky, and A. S. Wein. Computational hardness of certifying bounds on constrained PCA problems. In 11th Innovations in Theoretical Computer Science Conference (ITCS), volume 151. Schloss Dagstuhl-Leibniz-Zentrumfűr Informatik, 2020.

- [3] A. S. Bandeira, A. E. Alaoui, S. Hopkins, T. Schramm, A. S. Wein, and I. Zadik. The Franz-Parisi criterion and computational trade-offs in high dimensional statistics. In Advances in Neural Information Processing Systems (NIPS), volume 35, pages 33831–33844. Curran Associates, Inc., 2022.
- [4] B. Barak, C. N. Chou, Z. Lei, T. Schramm, and Y. Sheng. (Nearly) efficient algorithms for the graph matching problem on correlated random graphs. In Advances in Neural Information Processing Systems (NIPS), volume 32, pages 9190–9198. Curran Associates, Inc., 2019.
- [5] B. Barak, S. Hopkins, J. Kelner, P. K. Kothari, A. Moitra, and A. Potechin. A nearly tight sum-of-squares lower bound for the planted clique problem. In SIAM Journal on Computing, 48(2):687–735, 2019.
- [6] G. Bresler and B. Huang. The algorithmic phase transition of random k-SAT for low degree polynomials. In *IEEE 62nd Annual Symposium on Foundations of Computer Science (FOCS)*, pages 298–309. IEEE, 2022.
- [7] R. E. Burkard, E. Cela, P. M. Pardalos, and L. S. Pitsoulis. The quadratic assignment problem. In Handbook of combinatorial optimization, pages 1713–1809. Springer, 1998.
- [8] S. Chai and M. Z. Racz. Efficient graph matching for correlated stochastic block models. In Advances in Neural Information Processing Systems (NIPS), volume 37. Curran Associates, Inc., 2024.
- [9] G. Chen, J. Ding, S. Gong, and Z. Li. A computational transition for detecting correlated stochastic block models by low-degree polynomials. arXiv preprint, arXiv:2409.00966.
- [10] D. Cullina and N. Kiyavash. Exact alignment recovery for correlated Erdős-Rényi graphs. arXiv preprint, arXiv:1711.06783.
- [11] O. E. Dai, D. Cullina, N. Kiyavash, and M. Grossglauser. Analysis of a canonical labeling algorithm for the alignment of correlated Erdős-Rényi graphs. In *Proceedings of the ACM on Measurement and Analysis of Computing Systems*, 3(2):1–25, 2019.
- [12] A. Dhawan, C. Mao, and A. S. Wein. Detection of dense subhypergraphs by low-degree polynomials. to appear in *Random Structures and Algorithms*.
- [13] J. Ding and H. Du. Detection threshold for correlated Erdős-Rényi graphs via densest subgraph. In *IEEE Transactions on Information Theory*, 69(8):5289–5298, 2023.
- [14] J. Ding and H. Du. Matching recovery threshold for correlated random graphs. In Annals of Statistics, 51(4): 1718–1743, 2023.
- [15] J. Ding, H. Du, and Z. Li. Low-degree hardness of detection for correlated Erdős-Rényi Graphs. arXiv preprint, arXiv:2311.15931.
- [16] J. Ding and Z. Li. A polynomial time iterative algorithm for matching Gaussian matrices with non-vanishing correlation. To appear in *Foundations of Computational Mathematics*.
- [17] J. Ding and Z. Li. A polynomial-time iterative algorithm for random graph matching with non-vanishing correlation. arXiv preprint, arXiv:2306.00266.
- [18] J. Ding, Z. Ma, Y. Wu, and J. Xu. Efficient random graph matching via degree profiles. In Probability Theory and Related Fields, 179(1-2):29–115, 2021.

- [19] Y. Ding, D. Kunisky, A. S. Wein, and A. S. Bandeira. Subexponential-time algorithms for sparse PCA. In Foundations of Computational Mathematics, 22(1):1–50, 2022.
- [20] Z. Fan, C. Mao, Y. Wu, and J. Xu. Spectral graph matching and regularized quadratic relaxations I: Algorithm and theory. In *Foundations of Computational Mathematics*, 23(5):1511– 1565, 2023.
- [21] Z. Fan, C. Mao, Y. Wu, and J. Xu. Spectral graph matching and regularized quadratic relaxations II: Erdős-Rényi graphs and universality. In *Foundations of Computational Mathematics*, 23(5):1567–1617, 2023.
- [22] D. Gamarnik. The overlap gap property: A topological barrier to optimizing over random structures. In Proceedings of the National Academy of Sciences of the United States of America, 118(41):e2108492118, 2021.
- [23] D. Gamarnik, A. Jagannath and A. S. Wein. Low-Degree Hardness of Random Optimization Problems. In *IEEE 61st Annual Symposium on Foundations of Computer Science (FOCS)*, pages 131–140. IEEE, 2020.
- [24] L. Ganassali and L. Massoulié. From tree matching to sparse graph alignment. In Proceedings of 33rd Conference on Learning Theory (COLT), pages 1633–1665. PMLR, 2020.
- [25] L. Ganassali, L. Massoulie, and M. Lelarge. Impossibility of partial recovery in the graph alignment problem. In *Proceedings of 34th Conference on Learning Theory (COLT)*, pages 2080–2102. PMLR, 2021.
- [26] L. Ganassali, L. Massoulié, and M. Lelarge. Correlation detection in trees for planted graph alignment. In Annals of Applied Probability, 34(3):2799–2843, 2024.
- [27] L. Ganassali, L. Massoulié, and G. Semerjian. Statistical limits of correlation detection in trees. In Annals of Applied Probability, 34(4):3701–3734, 2024.
- [28] J. Gaudio, M. Z. Racz, and A. Sridhar. Exact community recovery in correlated stochastic block models. In *Conference on Learning Theory (COLT)*, pages 2183–2241. PMLR, 2022.
- [29] G. Hall and L. Massoulié. Partial recovery in the graph alignment problem. In Operations Research, 71(1):259–272, 2023.
- [30] S. Hopkins. Statistical inference and the sum of squares method. PhD thesis, Cornell University, 2018.
- [31] S. Hopkins, P. K. Kothari, A. Potechin, P. Raghavendra, T. Schramm, and D. Steurer. The power of sum-of-squares for detecting hidden structures. In *IEEE 58th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 720–731. IEEE, 2017.
- [32] S. Hopkins and D. Steurer. Efficient Bayesian estimation from few samples: community detection and related problems. In *IEEE 58th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 379–390. IEEE, 2017.
- [33] H. Kesten and B. P. Stigum. Additional limit theorems for indecomposable multidimensional Galton-Watson processes. In Annals of Mathematical Statistics, 37:1463–1481, 1966.
- [34] P. K. Kothari, S. S. Vempala, A. S. Wein, and J. Xu. Is planted coloring easier than planted clique? In *Proceedings of 36th Conference on Learning Theory (COLT)*, pages 5343–5372. PMLR, 2023.

- [35] D. Kunisky. Hypothesis testing with low-degree polynomials in the Morris class of exponential families. In Proceedings of 34th Conference on Learning Theory (COLT), pages 2822–2848. PMLR, 2021.
- [36] D. Kunisky, C. Moore, and A. S. Wein. Tensor cumulants for statistical inference on invariant distributions. In *IEEE 65th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 1007–1026. IEEE, 2024.
- [37] D. Kunisky, A. S. Wein, and A. S. Bandeira. Notes on computational hardness of hypothesis testing: Predictions using the low-degree likelihood ratio. In *Mathematical Analysis, its Applications and Computation: ISAAC*, pages 1–50. Springer, 2022.
- [38] K. Makarychev, R. Manokaran, and M. Sviridenko. Maximum quadratic assignment problem: Reduction from maximum label cover and lp-based approximation algorithm. In *International Colloquium on Automata, Languages, and Programming (ICALP)*, pages 594–604. Springer, 2010.
- [39] C. Mao, M. Rudelson, and K. Tikhomirov. Random graph matching with improved noise robustness. In *Proceedings of 34th Conference on Learning Theory (COLT)*, pages 3296–3329. PMLR, 2021.
- [40] C. Mao, M. Rudelson, and K. Tikhomirov. Exact matching of random graphs with constant correlation. In *Probability Theory and Related Fields*, 186(2):327–389, 2023.
- [41] C. Mao, Y. Wu, J. Xu, and S. H. Yu. Random graph matching at Otter's threshold via counting chandeliers. In *Proceedings of the 55th Annual ACM Symposium on Theory of Computing* (STOC), pages 1345–1356. ACM, 2023.
- [42] C. Mao, Y. Wu, J. Xu, and S. H. Yu. Testing network correlation efficiently via counting trees. In Annals of Statistics, 52(6):2483–2505, 2024.
- [43] A. Montanari and A. S. Wein. Equivalence of approximate message passing and low-degree polynomials in rank-one matrix estimation. to appear in *Probability Theory and Related Fields*.
- [44] A. Moitra and A. S. Wein. Precise error rates for computationally efficient testing. to appear in Annals of Statistics.
- [45] A. Narayanan and V. Shmatikov. Robust de-anonymization of large sparse datasets. In *IEEE Symposium on Security and Privacy*, pages 111–125. IEEE, 2008.
- [46] R. Otter. The number of trees. In Annals of Mathematics, pages 583–599, 1948.
- [47] M. Z. Racz and A. Sridhar. Correlated stochastic block models: exact graph matching with applications to recovering communities. In Advances in Neural Information Processing Systems (NIPS), volume 34, pages 22259–22273. Curran Associates, Inc., 2021.
- [48] P. Raghavendra, T. Schramm, and D. Steurer. High-dimensional estimation via sum-of-squares proofs. In *Proceedings of the International Congress of Mathematicians (ICM 2018)*, pages 3389–3423, 2019.
- [49] C. Rush, F. Skerman, A. S. Wein, and D. Yang. Is it easier to count communities than find them? In 14th Innovations in Theoretical Computer Science Conference (ITCS), Schloss Dagstuhl-Leibniz-Zentrumfűr Informatik, 2023.

- [50] T. Schramm and A. S. Wein. Computational barriers to estimation from low-degree polynomials. In Annals of Statistics, 50(3):1833–1858, 2022.
- [51] R. Singh, J. Xu, and B. Berger. Global alignment of multiple protein interaction networks with application to functional orthology detection. In *Proceedings of the National Academy of Sciences of the United State of America*, 105(35):12763–12768, 2008.
- [52] Y. Sohn and A. S. Wein. Sharp phase transitions in estimation with low-degree polynomials. To appear in STOC 2025.
- [53] A. S. Wein. Optimal low-degree hardness of maximum independent set. In *Mathematical Statistics and Learning*, pages 221–251, 2022.
- [54] Y. Wu, J. Xu, and S. H. Yu, Settling the sharp reconstruction thresholds of random graph matching. In *IEEE Transactions on Information Theory*, 68(8):5391–5417, 2022.
- [55] Y. Wu, J. Xu, and S. H. Yu, Testing correlation of unlabeled random graphs. In Annals of Applied Probability, 33(4): 2519–2558, 2023.
- [56] I. Zadik, M. Jae Song, A. S. Wein, and J. Bruna. Lattice-based methods surpass sum-ofsquares in clustering. In *Proceedings of 35th Conference on Learning Theory (COLT)*, pages 1247–1248. PMLR, 2022.
- [57] L. Zdeborová and F. Krzakala. Statistical physics of inference: thresholds and algorithms. In Advances in Physics, 65(5):453–552, 2016.