

Zero patterns in multi-way binary contingency tables with uniform margins

Roberto Fontana^a, Elisa Perrone^b, Fabio Rapallo^c

^a Department of Mathematical Sciences, Politecnico di Torino, Corso Duca degli Abruzzi 24, 10129 Torino, Italy

^b Department of Mathematics and Computer Science, Eindhoven University of Technology, Groene Loper 3, 5612 AE Eindhoven, The Netherlands

^c Department of Economics, Università di Genova, via Vivaldi 5, 16126 Genova, Italy

Abstract

We study the problem of transforming a multi-way contingency table into an equivalent table with uniform margins and same dependence structure. This is an old question which relates to recent advances in copula modeling for discrete random vectors. In this work, we focus on multi-way binary tables and develop novel theory to show how the zero patterns affect the existence and uniqueness of the transformation as well as its statistical interpretability in terms of dependence structure. The implementation of the theory relies on combinatorial and linear programming techniques. Several examples are described to illustrate the approach and point to interesting future research directions.

Keywords: Categorical data analysis, odds ratios, discrete copulas, multivariate Bernoulli, Iterative Proportional Fitting.

1. Introduction

Largely employed across numerous domains such as healthcare, biology, and social sciences, contingency tables serve to display data in tabular format. Contingency tables have been extensively analyzed within the field of statistics, primarily with the objective of developing methods to understand the dependence between variables (see, for example, [1]). Recent work presented in [2] highlights fascinating connections between the analysis of two-way contingency tables and copulas. The essence of copula theory is centered on distinguishing between the influence of separate variables and their mutual dependency within the model. This separation facilitates the use of *ad hoc* dependence modeling techniques by transforming the initial joint probability distribution into one with uniform margins on $[0, 1]$, known as the copula. When the marginal distributions are continuous, the copula associated with the original distribution is uniquely defined ([3]). For discrete random variables, it is not possible to transform marginal distributions into uniform distributions using the Probability Integral Transform (PIT). The PIT identifies the copula solely within a specific subdomain. As a result, numerous copula models align with this specific subdomain. The adaptation of copula theory to utilize an analogous notion for establishing a margin-free model within a discrete framework has been investigated in [2, 4]. In that study and the referenced literature, the authors explore the concept of transforming a given two-way contingency table into a new one that has uniform margins. This type of transformation clarifies the underlying relationships in the table, which could be obscured by notably uneven margins, as explained in the example below.

Motivating example. For 2×2 tables, the process entails converting the original table into a new one in which each marginal probability equals $1/2$, indicating a uniform distribution across the subdomain. In Table 1, we provide a classic instance from [5] that demonstrates smallpox cases documented at Sheffield Hospital. The dataset categorizes patients based on their vaccination status (yes or no) and their recovery outcome (yes or no). The odds ratio, frequently used to evaluate associations in contingency table analysis, is notable with a value of 19.47. Despite this, the unevenness of the marginals could potentially




Figure 1: Bubble plot of original data (a) and transformed data (b) for the Sheffield smallpox epidemic data in Table 1. Part (c) represents a possible zero pattern in the observed table that motivates the investigations in this paper.

obscure the link within the original dataset. The connection becomes apparent once the data is altered to show marginal probabilities of $1/2$, as illustrated in Fig. 1 (b), while maintaining an odds ratio of 19.47.

Transforming an original table into a new table with same odds ratios and uniform margins becomes challenging when the original table contains zero entries. This is not only due to the vanishing odds ratios but also to the possible non-existence of a transformed table which maintains the zero-structure and has uniform margins. To clarify this, we consider a modification of the above example where the probability mass p_{00} is zero. A generic table with this structure is represented in Fig. 1 (c). It is straightforward to check that a table that preserves such a zero-structure and has uniform margins does not exist. Indeed, if such a table would exist, it would be the solution of the following system of equations

$$\begin{cases} p_{00} + p_{01} + p_{10} + p_{11} = 1 \\ p_{00} + p_{01} - p_{10} - p_{11} = 0 \\ p_{00} - p_{01} + p_{10} - p_{11} = 0 \\ p_{00} = 0 \end{cases} \quad (1)$$

However, the system in Eq. (1) has no solutions unless p_{11} is also set to zero, which means the original zero pattern is not maintained.

Research problem and paper's contribution. The situation described in Fig. 1 (c) is not artificial as zero entries are often observed in real datasets. Being an issue in applications, the problem of zero entries in contingency table analysis has recently caught the attention of researchers, especially in algebraic statistics, see for instance [6, Chap. 9]. When working with sampling distributions, zeros can appear both as structural zeros and sampling zeros, a far more common situation than the structural zeros: As the number of cells increases in multi-way tables, so does the probability of sampling zeros even when the sample size is moderate or large. In both cases, some of the odds ratios become zero or undefined, leading to the removal of the corresponding odds ratio equation in the transformation. As a consequence, odds ratios become uninformative of the relationship between the variables and new issues on the uniqueness of a

Vaccination (X_1)	Recovery (X_2)	\tilde{n}	\tilde{p}
no	no	274	0.058
no	yes	278	0.059
yes	no	200	0.043
yes	yes	3951	0.840

Table 1: Sheffield smallpox epidemic reported in [5].

transformed table that preserves the dependence structure arise.

In this work, we further develop the initial investigations of [7, 8] to address the case of zeros in binary multi-way contingency tables. Our *problem statement* is the following: Given a table T (a probability table or a contingency table of observed frequencies), find a table \tilde{T} with the uniform margins and the same odds ratios structure as T . In particular, we provide conditions to ensure the existence of a transformed table with uniform margins, prescribed odds ratios, and zero pattern. We further investigate the uniqueness of the transformed table when it exists in relation to the zero pattern.

The paper is organized as follows: In Section 2, we describe the mathematical context of this work, introduce the notation, and connect our research with the state-of-the-art. Section 3 presents novel theoretical results on the existence of a transformed table with certain zero patterns. In Section 4, we show that in some degenerate cases, there might be a wider class of tables that satisfy the odds ratio constraints and have uniform margins. We conclude the paper with a discussion in Section 5.

2. Background, notation, and relevance to the state-of-the-art

In this work, we consider the case of d -dimensional multi-way binary contingency tables with zero entries and study how the presence of zeros affects the possibility of transforming a given table into one with uniform margins. In particular, we analyze all possible zero patterns where a zero pattern represents the set of the table cells whose corresponding frequencies are zero.

Here, we consider d binary factors, X_1, \dots, X_d , i.e., a d -dimensional random vector (X_1, \dots, X_d) . We refer to a specific cell of a table T with the binary vector $\alpha = (\alpha_1, \dots, \alpha_d) \in \{0, 1\}^d$. We assume the set $\{0, 1\}^d$ in lexicographic order and then we can refer to the k -th cell of T , $k = 1, \dots, 2^d$. We observe that the k -th cell of a table is the one corresponding to $\alpha_k = (\alpha_{k_1}, \dots, \alpha_{k_d})$ such that $\sum_{j=1}^d \alpha_{k_j} 2^{d-j} = k - 1$, $k = 1, \dots, 2^d$, i.e., α_k is the binary representation of $k - 1$. For each cell α , $\alpha \in \{0, 1\}^d$, we define p_α , the relative frequency of the cell. Thus, a table T is described by the 2^d -vector $p = (p_\alpha, \alpha \in \{0, 1\}^d)$. In the d -dimensional case, there are 2^d possible zero patterns. We can represent each zero pattern \mathcal{Z} with a binary 2^d vector, i.e., $\mathcal{Z} = (z_1, \dots, z_{2^d})$, where $z_i = 0$ ($z_i = 1$) indicates that the corresponding cell α_i has $p_{\alpha_i} = 0$ ($p_{\alpha_i} > 0$, respectively). We write that a zero pattern \mathcal{Z} is a k -zero pattern if the number of zeros in \mathcal{Z} is k , with $k = 0, \dots, 2^d$. Finally, we say that a table T with uniform margin is \mathcal{Z} -compatible if the corresponding probability mass function (pmf) p satisfies that $p_{\alpha_i} = 0$ ($p_{\alpha_i} > 0$) when $z_i = 0$ ($z_i = 1$, respectively), $i = 1, \dots, 2^d$.

We now introduce the notion of conditional odds ratios, which is key to this work. In particular, we define the conditional odds ratios constraints on each 2×2 sub-tables by fixing $d - 2$ variables. When $d = 2$, there is only one odds ratio given by

$$\omega^{12} = \frac{p_{11}p_{00}}{p_{10}p_{01}}.$$

In the case $d = 3$ there are six odds ratios given by

$$\begin{aligned} \omega_0^{23} &= \frac{p_{000}p_{011}}{p_{001}p_{010}} & \omega_1^{23} &= \frac{p_{100}p_{111}}{p_{101}p_{110}} \\ \omega_0^{13} &= \frac{p_{000}p_{101}}{p_{001}p_{100}} & \omega_1^{13} &= \frac{p_{010}p_{111}}{p_{011}p_{110}} \\ \omega_0^{12} &= \frac{p_{000}p_{110}}{p_{010}p_{100}} & \omega_1^{12} &= \frac{p_{001}p_{111}}{p_{011}p_{101}} \end{aligned} \quad (2)$$

The notation ω_0^{23} denotes the odds ratio for the variables X_2 and X_3 given the value 0 for X_1 . The conditions in Eq. (2) are not independent, and only four of them are independent.

In arbitrary dimension $d > 3$, the generic conditional odds ratio is defined in the same way: for each pair of variables X_i and X_j we consider the conditional odds ratio for given values of the other $(d - 2)$ variables. If we denote with $\alpha' \in \{0, 1\}^{d-2}$ the values of the fixed $(d - 2)$ variables, the odds ratio is denoted as $\omega_{\alpha'}^{ij}$ and defined by:

$$\omega_{\alpha'}^{ij} = \frac{p_{\alpha_1 p_{\alpha_2}}}{p_{\alpha_3 p_{\alpha_4}}}, \quad (3)$$

where $\alpha_1, \alpha_2, \alpha_3, \alpha_4$ are equal to $\alpha' \in \{0, 1\}^{d-2}$ in the entries $\{1, \dots, d\} - \{i, j\}$ and are equal to $(1, 1), (0, 0), (1, 0), (0, 1)$ respectively in the entries (i, j) . In the d -way case there are $\binom{d}{2} 2^{d-2}$ equations, which are not independent. In the following, we draw connections with previous research on discrete copulas and classical results in the context of log-linear models.

2.1. Table transformations and discrete copulas

In this section, we related our research problems to the theory of discrete copulas. For the sake of clarity, we first restrict to the two-dimensional case. We consider $R \in \mathbb{Z}_{>0}$ and denote $I_R = \{0, 1/R, \dots, (R-1)/R, 1\}$, $[R] = \{1, \dots, R\}$, and $\langle R \rangle = \{0, \dots, R\}$. Given R and S as positive integers, we define $U_R = \{u_0 = 0, u_1, \dots, u_{R-1}, u_R = 1\}$, with $u_0 < \dots < u_R$, and $V_S = \{v_0 = 0, v_1, \dots, v_{S-1}, v_S = 1\}$, with $v_0 < \dots < v_S$, as two discrete grid partitions over the unit interval. A discrete copula C_{U_R, V_S} is defined on the set $U_R \times V_S$ and retains the characteristic properties of a copula function over the grid domain $U_R \times V_S$. As noted in [9, 10], interesting relationships exist between the domain of discrete copulas and certain convex polytopes known as *transportation polytopes*. These polytopes are also associated with the contingency table analysis and the existence of a table with prescribed margins [11]. Namely, considering two vectors $\tilde{u} = (\tilde{u}_1, \dots, \tilde{u}_R) \in \mathbb{R}_{>0}^R$ and $\tilde{v} = (\tilde{v}_1, \dots, \tilde{v}_S) \in \mathbb{R}_{>0}^S$, we can define the transportation polytope $\mathcal{T}(\tilde{u}, \tilde{v})$ as the convex polytope in the RS variables $x_{i,j}$ satisfying, for all $i \in [R]$ and $j \in [S]$, the following conditions: $x_{i,j} \geq 0$, $\sum_{h=1}^S x_{i,h} = \tilde{u}_i$, $\sum_{\ell=1}^R x_{\ell,j} = \tilde{v}_j$. The two vectors \tilde{u} and \tilde{v} are called the margins of $\mathcal{T}(\tilde{u}, \tilde{v})$. According to [12], each discrete copula C_{U_R, V_S} is associated with a matrix in a transportation polytope $\mathcal{T}(\tilde{u}, \tilde{v})$, and conversely. The transportation matrix is connected to the probability mass function of the discrete random vector, whereas the associated discrete copula corresponds to the cumulative distribution function.

We now show how to derive the discrete copula associated with a given contingency table. We consider the motivating example presented in Table 1. There, we get $N = 4703$ total observations. In this example, $R = S = 2$, the vectors \tilde{u} and \tilde{v} are the margins of the contingency table, i.e., $\tilde{u} = (552, 4151)$ and $\tilde{v} = (474, 4229)$, while the defining grids of the corresponding discrete copula are $U_2 = \frac{1}{N}\{0, \tilde{u}_1, \tilde{u}_1 + \tilde{u}_2\} = \{0, 0.12, 1\}$ and $V_2 = \{0, 0.1, 1\}$. The entries of the discrete copula $C_1 = C_{U_2, V_2} = (c_{i,j})$, $i \in [2]$ and $j \in [2]$ are computed from the entries of the contingency table $(x_{i,j})$ by summing up and normalizing, i.e., $c_{i,j} = \frac{1}{N} \sum_{\ell=1}^i \sum_{h=1}^j x_{\ell,h}$, while $c_{0,0} = c_{i,0} = c_{0,j} = 0$, for $i \in [2]$ and $j \in [2]$. Namely, C_1 is as follows:

$$C_1 = \begin{pmatrix} 0 & 0.00 & 0.00 \\ 0 & 0.058 & 0.12 \\ 0 & 0.1 & 1.00 \end{pmatrix}.$$

As discussed in the introduction, in [2], the author emphasizes the challenge of inferring dependence from tables with non-uniform margins, such as Table 1. In line with copula theory for continuous random variables, the author proposes finding a representative \mathbf{p} for all $(R \times S)$ probability distributions that (1) maintains the inter-dependencies in a contingency table using odds ratios, and (2) has uniform margins of $1/R$ and $1/S$. In discrete copulas terms, this corresponds to finding a discrete copula defined on the rectangular grid $I_R \times I_S$ that maintains the dependence structure of the initial discrete copula derived from a specific contingency table. In the example above, the problem is to identify a discrete copula C_1 with its domain on $I_2 \times I_2$ and margins I_2 , which is somehow compatible with the original discrete copula C_1 .

A natural question that arises is whether or not such an element exists and is unique. For two-dimensional tables, the answer to this question is given in a theorem presented in [2], which we report below in a simplified version.

Theorem 2.1. *Let \mathbf{p} be in the set $\mathcal{P}_{R \times S}$ of all $(R \times S)$ probability distributions. We define $\text{Supp}(\mathbf{p}) = \{(i, j) \in [R] \times [S] \text{ s.t. } p_{i,j} > 0\}$ and $N(\mathbf{p}) = \{(v_{X_1} \times v_{X_2}) : v_{X_1} \subset [R], v_{X_2} \subset [S] \text{ s.t. } \sum_{(i,j) \in v_{X_1} \times v_{X_2}} p_{i,j} = 0\}$, the set of rectangular subset of $[R] \times [S]$ where \mathbf{p} is null. The cardinality of a set A is denoted by $|A|$.*

1. *Suppose that for all $(v_{X_1} \times v_{X_2}) \in N(\mathbf{p})$, $\frac{|v_{X_1}|}{R} + \frac{|v_{X_2}|}{S} < 1$, then there exists a unique \mathbf{p} with uniform margins associated with a discrete copula $C_{I_R \times I_S}$.*

2. Suppose that for all $(v_{X_1} \times v_{X_2}) \in N(\mathbf{p})$, $\frac{|v_{X_1}|}{R} + \frac{|v_{X_2}|}{S} \leq 1$ with $\frac{|\tilde{v}_{X_1}|}{R} + \frac{|\tilde{v}_{X_2}|}{S} = 1$ for some $(\tilde{v}_{X_1} \times \tilde{v}_{X_2}) \in N(\mathbf{p})$.
 - (i) If, for all $(\tilde{v}_{X_1} \times \tilde{v}_{X_2}) \in N(\mathbf{p})$ such that $\frac{|\tilde{v}_{X_1}|}{R} + \frac{|\tilde{v}_{X_2}|}{S} = 1$, $([R] \setminus \tilde{v}_{X_1} \times [S] \setminus \tilde{v}_{X_2}) \in N(\mathbf{p})$, then there exists a unique $\tilde{\mathbf{p}}$ with uniform margins associated with a discrete copula $C_{I_R \times I_S}$.
 - (ii) If, there exists $(\tilde{v}_{X_1}^* \times \tilde{v}_{X_2}^*) \in N(\mathbf{p})$ such that $\frac{|\tilde{v}_{X_1}^*|}{R} + \frac{|\tilde{v}_{X_2}^*|}{S} = 1$, and $([R] \setminus \tilde{v}_{X_1}^* \times [S] \setminus \tilde{v}_{X_2}^*) \notin N(\mathbf{p})$, then there is no element $\tilde{\mathbf{p}}$ with uniform margins and same support of the original table, but there is one element $\tilde{\mathbf{p}}$ with modified support.
3. Suppose that there exists $\tilde{v}_{X_1} \times \tilde{v}_{X_2} \in N(\mathbf{p})$ such that $\frac{|\tilde{v}_{X_1}|}{R} + \frac{|\tilde{v}_{X_2}|}{S} > 1$. Then there is no element $\tilde{\mathbf{p}}$ with uniform margins such that it has same odds ratio structure of \mathbf{p} and is associated with a discrete copula $C_{I_R \times I_S}$, not even with a modified support.

We can interpret the example of Table 1 in light of Thm. 2.1. For a broader discussion of the two-dimensional binary case, we refer the reader to [2]. The case reported in Table 1 does not show any zero pattern, since all entries are strictly positive. Therefore, the transformed table exists and is unique in agreement with Part 1 of Thm. 2.1. We now analyze the situation depicted in Fig. 1 (c) where one entry is zero while the others are non-zero. This case falls under point 2 (ii) of Thm. 2.1. Specifically, $|v_{X_1}| = |v_{X_2}| = 1$. Hence, $\frac{|\tilde{v}_{X_1}^*|}{2} + \frac{|\tilde{v}_{X_2}^*|}{2} = 1$ and a solution does not exist in agreement with the conclusion of the introduction. If we move to the boundary of the probability space by adding an extra zero in the zero pattern (that is, allowing p_{11} to be set equal to zero), a solution exists and is unique. When it exists, the element $\tilde{\mathbf{p}}$ can be obtained using the Iterative Proportional Fitting Procedure (IPFP), which is a standard method in contingency table analysis for a meaningful comparison of tables with different margins and same dependence structure in terms of corresponding odds ratios [2, 1]. In the following, we further discuss how our problem relates to standard theory in log-linear models.

2.2. Relations with the classical theory of log-linear models

In log-linear models, there is a well established link between IPFP and Maximum Likelihood Estimation (MLE). The use of IPFP to find MLE in complete tables dates back to [13] (see also [14]). IPFP works in log-linear models as follows: First, define a table satisfying all the desired independence and/or conditional independence statements, and then, run the IPFP to adjust the table with respect to the relevant marginal distributions of the sufficient statistic. When there are zeros in the table, the existence of the MLE is generally not guaranteed. Important developments in this case have been achieved with the use of algebraic Statistics: In most cases there are criteria to identify zero patterns which preserve the existence of the MLE, see [15]. In large classes of statistical models, including hierarchical log-linear models and staged trees, recent results have been introduced in [16]. Despite these connections, our problem is different because the margins are constrained to be uniform, while in log-linear models the margins are determined by suitable margins of the observed table. To clarify this key difference, we can refer again to the example depicted in Fig. 1 (c). In that situation, it is clear that there is no table with uniform margins and the prescribed support. However, if we start with unbalanced margins, i.e., $(1/3, 2/3)$, and relax the constraints of uniform margins, we can easily obtain tables that maintain the zero pattern.

3. Impact of zeros in the existence of the transformed table - binary case

As mentioned in the previous sections, the presence of zeros in the table affects the construction of a table with uniform margins in different ways. First, some odds ratio equations in Eq. (3) become meaningless and can thus be removed from the system. Additionally, the presence of zeros reduces the support of the probability distribution, reducing the number of free probabilities in the table.

As a prototype, let us consider again the 2×2 case. Given a 2×2 contingency table $(\tilde{n}_{ij}, i, j = 0, 1)$ let $N = \tilde{n}_{00} + \tilde{n}_{11} + \tilde{n}_{01} + \tilde{n}_{10}$ be the grand total and $\omega^{12} = \frac{\tilde{n}_{00}\tilde{n}_{11}}{\tilde{n}_{01}\tilde{n}_{10}}$ the odds ratio. We examine the table in terms of the relative frequencies $\tilde{p}_{ij} = \frac{\tilde{n}_{ij}}{N}$ rather than the counts \tilde{n}_{ij} . As shown in the motivating example in Sect. 1, the goal is to determine a new table $(p_{ij} \geq 0, i, j = 0, 1)$ such that the marginals are still $1/2$ and the

original table's odds ratio ω^{12} is maintained. Such a table can be obtained by solving the following system of equations:

$$\begin{cases} \frac{p_{00}p_{11}}{p_{01}p_{10}} = \omega^{12} \\ p_{00} + p_{01} + p_{10} + p_{11} = 1 \\ p_{00} + p_{01} - p_{10} - p_{11} = 0 \\ p_{00} - p_{01} + p_{10} - p_{11} = 0 \end{cases}, \quad (4)$$

whose unique solution is given as follows:

$$p_{00} = p_{11} = \frac{\sqrt{\omega^{12}}}{2(1 + \sqrt{\omega^{12}})}, \quad p_{01} = p_{10} = \frac{1}{2(1 + \sqrt{\omega^{12}})}. \quad (5)$$

In the described 2^2 scenario, addressing the issue of zeros in the table is straightforward, with two possible situations: (a) The table is entirely filled, or (b) The table might contain one or more zero entries. As discussed in Sec. 2, if the table is complete, there exists a unique table with identical odds ratios and uniform margins, obtainable through the IPFP. Referring again to [2], in scenario (b), odds ratios are absent, simplifying the task of identifying a table with the same support as the original and uniform margins. This scenario is only achievable when there are two zeros located at $(0, 0)$, $(1, 1)$ or $(1, 0)$, $(0, 1)$. In an intermediate scenario with a single zero present in the table as in Fig. 1 (c), the IPFP leads to a boundary solution, resulting in another cell probability approaching zero. In this case, there are no tables which are zero-compatible with the original table and also have uniform margins.

While in the two-way case, zeros only affect the existence of a table with uniform margins and desired odds ratios as discussed in Section 2, in the multi-way case the presence of zeros gives rise to a number of different situations. In the next section, we provide existence and uniqueness results for arbitrary dimensions and zero patterns.

3.1. d -way binary tables

We now move to the general d -way case. We denote $\{0, 1\}^k$ by V_k , $k = 1, \dots, d$. To simplify the notation, sometimes we write \mathcal{D} instead of $V_d = \{0, 1\}^d$. Here, we are interested in tables with uniform margins. More specifically, there are d margins m_i , $i = 1, \dots, d$ defined as

$$m_i = (m_{i,0}, m_{i,1}) = \left(\sum_{\alpha \in \mathcal{D}, \alpha_i=0} p_\alpha, \sum_{\alpha \in \mathcal{D}, \alpha_i=1} p_\alpha \right),$$

where $\mathcal{D} = \{0, 1\}^d$. We observe that $m_{i,1} = E[X_i]$ and $m_{i,0} + m_{i,1} = 1$, $i = 1, \dots, d$. To obtain uniform margins, the pmf of the table must satisfy the following constraints

$$m_{i,0} = m_{i,1} = \frac{1}{2}, \quad i = 1, \dots, d.$$

The next proposition 3.1 states a general result on the structure of the tables with uniform margins that are compatible with a class of zero patterns.

Proposition 3.1. *Let us consider a table T , with relative frequencies $p = (p_\alpha, \alpha \in V_d)$ and uniform margins. Let us suppose that $p_\alpha = 0$ for $\alpha \in \{(\alpha_1, \dots, \alpha_d) \in V_d : \alpha_{i_1} = y_{i_1}, \alpha_{i_2} = y_{i_2}, i_1, i_2 \in \{1, \dots, d\}, i_1 \neq i_2, y_{i_1}, y_{i_2} \in \{0, 1\}\}$. Then $p_\alpha = 0$ for $\alpha \in \{(\alpha_1, \dots, \alpha_d) \in V_d : \alpha_{i_1} = 1 - y_{i_1}, \alpha_{i_2} = 1 - y_{i_2}\}$.*

Proof. Without loss of generality, we can take $i_1 = 1$, $i_2 = 2$, $y_{i_1} = y_{i_2} = 0$. The hypothesis on p can be written as $p_{0\alpha} = 0$, $\alpha \in V_{d-2}$. We compute $m_{1,0}$, the first component of the first margin:

$$m_{1,0} = \sum_{\alpha' \in V_{d-1}} p_{0\alpha'} = \sum_{\alpha'' \in V_{d-2}} p_{00\alpha''} + \sum_{\alpha'' \in V_{d-2}} p_{01\alpha''} = \sum_{\alpha'' \in V_{d-2}} p_{01\alpha''}.$$

The last equality in the above equation follows by $p_{00\alpha} = 0$, $\alpha \in V_{d-2}$. The table has uniform margins, in particular $m_{1,0} = \frac{1}{2}$, and as a consequence,

$$\sum_{\alpha'' \in V_{d-2}} p_{01\alpha''} = \frac{1}{2}.$$

Let us now consider the second component of the second margin, $m_{2,1}$. The table T has uniform margins and then $m_{2,1} = \frac{1}{2}$. Thus, we obtain the following chain of equalities:

$$m_{2,1} = \sum_{\alpha_1 \in \{0,1\}, \alpha'' \in V_{d-2}} p_{\alpha_1,1,\alpha''} = \sum_{\alpha'' \in V_{d-2}} p_{01\alpha''} + \sum_{\alpha'' \in V_{d-2}} p_{11\alpha''} = \frac{1}{2} + \sum_{\alpha'' \in V_{d-2}} p_{11\alpha''}.$$

Consequently, we obtain $\sum_{\alpha'' \in V_{d-2}} p_{11\alpha''} = 0$, that is $p_{11\alpha''} = 0$, $\alpha'' \in V_{d-2}$.

To complete the proof, we show that there exists a table T with uniform margins and $p_{00\alpha''} = p_{11\alpha''} = 0$, $\alpha'' \in V_{d-2}$. Namely, we consider

$$p_{\alpha} = \begin{cases} 0 & \alpha = (0, 0, \alpha'') \text{ or } \alpha = (1, 1, \alpha''), \alpha'' \in V_{d-2} \\ \frac{1}{2^{d-1}} & \text{elsewhere} \end{cases}.$$

Since all margins have a similar structure, it is enough to compute $m_{1,0}$ and $m_{3,0}$, which are:

$$m_{1,0} = \sum_{\alpha'' \in V_{d-2}} p_{01\alpha''} = \sum_{\alpha'' \in V_{d-2}} \frac{1}{2^{d-1}} = \frac{2^{d-2}}{2^{d-1}} = \frac{1}{2},$$

$$m_{3,0} = \sum_{\alpha \in \mathcal{D}, \alpha_3=0} p_{\alpha} = \sum_{\alpha \in V_{d-3}} p_{010\alpha} + \sum_{\alpha \in V_{d-3}} p_{100\alpha} = \frac{1}{2^{d-1}} 2^{d-3} + \frac{1}{2^{d-1}} 2^{d-3} = \frac{1}{2}.$$

Finally, the condition $\sum_{\alpha \in \mathcal{D}} p_{\alpha} = 1$ can be verified by observing, for example, that

$$\sum_{\alpha \in \mathcal{D}} p_{\alpha} = m_{1,0} + m_{1,1} = \frac{1}{2} + \frac{1}{2} = 1.$$

□

To further explore the existence of a table with uniform margins, we use the following lemma. The lemma is based on a rewriting of the multidimensional case Th. 3 in [17], and an adaptation of the notation to binary tables.

Lemma 3.2. *Let $T_1 = (p_{\alpha})$ and $T_2 = (q_{\alpha})$ be two d -dimensional binary probability tables with the same zero pattern, and assume that T_2 has uniform margins. Then there exist positive vectors $w_1 = (w_{10}, w_{11}), \dots, w_d = (w_{d0}, w_{d1})$ such that the table T_3 with generic element*

$$\tilde{p}_{\alpha_1 \dots \alpha_d} = p_{\alpha_1 \dots \alpha_d} w_{1\alpha_1} \dots w_{d\alpha_d}$$

has uniform margins.

We now highlight two key points on the use of Lemma 3.2. First, for the existence of a solution to our problem, it is enough to check the conditions on margins, since the conditions on the odds ratios are automatically adjusted by the IPFP. Second, it is enough to prove the existence of a table with the same zero pattern and uniform margins, and it is not even necessary to actually determine such a table.

To do this, we propose two methods. The first method is based on combinatorial objects, while the second method is based on integer programming.

We point out that to have uniform margins the pmf p of a table T must satisfy the following constraints

$$m_{i0} = m_{i1} = \frac{1}{2} \Leftrightarrow m_{i0} - m_{i1} = 0 \Leftrightarrow \sum_{\alpha \in \mathcal{D}, \alpha_i=0} p_{\alpha} - \sum_{\alpha \in \mathcal{D}, \alpha_i=1} p_{\alpha} = 0, \quad i = 1 \dots, d$$

that can be rewritten as

$$(\mathbf{1}_{\{\alpha \in \mathcal{D}, \alpha_i=0\}} - \mathbf{1}_{\{\alpha \in \mathcal{D}, \alpha_i=1\}})^T p = 0, \quad i = 1, \dots, d, \quad (6)$$

where $\mathbf{1}_A$ is the indicator vector of A , b^T denotes the transpose of b , and p is the vector $p = (p_\alpha, \alpha \in \mathcal{D})$. Eq. (6) can be rewritten in compact form by denoting as H_d its matrix of coefficients. Namely, Eq. (6) becomes:

$$H_d p = 0. \quad (7)$$

For example, for $d = 3$ we obtain

$$H_3 p = \begin{pmatrix} 1 & 1 & 1 & 1 & -1 & -1 & -1 & -1 \\ 1 & 1 & -1 & -1 & 1 & 1 & -1 & -1 \\ 1 & -1 & 1 & -1 & 1 & -1 & 1 & -1 \end{pmatrix} p = 0.$$

The extreme rays of the system defined by $H_d y = 0$ with $y = (y_\alpha, \alpha \in \mathcal{D}, y_\alpha \geq 0)$ can be computed using software for commutative algebra. We used 4ti2, [18]. Then the corresponding extreme pmfs are computed by simple normalization $p_\alpha = \frac{y_\alpha}{\sum_{\alpha \in \mathcal{D}} y_\alpha}$. For the 3-dimensional case we obtain the extreme pmfs that are shown in Table 2.

α_1	α_2	α_3	r_1	r_2	r_3	r_4	r_5	r_6
0	0	0	$\frac{1}{2}$	0	0	0	$\frac{1}{4}$	0
1	0	0	0	$\frac{1}{2}$	0	0	0	$\frac{1}{4}$
0	1	0	0	0	$\frac{1}{2}$	0	0	$\frac{1}{4}$
1	1	0	0	0	0	$\frac{1}{2}$	$\frac{1}{4}$	0
0	0	1	0	0	0	$\frac{1}{2}$	0	$\frac{1}{4}$
1	0	1	0	0	$\frac{1}{2}$	0	$\frac{1}{4}$	0
0	1	1	0	$\frac{1}{2}$	0	0	$\frac{1}{4}$	0
1	1	1	$\frac{1}{2}$	0	0	0	0	$\frac{1}{4}$

Table 2: Extreme pmfs with uniform margins $d = 3$.

It is worth noting that, given the extreme pmfs $r_i, i = 1, \dots, n_d$ where n_d depends on the dimension d of the multi-way table (e.g., for the three-dimensional case we have $n_3 = 6$) any pmfs p with uniform margins can be written as $p = \sum_{i=1}^{n_d} \lambda_i r_i$ with $\lambda_i \geq 0, i = 1, \dots, n_d$ and $\sum_{i=1}^{n_d} \lambda_i = 1$. In Thm. 3.3, we provide conditions that a zero pattern must satisfy to be compatible with the existence of a table with uniform margins.

Theorem 3.3. *Let \mathcal{Z} be a zero pattern with $\mathcal{Z} = (z_i, i = 1, \dots, 2^d)$. Let us denote by \mathcal{A} the set of indices corresponding to zero cells, $\mathcal{A} = \{i \in \{0, \dots, 2^d\} : z_i = 0\}$ and $\bar{\mathcal{A}}$ the set of the indices corresponding to non-zero cells, $\bar{\mathcal{A}} = \{i \in \{0, \dots, 2^d\} : z_i = 1\}$. Let $\{r_i : i = 1, \dots, n_d\}$ be the set of extreme pmfs of the polytope defined by $H_d p = 0$.*

Let S_1 be the subset of $\{1, \dots, n_d\}$ defined as $S_1 = \{k \in \{1, \dots, n_d\} : r_k(\alpha_i) = 0, i \in \mathcal{A}\}$. If S_1 is not empty we consider the set $S_2 \subseteq \bar{\mathcal{A}}$ that contains all $i \in \bar{\mathcal{A}}$ such that $\sum_{j \in S_1} r_j(\alpha_i) > 0$.

If $S_2 = \mathcal{B}$ then $p = \sum_{i \in S_1} \frac{1}{|S_1|} r_i$ is a pmf of a \mathcal{Z} -compatible table with uniform margins. If $S_1 = \emptyset$ or $S_2 \subsetneq \mathcal{B}$ no \mathcal{Z} -compatible table with uniform margins exists.

Proof. We know that any p associated with a table T with uniform margins can be represented as $p = \sum_{k=1}^{n_d} \lambda_k r_k$ where $\lambda_k \geq 0, \sum_{k=1}^{n_d} \lambda_k = 1$, and $r_k, k = 1, \dots, n_d$ are the extreme pmfs of the polytope defined by $H_d p = 0$. We must have

$$p(\alpha_i) = \sum_{k=1}^{n_d} \lambda_k r_k(\alpha_i) = 0, \quad i \in \mathcal{A}.$$

We can choose a strictly positive value for λ_k only if the extreme pmf r_k satisfies the condition $r_k(\alpha_i) = 0, i \in \mathcal{A}$. We define S_1 as the subset of $\{1, \dots, n_d\}$ that contains the indexes k such that $r_k(\alpha_i) = 0, i \in \mathcal{A}$. If

S_1 is empty, no \mathcal{Z} -compatible pmf can be built. If S_1 is not empty, we have

$$p = \sum_{k \in S_1} \lambda_k r_k, \lambda_k \geq 0, \sum_{k \in S_1} \lambda_k = 1.$$

The pmf p must be positive for $\alpha_i, i \in \bar{\mathcal{A}}$:

$$p(\alpha_i) = \sum_{k \in S_1} \lambda_k r_k(\alpha_i) > 0, i \in \bar{\mathcal{A}}.$$

The coefficients $\lambda_k, k \in S_1$ are positive or zero. Then we can consider $\sum_{k \in S_1} r_k(\alpha_i)$. Let us define S_2 as the subset of $\bar{\mathcal{A}}$ that contains all the indexes $i \in \bar{\mathcal{A}}$ for which this sum is strictly positive, $\sum_{k \in S_1} r_k(\alpha_i) > 0$. If $S_2 = \bar{\mathcal{A}}$ it would be enough to choose $\lambda_k = \frac{1}{|\bar{\mathcal{A}}|}$: the pmf $p = \sum_{k \in \bar{\mathcal{A}}} \frac{1}{|\bar{\mathcal{A}}|} r_k$ has uniform margins and is \mathcal{Z} -compatible by construction. If S_2 is a proper subset of $\bar{\mathcal{A}}, S_2 \subset \bar{\mathcal{A}}$ no \mathcal{Z} -compatible pmf can be built. \square

Corollary 3.4. *Let \mathcal{Z} be a zero pattern with $\mathcal{Z} = (z_i, i = 1, \dots, 2^d)$. If S_1 is empty then no \mathcal{Z}' -compatible table with uniform margins exists for any $\mathcal{Z}' \leq \mathcal{Z}$, where $\mathcal{Z}' \leq \mathcal{Z}$ means $z'_i \leq z_i, i = 1, \dots, 2^d, \mathcal{Z}' = (z'_1, \dots, z'_{2^d})$.*

Example 3.5. *We consider some examples for $d = 3$ with zero patterns reported in the last columns of Table 3.*

Case 1. *The first case, \mathcal{Z}_1 , is a 7-zero pattern. For this case, we get $S_1 = \emptyset$ and then no table with uniform margins exists for this case.*

Case 2. *The second case, \mathcal{Z}_2 , is a 6-zero pattern. Here, we get $S_1 = \emptyset$. Thus, in such a case, no table with uniform margins exists. Using Cor. (3.4) we can conclude that for any zero pattern $\mathcal{Z}', z' = (z_1, z_2, 0, 0, 0, 0, 0, 0), z_1, z_2 \in \{0, 1\}$ no \mathcal{Z}' -compatible table with uniform margins exists.*

Case 3. *The third case, \mathcal{Z}_3 , is a 2-zero pattern. We get $S_1 = \{2, 4\}$. In this case, $\mathcal{A} = \{1, 3\}, \bar{\mathcal{A}} = \{2, 4, 5, 6, 7, 8\}$, and $S_2 = \{2, 4, 5, 7\} \subseteq \bar{\mathcal{A}}$. Then no table with uniform margins exists for this case.*

Case 4. *The fourth case, \mathcal{Z}_4 , is a 4-zero pattern. We obtain $S_1 = \{2, 4\}$. Here, $\mathcal{A} = \{1, 3, 6, 8\}, \bar{\mathcal{A}} = \{2, 4, 5, 7\}$, and $S_2 = \{2, 4, 5, 7\} = \bar{\mathcal{A}}$. Then $p = \sum_{k \in \{2, 4\}} \frac{1}{2} r_k = (0.0.25 \ 0.0.25 \ 0.25 \ 0.0.25 \ 0)$ has uniform margin and is compatible with \mathcal{Z}_4 .*

As previously discussed, given the extreme pmfs $r_i, i = 1, \dots, n_d$, any pmf with uniform margins can be expressed as follows:

$$p = \sum_{i=1}^{n_d} \lambda_i r_i, \lambda_i \geq 0, \sum_{i=1}^{n_d} \lambda_i = 1.$$

As a consequence, we have that each extreme pmf r_i itself represents a table with uniform margins that is compatible with $\mathcal{Z} = (z_1, \dots, z_{2^d})$, where $z_k = 0$ (or $z_k = 1$) if $r_i(\alpha_k) = 0$ (or $r_i(\alpha_k) > 0$, respectively), for $k = 1, \dots, 2^d$ and $i = 1, \dots, n_d$. Moreover, it is possible to sample from the set of tables with uniform margins by selecting $\lambda_i, i = 1, \dots, n_d$, according to some criterion within the $(n_d - 1)$ -dimensional simplex.

α_1	α_2	α_3	r_1	r_2	r_3	r_4	r_5	r_6	\mathcal{Z}_1	\mathcal{Z}_2	\mathcal{Z}_3	\mathcal{Z}_4
0	0	0	$\frac{1}{2}$	0	0	0	$\frac{1}{4}$	0	1	1	0	0
1	0	0	0	$\frac{1}{2}$	0	0	0	$\frac{1}{4}$	0	1	1	1
0	1	0	0	0	$\frac{1}{2}$	0	0	$\frac{1}{4}$	0	0	0	0
1	1	0	0	0	0	$\frac{1}{2}$	$\frac{1}{4}$	0	0	0	1	1
0	0	1	0	0	0	$\frac{1}{2}$	0	$\frac{1}{4}$	0	0	1	1
1	0	1	0	0	$\frac{1}{2}$	0	$\frac{1}{4}$	0	0	0	1	0
0	1	1	0	$\frac{1}{2}$	0	0	$\frac{1}{4}$	0	0	0	1	1
1	1	1	$\frac{1}{2}$	0	0	0	0	$\frac{1}{4}$	0	0	1	0

Table 3: $d = 3$: Extreme pmfs with uniform margins and four zero patterns.

On the other hand, it is important to note that the cardinality n_d of the set of extreme pmfs grows rapidly with d . For instance, when $d = 6$, there are $n_6 = 707,264$ extreme pmfs, which poses a significant computational challenge for this approach.

An alternative way to tackle our problem would be through integer programming. Since our problem reduces to linear constraints through the Lemma 3.2, we can set up a linear programming problem as follows. We start by considering the matrix C_d with dimensions $(d + 1) \times 2^d$ with: A first row of 1's; then block-wise repetitions of 1's and -1 's (as in a classical full-factorial 2^d design). C_d is the matrix H_d above except for the first row. Also, let us define the vector with length $d + 1$ as $b_d = (1, 0, \dots, 0)$. For a given zero pattern \mathcal{Z} , we introduce the matrix $C_d(\mathcal{Z})$ obtained from C_d by removing the columns that belong to the cells in \mathcal{Z} . The problem of the existence of a strictly positive probability distribution with uniform margins can be stated as follows:

$$\begin{aligned} & \max_p \delta \\ \text{subject to } & C_d(\mathcal{Z})p = b_d \\ & p_\alpha \geq \delta, \quad \alpha \in \mathcal{D} - \mathcal{Z}. \end{aligned} \tag{8}$$

If the maximum is $\delta^* = 0$, then there is no strictly positive solution. If the maximum δ^* is strictly positive, the solution exists and can be found through a standard algorithm based on the simplex method which yields a strictly positive probability table together with the value of δ^* . The algorithm above can be easily implemented with the `lpsolve` package, see [19] and [20], in R[21]. An alternative version of the optimization problem can be stated using separation theorems like the Farkas lemma, see, e.g., [22], but the refinement of the optimization problem is outside the scope of this work.

The approach via `lpsolve` is quite efficient and allows finding a solution even for large binary tables. For example, the analysis for $d = 12$ is performed in less than 1 sec. on standard hardware. For illustrative purposes, we conclude with two examples in the case $d = 4$. They are depicted in Fig. 2. In the upper panel, Fig. 2 shows a table with 4 zeros on a two-dimensional sub-table as zero pattern. According to Prop. 3.1, there is no strictly positive solution, and the linear programming algorithm yields a value of $\delta^*=0$. In the lower panel, Fig. 2 shows a different zero pattern, which is a table with again 4 zeros but in different positions. In this case, there is a strictly positive solution, and the output of the optimization procedure is $\delta^* = 0.07143$.

In summary, a key advantage of the linear programming-based method is its computational efficiency. Conversely, the method based on extreme pmfs offers valuable insight into the structure of zero patterns that are compatible with the existence of tables with uniform margins. The computation of extreme pmfs for the class of multivariate Bernoulli distributions with given margins, particularly in high dimensions, can also benefit from recent work (e.g., see [23]). We plan to further develop this method in the future by exploring this connection further.

3.2. Examples in the 2^3 case

In this section, we show how some 2^3 tables with only a few zeros can behave differently. Specifically, we give explicit configurations where the existence and uniqueness of the solution are not guaranteed. For our goal, it is sufficient to analyze 2^3 tables with one or two zeros, leading (up to isomorphism) to four possible configurations. We recall that, in our context, two tables are isomorphic if one can be obtained from the other by permuting the levels and relabeling the variables.

1. The table in Fig. 3 (a) has one zero. In this configuration, there are still four independent equations from marginal constraints and three independent odd ratios equations from the system in Eq. (2). Thus, the number of equations is equal to the number of positive probabilities.
2. The table in Fig. 3 (b) has two zero on the same edge, specifically in the entries $(0, 1, 0)$ and $(1, 1, 0)$. According to Prop. 3.1, there is no solution unless we also force the two opposite cells in $(0, 0, 1)$ and $(1, 0, 1)$ to be zero.
3. The table in Fig. 3 (c) has two zeros on the opposite cells of a face. In this configuration, there are four independent equations from marginal constraints and one independent odds ratios equation




Figure 2: Two 2^4 tables with zero patterns. The first table in the upper panel, the second table in the lower panel. The variable X_1, X_2, X_3 are the coordinates of the cubes, the variable X_4 divides the table into two cubes. Red dots denote zero probabilities. To ease readability, vertex labels are omitted.

from the system in Eq. (2). Since the positive probabilities are six, in this case there is no uniqueness. We discuss this example in full detail in Sect. 4.

4. The table in Fig. 3 (d) has two zeros on two different faces. Here, there are no valid constraints on the odds ratios. The problem reduces to the search for a table with uniform margins, and the number of positive probabilities is two more than the number of marginal constraints.

4. Uniqueness

In the previous section we focused on the existence of the solution. However, also uniqueness is an issue in the multidimensional case, as mentioned in the illustration of the 3-dimensional examples above. Here, we consider only an example where uniqueness does not hold and provide a description of the set of solutions. Although IPFP is known to provide a solution with good theoretical properties, it can be used to emphasize that the solution of our problem is not unique and there are zero patterns for which there are infinite probability tables with uniform margins and given odds ratio structure.

We consider a 2^3 table with zeros in the entries $(1, 0, 0)$ and $(0, 1, 0)$. In this case we have one non-trivial condition on the odds ratios given by

$$\omega_1^{12} = \frac{p_{001}p_{111}}{p_{011}p_{101}}. \quad (9)$$

Additionally, we have four independent conditions for the margins. Again using the lexicographic order for the entries, the kernel of the model matrix is generated by the two vectors:

$$(-1, 1, 0, 0, 1, -1), \quad (0, 1, -1, -1, 0, 1).$$

Given a particular solution \tilde{p} , for instance the solution of the IPFP which is strictly positive, the general solution to the problem with fixed margins is given by

$$(p_{000} = \tilde{p}_{000} - \alpha, p_{001} = \tilde{p}_{001} + \alpha + \beta, p_{011} = \tilde{p}_{011} - \beta, \\ p_{101} = \tilde{p}_{101} - \beta, p_{110} = \tilde{p}_{110} + \alpha, p_{111} = \tilde{p}_{111} - \alpha + \beta) \quad (10)$$




Figure 3: Configurations with 1 or 2 zeros in the 2^3 table. Red dots denote zero probabilities. (a): one zero; (b): two zeros on the same edge; (c): two zeros in diagonal position on a face; (d): two zeros in different faces.

for suitable values of (α, β) . This distribution is shown in Fig. 4. By substitution of the generic probability distribution in Eq. (10) into the odds ratio equation in Eq. (9), we get:

$$f(\alpha, \beta) = (\tilde{p}_{111} - \tilde{p}_{001})\alpha + (\tilde{p}_{111} + \tilde{p}_{001} + \omega_1^{12}(\tilde{p}_{011} + \tilde{p}_{101}))\beta - \alpha^2 + (1 - \omega_1^{12})\beta^2 = 0. \quad (11)$$

The equation above introduces an implicit function $\beta(\alpha)$, whose explicit expression is somewhat complicated and not relevant for our application, having quadratic terms in both α and β . However, we can see that, at least locally, we have infinite solutions in a neighborhood of the point $(0, 0)$. The point $(0, 0)$ is a solution because \tilde{p} is the probability distribution given by the IPFP. Since \tilde{p} is strictly positive, there is a neighborhood of $(0, 0)$ in the strictly positive simplex of the probabilities. The local behavior of $\beta(\alpha)$ can be studied by means of the implicit functions theorem. The partial derivatives are

$$\frac{\partial f}{\partial \alpha} = \tilde{p}_{111} - \tilde{p}_{001} - 2\alpha, \quad \frac{\partial f}{\partial \beta} = \tilde{p}_{111} + \tilde{p}_{001} + \omega_1^{12}(\tilde{p}_{011} + \tilde{p}_{101}) + 2(1 - \omega_1^{12})\beta,$$

leading to

$$\beta(0) = 0, \quad \beta'(0) = -\frac{\tilde{p}_{111} - \tilde{p}_{001}}{\tilde{p}_{111} + \tilde{p}_{001} + \omega_1^{12}(\tilde{p}_{011} + \tilde{p}_{101})}.$$




Figure 4: The general probability table in the 2^3 case with two zeros in the diagonal of a face.

A special case arises for conditional independence, i.e., $\omega_1^{12} = 1$ where the function in Eq. (11) reduces to

$$f(\alpha, \beta) = (\tilde{p}_{111} - \tilde{p}_{001})\alpha + \frac{1}{2}\beta - \alpha^2 = 0$$

from which

$$\beta(\alpha) = 2\alpha(\alpha - (\tilde{p}_{111} - \tilde{p}_{001})).$$

A more general approach would involve the use of Markov bases for the complete table and Markov moves in the continuous setting, as done in [24]. Indeed, Markov moves allow for considering tables in the closed simplex. However, a systematic analysis of the uniqueness issues with Algebraic Statistics is outside the aims of the present paper.

From the previous example and the discussion in Sect. 3.2, it is evident that there are cases where the solution space has an even higher dimension. For example, in the table drawn in panel (d) of Fig. 3, the difference in dimension is 2. However, this case is less relevant in our context, as none of the odds ratio equations remains. As a result, the problem reduces to a problem that involves only the search for a probability table with fixed uniform margins. Although this situation may still be of interest in broader statistical analysis, it does not pose additional challenges within our framework. Therefore, we primarily focus on cases where odds ratio constraints play a fundamental role in shaping the structure of the solution space.

5. Conclusions

In this paper, we introduce new theoretical insights that demonstrate the impact of zero patterns on both the existence and uniqueness of a transformed multi-way binary table with prescribed dependence structure in terms of odds ratios. While working on this project, we identified interesting research directions for future investigations. A natural question relates the extension of the proposed approach to generic d -way tables. In this regard, it is worth noting that both computational methods described in Sect. 3.1 can be generalized to any d -way table, where each variable X_i takes values in $0, 1, \dots, x_i - 1$ for $x_i \geq 2$, with $i = 1, \dots, d$. For this, it is sufficient to modify H_d and C_d to express the constraints

$$m_{i,0} = \dots = m_{i,x_i-1} = \frac{1}{x_i}, \quad i = 1, \dots, d.$$

From a theoretical viewpoint, the combinatorial approach is more powerful in obtaining necessary and sufficient conditions for the existence of a solution. However, such an approach needs further investigations from the combinatorial side, namely the characterization of the extreme rays to avoid the computational step. Some advances in this direction have recently been made in the context of multivariate Bernoulli distributions [23]. The possible adaptation of these results in the framework of discrete copulas is worth investigating. With these new insights, it would be possible to prove if the condition on the opposite facets of Prop. 3.1 is also sufficient and if it can be generalized to multi-way tables with an arbitrary number of levels. Moreover, it would be interesting to characterize the entire class solutions when uniqueness does not hold in terms of the associated algebraic variety. An algebraic approach would make it possible to identify extreme tables in this variety and study if there are valuable alternatives with respect to the solution provided by the IPFP. Finally, we could investigate how the approach adapts to different dependence conditions. For instance, we might consider suitable marginal odds ratios to have a closer connection with the classical log-linear models of independence and conditional independence for multi-way contingency tables.

References

- [1] T. Rudas, Lectures on categorical data analysis, Springer, 2018.
- [2] G. Geenens, Copula modeling for discrete random vectors, *Depend. Model.* 8 (1) (2020) 417–440.
- [3] A. Sklar, Fonctions de répartition à n dimensions et leurs marges, *Publications de l'Institut de Statistique de Paris* 8 (1959) 229–231.

- [4] I. Kojadinovic, T. Martini, Copula-like inference for discrete bivariate distributions with rectangular supports, *Electron. J. Stat.* 18 (1) (2024) 2571–2619. doi:10.1214/24-EJS2261.
URL <https://doi.org/10.1214/24-EJS2261>
- [5] G. U. Yule, On the methods of measuring association between two attributes, *Journal of the Royal Statistical Society* 75 (6) (1912) 579–652.
- [6] S. Sullivant, *Algebraic Statistics*, Vol. 194 of Graduate Studies in Mathematics, American Mathematical Society, Providence, RI, 2018.
- [7] E. Perrone, R. Fontana, F. Rapallo, Multi-way contingency tables with uniform margins, in: J. Ansari, S. Fuchs, W. Trutschnig, M. A. Lubiano, M. Á. Gil, P. Grzegorzewski, O. Hryniewicz (Eds.), *Combining, Modelling and Analyzing Imprecision, Randomness and Dependence*, Springer Nature Switzerland, Cham, 2024, pp. 349–356.
- [8] R. Fontana, E. Perrone, F. Rapallo, Contingency tables with structural zeros and discrete copulas, in: *Book of the short papers - SIS 2023*, June 21–23, 2023, Ancona (Italy), Pearson, 2023, pp. 713–718.
- [9] E. Perrone, F. Durante, Extreme points of polytopes of discrete copulas, in: *Joint Proceedings of the 19th World Congress of the International Fuzzy Systems Association (IFSA), the 12th Conference of the European Society for Fuzzy Logic and Technology (EUSFLAT), and the 11th International Summer School on Aggregation Operators (AGOP)*, Atlantis Press, 2021, pp. 596–601.
- [10] E. Perrone, Polytopes of discrete copulas and applications, in: L. A. García-Escudero, A. Gordaliza, A. Mayo, M. A. Lubiano Gomez, M. A. Gil, P. Grzegorzewski, O. Hryniewicz (Eds.), *Building Bridges between Soft and Statistical Methodologies for Data Science*, Springer International Publishing, Cham, 2023, pp. 319–325.
- [11] J. A. De Loera, E. D. Kim, Combinatorics and geometry of transportation polytopes: An update, in: *Discrete Geometry and Algebraic Combinatorics*, American Mathematical Society, Providence, RI, 2014, pp. 37–76.
- [12] E. Perrone, L. Solus, C. Uhler, Geometry of discrete copulas, *J. Multivar. Anal.* 172 (2019) 162–179.
- [13] S. E. Fienberg, An iterative procedure for estimation in contingency tables, *Ann. Math. Stat.* 41 (3) (1970) 907–917. doi:10.1214/aoms/1177696968.
- [14] S. E. Fienberg, M. M. Meyer, *Iterative Proportional Fitting*, John Wiley & Sons, Ltd, 2006. doi:<https://doi.org/10.1002/0471667196.ess1297.pub2>.
- [15] S. E. Fienberg, A. Rinaldo, Maximum likelihood estimation in log-linear models, *Ann. Stat.* 40 (2) (2012) 996–1023. doi:10.1214/12-AOS986.
URL <https://doi.org/10.1214/12-AOS986>
- [16] J. I. Coons, C. Langer, M. Ruddy, Classical iterative proportional scaling of log-linear models with rational maximum likelihood estimator, *Int. J. Approx. Reason.* 164 (2024) 109043. doi:<https://doi.org/10.1016/j.ijar.2023.109043>.
URL <https://www.sciencedirect.com/science/article/pii/S0888613X23001743>
- [17] J. Franklin, J. Lorenz, On the scaling of multidimensional matrices, *Linear Algebra Its Appl.* 114/115 (1989) 717–735.
- [18] 4ti2 team, 4ti2—a software package for algebraic, geometric and combinatorial problems on linear spaces.
URL <https://4ti2.github.io>
- [19] G. Csárdi, M. Berkelaar, lpSolve: Interface to 'Lp_solve' v. 5.5 to Solve Linear/Integer Programs, R package version 5.6.23 (2024).
URL <https://CRAN.R-project.org/package=lpSolve>
- [20] S. E. Buttrey, Calling the lp_solve linear program software from R, S-PLUS and Excel, *J. Stat. Softw.* 14 (4) (2005) 1–13. doi:10.18637/jss.v014.i04.
- [21] R Core Team, *R: A Language and Environment for Statistical Computing*, R Foundation for Statistical Computing, Vienna, Austria (2024).
URL <https://www.R-project.org/>
- [22] O. L. Mangasarian, *Nonlinear Programming*, Vol. 10 of Classics in Applied Mathematics, Society for Industrial and Applied Mathematics, Philadelphia, 1994. doi:10.1137/1.9781611971255.
- [23] R. Fontana, P. Semeraro, High dimensional bernoulli distributions: Algebraic representation and applications, *Bernoulli* 30 (1) (2024) 825–850.
- [24] G. Pistone, F. Rapallo, M. P. Rogantin, Finite space kantorovich problem with an MCMC of table moves, *Electron. J. Stat.* 15 (1) (2021) 880–907. doi:10.1214/21-EJS1804.