# Non Local Mixed Systems with Neumann Boundary Conditions

Rinaldo M. Colombo<sup>1</sup>

Elena Rossi<sup>2</sup>

Abraham Sylla<sup>1</sup>

#### Abstract

We prove well posedness and stability in  $\mathbf{L}^1$  for a class of mixed hyperbolic-parabolic non linear and non local equations in a bounded domain with *no flow* along the boundary. While the treatment of boundary conditions for the hyperbolic equation is standard, the extension to  $\mathbf{L}^1$  of classical results about parabolic equations with Neumann conditions is here achieved.

2020 Mathematics Subject Classification: 35M30, 35L04, 35K20

Keywords: Mixed Hyperbolic-Parabolic Initial Boundary Value Problems; Parabolic Problems with Neumann Boundary Conditions in  $\mathbf{L}^1$ ; Non Local Mixed Boundary Value Problems.

### **1** Introduction

We consider the following non linear and non local problem on a bounded domain  $\Omega \subset \mathbb{R}^n$ 

$$\begin{cases} \partial_t u + \nabla \cdot \left( u \, v \left( t, w(t) \right) \right) = \alpha \left( t, x, w(t) \right) \, u + a(t, x) \,, \\ \partial_t w - \mu \, \Delta w = \beta \left( t, x, u(t), w(t) \right) \, w + b(t, x) \,, \end{cases} \quad (t, x) \in [0, T] \times \Omega \,. \tag{1.1}$$

When n = 2, 3, this mixed system is motivated by a variety of predator-prey models. Indeed, for instance, u = u(t, x) can be the density of a population that *chases* the other population w = w(t, x). This chase is described through the non local operator v, able to model the movement of u towards regions where the concentration of w is higher. A change in the sign of v allows to model the case where the population u escapes from w. The w population diffuses isotropically in all directions. The source terms  $\alpha$  and  $\beta$  account for natality, mortality or predation, while a and b may describe controls acting on the system, which consist in the introduction of the two species at desired times and locations.

We equip problem (1.1) with the following initial data and conditions at the boundary:

$$\begin{cases} u(0,x) = u_o(x) \\ w(0,x) = w_o(x) \end{cases} \quad x \in \Omega, \qquad \begin{cases} u(t,\xi) = 0 \\ \nabla w(t,\xi) \cdot \nu(\xi) = 0 \end{cases} \quad (t,\xi) \in \left]0, T\right[ \times \partial\Omega. \quad (1.2) \end{cases}$$

<sup>&</sup>lt;sup>1</sup>INdAM Unit & Department of Information Engineering, University of Brescia, via Branze, 38, 25123 Brescia, Italy. Email: rinaldo.colombo@unibs.it and abraham.sylla@unibs.it

<sup>&</sup>lt;sup>2</sup>University of Modena and Reggio Emilia, INdAM Unit & Department of Sciences and Methods for Engineering, Via Amendola 2 – Pad. Morselli, 42122 Reggio Emilia, Italy. Email: elena.rossi13@unimore.it

The present choice of Neumann boundary conditions for the parabolic equation is motivated by the *no flow* requirement typically suited to the physical setting considered. Recall that boundary conditions for the hyperbolic equation bear an entirely different meaning, since they can be essentially neglected whenever characteristics exit the domain, see [1, 19, 21, 25]. Thus, the first boundary condition in (1.2) does not prevent u from exiting  $\Omega$ .

The analytical treatment of (1.1) relies on that of the separate hyperbolic and parabolic equations

$$\partial_t u + \nabla \cdot \left( u c(t, x) \right) = A(t, x) u + a(t, x) \quad \text{and} \quad \partial_t w - \mu \Delta w = B(t, x) w + b(t, x) \quad (1.3)$$

together with the initial and boundary conditions (1.2). While general well posedness results related to the former equation are available in the literature, the treatment in  $\mathbf{L}^1$  of the latter equation with Neumann boundary conditions has received far less attention. Indeed, since the classical books [11, 17], the literature offers a variety of results in  $\mathbf{L}^2$ , a choice which is hardly justifiable in the present physical setting. Here, on the contrary, the  $\mathbf{L}^1$  norm has a clear meaning but  $\mathbf{L}^1$  stability estimates were not available, not even in [11, 17], especially in the case of Neumann boundary conditions. Also the general mixed hyperbolic-parabolic setting in [10] does not comprise the well posedness of (1.1). Therefore, we provide a definition of weak solution to the parabolic equation and, correspondingly, we develop a well posedness and stability theory in the  $\mathbf{L}^1$  norm. In this procedure, clearly, the many properties that follow from reflexivity can not be used. However, it is remarkable that the weak completeness of  $\mathbf{L}^1$  plays a key role in our treatment of the parabolic equation.

Note that in (1.1) the dependence of v,  $\alpha$  and  $\beta$  on u(t) and w(t) is of a functional nature, allowing for non local dependencies. Indeed, u(t) and w(t) in (1.1) denote the functions  $x \mapsto u(t, x)$  and  $x \mapsto w(t, x)$ , both defined on all  $\Omega$ .

Systems of this form arise, for instance, in predator-prey models [5] and can be used in the control of parasites, see [7, 20]. A similar mixed hyperbolic-parabolic system is considered, in one space dimension, in [14], where Euler equations substitute the balance law in (1.1). A mixed ODE – parabolic PDE predator-prey model with Neumann boundary conditions is presented in [24]: here the predators' movement is the superposition of a directed hunting and a random dispersion. The present model (1.1)-(1.2) is applicable also to the setting of pursuit-evasion games, similarly to [12]. Here, however, the movement of the pursuer is not purely diffusive, but it is directed towards the average gradient of the evaders' density.

A mixed hyperbolic-parabolic system motivated by population dynamics is considered in [15], where the theoretical framework is set in  $\mathbf{L}^2$  and the theory of *m*-accretive operators is the key analytic tool. A more applied result is [16], where the description of an aneurysm leads to a mixed hyperbolic-parabolic system in 1 space dimension. Global classical solutions to a parabolic predator-prey system, under Neumann boundary conditions, are exhibited in [18], motivated by the dynamics of competing populations with repulsive chemotaxis. In the  $\mathbf{L}^2$  framework, local in time well posedness of a mixed hyperbolic-parabolic system is obtained in [9], by means of a Cauchy sequence of approximate solutions. A mixed ellipticparabolic problem also with biological motivations is studied in [23].

The next section presents the main analytical results. System (1.1) is split in the 2 equations (1.3). The former one is dealt with by means of results mainly coming from the literature, see § 2.1. On the contrary, in § 2.2 we present results on the parabolic part developed *ad hoc* in  $\mathbf{L}^1$  for the purposes of the present work. Finally, all proofs are deferred to Section 3.

# 2 Main Results

Throughout, the following notation is used.  $\mathbb{R}_+ = [0, +\infty[$ . If  $A \subseteq \mathbb{R}^n$ , the characteristic function  $\chi_A$  is defined by  $\chi_A(x) = 1$  if and only if  $x \in A$  and  $\chi_A(x) = 0$  if and only if  $x \in \mathbb{R}^n \setminus A$ . For  $x_o \in \mathbb{R}^n$  and r > 0,  $B(x_o, r)$  is the open sphere centered at  $x_o$  with radius r.

Fix T > 0. We pose the following assumptions on  $\Omega$  and on the functions appearing in problem (1.1):

- ( $\Omega$ )  $\Omega$  is a non empty, bounded and connected open subset of  $\mathbb{R}^n$ , with  $\mathbf{C}^{1,\gamma}$  boundary, for a  $\gamma \in [0,1]$ .
- (v)  $v: [0,T] \times \mathbf{L}^{1}(\Omega; \mathbb{R}) \to (\mathbf{C}^{2} \cap \mathbf{W}^{1,\infty})(\Omega; \mathbb{R}^{n})$  is such that for a constant  $K_{v} > 0$  and for a map  $k_{v} \in \mathbf{L}^{\infty}_{\mathbf{loc}}([0,T] \times \mathbb{R}_{+}; \mathbb{R}_{+})$  non decreasing in each argument, for all  $t, t_{1}, t_{2} \in [0,T]$  and  $w, w_{1}, w_{2} \in \mathbf{L}^{1}(\Omega; \mathbb{R}),$

$$\begin{aligned} \|v(t,w)\|_{\mathbf{L}^{\infty}(\Omega;\mathbb{R}^{n})} &\leq K_{v} \|w\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} \\ \|D_{x}v(t,w)\|_{\mathbf{L}^{\infty}(\Omega;\mathbb{R}^{n})} &\leq K_{v} \|w\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} \\ \|v(t_{1},w_{1})-v(t_{2},w_{2})\|_{\mathbf{L}^{\infty}(\Omega;\mathbb{R}^{n})} &\leq K_{v} \left(|t_{1}-t_{2}|+\|w_{1}-w_{2}\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})}\right) \\ \|D_{x}^{2}v(t,w)\|_{\mathbf{L}^{1}(\Omega;\mathbb{R}^{n\times n})} &\leq k_{v} \left(t,\|w\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})}\right) \|w\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} \\ \|\nabla \cdot \left(v(t_{1},w_{1})-v(t_{2},w_{2})\right)\|_{\mathbf{L}^{\infty}(\Omega;\mathbb{R})} &\leq k_{v} \left(t,\max_{i=1,2}\|w_{i}\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})}\right) \|w_{1}-w_{2}\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})}. \end{aligned}$$

( $\boldsymbol{\alpha}$ )  $\boldsymbol{\alpha} : [0,T] \times \Omega \times \mathbf{W}^{1,1}(\Omega; \mathbb{R}) \to \mathbb{R}$  admits a constant  $K_{\boldsymbol{\alpha}} > 0$  such that, for a.e.  $t \in [0,T]$ , for a.e.  $x \in \Omega$  and all  $w_1, w_2 \in \mathbf{W}^{1,1}(\Omega; \mathbb{R})$ ,

$$\sup_{x \in \Omega} \left| \alpha(t, x, w_1) - \alpha(t, x, w_2) \right| \le K_\alpha \| w_1 - w_2 \|_{\mathbf{L}^1(\Omega; \mathbb{R})}$$

and there exists  $k_{\alpha} \in \mathbf{L}^1([0,T]; \mathbb{R}_+)$  such that for all  $w \in \mathbf{W}^{1,1}(\Omega; \mathbb{R})$  and a.e.  $t \in [0,T]$ ,

$$\operatorname{TV}\left(\alpha(t,\cdot,w)\right) \leq K_{\alpha}\left(1+\|\nabla w\|_{\mathbf{L}^{1}(\Omega;\mathbb{R}^{n})}\right)$$
  
 
$$\operatorname{ess\,sup}_{x\in\Omega}\left|\alpha(t,x,w)\right| \leq k_{\alpha}(t)\left(1+\|w\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})}\right).$$

(a)  $a \in \mathbf{L}^1([0,T]; \mathbf{L}^{\infty}(\Omega; \mathbb{R}))$  and  $t \mapsto \mathrm{TV}(a(t))$  is in  $\mathbf{L}^1([0,T]; \mathbb{R})$ .

( $\beta$ )  $\beta$ :  $[0,T] \times \Omega \times \mathbf{L}^{1}(\Omega; \mathbb{R}) \times \mathbf{L}^{1}(\Omega; \mathbb{R}) \to \mathbb{R}$  admits a constant  $K_{\beta} > 0$  and a  $k_{\beta} \in \mathbf{L}^{1}([0,T]; \mathbb{R}_{+})$  such that, for a.e.  $t \in [0,T]$ , all  $x \in \Omega$  and  $u, u_{1}, u_{2}, w, w_{1}, w_{2} \in \mathbf{L}^{1}(\Omega; \mathbb{R})$ ,

(b)  $b \in \mathbf{L}^1([0,T] \times \Omega; \mathbb{R}).$ 

Since (1.1) is the coupling of a hyperbolic and a parabolic problem, the following definition of solution to (1.1) is a sort of gluing of the definitions of solutions to the equations in (1.3)

**Definition 2.1.** A pair  $(u, w) \in \mathbb{C}^0([0, T]; \mathbb{L}^1(\Omega; \mathbb{R}^2))$  is a solution to problem (1.1)–(1.2) if, setting

$$c(t,x) = v\left(t,w(t)\right)(x), \qquad A(t,x) = \alpha\left(t,x,w(t)\right), \qquad B(t,x) = \beta\left(t,x,u(t),w(t)\right),$$

the function u solves, according to Definition 2.3, the problem

$$\begin{aligned}
\partial_t u + \nabla \cdot (u \ c(t, x)) &= A(t, x)u + a(t, x) & (t, x) \in [0, T] \times \Omega \\
u(t, \xi) &= 0 & (t, \xi) \in ]0, T[\times \partial\Omega & (2.1) \\
u(0, x) &= u_o(x) & x \in \Omega
\end{aligned}$$

and the function w solves, according to Definition 2.5, the problem

$$\partial_t w - \mu \,\Delta w = B(t, x)w + b(t, x) \qquad (t, x) \in [0, T] \times \Omega$$
  

$$\nabla w(t, \xi) \cdot \nu(\xi) = 0 \qquad (t, \xi) \in ]0, T[\times \partial \Omega \qquad (2.2)$$
  

$$w(0, x) = w_o(x) \qquad x \in \Omega.$$

We now state the main result of this work: it ensures the well posedness in  $\mathbf{L}^1$  of (1.1) and provides stability estimates to be used, for instance, in control problems based on (1.1).

**Theorem 2.2.** Fix T > 0. Assume that  $(\Omega)$ , (v),  $(\alpha)$ ,  $(\beta)$  and (b) hold. Let  $(u_o, w_o) \in (\mathbf{L}^1 \cap \mathbf{BV})(\Omega; \mathbb{R}) \times \mathbf{L}^1(\Omega; \mathbb{R})$ . Then:

- (M1) Problem (1.1) admits a unique solution (u, w) in the sense of Definition 2.1.
- (M2) The map  $t \mapsto (u, w)(t)$  is in  $\mathbf{C}^0([0, T]; \mathbf{L}^1(\Omega; \mathbb{R}^2))$ .
- (M3) If  $(u_o^1, w_o^1), (u_o^2, w_o^2) \in (\mathbf{L}^1 \cap \mathbf{BV})(\Omega; \mathbb{R}) \times \mathbf{L}^1(\Omega; \mathbb{R})$ , the corresponding solutions  $(u_1, w_1)$ and  $(u_2, w_2)$  satisfy the estimate:

$$\left\| (u_1, w_1)(t) - (u_2, w_2)(t) \right\|_{\mathbf{L}^1(\Omega; \mathbb{R}^2)} \le C(t) \left\| (u_o^1, w_o^1) - (u_o^2, w_o^2) \right\|_{\mathbf{L}^1(\Omega; \mathbb{R}^2)}$$

where  $C \in \mathbf{L}^{\infty}([0,T]; \mathbb{R}_{+})$  depends on  $\mu$ ,  $(\Omega)$ ,  $(\boldsymbol{v})$ ,  $(\boldsymbol{\alpha})$ ,  $(\boldsymbol{\beta})$ ,  $(\boldsymbol{a})$ ,  $(\boldsymbol{b})$  and on  $\mathrm{TV}(u_{o}^{1})$ ,  $\mathrm{TV}(u_{o}^{2})$ ,  $\left\|(u_{o}^{1}, w_{o}^{1})\right\|_{\mathbf{L}^{1}(\Omega; \mathbb{R}^{2})}$ ,  $\left\|(u_{o}^{2}, w_{o}^{2})\right\|_{\mathbf{L}^{1}(\Omega; \mathbb{R}^{2})}$ .

(M4) If  $a_1, a_2$  satisfy (a) and  $b_1, b_2$  satisfy (b), the corresponding solutions  $(u_1, w_1)$  and  $(u_2, w_2)$  satisfy the estimate:

$$\begin{aligned} & \left\| (u_1, w_1)(t) - (u_2, w_2)(t) \right\|_{\mathbf{L}^1(\Omega; \mathbb{R}^2)} \le C(t) \left( \|a_1 - a_2\|_{\mathbf{L}^1([0,t] \times \Omega; \mathbb{R})} + \|b_1 - b_2\|_{\mathbf{L}^1([0,t] \times \Omega; \mathbb{R})} \right) \\ & \text{where } C \in \mathbf{L}^{\infty}([0,T]; \mathbb{R}_+) \text{ depends on } \mu, \ (\mathbf{\Omega}), \ (\mathbf{v}), \ (\mathbf{\alpha}), \ (\boldsymbol{\beta}), \ (\mathbf{a}), \ (\mathbf{b}) \text{ and on } \mathrm{TV}(u_o), \\ & \left\| (u_o, w_o) \right\|_{\mathbf{L}^1(\Omega; \mathbb{R}^2)}. \end{aligned}$$

(M5) If  $k_{\beta}$  in ( $\beta$ ) is bounded and  $u_o \ge 0$ ,  $w_o \ge 0$ ,  $a \ge 0$  and  $b \ge 0$ , then for a.e.  $t \in [0,T]$ , the solution (u, w) satisfies  $u(t) \ge 0$  and  $w(t) \ge 0$ .

Remark that if the various assumptions hold on the time interval  $\mathbb{R}_+$ , then Theorem 2.2 ensures well posedness on  $\mathbb{R}_+$ , with the function *C* appearing in (M3) and (M4) belonging to  $\mathbf{L}_{\mathbf{loc}}^{\infty}(\mathbb{R}_+, \mathbb{R}_+)$ .

To deal with non local operators on a bounded domain  $\Omega$ , the modified convolution introduced in [6, § 3] is an adequate tool. For a function  $\rho \in \mathbf{L}^1(\Omega; \mathbb{R})$  and a smooth kernel  $\eta$ , it reads

$$(\rho *_{\Omega} \eta)(x) = \frac{\int_{\Omega} \rho(y) \eta(x-y) \,\mathrm{d}y}{\int_{\Omega} \eta(x-y) \,\mathrm{d}y}.$$
(2.3)

The quantity  $(\rho *_{\Omega} \eta)(x)$  is an average of the values attained by  $\rho$  in  $\Omega$  around x as soon as the kernel  $\eta$  satisfies, for instance,

( $\eta$ )  $\eta(x) = \tilde{\eta}(||x||)$ , where  $\tilde{\eta} \in \mathbf{C}^3(\mathbb{R}_+; \mathbb{R})$ , spt  $\tilde{\eta} = [0, \ell], \ \ell > 0, \ \tilde{\eta}' \le 0, \ \tilde{\eta}'(0) = \tilde{\eta}''(0) = 0$  and  $\int_{\mathbb{R}^n} \eta(\xi) \, d\xi = 1.$ 

It is often reasonable to assume that u hunts w moving towards areas with higher density of w, or else that u escapes from w towards regions with lower w density. Thus, v is parallel to the average gradient of w in  $\Omega$ , such as

$$v(t,w) = k(t) \frac{\nabla (w *_{\Omega} \eta)}{\sqrt{1 + \left\|\nabla (w *_{\Omega} \eta)\right\|^2}}$$
(2.4)

where, for instance,

$$\eta(x) = \overline{\eta} \left( 1 - (\|x\|/\ell)^4 \right)^4 \chi_{B(0,\ell)}(x) \,. \tag{2.5}$$

Here,  $\ell$  has the clear physical meaning of the distance, or *horizon*, at which individuals of the u population *feel* the presence of the w population. The normalization parameter  $\overline{\eta}$  is chosen so that  $\int_{\mathbb{R}^n} \eta(x) \, \mathrm{d}x = 1$ . A choice like (2.4) is consistent with the requirements  $(\boldsymbol{v})$ , as proved in [6, Lemma 3.2].

### 2.1 The Hyperbolic Problem

We focus on the hyperbolic problem (2.1). For completeness, we present the standard definition of solution and a detailed well posedness result based on the current literature. Precise references are provided in § 3.1.

**Definition 2.3.** A function  $u \in \mathbf{L}^{\infty}([0,T] \times \Omega; \mathbb{R})$  is a solution to (2.1) if for any test function  $\varphi \in \mathbf{C}_{c}^{1}(]-\infty, T[\times \Omega; \mathbb{R}),$ 

$$\int_0^T \int_\Omega \left( u \left( \partial_t \varphi + c \cdot \nabla \varphi \right) + (A \, u + a) \varphi \right) \mathrm{d}x \, \mathrm{d}t + \int_\Omega u_o(x) \, \varphi(0, x) \, \mathrm{d}x = 0. \tag{2.6}$$

**Theorem 2.4.** Fix T > 0. Assume (a) and

- (c)  $c \in (\mathbf{C}^0 \cap \mathbf{L}^\infty) ([0,T] \times \Omega; \mathbb{R}^n), c(t) \in \mathbf{C}^1(\Omega; \mathbb{R}^n) \text{ for all } t \in [0,T], D_x c \in \mathbf{L}^\infty([0,T] \times \Omega; \mathbb{R}^{n \times n}).$
- (A)  $A \in \mathbf{L}^1([0,T]; \mathbf{L}^{\infty}(\Omega; \mathbb{R}))$  and  $t \mapsto \mathrm{TV}(A(t))$  is in  $\mathbf{L}^1([0,T]; \mathbb{R})$ .

Then, for all  $u_o \in \mathbf{L}^1(\Omega; \mathbb{R})$ , problem (2.1) admits a unique solution in the sense of Definition 2.3. Moreover

(H1) For all  $t \in [0, T]$ ,

$$\left\| u(t) \right\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} \leq \left( \left\| u_{o} \right\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} + \left\| a \right\|_{\mathbf{L}^{1}([0,t]\times\Omega;\mathbb{R})} \right) \exp \left( \left\| A \right\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}))} \right) .$$
(2.7)

(H2) For all  $t \in [0,T]$ , if  $u_o \in \mathbf{L}^{\infty}(\Omega; \mathbb{R})$ ,

$$\begin{aligned} \|u(t)\|_{\mathbf{L}^{\infty}(\Omega;\mathbb{R})} &\leq \left( \|u_o\|_{\mathbf{L}^{\infty}(\Omega;\mathbb{R})} + \|a\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}))} \right) \\ &\times \exp\left( \|A\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}))} + \|\nabla \cdot c\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}))} \right) . \end{aligned}$$

$$(2.8)$$

**(H3)** If  $u_o \ge 0$  and  $a \ge 0$ , then  $u \ge 0$ .

**(H4)** If  $u_o^1, u_o^2 \in \mathbf{L}^1(\Omega; \mathbb{R})$ , then the corresponding solutions  $u_1, u_2$  satisfy for all  $t \in [0, T]$ ,

$$\left\| u_{1}(t) - u_{2}(t) \right\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} \leq \left\| u_{o}^{1} - u_{o}^{2} \right\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} \exp\left( \|A\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}))} \right).$$
(2.9)

(H5) If  $A_1, A_2$  satisfy (A) and  $a_1, a_2$  satisfy (a), the corresponding solutions  $u_1$  and  $u_2$  satisfy for all  $t \in [0, T]$ 

$$\begin{aligned} & \left\| u_{1}(t) - u_{2}(t) \right\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} \\ & \leq \exp\left( \max\left\{ \|A_{1}\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}))}, \|A_{2}\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}))} \right\} \right) \\ & \times \left( \|u_{o}\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} + \|a_{1}\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}))} \right) \|A_{2} - A_{1}\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}))} \\ & + \exp\left( \|A_{2}\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}))} \right) \|a_{1} - a_{2}\|_{\mathbf{L}^{1}([0,t]\times\Omega;\mathbb{R})} . \end{aligned}$$

- (H6) If  $u_o \in \mathbf{BV}(\Omega; \mathbb{R})$ ,  $c(t) \in \mathbf{C}^2(\Omega; \mathbb{R}^n)$  for all  $t \in [0, T]$  and  $\nabla \nabla \cdot c \in \mathbf{L}^1([0, T] \times \Omega; \mathbb{R}^n)$ , then the map  $t \to u(t)$  is locally Lipschitz continuous in  $\mathbf{L}^1(\Omega; \mathbb{R})$ .
- (H7) If  $u_o \in \mathbf{BV}(\Omega; \mathbb{R})$ ,  $c_1, c_2$  satisfy (c) and  $c_1(t), c_2(t) \in \mathbf{C}^2(\Omega; \mathbb{R}^n)$  for all  $t \in [0, T]$  and  $\nabla \nabla \cdot c_1, \nabla \nabla \cdot c_2 \in \mathbf{L}^1([0, T] \times \Omega; \mathbb{R}^n)$ , then the corresponding solutions  $u_1, u_2$  satisfy

$$\begin{aligned} &\|u_{2}(t) - u_{1}(t)\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} \\ \leq & C \left( \|D_{x}c_{1}\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}^{n\times n}))}, \|\nabla\nabla\cdot c_{1}\|_{\mathbf{L}^{1}([0,t]\times\Omega;\mathbb{R}^{n})}, \|A\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}))}, \int_{0}^{t} \mathrm{TV}A(\tau) \,\mathrm{d}\tau \right) \\ & \times \left( \|c_{1} - c_{2}\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}^{n}))} + \|\nabla\cdot (c_{1} - c_{2})\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}))} \right) \end{aligned}$$

where C also depends on  $\|a\|_{\mathbf{L}^1([0,t];\mathbf{L}^\infty(\Omega;\mathbb{R}))}$ ,  $\int_0^t \mathrm{TV}(a(\tau)) \,\mathrm{d}\tau$  and  $\|u_o\|_{\mathbf{L}^1(\Omega;\mathbb{R})}$ ,  $\|u_o\|_{\mathbf{L}^\infty(\Omega;\mathbb{R})}$ ,  $\mathrm{TV}(u_o)$ .

The proof is based on results from the literature detailed in § 3.1.

### 2.2 The Parabolic Problem

We focus on the  $L^1$  well posedness for the parabolic problem (2.2), first adapting the classical definition of weak solution, see for instance [3, § 2.3], to the case of interest here.

**Definition 2.5.** A function  $w \in \mathbf{L}^{\infty}([0,T]; \mathbf{L}^{1}(\Omega; \mathbb{R}))$  with  $w(t) \in \mathbf{W}^{1,1}(\Omega; \mathbb{R})$  for a.e.  $t \in [0,T]$  is a weak solution to (2.2) if for all test functions  $\varphi \in \mathbf{C}^{\infty}_{c}([0,T] \times \overline{\Omega}; \mathbb{R})$ 

$$\int_{0}^{T} \int_{\Omega} w \,\partial_{t} \varphi \,\mathrm{d}x \,\mathrm{d}t - \mu \int_{0}^{T} \int_{\Omega} \nabla w \cdot \nabla \varphi \,\mathrm{d}x \,\mathrm{d}t + \int_{0}^{T} \int_{\Omega} B \,w \,\varphi \,\mathrm{d}x \,\mathrm{d}t + \int_{0}^{T} \int_{\Omega} b \,\varphi \,\mathrm{d}x \,\mathrm{d}t + \int_{\Omega} w_{o}(x) \,\varphi(0,x) \,\mathrm{d}x = 0.$$

$$(2.10)$$

A relevant consequence of the definition chosen above is the following convergence result where the weak completeness of  $\mathbf{L}^1$  is essential. This Lemma is of use in a few key points in the proof of Theorem 2.7.

**Lemma 2.6.** For  $h \in \mathbb{N}$ , let  $b_h \in \mathbf{L}^1([0,T] \times \Omega; \mathbb{R})$ ,  $B_h \in \mathbf{L}^1([0,T]; \mathbf{L}^{\infty}(\Omega; \mathbb{R}))$  and  $w_o^h \in \mathbf{L}^1(\Omega; \mathbb{R})$  be such that (2.2) admits the solution  $w_h$  in the sense of Definition 2.5. Moreover, assume that

$$\lim_{h \to +\infty} b_h = b \qquad in \ \mathbf{L}^1([0,T] \times \Omega; \mathbb{R}) ,\qquad \lim_{h \to +\infty} B_h = B \qquad in \ \mathbf{L}^1\left([0,T]; \mathbf{L}^\infty(\Omega; \mathbb{R})\right) ,\\ \lim_{h \to +\infty} w_o^h = w_o \qquad in \ \mathbf{L}^1(\Omega; \mathbb{R}) ,\qquad \qquad \lim_{h \to +\infty} w_h = w \qquad in \ \mathbf{L}^\infty\left([0,T]; \mathbf{L}^1(\Omega; \mathbb{R})\right) .$$

Then:

- (1) For a.e.  $t \in [0, T], w(t) \in \mathbf{W}^{1,1}(\Omega; \mathbb{R}).$
- (2)  $\lim_{h\to+\infty} \nabla w_h = \nabla w$  weakly in  $\mathbf{L}^1([0,T] \times \Omega; \mathbb{R})$ .
- (3) w is a solution to (2.2) in the sense of Definition 2.5.

The following result differs from others found in the literature in its being set in  $\mathbf{L}^1$  and in its referring to Neumann boundary conditions.

**Theorem 2.7.** Let  $\Omega$  satisfy  $(\Omega)$ . Fix  $T, \mu > 0$ . Assume  $B \in \mathbf{L}^1([0,T]; \mathbf{L}^{\infty}(\Omega; \mathbb{R}))$  and  $b \in \mathbf{L}^1([0,T] \times \Omega; \mathbb{R})$ . Then, for all  $w_o \in \mathbf{L}^1(\Omega; \mathbb{R})$ , problem (2.2) admits a unique solution in the sense of Definition 2.5. Moreover:

(P1) It also holds that  $w \in \mathbf{C}^0([0,T]; \mathbf{L}^1(\Omega; \mathbb{R})), \nabla w \in \mathbf{L}^1([0,T] \times \Omega; \mathbb{R}^n)$  and

$$\|\nabla w\|_{\mathbf{L}^{1}([0,T]\times\Omega;\mathbb{R}^{n})} \leq \frac{1}{\mu} \|B\|_{\mathbf{L}^{1}([0,T];\mathbf{L}^{\infty}(\Omega;\mathbb{R}))} \|w\|_{\mathbf{L}^{\infty}([0,T];\mathbf{L}^{1}(\Omega;\mathbb{R}))} + \frac{1}{\mu} \|b\|_{\mathbf{L}^{1}([0,T]\times\Omega;\mathbb{R})} + \frac{1}{\mu} \|w_{o}\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})}.$$
(2.11)

**(P2)** The following implicit representation formula holds:

$$w(t,x) = \int_{\Omega} N(t,x,0,y) w_o(y) dy + \int_0^t \int_{\Omega} N(t,x,s,y) (B(s,y) w(s,y) + b(s,y)) dy ds$$
(2.12)

and the Neumann function N is defined in Proposition 3.3.

(P3) There exists a positive K depending on  $\mu$ ,  $\Omega$  — hence on n — such that the following a priori bound holds for all  $t \in [0,T]$ :

$$\left\|w(t)\right\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} \leq K\left(\left\|w_{o}\right\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} + \left\|b\right\|_{\mathbf{L}^{1}([0,t]\times\Omega;\mathbb{R})}\right) \exp\left(K\left\|B\right\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}))}\right).$$
 (2.13)

(P4) If  $w_o^1, w_o^2 \in \mathbf{L}^1(\Omega; \mathbb{R})$ , the corresponding solutions  $w_1$  and  $w_2$  satisfy for all  $t \in [0, T]$ ,

$$\left\| w_{1}(t) - w_{2}(t) \right\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} \leq K \left\| w_{o}^{1} - w_{o}^{2} \right\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} \exp\left( K \left\| B \right\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}))} \right).$$
(2.14)

(P5) If  $B_1, B_2 \in \mathbf{L}^1([0,T]; \mathbf{L}^{\infty}(\Omega; \mathbb{R}))$ , the corresponding solutions  $w_1$  and  $w_2$  satisfy for all  $t \in [0,T]$ ,

$$\|w_{1}(t) - w_{2}(t)\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})}$$

$$\leq K^{2} \exp\left(K(\|B_{1}\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}))} + \|B_{2}\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}))})\right)$$

$$\times \left(\|w_{o}\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} + \|b\|_{\mathbf{L}^{1}([0,t]\times\Omega;\mathbb{R})}\right) \|B_{1} - B_{2}\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}))}.$$

$$(2.15)$$

(P6) If  $b_1, b_2 \in \mathbf{L}^1([0,T] \times \Omega; \mathbb{R})$ , the corresponding solutions  $w_1, w_2$  satisfy for all  $t \in [0,T]$ ,

$$\left\|w_{1}(t) - w_{2}(t)\right\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} \leq K \exp\left(K \left\|B\right\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}))}\right) \left\|b_{1} - b_{2}\right\|_{\mathbf{L}^{1}([0,t]\times\Omega;\mathbb{R})}.$$
 (2.16)

(P7) Assume  $B \in \mathbf{L}^{\infty}([0,T] \times \Omega; \mathbb{R})$ . If  $b \ge 0$  and  $w_o \ge 0$ , then  $w \ge 0$ .

The proof is deferred to  $\S$  3.2.

## 3 Analytical Proofs

#### 3.1 Hyperbolic Problem

**Proof of Theorem 2.4.** The existence and uniqueness of u follow from [8, Proposition 3.9]. The *a priori*  $\mathbf{L}^1$  and  $\mathbf{L}^\infty$  bounds (**H1**) and (**H2**) are obtained in [4, Lemma 4.2]. Positivity in (**H3**) is proved as in [8, Lemma 3.12]. The Lipschitz continuous dependence on the initial datum (**H4**) follows from (**H1**) by linearity. The stability estimate (**H5**) is proved through the same computations as in [4, Lemma 4.3], taking advantage of linearity and of (**H1**). The continuity (**H6**) follows from [8, Lemma 3.13]. The stability in (**H7**) is proved in [8, Lemma 3.1].

#### 3.2 Parabolic Problem

**Proof of Lemma 2.6.** Using Definition 2.5, pass to the limit  $h \to +\infty$  in

$$\mu \int_0^T \int_\Omega \nabla w_h \cdot \nabla \varphi \, \mathrm{d}x \, \mathrm{d}t = \int_0^T \int_\Omega w_h \, \partial_t \varphi \, \mathrm{d}x \, \mathrm{d}t + \int_0^T \int_\Omega B_h \, w_h \, \varphi \, \mathrm{d}x \, \mathrm{d}t + \int_0^T \int_\Omega b_h \, \varphi \, \mathrm{d}x \, \mathrm{d}t + \int_\Omega w_o^h(x) \, \varphi(0, x) \, \mathrm{d}x \; .$$

By the Dominated Convergence Theorem, for all test functions  $\varphi \in \mathbf{C}_c^{\infty}([0, T[\times \overline{\Omega}; \mathbb{R})$ 

$$\int_{0}^{T} \int_{\Omega} w_{h} \partial_{t} \varphi \, \mathrm{d}x \, \mathrm{d}t = \int_{0}^{T} \int_{\Omega} w \, \partial_{t} \varphi \, \mathrm{d}x \, \mathrm{d}t \qquad [\text{Since } w_{h} \to w \text{ in } \mathbf{L}^{1}]$$

$$\int_{0}^{T} \int_{\Omega} B_{h} w_{h} \varphi \, \mathrm{d}x \, \mathrm{d}t = \int_{0}^{T} \int_{\Omega} B w \varphi \, \mathrm{d}x \, \mathrm{d}t \qquad [\text{Since } w_{h} \to w \text{ and } B_{h} \to B]$$

$$\int_{0}^{T} \int_{\Omega} b_{h} \varphi \, \mathrm{d}x \, \mathrm{d}t = \int_{0}^{T} \int_{\Omega} b \varphi \, \mathrm{d}x \, \mathrm{d}t \qquad [\text{Since } w_{h} \to w \text{ and } B_{h} \to B]$$

$$\int_{\Omega} w_{o}^{h}(x) \varphi(0, x) \, \mathrm{d}x = \int_{\Omega} w_{o}(x) \varphi(0, x) \, \mathrm{d}x \qquad [\text{Since } w_{o}^{h} \to w_{o} \text{ in } \mathbf{L}^{1}]$$

As a consequence, we obtain that

$$\lim_{h \to +\infty} \int_0^T \int_\Omega \nabla w_h \cdot \nabla \varphi \, \mathrm{d}x \, \mathrm{d}t = c_\varphi$$

for a real number  $c_{\varphi}$ , so that by the weak completeness of  $\mathbf{L}^1$ , see [26, Corollary 14], there exists a map  $z \in \mathbf{L}^1([0,T] \times \Omega; \mathbb{R}^n)$  such that  $\nabla w_h \xrightarrow[h \to +\infty]{} z$  in  $\mathbf{L}^1$ . Choose now  $\psi \in \mathbf{C}^{\infty}_c(]0, T[\times \Omega; \mathbb{R})$ :

$$\int_{0}^{T} \int_{\Omega} w \,\nabla\psi \,\mathrm{d}x \,\mathrm{d}t = \lim_{h \to +\infty} \int_{0}^{T} \int_{\Omega} w_{h} \,\nabla\psi \,\mathrm{d}x \,\mathrm{d}t \qquad \qquad [\text{Since } w_{h} \stackrel{\rightharpoonup}{\to} w \text{ in } \mathbf{L}^{1}]$$
$$= -\lim_{h \to +\infty} \int_{0}^{T} \int_{\Omega} \nabla w_{h} \,\psi \,\mathrm{d}x \,\mathrm{d}t \qquad \qquad [\text{Since } w_{h} \in \mathbf{W}^{1,1}]$$
$$= -\int_{0}^{T} \int_{\Omega} z \,\psi \,\mathrm{d}x \,\mathrm{d}t \qquad \qquad [\text{Since } \nabla w_{h} \stackrel{\rightharpoonup}{\to} z \text{ in } \mathbf{L}^{1}]$$

proving that  $w(t) \in \mathbf{W}^{1,1}(\Omega; \mathbb{R})$  for a.e.  $t \in [0,T]$  and  $\nabla w \in \mathbf{L}^1([0,T] \times \Omega; \mathbb{R}^n)$ . Hence, w satisfies the regularity requirements in Definition 2.5. Since it also satisfies (2.10), w is a weak solution to (2.2) in the sense of Definition 2.5.

We first consider problem (2.2) with B = 0, namely

$$\partial_t w - \mu \,\Delta w = b(t, x) \qquad (t, x) \in [0, T] \times \Omega$$
  

$$\nabla w(t, \xi) \cdot \nu(\xi) = 0 \qquad (t, \xi) \in ]0, T[\times \partial \Omega \qquad (3.1)$$
  

$$w(0, x) = w_o(x) \qquad x \in \Omega$$

under condition  $(\mathbf{\Omega})$  on  $\Omega$ .

**Lemma 3.1.** Fix  $w_o \in \mathbf{L}^1(\Omega; \mathbb{R})$ ,  $b \in \mathbf{L}^1([0, T] \times \Omega; \mathbb{R})$  and let w solve (3.1) in the sense of Definition 2.5. Then, for all  $\eta \in \mathbf{C}^{\infty}(\overline{\Omega}; \mathbb{R})$ 

$$\forall t \in ]0, T[: \qquad \lim_{h \to 0} \int_{\Omega} \left( w(t+h, x) - w(t, x) \right) \ \eta(x) \, \mathrm{d}x = 0 t = 0: \qquad \qquad \lim_{h \to 0+} \int_{\Omega} \left( w(h, x) - w_o(x) \right) \ \eta(x) \, \mathrm{d}x = 0.$$

$$(3.2)$$

**Proof of Lemma 3.1.** Fix  $t \in [0, T]$  and introduce the sequence of functions

$$\chi_k \in \mathbf{C}^\infty_c(\mathbb{R}; [0,1])\,, \quad \operatorname{spt} \chi_k \subseteq [-1,t] \quad \text{ and } \quad \chi_k \xrightarrow[k \to +\infty]{} \mathbf{1}_{[0,t]} \text{ pointwise a.e. on } [0,T]\,.$$

Then use (2.10) with  $B \equiv 0$ , first  $\varphi(s, x) = \chi_k(s+h) \eta(x)$  and then  $\varphi(s, x) = \chi_k(s) \eta(x)$ , for a suitable  $\eta \in \mathbb{C}^{\infty}(\overline{\Omega}; \mathbb{R})$ , with  $t \in [0, T[$  and h sufficiently small. Taking the difference of the resulting expressions, we have:

$$\int_0^T \int_\Omega w(s,x) \left(\partial_t \chi_k(s+h) - \partial_t \chi_k(s)\right) \eta(x) \, \mathrm{d}x \, \mathrm{d}s$$
  
=  $\mu \int_0^T \int_\Omega \left(\chi_k(s+h) - \chi_k(s)\right) \nabla w(s,x) \cdot \nabla \eta(x) \, \mathrm{d}x \, \mathrm{d}s$   
 $- \int_0^T \int_\Omega b(s,x) \left(\chi_k(s+h) - \chi_k(s)\right) \eta(x) \, \mathrm{d}x \, \mathrm{d}s + \int_\Omega w_o(x) \left(\chi_k(0) - \chi_k(h)\right) \eta(x) \, \mathrm{d}x \, \mathrm{d}s$ 

If t > 0, the latter term above vanishes and in the limit  $k \to +\infty$  the first equality in (3.2) follows.

When t = 0, the above terms reduce to

$$\int_0^T \int_\Omega w(s,x) \,\partial_t \chi_k(s+h) \,\eta(x) \,\mathrm{d}x \,\mathrm{d}s$$
  
=  $\mu \int_0^T \int_\Omega \chi_k(s+h) \,\nabla w(s,x) \cdot \nabla \eta(x) \,\mathrm{d}x \,\mathrm{d}s$   
 $- \int_0^T \int_\Omega b(s,x) \,\chi_k(s+h) \,\eta(x) \,\mathrm{d}x \,\mathrm{d}s - \int_\Omega w_o(x) \,\chi_k(h) \,\eta(x) \,\mathrm{d}x$ 

and as  $k \to +\infty$  the second equality in (3.2) follows.

When the initial datum  $w_o$  is in  $\mathbf{L}^2(\Omega; \mathbb{R})$  and the source b is in  $\mathbf{L}^2([0, T] \times \Omega; \mathbb{R})$ , strong  $\mathbf{L}^2$  continuity in time is available.

**Lemma 3.2.** If  $b \in \mathbf{L}^2([0,T]; \mathbf{L}^2(\Omega; \mathbb{R}))$  and  $w_o \in \mathbf{L}^2(\Omega; \mathbb{R})$ , then problem (3.1) admits a unique solution  $w \in \mathbf{L}^2([0,T]; \mathbf{W}^{1,2}(\Omega; \mathbb{R}))$  in the sense of Definition 2.5 and  $\partial_t w \in \mathbf{L}^2([0,T]; \mathbf{W}^{1,2}(\Omega; \mathbb{R})^*)$ , so that  $w \in \mathbf{C}^0([0,T]; \mathbf{L}^2(\Omega; \mathbb{R}))$ .

**Proof of Lemma 3.2.** Use [22, Problem (10.35)] and apply [22, point b) in Theorem 7.104]. Note that the definition of weak solution in [22, p. 592] implies Definition 2.5 due to the density of  $\mathbf{C}_c^{\infty}(\Omega; \mathbb{R})$  in  $\mathbf{W}^{1,2}(\Omega; \mathbb{R})$  and by the Dominated Convergence Theorem.

We introduce for later use the space  $\mathcal{V}$  as the closure of  $\mathbf{W}^{1,2}(\mathbb{R} \times \Omega; \mathbb{R})$  with respect to the norm

$$\|w\|_{\mathcal{V}} = \|\nabla w\|_{\mathbf{L}^{2}(\mathbb{R}\times\Omega;\mathbb{R})} + \operatorname{ess\,sup}_{t\in\mathbb{R}} \|w(t)\|_{\mathbf{L}^{2}(\Omega;\mathbb{R})}.$$

**Proposition 3.3.** Let  $\Omega$  satisfy  $(\Omega)$  and  $\mu > 0$ . Then, there exists a function

$$N \in \mathbf{C}^0\left(\left\{(t, x, s, y) \in (\mathbb{R} \times \Omega)^2 \colon (t, x) \neq (s, y)\right\}; \mathbb{R}\right)$$

such that

**(N1)** For all  $(s, y) \in \mathbb{R} \times \Omega$ ,

$$(t,x) \mapsto N(t,x,s,y) \in \mathbf{L}^{1}_{\mathbf{loc}}(\mathbb{R} \times \Omega; \mathbb{R}) \text{ and } (t,x) \mapsto \nabla_{x} N(t,x,s,y) \in \mathbf{L}^{1}_{\mathbf{loc}}(\mathbb{R} \times \Omega; \mathbb{R}^{n}).$$

**(N2)** There exist positive C,  $\kappa$  and K such that for all  $(t, x, s, y) \in (\mathbb{R}_+ \times \Omega)^2$  with t > s

$$|N(t,x,s,y)| \leq C \left(1 + \frac{1}{(t-s)^{n/2}}\right) \exp\left(-\frac{\kappa ||x-y||^2}{t-s}\right);$$
 (3.3)

$$\int_{\Omega} |N(t,x,s,y)| \, \mathrm{d}x \leq K.$$
(3.4)

**(N3)** Fix T > 0. For every  $w_o \in \mathbf{L}^2(\Omega; \mathbb{R})$  and all  $b \in \mathbf{C}^{\infty}([0,T] \times \overline{\Omega}; \mathbb{R})$ , the map

$$w(t,x) = \int_{\Omega} N(t,x,0,y) \, w_o(y) \, \mathrm{d}y + \int_0^t \int_{\Omega} N(t,x,s,y) \, b(s,y) \, \mathrm{d}y \, \mathrm{d}s \tag{3.5}$$

is the unique function in  $\mathcal{V}$  satisfying (2.10), with  $B \equiv 0$ , for all test function  $\varphi \in \mathbf{C}_{c}^{\infty}([0,T[\times\overline{\Omega};\mathbb{R}).$ 

**Proof of Proposition 3.3.** Here we apply [3, Theorem 3.9], thus we need to verify conditions (A1)–(A2) in [3]. (A1) holds by [3, (1) in Examples 4.1.1] thanks to [2, Theorem 9.7] which can be applied thanks to  $(\Omega)$ ; (A2) holds by [3, (1) in Examples 4.1.2], that applies in the present scalar case. We can apply [3, Theorem 3.21] since also (A3) in [3] holds by [3, (3) in Examples 4.1.3], since the coefficients in (3.1) are constant and  $(\Omega)$  holds.

The regularity of N and the proof of (N1) directly follow from [3, Theorem 3.9]. To prove (3.3), start from the last line in the proof of [3, Theorem 3.21]:

$$\left|N(t,x,s,y)\right| \le C \max\left\{1, \frac{t-s}{\operatorname{diam}(\Omega)}\right\}^{n/2} (t-s)^{-n/2} \exp\left(-\frac{\kappa \|x-y\|^2}{t-s}\right)$$

which implies (3.3), up to relabeling C. Now, (3.4) is a direct consequence with  $K = C \left( \max(\Omega) + (\pi/\kappa)^{n/2} \right)$ .

Consider (N3). The proof that the expression (3.5) solves (3.1) in the sense of Definition 2.5 follows from [3, Theorem 3.9], as well as uniqueness.

**Remark 3.4.** A close look at the proof of [3, Theorem 3.9] shows also that  $N \ge 0$ . This positivity is not explicitly considered in [3] since the object of that work is a system of parabolic equations and thus N results to be matrix valued. This property, though basic, is not necessary in the sequel, hence we omit its proof whose rigorous exposition might significantly lengthen this work. However,  $N \ge 0$  ensures that

$$w_o \in \mathbf{L}^2(\Omega; \mathbb{R}_+)$$
 and  $b \in \mathbf{C}^{\infty}([0, T] \times \overline{\Omega}; \mathbb{R}_+) \implies w \ge 0$ .

**Proposition 3.5.** Let  $\Omega$  satisfy  $(\Omega)$ . Fix  $T, \mu > 0$ . For every  $w_o \in \mathbf{L}^1(\Omega; \mathbb{R})$  and any  $b \in \mathbf{L}^1([0,T] \times \Omega; \mathbb{R})$ , problem (3.1) admits a unique weak solution w in the sense of Definition 2.5. Moreover,

- (1) w admits the representation (3.5) with N as defined in Proposition 3.3.
- (2)  $w \in \mathbf{C}^0([0,T]; \mathbf{L}^1(\Omega; \mathbb{R}))$  and  $\nabla w \in \mathbf{L}^1([0,T] \times \Omega; \mathbb{R}^n)$ .

Proof of Proposition 3.5. We split the proof in a few steps.

**Step 1: Uniqueness.** Assume  $w_1, w_2$  solve (3.1) in the sense of Definition 2.5. Then, their difference w satisfies, for all test function  $\varphi \in \mathbf{C}_c^{\infty}([0, T[\times \overline{\Omega}; \mathbb{R}),$ 

$$\int_0^T \int_\Omega w \,\partial_t \varphi \,\mathrm{d}x \,\mathrm{d}t - \mu \int_0^T \int_\Omega \nabla w \cdot \nabla \varphi \,\mathrm{d}x \,\mathrm{d}t = 0.$$
(3.6)

First apply (3.6) with a test function depending only on time to obtain, thanks to Lemma 3.1, that for all  $t \in [0, T]$ ,

$$\int_{\Omega} w(t,x) \,\mathrm{d}x = 0\,. \tag{3.7}$$

Then, fix arbitrary  $a, b \in [0, T[$  with a < b. Introduce a sequence of test functions

$$\chi_h \in \mathbf{C}_c^{\infty}(]0, T[; [0, 1]), \quad \operatorname{spt} \chi_h \subseteq [a, b] \quad \text{and} \quad \chi_h \to \mathbf{1}_{[a, b]} \text{ pointwise a.e. on } [0, T].$$

Let  $\psi \in \mathbf{C}^{\infty}(\overline{\Omega}; \mathbb{R})$  and apply (3.6) with  $\varphi = \chi_h \psi$ , obtaining

$$0 = \int_{a}^{b} \int_{\Omega} w \,\partial_{t} \chi_{h} \,\psi \,\mathrm{d}x \,\mathrm{d}t - \mu \int_{a}^{b} \int_{\Omega} \chi_{h} \,\nabla w \cdot \nabla \psi \,\mathrm{d}x \,\mathrm{d}t$$
$$= \int_{\Omega} w(b,x) \,\mathrm{d}x - \int_{\Omega} w(a,x) \,\mathrm{d}x - \mu \int_{a}^{b} \int_{\Omega} \nabla w \cdot \nabla \psi \,\mathrm{d}x \,\mathrm{d}t$$
$$= -\mu \int_{a}^{b} \int_{\Omega} \nabla w \cdot \nabla \psi \,\mathrm{d}x \,\mathrm{d}t$$

where we used (3.7). Hence, there exists a  $c \in \mathbb{R}^n$  such that for a.e.  $x \in \Omega$ ,

$$\int_{a}^{b} \nabla w(t,x) \,\mathrm{d}t = c \tag{3.8}$$

and, by the arbitrariness of a and b, it must be c = 0. From (3.8) we thus obtain that for a.e.  $x \in \Omega$  and for all  $g \in \mathbf{L}^{\infty}([0,T]; \mathbb{R}^n)$ 

$$\int_0^T \nabla w(t, x) \cdot g(t) \, \mathrm{d}t = 0$$

so that  $\nabla w(t, x) = 0$  for all  $t \in [0, T]$  and a.e.  $x \in \Omega$ . Together with (3.7) and the connectedness of  $\Omega$ , this implies that w = 0, proving uniqueness.

**Step 2: Approximation.** By [13, (1.8) in § 1.14], there exists a sequence  $b_h \in \mathbf{C}_c^{\infty}(\mathbb{R}^{1+n};\mathbb{R})$ such that  $b_h \to b$  in  $\mathbf{L}^1([0,T] \times \Omega;\mathbb{R})$  as  $h \to +\infty$ . Similarly, there exists a sequence  $w_o^h \in \mathbf{C}_c^{\infty}(\mathbb{R}^n,\mathbb{R})$  such that  $w_o^h \to w_o$  in  $\mathbf{L}^1(\Omega;\mathbb{R})$  as  $h \to +\infty$ . Note that  $w_o^h \in \mathbf{L}^2(\Omega;\mathbb{R})$  for all h.

For all  $h \in \mathbb{N}$ , by **(N3)** in Proposition 3.3 there is a unique solution  $w_h \in \mathcal{V}$  to (3.1) with source  $b_h$  and initial datum  $w_o^h$ , which is given by the representation

$$w_h(t,x) = \int_{\Omega} N(t,x,0,y) \ w_o^h(y) \, \mathrm{d}y + \int_0^t \int_{\Omega} N(t,x,s,y) \ b_h(s,y) \, \mathrm{d}y \, \mathrm{d}s \ . \tag{3.9}$$

Then, for  $h, k \in \mathbb{N}$  with k > h, thanks to (3.4) we have

$$\left\|w_k(t) - w_h(t)\right\|_{\mathbf{L}^1(\Omega;\mathbb{R})} \leq \int_{\Omega} \int_{\Omega} \left|N(t,x,0,y)\right| \left|w_o^k(y) - w_o^h(y)\right| \mathrm{d}y \,\mathrm{d}x$$

$$\begin{aligned} &+ \int_{\Omega} \int_{0}^{t} \int_{\Omega} \left| N(t,x,s,y) \right| \left| b_{k}(s,y) - b_{h}(s,y) \right| \mathrm{d}y \, \mathrm{d}s \, \mathrm{d}x \\ &= \int_{\Omega} \left( \int_{\Omega} \left| N(t,x,0,y) \right| \, \mathrm{d}x \right) \, \left| w_{o}^{k}(y) - w_{o}^{h}(y) \right| \, \mathrm{d}y \\ &+ \int_{\Omega} \int_{0}^{t} \left( \int_{\Omega} \left| N(t,x,s,y) \right| \, \mathrm{d}x \right) \left| b_{k}(s,y) - b_{h}(s,y) \right| \, \mathrm{d}s \, \mathrm{d}y \\ &\leq K \left( \left\| w_{o}^{k} - w_{o}^{h} \right\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} + \left\| b_{k} - b_{h} \right\|_{\mathbf{L}^{1}([0,T] \times \Omega;\mathbb{R})} \right) \end{aligned}$$

proving that there exists a function  $w \in \mathbf{L}^{\infty}([0,T];\mathbf{L}^{1}(\Omega;\mathbb{R}))$  such that

$$\lim_{h \to +\infty} \|w_h - w\|_{\mathbf{L}^{\infty}\left([0,T];\mathbf{L}^1(\Omega;\mathbb{R})\right)} = 0.$$
(3.10)

Step 3: Existence in  $\mathbf{L}^{\infty}([0,T]; \mathbf{L}^{1}(\Omega; \mathbb{R}))$ . To prove that w solves (3.1) in the sense of Definition 2.5 it is now sufficient to apply Lemma 2.6 with  $B_h \equiv 0$ .

Hence, w satisfies the regularity requirements in Definition 2.5. Since it also satisfies (2.10), w is a weak solution to (3.1) in the sense of Definition 2.5, with  $B \equiv 0$ .

Step 4:  $\mathbf{L}^{1}(\Omega; \mathbb{R})$  Continuity in Time. Note that the sequence  $w_{h}$  defined above also satisfies  $w_{h} \in \mathbf{L}^{2}([0,T]; \mathbf{W}^{1,2}(\Omega; \mathbb{R}))$  and  $\partial_{t}w_{h} \in \mathbf{L}^{2}([0,T]; \mathbf{W}^{1,2}(\Omega; \mathbb{R})^{*})$ , so that  $w_{h} \in \mathbf{C}^{0}([0,T]; \mathbf{L}^{2}(\Omega; \mathbb{R}))$  by Lemma 3.2. Recall that  $\mathbf{C}^{0}([0,T]; \mathbf{L}^{2}(\Omega; \mathbb{R})) \subseteq \mathbf{C}^{0}([0,T]; \mathbf{L}^{1}(\Omega; \mathbb{R}))$ , hence the convergence (3.10) then ensures that  $w \in \mathbf{C}^{0}([0,T]; \mathbf{L}^{1}(\Omega; \mathbb{R}))$ .

Step 5: Representation Formula. It is now sufficient to pass to the limit  $h \to +\infty$  in (3.9) to prove that w admits the representation (3.5).

The proof is completed.

**Corollary 3.6.** Let  $(\Omega)$  hold. Fix  $T, \mu > 0$ . Let  $w_o^1, w_o^2 \in \mathbf{L}^1(\Omega; \mathbb{R})$  and  $b_1, b_2 \in \mathbf{L}^1([0,T] \times \Omega; \mathbb{R})$ . Call  $w_1, w_2$  the corresponding solutions to (3.1). Then, for all  $t \in [0,T]$ ,

$$\left\|w_{1}(t) - w_{2}(t)\right\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} \leq K\left(\left\|w_{o}^{1} - w_{o}^{2}\right\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} + \|b_{1} - b_{2}\|_{\mathbf{L}^{1}([0,t]\times\Omega;\mathbb{R})}\right)$$
(3.11)

where K is as in (N2) of Proposition 3.3.

Simply apply (1) of Proposition 3.5 to the difference  $w_2 - w_1$  and exploit the linearity of (3.1).

**Proof of Theorem 2.7.** We split the proof in a few steps.

Step 1: Problem (2.2) admits a unique solution on [0,T] satisfying (P1) and (P2). Consider the operators

$$\Lambda : \begin{bmatrix} \mathbf{C}^{0} \left( [0,T]; \mathbf{L}^{1}(\Omega; \mathbb{R}) \right) & \longrightarrow & \mathbf{L}^{1}([0,T] \times \Omega; \mathbb{R}) \\ w & \longmapsto & \Lambda w \quad \text{where } (\Lambda w)(t,x) = B(t,x) w(t,x) + b(t,x) \end{bmatrix}$$
(3.12)

and, with reference to Definition 2.5,

$$\Phi: \begin{bmatrix} \mathbf{L}^{1}([0,T] \times \Omega; \mathbb{R}) & \longrightarrow & \mathbf{C}^{0}\left([0,T]; \mathbf{L}^{1}(\Omega; \mathbb{R})\right) \\ \beta & \longmapsto & w \quad \text{where} \begin{cases} \partial_{t}w - \mu \,\Delta w = \beta(t,x) & (t,x) \in [0,T] \times \Omega \\ \nabla w(t,\xi) \cdot \nu(\xi) = 0 & (t,\xi) \in ]0, T[ \times \partial \Omega \\ w(0,x) = w_{o}(x) & x \in \Omega \,. \end{cases}$$

Let us precise that  $\Lambda$  is well defined, meaning that if  $w \in \mathbf{C}^0([0,T]; \mathbf{L}^1(\Omega; \mathbb{R}))$  then, by the assumptions on B and b,  $\Lambda(w) \in \mathbf{L}^1([0,T] \times \Omega; \mathbb{R})$ . Similarly,  $\Phi$  is well defined by Proposition 3.5.

By the assumption on B, there exist times  $t_i$  for i = 0, ..., m such that  $0 = t_0 < t_1 < \cdots < t_i < t_{i+1} < \cdots < t_m = T$  and

$$\int_{t_i}^{t_{i+1}} \left\| B(s) \right\|_{\mathbf{L}^{\infty}(\Omega;\mathbb{R})} \mathrm{d}s < \frac{1}{2K}$$
(3.13)

with K as in (3.4) in Proposition 3.3. Clearly, thanks to the continuity proved in Proposition 3.5, w in  $\mathbf{L}^{\infty}([0,T]; \mathbf{L}^{1}(\Omega; \mathbb{R}))$  with  $w(t) \in \mathbf{W}^{1,1}(\Omega; \mathbb{R})$  for a.e.  $t \in [0,T]$  is a weak solution to (2.2) if and only if it is a fixed point of  $\Phi \circ \Lambda$  in  $\mathbf{C}^{0}([0,T]; \mathbf{L}^{1}(\Omega; \mathbb{R}))$ . To construct such fixed point we apply Banach Fixed Point Theorem iteratively in each of the spaces  $\mathbf{C}^{0}([t_{i}, t_{i+1}]; \mathbf{L}^{1}(\Omega; \mathbb{R}))$  for  $i = 0, \ldots, m-1$ . Notice that for all  $w_{1}, w_{2} \in \mathbf{C}^{0}([t_{i}, t_{i+1}]; \mathbf{L}^{1}(\Omega; \mathbb{R}))$  with  $w_{1}(t_{i}) = w_{2}(t_{i})$ , for all  $t \in$ 

Notice that for all  $w_1, w_2 \in \mathbf{C}^0([t_i, t_{i+1}]; \mathbf{L}^1(\Omega; \mathbb{R}))$  with  $w_1(t_i) = w_2(t_i)$ , for all  $t \in [t_i, t_{i+1}]$ ,

$$\begin{split} \|\Phi \circ \Lambda(w_{1})(t) - \Phi \circ \Lambda(w_{2})(t)\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} \\ &\leq K \int_{t_{i}}^{t} \|\Lambda(w_{1})(s) - \Lambda(w_{2})(s)\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} \, \mathrm{d}s \qquad \text{[By Corollary 3.6]} \\ &= K \int_{t_{i}}^{t} \int_{\Omega} |B(s,y)| \, |w_{1}(s,y) - w_{2}(s,y)| \, \mathrm{d}y \, \mathrm{d}s \qquad \text{[By (3.12)]} \\ &\leq K \int_{t_{i}}^{t} \|B(s)\|_{\mathbf{L}^{\infty}(\Omega;\mathbb{R})} \int_{\Omega} |w_{1}(s,y) - w_{2}(s,y)| \, \mathrm{d}y \, \mathrm{d}s \\ &\leq K \int_{t_{i}}^{t} \|B(s)\|_{\mathbf{L}^{\infty}(\Omega;\mathbb{R})} \, \mathrm{d}s \, \|w_{1} - w_{2}\|_{\mathbf{C}^{0}([t_{i},t_{i+1}];\mathbf{L}^{1}(\Omega;\mathbb{R}))} \\ &\leq \frac{1}{2} \, \|w_{1} - w_{2}\|_{\mathbf{C}^{0}([t_{i},t_{i+1}];\mathbf{L}^{1}(\Omega;\mathbb{R}))} \,. \end{aligned}$$

An iterated application of Banach Fixed Point Theorem ensures the existence of  $w_* \in \mathbf{C}^0([0,T]; \mathbf{L}^1(\Omega; \mathbb{R}))$  such that  $w_* = \Phi \circ \Lambda(w_*)$ .

Define  $\tilde{b} = B w_* + b$ , so that  $\tilde{b} \in \mathbf{L}^1([0,T] \times \Omega; \mathbb{R})$ . By construction,  $w_*$  solves

$$\partial_t w - \mu \ \Delta w = \tilde{b}(t, x) \qquad (t, x) \in [0, T] \times \Omega$$
$$\nabla w(t, \xi) \cdot \nu(\xi) = 0 \qquad (t, \xi) \in ]0, T[\times \partial \Omega$$
$$w(0, x) = w_o(x) \qquad x \in \Omega$$

in the sense of Definition 2.5. Hence,  $w_*(t) \in \mathbf{W}^{1,1}(\Omega; \mathbb{R})$  for a.e.  $t \in [0,T]$  and  $\nabla w \in \mathbf{L}^1([0,T] \times \Omega; \mathbb{R}^n)$  by (2) in Proposition 3.5. This shows that  $w_*$  is a weak solution to (2.2) in the sense of Definition 2.5 on [0,T] and (P1) holds.

To prove the bound (2.11), by (2.10) we have

$$\begin{aligned} & \left| \int_{0}^{T} \int_{\Omega} [w - \mu \nabla w] \begin{bmatrix} \partial_{t} \varphi \\ \nabla \varphi \end{bmatrix} \mathrm{d}x \, \mathrm{d}t \right| \\ & \leq \left| \int_{0}^{T} \int_{\Omega} (B \, w + b) \varphi \, \mathrm{d}x \, \mathrm{d}t \right| + \left| \int_{\Omega} w_{o} \, \varphi(0, \cdot) \, \mathrm{d}x \right| \\ & \leq \left( \|B\|_{\mathbf{L}^{1}([0,T];\mathbf{L}^{\infty}(\Omega;\mathbb{R}))} \|w\|_{\mathbf{L}^{\infty}([0,T];\mathbf{L}^{1}(\Omega;\mathbb{R}))} + \|b\|_{\mathbf{L}^{1}([0,T]\times\Omega;\mathbb{R})} + \|w_{o}\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} \right) \|\varphi\|_{\mathbf{L}^{\infty}([0,T]\times\Omega;\mathbb{R})} \end{aligned}$$

which together with

$$\left\|\nabla w\right\|_{\mathbf{L}^{1}\left([0,T]\times\Omega;\mathbb{R}^{n}\right)} \leq \frac{1}{\mu} \left\|\left[w - \mu\,\nabla w\right]\right\|_{\mathbf{L}^{1}\left([0,T]\times\Omega;\mathbb{R}^{n+1}\right)}$$

completes the proof of (2.11).

By construction, (P2) holds. Step 1 is proved.

**Step 2: (P3) holds.** Using (2.12), for all  $t \in [0, T]$ , by (3.4)

$$\begin{split} \left\|w(t)\right\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} &\leq K\left(\left\|w_{o}\right\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} + \int_{0}^{t} \int_{\Omega} \left(\left|B(s,y)\right| \left|w(s,y)\right| + \left|b(s,y)\right|\right) \, \mathrm{d}y \, \mathrm{d}s\right) \\ &\leq K\left(\left\|w_{o}\right\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} + \left\|b\right\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{1}(\Omega;\mathbb{R}))} + \int_{0}^{t} \left\|B(s)\right\|_{\mathbf{L}^{\infty}(\Omega;\mathbb{R})} \left\|w(s)\right\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} \, \mathrm{d}s\right). \end{split}$$

An application of Gronwall Lemma leads to (2.13).

Step 3: (P4) holds. This is an immediate consequence of (P3) by linearity.

Step 4: (P5) and (P6) hold. Using (2.12), compute:

$$\begin{split} &\|w_{1}(t) - w_{2}(t)\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} \\ &\leq K \int_{0}^{t} \int_{\Omega} \left|B_{1}(s,y) w_{1}(s,y) - B_{2}(s,y) w_{2}(s,y)\right| \mathrm{d}y \, \mathrm{d}s + K \int_{0}^{t} \int_{\Omega} \left|b_{1}(s,y) - b_{2}(s,y)\right| \mathrm{d}y \, \mathrm{d}s \\ &\leq K \int_{0}^{t} \int_{\Omega} \left|B_{1}(s,y) w_{1}(s,y) - B_{1}(s,y) w_{2}(s,y)\right| \mathrm{d}y \, \mathrm{d}s \\ &+ K \int_{0}^{t} \int_{\Omega} \left|B_{1}(s,y) w_{2}(s,y) - B_{2}(s,y) w_{2}(s,y)\right| \mathrm{d}y \, \mathrm{d}s + K \int_{0}^{t} \int_{\Omega} \left|b_{1}(s,y) - b_{2}(s,y)\right| \mathrm{d}y \, \mathrm{d}s \\ &\leq K \int_{0}^{t} \left\|B_{1}(s)\right\|_{\mathbf{L}^{\infty}(\Omega;\mathbb{R})} \left\|w_{1}(s) - w_{2}(s)\right\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} \, \mathrm{d}s \\ &+ K \left\|B_{1} - B_{2}\right\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}))} \left\|w_{2}\right\|_{\mathbf{C}^{0}([0,t];\mathbf{L}^{1}(\Omega;\mathbb{R}))} + K \left\|b_{1} - b_{2}\right\|_{\mathbf{L}^{1}([0,t]\times\Omega;\mathbb{R})}. \end{split}$$

By Gronwall Lemma and using (P3),

$$\begin{aligned} & \left\| w_{1}(t) - w_{2}(t) \right\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} \\ \leq & K \left( \|B_{1} - B_{2}\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}))} \|w_{2}\|_{\mathbf{C}^{0}([0,t];\mathbf{L}^{1}(\Omega;\mathbb{R}))} + \|b_{1} - b_{2}\|_{\mathbf{L}^{1}([0,t]\times\Omega;\mathbb{R})} \right) \\ & \times \exp\left( K \|B_{1}\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}))} \right) \\ \leq & K \|b_{1} - b_{2}\|_{\mathbf{L}^{1}([0,t]\times\Omega;\mathbb{R})} \exp\left( K \|B_{1}\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}))} \right) \\ & + K^{2} \|B_{1} - B_{2}\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}))} \left( \|w_{o}\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} + \|b_{2}\|_{\mathbf{L}^{1}([0,t]\times\Omega;\mathbb{R})} \right) \\ & \times \exp\left( K \left( \|B_{1}\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}))} + \|B_{2}\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}))} \right) \right), \end{aligned}$$

which implies (2.15) and (2.16).

Step 5: (P7) holds with  $B \in \mathbf{L}^{\infty}([0,T] \times \Omega; \mathbb{R})$ . By [13, (1.8) in § 1.14], there exists a sequence  $b_h \in \mathbf{C}^{\infty}_c(\mathbb{R}^{1+n}; \mathbb{R}_+)$  such that  $b_h \to b$  in  $\mathbf{L}^1([0,T] \times \Omega; \mathbb{R})$  as  $h \to +\infty$ . Similarly, there exists a sequence  $w_o^h \in \mathbf{C}^{\infty}_c(\mathbb{R}^n, \mathbb{R}_+)$  such that  $w_o^h \to w_o$  in  $\mathbf{L}^1(\Omega; \mathbb{R})$  as  $h \to +\infty$ . Note that  $w_o^h \in \mathbf{L}^2(\Omega; \mathbb{R})$  for all h. Call  $w_h$  the solution to

$$\partial_t w_h - \mu \,\Delta w_h = B(t, x)w_h + b_h(t, x) \qquad (t, x) \in [0, T] \times \Omega$$
$$\nabla w_h(t, \xi) \cdot \nu(\xi) = 0 \qquad (t, \xi) \in ]0, T[\times \partial \Omega$$
$$w(0, x) = w_o^h(x) \qquad x \in \Omega.$$

in the sense of the definition given in [22, p. 592]. By [22, Remark 10.19],  $w_h \ge 0$ .

Using (P4) and (P6), we have

$$\left\| w_{h}(t) - w_{k}(t) \right\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} \leq K e^{K \|B\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}))}} \left( \left\| w_{o}^{h} - w_{o}^{k} \right\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} + \|b_{h} - b_{k}\|_{\mathbf{L}^{1}([0,t]\times\Omega;\mathbb{R})} \right)$$

hence there exists a  $w \in \mathbf{L}^{\infty}([0,T];\mathbf{L}^{1}(\Omega;\mathbb{R}))$  such that  $w_{h} \to w$  in  $\mathbf{L}^{\infty}([0,T];\mathbf{L}^{1}(\Omega;\mathbb{R}))$ .

Apply now Lemma 2.6 to prove that w is a solution to (2.2) in the sense of Definition 2.5. By construction,  $w \ge 0$ .

The proof is completed.

### 3.3 Mixed Problem

**Proof of Theorem 2.2.** Define  $u_1$  and  $w_1$  as solutions to

$$\begin{cases} \partial_t u_1 = a(t,x) & (t,x) \in [0,T] \times \Omega \\ u_1(t,\xi) = 0 & (t,\xi) \in ]0, T[\times \partial \Omega \\ u_1(0,x) = u_o(x) & x \in \Omega \end{cases} \qquad \begin{cases} \partial_t w_1 - \mu \,\Delta w_1 = b(t,x) & (t,x) \in [0,T] \times \Omega \\ \nabla w_1(t,\xi) \cdot \nu(\xi) = 0 & (t,\xi) \in ]0, T[\times \partial \Omega \\ w_1(0,x) = w_o(x) & x \in \Omega \end{cases}$$

in the sense of Definition 2.3 and of Definition 2.5. Clearly,  $u_1$  and  $w_1$  are well defined by Theorem 2.3 and Theorem 2.7. Define recursively, for  $i \in \mathbb{N} \setminus \{0\}$ ,  $u_{i+1}$  as solution to

$$\partial_t u_{i+1} + \nabla \cdot \left( u_{i+1} c_i(t, x) \right) = A_i(t, x) u_{i+1} + a(t, x) \qquad (t, x) \in [0, T] \times \Omega$$
$$u_{i+1}(t, \xi) = 0 \qquad (t, \xi) \in ]0, T[\times \partial \Omega \qquad (3.14)$$
$$u_{i+1}(0, x) = u_o(x) \qquad x \in \Omega$$

in the sense of Definition 2.3 and  $w_{i+1}$  as solution to

$$\partial_t w_{i+1} - \mu \Delta w_{i+1} = B_i(t, x) w_{i+1} + b(t, x) \qquad (t, x) \in [0, T] \times \Omega$$
  

$$\nabla w_{i+1}(t, \xi) \cdot \nu(\xi) = 0 \qquad (t, \xi) \in ]0, T[\times \partial \Omega \qquad (3.15)$$
  

$$w_{i+1}(0, x) = w_o(x) \qquad x \in \Omega$$

in the sense of Definition 2.5, where for  $i \in \mathbb{N} \setminus \{0\}$ ,

$$c_i(t,x) = v(t,w_i(t))(x)$$
  

$$A_i(t,x) = \alpha(t,x,w_i(t))$$
  

$$B_i(t,x) = \beta(t,x,u_i(t),w_i(t)).$$
(3.16)

We aim to prove that  $(u_i, w_i)$  are well defined for all  $i \in \mathbb{N} \setminus \{0, 1\}$  and constitute a Cauchy sequence with respect to the  $\mathbf{C}^0([0, T]; \mathbf{L}^1(\Omega; \mathbb{R}^2))$  distance as soon as T is sufficiently small.

Step 1:  $c_i$  satisfies (c) in Theorem 2.4. We first check the continuity. By (3.16) and (v) we have

$$\begin{aligned} & \left\|c_{i}(t,x) - c_{i}(t_{o},x_{o})\right\| \\ \leq & \left\|c_{i}(t,x) - c_{i}(t_{o},x)\right\| + \left\|c_{i}(t_{o},x) - c_{i}(t_{o},x_{o})\right\| \\ = & \left\|v\left(t,w_{i}(t)\right)(x) - v\left(t_{o},w_{i}(t_{o})\right)(x)\right\| + \left\|v\left(t_{o},w_{i}(t_{o})\right)(x) - v\left(t_{o},w_{i}(t_{o})\right)(x_{o})\right\| \\ \leq & K_{v}\left(\left|t - t_{o}\right| + \left\|w_{i}(t) - w_{i}(t_{o})\right\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})}\right) + \left\|v\left(t_{o},w_{i}(t_{o})\right)(x) - v\left(t_{o},w_{i}(t_{o})\right)(x_{o})\right\| \end{aligned}$$

where the terms  $\|w_i(t) - w_i(t_o)\|_{\mathbf{L}^1(\Omega;\mathbb{R})}$  and  $\|v(t_o, w_i(t_o))(x_o) - v(t_o, w_i(t_o))(x_o)\|$  vanish as  $t \to t_o$  and  $x \to x_o$  by the continuity of  $w_i$  and that of  $v(t_o, w_i(t_o))$ , ensured by  $(\boldsymbol{v})$ .

To prove the  $\mathbf{L}^{\infty}$  boundedness, observe that by (v)

$$\left\|c_{i}(t,x)\right\| \leq \left\|v\left(t,w_{i}(t)\right)\right\|_{\mathbf{L}^{\infty}(\Omega;\mathbb{R}^{n})} \leq K_{v}\left\|w_{i}(t)\right\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} \leq K_{v}\left\|w_{i}\right\|_{\mathbf{C}^{0}([0,T];\mathbf{L}^{1}(\Omega;\mathbb{R}))}.$$
 (3.17)

The regularity  $c_i(t) \in \mathbf{C}^1(\Omega; \mathbb{R}^n)$  directly follows from  $(\boldsymbol{v})$ . Concerning the boundedness of  $D_x c_i$ , observe that by  $(\boldsymbol{v})$ ,  $\|D_x c_i\|_{\mathbf{L}^{\infty}([0,T] \times \Omega; \mathbb{R}^{n \times n})} \leq K_v \|w_i\|_{\mathbf{C}^0([0,T]; \mathbf{L}^1(\Omega; \mathbb{R}))}$ .

Step 2:  $A_i$  satisfies (A) in Theorem 2.3. Compute:

$$\|A_i\|_{\mathbf{L}^1([0,T];\mathbf{L}^\infty(\Omega;\mathbb{R}))} = \int_0^T \operatorname{ess\,sup}_{x\in\Omega} \left| \alpha\left(t, x, w_i(t)\right) \right| \mathrm{d}t$$
[By (3.16)]

$$\leq \int_{0}^{T} k_{\alpha}(t) \left( 1 + \left\| w_{i}(t) \right\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} \right) \mathrm{d}t$$

$$\leq \|k_{\alpha}\|_{\mathbf{L}^{1}(0,T];\mathbb{R})} \left( 1 + \|w_{i}\|_{\mathbf{L}^{\infty}([0,T];\mathbf{L}^{1}(\Omega;\mathbb{R}))} \right)$$
[By (\alpha)]

proving that  $A_i \in \mathbf{L}^1([0,T]; \mathbf{L}^{\infty}(\Omega; \mathbb{R}))$ . Moreover,

$$\int_{0}^{T} \operatorname{TV}\left(A_{i}(t)\right) \mathrm{d}t \leq \int_{0}^{T} K_{\alpha} \left(1 + \left\|\nabla w_{i}(t)\right\|_{\mathbf{L}^{1}(\Omega;\mathbb{R}^{n})}\right) \mathrm{d}t \qquad [By (\boldsymbol{\alpha})]$$
$$= K_{\alpha} \left(T + \left\|\nabla w_{i}\right\|_{\mathbf{L}^{1}([0,T]\times\Omega;\mathbb{R}^{n})}\right), \qquad [By (\mathbf{2}) \text{ in Proposition 3.5}]$$

proving that (A) holds.

**Step 3:**  $B_i \in \mathbf{L}^1([0,T]; \mathbf{L}^{\infty}(\Omega; \mathbb{R}))$ . Using  $(\boldsymbol{\beta})$  compute:

$$\|B_i\|_{\mathbf{L}^1([0,T];\mathbf{L}^\infty(\Omega;\mathbb{R}))} = \int_0^T \operatorname{ess\,sup}_{x\in\Omega} \left|\beta\left(\tau, x, u_i(\tau), w_i(\tau)\right)\right| \mathrm{d}\tau \le \|k_\beta\|_{\mathbf{L}^1([0,T];\mathbb{R})} \tag{3.18}$$

proving Step 3.

Hence Theorem 2.4 and Theorem 2.7 apply, ensuring that the sequence  $(u_i, w_i)$  can be recursively defined. The next steps aim at proving that  $(u_i, w_i)$  is a Cauchy sequence.

Step 4:  $w_i$  is bounded in  $\mathbf{L}^{\infty}([0,T]; \mathbf{L}^1(\Omega; \mathbb{R}))$  uniformly in *i*. By (2.13) and (3.18) we have

$$\begin{aligned} \left\| w_{i}(t) \right\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} &\leq K \left( \left\| w_{o} \right\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} + \left\| b \right\|_{\mathbf{L}^{1}([0,t]\times\Omega;\mathbb{R})} \right) \exp \left( K \left\| B_{i-1} \right\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}))} \right) \\ &\leq K_{w} \end{aligned}$$

where we set

$$K_{w} = K\left(\|w_{o}\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} + \|b\|_{\mathbf{L}^{1}([0,T]\times\Omega;\mathbb{R})}\right) \exp\left(K\|k_{\beta}\|_{\mathbf{L}^{1}([0,T];\mathbb{R})}\right).$$
(3.19)

Step 5: Bounds uniform in *i*. Note that  $c_i(t) \in \mathbf{C}^2(\Omega; \mathbb{R}^n)$  by  $(\boldsymbol{v})$  and (3.16). Moreover,

$$\|c_i(t)\|_{\mathbf{L}^{\infty}(\Omega;\mathbb{R}^n)} \le K_v \|w_i\|_{\mathbf{C}^0([0,t];\mathbf{L}^1(\Omega;\mathbb{R}))}$$
[By (3.17)]

$$\leq K_v K_w \qquad [By (3.19)]$$

$$\begin{aligned} \|D_x c_i\|_{\mathbf{L}^{\infty}([0,T] \times \Omega; \mathbb{R}^{n \times n})} &\leq K_v \|w_i\|_{\mathbf{C}^0([0,T]; \mathbf{L}^1(\Omega; \mathbb{R}))} \\ &\leq K_v K_w \end{aligned} \tag{By (v)} \\ \end{aligned}$$

$$\begin{aligned} \|\nabla\nabla \cdot c_i\|_{\mathbf{L}^1([0,T]\times\Omega;\mathbb{R}^n)} &= \int_0^T \left\|\nabla\nabla \cdot v\left(t, w_i(t)\right)\right\|_{\mathbf{L}^1(\Omega;\mathbb{R}^n)} \mathrm{d}t \\ &\leq \int_0^T \left\|D_x^2 v\left(t, w_i(t)\right)\right\|_{\mathbf{L}^1(\Omega;\mathbb{R}^n)} \mathrm{d}t \end{aligned} \tag{By (3.16)}$$

$$\leq \int_{0}^{T} k_{v} \left( t, \left\| w_{i}(t) \right\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} \right) \left\| w_{i}(t) \right\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} \mathrm{d}t \qquad [\mathrm{By} (\boldsymbol{v})]$$

which is bounded since  $k_v \in \mathbf{L}^{\infty}_{\mathbf{loc}}([0,T] \times \mathbb{R}_+; \mathbb{R}_+)$ . Using (3.19) and  $(\boldsymbol{v})$ , prepare for later use:

$$\begin{aligned} \|c_{i+1} - c_{i}\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}^{n}))} &\leq \int_{0}^{t} \left\| v\left(\tau, w_{i+1}(\tau)\right) - v\left(\tau, w_{i}(\tau)\right) \right\|_{\mathbf{L}^{1}(\Omega;\mathbb{R}^{n})} \mathrm{d}\tau \\ &\leq K_{v} \|w_{i+1} - w_{i}\|_{\mathbf{L}^{1}([0,t]\times\Omega;\mathbb{R})} \\ \|\nabla \cdot (c_{i+1} - c_{i})\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}^{n}))} &\leq \int_{0}^{t} \left\| \nabla \cdot \left( v\left(\tau, w_{i+1}(\tau)\right) - v\left(\tau, w_{i}(\tau)\right) \right) \right\|_{\mathbf{L}^{\infty}(\Omega;\mathbb{R})} \mathrm{d}\tau \\ &\leq \int_{0}^{t} k_{v}(\tau, K_{w}) \|w_{i+1}(\tau) - w_{i}(\tau)\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} \mathrm{d}\tau \\ &= \|k_{v}\|_{\mathbf{L}^{\infty}([0,t]\times[0,K_{w}];\mathbb{R})} \|w_{i+1} - w_{i}\|_{\mathbf{L}^{1}([0,t]\times\Omega;\mathbb{R})} .\end{aligned}$$

Passing to the  $A_i$ :

$$\|A_i\|_{\mathbf{L}^1([0,t];\mathbf{L}^{\infty}\Omega;\mathbb{R}))} = \int_0^t \operatorname{ess\,sup}_{x\in\Omega} \left| \alpha \left(\tau, x, w_i(\tau)\right) \right| \mathrm{d}\tau$$
[By (3.16)]

$$\leq \int_{0}^{t} k_{\alpha}(\tau) \left( 1 + \left\| w_{i}(\tau) \right\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} \right) \mathrm{d}\tau$$
[By (\alpha)]

$$\leq (1 + K_w) \|k_a\|_{\mathbf{L}^1([0,t];\mathbb{R})}; \qquad [By (3.19)]$$

$$\int_{0}^{} \operatorname{TV}(A_{i}(\tau)) \,\mathrm{d}\tau$$

$$\leq K_{a}\left(1 + \|\nabla w_{i}\|_{\mathbf{L}^{1}([0,t]\times\Omega;\mathbb{R}^{n})}\right)$$
[By Step 2]

$$\leq K_{a} + \frac{K_{\alpha}}{\mu} \|B_{i-1}\|_{\mathbf{L}^{1}([0,T];\mathbf{L}^{\infty}(\Omega;\mathbb{R}))} \|w_{i}\|_{\mathbf{L}^{\infty}([0,T];\mathbf{L}^{1}(\Omega;\mathbb{R}))} + \frac{K_{\alpha}}{\mu} \|b\|_{\mathbf{L}^{1}([0,T]\times\Omega;\mathbb{R})} + \frac{K_{\alpha}}{\mu} \|w_{o}\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})}$$
[By (2.11)]

$$\leq K_{a} + \frac{K_{\alpha}}{\mu} K_{w} \|k_{\beta}\|_{\mathbf{L}^{1}([0,T];\mathbb{R})}$$

$$+ \frac{K_{\alpha}}{\mu} \|b\|_{\mathbf{L}^{1}([0,T]\times\Omega;\mathbb{R})} + \frac{K_{\alpha}}{\mu} \|w_{o}\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})};$$
[By (3.18), (3.19)]

$$\begin{aligned} &\|A_{i+1} - A_i\|_{\mathbf{L}^1([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}))} \\ &= \int_0^t \|A_{i+1}(\tau) - A_i(\tau)\|_{\mathbf{L}^{\infty}(\Omega;\mathbb{R})} \,\mathrm{d}\tau \\ &= \int_0^t \|\alpha\left(\tau, \cdot, w_{i+1}(\tau)\right) - \alpha\left(\tau, \cdot, w_i(\tau)\right)\|_{\mathbf{L}^{\infty}(\Omega;\mathbb{R})} \,\mathrm{d}\tau \qquad \qquad [By (3.16)] \\ &\leq K_a \|w_{i+1} - w_i\|_{\mathbf{L}^1([0,t]\times\Omega;\mathbb{R})} \,. \end{aligned}$$

Step 6:  $(u_i, w_i)$  is a Cauchy sequence in  $\mathbf{C}^0([0, \Delta T]; \mathbf{L}^1(\Omega; \mathbb{R}^2))$  for  $\Delta T$  small. Thanks to the above bounds uniform in i, all the constants appearing in (H5) and (H7) are bounded uniformly by quantities depending on  $\Omega$ ,  $u_o$ , a, b and by the constants in the assumptions ( $\Omega$ ),  $(v), (\alpha), (a), (\beta), (b)$ . Hence, (H5) and (H7) yield

$$\left\|u_{i+1}(t) - u_i(t)\right\|_{\mathbf{L}^1(\Omega;\mathbb{R})}$$

$$\leq \mathcal{O}(1) \left( \|c_{i} - c_{i-1}\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}^{n}))} + \|\nabla \cdot (c_{i} - c_{i-1}\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}^{n}))} \right) \\ + \mathcal{O}(1) \left( \|A_{i} - A_{i-1}\|_{\mathbf{L}^{1}([0,t];\mathbf{L}^{\infty}(\Omega;\mathbb{R}))} \right) \\ \leq \mathcal{O}(1) \|w_{i} - w_{i-1}\|_{\mathbf{L}^{1}([0,t]\times\Omega;\mathbb{R})} .$$

To compute the distance  $\|w_{i+1}(t) - w_i(t)\|_{\mathbf{L}^1(\Omega:\mathbb{R})}$ , apply (P5) thanks to Step 3 and get

$$\leq K^{2} K_{\beta} \left( \|w_{o}\|_{\mathbf{L}^{1}(\Omega;\mathbb{R})} + \|b\|_{\mathbf{L}^{1}([0,t]\times\Omega;\mathbb{R})} \right) e^{2K \|k_{\beta}\|_{\mathbf{L}^{1}([0,t];\mathbb{R})}}$$
[By (3.18)]

$$\times \left( \|u_{i} - u_{i-1}\|_{\mathbf{L}^{1}([0,t] \times \Omega;\mathbb{R})} + \|w_{i} - w_{i-1}\|_{\mathbf{L}^{1}([0,t] \times \Omega;\mathbb{R})} \right) \,. \tag{By } (\boldsymbol{\beta})$$

As a consequence,

ш

$$\|u_{i+1} - u_i\|_{\mathbf{C}^0([0,\Delta T];\mathbf{L}^1(\Omega;\mathbb{R}))} + \|w_{i+1} - w_i\|_{\mathbf{C}^0([0,\Delta T];\mathbf{L}^1(\Omega;\mathbb{R}))}$$

$$\leq \mathcal{O}(1) \left( \|u_i - u_{i-1}\|_{\mathbf{L}^1([0,\Delta T]\times\Omega;\mathbb{R})} + \|w_i - w_{i-1}\|_{\mathbf{L}^1([0,\Delta T]\times\Omega;\mathbb{R})} \right)$$

$$\leq \mathcal{O}(1) \Delta T \left( \|u_i - u_{i-1}\|_{\mathbf{C}^0([0,\Delta T];\mathbf{L}^1(\Omega;\mathbb{R}))} + \|w_i - w_{i-1}\|_{\mathbf{C}^0([0,\Delta T];\mathbf{L}^1(\Omega;\mathbb{R}))} \right)$$

proving that for  $\Delta T$  small,  $(u_i, w_i)$  is a Cauchy sequence in  $\mathbf{C}^0([0, \Delta T]; \mathbf{L}^1(\Omega; \mathbb{R}))$ . Call (u, w)the corresponding limit.

Step 7: Problem (1.1) admits a global solution in the sense of Definition 2.1. Note that by  $(\boldsymbol{v}), (\boldsymbol{\alpha})$  and  $(\boldsymbol{\beta})$ , we can pass to the  $\mathbf{C}^0([0, \Delta T]; \mathbf{L}^1(\Omega; \mathbb{R}))$  limit also in (3.16). Hence, the Dominated Convergence Theorem allows to pass to the limit in (2.6), so that Definition 2.3 applies to u. A further use of Lemma 2.6 allows to pass to the limit also in (2.10), proving the existence of a solution on the time interval  $[0, \Delta T]$ .

Further iterations of the above procedure yield a solution, say (u, w), to (1.1) in the sense of Definition 2.1. Call  $[0, T_*]$ , for a  $T_* > \Delta T$ , the biggest time interval on which (u, w) can be extended.

Define  $B(t,x) = \beta(t,x,u(t),w(t))$ . By  $(\beta), B \in \mathbf{L}^1([0,T_*];\mathbf{L}^\infty(\Omega;\mathbb{R}))$ . Hence, Theorem 2.7 ensures that problem (2.2) admits a solution w on  $[0, T_*]$  in the sense of Definition 2.5. By (P1),  $w \in \mathbf{C}^0([0, T_*]; \mathbf{L}^1(\Omega; \mathbb{R})).$ 

Define c(t, x) = v(t, w(t))(x),  $A(t, x) = \alpha(t, x, w(t))$  and repeat the same computations as in Step 1 and Step 2 to obtain that problem (2.1) admits as solution u on  $[0, T_*]$  in the sense of Definition 2.3. By (H6), u is continuous on  $[0, T_*]$ .

Thus, (u, w) is extended up to time  $T_*$ . If  $T_* < T$ , the above procedure can be repeated with reference to problem (1.1) with initial datum  $(u, w)(T_*)$  assigned at time  $T_*$ , obtaining an extension of (u, w) beyond time  $T_*$ , hence contradicting the maximality of  $T_*$ , unless  $T_* = T$ . This proves the global existence stated in (M1).

The continuity at (M2) follows from (H6) and (P1).

The continuous dependence and stability estimates (M3), and (M4) follow through long and tedious computations based on the estimates obtained so far. More precisely, to prove (M3), use (H4), (H5), (H7), (P4), (P5) and repeat the computations in Step 3, 4, 5 and 6. To prove (M4) the procedure is entirely similar, also using (P6).
The proof of (M5) is a direct consequence of (H3) and (P7).

**Acknowledgment:** The authors acknowledge the PRIN 2022 project *Modeling, Control* and Games through Partial Differential Equations (D53D23005620006), funded by the European Union - Next Generation EU. The first two authors also acknowledge the 2024 GNAMPA project *Modelling and Analysis through Conservation Laws*.

**Data Availability Statement:** Data sharing not applicable to this article as no data set were generated nor analysed during the current study.

### References

- C. Bardos, A. Y. le Roux, and J.-C. Nédélec. First order quasilinear equations with boundary conditions. Comm. Partial Differential Equations, 4(9):1017–1034, 1979.
- [2] H. Brezis. Functional analysis, Sobolev spaces and partial differential equations. Universitext. Springer, New York, 2011.
- J. Choi and S. Kim. Green's function for second order parabolic systems with Neumann boundary condition. J. Differential Equations, 254(7):2834–2860, 2013.
- [4] R. M. Colombo, M. Garavello, F. Marcellini, and E. Rossi. General renewal equations motivated by biology and epidemiology. J. Differential Equations, 354:133–169, 2023.
- [5] R. M. Colombo and E. Rossi. Hyperbolic predators vs. parabolic prey. Commun. Math. Sci., 13(2):369–400, 2015.
- [6] R. M. Colombo and E. Rossi. Nonlocal conservation laws in bounded domains. SIAM J. Math. Anal., 50(4):4041–4065, 2018.
- [7] R. M. Colombo and E. Rossi. Well-posedness and control in a hyperbolic-parabolic parasitoidparasite system. *Stud. Appl. Math.*, 147(3):839–871, 2021.
- [8] R. M. Colombo and E. Rossi. Non linear hyperbolic-parabolic systems with Dirichlet boundary conditions. *Diff. Int. Eq.*, 2024.
- [9] W. Dan. On a local in time solvability of the Neumann problem of quasilinear hyperbolic parabolic coupled systems. *Math. Methods Appl. Sci.*, 18(13):1053–1082, 1995.
- [10] S. Ebenfeld. Non-linear initial boundary value problems of hyperbolic-parabolic type. A general investigation of admissible couplings between systems of higher order. II. Abstract quasilinear theory. *Math. Methods Appl. Sci.*, 25(3):213–240, 2002.
- [11] A. Friedman. Partial differential equations of parabolic type. Prentice-Hall Inc., Englewood Cliffs, N.J., 1964.
- [12] M. Fuest. Global weak solutions to fully cross-diffusive systems with nonlinear diffusion and saturated taxis sensitivity. *Nonlinearity*, 35(1):608–657, 2022.
- [13] E. Giusti. Minimal surfaces and functions of bounded variation, volume 80 of Monographs in Mathematics. Birkhäuser Verlag, Basel, 1984.
- [14] G. Hong, H. Peng, Z.-A. Wang, and C. Zhu. Nonlinear stability of phase transition steady states to a hyperbolic-parabolic system modeling vascular networks. J. Lond. Math. Soc. (2), 103(4):1480–1514, 2021.

- [15] M. Iannelli and G. Marinoschi. Well posedness for a hyperbolic-parabolic Cauchy problem arising in population dynamics. *Diff. Int. Eq.*, 21(9-10):917–934, 2008.
- [16] V. A. Kozlov and S. A. Nazarov. A simple one dimensional model of a false aneurysm in the femoral artery. J. Math. Sci. (N.Y.), 214(3):287–301, 2016.
- [17] O. A. Ladyženskaja, V. A. Solonnikov, and N. N. Uralceva. *Linear and quasilinear equations of parabolic type*. Translations of Mathematical Monographs, Vol. 23. American Mathematical Society, Providence, R.I., 1968. Translated from the Russian by S. Smith.
- [18] G. Li and Y. Yao. Two-species competition model with chemotaxis: well-posedness, stability and dynamics. *Nonlinearity*, 35(3):1329–1359, 2022.
- [19] S. Martin. First order quasilinear equations with boundary conditions in the  $L^{\infty}$  framework. J. Differential Equations, 236(2):375–406, 2007.
- [20] F. Pfab, M. Stacconi, G. Anfora, A. Grassi, V. Walton, and A. Pugliese. Optimized timing of parasitoid release: a mathematical model for biological control of Drosophila suzukii. *Theoretical Ecology*, 11(4):489–501, 2018.
- [21] E. Rossi. Definitions of solutions to the IBVP for multi-dimensional scalar balance laws. J. Hyperbolic Differ. Equ., 15(2):349–374, 2018.
- [22] S. Salsa. *Partial differential equations in action*, volume 86 of *Unitext*. Springer, Cham, second edition, 2015. From modelling to theory.
- [23] S. Sastre-Gomez and J. I. Tello. On the existence of solutions for a parabolic-elliptic chemotaxis model with flux limitation and logistic source. *Math. Methods Appl. Sci.*, 46(8):9252–9267, 2023.
- [24] J. I. Tello and D. Wrzosek. Predator-prey model with diffusion and indirect prey-taxis. Math. Models Methods Appl. Sci., 26(11):2129–2162, 2016.
- [25] J. Vovelle. Convergence of finite volume monotone schemes for scalar conservation laws on bounded domains. Numer. Math., 90(3):563–596, 2002.
- [26] P. Wojtaszczyk. Banach spaces for analysts, volume 25 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 1991.