On the relevance of GIKS instability of charged $\mathcal{N} = 4$ SYM plasma

Alex Buchel

Department of Physics and Astronomy University of Western Ontario London, Ontario N6A 5B7, Canada Perimeter Institute for Theoretical Physics Waterloo, Ontario N2J 2W9, Canada

Abstract

Recently Gladden, Ivo, Kovtun, and Starinets (GIKS) identified a novel instability of $\mathcal{N} = 4$ supersymmetric Yang-Mills plasma with a diagonal U(1) *R*-symmetry chemical potential. Since $\mathcal{N} = 4$ SYM plasma has *R*-charged matter fields, it might suffer from the standard superconducting instability. We show that while the superconducting instability is indeed present, it happens at a lower critical temperature compare to that of the GIKS instability.

February 16, 2025

Strongly coupled $\mathcal{N} = 4 SU(N_c)$ supersymmetric Yang-Mills (SYM) theory has a holographic dual as type IIB string theory in $AdS_5 \times S^5$ [1]. In the planar limit, and at large N_c , the thermal states of the SYM are represented by black branes in type IIB supergravity. This gauge theory has SU(4) *R*-symmetry, and one can study its charged plasma, with the same chemical chemical potential μ for all U(1) factors of the maximal Abelian subgroup $U(1)^3 \subset SU(4)$. In this case the gravitational dual is represented by a Reissner-Nordstrom (RN) black brane in asymptotically AdS_5 spacetime. One reason why this model is interesting, is that it allows to study extremal horizons (where it is possible to take a limit of vanishing temperature) in the best explored holographic example.

In [2] (GIKS) the authors argued that the extremal limit is never achieved¹: there is a critical temperature, more precisely the critical value of $\frac{T}{\mu}$, below which the charged SYM plasma is unstable to fluctuations of the off-diagonal gauge fields and the neutral scalars of the bulk gravitational description, when interpreted within STU consistent truncation [4–6] of type IIB supergravity. However, $\mathcal{N} = 4$ SYM also has *R*-charged matter fields, which could condense at sufficiently low values of $\frac{T}{\mu}$ [7]. STU truncation does not provide gravitational description of such charged matter.

The question we address in this note is whether the GIKS instability is physically relevant: *i.e.*, whether the standard superconducting instability in charged SYM plasma occurs before the former can be triggered. Jumping to the result, we find that

$$\frac{T_c}{\mu}\Big|_{\text{superconducting}} = 0.57186(4) \left.\frac{T_c}{\mu}\right|_{\text{GIKS}},\tag{0.1}$$

i.e., the superconducting instability actually occurs at a lower temperature compare to the onset of the instability reported in [2]. Our result highlights the importance of understanding the end-point of the linearized GIKS instability.

The rest of this note explains how we reached the conclusion (0.1). As we already alluded to, it is important to enlarge the dual gravitational description beyond STU.

¹Identical phenomenon occurs in other holographic models as well [3].

Luckily, such a model is known [8]:

$$S_{5} = \frac{1}{4\pi G_{5}} \int_{\mathcal{M}_{5}} d^{5}\xi \sqrt{-g} \left[\frac{R}{4} - \frac{1}{4} \left(\rho^{4} \nu^{-4} F_{\mu\nu}^{(1)} F^{(1)\mu\nu} + \rho^{4} \nu^{4} F_{\mu\nu}^{(2)} F^{(2)\mu\nu} + \rho^{-8} F_{\mu\nu}^{(3)} F^{(3)\mu\nu} \right) \right. \\ \left. - \frac{1}{2} \sum_{j=1}^{4} \left(\partial_{\mu} \phi_{j} \right)^{2} - 3 \left(\partial_{\mu} \alpha \right)^{2} - \left(\partial_{\mu} \beta \right)^{2} - \frac{1}{8} \sinh^{2} (2\phi_{1}) \left(\partial_{\mu} \theta_{1} + \left(A_{\mu}^{(1)} + A_{\mu}^{(2)} - A_{\mu}^{(3)} \right) \right)^{2} \right. \\ \left. - \frac{1}{8} \sinh^{2} (2\phi_{2}) \left(\partial_{\mu} \theta_{2} + \left(A_{\mu}^{(1)} - A_{\mu}^{(2)} + A_{\mu}^{(3)} \right) \right)^{2} - \frac{1}{8} \sinh^{2} (2\phi_{3}) \left(\partial_{\mu} \theta_{3} + \left(-A_{\mu}^{(1)} + A_{\mu}^{(2)} + A_{\mu}^{(3)} \right) \right)^{2} - \frac{1}{8} \sinh^{2} (2\phi_{4}) \left(\partial_{\mu} \theta_{4} - \left(A_{\mu}^{(1)} + A_{\mu}^{(2)} + A_{\mu}^{(3)} \right) \right)^{2} - \mathcal{P} \right].$$

$$(0.2)$$

Here the $F^{(J)}$ are the field strengths of the U(1) gauge fields $A^{(J)}$ (dual to $U(1)^3 \subset SU(4)$ global *R*-symmetry currents), α and β are the STU-truncation bulk scalars, ϕ_i are the bulk scalars dual to dimension-3 fermion bilinears of the boundary SYM, and \mathcal{P} is the scalar potential. We introduced

$$\rho \equiv e^{\alpha} \,, \qquad \nu \equiv e^{\beta} \,. \tag{0.3}$$

The scalar potential, \mathcal{P} , is given in terms of a superpotential

$$\mathcal{P} = \frac{g^2}{8} \left[\sum_{j=1}^{4} \left(\frac{\partial W}{\partial \phi_j} \right)^2 + \frac{1}{6} \left(\frac{\partial W}{\partial \alpha} \right)^2 + \frac{1}{2} \left(\frac{\partial W}{\partial \beta} \right)^2 \right] - \frac{g^2}{3} W^2, \qquad (0.4)$$

where

$$W = -\frac{1}{4\rho^{2}\nu^{2}} \left[\left(1 + \nu^{4} - \nu^{2}\rho^{6} \right) \cosh(2\phi_{1}) + \left(-1 + \nu^{4} + \nu^{2}\rho^{6} \right) \cosh(2\phi_{2}) + \left(1 - \nu^{4} + \nu^{2}\rho^{6} \right) \cosh(2\phi_{3}) + \left(1 + \nu^{4} + \nu^{2}\rho^{6} \right) \cosh(2\phi_{4}) \right].$$

$$(0.5)$$

In what follows we set the gauged supergravity coupling g = 1, this corresponds to setting the asymptotic AdS_5 radius to L = 2. The five dimensional gravitational constant G_5 is related to the rank of the supersymmetric $\mathcal{N} = 4 SU(N_c)$ UV fixed point as

$$G_5 = \frac{4\pi}{N^2}.$$
 (0.6)

Different consistent truncations of (0.5) were discussed in [9]. For this paper we care about²:

²While we do not present the details here, we did verify that other then the instabilities of (**Tr-II**) and (**Tr-III**) discussed below, the RN black branes of (**Tr-I**) do not have additional spatially homogeneous and isotropic instabilities, when described within the gravitational dual (0.2).

• (Tr-I), the minimal holographic dual to the SYM with a diagonal U(1) *R*-symmetry is

$$\alpha = \beta = \phi_j = \theta_j = 0, \qquad A_{\mu}^{(1)} = A_{\mu}^{(2)} = A_{\mu}^{(3)}; \qquad (0.7)$$

• (Tr-II), the STU model,

$$\phi_j = \theta_j = 0; \tag{0.8}$$

• (Tr-III), the truncation describing the superconducting phase of the charged $\mathcal{N} = 4$ SYM plasma,

$$\alpha = \beta = \phi_{1,2,3} = \theta_{1,2,3} = 0, \qquad A_{\mu}^{(1)} = A_{\mu}^{(2)} = A_{\mu}^{(3)}. \tag{0.9}$$

(Tr-I): The RN black brane is a solution of the effective action (0.2) with (0.7):

$$ds_5 = \frac{4}{r} \left(-f \ dt^2 + d\boldsymbol{x}^2 \right) + \frac{dr^2}{r^2 f}, \qquad f = 1 - (p^2 + 1) \ r^2 + p^2 \ r^3$$

$$A^{(j)}_{\mu} = p \ (1 - r) \ \delta^t_{\mu}, \qquad (0.10)$$

where the radial coordinate $r \in (0, 1]$ (from the asymptotic AdS_5 boundary to the regular Schwarzschild horizon), and parameter p determines

$$\frac{T}{\mu} = \frac{2 - p^2}{2\pi p},$$
(0.11)

with the extremal limit achieved as $p \to \sqrt{2}$.

<u>(Tr-II)</u>: The RN black brane (0.10), is unstable in grand canonical ensemble, within the STU model (0.8), to an ordered phase [10, 11] below³

$$\frac{T_c}{\mu} = \frac{1}{2\pi} \equiv \frac{T_c}{\mu} \Big|_{\text{GIKS}}, \qquad (0.12)$$

as it turns out that (0.12) is also the onset of the GIKS instability⁴. The ordered phase of the charged SYM plasma is however thermodynamically unstable [2, 11], and thus is expected [2, 14, 15] to suffer from the same charge clumping instabilities as that of the dual to the RN black brane (0.10).

³See [12, 13] for earlier related work.

⁴The $\sqrt{2}$ difference in the critical value of the temperature is due to a different normalization of the gauge fields in [2] and in (0.5).

(Tr-III): Given (0.9), the effective action (0.5) takes the form:

$$S_{5} = \frac{1}{16\pi G_{5}} \int_{\mathcal{M}_{5}} d^{5}\xi \sqrt{-g} \left[R - 3F_{\mu\nu}F^{\mu\nu} - 2\left(\partial_{\mu}\phi\right)^{2} - \frac{1}{2}\sinh^{2}(2\phi)\left(\partial_{\mu}\theta - 3A_{\mu}\right)^{2} - 4\mathcal{P} \right],$$
(0.13)

where we introduced $\phi_4 \equiv \phi$, $\theta_4 \equiv \theta$, and $A^{(j)}_{\mu} \equiv A_{\mu}$. Following [7], to have a properly normalized *R*-charge 2 superpotential of the boundary SYM we need to rescale the bulk gauge field as

$$A_{\mu} = \frac{2}{3}\hat{A}_{\mu} \,. \tag{0.14}$$

Further introducing $\phi \equiv \frac{\eta}{2}$, the effective action (0.13) becomes identical to that of [7]. The superconducting phase transition in [7] was reported at

$$\frac{T_c}{\hat{\mu}} = 0.060676(6) \,, \tag{0.15}$$

where $\hat{\mu}$ is the chemical potential of the \hat{A} gauge field. Thus, given (0.14), the phase transition in the effective action (0.13) is at⁵

$$\frac{T_c}{\mu}\Big|_{\text{superconducting}} = \frac{3}{2} \cdot 0.060676(6) = 0.091014(9)
= 0.57186(4) \cdot \frac{1}{2\pi} = 0.57186(4) \cdot \frac{T_c}{\mu}\Big|_{\text{GIKS}},$$
(0.16)

where we used (0.12).

Acknowledgments

Research at Perimeter Institute is supported by the Government of Canada through Industry Canada and by the Province of Ontario through the Ministry of Research & Innovation. This work was further supported by NSERC through the Discovery Grants program.

References

 J. M. Maldacena, The Large N limit of superconformal field theories and supergravity, Int. J. Theor. Phys. 38 (1999) 1113–1133, [hep-th/9711200].

⁵We recovered this result directly from the effective action (0.13).

- [2] L. Gladden, V. Ivo, P. Kovtun and A. O. Starinets, *Instability in* $\mathcal{N} = 4$ supersymmetric Yang-Mills theory at finite density, 2412.12353.
- [3] A. Buchel, Instability of Baryonic Black Branes, 2502.05971.
- [4] K. Behrndt, M. Cvetic and W. A. Sabra, Nonextreme black holes of five-dimensional N=2 AdS supergravity, Nucl. Phys. B553 (1999) 317-332, [hep-th/9810227].
- [5] M. Cvetic, M. J. Duff, P. Hoxha, J. T. Liu, H. Lu, J. X. Lu et al., Embedding AdS black holes in ten-dimensions and eleven-dimensions, Nucl. Phys. B 558 (1999) 96-126, [hep-th/9903214].
- [6] M. Cvetic, H. Lu, C. N. Pope, A. Sadrzadeh and T. A. Tran, Consistent SO(6) reduction of type IIB supergravity on S**5, Nucl. Phys. B 586 (2000) 275–286, [hep-th/0003103].
- [7] S. S. Gubser, C. P. Herzog, S. S. Pufu and T. Tesileanu, Superconductors from Superstrings, Phys. Rev. Lett. 103 (2009) 141601, [0907.3510].
- [8] N. Bobev, A. Kundu, K. Pilch and N. P. Warner, Supersymmetric Charged Clouds in AdS₅, JHEP 03 (2011) 070, [1005.3552].
- [9] A. Buchel and B. Umbert, QGP universality in a magnetic field?, JHEP 06 (2020) 149, [2004.01583].
- [10] A. Buchel, The ordered phase of charged N=4 SYM plasma, 2501.01856.
- [11] A. Buchel, The ordered phase of charged N=4 SYM plasma from STU, 2501.12403.
- [12] O. Henriksson, C. Hoyos and N. Jokela, Novel color superconducting phases of $\mathcal{N} = 4$ super Yang-Mills at strong coupling, JHEP **09** (2019) 088, [1907.01562].
- [13] A. Anabalon and J. Oliva, Plasma-Plasma Third Order Phase Transition from Type IIB Supergravity, Phys. Rev. Lett. 133 (2024) 121601, [2405.04611].
- [14] A. Buchel, A Holographic perspective on Gubser-Mitra conjecture, Nucl. Phys. B731 (2005) 109–124, [hep-th/0507275].

 [15] A. Buchel, Critical phenomena in N=4 SYM plasma, Nucl. Phys. B 841 (2010) 59–99, [1005.0819].