NON-UNIQUENESS OF REGULAR SOLUTIONS FOR INCOMPRESSIBLE STATIC EULER EQUATIONS WITH GIVEN BOUNDARY CONDITIONS AND TURBULENT GLOBAL SOLUTIONS OF INCOMPRESSIBLE NAVIER-STOKES EQUATIONS

YONGQIAN HAN

ABSTRACT. The incompressible Navier-Stokes equations and static Euler equations are considered. We find that there exist infinite non-trivial regular solutions of incompressible static Euler equations with given boundary conditions. Moreover there exist random solutions of incompressible static Euler equations. Provided Reynolds number is large enough and time variable t goes to infinity, these random solutions of static Euler equations are the path limits of corresponding Navier-Stokes flows. But the double limits of these Navier-Stokes flows do not exist. These phenomena reveal randomness and turbulence of incompressible fluids. Therefore these solutions are called turbulent solutions. Here some typing models without Prandtl layer are given.

1. INTRODUCTION

The incompressible Navier-Stokes equations in \mathbb{R}^3 are given by

(1.1)
$$u_t - \nu \Delta u + (u \cdot \nabla)u + \nabla P = 0,$$

(1.2)
$$\nabla \cdot u = 0,$$

where $u = u(t, x) = (u^1(t, x), u^2(t, x), u^3(t, x))$ and P = P(t, x) stand for the unknown velocity vector field of fluid and its pressure, $\nu > 0$ is the coefficient of viscosity. Here $x = (x_1, x_2, x_3), \nabla = (\partial_1, \partial_2, \partial_3)$ and $\partial_j = \frac{\partial}{\partial x_j}$ (j = 1, 2, 3).

Taking curl with equation (1.1), we have

(1.3)
$$\omega_t - \nu \Delta \omega + (u \cdot \nabla) \omega - (\omega \cdot \nabla) u = 0.$$

Here vorticity $\omega = (\omega^1, \omega^2, \omega^3)$ and

(1.4)
$$\omega(t,x) = \nabla \times u(t,x).$$

The incompressible static Euler equations in \mathbb{R}^3 are given by

(1.5)
$$(u \cdot \nabla)u + \nabla P = 0,$$

(1.6)
$$\nabla \cdot u = 0.$$

By taking curl, the equation (1.5) is equivalent to

(1.7)
$$(u \cdot \nabla)\omega - (\omega \cdot \nabla)u = 0.$$

²⁰⁰⁰ Mathematics Subject Classification. 35Q30, 76D05, 76F02, 37L20.

Key words and phrases. Incompressible Navier-Stokes Equations, Incompressible Static Euler Equations, Randomness and Turbulence of Incompressible Fluids, Typing Models without Prandtl Layer.

The velocity vector u is solution of equations (1.1) (1.2) provided (u, ω) satisfies equation (1.3). The velocity vector u is solution of equations (1.5) (1.6) provided (u, ω) satisfies equation (1.7).

The Fourier transformation of u(t, x) with respect to x denotes by $\hat{u}(t, \xi)$. Then equation (1.2) and (1.6) are equivalent to

(1.8)
$$\xi_j \hat{u}^j = \xi_1 \hat{u}^1 + \xi_2 \hat{u}^2 + \xi_3 \hat{u}^3 = 0.$$

It is that $\hat{u}(t,\xi)$ is perpendicular to $\xi \in \mathbb{R}^3$. Denote by $\xi \perp \hat{u}$ for any $\xi \in \mathbb{R}^3$.

There is a large literature (for instance, see [1, 2, 3, 5, 8, 9, 11] and therein references) studying the incompressible Navier-Stokes and steady Euler equations.

To construct solutions of Navier-Stokes equations (1.1) (1.2) and Euler equations (1.5) (1.6), we define some linear operators which are perpendicular to operator ∇ .

Definition 1.1. Let domain $D \subset \mathbb{R}^3$. Assume that a vector $v(x) = (v_1(x), v_2(x), v_3(x))$ satisfies equation (1.2) for any $x \in D$ and the Fourier transformation $\hat{v}(\xi)$ is well defined.

First provided D is bounded, we define a linear operator $v(\nabla)$ as follows

(1.9)
$$(v(\nabla)f)^{\hat{}}(\xi) = \hat{v}(\xi)\hat{f}(\xi), \quad \forall f \in C(D),$$

where $\hat{f}(\xi)$ is the Fourier transformation of f(x).

Next provided $D = \mathbb{R}^3$, the linear operator $v(\nabla)$ can be defined in space $C_c^{\infty}(\mathbb{R}^3)$

(1.10)
$$(v(\nabla)f)^{\hat{}}(\xi) = \hat{v}(\xi)\hat{f}(\xi), \quad \forall f \in C_c^{\infty}(\mathbb{R}^3).$$

Especially let a vector $P = P(x) = (p_1(x), p_2(x), p_3(x))$ and $p_j(x)$ (j = 1, 2, 3) be polynomial. Provided polynomial vector P satisfies

(1.11)
$$x \cdot P(x) = x_j p_j(x) = x_1 p_1(x) + x_2 p_2(x) + x_3 p_3(x) \equiv 0, \quad \forall x \in \mathbb{R}^3.$$

Then we define linear operator $P(\nabla) = (p_1(\nabla), p_2(\nabla), p_3(\nabla))$. Now we have

(1.12)
$$(P(\nabla)f)^{\hat{}}(\xi) = P(i\xi)f(\xi) = (p_1(i\xi)\hat{f}(\xi), p_2(i\xi)\hat{f}(\xi), p_3(i\xi)\hat{f}(\xi)), \quad \forall f \in C_c^{\infty}(\mathbb{R}^3),$$

where $\hat{f}(\xi)$ is the Fourier transformation of f(x).

It is obvious that any linear operator $A(\nabla)$ defined in Definition 1.1 is perpendicular to operator ∇ . Let velocity vector u(x) be the solution of static Euler equations (1.5) (1.6) and regular enough, then we can define a linear operator $u(\nabla)$ by using Definition 1.1.

Choose a real linear operator $A = A(\nabla) = (a_1(\nabla), a_2(\nabla), a_3(\nabla))$ be defined in Definition 1.1, then $A(\nabla)$ is perpendicular to ∇ , i.e. $\nabla \cdot A(\nabla) = 0$. Since $\nabla, A(\nabla)$ and $A(\nabla) \times \nabla$ are vertical two by two, we can write velocity vector u as follows

$$u(t,x) = A(\nabla)\phi(t,x) + \{A(\nabla) \times \nabla\}\psi(t,x)$$

$$(1.13) = \left(a_1(\nabla)\phi + (a_2(\nabla)\partial_3 - a_3(\nabla)\partial_2)\psi, a_2(\nabla)\phi + (a_3(\nabla)\partial_1 - a_1(\nabla)\partial_3)\psi, a_3(\nabla)\phi + (a_1(\nabla)\partial_2 - a_2(\nabla)\partial_1)\psi\right)(t,x),$$

where ϕ and ψ are real scalar functions. Then u satisfies the equation (1.2) and (1.6). We call that the formulation (1.13) is symplectic representation of velocity vector u.

 $\mathbf{2}$

Now the curl of velocity vector u can be rewritten as follows

$$(1.14) \begin{aligned} \omega(t,x) &= \nabla \times u(t,x) \\ &= -\{(A(\nabla) \times \nabla) \phi(t,x) - \{(A(\nabla) \times \nabla) \times \nabla\} \psi(t,x) \\ &= -\{(A(\nabla) \times \nabla) \phi(t,x) + A(\nabla) \Delta \psi(t,x) \\ &= \left(-(a_2(\nabla)\partial_3 - a_3(\nabla)\partial_2) \phi + a_1(\nabla) \Delta \psi, \\ &- (a_3(\nabla)\partial_1 - a_1(\nabla)\partial_3) \phi + a_2(\nabla) \Delta \psi, \\ &- (a_1(\nabla)\partial_2 - a_2(\nabla)\partial_1) \phi + a_3(\nabla) \Delta \psi \right)(t,x). \end{aligned}$$

Some turbulent global solutions of equations (1.1) (1.2) have been constructed in [4] by using (1.13) (1.14) with special linear operator $A(\nabla) = e_3 \times \nabla$, provided (ϕ, ψ) is radial symmetric and cylindrical symmetric respectively.

By employing equations (1.3) (1.7) (1.13) (1.14), some new interesting results are obtained in this paper. The main result is as follows

Theorem 1.2 (Global Turbulent Solution). Assume that $A(\nabla)$ is any linear operator defined by Definition 1.1, λ is constant, $\psi(x)$ is regular enough and satisfies the following equation

(1.15)
$$-\Delta\psi(x) = \lambda^2\psi(x),$$

and $\phi(x) = \lambda \psi(x)$. Let

(1.16)
$$u_e(x) = A(\nabla)\phi(x) + \{A(\nabla) \times \nabla\}\psi(x) \\ = \lambda A(\nabla)\psi(x) + \{A(\nabla) \times \nabla\}\psi(x),$$

then $u = u_e(x)$ is the solution of static incompressible Euler equations (1.5) (1.6). Moreover let

(1.17)
$$u_{ns}(t,x) = e^{-\nu\lambda^{2}t}A(\nabla)\phi(x) + e^{-\nu\lambda^{2}t}\{A(\nabla)\times\nabla\}\psi(x)$$
$$= \lambda e^{-\nu\lambda^{2}t}A(\nabla)\psi(x) + e^{-\nu\lambda^{2}t}\{A(\nabla)\times\nabla\}\psi(x),$$

then $u = u_{ns}$ is the solution of incompressible Navier-Stokes equations (1.1) (1.2).

It is easy to prove this Theorem 1.2. In fact, inserting $\phi(x) = \lambda \psi(x)$ and equation (1.15) into (1.13) (1.14), we find $\omega = -\lambda u$. Inserting $\omega = -\lambda u$ into equations (1.3) (1.7), we can prove Theorem 1.2.

By taking samples of $(A(\nabla), \lambda, \phi, \psi)$ in Theorem 1.2, some new interesting results, such as non-uniqueness of static incompressible Euler equations (1.5) (1.6) with given boundary conditions, can be shown clearly.

Let us solve the equation (1.15) with periodic boundary condition

(1.18)
$$\psi(x+2\pi) = \psi(x), \quad \forall x \in \mathbb{R}^3$$

This is eigenvalue problem. By spectrum theory of compact operator $(-\Delta)^{-1}$, there exist countable infinite eigenvalues

 $0 \le \lambda_1 \le \dots \le \lambda_n \le \dots, \quad n = 1, 2, \dots$

Each eigenvalue is real, and may be geometrically simple or finite.

For eigenvalue λ_n of problem (1.15) (1.18), we define linear spaces E_n and E_n^* as follows

$$(1.19) \qquad E_n = \{\psi_n(x) | \text{any eigenfunction } \psi_n \text{ corresponding to eigenvalue } \lambda_n \}, \\ E_n^* = \text{span}\{ \sin(k_1 x_1) \sin(k_2 x_2) \sin(k_3 x_3), \sin(k_1 x_1) \sin(k_2 x_2) \cos(k_3 x_3), \\ \sin(k_1 x_1) \cos(k_2 x_2) \sin(k_3 x_3), \sin(k_1 x_1) \cos(k_2 x_2) \cos(k_3 x_3), \\ (1.20) \qquad \cos(k_1 x_1) \sin(k_2 x_2) \sin(k_3 x_3), \cos(k_1 x_1) \sin(k_2 x_2) \cos(k_3 x_3), \\ \cos(k_1 x_1) \cos(k_2 x_2) \sin(k_3 x_3), \cos(k_1 x_1) \cos(k_2 x_2) \cos(k_3 x_3), \\ \lambda_n^2 = k_1^2 + k_2^2 + k_3^2, \text{ integer } k_j \ge 0, \quad j = 1, 2, 3 \}.$$

It is obvious that $E_n^* \subseteq E_n$.

As the consequence of Theorem 1.2, we have

Theorem 1.3 (Global Turbulent Periodic Solution). Let $A(\nabla)$ be any linear operator which is defined in Definition 1.1 by using polynomials. Provided that λ_n is eigenvalue and ψ_n is eigenfunction of problem (1.15) (1.18), $\psi_n \in E_n$, $\phi_n(x) = \lambda_n \psi_n(x)$ ($n = 1, 2, \cdots$). Let

(1.21)
$$u_e^n(x) = A(\nabla)\phi_n(x) + \{A(\nabla) \times \nabla\}\psi_n(x) \\ = \lambda_n A(\nabla)\psi_n(x) + \{A(\nabla) \times \nabla\}\psi_n(x)$$

then $u = u_e^n(x)$ $(n = 1, 2, \dots)$ is the solution of static incompressible Euler equations (1.5) (1.6) with periodic boundary condition

(1.22)
$$u(x+2\pi) = u(x), \quad \forall x \in \mathbb{R}^3$$

Moreover let

(1.23)
$$u_{ns}^{n}(t,x) = e^{-\nu\lambda_{n}^{2}t}A(\nabla)\phi_{n}(x) + e^{-\nu\lambda_{n}^{2}t}\{A(\nabla)\times\nabla\}\psi_{n}(x)$$
$$= \lambda_{n}e^{-\nu\lambda_{n}^{2}t}A(\nabla)\psi_{n}(x) + e^{-\nu\lambda_{n}^{2}t}\{A(\nabla)\times\nabla\}\psi_{n}(x),$$

then $u = u_{ns}^n(t,x)$ $(n = 1, 2, \dots)$ is the solution of incompressible Navier-Stokes equations (1.1) (1.2) with periodic boundary condition

(1.24)
$$u(t, x + 2\pi) = u(t, x), \quad \forall t \ge 0, \ x \in \mathbb{R}^3$$

and initial data

(1.25)
$$u(t,x)|_{t=0} = u_e^n(x)$$

It is obvious that the following corollaries are the consequences of Theorem 1.3.

Corollary 1.4 (Non-unique Periodic Solutions of Euler Equations). The solution of static incompressible Euler equations (1.5) (1.6) with periodic boundary condition (1.22) is non-unique.

Corollary 1.5 (Global Well-posedness of NS Periodic Solution). The incompressible Navier-Stokes equations (1.1) (1.2) with periodic boundary condition (1.24) and initial data

(1.26)
$$u(t,x)\Big|_{t=0} = u_0(x)$$

is global well-posed, provided $u_0 \in B$. Here subset B of C_{per}^{∞} is defined by

(1.27)
$$B = \bigcup_{A(\nabla)} \bigcup_{n=1,2,\cdots} \{ u(x) | u = \lambda_n A(\nabla) \psi_n(x) + \{ A(\nabla) \times \nabla \} \psi_n(x), \forall \psi_n \in E_n \}.$$

4

 $A(\nabla)$ is any linear operator which is defined in Definition 1.1 by using polynomials.

To show randomness and turbulence of incompressible fluids, we consider limits of solution u_{ns}^n of Navier-Stokes equations as $\nu \to 0$ and $t \to \infty$.

Let μ be random number and Γ_{μ} be a path. In the path Γ_{μ} , we have

$$(1.28) t\nu = \mu$$

for any $(\nu, t) \in \Gamma_{\mu}$.

For the solution $u_{ns}^n(t,x)$ of Navier-Stokes equations constructed in (1.23), we have path limit

(1.29)
$$u_{e}^{n,\mu}(x) = \lim_{(\nu,t)\in\Gamma_{\mu},(\nu,1/t)\to(0,0)} u_{ns}^{n}(t,x)$$
$$= \lambda_{n}e^{-\mu\lambda_{n}^{2}}A(\nabla)\psi_{n}(x) + e^{-\mu\lambda_{n}^{2}}\{A(\nabla)\times\nabla\}\psi_{n}(x)$$
$$= \lambda_{n}A(\nabla)\{e^{-\mu\lambda_{n}^{2}}\psi_{n}(x)\} + \{A(\nabla)\times\nabla\}\{e^{-\mu\lambda_{n}^{2}}\psi_{n}(x)\}.$$

It is obvious that random $u_e^{n,\mu}$ is the solution of static incompressible Euler equations (1.5) (1.6) with periodic boundary condition (1.22) since E_n is linear space and $e^{-\mu\lambda_n^2}\psi_n(x) \in E_n$. On the other hand double limit $\lim_{(\nu,1/t)\to(0,0)} u_{ns}^n(t,x)$ does not exist. These phenomena reveal randomness and turbulence of incompressible fluids.

Now we give another typification to show that Prandtl layer dose not appear.

For any bounded domain $D \subset \mathbb{R}^3$ and $\partial D \in C^{\infty}$, we consider equation (1.15) with Dirichlet boundary condition

(1.30)
$$\psi(x)\big|_{x\in\partial D} = 0.$$

The equation (1.15) with Dirichlet boundary condition (1.30) is so-called Dirichlet eigenvalue problem. The eigenvalue λ of problem (1.15) (1.30) is so-called Dirichlet eigenvalue. By spectrum theory of compact operator $(-\Delta)^{-1}$, there exist countable infinite Dirichlet eigenvalues

$$0 < \lambda_1 \leq \cdots \leq \lambda_n \leq \cdots, n = 1, 2, \cdots$$

Each Dirichlet eigenvalue is real, and may be geometrically simple or finite. We define linear space F_n as follows

(1.31) $F_n = \{\psi_n(x) | \text{any eigenfunction } \psi_n \text{ corresponding to eigenvalue } \lambda_n \}.$

By the regular theory of elliptic equations, we have $\psi_n \in H^{\infty}(D)$ for any $\psi_n \in F_n$ and $n = 1, 2, \cdots$

As corollary of Theorem 1.2, we derive

Theorem 1.6 (Global Turbulent Dirichlet Solution). Let $A(\nabla)$ be any linear operator which is defined in Definition 1.1. Provided that λ_n is eigenvalue and ψ_n is eigenfunction of problem (1.15) (1.30), $\psi_n \in F_n$, $\phi_n(x) = \lambda_n \psi_n(x)$ $(n = 1, 2, \cdots)$. Let

(1.32)
$$u_e^{D,n}(x) = A(\nabla)\phi_n(x) + \{A(\nabla) \times \nabla\}\psi_n(x) \\ = \lambda_n A(\nabla)\psi_n(x) + \{A(\nabla) \times \nabla\}\psi_n(x),$$

then $u = u_e^{D,n}(x)$ $(n = 1, 2, \dots)$ is the solution of static incompressible Euler equations (1.5) (1.6) with boundary condition

(1.33)
$$\sigma \cdot u(x)\big|_{x \in \partial D} = \left\{\lambda_n \sigma \cdot A(\nabla)\psi_n(x) + \sigma \cdot \left\{A(\nabla) \times \nabla\right\}\psi_n(x)\right\}\big|_{x \in \partial D},$$

where σ is the outer normal of ∂D .

Moreover let

(1.34)
$$u_{ns}^{D,n}(t,x) = e^{-\nu\lambda_n^2 t} A(\nabla)\phi_n(x) + e^{-\nu\lambda_n^2 t} \{A(\nabla) \times \nabla\}\psi_n(x)$$
$$= \lambda_n e^{-\nu\lambda_n^2 t} A(\nabla)\psi_n(x) + e^{-\nu\lambda_n^2 t} \{A(\nabla) \times \nabla\}\psi_n(x),$$

then $u = u_{ns}^{D,n}(t,x)$ $(n = 1, 2, \dots)$ is the solution of incompressible Navier-Stokes equations (1.1) (1.2) with Dirichlet boundary condition

(1.35)
$$u(t,x)\big|_{x\in\partial D} = \{\lambda_n e^{-\nu\lambda_n^2 t} A(\nabla)\psi_n(x) + e^{-\nu\lambda_n^2 t} \{A(\nabla)\times\nabla\}\psi_n(x)\}\big|_{x\in\partial D}$$

and initial data

(1.36)
$$u(t,x)\big|_{t=0} = u_e^{D,n}(x).$$

Indeed the solution of Navier-Stokes equations (1.1) (1.2) with Dirichlet boundary condition (1.35) and initial data (1.36) converges to the solution of Euler equations (1.5) (1.6) with boundary condition (1.33) as $\nu \to 0$, i.e.

$$\lim_{\nu \to 0} u_{ns}^{D,n} = u_e^{D,n}.$$

Here Prandtl layer dose not appear.

Similarly we have path limit

(1.37)
$$u_{e}^{D,n,\mu}(x) = \lim_{(\nu,t)\in\Gamma_{\mu},(\nu,1/t)\to(0,0)} u_{ns}^{D,n}(t,x)$$
$$=\lambda_{n}e^{-\mu\lambda_{n}^{2}}A(\nabla)\psi_{n}(x) + e^{-\mu\lambda_{n}^{2}}\{A(\nabla)\times\nabla\}\psi_{n}(x)$$
$$=\lambda_{n}A(\nabla)\{e^{-\mu\lambda_{n}^{2}}\psi_{n}(x)\} + \{A(\nabla)\times\nabla\}\{e^{-\mu\lambda_{n}^{2}}\psi_{n}(x)\}.$$

Since F_n is linear space and $e^{-\mu\lambda_n^2}\psi_n(x) \in F_n$, it is obvious that random $u_e^{D,n,\mu}$ is the solution of static incompressible Euler equations (1.5) (1.6) with boundary condition

(1.38)
$$\begin{aligned} \sigma \cdot u(x)\big|_{x \in \partial D} \\ = \big(\lambda_n \sigma \cdot A(\nabla) \{e^{-\mu \lambda_n^2} \psi_n(x)\} + \sigma \cdot \{A(\nabla) \times \nabla\} \{e^{-\mu \lambda_n^2} \psi_n(x)\}\big)\big|_{x \in \partial D} \end{aligned}$$

On the other hand double limit $\lim_{(\nu,1/t)\to(0,0)} u_{ns}^{D,n}(t,x)$ does not exist. These phenomena also reveal randomness and turbulence of incompressible fluids.

Assume that $A(\nabla) = e_3 \times \nabla$, $\phi = 0$ and $\psi(t, x) = \psi(t, r, x_3)$ $(r^2 = x_1^2 + x_2^2)$ in (1.13). Inserting this special (1.13) into Navier-Stokes equations (1.1) (1.2), the equations reduce to axially symmetric Navier-Stokes equations without swirl. The global well-posedness of axially symmetric Navier-Stokes equations without swirl was first established by Ladyzhenskaya [6] and Ukhovskii-Yudovich [10]. Leonardi et al. [7] give a simpler proof to this result by using a priori weight estimates. Recently a different proof of the global well-posed result was given in [5].

Provided that $A(\nabla) = e_3 \times \nabla = (-\partial_2, \partial_1, 0)$, $\phi(t, x) = \phi(x)$ and $\psi(t, x) = 0$ in (1.13). Inserting this special (1.13) and (1.14) into static Euler equation (1.7), we derive

Theorem 1.7 (Non-unique Solutions of Periodic Pipeline Euler Flow). For any function $\phi(\xi, \eta) \in C^{\infty}(\mathbb{R} \times (L_1, L_2))$ with

(1.39)
$$\phi(\xi + 2\pi, \eta) = \phi(\xi, \eta), \quad \forall \xi \in \mathbb{R}, \ \eta \in (L_1, L_2),$$

6

we define

(1.40)
$$u_e^{pp}(x) = \left(-\partial_2\phi(x_1 + x_2, x_3), \partial_1\phi(x_1 + x_2, x_3), 0\right),$$

then $u = u_e^{pp}(x)$ is the solution of static incompressible Euler equations (1.5) (1.6) with boundary condition

$$u(x_1 + 2\pi, x_2, x_3) = u(x_1, x_2, x_3), \ \forall (x_1, x_2) \in \mathbb{R}^2, \ x_3 \in (L_1, L_2),$$

(1.41)
$$u(x_1, x_2 + 2\pi, x_3) = u(x_1, x_2, x_3), \ \forall (x_1, x_2) \in \mathbb{R}^2, \ x_3 \in (L_1, L_2), u_3(x_1, x_2, x_3)\big|_{x_3 = L_1} = u_3(x_1, x_2, x_3)\big|_{x_3 = L_2} = 0, \ \forall (x_1, x_2) \in \mathbb{R}^2.$$

Here $\omega \perp u$ and

(1.42)
$$\begin{aligned} \omega &= curl \ u = \nabla \times u \\ &= \left(-\partial_1 \partial_3 \phi(x_1 + x_2, x_3), -\partial_2 \partial_3 \phi(x_1 + x_2, x_3), \{\partial_1^2 + \partial_2^2\} \phi(x_1 + x_2, x_3) \right). \end{aligned}$$

Indeed the solution of static incompressible Euler equations (1.5) (1.6) with boundary condition (1.41) is non-unique.

This Theorem 1.7 is proved in Section 2.

Assume that $A(\nabla) = e_3 \times \nabla = (-\partial_2, \partial_1, 0)$ and $\psi(t, x) = 0$ in (1.13). Inserting this special (1.13) and (1.14) into Navier-Stokes equations (1.3), we obtain

Theorem 1.8 (Global Solution of Periodic Pipeline NS Flow). Provided regular function $\phi(t, \xi, \eta)$ satisfies the following equations

(1.43)
$$\begin{aligned} \phi_t - \nu (2\partial_{\xi}^2 + \partial_{\eta}^2)\phi &= 0, \\ \phi(t, \xi + 2\pi, \eta) &= \phi(t, \xi, \eta), \quad \forall t \ge 0, \ \xi \in \mathbb{R}, \ \eta \in (L_1, L_2), \\ \phi(t, \xi, \eta)\big|_{\eta = L_1} &= \phi(t, \xi, \eta)\big|_{\eta = L_2} = 0, \quad \forall t \ge 0, \ \xi \in \mathbb{R}, \\ \phi(t, \xi, \eta)\big|_{t=0} &= \phi_0(\xi, \eta). \end{aligned}$$

Let

(1.44)
$$u_{ns}^{pp}(t,x) = \left(-\partial_2\phi(t,x_1+x_2,x_3),\partial_1\phi(t,x_1+x_2,x_3),0\right),$$

then $u = u_{ns}^{pp}$ is the solution of incompressible Navier-Stokes equations (1.1) (1.2) with boundary condition

$$\begin{aligned} u(t, x_1 + 2\pi, x_2, x_3) &= u(t, x_1, x_2, x_3), \quad \forall t \ge 0, \ (x_1, x_2) \in \mathbb{R}^2, \ x_3 \in (L_1, L_2), \\ (1.45) \quad u(t, x_1, x_2 + 2\pi, x_3) &= u(t, x_1, x_2, x_3), \quad \forall t \ge 0, \ (x_1, x_2) \in \mathbb{R}^2, \ x_3 \in (L_1, L_2), \\ u(t, x_1, x_2, x_3)\big|_{x_3 = L_1} &= u(t, x_1, x_2, x_3)\big|_{x_3 = L_2} = 0, \quad \forall t \ge 0, \ (x_1, x_2) \in \mathbb{R}^2, \end{aligned}$$

and initial data

(1.46)
$$u(t,x)\big|_{t=0} = \big(-\partial_2\phi_0(x_1+x_2,x_3),\partial_1\phi_0(x_1+x_2,x_3),0\big).$$

This Theorem 1.8 is easy to prove by using Theorem 1.7. As a consequence of Theorem 1.8, we get

Corollary 1.9 (Typification without Prandtl Layer). Under the condition of Theorem 1.8, we derive that

$$\lim_{\nu \to 0} u_{ns}^{pp}(t,x) = \left(-\partial_2 \phi_0(x_1 + x_2, x_3), \partial_1 \phi_0(x_1 + x_2, x_3), 0 \right) = u_{e0}^{pp}(x),$$

and $u = u_{e0}^{pp}(x)$ is the solution of static incompressible Euler equations (1.5) (1.6) with boundary condition (1.41).

Here Prandtl layer dose not appear.

Let us give an interesting exemplification of Theorem 1.8 by taking

(1.47)
$$\phi_0(\xi,\eta) = \sum_{j\in J} \sum_{k\in K} \{\alpha_{jk}\sin(j\xi) + \beta_{jk}\cos(j\xi)\}\sin\left(\pi k(\eta - L_1)/(L_2 - L_1)\right)$$

where L_1 , L_2 , α_{jk} and β_{jk} are constants, J and K are any finite nonempty subsets of $\{1, 2, \dots\}$. For the problem (1.43), there exists a unique solution

(1.48)
$$\phi(t,\xi,\eta) = \sum_{j\in J} \sum_{k\in K} e^{-t\nu\{2j^2 + \pi^2k^2/(L_2 - L_1)^2\}} \{\alpha_{jk}\sin(j\xi) + \beta_{jk}\cos(j\xi)\}\sin\left(\pi k(\eta - L_1)/(L_2 - L_1)\right).$$

Inserting (1.48) into (1.44), we have path limit

(1.49)
$$u_{ef}^{pp\mu}(x) = \lim_{(\nu,t)\in\Gamma_{\mu},(\nu,1/t)\to(0,0)} u_{ns}^{pp}(t,x)$$
$$= \sum_{j\in J} \sum_{k\in K} e^{-\mu\{2j^2+\pi^2k^2/(L_2-L_1)^2\}}$$
$$\left(-\partial_2\{\alpha_{jk}\sin\left(j(x_1+x_2)\right)+\beta_{jk}\cos\left(j(x_1+x_2)\right)\}\sin\frac{\pi k(x_3-L_1)}{L_2-L_1},\right.$$
$$\partial_1\{\alpha_{jk}\sin\left(j(x_1+x_2)\right)+\beta_{jk}\cos\left(j(x_1+x_2)\right)\}\sin\frac{\pi k(x_3-L_1)}{L_2-L_1},0\right).$$

By Theorem 1.7, random $u = u_{ef}^{pp\mu}$ is the solution of static incompressible Euler equations (1.5) (1.6) with boundary condition (1.41). On the other hand double limit $\lim_{(\nu,1/t)\to(0,0)} u_{ns}^{pp}(t,x)$ does not exist. These phenomena reveal randomness and turbulence of incompressible fluids again.

Besides the non-uniqueness of solutions of static Euler equations in Corollary 1.4 and Theorem 1.7, the following non-unique theorem is well known (see [3, 4] for instance)

Theorem 1.10 (Non-unique Solutions of 2d Euler Equations). Provided that $r^2 = x_1^2 + x_2^2$, disc $B_R = \{(x_1, x_2) | r < R\}$, function $\phi(x) = \phi(r)$. For any radial function $\phi(r) \in C_c^{\infty}(B_R)$, let

(1.50)
$$u_e^{2d}(x_1, x_2) = \left(-\partial_2 \phi(r), \partial_1 \phi(r)\right),$$

then $u = u_e^{2d}(x_1, x_2)$ is the solution of two dimensional static incompressible Euler equations with boundary condition

(1.51)
$$\sigma \cdot u(x_1, x_2)\big|_{(x_1, x_2) \in \partial B_R} = 0,$$

where σ is the outer normal of ∂B_R .

Indeed the solution of two dimensional static incompressible Euler equations with boundary condition (1.51) is non-unique too.

8

2. Proof of Theorem 1.7

Provided that $A(\nabla) = e_3 \times \nabla = (-\partial_2, \partial_1, 0)$, $\phi(t, x) = \phi(x)$ and $\psi(t, x) = 0$ in (1.13). Theorem 1.7 can be proved by inserting this special (1.13) (1.14) into static Euler

equation (1.7).

Proof. Inserting $a_1(\nabla) = -\partial_2$, $a_2(\nabla) = \partial_1$, $a_3(\nabla) = 0$ and $\psi = 0$ into (1.13) (1.14), we have

(2.1)
$$u = (-\partial_2 \phi, \partial_1 \phi, 0),$$
$$\omega = \nabla \times u = (-\partial_1 \partial_3 \phi, -\partial_2 \partial_3 \phi, (\partial_1^2 + \partial_2^2) \phi).$$

It is well-known that the equation (1.7) is equivalent to

(2.2)
$$(u \cdot \nabla)\omega - (\omega \cdot \nabla)u = \sum_{j=1}^{3} \partial_j (u^j \omega - \omega^j u) = 0.$$

To verify that $u = u_e$ is the solution of Euler equations (1.5) (1.6), it is necessary to calculate $u^j \omega^k - \omega^j u^k$. By observation

(2.3)
$$\begin{aligned} u^{j}\omega^{k} - \omega^{j}u^{k} &= 0, \quad j = k, \\ u^{j}\omega^{k} - \omega^{j}u^{k} &= -(u^{k}\omega^{j} - \omega^{k}u^{j}) \end{aligned}$$

only three terms $u^2\omega^1 - \omega^2 u^1$, $u^3\omega^1 - \omega^3 u^1$ and $u^3\omega^2 - \omega^3 u^2$ have to calculate. Using (2.1), we derive

(2.4)
$$u^2\omega^1 - \omega^2 u^1 = -\partial_1 \phi \partial_1 \partial_3 \phi - \partial_2 \phi \partial_2 \partial_3 \phi,$$

(2.5)
$$u^3\omega^1 - \omega^3 u^1 = \partial_2 \phi (\partial_1^2 \phi + \partial_2^2 \phi),$$

(2.6)
$$u^3\omega^2 - \omega^3 u^2 = -\partial_1 \phi (\partial_1^2 \phi + \partial_2^2 \phi).$$

Therefore we have

$$(2.7) \qquad \begin{aligned} (u \cdot \nabla)\omega^{1} - (\omega \cdot \nabla)u^{1} &= \partial_{k}(u^{k}\omega^{1} - \omega^{k}u^{1}) \\ &= \partial_{2}(u^{2}\omega^{1} - \omega^{2}u^{1}) + \partial_{3}(u^{3}\omega^{1} - \omega^{3}u^{1}) \\ &= \partial_{2}(-\partial_{1}\phi\partial_{1}\partial_{3}\phi - \partial_{2}\phi\partial_{2}\partial_{3}\phi) + \partial_{3}\{\partial_{2}\phi(\partial_{1}^{2}\phi + \partial_{2}^{2}\phi)\} \\ &= -\partial_{1}\partial_{2}\phi\partial_{1}\partial_{3}\phi - \partial_{1}\phi\partial_{1}\partial_{2}\partial_{3}\phi + \partial_{2}\partial_{3}\phi\partial_{1}\partial_{1}\phi + \partial_{2}\phi\partial_{1}\partial_{1}\partial_{3}\phi, \end{aligned}$$

(2.8)
$$(u \cdot \nabla)\omega^{2} - (\omega \cdot \nabla)u^{2} = \partial_{k}(u^{k}\omega^{2} - \omega^{k}u^{2})$$
$$= -\partial_{1}(u^{2}\omega^{1} - \omega^{2}u^{1}) + \partial_{3}(u^{3}\omega^{2} - \omega^{3}u^{2})$$
$$= \partial_{1}(\partial_{1}\phi\partial_{1}\partial_{3}\phi + \partial_{2}\phi\partial_{2}\partial_{3}\phi) + \partial_{3}\{-\partial_{1}\phi(\partial_{1}^{2}\phi + \partial_{2}^{2}\phi)\}$$
$$= \partial_{1}\partial_{2}\phi\partial_{2}\partial_{3}\phi + \partial_{2}\phi\partial_{1}\partial_{2}\partial_{3}\phi - \partial_{1}\partial_{3}\phi\partial_{2}\partial_{2}\phi - \partial_{1}\phi\partial_{2}\partial_{2}\partial_{3}\phi,$$

(2.9)

$$(u \cdot \nabla)\omega^{3} - (\omega \cdot \nabla)u^{3} = \partial_{k}(u^{k}\omega^{3} - \omega^{k}u^{3})$$

$$= -\partial_{1}(u^{3}\omega^{1} - \omega^{3}u^{1}) - \partial_{2}(u^{3}\omega^{2} - \omega^{3}u^{2})$$

$$= -\partial_{1}\{\partial_{2}\phi(\partial_{1}^{2}\phi + \partial_{2}^{2}\phi)\} + \partial_{2}\{\partial_{1}\phi(\partial_{1}^{2}\phi + \partial_{2}^{2}\phi)\}$$

$$= \partial_{1}\phi\partial_{2}(\partial_{1}^{2}\phi + \partial_{2}^{2}\phi) - \partial_{2}\phi\partial_{1}(\partial_{1}^{2}\phi + \partial_{2}^{2}\phi).$$

For any smooth function $f(\xi, \eta)$, let $\phi(x_1, x_2, x_3) = f(x_1 + x_2, x_3)$. Then

(2.10)
$$\partial_1 \phi = \partial_{\xi} f, \ \partial_2 \phi = \partial_{\xi} f, \ \partial_3 \phi = \partial_{\eta} f$$

Applying (2.10) in (2.7) (2.8) (2.9), we obtain that

(2.11)
$$(u \cdot \nabla)\omega^1 - (\omega \cdot \nabla)u^1 = (u \cdot \nabla)\omega^2 - (\omega \cdot \nabla)u^2 = (u \cdot \nabla)\omega^3 - (\omega \cdot \nabla)u^3 = 0.$$

Equation (2.11) means that $u = u_e^{pp}(x)$ is the solution of static incompressible Euler equations (1.5) (1.6) with boundary condition (1.41).

Acknowledgments

This work is supported by National Natural Science Foundation of China–NSF, Grant No.11971068 and No.11971077.

References

- P. Constantin. Some open problems and research dierections in the mathematical study of fluid dynamics. In Mathematics Unlimited-2001 and Beyond, Springer Verlag, Berlin, 2002, 353–360.
- [2] Y. Guo and S. Iyer. Validity of steady Prandtl layer expansions. Comm. Pure Appl. Math. 76 (11), 2023, 3150–3232.
- [3] Yongqian Han. Symplectic Symmetry and Radial Symmetry Either Persistence or Breaking of Incompressible Fluid. ArXiv: 2305.13737v3.
- [4] Yongqian Han. Symplectic Representation and Turbulent Global Solutions of Incompressible Navier-Stokes Equations in ℝ³. ArXiv: 2311.18308v2.
- [5] Yongqian Han. Cylindrical Symplectic Representation and Global Regular Solution of Incompressible Navier-Stokes Equations in R³. ArXiv: 2311.18315v1.
- [6] O. A. Ladyzhenskaya. Unique global solvability of the three-dimensional Cauchy problem for the Navier-Stokes equations in the presence of axial symmetry. Zap. Naucn. Sem. Leningrad. Otdel. Math. Inst. Steklov. (LOMI) 7, 1968, 155–177 (Russian).
- [7] S. Leonardi, J. Malek, J. Necas and M. Porkorny. On axially symmetric flows in ℝ³. Z. Anal. Anwendungen 18, 1999, 639–649.
- [8] K. Li and B. Wang. Blowup criterion for Navier-Stokes equation in critical Besov space with spatial dimensions $d \ge 4$. Ann. Inst. Henri Poincare (C) Anal. Nonlinear 36(6), 2019, 1679–1707.
- R. Temam. Navier-Stokes Equations. Theory and numerical analysis. (Reprint of the 1984 edition). AMS Chelsea Publishing, Providence, RI, 2001.
- [10] M. R. Ukhovskii and V. I. Yudovich. Axially symmetric flows of ideal and viscous fluids filling the whole space. J. Appl. Math. Mech. 32, 1968, 52–61.
- [11] Qi S. Zhang. A review of results on axially symmetric Navier-Stokes equations, with addendum by X. Pan and Q. S. Zhang. ArXiv: 2102.04905v2.

INSTITUTE OF APPLIED PHYSICS AND COMPUTATIONAL MATHEMATICS, BEIJING 100088, CHINA

NATIONAL KEY LABORATORY OF COMPUTATIONAL PHYSICS, BEIJING 100088, CHINA *Email address:* han_yongqian@iapcm.ac.cn