ON THE MOUTARD TRANSFORMATION SINGULARITY FOR THE DAVEY-STEWARTSON II EQUATION

YI C. HUANG

ABSTRACT. We indicate explicitly how to obtain the Moutard transformation singularity for the Davey–Stewartson II equation with two indeterminancy points. We also justify an arbitrary order "mass drop" phenomenon via this singularity formation scheme.

1. INTRODUCTION: THE DAVEY-STEWARTSON II EQUATION

In this Letter, following Taĭmanov [Taĭ21, Taĭ24], we shall consider the Moutard transformation solutions of the following Davey–Stewartson II equation [DS74]:

$$U_t = i \left(U_{zz} + U_{\overline{zz}} + 2(V + \overline{V})U \right), \quad V_{\overline{z}} = (|U|^2)_z, \tag{1.1}$$

which is the compatibility condition for the linear problems

$$\mathcal{D}\Psi = 0, \quad \partial_t \Psi = \mathcal{A}\Psi,$$

where

$$\mathcal{D} = \left(\begin{array}{cc} 0 & \partial \\ -\overline{\partial} & 0 \end{array}\right) + \left(\begin{array}{cc} U & 0 \\ 0 & \overline{U} \end{array}\right)$$

is a two-dimensional Dirac operator with a complex-valued potential U with

$$\partial = \frac{1}{2} \left(\frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right), \quad \overline{\partial} = \frac{1}{2} \left(\frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right),$$

the time-dependent generator \mathcal{A} is given by

$$\mathcal{A} = i \begin{pmatrix} -\partial^2 - V & \overline{U} \overline{\partial} - \overline{U}_{\overline{z}} \\ U \partial - U_z & \overline{\partial}^2 + \overline{V} \end{pmatrix},$$

and the solution Ψ is taken as the following matrix form

$$\Psi = \left(\begin{array}{cc} \psi_1 & -\overline{\psi}_2 \\ \psi_2 & \overline{\psi}_1 \end{array}\right).$$

Note that if (U_0, V_0) solves (1.1), then (\overline{U}_0, V_0) solves

$$U_t = -i \left(U_{zz} + U_{\overline{z}\overline{z}} + 2(V + \overline{V})U \right), \quad V_{\overline{z}} = (|U|^2)_z.$$

Hence, (1.1) is also the compatibility condition of the "conjugate" problems

$$\mathcal{D}^{\vee}\Phi=0, \ \Phi_t=\mathcal{A}^{\vee}\Phi,$$

Date: February 18, 2025.

²⁰²⁰ Mathematics Subject Classification. Primary: 35Q53; Secondary: 37K35, 35B44, 49Q05.

Key words and phrases. Davey–Stewartson II equation, Dirac operators, Moutard transformation, singularity formation, spinorial representations, surface deformation.

Research of the author is partially supported by the National NSF grant of China (no. 11801274), the JSPS Invitational Fellowship for Research in Japan (no. S24040), and the Open Projects from Yunnan Normal University (no. YNNUMA2403) and Soochow University (no. SDGC2418). This text is written during a visit to Novosibirsk and the author thanks sincerely Iskander A. Taĭmanov for enlightening discussions.

where

$$\mathcal{D}^{\vee} = \left(\begin{array}{cc} 0 & \partial \\ -\overline{\partial} & 0 \end{array}\right) + \left(\begin{array}{cc} \overline{U} & 0 \\ 0 & U \end{array}\right),$$

and

$$\mathcal{A}^{\vee} = -i \left(\begin{array}{cc} -\partial^2 - V & U \overline{\partial} - U_{\overline{z}} \\ \overline{U} \partial - \overline{U}_z & \overline{\partial}^2 + \overline{V} \end{array} \right).$$

Analogously, the solution Φ is taken as the following matrix form

$$\Phi = \left(\begin{array}{cc} \varphi_1 & -\overline{\varphi}_2 \\ \varphi_2 & \overline{\varphi}_1 \end{array} \right).$$

If U = 0, (ψ_1, φ_1) is pair of holomorphic functions, while (ψ_2, φ_2) anti-holomorphic. When $U = \overline{U}$, the "conjugate" problems are irrelevant and one regards $\Psi = \Phi$. The pair (Ψ, Φ) serves as a Davey–Stewartson II deformation of surfaces in \mathbb{R}^4 , see e.g. [Taĭ22].

2. MOUTARD SINGULARITY ON TWO POINTS OR MULTIPLE ZEROES

Using the Moutard transformation of the solution U = V = 0 with spinors

$$\Psi = \left(\begin{array}{cc} g' & 0\\ 0 & g' \end{array}\right), \ \Phi = \left(\begin{array}{cc} f' & i\\ i & f' \end{array}\right),$$

the following two-parameter family of solutions were found in [Taĭ24, Theorem 4.1].

Theorem 2.1 (Taĭmanov, 2024). Let f(z,t) and g(z,t) be two functions which are holomorphic in z and satisfy the equations

$$\frac{\partial f}{\partial t} = i \frac{\partial^2 f}{\partial z^2}, \quad \frac{\partial g}{\partial t} = -i \frac{\partial^2 g}{\partial z^2}.$$
(2.1)

Let h be a function which is holomorphic in z and satisfies the relations

$$h' = f'g', \quad ih_t = g''f' - g'f''.$$

Then

$$U = i\overline{g'}\frac{f'g - h}{|g|^2 + |h|^2}$$
(2.2)

satisfies the Davey–Stewartson II equation (1.1).

It is noteworthy to point out that in the singularity formation scheme of Moutard transformation, the singularity points (z_0, t_0) are encoded in the zero set of $|g|^2 + |h|^2$.

Remark 2.2. Let us first write down the example given in [Taĭ24]. Take

$$f(z,t) = z^2 + 2it, \quad g(z,t) = z^3 - 6itz,$$

and choose

$$h = \frac{3}{2}z^4 - 6itz^2 + 6t^2.$$

We derive the following solution

$$U = \frac{i(3\overline{z}^{2} + 6it)(\frac{1}{2}z^{4} - 6itz^{2} - 6t^{2})}{|z^{3} - 6itz|^{2} + |\frac{3}{2}z^{4} - 6itz^{2} + 6t^{2}|^{2}}$$
$$= O(|z|^{-2}), \qquad |z| \to +\infty.$$

For t = 0 it has the indeterminacy type singularity $\sim \frac{3}{2}i\frac{z^2}{|z|^2}$ as $|z| \to 0$.

The first observation of this Letter can be summarized as follows:

Observation 2.3. In above example, we modify the data set (f,g) in the following way

$$f(z,t) = (z^2 + 2it) - z, \quad g(z,t) = (z^3 - 6itz) - z.$$

The natural observation (or rather, motivation) here is that the lower order modification terms is set to solve the *linear* equations for f and g. Thus it remains to compute

$$f'(z,t) = 2z - 1, \quad g'(z,t) = 3z^2 - (1 + 6it)$$

and

$$h'(z,t) = f'(z,t)g'(z,t) = 6z^3 - 3z^2 - 2(1+6it)z + (1+6it).$$

We choose

$$h(z,t) = \frac{3}{2}z^4 - z^3 - (1+6it)z^2 + (1+6it)z - i(t^2 + 4it^3) - \frac{1}{2}.$$

Here, by adding the temporal term $-i(t^2 + 4it^3)$ we meet the constraint $ih_t = g''f' - g'f''$, and by adding the constant $-\frac{1}{2}$, we see that $h|_{t=0}$ vanishes at $z = \pm 1$. By Theorem 2.1, the data triple (f, g, h) yields the following solution for the Davey–Stewartson II equation

$$U = i\overline{g'} \frac{f'g - h}{|g|^2 + |h|^2}$$

= $\frac{i(3\overline{z}^2 - (1 - 6it))(\frac{1}{2}z^4 - (1 + 6it)z^2 + i(t^2 + 4it^3) + \frac{1}{2})}{|(z^3 - 6itz) - z|^2 + |\frac{3}{2}z^4 - z^3 - (1 + 6it)z^2 + (1 + 6it)z - i(t^2 + 4it^3) - \frac{1}{2}|^2}$

Note that

$$h|_{t=0} = \frac{3}{2}z^4 - z^3 - z^2 + z - \frac{1}{2} = (z^2 - 1)\left(\frac{3}{2}z^2 - z + \frac{1}{2}\right),$$
$$g|_{t=0} = z(z^2 - 1), \quad g'|_{z=\pm 1} \neq 0,$$

and

$$(f'g-h)|_{t=0} = \frac{(z^2-1)^2}{2}$$

Thus the solution U at t = 0 has a two-point indeterminancy type singularity

$$U|_{t=0} \sim C_{\pm 1} \frac{(z^2 - 1)^2}{|z^2 - 1|^2}, \quad z \to \pm 1,$$

The involved implicit constants $C_{\pm 1}$ can be computed directly, and we omit this step.

Remark 2.4. For higher order polynomials one can construct in the similar manner the multiple-point indeterminancy singularity. We choose not to dwell on that issue.

Note that the solution in Observation 2.3 has mass $||U(\cdot,t)||_2^2 \equiv 4\pi$, except at the singular time t = 0, it has a "mass drop" with magnitude 2π (with 2 being the number of indeterminancy points). The second observation of this Letter can be summarized as follows:

Observation 2.5. According to (2.1), we construct for $2 \le m \in \mathbb{N}$ and $1 \le n \in \mathbb{N}$ that

$$f_m = z^m + im(m-1)tz^{m-2} + \cdots,$$

$$g_n = z^n - in(n-1)tz^{n-2} - \cdots,$$

and for convenience, with the normalization $f_m(0,0) = g_n(0,0) = 0$. For example, we have

$$f_2(z,t) = z^2 + 2it,$$

$$f_3(z,t) = z^3 + 6itz,$$

$$f_4(z,t) = z^4 + 12itz^2 - 12t^2,$$

$$f_5(z,t) = z^5 + 20itz^3 - 60t^2z$$

and

$$g_{1}(z,t) = z,$$

$$g_{2}(z,t) = z^{2} - 2it,$$

$$g_{3}(z,t) = z^{3} - 6itz,$$

$$g_{4}(z,t) = z^{4} - 12itz^{2} - 12t^{2},$$

$$g_{5}(z,t) = z^{5} - 20itz^{3} - 60t^{2}z.$$

In constructing the function *h* we also assume h(0,0) = 0, thus the solution *U* in (2.2) with data (f_m, g_n) has mass $||U(\cdot, t)||_2^2 \equiv (m+n-1)\pi$, except at the singular time t = 0, it has a "mass drop" with magnitude $n\pi$ (with *n* being the order of *g*) to $(m-1)\pi$. The normalization on *h* can not be fulfilled when m = 1 unless *U* is the trivial zero solution.

COMPLIANCE WITH ETHICAL STANDARDS

Conflict of interest The author has no known competing financial interests or personal relationships that could have appeared to influence this reported work.

Availability of data and material Not applicable.

REFERENCES

- [DS74] A. Davey and K. Stewartson. On three-dimensional packets of surface waves. Proc. Roy. Soc. London Ser. A, 338:101–110, 1974.
- [Taĭ21] I. A. Taĭmanov. The Moutard transformation for the Davey-Stewartson II equation and its geometrical meaning. *Mat. Zametki*, 110(5):751–765, 2021.
- [Taĭ22] I. A. Taĭmanov. Surfaces via spinors and soliton equations. In Proc. Int. Cong. Math, volume 4, pages 2638–2654, 2022.
- [Taĭ24] I. A. Taĭmanov. On a formation of singularities of solutions to soliton equations represented by *L*,*A*,*B*-triples. *Acta Math. Sin. (Engl. Ser.)*, 40(1):406–416, 2024.

SCHOOL OF MATHEMATICAL SCIENCES, NANJING NORMAL UNIVERSITY, NANJING 210023, PEO-PLE'S REPUBLIC OF CHINA

E-mail: Yi.Huang.Analysis@gmail.com Homepage: https://orcid.org/0000-0002-1297-7674