

Multipoint conformal integrals in D dimensions. Part I

Bipartite Mellin-Barnes representation and reconstruction

Konstantin Alkalaev and Semyon Mandrygin

*I.E. Tamm Department of Theoretical Physics,
P.N. Lebedev Physical Institute, 119991 Moscow, Russia*

E-mail: alkalaev@lpi.ru, semyon.mandrygin@gmail.com

ABSTRACT: We propose a systematic approach to calculating n -point one-loop parametric conformal integrals in D dimensions which we call the reconstruction procedure. It relies on decomposing a conformal integral over basis functions which are generated from a set of master functions by acting with the cyclic group \mathbb{Z}_n . In order to identify the master functions we introduce a bipartite Mellin-Barnes representation by means of splitting a given conformal integral into two additive parts, one of which can be evaluated explicitly in terms of multivariate generalized hypergeometric series.

For the box and pentagon integrals (i.e. $n = 4, 5$) we show that a computable part of the bipartite representation contains all master functions. In particular, this allows us to evaluate the parametric pentagon integral as a sum of ten basis functions generated from two master functions by the cyclic group \mathbb{Z}_5 . The resulting expression can be tested in two ways. First, when one of propagator powers is set to zero, the pentagon integral is reduced to the known box integral, which is also rederived through the reconstruction procedure. Second, going to the non-parametric case, we reproduce the known expression for the pentagon integral given in terms of logarithms derived earlier within the geometric approach to calculating conformal integrals.

We conclude by considering the hexagon integral ($n = 6$) for which we show that those basis functions which follow from the computable part of the bipartite representation are not enough and more basis functions are required. In the second part of our project we will describe a method of constructing a complete set of master/basis functions in the n -point case.

Contents

1	Introduction	2
2	Multipoint parametric conformal integrals	4
2.1	Review of the Symanzik approach	4
2.2	Bipartite Mellin-Barnes representation	7
3	Constraints from the permutation invariance	12
3.1	Analytic continuation	13
3.2	Box integral	15
3.2.1	Kinematic group	16
3.2.2	Cyclic group	17
3.2.3	Consistency checks	19
3.2.4	Summary of the box integral reconstruction	21
3.3	Pentagon integral	22
3.3.1	Cyclic group	23
3.3.2	Reduction to the box integral	24
3.4	Hexagon integral	25
4	Reconstruction conjecture	27
5	Conclusion	30
A	Notation and conventions	31
B	Generalized hypergeometric functions	31
B.1	General definitions	32
B.2	Integral representations	34
B.3	Analytic continuation	35
C	One more Mellin-Barnes representation of the conformal integral	37
D	Pentagon integral	37
D.1	Basis functions	38
D.2	Non-parametric integral	40
E	Kinematic group extensions	43

1 Introduction

Conformal integrals are an interesting specific class of integrals that arise in various areas of quantum field theory [1]. E.g. they appear in the study of scattering multi-loop amplitudes, where the corresponding Feynman integrals respect a dual conformal invariance [2, 3] as well as in the Fishnet CFT models [4–9] (see [10, 11] for reviews). There is a number of methods of calculating conformal integrals among which the most fruitful are the geometric approach which treats these integrals as volumes of simplices [12–17], Yangian bootstrap [18–21], GKZ differential equations [22–24] as well as other approaches which combine various techniques [25–32]. Nonetheless, despite this recent remarkable progress in calculating conformal integrals, explicit closed-form expressions for general conformal integrals are not yet known.

On the other hand, a more traditional way to come across conformal integrals is to use the shadow formalism in CFT which represents correlation functions as particular integrals possessing conformal invariance by construction [33–39]. In particular, using the shadow formalism allows one to take advantage of various Feynman integral techniques for CFT calculations.¹ In this paper, we instead are inspired by CFT methods for dealing with conformal integrals. Namely, we develop the idea of asymptotic analysis of conformal integrals which is borrowed from the conformal block decomposition of correlation functions in CFT. Recall that the conformal blocks can be defined by doing OPEs between pairs of primary operators inside a correlation function. This results in choosing of a particular channel, i.e. one fixes an order and closeness of points which is known as the OPE limit. This eventually defines a domain of convergence of the conformal block in the coordinate space. Now when the conformal block is represented in terms of conformal integrals, the latter are calculated in the OPE limit to have correct asymptotic behaviour. Other asymptotics of the respective conformal integral describe the so-called shadow blocks [38, 39]. The conformal blocks in other coordinate domains, i.e. in other possible channels, can be obtained by means of analytic continuation formulas.

Relying on this relation between conformal blocks and conformal integrals we can formulate the first point of our approach to calculating n -point conformal integrals in D dimensions: one fixes a convenient coordinate domain where a conformal integral can be represented in terms of a multivariate power series and then analytically continued onto other domains. The second point is that there is no need to evaluate conformal integral entirely, but only partially. To this end, one decomposes a conformal integral into two additive parts which are given by particular Mellin-Barnes integrals. This is a *bipartite* representation. The first term can be evaluated explicitly in some coordinate domain, while the second one is quite complicated and has not been computed yet. Here, we put forward our main idea that the second term can be reconstructed from the first one by invoking permutation invariance of the full conformal integral. Note that such a partition of conformal integral in no way is invariant against transformations from the symmetric group \mathcal{S}_n . Thus, the third point of our approach is to introduce a subgroup of the symmetric group which is claimed to generate the

¹Conformal integrals are also instrumental in CFT on non-trivial manifolds [40–47].

full conformal integral by acting on a set of *master* functions. We call this calculation scheme a reconstruction.

The most intricate part of this approach is to identify both a generating group and master functions. It turns out that a generating group can be chosen as the cyclic group $\mathbb{Z}_n \subset \mathcal{S}_n$ though this is not the only choice. A generating group is required to act on master functions to generate a complete set of *basis* functions which sum up to the full conformal integral. It follows that the generating group acts on the basis functions by reshuffling them. To single out an independent set of master functions we claim that they are not related by cyclic permutations.² In other words, a set of basis functions decomposes into subsets each of which carries a representation of \mathbb{Z}_n .³

In the cases of the box and pentagon conformal integrals the corresponding master functions can be naturally identified by explicitly calculating a first term of the bipartite representation and then analytically continuing the resulting expression onto some coordinate domain around a specified point. This procedure yields the first term as a sum of generalized hypergeometric multivariate functions from which a number of functions can be selected as master functions. The cyclic group generates a set of basis functions which define the second part of the bipartite representation thereby giving the full conformal integral.

However, when considering the hexagon integral we find out that those basis functions which come from the first part of the bipartite representations are not enough to build the full conformal integral. Nonetheless, we put forward a conjecture how to build a complete set of basis functions which is the subject of our forthcoming paper [48]. Essentially, there we claim that the basis functions which we are able to identify explicitly from the bipartite representation have a number of systematic properties that will eventually allow us to build a desired complete set.

This paper is organized as follows. In section 2 we introduce (non)-parametric n -point conformal integrals and review the standard calculation method by Symanzik which is based on using the Mellin-Barnes integrals. Then, in section 2.2 we propose a bipartite Mellin-Barnes representation which splits the original conformal integral into two additive parts. Here, we explicitly calculate the first part in some coordinate domain and discuss the second one. In section 3 we describe the action of symmetric group and consider the analytic continuation to the other coordinate domain. Section 3.2 considers the parametric box integral which has been studied in detail in the literature and examines our reconstruction approach using this example. Summarizing our study of the box integral, in section 3.2.4 we outline a heuristic procedure of obtaining the full asymptotic expansions in the n -point case. Then, in section 3.3 we apply this reconstruction machinery to the 5-point one-loop conformal integral. Here, we find the parametric pentagon integral as a sum of ten basis functions and then perform a number of consistency checks. In section 3.4 we consider the hexagon integral and discover that the way of choosing master functions used in the box and pentagon cases

²In this paper, a cyclic permutation will be understood only as a cycle of maximum length n , which generates the cyclic group $\mathbb{Z}_n \subset \mathcal{S}_n$.

³In general, these are inequivalent representations of \mathbb{Z}_n .

is insufficient to construct a complete asymptotic expression because there should be more master functions. Finally, in section 4, based on the considered examples, we formulate and discuss a conjecture regarding a reconstruction of n -point one-loop conformal integral. In the concluding section 5 we summarize our results and elaborate on some perspectives for future work.

A few appendices contain numerous technical aspects of calculations. Appendix A collects our notation and conventions used throughout the paper. Appendix B describes various generalized hypergeometric multivariate functions, their integral representations and analytic continuation formulas. In appendix C we suggest one more Mellin-Barnes representation of the conformal integral which can be useful in practice, e.g. for finding the conformal integral using the computer algebra methods. Appendix D collects explicit expressions for the basis functions which define the parametric pentagon conformal integral. In particular, in appendix D.2 we examine our representation of the non-parametric (i.e. with unit propagator powers) pentagon integral. Appendix E discusses an extended kinematic group which is a particular subgroup in the symmetric group.

2 Multipoint parametric conformal integrals

The following integral over the D -dimensional Euclidean space \mathbb{R}^D

$$I_n^{\mathbf{a}}(\mathbf{x}) = \int_{\mathbb{R}^D} \frac{d^D x_0}{\pi^{\frac{D}{2}}} \prod_{i=1}^n X_{0i}^{-a_i}, \quad X_{ij} \equiv X_{i,j} = (x_i - x_j)^2, \quad x_i \in \mathbb{R}^D, \quad (2.1)$$

where $\mathbf{a} = \{a_1, a_2, \dots, a_n\}$, $\mathbf{x} = \{x_1, x_2, \dots, x_n\}$, is called the n -point one-loop conformal integral [1] provided that the propagator powers $a_i \in \mathbb{R}$ obey the constraint

$$\sum_{i=1}^n a_i = D, \quad (2.2)$$

which guarantees that $I_n^{\mathbf{a}}(\mathbf{x})$ transforms covariantly under $O(D+1, 1)$ conformal transformations (see e.g. [11] for details). The integrals with arbitrary a_i subjected to (2.2) are called *parametric*, while those with all $a_i = 1$ in $D = n$ dimensions are called *non-parametric*.

Since the paper of Symanzik [1], the conformal integrals are traditionally represented in terms of the Mellin-Barnes integrals. In this section we first recall the Symanzik approach and then introduce a bipartite Mellin-Barnes representation.

2.1 Review of the Symanzik approach

In order to calculate the conformal integral (2.1) one can use the Schwinger parametrization

$$\frac{1}{X_{0i}^{a_i}} = \frac{1}{\Gamma(a_i)} \int_0^{+\infty} \frac{d\lambda_i}{\lambda_i} \lambda_i^{a_i} \exp(-\lambda_i X_{0i}), \quad a_i > 0. \quad (2.3)$$




Figure 1. The parametric conformal integral $I_n^{\mathbf{a}}(\mathbf{x})$ can be depicted as an n -valent vertex. The i -th leg denotes the propagator $X_{0i}^{-a_i}$ which is characterized by i -th position x_i and propagator power a_i , the central dot denotes integration over x_0 .

This trick is standard and allows one to evaluate the D -dimensional integral over x_0 by obtaining a set of one-dimensional integrals over λ_i . Then, the n -point conformal integral is represented as

$$I_n^{\mathbf{a}}(\mathbf{x}) = N_n^{\mathbf{a}} \int_0^{+\infty} \prod_{i=1}^n \left(\frac{d\lambda_i}{\lambda_i} \lambda_i^{a_i} \right) \frac{1}{|\boldsymbol{\lambda}_{1,n}|^{\frac{D}{2}}} \exp \left(- \frac{1}{|\boldsymbol{\lambda}_{1,n}|} \sum_{1 \leq i < j \leq n} \lambda_i \lambda_j X_{ij} \right), \quad (2.4)$$

where⁴ $|\boldsymbol{\lambda}_{i,j}| = \sum_{l=i}^j \lambda_l$ and for brevity we defined the prefactor

$$N_j^{\mathbf{a}} = \frac{1}{\Gamma(a_1, \dots, a_j)}, \quad j = 1, \dots, n, \quad (2.5)$$

for which the standard notation for the product of Γ -functions (A.3) is used.

The key observation [1] is that in (2.4) one can substitute $|\boldsymbol{\lambda}_{1,n}| \rightarrow \lambda_n$ as a consequence of the conformality condition (2.2). Representing the exponential functions by means of the Mellin-Barnes formula (B.20)

$$\begin{aligned} \exp \left(- \frac{\lambda_i \lambda_j X_{ij}}{\lambda_n} \right) &= \int_{-i\infty}^{+i\infty} \widehat{ds}_{ij} \left(\frac{\lambda_i \lambda_j X_{ij}}{\lambda_n} \right)^{s_{ij}}, \quad 1 \leq i < j \leq n-2, \\ \exp \left(- \frac{\lambda_k \lambda_{n-1} X_{k,n-1}}{\lambda_n} \right) &= \int_{-i\infty}^{+i\infty} \widehat{dt}_k \left(\frac{\lambda_k \lambda_{n-1} X_{k,n-1}}{\lambda_n} \right)^{t_k}, \quad 1 \leq k \leq n-3, \end{aligned} \quad (2.6)$$

one can successively calculate the integrals over $\lambda_n, \lambda_{n-1}, \dots, \lambda_1$ (the integration measures are defined in (A.4)). As a result, the conformal integral can be represented as a product

$$I_n^{\mathbf{a}}(\mathbf{x}) = L_n^{\mathbf{a}}(\mathbf{x}) \mathcal{I}_n^{\mathbf{a}}(\boldsymbol{\eta}), \quad (2.7)$$

where the *leg-factor*

$$L_n^{\mathbf{a}}(\mathbf{x}) = (Z_n^{n-2, n-1})^{-a'_n} \prod_{i=1}^{n-1} \left(X_{i,n}^{-a_i} \right), \quad a'_i = \frac{D}{2} - a_i, \quad (2.8)$$

⁴The most frequently used notation and conventions are collected in appendix A.

depends on the following combinations of squared distances,

$$Z_k^{ij} \equiv Z_k^{i,j} = \frac{X_{ij}}{X_{ik}X_{jk}}, \quad i, j \neq k, \quad (2.9)$$

which conveniently define the cross-ratios as⁵

$$(\eta_k)_{ml}^{ij} \equiv (\eta_k)_{m,l}^{i,j} = \frac{Z_k^{ij}}{Z_k^{ml}} = \frac{X_{ij}X_{mk}X_{lk}}{X_{ik}X_{jk}X_{ml}}. \quad (2.10)$$

The splitting (2.7) is the consequence of the conformal covariance of (2.1): the leg-factor encodes the conformal transformation properties, while the function $\mathcal{I}_n^\alpha(\boldsymbol{\eta})$, called a *bare conformal integral*, depends only on the cross-ratios (2.10). The bare conformal integral is represented as the following $n(n-3)/2$ -folded Mellin-Barnes integral⁶

$$\begin{aligned} \mathcal{I}_n^\alpha(\boldsymbol{\eta}) &= N_n^\alpha \int_{-\infty}^{+\infty} \prod_{1 \leq i < j \leq n-2} \left(\widehat{ds}_{ij} \left((\eta_n)_{n-2, n-1}^{i,j} \right)^{s_{ij}} \right) \prod_{k=1}^{n-3} \left(\widehat{dt}_k \left((\eta_n)_{n-2, n-1}^{k, n-1} \right)^{t_k} \right) \\ &\times \Gamma \left(|\mathbf{t}_{1, n-3}| + \sum_{1 \leq i < j \leq n-2} s_{ij} + a'_n \right) \prod_{l=1}^{n-3} \Gamma \left(a_l + t_l + \sum_{j=1}^{l-1} s_{jl} + \sum_{j=l+1}^{n-2} s_{lj} \right) \\ &\times \Gamma \left(\alpha_{n-2, n} - \sum_{1 \leq i < j \leq n-3} s_{ij} - |\mathbf{t}_{1, n-3}| \right) \Gamma \left(\alpha_{n-1, n} - \sum_{1 \leq i < j \leq n-2} s_{ij} \right). \end{aligned} \quad (2.11)$$

Here, $\alpha_{i,j}$ is defined in (A.2) and the number of cross-ratios $\boldsymbol{\eta}$ equals the number of integrals on the right-hand side of (2.11). By construction, the integrals over each variable are *balanced* (see e.g. [49]), which means that the contours can be closed either to the left or to the right. In fact, the choice of a contour is determined by a domain of convergence in which one wants to obtain the asymptotic expansion of the conformal integral. In its turn, this determines a specific set of cross-ratios $\boldsymbol{\eta}$.

As n increases, the evaluation of the integral (2.11) quickly becomes difficult due to the complex pole structure of the integrand. In fact, the only well-understood example is given by the box ($n=4$) conformal integral which has been calculated by many authors in various contexts, see e.g. [35] and references therein. Recently, however, significant progress has been made in the computation of higher-folded Mellin-Barnes integrals involving various computer algebra methods [50, 51], in particular, the hexagon ($n=6$) conformal integral was calculated in [52, 53].

We note finally that for some applications it is not necessary to know the whole integral (2.11), but only that part of it which can be obtained by evaluating over a particular subset of poles. E.g. this happens when calculating conformal blocks within the shadow formalism of CFT_D [38, 39, 44].

⁵ The cross-ratios here are generally defined as ratios of cubic combinations of the squared distances. When any two of upper and lower indices coincide they reduce to ratios of quadratic combinations. With a slight abuse of terminology we will call them as quadratic and cubic cross-ratios.

⁶ Appendix B illustrates the formalism of Mellin-Barnes integrals using examples of hypergeometric-type functions.

2.2 Bipartite Mellin-Barnes representation

In what follows, we suggest a different way of representing conformal integrals (2.1) in terms of Mellin-Barnes integrals which does not use the Symanzik conformal trick. The advantage of our approach is that a part of the n -point conformal integral can be computed effortlessly, while the remaining part is conjectured to be recovered by using permutation invariance of the whole conformal integral (2.1) (see section 3).

The first step is to remove $1/|\lambda_{1,n}|$ from the exponential in (2.4) by changing integration variables $\lambda_i \rightarrow |\lambda_{1,n}|\lambda_i$ and obtaining

$$I_n^{\mathbf{a}}(\mathbf{x}) = 2N_n^{\mathbf{a}} \int_0^{+\infty} \prod_{i=1}^n \left(\frac{d\lambda_i}{\lambda_i} \lambda_i^{a_i} \right) \frac{1}{|\lambda_{1,n}|^{D-|\mathbf{a}_{1,n}|}} \exp \left(- \sum_{1 \leq i < j \leq n} \lambda_i \lambda_j X_{ij} \right). \quad (2.12)$$

Then, the contribution of $|\lambda_{1,n}|$ cancels due to the conformality constraint (2.2). Evaluating the integral over λ_n yields

$$I_n^{\mathbf{a}}(\mathbf{x}) = 2N_{n-1}^{\mathbf{a}} \int_0^{+\infty} \prod_{i=1}^{n-1} \left(\frac{d\lambda_i}{\lambda_i} \lambda_i^{a_i} \right) \left(\sum_{i=1}^{n-1} \lambda_i X_{in} \right)^{-a_n} \exp \left(- \sum_{1 \leq i < j \leq n-1} \lambda_i \lambda_j X_{ij} \right). \quad (2.13)$$

Now, changing integration variables as $\lambda_i \rightarrow X_{in}^{-1} \lambda_i$ (no summation over i) one finds

$$I_n^{\mathbf{a}}(\mathbf{x}) = 2N_{n-1}^{\mathbf{a}} \prod_{i=1}^{n-1} X_{i,n}^{-a_i} \int_0^{+\infty} \prod_{i=1}^{n-1} \left(\frac{d\lambda_i}{\lambda_i} \lambda_i^{a_i} \right) \frac{1}{|\lambda_{1,n-1}|^{a_n}} \exp \left(- \sum_{1 \leq i < j \leq n-1} \lambda_i \lambda_j Z_n^{ij} \right), \quad (2.14)$$

where Z_n^{ij} are given by (2.9). Rescaling $\lambda_i \rightarrow \lambda_i \sqrt{\lambda_{n-1}}$ for $i = 1, \dots, n-2$ and $\lambda_{n-1} \rightarrow \sqrt{\lambda_{n-1}}$ one can evaluate the integral over λ_{n-1} :

$$I_n^{\mathbf{a}}(\mathbf{x}) = N_{n-1}^{\mathbf{a}} \Gamma(a'_n) \prod_{i=1}^{n-1} X_{i,n}^{-a_i} \int_0^{+\infty} \prod_{j=1}^{n-2} \left(\frac{d\lambda_j}{\lambda_j} \lambda_j^{a_j} \right) \times \frac{\left(1 + |\lambda_{1,n-2}| \right)^{-a_n}}{\left(\sum_{1 \leq l < m \leq n-2} \lambda_l \lambda_m Z_n^{lm} + \sum_{k=1}^{n-2} \lambda_k Z_n^{k,n-1} \right)^{a'_n}}. \quad (2.15)$$

Introducing new variables by $\lambda_i = s\sigma_i$ for $i = 1, \dots, n-3$ and $\lambda_{n-2} = s(1 - |\sigma|)$, where $|\sigma| \equiv |\sigma_{1,n-3}|$, the integral over s is recognized as the integral representation of the Gauss hypergeometric function (B.18). As a result, the conformal integral is again represented as the product (2.7) with the same leg-factor $L_n^{\mathbf{a}}(\mathbf{x})$ (2.8), but with the bare integral $\mathcal{I}_n^{\mathbf{a}}(\boldsymbol{\eta})$ written in the form

$$\mathcal{I}_n^{\mathbf{a}}(\boldsymbol{\eta}) = N_{n-2}^{\mathbf{a}} \Gamma \left[\begin{matrix} a'_{n-1}, a'_n \\ \frac{D}{2} \end{matrix} \right] \int_{\Omega} d^{n-3} \sigma \frac{\prod_{i=1}^{n-3} \sigma_i^{a_i-1} (1 - |\sigma|)^{a_{n-2}-1}}{(1 - \boldsymbol{\sigma} \cdot \boldsymbol{\xi})^{a'_n}} {}_2F_1 \left[\begin{matrix} a'_{n-1}, a'_n \\ \frac{D}{2} \end{matrix} \middle| 1 - \xi(\boldsymbol{\sigma}) \right], \quad (2.16)$$

where the integration domain is the standard orthogonal simplex $\Omega = \{\boldsymbol{\sigma} \in \mathbb{R}^{n-3} : \sigma_j \geq 0, j = 1, \dots, n-3; |\boldsymbol{\sigma}| \leq 1\}$, and

$$\xi(\boldsymbol{\sigma}) = \frac{\sum_{1 \leq i < j \leq n-3} \sigma_i \sigma_j (\eta_n)_{n-2, n-1}^{ij} + (1 - |\boldsymbol{\sigma}|) \sum_{l=1}^{n-3} \sigma_l (\eta_n)_{n-2, n-1}^{l, n-2}}{1 - \boldsymbol{\sigma} \cdot \boldsymbol{\xi}}, \quad (2.17)$$

$$\boldsymbol{\sigma} \cdot \boldsymbol{\xi} = \sum_{l=1}^{n-3} \sigma_l \xi_l, \quad \xi_i = 1 - (\eta_n)_{n-2, n-1}^{i, n-1}, \quad 1 \leq i \leq n-3,$$

where $(\eta_n)_{n-2, n-1}^{ij}$ are the cross-ratios (2.10). Note that there are $(n-3)(n-4)/2 + (n-3) = (n-3)(n-2)/2$ terms in the numerator of $\xi(\boldsymbol{\sigma})$, each of which contains one cross-ratio. Adding $(n-3)$ cross-ratios from the denominator one concludes that the bare conformal integral (2.16) does depend on $n(n-3)/2$ cross-ratios of the type (2.10), as discussed below (2.11).

Let us now express the bare integral (2.16) through the Mellin-Barnes integrals. For this purpose, it turns out to be convenient to use the analytic continuation formula for the Gauss hypergeometric function (B.25) that results in splitting the bare integral into two terms

$$\mathcal{I}_n^{\mathbf{a}}(\boldsymbol{\eta}) = \mathcal{I}_n^{(1), \mathbf{a}}(\boldsymbol{\eta}) + \mathcal{I}_n^{(2), \mathbf{a}}(\boldsymbol{\eta}), \quad (2.18)$$

where the terms on the right-hand side which we refer to as the *first and second bare integrals* are given by

$$\mathcal{I}_n^{(1), \mathbf{a}}(\boldsymbol{\eta}) = N_n^{\mathbf{a}} \Gamma(\alpha_{n-1, n}, a'_{n-1}, a'_n) \int_{\Omega} d^{n-3} \sigma \frac{\prod_{i=1}^{n-3} \sigma_i^{a_i-1} (1 - |\boldsymbol{\sigma}|)^{a_{n-2}-1}}{(1 - \boldsymbol{\sigma} \cdot \boldsymbol{\xi})^{a'_n}} \times {}_2F_1 \left[\begin{matrix} a'_{n-1}, a'_n \\ 1 - \alpha_{n-1, n} \end{matrix} \middle| \xi(\boldsymbol{\sigma}) \right], \quad (2.19)$$

$$\mathcal{I}_n^{(2), \mathbf{a}}(\boldsymbol{\eta}) = N_n^{\mathbf{a}} \Gamma(-\alpha_{n-1, n}, a_{n-1}, a_n) \int_{\Omega} d^{n-3} \sigma \frac{\prod_{i=1}^{n-3} \sigma_i^{a_i-1} (1 - |\boldsymbol{\sigma}|)^{a_{n-2}-1}}{(1 - \boldsymbol{\sigma} \cdot \boldsymbol{\xi})^{a_n}} \times (\xi(\boldsymbol{\sigma}))^{\alpha_{n-1, n}} {}_2F_1 \left[\begin{matrix} a_{n-1}, a_n \\ 1 + \alpha_{n-1, n} \end{matrix} \middle| \xi(\boldsymbol{\sigma}) \right]. \quad (2.20)$$

This is a *bipartite representation* of the n -point (bare) conformal integral. A few comments are in order.

- Using a certain analytic continuation formula is a formal trick because we cannot precisely describe a domain of the functional argument (2.17). Nonetheless, this procedure

will eventually lead to representing the conformal integral as a multivariate powers series in $n(n-3)/2$ cross-ratios which is valid for a particular arrangement of points x_i .

- We note that one could directly represent the Gauss hypergeometric function in (2.16) by means of one of its Mellin-Barnes representations (B.24). This yields a representation of the conformal integral in terms of the Mellin-Barnes integrals given in appendix C. However, this particular representation turns out to be less computationally efficient than that one we derive in this section.
- Similarly, another analytic continuation of the Gauss hypergeometric function could be used that would change the form of two terms in (2.18). Nonetheless, the convenience of choosing this particular partitioning of the conformal integral can be seen, in particular, in the $n=3$ case, when the second bare integral vanishes since the numerator of (2.17) has no terms, while the first bare integral leads to the star-triangle relation [1]

$$I_3^a(\mathbf{x}) = S_3^{(123)} X_{12}^{-a'_3} X_{13}^{-a'_2} X_{23}^{-a'_1}, \quad \text{where} \quad S_3^{(123)} \equiv \Gamma \left[\begin{matrix} a'_1, a'_2, a'_3 \\ a_1, a_2, a_3 \end{matrix} \right], \quad (2.21)$$

which is commonly represented as the following diagrammatic equality

- In what follows we restrict $n \geq 3$ so that the 1-point and 2-point conformal integrals fall out of our consideration. These lower-point integrals diverge and require careful regularizations which have been extensively studied, see e.g. [54–57]. Notably, the 2-point conformal integral being proportional to $\delta(x_1 - x_2)$ is the cornerstone of the shadow formalism of CFT_D [33, 58].

The term $\mathcal{I}_n^{(1),a}$. Let us now explicitly calculate the first bare integral (2.19). To integrate over σ_i we use the Mellin-Barnes representation for the Gauss hypergeometric function (B.19):

$${}_2F_1 \left[\begin{matrix} a'_{n-1}, a'_n \\ 1 - \alpha_{n-1,n} \end{matrix} \middle| \xi(\boldsymbol{\sigma}) \right] = \Gamma \left[\begin{matrix} 1 - \alpha_{n-1,n} \\ a'_{n-1}, a'_n \end{matrix} \right] \int_{-i\infty}^{+i\infty} \widehat{d}t \Gamma \left[\begin{matrix} a'_{n-1} + t, a'_n + t \\ 1 - \alpha_{n-1,n} + t \end{matrix} \right] (-\xi(\boldsymbol{\sigma}))^t. \quad (2.22)$$

When $n=4$, the numerator of (2.17) contains only one term, and thus we are left with the 1-dimensional integral over σ_1 which can be transformed into the Gauss hypergeometric function

(B.18). At $n > 4$ the functional argument (2.17) is more complicated and its numerator can be converted into the $(n-3)(n-2)/2 - 1$ Mellin-Barnes integrals by means of the Mellin-Barnes expansion (B.21):

$$\begin{aligned}
& \left(\sum_{1 \leq i < j \leq n-3} \sigma_i \sigma_j (\eta_m)_{n-2, n-1}^{ij} + (1 - |\boldsymbol{\sigma}|) \sum_{l=1}^{n-3} \sigma_l (\eta_m)_{n-2, n-1}^{l, n-2} \right)^t = \frac{1}{\Gamma(-t)} \\
& \times \int_{-i\infty}^{+i\infty} \prod_{l=1}^{n-4} \widehat{d}t_l \int_{-i\infty}^{+i\infty} \prod_{1 \leq k < m \leq n-3} \widehat{d}s_{km} \Gamma \left(|\mathbf{t}_{1, n-4}| + \sum_{1 \leq i < j \leq n-3} s_{ij} - t \right) \\
& \times \prod_{l=1}^{n-4} \left((1 - |\boldsymbol{\sigma}|) \sigma_l (\eta_m)_{n-2, n-1}^{l, n-2} \right)^{t_l} \prod_{1 \leq i < j \leq n-3} \left(\sigma_i \sigma_j (\eta_m)_{n-2, n-1}^{ij} \right)^{s_{ij}} \\
& \times \left((1 - |\boldsymbol{\sigma}|) \sigma_{n-3} (\eta_m)_{n-2, n-1}^{n-3, n-2} \right)^{t - |\mathbf{t}_{1, n-4}| - \sum_{1 \leq i < j \leq n-3} s_{ij}}.
\end{aligned} \tag{2.23}$$

The integral over σ_i in (2.19) is recognized as the integral representation of the Lauricella function $F_D^{(n-3)}$ (B.15). Thus, we obtain the first bare integral (2.19) in the following form

$$\begin{aligned}
\mathcal{I}_n^{(1), \mathbf{a}} &= N_n^{\mathbf{a}} \Gamma(1 - \alpha_{n-1, n}, \alpha_{n-1, n}) \int_{-i\infty}^{+i\infty} \frac{dt}{2\pi i} \Gamma \left[\begin{matrix} a'_{n-1} + t, a'_n + t \\ 1 - \alpha_{n-1, n} + t \end{matrix} \right] \left(-(\eta_m)_{n-2, n-1}^{n-3, n-2} \right)^t \\
& \times \int_{-i\infty}^{+i\infty} \prod_{l=1}^{n-4} \left(\widehat{d}t_l \left((\eta_m)_{n-3, n-2}^{l, n-2} \right)^{t_l} \right) \int_{-i\infty}^{+i\infty} \prod_{1 \leq i < j \leq n-3} \left(\widehat{d}s_{ij} \left((\eta_m)_{n-3, n-2}^{ij} \right)^{s_{ij}} \right) \\
& \times \Gamma \left(|\mathbf{t}_{1, n-4}| + \sum_{1 \leq i < j \leq n-3} s_{ij} - t \right) \Gamma \left[\begin{matrix} C^{(1)} - |\mathbf{B}^{(1)}|, \mathbf{B}^{(1)} \\ C^{(1)} \end{matrix} \right] F_D^{(n-3)} \left[\begin{matrix} A^{(1)}, \mathbf{B}^{(1)} \\ C^{(1)} \end{matrix} \middle| \boldsymbol{\xi} \right],
\end{aligned} \tag{2.24}$$

where $\boldsymbol{\xi} = \{\xi_1, \dots, \xi_{n-3}\}$ are given in (2.17), and the parameters here, including $\Gamma(\mathbf{B}^{(1)}) \equiv \Gamma(B_1^{(1)}, \dots, B_{n-3}^{(1)})$, are expressed in terms of propagator powers a_i as

$$\begin{aligned}
A^{(1)} &= a'_n + t, & C^{(1)} &= |\mathbf{a}_{1, n-2}| + 2t, & B_{n-3}^{(1)} &= a_{n-3} + t - |\mathbf{t}_{1, n-4}| - \sum_{1 \leq i < j \leq n-4} s_{ij}, \\
B_l^{(1)} &= a_l + t_l + \sum_{j=1}^{l-1} s_{jl} + \sum_{j=l+1}^{n-3} s_{lj}, & & & & l = 1, \dots, n-4.
\end{aligned} \tag{2.25}$$

By construction, all the contours in (2.24) for all integrals can be closed both to the left and right. When the contour over t is closed to the right, only the poles coming from $\Gamma \left(|\mathbf{t}_{1, n-4}| + \sum_{1 \leq i < j \leq n-3} s_{ij} - t \right)$ contribute to the integral (see the third line in (2.24)). Calculating the integral over t then leads to the fact that closing contours to the right in all the remaining integrals over t_i and s_{ij} only one set of poles contributes for each integration

variable, namely, these are the poles coming from $\Gamma(-t_i)$ and $\Gamma(-s_{ij})$ (cf. the integration measure (A.4)). In this way, the whole integral (2.24) can be explicitly calculated as

$$\begin{aligned} \mathcal{I}_n^{(1),\mathbf{a}} &= N_n^{\mathbf{a}} \Gamma(1 - \alpha_{n-1,n}, \alpha_{n-1,n}) \prod_{l=1}^{n-3} v_l^{\alpha_l} \sum_{\{k_l, p_{ij}\}=0}^{\infty} \prod_{l=1}^{n-3} \frac{u_l^{k_l}}{k_l!} \prod_{1 \leq i < j \leq n-3} \frac{w_{ij}^{p_{ij}}}{p_{ij}!} \\ &\times \Gamma \left[\begin{matrix} a_{n-2} + |\mathbf{k}|, a'_n + |\mathbf{k}| + |\mathbf{p}|, \mathcal{A}^{(1)}, \mathcal{B}^{(1)} \\ 1 - \alpha_{n-1,n} + |\mathbf{k}| + |\mathbf{p}|, \mathcal{C}^{(1)} \end{matrix} \right] F_D^{(n-3)} \left[\begin{matrix} \mathcal{A}^{(1)}, \mathcal{B}^{(1)} \\ \mathcal{C}^{(1)} \end{matrix} \middle| \mathbf{1} - \mathbf{v} \right], \end{aligned} \quad (2.26)$$

where we used the transformation formula for the Lauricella function (B.17); the parameters here are related to propagator powers and summation indices as follows

$$\mathcal{A}^{(1)} = a'_{n-1} + |\mathbf{k}| + |\mathbf{p}|, \quad \mathcal{B}_l^{(1)} = a_l + k_l + 2p_l, \quad \mathcal{C}^{(1)} = |\mathbf{a}_{1,n-2}| + 2(|\mathbf{k}| + |\mathbf{p}|), \quad (2.27)$$

where $l = 1, \dots, n-3$; the sums are denoted as

$$|\mathbf{p}| = \sum_{1 \leq i < j \leq n-3} p_{ij} = \sum_{l=1}^{n-3} p_l, \quad p_l = \frac{1}{2} \left(\sum_{j=1}^{l-1} p_{jl} + \sum_{j=l+1}^{n-3} p_{lj} \right), \quad l = 1, \dots, n-3. \quad (2.28)$$

The cross-ratios in (2.26) are given by

$$\begin{aligned} u_i &= (\eta_{n-1})_{n-2,n}^{i,n-2}, & i &= 1, \dots, n-3, \\ w_{ij} &= (\eta_{n-1})_{n-2,n}^{i,j}, & 1 \leq i < j \leq n-3, \\ v_i &= (\eta_{n-1})_{n-2,n}^{i,n}, & i &= 1, \dots, n-3, \end{aligned} \quad (2.29)$$

where u_i, v_j , and w_{kl} are quadratic and cubic, respectively (see footnote 5). Given the multivariate power series (2.26) one can find its region of convergence, that, however, can be a rather complicated problem. For simplicity, we suppose that such a series converges at least when the cross-ratios (2.29) tend to specific points. In particular, since the Lauricella function F_D (B.4) is supported on the polydisk (B.5), we can claim that the multivariate power series (2.26) converges when

$$u_i, w_{kl} \rightarrow 0 \quad \text{and} \quad v_j \rightarrow 1. \quad (2.30)$$

Clearly, a range of the cross-ratios is determined by a particular arrangement of points x_i and (2.30) corresponds to the coincidence limit

$$x_n \rightarrow x_{n-1}. \quad (2.31)$$

The term $\mathcal{I}_n^{(2),\mathbf{a}}$. Consider the second bare integral (2.20). Repeating the same steps as in the previous paragraph one shows that

$$\begin{aligned} \mathcal{I}_n^{(2),\mathbf{a}} &= N_n^{\mathbf{a}} \Gamma(1 + \alpha_{n-1,n}, -\alpha_{n-1,n}) \int_{-i\infty}^{+i\infty} \widehat{d}t \Gamma \left[\begin{matrix} a_{n-1} + t, a_n + t \\ 1 + \alpha_{n-1,n} + t, -\alpha_{n-1,n} - t \end{matrix} \right] \left((\eta_n)_{n-2, n-1}^{n-3, n-2} \right)^{\alpha_{n-1,n} + t} \\ &\times \int_{-i\infty}^{+i\infty} \prod_{l=1}^{n-4} \left(\widehat{d}t_l \left((\eta_n)_{n-3, n-2}^{l, n-2} \right)^{t_l} \right) \int_{-i\infty}^{+i\infty} \prod_{1 \leq i < j \leq n-3} \left(\widehat{d}s_{ij} \left((\eta_n)_{n-3, n-2}^{i, j} \right)^{s_{ij}} \right) \\ &\Gamma \left(|\mathbf{t}_{1, n-4}| + \sum_{1 \leq i < j \leq n-3} s_{ij} - t - \alpha_{n-1,n} \right) \Gamma \left[\begin{matrix} C^{(2)} - |\mathbf{B}^{(2)}|, \mathbf{B}^{(2)} \\ C^{(2)} \end{matrix} \right] F_D^{(n-3)} \left[\begin{matrix} A^{(2)}, \mathbf{B}^{(2)} \\ C^{(2)} \end{matrix} \middle| \boldsymbol{\xi} \right], \end{aligned} \quad (2.32)$$

where

$$\begin{aligned} A^{(2)} &= a_{n-1} + t, \quad C^{(2)} = |\mathbf{a}_{1, n-2}| + 2\alpha_{n-1,n} + 2t, \\ B_l^{(2)} &= a_l + t_l + \sum_{j=1}^{l-1} s_{jl} + \sum_{j=l+1}^{n-3} s_{lj}, \quad l = 1, \dots, n-4, \\ B_{n-3}^{(2)} &= a_{n-3} + \alpha_{n-1,n} + t - |\mathbf{t}_{1, n-4}| - \sum_{1 \leq i < j \leq n-4} s_{ij}; \end{aligned} \quad (2.33)$$

the arguments of the Lauricella function $F_D^{(n-3)}$ are given in (2.17).

The pole structure of the second bare integral $\mathcal{I}_n^{(2),\mathbf{a}}(\mathbf{x})$ is more complicated than that of the first bare integral $\mathcal{I}_n^{(1),\mathbf{a}}(\mathbf{x})$ so we cannot immediately calculate the integrals in (2.32). The reason for this difference is the use of particular analytic continuation formula for the Gauss hypergeometric function in the integral (2.16), which makes the two terms (2.19) and (2.20) different due to the additional factor of $\xi(\boldsymbol{\sigma})$ in (2.20). In turn, when closing contours to the right, this gives an additional set of poles over t in (2.32). Nonetheless, the problem of computing the second bare integral $\mathcal{I}_n^{(2),\mathbf{a}}(\mathbf{x})$ can be addressed by invoking invariance of the whole conformal integral under the symmetric group.

3 Constraints from the permutation invariance

The symmetric group \mathcal{S}_n naturally acts on functions of coordinates and propagator powers by permuting elements from $\mathbf{x} = (x_1, \dots, x_n)$ and $\mathbf{a} = (a_1, \dots, a_n)$ as $x_i \rightarrow x_{\pi(i)}$ and $a_i \rightarrow a_{\pi(i)}$. The conformal integral (2.1) is invariant under the symmetric group:

$$\forall \pi \in \mathcal{S}_n : \quad \mathbf{R}(\pi) \circ I_n^{\mathbf{a}}(\mathbf{x}) = I_n^{\mathbf{a}}(\mathbf{x}), \quad (3.1)$$

where $\mathbf{R}(\pi)$ stands for the representation of π .⁷ This invariance property is manifestly seen on the conformal graph on fig. 1 which remains invariant with respect to any permutation of legs which are labelled by (x_i, a_i) .

⁷We will use the one-line notation for elements of the symmetric group. The identical permutation will be denoted as e and $\mathbf{R}(e) = \mathbf{1}$.

On the other hand, a splitting of the conformal integral into two terms provided by the bipartite representation is not invariant with respect to the symmetric group. In this way, having found the first term explicitly, we expect to generate the unknown second term by acting on the first one with particular permutations.

3.1 Analytic continuation

To simplify the study of permutation invariance, it is useful to analytically continue the known first bare integral, which is originally defined at $\mathbf{u}, \mathbf{w} \rightarrow 0$ and $\mathbf{v} \rightarrow 1$, into the other domain, where all cross-ratios are near zero:

$$\mathbf{u}, \mathbf{w}, \mathbf{v} \rightarrow 0. \quad (3.2)$$

Recalling the explicit expression (2.26) we notice that this can be achieved by exploiting the analytic continuation formulas for the Lauricella function $F_D^{(n-3)}$ derived by Bezrodnykh in [59–63] (see also appendix B). Indeed, by applying the analytic continuation formula (B.26) one can split (2.26) into $n - 2$ terms thereby obtaining the first bare integral in the domain (3.2) as the following sum⁸

$$\mathcal{I}_n^{(1),\mathbf{a}}(\boldsymbol{\eta}) = \sum_{q=0}^{n-3} \mathcal{U}_n^{(q),\mathbf{a}}(\boldsymbol{\eta}), \quad (3.3)$$

where the $q = 0$ term:

$$\begin{aligned} \mathcal{U}_n^{(0),\mathbf{a}} &= S_n^{(n-2,n-1,n)} \prod_{i=1}^{n-3} v_i^{a_i} \sum_{\{k_i, p_{ij}, m_i\}=0}^{\infty} \prod_{i=1}^{n-3} \frac{u_i^{k_i}}{k_i!} \prod_{1 \leq i < j \leq n-3} \frac{w_{ij}^{p_{ij}}}{p_{ij}!} \prod_{l=1}^{n-3} \frac{(v_l)^{m_l}}{m_l!} \\ &\quad \times \frac{\prod_{l=1}^{n-3} (a_l)_{k_l+2p_l+m_l} (a'_{n-1})_{|\mathbf{k}|+|\mathbf{p}|+|\mathbf{m}|} (-)^{|\mathbf{p}|}}{(1 - \alpha_{n-1,n})_{|\mathbf{k}|+|\mathbf{p}|} (1 - \alpha_{n-2,n-1})_{|\mathbf{p}|+|\mathbf{m}|}}, \end{aligned} \quad (3.4)$$

the terms with $q = 1, \dots, n - 3$:

$$\begin{aligned} \mathcal{U}_n^{(q),\mathbf{a}} &= S_n^{(q,n-1,n)} v_q^{a'_n - |\mathbf{a}_{1,q-1}|} \prod_{l=1}^{q-1} v_l^{a_l} \sum_{\{k_i, p_{ij}, m_i\}=0}^{\infty} \prod_{l=1}^{n-3} \frac{(\mathcal{U}_l^{(q)})^{k_l}}{k_l!} \prod_{1 \leq i < j \leq n-3} \frac{(\mathcal{W}_{ij}^{(q)})^{p_{ij}}}{p_{ij}!} \prod_{l=1}^{n-3} \frac{(\mathcal{V}_l^{(q)})^{m_l}}{m_l!} \\ &\quad \times \frac{(a'_n - |\mathbf{a}_{1,q-1}|)_{|\mathbf{m}_q|+|\mathbf{k}_{q,n-3}|+|\mathbf{p}_q|} (-)^{|\mathbf{k}_{q+1,n-3}|+|\mathbf{p}_{q+1}|}}{(1 + a'_n - |\mathbf{a}_{1,q}|)_{|\mathbf{m}_q|+|\mathbf{k}_{q+1,n-3}|+|\mathbf{p}_{q+1}|}} \\ &\quad \times \frac{\prod_{l=1}^{q-1} (a_l)_{k_l+2p_l+m_l} (a_{n-2})_{|\mathbf{k}|+m_q} \prod_{l=q+1}^{n-3} (a_l)_{k_l+2p_l+m_l}}{(1 - \alpha_{n-1,n})_{|\mathbf{k}|+|\mathbf{p}|}}. \end{aligned} \quad (3.5)$$

⁸Here, the q -th term $\mathcal{U}_n^{(q),\mathbf{a}}$ arises from the q -th term D_q (B.28).

Here, $|\mathbf{p}|$ is defined in (2.28), other notations are given in appendix A. The new cross-ratios in (3.5) are expressed in terms of the previously introduced cross-ratios (2.29) as follows

$$\mathcal{U}_l^{(q)} = \begin{cases} u_l, & l = 1, \dots, q, \\ \frac{u_l v_q}{v_l}, & l = q + 1, \dots, n - 3, \end{cases} \quad \mathcal{V}_l^{(q)} = \begin{cases} \frac{v_l}{v_q}, & l = 1, \dots, q - 1, \\ v_q, & l = q, \\ \frac{v_q}{v_l}, & l = q + 1, \dots, n - 3, \end{cases} \quad (3.6)$$

$$\mathcal{W}_{ij}^{(q)} = \begin{cases} \frac{w_{ij}}{v_q}, & 1 \leq i < j \leq q, \\ \frac{w_{ij}}{v_j}, & i = 1, \dots, q, \quad j = q + 1, \dots, n - 3, \\ \frac{w_{ij} v_q}{v_i v_j}, & q + 1 \leq i < j \leq n - 3. \end{cases} \quad (3.7)$$

The overall coefficients in (3.4)-(3.5) are defined as

$$S_n^{(ijk)} = \Gamma \left[\begin{matrix} |\mathbf{a}_{i,j}| - \frac{D}{2}, |\mathbf{a}_{j,k}| - \frac{D}{2}, |\mathbf{a}_{k,i}| - \frac{D}{2} \\ a_i, \quad a_j, \quad a_k \end{matrix} \right], \quad 1 \leq i < j < k \leq n. \quad (3.8)$$

These generalize the prefactor in the star-triangle relation (2.21). Note that the last multiplier in the numerator contains $|\mathbf{a}_{k,i}|$ with $k > i$, which is defined in (A.2).

It is important to stress that having the expansion (3.3) near the origin in the space of cross-ratios, i.e. at $u_i, w_{kl}, v_j \rightarrow 0$, means that other arguments must also tend to zero, i.e.

$$\mathcal{U}_i^{(q)}, \mathcal{W}_{kl}^{(q)}, \mathcal{V}_j^{(q)} \rightarrow 0. \quad (3.9)$$

Since $\mathcal{U}_i^{(q)}, \mathcal{W}_{kl}^{(q)}, \mathcal{V}_j^{(q)}$ are (rational) functions of u_i, w_{kl}, v_j (3.6), then the condition (3.9) determines a specific way u_i, w_{kl}, v_j tend to zero.⁹

For future purposes, we note here that there are only two terms in the expansion (3.3) which can be related by the action of a longest cycle $C_n \equiv (12\dots n) \in \mathcal{S}_n$. Namely,

$$C_n \circ \left(L_n^{\mathbf{a}}(\mathbf{x}) U_n^{(0), \mathbf{a}}(\boldsymbol{\eta}) \right) = L_n^{\mathbf{a}}(\mathbf{x}) U_n^{(1), \mathbf{a}}(\boldsymbol{\eta}), \quad (3.10)$$

where by $C_n = R(C_n)$ we denote the action of C_n on functions of \mathbf{x} and \mathbf{a} . The proof is fairly straightforward if we consider how C_n changes both the leg-factor (2.8):

$$C_n \circ L_n^{\mathbf{a}}(\mathbf{x}) = v_1^{a'_n} u_1^{-a_n - 2} \prod_{j=2}^{n-3} \left(\frac{w_{1j}}{v_j} \right)^{-a_j} L_n^{\mathbf{a}}(\mathbf{x}), \quad (3.11)$$

⁹E.g. there are two variables such that $x, y \rightarrow 0$ and $x/y \rightarrow 0$ that means that $x \rightarrow 0$ faster than $y \rightarrow 0$.

and the cross-ratios (2.29):

$$\begin{aligned} C_n \circ u_i &= \frac{v_1}{v_{i+1}}, & C_n \circ u_{n-3} &= v_1, & C_n \circ w_{ij} &= \frac{w_{i+1,j+1}v_1}{v_{i+1}v_{j+1}}, \\ C_n \circ v_i &= \frac{w_{1,i+1}}{v_{i+1}}, & C_n \circ v_{n-3} &= u_1, & C_n \circ w_{i,n-3} &= \frac{u_{i+1}v_1}{v_{i+1}}, \end{aligned} \quad (3.12)$$

where $i, j = 1, \dots, n-4$ and $i < j$. The relation (3.10) stems directly from the fact that both power series $U_n^{(0),\mathbf{a}}$, $U_n^{(1),\mathbf{a}}$ arise from the Lauricella function $F_D^{(n-3)}$ (see (B.11) and (B.28)) by means of the analytic continuation formula (B.26). Note that all other terms ($q \neq 0, 1$) in (3.3) arise from the generalized hypergeometric series $G^{(n-3,q)}$ (see (B.28)) which are functions of different types for different q . Hence, other possible relations among $U_n^{(q),\mathbf{a}}$ involving permutations different from the longest cycle cannot be obtained in such an obvious way.

In the subsequent sections we will study the bipartite representation (2.18) supplemented with the analytic continuation formula (3.3) and the symmetric group action on the particular examples of the box, pentagon, and hexagon conformal integrals. This will allow us to have a number of observations, conclusions, and exact results which will eventually be summarized in section 4 in the form of the reconstruction conjecture.

3.2 Box integral

In this case, the first bare integral $\mathcal{I}_4^{(1),\mathbf{a}}(u, v)$ is supported on some domain on the (u, v) -plane of cross-ratios

$$u = \frac{X_{12}X_{34}}{X_{13}X_{24}}, \quad v = \frac{X_{14}X_{23}}{X_{13}X_{24}}. \quad (3.13)$$

Note that there are no cubic cross-ratios in this case. According to the general formula (2.26), it can be explicitly calculated as the asymptotic expansion at $(u, v) \rightarrow (0, 1)$:

$$\begin{aligned} \mathcal{I}_4^{(1),\mathbf{a}}(u, v) &= N_4^{\mathbf{a}} \Gamma(1 - \alpha_{3,4}, \alpha_{3,4}) v^{a_1} \\ &\times \sum_{k=0}^{\infty} \frac{u^k}{k!} \Gamma \left[\begin{matrix} a_1 + k, a_2 + k, a'_3 + k, a'_4 + k \\ a_1 + a_2 + 2k, 1 - \alpha_{3,4} + k \end{matrix} \right] {}_2F_1 \left[\begin{matrix} a_1 + k, a'_3 + k \\ a_1 + a_2 + 2k \end{matrix} \middle| 1 - v \right], \end{aligned} \quad (3.14)$$

We note that an arbitrary permutation from the symmetric group \mathcal{S}_4 acting on the first bare integral $\mathcal{I}_4^{(1),\mathbf{a}}$ (3.14) produces a power series whose domain of convergence is generally different from the original domain. For instance, a transposition $(23) \in \mathcal{S}_4$ acts on the cross-ratios as

$$\sigma_{23} \circ u = \frac{1}{u}, \quad \sigma_{23} \circ v = \frac{v}{u}, \quad (3.15)$$

where $\sigma_{ij} \equiv \mathbb{R}((ij))$ denotes the action of (ij) on functions of \mathbf{x} and \mathbf{a} . Obviously, such a permutation sends the original domain $(u, v) \rightarrow (0, 1)$ to $(u, v) \rightarrow \infty$. In order to return back to the original domain, the action (3.15) must be accompanied by making a suitable analytic continuation of the resulting power series for $\mathcal{I}_4^{(1),\mathbf{a}}$. However, analytic continuation formulas

are known only for a limited number of generalized hypergeometric functions. Thus, to avoid the need to use analytical continuation we first try permutations which leave the cross-ratios invariant thereby keeping any domain of convergence intact.

3.2.1 Kinematic group

Those elements of the symmetric group \mathcal{S}_4 which leave the cross-ratios invariant,

$$\exists \pi \in \mathcal{S}_4 : \quad \mathbf{R}(\pi) \circ u = u, \quad \mathbf{R}(\pi) \circ v = v, \quad (3.16)$$

form a subgroup called *kinematic* [64]:

$$\mathcal{S}_4^{\text{kin}} = \left\{ e, (12)(34), (13)(24), (23)(14) \right\} \equiv \mathbb{Z}_2 \times \mathbb{Z}_2 \subset \mathcal{S}_4. \quad (3.17)$$

By construction, a given element of the kinematic group generates from the first bare integral a power series in the same domain $(u, v) \rightarrow (0, 1)$. E.g. let us consider a permutation (23)(14) and examine its action on $\mathcal{I}_4^{(1),\mathbf{a}}(u, v)$. Note that despite the cross-ratios are invariant such an action is not trivial since the permutation also interchanges $a_2 \leftrightarrow a_3$ and $a_1 \leftrightarrow a_4$. To this end, one considers the invariance condition (3.1) which in this case takes the form

$$\sigma_{23}\sigma_{14} \circ I_4^{\mathbf{a}}(\mathbf{x}) = I_4^{\mathbf{a}}(\mathbf{x}). \quad (3.18)$$

Recalling the double product representation of the conformal integral (2.7) and using that the 4-point leg-factor (2.8) transforms as

$$\sigma_{23}\sigma_{14} \circ L_4^{\mathbf{a}}(\mathbf{x}) = u^{\alpha_{3,4}} v^{a_1 - a_4} L_4^{\mathbf{a}}(\mathbf{x}), \quad \text{where } L_4^{\mathbf{a}}(\mathbf{x}) = X_{14}^{-a_1} X_{23}^{-a_2} X_{24}^{-\alpha_{2,4}} X_{34}^{-\alpha_{3,4}}, \quad (3.19)$$

one obtains that the bare conformal integrals are related as

$$u^{\alpha_{3,4}} v^{a_1 - a_4} \left(\sigma_{23}\sigma_{14} \circ \mathcal{I}_4^{(1),\mathbf{a}}(u, v) + \sigma_{23}\sigma_{14} \circ \mathcal{I}_4^{(2),\mathbf{a}}(u, v) \right) = \mathcal{I}_4^{(1),\mathbf{a}}(u, v) + \mathcal{I}_4^{(2),\mathbf{a}}(u, v). \quad (3.20)$$

It can be seen that the asymptotics of the first term on the left-hand side at $(u, v) \rightarrow (0, 1)$ is different from that of $\mathcal{I}_4^{(1),\mathbf{a}}$ (3.14). It is the prefactor $u^{\alpha_{3,4}} v^{a_1 - a_4}$ which changes the asymptotic behaviour since by construction both $\mathcal{I}_4^{(1),\mathbf{a}}$ and $\sigma_{23}\sigma_{14} \circ \mathcal{I}_4^{(1),\mathbf{a}}$ have the same asymptotics in the considered domain. Thus, the first term on the left-hand side must coincide with the second term on the right-hand side which is the second bare integral $\mathcal{I}_4^{(2),\mathbf{a}}$, i.e.

$$\mathcal{I}_4^{(2),\mathbf{a}}(u, v) = u^{\alpha_{3,4}} v^{a_1 - a_4} \sigma_{23}\sigma_{14} \circ \mathcal{I}_4^{(1),\mathbf{a}}(u, v). \quad (3.21)$$

Explicitly, this gives the second bare integral as the asymptotic expansion at $(u, v) \rightarrow (0, 1)$:

$$\begin{aligned} \mathcal{I}_4^{(2),\mathbf{a}}(u, v) &= N_4^{\mathbf{a}} \Gamma(1 - \alpha_{1,2}, \alpha_{1,2}) u^{\alpha_{3,4}} v^{a_1} \\ &\times \sum_{k=0}^{\infty} \frac{u^k}{k!} \Gamma \left[\begin{matrix} a_1' + k, a_2' + k, a_3 + k, a_4 + k \\ a_3 + a_4 + 2k, 1 - \alpha_{1,2} + k \end{matrix} \right] {}_2F_1 \left[\begin{matrix} a_2' + k, a_4 + k \\ a_3 + a_4 + 2k \end{matrix} \middle| 1 - v \right], \end{aligned} \quad (3.22)$$

cf. (3.14). We conclude that the second bare integral can be reconstructed from the first one using a certain permutation. On the other hand, going back to the Mellin-Barnes representation of $\mathcal{I}_4^{(2),\mathbf{a}}$ (2.32), we note that in this case the integrals can be evaluated explicitly and the resulting function does coincide with (3.22).

The remaining permutations from $\mathcal{S}_4^{\text{kin}}$ can be considered along the same lines. One finds out that acting with (13)(24) yields the same result, i.e. this permutation also reconstructs the second bare integral $\mathcal{I}_4^{(2),\mathbf{a}}$, while (12)(34) keeps each bare integral invariant, i.e.

$$\sigma_{12}\sigma_{34} \circ \left(L_4^\mathbf{a}(\mathbf{x}) \mathcal{I}_4^{(m),\mathbf{a}}(u, v) \right) = L_4^\mathbf{a}(\mathbf{x}) \mathcal{I}_4^{(m),\mathbf{a}}(u, v), \quad m = 1, 2. \quad (3.23)$$

Thus, the conformal integral can be represented as

$$\begin{aligned} I_4^\mathbf{a}(\mathbf{x}) &= \left(\mathbb{1} + \sigma_{23}\sigma_{14} \right) \circ L_4^\mathbf{a}(\mathbf{x}) \mathcal{I}_4^{(1),\mathbf{a}}(\boldsymbol{\eta}) \\ &= \left(\mathbb{1} + \sigma_{13}\sigma_{24} \right) \circ L_4^\mathbf{a}(\mathbf{x}) \mathcal{I}_4^{(1),\mathbf{a}}(\boldsymbol{\eta}). \end{aligned} \quad (3.24)$$

Here, a key concept is that the conformal integral in an appropriate coordinate domain can be generated by acting with particular permutations on a single function which will be called a *master function*.¹⁰

It should be noted that for $n > 4$ the kinematic groups are all trivial, $\mathcal{S}_n^{\text{kin}} = \{e\}$ [64]. This motivates the search for other subgroups of \mathcal{S}_n which also reconstruct the conformal integrals from a given set of master functions. For any n the problem looks quite complicated. Nonetheless, going back to the $n = 4$ case and bearing in mind that then all the Mellin-Barnes integrals involved into the bipartite representation can be calculated explicitly, an educated guess is to consider the cyclic group $\mathbb{Z}_4 \subset \mathcal{S}_4$. In the higher-point case, this naturally generalizes to picking $\mathbb{Z}_n \subset \mathcal{S}_n$ as a *generating group*: a subgroup of \mathcal{S}_n which allows reconstructing the full conformal integral from a given set of master functions.

3.2.2 Cyclic group

The cyclic group \mathbb{Z}_4 is

$$\mathbb{Z}_4 = \left\{ e, C_4, (C_4)^2, (C_4)^3 \right\} \subset \mathcal{S}_4, \quad (3.25)$$

where $C_4 = (1234)$ is a longest cycle. The longest cycle permutes the two cross-ratios

$$C_4 \circ u = v, \quad C_4 \circ v = u, \quad (3.26)$$

cf. (3.15). Hence, the cycles interchange domains $(u, v) \rightarrow (0, 1)$ and $(u, v) \rightarrow (1, 0)$. Note, however, that $u, v = 0$ is a *fixed point* of \mathbb{Z}_4 that suggests that we first analytically continue the first bare integral (3.14) from $(u, v) \rightarrow (0, 1)$ to $(u, v) \rightarrow (0, 0)$ and then consider cycles

¹⁰This observation is known in the literature on (non-)conformal integrals. E.g. a similar permutation generated representation is valid for the non-parametric hexagon integral with three massive corners [25] or for the non-parametric pentagon integral [15] (see also appendix D.2).

as generating permutations which reconstruct the full conformal integral near the origin on the (u, v) -plane.

Thus, we can apply the general analytic continuation formula (3.3) which in this case takes the form

$$\mathcal{I}_4^{(1),\mathbf{a}}(u, v) = U_4^{(0),\mathbf{a}}(u, v) + U_4^{(1),\mathbf{a}}(u, v), \quad (3.27)$$

where

$$\begin{aligned} U_4^{(0),\mathbf{a}}(u, v) &= S_4^{(234)} v^{a_1} F_4 \left[\begin{matrix} a_1, a'_3 \\ 1 - \alpha_{3,4}, 1 + \alpha_{1,4} \end{matrix} \middle| u, v \right], \\ U_4^{(1),\mathbf{a}}(u, v) &= S_4^{(134)} v^{a'_4} F_4 \left[\begin{matrix} a_2, a'_4 \\ 1 - \alpha_{3,4}, 1 - \alpha_{1,4} \end{matrix} \middle| u, v \right]. \end{aligned} \quad (3.28)$$

Here, F_4 is the fourth Appell function (B.13) which resulted from using the splitting identity (B.14), the coefficient $S_n^{(ijk)}$ is defined in (3.8).

Let us now consider how \mathbb{Z}_4 acts on $U_4^{(0),\mathbf{a}}$ and $U_4^{(1),\mathbf{a}}$. It turns out that these two terms are related by the cyclic permutation as follows

$$C_4 \circ \left(L_4^{\mathbf{a}}(\mathbf{x}) U_4^{(0),\mathbf{a}}(u, v) \right) = L_4^{\mathbf{a}}(\mathbf{x}) U_4^{(1),\mathbf{a}}(u, v), \quad (3.29)$$

see the general statement (3.10). Practically, this relation implies that there is only one *master* function which we choose to be $L_4^{\mathbf{a}}(\mathbf{x}) U_4^{(0),\mathbf{a}}(u, v)$. By construction, this new master function is supported near the origin $(u, v) = 0$. The cyclic group \mathbb{Z}_4 is of order four and hence its elements generate from the master function three more functions which are also the fourth Appell functions supported on the same domain. Introducing the collective notation one can represent the resulting four functions (including the master one) as follows

$$\begin{aligned} \varphi_4^{(234)}(\mathbf{x}) &:= L_4^{\mathbf{a}}(\mathbf{x}) U_4^{(0),\mathbf{a}}(u, v) = S_4^{(234)} V_4^{(234)}(\mathbf{x}) F_4 \left[\begin{matrix} a'_3, a_1 \\ 1 - \alpha_{3,4}, 1 - \alpha_{2,3} \end{matrix} \middle| u, v \right], \\ \varphi_4^{(134)}(\mathbf{x}) &:= (C_4)^1 \circ \varphi_4^{(234)}(\mathbf{x}) = S_4^{(134)} V_4^{(134)}(\mathbf{x}) F_4 \left[\begin{matrix} a_2, a'_4 \\ 1 - \alpha_{3,4}, 1 - \alpha_{1,4} \end{matrix} \middle| u, v \right], \\ \varphi_4^{(124)}(\mathbf{x}) &:= (C_4)^2 \circ \varphi_4^{(234)}(\mathbf{x}) = S_4^{(124)} V_4^{(124)}(\mathbf{x}) F_4 \left[\begin{matrix} a'_1, a_3 \\ 1 - \alpha_{1,2}, 1 - \alpha_{1,4} \end{matrix} \middle| u, v \right], \\ \varphi_4^{(123)}(\mathbf{x}) &:= (C_4)^3 \circ \varphi_4^{(234)}(\mathbf{x}) = S_4^{(123)} V_4^{(123)}(\mathbf{x}) F_4 \left[\begin{matrix} a'_2, a_4 \\ 1 - \alpha_{1,2}, 1 - \alpha_{2,3} \end{matrix} \middle| u, v \right], \end{aligned} \quad (3.30)$$

where $S_n^{(ijk)}$ are defined in (3.8), the leg-factors $V_4^{(ijk)}(\mathbf{x})$ are given by

$$\begin{aligned} V_4^{(234)}(\mathbf{x}) &= X_{13}^{-a_1} X_{23}^{\alpha_{14}} X_{34}^{\alpha_{12}} X_{24}^{-a'_3}, & V_4^{(134)}(\mathbf{x}) &= X_{24}^{-a_2} X_{34}^{\alpha_{12}} X_{14}^{\alpha_{23}} X_{13}^{-a'_4}, \\ V_4^{(124)}(\mathbf{x}) &= X_{13}^{-a_3} X_{14}^{\alpha_{23}} X_{12}^{\alpha_{34}} X_{24}^{-a'_1}, & V_4^{(123)}(\mathbf{x}) &= X_{24}^{-a_4} X_{12}^{\alpha_{34}} X_{23}^{\alpha_{14}} X_{13}^{-a'_2}, \end{aligned} \quad (3.31)$$

and any two of them are related by a cyclic permutation, e.g. $V_4^{(134)} = C_4 \circ V_4^{(234)}$ (this also holds true for $S_4^{(ijk)}$, e.g. $S_4^{(134)} = C_4 \circ S_4^{(234)}$). The functions $\varphi_4^{(ijk)}(\mathbf{x})$ will be called *basis* functions.

One can show that in terms of the basis functions both the first and second contributions to the conformal integral can be represented as

$$L_4^{\mathbf{a}}(\mathbf{x}) \mathcal{I}_4^{(1),\mathbf{a}}(u, v) = \varphi_4^{(234)}(\mathbf{x}) + \varphi_4^{(134)}(\mathbf{x}), \quad (3.32)$$

$$L_4^{\mathbf{a}}(\mathbf{x}) \mathcal{I}_4^{(2),\mathbf{a}}(u, v) = \varphi_4^{(124)}(\mathbf{x}) + \varphi_4^{(123)}(\mathbf{x}). \quad (3.33)$$

Here, the first line is just (3.27) in the new notation, while the second line results from analytically continuing the second bare integral (3.22) to the domain $(u, v) \rightarrow 0$ using the same continuation formula (B.25). In this way, one obtains yet another representation of the full conformal integral in terms of the generalized hypergeometric series generated from the master function by the cyclic group \mathbb{Z}_4 :

$$\begin{aligned} I_4^{\mathbf{a}}(\mathbf{x}) &= \left(\mathbb{1} + (C_4)^1 + (C_4)^2 + (C_4)^3 \right) \circ \varphi_4^{(234)}(\mathbf{x}) \\ &= \varphi_4^{(234)}(\mathbf{x}) + \varphi_4^{(134)}(\mathbf{x}) + \varphi_4^{(124)}(\mathbf{x}) + \varphi_4^{(123)}(\mathbf{x}). \end{aligned} \quad (3.34)$$

This expansion in basis functions is valid within the domain of convergence of the fourth Appell function $\sqrt{u} + \sqrt{v} < 1$ and reproduces the 4-point conformal integral in the form known in the literature.¹¹ Finally, note that the kinematic group considered in the previous section also acts on the basis functions (3.30) so that their sum (3.34) remains invariant, see appendix E.

3.2.3 Consistency checks

There are at least two possible non-trivial ways to test the resulting expressions. First, the most direct way to check the n -point parametric conformal integral is to reduce a number of points by one, i.e. $n \rightarrow n - 1$ along with $a_j \rightarrow 0$ for some j from 1 to n and then constrain the remaining propagator powers by the $(n - 1)$ -point conformality condition

$$\sum_{\substack{i=1 \\ i \neq j}}^n a_i = D, \quad (3.35)$$

¹¹The explicit calculation of the 4-point conformal integral has a long history which is discussed in detail in e.g. [35].

cf. (2.2). The result must be given by the $(n - 1)$ -point parametric conformal integral. Second, one can consider the non-parametric case i.e. choose all $a_i = 1$, and then compare the resulting expression with those known in the literature which were obtained by other methods. These two checks are described in detail below.¹²

Reduction to the star-triangle. From the very definition of the 4-point integral (2.1) it follows that if one of propagator powers equals zero, e.g. $a_4 = 0$, then the 4-point conformal integral reduces to the 3-point conformal integral:

$$I_4^a(\mathbf{x})\Big|_{a_4=0} = I_3^{a_1, a_2, a_3}(x_1, x_2, x_3). \quad (3.36)$$

Given the 4-point conformal integral calculated in the form (3.34) one can see from (3.30) that by setting $a_4 = 0$ all basis functions except $\varphi_4^{(123)}$ vanish due to $\Gamma(a_4 \rightarrow 0) \rightarrow \pm\infty$ in their denominators. The only non-vanishing contribution then leads to¹³

$$I_4^a(\mathbf{x})\Big|_{a_4=0} = \left(S_4^{(123)} V_4^{(123)}(\mathbf{x}) \right)\Big|_{a_4=0}, \quad (3.37)$$

where we used that the Appell function F_4 equals 1 if one of the upper parameters is 0, see (B.13). Substituting (3.8) and (3.31) one finds out that the left-hand side of (3.37) reproduces the star-triangle relation (2.21) provided that $a_1 + a_2 + a_3 = D$. The remaining limiting cases $a_i = 0$, $i = 1, 2, 3$, can be analyzed along the same lines. The only difference is which of the series (3.30) is non-zero in this limit. However, the numbering of basis functions is designed in such a way that when $a_l = 0$ is imposed, a basis function $\varphi_4^{(ijk)}$ survives which does not have l among upper indices, i.e. $i, j, k \neq l$.

Non-parametric box. For the 4-point conformal integral, the transition from the parametric to non-parametric case was considered in [35, 67]. Here, we reproduce a part of this analysis for completeness. The subtlety is that when choosing $a_i = 1$ in $D = 4$ (cf. (2.2)) the resulting integral diverges because the basis functions (3.30) contain divergent Γ -function prefactors. Introducing a cutoff parameter $\epsilon \rightarrow 0$ as

$$a_1 = a_2 = a_3 = 1, \quad a_4 = 1 - 2\epsilon \quad \Rightarrow \quad D = 4 - 2\epsilon, \quad (3.38)$$

and expanding the 4-point conformal integral (3.34) around $\epsilon = 0$ one finds

$$I_4^{1,1,1,1-2\epsilon}(\mathbf{x}) = \frac{1}{X_{13}X_{24}} \frac{\Phi(u, v)}{(1 - u - v)^2 - 4uv} + O(\epsilon), \quad (3.39)$$

¹²One may also consider conformal integrals as one of points goes to infinity, e.g. $x_n \rightarrow \infty$, that partially breaks conformal invariance. In this limit, the 4-point conformal integral is a particular case of the triangle integral evaluated by Boos and Davydychev [65, 66]. Note that conformal integrals with partially broken conformal invariance are instrumental within the shadow formalism for CFT on nontrivial backgrounds [44].

¹³Note that the product $V_4^{(ijk)} S_4^{(ijk)}$ can then be thought of as a generalization of the right-hand side of the star-triangle relation (2.21).

where the Bloch-Wigner function $\Phi(u, v)$ is expressed in terms of polylogarithms [67]:

$$\begin{aligned} \Phi(u, v) &= \frac{\pi^2}{3} + \ln u \ln v + \ln \frac{1+u-v-\lambda(u, v)}{2u} \ln \frac{1-u+v-\lambda(u, v)}{2v} \\ &+ 2 \ln \frac{1+u-v-\lambda(u, v)}{2u} + 2 \ln \frac{1-u+v-\lambda(u, v)}{2v} \\ &- 2 \operatorname{Li}_2 \frac{1+u-v-\lambda(u, v)}{2} - 2 \operatorname{Li}_2 \frac{1-u+v-\lambda(u, v)}{2}, \end{aligned} \quad (3.40)$$

where Li_2 is a dilogarithm. Introducing variables $u = z(1-y)$ and $v = y(1-z)$ and using the identity $\operatorname{Li}_2 z + \operatorname{Li}_2(1-z) = \pi^2/6 - \ln z \ln(1-z)$ the Bloch-Wigner function (3.40) can be cast into the form [35]

$$\Phi(z, y) = \ln(y(1-z)) \ln \frac{z}{1-y} + 2 \operatorname{Li}_2(1-z) - 2 \operatorname{Li}_2 y. \quad (3.41)$$

Then, the non-parametric box integral (3.39) can be given in a standard form [35, 67]

$$I_4^{1,1,1,1}(\mathbf{x}) = \frac{1}{X_{13} X_{24}} \frac{\Phi(z, y)}{1-z-y}. \quad (3.42)$$

3.2.4 Summary of the box integral reconstruction

Let us briefly formulate an emerging strategy of reconstructing a given conformal integral in some coordinate domain. We will keep n arbitrary assuming that the box integral example can be directly generalized to the higher-point case.

- Having calculated the first bare integral in the form of multivariate hypergeometric series (2.26), we analytically continue the resulting function from the original region to another region around the origin in the space of cross-ratios ($\mathbf{u}, \mathbf{w}, \mathbf{v} = 0$) by means of the analytic continuation formula (3.3) which splits $\mathcal{I}_n^{(1), \mathbf{a}}$ into $n-2$ additive terms $U_n^{(q), \mathbf{a}}$, $q = 0, 1, \dots, n-3$.
- Among these $n-2$ terms we single out $n-3$ terms which are not related through the action of the cyclic group (see (3.10))

$$\mathbb{Z}_n = \{e, C_n, (C_n)^2, \dots, (C_n)^{n-1}\} \subset \mathcal{S}_n, \quad (3.43)$$

where $C_n = (123\dots n) \in \mathcal{S}_n$ is a longest cycle.

- Combining cyclically independent functions with the leg-factor (2.8) one defines $n-3$ master functions

$$\begin{aligned} \varphi_n^{(q, n-1, n)}(\mathbf{x}) &= L_n^{\mathbf{a}}(\mathbf{x}) U_n^{(q), \mathbf{a}}, \quad q = 2, 3, \dots, n-3, \\ \varphi_n^{(n-2, n-1, n)}(\mathbf{x}) &= L_n^{\mathbf{a}}(\mathbf{x}) U_n^{(0), \mathbf{a}}. \end{aligned} \quad (3.44)$$

- The idea of reconstruction: the conformal integral $I_n^{\mathbf{a}}(\mathbf{x})$ is supposed to be given by a sum of functions obtained by acting with all elements of \mathbb{Z}_n on the master functions.

- The resulting expression can be checked in at least two ways:
 - The n -point conformal integral reduces to the $(n - 1)$ -point one upon setting one of propagator powers \mathbf{a} equal to zero

$$I_n^{a_1, \dots, a_{j-1}, a_j, a_{j+1}, \dots, a_n}(x_1, \dots, x_{j-1}, x_j, x_{j+1}, \dots, x_n) \Big|_{a_j=0} \quad (3.45)$$

$$= I_{n-1}^{a_1, \dots, a_{j-1}, a_{j+1}, \dots, a_n}(x_1, \dots, x_{j-1}, x_{j+1}, \dots, x_n),$$

for any $j = 1, \dots, n$. This reduction condition obviously follows from the very definition (2.1).

- Going to the non-parametric case, the resulting $I_n^{(1,1, \dots, 1)}(\mathbf{x})$ can be compared with expressions known in the literature, if any.

In the next sections 3.3 and 3.4 we examine how this scheme works for the pentagon and hexagon integrals and then in section 4 we will be able to formulate the general reconstruction conjecture.

3.3 Pentagon integral

For $n = 5$ the general formula (2.26) represents the first bare integral as the asymptotic expansion at $u_{1,2}, w_{12} \rightarrow 0$ and $v_{1,2} \rightarrow 1$:

$$\mathcal{I}_5^{(1), \mathbf{a}} = N_5^{\mathbf{a}} \Gamma(1 - \alpha_{4,5}, \alpha_{4,5}) v_1^{a_1} v_2^{a_2} \sum_{k_1, k_2, p_{12}=0}^{\infty} \frac{u_1^{k_1}}{k_1!} \frac{u_2^{k_2}}{k_2!} \frac{w_{12}^{p_{12}}}{p_{12}!}$$

$$\times \Gamma \left[\begin{matrix} a_1 + k_1 + p_{12}, a_2 + k_2 + p_{12}, a_3 + |\mathbf{k}|, a'_4 + |\mathbf{k}| + p_{12}, a'_5 + |\mathbf{k}| + p_{12} \\ a_1 + a_2 + a_3 + 2(|\mathbf{k}| + p_{12}), \quad 1 - \alpha_{4,5} + |\mathbf{k}| + p_{12} \end{matrix} \right] \quad (3.46)$$

$$\times F_1 \left[\begin{matrix} a'_4 + |\mathbf{k}| + p_{12}, a_1 + k_1 + p_{12}, a_2 + k_2 + p_{12} \\ a_1 + a_2 + a_3 + 2(|\mathbf{k}| + p_{12}) \end{matrix} \middle| 1 - v_1, 1 - v_2 \right],$$

where F_1 is the first Appell function (B.8). Among the respective cross-ratios, one is cubic and the others are quadratic:

$$u_1 = \frac{X_{13} X_{45}}{X_{14} X_{35}}, \quad u_2 = \frac{X_{23} X_{45}}{X_{24} X_{35}}, \quad w_{12} = \frac{X_{12} X_{34} X_{45}}{X_{14} X_{24} X_{35}}, \quad (3.47)$$

$$v_1 = \frac{X_{15} X_{34}}{X_{14} X_{35}}, \quad v_2 = \frac{X_{25} X_{34}}{X_{24} X_{35}}.$$

According to the general strategy outlined in the previous section, we can try to reconstruct the conformal integral from a set of master functions which can be found by particular analytic continuation of the first bare integral. The analytic continuation formula (3.3)-(3.5) splits the first bare integral (3.46) into three terms

$$\mathcal{I}_5^{(1), \mathbf{a}}(\boldsymbol{\eta}) = U_5^{(0), \mathbf{a}}(\boldsymbol{\eta}) + U_5^{(1), \mathbf{a}}(\boldsymbol{\eta}) + U_5^{(2), \mathbf{a}}(\boldsymbol{\eta}), \quad (3.48)$$

where

$$U_5^{(0),\mathbf{a}}(\boldsymbol{\eta}) = S_5^{(345)} v_1^{a_1} v_2^{a_2} \sum_{k_1, k_2, p_{12}, m_1, m_2=0}^{\infty} \frac{u_1^{k_1} u_2^{k_2} w_{12}^{p_{12}} v_1^{m_1} v_2^{m_2}}{k_1! k_2! p_{12}! m_1! m_2!} \times \frac{(a_1)_{k_1+p_{12}+m_1} (a_2)_{k_2+p_{12}+m_2} (a'_4)_{|\mathbf{k}|+p_{12}+|\mathbf{m}|} (-)^{p_{12}}}{(1-\alpha_{4,5})_{|\mathbf{k}|+p_{12}} (1-\alpha_{3,4})_{p_{12}+|\mathbf{m}|}}, \quad (3.49)$$

$$U_5^{(1),\mathbf{a}}(\boldsymbol{\eta}) = S_5^{(145)} v_1^{a'_5} \sum_{k_1, k_2, p_{12}, m_1, m_2=0}^{\infty} \frac{u_1^{k_1} (u_2 v_1 / v_2)^{k_2} (w_{12} / v_2)^{p_{12}} v_1^{m_1} (v_1 / v_2)^{m_2}}{k_1! k_2! p_{12}! m_1! m_2!} \times \frac{(a_2)_{k_2+p_{12}+m_2} (a_3)_{|\mathbf{k}|+m_1} (a'_5)_{|\mathbf{k}|+p_{12}+|\mathbf{m}|} (-)^{k_2}}{(1-\alpha_{4,5})_{|\mathbf{k}|+p_{12}} (1-\alpha_{1,5})_{k_2+|\mathbf{m}|}}, \quad (3.50)$$

$$U_5^{(2),\mathbf{a}}(\boldsymbol{\eta}) = S_5^{(245)} v_1^{a_1} v_2^{-\alpha_{15}} \sum_{k_1, k_2, p_{12}, m_1, m_2=0}^{\infty} \frac{u_1^{k_1} u_2^{k_2} (w_{12} / v_2)^{p_{12}} (v_1 / v_2)^{m_1} v_2^{m_2}}{k_1! k_2! p_{12}! m_1! m_2!} \times \frac{(a_1)_{k_1+p_{12}+m_1} (a_3)_{|\mathbf{k}|+m_2} (-\alpha_{1,5})_{m_2+k_2-m_1} (-)^{p_{12}}}{(1-\alpha_{4,5})_{|\mathbf{k}|+p_{12}} (1+\alpha_{3,4})_{m_2-m_1-p_{12}}}. \quad (3.51)$$

These functions are supported near the origin of coordinates (3.2) such that the cross-ratios satisfy the condition (3.9).

3.3.1 Cyclic group

The cyclic group is $\mathbb{Z}_5 = \{e, C_5, (C_5)^2, (C_5)^3, (C_5)^4\} \subset \mathcal{S}_5$, where $C_5 = (12345)$ is a longest cycle. The general statement (3.10) about the relation of two terms in the analytic continuation formula for the first bare integral now reads as

$$C_5 \circ \left(L_5^{\mathbf{a}}(\mathbf{x}) U_5^{(0),\mathbf{a}}(\boldsymbol{\eta}) \right) = L_5^{\mathbf{a}}(\mathbf{x}) U_5^{(1),\mathbf{a}}(\boldsymbol{\eta}), \quad (3.52)$$

where the 5-point leg-factor (2.8) is given by

$$L_5^{\mathbf{a}}(\mathbf{x}) = X_{15}^{-a_1} X_{25}^{-a_2} X_{34}^{-a'_5} X_{35}^{-\alpha_{3,5}} X_{45}^{-\alpha_{4,5}}. \quad (3.53)$$

It means that among three functions in (3.48) there are two which are not related by cyclic permutations, e.g. these are $U_5^{(0),\mathbf{a}}$ and $U_5^{(2),\mathbf{a}}$. Then, adding the leg-factor we introduce two master functions as

$$\begin{aligned} \varphi_5^{(345)}(\mathbf{x}) &:= L_5^{\mathbf{a}}(\mathbf{x}) U_5^{(0),\mathbf{a}}(\boldsymbol{\eta}) = S_5^{(345)} V_5^{(345)}(\mathbf{x}) P_1 \left[\begin{array}{c|c} a_1, a_2, a'_4 & u_1, w_{12}, u_2, v_1, v_2 \\ \hline 1 - \alpha_{45}, 1 - \alpha_{34} & \end{array} \right], \\ \varphi_5^{(245)}(\mathbf{x}) &:= L_5^{\mathbf{a}}(\mathbf{x}) U_5^{(2),\mathbf{a}}(\boldsymbol{\eta}) = S_5^{(245)} V_5^{(245)}(\mathbf{x}) P_2 \left[\begin{array}{c|c} a_1, a_3, -\alpha_{15} & u_1, \frac{w_{12}}{v_2}, u_2, \frac{v_1}{v_2}, v_2 \\ \hline 1 - \alpha_{45}, 1 + \alpha_{34} & \end{array} \right], \end{aligned} \quad (3.54)$$

where we defined two generalized hypergeometric functions P_1 (D.1) and P_2 (D.2) as well as two leg-factors

$$\begin{aligned} V_5^{(345)}(\mathbf{x}) &= X_{14}^{-a_1} X_{24}^{-a_2} X_{45}^{-\alpha_{45}} X_{34}^{-\alpha_{34}} X_{35}^{-a'_4}, \\ V_5^{(245)}(\mathbf{x}) &= X_{14}^{-a_1} X_{35}^{-a_3} X_{24}^{-\alpha_{15}} X_{45}^{-\alpha_{45}} X_{25}^{\alpha_{34}}, \end{aligned} \tag{3.55}$$

cf. (3.31).

The cyclic group \mathbb{Z}_5 is of order five so that one can construct four more basis functions from each of the master functions. In this way, we obtain *ten* basis functions which are listed in appendix D. Note that our notation for the basis functions is designed in such a way that the action of a cyclic permutation on the master function $\varphi_5^{(ijk)}$ cyclically permutes the upper indices i, j, k . In particular, it follows that the two master functions (3.54) as well as prefactors (3.55) are not related to each other since there are no cyclic permutations which turn 345 into 245. Thus, the set of basis functions is split into two subsets of cyclically related functions, i.e. into two representations of \mathbb{Z}_5 .

Then, the 5-point parametric conformal integral is represented as the \mathbb{Z}_5 -invariant sum of ten basis functions:

$$\begin{aligned} I_5^{\mathbf{a}}(\mathbf{x}) &= \sum_{j=0}^4 (C_5)^j \circ \left(\varphi_5^{(345)}(\mathbf{x}) + \varphi_5^{(245)}(\mathbf{x}) \right) \\ &= \varphi_5^{(345)}(\mathbf{x}) + \varphi_5^{(145)}(\mathbf{x}) + \varphi_5^{(125)}(\mathbf{x}) + \varphi_5^{(123)}(\mathbf{x}) + \varphi_5^{(234)}(\mathbf{x}) \\ &\quad + \varphi_5^{(245)}(\mathbf{x}) + \varphi_5^{(135)}(\mathbf{x}) + \varphi_5^{(124)}(\mathbf{x}) + \varphi_5^{(235)}(\mathbf{x}) + \varphi_5^{(134)}(\mathbf{x}). \end{aligned} \tag{3.56}$$

The expansion is defined in a domain near the origin in the space of cross-ratios. By construction, both sides here are invariant under cyclic permutations.

Note that one can define a generalization of the kinematic group which we call an *extended* kinematic group. Given two master functions (3.54), such a group generates the same set of basis functions thereby providing an equivalent way of reconstructing the conformal pentagon integral. Details are given in appendix E.

3.3.2 Reduction to the box integral

In appendix D.2 we check that going to the non-parametric case we indeed do reproduce the formula for the non-parametric pentagon integral known in the literature [15]. Of course, we can compare only in a given coordinate domain. Below we examine the reduction to the box integral.

Setting one of propagator powers equal to zero, e.g. $a_5 = 0$, the pentagon integral has to reduce to the box integral:

$$I_5^{\mathbf{a}}(\mathbf{x}) \Big|_{a_5=0} = I_4^{a_1, a_2, a_3, a_4}(x_1, x_2, x_3, x_4). \tag{3.57}$$

In fact, it is quite easy to verify this relation provided that $I_5^{\mathbf{a}}(\mathbf{x})$ is represented as (3.56) At $a_5 = 0$ only those basis functions $\varphi_5^{(ijk)}$ survive which upper indices $i, j, k \neq 5$, while

the remaining basis functions vanish due to the presence of $\Gamma(a_5 \rightarrow 0) \rightarrow \pm\infty$ in their denominators, see (3.8). As a result,

$$I_5^{\mathbf{a}} \Big|_{a_5=0} = \left(\varphi_5^{(123)}(\mathbf{x}) + \varphi_5^{(234)}(\mathbf{x}) + \varphi_5^{(124)}(\mathbf{x}) + \varphi_5^{(134)}(\mathbf{x}) \right) \Big|_{a_5=0}. \quad (3.58)$$

The basis function are written in terms of functions (D.1) and (D.2) which are reduced to the fourth Appell function (B.13) when one of the upper parameters equals zero, i.e.

$$P_1 \left[\begin{matrix} 0, A, B \\ C_1, C_2 \end{matrix} \middle| \boldsymbol{\xi} \right] = P_2 \left[\begin{matrix} 0, A, B \\ C_1, C_2 \end{matrix} \middle| \boldsymbol{\xi} \right] = F_4 \left[\begin{matrix} A, B \\ C_1, C_2 \end{matrix} \middle| \xi_3, \xi_5 \right]. \quad (3.59)$$

On the other hand, the prefactors $S_5^{(ijk)} V_5^{(ijk)}(\mathbf{x})$ are reduced as

$$\begin{aligned} S_5^{(123)} V_5^{(123)} \Big|_{a_5=0} &= S_4^{(123)} V_4^{(123)}, & S_5^{(234)} V_5^{(234)} \Big|_{a_5=0} &= S_4^{(234)} V_4^{(234)}, \\ S_5^{(124)} V_5^{(124)} \Big|_{a_5=0} &= S_4^{(124)} V_4^{(124)}, & S_5^{(134)} V_5^{(134)} \Big|_{a_5=0} &= S_4^{(134)} V_4^{(134)}. \end{aligned} \quad (3.60)$$

Gathering (3.60) and (3.59) together and taking into account the symmetry properties of $P_{1,2}$ (D.3)-(D.4), one can see that the right-hand side of the reduced pentagon integral (3.58) reproduces the right-hand side of the box integral (3.34) so that the desired reduction condition (3.57) holds true.

The other cases $a_l = 0$ for $l = 1, 2, 3, 4$ can be analyzed in a similar way. The only difference is which of basis functions $\varphi_5^{(ijk)}$ survive at $a_l = 0$, but our way of numbering basis functions immediately indicates that these are functions with upper indices $i, j, k \neq l$. Thus, our expression for the pentagon integral (3.56) consistently reduces to the box integral formula (3.34).

3.4 Hexagon integral

Let us finally examine the reconstruction idea for the 6-point parametric conformal integral. The first bare integral (2.26) is given by

$$\begin{aligned} \mathcal{I}_6^{(1),\mathbf{a}} &= N_6^{\mathbf{a}} \Gamma(1 - \alpha_{5,6}, \alpha_{5,6}) v_1^{a_1} v_2^{a_2} v_3^{a_3} \sum_{\{k_l, p_{ij}\}=0}^{\infty} \frac{u_1^{k_1}}{k_1!} \frac{u_2^{k_2}}{k_2!} \frac{u_3^{k_3}}{k_3!} \frac{w_{12}^{p_{12}}}{p_{12}!} \frac{w_{13}^{p_{13}}}{p_{13}!} \frac{w_{23}^{p_{23}}}{p_{23}!} \\ &\times \Gamma \left[\begin{matrix} a_4 + |\mathbf{k}|, a'_5 + |\mathbf{k}| + |\mathbf{p}|, a'_6 + |\mathbf{k}| + |\mathbf{p}|, \mathcal{B}_1, \mathcal{B}_2, \mathcal{B}_3 \\ 1 - \alpha_{5,6} + |\mathbf{k}| + |\mathbf{p}|, \mathcal{C} \end{matrix} \right] \\ &\times F_D^{(3)} \left[\begin{matrix} a'_5 + |\mathbf{k}| + |\mathbf{p}|, \mathcal{B}_1, \mathcal{B}_2, \mathcal{B}_3 \\ \mathcal{C} \end{matrix} \middle| 1 - v_1, 1 - v_2, 1 - v_3 \right], \end{aligned} \quad (3.61)$$

where $|\mathbf{k}| = k_1 + k_2 + k_3$, $|\mathbf{p}| = p_{12} + p_{13} + p_{23}$, and the parameters are encoded as

$$\begin{aligned} \mathcal{B}_1 &= a_1 + k_1 + p_{12} + p_{13}, & \mathcal{B}_2 &= a_2 + k_2 + p_{12} + p_{23}, & \mathcal{B}_3 &= a_3 + k_3 + p_{13} + p_{23}, \\ \mathcal{C} &= a_1 + a_2 + a_3 + a_4 + 2(|\mathbf{k}| + |\mathbf{p}|). \end{aligned} \quad (3.62)$$

The nine cross-ratios in (3.61) are

$$\begin{aligned} u_1 &= \frac{X_{14}X_{56}}{X_{15}X_{46}}, & u_2 &= \frac{X_{24}X_{56}}{X_{25}X_{46}}, & u_3 &= \frac{X_{34}X_{56}}{X_{35}X_{46}}, \\ w_{12} &= \frac{X_{12}X_{45}X_{56}}{X_{15}X_{25}X_{46}}, & w_{13} &= \frac{X_{13}X_{45}X_{56}}{X_{15}X_{35}X_{46}}, & w_{23} &= \frac{X_{23}X_{45}X_{56}}{X_{25}X_{35}X_{46}}, \\ v_1 &= \frac{X_{16}X_{45}}{X_{15}X_{46}}, & v_2 &= \frac{X_{26}X_{45}}{X_{25}X_{46}}, & v_3 &= \frac{X_{36}X_{45}}{X_{35}X_{46}}. \end{aligned} \quad (3.63)$$

In order to reconstruct the full hexagon conformal integral we represent the first bare integral (3.61) near $\boldsymbol{\eta} = \mathbf{0}$ by means of the analytic continuation formula (3.3):

$$\mathcal{I}_6^{(1),\mathbf{a}}(\boldsymbol{\eta}) = \mathcal{U}_6^{(0),\mathbf{a}}(\boldsymbol{\eta}) + \mathcal{U}_6^{(1),\mathbf{a}}(\boldsymbol{\eta}) + \mathcal{U}_6^{(2),\mathbf{a}}(\boldsymbol{\eta}) + \mathcal{U}_6^{(3),\mathbf{a}}(\boldsymbol{\eta}), \quad (3.64)$$

where each term can be read off from (3.4)-(3.5). According to the general formula (3.10) two of four terms here are related as

$$C_6 \circ \left(L_6^{\mathbf{a}}(\mathbf{x}) \mathcal{U}_6^{(0),\mathbf{a}}(\boldsymbol{\eta}) \right) = L_6^{\mathbf{a}}(\mathbf{x}) \mathcal{U}_6^{(1),\mathbf{a}}(\boldsymbol{\eta}), \quad (3.65)$$

where $C_6 = (123456) \in \mathcal{S}_6$. Again, $\mathcal{U}_6^{(0),\mathbf{a}}$, $\mathcal{U}_6^{(2),\mathbf{a}}$, $\mathcal{U}_6^{(3),\mathbf{a}}$ can be chosen as master functions, while the corresponding basis functions $\varphi_6^{(ijk)}$ are generated from them by cycle permutations. Since the respective cyclic group \mathbb{Z}_6 is of order six, then the three master functions generate a set of 18 basis functions.

At this point, we could suggest that the 6-point conformal integral is to be constructed as the \mathbb{Z}_6 -invariant sum of 18 basis functions. If so, one can consider a reduction to the pentagon integral which in its turn has 10 terms (3.56). However, it can be shown that 18 basis functions of the hexagon integral will suffice to reproduce only 9 basis functions of the pentagon integral. This happens if any of the hexagon propagator powers is set to zero, $a_l = 0$ for some $l = 1, \dots, 6$.

Assuming that we are correct in our conjecture of reconstruction this means that there should be additional master functions which are not directly seen from the analytic continuation formula for the first bare integral (3.64). Indeed, one can show that available master and basis functions turn out to possess a number of structural properties which allow one to introduce and systematically describe a complete set of functions needed for reconstruction [48]. This set includes both already known functions and those that are missing, as in the hexagon case discussed above.

4 Reconstruction conjecture

Summing up our discussion in the previous sections we can now describe the reconstruction procedure in more detail.

- *Bipartite representation.* By formal identical transformations the conformal integral $I_n^\alpha(\mathbf{x})$ is split into two integral parts that defines the bipartite Mellin-Barnes representation:

$$I_n^\alpha(\mathbf{x}) = L_n^\alpha(\mathbf{x}) \mathcal{I}_n^{(1),\alpha}(\boldsymbol{\eta}) + L_n^\alpha(\mathbf{x}) \mathcal{I}_n^{(2),\alpha}(\boldsymbol{\eta}). \quad (4.1)$$

The first term here is found in the form of multivariate generalized hypergeometric series (2.26), while the second term is represented through the $(n-3)(n-2)/2$ -folded Mellin-Barnes integral (2.32) with a complex structure of poles, which currently is not known in analytic form. It is the reconstruction procedure that suggests avoiding any further integration because the second bare integral can be recovered from the first one by invoking the permutation invariance of the full conformal integral.

- *Coordinate domains.* The function $I_n^\alpha(\mathbf{x})$ is defined on a domain $D_n \subset (\mathbb{R}^D \times)^n$, which can be found by first calculating the conformal integral explicitly in a particular coordinate domain and then using analytic continuation formulas.¹⁴

The two terms in (4.1) are supported on their own domains which are generally wider than D_n . The domain of the first term $D_n^{(1)}$ is such that $D_n \subset D_n^{(1)}$, it is also hard to identify explicitly. We denote the respective domain in the space of cross-ratios \mathcal{H}_n as $\widehat{D}_n^{(1)} \subset \mathcal{H}_n$, i.e. $\widehat{D}_n^{(1)} = \boldsymbol{\eta}(D_n^{(1)})$.

- There is a smaller domain \widehat{A}_n which is the convergence region of $\mathcal{I}_n^{(1),\alpha}$ calculated as a multivariate power series near a particular point:

$$\widehat{D}_n^{(1)} \supset \widehat{A}_n \ni (\mathbf{u} = 0, \mathbf{w} = 0, \mathbf{v} = 1). \quad (4.2)$$

- Using the known analytic continuation formulas one can define $\mathcal{I}_n^{(1),\alpha}$ on a different domain \widehat{B}_n containing the origin of coordinates in \mathcal{H}_n :

$$\widehat{D}_n^{(1)} \supset \widehat{B}_n \ni (\mathbf{u} = 0, \mathbf{w} = 0, \mathbf{v} = 0). \quad (4.3)$$

- In fact, an exact shape of either domain remains unclear. Instead, we operate with small enough neighbourhoods of concrete points in \mathcal{H}_n like $0, 1, \infty$.

- *Symmetric group.* By construction, the conformal integral is invariant under action of the symmetric group

$$\forall \pi \in \mathcal{S}_n : \quad \mathbf{R}(\pi) \circ I_n^\alpha(\mathbf{x}) = I_n^\alpha(\mathbf{x}). \quad (4.4)$$

¹⁴This can be contrasted with other possible calculation schemes such as the geometric approach where the conformal integral is given by volume of a simplex in a space of constant curvature which sign depends on a particular kinematic regime, i.e. on a particular coordinate domain, see e.g. [16].

There are maps of the (co)domains induced by elements of the symmetric group \mathcal{S}_n acting on $\mathbf{x} = (x_1, \dots, x_n)$. In the space of cross-ratios \mathcal{H}_n the symmetric group acts by transformations

$$\forall \pi \in \mathcal{S}_n : \quad \boldsymbol{\eta} \rightarrow \boldsymbol{\eta}' = \mathbf{R}(\pi) \circ \boldsymbol{\eta} = H_\pi(\boldsymbol{\eta}), \quad (4.5)$$

where H_π are homogeneous rational functions. One can single out various subgroups of \mathcal{S}_n which act in \mathcal{H}_n in some special way.

- E.g., one can consider a stabilizer (kinematic) group,

$$\mathcal{S}_n^{\text{kin}} := \{\pi \in \mathcal{S}_n : \mathbf{R}(\pi) \circ \boldsymbol{\eta} = \boldsymbol{\eta}, \forall \boldsymbol{\eta} \in \mathcal{H}_n\} \subset \mathcal{S}_n. \quad (4.6)$$

It is clear that the stabilizer leaves any domain in \mathcal{H}_n invariant. However, one can show that the stabilizers are all trivial for $n > 4$ [64].

- Other subgroups change chosen domains. Given a domain $\widehat{X}_n \subset \mathcal{H}_n$ one can consider a subgroup

$$\mathcal{G}_n \subset \mathcal{S}_n : \quad \widehat{X}'_n = \mathbf{R}(\mathcal{G}_n) \circ \widehat{X}_n \quad \text{such that} \quad \widehat{X}_n \cap \widehat{X}'_n \neq \emptyset. \quad (4.7)$$

E.g., there is a cyclic subgroup generated by $C_n = (123\dots n) \in \mathcal{S}_n$, i.e.

$$\mathbb{Z}_n = \{e, C_n, (C_n)^2, \dots, (C_n)^{n-1}\} \subset \mathcal{S}_n. \quad (4.8)$$

- One may consider extended kinematic groups $\widehat{\mathcal{S}}_n^{\text{kin}}$ which are nontrivial for $n > 4$ and coincide with $\mathcal{S}_4^{\text{kin}}$ when $n = 4$ (see appendix E). However, at our current level of understanding, $\widehat{\mathcal{S}}_n^{\text{kin}}$ are derived objects, i.e. these subgroups can be defined only in terms of the cyclic group and its action on basis functions that is not useful in practice.
- *Basis functions.* One can choose a set of *basis* functions $\varphi_n^{(ijk)}$ supported on a common domain $\widehat{X}_n \subset \widehat{D}_n$. The considered cases $n = 4, 5, 6$ allow us to emphasize that the basis functions share a number of common features:
 - Γ -function prefactor $S_n^{(ijk)}$;
 - Leg-factor $V_n^{(ijk)}(\mathbf{x})$;
 - Hypergeometric-type power series in $n(n-3)/2$ cross-ratios.

Moreover, each of these elements and, therefore, a given basis function is determined by choosing 3 pairs (x_i, a_i) out of n such pairs.

- *Master functions.* The conformal integral $I_n^{\mathbf{a}}(\mathbf{x})$ must be invariant under the symmetric group \mathcal{S}_n . However, the basis functions $\varphi_n^{(ijk)}(\mathbf{x})$ are supported on a particular coordinate domain that implies that the full symmetric group is broken down to a particular *generating* subgroup of the type (4.7) which acts on the basis functions in a specific

way: among all basis functions one can single out a number of *master* functions which generate all other basis functions by acting with elements of the generating group. By construction, the generating group reshuffles a set of basis functions. It is important to stress that a set of basis functions depends on the choice of a coordinate domain that in its turn defines a suitable generating group.

- *Consistency checks.* In order to check the resulting expressions for the conformal integrals obtained by the reconstruction method one can compare them with those known in the literature. There are two possible checks: (1) reducing a number of points, $n \rightarrow n - 1$; (2) going to the non-parametric case, $a_i \rightarrow 1$.

If one chooses the domain as $\widehat{X}_n = \widehat{B}_n$ and the generating group as \mathbb{Z}_n , then the conformal integral can explicitly be evaluated by means of the following

Reconstruction conjecture. *The n -point conformal integral (2.1) is the \mathbb{Z}_n -invariant sum of master functions*

$$I_n^\alpha(\mathbf{x}) \stackrel{\text{nr}}{=} \sum_{m=0}^{n-1} (C_n)^m \circ \sum_{(ijk) \in \mathbb{T}_n} \varphi_n^{(ijk)}(\mathbf{x}), \quad (4.9)$$

where \mathbb{T}_n is a set of index triples (ijk) which are not related to each other by cyclic permutations, $1 \leq i < j < k \leq n$. A cardinal number of \mathbb{T}_n equals the number of master functions,

$$|\mathbb{T}_n| = \begin{cases} \frac{(n-2)(n-1)}{3!}, & \text{if } \frac{n}{3} \notin \mathbb{N}, \\ \frac{(n-2)(n-1) + 4}{3!}, & \text{if } \frac{n}{3} \in \mathbb{N}. \end{cases} \quad (4.10)$$

A total number of basis functions is equal to the binomial coefficient

$$\binom{n}{3} = \frac{n(n-1)(n-2)}{3!}, \quad (4.11)$$

which determines a number of all possible index triples. The symbol $\stackrel{\text{nr}}{=}$ in (4.9) means that among all basis functions produced by acting with cycles on master functions one keeps only non-repeating ones which number equals (4.11).

As drafted, this conjecture shifts the focus to finding master functions as well as describing their properties. A few comments are in order.

- Some of master functions are contained directly in the calculated part $L_n^\alpha(\mathbf{x}) \mathcal{I}_n^{(1),\alpha}$. Since among $n - 2$ candidate master functions delivered by the analytic continuation formula (3.3) the first two functions are cyclically dependent (see (3.10)), there remain $n - 3$ master functions which we explicitly know from evaluating the first bare integral. However, a number of master functions is conjectured to grow quadratically (4.10) and $n = 6$ is the first time when the $n - 3$ functions turn out to be insufficient for the full reconstruction.

- The number of master and basis functions (4.10) and (4.11) in the considered cases $n = 4, 5, 6$ equals 1, 2, 4 and 4, 10, 20, respectively. In the hexagon case we see that in addition to 18 basis functions coming from the first bare integral there are two more basis functions produced from the additional master function¹⁵ and now their total number is sufficient to reproduce the pentagon integral when checking $n \rightarrow n - 1$ reduction.
- We claim that unknown part of master functions can be derived from already known basis functions. The point is that the proposed parameterization by index triples allows one to reveal a number of remarkable properties of master/basis functions which can be effectively described by means of some diagrammatics. This will enable us eventually to build a complete set of functions explicitly, thereby reconstructing the full conformal integral in a given coordinate domain [48].

5 Conclusion

In this paper we have proposed a new method of calculating n -point parametric one-loop conformal integrals and elaborated two examples of the box and pentagon (non)-parametric integrals. Essentially, there are three main points: (1) a bipartite Mellin-Barnes representation which allows one to evaluate an additive part of the conformal integral explicitly; (2) using asymptotic expansions near particular points and analytic continuation on other domains; (3) permutation invariance can be used to recover the remaining unknown part.

This reconstruction procedure operates with a set of master functions which generate a wider set of basis functions by means of permutations from the cyclic group. The resulting expression for the conformal integral supported on a particular coordinate domain is given simply by summing all basis functions. For the box and pentagon integrals we have shown that the basis functions are directly defined from the explicitly found part of the bipartite representation. For the hexagon and higher-point integrals a complete set of basis functions will be described in our forthcoming paper [48] in which will finalize the reconstruction method.

It would be natural to find an appropriate modification of the reconstruction method for at least two types of Feynman integrals possessing conformal symmetry. These are multi-loop higher-point conformal integrals which are currently being considered by other methods, see e.g. [28, 68–71] as well as massive conformal integrals studied in [16, 72, 73]. The latter have recently been shown to play an essential role when calculating contact Witten diagrams [74].

Besides possible generalizations, the already obtained results can be immediately applied in CFT_D by means of the shadow formalism (see our discussion in the Introduction). Especially, the one-loop conformal integrals are useful in CFT_D on non-trivial backgrounds, where one can show that the problem of calculating n -point thermal conformal blocks boils down to knowing $(2n + 2)$ -point one-loop conformal integrals explicitly (see [44] for 1-point thermal blocks).

¹⁵This means that the respective representation of the cyclic group \mathbb{Z}_6 on basis functions is two-dimensional, and not six-dimensional as in the case of other hexagon master functions. Such a shortening happens every time when $n/3 \in \mathbb{N}$.

Acknowledgements. We would like to thank Leonid Bork, Sergey Derkachev, Alexey Isaev for discussions about related topics and Marcus Spradlin for correspondence. S.M. is also very grateful to Ekaterina Semenova for her support at all stages of this project as well as for her help in drawing diagrams.

Our work was supported by the Foundation for the Advancement of Theoretical Physics and Mathematics “BASIS”.

A Notation and conventions

Let $\mathbf{g} = \{g_1, \dots, g_N\}$ denote a set of elements such as parameters of functions, powers, integration variables, etc. Their various sums will be denoted as

$$|\mathbf{g}_{i,j}| = \sum_{l=i}^j g_l, \quad |\mathbf{g}| = \sum_{l=1}^N g_l, \quad |\mathbf{g}_q| = |\mathbf{g}_{q,N}| - |\mathbf{g}_{1,q-1}|. \quad (\text{A.1})$$

In the case of propagator powers $\mathbf{a} = (a_1, \dots, a_n)$ we also use

$$|\mathbf{a}_{i,j}| = \begin{cases} \sum_{l=i}^j a_l, & i < j, \\ \sum_{l=i}^n a_l + \sum_{l=1}^j a_l, & i > j, \end{cases} \quad a'_i = \frac{D}{2} - a_i, \quad \alpha_{i,j} = a_i + a_j - \frac{D}{2} \equiv a_i - a'_j, \quad (\text{A.2})$$

where D is the dimension of Euclidean space \mathbb{R}^D .

The products of Γ -functions are denoted as

$$\Gamma \left[\begin{matrix} a_1, \dots, a_M \\ b_1, \dots, b_K \end{matrix} \right] = \frac{\Gamma(a_1, \dots, a_M)}{\Gamma(b_1, \dots, b_K)}, \quad \Gamma(a_1, \dots, a_M) = \prod_{i=1}^M \Gamma(a_i). \quad (\text{A.3})$$

In the Mellin-Barnes integrals it is convenient to use the following modified measure:

$$\widehat{dt} = \frac{dt}{2\pi i} \Gamma(-t). \quad (\text{A.4})$$

B Generalized hypergeometric functions

In this appendix we discuss various generalized hypergeometric functions focusing on their integral representations, identities as well as analytic continuation formulas. We closely follow the review [63]. See also [49, 62, 75, 76].

B.1 General definitions

Following the Horn's classification scheme, a N -point multivariate power series

$$\sum_{m_1, \dots, m_N = -\infty}^{+\infty} A(m_1, \dots, m_N) \xi_1^{m_1} \dots \xi_N^{m_N} \equiv \sum_{\mathbf{m} \in \mathbb{Z}^N} A(\mathbf{m}) \boldsymbol{\xi}^{\mathbf{m}} \quad (\text{B.1})$$

is a generalized hypergeometric function if

$$\frac{A(\mathbf{m} + \mathbf{e}_j)}{A(\mathbf{m})} = \frac{P_j(\mathbf{m})}{Q_j(\mathbf{m})}, \quad \forall j = 1, \dots, n, \quad (\text{B.2})$$

where $P_j(\mathbf{m})$ and $Q_j(\mathbf{m})$ are some polynomials in $\mathbf{m} = \{m_1, \dots, m_N\}$ and $\mathbf{e}_j = \{0, \dots, 1, \dots, 0\}$ is a vector with all components equal to zero except j -th one. In other words, a given generalized hypergeometric function (B.1) have expansion coefficients such that their ratios are rational functions of parameters.

The relation (B.2) treated as an equation for $A(m_1, \dots, m_N)$ admits many solutions. In particular, when

$$P_j(\mathbf{m}) = (a + |\mathbf{m}|)(b_j + m_j), \quad Q_j(\mathbf{m}) = (c + |\mathbf{m}|)(1 + m_j), \quad (\text{B.3})$$

where $\mathbf{m}, \mathbf{b}, a, c$ are constant complex-valued parameters (see (A.1) for notation), the solution to (B.2) is the N -point Lauricella function D :

$$F_D^{(N)} \left[\begin{matrix} a, \mathbf{b} \\ c \end{matrix} \middle| \boldsymbol{\xi} \right] = \sum_{|\mathbf{m}|=0}^{\infty} \frac{(a)_{|\mathbf{m}|} (\mathbf{b})_{\mathbf{m}} \boldsymbol{\xi}^{\mathbf{m}}}{(c)_{|\mathbf{m}|} m!}, \quad (\text{B.4})$$

which is convergent in the unit polydisk

$$\mathbb{U}^N = \{ \boldsymbol{\xi} \in \mathbb{C}^N : |\xi_j| < 1, j = 1, \dots, N \}. \quad (\text{B.5})$$

The (rising) factorials in (B.4) are defined as

$$(a)_m = \frac{\Gamma(a+m)}{\Gamma(a)}, \quad (\mathbf{b})_{\mathbf{m}} = (b_1)_{m_1} \dots (b_N)_{m_N}, \quad \mathbf{m}! = m_1! \dots m_N!. \quad (\text{B.6})$$

At $N = 1$ and $N = 2$ the Lauricella function D (B.4) becomes the Gauss hypergeometric function ${}_2F_1$ and the first Appell function F_1 , respectively:

$${}_2F_1 \left[\begin{matrix} a, b \\ c \end{matrix} \middle| \xi \right] = \sum_{m=0}^{\infty} \frac{(a)_m (b)_m \xi^m}{(c)_m m!}, \quad (\text{B.7})$$

$$F_1 \left[\begin{matrix} a, b_1, b_2 \\ c \end{matrix} \middle| \xi_1, \xi_2 \right] = \sum_{m_1, m_2=0}^{\infty} \frac{(a)_{m_1+m_2} (b_1)_{m_1} (b_2)_{m_2} \xi_1^{m_1} \xi_2^{m_2}}{(c)_{m_1+m_2} m_1! m_2!}. \quad (\text{B.8})$$

The Lauricella function $F_D^{(N)}$ belongs to the family of generalized hypergeometric functions $G^{(N,j)}$ labelled by $j = 1, \dots, N + 1$ (see (A.1) for notation):

$$G^{(N,j)} \left[\begin{matrix} a, \mathbf{b} \\ c \end{matrix} \middle| \boldsymbol{\xi} \right] = \sum_{|\mathbf{m}|=0}^{\infty} \frac{(a)_{|\mathbf{m}_j|} (\mathbf{b})_{\mathbf{m}} \boldsymbol{\xi}^{\mathbf{m}}}{(c)_{|\mathbf{m}_j|} \mathbf{m}!}, \quad (\text{B.9})$$

which converge in the unit polydisk (B.5). The subscripts of the Pochhammer symbols in (B.9) can be negative, in which case the Pochhammer symbol reads as

$$(a)_{-m} = \frac{(-1)^m}{(1-a)_m}, \quad m \in \mathbb{Z}^+. \quad (\text{B.10})$$

The Lauricella function $F_D^{(N)}$ follows from (B.9) at $j = 1$ and $j = N + 1$:

$$F_D^{(N)} \left[\begin{matrix} a, \mathbf{b} \\ c \end{matrix} \middle| \boldsymbol{\xi} \right] = G^{(N,1)} \left[\begin{matrix} a, \mathbf{b} \\ c \end{matrix} \middle| \boldsymbol{\xi} \right], \quad F_D^{(N)} \left[\begin{matrix} 1-c, \mathbf{b} \\ 1-a \end{matrix} \middle| \boldsymbol{\xi} \right] = G^{(N,N+1)} \left[\begin{matrix} a, \mathbf{b} \\ c \end{matrix} \middle| \boldsymbol{\xi} \right]. \quad (\text{B.11})$$

The Horn function G_2 follows from (B.9) at $j = 2$ and $N = 2$:

$$\begin{aligned} G^{(2,2)} \left[\begin{matrix} a, b_1, b_2 \\ c \end{matrix} \middle| \xi_1, \xi_2 \right] &= G_2 (b_1, b_2, a, 1-c | -\xi_1, -\xi_2) \\ &= \sum_{m_1, m_2=0}^{\infty} (b_1)_{m_1} (b_2)_{m_2} (a)_{m_2-m_1} (1-c)_{m_1-m_2} \frac{(-\xi_1)^{m_1}}{m_1!} \frac{(-\xi_2)^{m_2}}{m_2!}. \end{aligned} \quad (\text{B.12})$$

Finally, note that given a N -point generalized hypergeometric function one can construct a $(N + m)$ -point hypergeometric function through the so-called splitting identities. E.g., the fourth Appell function F_4 defined by the double power series

$$F_4 \left[\begin{matrix} a_1, a_2 \\ c_1, c_2 \end{matrix} \middle| \xi_1, \xi_2 \right] = \sum_{m_1, m_2=0}^{\infty} \frac{(a_1)_{m_1+m_2} (a_2)_{m_2+m_1} \xi_1^{m_1} \xi_2^{m_2}}{(c_1)_{m_1} (c_2)_{m_2} m_1! m_2!}, \quad (\text{B.13})$$

and converging when $\sqrt{\xi_1} + \sqrt{\xi_2} < 1$ can be represented in terms of the Gauss hypergeometric function (B.7) by means of the following splitting identity

$$F_4 \left[\begin{matrix} a_1, a_2 \\ c_1, c_2 \end{matrix} \middle| \xi_1, \xi_2 \right] = \sum_{m_1=0}^{\infty} \frac{\xi_1^{m_1}}{m_1!} \frac{(a_1)_{m_1} (a_2)_{m_1}}{(c_1)_{m_1}} {}_2F_1 \left[\begin{matrix} a_1 + m_1, a_2 + m_2 \\ c_2 \end{matrix} \middle| \xi_2 \right]. \quad (\text{B.14})$$

B.2 Integral representations

Generalized hypergeometric functions can also be defined through the Euler-type integrals. E.g., for $\text{Re } b_i > 0, \text{Re}(c - |\mathbf{b}|) > 0$ the Lauricella function $F_D^{(N)}$ (B.4) can be represented as

$$F_D^{(N)} \left[\begin{matrix} a, \mathbf{b} \\ c \end{matrix} \middle| \boldsymbol{\xi} \right] = \Gamma \left[\begin{matrix} c \\ c - |\mathbf{b}|, \mathbf{b} \end{matrix} \right] \int_{\Omega} d^N \sigma \frac{\boldsymbol{\sigma}^{b-1} (1 - |\boldsymbol{\sigma}|)^{c-|\mathbf{b}|-1}}{(1 - \boldsymbol{\sigma} \cdot \boldsymbol{\xi})^a}, \quad (\text{B.15})$$

where $\boldsymbol{\sigma} \cdot \boldsymbol{\xi} = \sigma_1 \xi_1 + \dots + \sigma_N \xi_N$, and the integration domain is the standard orthogonal simplex $\Omega = \{\boldsymbol{\sigma} \in \mathbb{R}^N : \sigma_j \geq 0, j = 1, \dots, N; |\boldsymbol{\sigma}| \leq 1\}$ (see (A.3) for notation).

Euler-type integrals are useful in finding transformation formulas for generalized hypergeometric functions. This can be illustrated by another integral representation of the Lauricella function $F_D^{(N)}$:

$$F_D^{(N)} \left[\begin{matrix} a, \mathbf{b} \\ c \end{matrix} \middle| \boldsymbol{\xi} \right] = \Gamma \left[\begin{matrix} c \\ c - a, a \end{matrix} \right] \int_0^{\infty} dt t^{a-1} (1-t)^{c-a-1} \prod_{j=1}^N (1-t\xi_j)^{-b_j}, \quad (\text{B.16})$$

valid when $\text{Re } a > 0, \text{Re}(c - a) > 0$ and $|\arg(1 - \xi_j)| < \Pi, \forall j = 1, \dots, N$. Changing the integration variable as $t \rightarrow 1 - t$ one can establish the following transformation

$$F_D^{(N)} \left[\begin{matrix} a, \mathbf{b} \\ c \end{matrix} \middle| \boldsymbol{\xi} \right] = (1 - \boldsymbol{\xi})^{-b} F_D^{(N)} \left[\begin{matrix} c - a, \mathbf{b} \\ c \end{matrix} \middle| \mathcal{P}(\boldsymbol{\xi}) \right], \quad \mathcal{P}(\boldsymbol{\xi}) = \left\{ \frac{\xi_1}{\xi_1 - 1}, \dots, \frac{\xi_N}{\xi_N - 1} \right\}. \quad (\text{B.17})$$

At $N = 1$ it is the Pfaff transformation of the Gauss hypergeometric function.

Moreover, the integral representations can be useful in finding analytic continuation formulas. E.g. the Gauss hypergeometric function ${}_2F_1$ has the following integral representation

$${}_2F_1 \left[\begin{matrix} a, b \\ 1 + a + b - c \end{matrix} \middle| 1 - \xi \right] = \Gamma \left[\begin{matrix} 1 + a + b - c \\ a, b \end{matrix} \right] \int_0^{\infty} ds \frac{s^{b-1} (1+s)^{c-b-1}}{(1+s\xi)^a}, \quad (\text{B.18})$$

which extends the original power series definition to a domain around 1.

Mellin-Barnes integrals. In order to deal with the problem of analytic continuation of generalized hypergeometric functions, it is useful to use the Mellin-Barnes integral technique. E.g. the Mellin-Barnes representation for the Gauss hypergeometric function is given by

$${}_2F_1 \left[\begin{matrix} a, b \\ c \end{matrix} \middle| \xi \right] = \Gamma \left[\begin{matrix} c \\ a, b \end{matrix} \right] \int_{-i\infty}^{+i\infty} \frac{ds}{2\pi i} \Gamma \left[\begin{matrix} a + s, b + s, -s \\ c + s \end{matrix} \right] (-\xi)^s, \quad (\text{B.19})$$

where the integration contour is chosen such that the poles of the Γ -functions $s^{(1)} = -a - k$ and $s^{(2)} = -b - k$ lie to the left of the contour and the poles $s^{(3)} = k$ lie to the right ($k \in \mathbb{Z}^+$).

The Mellin-Barnes integrals are also useful in representing elementary functions, e.g. the exponential function

$$\exp(-X) = \frac{1}{2\pi i} \int_{-i\infty}^{+i\infty} ds \Gamma(-s) X^s, \quad (\text{B.20})$$

and the power function

$$\begin{aligned} \frac{1}{(\lambda_1 + \dots + \lambda_N)^a} &= \frac{1}{\Gamma(a)} \frac{1}{(2\pi i)^{N-1}} \int_{-i\infty}^{+i\infty} ds_1 \cdots \int_{-i\infty}^{+i\infty} ds_{N-1} \Gamma(-s_1) \cdots \Gamma(-s_{N-1}) \\ &\quad \times \Gamma(s_1 + \dots + s_{N-1} + a) \lambda_1^{s_1} \cdots \lambda_{N-1}^{s_{N-1}} \lambda_N^{-s_1 - \dots - s_{N-1} - a}. \end{aligned} \quad (\text{B.21})$$

In the last formula the integration contours for s_i are chosen such that poles $s_i^{(1)} = -S_i - k_i$ are to the left, where $S_i = s_1 + \dots + s_N + a - s_i$ and poles $s_i^{(2)} = -k_i$ are to the right, $k_i \in \mathbb{Z}^+$, $\forall i = 1, \dots, N-1$.

B.3 Analytic continuation

When calculating Mellin-Barnes integrals, e.g. (B.19), there is an ambiguity in closing the contour to the left or right. However, it can be fixed by specifying domains of argument ξ and parameters a, b, c . More precisely, when closing a contour to the right, the integral (B.19) can be calculated as a sum of residues at the poles $s^{(3)} = k \in \mathbb{Z}^+$, yielding the Gauss hypergeometric series (B.7) which converges for $|\xi| < 1$. On the other hand, when closing a contour to the left, the same integral is a sum of residues at the poles $s^{(1)} = -k - a$ and $s^{(2)} = -k - b$ with $k \in \mathbb{Z}^+$. In this way, one obtains the analytic continuation formula for the Gauss hypergeometric series (B.7) in the domain $|\xi| > 1$:

$$\begin{aligned} {}_2F_1 \left[\begin{matrix} a, b \\ c \end{matrix} \middle| \xi \right] &= \Gamma \left[\begin{matrix} c, b-a \\ b, c-a \end{matrix} \right] (-\xi)^{-a} {}_2F_1 \left[\begin{matrix} a, 1+a-c \\ 1+a-b \end{matrix} \middle| \frac{1}{\xi} \right] \\ &\quad + \Gamma \left[\begin{matrix} c, a-b \\ a, c-b \end{matrix} \right] (-\xi)^{-b} {}_2F_1 \left[\begin{matrix} b, 1+b-c \\ 1+b-a \end{matrix} \middle| \frac{1}{\xi} \right]. \end{aligned} \quad (\text{B.22})$$

Another analytic continuation formula can be derived by means of the first Barnes lemma:

$$\int_{-i\infty}^{+i\infty} \frac{dt}{2\pi i} \Gamma(a+t, b+t, s-t, c-a-b-t) = \Gamma \left[\begin{matrix} a+s, b+s, c-b, c-a \\ c+s \end{matrix} \right]. \quad (\text{B.23})$$

Substituting this relation into (B.19) one obtains the following integral representation

$${}_2F_1 \left[\begin{matrix} a, b \\ c \end{matrix} \middle| \xi \right] = \Gamma \left[\begin{matrix} c \\ a, b, c-a, c-b \end{matrix} \right] \int_{-i\infty}^{+i\infty} \frac{dt}{2\pi i} \Gamma(a+t, b+t, c-a-b-t, -t) (1-\xi)^t, \quad (\text{B.24})$$

where we also used (B.21) to evaluate the integral over s . Computing the integral as a sum over residues at the poles $t^{(1)} = k \in \mathbb{Z}^+$ and $t^{(2)} = c - a - b + k$, we find the following analytic continuation formula for the Gauss hypergeometric function

$$\begin{aligned} {}_2F_1 \left[\begin{matrix} a, b \\ c \end{matrix} \middle| \xi \right] &= \Gamma \left[\begin{matrix} c, c - a - b \\ c - a, c - b \end{matrix} \right] {}_2F_1 \left[\begin{matrix} a, b \\ 1 + a + b - c \end{matrix} \middle| 1 - \xi \right] \\ &+ \Gamma \left[\begin{matrix} c, a + b - c \\ a, b \end{matrix} \right] (1 - \xi)^{c-a-b} {}_2F_1 \left[\begin{matrix} c - a, c - b \\ 1 + c - a - b \end{matrix} \middle| 1 - \xi \right]. \end{aligned} \quad (\text{B.25})$$

By extending this procedure to generalized hypergeometric functions we also find the analytic continuation formula for the Lauricella function $F_D^{(N)}$ (B.4):¹⁶

$$F_D^{(N)} \left[\begin{matrix} a, \mathbf{b} \\ c \end{matrix} \middle| \boldsymbol{\xi} \right] = \sum_{q=0}^N A_q D_q \left[\begin{matrix} a, \mathbf{b} \\ c \end{matrix} \middle| \boldsymbol{\xi} \right], \quad (\text{B.26})$$

with a new domain of convergence

$$\mathbb{K}^N = \{ \boldsymbol{\xi} \in \mathbb{C}^N : 0 < |1 - \xi_1| < \dots < |1 - \xi_N| < 1; |\arg(1 - \xi_j)| < \pi, j = 1, \dots, N \}, \quad (\text{B.27})$$

cf. (B.5). The functions D_q are given by

$$\begin{aligned} D_0 \left[\begin{matrix} a, \mathbf{b} \\ c \end{matrix} \middle| \boldsymbol{\xi} \right] &= F_D^{(N)} \left[\begin{matrix} a, \mathbf{b} \\ 1 + a + |\mathbf{b}| - c \end{matrix} \middle| \mathbf{1} - \boldsymbol{\xi} \right], \\ D_q \left[\begin{matrix} a, \mathbf{b} \\ c \end{matrix} \middle| \boldsymbol{\xi} \right] &= (1 - \xi_q)^{c-a-|\mathbf{b}_{1,q}|} \left(\prod_{l=q+1}^N (1 - \xi_l)^{-b_l} \right) G^{(N,q)} \left[\begin{matrix} c - a - |\mathbf{b}_{1,q-1}|, \tilde{\mathbf{b}}_q \\ 1 + c - a - |\mathbf{b}_{1,q}| \end{matrix} \middle| \boldsymbol{\Xi}^{(q)} \right], \end{aligned} \quad (\text{B.28})$$

where $q = 1, \dots, N$ and $G^{(N,q)}$ is defined in (B.9). The parameters and arguments in (B.28) are packed into the following combinations:

$$\begin{aligned} \tilde{\mathbf{b}}_q &= \{ b_1, \dots, b_{q-1}, c - |\mathbf{b}|, b_{q+1}, \dots, b_N \}, \\ \boldsymbol{\Xi}^{(q)} &= \left\{ \frac{1 - \xi_1}{1 - \xi_q}, \dots, \frac{1 - \xi_{q-1}}{1 - \xi_q}, 1 - \xi_q, \frac{1 - \xi_q}{1 - \xi_{q+1}}, \dots, \frac{1 - \xi_q}{1 - \xi_N} \right\}. \end{aligned} \quad (\text{B.29})$$

The coefficients in (B.26) are given by

$$A_0 = \Gamma \left[\begin{matrix} c, c - a - |\mathbf{b}| \\ c - |\mathbf{b}|, c - a \end{matrix} \right], \quad A_q = \Gamma \left[\begin{matrix} c, c - |\mathbf{b}_{1,q-1}| - a, a + |\mathbf{b}_{1,q}| - c \\ a, b_q, c - a \end{matrix} \right]. \quad (\text{B.30})$$

¹⁶For $N = 2$, the formula (B.26) which was first derived in [76], provides an analytic continuation of the first Appell function F_1 .

It is worth noting that the right-hand side of (B.26) contains two Lauricella functions. The first one is given by D_0 , while the term D_1 is proportional to the Lauricella function by virtue of (B.11).

C One more Mellin-Barnes representation of the conformal integral

In this section, we describe an alternative way of representing the conformal integral in terms of Mellin-Barnes integrals. To this end, one steps back to (2.16) and represents the Gauss hypergeometric function ${}_2F_1$ therein by means of the analytic continuation formula (B.24):

$${}_2F_1 \left[\begin{matrix} a'_{n-1}, a'_n \\ \frac{D}{2} \end{matrix} \middle| 1 - \xi(\boldsymbol{\sigma}) \right] = \int_{-i\infty}^{+i\infty} \widehat{dt} \Gamma \left[\begin{matrix} \frac{D}{2}, a'_{n-1} + t, a'_n + t, \alpha_{n-1, n} - t \\ a_{n-1}, a_n, a'_{n-1}, a'_n \end{matrix} \right] (\xi(\boldsymbol{\sigma}))^t, \quad (\text{C.1})$$

where $\xi(\boldsymbol{\sigma})$ is given in (2.17). Applying to the numerator of $\xi(\boldsymbol{\sigma})$ the same procedure as in (2.23), one can obtain the following representation of the n -point conformal integral:

$$\begin{aligned} \mathcal{I}_n^{\mathbf{a}} &= N_n^{\mathbf{a}} \int_{-i\infty}^{+i\infty} \frac{dt}{2\pi i} \Gamma(a'_{n-1} + t, a'_n + t, \alpha_{n-1, n} - t) \left((\eta_n)_{n-2, n-1}^{n-3, n-2} \right)^t \\ &\times \int_{-i\infty}^{+i\infty} \prod_{i=1}^{n-4} \left(\widehat{dt}_i \left((\eta_n)_{n-3, n-2}^{i, n-2} \right)^{t_i} \right) \int_{-i\infty}^{+i\infty} \prod_{1 \leq i < j \leq n-3} \left(\widehat{ds}_{ij} \left((\eta_n)_{n-3, n-2}^{ij} \right)^{s_{ij}} \right) \\ &\times \Gamma \left(|\mathbf{t}_{1, n-4}| + \sum_{1 \leq i < j \leq n-3} s_{ij} - t \right) \Gamma \left[\begin{matrix} C^{(1)} - |\mathbf{B}^{(1)}|, \mathbf{B}^{(1)} \\ C^{(1)} \end{matrix} \right] F_D^{(n-3)} \left[\begin{matrix} A^{(1)}, \mathbf{B}^{(1)} \\ C^{(1)} \end{matrix} \middle| \boldsymbol{\xi} \right], \end{aligned} \quad (\text{C.2})$$

where $A^{(1)}, \mathbf{B}^{(1)}$ and $C^{(1)}$ are defined in (2.25). Although the parameters here are the same as in the first bare integral (2.24), the pole structure is more similar to that of the second bare integral (2.32). In particular, when closing a contour over t to the right, there are two sets of poles coming from $\Gamma(\alpha_{n-1, n} - t)$ and $\Gamma(|\mathbf{t}_{1, n-4}| + \sum_{1 \leq i < j \leq n-3} s_{ij} - t)$. Therefore, in the main text we focus on the bipartite representation of the conformal integral which allows one to evaluate the first bare integral explicitly.

D Pentagon integral

Here, we collect the corresponding master and basis functions and then verify that the obtained expression for the parametric pentagon integral correctly reproduces the known expression in the non-parametric case.

D.1 Basis functions

A basis function for the pentagon integral can be expressed in terms of two generalized hypergeometric functions which are defined as¹⁷

$$P_1 \left[\begin{matrix} A_1, A_2, B \\ C_1, C_2 \end{matrix} \middle| \boldsymbol{\xi} \right] = \sum_{l_i=0} (-)^{l_2} \frac{(A_1)_{l_1+l_2+l_4} (A_2)_{l_2+l_3+l_5} (B)_{l_1+l_2+l_3+l_4+l_5}}{(C_1)_{l_1+l_2+l_3} (C_2)_{l_2+l_4+l_5}} \prod_{i=1}^5 \frac{\xi_i^{l_i}}{l_i!}, \quad (\text{D.1})$$

$$P_2 \left[\begin{matrix} A_1, A_2, B \\ C, E \end{matrix} \middle| \boldsymbol{\xi} \right] = \sum_{l_i=0} (-)^{l_2} \frac{(A_1)_{l_1+l_2+l_4} (A_2)_{l_1+l_3+l_5} (B)_{l_5+l_3-l_4}}{(C)_{l_1+l_2+l_3} (E)_{l_5-l_4-l_2}} \prod_{i=1}^5 \frac{\xi_i^{l_i}}{l_i!}. \quad (\text{D.2})$$

The functions obey obvious symmetry relations

$$\begin{aligned} P_1 \left[\begin{matrix} A_1, A_2, B \\ C_1, C_2 \end{matrix} \middle| \xi_1, \xi_2, \xi_3, \xi_4, \xi_5 \right] &= P_1 \left[\begin{matrix} A_2, A_1, B \\ C_1, C_2 \end{matrix} \middle| \xi_1, \xi_4, \xi_5, \xi_2, \xi_3 \right] \\ &= P_1 \left[\begin{matrix} A_1, A_2, B \\ C_2, C_1 \end{matrix} \middle| \xi_4, \xi_2, \xi_5, \xi_1, \xi_3 \right], \end{aligned} \quad (\text{D.3})$$

$$P_2 \left[\begin{matrix} A_1, A_2, B \\ C, E \end{matrix} \middle| \xi_1, \xi_2, \xi_3, \xi_4, \xi_5 \right] = P_2 \left[\begin{matrix} A_2, A_1, 1-E \\ C, 1-B \end{matrix} \middle| \xi_1, \xi_3, \xi_2, \xi_5, \xi_4 \right]. \quad (\text{D.4})$$

To prove (D.4) one uses (B.10).

The action of $C_5 \in \mathbb{Z}_5$ on the cross-ratios (3.47) is summarized in the table

	u_1	u_2	w_{12}	v_1	v_2
C_5	$\frac{v_1}{v_2}$	v_1	$\frac{u_2 v_1}{v_2}$	$\frac{w_{12}}{v_2}$	u_1
$(C_5)^2$	$\frac{w_{12}}{u_1 v_2}$	$\frac{w_{12}}{v_2}$	$\frac{w_{12} v_1}{u_1 v_2}$	$\frac{u_2 v_1}{u_1 v_2}$	$\frac{v_1}{v_2}$
$(C_5)^3$	$\frac{u_2}{u_1}$	$\frac{u_2 v_1}{u_1 v_2}$	$\frac{w_{12} u_2}{u_1 v_2}$	$\frac{w_{12}}{u_1}$	$\frac{w_{12}}{u_1 v_2}$
$(C_5)^4$	v_2	$\frac{w_{12}}{u_1}$	$\frac{u_2 v_1}{u_1}$	u_2	$\frac{u_2}{u_1}$

The pentagon integral $I_5^a(\mathbf{x})$ is the sum of 10 terms (3.56) which are divided into two groups depending on two master function $\varphi_5^{(345)}(\mathbf{x})$ and $\varphi_5^{(245)}(\mathbf{x})$ (3.54) used to generate them through the cyclic permutations.

¹⁷Note that P_1 is the Srivastava-Daoust hypergeometric function [77, 78], which also arises in the analysis of the hexagon integral [18]. The pentagon example shows that there is also another type of function, i.e. P_2 , which to the best of our knowledge has not been considered in the literature.

Basis functions generated from $\varphi_5^{(345)}(\mathbf{x})$:

$$\varphi_5^{(345)}(\mathbf{x}) = S_5^{(345)} V_5^{(345)}(\mathbf{x}) P_1 \left[a_1, a_2, a'_4 \left| \begin{array}{l} u_1, w_{12}, u_2, v_1, v_2 \\ 1 - \alpha_{45}, 1 - \alpha_{34} \end{array} \right. \right], \quad (\text{D.5})$$

$$\varphi_5^{(145)}(\mathbf{x}) = S_5^{(145)} V_5^{(145)}(\mathbf{x}) P_1 \left[a_2, a_3, a'_5 \left| \begin{array}{l} \frac{v_1}{v_2}, \frac{u_2 v_1}{v_2}, v_1, \frac{w_{12}}{v_2}, u_1 \\ 1 - \alpha_{15}, 1 - \alpha_{45} \end{array} \right. \right], \quad (\text{D.6})$$

$$\varphi_5^{(125)}(\mathbf{x}) = S_5^{(125)} V_5^{(125)}(\mathbf{x}) P_1 \left[a_3, a_4, a'_1 \left| \begin{array}{l} \frac{w_{12}}{u_1 v_2}, \frac{w_{12} v_1}{u_1 v_2}, \frac{w_{12}}{v_2}, \frac{u_2 v_1}{u_1 v_2}, \frac{v_1}{v_2} \\ 1 - \alpha_{12}, 1 - \alpha_{15} \end{array} \right. \right], \quad (\text{D.7})$$

$$\varphi_5^{(123)}(\mathbf{x}) = S_5^{(123)} V_5^{(123)}(\mathbf{x}) P_1 \left[a_4, a_5, a'_2 \left| \begin{array}{l} \frac{u_2}{u_1}, \frac{w_{12} u_2}{u_1 v_2}, \frac{u_2 v_1}{u_1 v_2}, \frac{w_{12}}{u_1}, \frac{w_{12}}{u_1 v_2} \\ 1 - \alpha_{23}, 1 - \alpha_{12} \end{array} \right. \right], \quad (\text{D.8})$$

$$\varphi_5^{(234)}(\mathbf{x}) = S_5^{(234)} V_5^{(234)}(\mathbf{x}) P_1 \left[a_5, a_1, a'_3 \left| \begin{array}{l} v_2, \frac{u_2 v_1}{u_1}, \frac{w_{12}}{u_1}, u_2, \frac{u_2}{u_1} \\ 1 - \alpha_{34}, 1 - \alpha_{23} \end{array} \right. \right]. \quad (\text{D.9})$$

Basis functions generated from $\varphi_5^{(245)}(\mathbf{x})$:

$$\varphi_5^{(245)}(\mathbf{x}) = S_5^{(245)} V_5^{(245)}(\mathbf{x}) P_2 \left[a_1, a_3, -\alpha_{15} \left| \begin{array}{l} u_1, \frac{w_{12}}{v_2}, u_2, \frac{v_1}{v_2}, v_2 \\ 1 - \alpha_{45}, 1 + \alpha_{34} \end{array} \right. \right], \quad (\text{D.10})$$

$$\varphi_5^{(135)}(\mathbf{x}) = S_5^{(135)} V_5^{(135)}(\mathbf{x}) P_2 \left[a_2, a_4, -\alpha_{12} \left| \begin{array}{l} \frac{v_1}{v_2}, \frac{u_2 v_1}{u_1 v_2}, v_1, \frac{w_{12}}{u_1 v_2}, u_1 \\ 1 - \alpha_{15}, 1 + \alpha_{45} \end{array} \right. \right], \quad (\text{D.11})$$

$$\varphi_5^{(124)}(\mathbf{x}) = S_5^{(124)} V_5^{(124)}(\mathbf{x}) P_2 \left[a_3, a_5, -\alpha_{23} \left| \begin{array}{l} \frac{w_{12}}{u_1 v_2}, \frac{w_{12}}{u_1}, \frac{w_{12}}{v_2}, \frac{u_2}{u_1}, \frac{v_1}{v_2} \\ 1 - \alpha_{12}, 1 + \alpha_{15} \end{array} \right. \right], \quad (\text{D.12})$$

$$\varphi_5^{(235)}(\mathbf{x}) = S_5^{(235)} V_5^{(235)}(\mathbf{x}) P_2 \left[a_4, a_1, -\alpha_{34} \left| \begin{array}{l} \frac{u_2}{u_1}, u_2, \frac{u_2 v_1}{u_1 v_2}, v_2, \frac{w_{12}}{u_1 v_2} \\ 1 - \alpha_{23}, 1 + \alpha_{12} \end{array} \right. \right], \quad (\text{D.13})$$

$$\varphi_5^{(134)}(\mathbf{x}) = S_5^{(134)} V_5^{(134)}(\mathbf{x}) P_2 \left[a_5, a_2, -\alpha_{45} \left| \begin{array}{l} v_2, v_1, \frac{w_{12}}{u_1}, u_1, \frac{u_2}{u_1} \\ 1 - \alpha_{34}, 1 + \alpha_{23} \end{array} \right. \right]. \quad (\text{D.14})$$

D.2 Non-parametric integral

In what follows we compare our asymptotic expansion of the non-parametric pentagon integral with the exact formula proposed in [15].

Geometric approach. Within the ambient space approach the integral is defined as

$$I^{(5)} = \int \frac{d^5 x_0}{i\pi^{5/2}} \prod_{i=1}^5 \frac{1}{(x_i - x_0)^2} = \int \frac{d^5 Q}{i\pi^{d/2}} \prod_{i=1}^5 \frac{1}{(-2P_i \cdot Q)}, \quad (\text{D.15})$$

where $P_i^A = P^A(x_i)$, $Q^A = Q^A(x_0)$, $A = 1, \dots, d+2$, are $n+1$ null vectors in $\mathbb{R}^{1,d+1}$ such that local coordinates $x^\mu \in \mathbb{R}^d$ are introduced as $X^A(x) = (1, x^2, x^\mu)$. The volume of a particular 4-simplex in a constant curvature 4-dimensional space calculates the pentagon integral (D.15) [12–14, 16]:

$$I^{(5)} = \frac{2^{\frac{5}{2}} \Gamma(\frac{5}{2})}{\sqrt{|\det P_{ij}|}} V^{(4)}, \quad (\text{D.16})$$

where $P_{ij} \equiv -2P_i \cdot P_j = (x_i - x_j)^2$ and the volume can be calculated by means of the Schläfli formula given by

$$V_{(4)} = \frac{\pi}{6} \sum_{1 \leq i < j \leq 5} \log \frac{W_i \cdot W_j - \sqrt{(W_i \cdot W_j)^2 - W_i^2 W_j^2}}{W_i \cdot W_j + \sqrt{(W_i \cdot W_j)^2 - W_i^2 W_j^2}}. \quad (\text{D.17})$$

In this formula n vectors W_i are uniquely defined through $W_i \cdot P_j = \delta_{ij}$ (for review see e.g. [79]). Introducing the rescaled integral

$$I^{(5)} = (P_{13} P_{14} P_{24} P_{25} P_{35})^{-1/2} \tilde{I}^{(5)}, \quad (\text{D.18})$$

after identical transformations one finds that [15]

$$\tilde{I}^{(5)} = \frac{\pi^{\frac{3}{2}}}{2\sqrt{-\Delta^{(5)}}} (1 + g + g^2 + g^3 + g^4) \left\{ \log \left(\frac{r - \sqrt{-\Delta^{(5)}}}{r + \sqrt{-\Delta^{(5)}}} \right) \left(\frac{s - \sqrt{-\Delta^{(5)}}}{s + \sqrt{-\Delta^{(5)}}} \right) \right\}, \quad (\text{D.19})$$

where

$$r = \frac{(1 - t_2)(1 - t_5) - t_1(2 - t_3 - t_4) - t_3 t_5 - t_2 t_4 + t_1 t_3 t_4}{2\sqrt{t_1}}, \quad (\text{D.20})$$

$$s = \frac{(1 - t_5)(1 - t_2 t_5) - t_1(1 + t_5 - 2t_3 t_5 + t_4 + t_2 t_4 t_5 - t_1 t_4)}{2\sqrt{t_1 t_5}},$$

$$\Delta^{(5)} = 1 - [t_1(1 - t_3(1 + t_4) + t_2 t_4^2) + \text{cyclic}] + t_1 t_2 t_3 t_4 t_5, \quad (\text{D.21})$$

the cross-ratios are chosen as

$$t_1 = \frac{P_{14} P_{23}}{P_{13} P_{24}}, \quad t_2 = \frac{P_{25} P_{34}}{P_{24} P_{35}}, \quad t_3 = \frac{P_{13} P_{45}}{P_{14} P_{35}}, \quad t_4 = \frac{P_{15} P_{24}}{P_{14} P_{25}}, \quad t_5 = \frac{P_{12} P_{35}}{P_{13} P_{25}}, \quad (\text{D.22})$$

and $g : t_i \rightarrow t_{i+1}$ is a cyclic permutation which says that the pentagon formula (D.19) has a manifest cyclic permutation symmetry. The function $\Delta^{(5)} \geq 0$ depends on the choice a specific kinematical regime, i.e. on values of variables P_i . The resulting function (D.19) has 10 terms which number matches 10 terms in the asymptotic expansion (3.56).

Redefinitions. Introducing simplified notation, from (2.7) for the non-parametric pentagon integral one has

$$I_5 = LI_5, \quad (\text{D.23})$$

where the leg-factor is

$$L = (X_{15}X_{25}X_{34}X_{35}X_{45})^{-1} \quad (\text{D.24})$$

and \mathcal{I}_5 is expressed in terms of the cross-ratios (3.47):

$$\begin{aligned} u_1 &= \frac{X_{13}X_{45}}{X_{14}X_{35}}, & u_2 &= \frac{X_{23}X_{45}}{X_{24}X_{35}}, & w_{12} &= \frac{X_{12}X_{34}X_{45}}{X_{14}X_{24}X_{35}}, \\ v_1 &= \frac{X_{15}X_{34}}{X_{14}X_{35}}, & v_2 &= \frac{X_{25}X_{34}}{X_{24}X_{35}}. \end{aligned} \quad (\text{D.25})$$

Since $X_{ij} \equiv P_{ij}$, cf. (2.1), one finds that (D.22) and (D.25) are related as

$$\begin{aligned} t_1 &= \frac{u_2}{u_1}, & t_2 &= v_2, & t_3 &= u_1, & t_4 &= \frac{v_1}{v_2}, & t_5 &= \frac{w_{12}}{u_1v_2}; \\ u_1 &= t_3, & u_2 &= t_1t_3, & w_{12} &= t_2t_3t_5, & v_1 &= t_2t_4, & v_2 &= t_2. \end{aligned} \quad (\text{D.26})$$

On the other hand, the two definitions of the conformal integral (D.15) and (D.23) are related as $I_5 = iI^{(5)}$. Thus, one has the relation

$$\mathcal{I}_5 = iL^{-1}\tilde{L}\tilde{I}^{(5)} = it_2^{3/2}t_3^{-1/2}t_4\tilde{I}^{(5)}, \quad (\text{D.27})$$

where $\tilde{I}^{(5)}$ is given by (D.19) and the leg-factor $\tilde{L} = (X_{13}X_{14}X_{24}X_{25}X_{35})^{-1/2}$ in (D.18) is related to (D.24) as $\tilde{L} = t_2^{3/2}t_3^{-1/2}t_4L$.

Unit propagator powers. Contrary to the case of unit parameters in the box integral considered in section 3.2.3, the non-parametric pentagon integral has no divergent prefactors so that no regularization is required. Choosing $a_i = 1$ we introduce the simplified notation for two functions (D.1)-(D.2):

$$\begin{aligned} \mathbb{P}_1[z_1, z_2, z_3, z_4, z_5] &\equiv \mathbb{P}_1 \left[\begin{matrix} 1, 1, \frac{3}{2} \\ \frac{3}{2}, \frac{3}{2} \end{matrix} \middle| z_1, z_2, z_3, z_4, z_5 \right] = \sum_{\{n_i\}=0}^{\infty} \frac{z_1^{n_1} z_2^{n_2} z_3^{n_3} z_4^{n_4} z_5^{n_5}}{n_1! n_2! n_3! n_4! n_5!} \\ &\times (-)^{n_2} \frac{(n_1 + n_2 + n_4)!(n_2 + n_3 + n_5)! \left(\frac{3}{2}\right)_{n_1+n_2+n_3+n_4+n_5}}{\left(\frac{3}{2}\right)_{n_1+n_2+n_3} \left(\frac{3}{2}\right)_{n_2+n_4+n_5}}; \end{aligned} \quad (\text{D.28})$$

$$\begin{aligned} \mathbb{P}_2[z_1, z_2, z_3, z_4, z_5] &\equiv \mathbb{P}_2 \left[\begin{matrix} 1, 1, \frac{1}{2} \\ \frac{3}{2}, \frac{1}{2} \end{matrix} \middle| z_1, z_2, z_3, z_4, z_5 \right] = \sum_{\{n_i\}=0}^{\infty} \frac{z_1^{n_1} z_2^{n_2} z_3^{n_3} z_4^{n_4} z_5^{n_5}}{n_1! n_2! n_3! n_4! n_5!} \\ &\times (-)^{n_2} \frac{(n_1 + n_2 + n_4)!(n_1 + n_3 + n_5)! \left(\frac{1}{2}\right)_{n_3-n_4+n_5}}{\left(\frac{3}{2}\right)_{n_1+n_2+n_3} \left(\frac{1}{2}\right)_{n_5-n_2-n_4}}. \end{aligned} \quad (\text{D.29})$$

Then, the symmetry properties (D.3)-(D.4) take the form

$$\begin{aligned}\mathbb{P}_1[z_1, z_2, z_3, z_4, z_5] &= \mathbb{P}_1[z_1, z_4, z_5, z_2, z_3] = \mathbb{P}_1[z_4, z_2, z_5, z_1, z_3], \\ \mathbb{P}_2[z_1, z_2, z_3, z_4, z_5] &= \mathbb{P}_2[z_1, z_3, z_2, z_5, z_4].\end{aligned}\tag{D.30}$$

The asymptotic expansion of the pentagon integral \mathcal{I}_5 from (D.23) reads as

$$\begin{aligned}[2\pi^{\frac{3}{2}}]^{-1}\mathcal{I}_5 &= v_1 v_2 \mathbb{P}_1\left[u_1, w_{12}, u_2, v_1, v_2\right] - v_1 v_2^{\frac{1}{2}} \mathbb{P}_2\left[u_1, \frac{w_{12}}{v_2}, u_2, \frac{v_1}{v_2}, v_2\right] \\ &+ v_1^{\frac{3}{2}} \mathbb{P}_1\left[u_1, \frac{u_2 v_1}{v_2}, \frac{w_{12}}{v_2}, v_1, \frac{v_1}{v_2}\right] - v_1^{\frac{3}{2}} u_1^{-\frac{1}{2}} \mathbb{P}_2\left[\frac{v_1}{v_2}, \frac{v_1 u_2}{u_1 v_2}, v_1, \frac{w_{12}}{u_1 v_2}, u_1\right] \\ &+ v_1^{\frac{3}{2}} v_2^{-\frac{1}{2}} u_1^{-1} w_{12}^{\frac{1}{2}} \mathbb{P}_1\left[\frac{w_{12}}{u_1 v_2}, \frac{w_{12} v_1}{u_1 v_2}, \frac{w_{12}}{v_2}, \frac{v_1 u_2}{u_1 v_2}, \frac{v_1}{v_2}\right] - v_1 w_{12}^{\frac{1}{2}} u_1^{-1} \mathbb{P}_2\left[\frac{w_{12}}{u_1 v_2}, \frac{w_{12}}{u_1}, \frac{w_{12}}{v_2}, \frac{u_2}{u_1}, \frac{v_1}{v_2}\right] \\ &+ v_1 u_1^{-\frac{3}{2}} u_2^{\frac{1}{2}} w_{12}^{\frac{1}{2}} \mathbb{P}_1\left[\frac{u_2}{u_1}, \frac{w_{12} u_2}{u_1 v_2}, \frac{v_1 u_2}{u_1 v_2}, \frac{w_{12}}{u_1}, \frac{w_{12}}{u_1 v_2}\right] - v_1 v_2^{\frac{1}{2}} u_1^{-1} u_2^{\frac{1}{2}} \mathbb{P}_2\left[\frac{u_2}{u_1}, u_2, \frac{v_1 u_2}{u_1 v_2}, v_2, \frac{w_{12}}{u_1 v_2}\right] \\ &+ v_1 v_2 u_1^{-1} u_2^{\frac{1}{2}} \mathbb{P}_1\left[v_2, \frac{u_2 v_1}{u_1}, \frac{w_{12}}{u_1}, u_2, \frac{u_2}{u_1}\right] - v_1 v_2 u_1^{-\frac{1}{2}} \mathbb{P}_2\left[v_2, v_1, \frac{w_{12}}{u_1}, u_1, \frac{u_2}{u_1}\right].\end{aligned}\tag{D.31}$$

Comparing two functions by Wolfram Mathematica. The power series representation (D.31) is suitable for checking the asymptotic expansion of the analytic formula (D.19) at $\boldsymbol{\eta} \rightarrow 0$ (3.2) and (3.9). The issue is that a function of many variables can tend to its value in parametric way. This means that one can choose a particular path in the five-dimensional space of cross-ratios \mathcal{H}_5 and then compare expansions of (D.31) and (D.19).

- Consider how the multivariate power series (D.31) tends to zero. To this end, one collectively denotes the cross-ratios (D.25) as $\boldsymbol{\eta} = \{u_1, u_2, w_{12}, v_1, v_2\}$ and scales them near $\boldsymbol{\eta} = 0$ as $\eta_i \rightarrow \eta_i \lambda^{\alpha_i}$, where $\{\alpha_i > 0, i = 1, \dots, 5\} \equiv \boldsymbol{\alpha}$, the proper time parameter $\lambda \in [0, 1]$. This means that one approaches $\boldsymbol{\eta} = 0$ along a particular way defined by fixing powers $\boldsymbol{\alpha}$.
- Choose a set $\boldsymbol{\alpha}$ with not too large α_i (in this case the Mathematica computation takes less time) which is consistent with sending $\boldsymbol{\eta} \rightarrow 0$ (3.2) and (3.9).
- Single out the leading order in λ , i.e. $I_5 = a_0 + a_1 \lambda^\gamma + O(\lambda^{\gamma+\dots})$ with some power $\gamma = \gamma(\boldsymbol{\alpha}_i)$, where expansion coefficients $a_{0,1} = a_{0,1}(\boldsymbol{\eta})$ are some rational functions of cross-ratios.
- The resulting coefficients $a_{0,1}$ are to be compared with those ones arising when expanding the exact formula for $I^{(5)}$ using the same scaling pattern, i.e. $I^{(5)} = i(b_0 + b_1 \lambda^\gamma + O(\lambda^{\gamma+\dots}))$. To this end, one expresses the cross-ratios t_i (D.22) via $\boldsymbol{\eta}$ by means of relations (D.26) and then sends $\boldsymbol{\eta} \rightarrow 0$ along the same path in \mathcal{H}_5 , i.e. using the same powers $\boldsymbol{\alpha}$.




Figure 2. Action of the kinematic group S_4^{kin} (3.17) on the basis functions (3.30). Each node denotes a basis function $\varphi_4^{(ijk)}(\mathbf{x})$ which indices are shown inside a node. The color lines represent permutations listed on the right (the identity permutation acts on each node trivially). The diagram also reflects the group multiplication law, i.e. each triangle formed by color lines represents a multiplication $g_i \circ g_j = g_k$, where $g_{i,j,k} \in S_4^{\text{kin}}$.

- The exact formula and the asymptotic expansion of the pentagon integral may coincide in a given coordinate domain iff $a_{0,1} = b_{0,1}$. One can further expand in λ and compare higher-order expansion coefficients.

Following this procedure one fixes the powers e.g. as $\alpha = \{3, 1, 4, 1, 2\}$. Obviously, there are infinitely many choices of α_i , but here we are choosing those ones with not too high integer values that simplifies computations. Near $\eta = 0$ the function $\Delta^{(5)} > 0$ and the exact formula (D.19) has under the logarithm a unimodular complex-valued function, i.e. the logarithm is pure imaginary. However, the final expression for the conformal integral is real due to the prefactor $\sqrt{-\Delta^{(5)}}$ which is also pure imaginary in this case. Using simple Wolfram Mathematica functions we explicitly expand both functions up to $O(\lambda^{11/2})$ and find that the two resulting series have the same form.

E Kinematic group extensions

The kinematic group $S_4^{\text{kin}} = \mathbb{Z}_2 \times \mathbb{Z}_2$ also acts on the expansion in basis functions (3.34) obtained by means of the cyclic group \mathbb{Z}_4 . One can show that $\mathbb{Z}_2 \times \mathbb{Z}_2$ acts on the basis functions by rearranging them according to the diagram rule shown on fig. 2. Any two nodes are related by a group element from $\mathbb{Z}_2 \times \mathbb{Z}_2$ that means that there is a single master function. This observation allows one to give a complementary definition of the kinematic group S_4^{kin} which replaces the original definition as a stabilizer group of cross-ratios for $n > 4$. Thus, it is possible to overcome the difficulty that $S_n^{\text{kin}} = \{e\}$ for $n > 4$ since this new *extended* kinematic group $\widehat{S}_n^{\text{kin}}$ is non-trivial for $n > 4$.

One can introduce an extended kinematic group by following a number of steps: (0) one analytically continues the first bare integral to a domain around the origin of coordinates in the space of cross-ratios; (1) one acts with the cyclic group \mathbb{Z}_n on the master functions supported on the same domain; (2) one represents the conformal integral as a sum of basis functions; (3) one identifies all permutations $\notin \mathbb{Z}_n$ which rearrange basis functions while keeping their domains around the origin.




Figure 3. In the 5-point case, the ten basis functions (D.5)-(D.14) can be equivalently obtained by acting with $\widehat{\mathcal{S}}_5^{\text{kin}}$. These can be grouped into two pentagon diagrams. The identical permutation leaves each node invariant.

As can be seen, such a working definition is largely based on the cyclic group, and currently there are no guidelines that could help define an extended kinematic group independently. In the 4-point case we have $\mathcal{S}_4^{\text{kin}} = \widehat{\mathcal{S}}_4^{\text{kin}}$. In the 5-point case, one can show that an extended kinematic group $\widehat{\mathcal{S}}_5^{\text{kin}}$ is generated by the following elements

$$\{e, (14)(23), (15)(24), (25)(34), (12)(35), (13)(45)\} \subset \mathcal{S}_5. \quad (\text{E.1})$$

$\widehat{\mathcal{S}}_5^{\text{kin}}$ acts on the ten basis functions as shown on fig. 3.

References

- [1] K. Symanzik, *On Calculations in conformal invariant field theories*, *Lett. Nuovo Cim.* **3** (1972) 734–738.
- [2] J. M. Drummond, J. Henn, V. A. Smirnov and E. Sokatchev, *Magic identities for conformal four-point integrals*, *JHEP* **01** (2007) 064, [[hep-th/0607160](#)].
- [3] J. M. Drummond, J. Henn, G. P. Korchemsky and E. Sokatchev, *Dual superconformal symmetry of scattering amplitudes in $N=4$ super-Yang-Mills theory*, *Nucl. Phys. B* **828** (2010) 317–374, [[0807.1095](#)].
- [4] A. B. Zamolodchikov, *‘Fishnet’ diagrams as a completely integrable system*, *Phys. Lett. B* **97** (1980) 63–66.
- [5] O. Gürdoğan and V. Kazakov, *New Integrable 4D Quantum Field Theories from Strongly Deformed Planar $\mathcal{N} = 4$ Supersymmetric Yang-Mills Theory*, *Phys. Rev. Lett.* **117** (2016) 201602, [[1512.06704](#)].
- [6] J. a. Caetano, O. Gürdoğan and V. Kazakov, *Chiral limit of $\mathcal{N} = 4$ SYM and ABJM and integrable Feynman graphs*, *JHEP* **03** (2018) 077, [[1612.05895](#)].
- [7] V. Kazakov and E. Olivucci, *Biscalar Integrable Conformal Field Theories in Any Dimension*, *Phys. Rev. Lett.* **121** (2018) 131601, [[1801.09844](#)].
- [8] D. Chicherin, V. Kazakov, F. Loebbert, D. Müller and D.-l. Zhong, *Yangian Symmetry for Bi-Scalar Loop Amplitudes*, *JHEP* **05** (2018) 003, [[1704.01967](#)].

- [9] D. Chicherin, V. Kazakov, F. Loebbert, D. Müller and D.-l. Zhong, *Yangian Symmetry for Fishnet Feynman Graphs*, *Phys. Rev. D* **96** (2017) 121901, [[1708.00007](#)].
- [10] F. Loebbert, *Integrability for Feynman integrals*, *SciPost Phys. Proc.* **14** (2023) 008, [[2212.09636](#)].
- [11] L. Corcoran, *Conformal Feynman Integrals and Correlation Functions in Fishnet Theory*. PhD thesis, Humboldt U., Berlin, 2023. 10.18452/25602.
- [12] A. I. Davydychev and R. Delbourgo, *A Geometrical angle on Feynman integrals*, *J. Math. Phys.* **39** (1998) 4299–4334, [[hep-th/9709216](#)].
- [13] L. Mason and D. Skinner, *Amplitudes at Weak Coupling as Polytopes in AdS_5* , *J. Phys. A* **44** (2011) 135401, [[1004.3498](#)].
- [14] O. Schnetz, *The geometry of one-loop amplitudes*, [1010.5334](#).
- [15] D. Nandan, M. F. Paulos, M. Spradlin and A. Volovich, *Star Integrals, Convolutions and Simplices*, *JHEP* **05** (2013) 105, [[1301.2500](#)].
- [16] J. L. Bourjaily, E. Gardi, A. J. McLeod and C. Vergu, *All-mass n -gon integrals in n dimensions*, *JHEP* **08** (2020) 029, [[1912.11067](#)].
- [17] L. Ren, M. Spradlin, C. Vergu and A. Volovich, *One-loop integrals from volumes of orthoschemes*, *JHEP* **05** (2024) 104, [[2306.04630](#)].
- [18] F. Loebbert, D. Müller and H. Münkler, *Yangian Bootstrap for Conformal Feynman Integrals*, *Phys. Rev. D* **101** (2020) 066006, [[1912.05561](#)].
- [19] C. Duhr, A. Klemm, F. Loebbert, C. Nega and F. Porkert, *Yangian-Invariant Fishnet Integrals in Two Dimensions as Volumes of Calabi-Yau Varieties*, *Phys. Rev. Lett.* **130** (2023) 041602, [[2209.05291](#)].
- [20] C. Duhr, A. Klemm, F. Loebbert, C. Nega and F. Porkert, *Geometry from integrability: multi-leg fishnet integrals in two dimensions*, *JHEP* **07** (2024) 008, [[2402.19034](#)].
- [21] F. Loebbert and H. Mathur, *The Feyn-structure of Yangian symmetry*, *JHEP* **01** (2025) 112, [[2410.11936](#)].
- [22] A. Pal and K. Ray, *Conformal integrals in four dimensions*, *JHEP* **10** (2022) 087, [[2109.09379](#)].
- [23] A. Pal and K. Ray, *Conformal integrals in all dimensions as GKZ hypergeometric functions and Clifford groups*, [2303.17326](#).
- [24] F. Levkovich-Maslyuk and V. Mishnyakov, *Yangian symmetry, GKZ equations and integrable Feynman graphs in conformal variables*, [2412.19296](#).
- [25] V. Del Duca, L. J. Dixon, J. M. Drummond, C. Duhr, J. M. Henn and V. A. Smirnov, *The one-loop six-dimensional hexagon integral with three massive corners*, *Phys. Rev. D* **84** (2011) 045017, [[1105.2011](#)].
- [26] L. J. Dixon, J. M. Drummond and J. M. Henn, *The one-loop six-dimensional hexagon integral and its relation to MHV amplitudes in $N=4$ SYM*, *JHEP* **06** (2011) 100, [[1104.2787](#)].
- [27] M. F. Paulos, *Loops, Polytopes and Splines*, *JHEP* **06** (2013) 007, [[1210.0578](#)].
- [28] C. Duhr and F. Porkert, *Feynman integrals in two dimensions and single-valued hypergeometric functions*, *JHEP* **02** (2024) 179, [[2309.12772](#)].

- [29] C. Duhr, A. Klemm, F. Loebbert, C. Nega and F. Porkert, *The Basso-Dixon formula and Calabi-Yau geometry*, *JHEP* **03** (2024) 177, [[2310.08625](#)].
- [30] S. G. Prabhu, *Conformality of non-conformal correlators*, [2312.17067](#).
- [31] S. G. Prabhu, *Feynman Diagrams from Conformal Integrals*, [2501.00101](#).
- [32] S. He, X. Jiang, J. Liu and Y.-Q. Zhang, *Notes on conformal integrals: Coulomb branch amplitudes, magic identities and bootstrap*, [2502.08871](#).
- [33] S. Ferrara, A. F. Grillo, G. Parisi and R. Gatto, *The shadow operator formalism for conformal algebra. Vacuum expectation values and operator products*, *Lett. Nuovo Cim.* **4S2** (1972) 115–120.
- [34] S. Ferrara, A. F. Grillo, G. Parisi and R. Gatto, *Covariant expansion of the conformal four-point function*, *Nucl. Phys. B* **49** (1972) 77–98.
- [35] F. A. Dolan and H. Osborn, *Implications of $N=1$ superconformal symmetry for chiral fields*, *Nucl. Phys. B* **593** (2001) 599–633, [[hep-th/0006098](#)].
- [36] F. A. Dolan and H. Osborn, *Conformal four point functions and the operator product expansion*, *Nucl. Phys. B* **599** (2001) 459–496, [[hep-th/0011040](#)].
- [37] F. Dolan and H. Osborn, *Conformal Partial Waves: Further Mathematical Results*, [1108.6194](#).
- [38] D. Simmons-Duffin, *Projectors, Shadows, and Conformal Blocks*, *JHEP* **04** (2014) 146, [[1204.3894](#)].
- [39] V. Rosenhaus, *Multipoint Conformal Blocks in the Comb Channel*, *JHEP* **02** (2019) 142, [[1810.03244](#)].
- [40] V. Fateev and S. Ribault, *The Large central charge limit of conformal blocks*, *JHEP* **02** (2012) 001, [[1109.6764](#)].
- [41] A. C. Petkou, *Thermal one-point functions and single-valued polylogarithms*, *Phys. Lett. B* **820** (2021) 136467, [[2105.03530](#)].
- [42] M. Karydas, S. Li, A. C. Petkou and M. Vilatte, *Conformal Graphs as Twisted Partition Functions*, *Phys. Rev. Lett.* **132** (2024) 231601, [[2312.00135](#)].
- [43] K. Alkalaev and S. Mandrygin, *Torus shadow formalism and exact global conformal blocks*, *JHEP* **11** (2023) 157, [[2307.12061](#)].
- [44] K. Alkalaev and S. Mandrygin, *One-point thermal conformal blocks from four-point conformal integrals*, *JHEP* **10** (2024) 241, [[2407.01741](#)].
- [45] Y. Gobeil, A. Maloney, G. S. Ng and J.-q. Wu, *Thermal Conformal Blocks*, *SciPost Phys.* **7** (2019) 015, [[1802.10537](#)].
- [46] V. Belavin and J. Ramos Cabezas, *Global conformal blocks via shadow formalism*, *JHEP* **02** (2024) 167, [[2401.02580](#)].
- [47] V. Belavin, J. Ramos Cabezas and B. Runov, *Shadow formalism for supersymmetric conformal blocks*, *JHEP* **11** (2024) 048, [[2408.07684](#)].
- [48] K. Alkalaev and S. Mandrygin, *Multipoint conformal integrals in D dimensions. Part II, to appear* (2025).

- [49] I. Dubovyk, J. Gluza and G. Somogyi, *Mellin-Barnes Integrals: A Primer on Particle Physics Applications*, *Lect. Notes Phys.* **1008** (2022) pp., [[2211.13733](#)].
- [50] B. Ananthanarayan, S. Banik, S. Friot and S. Ghosh, *Multiple Series Representations of N-fold Mellin-Barnes Integrals*, *Phys. Rev. Lett.* **127** (2021) 151601, [[2012.15108](#)].
- [51] S. Banik and S. Friot, *Analytic Evaluation of Multiple Mellin-Barnes Integrals*, in *Loops and Legs in Quantum Field Theory*, 7, 2024. [2407.20120](#).
- [52] B. Ananthanarayan, S. Banik, S. Friot and S. Ghosh, *Double box and hexagon conformal Feynman integrals*, *Phys. Rev. D* **102** (2020) 091901, [[2007.08360](#)].
- [53] S. Banik and S. Friot, *Multiple Mellin-Barnes integrals and triangulations of point configurations*, *Phys. Rev. D* **110** (2024) 036002, [[2309.00409](#)].
- [54] S. G. Gorishnii and A. P. Isaev, *On an Approach to the Calculation of Multiloop Massless Feynman Integrals*, *Theor. Math. Phys.* **62** (1985) 232.
- [55] A. P. Isaev, *Multiloop Feynman integrals and conformal quantum mechanics*, *Nucl. Phys. B* **662** (2003) 461–475, [[hep-th/0303056](#)].
- [56] S. Derkachov, A. P. Isaev and L. Shumilov, *Conformal triangles and zig-zag diagrams*, *Phys. Lett. B* **830** (2022) 137150, [[2201.12232](#)].
- [57] S. E. Derkachov, A. P. Isaev and L. A. Shumilov, *Ladder and zig-zag Feynman diagrams, operator formalism and conformal triangles*, *JHEP* **06** (2023) 059, [[2302.11238](#)].
- [58] E. S. Fradkin and M. Y. Palchik, *Conformal quantum field theory in D-dimensions*. 1996.
- [59] S. I. Bezrodnykh, *Analytic continuation formulas and jacobi-type relations for Lauricella function*, *Doklady Mathematics* **93** (Mar., 2016) 129–134.
- [60] S. I. Bezrodnykh, *On the analytic continuation of the Lauricella function $F_D^{(N)}$* , *Mathematical Notes* **100** (July, 2016) 318–324.
- [61] S. I. Bezrodnykh, *Analytic continuation of the Appell function F_1 and integration of the associated system of equations in the logarithmic case*, *Computational Mathematics and Mathematical Physics* **57** (Apr., 2017) 559–589.
- [62] S. I. Bezrodnykh, *Analytic continuation of the Lauricella function $F_D^{(N)}$ with arbitrary number of variables*, *Integral Transforms Spec. Funct.* **29** (2018) 21–42.
- [63] S. I. Bezrodnykh, *The Lauricella hypergeometric function $F_D^{(N)}$, the Riemann–Hilbert problem, and some applications*, *Uspekhi Mat. Nauk* **73** (2018) 3–94.
- [64] P. Kravchuk and D. Simmons-Duffin, *Counting Conformal Correlators*, *JHEP* **02** (2018) 096, [[1612.08987](#)].
- [65] E. E. Boos and A. I. Davydychev, *A Method of the Evaluation of the Vertex Type Feynman Integrals*, *Moscow Univ. Phys. Bull.* **42N3** (1987) 6–10.
- [66] E. E. Boos and A. I. Davydychev, *A Method of evaluating massive Feynman integrals*, *Theor. Math. Phys.* **89** (1991) 1052–1063.
- [67] A. I. Davydychev, *Recursive algorithm for evaluating vertex-type Feynman integrals*, *J. Phys. A* **25** (1992) 5587.

- [68] M. F. Paulos, M. Spradlin and A. Volovich, *Mellin Amplitudes for Dual Conformal Integrals*, *JHEP* **08** (2012) 072, [[1203.6362](#)].
- [69] B. Basso and L. J. Dixon, *Gluing Ladder Feynman Diagrams into Fishnets*, *Phys. Rev. Lett.* **119** (2017) 071601, [[1705.03545](#)].
- [70] S. Derkachov, V. Kazakov and E. Olivucci, *Basso-Dixon Correlators in Two-Dimensional Fishnet CFT*, *JHEP* **04** (2019) 032, [[1811.10623](#)].
- [71] F. Loebbert and S. F. Stawinski, *Conformal four-point integrals: recursive structure, Toda equations and double copy*, *JHEP* **11** (2024) 092, [[2408.15331](#)].
- [72] F. Loebbert, J. Miczajka, D. Müller and H. Münkler, *Massive Conformal Symmetry and Integrability for Feynman Integrals*, *Phys. Rev. Lett.* **125** (2020) 091602, [[2005.01735](#)].
- [73] F. Loebbert, J. Miczajka, D. Müller and H. Münkler, *Yangian Bootstrap for Massive Feynman Integrals*, *SciPost Phys.* **11** (2021) 010, [[2010.08552](#)].
- [74] K. C. Rigatos and X. Zhou, *Yangian Symmetry in Holographic Correlators*, *Phys. Rev. Lett.* **129** (2022) 101601, [[2206.07924](#)].
- [75] H. Bateman and A. Erdélyi, *Higher transcendental functions*. California Institute of technology. Bateman Manuscript project. McGraw-Hill, New York, NY, 1953.
- [76] P. O. M. Olsson, *Integration of the Partial Differential Equations for the Hypergeometric Functions F_1 and F_D of Two and More Variables*, *Journal of Mathematical Physics* **5** (03, 1964) 420–430.
- [77] H. M. Srivastava and M. C. Daoust, *Nederl. Akad. Wetensch. Indag. Math.* **31** (1969) 449.
- [78] H. M. Srivastava and P. W. Karlsson, *Multiple gaussian hypergeometric series*, 1985.
- [79] E. B. Vinberg, *Volumes of non-Euclidean polyhedra*, *Russian Math. Surveys* **48** (1993) 15–45.