

#### **DOI:** DOIunavailable

#### Software

- Review 🗗
- Archive ♂

ditor: Pending Editor ♂ Reviewers:

• @Pending Reviewers

Submitted: N/A

#### License

Authors of papers retain copyright and release the work inder a Creative Commons Attribution 4.0 International ficense (CC BY 4.0).

# pylevin: efficient numerical integration of integrals containing up to three Bessel functions

### Robert Reischke\*

Argelander-Institut für Astronomie, Universität Bonn, Auf dem Hügel 71, D-53121 Bonn, Germany

Integrals involving highly oscillatory Bessel functions are notoriously challenging to compute using conventional integration techniques. While several methods are available, they predominantly cater to integrals with at most a single Bessel function, resulting in specialised yet highly optimised solutions. Here we present pylevin, a python package to efficiently compute integrals containing up to three Bessel functions of arbitrary order and arguments. The implementation makes use of Levin's method and allows for accurate and fast integration of these highly oscillatory integrals. In benchmarking pylevin against existing software for single Bessel function integrals, we find its speed comparable, usually within a factor of two, to specialised packages such as FFTLog. Furthermore, when dealing with integrals containing two or three Bessel functions, pylevin delivers performance up to four orders of magnitude faster than standard adaptive quadrature methods, while also exhibiting better stability for large Bessel function arguments. pylevin is available from source via github or directly from pypi.

## Introduction

Bessel functions commonly arise in physical systems exhibiting rotational symmetry. Consequently, theoretical predictions of observables often require evaluating integrals over these functions, which typically cannot be solved analytically and must be addressed numerically. Standard integration methods, such as quadrature, are generally inefficient and unreliable when handling these integrals due to the rapid oscillations characteristic of Bessel functions. Thus, it is essential to develop general tools capable of swiftly and accurately computing these challenging integrals. pylevin addresses this need by facilitating the calculation of the following frequently encountered types of integrals.

$$I_{\ell_1\ell_2\ell_3}(k_1,k_2,k_3) = \int_a^b \mathrm{d}x \, f(x) \prod_{i=1}^N \mathcal{J}_{\ell_i}(k_i x) \,, \quad N = 1, 2, 3 \,, \tag{1}$$

here  $\mathcal{J}_{\ell}(x)$  denotes a spherical,  $j_{\ell}$ , or cylindrical Bessel function,  $J_{\ell}$  of order  $\ell$  and f(x) can be any non-oscillatory function, i.e. that if the product,  $\prod_{i=1}^{n} \mathcal{J}_{\ell_i}(k_i x)$ , has a characteristic frequency  $\omega_{\mathcal{J}}$  then the characteristic frequency of f must satisfy  $\omega_f \ll \omega_{\mathcal{J}}$  over the integration domain. Otherwise, the function f can depend arbitrarily on the integration variable and the parameters  $k_i$ .

In the literature, there exist different methods<sup>1</sup> to solve integrals of the type of Equation (1), with public implementations solving the integral up to N = 2. Ogata (2005) proposed a method which has been for example implemented in the hankel<sup>2</sup> package Murray & Poulin (2019) to calculate Hankel integrals and transforms. These are integrals with the parameters: a = 0, N = 1 and  $\mathcal{J}_{\ell_1}(kx) = J_{\nu}(kx)$  being a cylindrical Bessel function of order  $\nu$ . The same problem is tackled using Fast Fourier Transforms (FFT) in logarithmic space (FFTLog Talman,

<sup>\*</sup>reischke@posteo.net, rreischke@astro.uni-bonn.de

 $<sup>^{1}</sup>$ The list of these excellent methods and packages here will be very biased and is driven by my own field of research.

<sup>&</sup>lt;sup>2</sup>https://hankel.readthedocs.io/en/latest/index.html



1978) implemented for example in pyfftlog package<sup>3</sup> (Hamilton, 2000), implementing discrete versions of the Hankel transformations as well. Another FFTLog implementation can be found in the in the hankl package<sup>4</sup> (Karamanis & Beutler, 2021). Schöneberg et al. (2018)<sup>5</sup> and Fang et al. (2020b)<sup>6</sup> used FFTLog as well to compute angular power spectra in the non-Limber projection, which requires evaluating Equation (1) over a compact range (which can formally extended to infinity) and for N = 1 and  $\mathcal{J}_{\ell_1}(kx) = j_{\ell}(kx)$ . Further applications of FFTLog were used, for example in Fang et al. (2017); Assassi et al. (2017); Grasshorn Gebhardt & Jeong (2018) while Campagne et al. (2017); Chiarenza et al. (2024) rely on Chebyshev polynomials.

FFTLog was further generalised to two dimensions (2DFFTLog) to compute integrals over products of two Bessel functions in Fang et al. (2020a)<sup>7</sup>. A completely different method was suggested by Levin (1996, 1997); Iserles & Nørsett (2005) (Levin's method) which relies on casting the integral into a solution of a differential equation. An adaptive version (see e.g. Chen et al., 2022, for details on convergence, especially for small  $\omega_{\mathcal{J}}$ ) of this method was used in a series of papers (Zieser & Merkel, 2016; Spurio Mancini et al., 2018a,b) to calculate integrals up to N = 2.

All the implementations presented are tailored to very specific applications and are highly optimised. However, they potentially lack flexibility and can only integrate a maximum of two Bessel functions. This is exactly where pylevin enters the stage. It allows for solving the type of integrals presented in Equation (1) and therefore includes all the integrals from the discussed methods and more. The code is written in C++ but is wrapped in python using pybind and is based on implementations used in Zieser & Merkel (2016) which is expanded and heavily optimised. We first discuss the implementation (for a description of the method, we refer to the appendix) and then do benchmarking against some popular codes.

## Implementation

As in Zieser & Merkel (2016), we implement the algorithm in C++ as an adaptive method, i.e. we calculate the estimate of Equation (1) by constructing a solution over the whole integral between [a, b] at using Equation (14) with n collocation points. This is followed by calculating a solution using n/2 collocation points. The interval [a, b] is then recursively bisected in that sub-interval where the relative error, i.e. the relative difference between n/2 and n, is largest. We truncate the recursion once convergence is achieved. The key improvements of pylevin compared to this previous implementation are:

- 1. The inclusion of more integral types, flexibility and an easy high-level interface via python.
- 2. Memory and speed optimisation as well as openmp parallelisation.
- 3. The possibility to reuse precomputed quantities. Specifically, the function f(x) appears solely within the inhomogeneity of the linear system of equations (see Equation 14). As a result, the homogeneous solution can always be established for a specific type of integral and a given bisection. In many practical applications, it is common to compute a fixed type of integral from Equation (1) multiple times, each time with slightly different f(x). A typical example is running a Monte Carlo Markov Chain (MCMC) in cosmology (see the comparison with hankl and pyCCL), where only the matter power spectrum and radial weight function changes, but the general structure of the integrand is kept the same. Hence, pylevin can update the function f(x) and reuse the solution to the homogeneous equation to construct the particular solution for the inhomogeneous

<sup>&</sup>lt;sup>3</sup>https://pyfftlog.readthedocs.io/en/latest/

<sup>&</sup>lt;sup>4</sup>https://github.com/minaskar/hankl

<sup>&</sup>lt;sup>5</sup>https://github.com/lesgourg/class\_public

<sup>&</sup>lt;sup>6</sup>https://github.com/xfangcosmo/FFTLog-and-beyond

<sup>&</sup>lt;sup>7</sup>https://github.com/xfangcosmo/2DFFTLog



solution from Equation (14). This speeds up calculations after the homogeneous solution is known by an order of magnitude.

A basic example of how to run the code can be seen in the code block below. For a comprehensive description, we refer to the readthedocs<sup>8</sup>. Here, we will briefly outline the purpose of the various steps involved:

```
import pylevin as levin
 1
 2
      import numpy as np
 3
     x = np.geomspace(1e-5,100,100)
 4
     number_integrands = 2
y = np.linspace(1,2, number_integrands)
f_of_x = x[:,None]**(3*y[None,:]) + (x**2 +x)[:, None]
 5
 6
 7
 8
     integral_type = 0
N_thread = 1
 9
10
11
      logx = True
      logy = True
12
13
      lp single = levin.pylevin(integral type, x, f of x, logx, logy, N thread)
14
15
     n\_sub~=~10
     n\_bisec\_max = 32
rel_acc = 1e-4
16
17
      boost_bessel = False
18
19
      verbose = False
20
     lp_single.set_levin(n_sub, n_bisec_max, rel_acc, boost_bessel, verbose)
21
     k = np.geomspace(1e-3,1e4,1000)
22
     ell = (5*np.ones_like(k)).astype(int)
a = x[0]*np.ones_like(k)
b = x[-1]*np.ones_like(k)
23
24
25
26
      diagonal = False
     result_levin = np.zeros((len(k), number_integrands))
lp_single.levin_integrate_bessel_single(a, b, k, ell, diagonal, result_levin)
27
28
29
     f_of_x_new = x[:,None]**(2.5*y[None,:]) + (x**1.5 +x)[:, None]
lp_single.update_integrand(x,f_of_x_new, logx, logy)
lp_single.levin_integrate_bessel_single(a, b, k, ell, diagonal, result_levin)
30
31
32
```

- Line 4 6: Defines the non-oscillatory part of the integrand, f(x). The *x*-range should include the minimum *a* and the maximum *b* you want to use. Note that the code allows to pass many integrands at the same time, therefore  $f_{of_x}$  has the shape  $(len(x), number_{integrands})$ . The second dimension always has to exist, even it is one.
- Line 9 13: Defines the type of the integral considered (here a single spherical Bessel function), the number of openmp threads and how the interpolation of the integrand should be carried out. In the last line, an instance of the class is created.
- Line 15 20: Defines the settings of the Levin collocation method. This is optional, as the class assumes default values for each of the parameters. Here we set the number of collocation points, the maximum number of bisections until the relative accuracy is reached, whether the Bessel functions should be computed using the GSL<sup>9</sup> or BOOST<sup>10</sup>.
- Line 22 28: First we set the parameters, a, b, l and k at which the integral should be calculated. These must all be one-dimensional arrays of the same length. In this case, we evaluate the integral at 10<sup>3</sup> different k values for the same values of a, b and l. If the number of integrands is equal to the number of parameters, one can choose to only calculate the diagonal of that square matrix. Here we calculate all values and therefore allocate a (len(k),number\_integrand) array in which the result is stored. If len(k) == number\_integrand and diagonal = True, the result will be an array of length number\_integrand.

<sup>&</sup>lt;sup>8</sup>A python notebook with examples can be found here  $\[mathbb{C}^3$ 

<sup>&</sup>lt;sup>9</sup>https://www.gnu.org/software/gsl/

<sup>&</sup>lt;sup>10</sup>https://www.boost.org



• Line 30 - 32: We can now define a new f(x) over the same x-range and simply update the integrand. By calling the integrator once more, it will utilise the previously constructed bisection, thereby significantly speeding up the calculation, as previously mentioned. However, there are a few caveats to consider: all parameters of the integral must remain unchanged. Furthermore, if the integrand alters sufficiently such that the previously employed bisection is no longer adequate for achieving convergence, the results may become erroneous. This issue may arise, for example, if the asymptotic behaviour of f(x) varies dramatically. Nonetheless, for most practical applications, this is typically not a concern.

## Comparison with various codes

In this section, we compare pylevin with several other, more specialised, codes who mostly solve integrals over single Bessel functions. All computing times presented here are averages over several runs. The comparison was done with an M3 processor with 8 cores (4 performance cores). For pylevin, we always measure the time with the homogeneous solution of Equation (14) known, as this is the case which occurs most in practice, and it is the most efficient way to run the code.

#### hankel

We calculate the following Hankel transformation:

integral(k) = 
$$\int_0^\infty \frac{x^2}{x^2 + 1} J_0(kx) \, \mathrm{d}x$$
, (2)

for 500 values of k logarithmically-spaced between 1 and  $10^4$ . The result is depicted on the left side of Figure 1. It can be seen that both methods agree very well and are roughly equally fast. While the Hankel transformation formally goes from zero to infinity,  $a = 10^{-5}$  and  $b = 10^8$  was used for pylevin. This choice of course depends on the specific integrand.



**Figure 1:** Comparison of pylevin with two methods to calculate a Hankel transformation. Dashed red is pylevin while solid blue is the alternative method. *Left*: Integral in Equation (2) evaluated with the Ogata method using the hankel package. *Right*: Integral in Equation (3) evaluated using the hankl package. Different lines refer to different redshifts.

#### hankl

Here, we follow the cosmology example provided in the hankl documentation to compute the monopole of the galaxy power spectrum:

$$\xi_0(s) = \int_0^\infty \left( b^2 + fb/3 + f^2/5 \right) P_{\rm lin}(k) J_0(ks) k^2 \, \mathrm{d}k \;, \tag{3}$$



where b is the galaxy bias, f the logarithmic growth rate and  $P_{\rm lin}(k)$  is the linear matter power spectrum, which is calculated using camb (Lewis et al., 2000; Lewis & Bridle, 2002; Howlett et al., 2012) at six redshifts. Since hankl is FFT based, it requires k to be discretised, the FFT-dual will then be calculated at the inverse grid points. For this comparison, we use  $2^{10}$ logarithmically-spaced points between  $k = 10^{-4}$  and k = 1 for the transformation to converge. For pylevin, the number of points where the transformation is evaluated is arbitrary. Here we use 100 points, which is more than enough to resolve all features in  $\xi_0$ . The results are shown on the right of Figure 1 and good agreement can be found between the two methods with hankl being roughly twice as fast as pylevin.

#### pyfftlog

For pyfftlog we use the following transformation:

$$FT(k) = \int_0^\infty r^5 e^{-r^2/2} J_4(kr) \, dr \,.$$
(4)

For pyfftlog,  $2^8$  logarithmically-spaced points between  $10^{-4}$  and  $10^4$  for r and hence also for k. pylevin is evaluated for the same number of points, this value could, however, be reduced due to the featureless transformation, thus increasing the speed. In the left panel of Figure 2, the result of this exercise is shown. Good agreement between the two methods is found, with both taken the same amount of time. The large relative error at large values of kis due to the small value of the integral, and hence purely numerical noise.



**Figure 2:** Comparison of pylevin with two other methods, the colour scheme is the same as in Figure 1. *Left*: Transformation defined in Equation (4) with the pyfftlog package. We show the relative difference between the two methods in the lower panel.

#### pyCCL

Here, we compare the implementation of the non-Limber projection for the angular power spectrum:

$$C_{\ell} = \frac{2}{\pi} \int d\chi_1 W(\chi_1) \int d\chi_2 W(\chi_2) \int k^2 dk \ P_{\rm m}(k,\chi_1,\chi_2) j_{\ell}(k\chi_1) j_{\ell}(k\chi_2) , \qquad (5)$$

for W, we assume a Gaussian shell in redshift with width  $\sigma_z = 0.01$  centred at z = 0.6,  $\chi(z)$  is the comoving distance. The matter power spectrum,  $P_{\rm m}$ , is again calculated with camb. The results from pylevin is compared to CCL (Chisari et al., 2019) which implements an FFTLog algorithm (Fang et al., 2020b; Leonard et al., 2023). This implementation first solves the two integrals over  $\chi_{1,2}$  using FFTLog and then carries out the remaining integral over k and assumes that the k and  $\chi_{1,2}$  dependence in the matter power spectrum is separable. To be consistent, we follow the same approach and solve the  $\chi_{1,2}$  integration using pylevin and



then calculate the remaining k integration with the composite Simpson's rule implemented in scipy. The right side of Figure 2 shows that the two methods agree very well with each other and that the method implemented in CCL is about a factor of 2 faster. If the power spectrum, however, would not be separable on small k, as it can be the case in modified gravity scenarios, the CCL method would need to split the integral up into sub-intervals where the separability holds, slowing down the computation by a factor equal to the number of sub-intervals. This assumption is not done in pylevin.

It should be noted that there is also an implementation in julia, which calculates Equation (5) solving the k integration first by expanding  $P_{\rm m}$  into Chebyshev polynomials and precomputing the resulting integrals (Chiarenza et al., 2024). This was shown to be an order of magnitude faster after pre-computation, but is also very tailored to this specific expression.

## Comparison with standard adaptive quadrature

Lastly, we calculate integrals over a product of two and three Bessel functions:

$$I_2 = \int_{10^{-5}}^{100} \mathrm{d}x \; (x^3 + x^2 + x) j_{10}(kx) j_5(kx) \;, \tag{6}$$

$$I_3 = \int_{10^{-5}}^{100} \mathrm{d}x \; (x^3 + x^2 + x) j_{10}(kx) j_5(kx) j_{15}(kx) \;, \tag{7}$$

for  $10^3 k$  values logarithmically spaced between  $10^{-2}$  and  $10^3$ . In Figure 3, the comparison between the adaptive quadrature of scipy and pylevin is shown with the integrals in Equations (6) and (7) on the left and the right respectively. In order for the quadrature to converge over an extended k-range, the number of maximum sub-intervals was increased to  $10^3$  (from the default of 50). The grey-shaded area indicates where the quadrature fails to reach convergence even after this change.



**Figure 3:** Comparison of pylevin with an adaptive quadrature (scipy.integrate.quad, the colour scheme is the same as in Figure 1. *Left*: Integral shown in Equation (6) with the relative difference in the lower panel. *Right*: Integral shown in Equation (7) with the relative difference in the lower panel. The grey shaded area indicates the region where the adaptive quadrature cannot reach convergence and catastrophically fails.

It is therefore clear that pylevin is more accurate and around three orders of magnitudes faster than standard integration routines.

## Summary

We have introduced pylevin, a python implementation of Levin's method designed for the efficient evaluation of integrals involving up to three Bessel functions. The code was compared



to different methods implementing the special case of a single Bessel function based on FFTLog or Ogata's method. Our tests show that pylevin is competitive with these more specialised codes and can reach comparable runtimes. For the case of integrals containing two and three Bessel functions, pylevin outperformed standard adaptive routines by almost four orders of magnitude. pylevin is completely general with respect to the order and the arguments of the Bessel functions in the integral, and therefore presents an efficient and easy to use method to compute these challenging and time-consuming integrals.

## Acknowledgements

I would like to thank Andrina Nicola and Paul Rogozenski for their help fixing issues with the CCL non-Limber comparison.

## References

Assassi V., Simonović M., Zaldarriaga M., 2017, Journal of Cosmology and Astroparticle Physics, 2017, 054

Campagne J. E., Neveu J., Plaszczynski S., 2017, Astronomy & Astrophysics, 602, A72

Chen S., Serkh K., Bremer J., 2022, arXiv e-prints, p. arXiv:2211.13400

Chiarenza S., Bonici M., Percival W., White M., 2024, The Open Journal of Astrophysics, 7, 112

Chisari N. E., et al., 2019, The Astrophysical Journal Supplement Series, 242, 2

Fang X., Blazek J. A., McEwen J. E., Hirata C. M., 2017, Journal of Cosmology and Astroparticle Physics, 2017, 030

Fang X., Eifler T., Krause E., 2020a, Monthly Notices of the Royal Astronomical Society, 497, 2699

Fang X., Krause E., Eifler T., MacCrann N., 2020b, Journal of Cosmology and Astroparticle Physics, 2020, 010

Grasshorn Gebhardt H. S., Jeong D., 2018, Physical Review D, 97, 023504

- Hamilton A. J. S., 2000, Monthly Notices of the Royal Astronomical Society, 312, 257
- Howlett C., Lewis A., Hall A., Challinor A., 2012, Journal of Cosmology and Astroparticle Physics, 4, 27
- Iserles A., Nørsett S. P., 2005, Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences, 461, 1383
- Karamanis M., Beutler F., 2021, hankl: Python implementation of the FFTLog algorithm for cosmology, Astrophysics Source Code Library, record ascl:2112.022

Leonard C. D., et al., 2023, The Open Journal of Astrophysics, 6, 8

Levin D., 1996, Journal of Computational and Applied Mathematics, 67, 95

Levin D., 1997, Journal of Computational and Applied Mathematics, 78, 131

Lewis A., Bridle S., 2002, Physical Review D, D66, 103511

Lewis A., Challinor A., Lasenby A., 2000, Astrophysical Journal, 538, 473

Murray S., Poulin F., 2019, The Journal of Open Source Software, 4, 1397

Ogata H., 2005, Publications of the Research Institute for Mathematical Sciences, 41, 949

Schöneberg N., Simonović M., Lesgourgues J., Zaldarriaga M., 2018, Journal of Cosmology and Astroparticle Physics, 2018, 047



- Spurio Mancini A., Taylor P. L., Reischke R., Kitching T., Pettorino V., Schäfer B. M., Zieser B., Merkel P. M., 2018a, Physical Reiew. D, 98, 103507
- Spurio Mancini A., Reischke R., Pettorino V., Schäfer B. M., Zumalacárregui M., 2018b, Monthly Notices of the Royal Astronomical Society, 480, 3725

Talman J. D., 1978, Journal of Computational Physics, 29, 35

Zieser B., Merkel P. M., 2016, Monthly Notices of the Royal Astronomical Society, 459, 1586



## Appendix: Levin's method

Levin's method (Levin, 1996, 1997) maps a quadrature problem of an oscillatory integral into the solution of a system of ordinary and linear differential equations. By assuming an orthonormal set of basis functions for the solution of the differential equation, the only thing which is required is to find the solution to a simple linear algebra problem. The method relies on integrals of the type:

$$I[\boldsymbol{f}(x,\{k_i\})] = \int_{x_1}^{x_2} \mathrm{d}x \left\langle \boldsymbol{f}(x,\{k_i\}), \mathbf{w}(\{k_i\},x) \right\rangle.$$
(8)

where  $\langle \cdot, \cdot \rangle$  denotes a scalar product. The non-oscillatory functions were bundled into a vector,  $f(x, \{k_i\})$  together with the oscillatory part denoted as  $w(x, \{k_i\})$ . Both depend on the integration variable and a set of external parameters. For the method to work, the oscillatory part, w, must satisfy:

$$\frac{\mathrm{d}\mathbf{w}(x)}{\mathrm{d}x} = \mathbf{A}(x)\mathbf{w}(x), \qquad (9)$$

with a matrix A. If one can find a vector p such that:

$$\langle \mathbf{p}, \mathbf{w} \rangle' = \left\langle \mathbf{p}' + \mathbf{A}^T \mathbf{p}, \mathbf{w} \right\rangle \approx \langle \mathbf{F}, \mathbf{w} \rangle ,$$
 (10)

the integral in Eq. 8 can be approximated by it. Thus, the condition  $\langle \mathbf{p}, \mathbf{w} \rangle' = \langle \mathbf{F}, \mathbf{w} \rangle$ , at n collocation points  $x_j, j = 1, 2, ..., n$ , leads to the following set of equations:

$$\left\langle \mathbf{p}' + \mathbf{A}^T \mathbf{p} - \mathbf{F}, \mathbf{w} \right\rangle (x_j) = 0,$$
 (11)

with the trivial solution:

$$\mathbf{p}'(x_j) + \mathbf{A}^T(x_j)\mathbf{p}(x_j) = \mathbf{F}(x_j).$$
(12)

Next, p is expanded into n differentiable basis function  $u_m$ :

$$\mathbf{p}(x) = \sum_{m=1}^{n} c^{(m)} u_m(x) , \qquad (13)$$

where  $c^{(m)}$  are the coefficients of p in the function basis. Inserting the expansion back into the equation yields

$$\sum_{m=1}^{n} \boldsymbol{c}^{(m)} u'_{m}(x_{j}) + \sum_{m=1}^{n} \mathbf{A}(x_{j}) \ \boldsymbol{c}^{(m)} u_{m}(x_{j}) = \mathbf{F}(x_{j}) , \qquad (14)$$

at the *n* collocation points. For the basis functions,  $u_m$ , we choose the Chebyshev polynomials of the first kind,  $T_n$  by first mapping the integral to the interval [-1,1]. This ensures more numerical stability than using simple polynomials, especially for low frequencies of the oscillatory part w.

The explicit form of  $\mathbf{w}$  and  $\mathbf{A}$  depends on the specific integral. In particular, we implement Equation (1), i.e. integrals with a single Bessel function, a product of two and a product of three Bessel functions. All types are implemented for spherical and cylindrical Bessel functions. We will now write down  $\mathbf{w}$  and  $\mathbf{A}$  explicitly for the cases of spherical Bessel functions by using their recurrence relation. These integrals can be evaluated by defining the following oscillatory



parts

$$\mathbf{w}_{\ell}^{j}(\mathbf{x}, \mathbf{k}) = \begin{pmatrix} j_{\ell}(xk) \\ j_{\ell+1}(xk) \end{pmatrix},$$

$$\mathbf{w}_{\ell_{1},\ell_{2}}^{j}(\mathbf{x}, \mathbf{k_{1}}, \mathbf{k_{2}}) = \begin{pmatrix} j_{\ell_{1}}(xk_{1})j_{\ell_{2}}(xk_{2}) \\ j_{\ell_{1}+1}(xk_{1})j_{\ell_{2}}(xk_{2}) \\ j_{\ell_{1}}(xk_{1})j_{\ell_{2}+1}(xk_{2}) \end{pmatrix},$$

$$\mathbf{w}_{\ell_{1},\ell_{2},\ell_{3}}^{j}(\mathbf{x}, \mathbf{k_{1}}, \mathbf{k_{2}}, \mathbf{k_{3}}) = \begin{pmatrix} j_{\ell_{1}}(xk_{1})j_{\ell_{2}}(xk_{2})j_{\ell_{3}}(xk_{3}) \\ j_{\ell_{1}+1}(xk_{1})j_{\ell_{2}}(xk_{2})j_{\ell_{3}}(xk_{3}) \\ j_{\ell_{1}}(xk_{1})j_{\ell_{2}+1}(xk_{2})j_{\ell_{3}}(xk_{3}) \\ j_{\ell_{1}}(xk_{1})j_{\ell_{2}+1}(xk_{2})j_{\ell_{3}}(xk_{3}) \\ j_{\ell_{1}+1}(xk_{1})j_{\ell_{2}+1}(xk_{2})j_{\ell_{3}+1}(kx_{3}) \\ j_{\ell_{1}+1}(xk_{1})j_{\ell_{2}+1}(xk_{2})j_{\ell_{3}+1}(k_{3}x) \\ j_{\ell_{1}+1}(xk_{1})j_{\ell_{2}+1}(xk_{2})j_{\ell_{3}+1}(k_{3}x) \\ j_{\ell_{1}+1}(xk_{1})j_{\ell_{2}+1}(xk_{2})j_{\ell_{3}+1}(k_{3}x) \\ j_{\ell_{1}+1}(xk_{1})j_{\ell_{2}+1}(xk_{2})j_{\ell_{3}+1}(k_{3}x) \end{pmatrix}.$$
(15)

These would be exactly the same for cylindrical Bessel functions,  $J_\ell$ , by just replacing  $j \to J$ . For the matrix,  $\mathbf{A}^{j/J}$ , one finds:

$$\begin{aligned} \mathbf{A}_{\ell}^{j}(x,k) &= \begin{pmatrix} \frac{\ell}{x} & -k \\ k & -\frac{\ell+2}{x} \end{pmatrix} , \\ \mathbf{A}_{\ell}^{J}(x,k) &= \begin{pmatrix} \frac{\ell}{x} & -k \\ k & -\frac{\ell+1}{x} \end{pmatrix} , \end{aligned}$$
(16)

for j and J respectively. These can now be applied successively to obtain the matrices **A** for the products of Bessel functions. From now on, we will only quote the results for j since the ones for J are easily obtained as well

$$\mathbf{A}_{\ell_1,\ell_2}^j(x,k_1,k_2) = \begin{pmatrix} \frac{\ell_1+\ell_2}{x} & -k_1 & -k_2 & 0\\ k_1 & \frac{\ell_2-\ell_1-2}{x} & 0 & -k_2\\ k_2 & 0 & \frac{\ell_1-\ell_2-2}{x} & -k_1\\ 0 & k_2 & k_1 & -\frac{\ell_1+\ell_2+4}{x} \end{pmatrix},$$
(17)

and

In principle, it is thus easy to generalise the method to any other oscillatory function with some kind of recurrence relation. The user just needs to provide new functions for the function  $\mathbf{w}$  and the matrix  $\mathbf{A}$  and define a new case in the code (that is, as long as the product of oscillatory functions does not extend beyond three).