

Energy spectrum of the long-range Lennard-Jones potential

Shahar Hod

The Ruppin Academic Center, Emeq Hefer 40250, Israel
and

The Hadassah Institute, Jerusalem 91010, Israel

(Dated: February 19, 2025)

The discrete energy spectra of composite inverse power-law binding potentials of the form $V(r; \alpha, \beta, n) = -\alpha/r^2 + \beta/r^n$ with $n > 2$ are studied *analytically*. In particular, using a functional matching procedure for the eigenfunctions of the radial Schrödinger equation, we derive a remarkably compact analytical formula for the discrete spectra of binding energies $\{E(\alpha, \beta, n; k)\}_{k=1}^{k=\infty}$ which characterize the highly-excited bound-state resonances of these long-range binding potentials. Our results are of practical importance for the physics of polarized molecules, the physics of composite polymers, and also for physical models describing the quantum interactions of bosonic particles.

I. INTRODUCTION

Inverse power-law potentials are ubiquitous in physics and chemistry. In particular, they play a central role in physical models describing the interactions of atoms and molecules (see [1–11] and references therein). Of particular interest is the effective Lennard-Jones potential [12]

$$V(r; \alpha, \beta, n_1, n_2)_{\text{LJ}} = -\frac{\alpha}{r^{n_1}} + \frac{\beta}{r^{n_2}} \quad \text{with} \quad n_2 > n_1 > 0, \quad (1)$$

which is composed of two inverse power-law interaction terms: the negative term $-\alpha/r^{n_1}$ describes an attractive interaction between the components of the system, whereas the positive term β/r^{n_2} describes an effective repulsive interaction [13]. The standard form of the Lennard-Jones potential is characterized by the inequality $n_2 > n_1$, in which case the radial function (1) has the form of an effective potential well whose minimum is located at $r^{\text{min}} = (\beta n_2 / \alpha n_1)^{\frac{1}{n_2 - n_1}}$.

The most studied form of the Lennard-Jones binding potential (1) is known as the 6-12 Lennard-Jones potential, which is characterized by an attractive Van der Waals interaction term with $n_1 = 6$. In this case (that is, $n_1 > 2$) one finds that the effective potential well, which is described by the composite inverse power-law functional relation (1), can only support a *finite* number of bound-state resonances [11].

On the other hand, there are known quantum mechanical systems which are characterized by attractive long-range inverse square potentials with $n_1 = 2$ [14–23]. For instance, the inverse square law potential naturally appears in physical models describing the interactions between polar molecules and laser-excited Rydberg atoms [16–19], in the Efimov effect describing the interactions of bosonic particles [20, 21], and also in theoretical models describing the physics of composite polymers [22, 23]. The attractive inverse square potential is especially interesting from physical as well as mathematical points of view since it sits on the boundary between long-range power-law potentials with $n_1 < 2$, which can support an infinite number of bound-state resonances [11], and short-range power-law potentials with $n_1 > 2$ which can only support a finite number of bound-state resonances [11].

The main goal of the present paper is to explore the physical properties of effective binding potentials of the Lennard-Jones form (1) with an attractive *long-range* inverse square law (that is, $n_1 = 2$) interaction term [14–23]. In particular, we shall analyze the discrete energy spectra $\{E(\alpha, \beta, n; k)\}_{k=1}^{k=\infty}$ [24] which characterize the family [25–28]

$$V(r; \alpha, \beta, n)_{\text{LR-LJ}} = -\frac{\alpha}{r^2} + \frac{\beta}{r^n} \quad \text{with} \quad n > 2 \quad (2)$$

of composed long-range inverse power-law binding potentials.

Interestingly, below we shall explicitly show that the Schrödinger differential equation with the effective radial potentials (2) is amenable to an analytical treatment in the regime of small binding energies. In particular, we shall derive a compact *analytical* formula [see Eq. (36) below] for the discrete energy spectra $\{E(\alpha, \beta, n; k)\}_{k=1}^{k=\infty}$ which characterize the highly-excited bound-state resonances of the composed *long-range* Lennard-Jones potentials (2).

II. DESCRIPTION OF THE SYSTEM

We shall study the stationary bound-state ($E < 0$) resonances of the Schrödinger differential equation [29]

$$\left[-\frac{\hbar^2}{2\mu} \frac{d^2}{dr^2} + V(r)_{\text{LR-LJ}} \right] \psi = E\psi , \quad (3)$$

where the spatial behavior of the binding potential $V(r)_{\text{LR-LJ}}$ is described by the composed inverse power-law (long-range Lennard-Jones) functional relation (2). The stationary eigenstates of the radial Schrödinger equation (3) are characterized by spatially bounded radial eigenfunctions:

$$\psi(r \rightarrow \infty) \sim e^{-\omega r} , \quad (4)$$

where the characteristic frequency is defined by the relation [30]

$$\omega^2 \equiv -\frac{2\mu}{\hbar^2} E \quad \text{with} \quad \omega \in \mathbb{R} . \quad (5)$$

In addition, we shall look for bound-state radial eigenfunctions which are spatially regular at the origin:

$$\psi(r \rightarrow 0) < \infty . \quad (6)$$

The Schrödinger differential equation (3), together with the physically motivated boundary conditions (4) and (6), determine the discrete spectra of binding energies $\{E(\alpha, r_0, n)\}$ which characterize the composed inverse power law binding potentials (2). As we shall explicitly show below, the qualitative behavior of the quantum system (3) is determined by the characteristic length-scale

$$r_0 \equiv \left(\frac{2\mu\beta}{\hbar^2} \right)^{1/(n-2)} . \quad (7)$$

In particular, in the next section we shall use a functional matching procedure for the characteristic eigenfunctions of the radial Schrödinger equation (3) in order to show that, in the regime [31]

$$\omega r_0 \ll 1 \quad (8)$$

of weakly bound-state resonances, the discrete energy spectra $\{E(\alpha, r_0, n)\}$ of the composite inverse power law binding potentials (2) [the long-range Lennard-Jones potentials (2)] can be determined *analytically*.

III. ASYMPTOTIC ANALYTICAL SOLUTIONS OF THE RADIAL SCHRÖDINGER EQUATION

In the present section we shall analyze the Schrödinger differential equation

$$\left\{ \frac{d^2}{dr^2} - \frac{1}{r^2} \left[(\omega r)^2 - \left(\frac{1}{4} + \nu^2 \right) + \left(\frac{r_0}{r} \right)^{n-2} \right] \right\} \psi(r; \omega, \nu, r_0, n) = 0 \quad (9)$$

for the long-range Lennard-Jones [26] binding potentials (2), where [32]

$$\frac{1}{4} + \nu^2 \equiv \frac{2\mu\alpha}{\hbar^2} \quad ; \quad \nu \in \mathbb{R} . \quad (10)$$

In particular, we shall explicitly show below that the radial Schrödinger equation (9), which determines the energy spectra of the composite inverse power-law potentials (2), is amenable to an analytical treatment in the asymptotic radial regions $r \ll 1/\omega$ and $r \gg r_0$. We shall then show that, by using a functional matching procedure for the characteristic radial solutions of (9) in the overlapping radial region $r_0 \ll r \ll 1/\omega$, one can determine *analytically* the discrete spectra of eigen-frequencies $\{\omega(\nu, r_0, n)\}$ which characterize the weakly-bound (highly-excited) resonances of the long-range Lennard-Jones potentials (2) [26].

We shall first obtain an analytical expression for the radial eigenfunction $\psi(r)$ in the radial region

$$r \ll 1/\omega , \quad (11)$$

in which case one may approximate the Schrödinger differential equation (9) by

$$\left(\frac{d^2}{dr^2} + \frac{1}{4} + \nu^2 - \frac{r_0^{n-2}}{r^n}\right)\psi = 0. \quad (12)$$

The general mathematical solution of the differential equation (12) can be expressed in terms of the Hankel functions of the first and second kinds (see Eq. 9.1.53 of [33]):

$$\psi(r) = A_1 r^{\frac{1}{2}} H_{\frac{2\bar{\nu}}{n-2}}^{(1)} \left[\frac{2i}{n-2} \left(\frac{r_0}{r}\right)^{(n-2)/2} \right] + A_2 r^{\frac{1}{2}} H_{\frac{2\bar{\nu}}{n-2}}^{(2)} \left[\frac{2i}{n-2} \left(\frac{r_0}{r}\right)^{(n-2)/2} \right], \quad (13)$$

where $\{A_1, A_2\}$ are normalization constants and $\bar{\nu} \equiv i\nu$.

Using the large-argument ($r_0/r \rightarrow \infty$) asymptotic behaviors

$$H_a^{(1)}(x \rightarrow \infty) = \frac{(1-i)e^{-ia\pi/2} e^{ix}}{\sqrt{\pi}} \cdot [1 + O(x^{-1})] \quad ; \quad H_a^{(2)}(x \rightarrow \infty) = \frac{(1+i)e^{ia\pi/2} e^{-ix}}{\sqrt{\pi}} \cdot [1 + O(x^{-1})] \quad (14)$$

of the Hankel functions, one finds from (13) the limiting behavior

$$\psi(r \rightarrow 0) = r_0^{\frac{1}{2}} \left(\frac{n-2}{\pi}\right)^{\frac{1}{2}} \left(\frac{r}{r_0}\right)^{\frac{n}{4}} \cdot \left\{ -iA_1 e^{\frac{\nu\pi}{n-2}} \exp\left[-\frac{2}{n-2} \left(\frac{r_0}{r}\right)^{(n-2)/2}\right] + A_2 e^{-\frac{\nu\pi}{n-2}} \exp\left[\frac{2}{n-2} \left(\frac{r_0}{r}\right)^{(n-2)/2}\right] \right\} \quad (15)$$

for the radial eigenfunctions near the origin. The physically acceptable bound-state resonances of the binding potentials (2) are characterized by regular radial eigenfunctions at the origin $r \rightarrow 0$ [see the boundary condition (6)]. One therefore concludes that the coefficient of the exploding exponent in (15) should vanish:

$$A_2 = 0. \quad (16)$$

Using the small-argument ($r_0/r \ll 1$) asymptotic behavior

$$H_a^{(1)}(x \rightarrow 0) = \left\{ \left[\frac{1}{\Gamma(1+a)} - i \frac{\cos(a\pi)\Gamma(-a)}{\pi} \right] \cdot (x/2)^a - i \frac{\Gamma(a)}{\pi} \cdot (x/2)^{-a} \right\} \cdot [1 + O(x^2)] \quad (17)$$

of the Hankel function, one finds from (13) the power-law behavior

$$\psi(r) = A_1 \left\{ r_0^{\frac{1}{2}} \left(\frac{i}{n-2}\right)^{\frac{2\bar{\nu}}{n-2}} \left[\frac{1}{\left(\frac{2\bar{\nu}}{n-2}\right)\Gamma\left(\frac{2\bar{\nu}}{n-2}\right)} - i \frac{\cos\left(\frac{2\bar{\nu}}{n-2}\pi\right)\Gamma\left(-\frac{2\bar{\nu}}{n-2}\right)}{\pi} \right] \cdot \left(\frac{r}{r_0}\right)^{\frac{1}{2}-\bar{\nu}} - i r_0^{\frac{1}{2}} \frac{\Gamma\left(\frac{2\bar{\nu}}{n-2}\right)}{\pi \left(\frac{i}{n-2}\right)^{\frac{2\bar{\nu}}{n-2}}} \cdot \left(\frac{r}{r_0}\right)^{\frac{1}{2}+\bar{\nu}} \right\} \quad (18)$$

for the characteristic radial eigenfunctions of the Schrödinger differential equation (9) in the intermediate radial region

$$r_0 \ll r \ll 1/\omega. \quad (19)$$

We shall next obtain an analytical expression for the radial eigenfunction $\psi(r)$ in the radial region

$$r \gg r_0, \quad (20)$$

in which case one may approximate the Schrödinger differential equation (9) by

$$\left(\frac{d^2}{dr^2} - \omega^2 + \frac{1}{4} + \nu^2\right)\psi = 0. \quad (21)$$

The general mathematical solution of the differential equation (21) can be expressed in terms of the Hankel functions of the first and second kinds (see Eq. 9.1.49 of [33])

$$\psi(r) = B_1 r^{\frac{1}{2}} H_{\bar{\nu}}^{(1)}(\bar{\omega}r) + B_2 r^{\frac{1}{2}} H_{\bar{\nu}}^{(2)}(\bar{\omega}r), \quad (22)$$

where $\{B_1, B_2\}$ is a normalization constants and $\bar{\omega} \equiv i\omega$.

Using the large-argument ($\omega r \rightarrow \infty$) asymptotic behaviors (14) of the Hankel functions, one finds from (22) the asymptotic behavior

$$\psi(r \rightarrow \infty) = B_1 \frac{-\sqrt{2}ie^{\nu\pi/2}}{\sqrt{\omega\pi}} e^{-\omega r} + B_2 \frac{\sqrt{2}e^{-\nu\pi/2}}{\sqrt{\omega\pi}} e^{\omega r} \quad (23)$$

for the radial eigenfunctions. The physically acceptable bound-state resonances of the effective radial potentials (2) are characterized by finite (exponentially decaying) radial eigenfunctions at spatial infinity $r \rightarrow \infty$ [see the boundary condition (4)]. One therefore concludes that the coefficient of the exploding exponent in (23) should vanish [34]:

$$B_2 = 0 . \quad (24)$$

Using the small-argument ($\omega r \ll 1$) asymptotic behavior (17) of the Hankel function, one finds from (22) the power-law behavior

$$\psi(r) = B_1 \left\{ (\bar{\omega}/2)^{\bar{\nu}} \left[\frac{1}{\bar{\nu}\Gamma(\bar{\nu})} - i \frac{\cos(\bar{\nu}\pi)\Gamma(-\bar{\nu})}{\pi} \right] r^{\frac{1}{2}+\bar{\nu}} - i \frac{\Gamma(\bar{\nu})}{\pi(\bar{\omega}/2)^{\bar{\nu}}} r^{\frac{1}{2}-\bar{\nu}} \right\} \quad (25)$$

for the radial eigenfunctions which characterize the Schrödinger differential equation (9) in the intermediate radial region

$$r_0 \ll r \ll 1/\omega . \quad (26)$$

IV. THE DISCRETE ENERGY SPECTRUM OF THE HIGHLY-EXCITED BOUND-STATE RESONANCES

In the previous section we have explicitly shown that, for small resonant eigen-frequencies in the regime $\omega r_0 \ll 1$ [35], there is an overlap radial region $r_0 \ll r \ll 1/\omega$ [see Eqs. (19) and (26)] in which the characteristic radial eigenfunctions $\psi(r)$ of the Schrödinger differential equation (9) are described by the two analytically derived mathematical expressions (13), (16), (22), and (24). In particular, inspecting the mathematical solutions (18) and (25) in the overlap region $r_0 \ll r \ll 1/\omega$ (where *both* solutions are valid), one immediately realizes that these two expressions for the radial eigenfunctions $\psi(r)$ are characterized by the same functional (power-law) behavior. Matching the coefficients of the power-law $r^{\frac{1}{2}+i\nu}$ terms in (18) and (25), one finds the dimensionless ratio [36]

$$\frac{B_1}{A_1} = \frac{\bar{\nu}\Gamma(\bar{\nu})\Gamma(\frac{2\bar{\nu}}{n-2})}{i\pi[1+i\cot(\bar{\nu}\pi)](\frac{i}{n-2})^{\frac{2\bar{\nu}}{n-2}}(\bar{\omega}r_0/2)^{\bar{\nu}}} . \quad (27)$$

Likewise, matching the coefficients of the power-law $r^{\frac{1}{2}-i\nu}$ terms in (18) and (25), one finds the dimensionless ratio [36]

$$\frac{B_1}{A_1} = \frac{i\pi[1+i\cot(\frac{2\bar{\nu}}{n-2}\pi)](\frac{i}{n-2})^{\frac{2\bar{\nu}}{n-2}}(\bar{\omega}r_0/2)^{\bar{\nu}}}{(\frac{2\bar{\nu}}{n-2})\Gamma(\bar{\nu})\Gamma(\frac{2\bar{\nu}}{n-2})} . \quad (28)$$

Taking cognizance of Eqs. (27) and (28), one finds the characteristic resonance equation

$$(\bar{\omega}r_0/2)^{2\bar{\nu}} = \frac{2}{(n-2)[1+i\cot(\bar{\nu}\pi)][1+i\cot(\frac{2\bar{\nu}}{n-2}\pi)]} \left[\frac{\nu\Gamma(\bar{\nu})\Gamma(\frac{2\bar{\nu}}{n-2})}{\pi(\frac{i}{n-2})^{\frac{2\bar{\nu}}{n-2}}} \right]^2 . \quad (29)$$

It is important to emphasize again that the analytically derived resonance equation (29) for the characteristic binding energies (resonant frequencies) of the effective long-range Lennard-Jones potentials (2) [26] is valid in the small frequency regime [see Eqs. (19) and (26)][37]

$$\omega r_0 \ll 1 . \quad (30)$$

From Eq. (29) we finally obtain the discrete resonance spectrum [38]

$$\omega r_0 = 2(n-2)^{\frac{2}{n-2}} \left\{ \frac{2e^{\frac{\pi\nu n}{n-2}}}{(n-2)[1+\coth(\nu\pi)][1+\coth(\frac{2\nu}{n-2}\pi)]} \left[\frac{\nu\Gamma(\bar{\nu})\Gamma(\frac{2\bar{\nu}}{n-2})}{\pi} \right]^2 \right\}^{\frac{1}{2\bar{\nu}}} \times e^{-\frac{\pi k}{\bar{\nu}}} ; \quad k \in \mathbb{Z} \quad (31)$$

which characterizes the weakly bound-state (highly-excited) resonances of the radial Schrödinger equation (3) with the effective long-range binding potentials (2). It is worth pointing out that the dimensionless expression on the r.h.s of Eq. (31) can be expressed in a more compact form. In particular, denoting

$$\phi_1 \equiv \arg[\Gamma(\bar{\nu})] ; \quad \phi_2 \equiv \arg\left[\Gamma\left(\frac{2\bar{\nu}}{n-2}\right)\right] , \quad (32)$$

and using the identity (see Eq. 6.1.43 of [33])

$$\Re[\Gamma(ix)] = \left[\frac{\pi}{x \sinh(\pi x)} \right]^{\frac{1}{2}}, \quad (33)$$

one can express the dimensionless resonance spectrum (31) in the remarkably compact form

$$\omega r_0 = 2(n-2)^{\frac{2}{n-2}} e^{\frac{\phi_1 + \phi_2}{\nu}} \times e^{-\frac{\pi k}{\nu}}; \quad k \in \mathbb{Z}. \quad (34)$$

V. SUMMARY

The discrete energy spectra of composite *long-range* Lennard-Jones potentials of the form (2) were studied *analytically*. Interestingly, using a functional matching procedure for the characteristic eigenfunctions of the radial Schrödinger equation (9), we have derived a remarkably compact analytical formula for the dimensionless binding energies

$$\mathcal{E}(\nu, n; k) \equiv \left(\frac{2\mu\beta_n^{\frac{2}{n}}}{\hbar^2} \right)^{\frac{n}{n-2}} \cdot E(\nu, n; k) \quad (35)$$

which characterize the highly-excited bound-state resonances of the long-range Lennard-Jones potentials (2) [26] [see Eqs. (5), (7), (32), and (34)]:

$$\mathcal{E}(\nu, n; k) = - \left[2(n-2)^{\frac{2}{n-2}} e^{\frac{\phi}{\nu}} \right]^2 \times e^{-\frac{2\pi k}{\nu}}; \quad k \in \mathbb{Z}, \quad (36)$$

where

$$\phi \equiv \arg[\Gamma(\bar{\nu})] + \arg\left[\Gamma\left(\frac{2\bar{\nu}}{n-2}\right)\right]. \quad (37)$$

Finally, it is worth noting that the long-range binding potentials (2) studied in the present paper are characterized by an attractive inverse square law interaction term which naturally appears in diverse quantum systems (see [14–23] and references therein). Thus, our results may be of practical importance in the physics of polarized molecules [16–19] and in the physics of composite polymers [22, 23] where this long-range interaction term naturally appears. In addition, the famous Efimov effect [20, 21] is characterized by the inverse square law potential term, and thus our results may also be of practical importance for physical models describing the quantum interactions of bosonic particles [39].

ACKNOWLEDGMENTS

This research is supported by the Carmel Science Foundation. I thank Yael Oren, Arbel M. Ongo, Ayelet B. Lata, and Alona B. Tea for stimulating discussions.

-
- [1] H. R. Thorsheim, J. Weiner, and P. S. Julienne, *Phys. Rev. Lett.* **58**, 2420 (1987).
 - [2] J. D. Miller, R. A. Cline, and D. J. Heinzen, *Phys. Rev. Lett.* **71**, 2204 (1993).
 - [3] R. A. Cline, J. D. Miller, and D. J. Heinzen, *Phys. Rev. Lett.* **73**, 632 (1994).
 - [4] C. J. Williams and P. S. Julienne, *J. Chem. Phys.* **101**, 2634 (1994).
 - [5] R. Cote, A. Dalgarno, and M. J. Jamieson, *Phys. Rev. A* **50**, 399 (1994).
 - [6] H. Wang, P. L. Gould, and W. C. Stwalley, *Phys. Rev. A* **53**, R1216 (1996).
 - [7] B. Gao, *Phys. Rev. A* **58**, 1728 (1998).
 - [8] J. Trost, C. Eltschka, and H. Friedrich, *J. Phys. B: At. Mol. Opt. Phys.* **31**, 361 (1998).
 - [9] B. Gao, *Phys. Rev. A* **59**, 2778 (1999).
 - [10] B. Gao, *Phys. Rev. Lett.* **83**, 4225 (1999).
 - [11] L. D. Landau and E. M. Lifshitz, *Quantum Mechanics. Non-relativistic theory* (Mir, Moscow, 1974).
 - [12] J. E. Lennard-Jones, *Proc. R. Soc. Lond. A*, **106**, 463 (1924).
 - [13] The repulsive interaction term in the Lennard-Jones potential (1) is usually attributed in the literature to the quantum effect of Pauli repulsion at short distances.
 - [14] Mott and Massey, *Theory of Atomic Collisions* (Clarendon Press, Oxford, 1949), p. 30.

- [15] K. M. Case, Phys. Rev. **80**, 797 (1950).
- [16] J.-M. Lévy-Leblond, Phys. Rev. **153**, 1 (1967).
- [17] O. H. Crawford, Proc. Phys. Soc. London **91**, 279 (1967).
- [18] C. Desfrancois, H. Abdoul-Carime, N. Khelifa, and J. P. Schermann, Phys. Rev. Lett. **73**, 2436 (1994).
- [19] H. E. Camblong, L. N. Epele, H. Fanchiotti, and C. A. Garcia Canal, Phys. Rev. Lett. **85**, 1590 (2000); H. E. Camblong, L. N. Epele, H. Fanchiotti, and C. A. Garcia Canal, Phys. Rev. Lett. **87** 220402 (2001).
- [20] V. Efimov, Phys. Lett. B **33**, 563 (1970); V. Efimov, Nucl. Phys. A **210**, 157 (1973).
- [21] T. Kraemer et al, Nature **440**, 315 (2006).
- [22] E. Marinari and G. Parisi, Europhys. Lett. **15**, 721 (1991).
- [23] C. Nisoli and B. R. Bishop, Phys. Rev. Lett. **112**, 070401 (2014).
- [24] Here k is the resonance parameter of the bound-state resonance [see Eq. (31) below].
- [25] Here the subscript LR-LJ stands for Long-Range Lennard-Jones potentials.
- [26] In the present paper we use the term ‘long-range Lennard-Jones potentials’ to describe composite inverse power-law binding potentials of the Lennard-Jones type (2) whose attractive interaction term decays asymptotically as an inverse square law with $n_1 = 2$ [this long-range attractive behavior should be contrasted with the standard Lennard-Jones potentials (1) whose attractive interaction terms decay asymptotically as inverse power laws with $n_1 > 2$].
- [27] It is important to emphasize that the physical case studied in the present work refers to eigenstates with square angular number l less than a critical value, determined by α , above which the total potential, which vanishes asymptotically as $1/r^2$, becomes repulsive due to the repulsive centrifugal force.
- [28] It is worth noting that one learns from the effective binding potential (2) that the effective radius \bar{r} of the radial eigenfunctions can be estimated by the functional expression $\bar{r} = n\beta/2\alpha$. It should be emphasized that, for physical systems which are made of molecular gases, our analytical treatment is valid provided the effective radius is less than the average half distance between the molecules.
- [29] Here μ stands for the effective (reduced) mass of the quantum system.
- [30] We shall assume, without loss of generality, that $\omega > 0$.
- [31] It is worth noting that the small frequency regime (8) that we shall explore in the present paper corresponds to small binding energies in the regime $(2\mu\beta^{2/n}/\hbar^2)^{n/(n-2)}E \ll 1$ [see Eqs. (5) and (7)].
- [32] We shall assume, without loss of generality, that $\nu > 0$.
- [33] M. Abramowitz and I. A. Stegun, *Handbook of Mathematical Functions* (Dover Publications, New York, 1970).
- [34] We recall that $0 < \omega \in \mathbb{R}$ [see Eq. (5) and [30]]. One therefore finds the physically acceptable asymptotic spatial behavior $\psi(r \rightarrow \infty) \sim e^{-\omega r} \rightarrow 0$ for $B_2 = 0$.
- [35] Note that this frequency regime corresponds to weakly bound-state (highly-excited) resonances of the radial Schrödinger equation (9).
- [36] Here we have used the functional identity $-x\Gamma(x)\Gamma(-x) = \pi/\sin(\pi x)$ (see Eq. 6.1.17 of [33]) in Eqs. (18) and (25).
- [37] Note, in particular, that the two strong inequalities $r_0 \ll r \ll 1/\omega$ in Eqs. (19) and (26) roughly correspond to an order-of-magnitude difference ($r_0/r \lesssim 10^{-1}$ and $\omega r \lesssim 10^{-1}$) between the relevant physical parameters. One therefore realizes that the analytically derived equation (29) for the characteristic resonant frequencies of the effective long-range Lennard-Jones potentials (2) [26] is expected to be valid in the regime $\omega r_0 \lesssim 10^{-2}$ of small binding energies.
- [38] Here we have used the relation $1 = e^{i2\pi k}$, where the resonance parameter k is an integer. In addition, we have used here the functional identity $i \cot(ix) = \coth(x)$.
- [39] Y. Nishida, Y. Kato, and C. D. Batista, Nature Physics **9**, 93 (2013).