Stability Bounds for Smooth Optimal Transport Maps and their Statistical Implications

Sivaraman Balakrishnan[†] and Tudor Manole[¢]

[†] Department of Statistics and Data Science Machine Learning Department Carnegie Mellon University siva@stat.cmu.edu

Statistics and Data Science Center Massachusetts Institute of Technology tmanole@mit.edu

February 19, 2025

Abstract

We study estimators of the optimal transport (OT) map between two probability distributions. We focus on plugin estimators derived from the OT map between estimates of the underlying distributions. We develop novel stability bounds for OT maps which generalize those in past work, and allow us to reduce the problem of optimally estimating the transport map to that of optimally estimating densities in the Wasserstein distance. In contrast, past work provided a partial connection between these problems and relied on regularity theory for the Monge-Ampère equation to bridge the gap, a step which required unnatural assumptions to obtain sharp guarantees. We also provide some new insights into the connections between stability bounds which arise in the analysis of plugin estimators and growth bounds for the semi-dual functional which arise in the analysis of Brenier potential-based estimators of the transport map. We illustrate the applicability of our new stability bounds by revisiting the smooth setting studied by Manole et al. [2024], analyzing two of their estimators under more general conditions. Critically, our bounds do not require smoothness or boundedness assumptions on the underlying measures. As an illustrative application, we develop and analyze a novel tuning parameter-free estimator for the OT map between two strongly log-concave distributions.

1 Introduction

The field of optimal transport (OT) is centered on the following question: Given two probability measures P and Q supported in \mathbb{R}^d , how can we transport P to Q while minimizing the transportation cost? For the squared Euclidean distance cost, an OT map T_0 from P to Q is any solution to the *Monge problem* [Monge, 1781],

$$\underset{T:T_{\#}P=Q}{\operatorname{argmin}} \int \|x - T(x)\|_2^2 \, dP(x), \tag{1}$$

where the minimizer is chosen from transport maps between P and Q, that is, the set of Borel-measurable functions $T : \mathbb{R}^d \to \mathbb{R}^d$ such that $T_{\#}P := P(T^{-1}(\cdot)) = Q$. Intuitively, a transport map T "transports" a random variable X with distribution P to a random variable T(X) which has distribution Q.

One of the central preoccupations of the field of *statistical* optimal transport [Chewi et al., 2024] is to understand the estimation of objects which arise from the OT framework—for instance, the optimal transport map T_0 or the optimal transport cost—when the two measures are unknown, but we have samples from them. Our focus in this paper is on estimating the OT map from samples. Concretely, given i.i.d. samples $X_1, \ldots, X_n \sim P$ and $Y_1, \ldots, Y_m \sim Q$ our goal is to construct an estimate \hat{T}_{nm} of the OT map T_0 between P and Q, for which the risk

$$R(\widehat{T}_{nm}, T_0) := \mathbb{E}\left[\int \|\widehat{T}_{nm}(x) - T_0(x)\|_2^2 dP(x)\right]$$
(2)

is small. Here, the outer expectation is taken over the randomness of the two samples. Two of the broad classes of estimators that have been studied in the literature are *plugin estimators*, which construct estimates of T_0 based on the optimal transport map between estimates \hat{P} and \hat{Q} of the underlying distribution [Chernozhukov et al., 2017, Deb and Sen, 2023, Ghosal and Sen, 2022, Gunsilius, 2022, Manole et al., 2024], and *dual estimators*, which are based on a characterization (see Theorems 1 and 2) of the optimal transport map as the gradient of a convex function φ_0 which solves the so-called *semi-dual* optimization problem:

$$\varphi_0 \in \operatorname*{argmin}_{\varphi \in L^1(P)} \int \varphi dP + \int \varphi^* dQ.$$
(3)

We refer to the quantity,

$$\mathcal{S}_{P,Q}(\varphi) := \int \varphi dP + \int \varphi^* dQ \tag{4}$$

as the semi-dual functional. Dual estimators approximate the solution to the program (3) building on standard ideas from the M-estimation literature [van de Geer, 2000, van der Vaart and Wellner, 1996], and then compute an estimate $\hat{T} = \nabla \hat{\varphi}$ [Ding et al., 2024, Divol et al., 2022, Hütter and Rigollet, 2021]. In contrast to plugin estimators, the computation of dual estimators is often impractical because computing the convex conjugate φ^* in the program above can be challenging. On the other hand, one can directly bring to bear powerful ideas from the analysis of M-estimation to sharply characterize the statistical performance of dual estimators and as a result they serve as a useful information-theoretic benchmark.

Central to the study of statistical properties of plugin estimators are stability bounds which aim to quantify the distance between the optimal transport map between \hat{P} and \hat{Q} and the one between P and Q, i.e. they quantify the stability of the optimal transport map between two distributions to perturbations of the distributions [Deb et al., 2021, Manole et al., 2024]. The study of dual estimators has on the other hand mirrored the study of M-estimators. As highlighted by the work of Hütter and Rigollet [2021], a key quantity in analyzing dual estimators is the so-called growth rate of the semi-dual functional, i.e. the rate at which the difference $S_{P,Q}(\varphi) - S_{P,Q}(\varphi_0)$ grows as a function of the distance between φ and φ_0 . Concretely, Hütter and Rigollet [2021] showed that under a certain curvature assumption on φ_0 , the growth rate is quadratic in the following sense:

$$\mathcal{S}_{P,Q}(\varphi) - \mathcal{S}_{P,Q}(\varphi_0) \asymp \int \|\nabla\varphi(x) - \nabla\varphi_0(x)\|_2^2 dP(x).$$
(5)

This quadratic growth, together with empirical process techniques from the analysis of Mestimators, has led to sharp statistical bounds for dual estimators [Divol et al., 2022, Hütter and Rigollet, 2021]. In our work, we shed some light on the connections between the growth rate of the semi-dual functional and stability bounds by showing that many existing and new stability bounds can be derived by studying the growth rate of appropriate semi-dual functionals.

Our Contributions: Our primary contribution (Theorem 3) is an improvement to the stability bounds of Manole et al. [2024]. We recover their one-sample stability bound and improved versions of their two-sample and empirical stability bounds as direct consequences of a single unified result. This new stability bound has immediate implications for the analysis of plugin estimators of the OT map. As an important example, we obtain new results for the estimation of the optimal transport map between distributions with smooth densities, in Section 4. Similar results were previously obtained by Manole et al. [2024] conditional on a uniform boundary regularity assumption holding. In order to obtain unconditional results, the authors made the unnatural assumption that the sampling distributions were supported on the d-dimensional flat torus. Using our new stability bound we provide an unconditional result with a much simpler proof, completely bypassing the regularity issues. We provide another illustration of the generality of our stability bounds by using them to provide guarantees for a new, practical estimator of the optimal transport map between two log-smooth and strongly log-concave distributions.

Organization: We begin with some background on optimal transport and introduce the semi-dual functional formally in Section 2. We state our new stability bound in Section 3 and discuss how it relates to primal and dual stability bounds developed in past work. In Section 4 we develop consequences of our stability bound in estimating the optimal transport map between two smooth distributions, highlighting how we are able to close some important gaps in this literature. In Section 5 we consider the problem of estimating the optimal transport map between two strongly smooth and strongly log-concave distributions. We conclude in Section 6.

Notation: For a domain $\Omega \subseteq \mathbb{R}^d$, we let $\mathcal{P}(\Omega)$ denote the set of Borel probability measures sures supported on Ω , and $\mathcal{P}_2(\Omega)$ be the subset of such probability measures with finite second moment. Similarly, let $\mathcal{P}_{ac}(\Omega)$ (resp. $\mathcal{P}_{2,ac}(\Omega)$) denote the set of probability measures in $\mathcal{P}(\Omega)$ (resp. $\mathcal{P}_2(\Omega)$) which are absolutely continuous with respect to the Lebesgue measure. We denote by $\mathcal{C}^s(\Omega)$ the standard Hölder space with real and positive exponent s > 0over Ω [Gilbarg and Trudinger, 2001], which is sometimes denoted $\mathcal{C}^{\lfloor s \rfloor, s - \lfloor s \rfloor}(\Omega)$. Given a vector field $T : \mathbb{R}^d \to \mathbb{R}^d$ and a measure μ on \mathbb{R}^d , we write $||T||^2_{L^2(\mu)} = \int_{\mathbb{R}^d} ||T(x)||^2_2 d\mu(x)$. For a pair of distributions $P, Q \in \mathcal{P}(\mathbb{R}^d)$, we denote by $\varphi_{P,Q}$ a *Brenier potential*, i.e. any potential which solves the semi-dual problem (3). For a convex function φ defined on Ω , we denote by φ^* its Fenchel conjugate. In general, φ^* takes values in the extended reals and is defined by $\varphi^*(y) := \sup_{x \in \Omega} [\langle x, y \rangle - \varphi(x)]$. For sequences $(a_n)_{n=1}^{\infty}$ and $(b_n)_{n=1}^{\infty}$, we write $a_n \leq b_n$ if there exists C > 0 such that $a_n \leq Cb_n$ for all $n \geq 1$. Following the convention in nonparametric statistics, the constant C above can depend on the dimension d, and other problem parameters when explicitly mentioned, but is otherwise universal.

2 Background

We work with the following statistical setup: We obtain i.i.d. samples $X_1, \ldots, X_n \sim P$ and $Y_1, \ldots, Y_m \sim Q$, where $P, Q \in \mathcal{P}_2(\Omega)$ and the distributions P, Q belong to some structured collection of distributions \mathcal{R} . In Section 4 our focus will be on the setting where \mathcal{R} is a collection of distributions with Hölder smooth densities, while in Section 5 we will consider the case where \mathcal{R} is the collection of strongly log-concave and log-smooth distributions. Our goal is to construct an estimator \widehat{T}_{nm} such that the risk in (2) is uniformly small over the collection \mathcal{R} , i.e. we are interested in constructing estimators for which the maximum risk:

$$\sup_{P,Q\in\mathcal{R}} R(\widehat{T}_{nm},T_0)$$

is small.

The Monge problem in (1) does not always have a solution, and the push-forward constraint is nonlinear in general. These aspects motivate the study of the Kantorovich relaxation [Kantorovich, 1942, 1948] of identifying an *optimal coupling* between P and Q, defined as:

$$\pi_0 \in \operatorname*{argmin}_{\pi \in \Pi(P,Q)} \int \|x - y\|_2^2 d\pi(x, y), \tag{6}$$

where the set $\Pi(P,Q)$ denotes the set of couplings of P and Q, i.e.

$$\Pi(P,Q) = \{ \pi \in \mathcal{P}_2(\Omega \times \Omega) : \pi(\cdot \times \Omega) = P, \pi(\Omega \times \cdot) = Q \}.$$

In contrast to the Monge problem (1), the Kantorovich problem (6) is always feasible, and is a linear program. From (6) we can derive the following intuitive optimality property: for any pair of random variables U with distribution P, and V with distribution Q we have that:

$$\mathbb{E}_{(X,Y)\sim\pi_0}\langle X,\,Y\rangle \ge \mathbb{E}\langle U,\,V\rangle,\tag{7}$$

i.e. the optimal coupling, among all pairs with given marginals, defines a pair of random variables with maximum correlation.

2.1 Brenier's Theorem and Alternatives

The Kantorovich problem is a relaxation of Monge's problem—every transport map between distributions defines a coupling of the distributions. In contrast to Monge's problem, the Kantorovich relaxation is always feasible, and is a linear program. A cornerstone result in OT theory, is Brenier's theorem which relates the solutions of the Monge and Kantorovich problems:

Theorem 1 (Brenier's Theorem). Let $P \in \mathcal{P}_{2,ac}(\Omega)$ and $Q \in \mathcal{P}_2(\Omega)$. Then, the following assertions hold.

- 1. There exists an optimal transport map T_0 pushing P forward onto Q which takes the form $T_0 = \nabla \varphi_0$ for a convex function $\varphi_0 : \mathbb{R}^d \to \mathbb{R}$ which solves the semi-dual problem (3). Furthermore, T_0 is uniquely determined P-almost everywhere.
- 2. If we further have $Q \in \mathcal{P}_{2,ac}(\Omega)$, then $S_0 := \nabla \varphi_0^*$ is the Q-almost everywhere uniquely determined optimal transport map pushing Q forward onto P. Furthermore, for Lebesguealmost every $x, y \in \Omega$,

$$\nabla \varphi_0^* \circ \nabla \varphi_0(x) = x, \quad \nabla \varphi_0 \circ \nabla \varphi_0^*(y) = y.$$

Our stability bounds also hold in some cases where Brenier's theorem does not apply. Indeed, we will generally only require the measures P and Q to be chosen such that the following assumption holds:

A0 There exists a strictly convex and differentiable function $\varphi_0 : \Omega \to \mathbb{R}$, such that $\varphi_0|_{\mathrm{supp}(P)}$ solves (3), and $T_0 := \nabla \varphi_0$ is a vector field from Ω into itself.

With a slight abuse of terminology, we will refer to the map $\varphi_0 : \Omega \to \mathbb{R}$ in condition **A0** as a Brenier potential. By Brenier's theorem, this condition is satisfied whenever $P \in \mathcal{P}_{2,\mathrm{ac}}(\Omega)$. However, it is also satisfied when P is an arbitrary measure, provided that Q is the pushforward of P under a map $\nabla \varphi_0 : \Omega \to \Omega$, with φ_0 a strictly convex and differentiable function. The following result (which can be deduced from Proposition 1.15 of Santambrogio [2015]) characterizes the optimal transport problem between P and Q when assumption **A0** holds:

Theorem 2. Let $P, Q \in \mathcal{P}_2(\Omega)$, and suppose that **A**0 holds. Then, there exists a unique optimal transport coupling between P and Q, which is induced by the optimal transport maps $T_0 := \nabla \varphi_0$ between P and Q, and $S_0 := \nabla \varphi_0^*$ between Q and P.

Theorems 1 and 2 provide sufficient conditions for the existence of a unique OT map between P and Q, and connects this unique map to the solution of the semi-dual program (3). These results are important as they allow us to study Monge's problem via its linear programming relaxation. This in turn allows us to bring tools from convex analysis and convex duality to bear on the OT problem. The dual perspective will play a key role in the development of stability bounds.

2.2 Wasserstein Distance and the Semi-Dual Functional

The optimal value of the Kantorovich problem in (6) defines a metric on distributions known as the 2-Wasserstein distance:

$$W_2^2(P,Q) := \inf_{\pi \in \Pi(P,Q)} \int \|x - y\|_2^2 d\pi(x,y).$$

The Wasserstein distance has a dual characterization via the identity:

$$W_{2}^{2}(P,Q) = \int \|\cdot\|_{2}^{2} dP + \int \|\cdot\|_{2}^{2} dQ - 2 \inf_{\varphi \in L^{1}(P)} \mathcal{S}_{P,Q}(\varphi),$$

where the semi-dual functional $S_{P,Q}$ is defined in (4). The semi-dual optimization problem (3) can be seen as a dual optimization problem to Kantorovich's linear program [Villani, 2003].

The semi-dual functional is central in our stability bounds. Of particular interest to us will be a characterization of the semi-dual growth function as an integrated Bregman divergence between transport maps (see Appendix B) [Muzellec et al., 2021, Vacher and Vialard, 2022]:

Lemma 1. Suppose that $P, Q, \widetilde{P}, \widetilde{Q} \in \mathcal{P}_2(\Omega)$ and that A0 holds. Let $T_0 := \nabla \varphi_0$, let $\widetilde{\pi}$ denote an optimal coupling of \widetilde{P} and \widetilde{Q} and let $\widetilde{\varphi}$ denote a Brenier potential between \widetilde{P} and \widetilde{Q} . Then:

$$\mathcal{S}_{\widetilde{P},\widetilde{Q}}(\varphi_0) - \mathcal{S}_{\widetilde{P},\widetilde{Q}}(\widetilde{\varphi}) = \int \left[\varphi_0^*(y) - \varphi_0^*(T_0(x)) - \langle \nabla \varphi_0^*(T_0(x)), y - T_0(x) \rangle \right] d\widetilde{\pi}(x,y).$$
(8)

In cases when the coupling $\tilde{\pi}$ is realized by a transport map \tilde{T} the expression (8) simplifies to:

$$\mathcal{S}_{\widetilde{P},\widetilde{Q}}(\varphi_0) - \mathcal{S}_{\widetilde{P},\widetilde{Q}}(\widetilde{\varphi}) = \int \left[\varphi_0^*(\widetilde{T}(x)) - \varphi_0^*(T_0(x)) - \langle \nabla \varphi_0^*(T_0(x)), \, \widetilde{T}(x) - T_0(x) \rangle \right] d\widetilde{P}(x).$$

Intuitively, this result shows that the growth of the semi-dual functional away from its minimizer effectively measures a "distance" between two transport maps, which in turn gives further insight into the crucial role played by the semi-dual functional in the estimation of transport maps.

2.3 Canonical Estimators of the OT Map

Having setup the necessary background on optimal transport we now describe the two classes of estimators that we study:

1. **Plugin Estimators:** We consider two types of plugin estimators. Smooth estimators of the OT map are obtained by first constructing estimates \hat{P}_n and \hat{Q}_m of the measures P and Q, and then defining \hat{T}_{nm} to be the optimal transport map between \hat{P}_n and \hat{Q}_m . For \hat{T}_{nm} to be well-defined, the estimators \hat{P}_n and \hat{Q}_m need to be proper, in the sense that they define probability measures in their own right, and are related by an optimal transport map (which is, for instance, the case if \hat{P}_n is absolutely continuous).

Empirical estimators are similar in spirit, but we first compute an optimal transport coupling between the empirical measures:

$$P_n = \frac{1}{n} \sum_{i=1}^n \delta_{X_i}, \text{ and } Q_m = \frac{1}{m} \sum_{j=1}^m \delta_{Y_j}.$$
 (9)

This empirical coupling, which is only defined at the sample points X_1, \ldots, X_n , can then be extended to the entire domain using ideas from nonparametric regression [Manole et al., 2024], or using the so-called barycentric projection [Deb et al., 2021]. We refer to empirical and smooth estimators as *plugin* estimators.

2. Dual Estimators: An alternative class of estimators is based on the fact that the optimal transport map is the gradient of a convex function φ_0 which solves the following *semi-dual* optimization problem:

$$\varphi_0 := \operatorname*{argmin}_{\varphi \in L^1(P)} \int \varphi dP + \int \varphi^* dQ, \tag{10}$$

where φ^* denotes the convex conjugate of φ . Given samples from the measures P and Q a natural class of estimators for φ_0 , is:

$$\widehat{\varphi} = \underset{\varphi \in \Phi}{\operatorname{argmin}} \int \varphi dP_n + \int \varphi^* dQ_m, \tag{11}$$

where Φ is an appropriate function class. Under some regularity conditions, solving the empirical semi-dual optimization problem yields estimators for the transport map $\widehat{T}_{nm} = \nabla \widehat{\varphi}$, which we refer to as dual estimators. Dual estimators have been proposed and analyzed in many prior works [Ding et al., 2024, Divol et al., 2022, Hütter and Rigollet, 2021, Vacher and Vialard, 2022].

3 Stability Bounds

We now present our main stability bound and discuss its consequences. In the analysis of plugin and dual transport map estimators the key technical condition, which endows the semidual optimization problem (3) with a desirable growth property (see (5)), is that there exists a Brenier potential $\varphi_0 : \Omega \to \mathbb{R}$ from P to Q in the sense of condition A0, which is additionally smooth and strongly convex:

A1(α) The function φ_0 is convex, continuously differentiable over Ω , and satisfies

$$\varphi_0(y) \ge \varphi_0(x) + \langle \nabla \varphi_0(x), y - x \rangle + \frac{\alpha}{2} \|y - x\|_2^2, \text{ for all } x, y \in \Omega.$$

 $A2(\beta)$ The function φ_0 is convex, continuously differentiable over Ω , and satisfies

$$\varphi_0(y) \le \varphi_0(x) + \langle \nabla \varphi_0(x), y - x \rangle + \frac{\beta}{2} ||y - x||_2^2, \text{ for all } x, y \in \Omega.$$

A standard fact in convex analysis is that β -smoothness of φ_0 implies $1/\beta$ -strong convexity of φ_0^* and α -strong convexity of φ_0 implies $1/\alpha$ -smoothness of φ_0^* [Hiriart-Urruty and Lemaréchal, 1993]. When assumptions **A0**, **A1**(α) and **A2**(β) hold then, by Theorem 2, there exists a *P*-almost everywhere uniquely defined transport map T_0 from *P* to *Q* which is bi-Lipschitz over Ω , and whose inverse is the *Q*-almost everywhere uniquely defined transport map from *Q* to *P*.

Our main contribution in this section is the following.

Theorem 3. Let $P, Q \in \mathcal{P}_2(\Omega)$, and assume A0, $A1(\alpha)$ and $A2(\beta)$ hold for some $\alpha, \beta > 0$. For any $\widehat{P}, \widehat{Q} \in \mathcal{P}_2(\Omega)$, let $\widehat{\pi}$ denote an optimal coupling of \widehat{P} and \widehat{Q} . Then,

$$\frac{1}{\beta} \mathbb{E}_{(X,Y)\sim\hat{\pi}} \|Y - T_0(X)\|_2^2 \le \frac{1}{\alpha} W_2^2(\widehat{Q}, Q) + \beta W_2^2(\widehat{P}, P) + 2W_2(\widehat{P}, P)W_2(\widehat{Q}, Q).$$
(12)

At a high-level, this inequality provides an upper bound on a notion of discrepancy between an estimated optimal coupling $\hat{\pi}$ and the target OT map T_0 in terms of the Wasserstein distances between \hat{P} and P, and \hat{Q} and Q. It is worth noting that this result holds even when the measures under consideration do not share the same support. This flexibility will be salient in Section 5.

Let us now highlight several interesting consequences of this general stability bound.

One-Sample Stability Bounds: An instructive special case to consider is when $P \in \mathcal{P}_{ac}(\Omega)$ is a known distribution, and we only obtain samples $Y_1, \ldots, Y_m \sim Q$. In this case, by Theorem 1, there is an optimal map \widehat{T} between P and \widehat{Q} , and our stability bound reduces to the following corollary:

Corollary 1. Let $P \in \mathcal{P}_{ac}(\Omega)$, $Q \in \mathcal{P}_{2}(\Omega)$, and assume **A0**, **A1**(α) and **A2**(β) hold for some $\alpha, \beta > 0$. For any $\widehat{Q} \in \mathcal{P}_{2}(\Omega)$, let \widehat{T} denote the OT map from P to \widehat{Q} . Then,

$$\frac{1}{\beta} \mathbb{E}_{X \sim P} \| \widehat{T}(X) - T_0(X) \|_2^2 \le \frac{1}{\alpha} W_2^2(\widehat{Q}, Q).$$
(13)

This recovers a result of Manole et al. [2024] (see Theorem 6). We could also consider the case where the source measure is sampled, and the target measure is known:

Corollary 2. Let $P, Q \in \mathcal{P}_2(\Omega)$, and assume A0 and $A2(\beta)$ hold for some $\beta > 0$. For any $\widehat{P} \in \mathcal{P}_2(\Omega)$, let $\widehat{\pi}$ denote the optimal coupling between \widehat{P} and Q. Then,

$$\frac{1}{\beta} \mathbb{E}_{(X,Y)\sim\hat{\pi}} \|Y - T_0(X)\|_2^2 \le \beta W_2^2(\hat{P}, P).$$
(14)

To our knowledge, this result is new, and surprisingly requires only smoothness of the Brenier potential φ_0 and not strong convexity.

Two-Sample Stability Bound: In the general setting, the stability bound (12) substantially strengthens the corresponding result in Manole et al. [2024] (see Proposition 13). In particular, under assumptions $A1(\alpha)$ and $A2(\beta)$, their work was only able to establish an upper bound on the semi-dual growth of the form:

$$\mathcal{S}_{\widehat{P},\widehat{Q}}(\varphi_0) - \mathcal{S}_{\widehat{P},\widehat{Q}}(\varphi_{\widehat{P},\widehat{Q}}) \leq \left[\beta + \frac{1}{\alpha}\right] \left[W_2(\widehat{P},P) + W_2(\widehat{Q},Q)\right]^2,$$

where $\varphi_{\widehat{P},\widehat{Q}}$ denotes any Brenier potential in the OT problem from \widehat{P} to \widehat{Q} . The above display does not yield useful bounds on the risk of the plugin transport map estimate. To remedy this deficiency, Manole et al. [2024] instead needed to assume *uniform bounds* on the smoothness and strong-convexity of the *estimated* Brenier potential $\varphi_{\widehat{P},\widehat{Q}}$. This step weakened their results considerably, requiring them to assume that the distributions under consideration were supported on the flat *d*-dimensional torus and that the target of inference was the torus OT map (the OT map with respect to a modified Euclidean metric). In contrast, building upon our improved stability bound we obtain sharp results for the usual plugin estimators of the OT map in the smooth setting, in Section 4.

Empirical Stability Bound: In the case when \widehat{P} and \widehat{Q} are empirical measures, we obtain the following corollary:

Corollary 3. Let $P, Q \in \mathcal{P}_2(\Omega)$, and assume A0, $A1(\alpha)$ and $A2(\beta)$ hold for some $\alpha, \beta > 0$. Let $\hat{\pi}$ denote an optimal coupling of the empirical distributions P_n and Q_m . Then,

$$\frac{1}{\beta} \mathbb{E}_{(X,Y)\sim\widehat{\pi}} \|Y - T_0(X)\|_2^2 \le \frac{1}{\alpha} W_2^2(Q_m, Q) + \beta W_2^2(P_n, P) + 2W_2(P_n, P)W_2(Q_m, Q).$$

This result strengthens Proposition 12 of Manole et al. [2024]. To build some intuition, it is worthwhile to consider the situation where n = m, in which case the optimal coupling between P_n and Q_n is realized by an optimal transport map \hat{T}_n . Corollary 3 then yields a bound on the *empirical* error of the map \hat{T}_n :

$$\frac{1}{n\beta}\sum_{i=1}^{n} \|\widehat{T}_n(X_i) - T_0(X_i)\|_2^2 \le \frac{1}{\alpha}W_2^2(Q_m, Q) + \beta W_2^2(P_n, P) + 2W_2(P_n, P)W_2(Q_m, Q).$$

The convergence of the empirical measure in the Wasserstein distance has been extensively studied [Fournier and Guillin, 2015, Lei, 2020, Weed and Bach, 2019], and these results provide sharp bounds on the right-hand side of the above display. In Section 5 we illustrate how the map \hat{T}_n can be extended to the domain Ω , providing guarantees on the $L^2(P)$ risk of the estimator \hat{T}_n .

Semi-Dual Quadratic Growth Bounds: The stability bound (12) is obtained by upper and lower bounding the growth of a semi-dual functional. As we highlighted earlier, past work analyzing dual estimators growth bounds by upper and lower bounding growth of the population semi-dual functional, i.e. by studying the growth of $S_{P,Q}(\varphi) - S_{P,Q}(\varphi_0)$ (see (5)). In contrast, when analyzing plugin estimators, it is fruitful to study the growth of the *estimated* semi-dual functional: $S_{\hat{P},\hat{Q}}(\varphi_0) - S_{\hat{P},\hat{Q}}(\varphi_{\hat{P},\hat{Q}})$. An important aspect of studying this latter quantity is that, due to the Bregman representation in Lemma 1, we are able to provide useful upper and lower bounds on the estimated semi-dual functional without requiring any regularity assumptions on the estimated Brenier potential $\varphi_{\widehat{P},\widehat{O}}$.

Stability Bounds for the OT Coupling: Theorem 3 can also be formulated as a stability bound over the space of couplings. We first define the Wasserstein distance between the estimated coupling $\hat{\pi}$ and $\pi_0 := (\mathrm{Id}, T_0)_{\#} P$, namely:

$$W_2^2(\hat{\pi}, \pi_0) = \inf_{\gamma \in \Pi(\hat{\pi}, \pi_0)} \int \|u - v\|_2^2 d\gamma(u, v).$$
(15)

Then, as a direct consquence of Corollary 3.9 of Li and Nochetto [2021], and our stability bound in (12) we obtain the following result:

Corollary 4. Under the conditions of Theorem 3, there exists a constant C > 0 depending only on α and β such that

$$W_2^2(\hat{\pi}, \pi_0) \le C \left(W_2^2(\hat{P}, P) + W_2^2(\hat{Q}, Q) \right).$$
(16)

This Corollary elucidates a certain sharpness of our stability bounds. Since the cost function $||u-v||_2^2$ in (15) depends additively on its coordinates, it must be the case that the Wasserstein distance between couplings is bounded from below by the Wasserstein distance between the corresponding marginal distributions, i.e.:

$$W_2^2(\hat{\pi}, \pi_0) \ge \max \{ W_2^2(\hat{P}, P), W_2^2(\hat{Q}, Q) \}.$$

It follows that the bound in Corollary 4 is sharp up to constants.

Finally, it is worth noting that in contrast to past work we obtain each of our stability bounds as direct consequences of a single unified result. We now briefly discuss a few highlights of the proof of Theorem 3 before providing some historical context for our work.

Proof Highlights for Theorem 3: We give a full proof of this result in Appendix C, but discuss some interesting aspects of the proof here. Despite the fact that our goal is to prove a stability bound, we proceed by establishing a semi-dual growth bound. Concretely, by Lemma 1 we have that,

$$\mathcal{S}_{\widehat{P},\widehat{Q}}(\varphi_0) - \mathcal{S}_{\widehat{P},\widehat{Q}}(\varphi_{\widehat{P},\widehat{Q}}) = \int \left[\varphi_0^*(y) - \varphi_0^*(T_0(x)) - \langle \nabla \varphi_0^*(T_0(x)), y - T_0(x) \rangle \right] d\widehat{\pi}(x,y).$$

Our goal will be to upper and lower bound this quantity by the two sides of the inequality (12), thus establishing the theorem. Noting that φ_0^* is $1/\beta$ strongly convex by A2(β) we have:

$$\frac{1}{\beta} \mathbb{E}_{(X,Y)\sim\widehat{\pi}} \|Y - T_0(X)\|_2^2 \le \mathcal{S}_{\widehat{P},\widehat{Q}}(\varphi_0) - \mathcal{S}_{\widehat{P},\widehat{Q}}(\varphi_{\widehat{P},\widehat{Q}}),$$

which immediately yields the desired lower bound.

In contrast to the proof of the lower bound which only uses pointwise bounds (i.e. bounds which apply to the integrand pointwise), the proof of the upper bound is more involved using instead average-case optimality properties of the OT coupling (7). To illustrate the main ideas, let us consider the case when \hat{P} is taken to be the true distribution P, and focus on upper bounding the difference $S_{P,\hat{Q}}(\varphi_0) - S_{P,\hat{Q}}(\varphi_{P,\hat{Q}})$. Let $\tilde{\pi}$ denote an optimal coupling of P and \hat{Q} , and let $(X, Y) \sim \tilde{\pi}$. Then,

$$\mathcal{S}_{P,\widehat{Q}}(\varphi_0) - \mathcal{S}_{P,\widehat{Q}}(\varphi_{P,\widehat{Q}}) = \mathbb{E}\left[\varphi_0^*(Y) - \varphi_0^*(T_0(X)) - \langle X, Y - T_0(X) \rangle\right],$$

using the fact that $\nabla \varphi_0^*(T_0(X)) = X$ almost surely. Now, given any random variable Z with marginal distribution \hat{Q} , we can equivalently write,

$$\mathcal{S}_{P,\widehat{Q}}(\varphi_0) - \mathcal{S}_{P,\widehat{Q}}(\varphi_{P,\widehat{Q}}) = \mathbb{E}\left[\varphi_0^*(Z) - \varphi_0^*(T_0(X)) - \langle X, Z - T_0(X) \rangle\right] - \mathbb{E}\langle X, Y - Z \rangle.$$

Recalling from (7) that

$$\mathbb{E}_{(X,Y)\sim\widetilde{\pi}}\langle X, Y\rangle \geq \mathbb{E}\langle X, Z\rangle,$$

we obtain,

$$\mathcal{S}_{P,\widehat{Q}}(\varphi_0) - \mathcal{S}_{P,\widehat{Q}}(\varphi_{P,\widehat{Q}}) \le \mathbb{E}\left[\varphi_0^*(Z) - \varphi_0^*(T_0(X)) - \langle X, Z - T_0(X) \rangle\right].$$

Using the $(1/\alpha)$ -smoothness of φ_0^* we obtain that,

$$\mathcal{S}_{P,\widehat{Q}}(\varphi_0) - \mathcal{S}_{P,\widehat{Q}}(\varphi_{P,\widehat{Q}}) \leq \frac{1}{\alpha} \mathbb{E} \|Z - T_0(X)\|_2^2.$$

The above display holds for any random variable $Z \sim \hat{Q}$, thus we are free to choose its joint distribution with X. Let $\pi_{Q,\hat{Q}}$ denote an optimal coupling of Q and \hat{Q} , and pick Z such that $(T_0(X), Z) \sim \pi_{Q,\hat{Q}}$. Then,

$$\mathbb{E}||Z - T_0(X)||_2^2 = \mathbb{E}_{(U,V) \sim \pi_{Q,\widehat{Q}}} ||V - U||_2^2 = W_2^2(Q,\widehat{Q}),$$

which yields the bound:

$$\mathcal{S}_{P,\widehat{Q}}(\varphi_0) - \mathcal{S}_{P,\widehat{Q}}(\varphi_{P,\widehat{Q}}) \leq \frac{1}{\alpha} W_2^2(Q,\widehat{Q}).$$

The proof of the general upper bound, when \widehat{P} is potentially different from P, requires a sharpening of the above argument (see Lemma 4) and we defer the details to Appendix C.

3.1 Related Work

With Theorem 3 in place we can now provide the appropriate context for our result. Stability bounds of the form we describe have a long history. For instance, a qualitative implication of the stability bound we derive in Corollary 1 is that the convergence of \hat{Q} to Q in the W_2 metric implies the convergence of the OT map \hat{T} to T_0 in the $L^2(P)$ metric. Qualitative results of this type are well-studied (for instance, see Villani [2003, Exercise 2.17], Panaretos and Zemel [2020, Proposition 1.7.11], and Segers [2022]), and hold under significantly weaker conditions than the ones we impose. On the other hand, qualitative results do not typically yield sharp statistical rates on the risk in (2).

To our knowledge, the earliest quantitative stability result was derived by Gigli [2011] (who attributed it to Ambrosio) under the strong convexity assumption $\mathbf{A1}(\alpha)$ (see also Theorem 3.2 of Ambrosio et al. [2019] and Theorem 3.5 of Li and Nochetto [2021]). Gigli showed that if $P \in \mathcal{P}_{2,\mathrm{ac}}(\mathbb{R}^d)$, \hat{T} is a transport map from P to \hat{Q} , and $T_0 = \nabla \varphi_0$ is the OT map from P to Q, with the potential φ_0 satisfying condition $\mathbf{A1}(\alpha)$, then:

$$||T_0 - \widehat{T}||^2_{L^2(P)} \lesssim \mathbb{E}_{X \sim P} ||\widehat{T}(X) - X||^2_2 - \mathbb{E}_{X \sim P} ||T_0(X) - X||^2_2.$$

This result suggests that the excess transport cost of a sub-optimal transport map grows in proportion to the $L^2(P)$ distance of the map from the OT map. This result is not directly useful for the statistical analysis of transport map estimates since it requires \hat{T} to be a valid transport map between P and \hat{Q} , a condition that is not always satisfied by estimates of the map. This deficiency inspired Hütter and Rigollet [2021] (see their Proposition 10) to develop the quadratic growth bound on the semi-dual functional $S_{P,Q}$, showing:

$$\|\nabla\varphi - T_0\|_{L^2(P)}^2 \lesssim \mathcal{S}_{P,Q}(\varphi) - \mathcal{S}_{P,Q}(\varphi_0) \lesssim \|\nabla\varphi - T_0\|_{L^2(P)}^2,$$

under the assumptions that φ , φ_0 are both smooth and strongly convex. Muzellec et al. [2021], Vacher and Vialard [2022] and Makkuva et al. [2020] observed that the proof of Hütter and Rigollet [2021] also applies when φ (alone) is smooth and strongly convex.

While the work of Hütter and Rigollet [2021] provides a semi-dual growth bound useful for the analysis of dual estimators, the works of Deb et al. [2021], Ghosal and Sen [2022] and Manole et al. [2024] provide stability bounds useful for the analysis of plugin estimators. We have already discussed the results of Manole et al. [2024], and highlighted how Theorem 3 provides useful improvements to their results. The stability bounds of Deb et al. [2021], Ghosal and Sen [2022] are quantitatively weaker than those in Manole et al. [2024], and replace the upper bound in (12), by an empirical process term involving the estimated measures \hat{P}, \hat{Q} and the corresponding Brenier potential $\varphi_{\hat{P},\hat{Q}}$. Analyzing this term to obtain risk bounds then necessitates strong assumptions to reason about the regularity of the Brenier potential $\varphi_{\hat{P},\hat{Q}}$ (akin to the torus assumption made by Manole et al. [2024]). A naïve analysis of the empirical process term has a further deficiency which can result in sub-optimal rates of convergence for transport map estimates. In rough terms, the sharper upper bounds of Theorem 3 behave as the square of an empirical process, yielding much faster rates of convergence.

Thus far, we have discussed quantitative stability bounds when one of conditions $A1(\alpha)$ and $A2(\beta)$ hold. In contrast, the works of Berman [2021], Delalande and Merigot [2023], Gallouët et al. [2022], Mérigot et al. [2020], and Letrouit and Mérigot [2024] provide various stability bounds under variations of the target measure, which hold without smoothness assumptions on the optimal transport maps involved. That is, given a sufficiently regular set $\Omega \subseteq \mathbb{R}^d$ and measure $P \in \mathcal{P}_2(\Omega)$, they derive bounds of the form:

$$\|\widehat{T} - T_0\|_{L^2(P)} \lesssim W_2^{\alpha}(\widehat{Q}, Q), \tag{17}$$

for any measures $\hat{Q}, Q \in \mathcal{P}(\Omega)$ satisfying appropriate tail conditions, and with corresponding Hölder exponents $\alpha \in [0, 1]$ which can be taken as high as $\alpha = 1/6$. Conversely, in this level of generality, it has been known since the work of Gigli [2011] that the exponent α cannot be made greater than 1/2, and it remains an open question to determine whether this threshold can be achieved. Our stability bounds show that the much more favorable Hölder exponent $\alpha = 1$ is achievable when one of the optimal transport maps is smooth.

Beyond the statistical applications that we have in mind, bounds of type (17) are highly sought after, as the quantity $\|\hat{T} - T_0\|_{L^2(P)}$ is itself a metric between \hat{Q} and Q, which can be viewed as a proxy for the Wasserstein distance. This quantity is sometimes known as the linearized Wasserstein distance [Wang et al., 2013], and is widely-used in applications due to its Hilbertian structure, and its favorable computational properties (e.g. Cai et al. [2020]). Corollary 1 provides compelling motivation for the use of linearized Wasserstein distance, by showing that it is in fact equivalent to the original Wasserstein distance under appropriate smoothness assumptions.

4 Estimating OT Maps Between Smooth Distributions

As a first application, we will now show how Theorem 3 can be used to characterize the risk of plugin estimators of optimal transport maps between smooth densities. Let P and Q denote two absolutely continuous distributions over Ω with respective densities p and q. To define our class of plugin estimators, let \hat{P}_n denote an estimator of the distribution P based on the i.i.d. sample $X_1, \ldots, X_n \sim P$. We assume that \hat{P}_n is a *proper* estimator, in the sense that it almost surely defines a probability measure in its own right. We further assume that \hat{P}_n is absolutely continuous with respect to the Lebesgue measure on Ω , almost surely. Finally, we denote by \hat{Q}_m any proper estimator of Q based on the i.i.d. sample $Y_1, \ldots, Y_m \sim Q$. Due to the absolute continuity of \hat{P}_n . Theorem 1 implies that there exists a unique optimal transport coupling between \hat{P}_n and \hat{Q}_m which is induced by an optimal transport map \hat{T}_{nm} . This uniquely defined map \hat{T}_{nm} is a natural plugin estimator of T_0 , and Theorem 3 implies the following bound on its $L^2(\hat{P}_n)$ risk:

$$\mathbb{E}\|\widehat{T}_{nm} - T_0\|_{L^2(\widehat{P}_n)}^2 \lesssim \mathbb{E}\Big[W_2^2(\widehat{P}_n, P) + W_2^2(\widehat{Q}_m, Q)\Big],$$

provided that T_0 satisfies assumptions $\mathbf{A1}(\alpha) - \mathbf{A2}(\beta)$. In order to characterize the $L^2(P)$ risk of \widehat{T}_{nm} from here, one requires two ingredients:

- 1. Risk of Distribution Estimation in Wasserstein Distance. First, one needs to provide upper bounds on the risk of \hat{P}_n and \hat{Q}_n as estimates of P and Q in the Wasserstein distance. Bounds of this type are well-studied in the literature for a variety of estimators, including smooth density estimators [Divol, 2022, Niles-Weed and Berthet, 2022], and the empirical measure (e.g. [Bobkov and Ledoux, 2019, Boissard and Le Gouic, 2014, Fournier and Guillin, 2015, Lei, 2020, Weed and Bach, 2019], and references therein).
- 2. Population and Empirical L^2 Norms. Second, one needs to relate the $L^2(\hat{P}_n)$ risk to the $L^2(P)$ risk. The simplest approach is to work with an estimator \hat{P}_n for which the density ratio $dP/d\hat{P}_n$ is almost surely bounded over Ω , in which case the $L^2(P)$ risk is trivially bounded above by the $L^2(\hat{P}_n)$ risk:

$$\mathbb{E}\|\widehat{T}_{nm} - T_0\|_{L^2(P)}^2 \lesssim \mathbb{E}\|\widehat{T}_{nm} - T_0\|_{L^2(\widehat{P}_n)}^2.$$

When such a condition is not met, one might instead try to bound the deviations

$$\left| \|\widehat{T} - T_0\|_{L^2(\widehat{P}_n)}^2 - \|\widehat{T} - T_0\|_{L^2(P)}^2 \right| = \left| \int \|\widehat{T} - T_0\|_2^2 d(\widehat{P}_n - P) \right|,$$

uniformly over the set of allowable transport maps \widehat{T} . If \widehat{P}_n were replaced by the empirical measure of X_1, \ldots, X_n , the above would simply be an empirical process indexed by the functions $\|\widehat{T} - T_0\|_2^2$. Similar quantities arise in the classical theory of nonparametric least squares regression, where localization and empirical process arguments are typically used to relate the $L^2(P_n)$ risk of least squares estimators to their $L^2(P)$ risk [van de Geer, 2000]. Such arguments cannot directly be translated to our setting since \widehat{P}_n cannot be taken to be the empirical measure, however for some choices of this estimator, such as linear smoothers, the above empirical process can in principle still be controlled using known bounds on suprema of smoothed empirical processes [Giné and Nickl, 2009, Radulović and Wegkamp, 2003]. In Section 5, we will illustrate a third method for relating the $L^2(P)$ and $L^2(\widehat{P}_n)$ norms, tailored to the nearest neighbor-based transport map estimator, which is based on characterizing the Voronoi cells generated by X_1, \ldots, X_n . Let us now provide an example of an estimator for which the above steps can be carried out. We adopt the same setting as Manole et al. [2024, Section 4.4], who derive upper bounds on a smooth plugin estimator *conditionally* on a boundary regularity condition (cf. condition (C2) therein). In contrast, we will derive an *unconditional* version of their result in this section, which is enabled by the fact that our new stability bound in Theorem 3 does not place any assumptions on the fitted coupling.

We adopt the same notation and conditions as Manole et al. [2024]. Concretely, assume that Ω is a *known* compact, convex subset of \mathbb{R}^d whose boundary is \mathcal{C}^{∞} . Without loss of generality, let Ω have unit volume. Assume that the true distributions P and Q have densities which lie in the following ball of Hölder continuous functions:

$$\mathcal{C}^{s}(\Omega; M, \gamma) = \Big\{ f \in \mathcal{C}^{s}(\Omega) : \|f\|_{\mathcal{C}^{s}(\Omega)} \le M, f \ge \gamma \text{ over } \Omega, D^{k}f = 0 \text{ on } \partial\Omega, k = 1, \dots, \lfloor s \rfloor \Big\},\$$

where $M, \gamma, s > 0$, and where differentiation along the boundary is to be understood in the weak sense (cf. Appendix J.1 of Manole et al. [2024]). Beyond the smoothness constraint, the class $C^s(\Omega; M, \gamma)$ requires the densities to be bounded away from zero over the domain, and to have vanishing derivatives at the boundary¹. It is well-known that the Wasserstein density estimation risk depends strongly on whether or not the densities are lower-bounded; we focus only on the lower-bounded case, for which the sharp minimax rate has been precisely characterized [Bobkov and Ledoux, 2019, Divol, 2022, Niles-Weed and Berthet, 2022]. On the other hand, the boundary condition allows us to construct density estimators which are exactly supported on Ω , which in turn is convenient for relating the Wasserstein density estimation risk to the simpler risk of estimating a density under the norm of a particular Hilbert space [Peyre, 2018]. These are certainly not the most general conditions under which Wasserstein density estimation can be studied, and improved results for Wasserstein density estimation will have direct implications for the induced optimal transport map estimators, due to the modular nature of Theorem 3.

Finally, in addition to the above conditions on the densities, we will assume that the T_0 satisfies the curvature conditions described in Section 3. That is, we will assume that the true densities lies in the class

$$\mathcal{F}^{s}(\Omega; M, \gamma, \alpha, \beta) = \{ (p,q) \in \mathcal{D}^{2} : p, q \in \mathcal{C}^{s}(\Omega; M, \gamma), \text{ there exists a Brenier potential } \varphi_{0} \\ \text{from } P \text{ to } Q \text{ satisfying } \mathbf{A1}(\alpha) - \mathbf{A2}(\beta) \},$$

where \mathcal{D} is the set of Lebesgue densities on \mathbb{R}^d . By a well-known result of Caffarelli [1996], the condition $p,q \in \mathcal{C}^s(\Omega; M, \gamma)$ implies the existence of a Brenier potential from P to Qwhich satisfies conditions $\mathbf{A1}(\widetilde{\alpha})-\mathbf{A2}(\widetilde{\beta})$ for some $\widetilde{\alpha}, \widetilde{\beta} > 0$. To the best of our knowledge, however, there is no known quantitative relation between the resulting parameters $\widetilde{\alpha}, \widetilde{\beta}$ and the original problem parameters M, d, γ, s , thus we prefer to formulate the above assumption on φ_0 directly. Similar considerations are discussed by Hütter and Rigollet [2021, Appendix E] and Manole et al. [2024, Section 4.3].

The main result of this section is:

Theorem 4. For any s > 0, there exist proper and absolutely continuous estimators \widehat{P}_n and \widehat{Q}_n such that for any $M, \gamma, \alpha, \beta > 0$, there exists a constant $C = C(M, \Omega, \gamma, s, \alpha, \beta) > 0$ for

¹The condition of vanishing derivatives can be relaxed to a Neumann boundary condition of sufficiently high order, as in the definition of the function class $C_N^s(\Omega)$ of Manole et al. [2024, Section 4.4].

which the unique optimal transport map \widehat{T}_n pushing \widehat{P}_n forward onto \widehat{Q}_m satisfies

$$\sup_{(p,q)\in\mathcal{F}^s(\Omega;M,\gamma,\alpha,\beta)} R(\widehat{T}_n,T_0) \le C\epsilon_{n\wedge m}, \quad where \ \epsilon_n := \begin{cases} 1/n, \quad d=1, \\ \log n/n, \quad d=2, \\ n^{-\frac{2(s+1)}{2s+d}}, \quad d\ge 3. \end{cases}$$
(18)

Proof. Assume without loss of generality that n = m. Let $(p,q) \in \mathcal{F}^s(\Omega; M, \gamma, \alpha, \beta)$. We consider the estimators \hat{P}_n and \hat{Q}_n defined in Section 4.4 of Manole et al. [2024]. By Lemma 64 therein, there exists a constant $C_1 = C_1(M, \Omega, \gamma, s)$ and an event A_n of probability content at least $1 - C_1/n^2$ such that the density \hat{p}_n of \hat{P}_n satisfies $\hat{p}_n \geq 1/C_1$ over Ω . Furthermore, since we have assumed that $p \leq M$ over Ω , we deduce that the density ratio $dP/d\hat{P}_n$ is bounded over Ω by a constant depending only on M, γ, d, s , with probability at least $1 - C_1/n^2$. Thus,

$$\begin{split} \mathbb{E} \| \widehat{T}_n - T_0 \|_{L^2(P)}^2 &\leq \mathbb{E} \Big[\| \widehat{T}_n - T_0 \|_{L^2(P)}^2 \, \Big| \, A_n \Big] + \mathbb{E} \Big[\| \widehat{T}_n - T_0 \|_{L^2(P)}^2 \, \Big| \, A_n^{\mathsf{c}} \Big] \mathbb{P}(A_n^{\mathsf{c}}) \\ &\lesssim \mathbb{E} \Big[\| \widehat{T}_n - T_0 \|_{L^2(\widehat{P}_n)}^2 \Big] + n^{-2}, \end{split}$$

where we obtained the final term by noting that $\|\hat{T}_n - T_0\|_{L^2(P)}^2$ is trivially bounded by a constant, due to the compactness of Ω . We are now in a position to apply Theorem 3, which leads to:

$$\mathbb{E}\|\widehat{T}_n - T_0\|_{L^2(P)}^2 \lesssim W_2^2(\widehat{P}_n, P) + W_2^2(\widehat{Q}_n, Q) + n^{-2}.$$

By Lemma 65 of Manole et al. [2024], the convergence rate of the density estimators \hat{P}_n and \hat{Q}_n is bounded above by ϵ_n under our conditions. The claim thus follows.

Theorem 4 is analogous to Theorem 18 of Manole et al. [2024], with the important difference that it does not rely on their assumption (C2), which they were unable to verify. This assumption was important for their work since it implied that their estimator \widehat{T}_{nm} has an induced potential $\widehat{\varphi}$ which satisfies the regularity conditions $A1(\alpha)-A2(\beta)$ with high probability. We are able to circumvent this route since our new stability bound in Theorem 3 places no smoothness assumptions on the fitted potential.

The convergence rate appearing in Theorem 4 matches known minimax lower bounds for estimating OT maps between Hölder-continuous densities [Hütter and Rigollet, 2021], up to a logarithmic factor when d = 2. To our knowledge, this is the first (unconditional) proof of minimax optimality of a two-sample plugin estimator for OT maps under Hölder regularity assumptions.

5 An Improved Analysis of the Nearest-Neighbor Estimator

As a further illustration of Theorem 3, we consider the case where \widehat{T}_{nm} is taken to be the empirical optimal transport map, extended to \mathbb{R}^d via one-nearest-neighbor extrapolation. This estimator has been studied in past work by Manole et al. [2024] (see also Pooladian et al. [2023]), however their analysis required P to be compactly-supported, and to admit a density bounded away from zero over its support. In what follows, we significantly weaken these assumptions, showing that the nearest-neighbor estimator is minimax optimal in estimating bi-Lipschitz optimal transport maps, subject only to mild moment constraints. Concretely, given i.i.d. samples $X_1, \ldots, X_n \sim P$ and $Y_1, \ldots, Y_m \sim Q$, let $\hat{\pi}_{nm}$ be (the probability mass function of) an optimal transport coupling between the empirical measures P_n and Q_m . That is,

$$\widehat{\pi}_{nm} \in \operatorname*{argmin}_{\pi} \sum_{i=1}^{n} \sum_{j=1}^{m} \pi_{ij} \|X_i - Y_j\|^2,$$

where the minimizer is over all matrices $\pi \in \mathbb{R}^{n \times m}$ with nonnegative entries, whose rows add up to 1/m, and whose columns add up to 1/n. Define the Voronoi partition induced by X_1, \ldots, X_n as

$$V_j = \{x \in \mathbb{R}^d : ||x - X_j|| \le ||x - X_i||, \forall i \ne j\}, \quad j = 1, \dots, n.$$

Then, we define the one-nearest neighbor estimator by

$$\widehat{T}_{nm}^{1\mathrm{NN}}(x) = \sum_{i=1}^{n} \sum_{j=1}^{m} (n\widehat{\pi}_{ij})I(x \in V_i)Y_j, \quad x \in \Omega.$$

Corollary 3 then immediately yields an empirical error bound for \widehat{T}_{nm}^{1NN} :

$$\frac{1}{n} \sum_{i=1}^{n} \|\widehat{T}_{nm}^{1NN}(X_i) - T_0(X_i)\|_2^2 = \frac{1}{n} \sum_{i=1}^{n} \left\| \sum_{j=1}^{m} n\widehat{\pi}_{ij}Y_j - T_0(X_i) \right\|_2^2$$
$$\leq \sum_{i=1}^{n} \sum_{j=1}^{m} \widehat{\pi}_{ij} \|Y_j - T_0(X_i)\|_2^2$$
$$\lesssim W_2^2(Q_m, Q) + W_2^2(P_n, P)$$
$$=: e_1.$$

In order to upper bound the $L^2(P)$ risk of T_{nm}^{1NN} we need to relate its in-sample error to the $L^2(P)$ error. We do so via a different recipe from that followed in Section 4. Concretely, suppose for any X we define nn(X) to be the nearest neighbor of X in the sample X_1, \ldots, X_n . Then we define the expected distance from X to its nearest neighbor as:

$$e_2 := \mathbb{E}_{X \sim P} \left[\|X - \operatorname{nn}(X)\|_2^2 | X_1, \dots, X_n \right].$$

We also define the maximum mass of any Voronoi cell by:

$$e_3 := \max_{i \in \{1, \dots, n\}} P(V_i).$$

With these quantities in place we show the following general result on the accuracy of the nearest neighbor map:

Lemma 2. Let $P, Q \in \mathcal{P}_2(\mathbb{R}^d)$, and suppose that assumptions A0, $A1(\alpha)$ and $A2(\beta)$ hold for some $\alpha, \beta > 0$. Then, there exists a constant C > 0 depending only on β such that

$$\|\widehat{T}_{nm}^{1NN} - T_0\|_{L^2(P)}^2 \le C(e_1 \times ne_3 + e_2).$$

We prove this result in Appendix D. To get some sense of the strength of this result we highlight that each of the terms in the above bound can be controlled under relatively mild moment assumptions on the distributions P and Q. Concretely, we suppose that for some r > 0, to be specified in the sequel, we have that:

$$\int \|x\|_2^r dP, \int \|x\|_2^r dQ \le M_r^r < \infty.$$
⁽¹⁹⁾

Lemma 3. Let d > 4, and let the measures $P, Q \in \mathcal{P}(\mathbb{R}^d)$ satisfy (19) for r > 4, and suppose that P is non-atomic. Then, it holds that, for implicit constants depending only on M_r, d, r :

$$\mathbb{E}[e_1^2] \lesssim \left[m^{-4/d} + n^{-4/d}\right]$$
$$\mathbb{E}[e_2] \lesssim n^{-2/d} \tag{20}$$

$$\mathbb{E}[e_3^2] \lesssim \left[\frac{\log n}{n}\right]^2.$$
(21)

The first claim above follows directly from past work (e.g. Fournier and Guillin [2015]). We prove the remaining claims in Appendix E. The restriction that d > 4 is not essential to our result, but different rates of convergence are obtained for e_1 and e_2 when $d \leq 4$. We do not aim to optimize the moment cutoff r > 4, but instead aim for concise and illustrative results and proofs. Under these mild conditions on the measures P and Q we obtain quantitative rates of convergence for a practical estimator of the transport map:

Corollary 5. Let d > 4, and let the measures $P, Q \in \mathcal{P}(\mathbb{R}^d)$ satisfy condition (19). Suppose P is non-atomic. Assume further that there exists a Brenier potential φ_0 from P to Q satisfying conditions A0, $A1(\alpha)$, and $A2(\beta)$. Then, there exists a constant C > 0 depending on M_r, α, β, d, r such that

$$R(\hat{T}_{nm}^{1NN}, T_0) \le C(n^{-2/d} + m^{-2/d})\log n.$$

Corollary 5 follows directly by combining Lemmas 2 and 3, and an application of the Cauchy-Schwarz inequality. This result shows that, up to a logarithmic factor, the nearest neighbor estimator achieves the convergence rate $(n \wedge m)^{-2/d}$ for estimating a bi-Lipschitz optimal transport map T_0 , and is therefore minimax optimal [Hütter and Rigollet, 2021]. Our result merely assumes that the measures admit $4+\epsilon$ moments for some $\epsilon > 0$; this stands in contrast to past analyses of OT map estimation, which either assume that P is compactly-supported (e.g. Gunsilius [2022], Manole et al. [2024]), has exponential tails [Deb et al., 2021, Divol et al., 2022], or yield suboptimal convergence rates when P has finitely many moments [Ding et al., 2024].

Transport between Log-Smooth and Strongly Log-Concave Distributions: As an illustration of Corollary 5, suppose we consider the case when P and Q, are log-smooth and strongly log-concave. Concretely, P and Q are supported on \mathbb{R}^d and admit Lebesgue densities of the form $P = \exp(-V)$ and $Q = \exp(-W)$, where V and W are twice-differentiable, and satisfy for all $x \in \mathbb{R}^d$:

$$\alpha_V I \preceq \nabla^2 V(x) \preceq \beta_V I$$
, and, $\alpha_W I \preceq \nabla^2 W(x) \preceq \beta_W I$. (22)

Under these assumptions, Caffarelli's contraction theorem [Caffarelli, 2000] implies that our regularity assumptions A0, A1(α) and A2(β) hold for $\Omega = \mathbb{R}^d$, with $\alpha = \sqrt{\alpha_V/\beta_W}$ and $\beta = \sqrt{\beta_V/\alpha_W}$. Furthermore, the tails of a log-concave distribution are sub-exponential [Ledoux, 2005], and these distributions satisfy the moment conditions of Lemma 3. We obtain as a direct consequence of Corollary 5 the following result for the nearest-neighbor estimator:

Corollary 6. Let $P = \exp(-V)$ and $Q = \exp(-W)$ be log-concave measures with potentials satisfying (22). Then, there exists a constant C > 0 depending only on $\alpha_V, \beta_V, \alpha_W, \beta_W$ such that

$$R(\widehat{T}_{nm}^{1NN}, T_0) \le C(n^{-2/d} + m^{-2/d})\log n.$$

This result improves (by some logarithmic factors) the result of Divol et al. [2022]. More importantly, and in contrast to past work, our estimator of the transport map between two log-smooth and log-strongly concave distributions is simple and practical; in particular, it is free of tuning parameters.

6 Discussion

Our results close an important gap in the literature on estimating smooth optimal transport maps via the plugin principle. The main contribution of our work was to derive a new twosample stability bound in Theorem 3, which does not rely on any regularity properties of the fitted optimal transport map. Stability bounds of this form are useful as they reduce the study of plugin transport map estimators to that of distribution estimation under the Wasserstein loss. The latter is a well-studied problem with sharp results available for smoothness classes and under mild moment conditions. There are, however other arguments that one might be able to exploit to more directly upper bound the semi-dual difference, and we expect this to be a fruitful direction for future investigation.

We have centered our discussion around two broad classes of OT map estimators: plugin estimators, and dual estimators. To our knowledge, these are the only methods which are known to be minimax optimal over typical classes of smooth optimal transport maps. Quantitative convergence rates have also been derived for several other methods, based on entropic OT with vanishing regularization [Divol et al., 2024, Eckstein and Nutz, 2024, Mordant, 2024, Pooladian and Niles-Weed, 2021], and sum-of-squares relaxations [Muzellec et al., 2021, Vacher et al., 2021]. Although these methods are not known to be minimax optimal in the same level of generality as plugin or dual estimators, they typically enjoy more favorable computational properties.

Our results relied on smoothness and strong convexity assumptions on the underlying Brenier potentials. These assumptions can be relaxed in various ways to study the estimation of nonsmooth optimal transport maps. This question has received some attention in the case when one of the measures P and Q is discrete [Pooladian et al., 2023, Sadhu et al., 2024a,b].

We have limited our attention to the $L^2(P)$ risk of OT map estimators, due to its close connection with the optimality property of the population OT map. It is nevertheless of interest to understand the risk of OT map estimators under other loss functions; for example, L^{∞} convergence results are useful in the study of Monge-Kantorovich ranks and quantiles [Ghosal and Sen, 2022, Hallin et al., 2021]. The recent works of Manole et al. [2023] and González-Sanz and Sheng [2024] have established quantitative stability bounds for the uniform risk of plugin estimators, however they require strong smoothness and boundary conditions on the underlying distributions. It would be interesting to explore if the ideas from our work, which relaxes these types of regularity assumptions in $L^2(P)$ estimation, have any implications in more challenging settings.

Acknowledgements

The authors are grateful to Jonathan Niles-Weed and Larry Wasserman for numerous helpful conversations about stability bounds, and to Shayan Hundrieser for discussions related to this work. The work of SB was supported by the NSF grant DMS-2310632. This work was done in part while SB was visiting the Simons Institute for the Theory of Computing. The work of TM is funded by a Norbert Wiener postdoctoral fellowship.

References

- L. Ambrosio, F. Glaudo, and D. Trevisan. On the optimal map in the 2-dimensional random matching problem. *Discrete and Continuous Dynamical Systems*, 39(12):7291–7308, 2019.
- S. Balakrishnan, S. Narayanan, A. Rinaldo, A. Singh, and L. Wasserman. Cluster trees on manifolds. In *Advances in Neural Information Processing Systems*, 2013.
- R. J. Berman. Convergence rates for discretized Monge–Ampère equations and quantitative stability of optimal transport. *Foundations of Computational Mathematics*, 21(4):1099–1140, 2021.
- S. Bobkov and M. Ledoux. One-dimensional empirical measures, order statistics, and Kantorovich transport distances. *Memoirs of the American Mathematical Society*, 261, 2019.
- E. Boissard and T. Le Gouic. On the mean speed of convergence of empirical and occupation measures in Wasserstein distance. Annales de l'Institut Henri Poincaré–Probabilités et Statistiques, 50:539–563, 2014.
- L. A. Caffarelli. Boundary Regularity of Maps with Convex Potentials–II. Annals of Mathematics, 144:453–496, 1996.
- L. A. Caffarelli. Monotonicity Properties of Optimal Transportation and the FKG and Related Inequalities. *Communications in Mathematical Physics*, 214:547–563, 2000.
- T. Cai, J. Cheng, N. Craig, and K. Craig. Linearized optimal transport for collider events. *Physical Review D*, 102(11):116019, 2020.
- K. Chaudhuri and S. Dasgupta. Rates of convergence for the cluster tree. In Advances in Neural Information Processing Systems 24, 2010.
- V. Chernozhukov, A. Galichon, M. Hallin, and M. Henry. Monge–Kantorovich depth, quantiles, ranks and signs. *The Annals of Statistics*, 45:223–256, 2017.
- S. Chewi, J. Niles-Weed, and P. Rigollet. Statistical optimal transport. Ecole d'Eté de Probabilités de Saint-Flour XLIX. arXiv:2407.18163, 2024.
- N. Deb and B. Sen. Multivariate Rank-Based Distribution-Free Nonparametric Testing Using Measure Transportation. Journal of the American Statistical Association, 118:192–207, 2023.
- N. Deb, P. Ghosal, and B. Sen. Rates of Estimation of Optimal Transport Maps using Plug-in Estimators via Barycentric Projections. Advances in Neural Information Processing Systems 34, 2021.
- A. Delalande and Q. Merigot. Quantitative stability of optimal transport maps under variations of the target measure. Duke Mathematical Journal, 172(17):3321–3357, 2023.
- Y. Ding, R. Li, and L. Xue. Statistical convergence rates of optimal transport map estimation between general distributions. arXiv preprint arXiv:2412.08064, 2024.
- V. Divol. Measure estimation on manifolds: An optimal transport approach. *Probability* Theory and Related Fields, 183(1):581–647, 2022.

- V. Divol, J. Niles-Weed, and A.-A. Pooladian. Optimal transport map estimation in general function spaces. *The Annals of Statistics (To appear). arXiv preprint arXiv:2212.03722*, 2022.
- V. Divol, J. Niles-Weed, and A.-A. Pooladian. Tight stability bounds for entropic brenier maps. arXiv preprint arXiv:2404.02855, 2024.
- S. Eckstein and M. Nutz. Convergence rates for regularized optimal transport via quantization. Mathematics of Operations Research, 49(2):1223–1240, 2024.
- N. Fournier and A. Guillin. On the rate of convergence in Wasserstein distance of the empirical measure. Probability Theory and Related Fields, 162:707–738, 2015.
- A. Gallouët, Q. Mérigot, and B. Thibert. Strong c-concavity and stability in optimal transport. arXiv preprint arXiv:2207.11042, 2022.
- P. Ghosal and B. Sen. Multivariate Ranks and Quantiles using Optimal Transport: Consistency, Rates, and Nonparametric Testing. *The Annals of Statistics*, 50:1012–4037, 2022.
- N. Gigli. On Hölder continuity-in-time of the optimal transport map towards measures along a curve. *Proceedings of the Edinburgh Mathematical Society*, 54:401–409, 2011.
- D. Gilbarg and N. S. Trudinger. *Elliptic Partial Differential Equations of Second Order*. Classics in Mathematics. Springer-Verlag, Berlin Heidelberg, 2 edition, 2001.
- E. Giné and R. Nickl. Uniform limit theorems for wavelet density estimators. The Annals of Probability, 37(4):1605–1646, 2009.
- A. González-Sanz and S. Sheng. Linearization of Monge-Ampère equations and data science applications. arXiv preprint arXiv:2408.06534, 2024.
- F. F. Gunsilius. On the convergence rate of potentials of Brenier maps. *Econometric Theory*, 38(2):381–417, 2022.
- L. Györfi, M. Kohler, A. Krzyzak, and H. Walk. A Distribution-Free Theory of Nonparametric Regression. Springer Science & Business Media, 2006.
- M. Hallin, E. del Barrio, J. Cuesta-Albertos, and C. Matrán. Distribution and quantile functions, ranks and signs in dimension d: A measure transportation approach. *The Annals of Statistics*, 49:1139–1165, 2021.
- J.-B. Hiriart-Urruty and C. Lemaréchal. Convex analysis and minimization algorithms II: Advanced theory and bundle methods. Springer-Verlag, 1993.
- J.-B. Hiriart-Urruty and C. Lemaréchal. *Fundamentals of Convex Analysis*. Springer Science & Business Media, 2004.
- J.-C. Hütter and P. Rigollet. Minimax rates of estimation for smooth optimal transport maps. The Annals of Statistics, 49:1166–1194, 2021.
- L. V. Kantorovich. On the translocation of masses. In Dokl. Akad. Nauk. USSR (NS), volume 37, pages 199–201, 1942.

- L. V. Kantorovich. On a problem of Monge. In CR (Doklady) Acad. Sci. URSS (NS), volume 3, pages 225–226, 1948.
- M. Ledoux. The concentration of measure phenomenon, volume 89 of Mathematical Surveys and Monographs. American Mathematical Society, 2005.
- J. Lei. Convergence and concentration of empirical measures under Wasserstein distance in unbounded functional spaces. *Bernoulli*, 26:767–798, 2020.
- C. Letrouit and Q. Mérigot. Gluing methods for quantitative stability of optimal transport maps. arXiv preprint arXiv:2411.04908, 2024.
- W. Li and R. H. Nochetto. Quantitative stability and error estimates for optimal transport plans. IMA Journal of Numerical Analysis, 41(3):1941–1965, 2021.
- A. Makkuva, A. Taghvaei, S. Oh, and J. Lee. Optimal transport mapping via input convex neural networks. In *International Conference on Machine Learning*, pages 6672–6681. PMLR, 2020.
- T. Manole, S. Balakrishnan, J. Niles-Weed, and L. Wasserman. Central limit theorems for smooth optimal transport maps. arXiv preprint arXiv:2312.12407, 2023.
- T. Manole, S. Balakrishnan, J. Niles-Weed, and L. Wasserman. Plugin estimation of smooth optimal transport maps. *The Annals of Statistics*, 52(3):966–998, 2024.
- Q. Mérigot, A. Delalande, and F. Chazal. Quantitative stability of optimal transport maps and linearization of the 2-Wasserstein space. In *International Conference on Artificial Intelligence and Statistics*, pages 3186–3196. PMLR, 2020.
- G. Monge. Mémoire sur la théorie des déblais et des remblais. *Histoire de l'Académie Royale des Sciences de Paris*, 1781.
- G. Mordant. The entropic optimal (self-)transport problem: Limit distributions for decreasing regularization with application to score function estimation. *arXiv preprint* arXiv:2412.12007, 2024.
- B. Muzellec, A. Vacher, F. Bach, F.-X. Vialard, and A. Rudi. Near-optimal estimation of smooth transport maps with kernel sums-of-squares. arXiv preprint arXiv:2112.01907, 2021.
- J. Niles-Weed and Q. Berthet. Minimax estimation of smooth densities in wasserstein distance. The Annals of Statistics, 50(3):1519–1540, 2022.
- V. M. Panaretos and Y. Zemel. An invitation to statistics in Wasserstein space. Springer Nature, 2020.
- R. Peyre. Comparison between W_2 distance and \dot{H}^{-1} norm, and localisation of Wasserstein distance. *ESAIM: Control, Optimisation and Calculus of Variations*, 24:1489–1501, 2018.
- A.-A. Pooladian and J. Niles-Weed. Entropic estimation of optimal transport maps. arXiv preprint arXiv:2109.12004, 2021.
- A.-A. Pooladian, V. Divol, and J. Niles-Weed. Minimax estimation of discontinuous optimal transport maps: The semi-discrete case. In *International Conference on Machine Learning*, pages 28128–28150. PMLR, 2023.

- D. Radulović and M. Wegkamp. Necessary and sufficient conditions for weak convergence of smoothed empirical processes. *Statistics and Probability Letters*, 61(3):321–336, 2003.
- R. Sadhu, Z. Goldfeld, and K. Kato. Approximation rates of entropic maps in semidiscrete optimal transport. arXiv preprint arXiv:2411.07947, 2024a.
- R. Sadhu, Z. Goldfeld, and K. Kato. Stability and statistical inference for semidiscrete optimal transport maps. *The Annals of Applied Probability*, 34(6):5694–5736, 2024b.
- F. Santambrogio. Optimal Transport for Applied Mathematicians: Calculus of Variations, PDEs, and Modeling, volume 87. Birkhäuser, 2015.
- J. Segers. Graphical and uniform consistency of estimated optimal transport plans. arXiv preprint arXiv:2208.02508, 2022.
- A. Vacher and F.-X. Vialard. Convex transport potential selection with semi-dual criterion. Advances in Neural Information Processing Systems 36, 2022.
- A. Vacher, B. Muzellec, A. Rudi, F. Bach, and F.-X. Vialard. A dimension-free computational upper-bound for smooth optimal transport estimation. In *Conference on Learning Theory*, pages 4143–4173. PMLR, 2021.
- S. van de Geer. Empirical Processes in M-Estimation, volume 3 of Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, 2000.
- A. W. van der Vaart and J. A. Wellner. Weak Convergence and Empirical Processes. Springer Series in Statistics. Springer-Verlag, New York, 1996.
- C. Villani. Topics in Optimal Transportation. American Mathematical Soc., 2003.
- C. Villani. Optimal Transport: Old and New, volume 338. Springer Science & Business Media, 2008.
- W. Wang, D. Slepčev, S. Basu, J. A. Ozolek, and G. K. Rohde. A linear optimal transportation framework for quantifying and visualizing variations in sets of images. *International Journal* of Computer Vision, 101:254–269, 2013.
- J. Weed and F. Bach. Sharp asymptotic and finite-sample rates of convergence of empirical measures in Wasserstein distance. *Bernoulli*, 25:2620–2648, 2019.

A The Semi-Dual Functional and Brenier Potentials

In this section, we recollect some well-known properties of the semi-dual functional, its optimizers, and some properties of convex conjugates that we use in our proofs. Recall that for a given pair of distributions $P, Q \in \mathcal{P}_2(\Omega)$ we define a Brenier potential $\varphi_{P,Q}$ as any solution to the optimization problem:

$$\varphi_{P,Q} \in \operatorname*{argmin}_{\varphi \in L^1(P)} \int \varphi dP + \int \varphi^* dQ.$$

If we denote by $\pi_{P,Q}$ an optimal coupling of P and Q, then we have that (see for instance Theorem 5.10 in Villani [2008]):

$$\varphi_{P,Q}(X) + \varphi_{P,Q}^*(Y) = \langle X, Y \rangle_{\mathcal{F}}$$

for $\pi_{P,Q}$ almost every (X, Y). Using this it is straightforward to see that,

$$\mathcal{S}_{P,Q}(\varphi_{P,Q}) = \mathbb{E}_{(X,Y)\sim\pi_{P,Q}}\langle X, Y \rangle.$$
(23)

We also recall some well-known facts about convex conjugates (see Hiriart-Urruty and Lemaréchal [2004]), and derive some implications of the regularity assumption A0. For a convex function φ defined on Ω , we always have the Fenchel-Young inequality for any $x, y \in \Omega$:

$$\varphi(x) + \varphi^*(y) \ge \langle x, y \rangle.$$

For a convex, differentiable function φ , the Fenchel-Young inequality holds with equality when $y = \nabla \varphi(x)$, i.e.:

$$\varphi(x) + \varphi^*(\nabla\varphi(x)) = \langle x, \, \nabla\varphi(x) \rangle.$$

If φ is a strictly convex, differentiable function, then φ^* is also convex and differentiable. In this case, we also have the identities:

$$\nabla \varphi^* \circ \nabla \varphi(x) = x, \quad \nabla \varphi \circ \nabla \varphi^*(y) = y.$$

B Proof of Lemma 1

Recall that $\tilde{\pi}$ denotes the optimal coupling between \tilde{P} and \tilde{Q} . We have that,

$$\mathcal{S}_{\widetilde{P},\widetilde{Q}}(\varphi_0) - \mathcal{S}_{\widetilde{P},\widetilde{Q}}(\widetilde{\varphi}) = \mathbb{E}_{(X,Y)\sim\widetilde{\pi}} \left[\varphi_0(X) + \varphi_0^*(Y) - \widetilde{\varphi}(X) - \widetilde{\varphi}^*(Y)\right].$$

Now, we recall that:

$$\widetilde{\varphi}(X) + \widetilde{\varphi}^*(Y) = \langle X, Y \rangle, \quad \widetilde{\pi} \text{ a.e.}$$

By Assumption A0 on φ_0 , and using the Fenchel-Young (in)equality, we have that:

$$\varphi_0(X) + \varphi_0^*(\nabla \varphi_0(X)) = \langle X, \, \nabla \varphi_0(X) \rangle.$$

Putting these together, we obtain that:

$$\mathcal{S}_{\widetilde{P},\widetilde{Q}}(\varphi_0) - \mathcal{S}_{\widetilde{P},\widetilde{Q}}(\widetilde{\varphi}) = \mathbb{E}_{(X,Y)\sim\widetilde{\pi}} \left[\varphi_0^*(Y) - \varphi_0^*(T_0(X)) - \langle X, Y - T_0(X) \rangle \right]$$

as desired.

C Proof of Theorem 3

As discussed in the main text, our proof proceeds by upper and lower bounding the semidual difference $S_{\widehat{P},\widehat{Q}}(\varphi_0) - S_{\widehat{P},\widehat{Q}}(\varphi_{\widehat{P},\widehat{Q}})$. The lower bound on this difference follows from the argument in the main text, and we provide a complete proof of the upper bound in this Appendix.

In the remainder of the proof, for any pair of distributions P, Q we let $\pi_{P,Q}$ denote the optimal coupling between P and Q. Then let us define the random variables

$$X \sim P, \quad Y \sim Q, \quad U_1, U_2, U_3 \sim \widehat{P}, \quad V_1, V_2, V_3 \sim \widehat{Q},$$

with the following joint distributions:

$$\begin{aligned} & (U_1, Y) \sim \pi_{\widehat{P}, Q}, \quad (X, V_1) \sim \pi_{P, \widehat{Q}} \\ & (X, U_2) \sim \pi_{P, \widehat{P}}, \quad (Y, V_2) \sim \pi_{Q, \widehat{Q}} \\ & (X, Y) \sim \pi_{P, Q}, \quad (U_3, V_3) \sim \pi_{\widehat{P}, \widehat{Q}}. \end{aligned}$$

These joint distributions are summarized in the following figure:



To begin we derive a technical result, giving one-sample stability bounds for the semi-dual functional:

Lemma 4. Let $P, Q \in \mathcal{P}_2(\Omega)$, and assume A0, $A1(\alpha)$ and $A2(\beta)$ hold for some $\alpha, \beta > 0$. For any $\widehat{P}, \widehat{Q} \in \mathcal{P}_2(\Omega)$,

$$\mathcal{S}_{P,\widehat{Q}}(\varphi_0) - \mathcal{S}_{P,\widehat{Q}}(\varphi_{P,\widehat{Q}}) \le \frac{1}{2\alpha} W_2^2(\widehat{Q}, Q) - \mathbb{E}\langle X, V_1 - V_2 \rangle$$
(24)

$$\mathcal{S}_{\widehat{P},Q}(\varphi_0) - \mathcal{S}_{\widehat{P},Q}(\varphi_{\widehat{P},Q}) \le \frac{\beta}{2} W_2^2(\widehat{P},P) - \mathbb{E}\langle Y, U_1 - U_2 \rangle.$$
⁽²⁵⁾

With this result in place we return to our main proof. The following identity follows directly from the definition of the semi-dual functional:

$$\mathcal{S}_{\widehat{P},\widehat{Q}}(\varphi_0) - \mathcal{S}_{\widehat{P},\widehat{Q}}(\varphi_{\widehat{P},\widehat{Q}}) = \mathcal{S}_{\widehat{P},Q}(\varphi_0) + \mathcal{S}_{P,\widehat{Q}}(\varphi_0) - \mathcal{S}_{P,Q}(\varphi_0) - \mathcal{S}_{\widehat{P},\widehat{Q}}(\varphi_{\widehat{P},\widehat{Q}}).$$
(26)

Then we observe that by (23),

$$\begin{split} \mathcal{S}_{\widehat{P},Q}(\varphi_{\widehat{P},Q}) &= \mathbb{E} \langle U_1, Y \rangle \\ \mathcal{S}_{P,\widehat{Q}}(\varphi_{P,\widehat{Q}}) &= \mathbb{E} \langle X, V_1 \rangle \\ \mathcal{S}_{\widehat{P},\widehat{Q}}(\varphi_{\widehat{P},\widehat{Q}}) &= \mathbb{E} \langle U_3, V_3 \rangle \\ \mathcal{S}_{P,Q}(\varphi_0) &= \mathbb{E} \langle X, Y \rangle. \end{split}$$

From these facts, and (26) we obtain,

$$\begin{split} \mathcal{S}_{\widehat{P},\widehat{Q}}(\varphi_{0}) - \mathcal{S}_{\widehat{P},\widehat{Q}}(\varphi_{\widehat{P},\widehat{Q}}) &= \mathcal{S}_{\widehat{P},Q}(\varphi_{0}) - \mathcal{S}_{\widehat{P},Q}(\varphi_{\widehat{P},Q}) + \mathcal{S}_{P,\widehat{Q}}(\varphi_{0}) - \mathcal{S}_{P,\widehat{Q}}(\varphi_{P,\widehat{Q}}) \\ &+ \mathbb{E}\left[\langle U_{1}, Y \rangle + \langle X, V_{1} \rangle - \langle X, Y \rangle - \langle U_{3}, V_{3} \rangle\right] \\ &\leq \frac{1}{2\alpha} W_{2}^{2}(\widehat{Q}, Q) + \frac{\beta}{2} W_{2}^{2}(\widehat{P}, P) + \mathbb{E}\left[\langle X, V_{2} - Y \rangle + \langle Y, U_{2} \rangle - \langle U_{3}, V_{3} \rangle\right], \end{split}$$

where the second inequality uses Lemma 4. Now, by (7), we know that, $\mathbb{E}\langle U_3, V_3 \rangle \geq \mathbb{E}\langle U_2, V_2 \rangle$ and so we obtain:

$$\mathcal{S}_{\widehat{P},\widehat{Q}}(\varphi_0) - \mathcal{S}_{\widehat{P},\widehat{Q}}(\varphi) \le \frac{1}{2\alpha} W_2^2(\widehat{Q},Q) + \frac{\beta}{2} W_2^2(\widehat{P},P) + \mathbb{E} \langle V_2 - Y, X - U_2 \rangle.$$

Applying the Cauchy-Schwarz inequality to the final term and noting that $\mathbb{E}||X - U_2||_2^2 = W_2^2(\hat{P}, P)$ and $\mathbb{E}||V_2 - Y||_2^2 = W_2^2(\hat{Q}, Q)$, we obtain the Theorem.

C.1 Proof of Lemma 4

The proofs of the two claims are similar, and we prove each in turn. For the first claim, notice that

$$\mathcal{S}_{P,\widehat{Q}}(\varphi_0) - \mathcal{S}_{P,\widehat{Q}}(\varphi_{P,\widehat{Q}}) = \mathbb{E}\left[\varphi_0(X) + \varphi_0^*(V_1) - \varphi_{P,\widehat{Q}}(X) - \varphi_{P,\widehat{Q}}^*(V_1)\right].$$

Furthermore, it holds almost surely that

$$\varphi_{P,\widehat{Q}}(X) + \varphi_{P,\widehat{Q}}^*(V_1) = \langle X, V_1 \rangle, \quad \varphi_0(X) + \varphi_0^*(Y) = \langle X, Y \rangle.$$

Now, since V_2 is equal in distribution to V_1 , we have

$$\begin{aligned} \mathcal{S}_{P,\widehat{Q}}(\varphi_0) - \mathcal{S}_{P,\widehat{Q}}(\varphi_{P,\widehat{Q}}) &= \mathbb{E}\left[\varphi_0^*(V_1) - \varphi_0^*(Y) - \langle X, V_1 - Y \rangle\right] \\ &= \mathbb{E}\left[\varphi_0^*(V_2) - \varphi_0^*(Y) - \langle \nabla \varphi_0^*(Y), V_2 - Y \rangle\right] + \mathbb{E}\langle X, V_2 - V_1 \rangle. \end{aligned}$$

Using this fact together with the $1/\alpha$ smoothness of φ_0 , we obtain the bound:

$$\mathcal{S}_{P,\widehat{Q}}(\varphi_0) - \mathcal{S}_{P,\widehat{Q}}(\varphi_{P,\widehat{Q}}) \le \frac{1}{2\alpha} \mathbb{E} \|V_2 - Y\|_2^2 - \mathbb{E} \langle X, V_1 - V_2 \rangle,$$

which gives the first claim. To prove the second claim, note again that

$$\mathcal{S}_{\widehat{P},Q}(\varphi_0) - \mathcal{S}_{\widehat{P},Q}(\varphi_{\widehat{P},Q}) = \mathbb{E}\left[\varphi_0(U_1) + \varphi_0^*(Y) - \varphi_{\widehat{P},Q}(U_1) - \varphi_{\widehat{P},Q}^*(Y)\right].$$

Furthermore, it holds almost surely that

$$\varphi_{\widehat{P},Q}(U_1) + \varphi_{\widehat{P},Q}^*(Y) = \langle U_1, Y \rangle, \quad \varphi_0(X) + \varphi_0^*(Y) = \langle X, Y \rangle.$$

Now, since U_2 is equal in distribution to U_1 , we have

$$\begin{aligned} \mathcal{S}_{\widehat{P},Q}(\varphi_0) - \mathcal{S}_{\widehat{P},Q}(\varphi_{\widehat{P},Q}) &= \mathbb{E}\left[\varphi_0(U_1) - \varphi_0(X) - \langle Y, U_1 - X \rangle\right] \\ &= \mathbb{E}\left[\varphi_0(U_2) - \varphi_0(X) - \langle Y, U_2 - X \rangle\right] - \mathbb{E}\langle Y, U_1 - U_2 \rangle. \end{aligned}$$

Using this fact together with the smoothness of φ_0 , we obtain the bound:

$$\mathcal{S}_{\widehat{P},Q}(\varphi_0) - \mathcal{S}_{\widehat{P},Q}(\varphi_{\widehat{P},Q}) \le \frac{\beta}{2} \mathbb{E} \|U_2 - X\|_2^2 - \mathbb{E} \langle Y, U_1 - U_2 \rangle,$$

which gives the desired claim.

D Proof of Lemma 2

For ease of notation we denote by \hat{T} the nearest neighbor OT map $\hat{T}_{nm}^{1\text{NN}}$. We observe that we can decompose the $L^2(P)$ error of the nearest neighbor map as:

$$\begin{split} \int \|\widehat{T} - T_0\|_2^2 dP &= \sum_{i=1}^n \int_{V_i} \|\widehat{T} - T_0\|_2^2 dP \\ &\leq 2 \sum_{i=1}^n \int_{V_i} \|\widehat{T}(X_i) - T_0(X_i)\|_2^2 dP + 2 \sum_{i=1}^n \int_{V_i} \|T_0(x) - T_0(X_i)\|_2^2 dP(x) \\ &\stackrel{(i)}{\leq} 2 \sum_{i=1}^n P(V_i) \|\widehat{T}(X_i) - T_0(X_i)\|_2^2 + 2\beta^2 \sum_{i=1}^n \int_{V_i} \|x - X_i\|_2^2 dP(x) \\ &\leq 2 \max_i P(V_i) \sum_{i=1}^n \|\widehat{T}(X_i) - T_0(X_i)\|_2^2 + 2\beta^2 \mathbb{E}_{X \sim P} \left[\|X - \operatorname{nn}(X)\|_2^2 |X_1, \dots, X_n \right], \end{split}$$

as desired. The inequality (i) follows from $A2(\beta)$, noting that $T_0 = \nabla \varphi_0$, and that for a differentiable convex function, smoothness is equivalent to Lipschitzness of the gradient [Hiriart-Urruty and Lemaréchal, 2004].

E Proof of Lemma 3

The first claim follows directly from past work, and we focus on the remaining claims. Similar results, albeit under much stronger boundedness assumptions, are classical in the analysis of k-NN regression (see, for instance, Chapter 6 of Györfi et al. [2006]).

Proof of Claim (20): Let $B_{x,r}$ denote the radius r ball in \mathbb{R}^d centered at x, and let N(S,r) denote the r-covering number of the set $S \subseteq \mathbb{R}^d$ with respect to the Euclidean distance. We partition \mathbb{R}^d into the sets:

$$S_0 = B_{0,1}, \quad S_j = B_{0,2^j} \setminus B_{0,2^{j-1}}, \quad j = 1, 2, \dots$$

Recalling that we denote by nn(x) the nearest neighbor of x in a sample X_1, \ldots, X_n , our goal is to bound:

$$\mathbb{E}[e_2] = \mathbb{E}_{X, X_1, \dots, X_n \sim P} \|X - \operatorname{nn}(X)\|_2^2 = \int_0^\infty \mathbb{P}(\|X - \operatorname{nn}(X)\|_2 \ge \sqrt{t}) dt.$$

For a fixed $x \in S_j$ we can write,

$$\mathbb{P}(\|x - nn(x)\|_2 \ge \sqrt{t}) = (1 - P(B_{x,\sqrt{t}}))^n,$$

so we obtain that,

$$\begin{split} \mathbb{P}(\|X - \operatorname{nn}(X)\|_2 \geq \sqrt{t}) &= \sum_j \int_{S_j} \mathbb{P}(\|x - \operatorname{nn}(x)\|_2 \geq \sqrt{t}) dP(x) \\ &= \sum_j \int_{S_j} (1 - P(B_{x,\sqrt{t}}))^n dP(x) \\ &\leq \sum_j \int_{S_j} \exp(-nP(B_{x,\sqrt{t}})) dP(x). \end{split}$$

Suppose we cover the set S_j with radius $\sqrt{t}/2$ balls, and use the following implication

$$x \in B_{x^c,\sqrt{t}/2} \implies B_{x^c,\sqrt{t}/2} \subseteq B_{x,\sqrt{t}},$$

where x^c is the center closest to x. Then, using the fact that $x \exp(-nx) \leq 1/n$, we get,

$$P(||X - nn(X)||_2 \ge \sqrt{t}) \le \sum_j \min\{P(S_j), N(S_j, \sqrt{t}/2)/n\}$$

Now, using Markov's inequality and the moment condition (19), we can upper bound the probability content of S_j as follows:

$$P(S_j) \le P(||x||_2 \ge 2^{j-1}) \le 2^{-r(j-1)}.$$

Furthermore, noting that $N(S_j, \sqrt{t}) \leq (2^j/\sqrt{t})^d$, we obtain

$$\mathbb{E} \|X - \operatorname{nn}(X)\|_{2}^{2} \lesssim \int_{0}^{\infty} \sum_{j} \min\left\{2^{-r(j-1)}, \frac{(2^{j}/\sqrt{t})^{d}}{n}\right\} dt$$
$$\lesssim \sum_{j} \left[2^{-rj} \frac{2^{2j(d+r)/d}}{n^{2/d}}\right] \lesssim n^{-2/d},$$

provided that $r \ge 2d/(d-2)$.

Proof of Claim (21): First, we note that by a standard uniform convergence argument (see, for instance, Lemma 7 in Chaudhuri and Dasgupta [2010] or Balakrishnan et al. [2013]), with probability at least $1 - \delta$, any ball B with $P(B) \gtrsim \log(n/\delta)/n$ must have at least one sample point in it. We condition on this event throughout the remainder of the proof.

We next use Lemma 6.2 from Györfi et al. [2006], which shows the following. Suppose, for any x and $a \ge 0$ we denote by $S_a(x)$ the set

$$S_a(x) := \{ y : P(B_{y, \|x-y\|_2}) \le a \}.$$

Then, since P is non-atomic, it holds for every $x \in \mathbb{R}^d$ that,

$$P(S_a(x)) \lesssim a. \tag{27}$$

For any sample point $x \in \{X_1, \ldots, X_n\}$ let us denote by V_x the Voronoi cell containing x. If $y \in V_x$, then $B_{y,||x-y||_2}$ must not contain any sample point other than x. This in turn means that $y \in S_a(x)$ for $a \simeq \log(n/\delta)/n$. Consequently, using the fact that P is non-atomic, we obtain from (27) that:

$$P(V_x) \lesssim \frac{\log(n/\delta)}{n}$$
.

Thus, letting $Z := \max_i P(V_i)$, we have shown that with probability at least $1 - \delta$,

$$Z \lesssim \frac{\log(n/\delta)}{n}.$$

From this we obtain that

$$\mathbb{E}[Z^2] \lesssim \delta + \left[\frac{\log(n/\delta)}{n}\right]^2 \lesssim \left[\frac{\log n}{n}\right]^2,$$

26

by choosing δ appropriately.