REGULARIZING EFFECT OF THE SPATIALLY HOMOGENEOUS LANDAU EQUATION WITH SOFT POTENTIAL

XIAODONG CAO & CHAOJIANG XU & YAN XU

ABSTRACT. This paper investigates the Cauchy problem of the spatially homogeneous Landau equation with soft potential under the perturbation framework to global equilibrium. We prove that the solution to the Cauchy problem exhibits analyticity in the time variable and the Gelfand-Shilov regularizing effect in the velocity variables.

1. INTRODUCTION

The Cauchy problem for the spatially homogeneous Landau equation is given by

$$\begin{cases} \partial_t F = Q(F, F), \\ F|_{t=0} = F_0, \end{cases}$$

$$\tag{1.1}$$

where $F = F(t, v) \ge 0$ represents the density distribution function at time $t \ge 0$, with the velocity variable $v \in \mathbb{R}^3$. The Landau bilinear collision operator is defined as follows:

$$Q(G,F)(v) = \sum_{j,k=1}^{3} \partial_j \bigg(\int_{\mathbb{R}^3} a_{jk} (v - v_*) [G(v_*)\partial_k F(v) - \partial_k G(v_*)F(v)] dv_* \bigg),$$
(1.2)

where the non-negative symmetric matrix (a_{jk}) is given by

$$a_{jk}(v) = (\delta_{jk}|v|^2 - v_j v_k)|v|^{\gamma}, \quad \gamma \ge -3.$$
 (1.3)

The parameter γ leads to the classification of hard potential if $\gamma > 0$, Maxwellian molecules if $\gamma = 0$, soft potential if $-3 < \gamma < 0$ and Coulombian potential if $\gamma = -3$.

The Landau equation is one of the fundamental kinetic equations obtained as a limit of the Boltzmann equation when grazing collisions prevail [23]. Extensive research has been conducted on the Landau equation with hard potential. Desvillettes-Villani [8] obtained the smoothness of solutions to the spatially homogeneous Landau equation.

For the hard potential case, the analytic smoothing effect for the spatially homogeneous Landau equation was established in [4], assuming an initial datum with finite mass and energy. Under the perturbation setting to the normalized global Maxwellian, the analytic smoothing effect to solution has been studied in [14], Gevrey regularity was treated in [5, 6], analytic Gelfand-Shilov smoothing effect was proven in [16]. For the Maxwellian molecules case, the existence, uniqueness and smoothness of solutions to the spatially homogeneous equation have been studied in [24], again under the assumption of an initial

Date: February 19, 2025.

²⁰¹⁰ Mathematics Subject Classification. 35B65,76P05,82C40.

Key words and phrases. Spatially homogeneous Landau equation, analytic and Gelfand-Shilov smoothing effect, soft potentials.

datum with finite mass and energy. Additionally, the analytic and Gelfand-Shilov smoothing effects were addressed in [13, 19, 20].

We are now interested in the spatially homogeneous Landau equation with soft potentials. In this work, we focus on studying the linearization of the Landau equation (1.1) around the Maxwellian distribution $\mu(v) = (2\pi)^{-\frac{3}{2}} e^{-\frac{|v|^2}{2}}$. Considering the fluctuation of the density distribution function

$$F(t,v) = \mu(v) + \sqrt{\mu(v)}f(t,v)$$

and noting that $Q(\mu, \mu) = 0$, the Cauchy problem (1.1) can be reformulated as

$$\begin{cases} \partial_t f + \mathcal{L}f = \Gamma(f, f), \\ f|_{t=0} = f_0, \end{cases}$$
(1.4)

where $F_0 = \mu + \sqrt{\mu} f_0$, and

$$\Gamma(f, f) = \mu^{-\frac{1}{2}} Q(\sqrt{\mu} f, \sqrt{\mu} f),$$

with the linear operator \mathcal{L} defined as

$$\mathcal{L} = \mathcal{L}_1 + \mathcal{L}_2$$
, with $\mathcal{L}_1 f = -\Gamma(\sqrt{\mu}, f)$, $\mathcal{L}_2 f = -\Gamma(f, \sqrt{\mu})$. (1.5)

The diffusion part \mathcal{L}_1 is expressed as

$$\mathcal{L}_1 f = -\nabla_v \cdot \left[A(v)\nabla_v f\right] + \left(A(v)\frac{v}{2} \cdot \frac{v}{2}\right)f - \nabla_v \cdot \left[A(v)\frac{v}{2}\right]f,\tag{1.6}$$

where $A(v) = (\bar{a}_{jk})_{1 \le j,k \le 3}$ is a symmetric matrix, and

$$\bar{a}_{jk} = a_{jk} * \mu = \int_{\mathbb{R}^3} \left(\delta_{jk} |v - v'|^2 - (v_j - v'_j)(v_k - v'_k) \right) |v - v'|^{\gamma} \mu(v') dv'.$$

Thus

$$\mathcal{L}_1 f \sim \langle v \rangle^{\gamma} (-\Delta_v + \frac{|v|^2}{4}),$$

and this operator is degenerate at $|v| = +\infty$ for the soft potential case $\gamma < 0$.

For soft potentials, existence and uniqueness results can be found in [23, 25]. Regarding smoothing properties, it was shown in [15] that solutions to the linear Landau equation with soft potentials exhibit analytic smoothness in both the velocity and time variables. The Gelfand-Shilov smoothing effect for moderately soft potentials $-2 < \gamma < 0$, was examined in [14]. Additionally, various regularity results in a near-equilibrium setting have been discussed in [2, 3, 22].

We mention that our discussion is based on the following function spaces. Let Ω be an open subset of \mathbb{R}^n . The analytical space $\mathcal{A}(\Omega)$, consists of smooth functions that satisfy the following condition: $f \in \mathcal{A}(\Omega)$ if there exists a constant C such that for all $\alpha \in \mathbb{N}^n$

$$\left\|\partial^{\alpha} f\right\|_{L^{2}(\Omega)} \leq C^{|\alpha|+1} \alpha!.$$

The Gelfand-Shilov spaces $S_{\nu}^{\sigma}(\mathbb{R}^n)$, with $\sigma, \nu > 0, \sigma + \nu \ge 1$, consist of smooth functions that satisfy the following condition: $f \in S_{\nu}^{\sigma}(\mathbb{R}^n)$ if there exists a constant C such that for all multi-indices $\alpha, \beta \in \mathbb{N}^n$,

$$\left\| v^{\beta} \partial^{\alpha} f \right\|_{L^{2}(\mathbb{R}^{n})} \leq C^{|\alpha| + |\beta| + 1}(\alpha!)^{\sigma}(\beta!)^{\nu}.$$

Before presenting the main result of the paper, we first introduce some notation. We denote the harmonic oscillators operator by $\mathcal{H} = -\Delta_v + \frac{|v|^2}{4}$ and the associated gradient by

$$\nabla_{\mathcal{H}_{\pm}} = (A_{\pm,1}, A_{\pm,2}, A_{\pm,3}),$$

where

$$A_{\pm,k} = \frac{1}{2} v_k \mp \partial_{v_k}, (1 \le k \le 3), \quad A_{\pm}^{\alpha} = A_{\pm,1}^{\alpha_1} A_{\pm,2}^{\alpha_2} A_{\pm,3}^{\alpha_3}, \ (\alpha \in \mathbb{N}^3)$$

We also define, for some constants $c_0 > 0$ and $0 < b \le 2$,

$$\omega_t(v) = e^{\frac{c_0}{1+t} \langle v \rangle^b}, \quad t \ge 0, \ v \in \mathbb{R}^3,$$

with $\langle v \rangle = (1 + |v|^2)^{\frac{1}{2}}$. Now, the main result of this paper is stated as follows.

Theorem 1.1. For the soft potential $-3 < \gamma < 0$, assume that there exist positive constants c_0 and $0 < b \leq 2$ such that for any $f_0 \in L^2(\mathbb{R}^3)$ satisfying $\|\omega_0 f_0\|_{L^2(\mathbb{R}^3)}$ small enough, then the Cauchy problem (1.4) admits a unique solution $\omega_t f \in \mathcal{A}(]0, \infty[; S^{\sigma}_{\sigma}(\mathbb{R}^3))$, where $\sigma = \max\{1, \frac{b-\gamma}{2b}\}$. Furthermore, for any T > 0, there exists a constant $C_T > 0$ which depends on γ , b, c_0 and T, such that

$$\left\| \omega_t \nabla_{\mathcal{H}_+}^m f(t) \right\|_{L^2(\mathbb{R}^3)} \le \left(\frac{C_T^{m+1} m!}{t^m} \right)^{\sigma}, \quad \forall \ m \in \mathbb{N}, \ \forall \ 0 < t \le T,$$
(1.7)

and

$$\|\partial_t^m f(t)\|_{L^2(\mathbb{R}^3)} \le \frac{C_T^{m+1}}{t^m} m!, \quad \forall \ m \in \mathbb{N}, \ \forall \ 0 < t \le T.$$

Remark 1.2. The results of reference [17] indicate that the solution of (1.4) belongs to $S_1^1(\mathbb{R}^3)$ when $-1 \leq \gamma < 0$, and belongs to $S_{\frac{1}{\gamma+2}}^{\frac{1}{\gamma+2}}(\mathbb{R}^3)$ when $-2 < \gamma < -1$. To overcome the difficulty of degeneracy, we introduce an exponential type weight $\omega_t(v) = e^{\frac{c_0}{1+t}\langle v \rangle^b}$ where the parameter *b* affects the analyticity index. Specifically, it implies that: the solution belongs to $\mathcal{A}(]0, \infty[; S_{\frac{b-\gamma}{2b}}^{\frac{b-\gamma}{2b}}(\mathbb{R}^3))$ for $-3 < \gamma < -b$. So that we can obtain the Gelfand-Shilov smoothing effect for all soft potential.

Remark 1.3. In [12], they consider the spatially inhomogeneous Boltzmann equation without angular cutoff for soft potentials and obtained the global Gevrey regularity with the Gevrey index $\max\{\frac{2-\gamma}{4s}, 1\}$, for initial data with some exponential weight(where the parameter b = 2).

Remark 1.4. In our main theorem 1.1, by Proposition 5.2 of [16] and Theorem 2.1 of [10], we can obtain

$$\left\| \omega_t v^\beta \partial_v^\alpha f(t) \right\|_{L^2(\mathbb{R}^3)} \le \left(\frac{C_T^{\alpha|+|\beta|+1} \beta! \alpha!}{t^{|\alpha|+|\beta|}} \right)^\sigma, \quad \forall \ \alpha, \beta \in \mathbb{N}^3, \ \forall \ 0 < t \le T.$$

Then we have

$$e^{\frac{c_0}{1+T}\langle v \rangle^b + c_1 t \langle v \rangle^{\frac{1}{\sigma}}} f \in L^2(\mathbb{R}^3), \quad 0 < t \le T.$$

(1) For the case of $-b \leq \gamma \leq 0$, we have proved that

$$e^{\frac{c_0}{1+T}\langle v \rangle^b + c_1 t \langle v \rangle} f \in L^2(\mathbb{R}^3), \quad 0 < t \le T,$$

Thus if $b \ge 1$, we have not improved the decreasing rate, and we obtain

$$e^{\frac{c_0}{1+T}\langle v \rangle^o} f \in L^2(\mathbb{R}^3), \quad 0 < t \le T,$$

on the other hand, if b < 1, we improve the decreasing rate, i. e.

$$e^{c_1 t \langle v \rangle} f \in L^2(\mathbb{R}^3), \quad 0 < t \le T$$

(2) For the case of $-3 < \gamma < -b$, we have proved that

$$e^{\frac{c_0}{1+T}\langle v \rangle^b + c_1 t \langle v \rangle^{\frac{2b}{b-\gamma}}} f \in L^2(\mathbb{R}^3), \quad 0 < t \le T.$$

Thus if $0 < \frac{2}{b-\gamma} \le 1$, the solution has the same decreasing rate as the initial data, i. e.

$$e^{\frac{c_0}{1+T}\langle v \rangle^b} f \in L^2(\mathbb{R}^3), \quad 0 < t \le T$$

On the other hand, once $\frac{2}{b-\gamma} > 1$, then we improve the decreasing rate, i. e.

$$e^{c_1 t \langle v \rangle^{\frac{2b}{b-\gamma}}} f \in L^2(\mathbb{R}^3), \quad 0 < t \le T.$$

The above analysis is easier to understand with the following figure: only in the red area (2), we improve the decreasing rate with respect to the initial date.



2. Schema of proof for the main theorem

We begin by recalling some preliminary results regarding Landau operators. For simplicity, with $r \in \mathbb{R}$, we denote the weighted Lebesgue spaces

$$\|\langle \cdot \rangle^r f\|_{L^p(\mathbb{R}^3)} = \|f\|_{p,r}, \quad 1 \le p \le \infty,$$

where the notation $\langle v \rangle = (1 + |v|^2)^{\frac{1}{2}}$. For the matrix A defined in (1.6), denote

$$\|f\|_{A}^{2} = \sum_{j,k=1}^{3} \int \left(\bar{a}_{jk}\partial_{j}f\partial_{k}f + \frac{1}{4}\bar{a}_{jk}v_{j}v_{k}f^{2}\right)dv = \frac{1}{2}\sum_{j,k=1}^{3} \int \bar{a}_{jk}\left(A_{-,j}fA_{-,k}f + A_{+,j}fA_{+,k}f\right)dv.$$

From corollary 1 in [9] and Proposition 2.3 in [16], for $\gamma > -3$, there exists a constant $C_1 > 0$ such that

$$\|f\|_{A}^{2} \geq C_{1}\left(\|\mathbf{P}_{v}\nabla f\|_{2,\frac{\gamma}{2}}^{2} + \|(\mathbf{I} - \mathbf{P}_{v})\nabla f\|_{2,1+\frac{\gamma}{2}}^{2} + \|f\|_{2,1+\frac{\gamma}{2}}^{2}\right),\tag{2.1}$$

and

$$\|f\|_{A}^{2} \geq C_{1}\left(\left\|\mathbf{P}_{v}\nabla_{\mathcal{H}_{\pm}}f\right\|_{2,\frac{\gamma}{2}}^{2} + \left\|(\mathbf{I}-\mathbf{P}_{v})\nabla_{\mathcal{H}_{\pm}}f\right\|_{2,1+\frac{\gamma}{2}}^{2}\right).$$
(2.2)

For any vector-valued function $G(v) = (G_1, G_2, G_3)$, defining the projection to the vector $v \in \mathbb{R}^3$ as

$$(\mathbf{P}_v G)_j = \sum_{k=1}^3 G_k v_k \frac{v_j}{|v|^2}, \quad 1 \le j \le 3.$$

Since $\nabla f = \mathbf{P}_v \nabla f + (\mathbf{I} - \mathbf{P}_v) \nabla f$, combining the inequality (2.1), we have

$$\|f\|_{A} \ge C_1 \left(\|\nabla f\|_{2,\frac{\gamma}{2}} + \|f\|_{2,1+\frac{\gamma}{2}} \right), \tag{2.3}$$

and

$$\|f\|_{A} \ge C_1 \left\|\nabla_{\mathcal{H}_{\pm}} f\right\|_{2,\frac{\gamma}{2}}.$$
(2.4)

We begin with a singular integration. Review that, for any $\gamma > -3$ and $\delta > 0$, we have

$$\int_{\mathbb{R}^3} |v - v'|^{\gamma} e^{-\delta |v'|^2} dv' \le C_{\gamma,\delta} \langle v \rangle^{\gamma}.$$
(2.5)

For later use, we need the following results for the linear Landau operator, which have been proved in [26]. Lemma 2.1. [26] Let $-3 < \gamma < 0$, then for any $0 < \epsilon_1 < 1$, there exists a constant $C_{\epsilon_1} > 0$ such that for any suitable function f

$$(1-\epsilon_1) \|f\|_A^2 \le (\mathcal{L}_1 f, f)_{L^2} + C_{\epsilon_1} \|f\|_{2,\frac{\gamma}{2}}^2.$$

Now, we construct the trilinear estimate of the nonlinear Landau operator.

Proposition 2.2. Let $-3 < \gamma < 0$, then there exists a constant $C_2 > 0$, only depends on γ , such that for any suitable functions F, G and H, we have

$$\left| (\Gamma(F,G),H)_{L^{2}(\mathbb{R}^{3})} \right| \leq C_{2} \, \|F\|_{2,\frac{\gamma}{2}} \, \|G\|_{A} \, \|H\|_{A} \, .$$

The proof of this Proposition is similar to that in [14], we just need to pay more attention to the parameter $\gamma < 0$.

Using the equation (1.4), we can obtain the following L^2 -estimate

$$\frac{1}{2} \frac{d}{dt} \left\| t^{m\sigma} \nabla^m_{\mathcal{H}_+} f(t) \right\|_{L^2(\mathbb{R}^3)}^2 + \left(t^{m\sigma} \nabla^m_{\mathcal{H}_+} \mathcal{L}f, t^{m\sigma} \nabla^m_{\mathcal{H}_+} f \right)_{L^2(\mathbb{R}^3)} \\
= m t^{2m\sigma - 1} \left\| \nabla^m_{\mathcal{H}_+} f(t) \right\|_{L^2(\mathbb{R}^3)}^2 + \left(t^{m\sigma} \nabla^m_{\mathcal{H}_+} \Gamma(f, f), t^{m\sigma} \nabla^m_{\mathcal{H}_+} f \right)_{L^2(\mathbb{R}^3)}.$$
(2.6)

Using Lemma 2.1, we have

$$\left(t^{m\sigma}\mathcal{L}_{1}\nabla_{\mathcal{H}_{+}}^{m}f, t^{m\sigma}\nabla_{\mathcal{H}_{+}}^{m}f\right)_{L^{2}(\mathbb{R}^{3})} \geq \frac{1}{2}\left\|t^{m\sigma}\nabla_{\mathcal{H}_{+}}^{m}f\right\|_{A}^{2} - C_{3}\left\|t^{m\sigma}\nabla_{\mathcal{H}_{+}}^{m}f\right\|_{2,\frac{\gamma}{2}}^{2}$$

then we can obtain

$$\begin{aligned} &\frac{1}{2} \frac{d}{dt} \left\| t^{m\sigma} \nabla_{\mathcal{H}_{+}}^{m} f(t) \right\|_{L^{2}(\mathbb{R}^{3})}^{2} + \frac{1}{2} \left\| t^{m\sigma} \nabla_{\mathcal{H}_{+}}^{m} f \right\|_{A}^{2} \\ &\leq \left| (m\sigma) t^{2m\sigma-1} \left\| \nabla_{\mathcal{H}_{+}}^{m} f(t) \right\|_{L^{2}(\mathbb{R}^{3})}^{2} \right| + C_{3} \left\| t^{m\sigma} \nabla_{\mathcal{H}_{+}}^{m} f \right\|_{2,\frac{\gamma}{2}}^{2} \\ &+ \left(t^{m\sigma} \nabla_{\mathcal{H}_{+}}^{m} \Gamma(f,f), t^{m\sigma} \nabla_{\mathcal{H}_{+}}^{m} f \right)_{L^{2}(\mathbb{R}^{3})} + \left| \left(t^{m\sigma} \nabla_{\mathcal{H}_{+}}^{m} \mathcal{L}_{2} f, t^{m\sigma} \nabla_{\mathcal{H}_{+}}^{m} f \right)_{L^{2}(\mathbb{R}^{3})} \right| \\ &+ \left| \left(t^{m\sigma} [\nabla_{\mathcal{H}_{+}}^{m}, \mathcal{L}_{1}] f, t^{m\sigma} \nabla_{\mathcal{H}_{+}}^{m} f \right)_{L^{2}(\mathbb{R}^{3})} \right|. \end{aligned}$$

The second term in the right hand side $\left\|t^{m\sigma}\nabla_{\mathcal{H}_{+}}^{m}f\right\|_{2,\frac{\gamma}{2}}^{2}$ can be treated by using Gronwall's inequality. The third-fifth term can be treated as the hard potential case using induction. However the first term $(m\sigma)t^{2m\sigma-1}\left\|\nabla_{\mathcal{H}_{+}}^{m}f(t)\right\|_{L^{2}(\mathbb{R}^{3})}^{2}$ cannot be bounded through induction. Since

$$(m\sigma)t^{2m\sigma-1} \left\| \nabla_{\mathcal{H}_{+}}^{m} f(t) \right\|_{L^{2}(\mathbb{R}^{3})}^{2} = (m\sigma)t^{2m\sigma-1} \left(\nabla_{\mathcal{H}_{+}}^{m} f(t), \nabla_{\mathcal{H}_{+}}^{m} f(t) \right)_{L^{2}(\mathbb{R}^{3})}$$
$$\leq (m\sigma)^{2}t^{2(m\sigma-1)} \left\| \nabla_{\mathcal{H}_{+}} \nabla_{\mathcal{H}_{+}}^{m-1} f(t) \right\|_{L^{2}(\mathbb{R}^{3})}^{2} + t^{2m\sigma} \left\| \nabla_{\mathcal{H}_{+}}^{m} f(t) \right\|_{L^{2}(\mathbb{R}^{3})}^{2},$$

and for the soft potential case $\gamma < 0$, due to (2.4), the term $\left\| \nabla_{\mathcal{H}_{+}} \nabla_{\mathcal{H}_{+}}^{m-1} f(t) \right\|_{L^{2}(\mathbb{R}^{3})}^{2}$ cannot be controlled by $\left\| \nabla_{\mathcal{H}_+}^{m-1} f(t) \right\|_A^2$.

On the other hand, by using the interpolation, for b > 0, $\gamma < 0$ and $\sigma = \max\{1, \frac{b-\gamma}{2b}\}$, we have that for $0 < t \leq T$,

$$(m\sigma)t^{2m\sigma-1} \left\| \nabla_{\mathcal{H}_{+}}^{m} f(t) \right\|_{L^{2}(\mathbb{R}^{3})}^{2}$$

$$= (m\sigma)t^{2m\sigma-1} \int_{\mathbb{R}^{3}} \left| \langle v \rangle^{\frac{b}{2}} \nabla_{\mathcal{H}_{+}}^{m} f(t,v) \right|^{2\cdot\frac{-\gamma}{b-\gamma}} \left| \langle v \rangle^{\frac{\gamma}{2}} \nabla_{\mathcal{H}_{+}}^{m} f(t,v) \right|^{2\cdot\frac{b}{b-\gamma}} dv$$

$$\leq (m\sigma)t^{\frac{2b\sigma}{b-\gamma}-1} \left\| \langle \cdot \rangle^{\frac{b}{2}} t^{m\sigma} \nabla_{\mathcal{H}_{+}}^{m} f(t) \right\|_{L^{2}(\mathbb{R}^{3})}^{\frac{-2\gamma}{b-\gamma}} \left(\frac{1}{C_{1}} \left\| t^{(m-1)\sigma} \nabla_{\mathcal{H}_{+}}^{m-1} f(t) \right\|_{A} \right)^{\frac{2b}{b-\gamma}}$$

$$\leq \epsilon \left\| \langle \cdot \rangle^{\frac{b}{2}} t^{m\sigma} \nabla_{\mathcal{H}_{+}}^{m} f(t) \right\|_{L^{2}(\mathbb{R}^{3})}^{2} + C_{\epsilon} \left(\frac{\tilde{T}_{0}m\sigma}{C_{1}} \right)^{\frac{b}{b-\gamma}} \left\| t^{(m-1)\sigma} \nabla_{\mathcal{H}_{+}}^{m-1} f(t) \right\|_{A}^{2}$$

$$\leq \epsilon \left\| \langle \cdot \rangle^{\frac{b}{2}} t^{m\sigma} \nabla_{\mathcal{H}_{+}}^{m} f(t) \right\|_{L^{2}(\mathbb{R}^{3})}^{2} + \tilde{C}_{\epsilon} (m-1)^{2\sigma} \left\| t^{(m-1)\sigma} \nabla_{\mathcal{H}_{+}}^{m-1} f(t) \right\|_{A}^{2},$$

here we use $\frac{2b\sigma}{b-\gamma} - 1 \ge 0$, and $\tilde{T}_0 = T^{\frac{2b\sigma}{b-\gamma}-1}$. Now, the second term on the right-hand side can be controlled by induction, but the first term on the right-hand side is with a positive weight $\langle v \rangle^{\frac{b}{2}}$. So we need to use the following exponential-type weighted functions

$$\omega_t(v) = e^{\frac{c_0}{1+t} \langle v \rangle^b},$$

with $c_0 > 0$, $0 < b \le 2$, then with the equation

$$\partial_t \left(\omega_t f \right) + \frac{c_0}{(1+t)^2} \langle v \rangle^b (\omega_t f) + \omega_t \mathcal{L} f = \omega_t \Gamma(f, f),$$

we can establish the following L^2 -estimate

$$\frac{1}{2} \frac{d}{dt} \left\| \omega_t t^{m\sigma} \nabla^m_{\mathcal{H}_+} f(t) \right\|_{L^2(\mathbb{R}^3)}^2 + \frac{c_0}{(1+t)^2} \left\| \langle \cdot \rangle^{\frac{b}{2}} \omega_t t^{m\sigma} \nabla^m_{\mathcal{H}_+} f(t) \right\|_{L^2(\mathbb{R}^3)}^2 + \left(\omega_t t^{m\sigma} \nabla^m_{\mathcal{H}_+} f(t) \right\|_{L^2(\mathbb{R}^3)}^2 \\
= (m\sigma) t^{2m\sigma-1} \left\| \omega_t \nabla^m_{\mathcal{H}_+} f(t) \right\|_{L^2(\mathbb{R}^3)}^2 + \left(\omega_t t^{m\sigma} \nabla^m_{\mathcal{H}_+} \Gamma(f,f), \ \omega_t t^{m\sigma} \nabla^m_{\mathcal{H}_+} f \right)_{L^2(\mathbb{R}^3)},$$

The primary distinction from (2.6) lies in the left-hand side above, where we gain a term

$$\left\|\langle\cdot\rangle^{\frac{b}{2}}\omega_t t^{m\sigma}\nabla^m_{\mathcal{H}_+}f(t)\right\|^2_{L^2(\mathbb{R}^3)}.$$

Consequently, the term $(m\sigma)t^{2m\sigma-1} \left\|\omega_t \nabla^m_{\mathcal{H}_+} f(t)\right\|^2_{L^2(\mathbb{R}^3)}$ can be bounded by induction.

Next, we will focus on the estimation of commutators of Landau operators with weight ω_t and $\nabla_{\mathcal{H}_+}^m$, as stated in the following Proposition.

Proposition 2.3. Let $-3 < \gamma < 0$, then for any suitable functions f and $m \in \mathbb{N}_+$, we have the following estimates with weights ω_t for $t \ge 0$

$$\left| \left(\omega_t \nabla_{\mathcal{H}_+}^m \Gamma(f, f), \ \omega_t \nabla_{\mathcal{H}_+}^m f \right)_{L^2(\mathbb{R}^3)} \right| \le C_4 \sum_{p=0}^m \binom{m}{p} \left\| \nabla_{\mathcal{H}_+}^p f \right\|_{2, \frac{\gamma}{2}} \left\| \omega_t \nabla_{\mathcal{H}_+}^{m-p} f \right\|_A \left\| \omega_t \nabla_{\mathcal{H}_+}^m f \right\|_A, \tag{2.7}$$

$$\left| \left(\omega_t \nabla_{\mathcal{H}_+}^m \mathcal{L}_2 f, \ \omega_t \nabla_{\mathcal{H}_+}^m f \right)_{L^2(\mathbb{R}^3)} \right| \le C_6 \sum_{p=0}^m \binom{m}{p} \left(\sqrt{C_5} \right)^{m-p} \sqrt{(m-p+3)!} \left\| \nabla_{\mathcal{H}_+}^p f \right\|_{2,\frac{\gamma}{2}} \left\| \omega_t \nabla_{\mathcal{H}_+}^m f \right\|_A,$$

$$(2.8)$$

$$\left| \left(\left[\omega_t \nabla_{\mathcal{H}_+}^m, \ \mathcal{L}_1 \right] f, \ \omega_t \nabla_{\mathcal{H}_+}^m f \right)_{L^2(\mathbb{R}^3)} \right| \leq \frac{1}{8} \left\| \omega_t \nabla_{\mathcal{H}_+}^m f \right\|_A^2 + C_7 \sum_{p=0}^{m-1} \binom{m}{p} \sqrt{\frac{(m-p+2)!}{2}} \left\| \omega_t \nabla_{\mathcal{H}_+}^p f \right\|_A \left\| \omega_t \nabla_{\mathcal{H}_+}^m f \right\|_A + C_7 \sum_{p=1}^m \binom{m}{p} p \sqrt{\frac{(m-p+2)!}{2}} \left\| \omega_t \nabla_{\mathcal{H}_+}^{p-1} f \right\|_A \left\| \omega_t \nabla_{\mathcal{H}_+}^m f \right\|_A,$$
(2.9)

where $C_4 - C_7$ are positive constants, which depend on γ , b and c_0 .

We will prove this Proposition in the next section.

3. Analysis of Landau operators

In this section, we give exact proofs of Proposition 2.3, which can be used directly in Section 6. In the following, we use the convention of implicit summation over repeated indices. We begin with proving (2.7). Reviewing that

Lemma 3.1. ([16]) For $f, g \in \mathcal{S}(\mathbb{R}^3_v)$, we have

$$\mathcal{L}_{1}f = A_{+,j} \left((a_{jk} * \mu)A_{-,k}f \right), \ \mathcal{L}_{2}f = -A_{+,j} \left(\sqrt{\mu}(a_{jk} * (\sqrt{\mu}A_{-,k}f)) \right),$$

$$\Gamma(f,g) = A_{+,j} \left((a_{jk} * (\sqrt{\mu}f))A_{+,k}g \right) - A_{+,j} \left((a_{jk} * (\sqrt{\mu}A_{+,k}f))g \right).$$

Lemma 3.2. ([16]) For any $m \in \mathbb{N}$, we have

$$\nabla_{\mathcal{H}_{+}}^{m}\left[\left(a_{jk}*\left(\sqrt{\mu}F\right)\right)G\right] = \sum_{p=0}^{m} \binom{m}{p} \left(a_{jk}*\left(\sqrt{\mu}\nabla_{\mathcal{H}_{+}}^{p}F\right)\right) \nabla_{\mathcal{H}_{+}}^{m-p}G.$$

In reviewing the above representations for Landau operators and the Leibniz-type formula, we need the following commutator between the nonlinear Landau operator Γ and weights ω_t .

Lemma 3.3. Let $-3 < \gamma < 0$, then there exists a constant $C_4 > 0$, which depends on γ , b and c_0 , such that for any suitable functions f, g and h

$$\left| \left(\omega_t \Gamma(f,g), \omega_t h \right)_{L^2(\mathbb{R}^3)} \right| \le C_4 \left\| f \right\|_{2,\frac{\gamma}{2}} \left\| \omega_t g \right\|_A \left\| \omega_t h \right\|_A$$

Proof. Since

 $(\omega_t \Gamma(f,g), \omega_t h)_{L^2(\mathbb{R}^3)} = (\omega_t \Gamma(f,g), \omega_t h)_{L^2(\mathbb{R}^3)} - (\Gamma(f,\omega_t g), \omega_t h)_{L^2(\mathbb{R}^3)} + (\Gamma(f,\omega_t g), \omega_t h)_{L^2(\mathbb{R}^3)},$ for the third term, it can be estimated from Proposition 2.2 directly, so we only need to consider estimates of the first two terms. Using integration by parts, we obtain that

$$\begin{split} &(\omega_t \Gamma(f,g),\omega_t h)_{L^2(\mathbb{R}^3)} - (\Gamma(f,\omega_t g),\omega_t h)_{L^2(\mathbb{R}^3)} \\ &= \int_{\mathbb{R}^3} \left(a_{jk} * (\sqrt{\mu}f) \right) \partial_k \omega_t \cdot g \cdot \partial_j (\omega_t h) dv - \int_{\mathbb{R}^3} \left(a_{jk} * (\sqrt{\mu}f) \right) \partial_j \omega_t \cdot \partial_k g \cdot (\omega_t h) dv \\ &+ \int_{\mathbb{R}^3} \left(a_{jk} * \left(\frac{v_j}{2} \sqrt{\mu}f \right) \right) \partial_k \omega_t \cdot g \cdot (\omega_t h) dv + \int_{\mathbb{R}^3} \left(a_{jk} * \left(\frac{v_k}{2} \sqrt{\mu}f \right) \right) \partial_j \omega_t \cdot g \cdot (\omega_t h) dv \\ &+ \int_{\mathbb{R}^3} \left(\partial_k a_{jk} * (\sqrt{\mu}f) \right) \partial_j \omega_t \cdot g \cdot (\omega_t h) dv = \Gamma_1 + \Gamma_2 + \Gamma_3 + \Gamma_4 + \Gamma_5. \end{split}$$

Since $\partial_j \omega_t(v) = \frac{bc_0}{1+t} v_j \langle v \rangle^{b-2} \omega_t(v)$ for $1 \le j \le 3$, then by using

$$\sum_{j} a_{jk} v_{j} = \sum_{k} a_{jk} v_{k} = \sum_{j,k} \partial_{l} a_{jk} v_{j} v_{k} = 0,$$
(3.1)

the following formulas hold:

$$\begin{split} \Gamma_1 &= \frac{bc_0}{1+t} \int_{\mathbb{R}^3} \left(a_{jk} * \left(v_k \sqrt{\mu} f \right) \right) \langle v \rangle^{b-2} \cdot \left(\omega_t g \right) \cdot \partial_j (\omega_t h), \\ \Gamma_2 &= -\frac{bc_0}{1+t} \int_{\mathbb{R}^3} \left(a_{jk} * \left(v_j \sqrt{\mu} f \right) \right) \langle v \rangle^{b-2} \cdot \partial_k (\omega_t g) \cdot (\omega_t h) \\ &+ \left(\frac{bc_0}{1+t} \right)^2 \int_{\mathbb{R}^3} \left(a_{jk} * \left(v_j v_k \sqrt{\mu} f \right) \right) \langle v \rangle^{2(b-2)} \cdot (\omega_t g) \cdot (\omega_t h), \\ \Gamma_3 &= \Gamma_4 = \frac{bc_0}{2(1+t)} \int_{\mathbb{R}^3} \left(a_{jk} * \left(v_j v_k \sqrt{\mu} f \right) \right) \langle v \rangle^{b-2} \cdot (\omega_t g) \cdot (\omega_t h), \\ \Gamma_5 &= \frac{bc_0}{1+t} \int_{\mathbb{R}^3} \left(\partial_k a_{jk} * \left(v_j \sqrt{\mu} f \right) \right) \langle v \rangle^{b-2} \cdot (\omega_t g) \cdot (\omega_t h) \\ &- \frac{bc_0}{1+t} \int_{\mathbb{R}^3} \left(a_{jk} * \left(\delta_{jk} \sqrt{\mu} f \right) \right) \langle v \rangle^{b-2} \cdot (\omega_t g) \cdot (\omega_t h). \end{split}$$

Since $0 < b \leq 2$, these terms can be treated as hard potential cases in the proof of the trilinear estimate to [14], then we have

$$|\Gamma_j| \lesssim \|f\|_{2,\frac{\gamma}{2}} \|\omega_t g\|_A \|\omega_t h\|_A, \quad j = 1, \cdots, 5.$$

Combining Proposition 2.2, there exists a constant $C_4 > 0$, which depends on γ , b and c_0 , such that

$$\left| (\omega_t \Gamma(f,g), \omega_t h)_{L^2(\mathbb{R}^3)} \right| \le C_4 \, \|f\|_{2,\frac{\gamma}{2}} \, \|\omega_t g\|_A \, \|\omega_t h\|_A \, .$$

Proof of (2.7). From the representation of Γ and Lemma 3.2, one has for all $m \in \mathbb{N}$

$$\nabla_{\mathcal{H}_{+}}^{m}\Gamma(f,f) = \sum_{p=0}^{m} \binom{m}{p} \Gamma\left(\nabla_{\mathcal{H}_{+}}^{p}f, \nabla_{\mathcal{H}_{+}}^{m-p}f\right).$$
(3.2)

Then applying Lemma 3.3, we can obtain that

$$\left| \left(\omega_t \nabla_{\mathcal{H}_+}^m \Gamma(f, f), \omega_t \nabla_{\mathcal{H}_+}^m f \right)_{L^2(\mathbb{R}^3)} \right| \leq \sum_{p=0}^m \binom{m}{p} \left| \left(\omega_t \Gamma \left(\nabla_{\mathcal{H}_+}^p f, \nabla_{\mathcal{H}_+}^{m-p} f \right), \omega_t \nabla_{\mathcal{H}_+}^m f \right)_{L^2(\mathbb{R}^3)} \right| \\ \leq C_4 \sum_{p=0}^m \binom{m}{p} \left\| \nabla_{\mathcal{H}_+}^p f \right\|_{2, \frac{\gamma}{2}} \left\| \omega_t \nabla_{\mathcal{H}_+}^{m-p} f \right\|_A \left\| \omega_t \nabla_{\mathcal{H}_+}^m f \right\|_A$$

Proof of (2.8). Since $\mathcal{L}_2 f = -\Gamma(f, \sqrt{\mu})$, then follows immediately from (2.7) that

$$\left| \left(\omega_t \nabla_{\mathcal{H}_+}^m \mathcal{L}_2 f, \omega_t \nabla_{\mathcal{H}_+}^m f \right)_{L^2(\mathbb{R}^3)} \right| \le C_4 \sum_{p=0}^m \binom{m}{p} \left\| \nabla_{\mathcal{H}_+}^p f \right\|_{2,\frac{\gamma}{2}} \left\| \omega_t \nabla_{\mathcal{H}_+}^{m-p} \sqrt{\mu} \right\|_A \left\| \omega_t \nabla_{\mathcal{H}_+}^m f \right\|_A$$

To estimate $\left\|\omega_t \nabla_{\mathcal{H}_+}^{m-p} \sqrt{\mu}\right\|_A^2$, we can write it as

$$\left\|\omega_{t}\nabla_{\mathcal{H}_{+}}^{m-p}\sqrt{\mu}\right\|_{A}^{2} = \sum_{|\alpha|=m-p} \frac{(m-p)!}{\alpha!} \left\|\omega_{t}A_{+}^{\alpha}\sqrt{\mu}\right\|_{A}^{2} = \sum_{|\alpha|=m-p} \frac{(m-p)!}{\alpha!} \left\|\omega_{t}\mu^{-\frac{1}{2}}\partial_{v}^{\alpha}\mu\right\|_{A}^{2}.$$

Since

$$\partial_v^{\alpha}\mu(v) = \prod_{q=1}^3 \sum_{l=0}^{\lceil \alpha_q/2 \rceil} \frac{\alpha_q!}{(\alpha_q - 2l)!l!} (-1)^{\alpha_q - l} \left(\frac{1}{2}\right)^l v_q^{\alpha_q - 2l} \mu(v),$$

this leads to

$$\begin{split} \left\|\omega_t \mu^{-\frac{1}{2}} \partial_v^{\alpha} \mu\right\|_A^2 &\lesssim \left\|\langle\cdot\rangle^{\gamma+2} \omega_t \mu^{-\frac{1}{2}} \partial_v^{\alpha+e_j} \mu\right\|_{L^2(\mathbb{R}^3)}^2 + \left\|\langle\cdot\rangle^{\gamma+4} \omega_t \mu^{-\frac{1}{2}} \partial_v^{\alpha} \mu\right\|_{L^2(\mathbb{R}^3)}^2 \\ &\lesssim 2^{m-p+1} \left[\frac{m-p+1}{2}\right]! \left\|\langle\cdot\rangle^{\gamma+3+m-p} \omega_t \mu^{\frac{1}{2}}\right\|_{L^2(\mathbb{R}^3)}^2 \\ &\quad + 2^{m-p} \left[\frac{m-p}{2}\right]! \left\|\langle\cdot\rangle^{\gamma+4+m-p} \omega_t \mu^{\frac{1}{2}}\right\|_{L^2(\mathbb{R}^3)}^2 \\ &\lesssim \left(\frac{2}{\frac{1}{4}-\frac{c_0}{1+t}}\right)^{m-p} (m-p+2)!, \end{split}$$

here $\frac{c_0}{1+t} \leq c_0 \leq \frac{1}{4}$, and we use the fact that $p!q! \leq (p+q)! \leq 2^{p+q}p!q!$, one has

$$\sum_{|\alpha|=m-p} \frac{(m-p)!}{\alpha!} \le 2^{m-p} \frac{(m-p+1)(m-p+2)}{2},$$

then, we can obtain that

$$\left\|\omega_t \nabla_{\mathcal{H}_+}^{m-p} \sqrt{\mu}\right\|_A^2 \le (C_5)^{m-p} (m-p+3)!.$$

Therefore, there exist positive constants C_5 and C_6 , depend on γ , b and c_0 , such that

$$\left| \left(\omega_t \nabla_{\mathcal{H}_+}^m \mathcal{L}_2 f, \omega_t \nabla_{\mathcal{H}_+}^m f \right)_{L^2(\mathbb{R}^3)} \right| \le C_6 \sum_{p=0}^m \binom{m}{p} \left(\sqrt{C_5} \right)^{m-p} \sqrt{(m-p+3)!} \left\| \nabla_{\mathcal{H}_+}^p f \right\|_{2,\frac{\gamma}{2}} \left\| \omega_t \nabla_{\mathcal{H}_+}^m f \right\|_A$$

Specially, for m = 0, we remark that

$$\left| \left(\omega_t \mathcal{L}_2 f, \omega_t f \right)_{L^2(\mathbb{R}^3)} \right| \le C_6 \left\| f \right\|_{2, \frac{\gamma}{2}} \left\| \omega_t f \right\|_A.$$

$$(3.3)$$

Proof of (2.9). For all $m \in \mathbb{N}_+$, by using (3.1) and Lemma 3.2, then follows from integration by parts that

$$\begin{split} &\left(\left[\omega_{t}\nabla_{\mathcal{H}_{+}}^{m},\ \mathcal{L}_{1}\right]f,\ \omega_{t}\nabla_{\mathcal{H}_{+}}^{m}f\right)_{L^{2}(\mathbb{R}^{3})} \\ &=\sum_{p=0}^{m-1}\binom{m}{p}\int_{\mathbb{R}^{3}}\left(a_{jk}*\left(\sqrt{\mu}\nabla_{\mathcal{H}_{+}}^{m-p}\sqrt{\mu}\right)\right)A_{-,k}\left(\omega_{t}\nabla_{\mathcal{H}_{+}}^{p}f\right)A_{-,j}\left(\omega_{t}\nabla_{\mathcal{H}_{+}}^{m}f\right) \\ &+\frac{bc_{0}}{1+t}\sum_{p=0}^{m}\binom{m}{p}\int_{\mathbb{R}^{3}}\left(a_{jk}*\left(v_{k}\sqrt{\mu}\nabla_{\mathcal{H}_{+}}^{m-p}\sqrt{\mu}\right)\right)\langle v\rangle^{b-2}\omega_{t}\nabla_{\mathcal{H}_{+}}^{p}f\cdot A_{-,j}\left(\omega_{t}\nabla_{\mathcal{H}_{+}}^{m}f\right) \\ &+\frac{bc_{0}}{1+t}\sum_{p=0}^{m}\binom{m}{p}\int_{\mathbb{R}^{3}}\left(a_{jk}*\left(v_{j}\sqrt{\mu}\nabla_{\mathcal{H}_{+}}^{m-p}\sqrt{\mu}\right)\right)\langle v\rangle^{b-2}A_{-,k}\left(\omega_{t}\nabla_{\mathcal{H}_{+}}^{p}f\right)\cdot\omega_{t}\nabla_{\mathcal{H}_{+}}^{m}f \\ &-\left(\frac{bc_{0}}{1+t}\right)^{2}\sum_{p=0}^{m}\binom{m}{p}\int_{\mathbb{R}^{3}}\left(a_{jk}*\left(v_{j}v_{k}\sqrt{\mu}\nabla_{\mathcal{H}_{+}}^{m-p}\sqrt{\mu}\right)\right)\langle v\rangle^{2(b-2)}\omega_{t}\nabla_{\mathcal{H}_{+}}^{p}f\cdot\omega_{t}\nabla_{\mathcal{H}_{+}}^{m}f \\ &-\sum_{p=1}^{m}\binom{m}{p}p\int_{\mathbb{R}^{3}}\left(a_{jk}*\left(\sqrt{\mu}\nabla_{\mathcal{H}_{+}}^{m-p}\sqrt{\mu}\right)\right)\omega_{t}\nabla_{\mathcal{H}_{+}}^{p-1}f\cdot\left(\omega_{t}\nabla_{\mathcal{H}_{+}}^{m}f\right) \\ &-\frac{bc_{0}}{1+t}\sum_{p=1}^{m}\binom{m}{p}p\int_{\mathbb{R}^{3}}\left(a_{jk}*\left(v_{j}\sqrt{\mu}\nabla_{\mathcal{H}_{+}}^{m-p}\sqrt{\mu}\right)\right)\langle v\rangle^{b-2}\omega_{t}\nabla_{\mathcal{H}_{+}}^{p-1}f\cdot\omega_{t}\nabla_{\mathcal{H}_{+}}^{m}f \\ &=I_{1}+I_{2}+I_{3}+I_{4}+I_{5}+I_{6}, \end{split}$$

here we use

$$\nabla_{\mathcal{H}_{+}}^{p} A_{-,k} = A_{-,k} \nabla_{\mathcal{H}_{+}}^{p} - p \nabla_{\mathcal{H}_{+}}^{p-1}, \ p \ge 1 \quad \text{and} \quad [\omega_{t}, A_{\pm,k}] = \pm \partial_{k} \omega_{t}, \ 1 \le k \le 3.$$

For the term I_1 , decomposing $\mathbb{R}^3 \times \mathbb{R}^3$ into

$$\{|v| \le 1\} \cup \{2|v'| \ge |v|, |v| \ge 1\} \cup \{2|v'| \le |v|, |v| \ge 1\} = \Omega_1 \cup \Omega_2 \cup \Omega_3.$$

In $\Omega_1 \cup \Omega_2$, we can deduce that

$$\left|a_{jk} * \left(\sqrt{\mu} \nabla_{\mathcal{H}_{+}}^{m-p} \sqrt{\mu}\right)\right| \lesssim \left\|\nabla_{\mathcal{H}_{+}}^{m-p} \sqrt{\mu}\right\|_{L^{2}(\mathbb{R}^{3})} \langle v \rangle^{\gamma},$$

where we use (2.5), the fact $|v| \leq 1$ in Ω_1 and $\sqrt{\mu}(v') \leq e^{-\frac{|v|^2}{16}}$ in Ω_2 . Then follows immediately from the Cauchy-Schwarz inequality and (2.4) that

$$\left|I_{1}\right|_{\Omega_{1}\cup\Omega_{2}}\right| \lesssim \sum_{p=0}^{m-1} \binom{m}{p} \left\|\nabla_{\mathcal{H}_{+}}^{m-p}\sqrt{\mu}\right\|_{L^{2}(\mathbb{R}^{3})} \left\|\omega_{t}\nabla_{\mathcal{H}_{+}}^{p}f\right\|_{A} \left\|\omega_{t}\nabla_{\mathcal{H}_{+}}^{m}f\right\|_{A}.$$

In Ω_3 , from (3.1) and Taylor's expansion

$$a_{jk}(v-v') = a_{jk}(v) + \sum_{1 \le l \le 3} \partial_l a_{jk}(v)v'_l + \frac{1}{2} \sum_{1 \le l, p \le 3} \int_0^1 \partial_{lp} a_{jk}(v-sv')dsv'_lv'_p,$$

we can get that

$$\begin{split} &\int_{\Omega_3} \left(a_{jk} * \left(\sqrt{\mu} \nabla_{\mathcal{H}_+}^{m-p} \sqrt{\mu} \right) \right) A_{-,k} \left(\omega_t \nabla_{\mathcal{H}_+}^p f \right) A_{-,j} \left(\omega_t \nabla_{\mathcal{H}_+}^m f \right) dv \\ &= \int_{\Omega_3} a_{jk}(v) \sqrt{\mu}(v') \nabla_{\mathcal{H}_+}^{m-p} \sqrt{\mu}(v') (\mathbf{I} - \mathbf{P}_v) A_{-,k} \left(\omega_t \nabla_{\mathcal{H}_+}^p f \right) (v) (\mathbf{I} - \mathbf{P}_v) A_{-,j} \left(\omega_t \nabla_{\mathcal{H}_+}^m f \right) (v) dv \\ &+ \sum_{1 \leq l \leq 3} \int_{\Omega_3} \partial_l a_{jk}(v) v_l' \sqrt{\mu}(v') \nabla_{\mathcal{H}_+}^{m-p} \sqrt{\mu}(v') \Big[\mathbf{P}_v A_{-,k} \left(\omega_t \nabla_{\mathcal{H}_+}^p f \right) (v) (\mathbf{I} - \mathbf{P}_v) A_{-,j} \left(\omega_t \nabla_{\mathcal{H}_+}^m f \right) (v) \\ &+ (\mathbf{I} - \mathbf{P}_v) A_{-,k} \left(\omega_t \nabla_{\mathcal{H}_+}^p f \right) (v) \mathbf{P}_v A_{-,j} \left(\omega_t \nabla_{\mathcal{H}_+}^m f \right) (v) \\ &+ (\mathbf{I} - \mathbf{P}_v) A_{-,k} \left(\omega_t \nabla_{\mathcal{H}_+}^p f \right) (v) (\mathbf{I} - \mathbf{P}_v) A_{-,j} \left(\omega_t \nabla_{\mathcal{H}_+}^m f \right) (v) \Big] dv \\ &+ \frac{1}{2} \sum_{1 \leq l, p \leq 3} \int_{\Omega_3} \int_0^1 \partial_{lp} a_{jk}(v - sv') ds v_l' v_p' \sqrt{\mu}(v') \nabla_{\mathcal{H}_+}^{m-p} \sqrt{\mu}(v') A_{-,k} \left(\omega_t \nabla_{\mathcal{H}_+}^p f \right) (v) A_{-,j} \left(\omega_t \nabla_{\mathcal{H}_+}^m f \right) (v) dv, \end{split}$$

noting that $|\partial_{lp}a_{jk}(v-sv')| \lesssim \langle v \rangle^{\gamma}$ in Ω_3 , then follows immediately from the Cauchy-Schwarz inequality and (2.2), (2.4) that

$$\left|I_{1}\right|_{\Omega_{3}}\right| \lesssim \sum_{p=0}^{m-1} \binom{m}{p} \left\|\nabla_{\mathcal{H}_{+}}^{m-p} \sqrt{\mu}\right\|_{L^{2}(\mathbb{R}^{3})} \left\|\omega_{t} \nabla_{\mathcal{H}_{+}}^{p} f\right\|_{A} \left\|\omega_{t} \nabla_{\mathcal{H}_{+}}^{m} f\right\|_{A}.$$

For the term I_2 , we can write it as

$$I_{2} = \frac{bc_{0}}{1+t} \sum_{p=0}^{m-1} {m \choose p} \int_{\mathbb{R}^{3}} \left(a_{jk} * \left(v_{k} \sqrt{\mu} \nabla_{\mathcal{H}_{+}}^{m-p} \sqrt{\mu} \right) \right) \langle v \rangle^{b-2} \omega_{t} \nabla_{\mathcal{H}_{+}}^{p} f \cdot A_{-,j} \left(\omega_{t} \nabla_{\mathcal{H}_{+}}^{m} f \right) \\ + \frac{bc_{0}}{1+t} \int_{\mathbb{R}^{3}} \left(a_{jk} * \left(v_{k} \mu \right) \right) \langle v \rangle^{b-2} \omega_{t} \nabla_{\mathcal{H}_{+}}^{m} f \cdot A_{-,j} \left(\omega_{t} \nabla_{\mathcal{H}_{+}}^{m} f \right) = I_{2,1} + I_{2,2},$$

Since $0 < b \leq 2$, then as discussion in I_1 , we can deduce that

$$|I_{2,1}| \lesssim \sum_{p=0}^{m-1} \binom{m}{p} \left\| \nabla_{\mathcal{H}_{+}}^{m-p} \sqrt{\mu} \right\|_{L^{2}(\mathbb{R}^{3})} \left\| \omega_{t} \nabla_{\mathcal{H}_{+}}^{p} f \right\|_{A} \left\| \omega_{t} \nabla_{\mathcal{H}_{+}}^{m} f \right\|_{A}.$$

To estimate $I_{2,2}$, using (3.1), we can write it as

$$I_{2,2} = \frac{bc_0}{1+t} \int_{\mathbb{R}^3} \left(a_{jk} * (v_k \mu) \right) \langle v \rangle^{b-2} \omega_t \nabla^m_{\mathcal{H}_+} f \cdot (\mathbf{I} - \mathbf{P}_v) A_{-,j} \left(\omega_t \nabla^m_{\mathcal{H}_+} f \right)$$

since $\gamma + 2 > -1$, from (2.5) we have

$$|a_{jk} * (v_k \mu)| \le \int_{\mathbb{R}^3} |v - v'|^{\gamma+2} \langle v \rangle \mu(v') dv' \le \| \langle \cdot \rangle \sqrt{\mu} \|_{L^{\infty}(\mathbb{R}^3)} \int_{\mathbb{R}^3} |v - v'|^{\gamma+2} e^{-\frac{|v'|^2}{4}} dv' \le C_0 \langle v \rangle^{\gamma+2}.$$

then follows immediately from (2.1), (2.2) and the Cauchy-Schwarz inequality that

$$|I_{2,2}| \leq \frac{bc_0}{1+t} \frac{C_7}{(C_1)^2} \left\| \omega_t \nabla_{\mathcal{H}_+}^m f \right\|_A^2 \leq \frac{1}{32} \left\| \omega_t \nabla_{\mathcal{H}_+}^m f \right\|_A^2,$$

if $\frac{bc_0\tilde{C}_7}{(C_1)^2} \leq \frac{1}{32}$. Similarly, we can deduce that

$$|I_3| \lesssim \sum_{p=0}^{m-1} \binom{m}{p} \left\| \nabla_{\mathcal{H}_+}^{m-p} \sqrt{\mu} \right\|_{L^2(\mathbb{R}^3)} \left\| \omega_t \nabla_{\mathcal{H}_+}^p f \right\|_A \left\| \omega_t \nabla_{\mathcal{H}_+}^m f \right\|_A + \frac{1}{32} \left\| \omega_t \nabla_{\mathcal{H}_+}^m f \right\|_A^2.$$

For term I_4 , we can write it as

$$I_{4} = -\left(\frac{bc_{0}}{1+t}\right)^{2} \sum_{p=0}^{m-1} {m \choose p} \int_{\mathbb{R}^{3}} \left(a_{jk} * \left(v_{j}v_{k}\sqrt{\mu}\nabla_{\mathcal{H}_{+}}^{m-p}\sqrt{\mu}\right)\right) \langle v \rangle^{2(b-2)} \omega_{t}\nabla_{\mathcal{H}_{+}}^{p} f \cdot \omega_{t}\nabla_{\mathcal{H}_{+}}^{m} f$$
$$-\left(\frac{bc_{0}}{1+t}\right)^{2} \int_{\mathbb{R}^{3}} \left(a_{jk} * \left(v_{j}v_{k}\mu\right)\right) \langle v \rangle^{2(b-2)} \omega_{t}\nabla_{\mathcal{H}_{+}}^{m} f \cdot \omega_{t}\nabla_{\mathcal{H}_{+}}^{m} f = I_{4,1} + I_{4,2},$$

by using (2.3) and (2.5), one shows

$$|I_{4,1}| \lesssim \sum_{p=0}^{m-1} \binom{m}{p} \left\| \nabla_{\mathcal{H}_{+}}^{m-p} \sqrt{\mu} \right\|_{L^{2}(\mathbb{R}^{3})} \int_{\mathbb{R}^{3}} \langle v \rangle^{\gamma+2} \left| \omega_{t} \nabla_{\mathcal{H}_{+}}^{p} f \cdot \omega_{t} \nabla_{\mathcal{H}_{+}}^{m} f \right| dv$$
$$\lesssim \sum_{p=0}^{m-1} \binom{m}{p} \left\| \nabla_{\mathcal{H}_{+}}^{m-p} \sqrt{\mu} \right\|_{L^{2}(\mathbb{R}^{3})} \left\| \omega_{t} \nabla_{\mathcal{H}_{+}}^{p} f \right\|_{A} \left\| \omega_{t} \nabla_{\mathcal{H}_{+}}^{m} f \right\|_{A},$$

as discuss in $I_{2,2}$, we have

$$I_{4,2} \leq \frac{1}{16} \left\| \omega_t \nabla^m_{\mathcal{H}_+} f \right\|_A^2.$$

It remains to consider I_5 and I_6 . By using (2.5), one has

$$\left|a_{jk}*\left(\sqrt{\mu}\nabla_{\mathcal{H}_{+}}^{m-p}\sqrt{\mu}\right)\right|+\left|a_{jk}*\left(v_{j}\sqrt{\mu}\nabla_{\mathcal{H}_{+}}^{m-p}\sqrt{\mu}\right)\right|\lesssim\left\|\nabla_{\mathcal{H}_{+}}^{m-p}\sqrt{\mu}\right\|_{L^{2}(\mathbb{R}^{3})}\langle v\rangle^{\gamma+2},$$

then from (2.1) and Cauchy inequality, we have

$$|I_6 + I_7| \lesssim \sum_{p=1}^m \binom{m}{p} p \left\| \nabla_{\mathcal{H}_+}^{m-p} \sqrt{\mu} \right\|_{L^2(\mathbb{R}^3)} \left\| \omega_t \nabla_{\mathcal{H}_+}^{p-1} f \right\|_A \left\| \omega_t \nabla_{\mathcal{H}_+}^m f \right\|_A.$$

To estimate $\|\nabla_{\mathcal{H}_+}^{m-p}\sqrt{\mu}\|_{L^2(\mathbb{R}^3)}$, we have

$$\left\|\nabla_{\mathcal{H}_{+}}^{m-p}\sqrt{\mu}\right\|_{L^{2}(\mathbb{R}^{3})}^{2} = \sum_{|\alpha|=m-p} \frac{(m-p)!}{\alpha!} \left\|A_{+}^{\alpha}\sqrt{\mu}\right\|_{L^{2}(\mathbb{R}^{3})}^{2} = \sum_{|\alpha|=m-p} (m-p)! \left\|\Psi_{\alpha}\right\|_{L^{2}(\mathbb{R}^{3})}^{2} = \frac{(m-p+2)!}{2}$$

where $\{\Psi_{\alpha}\}_{\alpha\in\mathbb{N}^3}$ is the orthonormal basis in $L^2(\mathbb{R}^3)$. Combining the above results, there exists a constant $C_7 > 0$, depends on γ , b and c_0 , such that

$$\begin{split} & \left| \left(\left[\omega_t \nabla_{\mathcal{H}_+}^m, \ \mathcal{L}_1 \right] f, \ \omega_t \nabla_{\mathcal{H}_+}^m f \right)_{L^2(\mathbb{R}^3)} \right| \\ & \leq \frac{1}{8} \left\| \omega_t \nabla_{\mathcal{H}_+}^m f \right\|_A^2 + C_7 \sum_{p=0}^{m-1} \binom{m}{p} \sqrt{\frac{(m-p+2)!}{2}} \left\| \omega_t \nabla_{\mathcal{H}_+}^p f \right\|_A \left\| \omega_t \nabla_{\mathcal{H}_+}^m f \right\|_A \\ & + C_7 \sum_{p=1}^m \binom{m}{p} p \sqrt{\frac{(m-p+2)!}{2}} \left\| \omega_t \nabla_{\mathcal{H}_+}^{p-1} f \right\|_A \left\| \omega_t \nabla_{\mathcal{H}_+}^m f \right\|_A. \end{split}$$

Specially, for m = 0, we remark that

$$\left| ([\omega_t, \mathcal{L}_1]f, \omega_t f)_{L^2(\mathbb{R}^3)} \right| \le \frac{1}{8} \|\omega_t f\|_A^2.$$
 (3.4)

4. EXISTENCE AND ENERGY ESTIMATE

Let c_0 be a positive constant, we will show the existence and L^2 -estimate concerning the following weighted function

$$\omega_t(v) = e^{\frac{c_0}{1+t} \langle v \rangle^b},$$

with $0 < b \leq 2$ and $t \geq 0$.

Proposition 4.1. Let $-3 < \gamma < 0$, assume that there exists $\varepsilon_0 > 0$ such that $e^{c_0 \langle \cdot \rangle^b} f_0 \in L^2(\mathbb{R}^3)$ satisfying

$$\left\| e^{c_0\langle \cdot \rangle^b} f_0 \right\|_{L^2(\mathbb{R}^3)} \le \varepsilon_0, \tag{4.1}$$

then the Cauchy problem (1.4) admits a weak solution $f \in L^{\infty}(]0, \infty[; L^{2}(\mathbb{R}^{3}))$, and for any T > 0

$$\|\omega_t f(t)\|_{L^2(\mathbb{R}^3)}^2 + \frac{2c_0}{(1+T)^2} \int_0^t \left\| \langle \cdot \rangle^{\frac{b}{2}} \omega_\tau f \right\|_{L^2(\mathbb{R}^3)}^2 d\tau + \int_0^t \|\omega_\tau f\|_A^2 d\tau \le C_T \varepsilon_0^2, \quad \forall \ 0 < t \le T.$$
(4.2)

Proof. Let $f^0 = \omega_0 f_0$, consider the following approximate equations

$$\partial_t f^n + \mathcal{L} f^n = \Gamma(f^{n-1}, f^n), \quad f^n|_{t=0} = f_0, \quad n \ge 1.$$

For n = 1, according to the general existence proof we have that the above Cauchy problem has a unique weak solution $f \in L^{\infty}([0, \infty[; L^2(\mathbb{R}^3)))$. With the equation

$$\left(\partial_t + \frac{c_0}{(1+t)^2} \langle v \rangle^b\right) \omega_t f^1 + \omega_t \mathcal{L} f^1 = \omega_t \Gamma(f^0, f^1),$$

taking the inner product with $\omega_t f^1$ on $L^2(\mathbb{R}^2)$, then it follows from Lemma 2.1, Lemma 3.3 and (3.3), (3.4) and (4.1), we have, for small ε_0 ,

$$\frac{d}{dt} \left\| \omega_t f^1 \right\|_{L^2(\mathbb{R}^3)}^2 + \frac{2c_0}{(1+t)^2} \left\| \langle \cdot \rangle^{\frac{b}{2}} \omega_t f^1 \right\|_{L^2(\mathbb{R}^3)}^2 + \|\omega_t f^1\|_A^2 \le C \left\| \omega_t f^1 \right\|_{2,\frac{\gamma}{2}}^2$$

then for all $0 < t \leq T$

$$\left\|\omega_{t}f^{1}(t)\right\|_{L^{2}(\mathbb{R}^{3})}^{2} + \frac{2c_{0}}{(1+T)^{2}}\int_{0}^{t}\left\|\langle\cdot\rangle^{\frac{b}{2}}\omega_{\tau}f^{1}\right\|_{L^{2}(\mathbb{R}^{3})}^{2}d\tau + \int_{0}^{t}\|\omega_{\tau}f\|_{A}^{2}d\tau \leq C(\varepsilon_{0})^{2}.$$
(4.3)

By iterating and (4.3), one can obtain that for all $n \ge 1$, the approximate equations admit a solution f^n satisfying

$$\|\omega_t f^n(t)\|_{L^2(\mathbb{R}^3)}^2 + \frac{2c_0}{(1+T)^2} \int_0^t \left\| \langle \cdot \rangle^{\frac{b}{2}} \omega_\tau f^n \right\|_{L^2(\mathbb{R}^3)}^2 d\tau + \int_0^t \|\omega_\tau f^n\|_A^2 d\tau \le C(\varepsilon_0)^2.$$

Setting $\zeta^n = f^{n+1} - f^n$ for $n \ge 1$, then

$$\partial_t \zeta^n + \mathcal{L} \zeta^n = \Gamma(f^n, \zeta^n) + \Gamma(\zeta^{n-1}, f^n), \quad \zeta^n|_{t=0} = 0.$$

Taking inner product with $\omega_t \zeta^n$ on both side of

$$\left(\partial_t + \frac{c_0}{(1+t)^2} \langle v \rangle^b\right) \omega_t \zeta^n + \omega_t \mathcal{L} \zeta^n = \omega_t \Gamma(f^n, \zeta^n) + \omega_t \Gamma(\zeta^{n-1}, f^n),$$
¹³

for any T > 0, by using Lemma 2.1, Lemma 3.3, (3.3) and (3.4), we can obtain that, for all $0 < t \leq T$

$$\|\omega_t \zeta^n(t)\|_{L^2(\mathbb{R}^3)}^2 + \frac{2c_0}{(1+T)^2} \int_0^t \left\| \langle \cdot \rangle^{\frac{b}{2}} \omega_\tau \zeta^n \right\|_{L^2(\mathbb{R}^3)}^2 d\tau + \int_0^t \|\omega_\tau \zeta^n\|_A^2 d\tau \le C \,\varepsilon_0^2 \left\|\omega_t \zeta^{n-1}\right\|_{L^\infty([0,T];L^2(\mathbb{R}^3))}^2,$$

this yields, for small ε_0 ,

$$\|\omega_t \zeta^n\|_{L^{\infty}([0,\infty[;L^2(\mathbb{R}^3)))}^2 \leq \frac{1}{2} \|\omega_t \zeta^{n-1}\|_{L^{\infty}([0,\infty[;L^2(\mathbb{R}^3)))}^2 \leq 2^{-n} \|\omega_t \zeta^0\|_{L^{\infty}([0,\infty[;L^2(\mathbb{R}^3)))}^2$$

Thus, $\{f^n\} \subset L^2(\mathbb{R}^3)$ is a Cauchy sequence, the limit function $f \in L^\infty([0,\infty[;L^2(\mathbb{R}^3)))$ is a unique solution to Cauchy problem (1.4), and satisfying (4.2).

With the same argument, we can also obtain the following global existence of weak solutions.

Corollary 4.2. Let $-3 < \gamma < 0$, and $f_0 \in L^2(\mathbb{R}^3)$ satisfying

$$\|f_0\|_{L^2(\mathbb{R}^3)} \le \varepsilon_0,\tag{4.4}$$

then the Cauchy problem (1.4) admits a global in time weak solution satisfying, for any T > 0, there exists $C_T > 0$ such that

$$\|f(t)\|_{L^{2}(\mathbb{R}^{3})}^{2} + \int_{0}^{t} \|f\|_{A}^{2} ds \leq C_{T} \varepsilon_{1}^{2}, \quad \forall \ 0 < t \leq T.$$

$$(4.5)$$

5. Analyticity in time variable

This section is devoted to proving time analyticity, for which the weight ω is not required. With the Cauchy problem (1.4), we can establish the following L^2 -esitmate.

Lemma 5.1. For $-3 < \gamma < 0$. Lef f be the solution of Cauchy problem (1.4) with $||f||_{L^{\infty}([0,\infty[:L^2)]}$ small enough. Then there exists a constant $C_8 > 0$ such that for any T > 0 and $t \in [0, T]$,

$$\|t\partial_t f\|_{L^2(\mathbb{R}^3)}^2 + \int_0^t \|s\partial_s f(s)\|_A^2 ds \le (C_8\epsilon)^2.$$
(5.1)

Proof. With the Cauchy problem (1.4), we have that for all $0 < t \le T$,

- *t*

$$\begin{aligned} &\frac{1}{2} \| t\partial_t f \|_{L^2(\mathbb{R}^3)}^2 + \int_0^t (\mathcal{L}_1(s\partial_s f), s\partial_s f)_{L^2(\mathbb{R}^3)} ds \\ &= \int_0^t s \| \partial_s f \|_{L^2(\mathbb{R}^3)}^2 ds - \int_0^t (\mathcal{L}_2(s\partial_s f), s\partial_s f)_{L^2(\mathbb{R}^3)} + \int_0^t (s\partial_s \Gamma(f, f), s\partial_s f)_{L^2(\mathbb{R}^3)} ds \\ &= D_1 + D_2 + D_3. \end{aligned}$$

It follows from (1.4) that

$$D_1 = \int_0^t (\Gamma(f, f), s\partial_s f)_{L^2(\mathbb{R}^3)} ds - \int_0^t (\mathcal{L}_1 f, s\partial_s f)_{L^2(\mathbb{R}^3)} ds - \int_0^t (\mathcal{L}_2 f, s\partial_s f)_{L^2(\mathbb{R}^3)} ds,$$

since $\mathcal{L}_1 f = -\Gamma(\sqrt{\mu}, f)$ and $\mathcal{L}_2 f = -\Gamma(f, \sqrt{\mu})$, then follows immediately from Lemma 2.2 and (2.3) that for all $0 < t \leq T$

$$D_{1} \leq C_{2}\varepsilon_{0} \int_{0}^{t} \|f(s)\|_{A} \|s\partial_{s}f\|_{A} ds + \tilde{C}_{2} \int_{0}^{t} \|f(s)\|_{A} \|s\partial_{s}f\|_{A} ds + \frac{\tilde{C}_{2}}{\sqrt{C_{1}}} \int_{0}^{t} \|f(s)\|_{A} \|s\partial_{s}f\|_{A} ds,$$

then from the Cauchy-Schwarz inequality, for any $0 < \delta < 1$, we have

$$D_{1} = \int_{0}^{t} s \|\partial_{s}f\|_{L^{2}(\mathbb{R}^{3})}^{2} ds \leq \delta \int_{0}^{t} \|s\partial_{s}f\|_{A}^{2} ds + C_{\delta} \int_{0}^{t} \|f(s)\|_{A}^{2} ds,$$
(5.2)

with C_{δ} depends on C_1, C_2 .

For D_2 , since $\mathcal{L}_2 f = -\Gamma(f, \sqrt{\mu})$, then by using Lemma 2.2 and (2.3), we have for all $0 < t \leq T$

$$|D_2| \le \epsilon_2 \int_0^t \|s\partial_s f\|_A^2 ds + C_{\epsilon_2} T \int_0^t s \|\partial_s f\|_{L^2(\mathbb{R}^3)}^2 ds,$$

by using (5.2) with $C_{\epsilon_2}T\delta \leq \epsilon_2$,

$$|D_2| \le 2\epsilon_2 \int_0^t \|s\partial_s f\|_A^2 ds + \tilde{C}_{\epsilon_2} \int_0^t \|f(s)\|_A^2 ds,$$

with \tilde{C}_{ϵ_2} depends on C_1, C_2 and T. Now, we trun to the term S_3 , since $\gamma < 0$, by using Lemma 2.2, (4.5) and Cauchy-Schwarz inequality, one has

$$|D_{3}| \leq \int_{0}^{t} s^{2} |(\Gamma(\partial_{s}f, f), \partial_{s}f)| ds + \int_{0}^{t} s^{2} |(\Gamma(f, \partial_{s}f), \partial_{s}f)| ds$$

$$\leq C_{2} \int_{0}^{t} ||s\partial_{s}f||_{L^{2}} ||f(s)||_{A} ||s\partial_{s}f||_{A} + ||f(s)||_{L^{2}} ||s\partial_{s}f||_{A}^{2} ds$$

$$\leq C_{2} \varepsilon_{1} ||s\partial_{s}f||_{L^{\infty}(]0,t];L^{2})}^{2} + 2C_{2} \varepsilon_{1} \int_{0}^{t} ||s\partial_{s}f||_{A}^{2} ds.$$

It remains to consider the operator \mathcal{L}_1 . Since $\gamma < 0$, by lemma 2.1, for all $0 < t \leq T$, we can conclude

$$\int_0^t (\mathcal{L}_1(s\partial_s f), s\partial_s f)_{L^2(\mathbb{R}^3)} ds \ge (1-\epsilon_1) \int_0^t \|s\partial_s f\|_A^2 ds - C_{\epsilon_1} \int_0^t \|s\partial_s f\|_{2,\frac{\gamma}{2}}^2 ds$$
$$\ge (1-\epsilon_1) \int_0^t \|s\partial_s f\|_A^2 ds - TC_{\epsilon_1} \int_0^t s\|\partial_s f\|_{L^2(\mathbb{R}^3)}^2 ds,$$

applying (5.2) with $TC_{\epsilon_1}\delta \leq \epsilon_1$, one has

$$\int_0^t (\mathcal{L}_1(s\partial_s f), s\partial_s f)_{L^2(\mathbb{R}^3)} ds \ge (1 - 2\epsilon_1) \int_0^t \|s\partial_s f\|_A^2 ds - \tilde{C}_{\epsilon_1} \int_0^t \|f(s)\|_A^2 ds$$

Combining these results and taking

$$2\epsilon_1 = \delta = 2\epsilon_2 = 2C_2\varepsilon_1 = \frac{1}{8},$$

then we can get that for all $0 < t \le T$

$$\|t\partial_t f\|_{L^2}^2 + \int_0^t \|s\partial_s f\|_A^2 ds \le 2\tilde{C}_8 \int_0^t \|f(s)\|_A^2 ds + 2(C_2\varepsilon_1)^2 \|s\partial_s f\|_{L^\infty([0,t];L^2)}^2,$$

here \tilde{C}_8 depends on C_1, C_2 and T. Now, we choose $2(C_2\varepsilon_1)^2 \leq \frac{1}{2}$, then

$$\frac{1}{2} \|t\partial_t f\|_{L^{\infty}(]0,T];L^2)}^2 \le 2\tilde{C}_8(\varepsilon_1)^2,$$

from this, one can deduce that for all $0 < t \leq T$

$$\|t\partial_t f\|_{L^2(\mathbb{R}^3)}^2 + \int_0^t \|s\partial_s f\|_A^2 ds \le 4\tilde{C}_8(\varepsilon_1)^2 \le (C_8\epsilon)^2,$$

here $C_8 \ge 2\sqrt{\tilde{C}_8}$ depends on C_1, C_2 and T.

Now, we turn to show the time analyticity.

Proposition 5.2. For $-3 < \gamma < 0$. Let f be the solution of Cauchy problem (1.4) with $||f||_{L^{\infty}([0,\infty[;L^2)})$ small enough. Then there exists a constant B > 0 such that for any T > 0, $t \in [0,T]$ and $k \in \mathbb{N}$,

$$\|t^k \partial_t^k f(t)\|_{L^2(\mathbb{R}^3)}^2 + \int_0^t \|s^k \partial_s^k f(s)\|_A^2 ds \le \left(B^{k-2}(k-1)!\right)^2.$$
(5.3)

Proof. We prove this proposition by induction on the index k. For k = 0, 1, it is enough to take in (4.5) and (5.1). Assume (5.3) holds true, for any $1 \le m \le k - 1$ with $k \ge 2$,

$$\|t^{m}\partial_{t}^{m}f(t)\|_{L^{2}(\mathbb{R}^{3})}^{2} + \int_{0}^{t} \|s^{m}\partial_{s}^{m}f(s)\|_{A}^{2}ds \leq \left(B^{m-1}m!\right)^{2}.$$
(5.4)

We shall prove (5.4) holds for m = k. Since μ is the function with respect to v, which implies

$$t^k \partial_t^k \mathcal{L}_1 f = \mathcal{L}_1(t^k \partial_t^k f), \quad t^k \partial_t^k \mathcal{L}_2 f = \mathcal{L}_2(t^k \partial_t^k f).$$

Then by (1.4), we have

$$\begin{split} \partial_t (t^k \partial_t^k f) &+ \mathcal{L}_1 (t^k \partial_t^k f) = k t^{k-1} \partial_t^k f - \mathcal{L}_2 (t^k \partial_t^k f) + \Gamma (f, t^k \partial_t^k f) \\ &+ \Gamma (t^k \partial_t^k f, f) + \sum_{1 \leq j \leq k-1} C_k^j \ \Gamma (t^j \partial_t^j f, t^{k-j} \partial_t^{k-j} f), \end{split}$$

where $C_k^j = \frac{k!}{j!(k-j)!}$. Then taking the $L^2(\mathbb{R}^3)$ inner product of both sides with respect to $t^k \partial_t^k f$, we have

$$\begin{aligned} \frac{1}{2} \frac{u}{dt} \| t^k \partial_t^k f \|_{L^2}^2 + (\mathcal{L}_1(t^k \partial_t^k f), t^k \partial_t^k f)_{L^2} &= k t^{2k-1} \| \partial_t^k f \|_{L^2}^2 - (\mathcal{L}_2(t^k \partial_t^k f), t^k \partial_t^k f)_{L^2} \\ &+ (\Gamma(f, t^k \partial_t^k f), t^k \partial_t^k f)_{L^2} + (\Gamma(t^k \partial_t^k f, f), t^k \partial_t^k f)_{L^2} \\ &+ \sum_{1 \le j \le k-1} C_k^j \, \Gamma(t^j \partial_t^j f, t^{k-j} \partial_t^{k-j} f), t^k \partial_t^k f)_{L^2}. \end{aligned}$$

For all $0 < t \le T$, integrating from 0 to t, since $\gamma < 0$, by using lemma 2.1, it follows that

$$\int_{0}^{t} (\mathcal{L}_{1}(s^{k}\partial_{s}^{k}f), s^{k}\partial_{s}^{k}f)_{L^{2}(\mathbb{R}^{3})}ds \geq (1-\epsilon_{1})\int_{0}^{t} \|s^{k}\partial_{s}^{k}f\|_{A}^{2}ds - TC_{\epsilon_{1}}\int_{0}^{t} s^{2k-1}\|\partial_{s}^{k}f\|_{L^{2}(\mathbb{R}^{3})}^{2}ds,$$

and since $\mathcal{L}_2 = -\Gamma(f, \sqrt{\mu})$, from Lemma 2.2, we have

$$\int_{0}^{t} |(\mathcal{L}_{2}(s^{k}\partial_{s}^{k}f), s^{k}\partial_{s}^{k}f)_{L^{2}}|ds \leq \epsilon_{2}\int_{0}^{t} \|s^{k}\partial_{s}^{k}f\|_{A}^{2}ds + TC_{\epsilon_{2}}\int_{0}^{t} s^{2k-1}\|\partial_{s}^{k}f\|_{L^{2}}^{2}ds.$$

Combining the results above and taking

$$\epsilon_1 = \epsilon_2 = \frac{1}{8},$$

we have for all $0 < t \leq T$,

$$\begin{aligned} |t^{k}\partial_{t}^{k}f(t)||_{L^{2}(\mathbb{R}^{3})}^{2} &+ \frac{3}{2}\int_{0}^{t} \|s^{k}\partial_{s}^{k}f\|_{A}^{2}ds \\ &\leq C_{9}\int_{0}^{t}s^{2k-1}\|\partial_{s}^{k}f\|_{L^{2}(\mathbb{R}^{3})}^{2}ds + 2\sum_{1\leq j\leq k-1\atop 16}C_{k}^{j}\int_{0}^{t}\Gamma(s^{j}\partial_{s}^{j}f,s^{k-j}\partial_{s}^{k-j}f),s^{k}\partial_{s}^{k}f)_{L^{2}}ds \end{aligned}$$

$$+ 2 \int_0^t (\Gamma(f, s^k \partial_s^k f), s^k \partial_s^k f)_{L^2} ds + 2 \int_0^t (\Gamma(s^k \partial_s^k f, f), s^k \partial_s^k f)_{L^2} ds \\= J_1 + J_2 + J_3 + J_4,$$

with C_9 depends on C_1, C_2 and T. For term J_2 , by using Lemma 2.2 and the fact $\gamma < 0$, one has

$$J_2 \le 2C_2 \sum_{1 \le j \le k-1} C_k^j \int_0^t \|s^j \partial_s^j f\|_{L^2} \|s^{k-j} \partial_s^{k-j} f\|_A \|s^k \partial_s^k f\|_A ds,$$

then by using Cauchy-Schwarz inequality and (5.4), it follows that

$$J_{2} \leq 2C_{2}B^{k-2} \sum_{1 \leq j \leq k-1} C_{k}^{j}(j-2)!(k-j-2)! \left(\int_{0}^{t} \|s^{k}\partial_{s}^{k}f\|_{A}^{2}ds\right)^{\frac{1}{2}}$$

$$\leq \frac{1}{8} \int_{0}^{t} \|s^{k}\partial_{s}^{k}f\|_{A}^{2}ds + 2 \left(2C_{2}B^{k-2} \sum_{1 \leq j \leq k-1} C_{k}^{j}(j-2)!(k-j-2)!\right)^{2}$$

$$\leq \frac{1}{8} \int_{0}^{t} \|s^{k}\partial_{s}^{k}f\|_{A}^{2}ds + 2 \left(64C_{2}B^{k-2}(k-2)!\right)^{2},$$

here we use

$$\sum_{\leq j \leq k-1} C_k^j (j-2)! (k-j-2)! \leq 32(k-2)!$$

 $1 \le j \le k-1$ For the term J_3 and J_4 , by using (2.2) and the fact $\gamma < 0$, one has

$$J_{3} + J_{4} \leq 2C_{2} \int_{0}^{t} \left(\|f(s)\|_{L^{2}} \|s^{k} \partial_{s}^{k} f\|_{A}^{2} + \|f(s)\|_{A} \|s^{k} \partial_{s}^{k} f\|_{L^{2}} \|s^{k} \partial_{s}^{k} f\|_{A} \right) ds$$

$$\leq 2C_{2} \varepsilon_{1} \int_{0}^{t} \|s^{k} \partial_{s}^{k} f\|_{A}^{2} ds + 2C_{2} \varepsilon_{1} \|s^{k} \partial_{s}^{k} f\|_{L^{\infty}(]0,t];L^{2})} \left(\int_{0}^{t} \|s^{k} \partial_{s}^{k} f\|_{A}^{2} ds \right)^{\frac{1}{2}}$$

$$\leq 4C_{2} \varepsilon_{1} \int_{0}^{t} \|s^{k} \partial_{s}^{k} f\|_{A}^{2} ds + \frac{1}{2}C_{2} \varepsilon_{1} \|s^{k} \partial_{s}^{k} f\|_{L^{\infty}(]0,t];L^{2})}.$$

Then taking $4C_2\varepsilon_1 \leq \frac{1}{8}$, we have

$$J_3 + J_4 \le \frac{1}{8} \int_0^t \|s^k \partial_s^k f\|_A^2 ds + \frac{1}{16} \|s^k \partial_s^k f\|_{L^{\infty}(]0,t];L^2)}^2$$

It remains to consider the term J_1 , since f is the solution of (1.4) and $k \ge 2$, then

$$\partial_t^k f = \partial_t^{k-1} \Gamma(f, f) - \mathcal{L}_1(\partial_t^{k-1} f) - \mathcal{L}_2(\partial_t^{k-1} f),$$

which implies

$$J_{1} = C_{9} \int_{0}^{t} (\Gamma(f, s^{k-1}\partial_{s}^{k-1}f), s^{k}\partial_{s}^{k}f)_{L^{2}} + (\Gamma(s^{k-1}\partial_{s}^{k-1}f, f), s^{k}\partial_{s}^{k}f)_{L^{2}}ds + C_{9} \sum_{1 \le j \le k-2} C_{k-1}^{j} \int_{0}^{t} (\Gamma(s^{j}\partial_{s}^{j}f, s^{k-1-j}\partial_{s}^{k-1-j}f), s^{k}\partial_{s}^{k}f)_{L^{2}}ds - C_{9} \int_{0}^{t} (\mathcal{L}_{1}(s^{k-1}\partial_{s}^{k-1}f), s^{k}\partial_{s}^{k}f)_{L^{2}(\mathbb{R}^{3})}ds - \int_{0}^{t} (\mathcal{L}_{2}(s^{k-1}\partial_{s}^{k-1}f), s^{k}\partial_{s}^{k}f)_{L^{2}(\mathbb{R}^{3})}ds 17$$

$$= J_{1,1} + J_{1,2} + J_{1,3} + J_{1,4}.$$

For $J_{1,1}$ and $J_{1,2}$, by using Lemma 2.2, (5.4), and the fact $\gamma < 0$, we have

$$|J_{1,1} + J_{1,2}| \le 2C_2C_9\varepsilon_0B^{k-2}(k-3)! \left(\int_0^t \|s^k\partial_s^k f\|_A^2 ds\right)^{\frac{1}{2}} + C_2C_9\sum_{1\le j\le k-2}C_{k-1}^jB^{k-3}(j-2)!(k-j-3)! \left(\int_0^t \|s^k\partial_s^k f\|_A^2 ds\right)^{\frac{1}{2}} \le 2\left(64C_2C_9B^{k-3}(k-2)!\right)^2 + \left((\varepsilon_1)^2 + \frac{1}{16}\right)\int_0^t \|s^k\partial_s^k f\|_A^2 ds$$

From Lemma 2.2, then applying (2.1) and (5.4), it follows that

$$\begin{aligned} |J_{1,3} + J_{1,4}| &\leq C_9 \left(\tilde{C}_2 + \frac{\tilde{C}_2}{\sqrt{C_1}} \right) B^{k-2} (k-3)! \left(\int_0^t \|s^k \partial_s^k f\|_A^2 ds \right)^{\frac{1}{2}} \\ &\leq \frac{1}{8} \int_0^t \|s^k \partial_s^k f\|_A^2 ds + 2 \left(C_9 \left(\tilde{C}_2 + \frac{\tilde{C}_2}{\sqrt{C_1}} \right) B^{k-2} (k-3)! \right)^2. \end{aligned}$$

So that, taking $(\varepsilon_1)^2 \leq \frac{1}{16}$, we can get that

$$J_1 \le \frac{1}{4} \int_0^t \|s^k \partial_s^k f\|_A^2 ds + 2\left(C_9\left(\tilde{C}_2 + \frac{\tilde{C}_2}{\sqrt{C_1}}\right)B^{k-2}(k-3)!\right)^2 + 2\left(64C_2C_9B^{k-3}(k-2)!\right)^2$$

Combining the results of $J_1 - J_4$, we have for all $0 < t \le T$,

$$\|t^k \partial_t^k f(t)\|_{L^2(\mathbb{R}^3)}^2 + \int_0^t \|s^k \partial_s^k f\|_A^2 ds \le \tilde{C}_9 \left(B^{k-2}(k-2)!\right)^2 + \frac{1}{2} \|t^k \partial_t^k f\|_{L^\infty(]0,T];L^2)}^2,$$

from this, we can get that for all T > 0

$$\frac{1}{2} \| t^k \partial_t^k f \|_{L^{\infty}(]0,T];L^2)}^2 \le \tilde{C}_9 \left(B^{k-2} (k-2)! \right)^2,$$

and therefore, we have for all $0 < t \leq T$,

$$\|t^k \partial_t^k f(t)\|_{L^2(\mathbb{R}^3)}^2 + \int_0^t \|s^k \partial_s^k f\|_A^2 ds \le 2\tilde{C}_9 \left(B^{k-2}(k-2)!\right)^2.$$

Finally, taking $B \ge \sqrt{2\tilde{C}_9}$, we can get that for all $0 < t \le T$,

$$\|t^k \partial_t^k f(t)\|_{L^2(\mathbb{R}^3)}^2 + \int_0^t \|s^k \partial_s^k f(s)\|_A^2 ds \le \left(B^{k-2}(k-1)!\right)^2.$$

6. Gelfand-Shilov Regularizing Effect

In this section, we will show the Gelfand-Shilov regularizing effect with the sub-exponential weighted functions.

Proposition 6.1. Assume that $-3 < \gamma < 0$. Let f be the solution of Cauchy problem (1.4) satisfying $\|\omega_t f\|_{L^{\infty}([0,\infty[;L^2(\mathbb{R}^3)))}$ small enough. Then there exists a positive constant L > 1, depends on γ , b, c_0 and T, such that

$$\begin{aligned} \left\| \omega_t t^{m\sigma} \nabla_{\mathcal{H}_+}^m f(t) \right\|_{L^2(\mathbb{R}^3)}^2 &+ \frac{c_0}{(1+T)^2} \int_0^t \left\| \omega_\tau \tau^{m\sigma} \langle \cdot \rangle^{\frac{b}{2}} \nabla_{\mathcal{H}_+}^m f(\tau) \right\|_{L^2(\mathbb{R}^3)}^2 d\tau \\ &+ \int_0^t \left\| \omega_\tau \tau^{m\sigma} \nabla_{\mathcal{H}_+}^m f(\tau) \right\|_A^2 d\tau \le \left(L^{m-1} (m-2)! \right)^{2\sigma}, \ \forall \ m \in \mathbb{N}, \ \forall \ t \in]0, T]. \end{aligned}$$

$$\tag{6.1}$$

Proof. We prove this proposition by induction on the index m. For m = 0, it has been proved in (4.2). From (1.4) and Lemma 2.1, we have

$$\frac{1}{2}\frac{d}{dt}\left\|\omega_{t}t^{m\sigma}\nabla_{\mathcal{H}_{+}}^{m}f(t)\right\|_{L^{2}(\mathbb{R}^{3})}^{2}+\frac{c_{0}}{(1+t)^{2}}\left\|\langle\cdot\rangle^{\frac{b}{2}}\omega_{t}t^{m\sigma}\nabla_{\mathcal{H}_{+}}^{m}f(t)\right\|_{L^{2}(\mathbb{R}^{3})}^{2}+\frac{7}{8}\left\|\omega_{t}t^{m\sigma}\nabla_{\mathcal{H}_{+}}^{m}f(t)\right\|_{A}^{2} \leq m\sigma t^{2m\sigma-1}\left\|\omega_{t}\nabla_{\mathcal{H}_{+}}^{m}f(t)\right\|_{L^{2}(\mathbb{R}^{3})}^{2}+\left|\left(\omega_{t}t^{m\sigma}\nabla_{\mathcal{H}_{+}}^{m}\Gamma(f,f),\omega_{t}t^{m\sigma}\nabla_{\mathcal{H}_{+}}^{m}f\right)_{L^{2}(\mathbb{R}^{3})}\right| \\ +\left|\left(\omega_{t}t^{m\sigma}\nabla_{\mathcal{H}_{+}}^{m}\mathcal{L}_{2}f,\omega_{t}t^{m\sigma}\nabla_{\mathcal{H}_{+}}^{m}f\right)_{L^{2}(\mathbb{R}^{3})}\right|+\left|\left(t^{m\sigma}[\omega_{t}\nabla_{\mathcal{H}_{+}}^{m},\mathcal{L}_{1}]f,\omega_{t}t^{m\sigma}\nabla_{\mathcal{H}_{+}}^{m}f\right)_{L^{2}(\mathbb{R}^{3})}\right|.$$

For all $t \in [0, T]$, integrating from 0 to t,

$$\begin{aligned} \left\| \omega_t t^{m\sigma} \nabla^m_{\mathcal{H}_+} f(t) \right\|_{L^2(\mathbb{R}^3)}^2 + \frac{2c_0}{(1+T)^2} \int_0^t \left\| \langle \cdot \rangle^{\frac{b}{2}} \omega_\tau \tau^{m\sigma} \nabla^m_{\mathcal{H}_+} f(\tau) \right\|_{L^2(\mathbb{R}^3)}^2 d\tau + \frac{3}{2} \int_0^t \left\| \omega_\tau \tau^{m\sigma} \nabla^m_{\mathcal{H}_+} f(\tau) \right\|_A^2 d\tau \\ &\leq 2m\sigma \int_0^t \tau^{2m\sigma-1} \left\| \omega_\tau \nabla^m_{\mathcal{H}_+} f(\tau) \right\|_{L^2(\mathbb{R}^3)}^2 d\tau + 2C_3 \int_0^t \left\| \omega_\tau \tau^{m\sigma} \nabla^m_{\mathcal{H}_+} f(\tau) \right\|_{2,\frac{\gamma}{2}}^2 d\tau + Q_1 + Q_2 + Q_3, \end{aligned}$$
with

W

$$Q_{1} = 2C_{7} \sum_{p=0}^{m-1} {m \choose p} \sqrt{\frac{(m-p+2)!}{2}} \int_{0}^{t} \tau^{(m-p)\sigma} \left\| \omega_{\tau} \tau^{p\sigma} \nabla_{\mathcal{H}_{+}}^{p} f(\tau) \right\|_{A} \left\| \omega_{\tau} \tau^{m\sigma} \nabla_{\mathcal{H}_{+}}^{m} f(\tau) \right\|_{A} d\tau$$
$$+ 2C_{7} \sum_{p=1}^{m} {m \choose p} p \sqrt{\frac{(m-p+2)!}{2}} \int_{0}^{t} \tau^{(m-p+1)\sigma} \left\| \omega_{\tau} \tau^{(p-1)\sigma} \nabla_{\mathcal{H}_{+}}^{p-1} f(\tau) \right\|_{A} \left\| \omega_{\tau} \tau^{m\sigma} \nabla_{\mathcal{H}_{+}}^{m} f(\tau) \right\|_{A} d\tau,$$

and

$$Q_2 = 2 \int_0^t \left| \left(\omega_\tau \tau^{m\sigma} \nabla^m_{\mathcal{H}_+} \mathcal{L}_2 f(\tau), \omega_\tau \tau^{m\sigma} \nabla^m_{\mathcal{H}_+} f(\tau) \right)_{L^2(\mathbb{R}^3)} \right| d\tau,$$

and

$$Q_3 = 2 \int_0^t \left| \left(\omega_\tau \tau^{m\sigma} \nabla^m_{\mathcal{H}_+} \Gamma(f, f), \omega_\tau \tau^{m\sigma} \nabla^m_{\mathcal{H}_+} f \right)_{L^2(\mathbb{R}^3)} \right| d\tau.$$

For m = 1, it follows immediately from Proposition 2.3, (4.2) that

$$\begin{split} \left\| \omega_{t} t^{\sigma} \nabla_{\mathcal{H}_{+}} f(t) \right\|_{L^{2}(\mathbb{R}^{3})}^{2} + \frac{2c_{0}}{(1+T)^{2}} \int_{0}^{t} \left\| \langle \cdot \rangle^{\frac{b}{2}} \omega_{\tau} \tau^{\sigma} \nabla_{\mathcal{H}_{+}} f(\tau) \right\|_{L^{2}(\mathbb{R}^{3})}^{2} d\tau + \int_{0}^{t} \left\| \omega_{\tau} \tau^{\sigma} \nabla_{\mathcal{H}_{+}} f(\tau) \right\|_{A}^{2} d\tau \\ &\leq 2\sigma \int_{0}^{t} \tau^{2\sigma-1} \left\| \omega_{\tau} \nabla_{\mathcal{H}_{+}} f(\tau) \right\|_{L^{2}(\mathbb{R}^{3})}^{2} d\tau + \left(2C_{3} + (4C_{6})^{2} \right) \int_{0}^{t} \left\| \omega_{\tau} \tau^{\sigma} \nabla_{\mathcal{H}_{+}} f(\tau) \right\|_{L^{2}(\mathbb{R}^{3})}^{2} d\tau \\ &+ \frac{1}{2} \left\| \omega_{t} t^{\sigma} \nabla_{\mathcal{H}_{+}} f \right\|_{L^{\infty}([0,c_{0}/2];L^{2}(\mathbb{R}^{3}))}^{2} + \left((4C_{6})^{2}C_{5} + (4C_{7})^{2} \right) (T+1)^{2\sigma} (\varepsilon_{1})^{2}, \end{split}$$

if we choose $C_4\varepsilon_1 \leq \frac{1}{4}$. For the first integral above, note that $0 < \frac{-\gamma}{b-\gamma}, \frac{b}{b-\gamma} < 1$, by applying Hölder's inequality, one has

$$\left\|\omega_t \nabla_{\mathcal{H}_+} f(\tau)\right\|_{L^2(\mathbb{R}^3)}^2 \le \left\|\langle\cdot\rangle^{\frac{b}{2}} \omega_t \nabla_{\mathcal{H}_+} f\right\|_{L^2(\mathbb{R}^3)}^{\frac{-2\gamma}{b-\gamma}} \left\|\langle\cdot\rangle^{\frac{\gamma}{2}} \omega_t \nabla_{\mathcal{H}_+} f\right\|_{L^2(\mathbb{R}^3)}^{\frac{2b}{b-\gamma}}$$

it follows from (2.3) and (2.4) that

$$\begin{split} \left\| \langle \cdot \rangle^{\frac{\gamma}{2}} \omega_t \nabla_{\mathcal{H}_+} f \right\|_{L^2(\mathbb{R}^3)} &\leq \left\| \langle \cdot \rangle^{\frac{\gamma}{2}} \nabla_{\mathcal{H}_+} \left(\omega_t f \right) \right\|_{L^2(\mathbb{R}^3)} + \left\| \langle \cdot \rangle^{\frac{\gamma}{2}} \left[\omega_t, \nabla_{\mathcal{H}_+} \right] f \right\|_{L^2(\mathbb{R}^3)} \\ &\leq \frac{1}{C_1} \left\| \omega_t f \right\|_A + \frac{bc_0}{1+t} \left\| \langle \cdot \rangle^{\frac{\gamma}{2}+b-1} \omega_t f \right\|_{L^2(\mathbb{R}^3)} \leq \frac{bc_0+1}{C_1} \left\| \omega_t f \right\|_A, \end{split}$$

since $\sigma = \max\{1, \frac{b-\gamma}{2b}\}$, then $\frac{2b}{b-\gamma}\sigma \ge 1$, by using Hölder's inequality and (4.2), we have

$$2\sigma \int_{0}^{t} \tau^{2\sigma-1} \left\| \omega_{t} \nabla_{\mathcal{H}_{+}} f(\tau) \right\|_{L^{2}(\mathbb{R}^{3})}^{2} d\tau$$

$$\leq 2\sigma \left(\frac{bc_{0}+1}{C_{1}} \right)^{\frac{2b}{b-\gamma}} \int_{0}^{t} \tau^{\frac{2b}{b-\gamma}\sigma-1} \left\| \langle \cdot \rangle^{\frac{b}{2}} \omega_{\tau} \tau^{\sigma} \nabla_{\mathcal{H}_{+}} f \right\|_{L^{2}(\mathbb{R}^{3})}^{\frac{-2\gamma}{b-\gamma}} \left\| \omega_{\tau} f(\tau) \right\|_{A}^{\frac{2b}{b-\gamma}} d\tau$$

$$\leq 2\sigma \left(\frac{b(c_{0}+1)^{\sigma+1}}{C_{1}} \right)^{\frac{2b}{b-\gamma}} \left(\int_{0}^{t} \left\| \langle \cdot \rangle^{\frac{b}{2}} \omega_{\tau} \tau^{\sigma} \nabla_{\mathcal{H}_{+}} f \right\|_{L^{2}(\mathbb{R}^{3})}^{2} d\tau \right)^{\frac{-\gamma}{b-\gamma}} \left(\int_{0}^{t} \left\| \omega_{\tau} f(\tau) \right\|_{A}^{2} d\tau \right)^{\frac{b}{b-\gamma}}$$

$$\leq \frac{c_{0}}{(1+T)^{2}} \int_{0}^{t} \left\| \langle \cdot \rangle^{\frac{b}{2}} \omega_{\tau} \tau^{\sigma} \nabla_{\mathcal{H}_{+}} f(\tau) \right\|_{L^{2}(\mathbb{R}^{3})}^{2} d\tau + \left(\frac{4C_{T} \varepsilon_{0} (b-\gamma) (c_{0}+1)^{2\sigma+1} (T+1)}{C_{1}} \right)^{2}.$$
(6.2)

From this, we can deduce that for all $0 < t \le T$

$$\begin{aligned} \left\|\omega_{t}t^{\sigma}\nabla_{\mathcal{H}_{+}}f(t)\right\|_{L^{2}(\mathbb{R}^{3})}^{2} + \frac{c_{0}}{(1+T)^{2}}\int_{0}^{t}\left\|\langle\cdot\rangle^{\frac{b}{2}}\omega_{\tau}\tau^{\sigma}\nabla_{\mathcal{H}_{+}}f(\tau)\right\|_{L^{2}(\mathbb{R}^{3})}^{2}d\tau + \int_{0}^{t}\left\|\omega_{\tau}\tau^{\sigma}\nabla_{\mathcal{H}_{+}}f(\tau)\right\|_{A}^{2}d\tau \\ \leq 2\left(2C_{3} + (4C_{6})^{2}\right)\int_{0}^{t}\left\|\omega_{\tau}\tau^{\sigma}\nabla_{\mathcal{H}_{+}}f(\tau)\right\|_{L^{2}(\mathbb{R}^{3})}^{2}d\tau + (\tilde{C}_{1}\varepsilon_{0})^{2}, \end{aligned}$$

where \tilde{C}_1 is a positive constant depends on c_0 , b, γ , T and $C_1 - C_7$. Then from Gronwall's inequality, we can get that for all $0 < t \le T$

$$\begin{aligned} \left\| \omega_{t} t^{\sigma} \nabla_{\mathcal{H}_{+}} f(t) \right\|_{L^{2}(\mathbb{R}^{3})}^{2} + \frac{c_{0}}{(1+T)^{2}} \int_{0}^{t} \left\| \langle \cdot \rangle^{\frac{b}{2}} \omega_{\tau} \tau^{\sigma} \nabla_{\mathcal{H}_{+}} f(\tau) \right\|_{L^{2}(\mathbb{R}^{3})}^{2} d\tau + \int_{0}^{t} \left\| \omega_{\tau} \tau^{\sigma} \nabla_{\mathcal{H}_{+}} f(\tau) \right\|_{A}^{2} d\tau \\ &\leq \left(1 + 2T \left(2C_{3} + (4C_{6})^{2} \right) \right)^{2} e^{2T \left(2C_{3} + (4C_{6})^{2} \right)} (\tilde{C}_{1} \varepsilon_{0})^{2}. \end{aligned}$$

Assume $m \ge 1$ and (6.1) holds for all $p \in \mathbb{N}$ with $1 \le p \le m - 1$

$$\begin{aligned} \left\| \omega_t t^{p\sigma} \nabla^p_{\mathcal{H}_+} f(t) \right\|_{L^2(\mathbb{R}^3)}^2 &+ \frac{c_0}{(1+T)^2} \int_0^t \left\| \omega_\tau \tau^{p\sigma} \langle \cdot \rangle^{\frac{b}{2}} \nabla^p_{\mathcal{H}_+} f(\tau) \right\|_{L^2(\mathbb{R}^3)}^2 d\tau \\ &+ \int_0^t \left\| \omega_\tau \tau^{p\sigma} \nabla^p_{\mathcal{H}_+} f(\tau) \right\|_A^2 d\tau \le \left(L^{p-1} (p-2)! \right)^{2\sigma}, \quad \forall \ 0 < t \le T. \end{aligned}$$

$$\tag{6.3}$$

Now, we shall prove (6.3) for p = m. We apply the Proposition 2.3 to analyse terms $Q_1 - Q_3$. For the term Q_1 , by using (4.2), the Cauchy-Schwarz inequality and hypothesis induction (6.3), one has

$$Q_1 \le (16C_7(m-2)!)^2(T+1)^{2m\sigma} + \left((\varepsilon_0(T+1))^{4\sigma} + \frac{1}{8} \right) \int_0^t \left\| \omega_\tau \tau^{m\sigma} \nabla_{\mathcal{H}_+}^m f(\tau) \right\|_A^2 d\tau$$

+ 128
$$\left(C_7 \sum_{p=1}^{m-1} {m \choose p} \sqrt{\frac{(m-p+2)!}{2}} (T+1)^{(m-p)\sigma} (L^{p-1}(p-2)!)^{\sigma}\right)^2$$
.

For terms Q_2 and Q_3 , noting that using (2.4), one has for all $0 < t \le T$

$$\left\| t^{p\sigma}\omega_t \nabla^p_{\mathcal{H}_+} f(t) \right\|_{2,\frac{\gamma}{2}} \le \frac{(T+1)^{\sigma}}{C_1} \left\| t^{(p-1)\sigma}\omega_t \nabla^{p-1}_{\mathcal{H}_+} f(t) \right\|_A, \quad \forall \ p \ge 1,$$

then follows from (2.8), (4.2), (6.3) and the Cauchy-Schwarz inequality that

$$Q_{2} \leq \frac{1}{8} \int_{0}^{t} \left\| \tau^{m\sigma} \omega_{\tau} \nabla_{\mathcal{H}_{+}}^{m} f(\tau) \right\|_{A}^{2} d\tau + T \left(C_{5}\right)^{m} (m+3)! + \left(\frac{C_{6}(T+1)^{\sigma}}{C_{1}} \sum_{p=1}^{m} {m \choose p} \left(\sqrt{C_{5}}\right)^{m-p} \sqrt{(m-p+3)!} \left(L^{p-2}(p-3)!\right)^{\sigma} \right)^{2},$$

from (2.7), (4.2), (6.3) and the Cauchy-Schwarz inequality, one has for all $0 < t \le T$

$$Q_{3} \leq \left(4C_{4}\varepsilon_{0} + \frac{\varepsilon_{0}C_{4}(T+1)^{\sigma}}{C_{1}} + \frac{1}{16}\right) \int_{0}^{t} \left\|\omega_{\tau}\tau^{m\sigma}\nabla_{\mathcal{H}_{+}}^{m}f(\tau)\right\|_{A}^{2} d\tau + C_{4}\varepsilon_{1} \left\|\omega_{t}t^{m\sigma}\nabla_{\mathcal{H}_{+}}^{m}f\right\|_{L^{\infty}([0,T];L^{2}(\mathbb{R}^{3}))}^{2} + \frac{2\varepsilon_{1}C_{4}(T+1)^{\sigma}}{C_{1}} \left(L^{m-2}(m-3)!\right)^{2\sigma} + 16 \left(\frac{C_{4}(T+1)^{\sigma}}{C_{1}}\sum_{p=2}^{m-1} \binom{m}{p} \left(L^{m-2}(p-2)!(m-p-2)!\right)^{\sigma}\right)^{2}.$$

Since $\sigma \ge 1$, then from $p!q! \le (p+q)!$, one can deduce

$$\sum_{p=2}^{m-1} \binom{m}{p} \left((p-2)!(m-p-2)!\right)^{\sigma} \le 4 \left((m-2)!\right)^{\sigma} \sum_{p=2}^{m-2} \frac{m^2}{p^2(m-p)^2} \le 64 \left((m-2)!\right)^{\sigma},$$

$$\sum_{p=2}^{m-1} \binom{m}{p} \sqrt{\frac{(m-p+2)!}{2}} \left((p-2)!\right)^{\sigma} \le 2 \sum_{p=2}^{m-1} \binom{m}{p} (m-p-2)! \left((p-2)!\right)^{\sigma} \le 128 \left((m-2)!\right)^{\sigma}, \quad (6.4)$$

$$\sum_{p=1}^{m} \binom{m}{p} \sqrt{(m-p+3)!} \left((p-3)!\right)^{\sigma} \le 6 \sum_{p=1}^{m} \binom{m}{p} (m-p-2)! \left((p-2)!\right)^{\sigma} \le 400 \left((m-2)!\right)^{\sigma},$$

$$(m-p+1)! = 1 \ge (m-1)!^{\sigma}, \quad (m-1)! = 1 \ge 0$$

then taking $L \ge (T+1)^{2\sigma}$ and $\varepsilon_0 > 0$ small enough such that

$$\varepsilon_0 \le \min\left\{\frac{1}{8(T+1)}, \frac{1}{128C_4}, \frac{C_1}{32C_4(T+1)^{\sigma}}\right\}, < T$$

we can get that for all $0 < t \leq T$

$$\begin{split} \left\| \omega_{t} t^{m\sigma} \nabla_{\mathcal{H}_{+}}^{m} f(t) \right\|_{L^{2}(\mathbb{R}^{3})}^{2} + \frac{2c_{0}}{(1+T)^{2}} \int_{0}^{t} \left\| \langle \cdot \rangle^{\frac{b}{2}} \omega_{\tau} \tau^{m\sigma} \nabla_{\mathcal{H}_{+}}^{m} f(\tau) \right\|_{L^{2}(\mathbb{R}^{3})}^{2} d\tau + \int_{0}^{t} \left\| \omega_{\tau} \tau^{m\sigma} \nabla_{\mathcal{H}_{+}}^{m} f(\tau) \right\|_{A}^{2} d\tau \\ &\leq 2m\sigma \int_{0}^{t} \tau^{2m\sigma-1} \left\| \omega_{\tau} \nabla_{\mathcal{H}_{+}}^{m} f(\tau) \right\|_{L^{2}(\mathbb{R}^{3})}^{2} d\tau + 2C_{3} \int_{0}^{t} \left\| \omega_{\tau} \tau^{m\sigma} \nabla_{\mathcal{H}_{+}}^{m} f(\tau) \right\|_{2,\frac{\gamma}{2}}^{2} d\tau \\ &+ \frac{1}{2} \left\| \omega_{t} t^{m\sigma} \nabla_{\mathcal{H}_{+}}^{m} f \right\|_{L^{\infty}(]0,c_{0}/2];L^{2}(\mathbb{R}^{3}))}^{2} + B_{0} \left(L^{m-2}(m-2)! \right)^{2\sigma}, \end{split}$$

$$\tag{6.5}$$

with

$$B_0 = 1000 \left(C_3 + \frac{(C_4 + C_6)(T+1)^{\sigma}}{C_1} + 1 \right).$$

It remains to consider

$$2m\sigma \int_0^t \tau^{2m\sigma-1} \left\| \omega_\tau \nabla^m_{\mathcal{H}_+} f(\tau) \right\|_{L^2(\mathbb{R}^3)}^2 d\tau.$$

Applying (6.2) with $f = \nabla_{\mathcal{H}_{+}}^{m-1} f$ and the hypothesis (6.3), we have

$$2m\sigma \int_{0}^{t} \tau^{2m\sigma-1} \left\| \omega_{\tau} \nabla_{\mathcal{H}_{+}}^{m} f(\tau) \right\|_{L^{2}(\mathbb{R}^{3})}^{2} d\tau \\ \leq \frac{c_{0}}{(1+T)^{2}} \int_{0}^{t} \left\| \langle \cdot \rangle^{\frac{b}{2}} \omega_{\tau} \tau^{m\sigma} \nabla_{\mathcal{H}_{+}}^{m} f \right\|_{L^{2}(\mathbb{R}^{3})}^{2} d\tau + \left(\frac{4(b-\gamma)(c_{0}+1)(T+1)}{C_{1}} L^{m-2}(m-3)! \right)^{2\sigma}.$$

Plugging it back into (6.5), we can get that for all $0 < t \le T$

$$\begin{aligned} \left\| \omega_t t^{m\sigma} \nabla_{\mathcal{H}_+}^m f(t) \right\|_{L^2(\mathbb{R}^3)}^2 + \frac{c_0}{(1+T)^2} \int_0^t \left\| \langle \cdot \rangle^{\frac{b}{2}} \omega_\tau \tau^{m\sigma} \nabla_{\mathcal{H}_+}^m f(\tau) \right\|_{L^2(\mathbb{R}^3)}^2 d\tau \\ + \int_0^t \left\| \omega_\tau \tau^{m\sigma} \nabla_{\mathcal{H}_+}^m f(\tau) \right\|_A^2 d\tau \\ \leq 4C_3 \int_0^t \left\| \omega_\tau \tau^{m\sigma} \nabla_{\mathcal{H}_+}^m f(\tau) \right\|_{2,\frac{\gamma}{2}}^2 d\tau + \frac{1}{2} \left\| \omega_t t^{m\sigma} \nabla_{\mathcal{H}_+}^m f \right\|_{L^\infty(]0,T];L^2(\mathbb{R}^3))}^2 \\ + \left(\tilde{B}_0 L^{m-2} (m-2)! \right)^{2\sigma}, \end{aligned}$$

$$(6.6)$$

where

$$\tilde{B}_0 = B_0 + \frac{4(b-\gamma)(c_0+1)(T+1)}{C_1},$$

this implies that for all $0 < t \leq T$

$$\frac{1}{2} \left\| \omega_t t^{m\sigma} \nabla^m_{\mathcal{H}_+} f \right\|_{L^{\infty}(]0,T];L^2(\mathbb{R}^3))}^2 \le 2C_3 \int_0^t \left\| \omega_\tau \tau^{m\sigma} \nabla^m_{\mathcal{H}_+} f(\tau) \right\|_{L^2(\mathbb{R}^3)}^2 d\tau + \left(\tilde{B}_0 L^{m-2} (m-2)! \right)^{2\sigma},$$

substituting it into (6.6) yields

$$\begin{aligned} \left\| \omega_t t^{m\sigma} \nabla_{\mathcal{H}_+}^m f(t) \right\|_{L^2(\mathbb{R}^3)}^2 + \frac{c_0}{(1+T)^2} \int_0^t \left\| \langle \cdot \rangle \omega_\tau \tau^{m\sigma} \nabla_{\mathcal{H}_+}^m f(\tau) \right\|_{L^2(\mathbb{R}^3)}^2 d\tau + \int_0^t \left\| \omega_\tau \tau^{m\sigma} \nabla_{\mathcal{H}_+}^m f(\tau) \right\|_A^2 d\tau \\ &\leq 4C_3 \int_0^t \left\| \omega_\tau \tau^{m\sigma} \nabla_{\mathcal{H}_+}^m f(\tau) \right\|_{L^2(\mathbb{R}^3)}^2 d\tau + 2 \left(\tilde{B}_0 L^{m-2} (m-2)! \right)^{2\sigma}. \end{aligned}$$

It follows from Gronwall inequality that for all $0 < t \leq T$

$$\left\|\omega_t t^{m\sigma} \nabla^m_{\mathcal{H}_+} f(t)\right\|^2_{L^2(\mathbb{R}^3)} \le 8C_3 e^{4TC_3} \left(\tilde{B}_0 L^{m-2}(m-2)!\right)^{2\sigma},$$

therefore, we can deduce that

$$\begin{aligned} \left\|\omega_{t}t^{m\sigma}\nabla_{\mathcal{H}_{+}}^{m}f(t)\right\|_{L^{2}(\mathbb{R}^{3})}^{2} + \frac{c_{0}}{(1+T)^{2}}\int_{0}^{t}\left\|\langle\cdot\rangle^{\frac{b}{2}}\omega_{\tau}\tau^{m\sigma}\nabla_{\mathcal{H}_{+}}^{m}f(\tau)\right\|_{L^{2}(\mathbb{R}^{3})}^{2}d\tau \\ + \int_{0}^{t}\left\|\omega_{\tau}\tau^{m\sigma}\nabla_{\mathcal{H}_{+}}^{m}f(\tau)\right\|_{A}^{2}d\tau \leq \left(L^{m-1}(m-2)!\right)^{2\sigma}, \quad \forall \ t \in]0,T], \end{aligned}$$

if we choose

$$L \ge \max\left\{4\tilde{B}_0\left(8C_3(T+1)+1\right)e^{4TC_3}, (T+1)^2\right\}.$$

Acknowledgements. This work was supported by the NSFC (No.12031006) and the Fundamental Research Funds for the Central Universities of China.

References

- H. M. Cao, W. X. Li, C. J. Xu, Analytic smoothing effect of the spatially inhomogeneous Landau equations for hard potentials, J. Math. Pures Appl., 176 (2023) 138-182.
- [2] K. Carrapatoso, I. Tristani and C. K. Wu, Cauchy problem and exponential stability for the inhomogeneous Landau equation. Arch. Ration. Mech. Anal., 221 (2016), 363-418.
- [3] K. Carrapatoso, S. Mischler, Landau equation for very soft and Coulomb potentials near Maxwellians. Ann. PDE, 3 (2017), Paper No. 1, 65 pp.
- [4] H. Chen, W. X. Li, C. J. Xu, Analytic smoothness effect of solutions for spatially homogeneous Landau equation. J. Differential Equations, 248 (2009), 77-94.
- [5] H. Chen, W. X. Li, C. j. Xu, Propagation of Gevrey regularity for solutions of Landau equations, *Kinet. Relat. Models*, 1(2008), 355-368.
- [6] H. Chen, W. X. Li, C. J. Xu, Gevrey regularity for solution of the spatially homogeneous Landau equation, Acta Math. Sci. Ser. B Engl. Ed., 29(3) (2009), 673-686.
- [7] Y. M. Chen, L. Desvillettes, L. B. He, Smoothing Effects for Classical Solutions of the Full Landau Equation. Arch. Ration. Mech. Anal., 193 (2009), 391-434.
- [8] L. Desvillettes, C. Villani, On the spatially homogeneous Landau equation for hard potentials part i: existence, uniqueness and smoothness. *Communications in Partial Differential Equations*, 25 (2000), 179-259.
- [9] Y. Guo., The Landau Equation in a Periodic Box. Comm. Math. Phys., 231 (2002), 391-434.
- [10] T. Gramchev, S. Pilipović and L. Rodino, Classes of degenerate elliptic operators in Gelfand- Shilov spaces, New Developments in Pseudo-Differential Operators. Birkhäuser Basel, (2009), 15-31.
- [11] C. Henderson, S. Snelson, C[∞] Smoothing for Weak Solutions of the Inhomogeneous Landau Equation. Arch. Ration. Mech. Anal., 236 (2020), 113-143.
- [12] L. B. He, J. Ji and W. X. Li., On the Boltzmann equation with soft potentials: Existence, uniqueness and smoothing effect of mild solutions. arXiv:2410.13205v1
- [13] H. G. Li, C. J. Xu, Cauchy problem for the spatially homogeneous Landau equation with Shubin class initial datum and Gelfand- Shilov smoothing effect. SIAM J. Math. Anal., 51 (2019), 532-564.
- [14] H. G. Li, C. J. Xu, Analytic smoothing effect of non-linear spatially homogeneous Landau equation with hard potentials, Sci. China Math., 65 (2022), 2079-2098.
- [15] H. G. Li, C. J. Xu, The analytic smoothing effect of linear Landau equation with soft potentials, Acta Math. Sci., 43(6), (2023).
- [16] H. G. Li, C. J. Xu, Analytic Gelfand-Shilov smoothing effect of the spatially homogeneous Landau equation with hard potentials, *Discrete and Continuous Dynamical Systems - B.*, 29(4), (2024), 1815-1840.
- [17] H. G. Li, C. J. Xu, Gelfand-Shilov smoothing effect of the spatially homogeneous Landau equation with moderately soft potential, *Math. Meth. Appl. Sci.*, 46(1), (2023), 14379-14406.
- [18] S. Liu, X. Ma, Regularizing effects for the classical solutions to the Landau equation in the whole space. J. Math. Anal. Appl., 417 (2014), 123-143.
- [19] Morimoto Y., Pravda-Starov K. and Xu C.-J., A remark on the ultra-analysis smoothing properties of the spatially homogeneous Landau equation. *Kinet. Relat. Models*, 6 (2013), 715-727.
- [20] Y. Morimoto, C.J. Xu, Ultra-analytic effect of Cauchy problem for a class of kinetic equations, J. Differ. Equ., 247(2) (2009), 596-617.
- [21] Y. Morimoto, C. J. Xu, Analytic smoothing effect of the nonlinear Landau equation of Maxwellian molecules. *Kinet. Relat. Models*, 13 (2020), 951-983.
- [22] R. M. Strain, Y. Guo, Exponential Decay for Soft Potentials near Maxwellian. Arch. Ration. Mech. Anal., 187 (2008), 287-339.
- [23] C. Villani, On a New Class of Weak Solutions to the Spatially Homogeneous Boltzmann and Landau Equations. Arch. Rational Mech. Anal., 143 (1998), 273-307.

- [24] C. Villani, On the spatially homogeneous Landau equation for Maxwellian molecules, Mathematical Methods and Methods in Applied Sciences, 8(6) (1998), 957-983.
- [25] K.C. Wu, Global in time estimates for the spatially homogeneous Landau equation with soft potentials, J. Funct. Anal., 266 (2014), 3134-3155.
- [26] C. J. Xu, Y. Xu, A remark about time-analyticity of the linear Landau equation with soft potential, Anal. Theory Appl., 40(1) (2024), 22-37.
- [27] C. J. Xu, Y. Xu, The analytic Gelfand-Shilov smoothing effect of the Landau equation with hard potential, J. Differ. Equ., 414 (2025), 645-681.

XIAO-DONG CAO, CHAO-JIANG XU

School of Mathematics and Key Laboratory of Mathematical MIIT,

NANJING UNIVERSITY OF AERONAUTICS AND ASTRONAUTICS, NANJING 210016, CHINA

Yan Xu

DEPARTMENT OF MATHEMATICAL SCIENCES, TSINGHUA UNIVERSITY, BEIJING 100084, CHINA

 ${\it Email\ address:\ \texttt{caoxiaodong@nuaa.edu.cn;\ xuchaojiang@nuaa.edu.cn;\ xuchaojiang@nuaa.edu$