# Form and function in biological filaments: A physicist's review

Jan Cammann,<sup>1,\*</sup> Hannah Laeverenz-Schlogelhofer,<sup>2,\*</sup> Kirsty Y. Wan,<sup>2,†</sup> and Marco G. Mazza<sup>1,‡</sup>

<sup>1</sup>Interdisciplinary Centre for Mathematical Modelling and Department of Mathematical Sciences,

Loughborough University, Loughborough, Leicestershire LE11 3TU, United Kingdom

<sup>2</sup>Living Systems Institute and Department of Mathematics and Statistics,

University of Exeter, Stocker Road, Exeter EX4 4QD, UK

(Dated: February 19, 2025)

Nature uses elongated shapes and filaments to build stable structures, generate motion, and allow complex geometric interactions. In this Review, we examine the role of biological filaments across different length scales. From the molecular scale, where cytoskeletal filaments provides a robust but dynamic cellular scaffolding, over the scale of cellular appendages like cilia and flagella, to the scale of filamentous microorganisms like cyanobacteria which are among the most successful genera on Earth, and even to the scale of elongated animals like worms and snakes, whose motility modes inspire robotic analogues. We highlight the general mechanisms that couple form and function. Physical principles, such as classical elasticity and the non-reciprocity of active matter can be used to trace unifying themes linking these systems spanning about six orders of magnitude in length.

# I. INTRODUCTION

The interconnectedness between form and function in biology has been quantified at least since D'Arcy Wentworth Thompson's influential work [1]. Because geometrical features influence and direct biological activity through physical forces, it seems worthwhile to explore the link between geometry and biological function. In this Review, we examine a simple geometrical shape – the filament – which is pervasive in the architecture of living organisms, and discuss the function of filamentous shapes and the physical advantages they offer.

Filamentous shapes permeate every length scale in the biological world. Natural evolution has found that the geometrical properties of long aspect-ratio shapes can be used in a multiplicity of useful ways. The biopolymers forming the cytoskeleton provide structural integrity to the cell and form an intracellular transportation system. Cytoskeletal networks generate cytokinesis [2], migration [3, 4], and apical contraction [5], and even generate cytoplasmic streaming in the *Drosophila* oocyte [6].

At length scales of a few microns, cilia and flagella can generate fluid flow and active motion due to their fluid dynamical interactions [7–9] and they can synchronize to generate metachronal waves [8, 10]. Microorganisms such as filamentous fungi form complex networks of hyphae following available water and resources, and can move cytoplasm along the network [11]. Filamentous fungi are also the group of fungi most often used in biotechnological processes and include perhaps the most important fungus in the history of humanity: *Penicillium rubens*, which was used by Alexander Fleming to extract penicillin [12–14]. The slime mold *Physarum polycephalum* performs cytoplasmic flow across its network akin to a generalised form of peristalsis, which is optimised when the wavelength of the peristaltic wave is of the order of the size of the network [15]; this filamentous network can even encode memory about nutrient location in its morphology [16]. In the prokaryotic domain, cyanobacteria were among the first microorganisms to straddle the transition from unicellular to multicellular organisms [17, 18], and include numerous filamentous species.

# Box I | Geometry of filaments

A filament can be described as a portion of a threedimensional curve, whose local orientation changes smoothly along its length. The curve needs to be sufficiently smooth, that is, it is at least thrice continuously differentiable. The Frenet frame or trihedron is a local, right-handed, orthogonal coordinate system with basis (t, n, b). We indicate the curve by r(s), where r is the position vector and s is the arc length of the curve. The tangent unit vector to the curve is defined as  $t \equiv \frac{dr}{ds}$ . The normal unit vector  $n \equiv \frac{dt}{ds} / |\frac{dt}{ds}|$ , and always satisfies  $n \cdot t = 0$ . Furthermore,  $\kappa \equiv |\frac{dt}{ds}|$  is the local curvature of the curve r(s). Finally, the binormal vector  $b \equiv t \times n$  together with t and n form an orthonormal basis for space. These three vectors obey the Frenet–Serret equations

$$rac{dm{t}}{ds} = \kappa m{n}\,, \quad rac{dm{n}}{ds} = -\kappa m{t} + au m{b}\,, \quad rac{dm{b}}{ds} = - au m{n}\,,$$

where  $\tau$  is the torsion. The Frenet trihedron can be useful in the description of the helical motion of freeswimming organisms [19–21], sperm cells [22, 23], chiral active particles [24], and for gliding motility [25].



At much larger length scales, organisms such as worms, snakes, and eels have elongated bodies that grant them ecological advantages. Worms, for example, can form en-

<sup>\*</sup> equal contribution

<sup>&</sup>lt;sup>†</sup> K.Y.Wan2@exeter.ac.uk

<sup>&</sup>lt;sup>‡</sup> M.G.Mazza@lboro.ac.uk



FIG. 1. Filamentous shapes in biology. (a) Growing complexity of filaments across orders of magnitude in length, from nucleic acids that store information, to biopolymers that provide structural integrity to cells and form networks spanning the entire cytoskeleton. Flagella and cilia self-assemble from polymers to generate motion. Bacterial cells such as cyanobacteria self-organize into filamentous shapes. Animals, from worms and nematodes to snakes and eels utilize elongated shapes to achieve locomotion and collective behaviour. (b) Hypothetical schematic diagram of the biological function emerging from the three main properties of biological filaments: their length, their rigidity (measured by, e.g., Young's modulus), and their structural complexity, such as twisting, cross-linking into a network, up until the tissue level of the muscle fibers or an entire body.

tangled 'blobs' that exhibit shear-thinning rheology [26] and reversible self-assembly [27]. Such advantages can be combined to create higher levels of complexity, such as in a network built from cross-linking filaments, as illustrated in Fig. 1.

Although the systems described above span about nine orders of magnitude in size (from nucleic acids in the nanometre range in diameter, to snakes and eels in the metre range), it is possible to identify and discuss general physical features of elongated shapes (from macromolecules to whole organisms) that depend on the geometrical form of filaments and, in turn, influence their biological activity.

How do cytoskeletal filaments generate structural integrity in the cell, how do groups of cilia generate fluid flow, or how do large elongated organisms generate undulatory locomotion? Understanding how physical systems interact via simple rules to generate interesting collective behaviour can and will have novel applications. The condensed-matter motto of 'More is Different' proposed by Philip Anderson [28] is a guiding principle in this field. For example, a self-organizing robot exhibits behaviour intermediate between solid and liquid with adaptive capabilities such that it can flow over obstacles [29]. This combination of flexibility and structural integrity is reminiscent of a cell's cytoskeleton, or even of snake locomotion. Each of these filamentous systems can serve as inspiration for new biosynthetic materials [30]. Examples of this principle are the realisation of strainstiffening metamaterials using PDMS slats that mimic cell behaviour [31], or the use of carbon nanotubes to study the effect of crowding and filament stiffness on their microscopic dynamics [32, 33].

Physically, the rigidity and length of different biofilaments determine the class of models used to describe their mechanical and dynamical properties. If fluctuations are not important (because of the filament's rigidity or because of the length scale of observation), the biofilaments can be effectively described as smooth lines. In this context, we can mathematically describe filaments as a curve in a three-dimensional (3D) space whose local orientation changes smoothly along their length. Box I describes the Frenet–Serret equations used in differential geometry to characterize a curve.

At smaller length scales, such as inside the cell, the polymeric nature of biofilaments is consequential and this character is amplified by the nonequilibrium nature of the fluctuations. For example, variations in polymer stiffness and connectivity of actomyosin networks by actin filament bundling and cross-linking proteins can systematically tune the mechanical response of actin networks from extensile to contractile [34, 35].

Another dimension, useful in classifying filaments in the natural world, is the complexity of their organisation. Single globular actin proteins (G-actins) assemble in filaments (F-actin), which together with cross-linking proteins and a myriad of other molecules form cytoskeletal networks. Multiple cells, in turn, can assemble into simple filaments, such as the Oscillatoria lutea cyanobacteria, or form even more complex filamentous organisms, such as nematodes, eels and snakes.

In this Review, we discuss the physical processes and properties of filamentous shapes in the biological world across length scales. Section II discusses the role of biofilaments inside the cell. Section III reviews the role of filamentous cellular appendages, such as the locomotor function of cilia and flagella. In Sec. IV we discuss the biological properties of filamentous bacteria. Section V deals with macroscopic elongated organisms, such as snakes and eels. Finally, in Sec. VI we outline a roadmap of future work and outstanding questions at this exciting frontier shared by biology and physics, to establish a common framework describing the physics of biological filaments (Fig. 1)

#### II. BIOPOLYMERS & CYTOSKELETAL FILAMENTS

The cytoskeleton is a 3D polymer-based scaffold that spans the entire interior of a cell and defines cell morphology [36]. The basic building blocks are few key proteins that assemble together into a broad range of filaments [37]. These filaments together with a host of proteins that bind to the filaments' sides or ends form a network that provides mechanical stability to the cell [38]. Using only a few building blocks allows great flexibility and economy of materials. For example, cytoskeletal polymers achieve the same rigidity (elastic modulus) of gel-like substances with two orders of magnitude less material [38, 39].

Biological filaments are subject to different types of deformations: extensional and compressional forces and moments that cause the filaments to bend and twist. Any deformation mode costs energy, which is continuously used in the cytoskeleton to maintain the cell in a nonequilibrium state and allows dynamic control and reconfiguration [35]. The theory of elasticity can help identify the associated bending energy  $E_b$  [40], see Box II.

Generally, one expects the bending energy to be proportional to the square of the deformation, i.e., of the curvature; if the symmetry of the system does not distinguish positive and negative curvature, linear terms in the curvature cannot be present. To lowest order, the energy of deformation is then quadratic

$$E_b = \frac{1}{2} \beta_{\rm f} \int_0^L \left( \frac{\partial^2 \boldsymbol{r}}{\partial s^2} \right)^2 ds \,, \tag{1}$$

which is known as the Kratky–Porod model [41].

At the molecular level, cytoskeletal filaments are linear polymers: monomers that are bound together as an unbranched chain. Physically, they can be distinguished by their stiffness. Thermal fluctuations induce random deformations of the filaments; thus, the thermal energy  $k_{\rm B}T$  is a natural scale to classify the ease of bending. The ratio  $\lambda_{\rm p} = \frac{\beta_t}{k_{\rm B}T}$  is the persistence length of the filament. By comparing  $\lambda_{\rm p}$  with the filament length we can classify filaments as follows.

### Box II | Elastodynamics

Let us consider a small section of a thin rod. We assume that bending is in the zx plane, where z is parallel to the unperturbed rod (deviations of a slightly bent curve from a plane, its torsion, are higher order than its curvature [40]). Because external forces are zero, it can be shown that a thin volume element of the rod deforms only by extension (or compression), although this amount varies from point to point along the rod. The relative extension of an element of length dz is  $x/R_c$ , where  $R_c$  is the local radius of curvature. The strain tensor component  $u_{zz} = x/R_c$ , and the stress tensor component  $\sigma_{zz} = E u_{zz}$  [40], where E is Young's modulus. The bending energy per unit length  $\boldsymbol{L}$  is then  $E_{\rm b}/L = \frac{1}{2}\sigma_{zz}u_{zz} = \frac{1}{2}Ex^2/R_c^2$ . Integrating now over the cross-sectional area yields  $E_{\rm b} = \frac{1}{2} \beta_{\rm f} \kappa^2 L$ , where  $\beta_{\rm f} \equiv EI$  is the flexural rigidity or bending stiffness of the filament,  $I \equiv \int x^2 dA$  the second area moment or the area moment of inertia. For a solid cylinder of radius R,  $\beta_{\rm f} = \frac{1}{4}\pi R^4 E$ , and for a hollow cylinder with inner radius  $R_i$ ,  $\beta_{\rm f} = \frac{1}{4}\pi(R^4 - R_i^4)E$ . In general, however, the radius of curvature of the bending need not be constant but might vary along the contour of the filament. For dynamical processes, the Euler-Lagrange equations for an energy function that includes kinetic energy, the elastic energy  $E_b$  and an external load q yield the Euler-Bernoulli dynamic beam equation [42, 43]

$$\frac{\partial^2}{\partial z^2} \left( EI \frac{\partial^2 w}{\partial z^2} \right) = -\rho A \frac{\partial^2 w}{\partial t^2} + q(z) \tag{2}$$

where w(z) is the deflection of the filament from the equilibrium along its length z, and q(z) the load. These equations can be used to characterize buckling, e.g. in microtubules observed in living cells [44, 45].

If  $\lambda_{\rm p} \ll L$  the filament is considered *flexible* (the filament will exhibit highly coiled configurations); if  $\lambda_{\rm p} \approx L$  the filament is termed *semiflexible*, coiling and knotting are not present but thermal fluctuations can induce bending [46]; if  $\lambda_{\rm p} \gg L$  the filament is *stiff*, and it appears as a stiff rod. The persistence length can be measured from the fluctuations of the filament. Under thermal fluctuations, the tangent t to different parts of the filament performs a random walk. The correlation between two different points will then exhibit an exponential decay  $\langle t(s + \Delta s)t(s) \rangle = e^{-\Delta s/\lambda_{\rm p}}$ .

The cytoskeleton is highly dynamic, because its constituents can be assembled and disassembled as the need arises due to permanent energy dissipation. Cytoskele-

Biofilament	Shear modulus $G$	Persistence length $\lambda_{\rm p}$	Young's modulus $E$	Bending stiffness $\beta_{\rm f}$
units	Pa	$\mu{ m m}$	kPa	${ m N}~{ m m}^2$
Actin	283 [ <b>47</b> ] (§)	(10 - 20) [48]	$(1-3) \times 10^6 \ [49]$	$7.3 \times 10^{-26}$ [50]
Fibrin	104 [ <b>47</b> ] (§)	(0.1 - 1) [48]	$(1-15) \times 10^3 \ [51]$	
Vimentin	32 [ <b>47</b> ] (§)			$(4-12) \times 10^{-27}$ [52]
Keratin			$6 \times 10^3 - 10^8$ [53, 54]	
Microtubules	34 <b>[47</b> ] (§)	$(4-6) \times 10^3 \ [48]$	$(1-50) \times 10^6 \ [50, \ 55]$	$2.2 \times 10^{-23}$ [50]
Cellulose			$138 \times 10^{6}$ [56]	
Bacterial flagella			$(0.1 - 380) \times 10^6 $ [57]	$(0.1 - 100) \times 10^{-23} $ [57]
Motile cilia			$(0.01 - 120) \times 10^6(\dagger)$	$(0.3-6) \times 10^{-21}$ [58, 59]
Filamentous cyanobacteria		$(0.4 - 11.2) \times 10^3 [60, 61]$	$(16 - 46) \times 10^3 \ [60]$	$(1-20)^{-17}$ [60, 62]
$C. \ elegans \ (adult)$			(80 - 140)(*)[63]	$\sim 10^{-13}$ [63]
Eels and snakes			100 - 1000 [64, 65]	$10^{-2} - 10^{-3}$ [66]

TABLE I. Physical parameters characterizing examples of biological filaments, from cytoskeletal structural components and flagella, to filamentous organisms, up to the scale of snakes. (§) For shear-modulus measurements, strains were imposed by rheological measurements ranged from 1 to 5% and frequencies from 1 to 10 rad/s [47]. (†) Estimated from  $\beta_f$ . See text for a discussion about this large range. (\*) Approximating *C. elegans* as uniform rods.

tal filaments are grouped into three main classes: microtubules, actin filaments, and intermediate filaments [37], see Fig. 2. The main differences among these three types of filaments are their mechanical stiffness, polarity, and the molecular motors associated with them [37]. We now discuss these three classes of biofilaments in turn.

• Microtubules are hollow and polar tubular structures, which grow by the addition of tubulin dimers (8) nm in length) bound to GTP (guanosine-5'-triphosphate) molecules [67, 68]. Microtubules are the stiffest cytoskeletal filaments, see Table I. They are formed by  $\alpha$ - and  $\beta$ tubulin protein dimers. The protein dimers polymerize into linear chains, called protofilaments, which arranged as a hollow cylinder constitute a microtubule; most eukaryotic cells contain microtubules with 13 protofilaments, characterised by an internal diameter of about 15 nm and an external diameter of about 25 nm [48, 69]. Microtubules perform crucial functions in the life of a cell. During cell division, microtubule nucleate from the centrioles that organize the mitotic spindle [70]. Microtubules are also the building blocks of eukaryotic cilia and flagella [71, 72].

The considerable stiffness of microtubules is reflected in a persistent length  $\lambda_{\rm p}$  of about 5 mm, which can span the entire length of an animal cell [37], and a Young's modulus ranging from one to 50 GPa, depending on their length, or on the formation of network structures [50, 55], see Table I. Classic experiments employed thermal fluctuations to measure the flexural rigidity of microtubules  $\beta_{\rm f} = 2.2 \times 10^{-23}$  N m<sup>2</sup> [50].

Microtubules are highly dynamic systems, where the cell-regulated nonequilibrium processes of polymerisation and depolymerisation [73, 74] of a microtubule coexist (known as 'dynamic instability' [75]) and extend to a large fraction of the microtubules in a cell. The tau protein (short for 'tubulin associated unit') maintain the stability of microtubules in the axons of neurons [76], however, when hyperphosphorylation occurs, the microtubule disintegrate and the tau proteins form aggregates (neurofibrillary tangles, which are tangles of tau filaments) that are associated with neurodegenerative diseases [77, 78].

A single microtubule never reaches a steady state length but persists in prolonged states of polymerisation and depolymerisation that interconvert infrequently [79]. The cell uses the free energy of GTP hydrolysis to drive the nonequilibrium polymerisation; this free energy is  $\approx 7.5$  kcal mol<sup>-1</sup> under standard conditions, or  $\approx 12.5$ kcal mol<sup>-1</sup> in vivo [79, 80].

Recent measurements have shown that as microtubules grow or shrink, they undergo conformational changes. The GTP-tubulin dimers are initially curved at an angle of  $\approx 12^{\circ}$ ; after binding they straighten but exert a stress on the rest of the filament structure; finally, after GTP hydrolysis the  $\alpha$ -tubulin shortens by 3 Å and releases the stress, which can be harnessed to perform work [67, 81, 82]. These conformational changes in the dimers generate long-range interactions along the microtubules that modulate their growth [67].

Perhaps the most important example of microtubule self-organisation is the mitotic spindle, which partitions the set of chromosomes from a cell into two copies for the daughter cells [83, 84]. The polarity of microtubules is essential in this process, as each half of the spindle contains uniformly oriented microtubules, with their minusends at the pole (towards the centrosome) and their plusends extending away. The initial growth of microtubules around centrosomes is based on their dynamic instability, with random phases of growth and shrinkage probing the cytoplasm for kinetochores (the so-called 'search-andcapture' model [85]). That many classes of kinesins play a role in the spindle formation is evidence that the mitotic spindle is a nonequilibrium self-organised structure [84].

• Actin filaments are composed of (globular) G-actin monomers which readily assemble into long strings called F-actin ("F" for filamentous) of about 8 nm in width [48]. Actin filaments are about 300 times less stiff than microtubules, with  $\beta_{\rm f} = 7.3 \times 10^{-26}$  N m<sup>2</sup> [50], a persistence length  $\lambda_{\rm p} \approx 18 \ \mu {\rm m}$  [50], and with a Young's modulus  $E \simeq (1-3)$  GPa, which is comparable with Young's modulus for bone (9 GPa) [49]. Interestingly, actin is less resistant to bending than microtubules only because it is thinner, and not because it is more compliant than tubulin [50]. It is also worth comparing the Young's moduli of actin and tubulin. The measurements of  $\beta_{\rm f}$  for a microtubule implies a Young's modulus  $E \sim 1.2$  GPa, while for actin  $E \sim 2.6$  GPa [50]. This is comparable with  $E \sim 2-8$  GPa of silk (*Bombyx mori*) [86–88], or with  $E \sim 2.5$  GPa of keratin (from birds' feathers) [89]. These are all proteins employed by the cells to generate tensile strength [50], so it is not so surprising after all that filaments with similar Young's modulus evolved for the cytoskeleton.

Both actin filaments and microtubules are polar filaments, that is, they have a positive and a negative end due to an asymmetry at the molecular level. Hence, they can function as train tracks along which molecular motors move. Dynein and kinesin are the motor proteins associated to microtubules; by means of ATP hydrolysis they can transport vesicles, mRNA, and other proteins. Myosins are the motor proteins associated to actin. They are a superfamily [90]: at least 35 classes of motor proteins associated to actin filaments are known [91] and used for vesicle transport; they are responsible for muscle contraction in myocytes [92], and are macromolecular filaments. Some members of this superfamily, the non-muscle myosin II (NM II) are even involved in cell adhesion and migration [93].

Actin is present in high concentrations in the actomyosin cortex of animal cells, a layer attached to the inner surface of the cell membrane, and its function is to generate tension and regulate force and cell shape [96–98]. Actin is the main constituent of lamellipodia, two-dimensional structures responsible for movement of e.g. epithelial cells [99], and is involved in cell adhesion [100, 101]. The versatility of actin is but an example of the fact that the molecular organisation of biofilaments is extremely flexible; as a further example, amyloid fibrils can change bending rigidity by four orders of magnitude due to varying backbone interactions in the cross- $\beta$  structure [102].

Actin plays a major role during embryonic development, as the cells have to undergo a specific sequence of physical and biological changes [103]. Physically, cell cortical forces, cell adhesion and motility together produce self-organisation both at the tissue level and within single cells, e.g., reorientation of the polarity plane [104]. Another physical factor involved in tissue morphogenesis is actin-myoosin flows (or cortical flow), that is, a cell-scale internal flow of actin components of the cortex [105]. Various aspects are at play here: the character of



(a)

FIG. 2. (a) Endothelial cells from cow lung with two components of the cytoskeleton: microtubules (green) and actin filaments (red) [94]. (b) Confocal microscope image of a mouse 3T3 fibroblast cell. The nucleus has been stained blue, and two components of the cytoskeleton, actin microfilaments and intermediate filaments, are stained red and green respectively [95].

the actin-myosin flow dynamics (pulsatile versus continuous) and direction (centripetal or anisotropic) influence force generation; additionally, to transmit forces to other cells, this flow needs to be coupled to adhesion complexes at the cell membrane [103]. Although much has been understood about force generation in cells and tissue, there are still many open questions about how the exquisitely complex self-organised process is controlled by the cell.

• Intermediate filaments (IFs) are the most flexible of the three classes of cytoskeletal filaments, but they are also a very diverse class of filaments [106]. Estimates of their persistence length range from  $\lambda_p = 415 \pm 45$  nm for human wild-type vimentin [107] to  $\lambda_p = 2.1 \pm 0.1 \ \mu \text{m}$ [108]. There are six classes of IFs [109], which include, e.g., keratin, desmin, vimentin, and nuclear lamin. Structurally, IFs exhibit a hierarchical structure [110]. The alpha-helical domains form a coiled-coil structural motif, composed of two parallel in-register chains. Two chains aggregate in parallel to generate an IF molecule, and then two such molecules assemble in an antiparallel fashion to form a four-chain structural unit [109]. IF proteins are highly charged, thus representing versatile polyampholytes with multiple functions due to their strong selfassembling ability [111]. IF is built as a complex network of ionic interactions formed by the acidic rod and the basic head domain, and, unlike microtubules and actin filaments, IFs are non-polar, and lack enzymatic activity [111]. IFs form networks of filaments with important mechanical properties: upon being mechanically stressed, the network becomes stiffer [47, 111, 112].

Cytoskeletal networks and viscoelastic properties. The viscoelastic properties of cytoskeletal filaments are closely linked to their biologic function [49], because their extreme aspect ratio and rod-like structure dominate the rheological behaviour of the cell, and changes in their structure may cause gel-sol transitions observed when cells are activated or begin to move [47]. Measurements of the storage modulus G' for F-actin and fibrin suggest that the molecular basis of the viscoelasticity of these two types of networks is similar and may be approximated by the steric hindrance of diffusion of long, interpenetrated filamentous polymers [47]. While the viscoelastic measurements of these biofilaments can be explained with isotropic models, there is ample evidence that microtubules and F-actin undergo a transition from isotropic to liquid crystalline solutions under the influence of shear stresses [113], other macromolecules [114], or even spontaneously [115–117], because of the strong thermodynamic drive for long rod-like filaments to form liquid-crystalline arrays [118, 119].

Cytoskeletal filaments play a fundamental role in the mechanical properties and the biological function of the cell. This functional importance is due to not just the physical properties of isolated filaments, but, perhaps more crucially, to their structural organisation. Most cytoskeletal filaments form networks, and by controlling their architecture and connectivity cells achieve remarkable material properties [120, 121].

Actin filaments are assembled by actin-binding proteins into branched, cross-linked networks; the list of such proteins is vast but includes, e.g.,  $\alpha$ -actinin,  $\beta$ -spectrin, filamin, and fimbrin. These complex, dynamicallyreassembling, dendritic networks generate force at the cell periphery that creates membrane protrusions (lamellipodia), that in turn produce the cell's directed motility, essential in processes such as embryonic development, wound healing, and the immune response [122].

The networks assembled with semi-flexible cytoskeletal filaments have unusual viscoelastic properties [38, 120, 121, 123–125]. Upon increasing stress, networks of actin filaments show stress stiffening behaviour up to a critical stress of  $\sigma_c \sim 270$  Pa, after which the networks enters a stress-softening regime that is reversible, that is, if the stress is reduced no hysteresis is observed [126]. A microscopic mechanism that explains such behaviour is based on the initial resistance to stretching due to entropic effects in the biofilaments, leading to the stress-stiffening regime. Above  $\sigma_c$ , some filaments under compressive stresses will buckle, which decreases the elasticity of the network and leads to the stress-softening regime [126]. This is at variance with the behaviour of flexible polymers networks, which exhibit stress softening only due to failure and is not reversible [49]. Furthermore, cytoskeletal filaments show negative normal stresses, that is, when sheared they tend to pull inward [127]. On the other hand, flexible-polymer networks exhibit positive normal stress. These nonlinear viscoelastic features of cytoskeletal filaments contribute to generate the structural integrity and strength of the cell.

Biofilaments exist also outside of the cell. In animal cells, collagen is the most important filament found in the extracellular matrix [128]. There are many types of collagen molecules, but e.g. in humans more than 90%are type I collagen, found in skin and tendons. Type I collagen forms fibers in a hierarchical structures. The amino acids chains form together a triple helix, called tropocollagen, with a diameter of  $\approx 1.5$  nm. Multiple tropocollagens bundle together into protofibrils (diameter  $\approx 6 - 25$  nm). Multiple protofibrils further bundle together and form collagen fibril (up to  $1\mu$ m in length,  $\approx 100$  nm in diameter). The fibrils can then form thicker fibers, which self-organize into a disordered network, the collagen matrix [128]. The collagen matrix exhibits nonlinear elasticity, with an elastic modulus depending on concentration, pH, and ionic strength, and ranging from few Pa up to hundreds of MPa [129, 130].

In plant cells, the most abundant and important filament is cellulose. This is a linear polysaccharide chain of glucose monomers. The numerous hydrogen bonds within and between chains gives them considerable stiffness and generates robust fibrils, which are used in the cell walls of plant cells. Depending on its microstructure, the Young's modulus of cellulose ranges from  $\approx 25$  GPa [131] (for microcrystalline cellulose) to 114 GPa (for bacterial cellulose) [132] or the classic measurement 138 GPa [56].

A review of cell biopolymers cannot be complete without mentioning nucleic acids: DNA and RNA. The physics of the DNA helix, or generally of nucleic acids, is surprisingly complex [133], and is inspiring a number of applications in nanotechnology [134–139]; but, unfortunately, we do not have the space to address it here. There are, however, numerous excellent reviews on DNA's and RNA's structure and biochemical interactions [140–144] and also on application of topology to DNA [145, 146]

#### III. CELLULAR APPENDAGES

In a letter sent to the Royal Society of London in 1676, van Leeuwenhoek reported the discovery of protozoa and the observation of cilia and flagella. He wrote: "I also discovered a second sort of animalcules, whose figure was an oval, ... provided with diverse incredibly thin little feet, or little legs [cilia], which were moved very nimbly ..., and wherewith they brought off incredibly quick motions" (cf. translation of letter 18 [147, 148]).

Today, our language has changed a bit, and we describe cilia and flagella as slender, hair-like appendages that extend from the main cell body and perform diverse functions, such as enabling cell movement [149, 150], generating flows [151–153], sensing environmental stimuli [154, 155] and attachment to surfaces [156, 157]. While there are a myriad of different filamentous appendages found in biology, here our focus is on those most widely studied, while also aiming to capture the diversity of their structures, properties, mechanisms and functionalities.

Across the tree of life, the most well studied cellular appendages are those used for motility, in particular the flagella in prokaryotes and cilia in eukaryotes. These filaments whip, wave, or rotate through the action of molecular motors that transform the cell's chemical energy into mechanical energy [8, 149]. The resulting nonreciprocal motion enables net propulsion within the cell's microscale, viscosity-dominated environment [9, 158].

# Box III | Drag anisotropy

Drag anisotropy refers to the directional dependence of drag forces acting on an object as it moves through a medium. Due to their elongated form, filaments experience less hydrodynamic drag when pulled along their length than when pulled perpendicular to it. Drag anisotropy can be quantified as the ratio

$$\xi = \frac{\zeta_{\perp}}{\zeta_{\parallel}},\tag{3}$$

where  $\zeta_{\perp}$  and  $\zeta_{\parallel}$  are the friction coefficients for motion perpendicular and parallel to the object's length. Typically, for filaments  $\zeta_{\perp} \approx 2\zeta_{\parallel}$ . The total drag force density (drag per unit length) opposing a filament's motion at low Reynolds number (i.e. viscosity-dominated regime) is given by  $\mathbf{F}_{d} = -(\zeta_{\parallel}\mathbf{v}_{\parallel} + \zeta_{\perp}\mathbf{v}_{\perp})$ , where  $\mathbf{v}_{\parallel}$  and  $\mathbf{v}_{\perp}$  are the projections of the local velocity into directions parallel and perpendicular to the segment [9, 159].

A low Reynolds number world. At the microscale in which these filaments operate, inertia is insignificant and viscous forces dominate. In fluid dynamics, this is the world of 'Low Reynolds number'. The Reynolds number describes the ratio of inertial to viscous forces and is defined as  $\text{Re} = \frac{\rho UL}{\eta}$ , where  $\rho$  is the fluid density,  $\eta$  its viscosity, and U and L are characteristic velocity and length scales, respectively. In the world of a planktonic cell,  $\text{Re} \approx 10^{-5} - 10^4$ , thus, in this viscosity-dominated

environment, cells overcome and even exploit drag to generate motion [9, 158]. Cells can achieve this due to the anisotropic drag (defined in Box III) of slender appendages, which allows propulsive forces to be generated perpendicular to the local velocity of the filament [9]. Additionally, drag-based thrust relies on the ability of filaments to perform time-periodic deformations that are also time-irreversible motions, crucial for creating a nonzero time-averaged propulsive force [9, 158]. This can be achieved via the rotation of helical flagella or the whiplike beating of cilia.

Multiscale dynamics. The activity of cellular filamentous appendages encompasses dynamics across multiple scales [8], offering physicists many opportunities to develop frameworks and provide insights into how these dynamics integrate and interact across scales. Nanoscale molecular motors drive shape changes and motions of individual filaments, while these shape changes drive fluid flows [8, 149]. Resistive force theory (RFT) and slender body theory (SBT) are two hydrodynamic models of fluid-structure interactions for slender bodies within a low Reynolds number environment (see reviews [9, 150]and references therein). Both are used to estimate the propulsive forces generated by slender appendages based on their waveforms and resulting microswimmer velocities. While RFT is a simpler approximation that neglects long-range hydrodynamic interaction, SBT is more accurate but also more computationally intensive. Mechanical coupling, both via hydrodynamic interactions and the internal cell cytoskeleton, enables coordination of multiple appendages [160, 161], and motivates a range of work that considers cellular appendages as coupled oscillators [162, 163]. Arrays of cilia can synchronise into metachronal waves [160, 164-166], while the bundling and unbundling dynamics of groups of bacterial flagella are key for their run-and-tumble motility behaviour [167, 168]. Large numbers of cilia can also group together to form compound ciliary structures, such as the leg-like 'cirri' the ciliate *Euplotes* uses for walking [169] and the comb plates of ctenophores that comprise tens of thousands of cilia [170]. The individual and collective activity of filamentous appendages drive fluid transport, self-propulsion and organism-environment interactions. Groups of microswimmers can display collective behaviours such as bacterial swarming [168, 171] and formation of bioconvection patterns that arise from hydrodynamic instabilities [172–176].

Considering the variety of mechanisms cells employ to actuate their filamentous appendages, these can be broadly classified into four different principles of activity: rotational actuation of passive helical propellors, linear motor actuation of flexible filaments to generate bending, assembly-disassembly dynamics, and pre-stressed springs; see Fig. 3 for the corresponding schematics and examples.

Rotationally actuated filaments. The rotation of passive filaments by cells involves a rotary complex embedded in the cell membrane at the base of the filament



FIG. 3. Four principles of activation in filamentous cellular appendages. (a) Rotationally actuated filaments are typically rigid helical filaments rotated at their base, such as bacterial flagella seen here in an image reproduced from [177] of a fluorescently labelled Escherichia coli cell. (b) Linearly actuated bending is the mode of activation found in the cilia of eukaryotes, with typical waveforms consisting of power and recovery strokes. The microalga Chlamydomonas uses two cilia for motility, here shown as three overlaid frames from a movie of its swimming behaviour. (c) Assembly-disassembly activity involves the dynamic extension and retraction of filaments, such as in pili used to generate twitching motility in bacteria and archaea. An example cryo-electron micrograph of a negatively stained wild-type Thermus thermophilus cell shows long flexible pili filaments [178]. (d) Pre-stressed springs respond to changes in intracellular calcium resulting in rapid movements, as seen in the time series of the contractile motion of the stalk of *Vorticella* [156] and the haptoneme of a haptophyte alga [179].

where stator and rotary components interact to generate torque that is transmitted to an inert rigid helical filament. There are two main examples of such filaments: the bacterial flagella and the archaea's archaella [149]. Their molecular structures differ. The bacterial flagellum is a hollow cylinder self-assembled from the flagellin protein into a a chiral structure [180, 181]. The archaea's archaella has a complex structure [182]. However, they are both ~10-20 nm wide and up to ~10  $\mu$ m in length. Archaella use an ATPase to generate torque via ATP

hydrolysis [183, 184], while the flagella's rotary complex takes a more structurally intricate form and is powered by the proton motive force [168, 185, 186]. In both cases, the torque generated by the embedded molecular machinery results in a rotation of the connected passive rigid filament that in turn acts as a propellor for cell motility [149].

Linearly actuated filaments. In contrast, cilia (sometimes referred to as eukaryotic flagella) are flexible filaments, typically  $\sim 250$  nm wide and up to  $\sim 100 \ \mu m$ long, and are actuated by linear stepper motors distributed along their entire length [149]. Cilia have a highly conserved structure across eukarvotes known as the axoneme, which comprises 9 microtubule doublets surrounding a central pair of microtubules [187]. Ciliary bending and twisting is powered by dynein motors that walk along the 9 outer doublets. Since the microtubules are usually fixed to the cell body at the base of the cilium, this linear forcing generates differential sliding between neighbouring doublets resulting in bending of the axoneme [188, 189]. The mechanism by which dynein motor activity generates ciliary beating is still under debate, with several biophysical models proposed that largely fall under two categories: (1) mechanical feedback within the axoneme regulates dynein activity [188, 190, 191], (2) continuous dynein activity along the flexible filament results in dynamic instabilities [192]. Dynein motors are typically distributed along the whole length of the cilium, giving a high degree of flexibility in the types of waveforms that are possible to produce by coordinating the dynein activity in different ways. Compared with the rotation of a rigid filament, which only has a small set of parameters that can be controlled (i.e. speed and direction of rotation), cilia have the potential to display a much more complex and nuanced diversity of waveforms, with calcium playing a central role in the regulation of ciliary activity [193]. The ability of cilia to change their waveform is critical for enabling ciliated organisms to switch locomotor gaits and perform complex motion [169, 194]. Although cilia are highly conserved, there are several examples of variations or additions to the standard 9+2 axoneme structure that exemplify how cilia can be adapted to perform specific functions [195]. For example, the cilium of the parasite Trypanosoma brucei has a paraflagellar rod, a lattice-like filament that runs along the length of the axoneme, thought to affect axonemal beating by imposing mechanical constraints and contributing to signaling pathways [196]. Excavates, species of phagotrophic flagellates, are characterised by possessing vaned cilia within a ventral groove whose confined beating is thought to be particularly efficient for generating feeding currents [197].

Assembly-disassembly dynamics. An entirely different principle of activity is that seen in the type IV pili of bacteria and archaea, in which cycles of extension, surface binding and retraction enable cells to perform functions such as twitching motility, surface sensing and DNA uptake [198]. This activity is enabled by ATP hydrolysis, which powers the rapid assembly and disassembly of pilin subunits [199].

Pre-stressed springs. Lastly, pre-stressed springs composed of filaments and tubules power many of the fastest cellular movements [200]. There are two main types: a microtubule-based system found in some microalgae (i.e. the haptoneme in haptophytes) [179] and a myoneme based system found in some ciliates (e.g. the spasmoneme of vorticella) [201, 202]. In both cases, rapid coiling is thought to be driven by an increase in  $Ca^{2+}$  concentration, which changes the energetically preferred conformation of the filament from elongated to coiled [202, 203]. For the haptoneme, some evidence suggests that this is driven by conformational changes in microtubule associated proteins that bind with  $Ca^{2+}$  [179], while for the spasmoneme it is hypothesised to be a combined electrostatic and entropic effect [200].

Elastodynamic properties. A key mechanical parameter for characterising the movements of cellular appendages is the bending stiffness (see Box II), which can be estimated from the deformation induced by forces and torques applied using techniques such as optical tweezers [204], microneedles [205], magnetic beads [206] or microfluidic flow devices [57]. Using such approaches the bending stiffness for the bacterial flagella has been estimated to be 1-1000 pN  $\mu$ m<sup>2</sup> (see table S1 in [57]), while estimates for eukaryotic motile cilia fall within the 300-6000 pN  $\mu m^2$  range [58, 59, 205, 206]. The Young's modulus E is generally estimated indirectly from experimentally measured values of the bending stiffness. This requires approximating the second area moment I, which is challenging when a filament is not easily defined as a solid or hollow cylinder, such as in the case for cilia. For motile cilia, two values are commonly used:  $I = 1.3 \times 10^{-31} \text{ m}^4$ [207] and  $I = 3 \times 10^{-29} \text{ m}^4$  [208], the latter assumed more relevant for tighter connections between the microtubulues of the axoneme [58]. This gives a large range of estimates  $E \sim (0.01 - 120)$  GPa.

Appendages as cellular antennae. In addition to their role in motility and fluid transport, appendages can also play important roles in sensory reception and signal transduction that benefit from the geometric and mechanical properties of their filamentous form [154, 155, 198, 209]. Cilia often possess many sensory structures such as chemical receptors and mechanically-sensitive ion channels that enable them to act as the cell's antenna [154, 210]. Filamentous appendages can extend far beyond the cell surface, reaching more well-mixed and therefore more 'representative' regions of the surrounding environment, making them effective as chemosensory probes [154]. Additionally, their high aspect-ratio and high surface area to volume ratio, improves their capture rate and chemosensory effectiveness [209]. Their elongated shape also means that filamentous appendages can be considered as extended cantilevers, facilitating their sensitivity to mechanical stimuli [155].



FIG. 4. Examples of cyanobacteria self-organisation. (a) Many cyanobacteria species grow into filaments made out of a chain of identical cells [211]. (b) These filaments are capable of organizing into intricate structures like reticulate mats [60], spirals [212] (c), and even ropes [213] (d).

#### **IV. FILAMENTOUS MICROBES**

Various microbial organisms have evolved a filamentous body shape. Examples of such organisms that have attracted the interest of physicists include various filamentous cyanobacteria [60, 62, 214–217], filamentous phages [218–220], and nematodes, [63, 221–223] that straddle the boundary between the microscopic and macroscopic world. As there are too many filamentous microorganisms to list, this section will focus on discussing more generally the implications of a filamentous body shape at the microscale for a few important classes of microorganisms.

Having a body with a large aspect-ratio like a filament comes with some advantages over microbes shaped as spheres or short rods. An elongated shape increases the chance of meeting one another in disordered environments [224]. Thanks to a filamentous shape, microbial organisms can self-propel without appendages. Rather than using flagella or cilia to generate motility, they can bend their whole body to generate thrust thanks to the different drag coefficients that a slender body experiences in the direction tangential and perpendicular to its motion in a viscous fluid [9, 159] (see Box III). A filamentous shape has been observed to be a viable strategy to avoid predation by protists by simply being too long to be ingested easily [225] (this approach is however futile at a certain size difference between devourer and the devoured, as is evident in the popularity of spaghetti as a human food source). The increased surface-to-volume

ratio of a filamentous shape allows for easier uptake of nutrients [226], but also allows stronger attachment to substrates [227] in bacteria, because of a larger contact

A filament's most crucial mechanical property is its bending stiffness. It is known one can draw conclusions about the forces at play from purely geometrical observations (see Box II). Techniques to measure the bending stiffness  $\beta_{\rm f}$  and Young's modulus E of gliding filaments with the help of microfluidics [60] and mircopipette techniques [62] have been developed (see Table I).

patch, compared to a spherical one.

Filamentous cyanobacteria often exist in a surfacebound but motile state, called gliding [228–230]. The underlying mechanism that allows the generation of propulsive forces in this state is still a matter of debate [231– 235], as no cyanobacteria species are known to possess flagella [229]. The main mechanisms that are debated are: type IV pili extension and retraction [229, 236], fibrils [232, 235], ejection of a jet of extracellular slime from junctional pores [233, 237–239] (at least for some species), and acoustic streaming [217, 234, 240]. By observing the self-buckling of a filament, the forces at play in its propulsion can be inferred via the bending stiffness [62]. Such observations combined with the independence of propulsion speed on filament length implies, that all cells within a cyanobacteria filament contribute roughly equal to its propulsion.

Cyanobacteria locomotion exhibits another interesting feature: path-tracking motion, where the body follows the path laid by the forward end of the filament [214]. This type of locomotion in worms is also known as metameric locomotion [241]. Physically, information about a filament's orientation persists over the time it takes a filament to traverse its own length [241]. Although their motility is induced by polar forces, filaments themselves are seemingly nematic, with no head or tail end. In fact, cyanobacteria have been observed to reverse their direction of motion frequently, as a strategy to navigate towards more favorable conditions, and even to form communities. [229, 242-244]. Given the delocalised nature of the filaments propulsive mechanism, a high degree of intercellular coordination is required to reverse the direction of motion, with the mechanisms of communication between cells not yet understood. Confined active filaments with low bending rigidity have been shown to curl up into tightly wound spirals (see Fig. 4(c)), effectively trapping themselves, greatly impeding their mobility [245–248]. Utilising reversals is a strategy to escape this trapped state [212].

Having a high aspect-ratio allows microbes to effectively sample changing conditions within space. While organisms with short bodies commonly have to move around to find favourable conditions, essentially sampling the environment at random to inform their movement through chemo- or phototaxis, elongated filaments can sample gradients along their body and react before they have wholly immersed themselves in an unfavourable environment. Cyanobacteria, for instance, have been observed to react to light shining only onto parts of their body [249, 250], showing scotophobic responses, well before the whole filament loses access to light by reversing its gliding direction [215].

Having an elongated body may even allow organisms to be in two places with different conditions at once. Cable bacteria are filamentous, multicellular bacteria belonging to the family *Desulfobulbaceae*. They can conduct electrons over centimetre scales [251, 252]. Through their filamentous shape they can couple sulfide oxidation in deeper, anoxic marine sediment with oxygen reduction in surface layers [253–255]. Electrical activity is also involved in a microbial feat of engineering: some electroactive bacteria produce nanowires to transfer electrons to extracellular electron acceptors [256]. These are, e.g.,the *Geobacter sulfurreducens* or the *Shewanella oneidensis* MR-1 bacteria [257–260].

Filamentous microbes can explore spaces inaccessible to plumper organisms with the same volume. Semiflexible active filaments have been shown to effectively explore porous environments [261]. Many soil dwelling microbes are filamentous in shape, in addition to their ability to explore a porous environment it has been speculated that their shape may enable them to fix their position in space, by wrapping around soil particles [226], allowing them to resist being washed away by percolating water. Entangled filamentous microbes have been observed to stabilize their habitats and protect them against erosion [262, 263]. Furthermore, nonmotile cells exploring a polymeric fluid can form long cables as they proliferate [264].

Their filamentous nature allows filamentous organisms such as cyanobacteria to organise into structures much larger than an individual filament [214, 216, 265, 266]. Their filamentous shape allows them to glide over one another without losing contact with the substrate. The dynamics of gliding filaments result in the formation of a network-like reticulate pattern of filament bundles [214, 216, 266] (see Fig. 4(b)). Under different conditions, filaments have been observed to form aster-like aggregates [244]. These relatively simple, mostly twodimensional structures may serve as a starting point for filaments to organize themselves into even more intricate three-dimensional structures, ranging from complex patterned biomats [265, 267] to the filaments weaving themselves into ropes [213] (see Fig. 4(d)), bridging large distances. The physical mechanisms behind this level of self-organisation and how the mechanical properties of these structures compare to that of a single filaments is yet to be understood, but can shed light on biofilm formation [268, 269]. It is plausible to presume that such complex assemblys could exhibit unusual rheological properties, just like those observed in cytosceletal networks (see Sec. II).

#### V. UNDULATORY LOCOMOTION AT THE MACROSCOPIC SCALE

At the organismal scale, animals with a slender and deformable body shape are also 'active filaments', and in some cases the analysis and framework presented in the previous sections could still apply. Diverse animals locomote by propagating undulatory waves. These body deformations are used to propel the organisms forward, although the actuation mechanisms may be very different across species and spatial scales (Figure 5). Well-studied examples include crawling worms [270, 271], swimming eels [272], and gliding snakes [273]. The study of both limbed and limbless movement has paved the way for understanding the origins and principles of neuromechanical control in diverse organisms, particularly the interactions between neural patterning and muscular activity [274]. Undulatory or sinusoidal locomotion strategies evolved across different phyla, often according to a basic neural circuit design involving mutually inhibitory halves with the left and right sides deforming alternately [275]. These insights into the neural basis of behaviour have led to the development of novel bio-inspired mechanisms for actuating synthetic systems including robots [276].

Since wave-like deformations are inherently nonreciprocal, they form the basis of diverse motility strategies through different media. In some cases, despite the large difference in scale compared to microscopic organisms such as flagellates and cyanobacteria (see Sec. III and IV), the undulatory locomotion of macroscopic animals such as nematodes and snakes can still be considered to operate in an overdamped regime where inertia is negligible, and where friction (instead of fluid viscosity) dominates. This is particularly true of locomotion in sand or granular media, where a resistive force theory based on drag anisotropy (see Box III) has been derived empirically, inspired by the movement of slender bodies through viscous fluids [277, 278].

At the individual organism scale, distinct undulatory behaviours can arise from the same underlying structure and neural circuitry. A given body shape can be decomposed into a small number of principal components or oscillation modes, reflecting the intrinsically lowdimensional nature of the posture space, and allowing diverse undulatory shape modes to be reconstructed with only a small number of parameters [283]. In the example of *C. elegans* crawling, its motion in 2D is stereotyped and well-captured by just six basis modes, or eigenworms [270, 271]. Frequency and amplitude thus uniquely define a gait, which in turn determines the distance traversed by the organism in one cycle. Notions such as geometric phase can help understand and predict the mapping between these gaits and eventual displacement in the lab frame [284]. These dynamics are not totally unconstrained however, undulatory locomotion across swimming sperm, nematodes, and even fish in overdamped environments may be subject to a universal dispersion re-



FIG. 5. Undulatory locomotion of active filaments across scales. (a) For *C. elegans* crawling movement on agar [279]. (b) Sidewinding locomotion of a snake, and (c) the timetrace of tracked points along the body of the snake as it moves on sand [280]. (d) Vinegar eels self-organise into metachronal waves [281]. (e) Topological entanglement in worm blobs [27]. (f) Bio-inspired snake-like robots with CPG-like control modules [282].

lationship [285, 286]. Thus, active filaments can achieve highly reconfigurable dynamics with minimal neurome-chanical coupling.

Collectively, filamentous organisms also engage in novel behaviours that may not be accessible by individuals. At sufficiently high densities, these filaments selfassemble to form adaptive active materials with both solid-like and liquid-like properties, that are capable of emergent functions such as directed locomotion or smart navigation. Macroscopic 'blobs' of the 1 cm-long California blackworm Lumbriculus variegatus undergo dynamic collective movements to avoid desiccation or thermal stress without the need for any centralised control [287]. The entanglement, or 'braiding', process enables fast and reversible tangling and untangling of worm collectives. Intriguingly, L. variegatus blobs can take minutes to assemble but only milliseconds to disentangle. A recent study proposed an explanation for this phenomenon by experimentally imaging the 3D trajectories of the worms and mapping the dynamics onto a 3D filament model of Kirchhoff rods [27]. The unique, activitydependent rheology of entangled blobs of *Tubifex tubifex*  worms was also studied and shown to be shear-thinning, but with a different scaling compared to polymer melts [26]. These properties can even be harnessed to enable activity-dependent separation (chromatography) of worm mixtures [288]. In another free-swimming nematode *Turbatrix aceti* (vingar eels), concentrated suspensions of these organisms display large-scale metachronal waves surprisingly similar to those found in ensembles of cilia and flagella [281, 289]. Understanding how these deformable filamentous bodies actively control the topology and rheology of the collective could inspire novel designs of functionally tunable synthetic materials [290].

More generally, undulatory or anguilliform locomotion in elongated bodies allows for direct interactions between an organism and the surrounding medium, which can yield novel principles of biomimetic/bioinspired locomotion reliant on embodied design with few mechanical components. Self-propulsion can also be easily adapted for movement through different environments, for example for snakes the same basic locomotion mechanism can be equally effective for locomotion on land as well as in water [291, 292]. In more extreme cases, 'flying' snakes from the genus *Chrysopelea* use body undulations to generate lift to glide stably over several metres through the air [293].

From a control theoretic perspective, undulatory locomotion is universal perhaps because it reduces the need for more complex neural control mechanisms and can be achieved simply by coordinating alternating waves of (usually muscle) contraction and relaxation. In some cases, oscillations emerge spontaneously as a result of instabilities, for example dynamics reminiscent of ciliary beating have been observed in macroscopic chains of selfpropelled microrobots, with the transition mapped to a supercritical Hopf bifurcation [294]. Indeed substrateattached, microscopic elastic filaments forced at one end also exhibit oscillations in different regimes [295–297]. The propagation of waves thus provides one of the simplest possible appendage or body actuation mechanisms, and naturally breaks time-reversal symmetry (particularly relevant for inertia-free swimming). This results in highly adaptable and flexible control of movement where both the speed and direction of travel can be readily changed in response to stimuli.

### VI. ROADMAP & KEY OPEN QUESTIONS

Filamentous structures are ubiquitous in biology from cytoskeletal filaments to propulsive appendages to whole organisms. Despite spanning several orders of magnitude in scale and complexity, surprising parallels can be observed in how biological form translates into physiological function in these different contexts. In this review, we have sought to outline the core physical principles that govern these relationships and common frameworks for studying filamentous systems. As cutting-edge experimental technologies continue to deliver ever finer resolution of the molecular composition and architecture of the constituent parts of organelles, cells, and organisms, there is increasing need to integrate these insights with functional assays and biophysical models. At the same time, we should also be mindful that in many cases the natural context or habitat in which organisms live could also contribute to any observed phenomena [298]. Therefore, there is also a growing need to study and understand the behaviour of living systems 'in the wild', not just under controlled laboratory settings.

Critical to many modelling and simulation approaches is detailed knowledge of the emergent material properties of the many types of filaments. Yet such measurements are not always available as they often rely on access to specialised technology, such as high-resolution imaging, atomic force microscopes, micropipette force sensors [63], or can only be performed on certain filaments, or in nonliving systems or systems that have been rendered passive. This apparent disconnect requires more holistic, physics-inspired understanding of the context-dependent material properties of a filament and filamentous network, particularly due to their out-of-equilibrium activity. As a concrete example, the measurements of physical parameters such as Young's modulus E or the bending stiffness  $\beta_{\rm f}$  of cytoskeletal filaments as shown in Table I strongly depends on the chemical and physical context.

Recently, theories of active polymers [299] and active nematics [300] have identified interesting collective behaviour (such as bands, asters, and topological defects [245, 246, 301–303]) that reproduce experiments with purified systems of microtubules and molecular motors [304–309]. While these systems shed light on the nonequilibrium dynamics of cytoskeletal filaments, the challenge that lies ahead is to develop these theories towards more realistic conditions like those inside the cell, which is a 3D crowded environment. Thus, these theoretical frameworks should be developed in tandem with experimental innovations. Identifying appropriate dimensionless quantities from filament properties, like the ones in Table I, will enable the development of theories describing common phenomena in the collective dynamics of filaments. Doing so will allow the development of experimentally convenient proxy systems. Thus, efforts in characterising filaments, completing and expanding the data in Table I is required to develop a unified understanding of filament dynamics across scales.

The study of active or self-propelled filaments in biology have led to advancement of new mathematical and physical theories, notably in the areas of topological braiding and entanglement, nonequilibrium statistical physics, and extreme mechanics such as supercoiling. However, one constraint in understanding filamentous systems is the practical challenge of acquiring good resolution data, and the lack of efficient computational tools to track filaments, particularly from 3D image stacks. There is also the open question of which metrics are most appropriate to describe the deformation of filaments and their collective organisation. Thus, devising universallyapplicable, relatively system-independent measures for such systems requires further innovation.

We speculate that, because living systems apparently increase entropy production [310, 311] while producing more complex structures (Fig. 1) during their evolution, a generalised form of entropy might be a useful quantity in the search of system-independent measures. Because configurational degrees of freedom (positions, orientations) can be measured, it is possible to define structural complexity (or entropy) and use it to characterize living systems. Consider a function  $\psi(\mathbf{x})$  representing the configuration of a system, for example, the atomic positions in a protein, the orientation of filaments in a network, or the muscle fibers in a vertebrate. The complexity of an ensemble can be quantified by means of a generalised Shannon–Boltzmann entropy [312–314]  $H = -\int D\psi P(\psi(\boldsymbol{x})) \ln P(\psi(\boldsymbol{x}))$ , where  $\int D\psi$  represents a path-integral sum over all possible configurations. We note that elongated shapes have five degrees of freedom

- [1] D'Arcy Wentworth Thompson, On Growth and Form (Cambridge University Press, 1917).
- [2] Ulrike S Eggert, Timothy J Mitchison, and Christine M Field, "Animal cytokinesis: from parts list to mechanisms," Annu. Rev. Biochem. 75, 543–566 (2006).
- [3] Ewa Paluch, Cécile Sykes, Jacques Prost, and Michel Bornens, "Dynamic modes of the cortical actomyosin gel during cell locomotion and division," Trends Cell Biol. 16, 5–10 (2006).
- [4] Tim Lämmermann, Bernhard L Bader, Susan J Monkley, Tim Worbs, Roland Wedlich-Söldner, Karin Hirsch, Markus Keller, Reinhold Förster, David R Critchley, Reinhard Fässler, et al., "Rapid leukocyte migration by integrin-independent flowing and squeezing," Nature 453, 51–55 (2008).
- [5] Adam C Martin, Matthias Kaschube, and Eric F Wieschaus, "Pulsed contractions of an actin–myosin network drive apical constriction," Nature 457, 495–499 (2009).
- [6] Sayantan Dutta, Reza Farhadifar, Wen Lu, Gokberk Kabacaoğlu, Robert Blackwell, David B Stein, Margot Lakonishok, Vladimir I Gelfand, Stanislav Y Shvartsman, and Michael J Shelley, "Self-organized intracellular twisters," Nat. Phys. 20, 1–9 (2024).
- [7] James Lighthill, "Flagellar hydrodynamics," SIAM Rev. 18, 161–230 (1976).
- [8] William Gilpin, Matthew Storm Bull, and Manu Prakash, "The multiscale physics of cilia and flagella," Nat. Rev. Phys. 2, 74–88 (2020).
- [9] Eric Lauga and Thomas R. Powers, "The hydrodynamics of swimming microorganisms," Rep. Prog. Phys. 72, 096601 (2009).
- [10] Andrej Vilfan and Frank Jülicher, "Hydrodynamic flow patterns and synchronization of beating cilia," Phys. Rev. Lett. 96, 058102 (2006).
- [11] D A Klein and M W Paschke, "Filamentous fungi: the indeterminate lifestyle and microbial ecology," Microb. Ecol. 47, 224–235 (2004).
- [12] Alexander Fleming, "On the antibacterial action of cul-

(compare with the three degrees of freedom for a spherically symmetric object); furthermore, a flexible elongated shape has 5 degrees of freedom per unit length. Such explosion of degrees of freedom might constitute an entropic incentive for biological systems to evolve and use biological filaments.

It is also instructive to ask why filaments are a recurring motif across living systems. Not only do single units admit complex shape deformations, filaments can also easily self-assemble to form higher-order structures, or be reconfigured into networks of increasing complexity. These active physical structures have the capacity to perform complex functions without sophisticated control or centralised computation [315, 316]. Future applications, particularly in the realm of synthetic biology or bioengineering, may seek to harness the mechanical intelligence inherent in filamentous structures, to create novel adaptive materials.

tures of a penicillium, with special reference to their use in the isolation of B. influenzae," Br. J. Exp. Pathol. **10**, 226 (1929).

- [13] Alexander Fleming, "Penicillin," Brit. Med. J. 2, 386 (1941).
- [14] Jos Houbraken, Jens C Frisvad, and Robert A Samson, "Fleming's penicillin producing strain is not *Penicillium chrysogenum* but *P. rubens*," IMA Fungus 2, 87–95 (2011).
- [15] Karen Alim, Gabriel Amselem, François Peaudecerf, Michael P Brenner, and Anne Pringle, "Random network peristalsis in physarum polycephalum organizes fluid flows across an individual," Proc. Natl. Acad. Sci. USA **110**, 13306–13311 (2013).
- [16] Mirna Kramar and Karen Alim, "Encoding memory in tube diameter hierarchy of living flow network," Proc. Natl. Acad. Sci. USA 118, e2007815118 (2021).
- [17] Richard K Grosberg and Richard R Strathmann, "The evolution of multicellularity: a minor major transition?" Annu. Rev. Ecol. Evol. Syst. 38, 621–654 (2007).
- [18] Dennis Claessen, Daniel E Rozen, Oscar P Kuipers, Lotte Søgaard-Andersen, and Gilles P Van Wezel, "Bacterial solutions to multicellularity: a tale of biofilms, filaments and fruiting bodies," Nat. Rev. Microbiol. **12**, 115–124 (2014).
- [19] Hugh C Crenshaw and Leah Edelstein-Keshet, "Orientation by helical motion-II. Changing the direction of the axis of motion," Bull. Math. Biol. 55, 213–230 (1993).
- [20] Benjamin M Friedrich and Frank Jülicher, "Steering chiral swimmers along noisy helical paths," Phys. Rev. Lett. 103, 068102 (2009).
- [21] Klaas Bente, Sarah Mohammadinejad, Mohammad Avalin Charsooghi, Felix Bachmann, Agnese Codutti, Christopher T Lefèvre, Stefan Klumpp, and Damien Faivre, "High-speed motility originates from cooperatively pushing and pulling flagella bundles in bilophotrichous bacteria," eLife 9, e47551 (2020).
- [22] Jan F Jikeli, Luis Alvarez, Benjamin M Friedrich, Lau-

rence G Wilson, René Pascal, Remy Colin, Magdalena Pichlo, Andreas Rennhack, Christoph Brenker, and U Benjamin Kaupp, "Sperm navigation along helical paths in 3d chemoattractant landscapes," Nat. Commun. **6**, 7985 (2015).

- [23] Benjamin M Friedrich and Frank Jülicher, "Chemotaxis of sperm cells," Proc. Natl. Acad. Sci. USA 104, 13256– 13261 (2007).
- [24] Francisco J Sevilla, "Diffusion of active chiral particles," Phys. Rev. E 94, 062120 (2016).
- [25] Leon Lettermann, Falko Ziebert, and Ulrich S Schwarz, "A geometrical theory of gliding motility based on cell shape and surface flow," Proc. Natl. Acad. Sci. USA 121, e2410708121 (2024).
- [26] A. Deblais, S. Woutersen, and D. Bonn, "Rheology of entangled active polymer-like *T. Tubifex* worms," Phys. Rev. Lett. **124**, 188002 (2020).
- [27] Vishal P Patil, Harry Tuazon, Emily Kaufman, Tuhin Chakrabortty, David Qin, Jörn Dunkel, and M Saad Bhamla, "Ultrafast reversible self-assembly of living tangled matter," Science **380**, 392–398 (2023).
- [28] Philip W Anderson, "More is Different: Broken symmetry and the nature of the hierarchical structure of science." Science 177, 393–396 (1972).
- [29] Baudouin Saintyves, Matthew Spenko, and Heinrich M Jaeger, "A self-organizing robotic aggregate using solid and liquid-like collective states," Sci. Robot. 9, eadh4130 (2024).
- [30] Magdalena Fladung, Alexander Berkes, Tim Alletzhaeusser, Yi Chen, Natalie Munding, Motomu Tanaka, Martin Wegener, and Martin Bastmeyer, "What lies beyond-insights into elastic microscaffolds with metamaterial properties for cell studies," Curr. Opin. Biomed. Eng., 100568 (2024).
- [31] Mohammadreza Taale, Målin Schmidt, Fereydoon Taheri, Michael Timmermann, and Christine Selhuber-Unkel, "A minimalistic, synthetic cell-inspired metamaterial for enabling reversible strain-stiffening," Adv. Mater. Technol. 8, 2201441 (2023).
- [32] Nikta Fakhri, Frederick C MacKintosh, Brahim Lounis, Laurent Cognet, and Matteo Pasquali, "Brownian motion of stiff filaments in a crowded environment," Science 330, 1804–1807 (2010).
- [33] Nikta Fakhri, Dmitri A Tsyboulski, Laurent Cognet, R Bruce Weisman, and Matteo Pasquali, "Diameterdependent bending dynamics of single-walled carbon nanotubes in liquids," Proc. Natl. Acad. Sci. USA 106, 14219–14223 (2009).
- [34] Samantha Stam, Simon L Freedman, Shiladitya Banerjee, Kimberly L Weirich, Aaron R Dinner, and Margaret L Gardel, "Filament rigidity and connectivity tune the deformation modes of active biopolymer networks," Proc. Natl. Acad. Sci. USA 114, E10037–E10045 (2017).
- [35] Shiladitya Banerjee, Margaret L Gardel, and Ulrich S Schwarz, "The actin cytoskeleton as an active adaptive material," Annu. Rev. Condens. Matter Phys. 11, 421– 439 (2020).
- [36] Florian Huber, Jörg Schnauß, Susanne Rönicke, Philipp Rauch, Karla Müller, Claus Fütterer, and Josef Käs, "Emergent complexity of the cytoskeleton: from single filaments to tissue," Adv. Phys. 62, 1–112 (2013).
- [37] Daniel A Fletcher and R Dyche Mullins, "Cell mechanics and the cytoskeleton," Nature 463, 485–492 (2010).
- [38] Adrian F Pegoraro, Paul Janmey, and David A Weitz,

"Mechanical properties of the cytoskeleton and cells," Cold Spring Harb. Perspect. Biol. **9**, a022038 (2017).

- [39] Philip Kollmannsberger and Ben Fabry, "Linear and nonlinear rheology of living cells," Annu. Rev. Mater. Res. 41, 75–97 (2011).
- [40] Lev Davidovich Landau, L P Pitaevskii, Arnol'd Markovich Kosevich, and Evgenii Mikhailovich Lifshitz, *Theory of elasticity: volume 7*, Vol. 7 (Elsevier, 2012).
- [41] Otto Kratky and Günther Porod, "Röntgenuntersuchung gelöster fadenmoleküle," Rec. Trav. Chim. 68, 1106–1122 (1949).
- [42] Ömer Civalek and Çiğdem Demir, "Bending analysis of microtubules using nonlocal Euler–Bernoulli beam theory," Appl. Math. Model. 35, 2053–2067 (2011).
- [43] Ondřej Kučera, Daniel Havelka, and Michal Cifra, "Vibrations of microtubules: Physics that has not met biology yet," Wave Motion 72, 13–22 (2017).
- [44] Mehrdad Mehrbod and Mohammad R K Mofrad, "On the significance of microtubule flexural behavior in cytoskeletal mechanics," PLoS One 6, e25627 (2011).
- [45] Carla Pallavicini, Alejandro Monastra, Nicolás González Bardeci, Diana Wetzler, Valeria Levi, and Luciana Bruno, "Characterization of microtubule buckling in living cells," Eur. Biophys. J. 46, 581–594 (2017).
- [46] Klaus Kroy and Erwin Frey, "Force-extension relation and plateau modulus for wormlike chains," Phys. Rev. Lett. 77, 306 (1996).
- [47] Paul A Janmey, Ursula Euteneuer, Peter Traub, and Manfred Schliwa, "Viscoelastic properties of vimentin compared with other filamentous biopolymer networks." J. Cell Biol. **113**, 155–160 (1991).
- [48] David H Boal, Mechanics of the Cell (Cambridge University Press, 2012).
- [49] Mohammad R K Mofrad, "Rheology of the cytoskeleton," Annu. Rev. Fluid Mech. 41, 433–453 (2009).
- [50] Frederick Gittes, Brian Mickey, Jilda Nettleton, and Jonathon Howard, "Flexural rigidity of microtubules and actin filaments measured from thermal fluctuations in shape." J. Cell Bio. **120**, 923–934 (1993).
- [51] Jean-Philippe Collet, Henry Shuman, Robert E Ledger, Seungtaek Lee, and John W Weisel, "The elasticity of an individual fibrin fiber in a clot," Proc. Natl. Acad. Sci. USA **102**, 9133–9137 (2005).
- [52] Nadine Walter, Tobias Busch, Thomas Seufferlein, and Joachim P Spatz, "Elastic moduli of living epithelial pancreatic cancer cells and their skeletonized keratin intermediate filament network," Biointerphases 6, 79–85 (2011).
- [53] Douglas S Fudge, Kenn H Gardner, V Trevor Forsyth, Christian Riekel, and John M Gosline, "The mechanical properties of hydrated intermediate filaments: insights from hagfish slime threads," Biophys. J. 85, 2015–2027 (2003).
- [54] Joanna McKittrick, P-Y Chen, S G Bodde, W Yang, E E Novitskaya, and M A Meyers, "The structure, functions, and mechanical properties of keratin," JOM 64, 449–468 (2012).
- [55] Abhimanyu Sharma and Michael Vershinin, "Length dependence of the rigidity of microtubules in small networks," Biochem. Biophys. Res. Commun. 529, 303–305 (2020).
- [56] Takashi Nishino, Kiyofumi Takano, and Katsuhiko

Nakamae, "Elastic modulus of the crystalline regions of cellulose polymorphs," J. Polym. Sci. B: Polym. Phys. **33**, 1647–1651 (1995).

- [57] Xinhui Shen, Phu N Tran, Benjamin Z Tay, et al., "Bending stiffness characterization of bacillus subtilis" flagellar filament," Biophysical Journal **121**, 1975–1985 (2022).
- [58] Makoto Okuno and Yukio Hiramoto, "Direct measurements of the stiffness of echinoderm sperm flagella," J. Exp. Biol. **79**, 235–243 (1979).
- [59] Sumio Ishijima and Yukio Hiramoto, "Flexural rigidity of echinoderm sperm flagella," Cell Structure and Function 19, 349–362 (1994).
- [60] Mixon K Faluweki and Lucas Goehring, "Structural mechanics of filamentous cyanobacteria," Journal of the Royal Society Interface 19, 20220268 (2022).
- [61] David Boal and Ray Ng, "Shape analysis of filamentous precambrian microfossils and modern cyanobacteria," Paleobiology 36, 555–572 (2010).
- [62] Maximilian Kurjahn, Antaran Deka, Antoine Girot, Leila Abbaspour, Stefan Klumpp, Maike Lorenz, Oliver Bäumchen, and Stefan Karpitschka, "Quantifying gliding forces of filamentous cyanobacteria by selfbuckling," eLife 12, RP87450 (2024).
- [63] Matilda Backholm, William S Ryu, and Kari Dalnoki-Veress, "Viscoelastic properties of the nematode caenorhabditis elegans, a self-similar, shearthinning worm," Proc. Natl. Acad. Sci. USA 110, 4528– 4533 (2013).
- [64] Zengcai V Guo and Lakshminarayanan Mahadevan, "Limbless undulatory propulsion on land," Proc. Natl. Acad. Sci. USA 105, 3179–3184 (2008).
- [65] Xiaotian Zhang, Noel Naughton, Tejaswin Parthasarathy, and Mattia Gazzola, "Friction modulation in limbless, three-dimensional gaits and heterogeneous terrains," Nat. Commun. 12, 6076 (2021).
- [66] John H Long Jr, "Muscles, elastic energy, and the dynamics of body stiffness in swimming eels," Am. Zool. 38, 771–792 (1998).
- [67] Gary J Brouhard and Luke M Rice, "Microtubule dynamics: an interplay of biochemistry and mechanics," Nat. Rev. Mol. Cell Biol. 19, 451–463 (2018).
- [68] Ju Zhou, Anhui Wang, Yinlong Song, Nan Liu, Jia Wang, Yan Li, Xin Liang, Guohui Li, Huiying Chu, and Hong-Wei Wang, "Structural insights into the mechanism of gtp initiation of microtubule assembly," Nat. Commun. 14, 5980 (2023).
- [69] Pierre Dustin, *Microtubules* (Springer Science & Business Media, 2012).
- [70] Juliette Azimzadeh and Wallace F Marshall, "Building the centriole," Curr. Biol. 20, R816–R825 (2010).
- [71] A V Grimstone and A Klug, "Observations on the substructure of flagellar fibres," J. Cell Sci. 1, 351–362 (1966).
- [72] Linda A Amos and A Klug, "Arrangement of subunits in flagellar microtubules," J. Cell Sci. 14, 523–549 (1974).
- [73] Marileen Dogterom and Stanislas Leibler, "Physical aspects of the growth and regulation of microtubule structures," Phys. Rev. Lett. 70, 1347 (1993).
- [74] Deborah Kuchnir Fygenson, Erez Braun, and Albert Libchaber, "Phase diagram of microtubules," Phys. Rev. E 50, 1579 (1994).
- [75] Tim Mitchison and Marc Kirschner, "Dynamic insta-

bility of microtubule growth," Nature **312**, 237–242 (1984).

- [76] Murray D Weingarten, Arthur H Lockwood, Shu-Ying Hwo, and Marc W Kirschner, "A protein factor essential for microtubule assembly." Proc. Natl. Acad. Sci. USA 72, 1858–1862 (1975).
- [77] Sebastian Palmqvist, Shorena Janelidze, Yakeel T Quiroz, Henrik Zetterberg, Francisco Lopera, Erik Stomrud, YI Su, Yinghua Chen, Geidy E Serrano, Antoine Leuzy, et al., "Discriminative accuracy of plasma phospho-tau217 for Alzheimer disease vs other neurodegenerative disorders," JAMA **324**, 772–781 (2020).
- [78] C Y Chen, Y H Tseng, and J P Ward, "A mathematical model on the propagation of tau pathology in neurodegenerative diseases," J. Math. Biol. 89, 4 (2024).
- [79] Arshad Desai and Timothy J Mitchison, "Microtubule polymerization dynamics," Annu. Rev. Cell Dev. Biol. 13, 83–117 (1997).
- [80] Albert L Lehninger, David L Nelson, and Michael M Cox, *Principles of Biochemistry*. (New York: Worth, 1993) 2nd ed.
- [81] Martine Coue, Vivian A Lombillo, and J Richard McIntosh, "Microtubule depolymerization promotes particle and chromosome movement in vitro," J. Cell. Biol. 112, 1165–1175 (1991).
- [82] Douglas E Koshland, T J Mitchison, and Marc W Kirschner, "Polewards chromosome movement driven by microtubule depolymerization in vitro," Nature 331, 499–504 (1988).
- [83] Torsten Wittmann, Anthony Hyman, and Arshad Desai, "The spindle: a dynamic assembly of microtubules and motors," Nat. Cell Biol. 3, E28–E34 (2001).
- [84] Sharat Gadde and Rebecca Heald, "Mechanisms and molecules of the mitotic spindle," Curr. Biol. 14, R797– R805 (2004).
- [85] Marc Kirschner and Tim Mitchison, "Beyond selfassembly: from microtubules to morphogenesis," Cell 45, 329–342 (1986).
- [86] Emiliano Lepore, Marco Isaia, Stefano Mammola, and Nicola Pugno, "The effect of ageing on the mechanical properties of the silk of the bridge spider *Larinioides cornutus* (Clerck, 1757)," Sci. Rep. 6, 24699 (2016).
- [87] Gabriele Greco, Hamideh Mirbaha, Benjamin Schmuck, Anna Rising, and Nicola M Pugno, "Artificial and natural silk materials have high mechanical property variability regardless of sample size," Sci. Rep. 12, 3507 (2022).
- [88] Stephen A Wainwright, Mechanical design in organisms (Princeton University Press, 1982).
- [89] Richard H C Bonser and Peter P Purslow, "The Young's modulus of feather keratin," J. Exp. Biol. 198, 1029– 1033 (1995).
- [90] M Amanda Hartman and James A Spudich, "The myosin superfamily at a glance," J. Cell Sci. 125, 1627– 1632 (2012).
- [91] Florian Odronitz and Martin Kollmar, "Drawing the tree of eukaryotic life based on the analysis of 2,269 manually annotated myosins from 328 species," Genome Biol. 8, 1–23 (2007).
- [92] H Lee Sweeney and Anne Houdusse, "Structural and functional insights into the myosin motor mechanism," Annu. Rev. Biophys. **39**, 539–557 (2010).
- [93] Miguel Vicente-Manzanares, Xuefei Ma, Robert S Adelstein, and Alan Rick Horwitz, "Non-muscle myosin ii

takes centre stage in cell adhesion and migration," Nat. Rev. Mol. Cell Biol. **10**, 778–790 (2009).

- [94] Tina Carvalho, "CIL:219," CIL Dataset, https://doi. org/doi:10.7295/W9CIL219 (2011).
- [95] David Becker, "CIL:39004," CIL Dataset, https:// doi.org/doi:10.7295/W9CIL39004 (2011).
- [96] Jocelyn Étienne, Jonathan Fouchard, Démosthene Mitrossilis, Nathalie Bufi, Pauline Durand-Smet, and Atef Asnacios, "Cells as liquid motors: Mechanosensitivity emerges from collective dynamics of actomyosin cortex," Proc. Natl. Acad. Sci. USA **112**, 2740–2745 (2015).
- [97] Arnab Saha, Masatoshi Nishikawa, Martin Behrndt, Carl-Philipp Heisenberg, Frank Jülicher, and Stephan W Grill, "Determining physical properties of the cell cortex," Biophys. J. 110, 1421–1429 (2016).
- [98] K Vijay Kumar, "The actomyosin cortex of cells: a thin film of active matter," J. Indian Inst. Sci. 101, 97–112 (2021).
- [99] Luca Cardamone, Alessandro Laio, Vincent Torre, Rajesh Shahapure, and Antonio DeSimone, "Cytoskeletal actin networks in motile cells are critically selforganized systems synchronized by mechanical interactions," Proc. Natl. Acad. Sci. USA 108, 13978–13983 (2011).
- [100] Ulrich S. Schwarz and Samuel A. Safran, "Physics of adherent cells," Rev. Mod. Phys. 85, 1327–1381 (2013).
- [101] Simon Garcia, Edouard Hannezo, Jens Elgeti, Jean-François Joanny, Pascal Silberzan, and Nir S Gov, "Physics of active jamming during collective cellular motion in a monolayer," Proc. Natl. Acad. Sci. USA 112, 15314–15319 (2015).
- [102] Tuomas P Knowles, Anthony W Fitzpatrick, Sarah Meehan, Helen R Mott, Michele Vendruscolo, Christopher M Dobson, and Mark E Welland, "Role of intermolecular forces in defining material properties of protein nanofibrils," Science **318**, 1900–1903 (2007).
- [103] Carl-Philipp Heisenberg and Yohanns Bellaïche, "Forces in tissue morphogenesis and patterning," Cell 153, 948– 962 (2013).
- [104] Benoît Aigouy, Reza Farhadifar, Douglas B Staple, Andreas Sagner, Jens-Christian Röper, Frank Jülicher, and Suzanne Eaton, "Cell flow reorients the axis of planar polarity in the wing epithelium of drosophila," Cell 142, 773–786 (2010).
- [105] D Bray and J G White, "Cortical flow in animal cells," Science 239, 883–888 (1988).
- [106] Sarah Köster, David A Weitz, Robert D Goldman, Ueli Aebi, and Harald Herrmann, "Intermediate filament mechanics in vitro and in the cell: from coiled coils to filaments, fibers and networks," Curr. Opin. Struct. Biol. **32**, 82–91 (2015).
- [107] Michael Schopferer, Harald Bär, Bernhard Hochstein, Sarika Sharma, Norbert Mücke, Harald Herrmann, and Norbert Willenbacher, "Desmin and vimentin intermediate filament networks: their viscoelastic properties investigated by mechanical rheometry," J. Mol. Biol. 388, 133–143 (2009).
- [108] Bernd Nöding and Sarah Köster, "Intermediate filaments in small configuration spaces," Phys. Rev. Lett. 108, 088101 (2012).
- [109] David A D Parry and Peter M Steinert, "Intermedi-

ate filament structure," Curr. Opin. Cell Biol. 4, 94–98 (1992).

- [110] Anastasia A Chernyatina, Dmytro Guzenko, and Sergei V Strelkov, "Intermediate filament structure: the bottom-up approach," Curr. Opin. Cell Biol. **32**, 65–72 (2015).
- [111] Harald Herrmann and Ueli Aebi, "Intermediate filaments: structure and assembly," Cold Spring Harb. Perspect. Biol. 8, a018242 (2016).
- [112] Qi Wen and Paul A Janmey, "Polymer physics of the cytoskeleton," Curr. Opin. Solid State Mater. Sci. 15, 177–182 (2011).
- [113] Andreas Kerst, Craig Chmielewski, Celeste Livesay, Robert E Buxbaum, and Steven R Heidemann, "Liquid crystal domains and thixotropy of filamentous actin suspensions." Proc. Natl. Acad. Sci. USA 87, 4241–4245 (1990).
- [114] Atsushi Suzuki, Masahito Yamazaki, and Tadanao Ito, "Osmoelastic coupling in biological structures: formation of parallel bundles of actin filaments in a crystalline-like structure caused by osmotic stress," Biochemistry 28, 6513–6518 (1989).
- [115] A L Hitt, A R Cross, and R C Williams Jr, "Microtubule solutions display nematic liquid crystalline structure." J. Biol. Chem. 265, 1639–1647 (1990).
- [116] Jorge Viamontes, Patrick W. Oakes, and Jay X. Tang, "Isotropic to nematic liquid crystalline phase transition of *f*-actin varies from continuous to first order," Phys. Rev. Lett. **97**, 118103 (2006).
- [117] H Hess and Jennifer L Ross, "Non-equilibrium assembly of microtubules: from molecules to autonomous chemical robots," Chem. Soc. Rev. 46, 5570–5587 (2017).
- [118] Lars Onsager, "The effects of shape on the interaction of colloidal particles," Ann. N. Y. Acad. Sci. 51, 627–659 (1949).
- [119] Paul J Flory, "Phase equilibria in solutions of rod-like particles," Proc. R. Soc. Lond. A 234, 73–89 (1956).
- [120] Bernd Wagner, Rainer Tharmann, Ilka Haase, Markus Fischer, and Andreas R Bausch, "Cytoskeletal polymer networks: the molecular structure of cross-linkers determines macroscopic properties," Proc. Natl. Acad. Sci. USA **103**, 13974–13978 (2006).
- [121] Margaret L Gardel, Karen E Kasza, Clifford P Brangwynne, Jiayu Liu, and David A Weitz, "Mechanical response of cytoskeletal networks," Methods Cell Biol. 89, 487–519 (2008).
- [122] Thomas D Pollard and Gary G Borisy, "Cellular motility driven by assembly and disassembly of actin filaments," Cell 112, 453–465 (2003).
- [123] Cornelis Storm, Jennifer J Pastore, Fred C MacKintosh, Tom C Lubensky, and Paul A Janmey, "Nonlinear elasticity in biological gels," Nature 435, 191–194 (2005).
- [124] Loïc Chaubet, Abdullah R Chaudhary, Hossein K Heris, Allen J Ehrlicher, and Adam G Hendricks, "Dynamic actin cross-linking governs the cytoplasm's transition to fluid-like behavior," Mol. Biol. Cell **31**, 1744–1752 (2020).
- [125] C Lorenz and Sarah Köster, "Multiscale architecture: Mechanics of composite cytoskeletal networks," Biophys. Rev. 3 (2022).
- [126] Ovijit Chaudhuri, Sapun H Parekh, and Daniel A Fletcher, "Reversible stress softening of actin networks," Nature 445, 295–298 (2007).
- [127] Paul A Janmey, Margaret E McCormick, Sebastian

Rammensee, Jennifer L Leight, Penelope C Georges, and Fred C MacKintosh, "Negative normal stress in semiflexible biopolymer gels," Nature Mater. **6**, 48–51 (2007).

- [128] Bo Sun, "The mechanics of fibrillar collagen extracellular matrix," Cell Rep. Phys. Sci. 2 (2021).
- [129] Ming-Thau Sheu, Ju-Chun Huang, Geng-Chang Yeh, and Hsiu-O Ho, "Characterization of collagen gel solutions and collagen matrices for cell culture," Biomater. 22, 1713–1719 (2001).
- [130] Pavel Dutov, Olga Antipova, Sameer Varma, Joseph P R O Orgel, and Jay D Schieber, "Measurement of elastic modulus of collagen type i single fiber," PloS One 11, e0145711 (2016).
- [131] S J Eichhorn and R J Young, "The young's modulus of a microcrystalline cellulose," Cellulose 8, 197–207 (2001).
- [132] Y-C Hsieh, H Yano, M Nogi, and S J Eichhorn, "An estimation of the young's modulus of bacterial cellulose filaments," Cellulose 15, 507–513 (2008).
- [133] Teresa E Strzelecka, Michael W Davidson, and Randolph L Rill, "Multiple liquid crystal phases of DNA at high concentrations," Nature 331, 457–460 (1988).
- [134] Graham Ramsay, "DNA chips: state-of-the art," Nat. Biotechnol. 16, 40–44 (1998).
- [135] Nadrian C Seeman, "Nanomaterials based on DNA," Annu. Rev. Biochem. 79, 65–87 (2010).
- [136] Barbara Saccà and Christof M Niemeyer, "DNA origami: the art of folding DNA," Angew. Chem. Int. Ed. 51, 58–66 (2012).
- [137] Nadrian C Seeman and Hanadi F Sleiman, "DNA nanotechnology," Nat. Rev. Mater. 3, 1–23 (2017).
- [138] Florian Praetorius, Benjamin Kick, Karl L Behler, Maximilian N Honemann, Dirk Weuster-Botz, and Hendrik Dietz, "Biotechnological mass production of DNA origami," Nature 552, 84–87 (2017).
- [139] Wouter Engelen and Hendrik Dietz, "Advancing biophysics using DNA origami," Annu. Rev. Biophys. 50, 469–492 (2021).
- [140] Alexei A Kornyshev, Dominic J Lee, Sergey Leikin, and Aaron Wynveen, "Structure and interactions of biological helices," Rev. Mod. Phys. 79, 943–996 (2007).
- [141] Gerard C L Wong and Lois Pollack, "Electrostatics of strongly charged biological polymers: ion-mediated interactions and self-organization in nucleic acids and proteins," Annu. Rev. Phys. Chem. 61, 171–189 (2010).
- [142] Vladimir B Teif and Klemen Bohinc, "Condensed DNA: condensing the concepts," Prog. Biophys. Mol. Biol. 105, 208–222 (2011).
- [143] Andrew Travers and Georgi Muskhelishvili, "DNA structure and function," FEBS J. 282, 2279–2295 (2015).
- [144] Davide Michieletto and Nick Gilbert, "Role of nuclear RNA in regulating chromatin structure and transcription," Curr. Opin. Cell Biol. 58, 120–125 (2019).
- [145] Dalibor Chevizovich, Davide Michieletto, Alain Mvogo, Farit Zakiryanov, and Slobodan Zdravković, "A review on nonlinear DNA physics," R. Soc. Open Sci. 7, 200774 (2020).
- [146] Davide Michieletto, "Kinetoplast DNA: a polymer physicist's topological Olympic dream," Nucleic Acids Res., gkae1206 (2024).
- [147] Clifford Dobell, Antony van Leeuwenhoek and his "Little Animals." (Russell & Russell New York, 1958).
- [148] Leah T Haimo and Joel L Rosenbaum, "Cilia, flagella,

and microtubules." J. Cell Biol. **91**, 125s–130s (1981).

- [149] Morgan Beeby, Josie L Ferreira, Patrick Tripp, Sonja-Verena Albers, and David R Mitchell, "Propulsive nanomachines: the convergent evolution of archaella, flagella and cilia," FEMS Microbiol. Rev. 44, 253–304 (2020).
- [150] Karen Grace Bondoc-Naumovitz, Hannah Laeverenz-Schlogelhofer, Rebecca N Poon, Alexander K Boggon, Samuel A Bentley, Dario Cortese, and Kirsty Y Wan, "Methods and measures for investigating microscale motility," Integrative and Comparative Biology 63, 1485–1508 (2023).
- [151] Gregor Eichele, Eberhard Bodenschatz, Zuzana Ditte, Ann-Kathrin Günther, Shoba Kapoor, Yong Wang, and Christian Westendorf, "Cilia-driven flows in the brain third ventricle," Philosophical Transactions of the Royal Society B 375, 20190154 (2020).
- [152] Brendan K Huang and Michael A Choma, "Microscale imaging of cilia-driven fluid flow," Cellular and molecular life sciences 72, 1095–1113 (2015).
- [153] Feng Ling, Tara Essock-Burns, Margaret McFall-Ngai, Kakani Katija, Janna C Nawroth, and Eva Kanso, "Flow physics guides morphology of ciliated organs," Nature Physics 20, 1679–1686 (2024).
- [154] Wallace F Marshall and Shigenori Nonaka, "Cilia: tuning in to the cell's antenna," Current Biology 16, R604– R614 (2006).
- [155] Rita R. Ferreira, Hajime Fukui, Renee Chow, Andrej Vilfan, and Julien Vermot, "The cilium as a force sensor- myth versus reality," Journal of cell science 132, jcs213496 (2019).
- [156] Sangjin Ryu and Paul Matsudaira, "Unsteady motion, finite reynolds numbers, and wall effect on vorticella convallaria contribute contraction force greater than the stokes drag," Biophys. J. 98, 2574–2581 (2010).
- [157] Ken F Jarrell, Yan Ding, Divya B Nair, and Sarah Siu, "Surface appendages of archaea: structure, function, genetics and assembly," Life 3, 86–117 (2013).
- [158] E. M. Purcell, "Life at low Reynolds number," Am. J. Phys. 45, 3–11 (1977).
- [159] Gregory J Hancock, "The self-propulsion of microscopic organisms through liquids," Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences 217, 96–121 (1953).
- [160] Kirsty Y Wan and Rebecca N Poon, "Mechanisms and functions of multiciliary coordination," Curr. Opin. Cell Biol. 86, 102286 (2024).
- [161] Kirsty Y Wan, "Coordination of eukaryotic cilia and flagella," Essays Biochem. 62, 829–838 (2018).
- [162] Nariya Uchida, Ramin Golestanian, and Rachel R Bennett, "Synchronization and collective dynamics of flagella and cilia as hydrodynamically coupled oscillators," J. Phys. Soc. Jpn. 86, 101007 (2017).
- [163] Nicolas Bruot and Pietro Cicuta, "Realizing the physics of motile cilia synchronization with driven colloids," Annu. Rev. Condens. Matter Phys. 7, 323–348 (2016).
- [164] Rebecca N Poon, Timothy A Westwood, Hannah Laeverenz-Schlogelhofer, Emelie Brodrick, Jamie Craggs, Eric E Keaveny, Gáspár Jékely, and Kirsty Y Wan, "Ciliary propulsion and metachronal coordination in reef coral larvae," Phys. Rev. Res. 5, L042037 (2023).
- [165] Fanlong Meng, Rachel R Bennett, Nariya Uchida, and Ramin Golestanian, "Conditions for metachronal coordination in arrays of model cilia," Proceedings of the

National Academy of Sciences  $\mathbf{118}$ ,  $\mathbf{e2102828118}$  (2021).

- [166] Jens Elgeti and Gerhard Gompper, "Emergence of metachronal waves in cilia arrays," Proc. Natl. Acad. Sci. USA 110, 4470–4475 (2013).
- [167] Marianne Grognot and Katja M Taute, "More than propellers: how flagella shape bacterial motility behaviors," Current Opinion in Microbiology 61, 73–81 (2021).
- [168] Navish Wadhwa and Howard C Berg, "Bacterial motility: machinery and mechanisms," Nat. Rev. Microbiol. 20, 161–173 (2022).
- [169] Hannah Laeverenz-Schlogelhofer and Kirsty Y Wan, "Bioelectric control of locomotor gaits in the walking ciliate euplotes," Current Biology 34, 697–709 (2024).
- [170] Kei Jokura, Yu Sato, Kogiku Shiba, and Kazuo Inaba, "Two distinct compartments of a ctenophore comb plate provide structural and functional integrity for the motility of giant multicilia," Current Biology **32**, 5144–5152 (2022).
- [171] Avraham Be'er and Gil Ariel, "A statistical physics view of swarming bacteria," Movement ecology 7, 1–17 (2019).
- [172] Martin A Bees, "Advances in bioconvection," Annual Review of Fluid Mechanics 52, 449–476 (2020).
- [173] Fabian Jan Schwarzendahl and Marco G Mazza, "Maximum in density heterogeneities of active swimmers," Soft Matter 14, 4666–4678 (2018).
- [174] Jérémy Vachier and Marco G Mazza, "Dynamics of sedimenting active brownian particles," Eur. Phys. J. E 42, 1–11 (2019).
- [175] Fabian Jan Schwarzendahl and Marco G Mazza, "Hydrodynamic interactions dominate the structure of active swimmers' pair distribution functions," J. Chem. Phys. 150 (2019).
- [176] Fabian Jan Schwarzendahl and Marco G Mazza, "Active percolation in pusher-type microswimmers," EPL 140, 47001 (2022).
- [177] Linda Turner, William S Ryu, and Howard C Berg, "Real-time imaging of fluorescent flagellar filaments," J. Bacteriol. 182, 2793–2801 (2000).
- [178] Alexander Neuhaus, Muniyandi Selvaraj, Ralf Salzer, Julian D Langer, Kerstin Kruse, Lennart Kirchner, Kelly Sanders, Bertram Daum, Beate Averhoff, and Vicki AM Gold, "Cryo-electron microscopy reveals two distinct type iv pili assembled by the same bacterium," Nature communications 11, 2231 (2020).
- [179] Mami Nomura, Kohei Atsuji, Keiko Hirose, Kogiku Shiba, Ryuji Yanase, Takeshi Nakayama, Ken-ichiro Ishida, and Kazuo Inaba, "Microtubule stabilizer reveals requirement of Ca2+-dependent conformational changes of microtubules for rapid coiling of haptonema in haptophyte algae," Biology Open 8, bio036590 (2019).
- [180] Michael Silverman and Melvin I Simon, "Bacterial flagella," Ann. Rev. Microbiol. 31, 397–419 (1977).
- [181] Daniel Coombs, Greg Huber, John O Kessler, and Raymond E Goldstein, "Periodic chirality transformations propagating on bacterial flagella," Phys. Rev. Lett. 89, 118102 (2002).
- [182] Virginija Cvirkaite-Krupovic, Ying Liu, Diana Baquero, Junfeng Liu, Ravi R Sonani, Chris R Calladine, Fengbin Wang, Mart Krupovic, and Edward Egelman, "The evolution of archaeal flagellar filaments," Proc. Natl. Acad. Sci. USA **120**, e2304256120 (2023).
- [183] Stefan Streif, Wilfried Franz Staudinger, Wolfgang Mar-

wan, and Dieter Oesterhelt, "Flagellar rotation in the archaeon halobacterium salinarum depends on atp," Journal of molecular biology **384**, 1–8 (2008).

- [184] Sonja-Verena Albers and Ken F Jarrell, "The archaellum: how archaea swim," Frontiers in Microbiology 6, 23 (2015).
- [185] Howard C Berg, "The rotary motor of bacterial flagella," Annual review of biochemistry 72, 19–54 (2003).
- [186] Michael D Manson, Pat Tedesco, Howard C Berg, Franklin M Harold, and Chris Van der Drift, "A protonmotive force drives bacterial flagella." Proceedings of the National Academy of Sciences 74, 3060–3064 (1977).
- [187] Daniela Nicastro, Cindi Schwartz, Jason Pierson, Richard Gaudette, Mary E Porter, and J Richard McIntosh, "The molecular architecture of axonemes revealed by cryoelectron tomography," Science **313**, 944– 948 (2006).
- [188] Charles B Lindemann and Kathleen A Lesich, "Flagellar and ciliary beating: the proven and the possible," Journal of cell science **123**, 519–528 (2010).
- [189] Peter Satir, Thomas Heuser, and Winfield S Sale, "A structural basis for how motile cilia beat," Bioscience 64, 1073–1083 (2014).
- [190] Ingmar H Riedel-Kruse, Andreas Hilfinger, Jonathon Howard, and Frank Jülicher, "How molecular motors shape the flagellar beat," HFSP journal 1, 192–208 (2007).
- [191] James F Cass and Hermes Bloomfield-Gadêlha, "The reaction-diffusion basis of animated patterns in eukaryotic flagella," Nat. Commun. 14, 5638 (2023).
- [192] Louis G Woodhams, Yenan Shen, and Philip V Bayly, "Generation of ciliary beating by steady dynein activity: the effects of inter-filament coupling in multi-filament models," J. R. Soc. Interface 19, 20220264 (2022).
- [193] Roger Eckert, "Bioelectric control of ciliary activity: Locomotion in the ciliated protozoa is regulated by membrane-limited calcium fluxes." Science 176, 473– 481 (1972).
- [194] Kirsty Y Wan and Raymond E Goldstein, "Time irreversibility and criticality in the motility of a flagellate microorganism," Phys. Rev. Lett. **121**, 058103 (2018).
- [195] Michael L Ginger, Neil Portman, and Paul G McKean, "Swimming with protists: perception, motility and flagellum assembly," Nature Reviews Microbiology 6, 838–850 (2008).
- [196] Gerasimos Langousis and Kent L Hill, "Motility and more: the flagellum of *Trypanosoma brucei*," Nat. Rev. Microbiol. **12**, 505–518 (2014).
- [197] Sei Suzuki-Tellier, Federica Miano, Seyed Saeed Asadzadeh, Alastair G B Simpson, and Thomas Kiørboe, "Foraging mechanisms in excavate flagellates shed light on the functional ecology of early eukaryotes," Proc. Natl. Acad. Sci. USA **121**, e2317264121 (2024).
- [198] Lisa Craig, Katrina T. Forest, and Berenike Maier, "Type IV pili: dynamics, biophysics and functional consequences," Nat. Rev. Microbiol. **17**, 429–440 (2019), number: 7 Publisher: Nature Publishing Group.
- [199] Courtney K Ellison, Gregory B Whitfield, and Yves V Brun, "Type iv pili: dynamic bacterial nanomachines," FEMS Microbiology Reviews 46, fuab053 (2022).
- [200] L. Mahadevan and P. Matsudaira, "Motility Powered by Supramolecular Springs and Ratchets," Science 288, 95–99 (2000), publisher: American Association for the Advancement of Science.

- [201] Sangjin Ryu, Rachel E. Pepper, Moeto Nagai, and Danielle C. France, "Vorticella: A Protozoan for Bio-Inspired Engineering," Micromachines 8, 4 (2017).
- [202] Carlos Floyd, Arthur T Molines, Xiangting Lei, Jerry E Honts, Fred Chang, Mary Williard Elting, Suriyanarayanan Vaikuntanathan, Aaron R Dinner, and M Saad Bhamla, "A unified model for the dynamics of atp-independent ultrafast contraction," Proc. Natl. Acad. Sci. USA **120**, e2217737120 (2023).
- [203] M. Kawachi and I. Inouye, "Ca2+-mediated induction of the coiling of the haptonema in Chrysochromulina hirta (Prymnesiophyta = Haptophyta)," Phycologia 33, 53–57 (1994).
- [204] Nicholas C Darnton and Howard C Berg, "Forceextension measurements on bacterial flagella: triggering polymorphic transformations," Biophys. J. 92, 2230– 2236 (2007).
- [205] Gang Xu, Kate S Wilson, Ruth J Okamoto, Jin-Yu Shao, Susan K Dutcher, and Philip V Bayly, "Flexural rigidity and shear stiffness of flagella estimated from induced bends and counterbends," Biophys. J. 110, 2759– 2768 (2016).
- [206] David B Hill, Vinay Swaminathan, Ashley Estes, Jeremy Cribb, E Timothy O'Brien, C William Davis, and R Superfine, "Force generation and dynamics of individual cilia under external loading," Biophys. J. 98, 57–66 (2010).
- [207] MEJ Holwill, "The motion of strigomonas oncopelti," Journal of Experimental Biology 42, 125–137 (1965).
- [208] Shoji A Baba, "Flexural rigidity and elastic constant of cilia," J. Exp. Biol. 56, 459–467 (1972).
- [209] David Hickey, Andrej Vilfan, and Ramin Golestanian, "Ciliary chemosensitivity is enhanced by cilium geometry and motility," Elife 10, e66322 (2021).
- [210] Robert A Bloodgood, "Sensory reception is an attribute of both primary cilia and motile cilia," J. Cell Sci. 123, 505–509 (2010).
- [211] Culture Collection of Algae, SAG, "Medium recipie vers. 10.2008," http://sagdb.uni-goettingen. de/culture\_media/01%20Basal%20Medium.pdf.
- [212] Xingting Gong and Manu Prakash, "Active dislocations and topological traps govern dynamics of spiraling filamentous cyanobacteria," arXiv preprint arXiv:2305.12572 (2023).
- [213] Ferran Garcia-Pichel and Martin F Wojciechowski, "The evolution of a capacity to build supra-cellular ropes enabled filamentous cyanobacteria to colonize highly erodible substrates," PloS one 4, e7801 (2009).
- [214] Mixon K Faluweki, Jan Cammann, Marco G Mazza, and Lucas Goehring, "Active spaghetti: collective organization in cyanobacteria," Phys. Rev. Lett. 131, 158303 (2023).
- [215] Maximilian Kurjahn, Leila Abbaspour, Franziska Papenfuß, Philip Bittihn, Ramin Golestanian, Benoît Mahault, and Stefan Karpitschka, "Collective selfcaging of active filaments in virtual confinement," arXiv preprint arXiv:2403.03093 (2024).
- [216] Jan Cammann, Mixon K Faluweki, Nayara Dambacher, Lucas Goehring, and Marco G Mazza, "Topological transition in filamentous cyanobacteria: from motion to structure," Commun. Phys. 7, 376 (2024).
- [217] Kurt M Ehlers, A D Samuel, Howard C Berg, and Richard Montgomery, "Do cyanobacteria swim using traveling surface waves?" Proc. Natl. Acad. Sci. USA

**93**, 8340–8343 (1996).

- [218] Zvonimir Dogic and Seth Fraden, "Smectic phase in a colloidal suspension of semiflexible virus particles," Phys. Rev. Lett. 78, 2417 (1997).
- [219] Zvonimir Dogic and Seth Fraden, "Ordered phases of filamentous viruses," Curr. Opin. Colloid Interface Sci. 11, 47–55 (2006).
- [220] Kirstin R Purdy and Seth Fraden, "Isotropic-cholesteric phase transition of filamentous virus suspensions as a function of rod length and charge," Phys. Rev. E 70, 061703 (2004).
- [221] Alejandro Bilbao, Eligiusz Wajnryb, Siva A Vanapalli, and Jerzy Blawzdziewicz, "Nematode locomotion in unconfined and confined fluids," Phys. Fluids 25 (2013).
- [222] Sean Gart, Dominic Vella, and Sunghwan Jung, "The collective motion of nematodes in a thin liquid layer," Soft Matter 7, 2444–2448 (2011).
- [223] Ernst Niebur and Paul Erdös, "Theory of the locomotion of nematodes: dynamics of undulatory progression on a surface," Biophysical journal 60, 1132–1146 (1991).
- [224] José-Agustín Arguedas-Leiva, Jonasz Słomka, Cristian C Lalescu, Roman Stocker, and Michael Wilczek, "Elongation enhances encounter rates between phytoplankton in turbulence," Proc. Natl. Acad. Sci. USA 119, e2203191119 (2022).
- [225] Jakob Pernthaler, "Predation on prokaryotes in the water column and its ecological implications," Nat. Rev. Microbiol. 3, 537–546 (2005).
- [226] Kevin D Young, "The selective value of bacterial shape," Microbiol. Mol. Biol. Rev. 70, 660–703 (2006).
- [227] Thomas H Weiss, Aaron L Mills, George M Hornberger, and Janet S Herman, "Effect of bacterial cell shape on transport of bacteria in porous media," Environ. Sci. Technol. 29, 1737–1740 (1995).
- [228] E Hoiczyk, "Gliding motility in cyanobacteria: observations and possible explanations," Archives of microbiology 174, 11–17 (2000).
- [229] Annegret Wilde and Conrad W Mullineaux, "Motility in cyanobacteria: polysaccharide tracks and Type IV pilus motors," Mol. Microbiol. 98, 998–1001 (2015).
- [230] Joël Tchoufag, Pushpita Ghosh, Connor B Pogue, Beiyan Nan, and Kranthi K Mandadapu, "Mechanisms for bacterial gliding motility on soft substrates," Proc. Natl. Acad. Sci. USA **116**, 25087–25096 (2019).
- [231] Mark J McBride, "Bacterial gliding motility: multiple mechanisms for cell movement over surfaces," Annu. Rev. Microbiol. 55, 49–75 (2001).
- [232] Nicholas Read, Simon Connell, and David G Adams, "Nanoscale visualization of a fibrillar array in the cell wall of filamentous cyanobacteria and its implications for gliding motility," J. Bacteriol. 189, 7361–7366 (2007).
- [233] Samia Dhahri, Michel Ramonda, and Christian Marliere, "In-situ determination of the mechanical properties of gliding or non-motile bacteria by atomic force microscopy under physiological conditions without immobilization," PLoS One 8, e61663 (2013).
- [234] Jair Koiller, Kurt M Ehlers, and Fabio Chalub, "Acoustic streaming, the "small invention" of cyanobacteria?" Arbor 186, 1089 (2010).
- [235] Lawrence N Halfen and Richard W Castenholz, "Gliding in a blue–green alga: a possible mechanism," Nature 225, 1163–1165 (1970).
- [236] Behzad Khayatan, John C Meeks, and Douglas D

Risser, "Evidence that a modified type IV pilus-like system powers gliding motility and polysaccharide secretion in filamentous cyanobacteria," Mol. Microbiol. **98**, 1021–1036 (2015).

- [237] EGBERT Hoiczyk and Wolfgang Baumeister, "Envelope structure of four gliding filamentous cyanobacteria," J. Bacteriol. 177, 2387–2395 (1995).
- [238] Egbert Hoiczyk and Wolfgang Baumeister, "The junctional pore complex, a prokaryotic secretion organelle, is the molecular motor underlying gliding motility in cyanobacteria," Curr. Biol. 8, 1161–1168 (1998).
- [239] Charles W Wolgemuth and George Oster, "The junctional pore complex and the propulsion of bacterial cells," J. Mol. Microbiol. Biotechnol. 7, 72–77 (2004).
- [240] Kurt M Ehlers and Jair Koiller, "Could cell membranes produce acoustic streaming? making the case for synechococcus self-propulsion," Math. Comput. Model. 53, 1489–1504 (2011).
- [241] Guangle Du, Sunita Kumari, Fangfu Ye, and Rudolf Podgornik, "Model of metameric locomotion in smooth active directional filaments with curvature fluctuations," EPL 136, 58003 (2022).
- [242] Jerko Rosko, Kelsey Cremin, Emanuele Locatelli, Mary Coates, Sarah JN Duxbury, Kieran Randall, Katie Croft, Chantal Valeriani, Marco Polin, Orkun S Soyer, *et al.*, "Cellular (de) coordination in gliding motility and plectoneme formation," bioRxiv (2024).
- [243] Leila Abbaspour, Ali Malek, Stefan Karpitschka, and Stefan Klumpp, "Effects of direction reversals on patterns of active filaments," Physical Review Research 5, 013171 (2023).
- [244] Ulrike Pfreundt, Jonasz Słomka, Giulia Schneider, Anupam Sengupta, Francesco Carrara, Vicente Fernandez, Martin Ackermann, and Roman Stocker, "Controlled motility in the cyanobacterium trichodesmium regulates aggregate architecture," Science **380**, 830–835 (2023).
- [245] Özer Duman, Rolf E Isele-Holder, Jens Elgeti, and Gerhard Gompper, "Collective dynamics of self-propelled semiflexible filaments," Soft matter 14, 4483–4494 (2018).
- [246] K R Prathyusha, Silke Henkes, and Rastko Sknepnek, "Dynamically generated patterns in dense suspensions of active filaments," Phys. Rev. E 97, 022606 (2018).
- [247] Rolf E Isele-Holder, Jens Elgeti, and Gerhard Gompper, "Self-propelled worm-like filaments: spontaneous spiral formation, structure, and dynamics," Soft matter 11, 7181–7190 (2015).
- [248] Szu-Ning Lin, Wei-Chang Lo, and Chien-Jung Lo, "Dynamics of self-organized rotating spiral-coils in bacterial swarms," Soft Matter 10, 760–766 (2014).
- [249] Wilhelm Nultsch and Klaus Wenderoth, "Partial irradiation experiments with anabaena variabilis (kütz)." Zeitschrift für Pflanzenphysiologie 111, 1–7 (1983).
- [250] Annegret Wilde and Conrad W Mullineaux, "Lightcontrolled motility in prokaryotes and the problem of directional light perception," FEMS Microbiol. Rev. 41, 900–922 (2017).
- [251] Christian Pfeffer, Steffen Larsen, Jie Song, Mingdong Dong, Flemming Besenbacher, Rikke Louise Meyer, Kasper Urup Kjeldsen, Lars Schreiber, Yuri A Gorby, Mohamed Y El-Naggar, *et al.*, "Filamentous bacteria transport electrons over centimetre distances," Nature **491**, 218–221 (2012).
- [252] Ugo Marzocchi, Daniela Trojan, Steffen Larsen, Rikke

Louise Meyer, Niels Peter Revsbech, Andreas Schramm, Lars Peter Nielsen, and Nils Risgaard-Petersen, "Electric coupling between distant nitrate reduction and sulfide oxidation in marine sediment," The ISME journal 8, 1682–1690 (2014).

- [253] Lars Peter Nielsen, Nils Risgaard-Petersen, Henrik Fossing, Peter Bondo Christensen, and Mikio Sayama, "Electric currents couple spatially separated biogeochemical processes in marine sediment," Nature 463, 1071–1074 (2010).
- [254] Jesper J Bjerg, Jamie J M Lustermans, Ian P G Marshall, Anna J Mueller, Signe Brokjær, Casper A Thorup, Paula Tataru, Markus Schmid, Michael Wagner, Lars Peter Nielsen, and Andreas Schramm, "Cable bacteria with electric connection to oxygen attract flocks of diverse bacteria," Nat. Commun. 14, 1614 (2023).
- [255] Nils Risgaard-Petersen, Michael Kristiansen, Rasmus B Frederiksen, Anders Lindequist Dittmer, Jesper Tataru Bjerg, Daniela Trojan, Lars Schreiber, Lars Riis Damgaard, Andreas Schramm, and Lars Peter Nielsen, "Cable bacteria in freshwater sediments," Appl. Environ. Microbiol. 81, 6003–6011 (2015).
- [256] Gemma Reguera, Kevin D McCarthy, Teena Mehta, Julie S Nicoll, Mark T Tuominen, and Derek R Lovley, "Extracellular electron transfer via microbial nanowires," Nature 435, 1098–1101 (2005).
- [257] Mohamed Y El-Naggar, Greg Wanger, Kar Man Leung, Thomas D Yuzvinsky, Gordon Southam, Jun Yang, Woon Ming Lau, Kenneth H Nealson, and Yuri A Gorby, "Electrical transport along bacterial nanowires from Shewanella oneidensis MR-1," Proc. Natl. Acad. Sci. USA **107**, 18127–18131 (2010).
- [258] Sahand Pirbadian, Sarah E Barchinger, Kar Man Leung, Hye Suk Byun, Yamini Jangir, Rachida A Bouhenni, Samantha B Reed, Margaret F Romine, Daad A Saffarini, Liang Shi, et al., "Shewanella oneidensis MR-1 nanowires are outer membrane and periplasmic extensions of the extracellular electron transport components," Proc. Natl. Acad. Sci. USA 111, 12883– 12888 (2014).
- [259] Gal Schkolnik, Matthias Schmidt, Marco G Mazza, Falk Harnisch, and Niculina Musat, "In situ analysis of a silver nanoparticle-precipitating shewanella biofilm by surface enhanced confocal Raman microscopy," PLoS One 10, e0145871 (2015).
- [260] Bruce E Logan, Ruggero Rossi, Ala'a Ragab, and Pascal E Saikaly, "Electroactive microorganisms in bioelectrochemical systems," Nat. Rev. Microbiol. 17, 307–319 (2019).
- [261] Zahra Mokhtari and Annette Zippelius, "Dynamics of active filaments in porous media," Phys. Rev. Lett. 123, 028001 (2019).
- [262] Harry D Kurtz Jr and Dennis I Netoff, "Stabilization of friable sandstone surfaces in a desiccating, windabraded environment of south-central utah by rock surface microorganisms," Journal of Arid Environments 48, 89–100 (2001).
- [263] J L A Pluis, "Algal crust formation in the inland dune area, Laarder Wasmeer, The Netherlands," Vegetatio 113, 41–51 (1994).
- [264] Sebastian Gonzalez La Corte, Corey A. Stevens, Gerardo Cárcamo-Oyarce, Katharina Ribbeck, Ned S. Wingreen, and Sujit S. Datta, "Morphogenesis of bacterial cables in polymeric environments," Science Ad-

vances **11**, eadq7797 (2025).

- [265] RN Shepard and DY Sumner, "Undirected motility of filamentous cyanobacteria produces reticulate mats," Geobiology 8, 179–190 (2010).
- [266] Carlos Tamulonis and Jaap Kaandorp, "A model of filamentous cyanobacteria leading to reticulate pattern formation," Life 4, 433–456 (2014).
- [267] Diana G Cuadrado and Jerónimo Pan, "Field observations on the evolution of reticulate patterns in microbial mats in a modern siliciclastic coastal environment," Journal of Sedimentary Research 88, 24–37 (2018).
- [268] Marco G Mazza, "The physics of biofilms-an introduction," J. Phys. D: Appl. Phys. 49, 203001 (2016).
- [269] Karin Sauer, Paul Stoodley, Darla M Goeres, Luanne Hall-Stoodley, Mette Burmølle, Philip S Stewart, and Thomas Bjarnsholt, "The biofilm life cycle: expanding the conceptual model of biofilm formation," Nat. Rev. Microbiol. 20, 608–620 (2022).
- [270] Antonio C Costa, Tosif Ahamed, David Jordan, and Greg J Stephens, "A markovian dynamics for caenorhabditis elegans behavior across scales," Proc. Natl. Acad. Sci. USA **121**, e2318805121 (2024).
- [271] Greg J. Stephens, Bethany Johnson-Kerner, William Bialek, and William S. Ryu, "Dimensionality and dynamics in the behavior of c. elegans," PLoS Comput. Biol. 4, e1000028 (2008).
- [272] Gary B. Gillis, "Environmental effects on undulatory locomotion in the american eel anguilla rostrata: Kinematics in water and on land," J. Exp. Biol. 201, 949–961 (1998).
- [273] Bruce C Jayne, "What defines different modes of snake locomotion?" Integr. Comp. Biol. 60, 156–170 (2020).
- [274] Sten Grillner, Peter Wallen, Lennart Brodin, and Anders Lansner, "Neuronal network generating locomotor behavior in lamprey: Circuitry, transmitters, membrane properties, and simulation," Annu. Rev. Neurosci. 14, 169–199 (1991).
- [275] Paul S. Katz, "Evolution of central pattern generators and rhythmic behaviours," Philos. Trans. R. Soc. B 371, 20150057 (2016).
- [276] Pavan Ramdya and Auke Jan Ijspeert, "The neuromechanics of animal locomotion: From biology to robotics and back," Sci. Robot. 8, eadg0279 (2023).
- [277] Ross L Hatton and Howie Choset, "Geometric motion planning: The local connection, stokes' theorem, and the importance of coordinate choice," Int. J. Robot. Res. 30, 988–1014 (2011).
- [278] Tingnan Zhang and Daniel I. Goldman, "The effectiveness of resistive force theory in granular locomotion," Phys. Fluids 26, 101308 (2014).
- [279] Venkat Padmanabhan, Zeina S Khan, Deepak E Solomon, Andrew Armstrong, Kendra P Rumbaugh, Siva A Vanapalli, and Jerzy Blawzdziewicz, "Locomotion of C. elegans: a piecewise-harmonic curvature representation of nematode behavior," PloS One 7, e40121 (2012).
- [280] Jennifer M Rieser, Tai-De Li, Jessica L Tingle, Daniel I Goldman, and Joseph R Mendelson III, "Functional consequences of convergently evolved microscopic skin features on snake locomotion," Proc. Natl. Acad. Sci. USA 118, e2018264118 (2021).
- [281] Anton Peshkov, Sonia McGaffigan, and Alice C Quillen, "Synchronized oscillations in swarms of nematode turbatrix aceti," Soft Matter 18, 1174–1182 (2022).

- [282] Xiaodong Wu and Shugen Ma, "Cpg-based control of serpentine locomotion of a snake-like robot," Mechatronics 20, 326–334 (2010).
- [283] Eviatar Yemini, Tadas Jucikas, Laura J Grundy, André E X Brown, and William R Schafer, "A database of caenorhabditis elegans behavioral phenotypes," Nat. Methods 10, 877–879 (2013).
- [284] Jennifer M. Rieser, Baxi Chong, Chaohui Gong, Henry C. Astley, Perrin E. Schiebel, Kelimar Diaz, Christopher J. Pierce, Hang Lu, Ross L. Hatton, Howie Choset, and Daniel I. Goldman, "Geometric phase predicts locomotion performance in undulating living systems across scales," Proc. Natl. Acad. Sci. USA 121, e2320517121 (2024).
- [285] Christopher J Pierce, Daniel Irvine, Lucinda Peng, Xuefei Lu, Hang Lu, and Daniel I Goldman, "Dispersion relations for active undulators in overdamped environments," arxiv.org/abs/2407.13037, arXiv-2407 (2024).
- [286] Jesús Sánchez-Rodríguez, Christophe Raufaste, and Médéric Argentina, "Scaling the tail beat frequency and swimming speed in underwater undulatory swimming," Nat. Commun. 14, 5569 (2023).
- [287] Yasemin Ozkan-Aydin, Daniel I Goldman, and M Saad Bhamla, "Collective dynamics in entangled worm and robot blobs," Proc. Natl. Acad. Sci. USA 118, e2010542118 (2021).
- [288] Tess Heeremans, Antoine Deblais, Daniel Bonn, and Sander Woutersen, "Chromatographic separation of active polymer–like worm mixtures by contour length and activity," Sci. Adv. 8, eabj7918 (2022).
- [289] AC Quillen, A Peshkov, Esteban Wright, and Sonia McGaffigan, "Metachronal waves in concentrations of swimming turbatrix aceti nematodes and an oscillator chain model for their coordinated motions," Phys. Rev. E 104, 014412 (2021).
- [290] Antoine Deblais, K. R. Prathyusha, Rosa Sinaasappel, Harry Tuazon, Ishant Tiwari, Vishal P. Patil, and M. Saad Bhamla, "Worm blobs as entangled living polymers: from topological active matter to flexible soft robot collectives," Soft Matter 19, 7057–7069 (2023).
- [291] Geoffrey Taylor, "Analysis of the swimming of long and narrow animals," Proc. R. Soc. A 214, 158–183 (1952).
- [292] Vincent Stin, Ramiro Godoy-Diana, Xavier Bonnet, and Anthony Herrel, "Form and function of anguilliform swimming," Biol. Rev. Camb. Philos. Soc. 99, 2190–2210 (2024).
- [293] Isaac J Yeaton, Shane D Ross, Grant A Baumgardner, and John J Socha, "Undulation enables gliding in flying snakes," Nat. Phys. 16, 974–982 (2020).
- [294] Ellen Zheng, Martin Brandenbourger, Louis Robinet, Peter Schall, Edan Lerner, and Corentin Coulais, "Selfoscillation and synchronization transitions in elastoactive structures," Phys. Rev. Lett. 130, 178202 (2023).
- [295] Gabriele De Canio, Eric Lauga, and Raymond E Goldstein, "Spontaneous oscillations of elastic filaments induced by molecular motors," J. R. Soc. Interface. 14, 20170491 (2017).
- [296] P V Bayly and S K Dutcher, "Steady dynein forces induce flutter instability and propagating waves in mathematical models of flagella," J. R. Soc. Interface. 13, 20160523 (2016).
- [297] Yi Man, Feng Ling, and Eva Kanso, "Cilia oscillations," Philos. Trans. R. Soc. B. 375, 20190157 (2020).
- [298] Nigel R Franks, Alan Worley, Katherine AJ Grant, Al-

ice R Gorman, Victoria Vizard, Harriet Plackett, Carolina Doran, Margaret L Gamble, Martin C Stumpe, and Ana B Sendova-Franks, "Social behaviour and collective motion in plant-animal worms," Proc. R. Soc. B **283**, 20152946 (2016).

- [299] Roland G Winkler and Gerhard Gompper, "The physics of active polymers and filaments," J. Chem. Phys. 153 (2020).
- [300] Amin Doostmohammadi, Jordi Ignés-Mullol, Julia M Yeomans, and Francesc Sagués, "Active nematics," Nat. Commun. 9, 3246 (2018).
- [301] T B Liverpool, A C Maggs, and A Ajdari, "Viscoelasticity of solutions of motile polymers," Physical review letters 86, 4171 (2001).
- [302] Valentino Bianco, Emanuele Locatelli, and Paolo Malgaretti, "Globulelike conformation and enhanced diffusion of active polymers," Phys. Rev. Lett. **121**, 217802 (2018).
- [303] Gerard A Vliegenthart, Arvind Ravichandran, Marisol Ripoll, Thorsten Auth, and Gerhard Gompper, "Filamentous active matter: Band formation, bending, buckling, and defects," Sci. Adv. 6, eaaw9975 (2020).
- [304] F J Nédélec, Thomas Surrey, Anthony C Maggs, and Stanislas Leibler, "Self-organization of microtubules and motors," Nature **389**, 305–308 (1997).
- [305] Tim Sanchez, Daniel T N Chen, Stephen J DeCamp, Michael Heymann, and Zvonimir Dogic, "Spontaneous motion in hierarchically assembled active matter," Nature 491, 431–434 (2012).
- [306] Karsten Kruse, Jean-François Joanny, Frank Jülicher, Jacques Prost, and Ken Sekimoto, "Asters, vortices, and rotating spirals in active gels of polar filaments," Phys. Rev. Lett. 92, 078101 (2004).
- [307] Yutaka Sumino, Ken H Nagai, Yuji Shitaka, Dan

Tanaka, Kenichi Yoshikawa, Hugues Chaté, and Kazuhiro Oiwa, "Large-scale vortex lattice emerging from collectively moving microtubules," Nature **483**, 448–452 (2012).

- [308] L Huber, R Suzuki, T Krüger, E Frey, and A R Bausch, "Emergence of coexisting ordered states in active matter systems," Science 361, 255–258 (2018).
- [309] Alfredo Sciortino and Andreas R Bausch, "Pattern formation and polarity sorting of driven actin filaments on lipid membranes," Proc. Natl. Acad. Sci. USA 118, e2017047118 (2021).
- [310] Leonid M Martyushev and Vladimir D Seleznev, "Maximum entropy production principle in physics, chemistry and biology," Phys. Rep. 426, 1–45 (2006).
- [311] Paul CW Davies, Elisabeth Rieper, and Jack A Tuszynski, "Self-organization and entropy reduction in a living cell," Biosyst. 111, 1–10 (2013).
- [312] Christoph Adami, Charles Ofria, and Travis C Collier, "Evolution of biological complexity," Proc. Natl. Acad. Sci. USA 97, 4463–4468 (2000).
- [313] Christoph Adami, "What is complexity?" BioEssays 24, 1085–1094 (2002).
- [314] Danail Bonchev and Gregory A Buck, "Quantitative Measures of Network Complexity," in *Complexity in Chemistry, Biology, and Ecology* (Springer, 2005) pp. 191–235.
- [315] Tianyu Wang, Christopher Pierce, Velin Kojouharov, Baxi Chong, Kelimar Diaz, Hang Lu, and Daniel I. Goldman, "Mechanical intelligence simplifies control in terrestrial limbless locomotion," Sci. Robot. 8, eadi2243 (2023).
- [316] Clément Moreau, Benjamin J Walker, Rebecca N Poon, Daniel Soto, Daniel I Goldman, Eamonn A Gaffney, and Kirsty Y Wan, "Minimal design of a synthetic cilium," Phys. Rev. Res. 6, L042061 (2024).