# Fundamental constraints on quantum fluctuations: Conservation laws, reality, and no-signaling

Thales A. B. Pinto Silva\* and David Gelbwaser-Klimovsky Schulich Faculty of Chemistry and Helen Diller Quantum Center, Technion-Israel Institute of Technology, Haifa 3200003, Israel

Quantum fluctuations and noise are fundamental in quantum technologies, affecting computing, sensing, cryptography, and thermodynamics. These include fluctuations in the variation of energy, charge, and other observables driven by interactions with lasers, amplifiers, and baths. Despite the precise rules quantum mechanics provides for measuring observables at single points in time, no standard framework exists for characterizing the fluctuations of their variations over time. This gap not only makes physical conclusions dependent on the chosen measurement protocol but also leads to inconsistencies in fluctuation predictions, impacting quantum technologies. We propose four basic criteria that any consistent measurement of these variations must satisfy, grounded in conservation laws, the no-signaling principle, and expected constraints on physical realism. We demonstrate that only one protocol fulfills all these criteria: the two-times quantum observables. This result enables the extension of key quantum information concepts, such as entanglement, steering, and Bell's inequalities, to processes rather than instantaneous observables. Beyond resolving ambiguities in quantum fluctuation measurements, our framework offers a foundation for improved fluctuation control in quantum devices, with potential applications in quantum computing, metrology, and thermodynamics.

arXiv:2502.12905v1 [quant-ph] 18 Feb 2025

Quantum fluctuations play an increasingly critical role in emerging devices [1], challenging the performance of quantum computers [2, 3], sensors [4], and cryptographic systems [5]. Central to control and predict these fluctuations is the ability to accurately characterize the *variation* of physical quantities (VPQs). For instance, in superconducting quantum computers, minimizing the impact of charge fluctuation is essential for ensuring reliable qubit performance [2, 3]. Likewise, in quantum thermodynamics, the statistical characterization of VPQs, such as variation of energy and particle number variations, is vital for refining fluctuation theorems and extending thermodynamic and conservation laws into the quantum realm [1, 6–13]. Remarkably, despite significant advancements, *there remains no universal and standard framework for characterizing the statistics of VPQs in quantum mechanics.* 

To illustrate this challenge, consider the simple task of describing the statistics of the variation of a quantum observable O for an arbitrary system S evolving under unitary dynamics  $U_t$  until an arbitrary time t. Quantum mechanics provides a standard protocol for measuring O at any time for any given initial quantum state  $\rho$ : by considering the eigenstates  $|o_n\rangle$  of O, we can define the probability of obtaining a measurement outcome  $o_k$  at time t as  $p(o_k, \rho, t) = \langle o_k | U_t \rho U_t^{\dagger} | o_k \rangle$ .  $p(o_k, \rho, t)$ fully characterizes the statistics of O at any given time. On the other hand, quantum mechanics provides no standard method for calculating the statistics of the variation of O over the interval [0, t]. This is because the variation of O is a quantity non-local in time, and it is unclear how to measure it without the measurement itself disturbing it. For instance, a common approach to measure such variations is the two-point measurement (TPM) protocol [12-17] (see Fig 1. for a theoretical example in trapped ions). Based on TPM, the system S, initially in state  $\rho$ , undergoes a projective measurement of O at t = 0, yielding an outcome  $o_n$ . After being measured, the system evolves under  $U_t$  until t, where a second measurement of O is done resulting in  $o_m$  (See the top of Figs. 1a and 1b). The variation  $\Delta o_{mn} = o_m - o_n$  is computed for this arbitrary run,

and repeating this protocol many times allows us to construct the probability of the variation of *O* of having any arbitrary value.

Although intuitive, the TPM approach has several drawbacks. Perhaps the most important one is that the initial measurement collapses any superpositions in  $\rho$  with respect to the eigenbasis of O. This can lead TPM to fail to respect conservation laws [15, 16, 19], as demonstrated in Fig. 1 and the "Methods" section. These and other limitations [20, 21] have prompted the development of alternative methodologies for describing the statistics of VPQs, including the well-known full-counting statistics [22], quasi-probabilities [23], Gaussian pointers [24], and two-time observables [19–21, 25–27], among others [16]. These approaches, however, often produce conflicting results, raising critical concerns for both foundational physics and practical applications. Indeed, because they provide contradictory predictions, it is unclear which approach, if any, accurately reflects the underlying physics. Also, in the absence of a standardized protocol, physical conclusions risk being influenced more by the measurement procedure chosen itself than by the process under investigation (e.g., the example in Fig. 1). Furthermore, the lack of a reliable standard has critical implications for quantum technologies. Predictions must be accurate and consistent across diverse scenarios to ensure quantum devices' functionality and resource efficiency-faulty or overly demanding predictions can undermine their practical feasibility.

Without a universally accepted standard methodology, grounding protocols in fundamental principles of physics offers a clear and reliable criterion to ensure their consistency and applicability across diverse scenarios [10, 15, 20, 28–30]. Motivated by this, we propose four fundamental properties that any consistent protocol for measuring VPQs should satisfy ensuring conservation laws, the no-signaling principle, and expected constraints on physical realism. By assuming these criteria, we prove that there is a unique measurement protocol able to meet all principles simultaneously: the two-



FIG. 1. Violation of energy conservation in TPM protocol. The system  $\Omega$  consists of a trapped Ca<sup>+</sup> ion model inspired in the Ref. [18]. The total energy of the system is  $H = H_{\rm H0} + H_e$ .  $H_{\rm H0} = \hbar\omega(N + 1/2) \otimes \mathbb{1}_s$  is the energy related with the center of mass (CM). The position of the CM is coupled to the spin of a covalent electron (red arrow) through a spin-dependent optical dipole force (ODF), with a coupling energy approximately  $H_e \approx \hbar(\omega_z/2 + \Delta_S k_{\rm SW} X/2) \otimes \sigma_z$  [18]. One round of the TPM measurement of the variations of  $H_{\rm H0}$  and  $H_e$  are represented in the above part of figures **a** and **b**, respectively. After collecting the results of many rounds of experiments, we represent in the lower graphs the two highest values of the probabilities  $p_{\rm TPM}(z, H_i, U_\tau, \rho)$  of the energies  $H_i \in \{H_{\rm H0}, H_e\}$  of varying *z* during the evolution  $U_\tau$  for the initial state  $\rho$ . Energy conservation requires the increase of  $H_{\rm H0}$  to be equal to the decrease of  $H_e$ . However, **a** and **b** reveal that TPM predicts  $H_{\rm H0}$  to be less likely to increase by  $\hbar\omega$  than  $H_e$  to decrease by  $-\hbar\omega$ , violating energy conservation. The calculations are done in the "Methods" section, considering an initial state  $\rho = |0\rangle \langle 0| \otimes |+\rangle \langle +|$ , with  $N |0\rangle = 0 |0\rangle$  and  $\sigma_z |+\rangle = |+\rangle$ , and parameters:  $\frac{\omega_L}{1.4} = \frac{\Delta S}{1.3} = 2\pi$  MHz,  $k_{\rm SW} = 2\pi/(280 \,\mathrm{nm})$ , and  $m = 6.68 \times 10^{-26}$  kg. The interval considered to compute the variation is  $[0, \tau]$ , where  $\tau = \pi/\omega$ .

times quantum observable protocol [19–21, 25–27]. This protocol can thus serve as the necessary standard measurement protocol to compute VPQs.

In our framework, we consider a general system  $\Omega$ , comprising arbitrary subsystems, prepared in an arbitrary quantum state  $\rho$  acting on a Hilbert space  $\mathcal{H}$ .  $\Omega$  evolves from time 0 to *t* under an arbitrary unitary operator *U*, with a countable eigenbasis. We focus on describing the measurement of the variation of *a specific part* of the total energy of the whole system  $\Omega$ , described by a time-independent Hermitian operator  $H_1$  acting on  $\mathcal{H}$  (see examples in Fig. 2). Our framework and results, however, readily extend to the variation of any other quantum observable, such as particle number, momentum, or angular momentum.

We first clarify what we mean by a measurement protocol. In quantum mechanics, measurements can be generally described by positive operator-valued measures (POVMs) [31– 33]. Following this perspective, we define a *measurement protocol*  $\mathbb{M}$  [15] to measure the *variation* of any arbitrary energy operator  $H_1$  under an arbitrary evolution U. For the protocol  $\mathbb{M}$  and for each  $(H_1, U)$  pair, the set  $\mathbb{M}(H_1, U) = \{M(z, H_1, U)\}$  defines a POVM whose operators satisfy  $\int_{-\infty}^{\infty} dz M(z, H_1, U) = \mathbb{1}$  and  $M(z, H_1, U) \ge 0$ . In this framework, the probability density of observing a variation z of  $H_1$  under U is  $\wp(z, H_1, U, \rho) = \text{Tr}[M(z, H_1, U)\rho]$ , for the initial state  $\rho$ . By the POVM properties, it follows that  $\int_{-\infty}^{\infty} dz \wp(z, H_1, U, \rho) = 1$  and  $\wp(z, H_1, U, \rho) \ge 0$ . This approach aims to keep the protocol as general as possible, so that  $\mathbb{M}$ applies universally.

An important example of a measurement protocol is the two-times observables (OBS) protocol, denoted by  $M_{OBS}$ . This protocol operates as follows: for any given energy operator  $H_1$  and unitary evolution U over time t, the variation of energy is defined by the two-time quantum observable [19–21]:

$$\Delta(H_1, U) = U^{\dagger} H_1 U - H_1, \qquad (1)$$

representing the difference between the Heisenberg picture operators  $U^{\dagger}H_1U$  and  $H_1$  at times t and 0.  $\Delta(H_1, U)$  is a Hermitian operator, with eigenvalues  $\{\delta_j(H_1, U)\}$  and corresponding eigenvectors  $\{|\delta_j(H_1, U)\rangle\}$ , as expressed in the decomposition  $\Delta(H_1, U) = \sum_j \delta_j(H_1, U) |\delta_j(H_1, U)\rangle \langle \delta_j(H_1, U)|$ . The probability of finding a specific eigenvalue  $\delta_j(H_1, U)$  is  $p_j(H_1, U, \rho) = \text{Tr}[P_j(H_1, U)\rho]$ , where  $P_j(H_1, U) = |\delta_j(H_1, U)\rangle \langle \delta_j(H_1, U)|$  and  $\rho$  is the initial state. Accordingly, the OBS protocol  $\mathbb{M}_{OBS}$  is defined by POVMs  $\mathbb{M}_{OBS}(H_1, U) = \{M_{OBS}(z, H_1, U)\}$ , where each element is given by  $M_{OBS}(z, H_1, U) = \sum_j \delta^{D}[z - \delta_j(H_1, U)]P_j(H_1, U)$ , and  $\delta^{D}$  is the Dirac's delta [19–21]. Consequently, the probability den-



FIG. 2. Illustration of some possible scenarios within our framework. (a) A general subsystem S with energy  $H_S \otimes \mathbb{1}_E$  interacting through an interaction term  $V_{SE}$  with an environment E, whose energy is  $\mathbb{1}_{S} \otimes H_{E}$ . This is the common scenario in deriving quantum open system evolution equations. The total energy of  $\Omega$  composed of S and E is given by  $H = H_S \otimes \mathbb{1}_E + \mathbb{1}_S \otimes H_E + V_{SE}$ , and the quantity of interest is the variation of  $H_1 = H_S \otimes \mathbb{1}_E$ . (b) Quantum computing scenario, where each qubit *i* has energy  $H_{ai}$ . The unitary evolution is determined by gate operations  $U_1$ ,  $U_2$ , and  $U_3$ , rather than by  $\exp[-iHt/\hbar]$ , where *H* is the sum of qubit energies. The quantity of interest is the variation of  $H_1$ , which can be restricted, for example, to the change of the last two qubit energies (green bracket and arrow). (c) Many-body system scenario, where the evolution follows  $\exp[-iHt/\hbar]$ , with the total energy described by  $H = \sum_i \hbar \Delta_i n_i + \sum_{ij} V_{ij}$ . The quantity of interest is the variation of  $H_1$ , where  $H_1 = \hbar \Delta_k n_k$  is the energy localized at one of the chain sites. (d) Two interacting particles, with kinetic energies  $K_1$  and  $K_2$ , coupled through a position-dependent potential  $V(X_2 - X_1)$ . The total energy is  $H = K_1 + K_2 + V(X_2 - X_1)$  and the evolution is unitary  $U = \exp[-iHt/\hbar]$ . The quantity of interest can be the variation of  $H_1 = K_1$ . These examples are just some among infinitely many possibilities for  $H_1$ , U and systems  $\Omega$ .

sity

$$\wp_{\text{OBS}}(z, H_1, U, \rho) = \text{Tr}[M_{\text{OBS}}(z, H_1, U)\rho]$$
(2)

provides the likelihood of measuring a variation of  $H_1$  equal to z under evolution U. An illustration of how the OBS protocol can be considered for a trapped ion system is provided in Fig. 3.

Another widely-used example is TPM protocol  $\mathbb{M}_{\text{TPM}}$ , briefly described in the introduction. For any  $(H_1, U)$  pair, we define [14, 15]  $\mathbb{M}_{\text{TPM}}(H_1, U) = \{M_{\text{TPM}}(z, H_1, U)\}$ , where  $M_{\text{TPM}}(z, H_1, U) = \sum_{jk} \delta^{\mathbf{p}}[z - (e_j - e_k)]|\langle e_j| U | e_k \rangle|^2 | e_k \rangle \langle e_k |$ , with  $| e_j \rangle$  and  $| e_k \rangle$  eigenvectors of  $H_1$  having eigenvalues  $e_j$ and  $e_k$ , respectively. The probability density is thus given by  $\varphi_{\text{TPM}}(z, H_1, U, \rho) = \text{Tr}[M_{\text{TPM}}(z, H_1, U)\rho]$ .

We now define the four fundamental conditions that we expect any consistent protocol  $\mathbb{M}$  for measuring energy variations to satisfy.

1. Conservation laws: For any preparation  $\rho$ , unitary evolution U, and energy operators  $H_1$  and  $H_2$  representing parts of the energy of any system  $\Omega$ , if  $[H_1 + H_2, U] = 0$ , then, for any z,  $\text{Tr}[M(z, H_1, U)\rho] =$  $\wp(z, H_1, U, \rho) = \wp(-z, H_2, U, \rho) = \text{Tr}[M(-z, H_2, U)\rho].$ 



FIG. 3. Schematic implementation of the OBS protocol for the trapped Ca<sup>+</sup> ion system described in Fig. 1. The variation of the center-of-mass energy,  $H_1 = H_{\rm HO} \otimes \mathbb{1}_s = \hbar\omega(N + 1/2) \otimes \mathbb{1}_s$ , is inferred entirely from measurements of  $X \otimes \sigma_z$ , an observable that commutes with  $\Delta(H_1, U)$ , with  $U = U_{\tau}$ . This highlights how OBS protocol can be applied via commuting observables. We consider an initial preparation  $\rho = |\alpha, -\rangle \langle \alpha, -|$ , where  $|\alpha\rangle$  is the coherent state and  $\sigma_z |-\rangle = -|-\rangle$ . The left graph below the scheme shows the expected probability distribution  $\wp(\bar{x}, \rho)$  of measuring  $X \otimes \sigma_z$  at time 0 and finding  $\bar{x} = k_{\rm sw} x$  for the dimensionless position. Eq. (2) together with  $\wp(\bar{x}, \rho)$  can be used to obtain the OBS probability distribution  $\wp_{\rm OBS}(\bar{z}, H_1, U, \rho)$  for the dimensionless variation  $\bar{z} = z/\hbar\omega$  (see "Methods" section).  $\wp_{\rm OBS}(\bar{z}, H_1, U, \rho)$  is shown in the right graph as a normalized Gaussian centered at  $\langle \Delta(H_1, U) \rangle / (\hbar\omega)$ , with variance  $\sigma_{\Delta(H_1, U)} / \hbar\omega = \sqrt{\langle \Delta^2(H_1, U) \rangle - \langle \Delta(H_1, U) \rangle^2} / \hbar\omega$ .

In other words, if the sum of energies  $H_1 + H_2$  is conserved under U, then the probability of  $H_1$  of increasing an amount z must equal the probability of  $H_2$  of *decreasing* the same amount. This condition ensures that conservation laws hold at the level of probability distributions instead of just on average. In light of the Wigner-Araki-Yanase (WAY) theorem [6, 7], it is interesting to note that this condition allows for the possibility that the measurement process may, in specific rounds, disturb the subsystem's energy or even not be repeatable. However, even with such single-shot disturbances, we assume that conservation laws remain preserved, ensuring no statistically detectable violation in the energy balance (see Fig. 4**a**).

- 2. **Reality**: Consider any system  $\Omega$ , operator  $H_1$  and evolution U. If the initial state is  $\rho = |e\rangle \langle e|$  such that  $|e\rangle$  is an eigenvector of both  $H_1$  and  $U^{\dagger}H_1U$  with respective eigenvalues e and  $\epsilon$ , then the POVM must result in the probabilities  $\wp(z, H_1, U, \rho) = \delta^p[z (\epsilon e)]$  for this specific  $\rho = |e\rangle \langle e|$ . In other words, if  $H_1$  is well-defined (or real, in an EPR-sense [34]) at both the start and end of the process, the variation should be precisely the difference between the initial and final eigenvalues (see Fig. 4b).
- 3. Independence of the initial state: For any system  $\Omega$ , operators  $H_1$ , and evolution operators U, the elements



FIG. 4. Schematic illustration of the four key conditions for a consistent measurement protocol of VPQs. (a) Energy Conservation (Condition 1): If  $[H_1 + H_2, U] = 0$ , the total energy  $H_1 + H_2$  must be conserved. This means the measurement protocols  $\mathbb{M}(H_1, U)$  and  $\mathbb{M}(H_2, U)$  should satisfy the relation  $\wp(z, H_1, U, \rho) = \wp(-z, H_2, U, \rho)$ for the same initial state  $\rho$ . (b) Reality Condition (Condition 2): If the whole system is in an eigenstate of the energy  $H_1$  at times 0 and t, with eigenvalues e and  $\epsilon$ , the measurement distribution for the variation of  $H_1$  should collapse to a delta function  $\delta^{\mathbf{D}}[z - (\epsilon - e)]$ , reflecting a well-defined energy difference. (c) Independence of the state (Condition 3): The measurement protocol must not depend on the initial state  $\rho$ , ensuring that the apparatus is set independently of prior knowledge about the system. (d) No-signaling (Condition 4): The measurement protocol must ensure that a local evolution U'on one part of a bipartite system does not affect the energy variation statistics of  $H_1$  in the other part, preserving the no-signaling principle during the evolution  $U \otimes U'$ .

of the POVM  $M(z, H_1, U)$  must not depend on the initial state  $\rho$ . This condition is generally satisfied in quantum mechanics [31–33] and ensures that the measurement apparatus is not adjusted based on the initial preparation of the system [10, 15, 16, 20] (see Fig. 4c). Notably, this condition is equivalent to requiring that the elements { $M(z, H_1, U)$ } of the POVM  $\mathbb{M}(H_1, U)$  are linear with respect to  $\rho$  [15].

4. No-signaling: Consider a system Ω evolving under an arbitrary bipartide unitary evolution U ⊗ U' acting on a bipartide Hilbert space H = H ⊗ H'. For any such system and every energy operator H<sub>1</sub> ⊗ 𝔅<sub>H'</sub> acting locally on H, M is such that its POVM elements satisfy M(z, H<sub>1</sub> ⊗ 𝔅<sub>H'</sub>, U ⊗ U') = M(z, H<sub>1</sub>, U) ⊗ 𝔅<sub>H'</sub> for every z. In other words, local statistics of H<sub>1</sub> should remain unaffected by changes (U') in another subspace, ensuring that no statistically detectable information is transmitted between different subspaces via the measurement process [35, 36] (see Fig. 4d). This condition should be crucial in setups where additional auxiliary subsystems are introduced to assist measurements [24, 37].

A measurement protocol that satisfies all four conditions (1-4) is referred to as an CRIN protocol. We are now in position to establish the main result of our work:

**Result 1.** The OBS protocol is the only protocol that satisfies the CRIN conditions.

The proof of this result is presented in the "Methods" section and further detailed in the Supplementary Material (SM) [38]. Importantly, the result is not confined to energy variations–it extends to the variation of any quantum observable, such as linear and angular momentum, or particle number. Additionally, we demonstrate in the SM that the result holds for explicitly time-dependent observables, with the CRIN conditions appropriately adapted for such cases. This broad applicability highlights the universal relevance of the OBS protocol in quantum mechanics.

The implications of result 1 offer valuable insights. First, because OBS is the only CRIN protocol, in order to measure variations in energy, charge, particle position, or other observables in quantum devices, then we must treat these variations as two-time observables. Any deviation from the OBS protocol leads to the violation of at least one of the CRIN conditions. For example, as shown in Fig. 1, the TPM protocol fails to satisfy condition 1, implying that energy conservation is not statistically preserved.

Interestingly, since the statistical framework for two-time observables mirrors that of conventional "one-time" quantum observables, the CRIN conditions naturally extend standard quantum phenomena to two-time observables. For instance, by assuming the CRIN conditions, entanglement, steering, or quantum superposition can be considered within the scope of two-time quantum observables (see Refs. [19, 21] for a discussion). Furthermore, we can deduce a two-time uncertainty relation for any initial state  $\rho$  [19]:

$$\sigma_{\Delta(H_j,U)}(\sigma_{U^{\dagger}H_jU} + \sigma_{H_j}) \ge |\langle [U^{\dagger}H_jU, H_j] \rangle|, \qquad (3)$$

where  $j \in \{1, 2\}$ ,  $\sigma_o = \sqrt{\langle O^2 \rangle - \langle O \rangle^2}$  is the variance of an observable *O* and  $\langle O \rangle = \text{Tr}[O\rho]$  its expectation value. Remarkably, this leads to scenarios where the variation  $\Delta(H_j, U)$  can become perfectly defined (i.e.,  $\sigma_{\Delta(H_j, U)} \rightarrow 0$ ), even when the individual energies  $H_j$  at 0 and  $U^{\dagger}H_jU$  at *t* cannot be completely determinate (i.e.,  $\sigma_{U^{\dagger}H_jU} + \sigma_{H_j} \rightarrow \infty$ ). The trapped ion example in Fig. 5 illustrates this behavior.

As demonstrated in Fig. 3, the OBS protocol is derived by measuring an observable commuting with  $\Delta(H_1, U)$ , allowing us to explicitly determine  $\wp_{OBS}(z, H_1, U, \rho)$ . However, in many experimental scenarios, direct measurements of certain quantities are impractical, and instead, indirect measurement schemes employing auxiliary probes are used [17, 37]. A natural question arises: can the same  $\wp_{OBS}(z, H_1, U, \rho)$  be obtained through the use of a probe to measure the energy? Our second key result addresses this challenge:

**Result 2.** For any U,  $H_1$  acting on  $\mathcal{H}$ , and eigenstate  $|\delta_i(H_1, U)\rangle$  of  $\Delta(H_1, U)$ , there exists a unitary U' acting on an auxiliary Hilbert space  $\mathcal{H}'$ , an additional Hamiltonian  $H_2$ 



FIG. 5. The role of commutation in Eq. (3) for the trapped Ca<sup>+</sup> ion system. (a) For  $H_1 = H_{HO}$ , we plot the probabilities  $\langle n, -|\rho|n, -\rangle$ (blue squares) and  $\langle n, -| U\rho U^{\dagger} | n, - \rangle$  (red circles) of the energy  $H_1$ being  $E_n = (n+1/2)\hbar\omega$  at t = 0 and  $t = \tau$ . The inset zooms into 395  $\leq$  $n \le 405$ , where small differences in the distributions cause small differences in the averages  $\langle U^{\dagger}H_1U\rangle - \langle H_1\rangle = \langle \Delta(H_1, U)\rangle$  correspond to the center of the normal distribution in Fig. 3. The variation of  $H_1$ has relatively low fluctuations, with  $\sigma_{\Delta(H_1,U)} \approx 0.243\hbar\omega$ . However, the uncertainties  $\sigma_{H_1} \approx \sigma_{U^{\dagger}H_1U} \approx |\Im(\alpha)|\hbar\omega = 20\hbar\omega$  are large, consistent with the lower bound in Eq. (3),  $|\langle [U^{\dagger}H_1U, H_1] \rangle| \approx 9.707\hbar^2\omega^2$ . (**b**) The probabilities distributions of finding  $H_2 = H - H_1 = H_e$ with the dimensionless value  $\bar{e} = e/(\hbar\omega)$  at t = 0 (solid blue line) and  $t = \tau$  (dashed red line) show low fluctuations. Here,  $[U^{\dagger}H_2U, H_2] = 0$ , allowing for  $\sigma_{H_2} = \sigma_{U^{\dagger}H_2U} \approx 0.121\hbar\omega$  and  $\sigma_{\Delta(H_2,U)} \approx \sigma_{\Delta(H_1,U)} \approx 0.243\hbar\omega$ . Although the absolute value of the energy variation is the same in both panels, the energy fluctuations in a are significantly larger than in b. This difference is consistent with Eq. (3): in **a**, the commutator  $[U^{\dagger}H_1U, H_1] \neq 0$  is non-zero, while in **b**,  $[U^{\dagger}H_2U, H_2] = 0$ . Notice that the figures illustrate that energy differences in  $H_1$  are more readily discernible when measuring  $H_2$ compared to  $H_1$ , supplementing Result 2.

acting on  $\mathcal{H} \otimes \mathcal{H}'$ , and a vector  $|v\rangle \in \mathcal{H}'$  such that:

$$[H_1 \otimes \mathbb{1}_{\mathcal{H}'} + H_2, U \otimes U'] = 0, \tag{4}$$

$$H_2 \left| \delta_i(H_1, U), v \right\rangle = E_i \left| \delta_i(H_1, U), v \right\rangle, \tag{5}$$

$$(U^{\dagger} \otimes U^{\dagger})H_2(U \otimes U^{\prime}) |\delta_i(H_1, U), v\rangle = E_i^{\prime} |\delta_i(H_1, U), v\rangle, (6)$$

where  $|\delta_i(H_1, U), v\rangle = |\delta_i(H_1, U)\rangle \otimes |v\rangle$ , and  $E_i$  and  $E'_i = E_i - \delta_i(H_1, U)$  are real numbers.

The proof of this result is outlined in the "Methods" section and detailed in the SM [38].

Result 2 reveals a significant insight into the interplay between the OBS protocol and the CRIN conditions, as examplified in Fig. 3. When employing directly the OBS protocol for an energy operator  $H_1$  without considering result 2, the measurement process projects the system into an eigenstate  $|\delta_i(H_1, U)\rangle$ , with the variation of energy inferred as  $\delta_i(H_1, U)$ . However, the variation  $\delta_i(H_1, U)$  is not directly measured. Instead, the protocol relies on measuring an auxiliary observable  $O^{\Delta}$  that commutes with  $\Delta(H_1, U)$ , as illustrated in Fig. 3. Considering result 2, an alternative perspective is offered, enabling a way to indirectly obtain the OBS statistics for  $\Delta(H_1, U)$  by means of a probe, while adhering to the CRIN conditions. Specifically, Result 2 guarantees the existence of a probe related with an auxiliary Hamiltonian  $H_2$  and dynamics U' such that the total energy  $H_1 \otimes \mathbb{1}_{\mathcal{H}'} + H_2$  is conserved under  $U \otimes U'$ . Moreover, once the combined state  $|\delta_i(H_1, U), v\rangle$ is prepared, it has well-defined values for  $H_2$  at both times

0 and *t*, enabling the variation of  $H_2$  to be precisely determined as  $-\delta_i(H_1, U)$ . By considering CRIN conditions 2 and 3, one can infer that the variation of  $H_2$  is  $-\delta_i(H_1, U)$  with 100% certainty (when disregarding practical experimental disturbances). Furthermore, conditions 1 and 4 ensure that this conclusion remains valid, as the conservation of  $H_1 \otimes \mathbb{1}_{\mathcal{H}'} + H_2$  enforces the same probability for  $H_1$  to vary by  $\delta_i(H_1, U)$ . This result applies universally to any observable  $O_1$  and  $O_2$  by substituting  $H_1, H_2 \rightarrow O_1, O_2$  in Result 2. While it may not always be practically feasible to design an experiment that implements the auxiliary Hamiltonian  $H_2$  and unitary operator U' described in Result 2, there are scenarios where the result can be directly applied. One case of this class of examples is the trapped ion case illustrated in Figs. 1, 3, and 5, as described in the SM [38].

To conclude, we have shown that the OBS protocol is unique in complying with the CRIN principles. Given the fundamental nature of these principles, we expect them to be fulfilled by any POVM protocol aiming to measure VPQs. In this sense, our results provide strong support for considering the OBS protocol as a standard for measuring variation of quantum observables.

Looking forward, several promising directions emerge. For instance, our findings open intriguing directions for exploring the observable properties of two-time observables. Can Bell's inequalities applied to two-time observables reveal new insights for quantum cryptography protocols or uncover deeper aspects of quantum nonlocality? How does this framework connect with relativity principles? Moreover, could entanglement and other quantum correlations enhance the capabilities of quantum machines when applied to processes rather than instantaneous (one-time) observables? Since the OBS protocol is the only measurement scheme consistent with the CRIN conditions, these questions are not merely speculative, as they emerge naturally within a physically framework consistent with the CRIN conditions.

Another avenue is to apply the OBS protocol in quantum computing platforms [2, 3, 39], to gain insights into charge fluctuations and energy dissipation in quantum processors. Is it possible to approximate to eigenstates like  $|\delta_i(H_1, U)\rangle$  with virtually no fluctuation in currents or energy variation in quantum devices? Our framework also opens new possibilities for advancing measurement techniques in complex quantum systems. For instance, using Result 2, we demonstrated that attaching an auxiliary system governed by  $H_2$  and U' can enable indirect measurement of variations of observables within a specific system. This approach could be extended to quantum many-body systems [40], where adding few extra degrees of freedom and leveraging conservation laws could provide insights into highly nontrivial observables. Moreover, our findings invite deeper questions about conservation laws and measurement precision. For example, the WAY theorem [6, 7] implies a fundamental tradeoff between energy conservation and the precise measurement of non-commuting observables. Could an auxiliary system be introduced, as in result 2, to restore energy conservation (by means of condition 1) while

enabling precise measurements?

It is important to remark that our framework is grounded in a universal perspective, with an open quantum system (OQS) part of a larger, closed universe treated as a special case (see Fig. 2a). Adapting the OBS protocol to account for scenarios involving noise and environmental interactions, perhaps using the adjoint Lindblad generator [41], is a necessary step. Within this line of research, we are currently investigating the relation between the statistics of variation of observables and the lack of detailed balance at equilibrium (nonreciprocity) and persistent quantum currents [50].

#### METHODS

#### **Proof of result 1**

We begin by proving that the OBS protocol is a CRIN protocol.

First, since the POVM  $\mathbb{M}_{OBS}(H_1, U)$  does not depend on the initial state  $\rho$  for any pair  $(H_1, U)$ , the condition 3 is satisfied. Second, for two energy operators  $H_1$  and  $H_2$  conserved under evolution U (i.e.,  $[H_1 + H_2, U] = 0$ ), it follows that  $\Delta(H_1, U) = -\Delta(H_2, U)$ . This implies the eigenvectors of both operators coincide,  $|\delta_i(H_1, U)\rangle = |\delta_i(H_2, U)\rangle$ , with eigenvalues related by  $\delta_i(H_1, U) = -\delta_i(H_2, U)$ . Consequently, for any z,  $M_{\text{OBS}}(z, H_1, U) = M_{\text{OBS}}(-z, H_2, U)$ , ensuring condition 1. Third, if  $\rho_1 = |e_1\rangle \langle e_1|$  is an eigenstate of both  $H_1$  and  $U^{\dagger}H_1U$ , with eigenvalues  $e_1$  and  $\epsilon_1$ , then  $|e_1\rangle$  is also an eigenstate of  $\Delta(H_1, U)$  with eigenvalue  $\epsilon_1 - e_1$ . Thus, the probability distribution  $\wp_{\text{OBS}}(z, H_1, U, \rho_1) = \delta^{\mathbf{D}}[z - (\epsilon_1 - e_1)]$  satisfies condition 2. At last, for a bipartite system  $\Omega$  evolving under  $U \otimes U'$  acting on a Hilbert space  $\mathcal{H} \otimes \mathcal{H}'$ , the local operator  $H_1 \otimes \mathbb{1}_{\mathcal{H}'}$  evolves as  $\Delta(H_1 \otimes \mathbb{1}_{\mathcal{H}'}, U \otimes U') = \sum_i \delta_i(H_1, U) P_i^{\Delta}(H_1, U) \otimes \mathbb{1}_{\mathcal{H}'}$ . This ensures that  $M_{\text{OBS}}(z, H_1 \otimes \mathbb{1}_2, U \otimes U') = M_{\text{OBS}}(z, H_1, U) \otimes \mathbb{1}_{\mathcal{H}'}$ , fulfilling condition 4.  $\mathbb{M}_{OBS}$  is indeed a CRIN protocol.

The final and most crucial task is to prove that the OBS protocol is the *only* one satisfying the CRIN conditions. Specifically, we aim to demonstrate that for any CRIN protocol  $\mathbb{M}'$ , and any  $H_1$  and U, any element  $M'(z, H_1, U)$  of the POVM  $\mathbb{M}'(H_1, U)$  satisfy

$$M'(z, H_1, U) = M_{\text{OBS}}(z, H_1, U),$$
(7)

for all *z*. To this end, consider an arbitrary CRIN protocol  $\mathbb{M}'$ . For simplicity, assume arbitrary  $H_1$  and U such that  $\Delta(H_1, U)$  is diagonal in a discrete non-degenerate basis { $|\delta_i(H_1, U)\rangle$ }. The general cases are addressed similarly in the SM [38]. We first establish that

$$\langle \delta_i(H_1, U) | M'(z, H_1, U) | \delta_i(H_1, U) \rangle = \delta^{\mathsf{p}}[z - \delta_i(H_1, U)] = = \langle \delta_i(H_1, U) | M_{\mathsf{OBS}}(z, H_1, U) | \delta_i(H_1, U) \rangle,$$
(8)

for all elements  $\{|\delta_i(H_1, U)\rangle\}$  of the eigenbasis. To prove this, consider operators U' and  $H_2$  acting on  $\mathcal{H}'$  and  $\mathcal{H} \otimes \mathcal{H}'$ , re-

spectively, along with a vector  $|v\rangle \in \mathcal{H}'$ , such that:

$$[H_1 \otimes \mathbb{1}_{\mathcal{H}'} + H_2, U \otimes U'] = 0, \tag{9}$$

$$H_2 \left| \delta_i(H_1, U), v \right\rangle = E_i \left| \delta_i(H_1, U), v \right\rangle, \tag{10}$$

$$(U^{\dagger} \otimes U^{\dagger})H_2(U \otimes U^{\prime}) |\delta_i(H_1, U), v\rangle = E_i^{\prime} |\delta_i(H_1, U), v\rangle (11)$$

where  $E'_i = E_i - \delta_i(H_1, U)$ . By Result 2, the existence of such  $H_2$ , U', and  $|v\rangle$  is guaranteed. Since  $\mathbb{M}'$  is CRIN, Conditions 1 and 3 imply for  $U \otimes U'$  in Eq. (9) and all *z* that (see section IC of the SM [38])

$$M'(z, H_1 \otimes \mathbb{1}_{\mathcal{H}'}, U \otimes U') = M'(-z, H_2, U \otimes U').$$
(12)

Moreover, considering Condition 4, we obtain

$$M'(z, H_1 \otimes \mathbb{1}_{\mathcal{H}'}, U \otimes U') = M'(z, H_1, U) \otimes \mathbb{1}_{\mathcal{H}'}.$$
 (13)

Considering the initial state  $\rho_i = |\delta_i(H_1, U), v\rangle \langle \delta_i(H_1, U), v|$ and condition 2, we obtain for any arbitrary z':

$$Tr[M'(z', H_2, U \otimes U')\rho_i] = \delta^{\mathbf{D}}[z' + \delta_i(H_1, U)].$$
(14)

Substituting  $z' \rightarrow -z$ , we have:

$$\operatorname{Tr}[M'(-z, H_2, U \otimes U')\rho_i] = \delta^{\mathbf{D}}[z - \delta_i(H_1, U)].$$
(15)

Combining Eqs. (12), (13), and (15), we obtain Eq. (8). This deduction is valid for arbitrary  $H_1$ , U, z, and  $|\delta_i(H_1, U)\rangle$ , demonstrating Eq. (8) for any CRIN protocol  $M'(z, H_1, U)$ .

Next, we prove that for any element  $M'(z, H_1, U)$  of an CRIN protocol  $\mathbb{M}'$  and for all  $i \neq j$ ,

$$\langle \delta_i(H_1, U) | M'(z, H_1, U) | \delta_j(H_1, U) \rangle = 0 = = \langle \delta_i(H_1, U) | M_{OBS}(z, H_1, U) | \delta_j(H_1, U) \rangle.$$
 (16)

To show this, first note that  $M'(z, H_1, U)$  is Hermitian, nonnegative, and can be diagonalized in a discrete basis  $\{|w_j\rangle\}$ (for the case of a continuous basis, see the SM [38]), such that  $M'(z, H_1, U) = \sum_j w_j |w_j\rangle \langle w_j|$ , with  $w_j \ge 0$ . Since  $\{|\delta_i(H_1, U)\rangle\}$  spans the Hilbert space  $\mathcal{H}$ , each  $|w_j\rangle$  can be expressed as  $|w_j\rangle = \sum_k \gamma_{jk} |\delta_k(H_1, U)\rangle$ , where  $\gamma_{jk} = \langle \delta_k(H_1, U)|w_j\rangle$ . As a result, we can write:

$$M'(z, H_1, U) = \sum_{j,k} w_j |\gamma_{jk}|^2 |\delta_k(H_1, U)\rangle \langle \delta_k(H_1, U)| + \sum_{j,k \neq k'} w_j \gamma^*_{jk'} \gamma_{jk} |\delta_k(H_1, U)\rangle \langle \delta_{k'}(H_1, U)|.$$
(17)

Two and only two cases arise: either there exists  $\bar{k}$  such that  $z = \delta_{\bar{k}}(H_1, U)$ , or  $z \neq \delta_k(H_1, U)$  for all k.

*Case 1*:  $z = \delta_{\bar{k}}(H_1, U)$ . From Eq. (8),  $\langle \delta_k(H_1, U) | M'(z, H_1, U) | \delta_k(H_1, U) \rangle = 0$  for all  $k \neq \bar{k}$ . Eq. (17) thus implies that  $\sum_j w_j |\gamma_{jk}|^2 = 0$  for all  $k \neq \bar{k}$ . Since  $w_j \geq 0$ , we must have  $w_j \gamma_{jk} = 0$  for all  $k \neq \bar{k}$ . Consequently, all off-diagonal terms in Eq. (17) vanish, proving Eq. (16) for all  $i \neq j$ .

*Case 2*:  $z \neq \delta_k(H_1, U)$  for all k. Eq. (8) implies  $\langle \delta_k(H_1, U) | M'(z, H_1, U) | \delta_k(H_1, U) \rangle = 0$  for all k. From Eq. (17), this leads to  $\sum_j w_j |\gamma_{jk}|^2 = 0$  for all k, and hence  $w_j \gamma_{jk} = 0$ . Thus, the off-diagonal terms vanish, and Eq. (16) holds for all  $i \neq j$ .

Combining Eq. (8) with Eq. (16), we find that for all *i* and *j*,  $\langle \delta_i(H_1, U) | M'(z, H_1, U) | \delta_j(H_1, U) \rangle =$  $\langle \delta_i(H_1, U) | M_{OBS}(z, H_1, U) | \delta_j(H_1, U) \rangle$ . Thus,  $M'(z, H_1, U) =$  $M_{OBS}(z, H_1, U)$  for all *z*, *H*<sub>1</sub>, and *U*. Therefore, any CRIN protocol M' must coincide with the OBS protocol  $\mathbb{M}_{OBS}$ , completing the proof.

## Outline of the proof of result 2

Since *U* is unitary and diagonalizable by a countable eigenbasis  $\{|u_i\rangle\}$ , we assume that for every basis element  $|u_i\rangle$ , the relations  $U |u_i\rangle = u_i |u_i\rangle$  and  $u_i = e^{i\theta_i}$  hold, where  $\theta_i \in \mathbb{R}$ . The set of all  $\theta_i$  is referred to as  $\Theta = \{\theta_i\}$ . Based on Lemma 1 in the SM [38], we show that for any such *U* and associated  $\Theta$ , we can define a *U'* acting on a Hilbert space  $\mathcal{H}'$  with countable basis  $\{|u'_i\rangle\}$  satisfying the following condition: (**U**) *Each basis element*  $|u'_i\rangle$  satisfies  $U' |u'_i\rangle = u'_i |u'_i\rangle$ , where  $u'_i = e^{i\theta'_i}$  and  $\theta'_i$  is real. The set of all  $\theta'_i$  associated with  $\{|u'_i\rangle\}$  is denoted by  $\Theta' = \{\theta'_i\}$  and is countably infinite set. Also, for any indexes *m*, *k*, and *j*, if  $\theta'_j \in \Theta'$  and  $\theta_m, \theta_k \in \Theta$ , there exists infinitely many  $\theta'_i \in \Theta'$  such that  $\theta_m + \theta'_i = \theta_k + \theta'_i \mod 2\pi$ .

Considering U' that satisfy (U), we are interested in construct  $H_2$  that satisfy Eq. (4), whose components with respect to the basis  $|u_i, u'_k\rangle$  can be written as

$$\langle u_m, u_j | [H_1 \otimes \mathbb{1}_{\mathcal{H}'} + H_2, U \otimes U'] | u_k, u_l \rangle = = (e^{i(\theta_k + \theta'_l)} - e^{i(\theta_m + \theta'_j)})((H_1)_{mk} \delta^{jl} + (H_2)^{jl}_{mk}) = 0.$$
 (18)

We considered the notation  $\langle u_m | H_1 | u_k \rangle = (H_1)_{mk}$ ,  $\langle u_m, u'_j | H_2 | u_k, u'_l \rangle = (H_2)^{jl}_{mk}$  and  $\delta^{jl} = \langle u'_j | \mathbb{1}_{\mathcal{H}'} | u'_l \rangle$ . The subindexes and superindexes refer, respectively, to the components of the basis of *U* and *U'*. As a result of property (**U**), it follows that for any *m*, *k* and *j*, there exist at least one *l* such that  $(e^{i(\theta_k + \theta'_l)} - e^{i(\theta_m + \theta'_j)}) = 0$ . For these cases, we can consider  $(H_2)^{jl}_{mk}$  as any arbitrary value and still the relation Eq. (18) holds. Therefore, we can define free complex variables  $h^{jl}_{mk}$ and assume the components of  $H_2$  to be

for either the indexes k = m and  $l \ge j$ , or the indexes k > m and any *l*. For the other indexes, we define

$$(H_2)_{mk}^{jl} = [(H_2)_{km}^{lj}]^*.$$
(20)

By taking into account that  $H_1$  is hermitian, these two equations guarantees that  $H_2$  is hermitian and satisfies Eq.(4) (see Lemma 2 of the SM [38] for more details). Therefore, we assume from this point on that  $H_2$  is defined by Eqs. (19) and (20) and all we need to do to prove the rest of the result is to characterize  $h_{mk}^{jl}$  such that Eqs. (5) and (6) are satisfied.

We begin by considering Eq. (5) and then show that  $H_2$ , satisfying Eqs. (5) and (4), also satisfies Eq. (6). For this, let  $|v\rangle \in \mathcal{H}'$  be a state whose components  $\beta_i = \langle u'_i | v \rangle$  satisfy  $\Re(\beta_j) \neq 0$  and  $\Im(\beta_j) \neq 0$ , i.e. non-vanishing real and imaginary parts. As shown in Lemma 3 of the SM [38], it is always possible to define such a  $|v\rangle$ . We analyze how to define the free variables  $h_{mk}^{jl}$  in Eq. (19) so that Eq. (5) is satisfied. For this, consider the components of Eq. (5) in the basis  $|u_m, u'_i\rangle$ :

$$E_i \alpha_{mi} \beta_j = \sum_{k,l} (H_2)^{jl}_{mk} \alpha_{ki} \beta_l, \qquad (21)$$

where  $\alpha_{mi} = \langle u_m | \delta_i(H_1, U) \rangle$ . Defining the set  $\mathbb{N}_{mj}$  of all pairs  $\{k, l\}$  satisfying  $(e^{i(\theta_k + \theta'_l)} - e^{i(\theta_m + \theta'_l)}) = 0$ , we can rewrite Eq. (21), using Eqs. (19) and (20), as:

$$E_i \alpha_{mi} \beta_j = -\sum_{\{k,l\} \notin \mathbb{N}_{mj}} (H_1)_{mk} \delta^{jl} \alpha_{ki} \beta_l + \sum_{\{k,l\} \in \mathbb{N}_{mj}} h_{mk}^{jl} \alpha_{ki} \beta_l.$$
(22)

By property (**U**), there always exists at least one pair  $\{k, l\}$ where  $\alpha_{ki} \neq 0$  and  $\mathbb{N}_{mj}$  is non-empty. Thus, for each *m* and *j*, at least one free variable  $h_{mk}^{jl}$  is available. In the section IIIB of the SM [38] we demonstrate that  $h_{mk}^{jl}$  can always be tuned to ensure Eq. (22) holds for any  $\alpha_{mi}$ , *m*, *j*,  $E_i$ , and  $(H_1)_{mk}$ , proving Eq. (5) for any U,  $H_1$ , and  $|\delta_i(H_1, U)\rangle$ .

To complete the proof, we show that the same  $H_2$  and  $|v\rangle \in \mathcal{H}'$  satisfying Eqs. (4) and (5) also satisfy Eq. (6). Applying  $U^{\dagger} \otimes U^{\dagger}$  to both sides of Eq. (4), we obtain:

$$\Delta(H_1, U) \otimes \mathbb{1}_{\mathcal{H}'} = -\Delta(H_2, U \otimes U').$$
<sup>(23)</sup>

Since  $\Delta(H_1, U) \otimes \mathbb{1}_{\mathcal{H}'} |\delta_i(H_1, U), v\rangle = \delta_i(H_1, U) |\delta_i(H_1, U), v\rangle$ , Eq. (23) implies:  $[(U^{\dagger} \otimes U'^{\dagger})H_2(U \otimes U') - H_2] |\delta_i(H_1, U), v\rangle = -\delta_i(H_1, U) |\delta_i(H_1, U), v\rangle$ . Using the fact that  $H_2$  satisfies Eq. (5) and the relation  $(U^{\dagger} \otimes U'^{\dagger})H_2(U \otimes U') = H_2 + \Delta(H_2, U \otimes U')$ , we deduce:  $(U^{\dagger} \otimes U'^{\dagger})H_2(U \otimes U') |\delta_i(H_1, U), v\rangle = (E_i - \delta_i(H_1, U)) |\delta_i(H_1, U), v\rangle$ , completing the proof.

#### **Example: Trapped ion**

The trapped ion system considered throughout the main text is similar to that in Ref. [18]. The total Hamiltonian is  $H = H_{\rm HO} + H_{\rm e}. \ H_{\rm HO} = \frac{P^2}{2m} \otimes \mathbb{1}_{\rm s} + \frac{m\omega^2 X^2}{2} \otimes \mathbb{1}_{\rm s} = \hbar\omega(N+1/2) \otimes \mathbb{1}_{\rm s}$ represents the center-of-mass energy of a Ca<sup>+</sup> ion, and  $H_{\rm e} \approx$  $\hbar(\omega_z/2 + \Delta_S k_{sw}X/2) \otimes \sigma_z$  describes the coupling between the ion's center-of-mass position and the spin of its covalent electron for small displacements, mediated by an optical dipole force (see Fig. 1). The system evolves in isolation under  $U_t = \exp(-itH/\hbar)$  until time t. Since  $[H, \sigma_z] = [X, \sigma_z] =$  $[P, \sigma_z] = 0$ , then  $U_t = U'_t \exp[-it(\hbar\omega_z\sigma_z - m\omega^2a^2)/2\hbar]$ , where we defined  $U'_t = \exp(-itH'/\hbar), H' = P^2/(2m) \otimes \mathbb{1}_s +$  $m\omega^2/2 (X \otimes \mathbb{1}_s + a(\mathbb{1}_{CM} \otimes \sigma_z))^2$ , and  $a = (\hbar\Delta_S k_{sw})/(2m\omega^2)$ . Defining  $X' = X \otimes \mathbb{1}_s + a(\mathbb{1}_{CM} \otimes \sigma_z)$ , we find that  $[X', P] = i\hbar \mathbb{1}$ and [X', X] = 0. Given the Heisenberg evolution of any operator  $O(t) = U_t^{\dagger} O U_t$  [42], it follows that  $X'(t) = U_t^{\dagger} X' U_t'$ and  $P(t) = U_t^{\dagger}(P \otimes \mathbb{1}_s)U_t^{\prime}$ . Differentiating these expressions with respect to t results in  $\partial_t X'(t) = P(t)/m$  and  $\partial_t P(t) =$  $-m\omega^2 X'(t)$ . These solve to  $X'(t) = X'(0)\cos(\omega t) + \frac{P(0)}{m\omega}\sin(\omega t)$ 

and  $P(t) = -m\omega X'(0) \sin(\omega t) + P(0) \cos(\omega t)$ , with initial conditions  $X'(0) = X \otimes \mathbb{1}_s + a(\mathbb{1}_{CM} \otimes \sigma_z)$  and  $P(0) = P \otimes \mathbb{1}_s$ . For  $\tau = \pi/\omega$ , we find:  $X'(\tau) = -X \otimes \mathbb{1}_s - a(\mathbb{1}_{CM} \otimes \sigma_z)$ ,  $P(\tau) = -P \otimes \mathbb{1}_s$ . Considering these expressions, we obtain:

$$H_{\rm HO}(\tau) = H_{\rm HO} + \Delta(H_{\rm HO}, U_{\tau}), \qquad (24)$$

$$H_{\rm e}(\tau) = H_{\rm e} + \Delta(H_{\rm e}, U_{\tau}), \qquad (25)$$

$$\Delta(H_{\rm HO}, U_{\tau}) = \hbar \Delta_S k_{\rm SW} \left( X \otimes \sigma_z + a \right) = -\Delta(H_{\rm e}, U_{\tau}).$$
(26)

Therefore,  $[\Delta(H_{\rm HO}, U_{\tau}), H_{\rm HO}] = [H_{\rm HO}(\tau), H_{\rm HO}] = i2\hbar\omega^2 aP \otimes \sigma_z$ ,  $[\Delta(H_{\rm e}, U_{\tau}), H_{\rm e}] = [H_{\rm e}(\tau), H_{\rm e}] = [\Delta(H_{\rm HO}, U_{\tau}), X \otimes \sigma_z] = -[\Delta(H_{\rm e}, U_{\tau}), X \otimes \sigma_z] = 0$ . Moreover, it follows that

$$\Delta(H_{\rm HO}, U_{\tau}) | x, \pm \rangle = \delta_{x,\pm} | \delta_{x,\pm} \rangle = -\Delta(H_{\rm e}, U_{\tau}) | x, \pm \rangle \qquad (27)$$

where  $\delta_{x,\pm} = \hbar \Delta_S k_{sw}(\pm x + a)$  and  $|\delta_{x,\pm}\rangle = |x,\pm\rangle$ . We are now ready to compute all results in Figs. 1, 3, and 5.

### Calculations for the Fig.1

In Fig. 1, we compute the TPM statistics for the variation of  $H_{\rm HO}$  and  $H_{\rm e}$  for a preparation  $\rho = |0, +\rangle \langle 0, +|$ , where  $N|0\rangle = 0|0\rangle$  and  $\sigma_z |+\rangle = |+\rangle$ . Let us first consider the TPM applied to the energy  $H_{\rm e}$  (Fig. 1b). In this case, we can use a result similar to Ref. [13] to show that if  $[H_{\rm e}(\tau), H_{\rm e}] =$ 0,  $\wp_{\rm TPM}(z, H_{\rm e}, U_{\tau}, \rho) = \wp_{\rm OBS}(z, H_{\rm e}, U_{\tau}, \rho)$  (see section V of the SM [38]). Thus, it suffices  $\wp_{\rm OBS}(z, H_{\rm e}, U_{\tau}, \rho)$  to describe  $\wp_{\rm TPM}(z, H_{\rm e}, U_{\tau}, \rho)$ . Using Eq. (27) and the OBS definition, we find:

$$M_{\rm OBS}(z, H_{\rm e}, U_{\tau}) = \sum_{s \in \{+, -\}} \int_{-\infty}^{\infty} dx \delta^{\rm D}[z - \delta_{x,s}(H_{\rm e}, U_{\tau})] P_{x,s}, \quad (28)$$

where  $P_{x,s} = |x, s\rangle \langle x, s|$  and the sum  $\sum_{s \in \{+,-\}}$  runs over the two possible values of  $s = \pm$ . Substituting  $M_{OBS}(z, H_e, U_\tau)$  into Eq. (2) and simplifying using Eq. (27), we find:

$$\mathcal{D}_{OBS}(z, H_e, U_\tau, \rho) = \frac{1}{\hbar \Delta_S k_{SW}} |\langle -a - \frac{z}{\hbar \Delta_S k_{SW}} |0\rangle|^2, \qquad (29)$$

where  $\langle -a - \frac{z}{\hbar \Delta_s k_{SW}} | 0 \rangle$  corresponds to  $\langle x | 0 \rangle$ , with  $x \to -a - \frac{z}{\hbar \Delta_s k_{SW}}$ . For the ground state  $| 0 \rangle$ , we have:

$$|\langle x|0\rangle|^2 = \mathcal{N}(x,0,\sigma^2), \tag{30}$$

where  $\sigma \equiv \sqrt{\hbar/(2m\omega)}$  and, from now on, we consider

$$\mathcal{N}(y, y_c, \sigma_y^2) = \frac{1}{\left(2\pi\sigma_y^2\right)^{1/2}} \exp\left[-\frac{(y-y_c)^2}{2\sigma_y^2}\right]$$
(31)

as a normal distribution of variable y, with mean  $y_c$  and variance  $\sigma_y^2$ . Substituting Eq. (30) into Eq. (29), and noting that  $\wp_{\text{TPM}}(z, H_e, U_\tau, \rho) = \wp_{\text{OBS}}(z, H_e, U_\tau, \rho)$ , we find:  $\wp_{\text{TPM}}(z, H_e, U_\tau, \rho) = \mathcal{N}(z, z_a, \sigma_z^2)$ , where  $z_a = -\hbar\Delta_S k_{\text{SW}}a$  and  $\sigma_z = \hbar\Delta_S k_{\text{SW}}\sigma$ . Finally, we compute the probability  $p_{\text{TPM}}(-n\hbar\omega, H_e, U_\tau, \rho) \coloneqq \int_{I_n^-} dz \, \wp_{\text{TPM}}(z, H_e, U_\tau, \rho)$  of  $H_e$  of varying by an amount z in intervals  $I_n^- = [-(n+1/2)\hbar\omega, -(n-1)/2)\hbar\omega$ 

 $1/2\hbar\omega$ ]. Figure 1b shows  $p_{\text{TPM}}(-n\hbar\omega, H_e, U_\tau, \rho)$  for n = 0 and n = 1.

Next, we use the main text expression for TPM protocol and Refs. [14, 15, 19, 20] to find:  $\wp_{\text{TPM}}(z, H_{\text{HO}} \otimes \mathbb{1}_{s}, U_{\tau}, \rho) = \sum_{m,n=0}^{\infty} \sum_{r,s \in \{+,-\}} \delta^{\text{D}}[z - (e_{m,r} - e_{n,s})]|\langle m, r| U_{\tau} | n, s \rangle|^{2} \langle n, s| \rho | n, s \rangle$ , where  $\sum_{r,s \in \{+,-\}}$  sums over  $r, s \in \{+,-\}$ , and  $|n, s \rangle$  are eigenvectors of  $H_{\text{HO}}$ satisfying  $H_{\text{HO}} | n, \pm \rangle = e_{n,\pm} | n, \pm \rangle = \hbar \omega (n + 1/2) | n, \pm \rangle$ . Since  $[U_{\tau}, \sigma_{z}] = 0$  and  $\langle n, s| \rho | n, s \rangle = 1$  for n = 0 and s = +, and 0 otherwise, it follows:  $\wp_{\text{TPM}}(z, H_{\text{HO}}, U_{\tau}, \rho) = \sum_{m=0}^{\infty} \delta^{\text{D}}[z - (e_{m,+} - e_{0,+})]| \langle m| U_{\tau}^{+} | 0 \rangle|^{2}$ , where  $U_{\tau}^{+} = \langle +| U_{\tau} | + \rangle = e^{-i\theta_{\tau}} \exp\left[-\frac{i\pi}{2} \left(\frac{p^{2}}{2m} + \frac{m\omega^{2}}{2} X_{+}^{2}\right)\right]$ ,  $X_{+} = \langle +| X' | + \rangle = X + a$ , and  $\theta_{\tau} = \frac{\hbar \omega_{zt}}{2\hbar} - \frac{m\omega^{2}a^{2}t}{2\hbar}$ . Analogously,  $p_{\text{TPM}}(n\hbar\omega, H_{\text{HO}}, U_{\tau}, \rho) :=$ 

Analogously,  $p_{\text{TPM}}(nh\omega, H_{\text{HO}}, U_{\tau}, \rho) = |\langle n| U_{\tau}^+ |0\rangle|^2$  is the probability of  $H_{\text{HO}}$  varying in intervals  $I_n^+ = [(n - 1/2)\hbar\omega, (n + 1/2)\hbar\omega]$ . We derive  $|\langle n| U_{\tau}^+ |0\rangle|^2$  analytically, but leave its details to the SM [38]. In Fig. 1, we show  $p_{\text{TPM}}(n\hbar\omega, H_{\text{HO}}, U_{\tau}, \rho)$  for n = 0 and n = 1.

#### Calculations for the Fig. 3 and 5

For Figs. 3 and 5, we consider an initial state  $\rho = |\alpha, -\rangle \langle \alpha, -|$ , where

$$|\alpha\rangle = e^{-|\alpha|^2/2} \sum_{n=0}^{\infty} \frac{\alpha^n}{\sqrt{n!}} |n\rangle, \qquad (32)$$

is the coherent state [43] and  $\sigma_z |-\rangle = -|-\rangle$ . For Fig. 3, we compute the OBS distribution:

$$\wp_{\text{OBS}}(z, H_{\text{HO}}, U_{\tau}, \rho) = \int_{-\infty}^{\infty} dx \delta^{\mathbf{p}}[z - \delta_{x, -}] |\langle x|\alpha \rangle|^2.$$
(33)

Using the coherent state probability  $|\langle x|\alpha \rangle|^2 = \mathcal{N}(x, \langle \alpha | X | \alpha \rangle, \sigma_X^2)$ , with  $\langle \alpha | X | \alpha \rangle = \sqrt{2\hbar/(m\omega)}\mathfrak{R}(\alpha)$ and  $\sigma_X = \sqrt{\hbar/(2m\omega)}$  [43], and introducing the dimensionless position  $\bar{x} = k_{sw}x$ , we get the probability  $\wp(\bar{x}, \rho) = \mathcal{N}(\bar{x}, k_{sw} \langle \alpha | X | \alpha \rangle, k_{sw}^2 \sigma_X^2)$  of the center of mass of the ion to be found at dimensionless position  $\bar{x}$  when initially prepared in the state  $\rho$ . The left-hand side of Fig. 3 shows this result.

For the right-hand side of Fig. 3, we compute  $\wp_{OBS}(\bar{z}, H_{HO}, U_{\tau}, \rho)$  with respect to the dimensionless variation  $\bar{z} = z/(\hbar\omega)$ . Substituting  $|\langle x|\alpha \rangle|^2 = \mathcal{N}(x, \langle \alpha|X|\alpha \rangle, \sigma_X^2)$  into Eq. (33), we find:  $\wp_{OBS}(\bar{z}, H_{HO}, U_{\tau}, \rho) = \mathcal{N}(x, \bar{z}_{\alpha}, \sigma_{\bar{z}}^2)$ , where  $\bar{z}_{\alpha} = -\frac{2m\omega^2 a \langle \alpha|X|\alpha \rangle}{\hbar} + \frac{2m\omega|a|^2}{\hbar}$  and  $\sigma_{\bar{z}} = \frac{2m\omega a \sigma_X}{\hbar}$ . The right-hand side of Fig. 3 displays  $\wp_{OBS}(\bar{z}, H_{HO}, U_{\tau}, \rho)$ , with parameters as described in the figure.

For Fig. 5, we compute the Heisenberg two-time uncertainty relations for  $H_{\rm HO}$  using Eqs. (24) and (26), and the coherent state properties from [43] to obtain:  $\sigma_{\bar{z}} := \sigma_{\Delta(H_{\rm HO},U_{\tau})}/(\hbar\omega) = \sqrt{\hbar\Delta_S^2 k_{SW}^2/\omega^3}$ ,  $\sigma_{H_{\rm HO}(\pi/\omega)}/(\hbar\omega) = \sqrt{(\Re(\alpha) - \sigma_{\bar{z}})^2 + \Im^2(\alpha)}$ ,  $\sigma_{H_{\rm HO}}/(\hbar\omega) = |\alpha|$ , and  $|\langle [H_{\rm HO}(\pi/\omega), H_{\rm HO}(0)] \rangle |/(\hbar^2 \omega^2) = 2\sigma_{\bar{z}} \Im(\alpha)$ .

Finally, we compute the probabilities for Fig. 5 for measurements of  $H_{\rm H0}(\pi/\omega)$  and  $H_{\rm H0}(0)$ . At t = 0,  $H_{\rm H0}(0) |n, -\rangle = \hbar\omega(n + 1/2) |n, -\rangle$ , and the probability  $p(n, H_{\rm H0}(0))$  is:  $p(n, H_{\rm H0}(0)) = |\langle n, -|\alpha, -\rangle|^2 = e^{-|\alpha|^2} \frac{|\alpha|^{2n}}{n!}$ . At  $t = \tau$ , we show in the SM [38] that  $p(n, H_{\rm H0}(\tau)) = e^{-|\alpha'|^2} \frac{|\alpha'|^{2n}}{n!}$ , where  $\alpha' = [\Re(\alpha) - 2a \sqrt{m\omega/(2\hbar)}] + i\Im(\alpha)$ . Fig. 5(a) shows  $p(n, H_{\rm H0}(0))$  (blue squares) and  $p(n, H_{\rm H0}(\tau))$  (red circles).

For the probability distribution  $\wp(\bar{e}, H_e(t))$  of  $H_e$  at t = 0 and  $t = \tau$ , using Eqs. (24) and (26), we compute:  $\wp(\bar{e}, H_e(0)) = \mathcal{N}(\bar{e}, (\hbar\omega)^{-1} \langle H_e(0) \rangle, (\hbar\omega)^{-2} \sigma_{H_e}^2)$  and  $\wp(\bar{e}, H_e(\tau)) = \mathcal{N}(\bar{e}, (\hbar\omega)^{-1} \langle H_e(\tau) \rangle, (\hbar\omega)^{-2} \sigma_{H_e}^2)$ , where  $\sigma_{H_e} = \hbar\Delta_S k_{sw} \sigma_X/2$ ,  $\langle H_e(0) \rangle = -(\hbar/2)(\omega_z - \Delta_S k_{sw} \langle \alpha | X | \alpha \rangle)$ , and  $\langle H_e(\tau) \rangle = (\hbar/2)(-\omega_z + \Delta_S k_{sw} \langle \alpha | X | \alpha \rangle - 2a)$ ). Fig. 5(b) shows  $\wp(\bar{e}, H_e(0))$  (blue solid line) and  $\wp(\bar{e}, H_e(\tau))$  (red dashed line).

#### ACKNOLEDGEMENTS

We thank Renato Angelo, Shay Blum, Časlav Brukner, Ohad Cremerman, Karen Hovhannisyan, Michael Iv, Gabriel Landi, Uri Peskin, Saar Rahav, and Ferdinand Schmidt-Kaler for fruitful conversations. T.A.B.P.S. was supported by the Helen Diller Quantum Center - Technion under Grant No. 86632417 and by the ISRAEL SCIENCE FOUNDATION (grant No. 2247/22). D.G.K. is supported by the ISRAEL SCIENCE FOUNDATION (grant No. 2247/22) and by the Council for Higher Education Support Program for Hiring Outstanding Faculty Members in Quantum Science and Technology in Research Universities.

\* pinto\_silva@campus.technion.ac.il

- Tesser, L. & Splettstoesser, J. Out-of-equilibrium fluctuationfissipation Bounds. *Phys. Rev. Lett.* 114, 186304 (2024).
- [2] Arute, F. et al. Quantum supremacy using a programmable superconducting processor. *Nature* 574, 505-510 (2019).
- [3] Neill, C. A path towards quantum supremacy with superconducting qubits. Ph.D. thesis, University of California, Santa Barbara (2017).
- [4] Zhao, N. et al. Atomic-scale magnetometry of distant nuclear spin clusters via nitrogen-vacancy spin in diamond. *Nat. Nan*otechnol 6, 242-246 (2011).
- [5] Bozzio, M. et al. Enhancing quantum cryptography with quantum dot single-photon sources. *Npj Quantum Inf.* 8, 104 (2022).
- [6] Loveridge, L. & Busch, P. 'Measurement of quantum mechanical operators' revisited. *Eur. Phys. J. D* 62, 297-307 (2011)
- [7] Gisin, N. & Cruzeiro, E. Z. Quantum Measurements, Energy Conservation and Quantum Clocks. Ann. Phys. 530, 1700388 (2018)
- [8] Aharonov, Y. et al. On conservation laws in quantum mechanics. PNAS 118, e1921529118 (2021)
- [9] Aamir, M.A. et al. Thermally driven quantum refrigerator autonomously resets a superconducting qubit. *Nat. Phys.* 21, 318-323 (2025)
- [10] Hovhannisyan, K. V. & Imparato, A. Energy conservation and fluctuation theorem are incompatible for quantum work. *Quantum* 8, 1336 (2024).

- [11] Hänggi, P. & Talkner, P. The other QFT. Nat. Phys. 11, 108 (2015).
- [12] Campisi, M. et al. *Colloquium*: Quantum fluctuation relations: Foundations and applications. *Rev. Mod. Phys.* 83, 771 (2011).
- [13] Talkner, P. et al. Fluctuation theorems: Work is not an observable, *Phys. Rev. E* 75, 050102(R) (2007).
- [14] Roncaglia, A. J. et al. Work measurement as a generalized quantum measurement. *Phys. Rev. Lett.* **113**, 250601 (2014).
- [15] Perarnau-Llobet, M. et al. No-go theorem for the characterization of work fluctuations in coherent quantum systems, *Phys. Rev. Lett.* **118**, 070601 (2017).
- [16] Bäumer, E. et al. Fluctuating work in coherent quantum systems: Proposals and limitations in *Thermodynamics in the quantum regime: Fundamental aspects and new directions* (Springer International, 2018), pp. 275-300.
- [17] Batalhão, T. B. et al. Experimental Reconstruction of Work Distribution and Study of Fluctuation Relations in a Closed Quantum System. *Phys. Rev. Lett.* **113**, 140601 (2014).
- [18] von Lindenfels, D. et al., Spin Heat Engine Coupled to a Harmonic-Oscillator Flywheel. *Phys. Rev. Lett.* **123**, 080602 (2019).
- [19] Pinto Silva, T. A. B. & Angelo, R. M. Quantum mechanical work. *Phys. Rev. A* **104**, 042215 (2021).
- [20] Pinto Silva, T. A. B. & Gelbwaser-Klimovsky, D. Quantum work: reconciling quantum mechanics and thermodynamics. *Phys. Rev. Research* 6, L022036 (2024).
- [21] Pinto Silva, T. A. B. & Angelo, R. M. Fluctuation theorems for genuine quantum mechanical regimes. *Phys. Rev. A* 107, 052211 (2023).
- [22] Esposito, M. et al. Nonequilibrium fluctuations, fluctuation theorems, and counting statistics in quantum systems. *Rev. Mod. Phys.* 81, 1665 (2009).
- [23] Gherardini, S. & De Chiara, G. Quasiprobabilities in Quantum Thermodynamics and Many-Body Systems, *PRX quantum* 5, 030201 (2024).
- [24] Talkner, P. & Hänggi, P. Aspects of quantum work, *Phys. Rev. E* **93**, 022131 (2016).
- [25] Allahverdyan, A. E. & Nieuwenhuizen, Th. M. Fluctuations of work from quantum subensembles: The case against quantum work-fluctuation theorems. *Phys. Rev. E* 71, 066102 (2005).
- [26] Maquedano et al. Two-time quantities as elements of physical reality. *Phys. Lett. A*, **525**, 129943 (2024)
- [27] Lindblad, G. Non-equilibrium Entropy and Irreversibility (D. Reidel Publishing Company, 1983).
- [28] Brandão, F. G. S. L. et al. Resource theory of quantum states out of thermal equilibrium. *Phys. Rev. Lett.* **111**, 250404 (2013)
- [29] Brandão, F. et al. The second laws of quantum thermodynamics. PNAS 112, 3275-3279 (2015).
- [30] Guryanova, Y. et al. Thermodynamics of quantum systems with multiple conserved quantities. *Nat. Commun.* 7, 12049 (2016).
- [31] Nielsen, M. A. & Chuang, I. L. Quantum Computation and Quantum Information Information (Cambridge University Press, 2010).
- [32] Heinosaari, T. A. & Ziman, M. *The Mathematical Language of Quantum Theory* (Cambridge University Press, 2012).
- [33] Busch, P. *The Quantum Theory of Measurement* (Springer-Verlag, 1996).
- [34] Einstein, A. et al. Can quantum-mechanical description of physical reality be considered complete? *Phys. Rev.* 47, 777 (1935).
- [35] Peres, A. & Terno, D. R. Colloquium: Quantum information and relativity theory, *Rev. Mod. Phys.* 76, 93 (2004).
- [36] Horodecki P. & Ramanathan, R. The relativistic causality versus no-signaling paradigm for multi-party correlations. *Nat. Commun.* 10, 1701 (2019).

- [38] See Supplemental Material at [URL will be inserted by publisher], which includes Refs. [13, 15, 19, 20, 32, 33, 42–49], for detailed proofs of results 1 and 2 and an in-depth analysis of the model described in the Figs. 1, 3, 5, and 6.
- [39] Kreppel, F. et al. Quantum Circuit Compiler for a Shuttling-Based Trapped-Ion Quantum Computer, *Quantum* 7, 1176 (2023).
- [40] J. Lukin, A. et al. Probing entanglement in a many-bodylocalized system, Science 364, 256 (2019).
- [41] Breuer, H.-P. & Petruccione, F. *The theory of open quantum systems*, (Oxford University Press, 2002).
- [42] Sakurai, J. J. *Modern quantum mechanics*, (Addison-Wesley Publishing Company, 1994).

- [43] Cohen-Tannoudji, C. et al. Quantum Mechanics, Volume 1: Basic Concepts, Tools, and Applications (Wiley-VCH Verlag GmbH & Co, 2020).
- [44] Athreya K. B. & Lahiri, S. N. Measure theory and probability theory (Springer, 2006).
- [45] Fristedt, B. E. & Gray, L. F. A modern approach to probability theory (Springer Science & Business Media, 2013).
- [46] Arfken, G. B. et al. *Mathematical methods for physicists: a comprehensive guide* (Academic press, 2011).
- [47] Freire, I. S. & Angelo, R. M. Quantifying continuous-variable realism. *Phys. Rev. A* 100, 022105 (2019).
- [48] Pinto Silva, T. A. B. A definition of quantum mechanical work. Master's dissertation, Universidade Federal do Paraná, Curitiba (2018).
- [49] Weisstein, E. W. Confluent hypergeometric function of the first kind. https://mathworld.wolfram.com/ (2003).
- [50] Alicki, et al. Violation of Detailed Balance in Quantum Open Systems. *Phys. Rev. Lett.* **131**, 040401 (2023).

# Supplemental Material for "Fundamental constraints on quantum fluctuations: Conservation laws, reality, and no-signaling"

Thales A. B. Pinto Silva<sup>\*</sup> and David Gelbwaser-Klimovsky Schulich Faculty of Chemistry and Helen Diller Quantum Center, Technion-Israel Institute of Technology, Haifa 3200003, Israel

# CONTENTS

I.	Preliminaries	2
	A. General two-time observables	2
	1. Degenerate observables	2
	2. Continuous observables	2
	B. Some formalities regarding POVMs	4
	C. Condition 1 and 3 imply $[H_1 + H_2, U] = 0 \implies M(z, H_1, U) = M(-z, H_2, U)$	5
II.	Completion of Result 1: degeneracies and continuous basis	7
	A. Discrete degeneracies	7
	B. Continuous, degenerate case	9
	C. Final remarks regarding result 1	12
III.	Complete proof of result 2	12
	A. Lemmas necessary for the deduction of Result 2	13
	B. The main proof of Result 2	16
IV.	Adapting to the time-dependent case	22
V.	Results needed for the ion trap examples in the figures	24
	A. OBS and TPM for the commuting continuous case	24
	B. Calculating the probabilities $ \langle \mathbf{n}   \mathbf{U}_{\tau}^+   0 \rangle ^2$ in the subsection "Calculations for the Fig.1" of Methods section	25
	C. Calculating the probabilities $p(n, H_{HO}(\tau))$ in subsection "Calculations for the Fig. 3 and 5" of the methods section	27
	D. Example of result 2 considering Trapped Ions	29
	References	30

<sup>\*</sup> pinto\_silva@campus.technion.ac.il

## I. PRELIMINARIES

Before presenting the main results and derivations of the paper, we introduce key formalities and preliminary results that will be essential throughout this Supplemental Material (SM). We primarily consider the same systems discussed in the main text. Specifically, we analyze a general quantum system,  $\Omega$ , composed of arbitrary subsystems, under minimal assumptions:  $\Omega$  is initially prepared in an arbitrary quantum state  $\rho$  in a Hilbert space  $\mathcal{H}$  and evolves from time 0 to *t* under an arbitrary unitary operator *U*, which has a countable eigenbasis. Our focus is on the measurement of the variation of *a part* of the total energy of  $\Omega$ , represented by a time-independent Hermitian operator  $H_1$  acting on  $\mathcal{H}$  (see Fig. 2 for examples).

# A. General two-time observables

In the Methods section of the main text, we presented Result 1 under the assumption that the variation observable  $\Delta(H_1, U)$  is diagonalizable in a discrete, countable and non-degenerate basis. Here, in this SM, we extend our analysis to cases where  $\Delta(H_1, U)$  may exhibit degeneracy and/or possess a continuous spectrum. In this subsection, we outline key details related to these more general scenarios.

## 1. Degenerate observables

We first consider the case where  $\Delta(H_1, U)$  is degenerate but still diagonalizable in a discrete, countable basis. For this, we assume the existence of a basis  $|\delta_i^j(H_1, U)\rangle$  such that

$$\Delta(H_1, U) |\delta_i^j(H_1, U)\rangle = \delta_i(H_1, U) |\delta_i^j(H_1, U)\rangle \tag{1}$$

where  $j \in \{1, 2, 3, ..., g(i)\}$ , meaning that each eigenvalue  $\delta_i(H_1, U)$  has a degeneracy g(i). This implies that the basis vectors satisfy the orthonormality condition

$$\langle \delta_{i'}^{j'}(H_1, U) | \delta_i^j(H_1, U) \rangle = \delta_{ii'}^{\mathsf{K}} \delta_{ii'}^{\mathsf{K}}, \tag{2}$$

where  $\delta_{ii'}^{\kappa}$  and  $\delta_{ii'}^{\kappa}$  are Kronecker's delta, and that the operator

$$P_i(H_1, U) = \sum_j |\delta_i^j(H_1, U)\rangle \langle \delta_i^j(H_1, U)|$$
(3)

is a projector onto the subspace  $\mathcal{E}_i$  associated with the eigenvalue  $\delta_i(H_1, U)$  [1]. We can then express  $\Delta(H_1, U)$  in terms of these projectors as

$$\Delta(H_1, U) = \sum_i \delta_i(H_1, U) P_i(H_1, U).$$
(4)

Using this decomposition, we define the OBS protocol POVM  $\mathbb{M}_{OBS}(H_1, U)$  for a given energy operator  $H_1$  and evolution U in terms of the operators

$$M_{\rm OBS}(z, H_1, U) = \sum_i \delta^{\mathbf{D}}[z - \delta_i(H_1, U)] P_i(H_1, U).$$
(5)

where  $\delta^{\mathbf{D}}$  is the Dirac's delta. Notice that the case where  $\Delta(H_1, U)$  has no degeneracies is simply a special case where g(i) = 1 and  $P_i(H_1, U) = |\delta_i(H_1, U)\rangle \langle \delta_i(H_1, U)|$  for all *i*.

#### 2. Continuous observables

The next step is to describe cases where  $\Delta(H_1, U)$  has a continuous spectrum. Although the unitary operator U remains diagonalizable in a discrete, countable basis, it is possible that not only  $\Delta(H_1, U)$  but also the energy operators  $H_1$  or  $U^{\dagger}H_1U$  exhibit continuous spectra.

When  $\Delta(H_1, U)$  has a continuous spectrum, we introduce the projectors  $|w(H_1, U), y\rangle \langle w(H_1, U), y| dw, dy$ , which satisfy

$$\Delta(H_1, U) |w(H_1, U), y\rangle \langle w(H_1, U), y| \, dw, dy = w(H_1, U) |w(H_1, U), y\rangle \langle w(H_1, U), y| \, dw, dy, \tag{6}$$

$$\Delta(H_1, U) = \int_{-\infty}^{\infty} dw w(H_1, U) P_w(H_1, U)$$
<sup>(7)</sup>

where the projectors onto the eigenspaces associated with  $w(H_1, U)$  are given by

$$P_{w}(H_{1}, U) = \int_{-\infty}^{\infty} dy |w(H_{1}, U), y\rangle \langle w(H_{1}, U), y|.$$
(8)

Here, we assume a general case where each eigenvalue  $w(H_1, U)$  has an uncountable degeneracy. However, degeneracy may also be discrete, in which case

$$P_{w}(H_{1}, U) = \sum_{y=1}^{g(w)} |w(H_{1}, U), y\rangle \langle w(H_{1}, U), y|, \qquad (9)$$

where g(w) denotes the degeneracy of each eigenvalue  $w(H_1, U)$ . If there is no degeneracy, the projectors reduce to

$$P_{w}(H_{1}, U) = |w(H_{1}, U)\rangle \langle w(H_{1}, U)|.$$
(10)

With this continuous formulation of  $\Delta(H_1, U)$ , we define the OBS POVM for  $H_1$  and U as

$$M_{\text{OBS}}(z, H_1, U) = \int_{-\infty}^{\infty} dw \,\delta^{\mathbf{p}}[z - w(H_1, U)] P_w(H_1, U) = P_z(H_1, U) = \int_{-\infty}^{\infty} dy \,|z, y\rangle \,\langle z, y| \,, \tag{11}$$

where  $|z, y\rangle$  is an eigenvector of  $\Delta(H_1, U)$  with eigenvalue z, satisfying  $\Delta(H_1, U) |z, y\rangle = z |z, y\rangle$ .

To extend our analysis to cases where  $H_1$  or  $U^{\dagger}H_1U$  have continuous spectra, we need to adapt the CRIN conditions accordingly. While Conditions 1, 3, and 4 remain well-defined for both discrete and continuous spectra, Condition 2 (the Reality Condition) is not generally well-posed in the continuous case.

To illustrate this issue, consider an energy operator  $H_1$  that is diagonalized in a continuous and unbounded basis, such as  $H_e$  in the main text. In such a scenario, there may exist an eigenvector  $|e\rangle$  satisfying

$$H_1 |e\rangle = e |e\rangle$$
 and  $U^{\dagger} H_1 U |e\rangle = \epsilon_e |e\rangle$  (12)

but  $|e\rangle$  is not *normalizable*. In this case, Condition 2 becomes ill-defined, as no *normalized* state  $\rho$  can simultaneously assign a well-defined energy value at both times. This raises the question: how should Condition 2 be adapted to accommodate such continuous scenarios?

To address this question, we first analyze the role of the Reality Condition in shaping the operator  $M(z, H_1, U)$  in the discrete case. In this setting, if a normalized state  $|e\rangle$  is an eigenvector of both  $H_1$  and  $U^{\dagger}H_1U$  with respective eigenvalues e and  $\epsilon_e$ , then it must also be an eigenstate  $|\delta_j^k(H_1, U)\rangle$  of  $\Delta(H_1, U)$  with eigenvalue  $\delta_j(H_1, U) = \epsilon_e - e$ . Consequently, the Reality Condition imposes the constraint:

$$\langle \delta_{i}^{k}(H_{1},U) | M(z,H_{1},U) | \delta_{i}^{k}(H_{1},U) \rangle = \delta^{\mathbf{D}}[z - \delta_{i}^{k}(H_{1},U)].$$
(13)

This implies that any  $M(z, H_1, U)$  satisfying the Reality Condition must take the form:

$$M(z, H_1, U) = \delta^{\mathbf{D}}[z - \delta^k_j(H_1, U)] |\delta^k_j(H_1, U)\rangle \langle \delta^k_j(H_1, U)| + M_n(z)$$
(14)

where  $M_n(z)$  is a Hermitian operator that satisfies  $\langle \delta_i^k(H_1, U) | M_n(z) | \delta_i^k(H_1, U) \rangle = 0$ .

Now, let us examine the continuous case. As in the discrete scenario, if Eq. (12) holds, then  $|e\rangle$  must also be an eigenstate of  $\Delta(H_1, U)$ . Given that  $\Delta(H_1, U)$  has a continuous and potentially degenerate spectrum, as described in Eq. (7), the state  $|e\rangle$  must take the form  $|w_e, y_e\rangle$ , where  $w_e = \epsilon_e - e$  represents the eigenvalue of  $\Delta(H_1, U)$ , and  $y_e$  is a real parameter accounting for the degeneracy.

We cannot simply assume for the continuous case that  $\langle w_e, y_e | M(z, H_1, U) | w_e, y_e \rangle = \delta^{\mathbf{p}}(z - w_e)$ , since  $|w_e, y_e \rangle$  is not normalized. However, we can adapt the reasoning from Eq. (14). Intuitively, given that  $|w_e, y_e\rangle$  is an eigenstate of both  $H_1$  and  $U^{\dagger}H_1U$ , we can assume:

$$M(z, H_1, U) = \delta^{\mathbf{D}}(z - w_e) |w_e, y_e\rangle \langle w_e, y_e| \, dwdy + M'_n(z) = |z, y_e\rangle \langle z, y_e| \, dwdy + M'_n(z) \text{ for } z = w_e \tag{15}$$

where  $dwdy \langle w_e, y_e | M'_n(z) | w_e, y_e \rangle dwdy = 0$ . This means that  $M(z, H_1, U)$  has its diagonal element  $|w_e, y_e \rangle \langle w_e, y_e | dwdy$  defined as  $\delta^{\mathbf{p}}(z - w_e)$ , while the remaining terms, represented by  $M'_n(z)$ , are arbitrary but must have a vanishing diagonal component.

Since the differentials dw and dy are not rigorously defined, expressing the reality condition in the form of Eq. (15) lacks precision. To formalize this further, we assume that there exist two open intervals  $I_{w_e}$  and  $I_{y_e}$  with bounded lengths  $d_{w_e} = \int_{I_{w_e}} dw$  and  $d_{y_e} = \int_{I_{w_e}} dy$ , such that:

$$M(z, H_1, U) = \int_{I_{y_e}} dy |z, y\rangle \langle z, y| + M_{nd}(z), \quad \text{for } z = w_e$$
(16)

where  $M_{nd}(z)$  is Hermitian and satisfies:

$$\langle w(H_1, U), y | M_{nd}(z) | w'(H_1, U), y' \rangle = 0, \quad \text{for all } w, w', y, y' \text{ such that } w, w' \in I_{w_e} \text{ and } y, y' \in I_{y_e}.$$

$$\tag{17}$$

By defining  $M(z, H_1, U)$  as in Eqs. (16) and (17), we ensure that the reality condition is well-defined for the continuous case. To see this, consider an arbitrary normalized state:

$$|\psi\rangle = \int_{I_1} dw \int_{I_2} dy \, |w(H_1, U), y\rangle \, \psi(w(H_1, U), y), \tag{18}$$

where  $\psi(w(H_1, U), y) = \langle w(H_1, U), y | \psi \rangle$  and  $I_1$  and  $I_2$  are real intervals. If  $I_1$  and/or  $I_2$  do not contain  $w_e$  and  $y_e$ , respectively, then the reality condition implemented via Eq. (16) imposes no restriction on  $\langle \psi | M(z, H_1, U) | \psi \rangle$ . However, if  $I_1 \subset I_w$  and  $I_2 \subset I_y$ , meaning they contain  $w_e$  and  $y_e$ , respectively, then:

$$\langle \psi | M(z, H_1, U) | \psi \rangle = \int_{I_2} dy | \psi(z, y) |^2.$$
 (19)

which leads to:

$$\langle \psi | \int_{I_1} dz M(z, H_1, U) | \psi \rangle = 1.$$
<sup>(20)</sup>

This implies that whenever the state  $|\psi\rangle$  is within the resolution range  $I_{w_e}$  and  $I_{y_e}$ , i.e., sufficiently close to an eigenstate of  $H_1$  and  $U^{\dagger}H_1U$ , its variation of energy is necessarily confined to the interval  $I_{w_e}$  around  $w_e$ .

We thus assume that Eqs. (16) and (17) will be taken to define the continuous analogue of the reality condition. Formally we then impose that:

• **Reality condition for the continuous case**: For any system  $\Omega$ , operators  $H_1$ , evolution U, if there is an eigenvector  $|w_e, y_e\rangle$  of  $\Delta(H_1, U)$ , such that  $H_1 |w_e, y_e\rangle = e |w_e, y_e\rangle$ ,  $U^{\dagger}H_1U |w_e, y_e\rangle = \epsilon_e |w_e, y_e\rangle$ , and  $\Delta(H_1, U) |w_e, y_e\rangle = w_e |w_e, y_e\rangle = (\epsilon_e - e) |w_e, y_e\rangle$  then there are intervals  $I_{w_e}$  and  $I_{y_e}$  with bounded lengths such that the operators  $\{M(z, H_1, U)\}$  satisfy Eqs. (16) and (17).

#### B. Some formalities regarding POVMs

Throughout the main text, we expressed the POVM elements as  $M(z, H_1, U)$ , ensuring that they satisfy  $\int_{-\infty}^{\infty} M(z, H_1, U) dz = 1$ and  $M(z, H_1, U) \ge 0$ . For an initial state  $\rho$ , the probability density was defined as  $\wp(z, H_1, U, \rho) = \text{Tr}[M(z, H_1, U)\rho]$ . However, for continuous variables, POVM elements are not strictly defined as  $M(z, H_1, U)$ . This notation is adopted throughout the main text and this SM for simplicity, but here we clarify its formal basis.

When considering energy variation as a continuous variable *z*, POVMs are formally defined as operators  $\mathcal{M}(I, H_1, U)$  acting on open intervals  $I \subset \mathbb{R}$  such that:

$$Tr[\mathcal{M}(I, H_1, U)\rho] = P(I, H_1, U, \rho)$$
(21)

where  $P(I, H_1, U, \rho)$  gives the probability that the energy variation falls within *I*. POVMs must satisfy  $\mathcal{M}((-\infty, \infty), H_1, U) = \mathbb{1}$ (as energy variation can take any real value) and  $\mathcal{M}(I, H_1, U) \ge 0$  for all  $I \subset \mathbb{R}$ .

The probability density function  $\wp(z, H_1, U, \rho)$  is thus defined as dP/dz (see page 54 of Ref. [2]), allowing us to express:

$$P(I, H_1, U, \rho) = \int_I dz \, \wp(z, H_1, U, \rho).$$
(22)

A key concept in measure theory [2, 3] is that of "almost everywhere in z" (*z*-a.e.), meaning a statement holds for all  $z \in \mathbb{R}$  except on subsets  $I_0$  of measure zero, i.e., open intervals  $I_0$  such that

$$\int_{I_0} dz = 0. \tag{23}$$

$$\int_{I} dz \, \wp(z, H_1, U, \rho) = \int_{I} dz \, \wp'(z, H_1, U, \rho).$$
<sup>(24)</sup>

For example, if  $\wp(z_0, H_1, U, \rho) \neq \wp'(z_0, H_1, U, \rho)$  for a single point  $z_0$  but  $\wp(z, H_1, U, \rho) = \wp'(z, H_1, U, \rho)$  everywhere else, they still define the same probability distribution since a single point has zero measure.

Similarly, it is common to define a "density operator"  $M(z, H_1, U)$  for POVM elements:

$$\mathcal{M}(I,H_1,U) = \int_I dz \, M(z,H_1,U). \tag{25}$$

This allows  $\text{Tr}[M(z, H_1, U)\rho]$  to represent the probability density. In both this SM and the main text, we referred to  $M(z, H_1, U)$  as POVM elements in  $\mathbb{M}(H_1, U)$  for simplicity. Formally, they are not exactly elements of a POVM set, and they are defined by the POVM map  $\mathcal{M}(I, H_1, U)$  in Eq. (21) (see [4, 5] for rigorous POVM definitions). The crucial point for our proofs is the equivalence relation: if  $M'(z, H_1, U) = M(z, H_1, U) z$ -a.e., they define the same POVM  $\mathcal{M}(I, H_1, U)$  for all interval of measure nonzero  $I \subset \mathbb{R}$ , since:

$$\mathcal{M}(I, H_1, U) = \int_I dz \, M(z, H_1, U) = \int_I dz \, M'(z, H_1, U).$$
(26)

Thus, to prove that two POVMs  $\mathcal{M}(I, H_1, U)$  and  $\mathcal{M}'(I, H_1, U)$  are identical for all  $I \subset \mathbb{R}$ , it suffices to show:

$$M'(z, H_1, U) = M(z, H_1, U)$$
 z-a.e. (27)

This formalism will be essential in extending the proof of Result 1 in this SM.

# **C.** Condition 1 and 3 imply $[H_1 + H_2, U] = 0 \implies M(z, H_1, U) = M(-z, H_2, U)$

In this section we show that for every POVM  $\mathbb{M}$  satisfying condition 1 and 3, it follows that  $M(z, H_1, U) = M(-z, H_2, U)$ . To prove this result, we first notice that, by Condition 1, whenever  $[H_1 + H_2, U] = 0$ , the following relation holds for any initial state  $\rho$ :

$$\wp(z, H_1, U, \rho) = \text{Tr}[M(z, H_1, U)\rho] = \text{Tr}[M(-z, H_2, U)\rho] = \wp(-z, H_2, U, \rho)$$
(28)

This leads to two possible scenarios:

- (i)  $M(z, H_1, U)$  and  $M(-z, H_2, U)$  act on a Hilbert space  $\mathcal{H}$  with countable basis  $\{|i\rangle\}$ , or
- (ii)  $M(z, H_1, U)$  and  $M(-z, H_2, U)$  act on a Hilbert space  $\mathcal{H}$  with continuous basis.

We consider both cases:

• Case (i): In this scenario,  $M(z, H_1, U)$  and  $M(-z, H_2, U)$  can only differ if there exist basis elements  $|j\rangle$  and  $|k\rangle$  in  $\{|i\rangle\}$  such that

$$\langle j| M(z, H_1, U) | k \rangle \neq \langle j| M(-z, H_2, U) | k \rangle.$$
<sup>(29)</sup>

However, this contradicts Eq. (28) in all possible cases. To see why, consider the following two scenarios:

- Case  $\mathbf{k} = \mathbf{j}$ : If Eq. (29) holds for k = j, then by Condition 3,  $M(z, H_1, U)$  and  $M(-z, H_2, U)$  must be independent of  $\rho$ , meaning we can freely choose  $\rho$  while still satisfying Eq. (28). Setting  $\rho = |j\rangle\langle j|$ , we obtain:

$$Tr[M(z, H_1, U)\rho] = \langle j | M(z, H_1, U) | j \rangle \neq \langle j | M(-z, H_2, U) | j \rangle = Tr[M(-z, H_2, U)\rho].$$
(30)

This directly contradicts Eq. (28).

- Case  $\mathbf{k} \neq \mathbf{j}$ : Suppose Eq. (29) holds for some  $k \neq \mathbf{j}$ , so that we define the difference:

$$\Phi_{jk} = \langle j | M(z, H_1, U) | k \rangle - \langle j | M(-z, H_2, U) | k \rangle, \qquad (31)$$

where  $|\Phi_{jk}| \neq 0$ . By Condition 3, we can consider any state  $\rho = |\psi\rangle \langle \psi|$  without altering  $M(z, H_1, U)$  or  $M(-z, H_2, U)$ . Choosing

$$|\psi\rangle = \frac{a\,|j\rangle + b\,|k\rangle}{|a|^2 + |b|^2},\tag{32}$$

where a = 1 and  $b = \Phi_{ik}^*$  (i.e., the conjugate of  $\Phi_{jk}$ ), we compute:

$$\operatorname{Tr}[M(z, H_1, U)\rho] - \operatorname{Tr}[M(-z, H_2, U)\rho] = \frac{ab^* \Phi_{jk}^* + a^* b \Phi_{jk}}{|a|^2 + |b|^2} = \frac{|\Phi_{jk}|^2 + |\Phi_{jk}|^2}{1 + |\Phi_{jk}|^2} > 0,$$
(33)

which contradicts Eq. (28).

Given that we consider arbitrary j and k, it follows that for all j and k it cannot be the case in which Eq. (29) holds. As a result, we conclude:

$$M(z, H_1, U) = M(-z, H_2, U).$$
(34)

• Case (ii): Since the space  $\mathcal{H}$  cannot be described by a countable basis, we consider arbitrary projectors  $|y\rangle \langle y| dy$ , with dy real, so that any operator A acting on  $\mathcal{H}$  can be written as:

$$A = \iint_{-\infty}^{\infty} dy \, dy' \, |y\rangle \, \langle y| \, A \, |y'\rangle \, \langle y'| \,. \tag{35}$$

Assuming that  $M(z, H_1, U) \neq M(-z, H_2, U)$ , and given that  $M(z, H_1, U)$  acts on and has its image in a space dense in  $L_2(-\infty, \infty)$  (i.e., the space of normalizable measurable functions defined by the 2-norm that is bounded. See Section 3.2 of [2]), then there must exist states  $|\psi\rangle$  and  $|\phi\rangle$  such that

$$\langle \phi | (M(z, H_1, U) - M(-z, H_2, U)) | \psi \rangle \neq 0.$$
 (36)

Thus, there must be intervals  $I_1$  and  $I_2$  with nonzero measure, i.e.,  $\int_{I_1} dy \neq 0$  and  $\int_{I_2} dy \neq 0$ , such that

$$|\psi_{1}\rangle = \frac{\int_{I_{1}} dy |y\rangle \psi(y)}{\sqrt{\int_{I_{1}} dy |\psi(y)|^{2}}}, \quad |\phi_{1}\rangle = \frac{\int_{I_{2}} dy |y\rangle \phi(y)}{\sqrt{\int_{I_{2}} dy |\phi(y)|^{2}}},$$
(37)

and

$$\bar{\Phi} := \langle \phi_1 | DM(z) | \psi_1 \rangle = \int_{I_2} dy \int_{I_1} dy' \phi_1^*(y) \langle y | DM(z) | y' \rangle \psi_1(y') \neq 0,$$
(38)

where we define, for simplicity,  $DM(z) = M(z, H_1, U) - M(-z, H_2, U)$ , as well as  $\langle y|\psi_1 \rangle = \psi_1(y)$  and  $\langle y|\phi_1 \rangle = \phi_1(y)$ , with  $|\bar{\Phi}| > 0$ . If, instead, Eq. (38) does not hold for all nonzero measure intervals  $I_1$  and  $I_2$ , then we obtain

$$\langle \phi | DM(z) | \psi \rangle = \langle \phi | (M(z, H_1, U) - M(-z, H_2, U)) | \psi \rangle = 0, \tag{39}$$

which contradicts our initial assumption in Eq. (36). There are only two possible cases: either (ii)a  $I_1 \cap I_2 \neq \emptyset$  or (ii)b  $I_1 \cap I_2 = \emptyset$ . We analyze both scenarios, demonstrating that assuming  $\langle \phi_1 | (M(z, H_1, U) - M(-z, H_2, U)) | \psi_1 \rangle \neq 0$  leads to a contradiction:

- Case (ii)a: Suppose first that  $I_1 = I_2$ . In this case, we consider the state  $\rho_{\psi} = |\psi\rangle \langle \psi|$ , where

$$|\psi\rangle = \frac{a\,|\psi_1\rangle + b\,|\phi_1\rangle}{|a|^2 + |b|^2}\tag{40}$$

with a and b being complex numbers. By Condition 3, it follows that

$$\langle \psi_1 | DM(z) | \psi_1 \rangle = \langle \phi_1 | DM(z) | \phi_1 \rangle = 0, \tag{41}$$

which, considering that both  $M(z, H_1, U)$  and  $M(-z, H_2, U)$  are Hermitian, results in

$$Tr[DM(z)\rho_{\psi}] = \frac{a^*b\bar{\Phi}^* + ab^*\bar{\Phi}}{|a|^2 + |b|^2}.$$
(42)

Choosing a = 1 and  $b = \overline{\Phi}$ , we obtain

$$Tr[DM(z)\rho_{\psi}] = \frac{2|\bar{\Phi}|^2}{1+|\bar{\Phi}|^2} > 0,$$
(43)

which contradicts Condition 3 and Eq. (28). Therefore, Eq. (38) cannot hold for  $I_1 = I_2$ . Next, consider the case where  $I_1 \cap I_2 = I'_3$ . We define the intervals  $I'_1 = I_1 \setminus I'_3$  and  $I'_2 = I_2 \setminus I'_3$ , where  $B \setminus A$  denotes the complement of A with respect to B. Thus, there is no intersection between  $\{I'_1, I'_2, I'_3\}$ . It must then follow that

$$\Phi_{jk} = \int_{I'_j} dy \int_{I'_k} dy' \phi_1^*(y) \langle y | DM(z) | y' \rangle \psi_1(y') \neq 0,$$
(44)

must hold for some  $j \neq k$ , with  $j \in \{2, 3\}$  and  $k \in \{1, 3\}$ . If this were not the case, then, considering the argument that  $\int_{I'_j} dy \int_{I'_j} dy' \phi_1^*(y) \langle y | DM(z) | y' \rangle \psi_1(y') = 0$ , Eq. (38) could not hold without the existence of  $j \neq k$  such that Eq. (44) is valid. Therefore, for the indices  $j \neq k$  for which Eq. (44) holds, we define the state  $\rho_{jk} = |\psi_{jk}\rangle \langle \psi_{jk}|$  such that

$$|\psi_{jk}\rangle = \frac{a\,|\psi_k\rangle + b\,|\psi_j\rangle}{|a|^2 + |b|^2},\tag{45}$$

with  $a = 1, b = \Phi_{ik}^*$ , and

$$|\psi_{j}\rangle = \frac{\int_{I'_{j}} |y\rangle \phi_{1}(y) dy}{\sqrt{\int_{I'_{j}} |\phi_{1}(y)|^{2} dy}}, \quad |\psi_{k}\rangle = \frac{\int_{I'_{k}} dy \psi_{1}(y)}{\sqrt{\int_{I'_{k}} |\psi_{1}(y)|^{2} dy}}.$$
(46)

By Condition 3,  $\langle \psi_k | DM(z) | \psi_k \rangle = \langle \psi_j | DM(z) | \psi_j \rangle$ , leading to

$$\operatorname{Tr}[DM(z)\rho_{jk}] = \frac{|\bar{\Phi}_{jk}|^2 + |\bar{\Phi}_{jk}|^2}{\sqrt{N'_j N'_k (1 + |\bar{\Phi}_{jk}|^2)}} > 0, \tag{47}$$

contradicting Eq. (28).

- Case (ii)b: The proof follows analogously by substituting  $I'_j \to I_1$ ,  $I'_k \to I_2$ ,  $|\psi_j\rangle \to |\phi_1\rangle$ , and  $|\psi_k\rangle \to |\psi_1\rangle$  in Eqs. (44)-(47).

By considering the only possibilities (ii)a and (ii)b for the continuous case (ii), we showed that there can be no intervals  $I_1$  and  $I_2$  such that Eq. (38) holds, and, therefore, it cannot be true that  $M(z, H_1, U) \neq M(-z, H_2, U)$ .

Since for the only possible cases (i) and (ii),  $M(z, H_1, U) = M(-z, H_2, U)$ , then we can only conclude that conditions 1 and 3 together with  $[H_1 + H_2, U] = 0$  imply  $M(z, H_1, U) = M(-z, H_2, U)$ .

# II. COMPLETION OF RESULT 1: DEGENERACIES AND CONTINUOUS BASIS

For the proof of Result 1 in the Methods section of the main text, we demonstrated that the OBS protocol satisfies the CRIN conditions for any energy operator  $H_1$  and unitary evolution U. This proof holds independently of whether  $\Delta(H_1, U)$  has degeneracies or a continuous spectrum. However, to establish that the OBS protocol is the *only* CRIN protocol, we considered the case where  $\Delta(H_1, U)$  has a discrete, non-degenerate basis. Here, we extend this proof by employing the formalism introduced in the Preliminaries section, completing the proof of Result 1 for cases where  $\Delta(H_1, U)$  exhibits degeneracies and/or has a continuous spectrum.

### A. Discrete degeneracies

To prove that the OBS protocol is the *only* CRIN protocol when  $\Delta(H_1, U)$  have discrete basis with degeneracies, we consider  $\Delta(H_1, U)$  as in Eq. (4), diagonalized by a basis  $\{|\delta_i^j(H_1, U)\rangle\}$ . Similar to the non-degenerate case, we aim here to demonstrate that for any CRIN protocol  $\mathbb{M}'$ , and any  $H_1$  and U, the POVM  $\mathbb{M}'(H_1, U)$  should coincide with the OBS protocol  $\mathbb{M}_{OBS}(H_1, U)$ . For that, as we discussed in the section IB of this SM, is enough for us to prove that

$$M'(z, H_1, U) = M_{OBS}(z, H_1, U), \quad z-a.e..$$
 (48)

Proving this relation establishes that there is no CRIN protocol  $\mathbb{M}'$  such that  $\mathbb{M}'(H_1, U) \neq \mathbb{M}_{OBS}(H_1, U)$ , completing the proof.

To this end, consider an arbitrary CRIN protocol  $\mathbb{M}'$ . We first establish that, because  $\mathbb{M}'$  is CRIN, then it follows:

$$\langle \delta_i^k(H_1, U) | M'(z, H_1, U) | \delta_i^k(H_1, U) \rangle = \delta^{\mathbf{D}}[z - \delta_i(H_1, U)] = \langle \delta_i^k(H_1, U) | M_{\text{OBS}}(z, H_1, U) | \delta_i^k(H_1, U) \rangle,$$
(49)

for all elements  $\{|\delta_i^k(H_1, U)\rangle\}$  of the eigenbasis. To prove this, we consider similar arguments as in the non-degenerate case. Consider operators U' and  $H_2$  acting on  $\mathcal{H}'$  and  $\mathcal{H} \otimes \mathcal{H}'$ , respectively, along with a vector  $|v\rangle \in \mathcal{H}'$ , such that:

$$[H_1 \otimes \mathbb{1}_{\mathcal{H}'} + H_2, U \otimes U'] = 0, \tag{50}$$

$$H_2 \left| \delta_i^k(H_1, U), v \right\rangle = E_i^k \left| \delta_i^k(H_1, U), v \right\rangle, \tag{51}$$

$$(U^{\dagger} \otimes U'^{\dagger})H_2(U \otimes U') |\delta_i(H_1, U), v\rangle = E_i^{\prime k} |\delta_i^k(H_1, U), v\rangle, \qquad (52)$$

where  $E_i^{k} = E_i^k - \delta_i(H_1, U)$ . By Result 2, the existence of such  $H_2$ , U', and  $|v\rangle$  is guaranteed. Since  $\mathbb{M}'$  satisfies all CRIN conditions for all arbitrary energy and evolution operators, Conditions 1, 3 and 4 (Conservation Laws, Independent of the initial state, No-Signaling) imply the following for the evolution  $U \otimes U'$  in Eq. (50) (see section IC of this SM):

$$M'(z, H_1 \otimes \mathbb{1}_{\mathcal{H}'}, U \otimes U') = M'(-z, H_2, U \otimes U'),$$
(53)

$$M'(z, H_1 \otimes \mathbb{1}_{\mathcal{H}'}, U \otimes U') = M'(z, H_1, U) \otimes \mathbb{1}_{\mathcal{H}'}.$$
(54)

Additionally, using the Reality Condition (Condition 2), we can consider the initial state  $\rho_i = |\delta_i(H_1, U), v\rangle \langle \delta_i(H_1, U), v|$ , which is an eigenstate of  $(U^{\dagger} \otimes U'^{\dagger})H_2(U \otimes U')$  and  $H_2$  with eigenvalues  $E_i'^k$  and  $E_i^k$ , respectively. For any arbitrary z', we then deduce:

$$Tr[M'(z', H_2, U \otimes U')\rho_i] = \delta^{\mathbf{D}}[z' + \delta_i(H_1, U)].$$
(55)

Substituting  $z' \rightarrow -z$ , we have:

$$\operatorname{Tr}[M'(-z, H_2, U \otimes U')\rho_i] = \delta^{\mathbf{D}}[z - \delta_i(H_1, U)].$$
(56)

Combining Eqs. (53), (54), (56), and the fact that  $\mathbb{M}'$  is state independent (according to the condition of Independence on state), we obtain Eq. (49). This deduction is valid for arbitrary *z*, and  $|\delta_i(H_1, U)\rangle$ , demonstrating Eq. (49) for any CRIN protocol  $M'(z, H_1, U)$ .

Our next goal is to prove that for any element  $M'(z, H_1, U)$  of an CRIN protocol  $\mathbb{M}'$ , it follows that

$$\langle \delta_i^k(H_1, U) | M'(z, H_1, U) | \delta_i^{k'}(H_1, U) \rangle = 0 = \langle \delta_i^k(H_1, U) | M_{\text{OBS}}(z, H_1, U) | \delta_i^{k'}(H_1, U) \rangle. \quad \text{z-a.e.}$$
(57)

for either  $i \neq j$  and any k and k' or for i = j and  $k \neq k'$ . To show this, first note that  $M'(z, H_1, U)$  is Hermitian, non-negative, and can be diagonalized as  $M'(z, H_1, U) = \int_{-\infty}^{\infty} w(z) |w(z)\rangle \langle w(z)| dw$ , with  $w(z) \ge 0$ . Notice that here we considered the continuous basis  $|w(z)\rangle \langle w(z)| dw$  seeking generality but we could consider the discrete in a similar way as in the main text. As a result, we get:

$$M'(z, H_1, U) = \int dw \sum_{ik} w(z) |\gamma_{ik}(z, w)|^2 |\delta_i^k(H_1, U)\rangle \langle \delta_i^k(H_1, U)| + \int dw \sum_{i \neq i', k, k'} w(z) \gamma_{ik'}^*(z, w) \gamma_{ik}(z, w) |\delta_i^k(H_1, U)\rangle \langle \delta_i^{k'}(H_1, U)| + \int dw \sum_{i \neq i', k, k'} w(z) \gamma_{i'k'}^*(z, w) \gamma_{ik}(z, w) |\delta_i^k(H_1, U)\rangle \langle \delta_{i'}^{k'}(H_1, U)| .$$
(58)

where the integrals are over  $(-\infty, \infty)$  and  $\gamma_{ik}(z, w) = \langle \delta_i^k(H_1, U) | w(z) \rangle$ . Two and only two cases arise: either (1)  $z \neq \delta_i(H_1, U)$  for all *i* or (2) there exists *j* such that  $z = \delta_i(H_1, U)$ . Let us consider both cases:

*Case* (1): In this case, Eq. (49) implies  $\langle \delta_i^k(H_1, U) | M'(z, H_1, U) | \delta_i^k(H_1, U) \rangle = 0$  for all *i* and *k*. From Eq. (58) and the fact that  $w(z) \ge 0$ , leads to  $\int dw w(z) |\gamma_{ik}(z, w)|^2 = 0$  for all *i* and *k*, and hence  $\int_I dw w(z) \gamma_{ik}(z, w) = \int_I dw w(z) \gamma_{ik}^*(z, w) = 0$  for any interval *I* of measure non-zero. Thus, all the off-diagonal terms of  $M'(z, H_1, U)$  with respect to the basis  $\{|\delta_i^k(H_1, U)\rangle\}$  vanish, and Eq. (57) holds for all  $i \ne j$  and *k* and *k'* and all i = j with  $k \ne k'$ .

*Case* 2: In this case, from Eq. (49),  $\langle \delta_i^k(H_1, U) | M'(z, H_1, U) | \delta_i^k(H_1, U) \rangle = 0$  for all  $j \neq i$ . From Eq. (58), this implies  $\int dw w(z) |\gamma_{ik}(z, w)|^2 = 0$  for all  $i \neq j$  and all k. Since  $w(z) \ge 0$  and real, we must have  $\int_I dww(z) \gamma_{ik}(z, w) = \int_I dww(z) \gamma_{ik}^*(z, w) = 0$  for any non-empty interval *I*, for all  $i \neq j$  and all k. Consequently, Eq. (57) holds for all  $i \neq j$  and any k and k'. All left is to prove that for i = j and  $k \neq k'$ , Eq. (57) holds for all  $i \neq j$  and k and k', it follows that, for the given j such that  $z = \delta_j(H_1, U)$  and any given  $k \neq k'$ ,

$$\langle \delta_{i}^{k}(H_{1},U) | M'(z,H_{1},U) | \delta_{i}^{k'}(H_{1},U) \rangle = \Gamma_{ikk'}(z).$$
(59)

where

$$\Gamma_{jkk'}(z) = \int dw w(z) \gamma_{jk'}^*(z, w) \gamma_{jk}(z, w).$$
(60)

Let us assume that there are two indexes  $l \neq l'$  and a non-empty interval  $I \subset \mathbb{R}$  such that  $\int_{I} \Gamma_{ill'}(z) dz = \mathcal{J} \neq 0$ , i.e. such that Eq. (57) does not hold for i = j and  $l \neq l'$ . Since  $\Gamma_{ill'}(z) = 0$  for  $i \neq j$ , then it is necessary that  $\delta_j(H_1, U) \in I$ . In this sense, consider an initial state  $\rho_{\psi} = |\psi\rangle \langle \psi|$  where

$$|\psi\rangle = \frac{\alpha |\delta_j^l(H_1, U)\rangle + \beta |\delta_j^{l'}(H_1, U)\rangle}{\sqrt{|\alpha|^2 + |\beta|^2}},\tag{61}$$

 $\alpha = 1$ , and  $\beta = \mathcal{J}^*$ . As a result, we get from Eqs. (49) and (59), and the fact that  $\delta_i(H_1, U) \in I$ , that

$$\int_{I} dz \operatorname{Tr}[M'(z, H_{1}, U)\rho_{\psi}] = \frac{|\alpha|^{2} + |\beta|^{2} + 2\Re[\int_{I} dz \Gamma_{\tilde{i}ll'}(z)\alpha^{*}\beta]}{|\alpha|^{2} + |\beta|^{2}} = 1 + \frac{2|\mathcal{J}|^{2}}{1 + |\mathcal{J}|^{2}} > 1,$$
(62)

This, however, contradicts the fact that  $M'(z, H_1, U)$  are elements of a POVM, since for any interval *I* and any state  $\rho$ , the POVM elements  $M'(z, H_1, U)$  should satisfy  $\int_I dz \operatorname{Tr}[M(z, H_X, U)\rho] \leq 1$ . Therefore, it cannot be the case that there is non-empty interval *I* and indexes  $l \neq l'$  such that  $\int_I \Gamma_{jll'}(z)dz \neq 0$ . In other words, it means that  $\Gamma_{jll'}(z) = 0$  *z*-a.e. and Eq. (57) is thus proved for all i = j and  $k \neq k'$ .

Combining Eq. (49) and Eq. (57) we find that for all i and j,

$$\langle \delta_{i}(H_{1}, U) | M'(z, H_{1}, U) | \delta_{i}(H_{1}, U) \rangle = \langle \delta_{i}(H_{1}, U) | M_{\text{OBS}}(z, H_{1}, U) | \delta_{i}(H_{1}, U) \rangle \quad z\text{-a.e.}.$$
(63)

Thus,

$$M'(z, H_1, U) = M_{\text{OBS}}(z, H_1, U)$$
 z-a.e. (64)

for all  $H_1$ , and U. Therefore, we conclude that any CRIN protocol  $\mathbb{M}'$  must be equivalent to the OBS protocol  $\mathbb{M}_{OBS}$ , completing the proof for the degenerate, discrete case.

#### B. Continuous, degenerate case

To prove that the OBS protocol is the *only* CRIN protocol when  $\Delta(H_1, U)$  have continuous, degenerate basis, we consider  $\Delta(H_1, U)$  as in Eq. (7), diagonalized in terms of the projectors { $|w(H_1, U), y\rangle \langle w(H_1, U), y|$ }. Similar to the non-degenerate case, we aim here to demonstrate that for any CRIN protocol  $\mathbb{M}'$ , and any  $H_1$  and U, the POVM  $\mathbb{M}'(H_1, U)$  should coincide with the OBS protocol  $\mathbb{M}_{ons}(H_1, U)$ . For that, as we discussed in the section IB of this SM, it is enough for us to prove that

$$M'(z, H_1, U) = M_{\text{OBS}}(z, H_1, U), \quad z-a.e..$$
 (65)

Proving this relation establishes that there is no CRIN protocol  $\mathbb{M}'$  such that  $\mathbb{M}'(H_1, U) \neq \mathbb{M}_{OBS}(H_1, U)$ , completing the proof. To this end, consider an arbitrary CRIN protocol  $\mathbb{M}'$  and its associated POVM  $\mathbb{M}'(H_1, U)$  and elements  $M(z, H_1, U)$ . Regarding  $H_1$  and U, consider operators U' and  $H_2$  acting on  $\mathcal{H}'$  and  $\mathcal{H} \otimes \mathcal{H}'$ , respectively, along with a vector  $|v\rangle \in \mathcal{H}'$ , such that:

$$[H_1 \otimes \mathbb{1}_{\mathcal{H}'} + H_2, U \otimes U'] = 0, \tag{66}$$

$$H_2 |w(H_1, U), y\rangle \otimes |v\rangle = E_w^y |w(H_1, U), y\rangle \otimes |v\rangle,$$
(67)

$$(U^{\dagger} \otimes U'^{\dagger})H_2(U \otimes U')|w(H_1, U), y\rangle \otimes |v\rangle = E_w^{y}|w(H_1, U), y\rangle \otimes |v\rangle,$$
(68)

where  $E'_w = E'_w - w(H_1, U)$ . By Result 2, the existence of such  $H_2$ , U', and  $|v\rangle$  is guaranteed. Since M' satisfies all CRIN conditions for all arbitrary energy and evolution operators, Conditions 1, 3 and 4 (Conservation Laws, Independent of the initial state, No-Signaling) imply the following for the evolution  $U \otimes U'$  in Eq. (66) (see section IC of this SM):

$$M'(z, H_1 \otimes \mathbb{1}_{\mathcal{H}'}, U \otimes U') = M'(-z, H_2, U \otimes U'), \tag{69}$$

$$M'(z, H_1 \otimes \mathbb{1}_{\mathcal{H}'}, U \otimes U') = M'(z, H_1, U) \otimes \mathbb{1}_{\mathcal{H}'}.$$
(70)

Additionally, since  $|w(H_1, U), y\rangle \otimes |v\rangle$  is a non-normalized eigenvector of  $H_2$  and  $(U^{\dagger} \otimes U'^{\dagger})H_2(U \otimes U')$ , with difference in the eigenvalues as  $-w(H_1, U)$ , then we can consider the Reality Condition (Condition 2), defined for the continuous case in section IA, to deduce that there exist open intervals  $I_w$  and  $I_y$  with bounded lengths  $d_w = \int_{I_w} dw$  and  $d_y = \int_{I_u} dy$ , such that

$$M'(z', H_2, U \otimes U') = \int_{I_y} dy \, |z, y, v\rangle \, \langle z, y, v| + M_{nd}(z', H_2, U \otimes U'), \quad \text{for } z' = -w(H_1, U)$$
(71)

where  $M_{nd}(z', H_2, U \otimes U')$  is hermitian and for  $z' = -w(H_1, U)$  satisfies

$$\langle w'(H_1, U), y', v | M_{nd}(z', H_2, U \otimes U') | w''(H_1, U), y'', v \rangle = 0,$$
(72)

for all w, w', y, y' such that  $w', w'' \in I_w$  and  $y', y'' \in I_y$ . Since z' here is arbitrary, we can redefine it as  $z' \to -z$ , to obtain:

$$M'(-z, H_2, U \otimes U') = \int_{I_y} dy \, |z, y, v\rangle \, \langle z, y, v| + M_{nd}(-z, H_2, U \otimes U') \quad \text{for } z = w(H_1, U)$$
(73)

Because  $|v\rangle$  is normalized, then we can combine Eqs. (69), (70), (73), to deduce that

$$\operatorname{Tr}_{\mathcal{H}'}[M'(-z, H_2, U \otimes U')(\mathbb{1}_{\mathcal{H}} \otimes |v\rangle \langle v|)] = \operatorname{Tr}_{\mathcal{H}'}[M'(z, H_1 \otimes \mathbb{1}, U \otimes U')(\mathbb{1}_{\mathcal{H}} \otimes |v\rangle \langle v|)]$$
  
= 
$$\operatorname{Tr}_{\mathcal{H}'}[M'(z, H_1, U) \otimes \mathbb{1}_{\mathcal{H}'}(\mathbb{1}_{\mathcal{H}} \otimes |v\rangle \langle v|)]$$
  
= 
$$M'(z, H_1, U).$$
(74)

Considering Eqs. (73) and (74), we then have that

$$M'(z, H_1, U) = \operatorname{Tr}_{\mathcal{H}'}[M'(-z, H_2, U \otimes U')(\mathbb{1}_{\mathcal{H}} \otimes |v\rangle \langle v|)] = M'_d(z) + M'_{nd}(z)$$
(75)

where

$$M'_{d}(z) = \int_{I_{y}} dy |z, y\rangle \langle z, y| \quad \text{for } z = w(H_{1}, U)$$
(76)

and  $M'_{nd}(z)$  is hermitian and satisfies

$$\langle w'(H_1, U), y' | M'_{nd}(z) | w''(H_1, U), y'' \rangle = 0,$$
(77)

for all w, w', y, y' such that  $w', w'' \in I_w$  and  $y', y'' \in I_y$ . Notice that the results deduced until now hold for arbitrary w and y, so that Eqs. (75)-(77) should be valid for any w and y. Therefore, for  $M'(z, H_1, U)$  to satisfy Eqs. (75)-(77) for all w and y, then it must have the form

$$M'(z, H_1, U) = M_{\text{OBS}}(z, H_1, U) + \bar{M}_{nd}(z)$$
(78)

where  $M_{OBS}(z, H_1, U)$  was defined in Eq. (11) and  $\overline{M}_{nd}(z)$  is hermitian and satisfy

$$\langle w'(H_1, U), y' | \bar{M}_{nd}(z) | w''(H_1, U), y'' \rangle = 0,$$
(79)

for all w, w', y, y' such that  $w', w'' \in I_z$ , where  $I_z$  is some interval with no-zero measure  $d_z = \int_{I_z} dw > 0$ . To end the deduction, all we need to prove is that

$$\bar{M}_{nd}(z) = 0, \quad z\text{-a.e.}.$$
 (80)

To do so, let us consider the opposite, so that there is some interval with nonzero measure I such that

$$\int_{I} \bar{M}_{nd}(z) \neq 0 \tag{81}$$

In this case, there must be two normalized vectors  $|\psi\rangle$  and  $|\phi\rangle$  such that

$$\langle \phi | \int_{I} \bar{M}_{nd}(z) | \psi \rangle \neq 0.$$
 (82)

Without loss of generality, let us assume that  $\psi(w, y) = \langle w(H_1, U), y | \psi \rangle$  and  $\psi(w, y) = \langle w(H_1, U), y | \psi \rangle$  are nonzero only when  $w \in I_2$  and  $w \in I_3$ , respectively, with  $I_2$  and  $I_3$  open intervals. Therefore,

$$|\psi\rangle = \int_{I_2} dw \int_{-\infty}^{\infty} dy \, |w(H_1, U), y\rangle \, \psi(w, y) \tag{83}$$

and

$$|\phi\rangle = \int_{I_3} dw \int_{-\infty}^{\infty} dy \, |w(H_1, U), y\rangle \, \phi(w, y). \tag{84}$$

$$\langle w'(H_1, U), y' | \bar{M}_{nd}(z) | w''(H_1, U), y'' \rangle = 0$$
(85)

for any  $w', w'' \in I$ , it cannot be the case that Eq.(82) holds. On the other hand, suppose that  $I_2 \cap I = I_3 \cap I = \emptyset$ , then since

$$\langle \psi | \int_{I_2} dz M'(z, H_1, U) | \psi \rangle = \langle \phi | \int_{I_3} dz M'(z, H_1, U) | \phi \rangle = 1,$$
(86)

it follows that

$$\langle \psi | \int_{I} dz M'(z, H_1, U) | \psi \rangle = \langle \phi | \int_{I} dz M'(z, H_1, U) | \phi \rangle = 0.$$
(87)

To deduce this, notice that  $M'(z, H_1, U)$  are elements of POVM and for any state  $|\psi'\rangle$  and interval  $I', \langle\psi'| \int_{I'} dz M'(z, H_1, U) |\psi'\rangle \ge 0, \langle\psi| \int_{-\infty}^{\infty} dz M'(z, H_1, U) |\psi\rangle = 1$ . Assuming that Eq.(82) holds, we can define

$$\Phi_{23} = \langle \phi | \int_{I} dz \bar{M}_{nd}(z) | \psi \rangle \neq 0$$
(88)

so that, by assuming the state

$$|\psi_{23}\rangle = \frac{a\,|\psi\rangle + b\,|\phi\rangle}{|a|^2 + |b|^2} \tag{89}$$

we get

$$\langle \psi_{23} | \int_{I} dz M'(z, H_1, U) | \psi_{23} \rangle = \frac{a b^* \Phi_{23} + b a^* \Phi_{23}^*}{|a|^2 + |b|^2}$$
(90)

Considering  $b = \Phi_{23}$  and a = -1, we get that

$$\langle \psi_{23} | \int_{I} dz M'(z, H_1, U) | \psi_{23} \rangle = -\frac{2|\Phi_{23}|^2}{1 + |\Phi_{23}|^2} < 0, \tag{91}$$

which cannot be the case for a POVM. Therefore, for  $I_2 \cap I = I_3 \cap I = \emptyset$ , Eq. (82) cannot hold.

Now, let us consider the case in which  $I_2 \subset I$  and  $I_3 \cap I = \emptyset$ . Then, since

$$\langle \phi | \int_{I_3} dz M'(z, H_1, U) | \phi \rangle = 1, \tag{92}$$

and, therefore,

$$\langle \phi | \int_{I} dz M'(z, H_1, U) | \phi \rangle = 0, \tag{93}$$

it follows that we can consider the state in the form  $|\psi_{23}\rangle$  once again to obtain

$$\langle \psi_{23} | \int_{I} dz M'(z, H_1, U) | \psi_{23} \rangle = \frac{|a|^2 + ab^* \Phi_{23} + ba^* \Phi_{23}^*}{|a|^2 + |b|^2}$$
(94)

Considering a = 1 and  $b = \Phi_{23}$ , we get

$$\langle \psi_{23} | \int_{I} dz M'(z, H_1, U) | \psi_{23} \rangle = \frac{1 + 2|\Phi_{23}|^2}{1 + |\Phi_{23}|^2} > 1,$$
(95)

which cannot hold for a POVM. Therefore, for  $I_2 \subset I$  and  $I_3 \cap I = \emptyset$ , Eq. (82) cannot hold. A similar argument can be given to deduce that for the case in which  $I_3 \subset I$  and  $I_2 \cap I = 0$ , Eq. (82) cannot hold as well.

With the previous arguments, we can thus conclude that for any vectors  $|\psi\rangle$  and  $|\phi\rangle$  with wave functions  $\langle w(H_1, U)|\psi\rangle$  and  $\langle w(H_1, U)|\psi\rangle$  with nonzero values for  $w \in I_2$  and  $w \in I_3$  such that  $I_2 \cap I = \emptyset$ ,  $I_2 \subset I$ ,  $I_3 \cap I = \emptyset$  or  $I_3 \subset I$ , Eq. (82) cannot hold.

Let us consider the general case in which  $I_2$  and  $I_3$  are not necessarily subsets of I and  $I_2 \cap I \neq \emptyset$  and  $I_3 \cap I \neq \emptyset$ . In this case, we can separate  $I'_2 = I_2 \cap I$  and  $I'_3 = I_3 \cap I$  and their complements  $I'_4 = I_2 \setminus I'_2$  and  $I'_5 = I_3 \setminus I'_3$ . Now, assuming Eq. (82) to hold implies that there is a  $k \in \{2, 4\}$  and  $j \in \{3, 5\}$ , such that

$$|\psi_{k}\rangle = \frac{1}{N'_{k}} \int_{I'_{k}} dw \int_{-\infty}^{\infty} dy \, |w(H_{1}, U)\rangle \, \psi(w, y), \quad N'_{k} = \int_{I'_{k}} dw \int_{-\infty}^{\infty} dy |\psi(w, y)|^{2}$$
(96)

and

$$|\phi_{j}\rangle = \frac{1}{N'_{j}} \int_{I'_{j}} dw \int_{-\infty}^{\infty} dy \,|w(H_{1}, U)\rangle \,\phi(w, y), \quad N'_{j} = \int_{I'_{j}} dw \int_{-\infty}^{\infty} dy |\phi(w, y)|^{2}$$
(97)

and

$$\langle \phi_j | \int_I \bar{M}_{nd}(z) | \psi_k \rangle \neq 0 \tag{98}$$

holds. If Eq. (98) does not hold for all  $k \in \{2, 4\}$  and  $j \in \{3, 5\}$ , then Eq.(82) cannot hold. However, notice that here either  $I'_j, I'_k \cap I = \emptyset$  or  $I'_j, I'_k \subset I$ . For these cases, we already concluded that Eq. (98) cannot hold in general. As a result, for general  $I_2$  and  $I_3$  and I, it cannot be the case that Eq. (82) holds. Consequently, there is no nonzero measure interval I such that

$$\int_{I} dz \bar{M}_{nd}(z) \neq 0.$$
<sup>(99)</sup>

As a result,  $M_{nd}(z) = 0$  z-a.e.. Considering this result, together with Eq.(78), we conclude that

$$M'(z, H_1, U) = M_{\text{OBS}}(z, H_1, U)$$
 z-a.e.. (100)

Given that we suppose arbitrary  $H_1$ , U, and arbitrary CRIN protocol  $\mathbb{M}'$ , then Eq.(100) is valid for any  $H_1$ , U and  $\mathbb{M}'$ . Therefore, there cannot be a CRIN protocol  $\mathbb{M}'$  different then a OBS protocol.

# C. Final remarks regarding result 1

By considering  $\Delta(H_1, U)$  to have continuous or discrete, degenerate or non-degenerate spectrum, we proved that there is no CRIN protocol different from the OBS protocol. As a result, for any  $\Delta(H_1, U)$  this result is valid and therefore, Result 1 is fully proved.

## **III. COMPLETE PROOF OF RESULT 2**

We derive in the present section result 2. As in the scope of result 1, we consider a unitary evolution U with countable basis and an energy operator of interest  $H_1$  acting on the same Hilbert space  $\mathcal{H}$  as U. We denote  $\{|\delta_j(H_1, U)\rangle\}$  as the basis that diagonalizes  $\Delta(H_1, U)$ , which can be discrete, continuous, degenerate or not. Considering this notation, we restate result 2 here for completeness:

**Result 2.** For any unitary evolution U, energy operator  $H_1$  acting on a Hilbert space  $\mathcal{H}$ , and eigenstate  $|\delta_i(H_1, U)\rangle$  of  $\Delta(H_1, U)$ , there exists a unitary U' acting on an auxiliary Hilbert space  $\mathcal{H}'$ , an additional Hamiltonian  $H_2$  acting on  $\mathcal{H} \otimes \mathcal{H}'$ , and a vector  $|v\rangle \in \mathcal{H}'$  such that the following equations are satisfied:

$$[H_1 \otimes \mathbb{1}_{\mathcal{H}'} + H_2, U \otimes U'] = 0, \tag{101}$$

$$H_2 \left| \delta_i(H_1, U), v \right\rangle = E_i \left| \delta_i(H_1, U), v \right\rangle, \tag{102}$$

$$(U^{\dagger} \otimes U'^{\dagger})H_2(U \otimes U') |\delta_i(H_1, U), v\rangle = E'_i |\delta_i(H_1, U), v\rangle, \qquad (103)$$

where  $|\delta_i(H_1, U), v\rangle = |\delta_i(H_1, U)\rangle \otimes |v\rangle$ , and  $E_i$  and  $E'_i = E_i - \delta_i(H_1, U)$  are real numbers.

To deduce this result, we separated the proof in many steps, considering a sequence of Lemmas. We present these Lemmas and then consider its main proof.

## A. Lemmas necessary for the deduction of Result 2

Throughout our main text, we assumed U to have a countable basis  $\{|u_i\rangle\}$ , then it follows that

$$U|u_i\rangle = u_i|u_i\rangle, \quad U^{\dagger}|u_i\rangle = u_i^{-1}|u_i\rangle, \quad u_i = e^{i\theta_i}$$
(104)

hold, where  $\theta_i \in \mathbb{R}$ . We define the countable set  $\Theta \subset \mathbb{R}$  containing all the arguments  $\theta_m$  defining the eigenvalues  $e^{i\theta_m}$  of U. Taking into account these definitions, we consider the following lemma:

**Lemma 1.** For any unitary U with countable basis, with the eigenvalues' arguments on  $\Theta$ , there exist at least one unitary operator U' acting on a Hilbert space H' such that:

1. U' is diagonalized by a countable basis  $\{|u'_i\rangle\}$ 

$$U' |u'_{j}\rangle = u'_{j} |u'_{j}\rangle, \quad U'^{\dagger} |u'_{j}\rangle = u'^{-1} |u'_{j}\rangle, \quad u'_{j} = e^{i\theta'_{j}}.$$
(105)

2. The countable set  $\Theta'$  containing all  $\theta'_j$  satisfying Eq. (105) is such that for any  $\theta'_j \in \Theta'$ , any  $\theta_k \in \Theta$ , and any  $\theta_l \in \Theta$ , there exist countably infinitely many  $\theta'_m \in \Theta'$  such that

$$\theta'_m + \theta_l = \theta'_i + \theta_k \mod 2\pi, \tag{106}$$

where mod  $2\pi$  denotes the sum modulo  $2\pi$ .

*Proof.* To prove our result, we only need to show that for an arbitrary U, there is at least one U' that satisfy the statement of the Lemma. In this sense, for an arbitrary U satisfying Eq. (104), we propose the following unitary:

$$U' = \sum_{n=0}^{\infty} e^{i\theta'_n} |n\rangle \langle n|$$
(107)

acting on a Hilbert space  $\mathcal{H}$  with countably infinite eigenbasis  $\{|n\rangle\}$  (e.g.  $|n\rangle$  can be the energy basis of an oscillator). Here,  $\theta'_n$  are real variables, each one related with a projector  $|n\rangle\langle n|$ . As a direct result of the definition (107), the set  $\{\theta'_n\} = \Theta'$  of all arguments  $\theta'_n$  is countable as well.

The definition of each  $\theta'_n$  is the most crucial so that condition 2 is satisfied. In this direction, we consider a procedure to define  $\theta'_i$  and then we prove that by using this procedure, all  $\theta'_i$  can be defined. For that consider an arbitrary natural  $l \ge 1$  and define

$$k_l = \left(\sum_{m=1}^{l-1} (2m+1)^m\right) + 1.$$
(108)

The  $\theta'_p$  for  $k_l \le p \le k_{l+1} - 1$  is defined using the following algorithm, which we call the permutation algorithm.

#### **Permutation algorithm**

Set  $k = k_l$  and then do the following:

• For  $i_1$  from -l to l do:

```
For i<sub>2</sub> from -l to l do:
∴
* For i<sub>l</sub> from -l to l do:
Define: θ'<sub>k</sub> = ∑<sup>l</sup><sub>j=1</sub> i<sub>j</sub>θ<sub>j</sub>, given that θ<sub>j</sub> ∈ Θ defined in Eq. (104);
Set k → k + 1;
* End i<sub>l</sub> loop.
∴
- End i<sub>2</sub> loop
• End i<sub>1</sub> loop
```

## End of the Algorithm

In other words, the set of  $\{\theta'_n\}$  defined in this procedure contains all the  $k_{l+1} - k_l = (2l+1)^l$  combinations of the type

$$\sum_{j=1}^{l} m_j \theta_j \tag{109}$$

with integers  $m_j$  satisfying  $|m_j| < l$ . Now, we show that using the permutation algorithm, we are able to define all  $\theta'_j$  in Eq. (107). First, notice that by considering l = 1, then we have, from Eq. (108) that  $k_1 = 1$  and, using the permutation algorithm:

$$\theta_1' = -\theta_1 \ \ \theta_2' = 0 \ \ \theta_3' = \theta_1 \tag{110}$$

Similarly, for l = 2, it follows that  $k_2 = 4$  and:

$$\begin{aligned} \theta'_{4} &= -2\theta_{2} - 2\theta_{1}, \ \theta'_{5} &= -2\theta_{2} - \theta_{1}, \ \theta'_{6} &= -2\theta_{2}, \ \theta'_{7} &= -2\theta_{2} + \theta_{1}, \ \theta'_{8} &= -2\theta_{2} + 2\theta_{1}, \ \theta'_{9} &= -\theta_{2} - 2\theta_{1}, \\ \theta'_{10} &= -\theta_{2} - \theta_{1}, \ \theta'_{11} &= -\theta_{2}, \ \theta'_{12} &= -\theta_{2} + \theta_{1}, \ \theta'_{13} &= -\theta_{2} + 2\theta_{1}, \ \theta'_{14} &= -2\theta_{1}, \ \theta'_{15} &= -\theta_{1}, \ \theta'_{16} &= 0, \\ \theta'_{17} &= \theta_{1}, \ \theta'_{18} &= 2\theta_{1}, \ \theta'_{19} &= \theta_{2} - 2\theta_{1}, \ \theta'_{20} &= \theta_{2} - \theta_{1}, \ \theta'_{21} &= \theta_{2}, \ \theta'_{22} &= \theta_{2} + \theta_{1}, \ \theta'_{23} &= \theta_{2} + 2\theta_{1}, \ \theta'_{24} &= 2\theta_{2} - 2\theta_{1}, \\ \theta'_{25} &= 2\theta_{2} - \theta_{1}, \ \theta'_{26} &= 2\theta_{2}, \ \theta'_{27} &= 2\theta_{2} + \theta_{1}, \ \theta'_{28} &= 2\theta_{2} + 2\theta_{1}, \end{aligned}$$
(111)

For l = 3, it follows that  $k_3 = 29$  and so on. Now, there are two options for the basis of U: either it is countably infinite or finite. Let us first consider that U is diagonalized by an countably infinite basis. Consider that  $\theta'_p$  is defined for  $1 \le p \le k_n - 1$ , where n is an arbitrary natural. Then, since the permutation algorithm used for the l = n, defines  $\theta'_p$  for all p such that  $k_n \le p \le k_{n+1} - 1$ , it follows that  $\theta'_p$  becomes defined for  $1 \le p \le k_{n+1} - 1$ . Therefore, by induction, we can repeat the procedure for arbitrary large l, and  $\theta'_p$  is defined for all  $p \ge 1$ . The set  $\Theta'$  containing all  $\theta'_p$  defined by this induction procedure is thus countably infinite.

In the case that *U* is diagonalized by a countably finite basis of dimension  $D < \infty$ , it follows that  $\theta_j$  is defined for  $1 \le j \ge D$ . We can define an extension  $\Theta_e = \Theta \bigcup_{k=1}^{\infty} \{\theta_{D+k}\}$  of  $\Theta$  by adding to it the extra variables  $\theta_{D+1} = \theta_1$ ,  $\theta_{D+2} = \theta_2$ ,  $\cdots$ ,  $\theta_{2D} = \theta_D$ ,  $\theta_{2D+1} = \theta_1$ ,  $\theta_{2D+2} = \theta_2$ , and inductively repeat this procedure. In this case,  $\theta'_j$  can be defined using the same procedure as in the case in which *U* is diagonalized by a countably infinite basis, but considering the elements  $\Theta_e$  instead of  $\Theta$ .

Now, we prove that all sums of the form

$$\sum_{j=1}^{N} m_j \theta_j \tag{112}$$

with finite natural  $N \ge 1$  and finite integers  $m_j$ , are inside  $\Theta'$ . To prove this, notice that, because  $m_j$  are finite, then there is an integer *s* such that  $|m_j| \le s$ . There are two possibilities: either  $s \ge N$  or s < N. Let us consider that  $s \ge N$ . In this case, if we consider the permutation algorithm for l = s, then the set all possible combinations

$$\sum_{j=1}^{s} i_j \theta_j \tag{113}$$

for  $|i_j| \le s$  will be contained in the set of all  $\theta'_k$  defined in the algorithm for l = s. Therefore, there is at least one  $\theta'_k$  defined in the algorithm for l = s equal to the sum in Eq. (112): the sum in Eq. (112) is contained in  $\Theta'$ . Now, consider that s < N. If we consider the permutation algorithm for l = N, then the set all possible combinations

$$\sum_{j=1}^{N} i_j \theta_j \tag{114}$$

for  $|i_j| \le N$  will be contained in the set of all  $\theta'_k$  defined in the algorithm for l = N. Therefore, there is at least one  $\theta'_k$  defined in the algorithm for l = N equal to the sum in Eq. (112). Since N is arbitrary as well as  $m_j$  then every combination as in Eq. (112) will be contained in  $\Theta'$ .

It is also straightforward to show (just by looking at the permutation algorithm) that all  $\theta'_j \in \Theta'$  are of the form of Eq. (112). As a result, all terms inside  $\Theta'$  are of the form of Eq. (112) and all the terms of the form of Eq. (112) are inside  $\Theta'$ .

To prove condition 2, select arbitrary  $\theta'_i \in \Theta'$ ,  $\theta_k \in \Theta$ , and  $\theta_q \in \Theta$ . Because  $\theta'_i$  is inside  $\Theta'$ , then it can be written as

$$\theta_j' = \sum_{n=1}^N m_n \theta_n \tag{115}$$

where N, and  $|m_n|$  are finite. As a result,

$$\theta'_j + \theta_k - \theta_q = \sum_{n=1}^N m_n \theta_n + \theta_k - \theta_q$$
(116)

Now, there are infinitely many  $\theta_m$  such that

$$\theta_m = \theta'_i + \theta_k - \theta_q. \tag{117}$$

To demonstrate this, let us define  $l_{\max} = \max\{N, k, q\}$ . For any finite natural  $l > l_{\max}$ , it is straightforward to check that, by applying the permutation algorithm for l, the element  $\theta'_j + \theta_k - \theta_q$  is defined. In other words, for every  $l > l_{\max}$ , there exists at least one *m* such that  $k_l \le m \le k_{l+1}$  and

$$\theta_m = \theta'_i + \theta_k - \theta_q. \tag{118}$$

Since the number of *l*'s such that  $l > l_{max}$  is countably infinite, then the set of all such  $\theta_m$  is also countably infinite. Given that *j*, *k*, and *n* are arbitrary, this establishes condition 2 of the lemma.

We call the conditions 1 and 2 of the Lemma as condition U from now on. Moreover, from this point on, we consider  $\Theta$  and  $\Theta'$  and U and U' as the sets and operators defined in Eq. (104), (105), and Lemma 1. Now, we prove the final two Lemmas needed for the proof of result 1.

**Lemma 2.** Consider a given  $H_1$  and U acting on a Hilbert space  $\mathcal{H}$ , such that the countable basis  $\{|u_i\rangle\}$  of U satisfy Eq. (104). Also, suppose that there is an hermitian operator  $H_2$  and an unitary U' acting on a Hilbert space  $\mathcal{H}'$ , and that the eigenvalues  $u'_i = e^{i\theta'_j}$  of U' satisfy Eq. (105). Moreover, suppose that

$$0 = \langle u_m, u'_i | [H_1 \otimes \mathbb{1}_{\mathcal{H}'} + H_2, U \otimes U'] | u_k, u'_i \rangle = (u_k u'_i - u_m u'_i) (\langle u_m | H_1 | u_k \rangle \delta_{jl} + \langle u_m, u'_i | H_2 | u_k, u'_l \rangle)$$
(119)

for every m, j, k, l such that

$$k = m \& l > j \quad or \quad k > m \& any l.$$
 (120)

Then it follows that

$$[H_1 \otimes \mathbb{1}_{\mathcal{H}'} + H_2, U \otimes U'] = 0.$$
(121)

*Proof.* To prove this result, all we need is to show that for m, j, k, l not satisfying Eq. (120), it can still be true that  $0 = \langle u_m, u'_j | [H_1 \otimes \mathbb{1}_{\mathcal{H}'} + H_2, U \otimes U'] | u_k, u'_l \rangle$ . For that, we first notice that if Eq. (119) is satisfied for m, j, k, l satisfying Eq. (120), then it follows that

$$(u_k u'_l - u_m u'_i)(\langle u_m | H_1 | u_k \rangle \,\delta_{jl} + \langle u_m, u'_j | H_2 | u_k, u'_l \rangle) = 0$$
(122)

which means that either (i)  $u_k u'_l - u_m u'_j = e^{i\theta_k + \theta'_l} - e^{i\theta_m + \theta'_j} = 0$ , or that (ii)  $\langle u_m | H_1 | u_k \rangle \delta_{jl} + \langle u_m, u'_j | H_2 | u_k, u'_l \rangle = 0$ , or both (i) and (ii) are simultaneously satisfied. If (i) is satisfied, then we can deduce that

$$\langle u_k, u'_l | [H_1 \otimes \mathbb{1}_{\mathcal{H}'} + H_2, U \otimes U'] | u_m, u'_j \rangle = (u_m u'_j - u_k u'_l) (\langle u_k | H_1 | u_m \rangle \delta_{jl} + \langle u_k, u'_l | H_2 | u_m, u'_j \rangle) = 0.$$
(123)

The same holds when (i) and (ii) are valid simultaneously. If, on the other hand, (i) is not valid, and (ii) holds, then  $\langle u_m | H_1 | u_k \rangle \delta_{jl} = -\langle u_m, u'_j | H_2 | u_k, u'_l \rangle$  and, given that  $H_1$  and  $H_2$  are hermitian, it follows that  $\langle u_k | H_1 | u_m \rangle^* \delta_{jl} = -\langle u_k, u'_l | H_2 | u_m, u'_j \rangle^*$ , which also implies in  $\langle u_k | H_1 | u_m \rangle \delta_{jl} = -\langle u_k, u'_l | H_2 | u_m, u'_j \rangle$ . As a result, it implies once again Eq. (123). Therefore, for any m, j, k, l such that  $\langle u_m, u'_j | [H_1 \otimes \mathbb{1}_{\mathcal{H}'} + H_2, U \otimes U'] | u_k, u'_l \rangle = 0$  it follows that  $\langle u_k, u'_l | [H_1 \otimes \mathbb{1}_{\mathcal{H}'} + H_2, U \otimes U'] | u_m, u'_j \rangle = 0$ . As a result, it follows that  $\langle u_m, u'_j | [H_1 \otimes \mathbb{1}_{\mathcal{H}'} + H_2, U \otimes U'] | u_k, u'_l \rangle = 0$  for any m, j, k, l satisfying

$$k = m & k \neq l \\ k = m & j < l \\ k < m & any l \\ k > m & any l. \end{cases}$$
(124)

The only case not considered to state that  $\langle u_m, u'_j | [H_1 \otimes \mathbb{1}_{\mathcal{H}'} + H_2, U \otimes U'] | u_k, u'_l \rangle = 0$  is the case where k = m and j = l. In this case, it is easy to show that  $(u_m u'_j - u_k u'_l) = (u_m u'_j - u_m u'_j) = 0$ . We can thus concluded that  $\langle u_k, u'_l | [H_1 \otimes \mathbb{1}_{\mathcal{H}'} + H_2, U \otimes U'] | u_m, u'_j \rangle = 0$  for every m, j, k, l, which results in  $[H_1 \otimes \mathbb{1}_{\mathcal{H}'} + H_2, U \otimes U'] = 0$ .

**Lemma 3.** Consider an arbitrary discrete and countable basis  $\{|n\rangle\}$  of an arbitrary discrete Hilbert space  $\mathcal{H}$ . We can always define a normalized vector  $|v\rangle \in \mathcal{H}$  such that  $\langle v|v\rangle = 1$  and the real and imaginary parts are non-null,  $\Re(\langle n|v\rangle) \neq 0$   $\Im(\langle n|v\rangle) \neq 0$ , for any of its components  $\langle n|v\rangle$ .

*Proof.* For a finite dimensional case, the result of the Lemma can be straightforwardly checked. Let us thus consider the infinite scenario. Since the basis is countable, then we can always assume a natural number *m* to identify the *m*-th element  $|m\rangle$  of the basis. Then, considering that  $m \ge 0$ , we can define  $\langle m|v\rangle$  as

$$\langle m|v\rangle = e^{i\eta_m} e^{-|\alpha|^2/2} \frac{(\alpha)^m}{\sqrt{(m)!}},\tag{125}$$

where  $\alpha \neq 0$  is a complex number and  $\eta_n$  is real. Notice that this state has the exactly same form as the components of the coherent state in the quantum oscillator for a complex  $\alpha$ , up to local phases  $\eta_m$ . Moreover,  $|v\rangle$  is normalized, since

$$\langle v|v\rangle = \langle v|\left(\sum_{m=0}^{\infty}|m\rangle\langle m|\right)|v\rangle = e^{-|\alpha|^2}\sum_{m=0}^{\infty}\frac{|\alpha|^{2m}}{m!} = 1.$$
(126)

Now, to show that  $\Re(\langle m|v\rangle)$ ,  $\Im(\langle m|v\rangle) \neq 0$ , first, notice that  $|\langle m|v\rangle| \neq 0$  for all *m*, given that  $\alpha \neq 0$ . Now, we can define  $\eta_m$  in the following form

$$\eta_m = 0 \quad \text{if } \Re(\alpha^m) \neq 0 \& \Im(\alpha^m) \neq 0$$
  

$$\eta_m = \pi/4 \text{ else.}$$
(127)

Since  $|\langle m|v\rangle| \neq 0$  for any natural *m*, then Eq. (127) ensure that  $\Re(\langle m|v\rangle), \Im(\langle m|v\rangle) \neq 0$ .

## B. The main proof of Result 2

Given all the Lemmas in the previous subsection, we are now in position to prove the result 2.

*Proof.* First, we consider a unitary U' that satisfies the U condition of Lemma 1. Therefore, it acts on a space  $\mathcal{H}'$  that satisfy Eq. (105), whose set  $\Theta' = \{\theta'_j\}$  of the arguments  $\theta'_j$  of its eigenvalues  $e^{i\theta'_j}$ . Because of Lemma 1, we know that such U' can always be defined. We thus consider the components  $[H_1 \otimes \mathbb{1}_{\mathcal{H}'} + H_2, U \otimes U']$ , in the basis  $|u_i, u'_k\rangle$  that diagonalizes  $U \otimes U'$  as follows:

$$\langle u_m, u_j | [H_1 \otimes \mathbb{1}_{\mathcal{H}'} + H_2, U \otimes U'] | u_k, u_l \rangle = (e^{i(\theta_k + \theta'_l)} - e^{i(\theta_m + \theta'_j)})(\langle u_m | H_1 | u_k \rangle \delta_{jl} + \langle u_m, u'_j | H_2 | u_k, u'_l \rangle)$$
(128)

To prove our result, we want to define  $H_2$  so as to make this expression to have a null value. In this sense, we recall that, because U' satisfy the U condition, then for any  $\theta'_i \in \Theta'$  and for any *m* and *k*, there exist infinitely many *l* such that  $\theta'_l \in \Theta'$  and

$$\theta_m + \theta'_i = \theta_k + \theta'_l \mod 2\pi. \tag{129}$$

As a result, it follows that for any *m*, *k* and *j*, there exist infinitely many *l* such that  $(e^{i(\theta_k + \theta'_l)} - e^{i(\theta_m + \theta'_j)}) = 0$ . Similarly, for any *m*, *k* and *j*, there exist infinitely many *l* such that  $(e^{i(\theta_k + \theta'_l)} - e^{i(\theta_m + \theta'_l)}) = 0$ . For the cases in which  $(e^{i(\theta_k + \theta'_l)} - e^{i(\theta_m + \theta'_l)}) = 0$ , we can consider any arbitrary value for  $\langle u_m, u'_i | H_2 | u_k, u'_l \rangle$  and still the following relation holds:

$$\langle u_m, u'_j | [H_1 \otimes \mathbb{1}_{\mathcal{H}'} + H_2, U \otimes U'] | u_k, u'_l \rangle = \left( e^{i(\theta_k + \theta'_l)} - e^{i(\theta_m + \theta'_j)} \right) (\langle u_m | H_1 | u_k \rangle \delta_{jl} + \langle u_m, u'_j | H_2 | u_k, u'_l \rangle) = 0.$$
(130)

As a consequence, we can define the free complex variables  $h_{mjkl}$  such that  $h_{klmj} = h_{mjkl}^*$ , so that the components of  $H_2$  are defined as

$$\langle u_m, u'_j | H_2 | u_k, u'_l \rangle = h_{mjkl} \quad \text{if} \quad (e^{i(\theta_k + \theta'_l)} - e^{i(\theta_m + \theta'_j)}) = 0 \langle u_m, u'_j | H_2 | u_k, u'_l \rangle = - \langle u_m | H_1 | u_k \rangle \, \delta_{jl} \quad \text{if} \quad (e^{i(\theta_k + \theta'_l)} - e^{i(\theta_m + \theta'_j)}) \neq 0$$

$$(131)$$

for the indexes

$$k = m \& l \ge j \quad \text{or} \quad k > m \& \text{ any } l, \tag{132}$$

and, by taking into account that  $H_1$  is hermitian so that  $\langle u_m | H_1 | u_k \rangle = \langle u_k | H_1 | u_m \rangle^*$ ,

$$\langle u_k, u'_l | H_2 | u_m, u'_j \rangle = \langle u_m, u'_j | H_2 | u_k, u'_l \rangle^*$$
(133)

for the indexes

$$k = m \& l < j \text{ or } k < m \& any l.$$
 (134)

Considering Eq. (131), we completely characterize the components  $\langle u_m, u'_j | H_2 | u_k, u'_l \rangle$  of  $H_2$  such as to ensure that  $H_2$  is hermitian and satisfy

$$0 = \langle u_m, u'_i | [H_1 \otimes \mathbb{1}_{\mathcal{H}'} + H_2, U \otimes U'] | u_k, u'_i \rangle = 0$$
(135)

for the indexes  $k = m \& l \ge j$  or k > m & any l. Considering Lemma 2, this means that by defining  $H_2$  as in Eqs. (131) and (133), it follows that

$$[H_1 \otimes \mathbb{1}_{\mathcal{H}'} + H_2, U \otimes U'] = 0, \tag{136}$$

so as to satisfy the first equation of result 2, *viz*. Eq. (101). We assume from this point on that  $H_2$  is defined by Eqs. (131) and (133).

Consider a state  $|v\rangle \in \mathcal{H}'$  whose only requirement is that its components  $\langle u'_j | v \rangle$  in the basis  $|u'_j\rangle$  satisfy  $\Re(\langle u'_j | v \rangle) \neq 0$  and  $\Im(\langle u'_j | v \rangle) \neq 0$ , i.e. the real and imaginary part of  $\langle u'_j | v \rangle$  are non-null. From Lemma 3 and the fact that the basis  $\{|u'_j\rangle\}$  is countable, we can always define such  $|v\rangle$ . Considering this  $|v\rangle$ , we can thus analyze how to define the free variables  $h_{klmj} = h^*_{mjkl}$  in (131) so that the eigenvalue/eigenvector equation

$$H_2 |\delta_i(H_1, U)\rangle \otimes |v\rangle = E_i |\delta_i(H_1, U)\rangle \otimes |v\rangle \equiv E_i |\delta_i(H_1, U), v\rangle$$
(137)

can be satisfied. For that, let us consider the components of (137) with respect to the specific element of the basis  $|u_m, u'_i\rangle$ :

$$\langle u_m, u'_j | E_i | \delta_i(H_1, U), v \rangle = E_i \alpha_{mi} \beta_j = \langle u_m, u'_j | H_2 | \delta_i(H_1, U), v \rangle = \sum_{k,l} \langle u_m, u'_j | H_2 | u_k, u'_l \rangle \alpha_{ki} \beta_l$$
(138)

where we named  $\alpha_{mi} = \langle u_m | \delta_i(H_1, U) \rangle$  and  $\beta_j = \langle u'_j | v \rangle$ . Consider the set  $\mathbb{N}_{mj}$  of all duple  $\{k, l\} \in \mathbb{N}_{mj}$  that, for the given component  $|u_m, u'_i\rangle$  satisfy  $(e^{i(\theta_k + \theta'_i)} - e^{i(\theta_m + \theta'_j)}) = 0$ . We can thus rewrite Eq. (138), considering Eqs. (131) and (133), as

$$E_{i}\alpha_{mi}\beta_{j} = -\sum_{\{k,l\}\notin\mathbb{N}_{mj}} \langle u_{m}|H_{1}|u_{k}\rangle \,\delta_{jl}^{\kappa}\alpha_{ki}\beta_{l} + \sum_{\{k,l\}\in\mathbb{N}_{mj}}h_{mjkl}\alpha_{ki}\beta_{l}.$$
(139)

Therefore, if we expect that  $H_2$  satisfy Eq. (137), then we need to properly define the free variables  $h_{mjkl}$  so that Eq. (139) holds. To do so, we first notice that there are only two possible alternatives: either  $\alpha_{mi} = 0$  and  $\alpha_{mi} \neq 0$ . We treat each of them separately.

In the case in which  $\alpha_{mi} = 0$ , for Eq. (139) for *m* and any arbitrary *j*, it is necessary that

$$\sum_{\{k,l\}\in\mathbb{N}_{mj}}h_{mjkl}\alpha_{ki}\beta_l = \sum_{\{k,l\}\notin\mathbb{N}_{mj}}\langle u_m|H_1|u_k\rangle\,\delta_{jl}^{\kappa}\alpha_{ki}\beta_l \tag{140}$$

holds. If  $|\delta_i(H_1, U)\rangle$  is a null vector so that  $\alpha_{ki} = 0$  for all  $k \neq m$ , then Eq. (137) holds directly for any value of  $h_{mjkl}$ . If, on the other hand,  $|\delta_i(H_1, U)\rangle$  is not a null vector, then there must be at least one  $k \neq m$  such that  $\alpha_{ki} \neq 0$ . Moreover, since  $\theta'_j \in \Theta'$ , then, from (129), for any m, k and j we can find infinitely many l such that  $\theta'_l \in \Theta'$  and  $\theta_k + \theta'_l = \theta_m + \theta'_j \mod 2\pi$ , i.e. such that  $\{k, l\} \in \mathbb{N}_{mj}$ . Therefore, there must exist a  $\{k, l\} \in \mathbb{N}_{mj}$  such that  $\alpha_{ki} \neq 0$ . Furthermore, since, by the definition of  $|v\rangle$ ,  $\beta_l \neq 0$  for all l, then for the specific m and j in Eq. (140) such that  $\alpha_{mi} = 0$ , there exist  $\{k, l\} \in \mathbb{N}_{mj}$  and  $k \neq m$ , such that  $\alpha_{ki}\beta_l \neq 0$ . As a result, the terms  $h_{mjkl}$ , being complex variables, can always be tuned to satisfy the Eq. (140) for all m and j such that  $\alpha_{mi} = 0$ . To show this, all we need is to show that there is at least one way of defining  $h_{mjkl}$  that make Eq. (140) to hold. In this sense, consider the set  $\mathbb{N}_{mj}^{\neq}$  of all the elements  $\{k, l\} \in \mathbb{N}_{mj}$  such that  $\alpha_{ki} \neq 0$ . Moreover, consider the indexes  $\bar{k}$  and  $\bar{l}$  as the smaller indexes in which  $\bar{k} + \bar{l} = \min_{\{k,l\} \in \mathbb{N}_{mj}} k + l$ . If there exist more then two duples  $\{\bar{k}', \bar{l}'\}$  and  $\{\bar{k}'', \bar{l}''\}$  such that  $\bar{k}' + \bar{l}' = \bar{k}'' + \bar{l}'' = \min_{\{k,l\} \in \mathbb{N}_{mj}} k + l$ , then we denote  $\{\bar{k}, \bar{l}\}$  as the duple with smallest from  $\bar{k}'$  and  $\bar{k}''$ . Let us define  $h_{mjkl}$  for all m and j such that  $\alpha_{mi} = 0$  as follows:

$$\begin{aligned} h_{mj\bar{k}\bar{l}} &= h_{\bar{k}\bar{l}mj}^* = \frac{1}{\alpha_{\bar{k}j}\beta_{\bar{l}}} \sum_{\{k,l\}\notin\mathbb{N}_{mj}} \langle u_m | H_1 | u_k \rangle \, \delta_{jl}^{\kappa} \alpha_{kj}\beta_l & \text{for } \{\bar{k},\bar{l}\} \\ h_{klmi}^* &= 0 & \text{for all } \{k,l\} \in \mathbb{N}_{mj}, \text{ such that } \{k,l\} \neq \{\bar{k},\bar{l}\}. \end{aligned}$$

Defining  $h_{mjkl}$  in this way, Eq. (140) will hold for any *m* and *j* such that  $\alpha_{mi} = 0$ . Importantly, notice that these definitions will not alter the other components of Eq. (137). To check that, we consider any other components  $|u_q, u'_r\rangle$  of Eq. (137), such that  $\{q, r\} \neq \{m, j\}$ :

$$\langle u_q, u_r' | H_2 | \delta_j(H_1, U) \rangle \otimes | v \rangle = E_i \alpha_{qi} \beta_r = -\sum_{sn \notin \mathbb{N}_{qr}} \langle u_q | H_1 | u_s \rangle \, \delta_{rn}^{\kappa} \alpha_{si} \beta_n + \sum_{sn \in \mathbb{N}_{qr}} h_{qrsn} \alpha_{si} \beta_n \tag{142}$$

The only way that Eq. (142) can be dependent on the terms  $h_{mjkl}$  for the specific *m*, *j* of Eq. (140) or their complex conjugate  $h_{mjkl}^* = h_{klmj}$  is either if, for the last sum in Eq. (142), we set (i) q = m, r = j, s = k, and n = l or (ii) q = k, r = l, s = m, and n = j. Since  $\{q, r\} \neq \{m, j\}$  then the case (i) is already not valid. On the other hand, in the case in which we consider (ii), then  $h_{qrsn}\alpha_{si} \rightarrow h_{klmj}\alpha_{mi}$  that will result in 0 for any  $h_{klmj}$ , since  $\alpha_{mi} = 0$ . So that, Eq. (142) will be independent of  $h_{mjkl}$  and  $h_{mjkl}^* = h_{klmj}$ . Since this holds for any  $\{q, r\} \neq \{m, j\}$ , then we proved that for any component  $|u_m, u'_j\rangle$  of Eq. (137) in which  $\alpha_{mi} = 0$ , we can define the independent variables  $h_{mjkl}$  as in Eq. (141) so as to make

$$\langle u_m, u'_j | H_2 | \delta_i(H_1, U), v \rangle = E_i \alpha_{mi} \beta_j = E_i \langle u_m, u'_j | \delta_i(H_1, U), v \rangle = 0 \quad \text{for } \alpha_{mi} = 0$$
(143)

to hold, in a way that does not interfere in the other components  $|u_q, u'_r\rangle \neq |u_m, u'_j\rangle$  of Eq. (137). We thus assume from now on  $h_{mikl}$  as in Eq. (141) for all *m* such that  $\alpha_{mi} = 0$ .

Now, let us consider the case in which  $\alpha_{mi} \neq 0$ . In this case, we rewrite Eq. (139) as

$$E_{i}\alpha_{mi}\beta_{j} = -\sum_{\substack{k\neq m, \{k,l\}\notin \mathbb{N}_{mj}}} \langle u_{m}|H_{1}|u_{k}\rangle \,\delta_{jl}^{\kappa}\alpha_{ki}\beta_{l} + \sum_{\substack{k\neq m, \{k,l\}\in \mathbb{N}_{mj}}} h_{mjkl}\alpha_{ki}\beta_{l} - \sum_{\substack{l\notin \mathbb{N}'_{j}}} \langle u_{m}|H_{1}|u_{m}\rangle \,\delta_{jl}^{\kappa}\alpha_{mi}\beta_{l} + \sum_{\substack{l\in \mathbb{N}'_{j}}} h_{mjkl}\alpha_{ki}\beta_{l} + \sum_{\substack{l\in \mathbb{N}'_{j}}} h_{mjkl}\alpha_{ki}\beta_{l} + \sum_{\substack{l\in \mathbb{N}'_{j}}} h_{mjml}\alpha_{mi}\beta_{l}.$$

$$(144)$$

where we denoted  $\mathbb{N}'_j$  as the set of indexes  $\{l\}$  such that  $\theta'_l = \theta'_j$  modulo  $2\pi$ . Therefore,  $l \in \mathbb{N}'_j$  and  $l \notin \mathbb{N}'_j$  means that  $e^{i\theta'_l} = e^{i\theta'_j}$ and  $e^{i\theta'_l} \neq e^{i\theta'_j}$ , respectively. Our goal, just as in the case in which  $\alpha_{mi} = 0$ , is to show that for each *m* and *j* there is at least one independent complex variable that can be tuned. Here, however, we have a difficulty since although  $h_{mjmj}$  is an independent variable, it must be *real* since  $H_2$  is hermitian. Moreover, the other terms  $h_{mjml}$  or  $h_{mjkl}$  for  $\{k, l\} \neq \{m, j\}$  cannot be considered as independent, since they can interfere in other components.

Our way to solve these problems is to consider an induction process. First, we consider

$$h_{mjml} = E_i \delta_{jl}^{\kappa} \tag{145}$$

for any *m* such that  $\alpha_{mi} \neq 0$  and any *j* and *l*. Moreover, consider that we order all indexes (because they are countable, we can always do this) and assume *m*(1) as the first index in which  $\alpha_{m(1)i} \neq 0$ . Consider the first index *j* = 1, such that Eq. (144) can be rewritten, considering Eq. (145), as:

$$E_{i}\alpha_{m(1)i}\beta_{1} = -\sum_{k \neq m(1), \{k\} \notin \mathbb{N}_{m(1)}^{\prime\prime}} \langle u_{m(1)} | H_{1} | u_{k} \rangle \alpha_{ki}\beta_{1} + \sum_{k \neq m(1), \{k,l\} \in \mathbb{N}_{m(1)1}} h_{m(1)1kl}\alpha_{ki}\beta_{l} + E_{i}\alpha_{m(1)i}\beta_{1}.$$
(146)

where  $\mathbb{N}''_{m(1)}$  is the set of indexes  $\{n\}$  such that  $\theta_n = \theta_{m(1)} \mod 2\pi$ . Now, let us consider the first k' such that  $\theta_{k'} \notin \mathbb{N}''_{m(1)}$  and  $\alpha_{k'i} \neq 0$ . By the definition of U' and of  $\Theta'$  as satisfying condition **U**, for all m(i), and k', there exist infinitely many l such that  $\theta_l' + \theta_{k'} = \theta_1' + \theta_{m(1)}$ , i.e. there are infinitely many l such that  $\{k', l\} \in \mathbb{N}_{m(1)1}$ . Let us define the smallest l such that  $\{k', l\} \in \mathbb{N}_{m(1)1}$  as l(k'). Moreover, from now on, let us denote, for any  $j, \beta_j = z_j e^{i\gamma_j}$ , where by the definition of  $\beta_j, z_j > 0$  is the real modulus and  $\gamma_j \neq 0, \pi/2 \mod 2\pi$ , is the argument. We then define the coefficients  $h_{m(1)1k'l}$ , with  $\{k', l\} \in \mathbb{N}_{m(1)1}$  as follows:

$$\begin{split} h_{m(1)lk'l} &= \langle u_{m(1)} | H_1 | u_{k'} \rangle e^{i(\gamma_1 - \gamma_l)} \frac{z_1}{z_l} & \text{if } l = l(k'), \{k', l\} \in \mathbb{N}_{m(1)1}, \text{ and } k' \notin \mathbb{N}'_{m(1)} \\ h_{m(1)lk'l} &= 0 & \text{if } l \neq l(k') \text{ and } \{k', l\} \in \mathbb{N}_{m(1)1}, \text{ or } k' \in \mathbb{N}'_{m(1)} \end{split}$$

$$\end{split}$$

$$(147)$$

Notice that, by definition, it follows that

$$\begin{array}{l} h_{k'lm(1)1} = \langle u_{k'} | H_1 | u_{m(1)} \rangle e^{i(\gamma_l - \gamma_1) \frac{z_1}{z_l}} & \text{if } l = l(k'), \{k', l\} \in \mathbb{N}_{m(1)1}, \text{ and } k' \notin \mathbb{N}'_{m(1)} \\ h_{k'lm(1)1} &= 0 & \text{if } l \neq l(k') \text{ and } \{k', l\} \in \mathbb{N}_{m(1)1}, \text{ or } k' \in \mathbb{N}'_{m(1)} \end{array}$$

$$(148)$$

By defining  $h_{m(1)1k'l}$  for  $\{k', l\} \in \mathbb{N}_{m(1)1}$  as in Eq. (147), we have that

$$\begin{split} h_{m(1)lk'l}\alpha_{k'i}\beta_l &= \langle u_{m(1)}|H_1|u'_k\rangle \,\alpha_{k'i}\beta_1 \quad \text{if } l = l(k'), \{k',l\} \in \mathbb{N}_{m(1)1}, \text{ and } k' \notin \mathbb{N}'_{m(1)} \\ h_{m(1)lk'l}\alpha_{k'i}\beta_l &= 0 \quad \text{if } l \neq l(k') \text{ and } \{k',l\} \in \mathbb{N}_{m(1)1}, \text{ or } k' \in \mathbb{N}'_{m(1)} \end{split}$$
(149)

Inserting this expression in Eq. (146), we have that

$$E_{i}\alpha_{m(1)i}\beta_{1} = -\sum_{k \neq m(1), k \neq k', \{k\} \notin \mathbb{N}'_{m(1)}} \langle u_{m(1)} | H_{1} | u_{k} \rangle \alpha_{ki}\beta_{1} + \sum_{k \neq m(1), k \neq k', \{k,l\} \in \mathbb{N}_{m(1)1}} h_{m(1)1kl}\alpha_{ki}\beta_{l} + E_{i}\alpha_{m(1)i}\beta_{1}.$$
(150)

In order words, we eliminated the first term k' of the first and second summation. Since the definition of  $h_{m(1)1k'l}$  does not alters the definition of  $h_{m(1)1kl}$  for other  $k \neq m(1)$ , then we can consider a similar definition for all the other  $k \neq m(1)$  in Eq.(152), as follows:

$$\begin{split} h_{m(1)1kl} &= \langle u_{m(1)} | H_1 | u_k \rangle \, \mathrm{e}^{i(\gamma_1 - \gamma_l)} \frac{z_1}{z_l} & \text{if } l = l(k), \, \{k, l\} \in \mathbb{N}_{m(1)1}, \, k \neq m(1), \, \alpha_{ki} \neq 0, \, k \notin \mathbb{N}'_{m(1)} \\ h_{m(1)1kl} &= 0 & \text{if } l \neq l(k), \, \{k, l\} \in \mathbb{N}_{m(1)1}, \, k \neq m(1), \, \text{and} \, \alpha_{ki} \neq 0, \, \text{or} \, k \in \mathbb{N}'_{m(1)} \end{split}$$
(151)

where l(k) is the smallest index l in which  $\{k, l\} \in \mathbb{N}_{m(1)1}$ . Using the definition in Eq. (151) for all  $k \neq m$  with  $\alpha_{ki} \neq 0$  in Eq. (152), we then obtain

$$\langle u_{m(1)}, u_1' | H_2 | \delta_i(H_1, U), v \rangle = E_i \alpha_{m(1)i} \beta_1$$
(152)

as we aimed.

Now, let us consider the second smallest index m(2) such that  $\alpha_{m(2)i} \neq 0$ . We then have the component  $|u_{m(2)}, u_1\rangle$  of Eq. (137), considering definition (145) can be written as

$$E_{i}\alpha_{m(2)i}\beta_{1} = -\sum_{k \neq m(2), \{k\} \notin \mathbb{N}'_{m(2)}} \langle u_{m(2)} | H_{1} | u_{k} \rangle \alpha_{ki}\beta_{1} + \sum_{k \neq m(2), \{k,l\} \in \mathbb{N}_{m(2)1}} h_{m(2)1kl}\alpha_{ki}\beta_{l} + E_{i}\alpha_{m(2)i}\beta_{1}$$
(153)

There are two possibilities: either  $\theta_{m(2)} = \theta_{m(1)}$ , so that  $m(2) \in \mathbb{N}'_{m(1)}$  or  $\theta_{m(2)} \neq \theta_{m(1)}$  and  $m(2) \notin \mathbb{N}'_{m(1)}$ . Let us first consider that  $m(2) \in \mathbb{N}'_{m(1)}$ . In this case, from the definition (151), it follows that  $h_{m(1)1m(2)l} = h^*_{m(2)lm(1)1} = h_{m(2)lm(1)1} = 0$  for any *l*. As a result, we can define as before, for any  $k \neq m(2)$  such that  $\alpha_{ki} \neq 0$ , the following:

$$\begin{split} h_{m(2)1kl} &= \langle u_{m(2)} | H_1 | u_k \rangle \, \mathrm{e}^{i(\gamma_1 - \gamma_l)} \frac{z_1}{z_l} & \text{if } l = l(k), \{k, l\} \in \mathbb{N}_{m(2)1}, \, k \neq m(2), \, \alpha_{ki} \neq 0, \, \text{and} \, k \notin \mathbb{N}'_{m(2)} \\ h_{m(2)1kl} &= 0 & \text{if } l \neq l(k), \, \{k, l\} \in \mathbb{N}_{m(2)1}, \, k \neq m(2), \, \text{and} \, \alpha_{ki} \neq 0, \, \text{or} \, k \in \mathbb{N}'_{m(2)} \end{split}$$
(154)

Notice that this condition does not contradicts the definition (154) for m(1), since in both cases we get  $h_{m(2)1m(1)1} = h_{m(1)1m(2)1} = 0$ when  $m(2) \in \mathbb{N}'_{m(1)}$  (and therefore  $m(1) \in \mathbb{N}'_{m(2)}$ ). Considering this definition in Eq. (153) we obtain again

$$\langle u_{m(2)}, u_1' | H_2 | \delta_i(H_1, U), v \rangle = E_i \alpha_{m(2)i} \beta_1.$$
(155)

Now, let us consider the case in which  $\theta_{m(2)} \neq \theta_{m(1)}$  so that  $m(2) \notin \mathbb{N}'_{m(1)}$ . In this case, for any  $k \neq m(1)$  and  $k \neq m(2)$  we can define  $h_{m(2)jkl}$  as in Eq. (154) and insert them in Eq. (153) to obtain

$$E_{i}\alpha_{m(2)i}\beta_{1} = -\langle u_{m(2)}|H_{1}|u_{m(1)}\rangle\alpha_{m(2)i}\beta_{1} + \sum_{\{m(1),l\}\in\mathbb{N}_{m(2)1}}h_{m(2)1m(1)l}\alpha_{m(1)i}\beta_{l} + E_{i}\alpha_{m(2)i}\beta_{1}.$$
(156)

where the summation  $\sum_{\{m(1),l\}\in\mathbb{N}_{m(2)1}}$  is over all l such that  $\{m(1),l\}\in\mathbb{N}_{m(2)1}$  for the fixed index m(1). Now, it cannot be the case that simultaneously  $m(2) \notin \mathbb{N}'_{m(1)}$  and 1 is equal to the smallest l such that  $\{m(2),l\}\in\mathbb{N}_{m(1)1}$ , since if  $m(2)\notin\mathbb{N}'_{m(1)}$  and therefore  $\theta_{m(1)}\neq\theta_{m(2)}$ , it cannot be the case that  $\theta'_1 + \theta_{m(1)} = \theta'_1 + \theta_{m(2)}$  and therefore  $\{m(1),1\}\notin\mathbb{N}_{m(2)1}$ . Therefore, for any l,  $h_{m(2)1m(1)l} = h^*_{m(1)lm(2)1}$  was not defined in the previous definitions in Eq. (151) for m(1). Thus, we can define  $h_{m(2)1m(1)l}$  as in Eq. (154) substituting  $k \to m(1)$  to obtain, considering Eq. (156),

$$\langle u_{m(2)}, u_1' | H_2 | \delta_i(H_1, U), v \rangle = E_i \alpha_{m(2)i} \beta_1.$$
(157)

This procedure can be repeated for all *m* in which  $\alpha_{mi} \neq 0$  for the same index j = 1 using the same reasoning, so that defining for all *m* 

$$\begin{aligned} h_{m1kl} &= \langle u_m | H_1 | u_k \rangle e^{i(\gamma_1 - \gamma_l) \frac{z_1}{z_l}} & \text{if } l = l(k), \{k, l\} \in \mathbb{N}_{m1}, k \neq m, \alpha_{ki} \neq 0, \text{ and } k \notin \mathbb{N}'_m \\ h_{m1kl} &= 0 & \text{if } l \neq l(k), \{k, l\} \in \mathbb{N}_{m1}, k \neq m, \text{ and } \alpha_{ki} \neq 0, \text{ or } k \in \mathbb{N}'_m, \end{aligned}$$

$$(158)$$

it follows that

$$\langle u_m, u_1' | H_2 | \delta_i(H_1, U), v \rangle = E_i \alpha_{mi} \beta_1 = E_i \langle u_m, u_1 | \delta_i(H_1, U), v \rangle.$$
(159)

Now, let us consider again m(1) but now we consider the second smallest index j = 2. We need to consider now the fact that it is possible that some  $h_{m(1)2k1} = h_{k1m(1)2}^*$  have already being defined in Eq. (158). Let us define the set  $\mathbb{O}_{12m(1)}$  as the set of indexes k such that  $h_{k1m(1)2}$  have already being defined in Eq. (158). For all  $k \notin \mathbb{O}_{12m(1)}$ , we can define

$$\begin{split} h_{m(1)2kl} &= \langle u_{m(1)} | H_1 | u_k \rangle e^{i(\gamma_2 - \gamma_l) \frac{z_2}{z_l}} & \text{if } l = l'(k), \{k, l\} \in \mathbb{N}_{m(1)2}, k \neq m(1), \, \alpha_{ki} \neq 0, \, \text{and} \, k \notin \mathbb{N}'_{m(1)} \\ h_{m(1)2kl} &= 0 & \text{if } l \neq l'(k), \{k, l\} \in \mathbb{N}_{m(1)2}, k \neq m(1), \, \text{and} \, \alpha_{ki} \neq 0, \, \text{or} \, k \in \mathbb{N}'_{m(1)}, \end{split}$$
(160)

where now l'(k) is the minimum l such that  $\{k, l\} \in \mathbb{N}_{m(1)2}$  and l > 1. Considering this definition, it follows that the eigenvectoreigenvalue equation for the component  $|u_{m(1)}, u'_2\rangle$ , taking into account (145), becomes

$$E_{i}\alpha_{m(1)i}\beta_{2} = -\sum_{\{k\}\notin\mathbb{N}_{m(1)}^{\prime}}^{\prime} \langle u_{m(1)}|H_{1}|u_{k}\rangle \alpha_{ki}\beta_{2} + \sum_{\{k,l\}\in\mathbb{N}_{m(1)2}}^{\prime} h_{m(1)2kl}\alpha_{ki}\beta_{l} + \sum_{l\in\mathbb{N}_{m(1)2}^{\prime}} h_{m(1)2m(1)l}\alpha_{m(1)l}\beta_{l}.$$

$$= -\sum_{\{k\}\notin\mathbb{N}_{m(1)}^{\prime}}^{\prime} \langle u_{m(1)}|H_{1}|u_{k}\rangle \alpha_{ki}\beta_{2} + \sum_{\{k,1\}\in\mathbb{N}_{m(1)2}}^{\prime} h_{m(1)2kl}\alpha_{ki}\beta_{1} + \sum_{l\neq1\{k,l\}\in\mathbb{N}_{m(1)2}}^{\prime} h_{m(1)2kl}\alpha_{ki}\beta_{l} + E_{i}\alpha_{m(1)i}\beta_{2}.$$
(161)

where we considered the notation

$$\sum_{k \neq m(1), k \in \mathcal{O}_{12m(1)}}^{\prime} = \sum_{k \neq m(1), k \in \mathcal{O}_{12m(1)}}^{\prime} .$$
(162)

Now, as can be straightforwardly checked, it is generally expected that

$$-\sum_{\{k\}\notin\mathbb{N}'_{m(2)}}^{\prime}\langle u_{m(1)}|H_{1}|u_{k}\rangle\alpha_{ki}\beta_{2} + \sum_{\{k,1\}\in\mathbb{N}_{m(1)2}}^{\prime}h_{m(1)2k1}\alpha_{ki}\beta_{1} \neq 0$$
(163)

i.e. the first and second sum in the second equality of Eq. (161) will not vanish in general. In this case, we have to define  $h_{m(1)2kl}$  appropriately so as to satisfy the eigenvalue-eigenvector equation. We then propose the following definition for all  $k \neq m(1)$ ,  $k \in \mathbb{O}_{12m(1)}$ :

$$\begin{split} h_{m(1)2kl} &= \langle u_{m(1)} | H_1 | u_k \rangle \, \mathrm{e}^{i(\gamma_2 - \gamma_l)} \frac{z_2}{z_l} - h_{m(1)2k1} \mathrm{e}^{i(\gamma_1 - \gamma_l)} \frac{z_1}{z_l} & \text{if } l = l''(k), \{k, l\} \in \mathbb{N}_{m(1)2}, \, \alpha_{ki} \neq 0, \, \text{and } k \notin \mathbb{N}'_{m(1)} \\ h_{m(1)2kl} &= 0 & \text{if } l \neq l''(k), \{k, l\} \in \mathbb{N}_{m(1)2}, \, \text{and } \alpha_{ki} \neq 0, \, \text{or } k \in \mathbb{N}'_{m(1)}. \end{split}$$
(164)

where l''(k) is the smallest l such that  $\{k, l\} \in \mathbb{N}_{m(1)2}$  with  $k \in \mathbb{O}_{12m(1)}$  and l > 1. Notice that because there are infinitely many l such that  $\{k, l\} \in \mathbb{N}_{m(1)2}$ , then we can always find such l''(k). Inserting the definition Eq. (164) in Eq. (161) for all  $k \neq m(1)$ ,  $k \in \mathbb{O}_{12m(1)}$ , we obtain

$$\langle u_{m(1)}, u'_2 | H_2 | \delta_i(H_1, U), v \rangle = E_i \alpha_{m(1)i} \beta_2 = E_i \langle u_{m(1)}, u_2 | \delta_i(H_1, U), v \rangle.$$
(165)

Let us now consider the index m(2) and j = 2. There are two possibilities: either  $m(2) \in \mathbb{N}'_{m(1)}$  or  $m(2) \notin \mathbb{N}'_{m(1)}$ . Let us first consider that  $m(2) \in \mathbb{N}'_{m(1)}$ . In this case, from the definition (160) or (164), it follows that  $h_{m(1)2m(2)l} = h^*_{m(2)lm(1)2} = h_{m(2)lm(1)2} = 0$  for any *l*. Moreover, we need to consider once again the possibility that for some  $k \neq m(1)$  and  $k \neq m(2)$ ,  $h_{m(2)2k1} = h^*_{k1m(2)2}$  have already being defined in Eq. (158). Defining analogously as for m(1) the set  $\mathbb{O}_{12m(2)}$  as the set of indexes *k* such that  $h_{k1m(2)2}$  have already being defined in Eq. (158), then we can define for every  $k \neq m(2)$  and  $k \neq m(1)$  the following:

$$\begin{split} h_{m(2)2kl} &= \langle u_{m(2)} | H_1 | u_k \rangle \, e^{i(\gamma_2 - \gamma_l) \frac{z_2}{z_l}} - h_{m(2)2k1} e^{i(\gamma_1 - \gamma_l) \frac{z_1}{z_l}} & \text{if } l = l''(k), \{k, l\} \in \mathbb{N}_{m(2)2}, \, \alpha_{ki} \neq 0, \, k \notin \mathbb{N}'_{m(2)}, \, \text{and } k \in \mathbb{O}_{12m(2)} \\ h_{m(2)2kl} &= \langle u_{m(2)} | H_1 | u_k \rangle \, e^{i(\gamma_2 - \gamma_l) \frac{z_2}{z_l}} & \text{if } l = l''(k), \{k, l\} \in \mathbb{N}_{m(2)2}, \, \alpha_{ki} \neq 0, \, k \notin \mathbb{N}'_{m(2)}, \, \text{and } k \notin \mathbb{O}_{12m(2)} \\ h_{m(2)2kl} &= 0 & \text{if } l \neq l''(k), \{k, l\} \in \mathbb{N}_{m(2)2}, \, \text{and } \alpha_{ki} \neq 0, \, \text{or } k \in \mathbb{N}'_{m(2)}. \end{split}$$

where l''(k) is the smallest *l* such that  $\{k, l\} \in \mathbb{N}_{m(2)2}$  with  $k \in \mathbb{O}_{12m(1)}$  and l > 1. Using the definition (166) it follows that in the component  $|u_{m(2)}, u'_{21}\rangle$  of the eigenvalue-eigenvector equation (137), we obtain

$$\langle u_{m(2)}, u'_{2} | H_{2} | \delta_{i}(H_{1}, U), v \rangle = E_{i} \alpha_{m(2)i} \beta_{2} = E_{i} \langle u_{m(2)}, u_{2} | \delta_{i}(H_{1}, U), v \rangle$$
(167)

as we wanted. If, on the other hand,  $\theta_{m(2)} \neq \theta_{m(1)}$  so that  $m(2) \notin \mathbb{N}'_{m(1)}$ , then it cannot be the case that simultaneously  $m(2) \notin \mathbb{N}'_{m(1)}$ and 2 is equal to the smallest *l* such that  $\{m(2), l\} \in \mathbb{N}_{m(1)2}$ , since if  $m(2) \notin \mathbb{N}'_{m(1)}$  and therefore  $\theta_{m(1)} \neq \theta_{m(2)}$ , it cannot be the case that  $\theta'_2 + \theta_{m(1)} = \theta'_2 + \theta_{m(2)}$  and therefore  $\{m(1), 2\} \notin \mathbb{N}_{m(2)2}$ . Therefore, for any l > 1,  $h_{m(2)2m(1)l} = h^*_{m(1)lm(2)2}$  was not defined in the previous definitions in Eq. (151). Thus, we can define  $h_{m(2)2m(1)l}$  as in Eq. (166) substituting  $k \to m(2)$  to obtain Eq. (167) when  $\theta_{m(2)} \neq \theta_{m(1)}$  as well.

Using the same arguments as for m(2), we can consider for any m the definition:

$$\begin{split} h_{m2kl} &= \langle u_m | H_1 | u_k \rangle e^{i(\gamma_2 - \gamma_l)} \frac{z_2}{z_l} - h_{m2k1} e^{i(\gamma_1 - \gamma_l)} \frac{z_1}{z_l} & \text{if } l = l''(k), \{k, l\} \in \mathbb{N}_{mj(2)}, \alpha_{ki} \neq 0, k \notin \mathbb{N}'_m, \text{ and } k \in \mathbb{O}_{12m} \\ h_{m2kl} &= \langle u_m | H_1 | u_k \rangle e^{i(\gamma_2 - \gamma_l)} \frac{z_2}{z_l} & \text{if } l = l''(k), \{k, l\} \in \mathbb{N}_{m2}, \alpha_{ki} \neq 0, k \notin \mathbb{N}'_m, \text{ and } k \notin \mathbb{O}_{12m} \\ h_{m2kl} &= 0 & \text{if } l \neq l''(k), \{k, l\} \in \mathbb{N}_{m2}, \alpha_{ki} \neq 0, \text{ or } k \in \mathbb{N}'_m. \end{split}$$
(168)

where  $\mathbb{O}_{12m}$  is the set of indexes k such that  $h_{m2k1}$  have already been defined, l''(k) is the smallest l > 1 such that  $\{k, l\} \in \mathbb{N}_{m2}$  with  $k \in \mathbb{O}_{12m}$ . Using this definition, it follows for any m that

$$\langle u_m, u_2 | H_2 | \delta_i(H_1, U), v \rangle = E_i \alpha_{mi} \beta_2 = E_i \langle u_m, u_2 | \delta_i(H_1, U), v \rangle.$$
(169)

Now, we want to prove a rule of induction considering the procedure that we did until now for all other indexes. Specifically, we want to define the component  $h_{m(1)jkl}$  of which  $\alpha_{m(1)i} \neq 0$  for any j, given that  $h_{mj'kl}$  was already defined using the same procedure for all k, l, m and j' = j - 1, j - 2,  $\cdots 1$  and that for all such m and j'

$$\langle u_m, u_{j'} | H_2 | \delta_i(H_1, U), v \rangle = E_i \alpha_{mi} \beta_{j'}.$$
(170)

Let us thus consider the indexes j and m(1). We define  $\mathbb{J}(j, m(1), k)$  as the set of all indexes  $\{j'\}$  such that  $j' \leq j$  and  $h_{m(1)jkj'} = h_{kj'm(1)j}$  was already defined in previous steps. Also, we define  $\mathbb{O}(j, m(1))$  of all the k indexes in which at least some j' < j, the term  $h_{m(1)jkj'}$  was already defined. Considering these sets, we then introduce the following definitions

$$h_{m(1)jkl} = \langle u_{m(1)} | H_1 | u_k \rangle \, \mathrm{e}^{i(\gamma_j - \gamma_l)} \frac{z_j}{z_l} - \sum_{j' \in \mathbb{J}(j,m(1),k)} h_{m(1)jkj'} \mathrm{e}^{i(\gamma_{j'} - \gamma_l)} \frac{z_{j'}}{z_l} \tag{171}$$

if l = l''(k), where l''(k) is the smallest l such that  $\{k, l\} \in \mathbb{N}_{m(1)j}$ , l > j,  $\alpha_{ki} \neq 0$ ,  $k \notin \mathbb{N}'_{m(1)}$ , and  $k \in \mathbb{O}(j, m(1))$ . If  $k \notin \mathbb{O}(j, m(1))$ , then we can consider the same definition l = l''(k), and consider

$$h_{m(1)jkl} = \langle u_{m(1)} | H_1 | u_k \rangle e^{i(\gamma_j - \gamma_l)} \frac{z_j}{z_l}$$
(172)

where  $\{k, l\} \in \mathbb{N}_{mj}, \alpha_{ki} \neq 0, k \notin \mathbb{N}'_{m(1)}$ . If  $l \neq l''(k), \{k, l\} \in \mathbb{N}_{m(1)j}, \alpha_{ki} \neq 0$ , and  $k \notin \mathbb{N}'_{m(1)}$  or in case that  $k \in \mathbb{N}'_{m(1)}$  then we define

$$h_{m(1)jkl} = 0. (173)$$

Notice that since for any k, j, and m(1), there are infinitely many l such that  $\{k, l\} \in \mathbb{N}_{m(1)j}$ , then l''(k) can always be defined.

Let us consider the eigenvalue-eigenvector equation for the component  $|u_{m(1)}, u'_{i}\rangle$ , considering (145):

$$E_{i}\alpha_{m(1)i}\beta_{j} = -\sum_{k \neq m(1), \{k\} \notin \mathbb{N}'_{m(1)}} \langle u_{m(1)} | H_{1} | u_{k} \rangle \alpha_{ki}\beta_{j} + \sum_{k \neq m(1), \{k,l\} \in \mathbb{N}_{m(1)j}} h_{m(1)jkl}\alpha_{ki}\beta_{l} + E_{i}\alpha_{m(1)i}\beta_{j}.$$
(174)

Now, considering the fact that for all  $k \in \mathbb{O}(j, m(1))$  and for such k, all indexes  $j' \in \mathbb{J}(j, m(1), k)$ ,  $h_{m(1)jkj'}$  was already defined for some j' < j, then Eq. (174) can be rewritten as

$$E_{i}\alpha_{m(1)i}\beta_{j} = -\sum_{k \neq m(1), \{k\} \notin \mathbb{N}'_{m(1)}} \langle u_{m(1)}| H_{1} | u_{k} \rangle \alpha_{ki}\beta_{j} + \sum_{k \in \mathbb{O}(j,m(1)), j' \in \mathbb{J}(j,m(1),k)} h_{m(1)jkj'}\alpha_{ki}\beta_{j'} + \sum_{k \in \mathbb{O}(j,m(1)), l} h_{m(1)jkl}\alpha_{ki}\beta_{l} + \sum_{k \notin \mathbb{O}(j,m(1)), l} h_{m(1)jkl}\alpha_{ki}\beta_{l} + E_{i}\alpha_{m(1)i}\beta_{j}.$$
(175)

where we consider the short notation:

$$\sum_{k \neq m(1), \{k,l\} \in \mathbb{N}_{m(1)j}} (176)$$

Substituting the definitions in Eqs. (171)-(173) in the third and second sum of Eq. (175), we can check that

$$0 = -\sum_{k \neq m(1), \{k\} \notin \mathbb{N}'_{m(1)}} \langle u_{m(1)} | H_1 | u_k \rangle \alpha_{ki} \beta_j + \sum_{k \in \mathbb{O}(j,m(1)), j' \in \mathbb{J}(j,m(1),k)} h_{m(1)jkj'} \alpha_{ki} \beta_{j'} + \sum_{k \in \mathbb{O}(j,m(1)), l} h_{m(1)jkl} \alpha_{ki} \beta_l + \sum_{k \notin \mathbb{O}(j,m(1)), l} h_{m(1)jkl} \alpha_{ki} \beta_l.$$
(177)

so that

$$\langle u_{m(1)}, u_j | H_2 | \delta_i(H_1, U), v \rangle = E_i \alpha_{m(1)i} \beta_j.$$
 (178)

Now, notice that for any r > 1, it follows that either  $m(r) \in \mathbb{N}'_{m(1)}$  or  $m(r) \notin \mathbb{N}'_{m(1)}$ . In the case in which  $m(r) \in \mathbb{N}'_{m(1)}$ , it follows that, in accordance with (173),  $h_{m(1)jm(r)l} = h^*_{m(r)lm(1)j} = h_{m(r)lm(1)j} = 0$  for any *l*. We can thus consider the same definitions as in Eqs. (171)-(173) substituting  $m(1) \to m(r)$  and this will not contradict the fact that  $h_{m(1)jm(r)j} = h^*_{m(r)jm(1)j} = 0$ . As

a result, it follows that using Eqs. (171)-(173), Eq. (179) will hold similarly substituting  $m(1) \rightarrow m(r)$ . On the other hand, if  $\theta_{m(r)} \neq \theta_{m(1)}$  and  $m(r) \notin \mathbb{N}'_{m(1)}$ , then it cannot be the case that simultaneously  $m(r) \notin \mathbb{N}'_{m(1)}$  and j is equal to the smallest l such that  $\{m(r), l\} \in \mathbb{N}_{m(1)j}$ . Therefore, for any l > j - 1,  $h_{m(r)jm(1)l} = h^*_{m(1)lm(r)j}$  was not yet defined. Thus, we can define  $h_{m(r)jkl}$  as in Eqs. (171)-(173) substituting  $m(1) \rightarrow m(r)$  to obtain Eq. (179) for any r. As a result, given that we assume for the induction process that Eq. (170) holds for any m and j' = j - 1, j - 2,  $\cdots$ , 1 and given the fact that for all  $r \ge 1$ , Eq. (179) holds substituting  $m(1) \rightarrow m(r)$ , then it follows that for all m and all j' = j, j - 1, j - 2,  $\cdots$ , 1, that

$$\langle u_m, u_{i'} | H_2 | \delta_i(H_1, U), v \rangle = E_i \alpha_{mi} \beta_{i'} = E_i \langle u_m, u_{i'} | \delta_i(H_1, U), v \rangle.$$
(179)

Therefore, by assuming that Eq. (170) holds for any *m* and all j' = j - 1, j - 2,  $\cdots 1$ , we proved that it also holds for all *m* and all j' = j, j - 1, j - 2,  $\cdots 1$ . Given that *j* is arbitrary here and that we proved Eqs. (159) and (169) for j = 1 and j = 2, respectively, then Eq. (170) holds for j = 1 and j = 2, and, by induction, it is valid for all  $j = 3, 4, 5, \cdots$ . As a result, for any *m* such that  $\alpha_{mi} \neq 0$  and any *j*,

$$\langle u_m, u'_i | H_2 | \delta_i(H_1, U), v \rangle = E_i \alpha_{mi} \beta_j.$$
(180)

Considering this result and (143), we can thus conclude that for the only two possible scenarios, in which either  $\alpha_{mi} \neq 0$  or  $\alpha_{mi} = 0$ , we can always tune  $h_{mjkl}$  that defines  $H_2$  in Eq. (131) to satisfy all the components  $|u_m, u'_j\rangle$  of Eq. (137). This means that we can always find a hermitian operator  $H_2$  that satisfy

$$[H_1 \otimes \mathbb{1}_{\mathcal{H}'} + H_2, U \otimes U'] = 0 \tag{181}$$

and

$$H_2 \left| \delta_i(H_1, U), v \right\rangle = E_i \left| \delta_i(H_1, U), v \right\rangle. \tag{182}$$

We conclude the proof of the result 2 by proving that the same  $H_2$  that satisfy Eqs. (181) and (182), also satisfy the relation

$$(U^{\dagger} \otimes U^{\dagger})H_2(U \otimes U')|\delta_i(H_1, U), v\rangle = E'_i|\delta_i(H_1, U), v\rangle, \qquad (183)$$

where  $E'_i = E_i - \delta_i(H_1, U)$ . To do so, we consider a  $H_2$  that satisfy Eqs. (181) and (182). Applying  $U^{\dagger} \otimes U'^{\dagger}$  on the left of both sides of Eq. (181), we obtain

$$\Delta(H_1, U) \otimes \mathbb{1}_{\mathcal{H}'} = U^{\dagger} H_1 U \otimes \mathbb{1}_{\mathcal{H}'} - H_1 \otimes \mathbb{1}_{\mathcal{H}'} = -[(U^{\dagger} \otimes U'^{\dagger})H_2(U \otimes U') - H_2] = -\Delta(H_2, U \otimes U').$$
(184)

Applying  $\Delta(H_2, U \otimes U')$  on  $|\delta_i(H_1, U), v\rangle$  considering the above equality, we get

$$[(U^{\dagger} \otimes U'^{\dagger})H_2(U \otimes U') - H_2]|\delta_i(H_1, U), v\rangle = \Delta(H_2, U \otimes U')|\delta_i(H_1, U), v\rangle = -\delta_i(H_1, U)|\delta_i(H_1, U), v\rangle$$
(185)

which means that  $|\delta_i(H_1, U), v\rangle$  is also an eigenvector of  $\Delta(H_2, U \otimes U')$  with eigenvalue  $-\delta_i(H_1, U)$ . Considering Eq. (185), the fact that  $H_2$  satisfy Eq. (182) and  $(U^{\dagger} \otimes U'^{\dagger})H_2(U \otimes U') = H_2 + \Delta(H_2, U \otimes U')$ , we deduce

$$(U^{\dagger} \otimes U^{\dagger})H_{2}(U \otimes U')|\delta_{i}(H_{1}, U), v\rangle = (H_{2} + \Delta(H_{2}, U \otimes U'))|\delta_{i}(H_{1}, U), v\rangle = (E_{i} - \delta_{i}(H_{1}, U))|\delta_{i}(H_{1}, U), v\rangle,$$
(186)

concluding the proof.

# IV. ADAPTING TO THE TIME-DEPENDENT CASE

It is interesting to notice that results 1 and 2 can be immediately adapted to any other quantum observalbe A substituting  $H_1 \rightarrow A$  in the results and deductions, given that U has a countable basis. Therefore, we can consider the substitution  $H_1 \rightarrow X, P, L, N, \cdots$ , i.e. position, linear and angular momentum, number of particles etc. In the present section, we show that we can adapt also our formalism for time-dependent cases, as long as we do adjustments in the CRIN conditions.

We then consider now that  $H_1$  can have a explicit time-dependence in the Schrödinger picture, so that  $H_1 \rightarrow H_1(t)$ , for any given time t. Similarly, we consider that such a system evolves according with a unitary U with countable basis. In this case, the two time observables defining the variation of  $H_1(t)$  on the interval is defined, for the discrete case, as

$$\Delta(H_1(0), H_1(t), U) = U^{\dagger} H_1(t) U - H_1(0) = \sum_i \delta_i(H_1(0), H_1(t), U) \left| \delta_i(H(0), H(t), U) \right\rangle,$$
(187)

where  $\Delta(H_1(0), H_1(t), U) |\delta_i(H(0), H(t), U)\rangle = \delta_i(H(0), H(t), U) |\delta_i(H(0), H(t), U)\rangle$ . An analogous definition can be made for the continuous and/or degenerate case. Notice that U in general will have the dependence of  $H_1(t')$  for all  $t' \in [0, t]$ . This dependence can thus be implicit in U and we take into account this fact in the notation of  $\Delta(H_1(0), H_1(t), U)$ .

Now, we define in analogy to the time-independent case, a time-dependent measurement protocol  $\mathbb{M}_t$  [6, 7] to measure the *variation* of any time-dependent arbitrary operator  $H_1(t)$  under an arbitrary evolution U; for the protocol  $\mathbb{M}_t$  and for each  $(H_1(0), H_1(t), U)$  triplet, the set  $\mathbb{M}_t(H_1(0), H_1(t), U) = \{M(z, H_1(0), H_1(t), U)\}$  defines a POVM whose operators satisfy the usual POVM properties, *viz.*  $\int_{-\infty}^{\infty} dz M(z, H_1(0), H_1(t), U) = \mathbb{1}$  and  $M(z, H_1(0), H_1(t), U) \ge 0$ . Given this definition, we redefine the CRIN conditions for the time-dependent case:

- 1. Conservation laws: For any preparation  $\rho$ , unitary evolution U, and time-dependent energy operators defined  $H_1(t)$  and  $H_2(t)$  representing parts of the energy of any system  $\Omega$ , if  $U^{\dagger}H_2(t)U + U^{\dagger}H_1(t)U = H_2(0) + H_1(0)$ , then, for any z,  $\text{Tr}[M(z, H_1(0), H_1(t), U)\rho] = \wp(z, H_1(0), H_1(t), U, \rho) = \wp(-z, H_2(0), H_2(t), U, \rho) = \text{Tr}[M(-z, H_2(0), H_2(t), U)\rho]$ .
- 2. **Reality**: For any system  $\Omega$ , operators  $H_1(0)$  and  $H_1(t)$ , and unitary evolution U, if there is an initial state  $\rho_1 = |e_1\rangle \langle e_1|$ such that  $|e_1\rangle$  is an eigenvector of both  $H_1(0)$  and  $U^{\dagger}H_1(t)U$  with respective eigenvalues  $e_1(0)$  and  $\epsilon_1(t)$ , then the POVM must result in the probabilities  $\wp(z, H_1(0), H_1(t), U) = \delta^{\wp}[z - (\epsilon_1(t) - e_1(0))]$ .
- 3. Independence of the initial state: For any system  $\Omega$ , operators  $H_1(0)$  and  $H_1(t)$ , and evolution operators U, the elements of the POVM  $M(z, H_1(0), H_1(t), U)$  must not depend on the initial state  $\rho$ .
- 4. No-signaling: For any system Ω, if Ω evolves under an arbitrary bipartide unitary evolution U ⊗ U' acting on a bipartide Hilbert space H = H<sub>1</sub> ⊗ H<sub>2</sub>, then, for every energy operators H<sub>1</sub>(0) ⊗ 1<sub>2</sub> and H<sub>1</sub>(t) ⊗ 1<sub>2</sub> acting locally on H<sub>1</sub>, M is such that its POVMs satisfy the following relation for every z: M(z, H<sub>1</sub>(0) ⊗ 1<sub>2</sub>, H<sub>1</sub>(t) ⊗ 1<sub>2</sub>, U ⊗ U') = M(z, H<sub>1</sub>(0), H<sub>1</sub>(t), U) ⊗ 1<sub>2</sub>.

Given this adaptation (and a similar adaptation of the reality condition for the continuous case), we now comment on how to adapt result 1 and 2 for the time-dependent case.

First, the following result is an immediate consequence of Result 2:

**Corollary 1.** For any unitary evolution U, energy operators  $H_1(0)$  and  $H_1(t)$  acting on a Hilbert space  $\mathcal{H}$ , and eigenstate  $|\delta_i(H_1(0), H_1(t), U)\rangle$  of  $\Delta(H_1(0), H_1(t), U)$ , there exists a unitary U' acting on an auxiliary Hilbert space  $\mathcal{H}'$ , an additional operators  $H_2(0)$  and  $H_2(t)$  acting on  $\mathcal{H} \otimes \mathcal{H}'$ , and a vector  $|v\rangle \in \mathcal{H}'$  such that the following equations are satisfied:

$$(U^{\dagger} \otimes U^{\dagger})H_1(t) \otimes \mathbb{1}_{\mathcal{H}'}(U \otimes U') + (U^{\dagger} \otimes U^{\dagger})H_2(U \otimes U') = H_1(0) \otimes \mathbb{1}_{\mathcal{H}'} + H_2(0) = 0,$$
(188)

$$H_2 |\delta_i(H_1(0), H_1(t), U), v\rangle = E_i |\delta_i(H_1(0), H_1(t), U), v\rangle,$$
(189)

$$(U^{\top} \otimes U^{\top})H_2(U \otimes U') |\delta_i(H_1(0), H_1(t), U), v\rangle = E'_i |\delta_i(H_1(0), H_1(t), U), v\rangle,$$
(190)

where  $|\delta_i(H_1(0), H_1(t), U), v\rangle = |\delta_i(H_1(0), H_1(t), U)\rangle \otimes |v\rangle$ , and  $E_i$  and  $E'_i = E_i - \delta_i(H_1(0), H_1(t), U)$  are real numbers.

*Proof.* By directly substituting  $H_1 \rightarrow H_1(0)$ ,  $|\delta_i(H_1, U)\rangle \rightarrow |\delta_i(H_1(0), H_1(t), U)\rangle$  in Result 2, we deduce that there is a  $H_2(0)$ , U', and  $|v\rangle$  that satisfy the following equations [?]:

$$[H_1(0) \otimes \mathbb{1}_{\mathcal{H}'} + H_2(0), U \otimes U'] = 0, \tag{191}$$

and

$$H_2(0)|\delta_i(H_1(0), H_1(t), U), v\rangle = E_i|\delta_i(H_1(0), H_1(t), U), v\rangle.$$
(192)

Applying  $U^{\dagger} \otimes U'^{\dagger}$  on both sides of Eq. (191), we get

$$(U^{\dagger} \otimes U^{\dagger})(H_1(0) \otimes \mathbb{1}_{\mathcal{H}'} + H_2(0))(U \otimes U') = H_1(0) \otimes \mathbb{1}_{\mathcal{H}'} + H_2(0).$$
(193)

Defining  $H_2(t) = H_1(0) \otimes \mathbb{1}_{\mathcal{H}'} + H_2(0) - H_1(t) \otimes \mathbb{1}_{\mathcal{H}'}$  and, considering Eqs. (191) and (193), we obtain

$$(U^{\dagger} \otimes U'^{\dagger})H_{2}(t)(U \otimes U') = (U^{\dagger} \otimes U'^{\dagger})(H_{1}(0) \otimes \mathbb{1}_{\mathcal{H}'} + H_{2}(0))(U \otimes U') - (U^{\dagger} \otimes U'^{\dagger})(H_{1}(t) \otimes \mathbb{1}_{\mathcal{H}'})(U \otimes U') = H_{1}(0) \otimes \mathbb{1}_{\mathcal{H}'} + H_{2}(0) - (U^{\dagger} \otimes U'^{\dagger})(H_{1}(t) \otimes \mathbb{1}_{\mathcal{H}'})(U \otimes U') = H_{1}(0) \otimes \mathbb{1}_{\mathcal{H}'} + H_{2}(0) - U^{\dagger}H_{1}(t)U \otimes \mathbb{1}_{\mathcal{H}'} = H_{2}(0) - (U^{\dagger}H_{1}(t)U \otimes \mathbb{1}_{\mathcal{H}'} - H_{1}(0) \otimes \mathbb{1}_{\mathcal{H}'}) = H_{2}(0) - (\Delta(H_{1}(0), H_{1}(t), U) \otimes \mathbb{1}_{\mathcal{H}'}).$$

$$(194)$$

As a result, it follows that

$$(U^{\dagger} \otimes U^{'\dagger})(H_2(t) + H_1(t) \otimes \mathbb{1}_{\mathcal{H}'})(U \otimes U') = H_2(0) + H_1(0) \otimes \mathbb{1}_{\mathcal{H}'}$$
(195)

and

$$(U^{\dagger} \otimes U^{\prime \dagger}) H_2(t) (U \otimes U^{\prime}) |\delta_i(H_1(0), H_1(t), U), v\rangle = (H_2(0) - (\Delta(H_1(0), H_1(t), U) \otimes \mathbb{1}_{\mathcal{H}^{\prime}})) |\delta_i(H_1(0), H_1(t), U), v\rangle$$

$$= E_i^{\prime} |\delta_i(H_1(0), H_1(t), U), v\rangle,$$
(196)

where  $E'_i = E_i - \delta_i(H_1(0), H_1(t), U)$ . Considering Eqs. (192), (195), and (196), the result is thus proved.

Corollary 1 is therefore the analogous of result 2, and follow almost immediately from it. Considering Corollary 1, an analogous of result 1 can be deduced substituting  $H_1 \rightarrow H_1(t)$ ,  $\Delta(H_1, U) \rightarrow \Delta(H_1(0), H_1(t), U)$ ,  $\mathbb{M} \rightarrow \mathbb{M}_t$ , and considering the adapted time-dependent CRIN conditions. As a consequence, the OBS protocol is also the only protocol that satisfies the time-dependent CRIN conditions for the time-dependent case.

# V. RESULTS NEEDED FOR THE ION TRAP EXAMPLES IN THE FIGURES

#### A. OBS and TPM for the commuting continuous case

In Ref. [8], it was shown that whenever a time-dependent Hamiltonian H(t) satisfies

$$[H(0), U^{\dagger}H(t)U] = 0, \tag{197}$$

then the TPM statistics for a thermal state  $\rho_{\beta} = Z^{-1} e^{-\beta H(0)}$  will be the same as the OBS statistics. The authors showed this result by taking into account the TPM characteristic function, defined as

$$G_{\text{TPM}}(u) = \int_{-\infty}^{\infty} dz e^{iuz} \wp_{\text{TPM}}(z, H(0), H(t), U, \rho_{\beta})$$
(198)

where, for the time-dependent case,

$$\varphi_{\text{TPM}}(z, H(0), H(t), U, \rho_{\beta}) = \text{Tr}[M_{\text{TPM}}(z, H(0), H(t), U)\rho_{\beta}],$$
(199)

and

$$M_{\text{TPM}}(z, H(0), H(t), U) = \sum_{jk} \delta^{\mathbf{p}}[z - (e_j(t) - e_k)] |\langle e_j(t) | U | e_k \rangle|^2 |e_k \rangle \langle e_k|.$$
(200)

Here,  $|e_j(t)\rangle$  and  $|e_k\rangle$  are eigenvectors of H(t) and H(0), respectively. Notice that  $M_{\text{TPM}}(z, H(0), H(t), U)$  is similar to the form defined in the main text, with adjustments due to the time-dependence. When H(t), H(0) and U satisfy Eq. (197), the authors of Ref. [8] proved that

$$G_{\text{TPM}}(u) = \int_{-\infty}^{\infty} dz e^{iuz} \varphi_{\text{TPM}}(z, H(0), H(t), U, \rho_{\beta}) = \sum_{jk} e^{iu(e_j(t) - e_k)} |\langle e_j(t)| U | e_k \rangle|^2 \langle e_k | \rho_{\beta} | e_k \rangle$$
  
= Tr[ $e^{iu(U^{\dagger}H(t)U - H(0))} \rho_{\beta}$ ], (201)

which will be equal to the OBS characteristic function:

$$G_{\rm OBS}(u) = \int_{-\infty}^{\infty} dz e^{iuz} \wp_{\rm OBS}(z, H(0), H(t), U, \rho) = \text{Tr}[e^{iu(\Delta(H(0), H(t), U))}\rho_{\beta}]$$
(202)

where, for the time-dependent case, we considered the two-time OBS statistics [7]

$$\wp_{\text{OBS}}(z, H(0), H(t), U) = \text{Tr}[M_{\text{OBS}}(z, H(0), H(t), U)\rho_{\beta}],$$
(203)

$$M_{\text{OBS}}(z, H(0), H(t), U) = \sum_{j} \delta^{\mathbf{D}}[z - \delta_{j}(H(0), H(t), U)] \left| \delta_{j}(H(0), H(t), U) \right\rangle \left\langle \delta_{j}(H(0), H(t), U) \right|,$$
(204)

and

$$\Delta(H(0), H(t), U) = U^{\dagger}H(t)U - H(0) = \sum_{j} \delta_{j}(H(0), H(t), U) \left| \delta_{j}(H(0), H(t), U) \right\rangle \left\langle \delta_{j}(H(0), H(t), U) \right|$$
(205)

Similarly, the same result can be deduced substituting in Eqs. (201) and (202) the following:  $H(0) \rightarrow H_1$ ,  $H(t) \rightarrow H_1$ ,  $\varphi_{OBS}(z, H(0), H(t), U) \rightarrow \varphi_{OBS}(z, H_1, U)$  and  $\varphi_{TPM}(z, H(0), H(t), U) \rightarrow \varphi_{TPM}(z, H_1, U)$  (defined in the main text), and  $\rho_{\beta} \rightarrow any \rho$ . Therefore, it follows, for the time-independent discrete basis case, that if  $[H_1, U^{\dagger}H_1U] = 0$ , then:

$$G_{\rm TPM}(u) = \int_{-\infty}^{\infty} dz e^{iuz} \wp_{\rm TPM}(z, H_1, U, \rho) = \int_{-\infty}^{\infty} dz e^{iuz} \wp_{\rm OBS}(z, H_1, U, \rho) = G_{\rm OBS}(u),$$
(206)

and, as a result,

$$\wp_{\text{TPM}}(z, H_1, U, \rho) = \wp_{\text{OBS}}(z, H_1, U, \rho).$$
 (207)

Now, for the deductions in [8] and the framework here, we considered discrete basis, so that TPM procedure is clear and needs no adaptation. This is not the case when  $H_1$  is diagonalized by a continuous basis  $\{|e\rangle\}$ , since after a first measurement of  $H_1$  of one round of a TPM procedure, the state is not normalizable after the measurement so that the statistics cannot be directly treated [9]. We can assume that after a first measurement, the state is a normalized state  $|\psi_{\mu}^e\rangle$ , such that  $\mu$  accounts for the precision of the experiment apparatus used and approximates an eigenstate  $|e\rangle$  whenever  $\mu \to 0$ . For instance,  $\langle e' | \psi_{\mu}^e \rangle$  can be a Gaussian state with its center located at e, such that  $\mu$  is the width of the gaussian. For such states, we expect that

$$\lim_{\mu \to 0} H_1^n |\psi_{\mu}^k\rangle = \lim_{\mu \to 0} e_k^n |\psi_{\mu}^k\rangle \quad \text{and} \quad \lim_{\mu \to 0} |\langle e|\psi_{\mu}^{e'}\rangle|^2 = \delta^{\mathbf{p}}(e'-e).$$
(208)

Whenever  $[H_1, U^{\dagger}H_1U] = 0$ , since and eigenvector  $|e\rangle$  is also eigenvector of  $U^{\dagger}H_1U$ , such that  $U^{\dagger}H_1U|e\rangle = \epsilon_e |e\rangle$  for some real  $\epsilon_e$ , then it also follows that

$$\lim_{\mu \to 0} U^{\dagger} H_1^n U |\psi_{\mu}^e\rangle = \lim_{\mu \to 0} \epsilon_e^n |\psi_{\mu}^e\rangle.$$
<sup>(209)</sup>

As a result, we can define the TPM POVM elements for the continuous case as

$$M_{\text{TPM}}(z, H_1, U) = \lim_{\mu \to 0} \iint_{-\infty}^{\infty} dede' \delta^{\mathbf{p}}[z - (e' - e)] |\langle e'| U |\psi_{\mu}^e \rangle|^2 |e\rangle \langle e|, \qquad (210)$$

So that, the characteristic function can be computed as

$$\begin{aligned} G_{\text{TPM}}(u) &= \int_{-\infty}^{\infty} dz e^{iuz} \wp_{\text{TPM}}(z, H_1, U, \rho) = \lim_{\mu \to 0} \iint_{-\infty}^{\infty} dede' e^{iu(\epsilon_{e'} - e)} |\langle \epsilon_{e'}| U | \psi_{\mu}^e \rangle|^2 \langle e|\rho|e \rangle \\ &= \lim_{\mu \to 0} \iint_{-\infty}^{\infty} dede' e^{iu(\epsilon_{e'} - e)} \langle e'| U | \psi_{\mu}^e \rangle \langle e|\rho|e \rangle \langle \psi_{\mu}^e | U^{\dagger}|e' \rangle, \\ &= \lim_{\mu \to 0} \iint_{-\infty}^{\infty} dede' \operatorname{Tr}[U^{\dagger}|e' \rangle e^{iu\epsilon_{e'}} \langle e'| U | \psi_{\mu}^e \rangle \langle e|e^{-iue\rho}\rho|e \rangle \langle \psi_{\mu}^e|] \\ &= \lim_{\mu \to 0} \iint_{-\infty}^{\infty} de \operatorname{Tr}[e^{iuU^{\dagger}H_1U} | \psi_{\mu}^e \rangle \langle e|e^{-iuH_1}\rho|e \rangle \langle \psi_{\mu}^e|] \\ &= \lim_{\mu \to 0} \iint_{-\infty}^{\infty} de \operatorname{Tr}[e^{iu\epsilon_{e'}} \langle e'|\psi_{\mu}^e \rangle \langle e|e^{-iuH_1}\rho|e \rangle \langle \psi_{\mu}^e|] \\ &= \lim_{\mu \to 0} \iint_{-\infty}^{\infty} de \int_{-\infty}^{\infty} de' e^{iu\epsilon_{e'}} \langle e|e^{-iuH_1}\rho|e \rangle \langle \psi_{\mu}^e|e' \rangle \\ &= \lim_{\mu \to 0} \iint_{-\infty}^{\infty} de \int_{-\infty}^{\infty} de' e^{iu\epsilon_{e'}} \langle e|e^{-iuH_1}\rho|e \rangle \langle \psi_{\mu}^e|e' \rangle|^2 \\ &= \lim_{\mu \to 0} \iint_{-\infty}^{\infty} de' \int_{-\infty}^{\infty} de' e^{iu\epsilon_{e'}} \langle e|e^{-iuH_1}\rho|e \rangle \langle \psi_{\mu}^e|e' \rangle|^2 \\ &= \int_{-\infty}^{\infty} de \int_{-\infty}^{\infty} de' e^{iu\epsilon_{e'}} \langle e|e^{-iuH_1}\rho|e \rangle \delta^{\mathbf{D}}(e - e') = \int_{-\infty}^{\infty} de \langle e|e^{iu(U^{\dagger}H_1U-H_1)}\rho|e \rangle = \operatorname{Tr}[e^{iu(U^{\dagger}H_1U-H_1)}\rho] = G_{OBS}(u) \end{aligned}$$

where we considered the properties of Eqs. (208) and (209) from the forth line on. As a result, in the limit in which  $\mu \rightarrow 0$ ,

$$\wp_{\rm TPM}(z, H_1, U, \rho) = \int_{-\infty}^{\infty} du e^{-iuz} G_{\rm TPM}(u) = \int_{-\infty}^{\infty} du e^{-iuz} G_{\rm OBS}(u) = \wp_{\rm OBS}(z, H_1, U, \rho).$$
(212)

# B. Calculating the probabilities $|\langle n| U_{\tau}^+ |0\rangle|^2$ in the subsection "Calculations for the Fig.1" of Methods section

We consider, as in the main text, that  $U_{\tau}^{+} = \langle + | U_{\tau} | + \rangle = e^{-i\theta_{\tau}} \exp\left[-\frac{i\tau}{\hbar} \left(\frac{P^{2}}{2m} + \frac{m\omega^{2}}{2}X_{+}^{2}\right)\right]$ ,  $X_{+} = \langle + | X' | + \rangle = X + a$ , and  $\theta_{\tau} = \frac{\hbar\omega_{\tau}\tau}{2\hbar} - \frac{m\omega^{2}a^{2}\tau}{2\hbar}$ . Considering  $|x_{+}\rangle$  as the basis that diagonalizes  $X_{+}$  and comparing with the basis  $|x\rangle$  that diagonalizes X, we have that  $|x_{+}\rangle_{x_{+} \text{ basis}} = |x_{+} - a\rangle_{x \text{ basis}}$ . Therefore, for any state  $|\psi\rangle$ , the following equality holds

$$\langle x_+ | \psi \rangle_{x_+ \text{ basis}} = \langle x_+ - a | \psi \rangle \tag{213}$$

so that

$$\langle x_{+}|0\rangle_{x_{+} \text{ basis}} = \frac{1}{(\pi 2\sigma^{2})^{1/4}} \exp\left[-\frac{(x_{+}-a)^{2}}{4\sigma^{2}}\right]$$
 (214)

and (see page 450 of [10] for the deduction of  $\langle x|n \rangle$  in the X basis)

$$\langle x_+|n\rangle_{x_+\text{ basis}} = (2^n n!)^{-1/2} \frac{1}{(2\pi\sigma^2)^{1/4}} \exp\left[-\frac{(x_+-a)^2}{4\sigma^2}\right] H_n\left(\frac{x_+-a}{\sqrt{2}\sigma}\right)$$
 (215)

where  $\sigma = \sqrt{\hbar/(2m\omega)}$  and  $H_n(y)$  are Hermite polynomials, that satisfy [11]

$$H_n(y) = \sum_{s=0}^{n/2} (-1)^s (2y)^{n-2s} \frac{n!}{(n-2s)!s!}$$
(216)

From this point on, we will consider all our calculations in the  $X_+$  basis. Notice that the term  $\frac{P^2}{2m} + \frac{m\omega^2}{2}X_+^2$  that appears inside the exponential defining  $U_{\tau}^+$  is just a simple harmonic oscillator energy term, with X displaced to  $X_+$ . Since  $[X_+, P] = [X, P] = i\hbar$ , then the same algebra rules for the oscillator can be applied here. As a result, we can write

$$\langle n|U_{\tau}^{+}|0\rangle = e^{-i\theta_{\tau}} \int_{-\infty}^{\infty} dx_{+} \langle n|x_{+}\rangle \langle x_{+}| \exp\left[-\frac{i\tau}{\hbar} \left(\frac{P^{2}}{2m} + \frac{m\omega^{2}}{2}X_{+}^{2}\right)\right] |0\rangle$$
(217)

The term

$$\langle x_{+}|\exp\left[-\frac{i\tau}{\hbar}\left(\frac{P^{2}}{2m}+\frac{m\omega^{2}}{2}X_{+}^{2}\right)\right]|0\rangle$$
 (218)

can be identified as the wave function evolved until time  $\tau$  of the state  $|0\rangle$ , which, in the  $X_+$  position basis, is a Gaussian centered around the phase point (a, 0) as described in Eq. (214). This gaussian evolves under a simple harmonic oscillator dynamics until time  $\tau$ . Using the formalism in [12, 13], we obtain

$$\langle x_{+} | \exp\left[-\frac{i\tau}{\hbar} \left(\frac{P^{2}}{2m} + \frac{m\omega^{2}}{2} X_{+}^{2}\right)\right] |0\rangle = \frac{1}{\left(2\pi\sigma_{t}^{2}\right)^{1/4}} \exp\left[-\frac{(x_{+} - a\cos(\omega t))^{2}}{\cos(\omega t)(4\sigma^{2}\cos(\omega t) + \frac{2i\hbar}{m\omega}\sin(\omega t))} - i\frac{m\omega\tan(\omega t)}{2\hbar} x_{+}^{2}\right]$$

$$= \frac{1}{\left(2\pi\sigma_{t}^{2}\right)^{1/4}} \exp\left[-\frac{(x_{+} - a\cos(\omega t))^{2}}{4\sigma^{2}\cos(\omega t)\left(\cos(\omega t) + \frac{i\hbar}{2m\omega\sigma^{2}}\sin(\omega t)\right)} - i\frac{m\omega\tan(\omega t)}{2\hbar} x_{+}^{2}\right]$$

$$= \frac{1}{\left(2\pi\sigma_{t}^{2}\right)^{1/4}} \exp\left[-\frac{(x_{+} - a\cos(\omega t))^{2}\left(\cos(\omega t) - \frac{i\hbar}{2m\omega\sigma^{2}}\sin(\omega t)\right)}{4\sigma_{t}^{2}\cos(\omega t)} - i\frac{m\omega\tan(\omega t)}{2\hbar} x_{+}^{2}\right]$$

$$(219)$$

where

$$\sigma_t = \sqrt{\sigma^2 \cos^2(\omega t) + \frac{\hbar^2}{4m^2 \omega^2 \sigma^2} \sin^2(\omega t)}$$
(220)

Now, inserting Eqs. (215) and (219) in Eq. (217), considering Eq. (216), we get

$$\langle n | U_{\tau}^{+} | 0 \rangle = (2^{n} n!)^{-1/2} \frac{e^{-i\theta_{\tau}}}{(\pi^{2} 4 \sigma^{2} \sigma_{\tau}^{2})^{1/4}} \int_{-\infty}^{\infty} dx_{+} \exp \left[ -\frac{(x_{+} - a)^{2}}{4\sigma^{2}} - \frac{(x_{+} - a\cos(\omega t))^{2} \left(\cos(\omega t) - \frac{i\hbar}{2m\omega\sigma^{2}}\sin(\omega t)\right)}{4\sigma_{\tau}^{2}\cos(\omega t)} - i\frac{m\omega\tan(\omega t)}{2\hbar}x_{+}^{2} \right] H_{n}\left(\frac{x_{+} - a}{\sqrt{2}\sigma}\right)$$

$$= (2^{n} n!)^{-1/2} \frac{e^{-i\theta_{\tau}}}{(\pi^{2} 4\sigma^{2} \sigma_{\tau}^{2})^{1/4}} \sum_{s=0}^{n/2} (-1)^{s} \frac{n!}{(n-2s)!s!} \times$$

$$\times \int_{-\infty}^{\infty} dx_{+} \exp \left[ -\frac{(x_{+} - a)^{2}}{4\sigma^{2}} - \frac{(x_{+} - a\cos(\omega t))^{2} \left(\cos(\omega t) - \frac{i\hbar}{2m\omega\sigma^{2}}\sin(\omega t)\right)}{4\sigma_{\tau}^{2}\cos(\omega t)} - i\frac{m\omega\tan(\omega t)}{2\hbar}x_{+}^{2} \right] (2\frac{x_{+} - a}{\sqrt{2}\sigma})^{n-2s}$$

$$(221)$$

Now, given that  $\tau = \pi/\omega$ , then  $\sigma_t = \sigma$  and, after some tedious calculations, we obtain

$$\langle n| U_{\tau}^{+} |0\rangle = (2^{n} n!)^{-1/2} \frac{e^{-\frac{a^{2}}{2\sigma^{2}} - i\theta_{\tau}}}{(2\pi\sigma^{2})^{1/2}} \sum_{s=0}^{n/2} \left(\frac{\sqrt{2}}{\sigma}\right)^{n-2s} (-1)^{s} \frac{n!}{(n-2s)!s!} f(n, s, a, \sigma),$$
(222)

where

$$f(n, s, a, \sigma) = \int_{-\infty}^{\infty} dx_{+} \exp\left[-\frac{(x_{+}+a)^{2}}{2\sigma^{2}}\right] (x_{+})^{n-2s}$$

$$= (-1)^{-2s} e^{-\frac{a^{2}}{4\sigma^{2}}} 2^{n-2s} |\sigma|^{n-2s-1} \times \\ \times \left(a|\sigma|\left((-1)^{n} - (-1)^{2s}\right) \Gamma\left(\frac{n}{2} - s + 1\right) {}_{1}F_{1}\left(\frac{n}{2} - s + 1; \frac{3}{2}; \frac{a^{2}}{4\sigma^{2}}\right) \\ + \sigma^{2}\left((-1)^{n} + (-1)^{2s}\right) \Gamma\left(\frac{1}{2}(n-2s+1)\right) {}_{1}F_{1}\left(\frac{1}{2}(n-2s+1); \frac{1}{2}; \frac{a^{2}}{4\sigma^{2}}\right)\right)$$
(223)

and  $\Gamma(z)$  is the Gamma function [11] and  ${}_{1}F_{1}\left(\frac{1}{2}(n-2s+1);\frac{1}{2};\frac{a^{2}}{4\sigma^{2}}\right)$  is the Kummer confluent hypergeometric function [14]. Using Eq. (222) with the parameters described in Fig. 1, we could compute the probabilities  $p_{\text{TPM}}(0, H_{\text{HO}}, U_{\tau}, \rho_{1}) = |\langle 0| U_{\tau}^{+} |0\rangle|^{2}$  and  $p_{\text{TPM}}(\hbar\omega, H_{\text{HO}}, U_{\tau}, \rho_{1}) = |\langle 1| U_{\tau}^{+} |0\rangle|^{2}$  presented in the Figure 1.

## C. Calculating the probabilities $p(n, H_{HO}(\tau))$ in subsection "Calculations for the Fig. 3 and 5" of the methods section

Our goal in this subsection is to compute the probability of the Harmonic oscillator part  $H_{\rm HO} = \frac{P^2}{2m} \otimes \mathbb{1}_{\rm s} + \frac{m\omega^2 X^2}{2} \otimes \mathbb{1}_{\rm s} = \hbar\omega(N + 1/2) \otimes \mathbb{1}_{\rm s}$  of the total energy to have some arbitrary value at time  $\tau$ . To do so, we first notice that since  $|n, \pm\rangle$  are eigenstates of  $H_{\rm HO}$  at time 0 with eigenvalues  $\hbar\omega(n + 1/2)$ , then the probability of the system of being  $\hbar\omega(n + 1/2)$  at time  $\tau$  is given by

$$p(n, H_{\rm H0}(\tau)) = |\langle n, -| U_{\tau} | \alpha, - \rangle|^2.$$
(224)

where we considered the fact that, since  $\sigma_z$  commutes with  $U_{\tau}$ , then

$$|\langle n, +| U_{\tau} | \alpha, - \rangle|^2 = 0.$$
 (225)

Our strategy to obtain  $|\langle n, -|U_{\tau}|\alpha, -\rangle|^2$  is to make a transition to the Heisenberg picture. For that, first notice that from the transition from the Schrödinger to the Heisenberg picture, we have that given that  $|n, \pm\rangle$  are eigenvectors of  $H_{\text{Ho}}$ , then  $U_{\tau}^{\dagger}|n, \pm\rangle = |n', \pm\rangle$  are eigenstates of the Heisenberg evolved version  $U_{\tau}^{\dagger}H_{\text{Ho}}U_{\tau}$ , since [10]

$$U_{\tau}^{\dagger}H_{\rm H0}U_{\tau}|n',\pm\rangle = U_{\tau}^{\dagger}H_{\rm H0}U_{\tau}U_{\tau}^{\dagger}|n,\pm\rangle = \left(n+\frac{1}{2}\right)\hbar\omega U^{\dagger}|n,\pm\rangle = \left(n+\frac{1}{2}\right)|n',\pm\rangle$$
(226)

Now, let us analyze  $\{|n'\rangle\}$ . We saw in the methods section, that the Heisenberg time-evolved version of the operator  $X' = X \otimes \mathbb{1}_s + a(\mathbb{1}_{CM} \otimes \sigma_z)$  is given by  $X'(\tau) = -X \otimes \mathbb{1}_s - a(\mathbb{1}_{CM} \otimes \sigma_z)$ . Therefore, given that  $\sigma_z$  does not change in time for the given evolution, it follows that  $X(\tau) = U_{\tau}^{\dagger}(X \otimes \mathbb{1}_s)U_{\tau} = -X \otimes \mathbb{1}_s - 2a(\mathbb{1}_{CM} \otimes \sigma_z)$ . Moreover, given that  $P(\tau) = -P \otimes \mathbb{1}_s$ , we have that

$$H'_{\rm HO} := \langle -|H_{\rm HO}(\tau)| - \rangle = \langle -|\left(\frac{P^2(\tau)}{2m} + \frac{m\omega^2 X^2(\tau)}{2}\right)| - \rangle = \frac{P^2(0)}{2m} + \frac{m\omega}{2}X'^2$$
(227)

and

$$\langle +|H_{\rm H0}(\tau)|-\rangle = \langle -|H_{\rm H0}(\tau)|+\rangle = 0 \tag{228}$$

where X' = X - 2a. Moreover, notice that  $[X', P] = i\hbar$  and [X', X'] = 0, so that, analogously as in the usual harmonic oscillator, by defining [10]

$$a' = \sqrt{\frac{m\omega}{2\hbar}} (X' + \frac{iP}{m\omega})$$
(229)

we have that

$$[a', a^{\dagger}] = 1, \quad N' = a^{\dagger} a' = \frac{H'_{S}}{\hbar \omega} - \frac{1}{2}, \quad [N', a^{\dagger}] = a^{\dagger}, \qquad [N', a'] = -a'.$$
(230)

So that

$$H'_{\rm HO} = \hbar\omega \left(N' + \frac{1}{2}\right) \tag{231}$$

and the same algebraic structure as the usual harmonic oscillator can be used to deduce that

$$N'|n'\rangle = n'|n'\rangle, \quad a^{\dagger}|n'\rangle = \sqrt{n'+1}|n'+1\rangle, \quad a^{\prime}|n'\rangle = \sqrt{n'}|n'-1\rangle.$$
(232)

As a result,

$$|n'\rangle = \frac{(a^{\prime\dagger})^n}{\sqrt{n!}} |0'\rangle \tag{233}$$

The difference from the usual harmonic oscillator appears in the wave-functions, since, considering that  $X' |x\rangle = (x - 2a) |x\rangle$ , then

$$\langle x|a'|0'\rangle = \left(x - 2a + x_0^2 \partial_x\right) \langle x|0\rangle = 0, \tag{234}$$

where  $x_0 = \sqrt{\frac{\hbar}{m\omega}}$ , resulting in

$$\langle x|0\rangle = \frac{1}{\pi^{\frac{1}{4}}\sqrt{x_0}} \exp\left[-\frac{1}{2}\left(\frac{x-2a}{x_0}\right)^2\right]$$
 (235)

From this, we obtain, using a method similar to the usual harmonic oscillator case [10], that

$$\langle x|n'\rangle = \left(\frac{1}{\pi^{\frac{1}{4}}\sqrt{2^{n}n!}}\right) \left(\frac{1}{x_{0}^{n+1/2}}\right) \left(x - 2a - x_{0}^{2}\partial_{x}\right)^{n} \exp\left[-\frac{1}{2}\left(\frac{x - 2a}{x_{0}}\right)^{2}\right]$$
(236)

These equations are the same as for the eigenvectors  $|n\rangle$  of  $\frac{p^2}{2m} + \frac{m\omega^2 X^2}{2} = \hbar\omega(N + 1/2)$ , but displaced by 2*a* (see section 2.3 of Ref. [10]), so that

$$\langle x - 2a|n \rangle = \langle x|n' \rangle \,. \tag{237}$$

Considering that the wave function of the coherent state is defined as [1]

$$\langle x|\alpha\rangle = e^{i\theta_{\alpha}} \left(\frac{m\omega}{\pi\hbar}\right)^{\frac{1}{4}} \exp\left[-\left(\frac{x-\langle \alpha|X|\alpha\rangle}{2\sigma_{X}^{2}}\right)^{2} + \frac{i\langle \alpha|P|\alpha\rangle x}{\hbar}\right]$$
(238)

with  $\sigma_X = \sqrt{\hbar/(2m\omega)}$  and  $e^{i\theta_\alpha} = e^{\alpha^{*2}-\alpha^2}$ , then we have that

$$\langle n'|\alpha\rangle = \int_{-\infty}^{\infty} dx \langle n'|x\rangle \langle x|\alpha\rangle = \int_{-\infty}^{\infty} dx \langle n'|x\rangle e^{i\theta_{\alpha}} \left(\frac{m\omega}{\pi\hbar}\right)^{\frac{1}{4}} \exp\left[-\left(\frac{x-\langle \alpha|X|\alpha\rangle}{2\sigma_{\chi}^{2}}\right)^{2} + \frac{i\langle \alpha|P|\alpha\rangle x}{\hbar}\right]$$

$$= \int_{-\infty}^{\infty} dx' \langle n'|x' + 2\alpha\rangle e^{i\theta_{\alpha}} \left(\frac{m\omega}{\pi\hbar}\right)^{\frac{1}{4}} \exp\left[-\left(\frac{x'+2\alpha-\langle \alpha|X|\alpha\rangle}{2\sigma_{\chi}^{2}}\right)^{2} + \frac{i\langle \alpha|P|\alpha\rangle (x'+2\alpha)}{\hbar}\right]$$

$$= e^{i\theta_{\alpha}+i2\alpha\langle \alpha|P|\alpha\rangle/\hbar} \int_{-\infty}^{\infty} dx' \langle n'|x' + 2\alpha\rangle \left(\frac{m\omega}{\pi\hbar}\right)^{\frac{1}{4}} \exp\left[-\left(\frac{x'+2\alpha-\langle \alpha|X|\alpha\rangle}{2\sigma_{\chi}^{2}}\right)^{2} + \frac{i\langle \alpha|P|\alpha\rangle x'}{\hbar}\right]$$

$$= e^{i\theta_{\alpha}+i2\alpha\langle \alpha|P|\alpha\rangle/\hbar} \int_{-\infty}^{\infty} dx' \langle n|x'\rangle \left(\frac{m\omega}{\pi\hbar}\right)^{\frac{1}{4}} \exp\left[-\left(\frac{x'+2\alpha-\langle \alpha|X|\alpha\rangle}{2\sigma_{\chi}^{2}}\right)^{2} + \frac{i\langle \alpha|P|\alpha\rangle x'}{\hbar}\right]$$

$$(239)$$

Defining

$$\alpha' = \sqrt{\frac{m\omega}{2\hbar}} (\langle \alpha | X | \alpha \rangle - 2a) + i \frac{1}{\sqrt{2m\hbar\omega}} \langle \alpha | P | \alpha \rangle, \qquad (240)$$

we can consider the coherent state  $|\alpha'\rangle$ , such that

$$\langle x|\alpha'\rangle = e^{i\theta_{\alpha'}} \left(\frac{m\omega}{\pi\hbar}\right)^{\frac{1}{4}} \exp\left[\left(\frac{x-\langle \alpha'|X|\alpha'\rangle}{2\sigma_X^2}\right)^2 + \frac{i\langle \alpha'|P|\alpha'\rangle x}{\hbar}\right],\tag{241}$$

where

$$\langle \alpha' | X | \alpha' \rangle = \sqrt{\frac{2\hbar}{m\omega}} \Re(\alpha') = \langle \alpha | X | \alpha \rangle - 2a, \quad \langle \alpha' | P | \alpha' \rangle = \sqrt{2m\hbar\omega} \Im(\alpha') = \langle \alpha | P | \alpha \rangle, \quad e^{i\theta_{\alpha'}} = e^{\frac{\alpha'^2 - \alpha'^2}{4}}.$$
 (242)

Therefore,

$$\langle n'|\alpha\rangle = e^{i(\theta_{\alpha}-\theta_{\alpha'})+i2a\langle\alpha|P|\alpha\rangle/\hbar} \int_{-\infty}^{\infty} dx' \langle n|x'\rangle \langle x'|\alpha'\rangle = \langle n|\alpha'\rangle e^{i(\theta_{\alpha}-\theta_{\alpha'})-i2a\langle\alpha|P|\alpha\rangle/\hbar}$$
(243)

Given that  $|\alpha'\rangle$  is a proper coherent state, then it follows that

$$|\alpha'\rangle = e^{-|\alpha'|^2/2} \sum_{n=0}^{\infty} \frac{\alpha'^n}{\sqrt{n!}} |n\rangle$$
(244)

and

$$\langle n|\alpha'\rangle = e^{-|\alpha'|^2/2} \frac{\alpha'^n}{\sqrt{n!}},\tag{245}$$

so that, from Eq. (243),

$$|\langle n'|\alpha\rangle|^2 = e^{-|\alpha'|^2} \frac{|\alpha'|^{2n}}{n!}.$$
(246)

As a result, it follows that

$$|\langle n', +|\alpha, -\rangle|^2 = 0, \quad |\langle n', -|\alpha, -\rangle|^2 = e^{-|\alpha'|^2} \frac{|\alpha'|^{2n}}{n!}$$
 (247)

The probability that  $H_{\rm HO}$  have an energy  $\hbar\omega(n+1/2)$  at time  $\tau$  is thus given by

$$p(n, H_{\rm Ho}(\tau)) = |\langle n', -|\alpha, -\rangle|^2 = e^{-|\alpha'|^2} \frac{|\alpha'|^{2n}}{n!}.$$
(248)

For values different from  $\hbar\omega(n + 1/2)$  for any *n*, the probability is null.

## D. Example of result 2 considering Trapped Ions

As mentioned in the main text, in specific scenarios, result 2 can be used to obtain the OBS statistics via a probe, an auxiliary energy operator  $H_2$ . However, as seen in the derivation of result 2, finding a physically intuitive operator  $H_2$  and a state  $|v\rangle$  such that result 2 can be directly applied is not always straightforward. Nevertheless, there exist special cases where  $H_2$  can be explicitly given without introducing an additional subspace ( $\mathcal{H}'$  in result 2). We analyze such cases here.

Consider a system  $\Omega$  whose total energy is given by

$$H = H_1 + H_2, (249)$$

where both  $H_1$  and  $H_2$  act on the same Hilbert space  $\mathcal{H}$ . The unitary evolution is specified as  $U = \exp[-iHt/\hbar]$ , and the following commutation relations hold:

$$[U^{\dagger}H_1U, H_1] \neq 0, \quad [U^{\dagger}H_2U, H_2] = 0.$$
(250)

Since

$$[\Delta(H_1, U), H_2] = [\Delta(H_1, U), U^{\dagger} H_2 U] = 0,$$
(251)

it follows that for any initial preparation  $\rho = |\delta_i(H_1, U)\rangle \langle \delta_i(H_1, U)|$  as an eigenstate of  $\Delta(H_1, U)$ , one can measure  $H_2$  at times 0 and *t*, obtaining with 100% certainty the energies  $E_i$  and  $E'_i = E_i - \delta_i(H_1, U)$ , where  $E_i$  is an eigenvalue of  $H_2$ . By energy conservation, the variation of  $H_1$  must necessarily be  $\delta_i(H_1, U)$ . This represents a special case of result 2, where  $H_2 \rightarrow H_2 \otimes \mathbb{I}_{\mathcal{H}'}$  and  $U' = \mathbb{I}_{\mathcal{H}'}$ .

An illustrative example of this scenario is provided in Fig. 1, corresponding to the trapped ion system analyzed in Figs. 1, 3, and 4 of the main text. In this case, the unitary evolution is given by

$$U_{\tau} = \exp[-iH\tau/\hbar],\tag{252}$$

where the Hamiltonian takes the form

$$H = H_{\rm HO} \otimes \mathbb{1}_s + H_{\rm e},\tag{253}$$

with

$$H_{\rm HO} = (N + \frac{1}{2})\hbar\omega, \quad H_{\rm e} = \hbar(\omega_z/2 + \Delta_S k_{\rm SW} X/2) \otimes \sigma_z. \tag{254}$$

By making the substitutions

$$H_1 \to H_{\rm H0} \otimes \mathbb{1}_s, \quad H_2 \to (H - H_1) = H_{\rm e}, \quad U \to U_{\tau},$$
(255)

we recover result 2, demonstrating that measuring  $H_2$  at times 0 and  $\tau$  yields the same statistics for the variation of  $H_1$  under  $U_{\tau}$  as obtained via the OBS protocol.



FIG. 1. Illustration of the application of result 2 to the trapped Ca<sup>+</sup> ion system. We consider the same system as in Figs. 1, 3, and 4 in the main text and define  $H_1 = H_{HO} \otimes \mathbb{1}_s$  and  $H_2 = H - H_1 = H_e$ . The variation of energy is analyzed over the interval  $[0, \tau]$ , where  $\tau = \pi/\omega$  and the unitary evolution is  $U \equiv U_\tau = \exp[-iH\tau/\hbar]$ . As detailed in the "Methods" section, the operator of variation of  $H_1$  for this interval is  $\Delta(H_1, U) = 2m\omega^2 aX \otimes \sigma_z + 2m\omega^2 a^2$ , where  $a = \hbar \Delta_s k_{sw}/(2m\omega^2)$ . The following conditions are satisfied  $[U^{\dagger}H_2U, H_2] = [U^{\dagger}H_2U, \Delta(H_1, U)] = [H_2, \Delta(H_1, U)] = 0$ . This scenario represents a special case of result 2. On the left, the OBS protocol is applied to measure  $\Delta(H_1, U)$ . The system's position X and spin  $\sigma_z$  are measured, yielding  $x_0$  and -1, respectively. This determines the variation of energy as  $\delta_{x_0}(H_1, U) = -2m\omega^2 ax_0 + 2m\omega^2 a^2$  which is the eigenvalue of  $\Delta(H_1, U)$  associated with the eigenvector  $|\delta_{x_0}(H_1, U)\rangle = |x_0, - \rangle$ . On the right, result 2 is applied to the same system, approximately prepared in the eigenstate  $|\delta_{x_0}(H_1, U)|$ . In this case,  $|\delta_{x_0}(H_1, U)\rangle$  is an eigenstate of both  $H_2$  and  $U^{\dagger}H_2U$ . Measurements of  $H_2$  at t = 0 and  $U^{\dagger}H_2U$  at  $t = \tau$ , via measurements of X and  $\sigma_z$ , yield the eigenvalues  $E_{x_0} = -\hbar(\omega_z/2 + \Delta_s k_{sw} x_0/2)$  and  $E'_{x_0} = E_{x_0} - \delta_{x_0}(H_1, U)$ , respectively. By the reality condition, the variation of  $H_2$  is  $-\delta_{x_0}(H_1, U)$ . By the conservation of energy (condition 2), the variation of  $H_1$  must be  $\delta_{x_0}(H_1, U)$ , thus showing the equivalence between the measurement results on the left and right side of the figure.

In this case, we arrive at a special instance of result 2 where no additional Hilbert space  $\mathcal{H}'$  or auxiliary vector  $|v\rangle$  is required to obtain the OBS statistics for the variation of  $H_1$ . However, to fully conform with result 2, we could introduce an auxiliary space  $\mathcal{H}'$  by redefining

$$H_1 \to H_{\text{HO}} \otimes \mathbb{1}_s, \quad H_2 \to (H - H_1) \otimes \mathbb{1}_{\mathcal{H}'} = H_{\circ} \otimes \mathbb{1}_{\mathcal{H}'}.$$
 (256)

Similarly, we set

$$U \otimes U' \to U_{\tau} \otimes \mathbb{1}_{\mathcal{H}'}.$$
(257)

With this formulation, the same result would be obtained for any normalized state  $|v\rangle \in \mathcal{H}'$ .

[5] Busch, P. The Quantum Theory of Measurement. (Springer-Verlag, 1996).

- [7] Pinto Silva, T. A. B. & Gelbwaser-Klimovsky, D. Quantum work: reconciling quantum mechanics and thermodynamics, Phys. Rev. Research 6, L022036 (2024).
- [8] Talkner, P. et al. Fluctuation theorems: Work is not an observable. Phys. Rev. E 75, 050102(R) (2007).
- [9] Pinto Silva, T. A. B. & Angelo, R. M. Quantum mechanical work, Phys. Rev. A 104, 042215 (2021).
- [10] Sakurai, J. J. Modern quantum mechanics, (Addison-Wesley Publishing Company, 1994).

<sup>[1]</sup> Cohen-Tannoudji, C. et al. *Quantum Mechanics, Volume I: Basic Concepts, Tools, and Applications* (Wiley-VCH Verlag GmbH & Co, 2020).

<sup>[2]</sup> Athreya K. B. & Lahiri, S. N. Measure theory and probability theory (Springer, 2006).

<sup>[3]</sup> Fristedt, B. E. & Gray, L. F. A modern approach to probability theory (Springer Science & Business Media, 2013).

<sup>[4]</sup> Heinosaari, T. A. & Ziman, M. The Mathematical Language of Quantum Theory (Cambridge University Press, 2012).

<sup>[6]</sup> Perarnau-Llobet, M. et al. No-go theorem for the characterization of work fluctuations in coherent quantum systems. *Phys. Rev. Lett.* **118**, 070601 (2017).

<sup>[11]</sup> Arfken, G. B. et al. Mathematical methods for physicists: a comprehensive guide (Academic press, 2011).

- [12] Freire, I. S. & Angelo, R. M. Quantifying continuous-variable realism. Phys. Rev. A 100, 022105 (2019).
- [13] Pinto Silva, T. A. B. A definition of quantum mechanical work. Master's dissertation, Universidade Federal do Paraná, Curitiba (2018).
- [14] Weisstein, E. W. Confluent hypergeometric function of the first kind. https://mathworld.wolfram.com/ (2003).