# A robust estimation and variable selection approach for sparse partially linear additive models

Alejandra Mercedes Martínez

CONICET and Universidad Nacional de Luján, Argentina

#### Abstract

In partially linear additive models the response variable is modelled with a linear component on a subset of covariates and an additive component in which the rest of the covariates enter to the model as a sum of univariate unknown functions. This structure is more flexible than the usual full linear or full nonparametric regression models, avoids the "curse of dimensionality", is easily interpretable and allows the user to include discrete or categorical variables in the linear part. On the other hand, in practice, the user incorporates all the available variables in the model no matter how they would impact on the response variable. For this reason, variable selection plays an important role since including covariates that has a null impact on the responses will reduce the prediction capability of the model. As in other settings, outliers in the data may harm estimations based on strong assumptions, such as normality of the response variable, leading to conclusions that are not representative of the data set.

In this work, we propose a family of robust estimators that estimate and select variables from both the linear and the additive part of the model simultaneously. This family considers an adaptive procedure on a general class of penalties in the regularization part of the objetive function that defines the estimators. We study the behaviour of the proposal againts its leastsquares counterpart under simulations and show the advantages of its use on a real data set.

### 1 Introduction

Partially linear additive models (PLAM) assume that  $(Y_i, \mathbf{Z}_i^{\mathrm{T}}, \mathbf{X}_i^{\mathrm{T}})^{\mathrm{T}} \in \mathbb{R}^{1+q+p}, 1 \leq i \leq n$ , are independent and identically distributed random vectors with the same distribution as  $(Y, \mathbf{Z}^{\mathrm{T}}, \mathbf{X}^{\mathrm{T}})^{\mathrm{T}}$  such that

$$Y = m(\mathbf{Z}^{\mathrm{T}}, \mathbf{X}^{\mathrm{T}}) + u = \mu + \boldsymbol{\beta}^{\mathrm{T}} \mathbf{Z} + \sum_{j=1}^{p} \eta_j(X_j) + \sigma\varepsilon$$
(1)

where the constant  $\mu \in \mathbb{R}$ , vector  $\beta \in \mathbb{R}^q$ , the univariate functions  $\eta_j : \mathbb{R} \to \mathbb{R}$ ,  $1 \leq j \leq p$ , and the scale parameter  $\sigma > 0$  are the objects to be estimated. In order to ensure identifiability of the additive components  $\eta_j$ , it is assumed that  $\int \eta_j(x) dx = 0$  and that  $\eta_j : \mathcal{I}_j \to \mathbb{R}$ , then  $\int_{\mathcal{I}} \eta_j(x) dx = 0$ , for  $1 \leq j \leq p$ . The errors  $\varepsilon$  are assumed to be independent from the vector of covariates  $(\mathbf{Z}^T, \mathbf{X}^T)^T$ . In the classical context, the errors  $\varepsilon$  are considered to have 0 mean and variance equals 1, while in the context of robustness they are considered to have a symmetric distribution  $F(\cdot)$  (that is, if they have a density, this density function has to be an even one) with scale parameter equals 1.

In Boente & Martinez (2023) a robust approach to estimate all the parameters in the model is developed. A *B*-spline basis is used to approximate each additive function  $\eta_j$  in (1), transforming the original model into a linear regression model. A robust regression approach is then applied to obtain the final estimators. In addition to having developed asymptotic theory for the proposal, a generalization of a BIC criteria was defined in order to robustly select the regularization parameters in the approximation procedure. However, this approach has its limitation. When many covariates are included, particularly in the additive part of the model, the estimation procedure becomes impractical, as it is either computationally very expensive or fails to produce unique solutions.

Usually, in practice and in a first step of modelling, researchers introduce all possible variables into the model based on their own experience. In this sense, some variables that have a small impact on the response variable will reduce the prediction capability of the model. For this reason, under a sparse model, variable selection plays an important role. Even though there exists different methods for selecting variables such as *stepwise* or *best subset*, these used to be computational expensive, their theoretical properties are hard to establish and, the most severe which is the lack of stability analyzed, for instance, by (Breiman, 1996). An alternative is to include a penalty term in the optimization problem that defines the estimators (see Efron & Hastie, 2016, for an overview of penalization methods). An example of these is the least absolute shrinkage and selection operator (LASSO) estimator, related to the  $\ell_1$  penalty, for linear regression models introduced by Tibshirani (1996). However, it has the disadvantages of selecting too many covariates and giving rise to biased solutions. To avoid the selection problem, Zou & Hastie (2005) considered a combination of the  $L_1$  and  $L_2$  penalties, named *Elastic Net*, while Fan & Li (2001) introduced the SCAD penalty (see Section 5 for more details). Other penalties are the adaptive LASSO, also called ADALASSO (Zou, 2006), and the Minimax Concave Penalty (MCP) introduced by Zhang (2010).

Similar to what happens with estimation procedures, simultaneous variable estimation and selection methods can be seriously affected if no robust procedures are used and there are even a few atypical observations presence in the data. In order to solve this problem, Smucler & Yohai (2017) considered sparse robust estimators for linear regression models. In generalized linear models, Avella-Medina & Ronchetti (2018) gave a robust proposal based on controlling the quasilikelihood for the variable selection procedure, and in Bianco et al. (2022) considered penalized robust estimators based on the deviance for the logistic model. On the other hand, Avella-Medina (2016) consideres robust approaches for generalized additive models.

As we have already mentioned, many researchers have studied the problem of estimating and selecting variables in linear and also in partly linear models by considering that the regression parameter vector is sparse (see, for instance, Desboulets, 2018, for a review). Specifically with respect to the simultaneous estimation and variable selection in PLAM, sparse models for the linear component were considered in Liu et al. (2011), Du et al. (2012) and Lian (2012) who developed a variable selection procedure based on the least squares regression estimator, spline approximation and penalty functions SCAD or ADALASSO for the regression parameters  $\beta$ . More resistant approaches to atypical data have been studied in Koenker (2011), Guo et al. (2013) and Sherwood & Wang (2016) who proposed methods based on quantile regression. However, as it is mentioned in Boente & Martinez (2023), the quantile estimators are related to an unbounded loss functions and, for this reason and as it happens in linear regression, it can be affected by high leverage points.

However, as far as we know, not many have studied the problem of estimating and simultaneously selecting variables from both the linear and the additive components of a PLAM. Among them, we can mention Kazemi et al. (2018) who proposed a two-step estimation and variable selection method in a high dimensional setting, used the SCAD penalty for regularizing and considered an optimization procedure that minimizes the sum the squared residuals. Besides, they proposed a BIC-type criteria for selecting the penalty parameters and a 2-fold cross-validation mechanism for determining the tuning parameters of the B-splines approximation. Hu et al. (2015) considered a hierarchical Bayesian estimation and model selection approach with quantile regression, while Banerjee & Ghosal (2014) proposed a Bayesian method for generalized partially linear additive models that uses group LASSO (see Yuan & Lin, 2006). When the response variable is a time series, Feng et al. (2018) studied an estimation and variable selection procedure using the adaptive group LASSO. In all these papers, the additive functions have been approximated by splines, except in Feng et al. (2018) where they used orthogonal series. However, all these approaches are not resistant to atypical data. A more robust approach has been given in Lv et al. (2017) by combining modal estimators based on B-splines with an adaptive SCAD penalty. Even though this proposal is more robust that the ones previously mentioned since can afford heavy-tailed distributions, these estimators are highly sensitive to high leverage points in the linear component of the model associated to large residuals.

The rest of the paper is organized as follows. In Section 2 the robust approach that estimates and selects variables from both the linear and the additive components of a PLAM is proposed. In Section 3, a robust BIC criterion is defined to select the penalty parameters, while in Section 4 it is mentioned a robust BIC criterion for selecting the parameters related to the B-splines approximation. Section 5 contains a brief review of the differences among some commonly used penalty functions. In Section 6, a general algorithm to obtain the penalized robust estimators is derived. The results of a simulation study carried out to compare the performance of the robust proposal against its least squares counterpart is shown in Section 7. Section 8 contains the analysis of the well-known plasma beta-carotene level data set where the advantages of using the penalized robust approach to simultaneously estimate and select variables from both linear and additive part of the PLAM are highlighted. Finally, some final comments can be found in Section 9.

## 2 The proposal

In order to estimate and select variables from both components of a PLAM, a B-spline basis is used to approximate each additive function  $\eta_j$  of (1). Similar to what is done in Boente & Martinez (2023),  $\eta_j$  is approximated as a  $\sum_{s=1}^{k_j-1} c_s^{(j)} B_s^{(j)}(t)$  with  $B_s^{(j)}$ , for  $1 \leq s \leq k_j - 1$  and  $1 \leq j \leq p$ , the sth element of the basis of centered B-splines of order  $\ell_j$ , that is,  $B_s^{(j)}$  is such that  $B_s^{(j)}(t) = \tilde{B}_s^{(j)}(t) - \int_{\mathcal{I}_j} \tilde{B}_s^{(j)}(t) dt$  with  $\{\tilde{B}_s^{(j)} : 1 \leq s \leq k_j\}$  a spline basis of order  $\ell_j$ . Besides, as it is explained in that paper, only  $k_j - 1$  elements of the basis are considered instead of the  $k_j$  total in order to allow the parameter  $\mu$  to be identifiable. Then, the PLAM in (1) is approximated by the following linear regression model

$$Y = \mu + \beta^{\mathrm{T}} \mathbf{Z} + \sum_{j=1}^{p} \sum_{s=1}^{k_j - 1} c_s^{(j)} B_s^{(j)}(X_j) + \sigma \varepsilon.$$
(2)

Let  $\boldsymbol{\beta} = (\beta_1, \dots, \beta_q)^{\mathrm{T}} \in \mathbb{R}^q$ ,  $\mathbf{c}^{(j)} = (c_1^{(j)}, \dots, c_{k_j-1}^{(j)})^{\mathrm{T}} \in \mathbb{R}^{k_j-1}$ ,  $\mathbf{c} = (\mathbf{c}^{(1)\mathrm{T}}, \dots, \mathbf{c}^{(p)\mathrm{T}}) \in \mathbb{R}^K$ ,  $K = \sum_{j=1}^p k_j - p$ , and  $\mathbf{V} = (\mathbf{V}^{(1)}(X_1)^{\mathrm{T}}, \dots, \mathbf{V}^{(p)}(X_p)^{\mathrm{T}})^{\mathrm{T}} \in \mathbb{R}^K$  where  $\mathbf{V}^{(j)}(t) = (B_1^{(j)}(t), \dots, B_{k_j-1}^{(j)}(t))^{\mathrm{T}} \in \mathbb{R}^{k_j-1}$ . Then, (2) can be written in a simplified form as

$$Y = \mu + \boldsymbol{\beta}^{\mathrm{T}} \mathbf{Z} + \mathbf{c}^{\mathrm{T}} \mathbf{V}_{i} + \sigma \varepsilon$$

Let  $\mathbf{b} = (b_1, \ldots, b_q)^{\mathrm{T}} \in \mathbb{R}^q$  and  $\mathbf{d}^{(j)} = (d_1^{(j)}, \ldots, d_{k_j-1}^{(j)})^{\mathrm{T}} \in \mathbb{R}^{k_j-1}$  two general vectors to approximate  $\boldsymbol{\beta}$  and  $\mathbf{c}^{(j)}$ , for  $1 \leq j \leq p$ , and  $\mathbf{d} = (\mathbf{d}^{(1)\mathrm{T}}, \ldots, \mathbf{d}^{(p)\mathrm{T}}) \in \mathbb{R}^K$ . Let  $\hat{\mu}$  be an estimator of  $\mu$  and  $\hat{\sigma}$  a scale estimator of  $\sigma$ , then we define the penalized robust estimators for the partially linear additive models as

$$(\hat{\boldsymbol{\beta}}, \hat{\mathbf{c}}) = \underset{\mathbf{b} \in \mathbb{R}^{q}, \mathbf{d} \in \mathbb{R}^{K}}{\operatorname{argmin}} PL_{n, \boldsymbol{\lambda}, \mathbf{k}}(\mathbf{b}, \mathbf{d}), \qquad (3)$$

where

$$PL_{n,\boldsymbol{\lambda},\mathbf{k}}(\mathbf{b},\mathbf{d}) = L_n(\widehat{\mu},\widehat{\sigma},\mathbf{b},\mathbf{d}) + \mathcal{J}_{\boldsymbol{\lambda},\mathbf{k}}(\mathbf{b},\mathbf{d}), \qquad (4)$$

with

$$L_n(a,\varsigma,\mathbf{b},\mathbf{d}) = \frac{1}{n} \sum_{i=1}^n \rho\left(\frac{r_i(a,\mathbf{b},\mathbf{d})}{\varsigma}\right), \qquad r_i(a,\mathbf{b},\mathbf{d}) = Y_i - a - \mathbf{b}^{\mathrm{T}} \mathbf{Z}_i - \mathbf{d}^{\mathrm{T}} \mathbf{V}_i$$

and  $\mathcal{J}_{\boldsymbol{\lambda},\mathbf{k}}(\mathbf{b},\mathbf{d})$  is an arbitrary penalty function (selected by the user) depending on a regularization parameter  $\boldsymbol{\lambda} = (\lambda_{1,1}, \ldots, \lambda_{1,q}, \lambda_{2,1}, \ldots, \lambda_{2,p})^{\mathrm{T}} \in \mathbb{R}^{q+p}$  that determines the model complexity, that is,  $\lambda_{1,s}$ ,  $1 \leq s \leq q$ , and  $\lambda_{2,j}$ ,  $1 \leq j \leq p$ , are the penalty parameters that control the sparsity of the parametric and nonparametric components, respectively, and  $\mathbf{k} = (k_1, \ldots, k_p)^{\mathrm{T}}$  is the parameter vector that controls the *B*-spline approximation of the additive parte of the model. For instance,  $\mathcal{J}_{\boldsymbol{\lambda},\mathbf{k}}(\mathbf{b},\mathbf{d})$  can be defined as

$$\mathcal{J}_{\lambda,\mathbf{k}}(\mathbf{b},\mathbf{d}) = \sum_{s=1}^{q} p_{\lambda_{1,s}}(|b_s|) + \sum_{j=1}^{p} p_{\lambda_{2,j}}(\|\mathbf{d}^{(j)}\|_{H_j})$$

with  $p_{\lambda}$  a univariate penalty function such as the SCAD or the MCP penalties. Additionally, the proposal includes, but it is not mandatory, the adaptive setting, as proposed in Zhao et al. (2014), as follows

$$\lambda_{1,s} = \frac{\lambda_1}{|\widetilde{\beta}_s|} \quad \text{and} \quad \lambda_{2,j} = \frac{\lambda_2}{\|\widetilde{\mathbf{c}}^{(j)}\|_{H_j}}, \quad (5)$$

where  $\tilde{\lambda}_1$  and  $\tilde{\lambda}_2$  are auxiliary penalty parameters. In addition,  $\|\mathbf{d}^{(j)}\|_{H_j} = (\mathbf{d}^{(j)T}H_j\mathbf{d}^{(j)})^{1/2}$  with  $H_j$  of dimension  $(k_j - 1) \times (k_j - 1)$  with its (s, s') element given by  $\int_0^1 B_s^{(j)}(t) B_{s'}^{(j)}(t) dt$ ,  $\rho$  is a bounded  $\rho$ -function as defined in Maronna et al. (2019). Besides,  $\tilde{\beta}_s$ , for  $1 \leq s \leq q$ , and  $\tilde{\mathbf{c}}^{(j)}$ , for  $1 \leq j \leq p$ , are unpenalized estimators of  $\beta_s$  and  $\mathbf{c}^{(j)}$ , respectively. A possible choice is to compute the robust estimators obtained with the non-penalized procedure proposed in Boente & Martinez (2023) from where it can also be obtained  $\hat{\mu}$  and  $\hat{\sigma}$ . It is worth noting that when the adaptive setting is considered, the penalty parameters  $\lambda_{1,s}$  and  $\lambda_{2,j}$ , for  $1 \leq s \leq q$ ,  $1 \leq j \leq p$ , associated to components  $b_s$  and  $\mathbf{d}^{(j)}$ , respectively, are given in (5) through the auxiliary penalty parameters  $\tilde{\lambda}_1$  and  $\tilde{\lambda}_2$ . In this context, the search for the "best" subset of parameters is reduced from dimension q + p to 2, as it is explained in Section 5.

*Remark.* When the objetive function  $PL_{n,\lambda,\mathbf{k}}$  is defined through auxiliary coefficients  $\lambda_1$  and  $\lambda_2$  as in (5), to determine the penality parameters for each component of the regression parameter vectors, say **b** and **d**, the resulting procedure is called *adaptive*. An example of this is the adaptive LASSO, called ADALASSO, defined in Zou (2006).

## **3** Selection of penalty parameters

Since any robust estimation and variable selection approach also needs robust methods for selecting the regularization and penalty parameters, we will consider a robust procedure for selecting the penalty parameters when the the number of terms to approximate the additive functions, that is,  $\mathbf{k}$ , is fixed. Following the ideas in Zhao et al. (2014) and Lv et al. (2017) for modal regression, for  $\mathbf{k}$  fixed, we propose the following robust BIC criterion

$$\operatorname{RBIC}_{\boldsymbol{\lambda}}(\boldsymbol{\lambda}) = \log\left(\widehat{\sigma}^2 \sum_{i=1}^n \rho\left(\frac{r_i(\widehat{\beta}_{\boldsymbol{\lambda}}, \widehat{\mathbf{c}}_{\boldsymbol{\lambda}})}{\widehat{\sigma}}\right)\right) + df_c \frac{\log(n)}{n} + df_n \frac{\log(n/K)}{n/K}$$
(6)

where  $K = \sum_{j=1}^{p} k_j - p$ ,  $(\widehat{\beta}_{\lambda}, \widehat{\mathbf{c}}_{\lambda})$  is the solution of (3) with penalty vector  $\boldsymbol{\lambda} = (\lambda_{1,1}, \ldots, \lambda_{1,q}, \lambda_{2,1}, \ldots, \lambda_{2,p})^{\mathrm{T}}$ ,  $df_c$  is the number of nonzero parametric components,  $df_n$  is the number of nonzero nonparametric components and  $\widehat{\sigma}$  is the preliminary *S*-estimator.

However, this proposal needs to select  $\lambda$  over a grid of dimension q + p which is very expensive if this number is very large. For this reason and taking into account the relation between  $\lambda_{1,s}$  and  $\tilde{\lambda}_1$  and between  $\lambda_{2,j}$  and  $\tilde{\lambda}_2$  as stated in (5), we will now propose a method for selecting only the penalty parameters  $\tilde{\lambda}_1$  and  $\tilde{\lambda}_2$  which implies a selection over a grid of dimension 2. The modified robust BIC criterion is defined in the following way:

$$\operatorname{RBIC}_{\boldsymbol{\lambda}}(\widetilde{\boldsymbol{\lambda}}) = \log\left(\widehat{\sigma}^2 \sum_{i=1}^n \rho\left(\frac{r_i(\widehat{\beta}_{\widetilde{\boldsymbol{\lambda}}}, \widehat{\mathbf{c}}_{\widetilde{\boldsymbol{\lambda}}})}{\widehat{\sigma}}\right)\right) + df_c \frac{\log(n)}{n} + df_n \frac{\log(n/K)}{n/K}$$
(7)

where  $K = \sum_{j=1}^{p} k_j - p$ ,  $df_c$ ,  $df_n$  and  $\widehat{\sigma}$  are defined as before, and where  $(\widehat{\beta}_{\widetilde{\lambda}}, \widehat{\mathbf{c}}_{\widetilde{\lambda}})$  is now the solution of (3) with  $\lambda$  given through  $\widetilde{\lambda} = (\widetilde{\lambda}_1, \widetilde{\lambda}_2)$  as in (5).

# 4 Selection of regularization parameters

Any B-spline approach requires the selection of a set of knots to define the basis functions. In this sense, since each additive function  $\eta_j$ ,  $1 \leq j \leq p$ , is approximated by a B-spline curve, p sequences of knots are necessary. Stone (1985) mentions that the number of knots is more important than their locations. For this reason, we will consider a generalization of a BIC criterion for the number of knots considered for each additive component, that is, for  $k_1, \ldots, k_p$ . However, regarding their locations, two possible choices are: the equally spaced knots and the quantile knots. Uniform knots are used when the function  $\eta_j$  does not present dramatic changes in its derivates, while non-uniform knots are preferred when the function shows very different behaviours in different regions. When

the latter is the case, quantile knots are used, that is, the knots are obtained as the quantiles of the observed explanatory variable with uniform percentile ranks.

Since, once again, a robust criterion should be used to obtain final robust estimators, in this paper, in order to determine  $\mathbf{k} = (k_1, \ldots, k_p)^T$ , we will use the robust BIC criterion defined in Boente & Martinez (2023) which is the following:

$$\operatorname{RBIC}_{\mathbf{k}}(\mathbf{k}) = \log\left(\widehat{\sigma}^2 \sum_{i=1}^n \rho\left(\frac{r_i(\widehat{\beta}_{\boldsymbol{\lambda},\mathbf{k}},\widehat{\mathbf{c}}_{\boldsymbol{\lambda},\mathbf{k}})}{\widehat{\sigma}}\right)\right) + \frac{\log(n)}{2n} \sum_{j=1}^p k_j \tag{8}$$

where  $\mathbf{k} = (k_1, \ldots, k_p)^{\mathrm{T}}$ ,  $r_i(\widehat{\beta}_{\boldsymbol{\lambda}, \mathbf{k}}, \widehat{\mathbf{c}}_{\boldsymbol{\lambda}, \mathbf{k}})$ , for  $1 \leq i \leq n$ , are the residuals obtained using a basis of dimension  $k_j$  to approximate the additive function  $\eta_j$  and  $\boldsymbol{\lambda}$  is given,  $\rho$  is the  $\rho$ -function used to compute the *M*-estimator and  $\widehat{\sigma}$  is a preliminary scale estimator of  $\sigma$ . When p = 1, that is, when it is considered a partially linear model with one covariate entering in the nonparametric part of the model, and  $\rho(t) = t^2$ , this criterion is reduced to the criterion proposed in He et al. (2003). When  $\boldsymbol{\lambda}$  in defined through  $\widetilde{\boldsymbol{\lambda}}$  as in (5), the dependence of the estimators can be hightlighted as  $\widehat{\beta}_{\widetilde{\boldsymbol{\lambda}}, \mathbf{k}}$  and  $\widehat{\mathbf{c}}_{\widetilde{\boldsymbol{\lambda}}, \mathbf{k}}$ .

Besides, when the number of elements to approximate each additive function is the same, that is, when  $k_j = k$  for  $1 \le j \le p$ , the robust BIC criterion reduces to

$$\operatorname{RBIC}_{\mathbf{k}}(k) = \log\left(\widehat{\sigma}^2 \sum_{i=1}^n \rho\left(\frac{r_i(\widehat{\beta}_{\mathbf{\lambda},\mathbf{k}},\widehat{\mathbf{c}}_{\mathbf{\lambda},\mathbf{k}})}{\widehat{\sigma}}\right)\right) + \frac{\log(n)}{2n}pk.$$
(9)

As it is mentioned in He et al. (2003) and also in Boente & Martinez (2023), since for cubic splines the smallest possible number of knots is 4, the *j*th component of **k**, that is,  $k_j$ , in (8) can be chosen in the interval  $[\max(n^{1/5}/2, 4); 8 + 2n^{1/5}]$  when cubic splines are considered. The numerical computations are even more reduced when the size of the basis is the same for all *j*. In this case, it is enough to look for the first local minimum of  $\text{RBIC}_{\mathbf{k}}(k)$  defined in (9) for  $\max(n^{1/5}/2, 4) \le k \le 8 + 2n^{1/5}$ .

## 5 The penalty functions

There are many univariate penalty functions  $p_{\lambda}$  that can be considered in the regularization component of (4), such as the  $L_1$  or, in general, the  $L_q$  penalties (see Tibshirani, 1996, 1997; Frank & Friedman, 1993; Fu, 1998), the hard thresholding (see Fan & Li, 2001; Pötscher & Leeb, 2009, for an overview and a comparison of penalty functions), the Minimax Concave Penalty (MCP) introduced by Zhang (2010) or the Smoothly Clipped Absolute Deviation Penalty (SCAD) proposed by Fan & Li (2001).

The hard thresholding penalty is defined as  $p_{\lambda}(\theta) = \lambda^2 - (|\theta| - \lambda)^2 I\{|\theta| < \lambda\}$ , while the  $L_q$  penalty is defined as  $p_{\lambda}(\theta) = \lambda |\theta|^q$ . When q = 1, we get the  $L_1$  penalty defined as  $p_{\lambda}(\theta) = \lambda |\theta|$  which is associated to the LASSO estimators. Additionally, the MCP is defined as

$$p_{\lambda}(\theta) = \begin{cases} \lambda |\theta| - \frac{\theta^2}{2\gamma} & |\theta| \le \lambda\gamma \\ \frac{\gamma\lambda^2}{2} & |\theta| > \lambda\gamma \end{cases}$$

where  $\gamma > 1$  is a tuning parameter, and the SCAD penalty is defined, for  $\lambda > 0$  (and some a > 2), as

$$p_{\lambda}(\theta) = \begin{cases} \lambda |\theta|, & |\theta| \le \lambda \\ -(\theta^2 - 2a\lambda|\theta| + \lambda^2)/(2(a-1)), & \lambda < |\theta| \le a\lambda \\ (a+1)\lambda^2/2, & |\theta| > a\lambda \end{cases}$$

The SCAD penalty has advantages over the  $L_q$  and the hard thresholding penalties such as its simultaneous unbiased, sparsity and continuity properties of the resulting estimators. For instace, the  $L_q$  penalties give rise to continuous solutions when  $q \ge 1$  but for q > 1 it does not produce sparse solutions, and when q = 1 the resulting estimator is shifted by a constant and so it is not unbiased; while the hard thresholding does not produce continuous estimatos (see Fan & Li, 2001, for more details). It can also be appreciated that the penalties hard thresholding, SCAD and MCP are bounded functions while the others are unbounded. Even though the similarities between the SCAD and the MCP penalties, as it is mentioned in Bianco et al. (2022), in some regression settings when considering classical estimadors, the first one outperforms the latter. In allow the reader some visual comparison, Figure 1 shows the plots of the penalty functions previously mentioned. In red solid line it is plotted the hard thresholding, in green line the  $L_1$  penalty, in pink the  $L_{0.5}$  penalty, in orange the MCP and in blue line the SCAD penalty. As previously mentioned, the  $L_2$  penalty function does not yield sparse solutions due to its differentiability at the origin, so its curve is represented by a dashed gray line.



Figure 1: Plot of penalty functions: the hard penalty in red, the  $L_1$  in green, the  $L_{0.5}$  in pink, the MCP in orange, the SCAD penalty in blue and the  $L_2$  in dashed gray line.

For all computational studies conducted in this article, we use the Smoothly Clipped Absolute Deviation (SCAD) penalty. As is customary, the parameter a in the SCAD penalty definition is fixed at 3.7, so it does not require selection.

# 6 General algorithm

In this section, we derive an algorithm to compute the penalized estimators for the general setting in which the penalized vector is given by  $\boldsymbol{\lambda} = (\lambda_{1,1}, \ldots, \lambda_{1,q}, \lambda_{2,1}, \ldots, \lambda_{2,p})^{\mathrm{T}}$  and the RBIC criterion for selecting the penalty parameters is defined in (6). However, when an adaptive procedure is used, that is, when the components of the penalty vector  $\boldsymbol{\lambda}$  are defined according to (5), the proposed algorithm can be easily modified by considering a grid for  $\boldsymbol{\lambda} = (\lambda_1, \lambda_2)^{\mathrm{T}}$  and minimizing the RBIC criterion defined in (7). Similarly, although we describe the general framework for selecting the regularization parameters  $\mathbf{k} = (k_1, \ldots, k_p)^{\mathrm{T}}$  where the RBIC criterion is defined in (9), a straightforward modification of the algorithm allows us to minimize the RBIC defined in (8) under the assumption that  $k_j = k$  for all  $j = 1, \ldots, p$ . In this sense, Algorithm 1 presents a general procedure for obtaining the proposed estimators.

#### Algorithm 1 General algorithm

- 1: Obtain preliminary robust estimators  $\hat{\mu}$ ,  $\hat{\beta}$ ,  $\hat{\eta}_j$ , for  $1 \leq j \leq p$ , and a robust scale estimator  $\hat{\sigma}$ . A possible choice is to compute the robust estimators obtained with the non-penalized procedure proposed in Boente & Martinez (2023).
- 2: For  $j = 1, \ldots, p$ , fix the spline order  $\ell_j$ .
- 3: Consider a grid for  $\mathbf{k} = (k_1, \dots, k_p)^T$  of M elements:  $\mathbf{k}_1, \dots, \mathbf{k}_M$ , and consider a grid for  $\boldsymbol{\lambda}$  of N elements:  $\boldsymbol{\lambda}_1, \dots, \boldsymbol{\lambda}_N$ .
- 4: for  $\ell = 1$  to M do
- 5: for j = 1 to N do
- 6: Solve the optimization problem

$$(\widehat{\beta}_{\lambda_j,\mathbf{k}_\ell}, \widehat{\mathbf{c}}_{\lambda_j,\mathbf{k}_\ell}) = \operatorname*{argmin}_{\mathbf{b}\in\mathbb{R}^q,\mathbf{d}\in\mathbb{R}^K} PL_{n,\lambda_j,\mathbf{k}_\ell}(\mathbf{b},\mathbf{d}).$$
(10)

Note that  $(\widehat{\beta}_{\lambda_j,\mathbf{k}_\ell}, \widehat{\mathbf{c}}_{\lambda_j,\mathbf{k}_\ell})$  is the argument that minimizes the optimization problem when  $\mathbf{k} = \mathbf{k}_\ell$  and  $\lambda = \lambda_j$  are fixed.

- Compute and save  $\operatorname{RBIC}_{\lambda}(\lambda_j)$ .
- 8: end for

7:

9: Solve

$$\boldsymbol{\lambda}_{\mathbf{k}_{\ell}} = \operatorname{argmin} \{ \operatorname{RBIC}_{\boldsymbol{\lambda}}(\boldsymbol{\lambda}_{j}) \}$$

That is,  $\lambda_{\mathbf{k}_{\ell}}$  is the optimal penalty parameter vector for  $\mathbf{k} = \mathbf{k}_{\ell}$  fixed.

- 10: Compute and save  $\operatorname{RBIC}_{\mathbf{k}}(\mathbf{k}_{\ell})$ .
- 11: end for
- 12: Obtain  ${\bf k}$  as

$$\widehat{\mathbf{k}} = \mathop{\mathrm{argmin}}_{\ell} \left\{ \mathrm{RBIC}_{\mathbf{k}}(\mathbf{k}_{\ell}) \right\}$$

13: The penalized estimators are  $(\widehat{\beta}_{\lambda_{\hat{k}},\hat{k}}, \widehat{c}_{\lambda_{\hat{k}},\hat{k}})$ .

As it can be appreciated, Algorithm 1 does not include how to solve the optimization problem (10), which we still need to describe. In Subsection 6.1, a method to solve the minimization task, for **k** and  $\tilde{\lambda}$  fixed, is developed. The procedure considers certain type of penalty functions, such as the SCAD and MCP penalties.

#### 6.1 Solving the optimization problem for k and $\lambda$ fixed

In what follows, we consider univariate penalties  $p_{\lambda}$  that are twice continuously differentiable on (0, M), with M > 0, and satisfy  $p_{\lambda}(0) = 0$ . These conditions allow for local approximation by quadratic functions near in a neighbourhood of the origin and include some common penalty functions such as the  $L_1$ , SCAD and MCP penalties.

For the sake of simplicity, let assume that  $\mu = 0$ . Then, for  $\lambda_j$  and  $\mathbf{k}_{\ell}$  fixed, we have

$$PL_{n,\boldsymbol{\lambda}_{j},\mathbf{k}_{\ell}}(\mathbf{b},\mathbf{d}) = \frac{1}{n} \sum_{i=1}^{n} \rho\left(\frac{r_{i}(\mathbf{b},\mathbf{d})}{\widehat{\sigma}}\right) + \mathcal{J}_{\boldsymbol{\lambda}_{j},\mathbf{k}_{\ell}}(\mathbf{b},\mathbf{d}).$$
(11)

with  $r(\mathbf{b}, \mathbf{d}) = Y_i - \mathbf{b}^{\mathrm{T}} \mathbf{Z}_i - \mathbf{d}^{\mathrm{T}} \mathbf{V}_i$ . If  $\mu \neq 0$ , obtain  $\hat{\mu}$  an estimator of  $\mu$ , for instance using the unpenalized robust estimator defined in Boente & Martinez (2023) or using the S- or the MM-Ridge estimators defined in Maronna (2011), and take the response variables  $Y_i - \hat{\mu}$  instead of  $Y_i$ .

Assume that the penalty function used  $p_{\lambda}$  is the same for both the linear and the additive parts of the model. To avoid burden notation, let  $p = p_{\lambda_{1,s}}$  and  $p = p_{\lambda_{2,j}}$  when it corresponds. The assumptions on p previously mentioned guarantee that, if  $t_0$  is a initial point close to 0, then there exists a quadratic function q(t) that is symmetric around 0,  $q(t_0) = p(|t_0|)$  and  $q'(t_0) =$  $p'(|t_0|) \operatorname{sgn}(t_0)$ . Using the symmetry, we have that  $q(t) = a + bt^2$ . Then, the conditions on  $q(t_0)$ and  $q'(t_0)$  imply that

$$\left\{ \begin{array}{rrrr} a+b|t_0|^2 &=& p(|t_0|) \\ 2bt_0 &=& p'(|t_0|)\operatorname{sgn}(t_0) \end{array} \Leftrightarrow \left\{ \begin{array}{rrrr} a+b|t_0|^2 &=& p(|t_0|) \\ 2b|t_0| &=& p'(|t_0|) \end{array} \right. \right.$$

Then, from the second equation we have that  $b = p'(|t_0|)/(2|t_0|)$  and replacing in the first equation we get that  $a = p(|t_0|) - p'(|t_0|)/(2|t_0|)t_0^2$ . Finally, the quadratic approximation is

$$\begin{aligned} q(t) &= p(|t_0|) - \frac{p'(|t_0|)}{2|t_0|} t_0^2 + \frac{p'(|t_0|)}{2|t_0|} t^2 \\ &= p(|t_0|) + \frac{p'(|t_0|)}{2|t_0|} (t^2 - t_0^2), \end{aligned}$$

that is, for t close to  $t_0$ , we have that

$$p(|t|) \approx p(|t_0|) + \frac{p'(|t_0|)}{2|t_0|} (t^2 - t_0^2).$$

In this way, given  $(\mathbf{b}_0, \mathbf{d}_0)$  close to the minimizer of (10) and  $(\mathbf{b}, \mathbf{d})$  close to  $(\mathbf{b}_0, \mathbf{d}_0)$  with  $|b_{0s}| > 0$ and  $\|\mathbf{d}_0^{(j)}\|_{H_j} > 0$ , for s = 1, ..., q and j = 1, ..., p, we have that

$$p(|b_s|) \approx p(|b_{0s}|) + \frac{p'(|b_{0s}|)}{2|b_{0s}|}(b_s^2 - b_{0s}^2)$$

and

$$p(\|\mathbf{d}^{(j)}\|_{H_j}) \approx p(\|\mathbf{d}_0^{(j)}\|_{H_j}) + \frac{p'(\|\mathbf{d}_0^{(j)}\|_{H_j})}{2\|\mathbf{d}^{(j)}\|_{H_j}} (\|\mathbf{d}^{(j)}\|_{H_j}^2 - \|\mathbf{d}_0^{(j)}\|_{H_j}^2)$$

Then, recalling that  $\|\mathbf{d}^{(j)}\|_{H_j}^2 = \mathbf{d}^{(j)\mathrm{T}} H_j \mathbf{d}^{(j)}$ , if we define

$$\boldsymbol{\Sigma}(\mathbf{b}_{0},\mathbf{d}_{0}) = \operatorname{diag}\left\{\frac{p'(|b_{01}|)}{2|b_{01}|},\ldots,\frac{p'(|b_{0q}|)}{2|b_{0q}|},\frac{p'(||\mathbf{d}_{0}^{(1)}||_{H_{1}})}{2||\mathbf{d}^{(1)}||_{H_{1}}}H_{1},\ldots,\frac{p'(||\mathbf{d}_{0}^{(p)}||_{H_{p}})}{2||\mathbf{d}^{(p)}||_{H_{p}}}H_{p}\right\} \in \mathbb{R}^{(q+K-p)\times(q+K-p)}$$

we get that the objective function given in (11) can be approximated by

$$\frac{1}{n}\sum_{i=1}^{n}\rho\left(\frac{r_{i}(\mathbf{b},\mathbf{d})}{\widehat{\sigma}}\right) + (\mathbf{b}^{\mathrm{T}},\mathbf{d}^{\mathrm{T}})\boldsymbol{\Sigma}(\mathbf{b}_{0},\mathbf{d}_{0})(\mathbf{b}^{\mathrm{T}},\mathbf{d}^{\mathrm{T}})^{\mathrm{T}}$$
(12)

and so the estimators obtained by minimizing (12) will be close to the minimizers of (10) with the penalized function given in (11).

Denote  $\boldsymbol{\theta} = (\mathbf{b}^{\mathrm{T}}, \mathbf{d}^{\mathrm{T}})^{\mathrm{T}}$  and  $\mathbf{W}_{i} = (\mathbf{Z}_{i}^{\mathrm{T}}, \mathbf{V}_{i}^{\mathrm{T}})^{\mathrm{T}}$ . The minimization of (12) can be obtained by a reweighted procedure as it is usual for M-estimators.

**Deduction of the iteration procedure**. Suppose now we have  $\boldsymbol{\theta}^{(m)} = (\mathbf{b}^{(m)T}, \mathbf{d}^{(m)T})^{T}$  the estimators obtained in the *m*-step and  $\hat{\sigma}$  is a preliminary robust estimator of  $\sigma$ . By first derivating (12) with respect to  $\boldsymbol{\theta}$  and then multiplying and dividing by  $(Y_i - \mathbf{W}_i^T \boldsymbol{\theta})/\hat{\sigma}$ , we obtain

$$\frac{1}{n} \sum_{i=1}^{n} \psi \left( \frac{Y_i - \mathbf{W}_i^{\mathrm{T}} \boldsymbol{\theta}}{\widehat{\sigma}} \right) \left( -\frac{\mathbf{W}_i}{\widehat{\sigma}} \right) + 2 \boldsymbol{\Sigma}(\boldsymbol{\theta}^{(m)}) \boldsymbol{\theta} = \mathbf{0}$$

$$\frac{1}{n} \sum_{i=1}^{n} \psi \left( \frac{Y_i - \mathbf{W}_i^{\mathrm{T}} \boldsymbol{\theta}}{\widehat{\sigma}} \right) \frac{\widehat{\sigma}}{Y_i - \mathbf{W}_i^{\mathrm{T}} \boldsymbol{\theta}} \frac{Y_i - \mathbf{W}_i^{\mathrm{T}} \boldsymbol{\theta}}{\widehat{\sigma}} \left( -\frac{\mathbf{W}_i}{\widehat{\sigma}} \right) + 2 \boldsymbol{\Sigma}(\boldsymbol{\theta}^{(m)}) \boldsymbol{\theta} = \mathbf{0}$$

$$\frac{1}{n} \sum_{i=1}^{n} w \left( \frac{r_i(\mathbf{b}, \mathbf{d})}{\widehat{\sigma}} \right) \frac{Y_i - \mathbf{W}_i^{\mathrm{T}} \boldsymbol{\theta}}{\widehat{\sigma}} \left( -\frac{\mathbf{W}_i}{\widehat{\sigma}} \right) + 2 \boldsymbol{\Sigma}(\boldsymbol{\theta}^{(m)}) \boldsymbol{\theta} = \mathbf{0}$$

where  $w(t) = \psi(t)/t$  and  $r_i(\mathbf{b}, \mathbf{d}) = Y_i - \mathbf{W}_i^{\mathrm{T}} \boldsymbol{\theta}$ . Then, the estimator  $\boldsymbol{\theta}^{(m+1)} = (\mathbf{b}^{(m+1)\mathrm{T}}, \mathbf{d}^{(m+1)\mathrm{T}})^{\mathrm{T}}$  of the (m+1)-step is defined as the solution of

$$\frac{1}{n}\sum_{i=1}^{n}w_{i,m}\frac{Y_{i}-\mathbf{W}_{i}^{\mathrm{T}}\boldsymbol{\theta}}{\widehat{\sigma}}\left(-\frac{\mathbf{W}_{i}}{\widehat{\sigma}}\right)+2\boldsymbol{\Sigma}(\boldsymbol{\theta}^{(m)})\boldsymbol{\theta}=\mathbf{0}$$

where  $w_{i,m} = W(r_i(\boldsymbol{\theta}^{(m)})/\hat{\sigma})$  are the weights obtained by replacing the estimator calculated in the previous step. Finally, the latter equation is equivalent to

$$-\frac{1}{n}\sum_{i=1}^{n}\frac{w_{i,m}}{\widehat{\sigma}^{2}}\mathbf{W}_{i}Y_{i} + \left(\frac{1}{n}\sum_{i=1}^{n}\frac{w_{i,m}}{\widehat{\sigma}^{2}}\mathbf{W}_{i}\mathbf{W}_{i}^{\mathrm{T}} + 2\boldsymbol{\Sigma}(\boldsymbol{\theta}^{(m)})\right)\boldsymbol{\theta} = \mathbf{0}$$

and so the iteration step is

$$\boldsymbol{\theta}^{(m+1)} = \left(\frac{1}{n}\sum_{i=1}^{n}\frac{w_{i,m}}{\widehat{\sigma}^2}\mathbf{W}_i\mathbf{W}_i^{\mathrm{T}} + 2\boldsymbol{\Sigma}(\boldsymbol{\theta}^{(m)})\right)^{-1}\frac{1}{n}\sum_{i=1}^{n}\frac{w_{i,m}}{\widehat{\sigma}^2}\mathbf{W}_iY_i.$$
 (13)

Finally, we obtain the following algorith to minimize (12).

The iteratively reweighted algorithm. Consider  $\epsilon > 0$  the stopping rule, then

- 1. Compute  $\hat{\sigma}$  and an initial  $\theta^{(0)}$  such as the one mentioned in Step 1 in Algorithm 1.
- 2. For m = 0, 1, 2, ... compute the weights  $w_{i,m}$ , for i = 1, ..., n, and then  $\theta^{(m+1)}$  in (13).
- 3. Stop when the incrementation is small enough, for instance, if  $\|\boldsymbol{\theta}^{(m+1)} \boldsymbol{\theta}^{(m)}\| / \|\boldsymbol{\theta}^{(m)}\| < \epsilon$ .

# 7 Monte Carlo Study

This section contains the results of a simulation study in which it is compared the robust proposal with the corresponding estimator based on least squares. All computations were carried out in R. The code used is available at https://github.com/alemermartinez/rplam-vs. The LS-estimator corresponds to use  $\rho(t) = t^2$  in (4). Observe that, in this case, no preliminary scale estimator is needed. For the robust estimator, the loss function considered is the Tukey's bisquare loss which is of the form  $\rho_c(t) = \min\{1 - (1 - (t/c)^2)^3, 1\}$ . The tuning constant c > 0 balances the robustness and efficiency properties of the associated estimators. For the simulation study, the tuning constant was fixed at c = 4.685, which is the constant necessary to obtain a 95% efficiency for linear regression estimators. Besides, an *S*-estimator is used for the preliminary scale estimator. From now on, the robust procedure proposed in this paper will be denoted as ROB and as LS the approach based on least squares.

The samples  $\{(Y_i, \mathbf{Z}_i^{\mathrm{T}}, \mathbf{X}_i^{\mathrm{T}})^{\mathrm{T}}\}_{i=1}^n$  are generated with the same distribution as  $(Y, \mathbf{Z}^{\mathrm{T}}, \mathbf{X}^{\mathrm{T}})^{\mathrm{T}}, \mathbf{Z} = (Z_1, \ldots, Z_q) \in \mathbb{R}^q, \mathbf{X} = (X_1, \ldots, X_p)^{\mathrm{T}} \in \mathbb{R}^p$ , with q = 10 and p = 10 and three different sample sizes n = 200,400 and 600. The covariates  $\mathbf{Z}_i = (Z_{i1}, \ldots, Z_{iq})^{\mathrm{T}}$  are generated from a multivariate normal distribution with zero mean and variance-covariance matrix  $\Sigma$  defined as  $(\Sigma)_{i,\ell} = \operatorname{Corr}(Z_{ik}, Z_{i\ell}) = 0.5^{|k-\ell|}$ , for  $1 \leq k, \ell \leq q$ , and  $X_{ij}$ , for  $j = 1, \ldots, p$ , are uniformly distributed on the interval [0; 1] and are independent from each other and from the vector  $\mathbf{Z}_i$ .

In all cases, the response and the covariates satisfy the partially linear additive model

$$Y = \mu + \boldsymbol{\beta}^{\mathrm{T}} \mathbf{Z} + \sum_{j=1}^{p} \eta_j(X_j) + u$$

where  $\beta = (3, 1.5, 2, -1.5, 0, 0, 0, 0, 0, 0) \in \mathbb{R}^{10}$ , the additive functions are

$$\eta_1(x) = 5x - \frac{5}{2}, \qquad \eta_2(x) = 3(2x - 1)^2 - 1,$$
  
$$\eta_3(x) = 60x^3 - 90x^2 + 30x, \qquad \eta_4(x) = 2\sin(\pi x) - \frac{4}{\pi}$$

 $\eta_j(x) \equiv 0$  for  $j = 5, \ldots, p$ ,  $u = \sigma \varepsilon$ ,  $\sigma = 1$  and  $\mu = 0$ . In all cases the additive functions are such that  $\mathbb{E}\eta_j(X_j) = 0$  for all  $j = 1, \ldots, 10$ . For clean samples, the error's distribution is  $\epsilon \sim N(0, 1)$ , denoted from now on as  $C_0$ .

In order to study the effect of atypical data on the estimators, seven different contamination schemes were considered:

- $C_1: u \sim t_3$
- $C_2: u \sim 0.9N(0, \sigma^2) + 0.1N(0, 25\sigma^2)$
- $C_3: u \sim 0.95N(0, \sigma^2) + 0.05N(0, 100\sigma^2)$
- $C_4: u \sim 0.95N(0, \sigma^2) + 0.05N(15, \sigma^2)$
- $C_5: U \sim 0.85N(0,\sigma^2) + 0.15N(15,\sigma^2)$

- $C_6$ : A 5% of covariates  $\mathbf{Z}_i$  were replaced by  $(20, \ldots, 20) \in \mathbb{R}^{10}$ .
- $C_7$ : A 10% of covariates  $\mathbf{Z}_i$  were replaced by  $(20, \ldots, 20) \in \mathbb{R}^{10}$ .

Contamination  $C_1$  is a heavy-tailed contamination setting where the errors are distributed as a Student's t-distribution with 3 degrees of freedom.  $C_2$  and  $C_3$  are variance contamination settings where a 10% and a 5%, respectively, of the errors are normaly distributed with enlarged variances.  $C_4$  and  $C_5$  are bias contamination schemes where 5% and 15% of the errors, respectively, have their location shifted to 15. These contamination schemes give rise to what are called *vertical outliers*.  $C_6$  and  $C_7$  are high-leverage contaminations where a 5% and 10%, respectively, of the covariates of the linear part were randomly chosen to take another value. In this case, the response variables and the errors are not modified. It is interesting to note that none of the contamination settings considered introduces outliers in the additive component of the model or in the response variable depending on a particular set of the additive components. The main reason for not considering outliers in the covariates of the additive part is that this would only generate high computational costs since the estimators would be defined using a larger number of terms to approximate the additive functions. With respect to contamination of the errors depending on the values taken by the covariates belonging to the additive part, similarly as what was done in (Boente et al., 2017), it would be needed a much larger number of sample sizes. For instance, if we want to contaminate errors with their 10 covariates in the interval [0, 0.1], since all the covariates are distributed as  $\mathcal{U}[0,1]$ , this would lead to  $10^{-10}\%$  of each sample with contaminated data, and even with the highest sample size of n = 600, this would represent an amount of  $6.10^{-8}$  observations.

With respect to the selection of the penalty and regularization parameters, both generalizations of the BIC criteria stated in Sections 3 and 4, respectively, were used. For the regularization parameters and for both robust and least squares estimators, equally spaced knots are used and the same number of terms to approximate each additive function were used, that is,  $k_j = k$  for all  $1 \leq j \leq 10$ , and so the selection problem consists of minimizing the simplified RBIC<sub>k</sub> defined in (9). As it was mentioned in Section 4, for n = 200, the grid for k was  $\{4; 5; \ldots; 13\}$ , for n = 400, tit was  $\{4; 5; \ldots; 14\}$  and, for n = 600, it was  $\{4; 5; \ldots; 15\}$ . For selecting the penalty parameters, an adaptive procedure was considered. The robust estimators proposed in (Boente & Martinez, 2023) were used as initial estimators for the robust proposal and the least-squares version for the LS-approach. Then, the RBIC<sub>k</sub> was minimized over the grid  $\{(\lambda_1, \lambda_2) : \lambda_1 \in \{0.05; 0.1; \ldots; 0.35\}, \lambda_2 \in \{0.20; 0.25; \ldots; 0.50\}\}$  for n = 200 and both robust and LS-estimators, for n = 400 and n = 600, the robust estimator was minimized over  $\{(\lambda_1, \lambda_2) : \lambda_1 \in \{0.15; 0.2; \ldots; 0.35\}, \lambda_2 \in \{0.05; 0.1; \ldots; 0.35\}, \lambda_3 \in \{0.05; 0.1; \ldots; 0.35\}, \lambda_4 \in \{0.05; 0.1; \ldots; 0.35\}$ 

Different measures were used for determining the variable selection results. For each sample, it is computed

C: the number of zero components correctly estimated to be zero,

IC: the number of zero components incorrectly estimated to be zero, and

CF: 1 if the true model is selected and 0 otherwise.

Then, they are averaged over the 500 replications to obtain the final values. All these three measures were calculated for both parametric and nonparametric components separately and also for the

complete model, and for both estimators. For instance, for both the parametric and nonparametric parts of the models, the closer it is the C measure to 6, the better. However, when the C measure is considered for the whole model, the closer to 12, the better. In all cases, the closer the IC measure is to 0 and the CF measure is to 1, the better.

To evaluate the performance of the parametric components, for each replication, we calculated the generalized mean square error (GMSE) defined as

$$GMSE = (\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta})^{\mathrm{T}} \Sigma (\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta})$$

where  $\Sigma \in \mathbb{R}^{q \times q}$  is the true variance-covariance matrix of **Z**. While for evaluating the performance of the estimator of the nonparametric components, we used the square root of the average square error (RASE) defined as

RASE = 
$$\sqrt{\frac{1}{m} \sum_{j=1}^{p} \sum_{k=1}^{m} \|\widehat{\eta}_{j}(t_{k}) - \eta_{j}(t_{k})\|^{2}}$$

where  $\{t_k\}_{k=1}^m$  is a grid of m = 1000 equidistant points in the interval [0, 1]. Then, once again, they are averaged over replications to obtain the final values and also, in order to visualize the variability, it is calculated the standard deviation for the GMSE and the RASE measures.

Both estimation measures, the GMSE for the parametric part and the RASE for the nonparametric components, were considered for the oracle estimators, that is, for the estimators obtained when using only the non-zero components.

Table 1 shows the measures obtained for the least squares and robust variable selection procedures for the parametric and nonparametric parts and for the complete model, under all contamination cases and the three sample sizes. When no contaminated data and for all sample sizes considered, both estimators throw similar values of the C, IC and CF measures. It can be appreciated that, in some cases, the robust approach performs slightly better than the LS-estimator, probably due to the more stable algorithm used to compute the robust estimator. For  $C_1$  to  $C_7$ , the least squares estimator throws much smaller values of the C and CF measures. By analizing the C measure, for instance, for the linear part of the model and under  $C_4$  and  $C_5$ , the C measure of the least squares estimator is reduced in more than a 42% with respect of the  $C_0$  for all sample sizes. A significant reduction in the C measure can also be observed for both the additive part and the complete model for the LS-estimator. For instance, for the complete model, under contamination  $C_4$ , the C values have been reduced to less than a 35% of the values obtained under the noncontaminated setting. In a more extreme way, for contamination  $C_7$ , the least squares estimator throws values of the C measure that are less than the 29% of the values obtained for noncontaminated data. On the other hand, across all the contamination scenarios and for all sample sizes, the robust approach shows a more stable behaviour, throwing values of the C measure close to the ones obtain under  $C_0$ . A similar analysis can be done for the CF measure for the three parts of the model. In this case, it is also even more notorious the bad performance of the LS-estimator across all the scenarios where, in some cases, the correct variables are never or almost never selected. With respect to the linear part of the model, when the sample size is 200, the correct variables are almost never selected under schemes  $C_4$  to  $C_7$ , and never or almost never under  $C_5$  to  $C_7$  for sample sizes 400 and 600. For the additive part, when n = 200 and under  $C_4$  to  $C_7$ , the values of the CF measure of the least squares estimator are close to 0. Moreover, under contaminations  $C_5$  to  $C_7$ , for the complete model, the values of the CF obtained by the LS-estimator under all the sample sizes considered are zero. This means that the components of whole model are never selected all at once. In contrast, the robust approach performs quite similarly across all the sample sizes and all contamination settings, yielding C measure values close to 6 for the linear and additive parts and to 12 for the complete model, and values of the CF measure close to 1, especially when the sample size increases. With respect to the IC measure, these values are 0 or close to 0 for both estimators and under all the different scenarios probably meaning that it is hard for the procedures to indicate as zero variables that are important to the model.

			Linear part		Additive part			Complete model			
n	Cont	Method	С	IC	CF	С	IC	CF	С	IC	CF
200	$C_0$	LS	5.12	0.00	0.76	5.95	0.00	0.95	11.07	0.00	0.73
200	$C_0$	ROB	5.55	0.00	0.88	5.92	0.00	0.93	11.47	0.00	0.84
200	$C_1$	LS	3.78	0.00	0.39	5.02	0.01	0.39	8.80	0.01	0.16
200	$C_1$	ROB	5.38	0.00	0.87	5.94	0.03	0.91	11.32	0.03	0.81
200	$C_2$	LS	3.31	0.00	0.29	4.82	0.01	0.29	8.13	0.01	0.10
200	$C_2$	ROB	5.51	0.00	0.90	5.94	0.01	0.95	11.45	0.01	0.86
200	$C_3$	LS	2.98	0.00	0.19	3.80	0.03	0.11	6.78	0.03	0.03
200	$C_3$	ROB	5.65	0.00	0.90	5.93	0.00	0.94	11.58	0.00	0.86
200	$C_4$	LS	2.96	0.01	0.07	2.11	0.06	0.01	5.07	0.07	0.00
200	$C_4$	ROB	5.79	0.00	0.94	5.95	0.00	0.96	11.74	0.00	0.90
200	$C_5$	LS	2.91	0.04	0.01	0.72	0.06	0.00	3.64	0.10	0.00
200	$C_5$	ROB	5.75	0.00	0.95	5.99	0.02	0.97	11.74	0.02	0.93
200	$C_6$	LS	1.37	0.01	0.00	2.95	0.03	0.01	4.32	0.04	0.00
200	$C_6$	ROB	5.59	0.00	0.90	5.91	0.00	0.93	11.50	0.00	0.86
200	$C_7$	LS	1.44	0.01	0.00	2.62	0.03	0.00	4.06	0.04	0.00
200	$C_7$	ROB	5.60	0.02	0.85	5.89	0.01	0.94	11.49	0.03	0.82
400	$C_0$	LS	5.64	0.00	0.92	6.00	0.00	1.00	11.64	0.00	0.92
400	$C_0$	ROB	5.79	0.00	0.95	5.99	0.00	1.00	11.79	0.00	0.95
400	$C_1$	LS	4.64	0.00	0.66	5.77	0.00	0.80	10.41	0.00	0.54
400	$C_1$	ROB	5.87	0.00	0.98	6.00	0.00	1.00	11.87	0.00	0.97
400	$C_2$	LS	4.37	0.00	0.56	5.62	0.00	0.65	9.99	0.00	0.37
400	$C_2$	ROB	5.94	0.00	0.99	6.00	0.00	1.00	11.94	0.00	0.99
400	$C_3$	LS	3.50	0.00	0.37	5.10	0.00	0.36	8.59	0.00	0.17
400	$C_3$	ROB	5.85	0.00	0.97	5.99	0.00	1.00	11.84	0.00	0.97
400	$C_4$	LS	2.87	0.00	0.17	3.86	0.02	0.04	6.73	0.02	0.01
400	$C_4$	ROB	5.95	0.00	0.99	5.99	0.00	1.00	11.94	0.00	0.99
400	$C_5$	LS	2.95	0.00	0.05	1.68	0.06	0.00	4.63	0.06	0.00
400	$C_5$	ROB	5.96	0.00	0.99	6.00	0.00	1.00	11.96	0.00	0.99
400	$C_6$	LS	0.79	0.00	0.00	4.73	0.01	0.18	5.52	0.01	0.00
400	$C_6$	ROB	5.93	0.00	0.98	5.98	0.00	0.99	11.92	0.00	0.98
400	$C_7$	LS	0.89	0.00	0.00	4.53	0.01	0.14	5.42	0.01	0.00
400	$C_7$	ROB	5.90	0.00	0.97	6.00	0.00	1.00	11.90	0.00	0.96
600	$C_0$	LS	5.88	0.00	0.97	6.00	0.00	1.00	11.88	0.00	0.97
600	$C_0$	ROB	5.94	0.00	0.99	6.00	0.00	1.00	11.94	0.00	0.99
600	$C_1$	LS	5.11	0.00	0.78	5.88	0.00	0.88	10.99	0.00	0.70
600	$C_1$	ROB	5.94	0.00	0.99	6.00	0.00	1.00	11.94	0.00	0.99
600	$C_2$	LS	4.97	0.00	0.75	5.89	0.00	0.89	10.86	0.00	0.67
600	$C_2$	ROB	5.95	0.00	0.99	6.00	0.00	1.00	11.95	0.00	0.99
600	$C_3$	LS	4.28	0.00	0.54	5.56	0.00	0.63	9.84	0.00	0.36
600	$C_3$	ROB	5.95	0.00	0.99	6.00	0.00	1.00	11.95	0.00	0.99
600	$C_4$	LS	3.14	0.00	0.27	4.66	0.01	0.16	7.80	0.01	0.05
600	$C_4$	ROB	5.92	0.00	0.98	5.99	0.00	1.00	11.91	0.00	0.98
600	$C_5$	LS	2.78	0.00	0.09	2.63	0.06	0.00	5.40	0.06	0.00
600	$C_5$	ROB	5.95	0.00	0.99	6.00	0.00	1.00	11.95	0.00	0.99
600	$C_6$	LS	0.43	0.00	0.00	5.36	0.00	0.49	5.78	0.00	0.00
600	$C_6$	ROB	5.92	0.00	0.98	6.00	0.00	1.00	11.92	0.00	0.98
600	$C_7$	LS	0.48	0.00	0.00	5.20	0.00	0.40	5.68	0.00	0.00
600	$C_7$	ROB	5.97	0.00	0.99	5.99	0.00	1.00	11.96	0.00	0.99

Table 1: C, IC, CF obtained for the least squares and robust approaches, for different sample sizes.

Table 2 shows the results of the estimation measures GMSE and RASE for the linear part and

the additive part, respectively, for the penalized and the oracle LS- and robust estimators, under different contamination settings and different sample sizes. For each estimator, it was computed the mean of the estimation measure, denoted MEAN, and also the standard deviation, denoted SD. As it is expected, under the noncontaminated scheme  $C_0$ , both the oracle and penalized robust estimators throw slightly larger values of these measures than their least squares counterparts under the different sample sizes considered, caused by the lack of efficiency of the robust estimators usually have. Besides, under this noncontaminted setting, the penalized approaches throw similar values to the ones obtained by the oracle versions since, as it was already mentioned when analysing Table 1, the correct model was highly selected. This behaviour can also be appreciated by observing the standard deviations obtained. For the contamination schemes  $C_1$  to  $C_7$ , similar values of the oracle estimator with respect to the penalized estimator can be still observe but only for the robust approach showing a more stable behaviour across all the contamination schemes and for both GMSE and RASE measures. However, despite of the greater stability of the robust approach, it can be appreciated that, for the  $C_7$  case and the GMSE measure, the robust proposal yields considerably higher results than the oracle estimator for all sample sizes, along with much larger standard deviations. It can be checked that a few atypical observations appeared in the data under this contamination setting (see, for instance Figures 2 and 3). For n = 200, a 7.4% of the GMSE values were detected as outliers by the adjusted boxplot, which is a boxplot adapted to asymmetric data. For n = 400 and n = 600, these percentages are reduced to 4% and 1.4%, respectively. Table 3 shows the results of the 10%-trimmed mean of the GMSE values for the oracle and penalized LS- and robust estimators. By computed the trimmed mean, the results obtained by the robust approach are quite similar to those obtained by its oracle version. Let us continue analysing the GMSE measure. When looking only the penalized estimators, the robust proposal shows a more stable behaviour across all the contamination settings than its least squares counterpart. On the contrary, the LS-estimator shows a more erratic behaviour under all sample sizes and all contamination settings, especially under  $C_5 \circ C_7$  where the mean of GMSE obtained are between 51 and 913 times larger than under the  $C_0$  case. In order to visualize the results obtained for the GMSE even more, Figures 2 and 3 show the adjusted boxplots for contaminations  $C_0, C_1, C_3, C_4$  and  $C_7$  and sample sizes n = 200 and n = 600, respectively. A similar analysis can be done for the RASE measure, that is, the measure used for the additive part. Under all sample sizes, when no contaminated data, both oracle estimators and both penalized estimators behave similarly between them, with also the robust approaches throwing slightly larger values than the least squares counterparts, as it was already mentioned. Across the contamination schemes  $C_1$  to  $C_7$ , both estimators based on least squares throw larger values of the RASE than the ones obtained under  $C_0$ , especially under  $C_4$  to  $C_7$ , achieving values up to 8 times larger. On the other hand, the robust proposal, and also its oracle version, shows a more stable behaviour across all the contamination scenarios. In order to visualize these behaviours, Figures 4 and 5 show the adjusted boxplots of the RASE values for contaminations  $C_0$ ,  $C_1$ ,  $C_3$ ,  $C_4$  and  $C_7$ , for the oracle and penalized estimators, and sample sizes n = 200 and n = 600, respectively.

In order to see how the penalized approaches estimates the additive functions, Figures 6 to 10 show the functional boxplots obtained for n = 200 under the contamination settings  $C_0$ ,  $C_1$ ,  $C_2$ ,  $C_4$  and  $C_7$ , respectively, for the least squares and robust estimators. Similar plots can be obtained for sample sizes n = 400 and n = 600. Functional boxplots were introduced by Sun & Genton (2011) and are a useful tool to visualize a collection of curves. The area in purple represents the 50% inner band of curves, and the solid blue and the dotted red lines correspond to the whiskers and outlying curves, respectively. Moreover, the black line indicates the deepest function. The true

		GMSE						RASE				
			Ora	cle	Pena	lized	Ora	acle	Penalized			
n	Cont	Method	MEAN	SD	MEAN	SD	MEAN	SD	MEAN	SD		
200	$C_0$	LS	0.022	0.016	0.026	0.021	0.270	0.064	0.278	0.082		
200	$C_0$	ROB	0.023	0.016	0.027	0.023	0.284	0.074	0.283	0.080		
200	$C_1$	LS	0.064	0.068	0.105	0.135	0.436	0.131	0.522	0.227		
200	$C_1$	ROB	0.036	0.025	0.044	0.037	0.355	0.103	0.369	0.128		
200	$C_2$	LS	0.076	0.062	0.129	0.102	0.481	0.140	0.557	0.179		
200	$C_2$	ROB	0.029	0.020	0.034	0.028	0.313	0.083	0.314	0.101		
200	$C_3$	LS	0.132	0.127	0.261	0.234	0.625	0.258	0.805	0.344		
200	$C_3$	ROB	0.026	0.018	0.028	0.021	0.302	0.086	0.298	0.092		
200	$C_4$	LS	0.249	0.198	0.527	0.384	0.859	0.226	1.252	0.418		
200	$C_4$	ROB	0.025	0.017	0.026	0.019	0.297	0.081	0.289	0.084		
200	$C_5$	LS	0.654	0.494	1.522	0.959	1.370	0.331	2.222	0.519		
200	$C_5$	ROB	0.027	0.018	0.029	0.023	0.299	0.099	0.307	0.106		
200	$C_6$	LS	12.172	0.445	6.708	0.368	0.962	0.236	1.027	0.354		
200	$C_6$	ROB	0.025	0.018	0.027	0.021	0.299	0.086	0.295	0.094		
200	$C_7$	LS	12.450	0.359	6.826	0.373	0.973	0.232	1.112	0.375		
200	$C_7$	ROB	0.050	0.526	0.745	2.790	0.304	0.144	0.353	0.243		
400	$C_0$	LS	0.011	0.008	0.012	0.009	0.186	0.041	0.184	0.041		
400	$C_0$	ROB	0.012	0.008	0.012	0.009	0.192	0.044	0.188	0.040		
400	$C_1$	LS	0.031	0.027	0.042	0.040	0.309	0.084	0.335	0.112		
400	$C_1$	ROB	0.016	0.012	0.017	0.012	0.234	0.057	0.229	0.057		
400	$C_2$	LS	0.036	0.026	0.048	0.036	0.325	0.077	0.362	0.104		
400	$C_2$	ROB	0.014	0.010	0.014	0.010	0.211	0.050	0.205	0.046		
400	$C_3$	LS	0.064	0.052	0.101	0.079	0.421	0.117	0.484	0.155		
400	$C_3$	ROB	0.012	0.008	0.013	0.009	0.201	0.048	0.194	0.041		
400	$C_4$	LS	0.125	0.089	0.223	0.159	0.598	0.155	0.720	0.204		
400	$C_4$	ROB	0.012	0.008	0.012	0.009	0.198	0.048	0.192	0.043		
400	$C_5$	LS	0.314	0.230	0.620	0.382	0.949	0.218	1.374	0.287		
400	$C_5$	ROB	0.013	0.009	0.013	0.009	0.206	0.049	0.199	0.042		
400	$C_6$	LS	12.005	0.273	6.402	0.230	0.648	0.135	0.590	0.153		
400	$C_6$	ROB	0.012	0.008	0.012	0.009	0.198	0.047	0.191	0.040		
400	$C_7$	LS	12.293	0.210	6.491	0.225	0.665	0.141	0.621	0.154		
400	$C_7$	ROB	0.012	0.009	0.319	1.890	0.201	0.046	0.209	0.101		
600	$C_0$	LS	0.007	0.005	0.007	0.005	0.149	0.031	0.147	0.029		
600	$C_0$	ROB	0.008	0.005	0.008	0.005	0.154	0.032	0.151	0.030		
600	$C_1$	LS	0.022	0.018	0.026	0.028	0.253	0.072	0.267	0.093		
600	$C_1$	ROB	0.011	0.008	0.011	0.008	0.190	0.042	0.186	0.041		
600	$C_2$	LS	0.023	0.017	0.028	0.020	0.265	0.061	0.285	0.086		
600	$C_2$	ROB	0.009	0.006	0.009	0.006	0.169	0.036	0.164	0.032		
600	$C_3$	LS	0.042	0.032	0.057	0.046	0.344	0.088	0.381	0.114		
600	$C_3$	ROB	0.008	0.006	0.008	0.006	0.160	0.036	0.155	0.031		
600	$C_4$	LS	0.077	0.063	0.129	0.096	0.494	0.118	0.568	0.150		
600	$C_4$	ROB	0.008	0.005	0.008	0.006	0.158	0.033	0.153	0.031		
600	$C_5$	LS	0.201	0.154	0.374	0.229	0.775	0.171	1.023	0.235		
600	$C_5$	ROB	0.009	0.006	0.009	0.006	0.164	0.035	0.161	0.036		
600	$C_6$	LS	11.970	0.221	6.300	0.177	0.525	0.108	0.462	0.116		
600	$C_6$	ROB	0.008	0.005	0.008	0.005	0.158	0.033	0.153	0.030		
600	$C_7$	LS	12.280	0.174	6.391	0.177	0.540	0.116	0.491	0.124		
600	$C_7$	ROB	0.008	0.006	0.132	1.162	0.161	0.034	0.162	0.061		

Table 2: GMSE and RASE obtained for the least squares and robust approaches and both oracle estimators, for different sample sizes.

functions are also plotted in solid green line. All y-limits were fixed across all the contamination scenarios for the ease of comparison. Recalling that Figure 6 shows the results obtained under  $C_0$ , it can be appreciated that both methods show similar estimates, for instance by noting that the black curves, corresponding to the deepest curve, and the purple boxes, corresponding to the 50% percent of the inner curves, are quite similar. Taking into account that only the first four additive functions are nonidentically zero functions and the last six are null functions, both estimators seem to be estimating the correct true functions (in green). In the last six functional boxplots, except for

		G	MSE
n	Method	Oracle	Penalized
200	LS	12.437	6.790
200	ROB	0.024	0.028
400	LS	12.292	6.471
400	ROB	0.011	0.012
600	LS	12.277	6.375
600	ROB	0.007	0.008

Table 3: 10%-trimmed mean of the GMSE values for the LS and robust approaches and both oracle estimators, for different sample sizes and under the  $C_7$  contamination setting.

the constant curves in green that are clearly visible, the boxes are almost indistinguishable and only the few outlying curves can be seen. It is also worth noting, especially for the first four additive functions, that near the boundaries, for both robust and least squares estimators, the boxes tend to be wider. The reason of this is the well-known bad behaviour of spline approximations in the boundaries due to the reduced number of data points. This can also be observed for the rest of the figures. To study the effect of the atypical data, Figures 7 to 10 show the same functional boxplots but for contamination squemes  $C_1, C_2, C_4$  and  $C_7$ . Recall that none of the contamination settings considered introduce outliers in the additive components of the model and, for this reason, the black curves representing the deepest curves are always close to the true one (in green). However, these functional boxplots allow us to see the general behaviour of the estimators by observing the width of the boxes and whiskers and the outlying curves. For instance, Figure 7, which shows the functional boxplots when considering the contamination scheme  $C_1$ , for the LS-estimator, boxes of the first four covariates are slightly larger than under  $C_0$  and present a few outlying curves. For the last six additive functions, the presence of outlying curves is quite remarkable. On the other hand, the robust estimator seems to show a behavior similar to that of  $C_0$ . An analogous analysis can be done for the contamination scenario  $C_2$ . However, for contamination scenarios  $C_4$  and  $C_7$ , the notorious poor performance of the least squares proposal can be observed in all additive functions. Furthermore and in contrast to the other settings, an increase of the size of the boxes and longer whiskers can be appreciated when the approach based on LS estimates the null functions. On the other hand, the robust approach seems to consistently estimate the ten additive functions very well, leading to estimations similar to those obtained under  $C_0$ . Interestingly and as it was already mentioned, under the most extreme contamination setting considered, that is, under  $C_7$  (see Figure 10), the robust approach presents some extra outlying curves that did not appear in the other plots. However, this unexpected behavior is not so bad when compared to the damage caused to the least squares-based estimator.



Figure 2: Adjusted boxplots for the GMSE values under  $C_0$ ,  $C_1$ ,  $C_3$ ,  $C_4$  and  $C_7$  for both oracle estimators and both penalized estimators for sample size n = 200.



Figure 3: Adjusted boxplots for the GMSE values under  $C_0$ ,  $C_1$ ,  $C_3$ ,  $C_4$  and  $C_7$  for both oracle estimators and both penalized estimators for sample size n = 600.



Figure 4: Adjusted boxplots for the RASE values under  $C_0$ ,  $C_1$ ,  $C_3$ ,  $C_4$  and  $C_7$  for both oracle estimators and both penalized estimators for sample size n = 200.



Figure 5: Adjusted boxplots for the RASE values under  $C_0$ ,  $C_1$ ,  $C_3$ ,  $C_4$  and  $C_7$  for both oracle estimators and both penalized estimators for sample size n = 600.

Figure 6: The first two rows contain the functional boxplots of the estimated additive functions using the least square-based estimator while the last two rows contain the functional boxplots of the estimated additive functions by the robust approach, for n = 200 when no contaminated data.





Figure 7: The first two rows contain the functional boxplots of the estimated additive functions using the least square-based estimator while the last two rows contain the functional boxplots of the estimated additive functions by the robust approach, for n = 200 and the contamination setting  $C_1$ .

0.0

5.0-

50

s.0 0.0 2.0-

Figure 8: The first two rows contain the functional boxplots of the estimated additive functions using the least square-based estimator while the last two rows contain the functional boxplots of the estimated additive functions by the robust approach, for n = 200 and the 8 . 8 . 8 8 3 0.1 8 0.1 contamination setting  $C_2$ . 8 . 8

0.0

5.0-

5.0



0

510 0.0 9.0-

Figure 9: The first two rows contain the functional boxplots of the estimated additive functions using the least square-based estimator while the last two rows contain the functional boxplots of the estimated additive functions by the robust approach, for n = 200 and the contamination setting  $C_4$ .





Figure 10: The first two rows contain the functional boxplots of the estimated additive functions using the least square-based estimator while the last two rows contain the functional boxplots of the estimated additive functions by the robust approach, for n = 200 and the contamination setting  $C_7$ .

0.0 2.0-

50

s.0 0.0 2.0-

It is interesting to see which are the  $\tilde{\lambda}_1$  and  $\tilde{\lambda}_2$  selected for each sample and each penalized estimator across the contamination settings. Figures 11 and 12 show the boxplots of the  $\tilde{\lambda}_1$  and  $\tilde{\lambda}_2$  show the boxplots of the selected when n = 200, respectively, while Figures 13 and 14 show the selected parameters under n = 400 and Figures 15 and 16 for n = 600. It can be appreciated that under all contamination scenarios, for a fixed sample size, the robust approach tends to select similar values of the parameters while the least squares estimators vary the selected values yielding to very different boxplots. When n = 200 and for selecting  $\tilde{\lambda}_1$ , except for contamination  $C_5$ , the sizes of the boxes of the robust approach are quite similar. All the boxplots present whiskers to both sides and the median value is located at the same number. A different behaviour is observed for contamination  $C_5$ , where the top whisker is much shorterr than in the other settings, the box is less than a half of the otherss, the median is located at a lower position and there are right-side outliers. On the other hand, the boxes of the boxplots of the selected  $\tilde{\lambda}_1$  by the least squares estimator are located at different values of y-axis: at the bottom of the plot ( $C_1$ ,  $C_2$ ,  $C_3$  and  $C_4$ ), in the middle of the plot ( $C_0$ ,  $C_6$  and  $C_7$ ) and at the top of the plot ( $C_5$ ), leading to different locations of the median values.

With respect to the selection of  $\lambda_2$ , Figure 12 shows a similar behavior. While the values selected by the robust approach are quite consistent, reflected in similar boxplots with comparable boxes, medians, and whisker lengths, the least squares estimator displays a different pattern. Although the boxes are of similar sizes (except in the  $C_5$  case), the medians vary slightly, and the whiskers range from shorter ones (in  $C_0$  to  $C_3$ ) to longer ones (in  $C_4$ ,  $C_6$ , and  $C_7$ ). Under contamination setting  $C_5$ , the LS-estimator seems to select values in a certian what that the box spans nearly the entire grid range for  $\lambda_2$ . For n = 400, shown in Figures 13 and 14, the more stable behavior of the robust estimator across scenarios is evident for both tuning parameters, particularly for  $\lambda_1$ , where all boxplots are positioned at the lower end of the plot. A distinct behavior appears under  $C_5$ , where the box is reduced to the value 0.05. For the selection of  $\lambda_2$ , the boxplots across contaminated settings are quite similar to that of the uncontaminated scenario  $C_0$ . In contrast, the values of  $\lambda_1$  under  $C_0$  for the LS-estimator show a small box with whiskers on both sides and some right-side outliers, while other settings show much larger boxes, differing medians, and different lengths of whiskers. A similarly unpredictable pattern is seen in the least-squares approach for selecting  $\lambda_2$ ; under  $C_0$ , the boxplot has its median at 0.3, centered with whiskers on both sides, but the other settings display larger medians and a range of box sizes, from narrower (e.g.,  $C_1$  or  $C_4$ ) to much larger ones (e.g.,  $C_5$ ). Finally, for n = 600 and selecting  $\lambda_1$  (see Figure 15), the robust estimator consistently selects the same value, 0.05, except for some outliers. However, the least squares approach shows greater variability, with larger boxes. As in the previous sample sizes, box sizes, median locations, and whisker lengths vary across contamination schemes. For selecting  $\lambda_2$ (Figure 16), the robust approach continues to show consistent boxplots, with boxes of similar size, whiskers on both sides, and medians at close values. The LS-based approach, in contrast, behaves differently. Under  $C_0$ , the box is centered on the y-axis with symmetrical whiskers and a median at 0.25. For other contamination scenarios, except  $C_4$ , the bottom whisker is slightly longer, and medians are equal or greater than 0.4. In  $C_3$  to  $C_7$ , the top whiskers are missing, in  $C_4$  the box is narrower and the left whisker shorter.

Figure 11: Boxplots of the  $\tilde{\lambda}_1$  selected under different contamination schemes for both least squares and robust estimators, when n = 200. ő ပဳ ഗ് Q ₽ ပဳ ပိ Ω ഗ്







 $_{\mathrm{LS}}$ 

Figure 12: Boxplots of the  $\tilde{\lambda}_2$  selected under different contamination schemes for both least squares and robust estimators, when n = 200. ő ပဳ ပိ - v ပဳ ပိ ပိ ۲.0 S.0





S.0

**Þ**.0

6.0

6.0

0

0

٢.0

0

0

0

0.0

ROB

Figure 13: Boxplots of the  $\tilde{\lambda}_1$  selected under different contamination schemes for both least squares and robust estimators, when n = 400.







0.1E

01.0

ο

0.20

0

ο

0

95.0

0

05.0

0

0.25

ROB

ů

ပဳ

ပိ

Q ₽

ပဳ

ΰ

Q

ပိ

-----

.....

. . . . . .

00.0

90<sup>.</sup>0





 $\mathbf{LS}$ 

S.0

6.0

6.0

**Þ**.0

ROB

ů

– ပိ

. م

0 - 3

ပဳ

- S

- ت

ပိ

0.0

0

0 0

0

0

0

٢.0

о

0 0 0

Figure 15: Boxplots of the  $\tilde{\lambda}_1$  selected under different contamination schemes for both least squares and robust estimators, when n = 600.

0	0	o	o	o	0	0 - 1
0	o	o	o	o	o	o – ,
0	o	o		0	o	0 - 0
0	0	o	o	o	o	0 - 0
0	0	0	o	o	0	0 - 0
0	o	o	o	o	o	o – ,
o	o	o	o	o	o	o – .
0	0	o	o	o	o	o – ,
6.0	0:30	0.25	0.20	٥r.0 ا	01.0	S0.0 00.0

ROB



 $\mathbf{LS}$ 



Figure 16: Boxplots of the  $\tilde{\lambda}_2$  selected under different contamination schemes for both least squares and robust estimators, when n = 600.

 $\mathbf{LS}$ 

S.0

٢.0

6.0

ROB

ů

- ပိ

- రో

- 0ţ

ပိ

ő

ū

ပိ

0.0

o

0

0

0 0

0

0

0

0 0

ŀ

6.0

**Þ**.0

# 8 A real data example

In this section, we analyze a real dataset: the well-known plasma beta-carotene level dataset collected by Nierenberg et al. (1989). This dataset is of significant interest as studies have shown a direct relationship between beta-carotene levels and certain types of cancer (see Fairfield & Fletcher, 2002). It contains 315 observations and 14 variables and is available in R as the dataset plasma within the library gamlss.data.

The aim is to explore the relationship between plasma beta-carotene levels and various regulatory factors, including age, sex, dietary intake, smoking status, and alcohol consumption. The response variable, plasma beta-carotene (measured in ng/ml) and denoted as BETAPLASMA, will be modeled using the following covariates:

AGE: age (years);

SEX: sex (1=male, 2=female);

SMOKSTAT: smoking status (1=never, 2=former, 3=current smoker);

QUET: quetelet  $(weight/(height)^2);$ 

VITUSE: vitamin use (1=yes, fairly often, 2= yes, not often, 3=no);

CALORIES: number of calories consumed per day;

FAT: grams of fat consumed per day;

FIBER: grams of fiber consumed per day;

ALCOHOL: number of alcoholic drinks consumed per week;

CHOL: cholesterol consumed (mg/day);

BETADIET: dietary beta-carotene consumed (mcg/day).

As it is pointed out by Liu et al. (2011) and Guo et al. (2013), there is one extremely high leverage point in ALCOHOL which, for obvious reasons, we will not remove. The categorical variables were divided into dummy variables before being included in the model. Thus, SMOKSTAT enters the model as SMOK1 (1=former smoker, 0=other) and SMOK2 (1=current smoker, 0=other) and VITUSE is separated into VIT1 (1=fairly often, 0=other) and VIT2 (1=not often, 0=other). Besides, all variables except binary ones are standarized using the MEDIAN as the center and the MAD as the deviation, assuming that some covariates may present high-leverage points.

This dataset has been widely studied by several authors, especially in the context of semiparametric models and, in particular, in partially linear additive models. For instance, in Liu et al. (2011), an approach based on least squares is used to model the logarithm of BETAPLASMA with nine covariates: QUET, CALORIES, FAT, FIBER, BETADIET, SEX, ALCOHOL, SMOK1 and SMOK2 entering in the linear part, and AGE and CHOL in the additive part. The same model was considered in Banerjee & Ghosal (2014) using a Bayesian model selection method, while Jeong et al. (2022) also used a Bayesian model selection approach but to study the specific effect of the CHOL covariate. On the other hand, Guo et al. (2013) proposed a composite quantile regression approach with a regularization procedure to select variables from the linear component, modeling BETAPLASMA without transforming it. They included ten variables in the linear part: SEX, SMOK1, SMOK2, QUET, VIT1, VIT2, CALORIES, FAT, ALCOHOL and BETADIET, while AGE, CHOL and FIBER were included in the additive part. Lv et al. (2017) considered a modal regression approach to model BETAPLASMA with ten covariates entering in the linear part: SEX, SMOKSTAT, QUET, VITUSE, CALORIES, FAT, ALCOHOL, BETADIET and two other covariates named RETDIET and RETPLASMA, and the same three covariates entering in the additive component of the model: AGE, FIBER and CHOL. While SMOKSTAT was treated as a discrete variable in their analysis, converting it into dummy variables might align better with standard modeling practices. Finally, Boente & Martinez (2024) considered the same data set and proposed the same model as in Guo et al. (2013), but with variable selection applied only to the linear component of the model.

For this reason, we adopt the same model as Guo et al. (2013), where AGE, CHOL and FIBER enter to the model in the additive part, while the remaining variables are considered to have a linear relationship with plasma beta-carotene levels. More precisely, the partially linear additive model considered is the following:

$$\begin{aligned} \text{BETAPLASMA} &= \mu + \beta_1 \text{SEX} + \beta_2 \text{SMOK1} + \beta_3 \text{SMOK2} + \beta_4 \text{QUET} + \beta_5 \text{VIT1} + \beta_6 \text{VIT2} \\ + \beta_7 \text{CALORIES} + \beta_8 \text{FAT} + \beta_9 \text{ALCOHOL} + \beta_{10} \text{BETADIET} + \eta_1 (\text{AGE}) \\ + \eta_2 (\text{CHOL}) + \eta_3 (\text{FIBER}) + \sigma \varepsilon, \end{aligned}$$

where the errors  $\varepsilon$  are assumed to be independent of the covariates, with symmetric distribution and scale 1. With the notation given in equation (1),  $\mathbf{Z} = (\text{SEX}, \text{SMOK1}, \text{SMOK2}, \text{QUET}, \text{VIT1}, \text{VIT2}, \text{CALORIES}, \text{FAT}, \text{ALCOHOL}, \text{BETADIET})^{\text{T}} \in \mathbb{R}^{10}$  and  $\mathbf{X} = (X_1, X_2, X_3)^{\text{T}} = (\text{AGE}, \text{CHOL}, \text{FIBER})^{\text{T}} \in \mathbb{R}^3$ .

For the analysis, two penalized estimators are computed: the robust estimator and the one based on least squares, in a similar way to what has been done in the simulation study. For the robust approach, it is used the Tukey's bisquare loss function as  $\rho$ -function with tuning constant c = 4.685, while for the LS-estimator, the quadratic function  $\rho(t) = t^2$ . Cubic *B*-splines are used to estimate the additive components. When estimating  $\eta_j$ , the internal knots are taken as the  $\ell/(k + 1)100\%$  quantiles,  $\ell = 1, \ldots, k$ , of the observed values of the covariate  $X_j$ . The number of internal knots is the same for all three covariates included in the nonparametric part of the model, i.e.  $k_j = k$  for j = 1, 2, 3. The regularization parameter k and the auxiliary penalty parameters  $\tilde{\lambda}_1$  and  $\tilde{\lambda}_2$  are automatically selected using the RBIC<sub>k</sub> and RBIC<sub> $\lambda$ </sub> criteria, respectively, as described in this work. For the robust estimators, the Tukey's bisquare loss function is used, while for the LS-estimator it is used  $\rho(t) = t^2$ . The grid considered for  $\tilde{\lambda} = (\tilde{\lambda}_1, \tilde{\lambda}_2)$  is  $\mathcal{G} = \{(\ell_1, \ell_2) : \ell_1 \in \{0; 0.01; 0.02; \ldots; 0.1\}, \ell_2 \in \{0; 0.1; 0.2; \ldots; 2\}\}$ , as this grid is used for both robust and least-squares approaches.

In order to compare the behaviour of the estimators, 100 observations are randomly selected for prediction purposes, while the remaining 215 observations are used for estimation and variable selection. This procedure is repeated 50 times. For each iteration, the prediction performance of an estimator is measured through the median absolute prediction error (MAPE), calculated as the median of  $\{|y_{\ell} - \hat{y}_{\ell}|, \ell = 1, ..., 100\}$  where  $(y_{\ell}, \mathbf{z}_{\ell}^{\mathrm{T}}, \mathbf{x}_{\ell}^{\mathrm{T}})^{\mathrm{T}}, 1 \leq \ell \leq 100$  represents the observations reserved for prediction and  $\hat{y}_{\ell}$  is the predicted value at  $(\mathbf{z}_{\ell}^{\mathrm{T}}, \mathbf{x}_{\ell}^{\mathrm{T}})^{\mathrm{T}}$ . The final number of this error is calculated as the mean of the 50 MAPE values. To evaluate the variable selection procedure, for each iteration and each estimator (robust and least-squares), the number of covariates identified as important and included in the model is recorded. These counts are averaged across the 50 replications to obtain the final number, named AV.SIZE.

All these measures are calculated for both the penalized robust approach and its least squares counterpart (named PENALIZED). Additionally, unpenalized estimators that do not select variables are computed. These correspond to the estimators defined in Boente & Martinez (2023) for the robust approach and to the least squares approach when using  $\rho(t) = t^2$  as  $\rho$ -function. Note that, since unpenalized estimators do not perform variable selection, their AV.SIZE are 13.

Table 4 summarizes the mean of the MAPE values obtained over the 50 replications together with the average number of covariates selected (AV.SIZE). Although the unpenalized robust estimator achieves the lowest MAPE value, the robust approach that selects variables slightly increases the prediction error while reducing the average number of selected variables to 5.32, compared to the 13 variables included in the model by the unpenalized approach. On the other hand, both LS-estimators yield higher MAPE values than their robust counterparts. Moreover, for the penalized LS-estimator, the average size of the selected model is 7.54, which is notably larger than that of penalized robust approach.

For better visualization, Figure 17 displays adjusted boxplots for the 50 MAPE values obtained from each of the four estimators. The two boxplots on the left correspond to the unpenalized estimators (those that do not perform variable selection), while the two on the right correspond to the penalized estimators that incorporate regularization. It can be appreciated that the boxplots for the robust approaches are consistenly lower than those of the least squares estimators. For instance, more than 75% of the MAPE values obtained by the robust estimators are smaller that at least 75% of the values obtained by their LS-counterparts. Additionally, as previously noted, the adjusted boxplot for the penalized robust estimator is spligthly higher than that of its unpenalized competitor.

METHOD	MAPE	AV.SIZE
PENALIZED LS	0.9113	7.54
PENALIZED ROB	0.6987	5.32
LS	0.8920	13
ROB	0.6365	13
PENALIZED LS <sup>(-OUT)</sup>	0.6598	5.96

Table 4: MAPE and AV.SIZE for the penalized least squares and robust approaches, for the unpenalized least squares and robust estimators in the first two rows and, in the last row, for the LS-estimator with the atypical observations removed.

To determine which variables are more frequently selected by each method, Tables 5 and 6 present the frequencies with which each variable is identified as non-zero. Table 5 corresponds to the ten variables in the parametric component, while Table 6 focuses on the three covariates in the nonparametric part. By considering a threshold of 0.5, the least squares estimator selects the variables SEX, SMOK2, QUET, VIT1, VIT2, FAT and BETADIET from the linear component, as well as FIBER from the additive component. In contrast, the robust approach selects SEX, SMOK2, QUET, VIT1 and BETADIET from the linear component, and none from the additive component.



Figure 17: Adjusted boxplots for the MAPE values obtained for the unpenalized estimators (on the left) and for the penalized ones (on the right).

PENALIZED	SEX	SMOK1	SMOK2	QUET	VIT1	VIT2	CALORIES	FAT	ALCOHOL	BETADIET
LS	0.78	0.22	0.90	1.00	1.00	0.80	0.32	0.56	0.14	0.94
ROB	0.66	0.40	0.62	0.88	0.54	0.42	0.32	0.24	0.02	0.72
LS <sup>(-OUT)</sup>	0.88	0.34	0.88	1.00	0.76	0.52	0.16	0.38	0.00	0.98

Table 5: Selected frequencies as non-zero for the ten covariates of the parametric component, for the least squares approach in the first row, the robust proposal in the second row and, in the last row, the least squares approach computed without the atypical observations.

PENALIZED	AGE	CHOL	FIBER
LS	0.24	0.04	0.60
ROB	0.06	0.04	0.40
LS <sup>(-OUT)</sup>	0.02	0.02	0.02

Table 6: Selected frequencies as non-zero for the three covariates of the additive component, for the least squares approach in the first row, the robust proposal in the second row and, in the last row, the least squares approach computed without the atypical observations.

Since outlying observations may influence the estimator based on least squares, resulting in higher MAPE values, the penalized robust approach is computed on the whole dataset to identify large residuals. The boxplot of residuals obtained using the robust method, shown in Figure 18, reveals the presence of 17 vertical outliers, that is, 17 observations detected with large residuals.

Subsequently, the procedure is repeated for the penalized LS-estimator after excluding the 17



Figure 18: Boxplot of the residuals obtained when fitting the complete data set using the penalized robust estimator.

vertical outliers and also the high leverage observation in ALCOHOL identified by Liu et al. (2011) and Guo et al. (2013). That is, for each of the 50 iterations, out of a total of 297 observations, 100 are randomly excluded for testing, while the 197 remaining are used for training the estimator. The MAPE and AV.SIZE values are then calculated. The results, shown in the last row of Table 4 (labeled with an upperscript (-OUT)), show the average MAPE and model size values. Figure 19 shows the adjusted boxplots of the MAPE values obtained by the same four estimators as before but with the recently computed LS-based estimator with fewer observations.

As it can be appreciated in Table 4, the averaged MAPE and AV.SIZE for this estimator are now quite similar to those obtained by the penalized robust approach. Additionally, Figure 19 displays adjusted boxplots for the MAPE values of the same four estimators as before, now including the LS-estimator computed on the reduced sample, which is in fact quite similar to those obtained by the robust estimators. Finally, last rows of Tables 5 and 6 indicate that the proportion of times each covariate is selected by this reduced-sample LS-estimator aligns closely with the selection frequencies of the robust method. Except for VIT2, both the robust approach and the LS-estimator on the reduced sample select the same covariates.

# 9 Concluding remarks

Partially linear additive models (PLAM) are a suitable choice for modeling a response variable using a set of covariates that may have either a linear or nonparametric relationship with the response. Even when the relationship of a covariate is unknown, the linear part of the model allows for the inclusion of a certain type of covariates, such as categorical or discrete ones, which cannot be easily accommodated by fully nonparametric models.

In practice, practitioners typically include all available variables in the model, assiging them to the linear or additive components based on their expertise or the intrinsic nature of the covariates.



Figure 19: Adjusted boxplots for the MAPE measures obtained for the estimators without penalization (on the left) and for the penalized (on the right) estimators. The one corresponding to the penalized least squares estimator computed without the outliers is labelled  $LS^{(-OUT)}$ .

However, including too many covariates can lead to issues such as lack of parsimonity or reduced predicted performance. Variable selection, therefore, becomes crucial, and much research has focused on it, especially on selecting variables from the linear part of the model. However, the excess of covariates can arise not only in linear component but also in the additive part of a PLAM, making this an interesting and relatively unexplored are of investigation.

On the other hand, since atypical data may arise, robust procedures are needed not only for accurately estimating the relationship between the response variable and the covariates but also for correctly identifying and selecting significant variables while discarding irrelevant ones.

To address these challenges, this paper introduces a robust method for estimating and selecting variables in sparse partially linear additive models. The proposal includes automatic selection of regularization and penalty parameters, achieved through generalized BIC criteria. A general algorithm is developed to compute the proposed estimator, with the option to incorporate an adaptive procedure for penalty parameters. A Monte Carlo study, conducted with three sample sizes and eight contamination scenarios (including a non-contaminated case), is used to compare the robust proposal with its least squares counterpart. Finally, the approach is evaluated using the well-known plasma beta-carotene level dataset. The results from both the simulation study and the real-data analysis highlight the advantages of the robust proposal, particularly in the presence of atypical data.

Acknowledgements. This research was partially supported by grants Proyecto Interno CD-CBLUJ 204/19 from the Departamento de Ciencias Básicas, Universidad Nacional de Luján, PICTO-2021-UNLU-

00016 from ANPCYT and Universidad Nacional de Luján, and PICT-2021-I-A-00260 from ANPCYT, Argentina.

# References

- Avella-Medina, M. (2016). Robust penalized M-estimators for generalized linear and additive models. PhD thesis, Geneva School of Economics and Management, University of Geneva, Switzerland (supervisor Ronchetti, E.).
- Avella-Medina, M. & Ronchetti, E. (2018). Robust and consistent variable selection in high-dimensional generalized linear models. *Biometrika*, 105:31–44.
- Banerjee, S. & Ghosal, S. (2014). Bayesian variable selection in generalized additive partial linear models. Stat, 3:363–378.
- Bianco, A., Boente, G., & Chebi, G. (2022). Penalized robust estimators in logistic regression with applications to sparse models. *Test*, 31:563–594.
- Boente, G. & Martinez, A. (2023). A robust spline approach in partially linear additive models. Computational Statistics and Data Analysis, 178:107611.
- Boente, G. & Martinez, A. (2024). Robust variable selection for partially linear additive models. *Statistics* and *Computing*, 34:201.
- Boente, G., Martinez, A., & Salibian-Barrera, M. (2017). Robust estimators for additive models using backfitting. *Journal of Nonparametric Statistics*, 29:744–767.
- Breiman, L. (1996). Heuristics of inestability and statibilization in model selection. *The Annals of Statistics*, 24:2350–2383.
- Desboulets, L. D. D. (2018). A review on variable selection in regression analysis. *Econometrics*, 6:45.
- Du, P., Cheng, G., & Liang, H. (2012). Semiparametric regression models with additive nonparametric components and high dimensional parametric components. *Computational Statistics and Data Analysis*, 56:2006–2017.
- Efron, B. & Hastie, T. (2016). Computer Age Statistical Inference. Cambridge University Press.
- Fairfield, K. M. & Fletcher, R. H. (2002). Vitamins for chronic disease prevention in adults. Journal of the American Medical Association, 287:3116–3226.
- Fan, J. & Li, R. (2001). Variable selection via non-concave penalized likelihood and its oracle properties. Journal of the American Statistical Association, 96:1348–1360.
- Feng, M., Chen, Z., & Cheng, X. (2018). Two-step variable selection in partially linear additive models with time series data. Communications in Statistics - Simulation and Computation, 47:661–671.
- Frank, L. E. & Friedman, J. H. (1993). A statistical view of some chemometrics regression tools. *Techno-metrics*, 35:109–135.
- Fu, W. J. (1998). Penalized regression: The bridge versus the LASSO. Journal of Computational and Graphical Statistics, 7:397–416.

- Guo, J., Tang, M., Tian, M., & Zhu, K. (2013). Variable selection in high-dimensional partially linear additive models for composite quantile regression. *Computational Statistics and Data Analysis*, 65:56– 67.
- He, X., Zhue, Z. Y., & Fung, W. K. (2003). Estimation in a semiparametric model for longitudinal data with unspecified dependence structure. *Biometrika*, 89:579–590.
- Hu, Y., Zhao, K., & Lian, H. (2015). Bayesian quantile regression for partially linear additive models. Statistics and Computing, 25:651–668.
- Jeong, S., Park, T., & van Dyk, D. A. (2022). Bayesian model selection in additive partial linear models via locally adaptive splines. *Journal of Computational and Graphical Statistics*, 31:324–336.
- Kazemi, M., Shahsavani, D., & Arashi, M. (2018). Variable selection and structure identification for ultrahigh-dimensional partially linear additive models with application to cardiomyopathy microarray data. Statistics, Optimization & Information Computing, 6:373–382.
- Koenker, R. (2011). Additive models for quantile regression: model selection and confidence bands. Brazilian Journal of Probability and Statistics, 25:239–262.
- Lian, H. (2012). Variable selection in high-dimensional partly linear additive models. Journal of Nonparametric Statistics, 24:825–839.
- Liu, X., Wang, L., & Wang, H. (2011). Estimation and variable selection for semiparametric additive partial linear models. *Statistica Sinica*, 21:1225–1248.
- Lv, J., Yang, H., & Guo, C. (2017). Variable selection in partially linear additive models for modal regression. Communications in Statistics - Simulation and Computation, 46:5646–5665.
- Maronna, R. (2011). Robust ridge regression for high-dimensional data. Technometrics, 53:44-53.
- Maronna, R., Martin, D., Yohai, V., & Salibián-Barrera, M. (2019). Robust Statistics: Theory and Methods (with R). John Wiley and Sons.
- Nierenberg, D., Stukel, T., Baron, J., Dain, B., & Greenberg, E. (1989). Determinants of plasma levels of beta-carotene and retinol. American Journal of Epidemiology, 130:511–521.
- Pötscher, B. M. & Leeb, H. (2009). On the distribution of penalized maximum likelihood estimators: The lasso, scad, and thresholding. *Journal of Multivariate Analysis*, 100:2065–2082.
- Sherwood, B. & Wang, L. (2016). Partially linear additive quantile regression in ultra-high dimension. Annals of Statistics, 44:288–317.
- Smucler, E. & Yohai, V. (2017). Robust and sparse estimators for linear regression models. Computational Statistics and Data Analysis, 111:116–130.
- Stone, C. (1985). Additive regression and other nonparametric models. Annals of Statistics, 13:689–705.
- Sun, Y. & Genton, M. G. (2011). Functional boxplots. Journal of Computational and Graphical Statistics, 20:316–334.
- Tibshirani, R. J. (1996). Regression shrinkage and selection via the LASSO. Journal of the Royal Statistical Society, Series B, 58:267–288.
- Tibshirani, R. J. (1997). The LASSO method for variable selection in cox model. *Statistics in Medicine*, 16:385–395.

- Yuan, M. & Lin, Y. (2006). Model selection and estimation in regression with grouped variables. *Journal* of the Royal Statistical Society: Series B (Statistical Methodology), 68:49–67.
- Zhang, C. H. (2010). Nearly unbiased variable selection under minimax concave penalty. Annals of Statistics, 38:894–942.
- Zhao, Y., Zhang, R., Liu, J., & Lv, Y. (2014). Robust and efficient variable selection for semiparametric partially linear varying coefficient model based on modal regression. Annals of the Institute of Statistical Mathematics, 66:165–191.
- Zou, H. (2006). The adaptive Lasso and its oracle properties. Journal of the American Statistical Association, 101:1418–1429.
- Zou, H. & Hastie, T. (2005). Regularization and variable selection via the elastic net. Journal of the Royal Statistical Society: Series B, 67:301–320.