

# Measurement-Device-Independent Certification of Schmidt Number

Saheli Mukherjee,<sup>1,\*</sup> Bivas Mallick,<sup>1,†</sup> Arun Kumar Das,<sup>1,‡</sup> Amit Kundu,<sup>1,§</sup> and Pratik Ghosal<sup>1,¶</sup>

<sup>1</sup>*S. N. Bose National Centre for Basic Sciences, Block JD, Sector III, Salt Lake, Kolkata 700 106, India*

Bipartite quantum states with higher Schmidt numbers have been shown to outperform those with lower Schmidt numbers in various information processing tasks. Therefore, to ensure the efficient use of resources in these tasks, it is essential to certify the Schmidt number of the resource states. Ideally, this certification should rely as little as possible on the certifying devices, ensuring robustness against potential imperfections. In this work, we explore the scope of fully and partially device-independent Schmidt number certification methods. We demonstrate the general impossibility of fully device-independent certification for all states. Specifically, in a restricted setting, we present a class of states with Schmidt number 3, for which it is impossible to certify that their Schmidt number is greater than 2. However, we show that the Schmidt number of all states can be certified in a measurement-device-independent manner via semi-quantum nonlocal games, which assume trust in the preparation devices. Finally, we present an explicit construction of a semi-quantum game for the measurement-device-independent certification of a class of states.

## I. INTRODUCTION

Entanglement [1] is a key resource in various quantum information processing tasks, such as quantum computation [2], quantum teleportation [3], superdense coding [4], secret sharing [5], quantum key distribution [6], and enhancing communication capacity [7], among others. In some of these tasks, entanglement offers a significant advantage over classical methods, while in others, it enables phenomena that have no classical counterpart. The degree of entanglement in a quantum system is therefore crucial when comparing the resources needed for a specific task.

Even in the simplest case of two subsystems, there are multiple independent and inequivalent ways to measure entanglement [8]. Each measure quantifies a different aspect of entanglement, making it important to choose the appropriate one for a given task. One particularly interesting way to quantify the degree of entanglement is by counting the number of degrees of freedom that actively contribute to the entanglement—the *dimension of entanglement*. For bipartite systems, this can be measured using the *Schmidt number of the density operator* representing the system's state [9]. Advantage of higher Schmidt number states has been established in various tasks, including channel discrimination [10], quantum communication [11], universal quantum computation [12], quantum control [13], experimental optics [14, 15], enhancing key rates and security in quantum key distribution [16–19], and many more. This highlights the practical importance of certifying and characterizing the

Schmidt number of a given state for the advancement of quantum technologies.

Consider a scenario where a task requires a minimum entanglement dimension, *i.e.*, entangled states with a Schmidt number greater than a specified threshold, say  $r$ . To determine whether a given state can be used as a resource for the task, one must ensure that the Schmidt number of the state exceeds  $r$ . Several methods have been proposed in the literature to achieve this, each exploiting different properties of quantum states [20–26]. Perhaps the simplest approach to separating states with a Schmidt number greater than  $r$  from those with a Schmidt number less than or equal to  $r$  is through the use of Schmidt number witnesses [27–29]. States with a Schmidt number less than or equal to  $r$  form a closed, convex, and compact set. For any state outside this set, there exists at least one Hermitian operator—called the Schmidt number witness operator—whose expectation value with respect to the state is strictly negative, while for all states within the set, the expectation value is positive semi-definite. However, in practice, this approach is not infallible due to imperfections in the implementation of measurement devices used to experimentally evaluate the expectation value of the witness operator. In such cases, it is possible to incorrectly certify a state's Schmidt number as greater than  $r$ , even when it is not. This prompts the search for certification methods that are fully or at least partially device-independent.

In this paper, we explore such methods. A fully device-independent (DI) approach relies solely on the correlation statistics obtained from local measurements on shared entangled states to certify the Schmidt number. Specifically, in this setup, a referee provides two classical inputs,  $x$  and  $y$ , selected from two independent finite index sets  $\mathcal{X}$  and  $\mathcal{Y}$ , to two spatially separated non-communicating experimenters (or parties) who share the state whose Schmidt number is to be certified. Each party then locally applies a measurement device based

\* [mukherjeesaheli95@gmail.com](mailto:mukherjeesaheli95@gmail.com)

† [bivasqic@gmail.com](mailto:bivasqic@gmail.com)

‡ [akd.qip@gmail.com](mailto:akd.qip@gmail.com)

§ [amit8967@gmail.com](mailto:amit8967@gmail.com)

¶ [ghoshal.pratik00@gmail.com](mailto:ghoshal.pratik00@gmail.com)

on its respective input, yielding classical outputs  $a$  and  $b$ , which are then sent back to the referee. This process is repeated multiple times to generate the correlation statistics  $p(a, b|x, y)$ . The referee assigns a payoff  $\mathcal{J}(a, b, x, y)$  based on the inputs  $(x, y)$  and outputs  $(a, b)$ , and calculates the average payoff. The parties can select a strategy (local measurement settings) to optimize this average payoff. This nonlocal game is said to certify the Schmidt number of the state if it is possible to find an appropriate payoff function  $\mathcal{J}(a, b, x, y)$ , such that the optimal average payoff is negative *only if* the Schmidt number of the shared state exceeds  $r$ , and positive semi-definite otherwise. Note that this method is independent of the specific measurements implemented, complete knowledge of the shared state, or any other devices used in the experiment. However, as we will discuss, DI certification method of the Schmidt number is *not faithful*, in the sense that, in general, it cannot certify the Schmidt number of all states (see also [30]). One simple argument for this is that DI certification of entanglement [31] for all states is not possible, due to the existence of entangled states that admit local- hidden-variable (LHV) models [32, 33]. Entanglement certification can be viewed as certifying whether a given bipartite state has Schmidt number greater than 1, which constitutes a special case of the more general Schmidt number certification. Still, it is reasonable to question whether DI certification of all states with Schmidt number greater than  $r$  is possible for  $r \neq 1$ . We provide an explicit example of a class of states with Schmidt number 3, where certifying that they have Schmidt number greater than 2 in a DI manner is not possible when the parties are restricted to only local projective measurements.

The faithfulness of Schmidt number certification can be restored by relaxing the demand for full device-independence and opting for partial device-independence. We propose a measurement-device-independent (MDI) method for certifying the Schmidt number of all bipartite states utilizing the framework of semi-quantum nonlocal games [34]. Semi-quantum games generalize the earlier Bell-nonlocal game setting by replacing the classical inputs  $(x, y)$  with classically indexed quantum inputs  $(\psi_x, \phi_y)$ . The parties are aware of the input ensembles  $\{\psi_x\}$  and  $\{\phi_y\}$ , but crucially, they are unaware of which particular states have been provided in a given run of the game. In this setting, the optimal average payoff generated by the correlation statistics  $p(a, b|\psi_x, \phi_y)$  certifies the Schmidt number by satisfying the same condition as before: it is negative *only if* the Schmidt number of the shared state exceeds  $r$ , and positive semi-definite otherwise. It is important to note that, while this method is independent of the specific measurement devices used in the experiment, it critically depends on the perfect preparation of the input states. Trusting the preparation devices is a more nat-

ural assumption than trusting the measurement devices, since the detection devices, in general, are open to the external environment. We demonstrate the feasibility of MDI certification of the Schmidt number for all states by explicitly constructing certification methods from standard Schmidt number witnesses. We also present an example illustrating the construction.

The framework of semi-quantum nonlocal games has previously been used for MDI certification [35] and characterization of entanglement [36–41], as well as for MDI certification of beyond-quantum states [42, 43]. This framework has also been invoked in the context of steering [44], non-classical teleportation [45], and classical simulation of quantum correlations [46, 47]. This MDI approach to certification has also been tested experimentally [48–50]. However, these works do not address the specific question of certifying whether a given state has a Schmidt number greater than  $r$ , which is the focus of this study. General MDI entanglement witnesses [35] certify whether a state is entangled, but they do not capture the distinct layers of entanglement present in different entangled states, as quantified by their Schmidt numbers.

The rest of the paper is organized as follows. In Sec. II, we formally introduce several preliminary concepts, such as the Schmidt number of bipartite states and the framework of semi-quantum nonlocal games. In Sec. III A, we elaborate on why fully device-independent Schmidt number certification is not possible for all states. Next, in Sec. III B, we demonstrate how the Schmidt number of all states can be certified through semi-quantum nonlocal games in a measurement-device-independent manner, with an explicit example illustrating the construction of a semi-quantum game that witnesses the Schmidt number of a class of states. Finally, we conclude with some discussions in Sec. IV.

## II. PRELIMINARIES

Throughout the paper, we will be considering finite-dimensional quantum systems, each of which is associated with a complex Hilbert space  $\mathcal{H} \cong \mathbb{C}^d$  where  $d = \dim(\mathcal{H}) < \infty$ . The set of all bounded linear operators on the Hilbert space is denoted by  $\mathcal{B}(\mathcal{H})$ . Pure states of the system are represented by vectors in  $\mathcal{H}$ , while mixed states are described by density operators, which are positive semi-definite operators with unit trace. We denote the set of all such density operators by  $\mathcal{D}(\mathcal{H}) \subset \mathcal{B}(\mathcal{H})$ .

### A. Schmidt number of density operators and its witness

Consider a bipartite quantum system consisting of two subsystems,  $A$  and  $B$ , with a composite Hilbert space




Figure 1: Illustration of the nested subset structure of sets of states with Schmidt number less than or equal to  $r$  and Schmidt number witnesses.

given by  $\mathcal{H}_A \otimes \mathcal{H}_B \cong \mathbb{C}^{d_A} \otimes \mathbb{C}^{d_B}$ . Any pure state  $|\psi\rangle_{AB}$  of the system can be always represented in Schmidt decomposition form [51, 52] as:

$$|\psi\rangle_{AB} = \sum_{i=1}^k \sqrt{\lambda_i} |u_i\rangle_A |v_i\rangle_B, \quad (1)$$

where the  $\sqrt{\lambda_i}$ 's are called Schmidt coefficients, with  $\lambda_i \geq 0$  and  $\sum_i \lambda_i = 1$ , and  $\{|u_i\rangle_A\}$  and  $\{|v_i\rangle_B\}$  are orthonormal bases for  $\mathbb{C}^{d_A}$  and  $\mathbb{C}^{d_B}$ , respectively. Here,  $1 \leq k \leq \min\{d_A, d_B\}$  represents the Schmidt rank (SR) of the state  $|\psi\rangle_{AB}$ , which indicates the number of non-vanishing Schmidt coefficients in the decomposition, *i.e.*,  $\text{SR}(|\psi\rangle_{AB}) = k$ . A bipartite state is product if and only if its  $\text{SR} = 1$ , and is entangled otherwise.

The Schmidt number (SN) is a generalization of the Schmidt rank (SR) for mixed states [9]. A mixed state  $\sigma_{AB} \in \mathcal{D}(\mathbb{C}^{d_A} \otimes \mathbb{C}^{d_B})$  admits a non-unique decomposition into pure states as  $\sigma_{AB} = \sum_k p_k |\psi_k\rangle_{AB} \langle \psi_k|$ , with  $|\psi_k\rangle_{AB} \in \mathbb{C}^{d_A} \otimes \mathbb{C}^{d_B}$ ,  $p_k \geq 0$  and  $\sum_k p_k = 1$ . The Schmidt number of  $\sigma_{AB}$  is defined as

$$\text{SN}(\sigma_{AB}) = \min_{\{p_k, |\psi_k\rangle_{AB}\}} \left[ \max_k \text{SR}(|\psi_k\rangle_{AB}) \right] \quad (2)$$

where the minimization is taken over all possible pure state decomposition of  $\sigma_{AB}$ . It follows that for any state  $\sigma_{AB}$ ,  $1 \leq \text{SN}(\sigma_{AB}) \leq \min\{d_A, d_B\}$ , and for separable states  $\sigma_{\text{sep}}$ ,  $\text{SN}(\sigma_{\text{sep}}) = 1$ .

Let us denote the set of all states in  $\mathcal{D}(\mathbb{C}^{d_A} \otimes \mathbb{C}^{d_B})$  with Schmidt number  $\leq r$  by  $\mathcal{S}_r$ . The set  $\mathcal{S}_r$  forms a closed, convex, and compact subset within the real vector space of all density operators, with the following subset relation:

$$\mathcal{S}_1 \subset \mathcal{S}_2 \subset \dots \subset \mathcal{S}_r.$$

Therefore, by the Hahn-Banach separation theorem [53], there always exists a real functional on this space, which

acts as a separating hyperplane, distinguishing states with Schmidt number greater than  $r$  from the set  $\mathcal{S}_r$  [27–29] (See Fig. 1). Specifically, for all states  $\rho_{AB}$  with  $\text{SN}(\rho_{AB}) > r$ , there exists at least one Hermitian operator  $\mathcal{W}$ , called a *Schmidt number witness*, such that

$$\begin{aligned} \text{Tr}[\mathcal{W}\rho_{AB}] &< 0, \\ \text{and } \text{Tr}[\mathcal{W}\sigma_{AB}] &\geq 0, \quad \forall \sigma_{AB} \in \mathcal{S}_r. \end{aligned} \quad (3)$$

## B. Semi-quantum nonlocal games

Semi-quantum nonlocal games [34] are a generalization of Bell-nonlocal games. The game involves two parties, say Alice and Bob. A referee chooses two classical indices,  $x$  and  $y$ , from two independent index sets  $\mathcal{X}$  and  $\mathcal{Y}$ , with respective probabilities  $p(x)$  and  $q(y)$ . Unlike in the Bell scenario, where the referee sends these classical variables directly to Alice and Bob, in the semi-quantum setting, the referee prepares two quantum systems in the states  $\psi_{A_0}^x \in \mathcal{D}(\mathcal{H}_{A_0})$  and  $\phi_{B_0}^y \in \mathcal{D}(\mathcal{H}_{B_0})$  and sends them to Alice and Bob, respectively. It is important to note that the states  $\psi_{A_0}^x$  (and similarly, the states  $\phi_{B_0}^y$ ) are non-orthogonal; otherwise, it would reduce to the Bell scenario. Consequently, the identities of the states sent in a given run of the game are known only to the referee, and not to Alice or Bob. Before the game starts, Alice and Bob can fix a strategy, like sharing classical shared randomness or, a bipartite state  $\rho_{AB} \in \mathcal{D}(\mathcal{H}_A \otimes \mathcal{H}_B)$ .

Upon receiving their respective states from the referee, each of the parties locally perform a generalized joint measurement on their received system and their respective subsystem of the pre-shared system. Mathematically, the generalized measurements are denoted by positive operator-valued measures (POVMs)  $\mathcal{E}_{AA_0} = \{\mathcal{E}_{AA_0}^a | \mathcal{E}_{AA_0}^a \in \mathbb{P}(\mathcal{H}_A \otimes \mathcal{H}_{A_0}) \forall a \text{ \& } \sum_a \mathcal{E}_{AA_0}^a = \mathbb{I}_{AA_0}\}$  and  $\mathcal{F}_{BB_0} = \{\mathcal{F}_{BB_0}^b | \mathcal{F}_{BB_0}^b \in \mathbb{P}(\mathcal{H}_B \otimes \mathcal{H}_{B_0}) \forall b \text{ \& } \sum_b \mathcal{F}_{BB_0}^b = \mathbb{I}_{BB_0}\}$ , with  $\mathbb{P}(\mathcal{H})$  denoting the set of positive operators on  $\mathcal{H}$ , and  $a$  and  $b$  denoting the measurement outcomes respectively. Repeating this multiple times generates a correlation statistics given by

$$p(a, b | \psi^x, \phi^y) = \text{Tr}[(\mathcal{E}_{AA_0}^a \otimes \mathcal{F}_{BB_0}^b)(\psi_{A_0}^x \otimes \rho_{AB} \otimes \phi_{B_0}^y)]. \quad (4)$$

Based on the inputs and outputs, the referee assigns a payoff function  $\mathcal{J}(a, b, x, y)$ . The goal of Alice and Bob is to optimize their average payoff, given by

$$\mathcal{J}_{\text{avg}}(\rho_{AB}) = \max_{a, b, x, y} \sum p(x)q(y) \mathcal{J}(a, b, x, y) p(a, b | \psi^x, \phi^y), \quad (5)$$

where the maximization is over all the POVM measurements performed by Alice and Bob, appearing in Eqn. (4).

In [34], it was shown that all entangled states can generate correlations  $p(a, b | \psi^x, \phi^y)$ , with a suitable choice of Alice and Bob's measurements, that cannot be simulated by classical shared randomness or separable states, thereby demonstrating the "nonlocality" of all entangled states, in contrast to Bell-nonlocal games. Later, in [35], it was shown that this semi-quantum game can be used to certify the entanglement of all entangled states in an MDI manner. In the following section, we will demonstrate that the semi-quantum game can not only be used to certify whether a given state is entangled, but also to certify whether the Schmidt number of the state is greater than  $r$ .

### III. CERTIFICATION OF SCHMIDT NUMBER

#### A. Scope of fully Device-Independent certification

Fully device-independent certification of entanglement for all entangled states is, in general, not possible. This is because there exists bipartite entangled states that admit a local-hidden-variable (LHV) model. In a Bell-nonlocal game, any correlation  $p(a, b | x, y)$  generated by these states can always be simulated by separable states, making it impossible to distinguish these states from separable ones based solely on the correlation.

Motivated by the above consideration, we present two explicit examples demonstrating that fully device-independent certification of Schmidt number is, in general, not possible for all states.

**Example 1:** Consider the Werner-class of states [32],

$$\rho_{AB}^W(p) = p|\phi^+\rangle_{AB}\langle\phi^+| + (1-p)\frac{\mathbb{I}_A}{2} \otimes \frac{\mathbb{I}_B}{2}, \quad (6)$$

where  $|\phi^+\rangle_{AB} = \frac{1}{\sqrt{2}}(|00\rangle_{AB} + |11\rangle_{AB})$  and  $p \in [0, 1]$ . This class of states is entangled for  $\frac{1}{3} < p \leq 1$ , and hence  $\rho_{AB}^W(p > \frac{1}{3}) \notin \mathcal{S}_1$ . However, for  $p \leq \frac{1}{2}$ ,  $\rho_{AB}^W(p)$  does not violate any Bell inequality for any choice of generalized POVM measurements of Alice and Bob [54, 55]. Therefore, for the class of states  $\rho_{AB}^W(p)$  with  $\frac{1}{3} < p \leq \frac{1}{2}$ , the condition  $\text{SN}(\rho_{AB}^W(p)) > 1$  cannot be certified in a fully device-independent (DI) manner.

Nevertheless, one may still question the possibility of DI certification of whether a given state  $\rho_{AB} \notin \mathcal{S}_r$ , for  $r \neq 1$  and for all  $\rho_{AB}$ . Our next example provides a negative answer under a restricted scenario. We show the existence of a class of states in  $\mathcal{D}(\mathbb{C}^8 \otimes \mathbb{C}^8)$  with Schmidt number 3, for which it is not possible to certify that their Schmidt number is greater than 2 in a DI manner, if Alice and Bob are restricted to implementing only local projective measurements.

**Example 2:** Consider the isotropic class of states  $\rho_{AB}^{\text{iso}}(p) \in \mathcal{D}(\mathbb{C}^d \otimes \mathbb{C}^d)$

$$\rho_{AB}^{\text{iso}}(p) = p|\Phi^+\rangle_{AB}\langle\Phi^+| + (1-p)\frac{\mathbb{I}_A}{d} \otimes \frac{\mathbb{I}_B}{d} \quad (7)$$

where  $|\Phi^+\rangle_{AB} = \frac{1}{\sqrt{d}}\sum_{i=0}^{d-1}|ii\rangle_{AB}$  and  $p \in [0, 1]$ . This class of states have the following properties:

1.  $\text{SN}(\rho_{AB}^{\text{iso}}(p)) \leq r$  if and only if  $0 \leq p \leq \frac{rd-1}{d^2-1}$ , [9, 56]
2.  $\rho_{AB}^{\text{iso}}(p)$  is entangled for  $\frac{1}{d+1} < p \leq 1$ ,
3.  $\rho_{AB}^{\text{iso}}(p)$  admits a LHV model for  $p \leq \frac{\sum_{k=2}^d k}{d-1}$  under local projective measurements [57].

Using properties 1 and 3, it can be verified that an admissible range of  $p$  exists such that states in this range have a Schmidt number greater than 2 as well as admit an LHV model, only if  $d \geq 8$ . For simplicity, we focus on the case of  $d = 8$ . In that case, for  $0.238 < p \leq 0.245$ ,  $\text{SN}(\rho_{AB}^{\text{iso}}(p)) = 3$ , and  $\rho_{AB}^{\text{iso}}(p)$  admits an LHV model under local projective measurements.

Now, consider another class of states in  $\mathcal{D}(\mathbb{C}^8 \otimes \mathbb{C}^8)$  given by

$$\tilde{\rho}_{AB}(\lambda) = \lambda \rho_{AB}^{\text{iso}}(p = 0.24) + (1-\lambda)|\phi_2^+\rangle_{AB}\langle\phi_2^+|, \quad (8)$$

where  $\rho_{AB}^{\text{iso}}(p = 0.24) \in \mathcal{D}(\mathbb{C}^8 \otimes \mathbb{C}^8)$ ,  $|\phi_2^+\rangle_{AB} = \frac{1}{\sqrt{2}}(|00\rangle_{AB} + |11\rangle_{AB})$ , and  $\lambda \in [0, 1]$ .

**Lemma 1.**  $\text{SN}(\tilde{\rho}_{AB}(\lambda)) = 3$  for  $\lambda \in (0, 1]$ .

*Proof.* Since  $\text{SR}(|\phi_2^+\rangle_{AB}) = 2$  and  $\text{SN}(\rho_{AB}^{\text{iso}}(p = 0.24)) = 3$ , it follows from the convexity of the set  $\mathcal{S}_3$  that  $\tilde{\rho}_{AB}(\lambda) \in \mathcal{S}_3$ . Now, consider the Schmidt number witness operator [29]

$$\mathcal{W}_r = \mathbb{I}_8 \otimes \mathbb{I}_8 - \frac{8}{r} \mathcal{P}, \quad (9)$$

where  $\mathcal{P}$  is the projector onto the maximally entangled state  $|\Phi^+\rangle = \frac{1}{\sqrt{8}}\sum_{i=1}^8|ii\rangle$ . The operator  $\mathcal{W}_r$  serves as a witness for whether a state has a Schmidt number greater than  $r$ . For  $\tilde{\rho}_{AB}(\lambda)$ , we get  $\text{Tr}[\mathcal{W}_2 \tilde{\rho}_{AB}] < 0$  for  $\lambda \in (0, 1]$ , which conclusively implies that  $\tilde{\rho}_{AB} \notin \mathcal{S}_2$  in this range. Therefore, we conclude that  $\text{SN}(\tilde{\rho}_{AB}) = 3$  for  $\lambda \in (0, 1]$ . ■

**Lemma 2.**  $\tilde{\rho}_{AB}(\lambda)$  is nonlocal under local projective measurements for  $\lambda \in [0, 0.312]$ .

*Proof.* Let Alice and Bob share the bipartite state  $\tilde{\rho}_{AB}(\lambda)$ . The goal is to find the range of  $\lambda$  such that the state exhibits nonlocal statistics under suitable choices of incompatible measurements on Alice's and Bob's sides. Let the inputs to Alice and Bob be represented by  $x$

and  $y$  respectively, where  $x, y \in \{0, 1\}$ . The respective outcomes for Alice and Bob are denoted by  $a$  and  $b$ , where  $a, b \in \{0, 1, 2\}$ . For  $x = 0$ , Alice performs the three-outcome projective measurement

$$M_0 = \{|0\rangle\langle 0|, |1\rangle\langle 1|, \mathbb{1} - |0\rangle\langle 0| - |1\rangle\langle 1|\}$$

where  $|0\rangle = (1\ 0\ 0\ 0\ 0\ 0\ 0\ 0)^T$  and  $|1\rangle = (0\ 1\ 0\ 0\ 0\ 0\ 0\ 0)^T$ . For  $x = 1$ , Alice performs the projective measurement

$$M_1 = \{|+\rangle\langle +|, |-\rangle\langle -|, \mathbb{1} - |+\rangle\langle +| - |-\rangle\langle -|\}$$

where  $|+\rangle = \left(\frac{1}{\sqrt{2}}\ \frac{1}{\sqrt{2}}\ 0\ 0\ 0\ 0\ 0\ 0\right)^T$  and  $|-\rangle = \left(\frac{1}{\sqrt{2}}\ \frac{-1}{\sqrt{2}}\ 0\ 0\ 0\ 0\ 0\ 0\right)^T$ . The corresponding measurements for Bob are  $N_0 = \{|\alpha_1\rangle\langle \alpha_1|, |\alpha_2\rangle\langle \alpha_2|, \mathbb{1} - |\alpha_1\rangle\langle \alpha_1| - |\alpha_2\rangle\langle \alpha_2|\}$  where  $|\alpha_1\rangle = \frac{1}{\sqrt{4+2\sqrt{2}}}\left((1+\sqrt{2})\ 1\ 0\ 0\ 0\ 0\ 0\ 0\right)^T$ ,  $|\alpha_2\rangle = \frac{1}{\sqrt{4-2\sqrt{2}}}\left((1-\sqrt{2})\ 1\ 0\ 0\ 0\ 0\ 0\ 0\right)^T$  and  $N_1 = \{|\beta_1\rangle\langle \beta_1|, |\beta_2\rangle\langle \beta_2|, \mathbb{1} - |\beta_1\rangle\langle \beta_1| - |\beta_2\rangle\langle \beta_2|\}$  where  $|\beta_1\rangle = \frac{1}{\sqrt{4+2\sqrt{2}}}\left((-1-\sqrt{2})\ 1\ 0\ 0\ 0\ 0\ 0\ 0\right)^T$ ,  $|\beta_2\rangle = \frac{1}{\sqrt{4-2\sqrt{2}}}\left((\sqrt{2}-1)\ 1\ 0\ 0\ 0\ 0\ 0\ 0\right)^T$ . It is known that corresponding to every Bell inequality, there exists a nonlocal game and vice-versa [58]. Let Alice and Bob play a game under these measurement settings and the state  $\tilde{\rho}_{AB}(\lambda)$ , with a payoff given by

$$\mathcal{J}(\rho) = \begin{cases} +1 & \text{for } a \oplus b = xy \text{ and } a, b \in \{0, 1\} \\ -1 & \text{for } a \oplus b = \bar{x}\bar{y} \text{ and } a, b \in \{0, 1\} \\ 0 & \text{for } a, b \notin \{0, 1\} \end{cases} \quad (10)$$

It can be clearly seen that the average payoff of this game reduces to the Clauser-Horne-Shimony-Holt (CHSH) inequality [59]. This inequality surpasses the local bound of 2 for  $\lambda \in [0, 0.312]$  indicating that the state  $\tilde{\rho}_{AB}(\lambda)$  is CHSH-nonlocal for  $\lambda \in [0, 0.312]$ . ■

Now, we state the following proposition.

**Proposition 1.** *For the class of states  $\tilde{\rho}_{AB}(\lambda)$  in Eqn. (8), DI certification of  $\tilde{\rho}_{AB}(\lambda) \notin \mathcal{S}_2$  is not possible in the range  $\lambda \in (0, 0.312)$ , if the parties are allowed to implement only local projective measurements.*

*Proof.* It follows from Lemma 1 and Lemma 2 that in the range  $\lambda \in (0, 0.312)$ ,  $\tilde{\rho}_{AB}(\lambda)$  is Bell-nonlocal and has Schmidt number  $\text{SN}(\tilde{\rho}_{AB}(\lambda)) = 3$ . Since  $\rho_{AB}^{\text{iso}}(p = 0.24)$  is local under projective measurements, any nonlocal contribution to the correlation of  $\tilde{\rho}_{AB}(\lambda)$  stems exclusively from the  $|\phi_2^+\rangle_{AB}\langle \phi_2^+|$  part. Therefore, any correlation generated by the state  $\tilde{\rho}_{AB}(\lambda)$  can be simulated by a Schmidt number 2 state:

$$\rho_{AB}^* = \lambda \rho_{AB}^{\text{sep}} + (1 - \lambda) |\phi_2^+\rangle_{AB}\langle \phi_2^+|,$$

where  $\rho_{AB}^{\text{sep}}$  is a separable state. This establishes that DI certification of  $\tilde{\rho}_{AB}(\lambda) \notin \mathcal{S}_2$  is not possible. ■

However, in the next subsection, we will demonstrate that MDI certification of Schmidt number is possible for all states.

## B. Measurement-Device-Independent certification

We propose our main theorem, which provides a prescription for constructing semi-quantum games that certify the Schmidt number of all states.

**Theorem 1.** *For every bipartite quantum state  $\rho_{AB} \in \mathcal{D}(\mathcal{H}_A \otimes \mathcal{H}_B)$  with  $\text{SN}(\rho_{AB}) > r$ , there exists a semi-quantum game such that the average payoff obtained using  $\rho_{AB}$ ,  $\mathcal{J}_{\text{avg}}(\rho_{AB}) < 0$ , whereas  $\mathcal{J}_{\text{avg}}(\sigma_{AB}) \geq 0$ ,  $\forall \sigma_{AB} \in \mathcal{S}_r$ .*

*Proof.* Consider that a referee provides Alice and Bob a state  $\rho_{AB} \in \mathcal{D}(\mathcal{H}_A \otimes \mathcal{H}_B)$  and wants to convince them that  $\rho_{AB}$  has a Schmidt number greater than  $r$ , i.e.,  $\rho_{AB} \notin \mathcal{S}_r$ .

To achieve this, referee, who knows the identity of  $\rho_{AB}$ , selects a witness operator  $\mathcal{W}_{AB}$  such that Eqn. (3) holds. Since the space of all Hermitian operators is spanned by the set of density operators, any witness operator, being Hermitian, can be written as

$$\mathcal{W}_{AB} = \sum_{x,y} \gamma_{x,y} (\psi_A^x)^\top \otimes (\phi_B^y)^\top, \quad (11)$$

where  $\gamma_{x,y}$ 's are real  $\forall x, y$ ,  $\{\psi^x\}$  and  $\{\phi^y\}$  form bases of the linear operator spaces  $\mathcal{B}(\mathcal{H}_A)$  and  $\mathcal{B}(\mathcal{H}_B)$ , respectively, and  $(\cdot)^\top$  denotes the transpose with respect to the computational basis of  $\mathcal{H}_A$  and  $\mathcal{H}_B$ . Note that this decomposition is, in general, non-unique.

Now, the referee plays a semi-quantum game with Alice and Bob, providing the quantum inputs from the ensembles  $\{\psi_{A_0}^x\}$  and  $\{\phi_{B_0}^y\}$ , appearing in Eqn. (11) with  $\mathcal{H}_{A_0} \cong \mathcal{H}_A$  and  $\mathcal{H}_{B_0} \cong \mathcal{H}_B$ , and  $a, b \in \{1, 2\}$ . Additionally, the referee ensures that the game parameters  $p(x)$ ,  $q(y)$ , and  $\mathcal{J}(a, b, x, y)$  satisfies the following conditions:

$$p(x)q(y)\mathcal{J}(a, b, x, y) = \gamma_{a,b,x,y} = \begin{cases} \gamma_{x,y} & \text{for } (a, b) = (1, 1) \\ 0 & \text{otherwise} \end{cases} \quad (12)$$

For certification, we need to show that:

- regardless of the measurement settings Alice and Bob choose, the average payoff of this game,  $\mathcal{J}_{\text{avg}}(\rho_{AB})$ , satisfies  $\mathcal{J}_{\text{avg}}(\rho_{AB}) \geq 0$  if  $\rho_{AB} \in \mathcal{S}_r$ , and
- if  $\rho_{AB} \notin \mathcal{S}_r$ , there exists a measurement setting such that  $\mathcal{J}_{\text{avg}}(\rho_{AB}) < 0$ .

The average payoff of this game is given by

$$\mathcal{J}_{\text{avg}}(\rho_{AB}) = \sum_{x,y} \gamma_{x,y} \text{Tr} \left[ (\mathcal{E}_{A_0A}^1 \otimes \mathcal{F}_{B_0B}^1) (\psi_{A_0}^x \otimes \rho_{AB} \otimes \phi_{B_0}^y) \right] \quad (13)$$

**Proof of the 1<sup>st</sup> condition:** Let us consider that  $\rho_{AB} \in \mathcal{S}_r$ . This implies that there exists at least one decomposition of  $\rho_{AB}$  in terms of pure states  $\rho_{AB} = \sum_k p_k |\eta_k\rangle_{AB} \langle \eta_k|$  such that  $\text{SR}(|\eta_k\rangle_{AB}) \leq r \ \forall k$ . Eqn. (13) can be re-expressed as

$$\mathcal{J}_{\text{avg}}(\rho_{AB}) = \sum_k p_k \sum_{x,y} \gamma_{x,y} \text{Tr}_{A_0B_0} [R_{A_0B_0}^k (\psi_{A_0}^x \otimes \phi_{B_0}^y)], \quad (14)$$

where

$$R_{A_0B_0}^k = \text{Tr}_{AB} [(\mathcal{E}_{A_0A}^1 \otimes \mathcal{F}_{B_0B}^1) (|\eta_k\rangle_{AB} \langle \eta_k| \otimes \mathbb{I}_{A_0B_0})]. \quad (15)$$

Note that  $R_{A_0B_0}^k$  is a positive operator that is proportional to the state obtained by locally implementing the measurements  $\{\mathcal{E}_{A_0A}^a\}$  and  $\{\mathcal{F}_{B_0B}^b\}$  on the state  $(|\eta_k\rangle_{AB} \langle \eta_k| \otimes \frac{\mathbb{I}_{A_0B_0}}{d_{A_0}d_{B_0}})$ , and filtering the outcome  $a = 1$  and  $b = 1$ .

It follows from the definition of the Schmidt number that it is invariant under rescaling of the state, *i.e.*, multiplying the state by a scalar factor. Since  $\text{SR}(|\eta_k\rangle_{AB}) \leq r$ , it follows that  $\text{SN}(|\eta_k\rangle_{AB} \otimes \mathbb{I}) \leq r$  for all  $k$  [9]. Moreover, the Schmidt number cannot increase under local filtering, as it is a monotone under a larger set of operations, namely stochastic local operations and classical communication (SLOCC) [60–64]. Consequently, we have  $\text{SN}(R_{A_0B_0}^k) \leq r \ \forall k$ . Following the convexity of the set  $\mathcal{S}_r$ , we also get  $\sum_k p_k R_{A_0B_0}^k \in \mathcal{S}_r$ .

Using Eqn. (11) in Eqn. (14), the average payoff can be written as

$$\begin{aligned} \mathcal{J}_{\text{avg}}(\rho_{AB}) &= \text{Tr}_{A_0B_0} \left[ \sum_k p_k R_{A_0B_0}^k \mathcal{W}_{A_0B_0}^\top \right] \\ &= \sum_k p_k \text{Tr}_{A_0B_0} \left[ (R_{A_0B_0}^k)^\top \mathcal{W}_{A_0B_0} \right]. \end{aligned} \quad (16)$$

The last step follows from the linearity of the trace operation and the self-duality of the transpose map. Additionally,

$$\sum_k p_k R_{A_0B_0}^k \in \mathcal{S}_r \implies \sum_k p_k (R_{A_0B_0}^k)^\top \in \mathcal{S}_r.$$

Using the properties of the Schmidt number witness, we obtain

$$\mathcal{J}_{\text{avg}}(\rho_{AB}) \geq 0. \quad (17)$$

**Proof of the 2<sup>nd</sup> condition:** Now, let us consider that  $\rho_{AB} \notin \mathcal{S}_r$  and Alice performs a two-outcome joint measurement  $\{\mathcal{E}_{A_0A}^1 = \mathcal{P}_{AA_0}, \mathcal{E}_{A_0A}^2 = \mathbb{I} - \mathcal{P}_{AA_0}\}$  and Bob performs  $\{\mathcal{F}_{B_0B}^1 = \mathcal{P}_{BB_0}, \mathcal{F}_{B_0B}^2 = \mathbb{I} - \mathcal{P}_{BB_0}\}$  where  $\mathcal{P}_{XX_0}$  is a projector on the maximally entangled state  $|\phi^+\rangle_{XX_0} = \frac{1}{\sqrt{d_X}} \sum_{i=0}^{d_X-1} |ii\rangle$  for  $X = A, B$ .

The average payoff, then, turns out to be

$$\begin{aligned} \mathcal{J}_{\text{avg}}(\rho_{AB}) &= \sum_{x,y} \gamma_{x,y} \text{Tr} \left[ (\mathcal{P}_{AA_0} \otimes \mathcal{P}_{BB_0}) (\psi_{A_0}^x \otimes \rho_{AB} \otimes \phi_{B_0}^y) \right] \\ &= \sum_{x,y} \gamma_{x,y} \text{Tr}_{AB} [(M_A^x \otimes M_B^y) \rho_{AB}], \end{aligned} \quad (18)$$

where

$$\begin{aligned} M_A^x &= \text{Tr}_{A_0} \left[ \mathcal{P}_{AA_0} (\psi_{A_0}^x \otimes \mathbb{I}_A) \right] \implies M_A^x = \frac{1}{d_A} [\psi_A^x]^\top, \\ \& \ M_B^y &= \text{Tr}_{B_0} \left[ \mathcal{P}_{BB_0} (\phi_{B_0}^y \otimes \mathbb{I}_B) \right] \implies M_B^y = \frac{1}{d_B} [\phi_B^y]^\top. \end{aligned}$$

Using Eqn. (11), we get

$$\mathcal{J}_{\text{avg}}(\rho_{AB}) = \frac{1}{d_A d_B} \text{Tr}_{AB} [\mathcal{W}_{AB} \rho_{AB}] < 0. \quad (19)$$

This completes the proof.  $\blacksquare$

Now, let us present an explicit example in support of Theorem 1.

**Example:** We illustrate the construction of a semi-quantum game corresponding to the witness operator [29]

$$\mathcal{W}_2 = \mathbb{I}_3 \otimes \mathbb{I}_3 - \frac{3}{2} |\phi^+\rangle \langle \phi^+| \in \mathcal{B}(\mathbb{C}^3 \otimes \mathbb{C}^3), \quad (20)$$

where  $|\phi^+\rangle = \frac{1}{\sqrt{3}}(|00\rangle + |11\rangle + |22\rangle)$ . This operator witnesses the Schmidt number of the class of states  $\rho_{AB}^{\text{iso}}(p) \in \mathcal{D}(\mathbb{C}^3 \otimes \mathbb{C}^3)$  [Eqn. (7)] having a Schmidt number greater than 2, for  $p \in (\frac{5}{8}, 1]$ .

$\mathcal{W}_2$  admits a decomposition of the form

$$\mathcal{W}_2 = \sum_{xy} \gamma_{x,y} \zeta^x \otimes \zeta^y,$$

where



- [3] C. H. Bennett, G. Brassard, C. Crépeau, R. Jozsa, A. Peres, and W. K. Wootters, Teleporting an unknown quantum state via dual classical and einstein-podolsky-rosen channels, *Phys. Rev. Lett.* **70**, 1895 (1993).
- [4] C. H. Bennett and S. J. Wiesner, Communication via one- and two-particle operators on einstein-podolsky-rosen states, *Phys. Rev. Lett.* **69**, 2881 (1992).
- [5] R. Cleve, D. Gottesman, and H.-K. Lo, How to share a quantum secret, *Phys. Rev. Lett.* **83**, 648 (1999).
- [6] A. K. Ekert, Quantum cryptography based on bell's theorem, *Phys. Rev. Lett.* **67**, 661 (1991).
- [7] C. Bennett, P. Shor, J. Smolin, and A. Thapliyal, Entanglement-assisted capacity of a quantum channel and the reverse shannon theorem, *IEEE Transactions on Information Theory* **48**, 2637 (2002).
- [8] M. B. Plenio and S. S. Virmani, An Introduction to Entanglement Theory, *Quant. Inf. Comput.* **7**, 001 (2007), [arXiv:quant-ph/0504163](https://arxiv.org/abs/quant-ph/0504163).
- [9] B. M. Terhal and P. Horodecki, Schmidt number for density matrices, *Physical Review A* **61**, 040301 (2000).
- [10] J. Bae, D. Chruściński, and M. Piani, More entanglement implies higher performance in channel discrimination tasks, *Physical Review Letters* **122**, 140404 (2019).
- [11] D. Cozzolino, B. Da Lio, D. Bacco, and L. K. Oxenløwe, High-dimensional quantum communication: benefits, progress, and future challenges, *Advanced Quantum Technologies* **2**, 1900038 (2019).
- [12] Y. Wang, Z. Hu, B. C. Sanders, and S. Kais, Qudits and high-dimensional quantum computing, *Frontiers in Physics* **8**, 589504 (2020).
- [13] M. Kues, C. Reimer, P. Roztocky, L. R. Cortés, S. Sciara, B. Wetzl, Y. Zhang, A. Cino, S. T. Chu, B. E. Little, *et al.*, On-chip generation of high-dimensional entangled quantum states and their coherent control, *Nature* **546**, 622 (2017).
- [14] M. Erhard, R. Fickler, M. Krenn, and A. Zeilinger, Twisted photons: new quantum perspectives in high dimensions, *Light: Science & Applications* **7**, 17146 (2018).
- [15] M. Erhard, M. Krenn, and A. Zeilinger, Advances in high-dimensional quantum entanglement, *Nature Reviews Physics* **2**, 365 (2020).
- [16] N. J. Cerf, M. Bourennane, A. Karlsson, and N. Gisin, Security of quantum key distribution using d-level systems, *Physical review letters* **88**, 127902 (2002).
- [17] L. Sheridan and V. Scarani, Security proof for quantum key distribution using qudit systems, *Physical Review A—Atomic, Molecular, and Optical Physics* **82**, 030301 (2010).
- [18] D. Cozzolino, D. Bacco, B. Da Lio, K. Ingerslev, Y. Ding, K. Dalgaard, P. Kristensen, M. Galili, K. Rottwitt, S. Ramachandran, *et al.*, Orbital angular momentum states enabling fiber-based high-dimensional quantum communication, *Physical Review Applied* **11**, 064058 (2019).
- [19] F. Bouchard, K. Heshami, D. England, R. Fickler, R. W. Boyd, B.-G. Englert, L. L. Sánchez-Soto, and E. Karimi, Experimental investigation of high-dimensional quantum key distribution protocols with twisted photons, *Quantum* **2**, 111 (2018).
- [20] J. Bavaresco, N. Herrera Valencia, C. Klöckl, M. Pivoluska, P. Erker, N. Friis, M. Malik, and M. Huber, Measurements in two bases are sufficient for certifying high-dimensional entanglement, *Nature Physics* **14**, 1032 (2018).
- [21] O. Lib, S. Liu, R. Shekel, Q. He, M. Huber, Y. Bromberg, and G. Vitagliano, Experimental certification of high-dimensional entanglement with randomized measurements, *arXiv preprint arXiv:2412.04643* (2024).
- [22] M. Dąbrowski, M. Mazelanik, M. Parniak, A. Leszczyński, M. Lipka, and W. Wasilewski, Certification of high-dimensional entanglement and einstein-podolsky-rosen steering with cold atomic quantum memory, *Physical Review A* **98**, 042126 (2018).
- [23] Z. Huang, L. Maccone, A. Karim, C. Macchiavello, R. J. Chapman, and A. Peruzzo, High-dimensional entanglement certification, *Scientific Reports* **6**, 27637 (2016).
- [24] S. Liu, Q. He, M. Huber, O. Gühne, and G. Vitagliano, Characterizing entanglement dimensionality from randomized measurements, *PRX Quantum* **4**, 020324 (2023).
- [25] A. Tavakoli and S. Morelli, Enhanced schmidt-number criteria based on correlation trace norms, *Physical Review A* **110**, 062417 (2024).
- [26] C. Zhang, S. Denker, A. Asadian, and O. Gühne, Analyzing quantum entanglement with the schmidt decomposition in operator space, *Physical Review Letters* **133**, 040203 (2024).
- [27] D. Chruściński and G. Sarbicki, Entanglement witnesses: construction, analysis and classification, *Journal of Physics A: Mathematical and Theoretical* **47**, 483001 (2014).
- [28] X. Shi, Families of schmidt-number witnesses for high dimensional quantum states, *Communications in Theoretical Physics* **76**, 085103 (2024).
- [29] A. Sanpera, D. Bruß, and M. Lewenstein, Schmidt-number witnesses and bound entanglement, *Physical Review A* **63**, 050301 (2001).
- [30] F. Hirsch and M. Huber, The schmidt number of a quantum state cannot always be device-independently certified, *arXiv preprint arXiv:2003.14189* (2020).
- [31] N. Brunner, D. Cavalcanti, S. Pironio, V. Scarani, and S. Wehner, Bell nonlocality, *Rev. Mod. Phys.* **86**, 419 (2014).
- [32] R. F. Werner, Quantum states with einstein-podolsky-rosen correlations admitting a hidden-variable model, *Physical Review A* **40**, 4277 (1989).
- [33] J. Barrett, Nonsequential positive-operator-valued measurements on entangled mixed states do not always violate a bell inequality, *Physical Review A* **65**, 042302 (2002).
- [34] F. Buscemi, All entangled quantum states are nonlocal, *Physical review letters* **108**, 200401 (2012).
- [35] C. Branciard, D. Rosset, Y.-C. Liang, and N. Gisin, Measurement-device-independent entanglement witnesses for all entangled quantum states, *Physical review letters* **110**, 060405 (2013).
- [36] Q. Zhao, X. Yuan, and X. Ma, Efficient measurement-device-independent detection of multipartite entanglement structure, *Phys. Rev. A* **94**, 012343 (2016).
- [37] F. Shahandeh, M. J. W. Hall, and T. C. Ralph, Measurement-device-independent approach to entanglement measures, *Phys. Rev. Lett.* **118**, 150505 (2017).
- [38] K. T. Goh, J.-D. Bancal, and V. Scarani, Measurement-device-independent quantification of entanglement for given hilbert space dimension, *New Journal of Physics* **18**, 045022 (2016).
- [39] D. Rosset, A. Martin, E. Verbanis, C. C. W. Lim, and R. Thew, Practical measurement-device-independent en-



- tanglement quantification, *Phys. Rev. A* **98**, 052332 (2018).
- [40] L.-L. Sun, X. Zhou, A. Tavakoli, Z.-P. Xu, and S. Yu, Bounding the amount of entanglement from witness operators, *Phys. Rev. Lett.* **132**, 110204 (2024).
- [41] I. Šupić, P. Skrzypczyk, and D. Cavalcanti, Measurement-device-independent entanglement and randomness estimation in quantum networks, *Phys. Rev. A* **95**, 042340 (2017).
- [42] E. P. Lobo, S. G. Naik, S. Sen, R. K. Patra, M. Banik, and M. Alimuddin, Certifying beyond quantumness of locally quantum no-signaling theories through a quantum-input bell test, *Phys. Rev. A* **106**, L040201 (2022).
- [43] B. Yu and M. Hayashi, Measurement-device-independent detection of beyond-quantum state, *New Journal of Physics* **26**, 103003 (2024).
- [44] E. G. Cavalcanti, M. J. W. Hall, and H. M. Wiseman, Entanglement verification and steering when alice and bob cannot be trusted, *Phys. Rev. A* **87**, 032306 (2013).
- [45] D. Cavalcanti, P. Skrzypczyk, and I. Šupić, All entangled states can demonstrate nonclassical teleportation, *Phys. Rev. Lett.* **119**, 110501 (2017).
- [46] D. Rosset, C. Branciard, N. Gisin, and Y.-C. Liang, Entangled states cannot be classically simulated in generalized bell experiments with quantum inputs, *New Journal of Physics* **15**, 053025 (2013).
- [47] M. Banik, Lack of measurement independence can simulate quantum correlations even when signaling can not, *Phys. Rev. A* **88**, 032118 (2013).
- [48] E. Verbanis, A. Martin, D. Rosset, C. Lim, R. Thew, and H. Zbinden, Resource-efficient measurement-device-independent entanglement witness, *Physical review letters* **116**, 190501 (2016).
- [49] S. Kocsis, M. J. Hall, A. J. Bennet, D. J. Saunders, and G. J. Pryde, Experimental measurement-device-independent verification of quantum steering, *Nature communications* **6**, 5886 (2015).
- [50] P. Xu, X. Yuan, L.-K. Chen, H. Lu, X.-C. Yao, X. Ma, Y.-A. Chen, and J.-W. Pan, Implementation of a measurement-device-independent entanglement witness, *Physical review letters* **112**, 140506 (2014).
- [51] A. Peres, *Quantum theory: concepts and methods*, Vol. 72 (Springer, 1997).
- [52] M. A. Nielsen and I. L. Chuang, *Quantum computation and quantum information* (Cambridge university press, 2010).
- [53] R. B. Holmes, *Geometric functional analysis and its applications*, Vol. 24 (Springer Science & Business Media, 2012).
- [54] M. J. Renner, Compatibility of generalized noisy qubit measurements, *Phys. Rev. Lett.* **132**, 250202 (2024).
- [55] Y. Zhang and E. Chitambar, Exact steering bound for two-qubit werner states, *Physical Review Letters* **132**, 250201 (2024).
- [56] B. Mallick, N. Ganguly, and A. S. Majumdar, On the characterization of schmidt number breaking and annihilating channels, *arXiv preprint arXiv:2411.19315* (2024).
- [57] R. Augusiak, M. Demianowicz, and A. Acín, Local hidden-variable models for entangled quantum states, *Journal of Physics A: Mathematical and Theoretical* **47**, 424002 (2014).
- [58] J. Silman, S. Machnes, and N. Aharon, On the relation between bell's inequalities and nonlocal games, *Physics Letters A* **372**, 3796 (2008).
- [59] J. F. Clauser, M. A. Horne, A. Shimony, and R. A. Holt, Proposed experiment to test local hidden-variable theories, *Physical review letters* **23**, 880 (1969).
- [60] S. Ishizaka, Bound entanglement provides convertibility of pure entangled states, *Physical review letters* **93**, 190501 (2004).
- [61] M. B. Plenio and S. Virmani, An introduction to entanglement measures, *Quantum Info. Comput.* **7**, 1–51 (2007).
- [62] W. Dür, G. Vidal, and J. I. Cirac, Three qubits can be entangled in two inequivalent ways, *Physical Review A* **62**, 062314 (2000).
- [63] C. Eltschka and J. Siewert, Quantifying entanglement resources, *Journal of Physics A: Mathematical and Theoretical* **47**, 424005 (2014).
- [64] C. Schmid, N. Kiesel, W. Laskowski, W. Wieczorek, . f. M. Żukowski, and H. Weinfurter, Discriminating multipartite entangled states, *Physical review letters* **100**, 200407 (2008).