Density estimates for a nonlocal variational model with a degenerate double-well potential via the Sobolev inequality

Serena Dipierro, Alberto Farina, Giovanni Giacomin and Enrico Valdinoci *

February 20, 2025

Abstract

We provide density estimates for level sets of minimizers of the energy

$$\frac{1}{2} \int_{\Omega} \int_{\Omega} \frac{|u(x) - u(y)|^p}{|x - y|^{n + sp}} \, dx \, dy + \int_{\Omega} \int_{\mathbb{R}^n \setminus \Omega} \frac{|u(x) - u(y)|^p}{|x - y|^{n + sp}} \, dx \, dy + \int_{\Omega} W(u(x)) \, dx$$

where $p \in (1, +\infty)$, $s \in (0, \frac{1}{p})$ and W is a double-well potential with polynomial growth $m \in [p, +\infty)$ from the minima.

These kinds of potentials are "degenerate", since they detach "slowly" from the minima, therefore they provide additional difficulties if one wishes to determine the relative sizes of the "layers" and the "pure phases". To overcome these challenges, we introduce new barriers allowing us to rely on the fractional Sobolev inequality and on a suitable iteration method.

The proofs presented here are robust enough to consider the case of quasilinear nonlocal equations driven by the fractional *p*-Laplacian, but our results are new even for the case p = 2.

Mathematics Subject Classification 2020: 47G10, 47B34, 35R11, 35B08.

Keywords: Nonlocal energies, fractional Laplacian, degenerate potential.

^{*}Serena Dipierro, Giovanni Giacomin and Enrico Valdinoci: Department of Mathematics and Statistics, University of Western Australia, 35 Stirling Highway, WA6009 Crawley, Australia.

Alberto Farina: LAMFA, UMR CNRS 7352, Université Picardie Jules Verne 33, rue St Leu, 80039 Amiens, France. serena.dipierro@uwa.edu.au, alberto.farina@u-picardie.fr giovanni.giacomin@research.uwa.edu.au, enrico.valdinoci@uwa.edu.au

SD, GG and EV are members of Australian Mathematical Society. Supported by the Australian Future Fellowship FT230100333 "New perspectives on nonlocal equations" and the Australian Laureate Fellowship FL190100081 "Minimal surfaces, free boundaries and partial differential equations". Helpful discussions with Francesco De Pas and Jack Thompson are acknowledged. AF and GG would like to thank the Institute Henri Poincaré where part of this research was performed.

1 Introduction

Phase coexistence models were introduced in physics to study the interfaces between regions associated with different values of a suitable state parameter, such as the density of a fluid in capillarity phenomena [Row79], the fraction of volume occupied by two different materials in a non homogeneous system [CH58], or the density of the superfluid component of helium [GP58]. Since these early works, phase coexistence models have been developed to describe a variety of physical systems characterized by the interaction of two or more components.

All these models are described by some region $\Omega \subset \mathbb{R}^n$ and a state parameter $u : \Omega \to [-1, 1]$, where the "pure phases" correspond in our setting to the values 1 and -1, and the phase separation is induced by the minimization of a suitable energy functional.

The prototype of such an energy consists of a potential and an interaction term. The role of the potential term is to push the state parameter towards the pure phases, while the interaction term discourages the production of unneeded interfaces by suitably "shaping" the portions of the domain in which the value of the state parameter is either 1 or -1 (or is "sufficiently close" to these values).

In the local setting, the interaction energy is proportional to a gradient term of L^p -type with $p \in (1, +\infty)$, see for instance [AC81, ACF84, Bou90, PV05a, PV05b]. In order to capture long-range interactions, in the nonlocal framework the L^p -norm of the gradient has been replaced by the Gagliardo seminorm, see for instance [PSV13, SV11, SV14]. For further references regarding local and nonlocal phase separation models we refer to the survey [DV23].

The potential energy is given by the L^1 -norm of a "double-well" function W whose absolute minima are the pure phases. More precisely, $W : [-1, 1] \to \mathbb{R}$ satisfies

$$W(t) > 0$$
 for every $t \in (-1, 1)$ and $W(\pm 1) = 0.$ (1.1)

In the study of phase coexistence models, it is often desirable to quantify the size of the transition layer with respect to that of the pure phases (or, more precisely, of the regions close to pure phases). This is important since, due to the complexity of the problem, it is often difficult, when not impossible, to have general descriptions of the interfaces and the best one can do is typically to describe the phase separation "in a measure theoretic sense" and conclude that, at a large scale, most of the space is occupied by state parameters close to pure phases, separated by "thin" interfaces. In this spirit, the study of density estimates, namely of obtaining sharp quantifications of the measure of the interface, happens to be quite a cross-disciplinary topic, involving, under various perspectives, mathematical analysis, statistical mechanics, and materials sciences.

In the framework of density estimates a common hypothesis on the potential is that its growth from the minima is comparable to a polynomial of degree m, where $m \in (0, p]$ and $p \in (1, +\infty)$ is the exponent of the interaction term, see for instance formula (1.10) in [PV05a] for the local case and formula (1.9) in [SV14] for the nonlocal one.

In the current paper, we consider the energy functional

$$\mathcal{E}_{s}^{p}(u,\Omega) := \frac{1}{2} \int_{\Omega} \int_{\Omega} \frac{|u(x) - u(y)|^{p}}{|x - y|^{n + sp}} \, dx \, dy + \int_{\Omega} \int_{\mathbb{R}^{n} \setminus \Omega} \frac{|u(x) - u(y)|^{p}}{|x - y|^{n + sp}} \, dx \, dy + \int_{\Omega} W(u(x)) \, dx \quad (1.2)$$

where W presents a polynomial growth from the minima -1 and 1 of degree $m \in [p, +\infty)$. We will refer to these potentials as "degenerate", since they have a "slow" growth rate from their minima (in particular, slower than the homogeneity of the associated interaction energy).

In particular, we prove density estimates for every $p \in (1, +\infty)$ and $s \in \left(0, \frac{1}{p}\right)$, see Theorem 1.3 below. The precise hypothesis on W, the statement of Theorem 1.3 and its corollaries are all discussed in Section 1.3. In what follows, we introduce the concept of density estimate and we present in more detail the implications of the presence of a degenerate double-well potential.

1.1 Density estimates

To establish a conceptual framework in a model case, let us consider the prototype energy functional

$$\mathcal{E}(u,\Omega) := \frac{1}{2} \int_{\Omega} |\nabla u(x)|^2 \, dx + \int_{\Omega} W(u(x)) \, dx, \tag{1.3}$$

with W satisfying the assumptions in (1.1) and "nondegenerate" (in the sense that W detaches at least quadratically from its minima). From one point of view, as expected from the shape of the potential W, a minimizer for \mathcal{E} will be more likely to attain values close to the pure phases, to reduce the overall contribution of W in Ω . Yet, from another point of view, the interaction term penalizes "big jumps" of the state parameter. As a consequence of this simultaneous action, the minimizer will be pushed to attain values close to the pure phases but in such a way that at large scales the portion of the domain where the state parameter jumps from one pure phase to the other is negligible.

A common question in physics and mathematics is related to the geometrical features of these interfaces as we "zoom out". So far, there have a been two approaches to this question: one exploiting Γ -convergence and the other relying on density estimates.

On the one hand, the idea behind Γ -convergence is to analyse the limit functional obtained by suitably rescaling the minimizers of \mathcal{E} and deduce qualitative information regarding the interfaces at large scales. In particular, if u minimizes \mathcal{E} in Ω , then for $\epsilon \in (0, 1)$ we consider the rescaled state parameter

$$u_{\epsilon}(x) := u\left(\frac{x}{\epsilon}\right).$$

In [MM77], and later in greater generality in [Bou90], it is proved that there exists some set $E \subset \mathbb{R}^n$ such that as $\epsilon \to 0^+$ the minimizers u_{ϵ} converge, up to a subsequence, in $L^1_{loc}(\Omega)$ to¹ a step function $u_0 := \chi_E - \chi_{E^c}$. Interestingly, the interface ∂E is a minimal surface, namely a minimizer of the perimeter functional.

On the other hand, the purpose of density estimates is to provide lower bounds for the measure of the portion of the domain characterized by values of the minimizers of \mathcal{E} close to the pure phases. The first result in this direction is due to L. Caffarelli and A. Córdoba. In [CC95] they prove that if $u: \Omega \to [-1, 1]$ minimizes \mathcal{E} , then, for any $\theta \in (-1, 1)$ and $r \in (0, +\infty)$ sufficiently large, one has that $\{|u| > \theta\} \cap B_r$ is comparable to r^n (i.e., to the volume of the full ball B_r), while $\{|u| \leq \theta\} \cap B_r$ is bounded from above by a constant times r^{n-1} (corresponding, for example, to the (n-1)-dimensional surface measure in B_r of a hyperplane passing through the origin). In other words, state parameters close to the pure phases occupy a considerable portion of the domain, while the interface is negligible in the sense of measure.

As a consequence, as the scale parameter $\epsilon \in (0, 1)$ approaches zero, the set $\{|u_{\epsilon}| \leq \theta\} \cap B_r$ converges in the Hausdorff distance to the hypersurface ∂E . More specifically, for any $\delta, R > 0$ there exists $\epsilon_0 := \epsilon_0(\delta, R) > 0$ such that, for any $\epsilon \in (0, \epsilon_0)$,

$$\{|u_{\epsilon}| \leq \theta\} \cap B_R \subset \bigcup_{x \in \partial E} B_{\delta}(x).$$

Since the pioneering work of L. Caffarelli and A. Córdoba, density estimates have been used to study the interface convergence of minimizers and quasiminimizers of a larger spectrum of energy

$$\chi_A(a) := \begin{cases} 1 & \text{if } a \in A, \\ 0 & \text{if } a \notin A. \end{cases}$$

¹As usual, we denote here by χ_A the characteristic function (or indicator function) of the set A, namely

functionals related to phase separation. In [PV05a] and [PV05b] gradient terms of L^p -type with $p \in$ $(1, +\infty)$ are considered, together also with χ -shaped potentials W, namely $W(x) := \chi_{(-1,1)}(x)Q(x)$ with Q uniformly bounded away from zero. This type of potentials are related to models for fluid jets and capillarity phenomena; see also [Val01] for the elliptic case. In [FV08] density estimates are also proved under the weaker assumption of quasiminimality.

As already mentioned, nonlocal counterparts of (1.3) have been dealt with in the literature. A consolidated choice consists in replacing the L^2 -norm of the gradient with the Gagliardo seminorm. In [SV12] and [SV14], for $s \in (0,1)$ and $\epsilon \in (0,1)$, nonlocal phase separation was examined for the energy

$$\mathcal{F}_{s,\epsilon}^{2}(u,\Omega) := \epsilon^{2s} \left(\frac{1}{2} \int_{\Omega} \int_{\Omega} \frac{|u(x) - u(y)|^{2}}{|x - y|^{n + 2s}} \, dx \, dy + \int_{\Omega} \int_{\mathbb{R}^{n} \setminus \Omega} \frac{|u(x) - u(y)|^{2}}{|x - y|^{n + 2s}} \, dx \, dy \right) + \int_{\Omega} W(u(x)) \, dx.$$
(1.4)

More precisely, in [SV12] it is proved that, by suitably scaling the energy in (1.4) by some factor depending on ϵ and s, as ϵ vanishes, the minimizers are compact and the rescaled energy Γ -converges (to a functional of geometric flavor which we discuss below). Such a gauge is proved to be ϵ^{-2s} , $\epsilon^{-1}|\log(\epsilon)|$ and ϵ^{-1} respectively for the cases $s \in (0, \frac{1}{2})$, $s = \frac{1}{2}$ and $s \in (\frac{1}{2}, 1)$. Interestingly, in Theorem 1.4 of [SV12], it is established that if $s \in (0, \frac{1}{2})$, then $\epsilon^{-2s} \mathcal{F}_{s,\epsilon}^2 \Gamma$ -converges

as $\epsilon \to 0^+$ to the nonlocal perimeter $\operatorname{Per}_s(\cdot, \Omega)$ in Ω , defined for every measurable set $E \subset \mathbb{R}^n$ as

$$\operatorname{Per}_{s}(E,\Omega) := \int_{E\cap\Omega} \int_{E^{c}\cap\Omega} \frac{dx\,dy}{|x-y|^{n+2s}} + \int_{E\cap\Omega^{c}} \int_{E^{c}\cap\Omega} \frac{dx\,dy}{|x-y|^{n+2s}} + \int_{E\cap\Omega} \int_{E^{c}\cap\Omega^{c}} \frac{dx\,dy}{|x-y|^{n+2s}} + \int_{E\cap\Omega^{c}} \int_{E^{c}\cap\Omega^{c}} \frac{dx\,dy}{|x-y|^{n+2s}} + \int_{E^{c}\cap\Omega^{c}} \int_{E^{c}$$

Additionally, in [SV12] it is proved that if $s \in \left[\frac{1}{2}, 1\right)$ the Γ -limit coincides with the classical perimeter. Surprisingly, this means that in the regime $s \in \left[\frac{1}{2}, 1\right)$ and at large scales the effect of the Gagliardo seminorm on the interfaces is similar to the one of the L^2 -norm of the gradient and the phase separation resembles the classical one.

In this sense, the case $s \in (0, \frac{1}{2})$ (when p = 2, or, more generally, the case $s \in (0, \frac{1}{2})$ when $p \in$ $(1, +\infty)$) is considered as "genuinely nonlocal", since the problem retains its nonlocal features at every scale.

For what concerns density estimates, the cases $s \in (0, \frac{1}{2})$ and $s \in [\frac{1}{2}, 1)$ can be treated separately. As a matter of fact, in [SV11] density estimates are proved for the functional in (1.4) for every $s \in$ $(0, \frac{1}{2})$ using the Sobolev inequality. This approach fails when $s \in \lfloor \frac{1}{2}, 1 \rfloor$ and in [SV14] a fine measure theoretic result is established, namely Theorem 1.6 in [SV14], to obtain density estimates for every $s \in$ (0, 1), see Theorem 1.4 in [SV14].

The Sobolev inequality will be employed also in the current paper to obtain density estimates for the energy in (1.2). Analogously to the case p = 2 dealt with in [SV11] and [SV14], this approach only works if $s \in \left(0, \frac{1}{p}\right)$. The case $s \in \left[\frac{1}{p}, 1\right)$ will be treated in a forthcoming article [DFGV].

1.2Degenerate double-well potentials

An interesting problem surrounding the topic of density estimates is how the growth from the minima of the double-well potential W might affect phase separation. For example, a slow growth might induce, in some cases, the formation of "intermediate phases". To picture this, we may consider the extreme situation where, for some $a \in (0, 1)$ and every $x \in (-1, 1)$, the potential is given by

$$W(x) := \chi_{[-a,a]}(x), \tag{1.5}$$

see Figure 1. In this case, the minimizer will occupy values close to -a and a to reduce the interaction, whether this is prompted by a nonlocal or local type of interaction term, while the state parameters with value in (-1, -a) and (a, 1) become empty, violating density estimates. In particular, this shows that to obtain density estimates some assumptions on W are necessary.

As already mentioned, a common hypothesis on the potential is that if $p \in (1, +\infty)$ is the exponent of the interaction term, then, for some $m \in (0, p]$, $\lambda \in (0, 1)$ and $\Lambda \in [1, +\infty)$, and every $x \in (-1, 1)$,

$$\lambda (1 - x^2)^m \leqslant W(x) \leqslant \Lambda (1 - x^2)^m.$$
(1.6)

This corresponds, in the framework of this paper, to a nondegenerate double-well potential, since the growth from the potential's minima is not larger than the homogeneity of the interaction term. In particular, in the classical case of a gradient term of L^2 -type this translates into a (sub)quadratic growth, see [CC95]. In such a case, a common double-well potential in the literature is

$$W(x) := \frac{(1-x^2)^2}{4}.$$
(1.7)

In view of (1.6) and (1.7), for $m \in [p, +\infty)$ and $p \in (1, +\infty)$ it is natural to define

$$W(x) := \frac{(1-x^2)^m}{2m}$$
(1.8)

as a prototype of a degenerate double-well potential. See Figure 1 for a comparison with the potential in (1.7).



Figure 1: Plots of $\chi_{[-1/2,1/2]}(x)$, $(1-x^2)^2$ and $(1-x^2)^8$ respectively in green, blue and red.

Of course, the conditions in (1.6) may be too restrictive, and density estimates may still hold for double-well potentials with a flatter profile close to the minima. This problem has been first addressed in the local setting in [DFV18], where the authors prove density estimates for quasiminimizers of a phase transition energy driven by the L^p -norm of the gradient and with a degenerate double-well potential of the type (1.8), as far as some condition on n, m and p are met, see equation (1.7) in [DFV18].

The aim of this paper is to obtain density estimates for minimizers of the energy in (1.2), when the potential W presents a polynomial growth from the minima comparable to the one of the degenerate potential in (1.8).

In the following section we introduce all the necessary mathematical notation, state precisely the main results and provide the structure of the paper.

1.3 Mathematical framework and main results

In what follows $n \in \mathbb{N}$, $\Omega \subset \mathbb{R}^n$ is open, $p \in (1, +\infty)$ and $s \in \left(0, \frac{1}{p}\right)$. We consider $W \in C^1([-1, 1], \mathbb{R}^+)$ such that for some $\theta \in (-1, 1)$, $\lambda \in (0, 1]$, $\Lambda \in [1, +\infty)$ and $m \in [p, +\infty)$, for every $x \in [-1, 1]$,

$$\lambda \chi_{(-\infty,\theta]}(x) |1+x|^m \leqslant W(x) \leqslant \Lambda |1+x|^m.$$
(1.9)

Moreover, we assume that, for some $q \in (0, \theta+1] \cap (0, 1)$ and $c_1 \in (0, +\infty)$, for every $x \in (-1, -1+q)$,

$$c_1 |1+x|^{m-1} \leq W'(x).$$
 (1.10)

Note that the prototype of a degenerate double-well potential in (1.8) satisfies (1.9) and (1.10).

Also, for every measurable function $u: \mathbb{R}^n \to \mathbb{R}$ we define

$$\mathcal{K}_{s}^{p}(u,\Omega) := \frac{1}{2} \int_{\Omega} \int_{\Omega} \frac{|u(x) - u(y)|^{p}}{|x - y|^{n + sp}} \, dx \, dy + \int_{\Omega} \int_{\mathbb{R}^{n} \setminus \Omega} \frac{|u(x) - u(y)|^{p}}{|x - y|^{n + sp}} \, dx \, dy \tag{1.11}$$

and rewrite the energy in (1.2) as

$$\mathcal{E}_s^p(u,\Omega) := \mathcal{K}_s^p(u,\Omega) + \int_{\Omega} W(u(x)) \, dx.$$
(1.12)

Furthermore, in analogy with [SV12], for every $\epsilon \in (0, 1)$, we consider the rescaled energy

$$\mathcal{F}^p_{s,\epsilon}(u,\Omega) := \mathcal{K}^p_s(u,\Omega) + \epsilon^{-sp} \int_{\Omega} W(u(x)) \, dx.$$
(1.13)

A convenient function space to study \mathcal{K}_s^p is

 $\widehat{H}^{s,p}(\Omega) := \left\{ u : \mathbb{R}^n \to \mathbb{R} \text{ measurable s.t. } \mathcal{K}^p_s(u,\Omega) < +\infty \right\}.$

Note that $W^{s,p}(\mathbb{R}^n) \subset \widehat{H}^{s,p}(\Omega)$. Since W is defined in the interval [-1,1], we consider the subspace of all those functions $u \in \widehat{H}^{s,p}(\Omega)$ such that $|u| \leq 1$ a.e., namely

$$X^{s,p}(\Omega) := \left\{ u \in \widehat{H}^{s,p}(\Omega) \text{ s.t. } \|u\|_{L^{\infty}(\mathbb{R}^n)} \leqslant 1 \right\}.$$
(1.14)

With these choices we have the following definition:

Definition 1.1 (Minimizer for \mathcal{E}_s^p and $\mathcal{F}_{s,\epsilon}^p$). If $\Omega \subset \mathbb{R}^n$ is bounded, we say that $u \in X^{s,p}(\Omega)$ is a minimizer for $\mathcal{E}_s^p(\cdot, \Omega)$ in $X^{s,p}(\Omega)$ if

$$\mathcal{E}^p_s(u,\Omega) \leqslant \mathcal{E}^p_s(v,\Omega)$$

for any $v \in X^{s,p}(\Omega)$ such that $v \equiv u$ in $\mathbb{R}^n \setminus \Omega$.

If $\Omega \subset \mathbb{R}^n$ is unbounded, we say that $u \in X^{s,p}(\Omega)$ is a minimizer for $\mathcal{E}^p_s(\cdot, \Omega)$ in $X^{s,p}(\Omega)$ if it minimizes $\mathcal{E}^p_s(\cdot, \Omega')$ in $X^{s,p}(\Omega')$ for every bounded and open set $\Omega' \subseteq \Omega$.

Similarly, one also defines a minimizer for $\mathcal{F}_{s,\epsilon}^p$.

Remark 1.2. Note that if u minimizes \mathcal{E}_s^p (or $\mathcal{F}_{s,\epsilon}^p$) in an open and bounded set $\Omega \subset \mathbb{R}^n$, then it minimizes \mathcal{E}_s^p (or $\mathcal{F}_{s,\epsilon}^p$) in every measurable subset $\Omega' \subset \Omega$.

The main result of this paper is the following density estimate:

Theorem 1.3 (Density estimates). Let $p \in (1, +\infty)$, $s \in \left(0, \frac{1}{p}\right)$ and $\theta_1, \theta_2 \in (-1, \theta]$.

Let also

$$\theta_* := \min \{\theta_1, \theta_2, -1 + q\} \quad and \quad \theta^* := \max \{\theta_1, \theta_2, -1 + q\}.$$
(1.15)

Assume that u is a minimizer for \mathcal{E}_s^p in Ω and that, for some $c_0, r_0 \in (0, +\infty)$ such that $B_{r_0} \subset \Omega$,

$$\mathcal{L}^{n}(B_{r_{0}} \cap \{u > \theta_{1}\}) > c_{0}.$$
(1.16)

Then, there exist $R^* := R^*_{s,n,p,m,\theta_*,r_0} \in [r_0, +\infty)$, $\tilde{c} := \tilde{c}_{s,n,p,m,\Lambda,c_1,\theta_*,r_0,c_0} \in (0,1)$, and $c := c_{m,\theta_*} \in (0, +\infty)$, such that, for any $r \in [R^*, +\infty)$ with $B_{\frac{3r}{2}} \subset \Omega$,

$$c \int_{B_r \cap \{\theta_* < u \leq \theta^*\}} |1 + u(x)|^m \, dx + \mathcal{L}^n(B_r \cap \{u > \theta_2\}) \geqslant \tilde{c} \, r^n.$$

$$(1.17)$$

Here above and in the rest of the paper, $\mathcal{L}^n(E)$ denotes the Lebesgue measure of a set $E \subset \mathbb{R}^n$.

The proof of the density estimates in Theorem 1.3 relies on the construction of a suitable barrier (see Theorem 3.1 below) that is inspired by Lemma 3.1 in [SV14], where a barrier is built in the case p = 2 and m = 2. The extension to the cases $p \neq 2$ and $m \in [p, +\infty)$ that we provide in this paper is nontrivial, due to the nonlinearity of the operator and the precence of a degenerate potential. In the case of the classical *p*-Laplacian, barriers have been built in [VSS06] in a different context. The case p = 2 in the nonlocal setting with a degenerate potential has been also considered in [DPDPV].

The presence of the first term in the left-hand side of (1.17) is a consequence of the weak lower bound in (1.9). More precisely, under some additional assumptions on the lower bound for W(see (1.19) below) it is possible to reabsorb such term in the right hand-side of the inequality, obtaining the "full density estimates". This is made possible thanks to the following upper bound on the energy \mathcal{E}_s^p . Its proof is contained in the forthcoming paper [DFGV].

Theorem 1.4 ([DFGV]). Let $p \in (1, +\infty)$, $s \in \left(0, \frac{1}{p}\right)$ and u be a minimizer of \mathcal{E}_s^p in $X^{s,p}(B_{R+2})$ with $R \ge 2$. Then, there exists $\overline{C} > 0$, depending only on s, n, p, m and Λ , such that

$$\mathcal{E}_s^p(u, B_R) \leqslant \bar{C}R^{n-sp}. \tag{1.18}$$

From Theorems 1.3 and 1.4 we obtain the following density estimate:

Corollary 1.5. Let θ_1 , $\theta_2 \in (-1, 1)$ and u be as in the statement of Theorem 1.3. Assume that, for every $\mu \in (-1, 1)$, there exists $\lambda_{\mu} \in (0, +\infty)$ such that, for every $x \in (-1, 1)$,

$$\lambda_{\mu}\chi_{(-\infty,\mu]}(x)\left|1+x\right|^{m} \leqslant W(x).$$
(1.19)

Moreover, let θ_* and θ^* be as in (1.15).

Then, there exist $\widetilde{R} := \widetilde{R}_{s,n,p,m,\theta_*,r_0,\lambda_{\theta^*}} \in [r_0,+\infty)$ and $\hat{c} := \hat{c}_{s,n,p,m,\Lambda,c_1,\theta_*,r_0,c_0} \in (0,1)$ such that, for any $r \in [\widetilde{R},+\infty)$ with $B_{\frac{3r}{2}} \subset \Omega$,

$$\mathcal{L}^n(B_r \cap \{u > \theta_2\}) \geqslant \hat{c} r^n. \tag{1.20}$$

As already mentioned, an important consequence of the density estimates is the uniform convergence of the interface of the minimizers of the rescaled energy functional to a hypersurface. To prove this result we require an additional condition on the potential W. In particular, we assume that there exists some $c_2 \in (0, +\infty)$ such that for every $x \in (-1, 1)$

$$W(x) \ge c_2(1-x^2)^m.$$
 (1.21)

Theorem 1.6 (Compactness). Let $p \in (1, +\infty)$, $s \in \left(0, \frac{1}{p}\right)$ and W satisfy (1.21). Assume that the sequence $\{u_{\epsilon}\}_{\epsilon} \subset X^{s,p}(\Omega)$ satisfies

$$\sup_{\epsilon \in (0,1)} \mathcal{F}^p_{s,\epsilon}(u_{\epsilon}, \Omega) < +\infty.$$
(1.22)

Then, there exists a measurable set $E \subset \mathbb{R}^n$ such that, up to a subsequence,

$$u_{\epsilon} \to u^* := \chi_E - \chi_{E^c} \quad in \quad L^1_{loc}(\Omega). \tag{1.23}$$

This compactness result is a consequence of the scaling chosen for $\mathcal{F}_{s,\epsilon}^p$. Indeed, it follows from (1.13) and (1.22) that the Gagliardo seminorm of u_{ϵ} is bounded on all compact subsets of Ω . Then, it is enough to use compact embeddings of $W^{s,p}$ to conclude. The case $s \in \left[\frac{1}{p}, 1\right)$ is more involved, and it will be dealt with in the forthcoming paper [DFGV].

As a byproduct of the density estimates in (1.20) and the compactness in Theorem 1.6 we obtain the Hausdorff convergence for the interfaces:

Corollary 1.7 (Hausdorff convergence). Let $p \in (1, +\infty)$, $s \in \left(0, \frac{1}{p}\right)$ and W satisfy (1.21). For $\epsilon \to 0^+$ let $\{u_{\epsilon}\}_{\epsilon} \subset X^{s,p}(\Omega)$ be such that u_{ϵ} is a minimizer of $\mathcal{F}_{s,\epsilon}^p$ in Ω and (1.22) holds. Moreover, let $E \subset \mathbb{R}^n$ be as in the statement of Theorem 1.6.

Then, for every $\Theta \in (0,1)$, $\delta \in (0,+\infty)$, and $R \in (0,+\infty)$ satisfying $B_R \Subset \Omega$, there exists $\epsilon_0 := \epsilon_0(\Theta, \delta, R) > 0$ such that, if $\epsilon \in (0, \epsilon_0)$,

$$\{|u_{\epsilon}| < \Theta\} \cap B_R \subset \bigcup_{x \in \partial E} B_{\delta}(x).$$

We remark that the results presented in this paper are stated and proved for all $p \in (1, +\infty)$, but they are new even for the case p = 2.

The rest of this paper is structured as follows. In Section 2 we recall some known results that will be used in the proofs of the main statements. In Section 3 we construct a suitable barrier, see Theorem 3.1 below, which will be used to prove Theorem 1.3. The construction of this barrier relies also on some technical results that are collected in Appendix A.

The proofs of Theorem 1.3 and Corollary 1.5 are contained in Section 4. Finally, in Section 5 we prove Theorem 1.6 and Corollary 1.7.

2 Some auxiliary results

In this section we recall some auxiliary results that will be used throughout the proofs of our main theorems.

2.1 A general estimate

The following statement is provided in Lemma 3.2. in [SV14]. We point out that by a careful inspection of the proof presented in [SV14] one can check that the dependance of the constants \tilde{c} and R_* is as claimed here below (this will be important in the forthcoming paper [DFGV]).

Lemma 2.1 (Lemma 3.2. in [SV14]). Let σ , $R_0 \in (0, +\infty)$, $\nu \in (\sigma, +\infty)$ and γ , μ , $C \in (1, +\infty)$.

Let $V: (0, +\infty) \to (0, +\infty)$ be a nondecreasing function. For any $r \in [R_0, +\infty)$, let

$$\alpha(r) := \min\left\{1, \frac{\log(V(r))}{\log(r)}\right\}$$

Also, suppose that

$$V(R_0) \geqslant \mu \tag{2.1}$$

and that, for any $r \in [R_0, +\infty)$,

$$r^{\sigma}\alpha(r)\big(V(r)\big)^{\frac{\nu-\sigma}{\nu}} \leqslant C V(\gamma r).$$
(2.2)

Then, there exist $\tilde{c} \in (0,1)$, depending on C, σ , μ , γ , ν and R_0 , and $R_* \in [R_0, +\infty)$, depending on R_0 and γ , such that, for any $r \in [R_*, +\infty)$,

$$V(r) \geqslant \tilde{c} r^{\nu}. \tag{2.3}$$

2.2 Hölder regularity for minimizers

In what follows, we let $F \in L^{\infty}(\mathbb{R}, \mathbb{R})$. Also, for every $u \in \widehat{H}^{s,p}(\Omega)$ we consider the renormalized energy

$$\bar{\mathcal{E}}_s^p(u,\Omega) := (1-s)\mathcal{K}_s^p(u,\Omega) + \int_{\Omega} F(u(x)) \, dx.$$

As customary, given some open set $B \subset \mathbb{R}^n$ and $\beta \in (0,1)$, we denote the Hölder seminorm of a function $f: B \to \mathbb{R}$ by

$$[f]_{C^{\beta}(B)} := \sup_{x,y \in B} \frac{|f(x) - f(y)|}{|x - y|^{\beta}}$$

If $\beta = 1$, we denote the Lipschitz seminorm of $f : B \to \mathbb{R}$ by

$$[f]_{C^{0,1}(B)} := \sup_{x,y \in B} \frac{|f(x) - f(y)|}{|x - y|}.$$

Also, given $x_0 \in \Omega$ and $R \in (0, \text{dist}(x_0, \partial \Omega))$, we set

$$\operatorname{Tail}(u, x_0, R) := \left[(1-s) R^{sp} \int_{\mathbb{R}^n \setminus B_{R(x_0)}} \frac{|u(y)|^{p-1}}{|y-x_0|^{n+sp}} \right]^{\frac{1}{p-1}}.$$

With this notation, we now recall the following result on the Hölder regularity for the minimizers of $\bar{\mathcal{E}}_s^p$. Such result is contained in [Coz17], where the Hölder regularity is established using fractional De Giorgi classes.

Theorem 2.2 (Theorem 6.4 and Proposition 7.5 in [Coz17]). Let $n \in \mathbb{N}$, $s_0 \in (0,1)$, $s \in [s_0,1)$ and $p \in (1, +\infty)$. Let $u \in \widehat{H}^{s,p}(\Omega)$ be a minimizer for $\overline{\mathcal{E}}_s^p$ in Ω .

Then, $u \in C_{loc}^{\alpha}(\Omega)$ for some $\alpha \in (0, 1)$. Moreover, for any $x_0 \in \Omega$ and $R \in \left(0, \frac{\text{dist}(x_0, \partial \Omega)}{8}\right)$,

$$[u]_{C^{\alpha}(B_{R}(x_{0}))} \leq \frac{C}{R^{\alpha}} \left(\|u\|_{L^{\infty}(B_{4R}(x_{0}))} + Tail(u; x_{0}, 4R) + R^{s} \|F(u)\|_{L^{\infty}(B_{8R}(x_{0}))}^{\frac{1}{p}} \right)$$

for some $C \ge 1$.

The constants α and C depend only on n, p, s₀ when $n \ge 2$, and also on s when n = 1.

3 Construction of a suitable barrier

The following Theorem 3.1 is the extension to the cases $p \neq 2$ and $m \in [p, +\infty)$ of Lemma 3.1 in [SV14].

We point out that, in our construction, the dependence of the constants with respect to s is explicit. In this way, as it will be discussed in detail in [DFGV], it is possible to obtain density estimates that are stable as $s \to 1^-$, recovering local density estimates from the nonlocal ones.

In what follows, for every $t \in \mathbb{R}$ we denote by [t] the integer part of t. Also, for every $z : \mathbb{R}^n \to \mathbb{R}$ measurable and $x \in \mathbb{R}^n$, we denote its *p*-fractional Laplacian at $x \in \mathbb{R}^n$ as

$$(-\Delta)_p^s z(x) := \int_{\mathbb{R}^n} \frac{(z(x) - z(y)) |z(x) - z(y)|^{p-2}}{|x - y|^{n+sp}} \, dy.$$

Throughout this section we will also make use of some technical results collected in Appendix A.

Theorem 3.1. For any $n \in \mathbb{N}$, $\tau \in (0, +\infty)$, $p \in (1, +\infty)$, $s \in (0, 1)$ and $m \in [p, +\infty)$ there exists $\widehat{C}_{n,p,m} \in (0, +\infty)$ such that if we define

$$\bar{R} = \bar{R}_{s,\tau,n,p,m} := \left(\frac{\widehat{C}_{n,p,m}}{s(1-s)\tau}\right)^{\frac{1}{ps}}$$
(3.1)

the following statement holds true.

For every $R \in [\bar{R}, +\infty)$ there exists a rotationally symmetric function $w \in C(\mathbb{R}^n, (-1, 1])$ such that

$$w \equiv 1 \quad in \quad \mathbb{R}^n \setminus B_R \tag{3.2}$$

and, for every $x \in B_R$,

$$-(-\Delta)_{p}^{s}w(x) \leqslant \tau (1+w(x))^{m-1}.$$
(3.3)

Also, if we set $q := \frac{p}{m-1}$ and

$$C_{s,\tau,n,p,m} := 2^{q+1} \max\left\{1, \bar{R}^{qs}\right\},$$
(3.4)

then, for every $x \in B_R$,

$$1 + w(x) \leqslant \frac{C_{s,\tau,n,p,m}}{(1 + R - |x|)^{qs}}.$$
(3.5)

The proof of Theorem 3.1 will be based on the following construction. We take a large $r \in [1, +\infty)$, to be conveniently chosen with respect to R and τ here below (see formula (3.41)).

We consider the integer quantity

$$k = k(s, p) := \begin{cases} 0 & \text{if } p \in (1, 2) \text{ and } s \in \left(0, \frac{p-1}{2p}\right), \\ \left[\frac{sp}{p-1}\right] + 1 & \text{if } p \in (1, 2) \text{ and } s \in \left[\frac{p-1}{2p}, 1\right), \\ 1 & \text{if } p \in [2, +\infty) \text{ and } s \in (0, 1). \end{cases}$$
(3.6)

We set $q := \frac{p}{m-1}$ and we define, for every $t \in (0, r)$,

$$g(t) := t^{-qs}$$

and $f(t) := g(r-t) + \sum_{i=0}^{k} \frac{(-1)^{i+1}}{i!} g^{(i)} \left(\frac{r}{2}\right) \left(t - \frac{r}{2}\right)^{i}.$ (3.7)

Furthermore, we also define the functions

$$h(t) := \begin{cases} 0 & \text{if } t \in (0, r/2], \\ \min\{f(t), 1\} & \text{if } t \in (r/2, r), \\ 1 & \text{if } t \in [r, +\infty) \end{cases}$$

$$v(x) := h(|x|) \quad \text{for every} \quad x \in \mathbb{R}^n.$$

$$(3.8)$$

We provide the following estimate for the function h:

and

Proposition 3.2. Let $s \in (0,1)$, $p \in (1,+\infty)$, $m \in [p,+\infty)$ and $q := \frac{p}{m-1}$. Let $h : (0,+\infty) \to [0,1]$ be as in (3.8).

Then, there exists $\bar{c}_{k,q,s} \in (1, +\infty)$ such that, for every $t \in (0, +\infty)$,

$$\min\left\{1, (r-t)^{-ps}\right\} \leqslant (h(t) + \bar{c}_{k,q,s}r^{-qs})^{m-1}.$$
(3.9)

Proof. We observe that if $t \in \left(\frac{r}{2}, r\right)$ and h(t) < 1, then

$$h(t) = f(t)$$

$$= g(r-t) + \sum_{i=0}^{k} \frac{(-1)^{i+1}}{i!} g^{(i)} \left(\frac{r}{2}\right) \left(t - \frac{r}{2}\right)^{i}$$

$$\geqslant (r-t)^{-qs} - \sum_{i=0}^{k} \frac{|g^{(i)} \left(\frac{r}{2}\right)|}{i!} \left(t - \frac{r}{2}\right)^{i}$$

$$\geqslant (r-t)^{-qs} - \sum_{i=0}^{k} c_{q,s,i} \left(\frac{r}{2}\right)^{-qs-i} \left(\frac{r}{2}\right)^{i}$$

$$= (r-t)^{-qs} - c_{k,q,s} \left(\frac{r}{2}\right)^{-qs},$$
(3.10)

where

$$c_{q,s,i} := \frac{\prod_{j=0}^{i-1} (qs+j)}{i!} \quad \text{and} \quad c_{k,q,s} := \sum_{i=0}^{k} c_{q,s,i}.$$
(3.11)

On this account, if $t \in \left(\frac{r}{2}, r\right)$ and h(t) < 1 and we adopt the notation

$$\tilde{c}_{k,q,s} := 2^{qs} c_{k,q,s},$$
(3.12)

we can use (3.10) to deduce that

$$(h(t) + \tilde{c}_{k,q,s}r^{-qs})^{\frac{p}{q}} \ge ((r-t)^{-qs})^{\frac{p}{q}} = (r-t)^{-ps} \ge \min\left\{1, (r-t)^{-ps}\right\}.$$
(3.13)

Moreover, if h(t) = 1, then we have that

$$(h(t) + \tilde{c}_{k,q,s}r^{-qs})^{m-1} = (1 + \tilde{c}_{k,q,s}r^{-qs})^{m-1} \ge 1 \ge \min\left\{1, (r-t)^{-ps}\right\}.$$
(3.14)

Finally, if $t \in \left(0, \frac{r}{2}\right]$,

$$(h(t) + 2^{qs}r^{-qs})^{m-1} = (2^{qs}r^{-qs})^{\frac{p}{q}} = 2^{ps}r^{-ps} \ge (r-t)^{-ps} \ge \min\left\{1, (r-t)^{-ps}\right\}.$$
(3.15)

Gathering these pieces of information, we obtain the estimate in (3.9), as desired.

In particular, from (3.13), (3.14) and (3.15) we evince that (3.9) holds true with $\bar{c}_{k,q,s}$ given by

$$\bar{c}_{k,q,s} := \max\left\{2^{qs}, \tilde{c}_{k,q,s}\right\}.$$
 (3.16)

This completes the proof of Proposition 3.2.

The next observation provides a uniform bound in s for $\overline{c}_{k,q,s}^{\frac{1}{q_s}}$. For this, we define the constant

$$\hat{c}_{p,m} := 2e^{c_{p,m}^{(1)}},\tag{3.17}$$

where

$$c_{p,m}^{(1)} := \left(\left[\frac{p}{p-1} \right] + 2 \right) \prod_{j=1}^{\left\lfloor \frac{p}{p-1} \right\rfloor + 1} \left(\frac{p}{m-1} + j \right).$$
(3.18)

Lemma 3.3. Let $p \in (1, +\infty)$, $m \in [p, +\infty)$ and $q := \frac{p}{m-1}$. Let $\bar{c}_{k,q,s}$ be given by Proposition 3.2. Then, for every $s \in (0, 1)$,

$$\overline{c}_{k,q,s}^{\frac{1}{qs}} \leqslant \hat{c}_{p,m}.$$

Proof. To show this, from (3.16) (and recalling also (3.11) and (3.12)), we evince that

$$\bar{c}_{k,q,s} = \max\left\{2^{qs}, \tilde{c}_{k,q,s}\right\} = 2^{qs} \max\left\{1, c_{k,q,s}\right\} \leqslant 2^{qs} \left(1 + c_{k,q,s}\right).$$
(3.19)

Moreover, we notice that, by (3.6),

$$k \leqslant \left[\frac{sp}{p-1}\right] + 1.$$

Therefore, recalling (3.11), we see that

$$c_{k,q,s} \leqslant (k+1) \prod_{j=0}^{k} (qs+j)$$

$$\leqslant qs \left(\left[\frac{p}{p-1} \right] + 2 \right) \prod_{j=1}^{\left[\frac{p}{p-1} \right] + 1} \left(\frac{p}{m-1} + j \right)$$

$$= qsc_{p,m}^{(1)}.$$
(3.20)

From this and (3.19) we obtain that, for every $s \in (0, 1)$,

$$\overline{c}_{k,q,s}^{\frac{1}{q_s}} \leqslant 2\left(1 + c_{k,q,s}\right)^{\frac{1}{q_s}} \leqslant 2\left(1 + qsc_{p,m}^{(1)}\right)^{\frac{1}{q_s}}.$$
(3.21)

Now, we define the function $z:(0,1)\to (0,+\infty)$ as

$$z(s) := \left(1 + qsc_{p,m}^{(1)}\right)^{\frac{1}{qs}}$$

and we notice that z is decreasing in (0, 1). Therefore, from (3.21), we find that

$$\overline{c}_{k,q,s}^{\frac{1}{q_s}} \leqslant 2z(s) \leqslant 2 \limsup_{s \to 0^+} z(s) = 2e^{c_{p,m}^{(1)}},$$

which, together with (3.17), gives the desired result.

The barrier w in Theorem 3.1 will be built by suitably rescaling and translating the function v in (3.8). Thus, as a preliminary step in this direction, we now estimate the *p*-fractional Laplacian of v, according to the following statement:

Proposition 3.4. Let $s \in (0,1)$, $p \in (1,+\infty)$, $m \in [p,+\infty)$, $q := \frac{p}{m-1}$ and $r \in [1,+\infty)$. Let $v : \mathbb{R}^n \to [0,1]$ be as in (3.8).

Then, there exists $\widetilde{C} \in (0, +\infty)$ depending only on n, m and p such that, for every $x \in B_r$,

$$-(-\Delta)_{p}^{s}v(x) \leqslant \frac{\widetilde{C}}{s(1-s)} \left(v(x) + \bar{c}_{k,q,s}r^{-qs}\right)^{m-1}, \qquad (3.22)$$

where $\bar{c}_{k,q,s}$ is given in (3.16).

Proof. In order to prove (3.22), we show that there exists $\overline{C} \in (0, +\infty)$, depending only on n, m and p, such that, for every $x \in B_r$,

$$-(-\Delta)_{p}^{s}v(x) \leqslant \frac{\bar{C}}{s(1-s)} \left((r-|x|)^{-qs(p-1)-ps} + (r-|x|)^{-ps} \right),$$
(3.23)

and also

$$-(-\Delta)_p^s v(x) \leqslant \frac{C}{s(1-s)}.$$
(3.24)

If equations (3.23) and (3.24) hold true, then using them together with the estimate in (3.9) we obtain that, for every $x \in B_r$,

$$-(-\Delta)_{p}^{s}v(x) \leq \frac{C}{s(1-s)} \min\left\{1, (r-|x|)^{-qs(p-1)-ps} + (r-|x|)^{-ps}\right\}$$
$$\leq \frac{2\bar{C}}{s(1-s)} \min\left\{1, (r-|x|)^{-ps}\right\}$$
$$\leq \frac{2\bar{C}}{s(1-s)} \left(h\left(|x|\right) + \bar{c}_{k,q,s}r^{-qs}\right)^{m-1}$$
$$= \frac{2\bar{C}}{s(1-s)} \left(v(x) + \bar{c}_{k,q,s}r^{-qs}\right)^{m-1},$$

and therefore (3.22) holds true with $\tilde{C} := 2\bar{C}$.

Hence, from now on, we focus on the proof of (3.23) and (3.24). To accomplish this, in what follows we denote by $C \in (0, +\infty)$ a constant depending only on p, m and n and possibly changing from line to line.

If $p \in (1,2)$ and $s \in \left[\frac{p-1}{2p}, 1\right)$, we write that

$$-(-\Delta)_{p}^{s}v(x) \leq \int_{B_{\rho}(x)} \frac{(v(y) - v(x)) |v(y) - v(x)|^{p-2}}{|x - y|^{n+sp}} dy + 2^{p-1} ||v||_{L^{\infty}(\mathbb{R}^{n})} \int_{\mathbb{R}^{n} \setminus B_{\rho}(x)} \frac{dy}{|x - y|^{n+sp}}.$$

As a result, applying Lemma A.5 with $\rho := \frac{r-|x|}{2}$ we find that

$$-(-\Delta)_{p}^{s}v(x) \leq \frac{C_{n,p,m}}{1-s}\rho^{-qs(p-1)-ps} + \frac{c_{n,p}}{s}\rho^{-ps}$$
$$\leq \frac{C}{s(1-s)}\left((r-|x|)^{-qs(p-1)-ps} + (r-|x|)^{-ps}\right),$$

which establishes (3.23), while making use of Corollary A.7 with $\rho = \mu := 1$ we obtain that

$$-(-\Delta)_p^s v(x) \leqslant \frac{C_{1,p,m,n}}{p-sp} + \frac{c_{n,p}}{s} \leqslant \frac{C}{s(1-s)},$$

which proves (3.24) in the case $p \in (1,2)$ and $s \in \left[\frac{p-1}{2p}, 1\right]$. Now we let $p \in (1,2)$ and $s \in \left(0, \frac{p-1}{2p}\right)$. In this setting, recalling (3.6) and (3.8), we have that k = 0 and that k = 0 and

$$v(y) := \begin{cases} 0 & \text{if } y \in B_{\frac{r}{2}}, \\ (r - |y|)^{-qs} - \left(\frac{r}{2}\right)^{-qs} & \text{if } y \in B_{\hat{r}} \setminus \overline{B_{\frac{r}{2}}}, \\ 1 & \text{if } y \in \mathbb{R}^n \setminus \overline{B}_{\hat{r}}, \end{cases}$$

where $\hat{r} \in \left(\frac{r}{2}, r\right)$ is defined as

$$\hat{r} := \min\left\{t \in \left(\frac{r}{2}, r\right) \text{ s.t. } f(t) \ge 1\right\}.$$
(3.25)

Accordingly,

$$\nabla v(y) = \begin{cases} 0 & \text{if } y \in B_{\frac{r}{2}}, \\ qs(r-|y|)^{-qs-1}\frac{y}{|y|} & \text{if } y \in B_{\hat{r}} \setminus \overline{B_{\frac{r}{2}}}, \\ 0 & \text{if } y \in \mathbb{R}^n \setminus \overline{B}_{\hat{r}}, \end{cases}$$
(3.26)

As a consequence, for a.e. $y \in \mathbb{R}^n$,

$$|\nabla v(y)| \leqslant qs \left| r - \left| y \right| \right|^{-qs-1} \chi_{B_{\hat{r}} \setminus \overline{B_{\frac{r}{2}}}}.$$
(3.27)

Also, if $y \in B_{\frac{r-|x|}{2}}(x)$ then

$$|y| \leq |y - x| + |x| \leq \frac{r - |x|}{2} + |x| \leq \frac{r + |x|}{2} < r,$$

and thus

$$|r - |y|| = r - |y| \ge \frac{r - |x|}{2}.$$
(3.28)

Hence, using (3.27) and (3.28) we obtain that, if $y \in B_{\frac{r-|x|}{2}}(x)$,

$$|\nabla v(y)| \leqslant qs 2^{qs+1} \left(r - |x|\right)^{-qs-1},$$

and therefore

$$\|\nabla v\|_{L^{\infty}\left(B_{\frac{r-|x|}{2}}(x)\right)} < q2^{q+1}(r-|x|)^{-qs-1}.$$
(3.29)

We thus apply Lemma A.1 with

$$\widehat{K} := \|\nabla v\|_{L^{\infty}\left(B_{\frac{r-|x|}{2}}\right)} \quad \text{and} \quad \rho := \frac{r-|x|}{2}$$

and, thanks to (3.29), we find that

$$\begin{aligned} \left| (-\Delta)_{p}^{s} v(x) \right| &\leq \frac{c_{n,p}}{s(p-1-ps)} \left(\left\| \nabla v \right\|_{L^{\infty} \left(B_{\frac{r-|x|}{2}} \right)}^{p-1} \left(\frac{r-|x|}{2} \right)^{-ps+p-1} + \left(\frac{r-|x|}{2} \right)^{-ps} \right) \\ &\leq \frac{C}{s(1-s)} \left((r-|x|)^{-qs(p-1)-ps} + (r-|x|)^{-ps} \right). \end{aligned}$$

This concludes the proof of (3.23) for the case $p \in (1,2)$ and $s \in \left(0, \frac{p-1}{2p}\right)$.

Furthermore, from (3.27) and Proposition A.6 we evince that, for every $y \in \mathbb{R}^n$,

$$|\nabla v(y)| \leqslant qs(r-\hat{r})^{-qs-1} \leqslant q\widehat{C}_{p,m}^{-qs-1} \leqslant C,$$

and so $\|\nabla v\|_{L^{\infty}(\mathbb{R}^n)} \leq C$. This allows us to use Lemma A.1 with $\widehat{K} := \|\nabla v\|_{L^{\infty}(\mathbb{R}^n)}$ and $\rho := 1$, obtaining that, for every $x \in B_r$,

$$-(-\Delta)_p^s v(x) \leqslant \frac{C}{s(1-s)}.$$

This concludes the proof of (3.24) for the case $p \in (1, 2)$ and $s \in \left(0, \frac{p-1}{2p}\right)$. Now, we let $p \in [2, +\infty)$ and $s \in (0, 1)$. In this case, recalling (3.6), we have that k = 1 and

$$v(y) := \begin{cases} 0 & \text{if } y \in \overline{B_{\frac{r}{2}}}, \\ (r - |y|)^{-qs} - \left(\frac{r}{2}\right)^{-qs} - qs\left(\frac{r}{2}\right)^{-qs-1}\left(|y| - \frac{r}{2}\right) & \text{if } y \in B_{\hat{r}} \setminus \overline{B_{\frac{r}{2}}} \\ 1 & \text{if } y \in \mathbb{R}^n \setminus \overline{B}_{\hat{r}} \end{cases}$$

where \hat{r} has been defined in (3.25). We can thereby compute that

$$\nabla v(y) = \begin{cases} 0 & \text{if } y \in \overline{B_{\frac{r}{2}}}, \\ \frac{y}{|y|} \left(qs(r-|y|)^{-qs-1} - qs\left(\frac{r}{2}\right)^{-qs-1} \right) & \text{if } y \in B_{\hat{r}} \setminus \overline{B_{\frac{r}{2}}}, \\ 0 & \text{if } y \in \mathbb{R}^n \setminus B_{\hat{r}}. \end{cases}$$
(3.30)

Consequently, for every $y \in B_r$,

$$|\nabla v(y)| \leqslant qs \left((r-|y|)^{-qs-1} - \left(\frac{r}{2}\right)^{-qs-1} \right) \chi_{B_{\hat{r}} \setminus \overline{B_{\frac{r}{2}}}} \leqslant C \left(r-|y|\right)^{-qs-1} \chi_{B_{\hat{r}} \setminus \overline{B_{\frac{r}{2}}}}.$$
 (3.31)

From this and (3.28) we infer that

$$\left\|\nabla v\right\|_{L^{\infty}\left(B_{\frac{r-|x|}{2}}(x)\right)} \leqslant C\left(r-|x|\right)^{-qs-1}.$$
(3.32)

Also, (3.31) and Proposition A.6 give that

$$\|\nabla v\|_{L^{\infty}(\mathbb{R}^n)} \leqslant C \left(r - \hat{r}\right)^{-qs-1} \leqslant C.$$
(3.33)

Moreover, for every $y \in B_{\hat{r}} \setminus B_{\frac{r}{2}}$ and $\mu, j \in \{1, \ldots, n\}$,

$$D_{j,\mu}^2 v(y) = qs \left(\frac{\delta_{\mu,j}}{|y|} - \frac{y_{\mu}y_j}{|y|^3}\right) \left((r - |y|)^{-qs-1} - \left(\frac{r}{2}\right)^{-qs-1}\right) + \frac{y_{\mu}y_j}{|y|^2} qs(qs+1)(r - |y|)^{-qs-2},$$

and thus, for every $y \in B_{\hat{r}} \setminus \overline{B}_{\frac{r}{2}}$,

$$\left|D^{2}v(y)\right| \leq C\left(\frac{\left(r-|y|\right)^{-qs-1}}{|y|} + \left(r-|y|\right)^{-qs-2}\right) \leq C\left(r-|y|\right)^{-qs-2}.$$
(3.34)

Accordingly, making use of this and (3.28),

$$\left\| D^{2} v \right\|_{L^{\infty} \left(B_{\frac{r-|x|}{2}}(x) \cap B_{\hat{r}} \right)} \leq C \left(r-|x| \right)^{-qs-2}.$$
(3.35)

In addition, we deduce from (3.34) and Proposition A.6 that

$$|D^2 v||_{L^{\infty}(B_{\hat{r}})} \leq C (r - \hat{r})^{-qs-2} \leq C.$$
 (3.36)

Now, we set

$$\Sigma(y) := \begin{cases} \nabla v(y) & \text{if } y \in B_{\hat{r}}, \\ 0 & \text{if } y \in \mathbb{R}^n \setminus B_{\hat{r}} \end{cases}$$

and we claim that there exists $C \in (0, +\infty)$ such that, for every $y \in B_{\frac{r-|x|}{2}}(x)$,

$$v(y) - v(x) - \Sigma(x) \cdot (y - x) \leq C \left(r - |x|\right)^{-qs-2} |x - y|^2$$
(3.37)

and, for every $y \in \mathbb{R}^n$,

$$v(y) - v(x) - \Sigma(x) \cdot (y - x) \leq C |x - y|^2.$$
 (3.38)

To check (3.37) and (3.38) we proceed as follows. If $x \in B_r \setminus B_{\hat{r}}$, the left-hand sides of both (3.37) and (3.38) are non positive, so the inequalities trivially hold. If instead $x, y \in B_{\hat{r}}$ the inequality in (3.38) follows from (3.36). Analogously, if $x \in B_{\hat{r}}$ and $y \in B_{\hat{r}} \cap B_{\frac{r-|x|}{2}}(x)$ the inequality in (3.37) follows from (3.35).

Now we consider a radial smooth extension \bar{v} of v outside $B_{\hat{r}}$ such that $1 \leq \bar{v} \leq 2$ and $D^2 \bar{v}(y) \leq C$ for every $y \in \mathbb{R}^n \setminus B_{\hat{r}}$. Then, from (3.36), we have that if $x \in B_{\hat{r}}$, for every $y \in \mathbb{R}^n \setminus B_{\hat{r}}$,

$$v(y) - v(x) - \Sigma(x) \cdot (y - x) = 1 - \overline{v}(x) - \nabla \overline{v}(x) \cdot (y - x)$$

$$\leqslant \overline{v}(y) - \overline{v}(x) - \nabla \overline{v}(x) \cdot (y - x)$$

$$\leqslant C |x - y|^2.$$

This concludes the proof of (3.38).

Similarly, if we define the extension

$$\hat{v}(y) := \begin{cases} v(y) & \text{if } y \in B_{\hat{r}}, \\ f(|y|) & \text{if } y \in B_r \setminus B_{\hat{r}}, \\ 1 & \text{if } y \in \mathbb{R}^n \setminus B_r, \end{cases}$$

we notice that, by (3.35) and the definition of f in (3.7), if $x \in B_{\hat{r}}$,

$$\left\|D^{2}\hat{v}\right\|_{L^{\infty}\left(B_{\frac{r-|x|}{2}}(x)\right)} \leqslant C\left(r-|x|\right)^{-qs-2}.$$

Then, for every $y \in B_{\frac{r-|x|}{2}}(x) \setminus B_{\hat{r}}$,

$$\begin{aligned} v(y) - v(x) - \Sigma(x) \cdot (y - x) &= 1 - \hat{v}(x) - \nabla \hat{v}(x) \cdot (y - x) \\ &\leq \hat{v}(y) - \hat{v}(x) - \nabla \hat{v}(x) \cdot (y - x) \\ &\leq C \left(r - |x|\right)^{-qs-2} |x - y|^2 \,. \end{aligned}$$

This concludes the proof of (3.37).

Now, we apply Lemma A.2 with

$$\rho := \frac{r - |x|}{2}, \quad \widehat{K} := \|\nabla v\|_{L^{\infty}\left(B_{\frac{r - |x|}{2}}(x)\right)} \quad \text{and} \quad \widetilde{C} := C \left(r - |x|\right)^{-qs-2}$$

and obtain, thanks to (3.32) and (3.37), that, if $x \in B_r$,

$$\begin{split} &-(-\Delta)_{p}^{s}v(x) \\ &\leqslant \frac{c_{n,p}}{s(1-s)} \left(\widetilde{C} \max\left\{ \left\| \nabla v \right\|_{L^{\infty}\left(B_{\frac{r-|x|}{2}}(x)\right)}, \left| \Sigma(x) \right| \right\}^{p-2} (r-|x|)^{-sp+p-qs-2} + (r-|x|)^{-sp} \right) \\ &\leqslant \frac{c_{n,p}}{1-s} \left(\widetilde{C} \left\| \nabla v \right\|_{L^{\infty}\left(B_{\frac{r-|x|}{2}}(x)\right)}^{p-2} (r-|x|)^{-sp+p-qs-2} + (r-|x|)^{-sp} \right) \\ &\leqslant \frac{C}{1-s} \left((r-|x|)^{-(qs+1)(p-2)-sp+p-qs-2} + (r-|x|)^{-sp} \right) \\ &= \frac{C}{1-s} \left((r-|x|)^{-qs(p-1)-sp} + (r-|x|)^{-sp} \right). \end{split}$$

This concludes the proof of (3.23) when $p \in [2, +\infty)$ and $s \in \left[\frac{p-1}{p}, 1\right)$.

Finally, applying Lemma A.2 with

$$\rho := 1 \quad \text{and} \quad \widehat{K} := \|\nabla v\|_{L^{\infty}(\mathbb{R}^n)},$$

and recalling also equations (3.33) and (3.38), we obtain that, if $x \in B_r$,

$$-(-\Delta)_p^s v(x) \leqslant \frac{C}{1-s}.$$

This concludes the proof of (3.24) when $p \in [2, +\infty)$ and $s \in (0, 1)$.

With the work done so far, we can now complete the construction of the barrier in Theorem 3.1. *Proof of Theorem 3.1.* Let $\bar{c}_{k,q,s}$ and \tilde{C} be as in Propositions 3.2 and 3.4 respectively. We define the constants

$$C_0 := \left(\frac{\widetilde{C}}{s(1-s)\tau}\right)^{\frac{1}{ps}} \quad \text{and} \quad \beta := 2\overline{c}_{k,q,s}r^{-qs}. \tag{3.39}$$

Also, we rescale and translate v as follows

$$w(x) := (2 - \beta)v\left(\frac{x}{C_0}\right) + \beta - 1.$$
(3.40)

Moreover, we set

$$r := \frac{R}{C_0} \tag{3.41}$$

and we define the constants

$$\widehat{C}_{n,p,m} := \widehat{c}_{p,m}^{p} \widetilde{C} \quad \text{and} \quad \bar{R}_{s,\tau,n,p,m} := \left(\frac{\widehat{C}_{n,p,m}}{s(1-s)\tau}\right)^{\frac{1}{ps}},$$
(3.42)

where $\hat{c}_{p,m} \in (1, +\infty)$ is provided in (3.17).

We prove that w, as defined in (3.40), satisfies the properties listed in the statement of Theorem 3.1.

We first observe that, for every $R \in [\bar{R}_{s,\tau,n,p,m}, +\infty)$, if $x \in \mathbb{R}^n \setminus B_R$ then

$$\frac{|x|}{C_0} \ge \frac{R}{C_0} = r$$

and so, recalling the definition of v in (3.8), we have that

$$w(x) = (2 - \beta)v\left(\frac{x}{C_0}\right) + \beta - 1 = 2 - \beta + \beta - 1 = 1.$$

This establishes (3.2).

We now check that (3.5) holds true for every $R \in [\bar{R}_{s,\tau,n,p,m}, +\infty)$. For this, we make use of Lemma 3.3 (recall also that $\hat{c}_{p,m} > 1$ thanks to (3.17)) to deduce that

$$R \geqslant \bar{R}_{s,\tau,n,p,m} = C_0 \hat{c}_{p,m}^{\frac{1}{s}} \geqslant C_0 \hat{c}_{p,m} \geqslant C_0 \bar{c}_{k,q,s}^{\frac{1}{q_s}}.$$

In light of this, (3.39) and (3.41) we obtain that

$$\beta = 2\bar{c}_{k,q,s} \left(\frac{C_0}{R}\right)^{qs} \leqslant 2$$

From this and (3.40) we infer that

$$-1 < w \leqslant 1. \tag{3.43}$$

Moreover, recalling the definitions in (3.7) and (3.8), for every $t \in \left(\frac{r}{2}, r\right)$, we have that

$$h(t) \leq f(t)$$

$$= g(r-t) + \sum_{i=0}^{k} \frac{(-1)^{i+1}}{i!} g^{(i)} \left(\frac{r}{2}\right) \left(t - \frac{r}{2}\right)^{i}$$

$$\leq (r-t)^{-qs} + \sum_{i=0}^{k} \frac{\left|g^{(i)}(r/2)\right|}{i!} \left(t - \frac{r}{2}\right)^{i}$$

$$\leq (r-t)^{-qs} + \sum_{i=0}^{k} c_{q,s,i} \left(\frac{r}{2}\right)^{-qs-i} \left(\frac{r}{2}\right)^{i}$$

$$= (r-t)^{-qs} + \tilde{c}_{k,q,s} r^{-qs},$$

where $c_{q,s,i}$ and $\tilde{c}_{k,q,s}$ have been defined respectively in (3.11) and (3.12).

From this we deduce that, for every $x \in B_r$,

$$v(x) \leqslant (r - |x|)^{-qs} + \tilde{c}_{k,q,s} r^{-qs}$$

For this reason, for any $x \in B_R$,

$$1 + w(x) \leq 2v \left(\frac{x}{C_0}\right) + \beta \leq 2C_0^{qs} (R - |x|)^{-qs} + 2\tilde{c}_{k,q,s} r^{-qs} + \beta$$

= $2C_0^{qs} (R - |x|)^{-qs} + 2\tilde{c}_{k,q,s} r^{-qs} + 2\bar{c}_{k,q,s} r^{-qs}.$ (3.44)

Since

$$r = \frac{R}{C_0} \geqslant \frac{R - |x|}{C_0},$$

we obtain from (3.44) that, for all $x \in B_R$,

$$1 + w(x) \leq 2C_0^{qs}(R - |x|)^{-qs} + 2\tilde{c}_{k,q,s}C_0^{qs}(R - |x|)^{-qs} + 2\bar{c}_{k,q,s}C_0^{qs}(R - |x|)^{-qs} \leq C_1(R - |x|)^{-qs},$$
(3.45)

where $C_1 := 2C_0^{qs} \max\{1, \tilde{c}_{k,q,s}, \bar{c}_{k,q,s}\} = 2C_0^{qs} \bar{c}_{k,q,s}$, in light of (3.16). Now, if $x \in B_{R-1}$ then $R - |x| \ge \frac{1+R-|x|}{2}$. Thus, by (3.45),

$$1 + w(x) \leqslant 2^{qs+1} C_0^{qs} \bar{c}_{k,q,s} (1 + R - |x|)^{-qs}.$$
(3.46)

Additionally, if $x \in B_R \setminus B_{R-1}$, then $R+1-|x| \leq 2$. Thus, by (3.43),

$$1 + w(x) \leq 2 = 2^{qs+1} 2^{-qs} \leq 2^{qs+1} (1 + R - |x|)^{-qs}.$$
(3.47)

From (3.46) and (3.47), and recalling Lemma 3.3, we see that, for every $x \in B_R$,

$$\begin{split} 1+w(x) &\leqslant 2^{qs+1} \max\left\{1, C_0^{qs} \bar{c}_{k,q,s}\right\} (1+R-|x|)^{-qs} \\ &\leqslant 2^{q+1} \max\left\{1, C_0^{qs} \hat{c}_{p,m}^{qs}\right\} (1+R-|x|)^{-qs} \\ &\leqslant 2^{q+1} \max\left\{1, C_0^{qs} \hat{c}_{p,m}^{q}\right\} (1+R-|x|)^{-qs} \\ &= 2^{q+1} \max\left\{1, \bar{R}_{s,\tau,n,p,m}^{qs}\right\} (1+R-|x|)^{-qs} \,, \end{split}$$

which proves (3.5).

Now we prove (3.3). For this, we use Lemma 3.4 and equation (3.39) and we see that, for every $x \in B_R$,

$$\begin{aligned} -(-\Delta)_{p}^{s}w(x) &= -(2-\beta)^{p-1}C_{0}^{-sp}(-\Delta)_{p}^{s}v\left(\frac{x}{C_{0}}\right) \\ &\leqslant (2-\beta)^{p-1}C_{0}^{-sp}\frac{\widetilde{C}}{s(1-s)}\left(v\left(\frac{x}{C_{0}}\right) + \bar{c}_{k,q,s}r^{-qs}\right)^{m-1} \\ &= \tau(2-\beta)^{p-1}\left(v\left(\frac{x}{C_{0}}\right) + \bar{c}_{k,q,s}r^{-qs}\right)^{m-1} \\ &\leqslant \tau(2-\beta)^{m-1}\left(v\left(\frac{x}{C_{0}}\right) + \bar{c}_{k,q,s}r^{-qs}\right)^{m-1} \\ &\leqslant \tau\left((2-\beta)v\left(\frac{x}{C_{0}}\right) + (2-\beta)\bar{c}_{k,q,s}r^{-qs}\right)^{m-1} \\ &\leqslant \tau\left((2-\beta)v\left(\frac{x}{C_{0}}\right) + 2\bar{c}_{k,q,s}r^{-qs}\right)^{m-1} \\ &= \tau\left(1+w(x)\right)^{m-1}. \end{aligned}$$

This establishes the claim in (3.3) and completes the proof of Theorem 3.1.

4 Proofs of Theorem 1.3 and Corollary 1.5

In the section we prove the density estimates in Theorem 1.3 and Corollary 1.5.

We begin with the following observation:

Remark 4.1. We notice that in (1.16) we can assume, without loss of generality, that $c_0 \in (1, +\infty)$. Indeed, if $c_0 \in (0, 1]$, we can choose some $\tilde{c}_0 \in (1, +\infty)$ and for $t := (c_0/\tilde{c}_0)^{\frac{1}{n}}$ we can consider the rescaled function

$$\tilde{u}(x) := u(tx).$$

Then, if we define $\tilde{r}_0 := r_0/t$, by scaling we obtain that

$$\mathcal{L}^n\left(B_{\tilde{r}_0}\cap\{\tilde{u}>\theta_1\}\right)=\frac{1}{t^n}\mathcal{L}^n\left(B_{r_0}\cap\{u>\theta_1\}\right)>\frac{c_0}{t^n}=\tilde{c}_0.$$

As a consequence, being \tilde{u} a minimizer² for $\mathcal{E}_{s}^{p}(\cdot, \Omega_{1/t})$ with potential $\widetilde{W} := t^{sp}W$, according to (1.17) we obtain the existence of some R^{*} and \tilde{c} , depending only on $n, s, p, c_{0}, W, r_{0}, \theta_{1}, \theta_{2}$ and \tilde{c}_{0} , such that, for any $r \geq R^{*}$ satisfying $B_{\frac{3}{2}r} \subset \Omega_{1/t}$,

$$\tilde{c}r^{n} < c_{m,\theta_{*}} \int_{B_{r} \cap \{\theta_{*} < \tilde{u} \le \theta^{*}\}} |1 + \tilde{u}(x)|^{m} dx + \mathcal{L}^{n}(B_{r} \cap \{\tilde{u} > \theta_{2}\})$$
$$= \frac{c_{m,\theta_{*}}}{t^{n}} \int_{B_{tr} \cap \{\theta_{*} < u \le \theta^{*}\}} |1 + u(x)|^{m} dx + \frac{1}{t^{n}} \mathcal{L}^{n}(B_{tr} \cap \{u > \theta_{2}\})$$

thus proving the density estimates for u with constants R^*t and \tilde{c} .

Proof of Theorem 1.3. Thanks to Remark 4.1, we can assume, without loss of generality, that

$$c_0 \in (1, +\infty). \tag{4.1}$$

Let $R \in (0, +\infty)$ such that $B_R \subset \Omega$ and $w \in C(\mathbb{R}^n, (-1, 1])$ such that $w \equiv 1$ in $\mathbb{R}^n \setminus B_R$. A more specific choice of w will be made later on in the proof (namely w will be the barrier constructed in Theorem 3.1 with a suitable choice of the parameter τ).

Moreover, we define

$$v := \min\left\{u, w\right\}$$

and we observe that

$$v = u \quad \text{in} \quad \mathbb{R}^n \setminus B_R. \tag{4.2}$$

To ease notation, we also set

$$D := (\mathbb{R}^n \times \mathbb{R}^n) \setminus ((\mathbb{R}^n \setminus B_R) \times (\mathbb{R}^n \setminus B_R)).$$

In this way, recalling the definition of the interaction energy \mathcal{K}_s^p in (1.11), we see that, for every $f : \mathbb{R}^n \to \mathbb{R}$,

$$\mathcal{K}_{s}^{p}(f, B_{R}) = \frac{1}{2} \int_{D} \frac{|f(x) - f(y)|^{p}}{|x - y|^{n + sp}} \, dx \, dy.$$
(4.3)

²In this paper, for any r > 0, we use the notation

$$\Omega_r := r\Omega := \big\{ rx \text{ with } x \in \Omega \big\}.$$

Now we introduce the following exponents

$$\hat{p} := \max\{2, p\}$$
 and $l := \min\{2, p\}$. (4.4)

We claim that, for $\tau := \frac{c_1}{p}$, $\bar{R}_{\tau} := \bar{R}_{s,\tau,n,p,m} \in (0, +\infty)$ given in (3.1) and $w \in C(\mathbb{R}^n, (-1, 1])$ given by Theorem 3.1, there exists $\hat{c}_p \in (0, +\infty)$ such that, for every $R \in [\bar{R}_{\tau}, +\infty)$ satisfying $B_R \subset \Omega$,

$$\mathcal{K}_{\frac{sl}{2}}^{\hat{p}}(u-v,B_R) \leqslant \frac{\Lambda + 2c_1}{\hat{c}_p} \int_{B_R \cap \{u > \max\{\theta_*,w\}\}} |1+w(x)|^{m-1} dx.$$
(4.5)

Here above, Λ is given in (1.9) and c_1 in (1.10).

To prove this, if $p \in [2, +\infty)$ we use formula (4.3) in [Lin90] to deduce that, for all $x, y \in \mathbb{R}^n$,

$$\frac{1}{2^{p-1}-1} |(u-v)(x) - (u-v)(y)|^p + |v(x) - v(y)|^p - |u(x) - u(y)|^p \\ \leqslant p(v(x) - v(y)) |v(x) - v(y)|^{p-2} (v(x) - v(y) - (u(x) - u(y))),$$
(4.6)

while if $p \in (1,2)$ we use formula (4.4) in [Lin90] to obtain that, for all $x, y \in \mathbb{R}^n$,

$$\frac{3p(p-1)}{4^2} \frac{|(u-v)(x) - (u-v)(y)|^2}{(|v(x) - v(y)| + |u(x) - u(y)|)^{2-p}} + |v(x) - v(y)|^p - |u(x) - u(y)|^p}{\leqslant p(v(x) - v(y)) |v(x) - v(y)|^{p-2} (v(x) - v(y) - (u(x) - u(y)))}.$$

Since $\|v\|_{L^{\infty}(\mathbb{R}^n)}$, $\|u\|_{L^{\infty}(\mathbb{R}^n)} \leq 1$, from the last equation we obtain that, for every $p \in (1, 2)$ and $x, y \in \mathbb{R}^n$,

$$\frac{3p(p-1)}{4^{4-p}} |(u-v)(x) - (u-v)(y)|^2 + |v(x) - v(y)|^p - |u(x) - u(y)|^p \\ \leqslant p(v(x) - v(y)) |v(x) - v(y)|^{p-2} (v(x) - v(y) - (u(x) - u(y))).$$

$$(4.7)$$

Therefore, we define the constant

$$\hat{c}_p := \begin{cases} \frac{1}{2^{p-1}} & \text{if } p \in [2, +\infty), \\ \frac{3p(p-1)}{4^{4-p}} & \text{if } p \in (1, 2) \end{cases}$$

and we infer from (4.6) and (4.7) that, for every $p \in (1, +\infty)$ and for every $x, y \in \mathbb{R}^n$,

$$\hat{c}_{p} |(u-v)(x) - (u-v)(y)|^{\hat{p}} + |v(x) - v(y)|^{p} - |u(x) - u(y)|^{p} \\ \leqslant p(v(x) - v(y)) |v(x) - v(y)|^{p-2} (v(x) - v(y) - (u(x) - u(y))).$$

$$(4.8)$$

Now, recalling (4.3), we obtain that

$$\hat{c}_{p}\mathcal{K}_{\frac{s_{l}}{2}}^{\hat{p}}(u-v,B_{R}) + \mathcal{K}_{s}^{p}(v,B_{R}) - \mathcal{K}_{s}^{p}(u,B_{R})$$

$$= \frac{\hat{c}_{p}}{2} \int_{D} \frac{\left|(u-v)(x) - (u-v)(y)\right|^{\hat{p}}}{\left|x-y\right|^{n+\hat{p}\frac{s_{l}}{2}}} dx dy$$

$$+ \frac{1}{2} \int_{D} \frac{\left|v(x) - v(y)\right|^{p}}{\left|x-y\right|^{n+sp}} dx dy - \frac{1}{2} \int_{D} \frac{\left|u(x) - u(y)\right|^{p}}{\left|x-y\right|^{n+sp}} dx dy.$$

We remark that, in light of (4.4),

$$\hat{p}\frac{sl}{2} = \frac{2sp}{2} = sp,$$

and therefore

$$\hat{c}_{p}\mathcal{K}_{\frac{sl}{2}}^{\hat{p}}(u-v,B_{R}) + \mathcal{K}_{s}^{p}(v,B_{R}) - \mathcal{K}_{s}^{p}(u,B_{R})$$

$$= \frac{\hat{c}_{p}}{2} \int_{D} \frac{|(u-v)(x) - (u-v)(y)|^{\hat{p}}}{|x-y|^{n+sp}} dx dy$$

$$+ \frac{1}{2} \int_{D} \frac{|v(x) - v(y)|^{p}}{|x-y|^{n+sp}} dx dy - \frac{1}{2} \int_{D} \frac{|u(x) - u(y)|^{p}}{|x-y|^{n+sp}} dx dy.$$

Employing (4.8) and then recalling (4.2) we thereby obtain that

$$\begin{aligned} \hat{c}_{p} \mathcal{K}_{\frac{sl}{2}}^{\hat{p}}(u-v,B_{R}) + \mathcal{K}_{s}^{p}(v,B_{R}) - \mathcal{K}_{s}^{p}(u,B_{R}) \\ \leqslant \quad \frac{p}{2} \int_{D} \frac{\left(v(x)-v(y)\right) |v(x)-v(y)|^{p-2} \left(v(x)-v(y)-(u(x)-u(y))\right)}{|x-y|^{n+sp}} \, dx \, dy \\ = \quad \frac{p}{2} \int_{\mathbb{R}^{n} \times \mathbb{R}^{n}} \frac{\left(v(x)-v(y)\right) |v(x)-v(y)|^{p-2} \left(v(x)-v(y)-(u(x)-u(y))\right)}{|x-y|^{n+sp}} \, dx \, dy \\ = \quad p \int_{\mathbb{R}^{n} \times \mathbb{R}^{n}} \frac{\left(v(y)-v(x)\right) |v(x)-v(y)|^{p-2} \left(u(x)-v(x)\right)}{|x-y|^{n+sp}} \, dx \, dy. \end{aligned}$$

As a result, making again use of (4.2) and of the fact that $w \equiv 1$ in $\mathbb{R}^n \setminus B_R$,

$$\hat{c}_{p}\mathcal{K}_{\frac{sl}{2}}^{\hat{p}}(u-v,B_{R}) + \mathcal{K}_{s}^{p}(v,B_{R}) - \mathcal{K}_{s}^{p}(u,B_{R}) \\
\leq p \int_{\mathbb{R}^{n}} \left((u(x)-v(x)) \int_{\mathbb{R}^{n}} \frac{(v(y)-v(x)) |v(x)-v(y)|^{p-2}}{|x-y|^{n+sp}} \, dy \right) \, dx \\
= p \int_{\{u>v=w\}} \left((u(x)-v(x)) \int_{\mathbb{R}^{n}} \frac{(v(y)-w(x)) |w(x)-v(y)|^{p-2}}{|x-y|^{n+sp}} \, dy \right) \, dx \\
= p \int_{B_{R} \cap \{u>v=w\}} \left((u(x)-v(x)) \int_{\mathbb{R}^{n}} \frac{(v(y)-w(x)) |w(x)-v(y)|^{p-2}}{|x-y|^{n+sp}} \, dy \right) \, dx.$$
(4.9)

Now, we define the function $\phi_p(t) := t|t|^{p-2}$ and we observe that ϕ_p is nondecreasing, since

$$\phi'_p(t) = |t|^{p-2} + t(p-2)|t|^{p-4}t = |t|^{p-2} + (p-2)|t|^{p-2} = (p-1)|t|^{p-2} \ge 0$$

As a consequence, since $v(y) \leq w(y)$, we see that

$$\phi_p(v(y) - w(x)) \leqslant \phi_p(w(y) - w(x)).$$

Plugging this information into (4.9), we thus find that

$$\hat{c}_{p}\mathcal{K}_{\frac{s_{l}}{2}}^{\hat{p}}(u-v,B_{R}) + \mathcal{K}_{s}^{p}(v,B_{R}) - \mathcal{K}_{s}^{p}(u,B_{R})$$

$$\leqslant p \int_{B_{R} \cap \{u > v = w\}} \left((u(x) - v(x)) \int_{\mathbb{R}^{n}} \frac{(w(y) - w(x)) |w(x) - w(y)|^{p-2}}{|x-y|^{n+sp}} \, dy \right) \, dx$$

$$= -p \int_{B_R \cap \{u > v = w\}} (u(x) - v(x)) (-\Delta)_p^s w(x) \, dx.$$

From this and the definition of the energy in (1.12) we obtain that

$$\hat{c}_p \mathcal{K}_{\frac{s_l}{2}}^{\hat{p}}(u-v, B_R) \leqslant \mathcal{E}(u, B_R) - \mathcal{E}(v, B_R) + \int_{B_R} W(v(x)) - W(u(x)) \, dx$$
$$- p \int_{B_R \cap \{u > v = w\}} (u(x) - v(x))(-\Delta)_p^s w(x) \, dx.$$

Thus, the minimality of u in B_R and (4.2) give that

$$\hat{c}_{p}\mathcal{K}_{\frac{sl}{2}}^{\hat{p}}(u-v,B_{R}) \\
\leqslant \int_{B_{R}} W(v(x)) - W(u(x)) \, dx - p \int_{B_{R} \cap \{u > v = w\}} (u(x) - v(x))(-\Delta)_{p}^{s} w(x) \, dx \\
= \int_{B_{R} \cap \{u > w\}} W(w(x)) - W(u(x)) \, dx - p \int_{B_{R} \cap \{u > v = w\}} (u(x) - v(x))(-\Delta)_{p}^{s} w(x) \, dx.$$
(4.10)

Now, we observe that, thanks to (1.10), if $x \in \{u > w\}$,

$$W(u(x)) - W(w(x)) = \int_{w(x)}^{u(x)} W'(\xi) \, d\xi \ge c_1 \int_{w(x)}^{u(x)} |1 + \xi|^{m-1} \, d\xi$$
$$\ge c_1 |1 + w(x)|^{m-1} (u(x) - w(x)).$$

Therefore, using also (1.9),

$$\int_{B_R \cap \{u > w\}} W(w(x)) - W(u(x)) \, dx$$

= $\int_{B_R \cap \{w < u \le \theta_*\}} W(w(x)) - W(u(x)) \, dx + \int_{B_R \cap \{u > \max\{\theta_*, w\}\}} W(w(x)) - W(u(x)) \, dx$
 $\leqslant -c_1 \int_{B_R \cap \{w < u \le \theta_*\}} |1 + w(x)|^{m-1} \left(u(x) - w(x)\right) \, dx + \Lambda \int_{B_R \cap \{u > \max\{\theta_*, w\}\}} |1 + w(x)|^m \, dx.$

Accordingly, using this fact into (4.10),

$$\begin{split} \hat{c}_{p} \mathcal{K}_{\frac{s_{l}}{2}}^{\hat{p}}(u-v,B_{R}) \\ &\leqslant -c_{1} \int_{B_{R} \cap \{w < u \leqslant \theta_{*}\}} |1+w(x)|^{m-1} (u(x)-w(x)) \, dx + \Lambda \int_{B_{R} \cap \{u > \max\{\theta_{*},w\}\}} |1+w(x)|^{m-1} \, dx \\ &- p \int_{B_{R} \cap \{u > v = w\}} (u(x)-v(x))(-\Delta)_{p}^{s} w(x) \, dx \\ &= -c_{1} \int_{B_{R} \cap \{u > w\}} |1+w(x)|^{m-1} (u(x)-w(x)) \, dx \\ &+ c_{1} \int_{B_{R} \cap \{u > \max\{w,\theta_{*}\}\}} |1+w(x)|^{m-1} (u(x)-w(x)) \, dx \\ &+ \Lambda \int_{B_{R} \cap \{u > \max\{\theta_{*},w\}\}} |1+w(x)|^{m-1} \, dx - p \int_{B_{R} \cap \{u > v = w\}} (u(x)-v(x))(-\Delta)_{p}^{s} w(x) \, dx. \end{split}$$

The computations done so far hold true for any function $w \in C(\mathbb{R}^n, (-1, 1])$ such that $w \equiv 1$ in $\mathbb{R}^n \setminus B_R$. We now choose w as in Theorem 3.1 with $\tau := \frac{c_1}{p}$. Then, if $\bar{R}_\tau := \bar{R}_{s,\tau,n,p,m} \in (0, +\infty)$ is given as in (3.1), thanks to Theorem 3.1 we obtain that, for every $R \in [\bar{R}_\tau, +\infty)$ such that $B_R \subset \Omega$,

$$\begin{split} \hat{c}_{p} \mathcal{K}_{\frac{s_{l}}{2}}^{\hat{p}}(u-v,B_{R}) \\ \leqslant & -c_{1} \int_{B_{R} \cap \{u>w\}} |1+w(x)|^{m-1} \left(u(x)-w(x)\right) dx \\ & +c_{1} \int_{B_{R} \cap \{u>\max\{w,\theta_{*}\}\}} |1+w(x)|^{m-1} \left(u(x)-w(x)\right) dx \\ & +\Lambda \int_{B_{R} \cap \{u>\max\{\theta_{*},w\}\}} |1+w(x)|^{m-1} dx + c_{1} \int_{B_{R} \cap \{u>v=w\}} \left(u(x)-v(x)\right) |1+w(x)|^{m-1} dx \\ = & \Lambda \int_{B_{R} \cap \{u>\max\{\theta_{*},w\}\}} |1+w(x)|^{m-1} dx + c_{1} \int_{B_{R} \cap \{u>\max\{w,\theta_{*}\}\}} \left(u(x)-v(x)\right) |1+w(x)|^{m-1} dx \\ \leqslant & (\Lambda+2c_{1}) \int_{B_{R} \cap \{u>\max\{\theta_{*},w\}\}} |1+w(x)|^{m-1} . \end{split}$$

This concludes the proof of the claim in (4.5).

Now, let $C_{\tau} = C_{s,\tau,n,p,m}$ be as in (3.4) with $\tau = \frac{c_1}{p}$, and define

$$k_0 := \left(\frac{2C_\tau}{1+\theta_*}\right)^{\frac{m-1}{ps}} - 1.$$

Note that by the definitions of C_{τ} and θ_* we have that

$$k_{0} \ge C_{\tau}^{\frac{m-1}{ps}} - 1 = \left(2^{\frac{p}{m-1}+1} \max\left\{1, \bar{R}_{\tau}^{\frac{ps}{m-1}}\right\}\right)^{\frac{m-1}{ps}} - 1$$

= $2^{\frac{1}{s} + \frac{m-1}{ps}} \max\left\{1, \bar{R}_{\tau}\right\} - 1 \ge \max\left\{1, \bar{R}_{\tau}\right\}.$ (4.11)

Thanks to this and (3.5), we have that if $R \in (k_0, +\infty)$ and $k \in (k_0, R)$, for all $x \in B_{R-k}$,

$$w(x) \leqslant -1 + \frac{C_{\tau}}{(1+R-|x|)^{\frac{ps}{m-1}}} \leqslant -1 + \frac{C_{\tau}}{(1+k)^{\frac{ps}{m-1}}} \leqslant -1 + \frac{1+\theta_*}{2}.$$

This gives that, for all $x \in B_{R-k} \cap \{u > \theta_*\},\$

$$|u(x) - v(x)| = u(x) - v(x) \ge u(x) - w(x) \ge \frac{1 + \theta_*}{2}.$$
(4.12)

Now, for every $t \in (0, +\infty)$ we denote by

$$V(t) := \mathcal{L}^n(B_t \cap \{u > \theta_*\}). \tag{4.13}$$

With this notation, using (4.12), we have that, for every $R \in (k_0, +\infty)$ and $k \in (k_0, R)$ satisfying $B_R \subset \Omega$,

$$\left(V(R-k)\right)^{\frac{n-sp}{n}} = \left(\int_{B_{R-k} \cap \{u > \theta_*\}} dx\right)^{\frac{n-sp}{n}}$$

$$\leqslant \left(\frac{2}{1+\theta_*}\right)^{\hat{p}} \left(\int_{B_{R-k}\cap\{u>\theta_*\}} |u(x)-v(x)|^{\frac{n\hat{p}}{n-sp}} dx\right)^{\frac{n-sp}{n}},$$

where \hat{p} has been defined in (4.4).

We also denote by $s' := \frac{sl}{2}$, with l given in (4.4), and we observe that

$$sp = \frac{2sp}{2} = \frac{\hat{p}sl}{2} = \hat{p}s'$$

Hence, we can write that

$$\left(V(R-k) \right)^{\frac{n-sp}{n}} \leq \left(\frac{2}{1+\theta_*} \right)^{\hat{p}} \left(\int_{B_{R-k} \cap \{u > \theta_*\}} |u(x) - v(x)|^{\frac{n\hat{p}}{n-\hat{p}s'}} dx \right)^{\frac{n-\hat{p}s'}{n}}$$

$$\leq \left(\frac{2}{1+\theta_*} \right)^{\hat{p}} \left(\int_{\mathbb{R}^n} |u(x) - v(x)|^{\frac{n\hat{p}}{n-\hat{p}s'}} dx \right)^{\frac{n-\hat{p}s'}{n}}.$$

$$(4.14)$$

Now, we recall the Sobolev inequality of Theorem 1 in [MS02], according to which there exists $c_{n,p} \in (0, +\infty)$ such that, for every $f \in W_0^{s,p}(\mathbb{R}^n)$,

$$\left(\int_{\mathbb{R}^n} |f(x)|^{\frac{pn}{n-sp}} dx\right)^{\frac{n-sp}{n}} \leqslant c_{n,p} \frac{s(1-s)}{(n-sp)^{p-1}} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|f(x)-f(y)|^p}{|x-y|^{n+sp}} dx dy.$$

From this inequality (used with f := u - v) and (4.14), we deduce that

$$\left(V(R-k)\right)^{\frac{n-sp}{n}} \leqslant c_{n,\hat{p}} \frac{s'(1-s')}{(n-s'\hat{p})^{\hat{p}-1}} \left(\frac{2}{1+\theta_*}\right)^{\hat{p}} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|(u-v)(x) - (u-v)(y)|^{\hat{p}}}{|x-y|^{n+s'\hat{p}}} \, dy \, dx.$$

Thus, making use of (4.2),

$$\left(V(R-k) \right)^{\frac{n-sp}{n}} \leq c_{n,\hat{p}} \frac{s'(1-s')}{(n-sp)^{\hat{p}-1}} \left(\frac{2}{1+\theta_*} \right)^{\hat{p}} \int_D \frac{\left| (u-v)(x) - (u-v)(y) \right|^{\hat{p}}}{|x-y|^{n+s'\hat{p}}} \, dy \, dx$$

$$= c_{n,\hat{p}} \frac{s'(1-s')}{(n-sp)^{\hat{p}-1}} \left(\frac{2}{1+\theta_*} \right)^{\hat{p}} \mathcal{K}_{s'}^{\hat{p}}(u-v, B_R).$$

This and (4.5) give that

$$\left(V(R-k)\right)^{\frac{n-sp}{n}} \leqslant c_{n,\hat{p}} \frac{s'(1-s')}{(n-sp)^{\hat{p}-1}} \left(\frac{2}{1+\theta_*}\right)^{\hat{p}} \frac{\Lambda + 2c_1}{\hat{c}_p} \int_{B_R \cap \{u > \max\{\theta_*, w\}\}} \left|1 + w(x)\right|^{m-1} dx.$$

Therefore, in light of (3.5),

$$\left(V(R-k) \right)^{\frac{n-sp}{n}} \leq c_{n,\hat{p}} \frac{s'(1-s')}{(n-sp)^{\hat{p}-1}} \left(\frac{2}{1+\theta_*} \right)^{\hat{p}} \frac{\Lambda + 2c_1}{\hat{c}_p} C_{\tau}^{m-1} \int_{B_R \cap \{u > \max\{\theta_*, w\}\}} (1+R-|x|)^{-ps} dx \\ \leq c_{n,\hat{p}} \frac{s'(1-s')}{(n-sp)^{\hat{p}-1}} \left(\frac{2}{1+\theta_*} \right)^{\hat{p}} \frac{\Lambda + 2c_1}{\hat{c}_p} C_{\frac{c_1}{p}}^{m-1} \int_{B_R \cap \{u > \theta_*\}} (1+R-|x|)^{-ps} dx.$$

As a result, using the coarea formula and setting

$$c = c_{s,n,p,m,\Lambda,c_1,\theta_*} := c_{n,\hat{p}} \frac{s'(1-s')}{(n-sp)^{\hat{p}-1}} \left(\frac{2}{1+\theta_*}\right)^{\hat{p}} \frac{\Lambda + 2c_1}{\hat{c}_p} C_{\frac{c_1}{p}}^{m-1},$$

we conclude that, for every $R \in (k_0, +\infty)$ and $k \in (k_0, R)$ satisfying $B_R \subset \Omega$,

$$\left(V(R-k)\right)^{\frac{n-sp}{n}} \leqslant c \int_0^R (1+R-|t|)^{-ps} V'(t) \, dt.$$
(4.15)

Now we integrate (4.15) in $R \in \left[\rho, \frac{3}{2}\rho\right]$ with $\rho \ge 4k$ and $B_{\frac{3}{2}\rho} \subset \Omega$. Also, we use the fact that $s \in \left(0, \frac{1}{p}\right)$ together with (4.11) and obtain that

$$\begin{split} \frac{\rho}{2} \big(V(\rho-k) \big)^{\frac{n-sp}{n}} &\leqslant \int_{\rho}^{\frac{3}{2}\rho} \big(V(R-k) \big)^{\frac{n-sp}{n}} dR \\ &\leqslant c \int_{\rho}^{\frac{3}{2}\rho} \left(\int_{0}^{R} (1+R-|t|)^{-sp} V'(t) \, dt \right) \, dR \\ &\leqslant c \int_{0}^{\frac{3}{2}\rho} \left(V'(t) \int_{t}^{\frac{3}{2}\rho} (1+R-|t|)^{-sp} \, dR \right) \, dt \\ &= \frac{c}{1-sp} \int_{0}^{\frac{3}{2}\rho} V'(t) \left[\left(1+\frac{3}{2}\rho-|t| \right)^{1-sp} - 1 \right] \, dt \\ &\leqslant \frac{c}{1-sp} 2^{1-sp} \rho^{1-sp} \int_{0}^{\frac{3}{2}\rho} V'(t) \, dt \\ &\leqslant \frac{2c}{1-sp} \rho^{1-sp} V\left(\frac{3}{2}\rho\right). \end{split}$$

In particular, setting $r := \rho - k$ and recalling that $\rho \ge 4k$ we deduce that

$$r^{sp}\left(V(r)\right)^{\frac{n-sp}{n}} \leqslant \hat{c}\,V(2r),\tag{4.16}$$

with

$$\hat{c} = \hat{c}_{s,n,p,m,\Lambda,c_1,\theta_*} := \frac{4c}{1-sp}$$

From now $k > k_0$ is fixed once and for all.

From (1.16), (4.1) and (4.16) we see that (2.1) and (2.2) in Lemma 2.1 hold true with V as in (4.13) and with the following choices

$$\sigma := sp, \quad \nu := n, \quad \gamma := 2, \quad \mu := c_0, \quad R_0 := \max\{3k, r_0\} \text{ and } C := \hat{c},$$

as far as $R \in [R_0, +\infty)$ is such that $B_{\frac{3R}{2}} \subset \Omega$.

Therefore, in virtue of (2.3) in Lemma 2.1, we obtain that there exist $\tilde{c} \in (0, 1)$, depending only on $s, n, p, m, \Lambda, c_1, \theta_*, c_0, R_0$ and k, and $R_* \in [R_0, +\infty)$, depending only on R_0 , such that, for every $r \in [R_*, +\infty)$ satisfying $B_{\frac{3r}{2}} \subset \Omega$,

$$V(r) \geqslant \tilde{c} r^n$$

As a result, for any $r \in [R_*, +\infty)$ such that $B_{\frac{3r}{2}} \subset \Omega$,

$$\tilde{c}r^n \leqslant \mathcal{L}^n(B_r \cap \{u > \theta_*\}) = \mathcal{L}^n(B_r \cap \{u > \theta^*\}) + \mathcal{L}^n(B_r \cap \{\theta_* < u \leqslant \theta^*\}).$$
(4.17)

Now we observe that

$$\mathcal{L}^{n}\left(B_{r} \cap \left\{\theta_{*} < u \leqslant \theta^{*}\right\}\right) = \int_{B_{r} \cap \left\{\theta_{*} < u \leqslant \theta^{*}\right\}} dx$$
$$\leqslant (1+\theta_{*})^{-m} \int_{B_{r} \cap \left\{\theta_{*} < u \leqslant \theta^{*}\right\}} |1+u(x)|^{m} dx.$$

Thus, it follows from this and (4.17) that, for every $r \in [R_*, +\infty)$ such that $B_{\frac{3r}{2}} \subset \Omega$,

$$\mathcal{L}^{n}(B_{r} \cap \{u > \theta^{*}\}) \ge c r^{n} - (1 + \theta_{*})^{-m} \int_{B_{r} \cap \{\theta_{*} < u \le \theta^{*}\}} |1 + u(x)|^{m} dx.$$

This entails the desired result in (1.17).

Proof of Corollary 1.5. Thanks to the claim (1.18) in Theorem 1.4 and the assumption in (1.19), we have that, for every $r \in (0, +\infty)$ such that $B_{r+2} \subset \Omega$,

$$\int_{B_r \cap \{\theta_* < u \leqslant \theta^*\}} |1 + u(x)|^m dx \leqslant \lambda_{\theta^*}^{-1} \int_{B_r \cap \{\theta_* < u \leqslant \theta^*\}} W(u(x)) dx
\leqslant \lambda_{\theta^*}^{-1} \mathcal{E}_s^p(u, B_r) \leqslant \lambda_{\theta^*}^{-1} \bar{C} r^{n-sp}.$$
(4.18)

Also, we observe that for every $r \ge 4$ it holds that $\frac{3}{2}r \ge r+2$. Hence, from (4.18) and (1.17) we deduce that, for every $r \ge \max\{4, R^*\}$ such that $B_{\frac{3}{2}r} \subset \Omega$,

$$\tilde{c} r^n \leqslant c_{m,\theta_*} \int_{B_r \cap \{\theta_* < u \leqslant \theta^*\}} |1 + u(x)|^m dx + \mathcal{L}^n(B_r \cap \{u > \theta_2\})$$
$$\leqslant \lambda_{\theta^*}^{-1} c_{m,\theta_*} \bar{C} r^{n-sp} + \mathcal{L}^n(B_r \cap \{u > \theta_2\}).$$

From this, the claim in (1.20) readily follows.

5 Proofs of Theorem 1.6 and Corollary 1.7

We begin this section by establishing the compactness result for sequences of functions $\{u_{\epsilon}\}_{\epsilon} \subset X^{s,p}(\Omega)$ satisfying the boundedness assumption (1.22), as stated in Theorem 1.6. As already mentioned in the introduction, the convergence is a straightforward consequence of the scaling of $\mathcal{F}_{s,\epsilon}^p$, the uniform bound in (1.22) and the compact embedding of $W^{s,p}(\Omega')$ into $L^1(\Omega')$ for $\Omega' \subseteq \Omega$. The details of the proof are as follows.

Proof of Theorem 1.6. As a consequence of (1.22), for every open and bounded set $\Omega' \subseteq \Omega$, we have that

$$\sup_{\epsilon \in (0,1)} \int_{\Omega'} \int_{\Omega'} \frac{\left| u_{\epsilon}(x) - u_{\epsilon}(y) \right|^p}{\left| x - y \right|^{n+sp}} \, dx \, dy < +\infty.$$
(5.1)

Moreover, since $u_{\epsilon} \in X^{s,p}(\Omega)$, we find that, for every $\epsilon \in (0,1)$,

$$\|u_{\epsilon}\|_{L^{p}(\Omega')} \leqslant |\Omega'|^{\frac{1}{p}}.$$
(5.2)

Г	1
	J

Thus, it follows from (5.1), (5.2) and Theorem 7.1 in [DNPV12] that there exists some $u_0 \in L^1(\Omega')$ such that, up to a subsequence, u_{ϵ} converges to u_0 in $L^1(\Omega')$ and pointwise a.e. in Ω' , as $\epsilon \to 0$.

Furthermore, using (1.21), (1.22) and Fatou's Lemma, we have that

$$0 = \lim_{\epsilon \to 0^+} \epsilon^{sp} \mathcal{F}_{\epsilon}(u_{\epsilon}, \Omega')$$

=
$$\lim_{\epsilon \to 0^+} \epsilon^{sp} \mathcal{K}_{s}^{p}(u_{\epsilon}, \Omega') + \int_{\Omega'} W(u_{\epsilon}(x)) dx$$

$$\geqslant c_{2} \lim_{\epsilon \to 0^+} \int_{\Omega'} (1 - u_{\epsilon}^{2}(x))^{m} dx$$

$$\geqslant c_{2} \int_{\Omega'} (1 - u_{0}^{2}(x))^{m} dx.$$

This yields that u_0 must be equal to -1 and 1 a.e. in Ω' . Accordingly, if we denote by

$$E := \{ x \in \Omega' \text{ s.t. } u_0(x) = 1 \}$$

we obtain that $u_0|_{\Omega'} = \chi_E - \chi_{E^c}$. The proof of Theorem 1.6 is thereby complete.

In what follows we prove the Hausdorff convergence of the interface stated in Corollary 1.7. This result follows from the density estimates in Corollary 1.5 together with the Hölder regularity result for minimizers of \mathcal{E}_s^p that we have recalled in Section 2.2.

Proof of Corollary 1.7. We argue by contradiction, and we assume that there exist $\Theta \in (0, 1), \delta \in (0, +\infty), R \in (0, +\infty)$ and a sequence $x_k \in \{|u_{\epsilon_k}| < \Theta\} \cap B_R$ such that either $B_{\delta}(x_k) \subset E$ or $B_{\delta}(x_k) \subset E^c$. Without loss of generality, we assume that for every k it holds that $B_{\delta}(x_k) \subset E^c$.

Then, according to Theorem 1.6, up to a subsequence we have that

$$0 = \lim_{k \to +\infty} \int_{B_{\delta}(x_k) \cap \Omega} |u_{\epsilon_k}(x) - \chi_E(x) + \chi_{E^c}(x)| dx$$

$$= \lim_{k \to +\infty} \int_{B_{\delta}(x_k) \cap \Omega} |u_{\epsilon_k}(x) + 1| dx.$$
(5.3)

Now, we define the rescaled set

$$\Omega_k := \left\{ \frac{x - x_k}{\epsilon_k} \text{ s.t. } x \in \Omega \right\},\,$$

and the sequence of functions

$$w_k(x) := u_{\epsilon_k} \left(x \epsilon_k + x_k \right).$$

In this way, w_k is a minimizer for \mathcal{E}_s^p in Ω_k . Moreover,

$$w_k(0) = u_{\epsilon_k}(x_k) > -\Theta. \tag{5.4}$$

Now, we apply Theorem 2.2 to w_k with the choice F(t) := (1 - s)W(t) and we obtain that $w_k \in C^{\alpha}_{loc}(\Omega_k)$ for some $\alpha \in (0, 1)$, depending only on n, s and p, and there exists $C \ge 1$, depending only

on n, s and p, such that, for every $R_0 \in \left(0, \frac{\operatorname{dist}(0, \partial \Omega_k)}{8}\right)$,

$$\begin{split} [w_k]_{C^{\alpha}(B_{R_0})} &\leqslant \frac{C}{R_0^{\alpha}} \left[\|w_k\|_{L^{\infty}(\mathbb{R}^n)} + \operatorname{Tail}(w_k; 0, 4R_0) + R_0^s (1-s)^{\frac{1}{p}} \|W\|_{L^{\infty}((-1,1))}^{\frac{1}{p}} \right] \\ &\leqslant \frac{C}{R_0^{\alpha}} \left[1 + \left((1-s)(4R_0)^{sp} \int_{\mathbb{R}^n \setminus B_{4R_0}} \frac{dy}{|y|^{n+sp}} \right)^{\frac{1}{p-1}} + R_0^s (1-s)^{\frac{1}{p}} \|W\|_{L^{\infty}((-1,1))}^{\frac{1}{p}} \right] \\ &= \frac{C}{R_0^{\alpha}} \left[1 + \left((1-s)\frac{\omega_{n-1}}{sp} \right)^{\frac{1}{p-1}} + R_0^s (1-s)^{\frac{1}{p}} \|W\|_{L^{\infty}((-1,1))}^{\frac{1}{p}} \right]. \end{split}$$
(5.5)

Furthermore, if we set $R_0 := \frac{\operatorname{dist}(\partial B_R, \partial \Omega)}{8} \in (0, +\infty)$, then $R_0 \in \left(0, \frac{\operatorname{dist}(0, \partial \Omega_k)}{8}\right)$ for every k. From this, (5.4) and (5.5), it follows that there exists r_0 such that $w_k(x) > -\Theta$ for every $x \in B_{r_0}$.

Therefore, we conclude that

$$\mathcal{L}^n(B_{r_0} \cap \{w_k > -\Theta\}) \geqslant \mathcal{L}^n(B_{r_0}) =: c_0.$$

We can thereby apply Corollary 1.5 and obtain that for any $\theta \in (-1, 1)$ there exist $\widetilde{R} := \widetilde{R}_{s,n,p,m,\theta_*,r_0,c_2} \in [r_0, +\infty)$ and $\hat{c} := \hat{c}_{s,n,p,m,\Lambda,c_1,\theta_*,r_0,c_0} \in (0, 1)$ such that, for every $r \in [\widetilde{R}, +\infty)$ satisfying $B_{\frac{3r}{2}} \subset \Omega_k$,

$$\mathcal{L}^n(B_r \cap \{w_k > \theta\}) \geqslant \hat{c} r^n.$$
(5.6)

Now, we choose $\theta := -\frac{1}{2}$ and $\delta_0 \in (0, \delta]$ such that $B_{\frac{3}{2}\delta_0}(x_k) \subset \Omega$, and we obtain from (5.3) and (5.6) that

$$0 = \lim_{k \to +\infty} \int_{B_{\delta}(x_k)} |1 + u_{\epsilon_k}(x)| dx$$

$$\geq \lim_{k \to +\infty} \int_{B_{\delta_0}(x_k)} |1 - u_{\epsilon_k}(x)| dx$$

$$= \lim_{k \to +\infty} \epsilon_k^n \int_{B_{\frac{\delta_0}{\epsilon_k}}} |1 + u_{\epsilon_k}(x\epsilon_k + x_k)| dx$$

$$= \lim_{k \to +\infty} \epsilon_k^n \int_{B_{\frac{\delta_0}{\epsilon_k}} \cap \{w_k > \theta\}} |1 + w_k(x)| dx$$

$$\geq \lim_{k \to +\infty} \epsilon_k^n \int_{B_{\frac{\delta_0}{\epsilon_k}} \cap \{w_k > \theta\}} |1 + w_k(x)| dx$$

$$\geq \frac{\hat{c}}{2} \delta_0^n,$$

thus providing a contradiction.

A Some technical results towards the proof of Theorem 3.1

In this section we collect some technical results that are used throughout Section 3 in order to prove Theorem 3.1.

We start with two estimates for the fractional p-Laplacian of a bounded and locally Lipschitz function.

Lemma A.1. Let $\rho \in (0, +\infty)$, $p \in (1, +\infty)$, $s \in \left(0, \frac{p-1}{p}\right)$ and $\psi \in L^{\infty}(\mathbb{R}^n)$.

Let $x \in \mathbb{R}^n$ and assume that there exists $\widehat{K} \in (0, +\infty)$ such that, for every $y \in B_{\rho}(x)$,

$$|\psi(y) - \psi(x)| \leqslant \widehat{K} |y - x|.$$

Then, there exists $c_{n,p} \in (0, +\infty)$ such that

$$\left| (-\Delta)_{p}^{s} \psi(x) \right| \leqslant \frac{c_{n,p}}{s(p-1-sp)} \left(\widehat{K}^{p-1} \rho^{-sp+p-1} + \|\psi\|_{L^{\infty}(\mathbb{R}^{n})}^{p-1} \rho^{-sp} \right).$$
(A.1)

Proof. We compute

$$\begin{split} & \left| (-\Delta)_{p}^{s} \psi(x) \right| \\ &= \left| \int_{B_{\rho}(x)} \frac{\left(\psi(y) - \psi(x) \right) |\psi(y) - \psi(x)|^{p-2}}{|x - y|^{n + sp}} \, dy + \int_{\mathbb{R}^{n} \setminus B_{\rho}(x)} \frac{\left(\psi(y) - \psi(x) \right) |\psi(y) - \psi(x)|^{p-2}}{|x - y|^{n + sp}} \, dy \right| \\ &\leqslant \widehat{K}^{p-1} \int_{B_{\rho}(x)} \frac{dy}{|x - y|^{n + sp - p + 1}} + 2^{p-1} \, \|\psi\|_{L^{\infty}(\mathbb{R}^{n})}^{p-1} \int_{\mathbb{R}^{n} \setminus B_{\rho}(x)} \frac{dy}{|x - y|^{n + sp}} \\ &= \frac{\widehat{K}^{p-1} \omega_{n-1}}{p - 1 - sp} \rho^{-sp + p - 1} + 2^{p-1} \, \|\psi\|_{L^{\infty}(\mathbb{R}^{n})}^{p-1} \frac{\omega_{n-1}}{sp} \rho^{-sp}. \end{split}$$

Now since sp < p-1 we conclude that

$$\begin{split} \left| (-\Delta)_{p}^{s} \psi(x) \right| &\leq \frac{(p-1)\widehat{K}^{p-1}\omega_{n-1}}{sp(p-1-sp)} \rho^{-sp+p-1} + 2^{p-1} \left\| \psi \right\|_{L^{\infty}(\mathbb{R}^{n})}^{p-1} \frac{(p-1)\omega_{n-1}}{sp(p-1-sp)} \rho^{-sp} \\ &\leq \frac{c_{n,p}}{s(p-1-sp)} \left(\widehat{K}^{p-1}\rho^{-sp+p-1} + \left\| \psi \right\|_{L^{\infty}(\mathbb{R}^{n})}^{p-1} \rho^{-sp} \right), \end{split}$$

which is the desired result.

Lemma A.2. Let $\rho \in (0, +\infty)$, $p \in [2, +\infty)_2$ $s \in (0, 1)$ and $\psi \in L^{\infty}(\mathbb{R}^n) \cap C(\mathbb{R}^n)$.

Let $x \in \mathbb{R}^n$ and assume that there exist $\widehat{K}, \widetilde{C} \in (0, +\infty)$ and $\Sigma(x) \in \mathbb{R}^n$ such that, for every $y \in B_{\rho}(x)$,

$$|\psi(x) - \psi(y)| \leq \widehat{K} |x - y| \quad and \quad \psi(y) - \psi(x) - \Sigma(x) \cdot (y - x) \leq \widetilde{C} |x - y|^2.$$
(A.2)

Then, there exists $c_{n,p} \in (0, +\infty)$ such that

$$-(-\Delta)_p^s\psi(x) \leqslant \frac{c_{n,p}}{s(1-s)} \left(\widetilde{C}\max\left\{\left|\Sigma(x)\right|, \widehat{K}\right\}^{p-2}\rho^{-sp+p} + \|\psi\|_{L^{\infty}(\mathbb{R}^n)}^{p-1}\rho^{-sp}\right)$$

Proof. We define the affine approximation of ψ at x as

$$\ell(y) := \psi(x) + \Sigma(x) \cdot (y - x).$$

Then, denoting $\phi_p(t) := t|t|^{p-2}$, we obtain that

$$P.V. \int_{B_{\rho}(x)} \frac{\phi_p \left(\ell(y) - \ell(x)\right)}{|x - y|^{n + sp}} \, dy = 0.$$

Consequently,

$$\begin{split} & \int_{\mathbb{R}^{n}} \frac{\phi_{p}\left(\psi(y) - \psi(x)\right)}{|x - y|^{n + sp}} \, dy \\ &= \int_{B_{\rho}(x)} \frac{\phi_{p}\left(\psi(y) - \psi(x)\right)}{|x - y|^{n + sp}} \, dy + \int_{\mathbb{R}^{n} \setminus B_{\rho}(x)} \frac{\phi_{p}\left(\psi(y) - \psi(x)\right)}{|x - y|^{n + sp}} \, dy \\ &= \int_{B_{\rho}(x)} \frac{\phi_{p}\left(\psi(y) - \psi(x)\right) - \phi_{p}\left(\ell(y) - \ell(x)\right)}{|x - y|^{n + sp}} \, dy + \int_{\mathbb{R}^{n} \setminus B_{\rho}(x)} \frac{\phi_{p}\left(\psi(y) - \psi(x)\right)}{|x - y|^{n + sp}} \, dy \\ &= : I + II. \end{split}$$

To estimate I, for every $y \in B_{\rho}(x)$ we denote by

$$m := \min \{ \psi(y) - \psi(x), \ell(y) - \ell(x) \} \quad \text{and} \quad M := \max \{ \psi(y) - \psi(x), \ell(y) - \ell(x) \}.$$
(A.3)

Then, thanks to the Mean Value Theorem and (A.2), we obtain that there exists some $\xi \in [m, M]$ such that

$$\phi_{p}(\psi(y) - \psi(x)) - \phi_{p}(\ell(y) - \ell(x)) = \phi_{p}'(\xi)(\psi(y) - \psi(x) - \ell(y) + \ell(x))
= \phi_{p}'(\xi)(\psi(y) - \ell(y))
= \phi_{p}'(\xi)(\psi(y) - \psi(x) - \Sigma(x) \cdot (y - x))
= (p - 1) |\xi|^{p-2} (\psi(y) - \psi(x) - \Sigma(x) \cdot (y - x))
\leqslant \widetilde{C}(p - 1) |\xi|^{p-2} |x - y|^{2}.$$
(A.4)

Also, according to (A.2) and (A.3),

$$|\xi| \leq \max\left\{ \left| \Sigma(x) \right|, \widehat{K} \right\} |x - y|,$$

and employing (A.4) we obtain that

$$\phi_p\left(\psi(y) - \psi(x)\right) - \phi_p\left(\ell(y) - \ell(x)\right) \leqslant \widetilde{C}(p-1) \max\left\{\left|\Sigma(x)\right|, \widehat{K}\right\}^{p-2} |x-y|^p$$

On this account, we have that

$$I \leqslant \widetilde{C}(p-1) \max\left\{ \left| \Sigma(x) \right|, \widehat{K} \right\}^{p-2} \int_{B_{\rho}(x)} \frac{dy}{|x-y|^{n+sp-p}}$$
$$= \widetilde{C}(p-1) \max\left\{ \left| \Sigma(x) \right|, \widehat{K} \right\}^{p-2} \frac{\omega_{n-1}}{p-sp} \rho^{-sp+p}.$$

We now estimate II, as follows

$$II \leqslant 2^{p-2} \|\psi\|_{L^{\infty}(\mathbb{R}^n)}^{p-1} \int_{\mathbb{R}^n} \frac{dy}{|x-y|^{n+sp}} = 2^{p-2} \|\psi\|_{L^{\infty}(\mathbb{R}^n)}^{p-1} \frac{\omega_{n-1}}{sp} \rho^{-sp}.$$

We thus conclude that

$$\begin{split} &-(-\Delta)_{p}^{s}\psi(x)\\ &\leqslant \widetilde{C}(p-1)\max\left\{\left|\Sigma(x)\right|,\widehat{K}\right\}^{p-2}\frac{\omega_{n-1}}{p-sp}\rho^{-sp+p}+2^{p-1}\left\|\psi\right\|_{L^{\infty}(\mathbb{R}^{n})}^{p-1}\frac{\omega_{n-1}}{sp}\rho^{-sp}\\ &=\widetilde{C}s(p-1)\max\left\{\left|\Sigma(x)\right|,\widehat{K}\right\}^{p-2}\frac{\omega_{n-1}}{sp(1-s)}\rho^{-sp+p}+2^{p-1}(1-s)\left\|\psi\right\|_{L^{\infty}(\mathbb{R}^{n})}^{p-1}\frac{\omega_{n-1}}{sp(1-s)}\rho^{-sp}\\ &\leqslant \frac{c_{n,p}}{s(1-s)}\left(\widetilde{C}\max\left\{\left|\Sigma(x)\right|,\widehat{K}\right\}^{p-2}\rho^{-sp+p}+\left\|\psi\right\|_{L^{\infty}(\mathbb{R}^{n})}^{p-1}\rho^{-sp}\right), \end{split}$$

as desired.

We now focus on the specific construction of the barrier in Theorem 3.1. We recall the setting in (3.6), (3.7) and (3.8) and we show the following:

Proposition A.3. Let $s \in (0,1)$ and $p \in (1,+\infty)$. Let k and f be respectively as in (3.6) and (3.7). Then, for every $i \in \{0,\ldots,k+1\}$ there exists $\widetilde{C}_{q,s,k,i} \in (0,+\infty)$ such that, if $\rho_0 \in (0,\frac{r}{2})$ and $t \in (\frac{r}{2}, r - \rho_0)$,

$$\widetilde{C}_{q,s,k,i}\left(\frac{r}{2}\right)^{-qs-k-1}\left(t-\frac{r}{2}\right)^{k+1-i} \leqslant f^{(i)}(t) \leqslant \widetilde{C}_{q,s,k,i}\,\rho_0^{-qs-k-1}\left(t-\frac{r}{2}\right)^{k+1-i}.$$
(A.5)

Also, for every $i \in \{0, \dots, k+1\}$ there exists $C_{i,q,s} \in (0, +\infty)$ such that, if $x \in \left(\frac{r}{2}, r\right)$ and $\rho \in \left(0, x - \frac{r}{2}\right)$, $\widetilde{C}_{q,s,k,i}(r - x + \rho)^{-qs-k-1}\rho^{k+1-i} \leq f^{(i)}(x) \leq C_{i,q,s}(r - x)^{-qs-i}$. (A.6)

$$(A.6)^{q_{s,k,i}}(r-x+\rho)^{-q_s-k-1}\rho^{k+1-i} \leqslant f^{(i)}(x) \leqslant C_{i,q,s}(r-x)^{-q_s-i}.$$

Proof. In order to prove (A.5) and (A.6), we notice that, owing to (3.7),

$$\begin{cases} f^{(i)}\left(\frac{r}{2}\right) = 0 & \text{for all } i \in \{0, 1, \dots, k\} \\ f^{(k+1)}(\eta) = (-1)^{(k+1)} g^{(k+1)}(r-\eta) & \text{for all } \eta \in \left(\frac{r}{2}, r\right]. \end{cases}$$
(A.7)

Furthermore, for every $\eta \in (0, +\infty)$ we have that

$$g^{(k+1)}(\eta) = (-1)^{k+1} \prod_{i=0}^{k} (qs+i)\eta^{-qs-k-1}.$$

From this, it follows that, given $0 < \rho_0 < \rho_1$, for all $\eta \in (\rho_0, \rho_1)$,

$$C_{q,s,k}\rho_1^{-qs-k-1} \leqslant (-1)^{(k+1)}g^{(k+1)}(\eta) \leqslant C_{q,s,k}\rho_0^{-qs-k-1}$$
(A.8)

where

$$C_{q,s,k} := \prod_{i=0}^{k} (qs+i).$$
 (A.9)

Gathering these pieces of information, we deduce that, for every $t \in \left(\frac{r}{2}, r - \rho_0\right)$,

$$C_{q,s,k}\left(\frac{r}{2}\right)^{-qs-k-1} \leqslant f^{(k+1)}(t) \leqslant C_{q,s,k}\rho_0^{-qs-k-1}.$$

From this we obtain that, for every $i \in \{0, \ldots, k\}$ and $y \in \left(\frac{r}{2}, r - \rho_0\right)$,

$$f^{(i)}(y) = \int_{\frac{r}{2}}^{y} f^{(i+1)}(t_1) dt_1$$

= $\int_{\frac{r}{2}}^{y} \int_{\frac{r}{2}}^{t_1} f^{(i+2)}(t_2) dt_2 dt_1$
= $\int_{\frac{r}{2}}^{y} \int_{\frac{r}{2}}^{t_1} \dots \int_{\frac{r}{2}}^{t_{k-i}} f^{(k+1)}(t_{k+1-i}) dt_{k+1-i} \dots dt_2 dt_1$
 $\geqslant \widetilde{C}_{q,s,k,i} \left(\frac{r}{2}\right)^{-qs-k-1} \left(y - \frac{r}{2}\right)^{k+1-i},$

where

$$\widetilde{C}_{q,s,k,i} := \frac{C_{q,s,k}}{(k+1-i)!}.$$
(A.10)

Similarly, one can show that for every $i \in \{0, ..., k\}$ and $y \in (\frac{r}{2}, r - \rho_0)$

$$f^{(i)}(y) \leqslant \widetilde{C}_{q,s,k,i} \rho_0^{-qs-k-1} \left(y - \frac{r}{2}\right)^{k+1-i}.$$

This concludes the proof of (A.5).

Now we prove (A.6). To do so, we make use of (A.7) and (A.8) to see that, for every $x \in \left(\frac{r}{2}, r\right)$ and $\rho \in \left(0, x - \frac{r}{2}\right)$,

$$C_{q,s,k}(r-x+\rho)^{-qs-k-1} \leq f^{(k+1)}(x) \leq C_{q,s,k}(r-x)^{-qs-k-1}.$$

Moreover, we point out that $f^{(i)}(x-\rho) \ge 0$ for all $i \in \{0, \ldots, k\}$. Therefore, for every $i \in \{0, \ldots, k\}$,

$$f^{(i)}(x) \ge \int_{x-\rho}^{x} f^{(i+1)}(t_1) dt_1$$

$$\ge \int_{x-\rho}^{x} \int_{x-\rho}^{t_1} f^{(i+2)}(t_2) dt_2 dt_1$$

$$\ge \int_{x-\rho}^{x} \int_{x-\rho}^{t_1} \cdots \int_{x-\rho}^{t_{k-i}} f^{(k+1)}(r-t_{k-i+1}) dt_{k-i+1} \dots dt_2 dt_1)$$

$$\ge \widetilde{C}_{q,s,k,i} (r-x+\rho)^{-qs-k-1} \rho^{k+1-i}.$$

This gives the lower bound in (A.6).

Additionally, recalling (3.7), we find that, for every $i \in \{0, 1, ..., k\}$ and $x \in \left(\frac{r}{2}, r\right)$,

$$f^{(i)}(x) = (-1)^{i} g^{(i)}(r-x) - (-1)^{i} g^{(i)}\left(\frac{r}{2}\right) + \sum_{j=i+1}^{k} \frac{(-1)^{j+1}}{(j-2)!} g^{(j)}\left(\frac{r}{2}\right) \left(x - \frac{r}{2}\right)^{j-2}$$

$$\leq (-1)^{i} g^{(i)}(r-x)$$

$$= \prod_{j=0}^{i-1} (qs+j)(r-x)^{-qs-i}.$$

This gives the upper bound in (A.6) with

$$C_{i,q,s} := \prod_{j=0}^{i-1} (qs+j), \tag{A.11}$$

as desired.

Remark A.4. We notice that if $p \in (1,2)$, $s \in \left[\frac{p-1}{2p}, 1\right)$ and $q := \frac{p}{m-1}$ it follows from (3.6) that

$$1 \leqslant k \leqslant \left[\frac{p}{p-1}\right] + 1.$$

Hence, if we recall (A.9) and we define

$$C_{p,m} := \prod_{i=0}^{\left[\frac{p}{p-1}\right]+1} \left(\frac{p}{m-1}+i\right),$$
(A.12)

we see that

$$\frac{p-1}{2(m-1)} \left(\frac{p-1}{2(m-1)} + 1\right) \leqslant C_{q,s,k} \leqslant C_{p,m}.$$
(A.13)

Thus, recalling also (A.10), for every $i \in \{0, \ldots, k+1\}$, we evince that

$$c_{p,m} := \frac{\frac{p-1}{2(m-1)} \left(\frac{p-1}{2(m-1)} + 1\right)}{\left(\left[\frac{p}{p-1}\right] + 2\right)!} \leqslant \widetilde{C}_{q,s,k,i} \leqslant C_{p,m}.$$

Hence, using also (A.5), we deduce that for every $i \in \{0, \ldots, k+1\}, \rho_0 \in (0, \frac{r}{2})$ and $t \in (\frac{r}{2}, r - \rho_0)$

$$c_{p,m}\left(\frac{r}{2}\right)^{-qs-k-1}\left(t-\frac{r}{2}\right)^{k+1-i} \leqslant f^{(i)}(t) \leqslant C_{p,m}\,\rho_0^{-qs-k-1}\left(t-\frac{r}{2}\right)^{k+1-i}.\tag{A.14}$$

Now we notice that, by (A.11) and (A.13),

$$C_{i,q,s} \leqslant C_{q,s,k} \leqslant C_{p,m}.\tag{A.15}$$

We infer from this and (A.6) that for every $x \in \left(\frac{2}{3}r, r\right)$, $\rho := \frac{r-x}{2}$ and $i \in \{0, \dots, k+1\}$

$$c_{p,m}3^{-qs-k-1}\rho^{-qs-i} \leqslant f^{(i)}(x) \leqslant C_{p,m}2^{-qs-i}\rho^{-qs-i}.$$

Thus, if we define the constant

$$\tilde{c}_{p,m} := c_{p,m} \left(\frac{1}{3}\right)^{\frac{p}{m-1} + \left[\frac{p}{p-1}\right] + 2}, \qquad (A.16)$$

we see that, for every $x \in \left(\frac{2}{3}r, r\right)$ and $i \in \{0, \dots, k+1\}$,

$$\tilde{c}_{p,m}\rho^{-qs-i} \leqslant f^{(i)}(x) \leqslant C_{p,m}\rho^{-qs-i},\tag{A.17}$$

where $\rho := \frac{r-x}{2}$.

Thanks to these observations, we now provide some auxiliary results that allow us to estimate the fractional *p*-Laplacian of the function v in (3.8), as stated in Proposition 3.4.

Lemma A.5. Let $p \in (1,2)$, $s \in \left[\frac{p-1}{2p}, 1\right)$ and $r \in (1,+\infty)$. Let $v : \mathbb{R}^n \to [0,1]$ be as in (3.8). Let $x \in B_r$ and

$$\rho := \frac{r - |x|}{2}.\tag{A.18}$$

Then, there exists $C_{n,p,m} \in (0, +\infty)$ such that

$$\int_{B_{\rho}(x)} \frac{\left(v(y) - v(x)\right) \left|v(y) - v(x)\right|^{p-2}}{\left|x - y\right|^{n+sp}} \, dy \leqslant \frac{C_{n,p,m}}{1 - s} \rho^{-qs(p-1)-sp}.$$
(A.19)

Proof. In what follows, for simplicity of notation, if $A, B \in (0, +\infty)$ we adopt the notation

$$A \lesssim B$$
 if $A \leqslant C_{n,p,m}B$

for some $C_{n,p,m} \in (0, +\infty)$ depending at most on n, p and m.

Moreover, for every $p \in (1,2), s \in \left[\frac{p-1}{2p}, 1\right)$ and k as in (3.6), we have that

$$(k+1)(p-1) - sp \ge (1-s)(p-1).$$
 (A.20)

Indeed,

$$(k+1)(p-1) - sp - (1-s)(p-1) = k(p-1) - s$$
$$= \left(\left[\frac{sp}{p-1} \right] + 1 \right) (p-1) - s \ge \frac{sp}{p-1}(p-1) - s = s(p-1) \ge 0,$$

as desired.

Now, we complete the proof of (A.19), which is based on a long and tedious computation, that we provide here below for the facility of the reader. We will deal separately with the cases $x \in \overline{B}_{\frac{r}{2}}$, $x \in B_{\hat{r}} \setminus \overline{B}_{\frac{r}{2}}$ and $x \in B_r \setminus B_{\hat{r}}$, where \hat{r} has been defined in (3.25).

(i) Case $x \in \overline{B}_{\frac{r}{2}}$. In this case, for every $y \in B_{\rho}(x)$ it holds that

$$|y| \le \rho + |x| = \frac{r - |x|}{2} + |x| = \frac{r + |x|}{2} \le \frac{3}{4}r$$

Hence, recalling (3.8), if $y \in B_{\rho}(x)$,

$$(v(y) - v(x)) |v(y) - v(x)|^{p-2} = (h(|y|) - h(|x|)) |h(|y|) - h(|x|)|^{p-2}$$

= $h(|y|) |h(|y|)|^{p-2} \le |f(|y|)|^{p-1} \chi_{\mathbb{R}^n \setminus B_{\frac{r}{2}}}(y).$

As a consequence,

$$\int_{B_{\rho}(x)} \frac{(v(y) - v(x)) |v(y) - v(x)|^{p-2}}{|x - y|^{n+sp}} \, dy \leqslant \int_{B_{\rho}(x) \setminus B_{\frac{r}{2}}} \frac{|f(|y|)|^{p-1}}{|x - y|^{n+sp}} \, dy.$$

We now use (A.14) with $\rho_0 := r/4$ and i := 0 and we obtain that

$$\int_{B_{\rho}(x)} \frac{\left(v(y) - v(x)\right) \left|v(y) - v(x)\right|^{p-2}}{\left|x - y\right|^{n+sp}} \, dy$$

$$\leqslant C_{p,m}^{p-1} \left(\frac{r}{4}\right)^{-(qs+k+1)(p-1)} \int_{B_{\rho}(x) \setminus B_{\frac{r}{2}}} \frac{\left(\left|y\right| - \frac{r}{2}\right)^{(p-1)(k+1)}}{\left|x - y\right|^{n+sp}} \, dy.$$

Thus, changing variable gives that

$$\int_{B_{\rho}(x)} \frac{\left(v(y) - v(x)\right) \left|v(y) - v(x)\right|^{p-2}}{\left|x - y\right|^{n+sp}} \, dy$$

$$\leqslant C_{p,m}^{p-1} \left(\frac{r}{4}\right)^{-(qs+k+1)(p-1)} \int_{B_{\rho} \setminus B_{\frac{r}{2}}(-x)} \frac{\left(|y + x| - \frac{r}{2}\right)^{(k+1)(p-1)}}{\left|y\right|^{n+sp}} \, dy$$

$$\leqslant C_{p,m}^{p-1} \left(\frac{r}{4}\right)^{-(qs+k+1)(p-1)} \int_{B_{\rho} \setminus B_{\frac{r}{2}}(-x)} \frac{\left(|y| + |x| - \frac{r}{2}\right)^{(k+1)(p-1)}}{\left|y\right|^{n+sp}} \, dy.$$

Since $|x| \leq r/2$, from this we find that

$$\int_{B_{\rho}(x)} \frac{\left(v(y) - v(x)\right) |v(y) - v(x)|^{p-2}}{|x - y|^{n+sp}} dy$$

$$\leq C_{p,m}^{p-1} \left(\frac{r}{4}\right)^{-(qs+k+1)(p-1)} \int_{B_{\rho}} \frac{|y|^{(k+1)(p-1)}}{|y|^{n+sp}} dy$$

$$= \frac{C_{p,m}^{p-1} \omega_{n-1}}{(k+1)(p-1) - sp} \left(\frac{r}{4}\right)^{-(qs+k+1)(p-1)} \rho^{(k+1)(p-1)-sp}.$$
(A.21)

Also, in light of (A.18) we have that $\rho \leqslant r/2$, and therefore

$$r^{-(qs+k+1)(p-1)}\rho^{(k+1)(p-1)-sp} \leqslant (2\rho)^{-(qs+k+1)(p-1)}\rho^{(k+1)(p-1)-sp} = 2^{-(qs+k+1)(p-1)}\rho^{-qs(p-1)-sp}.$$

Using this into (A.21) and recalling (A.20), we see that

$$\begin{split} \int_{B_{\rho}(x)} \frac{\left(v(y) - v(x)\right) \left|v(y) - v(x)\right|^{p-2}}{\left|x - y\right|^{n+sp}} \, dy &\leqslant \frac{C_{p,m}^{p-1} \,\omega_{n-1} 2^{(qs+k+1)(p-1)}}{(k+1)(p-1) - sp} \rho^{-qs(p-1) - sp} \\ &\leqslant \frac{C_{p,m}^{p-1} \,\omega_{n-1} 2^{(qs+k+1)(p-1)}}{(1 - s)(p-1)} \rho^{-qs(p-1) - sp}, \end{split}$$

which gives the desired estimate in (A.19).

(ii) Case $x \in B_{\hat{r}} \setminus \overline{B}_{\frac{r}{2}}$. If we denote by

$$B_{\rho}^{+} := \left\{ y \in B_{\rho} \text{ s.t. } y \cdot x \ge 0 \right\}$$

and
$$B_{\rho}^{-} := \left\{ y \in B_{\rho} \text{ s.t. } y \cdot x < 0 \right\},$$

we have that

$$|y+x| \ge |x|$$
 for every $y \in B^+_{\rho}$. (A.22)

Now, we write

$$\begin{split} &\int_{B_{\rho}} \frac{(v(y+x)-v(x))|v(y+x)-v(x)|^{p-2}}{|y|^{n+sp}} \, dy \\ &= \frac{1}{2} \int_{B_{\rho}} \frac{(v(y+x)-v(x))|v(y+x)-v(x)|^{p-2} + (v(x-y)-v(x))|v(x-y)-v(x)|^{p-2}}{|y|^{n+sp}} \, dy \\ &= \frac{1}{2} \int_{B_{\rho}^{+}} \frac{(v(y+x)-v(x))|v(y+x)-v(x)|^{p-2} + (v(x-y)-v(x))|v(x-y)-v(x)|^{p-2}}{|y|^{n+sp}} \, dy \\ &\quad + \frac{1}{2} \int_{B_{\rho}^{-}} \frac{(v(y+x)-v(x))|v(y+x)-v(x)|^{p-2} + (v(x-y)-v(x))|v(x-y)-v(x)|^{p-2}}{|y|^{n+sp}} \, dy. \end{split}$$

Since v is a radial function, changing variable in the last integral, we find that

$$\int_{B_{\rho}^{-}} \frac{(v(y+x)-v(x))|v(y+x)-v(x)|^{p-2} + (v(x-y)-v(x))|v(x-y)-v(x)|^{p-2}}{|y|^{n+sp}} \, dy$$

$$= \int_{B_{\rho}^{+}} \frac{(v(y+x)-v(x))|v(y+x)-v(x)|^{p-2} + (v(x-y)-v(x))|v(x-y)-v(x)|^{p-2}}{|y|^{n+sp}} \, dy,$$

and therefore

$$\begin{split} &\int_{B_{\rho}} \frac{(v(y+x)-v(x))|v(y+x)-v(x)|^{p-2}}{|y|^{n+sp}} \, dy \\ &= \int_{B_{\rho}^{+}} \frac{(v(y+x)-v(x))|v(y+x)-v(x)|^{p-2} + (v(x-y)-v(x))|v(x-y)-v(x)|^{p-2}}{|y|^{n+sp}} \, dy \\ &= \int_{B_{\rho}^{+}} \frac{(h(|y+x|)-h(|x|)) \left|h(|y+x|)-h(|x|)\right|^{p-2} + (h(|x-y|)-h(|x|)) \left|h(|x-y|)-h(|x|)\right|^{p-2}}{|y|^{n+sp}} \, dy. \end{split}$$

As a result, thanks to (A.22) and the fact that h is a non decreasing function,

$$\int_{B_{\rho}} \frac{(v(y+x) - v(x)) |v(y+x) - v(x)|^{p-2}}{|y|^{n+sp}} dy$$

$$= \int_{B_{\rho}^{+} \cap B_{|x|}(x) \cap B_{\frac{r}{2}}^{c}(x)} \frac{|h(|y+x|) - h(|x|)|^{p-1} - |h(|x|) - h(|x-y|)|^{p-1}}{|y|^{n+sp}} dy$$

$$+ \int_{B_{\rho}^{+} \cap B_{|x|}(x) \cap B_{\frac{r}{2}}(x)} \frac{|h(|y+x|) - h(|x|)|^{p-1} - |h(|x|) - h(|x-y|)|^{p-1}}{|y|^{n+sp}} dy$$

$$+ \int_{B_{\rho}^{+} \cap B_{|x|}^{c}(x)} \frac{|h(|y+x|) - h(|x|)|^{p-1} + |h(|x-y|) - h(|x|)|^{p-1}}{|y|^{n+sp}} dy.$$
(A.23)

Now, we claim that

$$\int_{B_{\rho}^{+}\cap B_{|x|}(x)\cap B_{\frac{r}{2}}^{c}(x)} \frac{|h(|y+x|) - h(|x|)|^{p-1} - |h(|x|) - h(|x-y|)|^{p-1}}{|y|^{n+sp}} dy \lesssim \frac{\rho^{-qs(p-1)-sp}}{1-s}, \qquad (A.24)$$

$$\int_{B_{\rho}^{+} \cap B_{|x|}(x) \cap B_{\frac{r}{2}}(x)} \frac{|h(|y+x|) - h(|x|)|^{p-1} - |h(|x|) - h(|x-y|)|^{p-1}}{|y|^{n+sp}} dy \lesssim \frac{\rho^{-qs(p-1)-sp}}{1-s}$$
(A.25)

and

$$\int_{B_{\rho}^{+} \cap B_{|x|}^{c}(x)} \frac{|h(|y+x|) - h(|x|)|^{p-1} + |h(|x-y|) - h(|x|)|^{p-1}}{|y|^{n+sp}} dy \lesssim \frac{\rho^{-qs(p-1)-sp}}{1-s}.$$
 (A.26)

We point out that if the claims in (A.24), (A.25) and (A.26) hold true, then from (A.23) the estimate in (A.19) for the case $x \in B_{\hat{r}} \setminus \overline{B}_{\frac{r}{2}}$ readily follows. Hence, from now on, we focus on the proofs of the claims in (A.24), (A.25) and (A.26).

Proof of (A.24): For this, we recall (3.8) and we use Proposition A.3 to find that, for every $y \in$ $B_{\rho}^+ \overline{\cap B_{|x|}(x)},$

$$\begin{aligned} h(|x+y|) - h(|x|) &\leq f(|x+y|) - f(|x|) = \int_{|x|}^{|x+y|} f'(t) \, dt \\ &= \int_{|x|}^{|x+y|} \left(f'(|x|) + \int_{|x|}^{t} f''(\xi) \, d\xi \right) \, dt \\ &\leq f'(|x|) \left(|y+x| - |x| \right) + f''(|x+y|) \int_{|x|}^{|x+y|} \left(\int_{|x|}^{t} d\xi \right) \, dt \\ &= f'(|x|) \left(|y+x| - |x| \right) + f''(|x+y|) \frac{(|x+y| - |x|)^2}{2}. \end{aligned}$$
(A.27)

Also, for every $y \in B^+_{\rho} \cap B_{|x|}(x) \cap B^c_{\frac{r}{2}}(x)$, we have that $|x-y| \in \left(\frac{r}{2}, \hat{r}\right)$, and so

$$h(|x|) - h(|x - y|) = f(|x|) - f(|x - y|) = \int_{|x - y|}^{|x|} f'(t) dt$$

$$= \int_{|x - y|}^{|x|} \left(f'(|x|) - \int_{t}^{|x|} f''(\xi) d\xi \right) dt$$

$$\ge f'(|x|) \left(|x| - |x - y| \right) - f''(|x|) \int_{|x - y|}^{|x|} \left(\int_{t}^{|x|} d\xi \right) dt$$

$$= f'(|x|) \left(|x| - |x - y| \right) - f''(|x|) \frac{\left(|x| - |x - y| \right)^{2}}{2}.$$
(A.28)

Now, we define

$$u(x) := 2\frac{f'(|x|)}{f''(|x|)} \tag{A.29}$$

and we notice that, for every $y \in B_{u(x)}$,

$$f'(|x|) - f''(|x|) \frac{(|x| - |x - y|)}{2} \ge f'(|x|) - f''(|x|) \frac{|y|}{2} \ge 0.$$

From this, (A.27) and (A.28) we deduce that, for every $y \in B^+_{\rho} \cap B_{|x|}(x) \cap B^c_{\frac{r}{2}}(x) \cap B_{u(x)}$,

$$\begin{split} &|h(|y+x|) - h(|x|)|^{p-1} - |h(|x|) - h(|x-y|)|^{p-1} \\ &\leq (|x+y| - |x|)^{p-1} \left| f'(|x|) + f''(|x+y|) \frac{|x+y| - |x|}{2} \right|^{p-1} \\ &- (|x| - |x-y|)^{p-1} \left| f'(|x|) - f''(|x|) \frac{(|x| - |x-y|)}{2} \right|^{p-1} \\ &= (|x+y| - |x|)^{p-1} \\ &\times \left(\left| f'(|x|) + f''(|x+y|) \frac{|x+y| - |x|}{2} \right|^{p-1} - \left| f'(|x|) - f''(|x|) \frac{(|x| - |x-y|)}{2} \right|^{p-1} \right) \\ &+ \left((|x+y| - |x|)^{p-1} - (|x| - |x-y|)^{p-1} \right) \left| f'(|x|) - f''(|x|) \frac{(|x| - |x-y|)}{2} \right|^{p-1} \\ &\leq |y|^{p-1} \left(\left| f'(|x|) + f''(|x+y|) \frac{|y|}{2} \right|^{p-1} - \left| f'(|x|) - f''(|x|) \frac{|y|}{2} \right|^{p-1} \right) \\ &+ \left((|x+y| - |x|)^{p-1} - (|x| - |x-y|)^{p-1} \right) \left| f'(|x|) - f''(|x|) \frac{|y|}{2} \right|^{p-1} . \end{split}$$

Applying Taylor's Theorem with the Lagrange remainder we obtain that

$$\begin{aligned} \left| f'(|x|) + f''(|x+y|) \frac{|y|}{2} \right|^{p-1} \\ &= |f'(|x|)|^{p-1} + (p-1) |f'(|x|)|^{p-2} f''(|x+y|) \frac{|y|}{2} \\ &+ \frac{(p-1)(p-2)}{8} |f'(|x|) + b|^{p-3} f''(|x+y|)^2 |y|^2 \\ &\leqslant |f'(|x|)|^{p-1} + (p-1) |f'(|x|)|^{p-2} f''(|x+y|) \frac{|y|}{2} \\ &+ \frac{(p-1)(2-p)}{8} |f'(|x|) + b|^{p-3} (f''(|x+y|))^2 |y|^2 \\ &\leqslant |f'(|x|)|^{p-1} + (p-1) |f'(|x|)|^{p-2} f''(|x+y|) \frac{|y|}{2} \\ &+ \frac{(p-1)(2-p)}{8} |f'(|x|)|^{p-3} (f''(|x+y|))^2 |y|^2, \end{aligned}$$
(A.31)

for some $b \in \left(0, f''(|x+y|)\frac{|y|}{2}\right)$. Also, for every $\xi \in \left(0, f''(|x|)\frac{|y|}{2}\right)$,

$$|f'(|x|) - \xi| \ge |f'(|x|)| - \xi \ge f'(|x|) - f''(|x|)\frac{|y|}{2} \ge 0.$$

Hence, thanks to Taylor's Theorem we find that, for some $a \in \left(0, f''(|x|)\frac{|y|}{2}\right)$,

$$\left| f'(|x|) - f''(|x|) \frac{|y|}{2} \right|^{p-1} = \left| f'(|x|) \right|^{p-1} - (p-1) \left| f'(|x|) \right|^{p-2} f''(|x|) \frac{|y|}{2} + \frac{(p-1)(p-2)}{8} \left| f'(|x|) - a \right|^{p-3} (f''(|x|))^2 |y|^2.$$

From this, (A.30) and (A.31) we conclude that, for every $y \in B_{\rho}^+ \cap B_{|x|}(x) \cap B_{\frac{r}{2}}^c(x) \cap B_{u(x)}$,

$$\begin{split} &|h(|y+x|) - h(|x|)|^{p-1} - |h(|x|) - h(|x-y|)|^{p-1} \\ \leqslant \frac{p-1}{2} |y|^p |f'(|x|)|^{p-2} \left(f''(|x+y|) + f''(|x|) \right) \\ &+ \frac{(p-1)(2-p)}{8} |y|^{p+1} \left(f'(|x|)^{p-3} (f''(|x+y|))^2 + \left| f'(|x|) - f''(|x|) \frac{|y|}{2} \right|^{p-3} (f''(|x|))^2 \right) \\ &+ \left((|x+y| - |x|)^{p-1} - (|x| - |x-y|)^{p-1} \right) |f'(|x|)|^{p-1}. \end{split}$$

Therefore, we deduce that, for every $w \in (0, u(x)) \cap (0, |x| - \frac{r}{2}) \cap (0, \rho)$,

$$\begin{split} &\int_{B_{\rho}^{+}\cap B_{|x|}(x)\cap B_{\frac{r}{2}}^{c}(x)} \frac{|h(|y+x|) - h(|x|)|^{p-1} - |h(|x|) - h(|x-y|)|^{p-1}}{|y|^{n+sp}} \, dy \\ &\leqslant \frac{p-1}{2} \int_{B_{w}^{+}\cap B_{|x|}(x)\cap B_{\frac{r}{2}}^{c}(x)} \frac{|f'(|x|)|^{p-2} \left(f''(|x+y|) + f''(|x|)\right)}{|y|^{n+sp-p}} \, dy \\ &+ \frac{(p-1)(2-p)}{8} \int_{B_{w}^{+}\cap B_{|x|}(x)\cap B_{\frac{r}{2}}^{c}(x)} \frac{f'(|x|)^{p-3} f''(|x+y|)^{2} + \left|f'(|x|) - f''(|x|)\frac{|y|}{2}\right|^{p-3} f''(|x|)^{2}}{|y|^{n+sp-p-1}} \, dy \\ &+ \int_{B_{w}^{+}\cap B_{|x|}(x)\cap B_{\frac{r}{2}}^{c}(x)} \frac{\left((|x+y| - |x|)^{p-1} - (|x| - |x-y|)^{p-1}\right) |f'(|x|)|^{p-1}}{|y|^{n+sp}} \, dy \\ &+ \int_{(B_{\rho}^{+}\setminus B_{w}^{+})\cap B_{|x|}(x)\cap B_{\frac{r}{2}}^{c}(x)} \frac{|h(|y+x|) - h(|x|)|^{p-1} - |h(|x|) - h(|x-y|)|^{p-1}}{|y|^{n+sp}} \, dy. \end{split}$$
(A.32)

Now we adopt the notation

$$\tilde{x} := |x| - \frac{r}{2}.\tag{A.33}$$

Also, we assume that $\hat{r} \in \left(\frac{2}{3}r, +\infty\right)$, the other case being similar. We notice that

$$\tilde{x} < \rho \quad \text{if } x \in B_{\frac{2}{3}r} \setminus \overline{B}_{\frac{r}{2}}$$
(A.34)

and
$$\tilde{x} \ge \rho$$
 if $x \in B_{\hat{r}} \setminus B_{\frac{2}{3}r}$. (A.35)

Moreover, we define

$$\alpha_{p,m} := \frac{\tilde{c}_{p,m}}{C_{p,m}}$$

and the function

$$w(x) := \begin{cases} \alpha_{p,m} \tilde{x} & \text{for } x \in B_{\frac{2}{3}r} \setminus \overline{B}_{\frac{r}{2}}, \\ \alpha_{p,m} \rho & \text{for } x \in B_{\hat{r}} \setminus B_{\frac{2}{3}r}, \end{cases}$$
(A.36)

where $\tilde{c}_{p,m}$ and $C_{p,m}$ are given respectively in (A.16) and (A.12).

We notice that $\alpha_{p,m} \leq 1$, thanks to (A.17). Moreover, recalling (A.29) and making use of (A.17),

$$u(x) = 2\frac{f'(|x|)}{f''(|x|)} \ge 2\frac{\tilde{c}_{p,m}\left(\frac{r-|x|}{2}\right)^{-qs-1}}{C_{p,m}\left(\frac{r-|x|}{2}\right)^{-qs-2}} = \alpha_{p,m}(r-|x|) = \alpha_{p,m}\left(\frac{r}{2} - \tilde{x}\right) \ge w(x).$$

Therefore, using also (A.34) and (A.35), we have that, if $x \in B_{\hat{r}} \setminus \overline{B}_{\frac{r}{2}}$,

$$w(x) \leqslant \min \left\{ u(x), \tilde{x}, \rho \right\}.$$
(A.37)

Accordingly, we can exploit (A.32) with w := w(x).

In this way, we deduce that, in order to show (A.24), it is enough to prove that, if $x \in B_{\hat{r}} \setminus B_{\frac{r}{2}}$,

$$\int_{B^{+}_{w(x)}\cap B_{|x|}(x)\cap B^{c}_{\frac{r}{2}}(x)} \frac{|f'(|x|)|^{p-2} \left(f''(|x+y|) + f''(|x|)\right)}{|y|^{n+sp-p}} dy \lesssim \frac{\rho^{-qs(p-1)-sp}}{1-s},$$
(A.38)

$$\int_{B^{+}_{w(x)}\cap B_{|x|}(x)\cap B^{c}_{\frac{r}{2}}(x)} \frac{f'(|x|)^{p-3}f''(|x+y|)^{2} + \left|f'(|x|) - f''(|x|)\frac{|y|}{2}\right|^{p-3}f''(|x|)^{2}}{|y|^{n+sp-p-1}} \, dy \tag{A.39}$$

$$\int_{B_{w(x)}^{+} \cap B_{|x|}(x) \cap B_{\frac{r}{2}}^{c}(x)} \frac{\left(\left(|x+y|-|x|\right)^{p-1} - \left(|x|-|x-y|\right)^{p-1}\right) |f'(|x|)|^{p-1}}{|y|^{n+sp}} dy \lesssim \frac{\rho^{-qs(p-1)-sp}}{1-s} \quad (A.40)$$

and

$$\int_{\left(B_{\rho}^{+}\setminus B_{w(x)}^{+}\right)\cap B_{|x|}(x)\cap B_{\frac{r}{2}}^{c}(x)}\frac{|h(|y+x|)-h(|x|)|^{p-1}-|h(|x|)-h(|x-y|)|^{p-1}}{|y|^{n+sp}}\,dy\lesssim\frac{\rho^{-qs(p-1)-sp}}{1-s}.$$
 (A.41)

Hence, from now on we focus on the proofs of these claims.

We first prove (A.38). To do so, if $x \in B_{\frac{2}{3}r} \setminus \overline{B}_{\frac{r}{2}}$, we observe that, in light of (A.14),

$$|f'(|x|)|^{p-2} \left(f''(|x+y|) + f''(|x|) \right)$$

$$\leq c_{p,m}^{p-2} \left(\frac{r}{2} \right)^{-(qs+k+1)(p-2)} \left(|x| - \frac{r}{2} \right)^{k(p-2)} C_{p,m} \left(\frac{r}{6} \right)^{-qs-k-1} \left(\left(|x+y| - \frac{r}{2} \right)^{k-1} + \left(|x| - \frac{r}{2} \right)^{k-1} \right)$$

$$\leq c_{p,m}^{p-2} C_{p,m} \left(\frac{r}{2} \right)^{-(qs+k+1)(p-2)} \left(\frac{r}{6} \right)^{-qs-k-1} \tilde{x}^{k(p-2)} \left((\tilde{x}+|y|)^{k-1} + \tilde{x}^{k-1} \right).$$

As a consequence,

$$\begin{split} &\int_{B_{w(x)}^{+}\cap B_{|x|}(x)\cap B_{\frac{r}{2}}^{c}(x)} \frac{\left|f'(|x|)\right|^{p-2} \left(f''(|x+y|) + f''(|x|)\right)}{|y|^{n+sp-p}} \, dy \\ &\leqslant c_{p,m}^{p-2} C_{p,m} \left(\frac{r}{2}\right)^{-(qs+k+1)(p-2)} \left(\frac{r}{6}\right)^{-qs-k-1} \int_{B_{w(x)}^{+}\cap B_{|x|}(x)\cap B_{\frac{r}{2}}^{c}(x)} \frac{\tilde{x}^{k(p-2)} \left(\left(\tilde{x}+|y|\right)^{k-1}+\tilde{x}^{k-1}\right)}{|y|^{n+sp-p}} \, dy \\ &\lesssim r^{-(p-1)(qs+k+1)} \int_{B_{w(x)}} \frac{\tilde{x}^{k(p-2)} \left(\left(\tilde{x}+|y|\right)^{k-1}+\tilde{x}^{k-1}\right)}{|y|^{n+sp-p}} \, dy. \end{split}$$

Thus, changing variable $z := y/\tilde{x}$, we obtain that

$$\begin{split} & \int_{B_{w(x)}^{+}\cap B_{|x|}(x)\cap B_{\frac{r}{2}}^{c}(x)} \frac{|f'(|x|)|^{p-2} \left(f''(|x+y|) + f''(|x|)\right)}{|y|^{n+sp-p}} \, dy \\ & \lesssim r^{-(p-1)(qs+k+1)} \tilde{x}^{(k+1)(p-1)-sp} \int_{B_{w(x)/\tilde{x}}} \frac{\left((1+|z|)^{k-1}+1\right)}{|z|^{n+sp-p}} \, dz \\ & \lesssim r^{-(p-1)(qs+k+1)} \tilde{x}^{(k+1)(p-1)-sp} \int_{B_{\alpha p,m}} \frac{\left((1+|z|)^{k-1}+1\right)}{|z|^{n+sp-p}} \, dz \\ & \lesssim \frac{r^{-(p-1)(qs+k+1)}}{p-sp} \tilde{x}^{(k+1)(p-1)-sp}. \end{split}$$

Now, we observe that $(k+1)(p-1) - sp \ge 0$ thanks to (A.20), and therefore we deduce from (A.34) that

$$\begin{split} \int_{B^+_{w(x)} \cap B_{|x|}(x) \cap B^c_{\frac{r}{2}}(x)} \frac{|f'(|x|)|^{p-2} \left(f''(|x+y|) + f''(|x|)\right)}{|y|^{n+sp-p}} \, dy \\ \lesssim \frac{r^{-(p-1)(qs+k+1)}}{1-s} \rho^{(k+1)(p-1)-sp} \leqslant \frac{\rho^{-qs(p-1)-sp}}{1-s}. \end{split}$$

This proves (A.38) when $x \in B_{\frac{2}{3}r} \setminus \overline{B}_{\frac{r}{2}}$.

If instead $x \in B_{\hat{r}} \setminus \overline{B}_{\frac{2}{3}r}$, we exploit (A.17) to obtain the estimate

$$|f'(x)|^{p-2} \left(f''(x+y) + f''(x)\right) \\ \leqslant \tilde{c}_{p,m}^{p-2} \left(\frac{r-|x|}{2}\right)^{(qs+1)(2-p)} C_{p,m} \left(\left(\frac{r-|x+y|}{2}\right)^{-qs-2} + \left(\frac{r-|x|}{2}\right)^{-qs-2}\right).$$
(A.42)

Also, if $y \in B_{w(x)}$ then $|x+y| \leq |x| + |y| \leq |x| + w(x) \leq |x| + \rho$, thanks to (A.37). Consequently,

$$\frac{r - |x + y|}{2} \ge \frac{r - |x| - \rho}{2} = \rho - \frac{\rho}{2} = \frac{\rho}{2}.$$
(A.43)

From this and (A.42), we infer that

$$|f'(x)|^{p-2} \left(f''(x+y) + f''(x)\right) \lesssim \rho^{(qs+1)(2-p)} \rho^{-qs-2} = \rho^{-qs(p-1)-p}.$$

As a result,

$$\int_{B_{w(x)}^{+}\cap B_{|x|}(x)\cap B_{\frac{r}{2}}^{c}(x)} \frac{|f'(x)|^{p-2} \left(f''(x+y)+f''(x)\right)}{|y|^{n+sp-p}} dy \lesssim \rho^{-qs(p-1)-p} \int_{B_{w(x)}} \frac{dy}{|y|^{n+sp-p}} \leqslant \rho^{-qs(p-1)-p} \int_{B_{\rho}} \frac{dy}{|y|^{n+sp-p}} \lesssim \frac{\rho^{-qs(p-1)-sp}}{1-s}.$$

This concludes the proof of (A.38).

Next, we show that (A.39) holds true. For this, we observe that, as a consequence of (A.14) and (A.16), if $x \in B_{\frac{2}{3}r} \setminus \overline{B}_{\frac{r}{2}}$ and $y \in B_{w(x)}$, then

$$f'(|x|) - f''(|x|)\frac{|y|}{2} \ge c_{p,m} \left(\frac{r}{2}\right)^{-qs-k-1} \tilde{x}^k - C_{p,m} \left(\frac{r}{6}\right)^{-qs-k-1} \tilde{x}^{k-1} \frac{|y|}{2} \ge c_{p,m} \left(\frac{r}{2}\right)^{-qs-k-1} \tilde{x}^k - C_{p,m} \left(\frac{r}{6}\right)^{-qs-k-1} \tilde{x}^{k-1} \frac{w(x)}{2} \ge c_{p,m} \left(\frac{r}{2}\right)^{-qs-k-1} \tilde{x}^k - \frac{\tilde{c}_{p,m}}{2} \left(\frac{r}{6}\right)^{-qs-k-1} \tilde{x}^k = c_{p,m} \left(\frac{r}{2}\right)^{-qs-k-1} \tilde{x}^k - \frac{c_{p,m}}{2} \left(\frac{1}{3}\right)^{q+\left[\frac{p}{p-1}\right]+1} \left(\frac{r}{6}\right)^{-qs-k-1} \tilde{x}^k \ge c_{p,m} \left(\frac{r}{2}\right)^{-qs-k-1} \tilde{x}^k - \frac{c_{p,m}}{2} \left(\frac{1}{3}\right)^{qs+k+1} \left(\frac{r}{6}\right)^{-qs-k-1} \tilde{x}^k = \frac{c_{p,m}}{2} \left(\frac{r}{2}\right)^{-qs-k-1} \tilde{x}^k.$$

Thus, making again use of (A.14),

$$f'(|x|)^{p-3}f''(|x+y|)^{2} + \left|f'(|x|) - f''(|x|)\frac{|y|}{2}\right|^{p-3}f''(|x|)^{2}$$

$$\leqslant c_{p,m}^{p-3}\left(\frac{r}{2}\right)^{-(qs+k+1)(p-3)}\left(|x| - \frac{r}{2}\right)^{k(p-3)}C_{p,m}^{2}\left(\frac{r}{6}\right)^{-2(qs+k+1)}\left(|x+y| - \frac{r}{2}\right)^{2(k-1)}$$

$$+ \frac{c_{p,m}^{p-3}}{2^{p-3}}\left(\frac{r}{2}\right)^{-(qs+k+1)(p-3)}\tilde{x}^{k(p-3)}C_{p,m}^{2}\left(\frac{r}{6}\right)^{-2(qs+k+1)}\left(|x| - \frac{r}{2}\right)^{2(k-1)}$$

$$\lesssim r^{-(p-1)(qs+k+1)} \tilde{x}^{k(p-3)} \left(\left(|x+y| - \frac{r}{2} \right)^{2(k-1)} + \tilde{x}^{2(k-1)} \right)$$

$$\leqslant r^{-(p-1)(qs+k+1)} \tilde{x}^{k(p-3)} \left((\tilde{x}+|y|)^{2(k-1)} + \tilde{x}^{2(k-1)} \right).$$

As a consequence, using also (A.34), we infer that

$$\begin{split} &\int_{B_{w(x)}^{+}\cap B_{|x|}(x)\cap B_{\frac{r}{2}}^{c}(x)} \frac{f'(|x|)^{p-3}f''(|x+y|)^{2} + \left|f'(|x|) - f''(|x|)\frac{|y|}{2}\right|^{p-3}f''(|x|)^{2}}{|y|^{n+sp-p-1}} \, dy \\ &\lesssim r^{-(p-1)(qs+k+1)}\tilde{x}^{k(p-3)} \left(\int_{B_{w(x)}} \frac{(\tilde{x}+|y|)^{2(k-1)}}{|y|^{n+sp-p-1}} \, dy + \int_{B_{w(x)}} \frac{\tilde{x}^{2(k-1)}}{|y|^{n+sp-p-1}} \, dy\right) \\ &\leqslant r^{-(p-1)(qs+k+1)}\tilde{x}^{(k+1)(p-1)-sp} \left(\int_{B_{\alpha p,m}} \frac{(1+|y|)^{2(k-1)}}{|y|^{n+sp-p-1}} \, dy + \int_{B_{\alpha p,m}} \frac{dy}{|y|^{n+sp-p-1}}\right) \\ &\leqslant r^{-(p-1)(qs+k+1)}\rho^{(k+1)(p-1)-sp} \int_{B_{\alpha p,m}} \frac{dy}{|y|^{n+sp-p-1}} \\ &\lesssim \frac{\rho^{-qs(p-1)-sp}}{1+p-sp} \\ &\leqslant \frac{\rho^{-qs(p-1)-sp}}{1-s}, \end{split}$$

which establishes (A.39) when $x \in B_{\frac{2}{3}r} \setminus \overline{B}_{\frac{r}{2}}$, Similarly, using (A.17), we find that, for all $x \in B_{\hat{r}} \setminus B_{\frac{2}{3}r}$ and $y \in B_{w(x)}$,

$$\begin{aligned} f'(|x|) - f''(|x|) \frac{|y|}{2} &\geq f'(|x|) - f''(|x|) \frac{w(x)}{2} \\ &= f'(|x|) - f''(|x|) \frac{\alpha_{p,m}\rho}{2} \\ &\geq \tilde{c}_{p,m} \rho^{-qs-1} - C_{p,m} \rho^{-qs-1} \frac{\alpha_{p,m}}{2} \\ &= \rho^{-qs-1} \frac{\tilde{c}_{p,m}}{2}. \end{aligned}$$

This and (A.17) give that

$$f'(|x|)^{p-3}f''(|x+y|)^{2} + \left|f'(|x|) - f''(|x|)\frac{|y|}{2}\right|^{p-3}f''(|x|)^{2}$$

$$\leqslant \quad \tilde{c}_{p,m}^{p-3}\rho^{-(qs+1)(p-3)}C_{p,m}^{2}\left(\frac{r-|x+y|}{2}\right)^{-2(qs+2)} + \rho^{-(qs+1)(p-3)}\frac{\tilde{c}_{p,m}^{p-3}}{2^{p-3}}C_{p,m}^{2}\rho^{-2(qs+2)}$$

Hence, recalling (A.43),

$$f'(|x|)^{p-3}f''(|x+y|)^2 + \left|f'(|x|) - f''(|x|)\frac{|y|}{2}\right|^{p-3}f''(|x|)^2 \lesssim \rho^{-(qs+1)(p-3)-2(qs+2)}.$$

From this we thereby find that

$$\begin{split} \int_{B_{w(x)}^+ \cap B_{|x|}(x) \cap B_{\frac{r}{2}}^-(x)} \frac{f'(|x|)^{p-3} f''(|x+y|)^2 + \left|f'(|x|) - f''(|x|)\frac{|y|}{2}\right|^{p-3} f''(|x|)^2}{|y|^{n+sp-p-1}} \, dy \\ \lesssim \rho^{(qs+1)(1-p)} \int_{B_{w(x)}} \frac{dy}{|y|^{n+sp-p-1}} \\ = \rho^{(qs+1)(1-p)} \rho^{-sp+p-1} \int_{B_{\alpha_{p,m}}} \frac{dy}{|y|^{n+sp-p-1}} \lesssim \frac{\rho^{-qs(p-1)-sp}}{1+p-sp} \leqslant \frac{\rho^{-qs(p-1)-sp}}{1-s}. \end{split}$$

Therefore the proof of (A.39) is complete.

Now we prove (A.40). To accomplish this goal, we prove first that there exists $C_p \in (0, +\infty)$ such that, for every $x \in B_{\hat{r}}$ and $y \in B^+_{\frac{w(x)}{|x|}} \cap B_1(x/|x|) \cap B^c_{\frac{r}{2|x|}}(x/|x|)$,

$$\left(\left|\frac{x}{|x|} + y\right| - 1\right)^{p-1} - \left(1 - \left|y - \frac{x}{|x|}\right|\right)^{p-1} \le C_p |y|^p.$$
(A.44)

To show the claim in (A.44), we assume without loss of generality that $x = |x|e_n$. Then, if we write $y = (y', y_n)$ with $y' := (y_1, \ldots, y_{n-1})$, we have that

$$\frac{d}{dy_n} \left| \frac{x}{|x|} + y \right| = \frac{d}{dy_n} \sqrt{|y'|^2 + y_n^2 + 2y_n + 1} = \frac{1 + y_n}{\sqrt{|y'|^2 + y_n^2 + 2y_n + 1}}$$

and similarly

$$\frac{d^2}{dy_n^2} \left| \frac{x}{|x|} + y \right| = \frac{\left(|y'|^2 + y_n^2 + 2y_n + 1 \right) - (1 + y_n)^2}{\left(|y'|^2 + y_n^2 + 2y_n + 1 \right)^{\frac{3}{2}}} = \frac{|y'|^2}{\left(|y'|^2 + y_n^2 + 2y_n + 1 \right)^{\frac{3}{2}}}.$$
(A.45)

We also compute that

$$\frac{d}{dy_n} \left| y - \frac{x}{|x|} \right| = \frac{d}{dy_n} \sqrt{|y'|^2 + y_n^2 - 2y_n + 1} = \frac{y_n - 1}{\sqrt{|y'|^2 + y_n^2 - 2y_n + 1}}$$

and

$$\frac{d^2}{dy_n^2} \left| y - \frac{x}{|x|} \right| = \frac{|y'|^2}{\left(|y'|^2 + y_n^2 - 2y_n + 1 \right)^{\frac{3}{2}}}.$$
(A.46)

Now we denote by $\pi_n : \mathbb{R}^n \to \mathbb{R}$ the projection along the *n*-th axis and we set

$$A := \pi_n \left(B^+_{\frac{w(x)}{|x|}} \cap B_1(x/|x|) \cap B^c_{\frac{r}{2|x|}}(x/|x|) \right).$$

Then, in light of (A.45) and (A.46),

$$\sup_{y_n \in A} \left| \frac{d^2}{dy_n^2} \left| \frac{x}{|x|} + y \right| \right| \leqslant \frac{|y'|^2}{\left(|y'|^2 + 1 \right)^{\frac{3}{2}}} \leqslant 1.$$

Also, since $x \in B_{\hat{r}}$,

$$\sup_{y_n \in A} \left| \frac{d^2}{dy_n^2} \left| y - \frac{x}{|x|} \right| \right| \leqslant \frac{1}{\left(\left| y' \right|^2 + y_n^2 - 2y_n + 1 \right)^{\frac{1}{2}}} \leqslant \frac{2|x|}{r} \leqslant \frac{2\hat{r}}{r} \leqslant 2.$$

By Taylor's Theorem we thereby deduce that

$$\left|\frac{x}{|x|} + y\right| - 1 = y_n + \frac{y_n^2}{2} \frac{d^2}{d\xi^2} \left|\frac{x}{|x|} - (y',\xi)\right| \le y_n + \frac{1}{2}y_n^2 \tag{A.47}$$

and that

$$1 - \left| y - \frac{x}{|x|} \right| = y_n - \frac{y_n^2}{2} \frac{d^2}{d\eta^2} \left| (y', \eta) - \frac{x}{|x|} \right| \ge y_n - y_n^2, \tag{A.48}$$

for some $\eta, \xi \in (0, y_n)$.

Furthermore, by (A.37), we have that, for every $y \in B^+_{\frac{w(x)}{|x|}} \cap B_1(x/|x|) \cap B^c_{\frac{r}{2|x|}}(x/|x|)$,

$$(1-y_n) \ge (1-|y|) \ge \left(1-\frac{w(x)}{|x|}\right) \ge \left(1-\frac{\min\left\{\tilde{x},\rho\right\}}{|x|}\right) \ge \frac{1}{2}.$$
(A.49)

Therefore, thanks to (A.47), (A.48) and (A.49), and applying Taylor's Theorem once again, we infer that, for every $y \in B^+_{\frac{w(x)}{|x|}} \cap B_1(x/|x|) \cap B^c_{\frac{r}{2|x|}}(x/|x|)$, there exist some $\xi, \eta \in (0, y_n)$ such that

$$\begin{pmatrix} \left| \frac{x}{|x|} + y \right| - 1 \end{pmatrix}^{p-1} - \left(1 - \left| \frac{x}{|x|} - y \right| \right)^{p-1} \\ = \left(\sqrt{|y'|^2 + y_n^2 + 2y_n + 1} - 1 \right)^{p-1} - \left(1 - \sqrt{|y'|^2 + y_n^2 - 2y_n + 1} \right)^{p-1} \\ \leqslant y_n^{p-1} \left(\left(1 + \frac{1}{2} y_n \right)^{p-1} - (1 - y_n)^{p-1} \right) \\ \leqslant y_n^{p-1} \left(1 + \frac{p-1}{2} \left(1 + \frac{1}{2} \xi \right)^{p-2} y_n - 1 + (p-1) (1 - \eta)^{p-2} y_n \right) \\ \leqslant y_n^{p-1} \left(\frac{p-1}{2} y_n + (p-1) (1 - y_n)^{p-2} y_n \right) \\ \leqslant y_n^{p-1} \left(\frac{p-1}{2} y_n + (p-1) 2^{2-p} y_n \right) \\ \leqslant C_p y_n^p \\ \leqslant C_p |y|^p$$

where we have defined $C_p := (p-1)(2^{-1}+2^{2-p})$. This concludes the proof of (A.44). As a consequence of a change of variable and (A.44), if $x \in B_{\frac{2}{3}r} \setminus \overline{B}_{\frac{r}{2}}$,

$$\int_{B_{w(x)}^{+} \cap B_{|x|}(x) \cap B_{\frac{r}{2}}^{c}(x)} \frac{\left(\left(|x+y|-|x|\right)^{p-1} - \left(|x|-|x-y|\right)^{p-1}\right) |f'(|x|)|^{p-1}}{|y|^{n+sp}} dy$$

$$= |x|^{p-1-sp} \int_{B_{\frac{w(x)}{|x|}}^{+} \cap B_{1}(x/|x|) \cap B_{\frac{r}{2|x|}}^{c}(x/|x|)} \frac{\left(\left(|x/|x|+y|-1\right)^{p-1} - \left(1-|x/|x|-y|\right)^{p-1}\right) |f'(|x|)|^{p-1}}{|y|^{n+sp}} dy$$

$$\leqslant C_p |x|^{p-1-sp} \int_{B^+_{\frac{w(x)}{|x|}} \cap B_1(x/|x|) \cap B^c_{\frac{r}{2|x|}}(x/|x|)} \frac{|f'(|x|)|^{p-1}}{|y|^{n+sp-p}} \, dy$$

From this and (A.14) we thus find that

$$\begin{split} & \int_{B_{w(x)}^{+} \cap B_{|x|}(x) \cap B_{\frac{r}{2}}^{c}(x)} \frac{\left((|x+y|-|x|)^{p-1} - (|x|-|x-y|)^{p-1} \right) |f'(|x|)|^{p-1}}{|y|^{n+sp}} \, dy \\ \leqslant & C_p C_{p,m}^{p-1} |x|^{p-1-sp} \left(\frac{r}{6}\right)^{-(p-1)(qs+k+1)} \left(|x| - \frac{r}{2} \right)^{k(p-1)} \int_{B_{\frac{w(x)}{|x|}}^{+} \cap B_1(x/|x|) \cap B_{\frac{r}{2}|x|}^{c}(x/|x|)} \frac{dy}{|y|^{n+sp-p}} \\ \lesssim & |x|^{p-1-sp} r^{-(p-1)(qs+k+1)} \tilde{x}^{k(p-1)} \int_{B_{\frac{w(x)}{|x|}}^{+} \cap B_1(x/|x|) \cap B_{\frac{r}{2}|x|}^{c}(x/|x|)} \frac{dy}{|y|^{n+sp-p}}. \end{split}$$

Now, we observe that $w(x) \leq \tilde{x} \leq |x|$, thanks to (A.33) and (A.37). Therefore,

$$\begin{split} & \int_{B^+_{w(x)}\cap B_{|x|}(x)\cap B^c_{\frac{r}{2}}(x)} \frac{\left(\left(|x+y|-|x|\right)^{p-1}-\left(|x|-|x-y|\right)^{p-1}\right)|f'(|x|)|^{p-1}}{|y|^{n+sp}} \, dy \\ \lesssim & |x|^{p-1-sp}r^{-(p-1)(qs+k+1)}\tilde{x}^{k(p-1)} \int_{B_1} \frac{dy}{|y|^{n+sp-p}} \\ \lesssim & \frac{|x|^{p-1-sp}r^{-(p-1)(qs+k+1)}\tilde{x}^{k(p-1)}}{p-sp}. \end{split}$$

Accordingly, recalling also (A.34),

$$\int_{B_{w(x)}^{+}\cap B_{|x|}(x)\cap B_{\frac{r}{2}}^{c}(x)} \frac{\left(\left(|x+y|-|x|\right)^{p-1}-\left(|x|-|x-y|\right)^{p-1}\right)|f'(|x|)|^{p-1}}{|y|^{n+sp}} \, dy$$
$$\lesssim \frac{r^{-sp-(p-1)(qs+k)}\rho^{k(p-1)}}{p-sp} \lesssim \frac{\rho^{-qs(p-1)-sp}}{1-s},$$

which proves (A.40) when $x \in B_{\frac{2}{3}r} \setminus \overline{B}_{\frac{r}{2}}$. If $x \in B_{\hat{r}} \setminus B_{\frac{2}{3}r}$, we argue in a similar way, exploiting a change of variable and (A.44), but using now (A.17) to estimate |f'(|x|)| and (A.35) to see that $w(x) \leq \rho$. In this way, we conclude that

$$\begin{split} &\int_{B_{w(x)}^{+}\cap B_{|x|}(x)\cap B_{\frac{r}{2}}^{c}(x)} \frac{\left(\left(|x+y|-|x|\right)^{p-1}-\left(|x|-|x-y|\right)^{p-1}\right)|f'(|x|)\right)^{p-1}}{|y|^{n+sp}} \, dy \\ &\lesssim |x|^{p-1-sp} \int_{B_{\frac{w(x)}{|x|}}^{+}\cap B_{1}(x/|x|)\cap B_{\frac{r}{2|x|}}^{c}(x/|x|)} \frac{|f'(|x|)|^{p-1}}{|y|^{n+sp-p}} \, dy \\ &\leqslant C_{p,m}^{p-1} |x|^{p-1-sp} \, \rho^{-(qs+1)(p-1)} \int_{B_{\frac{w(x)}{|x|}}} \frac{dy}{|y|^{n+sp-p}} \\ &\lesssim |x|^{p-1-sp} \, \rho^{-(qs+1)(p-1)} \frac{\rho^{p-sp}}{|x|^{p-sp}} \frac{1}{p-sp} \\ &\lesssim \frac{\rho^{-qs(p-1)-sp}}{1-s}. \end{split}$$

Thus, the claim in (A.40) holds true for every $x \in B_{\hat{r}} \setminus \overline{B}_{\frac{r}{2}}$. In order to complete the proof of (A.24) it is only left to show (A.41). To do so, we observe that, for every t > 0 and $y \in \mathbb{R}^n \setminus B_t$,

$$(1+|y|)^{k-1} - 1 \leq (1+|y|)^{k+1} \leq (t^{-1}+1)^{k+1} |y|^{k+1}.$$
(A.50)

Moreover, we see that

$$\begin{split} & \int_{\left(B_{\rho}^{+} \setminus B_{w(x)}^{+}\right) \cap B_{|x|}(x) \cap B_{\frac{r}{2}}^{c}(x)} \frac{\left|h(|y+x|) - h(|x|)\right|^{p-1} - \left|h(|x|) - h(|x-y|)\right|^{p-1}}{|y|^{n+sp}} \, dy \\ & \leqslant \int_{\left(B_{\rho}^{+} \setminus B_{w(x)}^{+}\right) \cap B_{|x|}(x) \cap B_{\frac{r}{2}}^{c}(x)} \frac{\left|h(|y+x|) - h(|x|)\right|^{p-1}}{|y|^{n+sp}} \, dy \\ & \leqslant \int_{B_{\rho}^{+} \setminus B_{w(x)}^{+}} \frac{\left|h(|y+x|) - h(|x|)\right|^{p-1}}{|y|^{n+sp}} \, dy \\ & \leqslant \int_{B_{\rho}^{+} \setminus B_{w(x)}^{+}} \frac{\left|f(|y+x|) - f(|x|)\right|^{p-1}}{|y|^{n+sp}} \, dy. \end{split}$$
(A.51)

Now, if $x \in B_{\frac{2}{3}r} \setminus \overline{B}_{\frac{r}{2}}$ and $y \in B_{\rho}$, we have that $(\tilde{x} + \frac{r}{2}, \tilde{x} + \frac{r}{2} + |y|) \subset (\frac{r}{2}, \frac{5r}{6})$. Therefore, we employ (A.14) with $\rho_0 := r/6$ and we obtain that

$$|f(|y+x|) - f(|x|)| \leq \left| f\left(|y| + \tilde{x} + \frac{r}{2}\right) - f\left(\tilde{x} + \frac{r}{2}\right) \right| = \int_{\tilde{x}}^{\tilde{x}+|y|} f'\left(t + \frac{r}{2}\right) dt$$

$$\leq C_{p,m} \left(\frac{r}{6}\right)^{-qs-k-1} \int_{\tilde{x}}^{\tilde{x}+|y|} t^k dt \leq r^{-qs-k-1} \left((\tilde{x}+|y|)^{k+1} - \tilde{x}^{k+1} \right).$$
(A.52)

Thus, if $x \in B_{\frac{2}{3}r} \setminus \overline{B}_{\frac{r}{2}}$, we employ a change of variable to deduce that

$$\begin{split} & \int_{\left(B_{\rho}^{+}\setminus B_{w(x)}^{+}\right)\cap B_{|x|}(x)\cap B_{\frac{r}{2}}^{c}(x)} \frac{\left|h(|y+x|)-h(|x|)\right|^{p-1}-\left|h(|x|)-h(|x-y|)\right|^{p-1}}{|y|^{n+sp}} \, dy \\ &\lesssim r^{-(p-1)(qs+k+1)} \int_{B_{\rho}\setminus B_{w(x)}} \frac{\left((\tilde{x}+|y|)^{k+1}-\tilde{x}^{k+1}\right)^{p-1}}{|y|^{n+sp}} \, dy \\ &\lesssim r^{-(p-1)(qs+k+1)} \tilde{x}^{(k+1)(p-1)-sp} \int_{B_{\rho/\tilde{x}}\setminus B_{\alpha_{p,m}}} \frac{\left((1+|y|)^{k+1}-1\right)^{p-1}}{|y|^{n+sp}} \, dy. \end{split}$$

Now, formula (A.50) (used here with $t := \alpha_{p,m}$) gives that

$$\begin{split} &\int_{\left(B_{\rho}^{+}\setminus B_{w(x)}^{+}\right)\cap B_{|x|}(x)\cap B_{\frac{r}{2}}^{c}(x)} \frac{|h(|y+x|)-h(|x|)|^{p-1}-|h(|x|)-h(|x-y|)|^{p-1}}{|y|^{n+sp}} \, dy \\ &\lesssim r^{-(p-1)(qs+k+1)}\tilde{x}^{(k+1)(p-1)-sp} \int_{B_{\rho/\tilde{x}}\setminus B_{\alpha_{p,m}}} \frac{dy}{|y|^{n+sp-(k+1)(p-1)}} \\ &\leqslant r^{-(p-1)(qs+k+1)}\tilde{x}^{(k+1)(p-1)-sp} \int_{B_{\rho/\tilde{x}}} \frac{dy}{|y|^{n+sp-(k+1)(p-1)}} \\ &\lesssim \frac{r^{-(p-1)(qs+k+1)}}{(k+1)(p-1)-sp} \rho^{(k+1)(p-1)-sp}. \end{split}$$

Thus, recalling (A.20),

$$\begin{split} \int_{\left(B_{\rho}^{+}\setminus B_{w(x)}^{+}\right)\cap B_{|x|}(x)\cap B_{\frac{r}{2}}^{c}(x)} \frac{|h(|y+x|)-h(|x|)|^{p-1}-|h(|x|)-h(|x-y|)|^{p-1}}{|y|^{n+sp}} \, dy \\ \lesssim \frac{\rho^{-qs(p-1)-sp}}{(k+1)(p-1)-sp} \leqslant \frac{\rho^{-qs(p-1)-sp}}{1-s}, \end{split}$$

which is (A.41) when $x \in B_{\frac{2}{3}r} \setminus \overline{B}_{\frac{r}{2}}$. Analogously, if $x \in B_{\hat{r}} \setminus B_{\frac{2}{3}r}$ we use (A.17) into (A.51) to obtain that

$$\begin{split} \int_{\left(B^+_{\rho}\setminus B^+_{w(x)}\right)\cap B_{|x|}(x)\cap B^c_{\frac{r}{2}}(x)} \frac{\left|h(|y+x|)-h(|x|)\right|^{p-1}-\left|h(|x|)-h(|x-y|)\right|^{p-1}}{|y|^{n+sp}} \, dy \\ \leqslant C^{p-1}_{p,m} \int_{B^+_{\rho}\setminus B^+_{w(x)}} \frac{\rho^{-(qs+1)(p-1)}}{|y|^{n+sp-p+1}} \, dy. \end{split}$$

Hence, changing variable and recalling (A.36),

This concludes the proof of (A.41) and thus of the claim in (A.24).

Proof of (A.25): Toward this objective, we use the notation for \tilde{x} in (A.33) and we observe that, for every $y \in B_{\frac{r}{2\tilde{x}}}\left(\frac{x}{\tilde{x}}\right)$,

$$(|y|+1)^{k+1} - 1 \leqslant (|y|+1)^{k+1} \leqslant 2^{k+1} |y|^{k+1}.$$
(A.53)

Indeed, the first inequality is obvious. For the second one, we use the triangular inequality to obtain that, if $y \in B_{\frac{r}{2\tilde{x}}}\left(\frac{x}{\tilde{x}}\right)$,

$$\frac{|x|}{\tilde{x}} - |y| \leqslant \left|y - \frac{x}{\tilde{x}}\right| \leqslant \frac{r}{2\tilde{x}}$$

from which it follows that

$$|y| \ge \left(|x| - \frac{r}{2}\right)\frac{1}{\tilde{x}} = 1,$$

and therefore the proof of (A.53) is complete.

Moreover, if $x \in B_{\frac{2}{3}r} \setminus \overline{B}_{\frac{r}{2}}$ and $y \in B_{\rho}$, we recall (A.52) and we see that

$$|h(|y+x|) - h(|x|)|^{p-1} - |h(|x|) - h(|x-y|)|^{p-1} \le |h(|y+x|) - h(|x|)|^{p-1}$$

$$\leq |f(|y+x|) - f(|x|)|^{p-1} \leq r^{-(qs+k+1)(p-1)} \big((\tilde{x}+|y|)^{k+1} - \tilde{x}^{k+1} \big)^{p-1},$$

and thus a change of variable gives that

$$\begin{split} & \int_{B_{\rho}^{+} \cap B_{\frac{r}{2}}(x)} \frac{|h(|y+x|) - h(|x|)|^{p-1} - |h(|x|) - h(|x-y|)|^{p-1}}{|y|^{n+sp}} \, dy \\ &\lesssim r^{-(p-1)(qs+k+1)} \int_{B_{\rho}^{+} \cap B_{\frac{r}{2}}(x)} \frac{\left((|y|+\tilde{x})^{k+1} - \tilde{x}^{k+1}\right)^{p-1}}{|y|^{n+sp}} \, dy \\ &= r^{-(p-1)(qs+k+1)} \tilde{x}^{(k+1)(p-1)-sp} \int_{B_{\frac{\rho}{x}}^{+} \cap B_{\frac{r}{2x}}\left(\frac{x}{\tilde{x}}\right)} \frac{\left((|y|+1)^{k+1} - 1\right)^{p-1}}{|y|^{n+sp}} \, dy. \end{split}$$

From this, and using also (A.20), (A.36) and (A.53) we obtain that

$$\begin{split} &\int_{B_{\rho}^{+}\cap B_{\frac{r}{2}}(x)} \frac{|h(|y+x|) - h(|x|)|^{p-1} - |h(|x|) - h(|x-y|)|^{p-1}}{|y|^{n+sp}} \, dy \\ &\lesssim r^{-(p-1)(qs+k+1)} \tilde{x}^{(k+1)(p-1)-sp} \int_{B_{\frac{\rho}{x}}} \frac{dy}{|y|^{n+sp-(k+1)(p-1)}} \, dy \\ &\lesssim \frac{r^{-(p-1)(qs+k+1)}}{(k+1)(p-1)-sp} \rho^{(k+1)(p-1)-sp} \\ &\lesssim \frac{\rho^{-qs(p-1)-sp}}{(k+1)(p-1)-sp} \\ &\leqslant \frac{\rho^{-qs(p-1)-sp}}{1-s}. \end{split}$$

This establishes (A.25) if $x \in B_{\frac{2}{3}r} \setminus \overline{B}_{\frac{r}{2}}$. If instead $x \in B_{\hat{r}} \setminus B_{\frac{2}{3}r}$ we argue as follows. We notice that

$$B_{\frac{r}{2}}(x) \subset \mathbb{R}^n \setminus B_{\tilde{x}},\tag{A.54}$$

since, if $y \in B_{\frac{r}{2}}(x)$ then, recalling (A.33),

$$|y| \ge |x| - |y - x| = \tilde{x} + \frac{r}{2} - |y - x| \ge \tilde{x}.$$

Thus, according to (A.35) and (A.54), we have that

$$B^+_{\rho} \cap B_{\frac{r}{2}}(x) \subset B^+_{\rho} \cap B^c_{\tilde{x}} = \varnothing.$$

As a consequence,

$$\int_{B_{\rho}^{+}\cap B_{|x|}(x)\cap B_{\frac{r}{2}}(x)} \frac{|h(|y+x|) - h(|x|)|^{p-1} - |h(|x|) - h(|x-y|)|^{p-1}}{|y|^{n+sp}} \, dy = 0.$$

This concludes the proof of claim (A.25).

Proof of (A.26): Making use of (A.6) we see that, for every $y \in B_{\rho}$,

$$\begin{aligned} |h(|y \pm x|) - h(|x|)| &\leq |f(|x \pm y|) - f(|x|)| \leq \int_{|x|}^{|x|+|y|} f'(t) \, dt \leq f'(|x|+|y|)|y| \\ &\lesssim (r - |x| - |y|)^{-qs-1} \, |y| = (2\rho - |y|)^{-qs-1} \, |y| \leq \rho^{-qs-1}|y|. \end{aligned}$$

Thanks to this, and also changing variable, we find that

$$\begin{split} & \int_{B_{\rho}^{+} \cap B_{|x|}^{c}(x)} \frac{|h(|y+x|) - h(|x|)|^{p-1} + |h(|x-y|) - h(|x|)|^{p-1}}{|y|^{n+sp}} \, dy \\ & \lesssim \rho^{-(p-1)(qs+1)} \int_{B_{\rho}^{+} \cap B_{|x|}^{c}(x)} \frac{dy}{|y|^{n+sp-p+1}} \\ & = \rho^{-(p-1)(qs+1)} \, |x|^{p-1-sp} \int_{B_{\frac{\rho}{|x|}}^{+} \cap B_{1}^{c}\left(\frac{x}{|x|}\right)} \frac{dy}{|y|^{n+sp-p+1}} \\ & \leqslant \rho^{-(p-1)(qs+1)} \, |x|^{p-1-sp} \int_{B_{\frac{\rho}{|x|}}^{+} \cap B_{1}^{c}\left(\frac{x}{|x|}\right)} \frac{dy}{|y'|^{n+sp-p+1}}. \end{split}$$

We suppose, without loss of generality, that $x = |x|e_n$, and so

$$\int_{B_{\rho}^{+} \cap B_{|x|}^{c}(x)} \frac{|h(|y+x|) - h(|x|)|^{p-1} + |h(|x-y|) - h(|x|)|^{p-1}}{|y|^{n+sp}} dy$$

$$\lesssim \rho^{-(p-1)(qs+1)} |x|^{p-1-sp} \int_{B_{\frac{\rho}{|x|}}^{+} \cap B_{1}^{c}(e_{n})} \frac{dy}{|y'|^{n+sp-p+1}}$$

Now, we remark that, if $x \in B_{\hat{r}} \setminus \overline{B}_{\frac{r}{2}}$,

$$\frac{\rho}{|x|} = \frac{r}{2|x|} - \frac{1}{2} \leqslant 1 - \frac{1}{2} = \frac{1}{2}.$$

As a consequence, denoting by $y = (y', y_n) \in \mathbb{R}^{n-1} \times \mathbb{R}$, we have that

$$B_{\frac{\rho}{|x|}}^+ \cap B_1^c(e_n) \subset \left\{ y \in \mathbb{R}^n \text{ s.t. } |y'| \leqslant \frac{\rho}{|x|} \text{ and } y_n \in \left(0, 1 - \sqrt{1 - |y'|^2}\right) \right\},$$

and therefore

$$\begin{split} &\int_{B_{\rho}^{+} \cap B_{|x|}^{c}(x)} \frac{|h(|y+x|) - h(|x|)|^{p-1} + |h(|x-y|) - h(|x|)|^{p-1}}{|y|^{n+sp}} \, dy \\ &\lesssim \rho^{-(p-1)(qs+1)} \, |x|^{p-1-sp} \int_{\left\{ |y'| \leqslant \frac{\rho}{|x|} \right\}} \frac{1}{|y'|^{n+sp-p+1}} \left(\int_{0}^{1-\sqrt{1-|y'|^{2}}} dy_{n} \right) \, dy' \\ &= \rho^{-(p-1)(qs+1)} \, |x|^{p-1-sp} \int_{\left\{ |y'| \leqslant \frac{\rho}{|x|} \right\}} \frac{1-\sqrt{1-|y'|^{2}}}{|y'|^{n+sp-p+1}} \, dy' \\ &\leqslant \rho^{-(p-1)(qs+1)} \, |x|^{p-1-sp} \int_{\left\{ |y'| \leqslant \frac{\rho}{|x|} \right\}} \frac{2}{\sqrt{3} \, |y'|^{n+sp-p-1}} \, dy' \\ &\lesssim \rho^{-(p-1)(qs+1)} \, |x|^{p-1-sp} \, \frac{\rho^{p-sp}}{|x|^{p-sp}} \frac{1}{p-sp} \\ &\lesssim \frac{\rho^{-qs(p-1)-sp}}{1-s}. \end{split}$$

This concludes the proof of claim (A.26).

(iii) Case $x \in B_r \setminus B_r$. In this case, for every $y \in \mathbb{R}^n$ it holds that

$$v(y) - v(x) = v(y) - 1 \leqslant 0.$$

From this, we deduce that

$$\int_{B_{\rho}(x)} \frac{(v(y) - v(x)) |v(y) - v(x)|^{p-2}}{|x - y|^{n+sp}} \, dy \leqslant 0,$$

and (A.19) trivially follows.

We now provide an estimate on $r - \hat{r}$, where \hat{r} has been defined in (3.25).

Proposition A.6. Let $p \in (1, +\infty)$, $m \in [p, +\infty)$, $s \in (0, 1)$ and $r \in (1, +\infty)$. Let f and \hat{r} be defined respectively as in (3.8) and (3.25).

Then, there exists $\widehat{C}_{p,m} \in (0, +\infty)$ such that

$$r - \hat{r} > \widehat{C}_{p,m}$$

Proof. From the computations in (3.10), we see that, for every $t \in \left(\frac{r}{2}, r\right)$,

$$f(t) \ge (r-t)^{-qs} - c_{k,q,s} \left(\frac{r}{2}\right)^{-qs},$$

where $c_{k,q,s}$ is given in (3.11).

Thus, if we define

$$r_0 := r - \left(c_{k,q,s} \left(\frac{r}{2}\right)^{-qs} + 1\right)^{-\frac{1}{qs}},\tag{A.55}$$

we obtain that $f(t) \ge 1$ for every $t \in (r_0, r)$.

Therefore, owing to (3.25), we obtain that $\hat{r} < r_0$. This gives that

$$r - \hat{r} \ge \left(c_{k,q,s}\left(\frac{r}{2}\right)^{-qs} + 1\right)^{-\frac{1}{qs}} \ge \left(c_{k,q,s}2^{qs} + 1\right)^{-\frac{1}{qs}} =: C_{q,s,k}$$

Hence, in order to complete the proof of Lemma A.6, it is only left to show that there exists $\widehat{C}_{p,m} \in (0, +\infty)$ such that

$$C_{q,s,k} \geqslant \widehat{C}_{p,m}.$$
 (A.56)

To prove (A.56), we make use of (3.20) and we obtain that

$$C_{q,s,k} \ge \left(qsc_{p,m}^{(1)}2^q + 1\right)^{-\frac{1}{qs}} = \left(1 - \frac{qsc_{p,m}^{(1)}2^q}{qsc_{p,m}^{(1)}2^q + 1}\right)^{\frac{1}{qs}}.$$
(A.57)

Now we consider the continuous function $l: (0,1] \to (0,+\infty)$ defined by

$$l(s) := \left(1 - \frac{qsc_{p,m}^{(1)}2^q}{qsc_{p,m}^{(1)}2^q + 1}\right)^{\frac{1}{qs}}.$$

_	L

We set $s_0 := \min\left\{1, \frac{1}{q2^{q+1}c_{p,m}^{(1)}}\right\}$ and we observe that, for every $s \in (0, s_0)$,

$$1 - \frac{qsc_{p,m}^{(1)}2^q}{qsc_{p,m}^{(1)}2^q + 1} \ge \frac{1}{2}$$

and therefore

$$l(s) \ge \left(1 - qsc_{p,m}^{(1)}2^q\right)^{\frac{1}{qs}}$$

Taking the limit as $s \to 0$, we obtain that

$$\liminf_{s \to 0} l(s) \ge e^{-c_{p,m}^{(1)}2^q}.$$

From this, the fact that $l \in C((0,1], (0, +\infty))$ and (A.57), we obtain (A.56), as desired. \Box Corollary A.7. Let $p \in (1,2), s \in \left[\frac{p-1}{2p}, 1\right), r \in (1, +\infty)$ and $\mu \in (0, +\infty)$. Let $v : \mathbb{R}^n \to [0,1]$ be

defined as in (3.8).

Then, there exists $C_{\mu,n,p,m} \in (0, +\infty)$ such that, for every $x \in B_r$,

$$\int_{B_{\mu}(x)} \frac{(v(y) - v(x)) |v(y) - v(x)|^{p-2}}{|x - y|^{n+sp}} \, dy \leqslant \frac{C_{\mu,n,p,m}}{1 - s}.$$
(A.58)

Proof. We recall that, if \hat{r} is given as in (3.25), then, for every $x \in B_r \setminus B_{\hat{r}}$ and every $y \in \mathbb{R}^n$,

$$v(y) - v(x) = v(y) - 1 \leqslant 0.$$

Consequently, for every $x \in B_r \setminus B_r$ and $\mu \in (0, +\infty)$,

$$\int_{B_{\mu}(x)} \frac{\left(v(y) - v(x)\right) \left|v(y) - v(x)\right|^{p-2}}{\left|x - y\right|^{n+sp}} \, dy \leqslant 0,$$

and (A.58) plainly follows.

Furthermore, in virtue of (A.19) and Lemma A.6, if $\rho := \frac{r-|x|}{2}$ we have that, for every $x \in B_{\hat{r}} \setminus \overline{B}_{\frac{r}{2}}$,

$$\int_{B_{\rho}(x)} \frac{\left(v(y) - v(x)\right) \left|v(y) - v(x)\right|^{p-2}}{\left|x - y\right|^{n+sp}} \, dy \leqslant C_{n,p,m} \frac{\rho^{-qs(p-1)-sp}}{p - sp} \\ \leqslant \frac{C_{n,p,m}}{p - sp} \left(\frac{r - \hat{r}}{2}\right)^{-qs(p-1)-sp} \leqslant \frac{C_{n,p,m}}{p - sp} \left(\frac{\widehat{C}_{p,m}}{2}\right)^{-qs(p-1)-sp}.$$

Moreover, if $\mu \in [\rho, +\infty)$,

$$\int_{B_{\mu}(x)\setminus B_{\rho}(x)} \frac{(v(y)-v(x))|v(y)-v(x)|^{p-2}}{|x-y|^{n+sp}} dy \leq \int_{B_{\mu}(x)\setminus B_{\rho}(x)} \frac{dy}{|x-y|^{n+sp}} \\ \leq \frac{\omega_{n-1}}{sp} \rho^{-sp} \leq \frac{1}{p-1} \frac{\omega_{n-1}}{p-sp} \rho^{-sp} \leq \frac{1}{p-1} \frac{\omega_{n-1}}{p-sp} \left(\frac{\widehat{C}_{p,m}}{2}\right)^{-sp}.$$

From the last two displays we thus obtain that, if $\mu \in [\rho, +\infty)$,

$$\int_{B_{\mu}(x)} \frac{(v(y) - v(x)) |v(y) - v(x)|^{p-2}}{|x - y|^{n + sp}} dy$$

$$\leq \frac{C_{n,m,p}}{p - sp} \left(\frac{\widehat{C}_{p,m}}{2}\right)^{-qs(p-1) - sp} + \frac{1}{p - 1} \frac{\omega_{n-1}}{p - sp} \left(\frac{\widehat{C}_{p,m}}{2}\right)^{-sp}$$

which gives the desired estimate.

Similarly, if $\mu \in (0, \rho)$,

$$\begin{split} &\int_{B_{\mu}(x)} \frac{\left(v(y) - v(x)\right) |v(y) - v(x)|^{p-2}}{|x - y|^{n + sp}} \, dy \\ &= \int_{B_{\rho}(x)} \frac{\left(v(y) - v(x)\right) |v(y) - v(x)|^{p-2}}{|x - y|^{n + sp}} \, dy - \int_{B_{\rho}(x) \setminus B_{\mu}(x)} \frac{\left(v(y) - v(x)\right) |v(y) - v(x)|^{p-2}}{|x - y|^{n + sp}} \, dy \\ &\leqslant \frac{C_{n,m,p}}{p - sp} \left(\frac{\widehat{C}_{p,m}}{2}\right)^{-qs(p-1) - sp} + 2^{p-1} \int_{B_{\rho}(x) \setminus B_{\mu}(x)} \frac{dy}{|x - y|^{n + sp}} \\ &\leqslant \frac{C_{n,m,p}}{p - sp} \left(\frac{\widehat{C}_{p,m}}{2}\right)^{-qs(p-1) - sp} + \frac{2^{p-1}}{p - 1} \frac{\omega_{n-1}}{p - sp} \mu^{-sp}, \end{split}$$

as desired.

References

[AC72] S. M. Allen and J. W. Cahn, Ground state structures in ordered binary alloys with second neighbor interactions, Acta Metallurgica 20 (1972), no. 3, 423–433, DOI https://doi.org/10.1016/0001-6160(72)90037-5.

- [AC81] H. W. Alt and L. A. Caffarelli, Existence and regularity for a minimum problem with free boundary, J. Reine Angew. Math. 325 (1981), 105–144. MR0618549
- [ACF84] Hans Wilhelm Alt, Luis A. Caffarelli, and Avner Friedman, Jets with two fluids. I. One free boundary, Indiana Univ. Math. J. 33 (1984), no. 2, 213–247, DOI 10.1512/iumj.1984.33.33011. MR0733897
- [Bou90] Guy Bouchitté, Singular perturbations of variational problems arising from a two-phase transition model, Appl. Math. Optim. 21 (1990), no. 3, 289–314, DOI 10.1007/BF01445167. MR1036589
- [CC95] Luis A. Caffarelli and Antonio Córdoba, Uniform convergence of a singular perturbation problem, Comm. Pure Appl. Math. 48 (1995), no. 1, 1–12, DOI 10.1002/cpa.3160480101. MR1310848
- [CH58] J. W. Cahn and J. E. Hilliard, Free energy of a nonuniform system. I. Interfacial free energy, The Journal of chemical physics 28 (1958), no. 2, 258–267.
- [Coz17] Matteo Cozzi, Regularity results and Harnack inequalities for minimizers and solutions of nonlocal problems: a unified approach via fractional De Giorgi classes, J. Funct. Anal. 272 (2017), no. 11, 4762–4837, DOI 10.1016/j.jfa.2017.02.016. MR3630640
- [DPDPV] F. De Pas, S. Dipierro, M. Piccinini, and E. Valdinoci, Heteroclinic connections for fractional Allen-Cahn equations with degenerate potentials, Preprint, available online at https://cvgmt.sns.it/paper/6925/.
- [DNPV12] E. Di Nezza, G. Palatucci, and E. Valdinoci, *Hitchhiker's guide to the fractional Sobolev spaces*, Bull. Sci. Math. **136** (2012), no. 5, 521–573, DOI 10.1016/j.bulsci.2011.12.004. MR2944369
 - [DFGV] S. Dipierro, A. Farina, G. Giacomin, and E. Valdinoci, *Density estimates for a nonlocal variational model* with a degenerate double-well potential driven by the Gagliardo norm, in preparation.
 - [DFV18] Serena Dipierro, Alberto Farina, and Enrico Valdinoci, Density estimates for degenerate double-well potentials, SIAM J. Math. Anal. 50 (2018), no. 6, 6333–6347, DOI 10.1137/17M114933X. MR3890785
 - [DV23] Serena Dipierro and Enrico Valdinoci, Some perspectives on (non)local phase transitions and minimal surfaces, Bull. Math. Sci. 13 (2023), no. 1, Paper No. 2330001, 77, DOI 10.1142/S1664360723300013. MR4581189
 - [FV08] Alberto Farina and Enrico Valdinoci, Geometry of quasiminimal phase transitions, Calc. Var. Partial Differential Equations 33 (2008), no. 1, 1–35, DOI 10.1007/s00526-007-0146-1. MR2413100

- [GP58] V. L. Ginzburg and L. P. Pitaevskiĭ, On the theory of superfluidity, Soviet Physics JETP 34(7) (1958), 858–861 (1240–1245 Ž. Eksper. Teoret. Fiz.) MR105929
- [Gur85] Morton E. Gurtin, On a theory of phase transitions with interfacial energy, Arch. Rational Mech. Anal. 87 (1985), no. 3, 187–212, DOI 10.1007/BF00250724. MR768066
- [Lin90] Peter Lindqvist, On the equation div $(|\nabla u|^{p-2}\nabla u) + \lambda |u|^{p-2}u = 0$, Proc. Amer. Math. Soc. **109** (1990), no. 1, 157–164, DOI 10.2307/2048375. MR1007505
- [MS02] V. Maz'ya and T. Shaposhnikova, On the Bourgain, Brezis, and Mironescu theorem concerning limiting embeddings of fractional Sobolev spaces, J. Funct. Anal. 195 (2002), no. 2, 230–238, DOI 10.1006/jfan.2002.3955. MR1940355
- [Mod87] Luciano Modica, The gradient theory of phase transitions and the minimal interface criterion, Arch. Rational Mech. Anal. 98 (1987), no. 2, 123–142, DOI 10.1007/BF00251230. MR0866718
- [MM77] Luciano Modica and Stefano Mortola, Un esempio di Γ⁻-convergenza, Boll. Un. Mat. Ital. B (5) 14 (1977), no. 1, 285–299 (Italian, with English summary). MR0445362
- [PSV13] Giampiero Palatucci, Ovidiu Savin, and Enrico Valdinoci, Local and global minimizers for a variational energy involving a fractional norm, Ann. Mat. Pura Appl. (4) 192 (2013), no. 4, 673–718, DOI 10.1007/s10231-011-0243-9. MR3081641
- [PV05a] Arshak Petrosyan and Enrico Valdinoci, Density estimates for a degenerate/singular phase-transition model, SIAM J. Math. Anal. 36 (2005), no. 4, 1057–1079, DOI 10.1137/S0036141003437678. MR2139200
- [PV05b] _____, Geometric properties of Bernoulli-type minimizers, Interfaces Free Bound. 7 (2005), no. 1, 55–77, DOI 10.4171/IFB/113. MR2126143
- [Row79] J. S. Rowlinson, Translation of J. D. van der Waals' "The thermodynamic theory of capillarity under the hypothesis of a continuous variation of density", J. Statist. Phys. 20 (1979), no. 2, 197–244, DOI 10.1007/BF01011513. MR0523642
- [SV11] Ovidiu Savin and Enrico Valdinoci, Density estimates for a nonlocal variational model via the Sobolev inequality, SIAM J. Math. Anal. 43 (2011), no. 6, 2675–2687, DOI 10.1137/110831040. MR2873236
- [SV12] _____, Γ-convergence for nonlocal phase transitions, Ann. Inst. H. Poincaré C Anal. Non Linéaire 29 (2012), no. 4, 479–500, DOI 10.1016/j.anihpc.2012.01.006. MR2948285
- [SV14] _____, Density estimates for a variational model driven by the Gagliardo norm, J. Math. Pures Appl. (9) 101 (2014), no. 1, 1–26, DOI 10.1016/j.matpur.2013.05.001 (English, with English and French summaries). MR3133422
- [TH13] D. Ter Haar, Collected papers of LD Landau, Elsevier, 2013.
- [Val01] Enrico Valdinoci, Plane-like minimizers in periodic media: Jet flows and Ginzburg-Landau, ProQuest LLC, Ann Arbor, MI, 2001. Thesis (Ph.D.)–The University of Texas at Austin. MR2703829
- [VSS06] Enrico Valdinoci, Berardino Sciunzi, and Vasile Ovidiu Savin, Flat level set regularity of p-Laplace phase transitions, Mem. Amer. Math. Soc. 182 (2006), no. 858, vi+144, DOI 10.1090/memo/0858. MR2228294
- [WZJL19] Christoph A Weber, D. Zwicker, F. Jülicher, and Chiu F. Lee, *Physics of active emulsions*, Reports on Progress in Physics 82 (2019), no. 6, 064601.