B-type anomaly coefficients for the D3-D7 domain wall

Georgios Linardopoulos^a

^a Asia Pacific Center for Theoretical Physics (APCTP)
Hogil Kim Memorial Building, #501 POSTECH
77 Cheongam-Ro Nam-gu, Pohang Gyeongsangbuk-do 37673, Korea.

E-mail: georgios.linardopoulos@apctp.org

ABSTRACT: We compute B-type Weyl anomaly coefficients for the two non-supersymmetric codimension-1 domain wall versions of $\mathcal{N} = 4$ super Yang-Mills theory which are holographically dual to the D3-D7 probe-brane system. We compute the two-point function of the (improved) energy-momentum tensor in the presence of the domain wall, to leading order in perturbation theory. From it we determine the two-point function of the corresponding displacement operator and the b_2 anomaly coefficient for both domain wall systems.

Contents

1	Introduction	1
2	The D3-D7 domain wall	3
3	Energy-momentum tensor	5
4	Anomaly coefficients	10
5	Conclusions	12
A	$\mathcal{N} = 4$ super Yang-Mills	13
в	Scalar propagators in the D3-D7 dCFT	14

1 Introduction

Even dimensional conformal field theories (CFTs) are known to develop conformal/Weyl (or trace) anomalies in curved spacetimes which make the trace of the energy-momentum (or stress) tensor acquire a non-vanishing expectation value. There is no Weyl anomaly in odd dimensions. Although flat space CFTs are generally free from Weyl anomalies, the corresponding (curved space) anomaly coefficients still show up in flat-space conformal data. For example, two and three-point functions of the energy-momentum tensor take the following forms in two spacetime dimensions (x_1, x_2) :

$$\langle T(\zeta_1) T(\zeta_2) \rangle = \frac{c/2}{(\zeta_1 - \zeta_2)^4}, \quad \langle T(\zeta_1) T(\zeta_2) T(\zeta_3) \rangle = \frac{c}{(\zeta_1 - \zeta_2)^2 (\zeta_2 - \zeta_3)^2 (\zeta_3 - \zeta_1)^2}, (1.1)$$

where $T \equiv T_{\zeta\zeta}$, and $\zeta \equiv x_1 + ix_2$, $\bar{\zeta} \equiv x_1 - ix_2$ are the holomorphic/anti-holomorphic coordinates, while c is the single (ambient) Weyl anomaly coefficient there is in two dimensions.

The presence of boundaries and defects gives rise to extra Weyl anomaly coefficients in both even and odd dimensions. The new anomaly coefficients are localized on the boundary. For example, the Weyl anomaly in the case of codimension-1 boundaries in 4-dimensional CFTs takes the following form (scheme-independent terms only) [1, 2] (see also [3]):

$$\left\langle T^{\mu}_{\mu} \right\rangle^{d=4} = \frac{1}{16\pi^2} \left(c W^2_{\mu\nu\rho\sigma} - a E_4 \right) + \frac{\delta \left(x_3 \right)}{16\pi^2} \left(a E_4^{(\text{bry})} - b_1 \operatorname{tr} \hat{K}^3 - b_2 h^{pq} \hat{K}^{rs} W_{pqrs} \right), \quad (1.2)$$

where $W_{\mu\nu\rho\sigma}$ is the Weyl tensor, E_4 is the ambient Euler density, $E_4^{(bry)}$ is the boundary term of the Euler characteristic, K_{pq} is the boundary extrinsic curvature ($\hat{K}_{pq} \equiv K_{pq} - Kh_{pq}/3$), and h_{pq} is the induced metric on the boundary at $x_3 = 0$. The Weyl anomaly (1.2) includes two main types of ambient/boundary terms, the A-term *a* and the B-terms *c*, b_1 , b_2 . Unlike 2d CFTs, where every correlation function of the energy-momentum tensor is determined by the single Weyl anomaly coefficient c, in four dimensions only two-point functions of the energy-momentum tensor are fully specified by the ambient B-type anomaly coefficient c. Three-point functions contain three independent unknown coefficients, so that only two of them can be fixed by the ambient anomaly coefficients a and c in (1.2) [4, 5].

On the other hand, the (B-type) boundary anomaly coefficients b_1 and b_2 of codimension-1 defects in four dimensions fully determine the two and three point functions of the so-called displacement operator. These are given by:

$$\left\langle \mathcal{D}\left(\mathbf{x}_{1}\right) \mathcal{D}\left(\mathbf{x}_{2}\right)\right\rangle = \frac{c_{\eta\eta}}{\mathbf{x}_{12}^{8}}, \qquad \left\langle \mathcal{D}\left(\mathbf{x}_{1}\right) \mathcal{D}\left(\mathbf{x}_{2}\right) \mathcal{D}\left(\mathbf{x}_{3}\right)\right\rangle = \frac{c_{\eta\eta\eta}}{\mathbf{x}_{12}^{4}\mathbf{x}_{23}^{4}\mathbf{x}_{31}^{4}}, \tag{1.3}$$

where $\mathbf{x}_{ij} \equiv |\mathbf{x}_i - \mathbf{x}_j|$ and the points $\mathbf{x}_{i,j}$ lie on the 3d boundary.¹ The displacement operator $\mathcal{D}(\mathbf{x})$ is a scalar operator that is localized on the codimension-1 boundary and quantifies the violation of translation invariance across it (see e.g. [6] for a relevant discussion). The boundary anomaly coefficients b_1 , b_2 are related to the two and three-point function structure constants of the displacement operator as follows [2, 7]:

$$b_1 = \frac{2\pi^3}{35} \cdot c_{\eta\eta\eta}, \qquad b_2 = \frac{2\pi^4}{15} \cdot c_{\eta\eta}.$$
 (1.4)

In free theories, the ambient B-type coefficient c is related to the boundary B-type coefficient b_2 as $b_2 = 8c$ [8, 9]. This relation generally ceases to hold in interacting theories [7].

In the present paper we compute the B-type boundary anomaly coefficient b_2 for two 4-dimensional, codimension-1 defect CFTs (dCFTs) which are holographically dual to the D3-D7 probe-brane system. The first of these systems has an $SU(2) \times SU(2)$ global symmetry, while the second system has an SO(5) global symmetry. Both dCFTs live in flat space and so they are anomaly-free. Yet we can compute the corresponding B-type anomaly coefficients from the correlation functions of the energy-momentum tensor and the displacement operator, as we have just seen. Our computations are perturbative in the 't Hooft coupling constant $\lambda \equiv g_{\rm YM}^2 N_c$, and we report the leading-order result for the b_2 anomaly coefficients. Recently, a similar computation was carried out in [10] for the codimension-1 dCFT₄ which is holographically dual to the SU(2) symmetric D3-probe-D5 brane system.

One important difference between the two (codimension-1, 4-dimensional, flat-space) dCFTs is that the dCFT which is dual to the D3-D5 system is half-BPS supersymmetric, while both dCFTs which are dual to the D3-D7 system are non-supersymmetric. The first direct consequence of this is that the D3-D7 system is unstable and needs to be stabilized by appropriate worldvolume fluxes [11–14]. The second direct consequence concerns the integrability of the D3-D7 system, for which it is known that the $SU(2) \times SU(2)$ symmetric case is not integrable [15], while the SO(5) symmetric case is only known to be integrable to leading order in perturbation theory [16–18] (see also [19]). What is more, supersymmetric localization methods which have recently become available for the supersymmetric D3-D5 dCFT [20–25], are not expected to be applicable to any of the non-supersymmetric dCFTs

¹We will be using roman typeface for the spacetime arguments of correlation functions in both CFTs and codimension-1 dCFTs, setting $\mathbf{x} = (\mathbf{x}, \mathbf{z})$ to denote the presence of a codimension-1 boundary at $\mathbf{z} = 0$.

which are dual to the D3-D7 probe-brane system.

Two-point functions of the energy-momentum tensor are very interesting observables in codimension-1 dCFTs [26, 27]. They are determined up to three (generally dependent) unknown coefficients which depend on the dCFT invariant ratio v, cf. (3.6). Because onepoint functions of the energy-momentum tensor vanish in codimension-1 dCFTs, two-point functions can be used together with the operator product expansion (OPE) to derive all higher-point correlation functions of the energy-momentum tensor. This is one of the main goals of the boundary conformal bootstrap program [28], which aims to determine all defect CFT correlation functions from a minimal set of scalar and spinorial data (such as for example one and two-point function structure constants of the energy-momentum tensor).

Our paper is organized as follows. In the following section 2, we introduce the two domain wall setups which are holographically dual to the $SU(2) \times SU(2)$ and the SO(5)symmetric D3-D7 probe-brane systems. In section 3 we compute the leading-order contribution to the energy-momentum tensor two-point function, for both versions of the D3-D7 domain wall. In section 4 we compute the corresponding two-point functions of the displacement operator and from it we determine the B-type boundary anomaly coefficients of the D3-D7 domain wall system (both versions). Our conclusions can be found in section 5.

2 The D3-D7 domain wall

The D3-D7 probe-brane system is made up from a large number of $N_c \to \infty$ coincident D3 branes which intersect a single (probe) D7 brane. The near-horizon geometry of the D3-branes is $AdS_5 \times S^5$. The relative orientation of the D-branes in this space is shown in table 1 below. The coordinates x_0, \ldots, x_3, z parametrize AdS_5 in the Poincaré frame, while the angles $\psi, \theta, \chi, \vartheta, \varrho$ parametrize the 5-sphere S^5 .

	x_0	x_1	x_2	x_3	z	ψ	θ	χ	θ	Q
D3	•	•	•	٠						
D7	•	٠	٠		٠		•	٠	•	•

Table 1. The D3-D7 intersection.

It can be shown that the geometry of the probe D7-brane inside $AdS_5 \times S^5$ can be either $AdS_4 \times S^2 \times S^2$ or $AdS_4 \times S^4$ [11–14]. The respective global bosonic symmetries are therefore $SO(3, 2) \times SO(3) \times SO(3)$ and $SO(3, 2) \times SO(5)$, breaking all (maximal) supersymmetries of the D3-brane system. There is also a tachyonic instability which causes the compact parts of the D7-brane ($S^2 \times S^2$ and S^4) to "slip off" either side of the S^5 equator. To lift the instability we add k_1 and k_2 units of magnetic flux on either of the two 2-spheres of the $AdS_4 \times S^2 \times S^2$ symmetric brane and $d_G \equiv (n+1)(n+2)(n+3)/6$ units of instanton flux on the 4-sphere of the $AdS_4 \times S^4$ symmetric brane. For more information, see e.g. [19, 29].

The D3-D7 probe-brane system is holographically dual to a 4-dimensional, codimension-1 dCFT. The ambient CFT₄ is just $\mathcal{N} = 4$ SYM. Refer to appendix A for the Lagrangian and our basic conventions regarding $\mathcal{N} = 4$ SYM. At $x_3 = 0$ there is a flat 2+1 dimensional boundary which may or may not host additional (mainly fermionic) degrees of freedom [12]. Defect CFT correlation functions can be computed at weak 't Hooft coupling λ by means of "interfaces" or domain walls [30, 31]. These are described by classical ("fuzzy funnel") solutions of the ambient equations of motion which share the global symmetries of the defect CFT [32, 33].

For the domain wall which is dual to the $SU(2) \times SU(2)$ symmetric D3-probe-D7 brane system, the classical solution of the scalar equations of motion of $\mathcal{N} = 4$ SYM reads [29]:

$$\varphi_{i} = \varphi_{i}^{\text{cl}}(x_{3}) = \frac{1}{x_{3}} \cdot \left[\begin{array}{c} \left(t_{i}^{(k_{1})} \otimes \mathbb{1}_{k_{2}} \right)_{k \times k} & 0_{k \times (N_{c} - k)} \\ 0_{(N_{c} - k) \times k} & 0_{(N_{c} - k) \times (N_{c} - k)} \end{array} \right]_{N_{c} \times N_{c}}, \quad i = 1, 2, 3, \tag{2.1}$$

$$\varphi_{i} = \varphi_{i}^{\text{cl}}(x_{3}) = \frac{1}{x_{3}} \cdot \left[\begin{array}{c} \left(\mathbb{1}_{k_{1}} \otimes t_{i}^{(k_{2})} \right)_{k \times k} & 0_{k \times (N_{c} - k)} \\ 0_{(N_{c} - k) \times k} & 0_{(N_{c} - k) \times (N_{c} - k)} \end{array} \right]_{N_{c} \times N_{c}}, \quad i = 4, 5, 6, \tag{2.2}$$

where $x_3 > 0$, and the fermionic/vector fields of $\mathcal{N} = 4$ SYM have been set to zero, i.e. $A_{\mu} = \psi_{\alpha} = 0$. For $k_1, k_2 = 0, 1, \ldots$ we have also set,

$$k \equiv k_1 \cdot k_2, \tag{2.3}$$

while the matrices $t_i^{(k_i)}$ furnish a $k_i \times k_i$ irreducible representation of SU(2) (where i = 1, 2):

$$\left[t_{i}^{(k_{i})}, t_{j}^{(k_{i})}\right] = i\epsilon_{ijl}t_{l}^{(k_{i})}, \qquad i, j, l = 1, 2, 3,$$
(2.4)

so that the domain wall solution (2.1)–(2.2) shares the global bosonic symmetry of its dual $SU(2) \times SU(2)$ symmetric D3-D7 probe-brane system.

For the domain wall that is dual to the SO(5) symmetric D3-D7 probe D-brane system, the classical solution of the scalar equations of motion of $\mathcal{N} = 4$ SYM is given by [29]:

$$\varphi_{i} = \varphi_{i}^{\text{cl}}(x_{3}) = \frac{1}{\sqrt{8}x_{3}} \cdot \begin{bmatrix} (G_{i})_{d_{G} \times d_{G}} & 0_{d_{G} \times (N_{c} - d_{G})} \\ 0_{(N_{c} - d_{G}) \times d_{G}} & 0_{(N_{c} - d_{G}) \times (N_{c} - d_{G})} \end{bmatrix}_{N_{c} \times N_{c}}, \quad i = 1, \dots, 5, \quad (2.5)$$

$$\varphi_{6} = 0, \quad (2.6)$$

where $x_3 > 0$, and the fermionic/vector fields of $\mathcal{N} = 4$ SYM have been set to zero as before $(A_{\mu} = \psi_{\alpha} = 0)$. We have also defined,

$$d_G \equiv \frac{1}{6} \cdot (n+1)(n+2)(n+3), \qquad n = 1, 2, \dots$$
(2.7)

The five $d_G \times d_G$ fuzzy S⁴ matrices (*G*-matrices) G_i are given by [34]:

$$G_{i} \equiv \left[\underbrace{\gamma_{i} \otimes \mathbb{1}_{4} \otimes \ldots \otimes \mathbb{1}_{4}}_{n \text{ terms}} + \mathbb{1}_{4} \otimes \gamma_{i} \otimes \ldots \otimes \mathbb{1}_{4} + \ldots + \mathbb{1}_{4} \otimes \ldots \otimes \mathbb{1}_{4} \otimes \gamma_{i}}_{n \text{ terms}}\right]_{\text{sym}}, \qquad (2.8)$$

where i = 1, ..., 5 and γ_i are the five 4×4 Euclidean Dirac matrices (in five dimensions):

$$\gamma_i = \begin{pmatrix} 0 & -i\sigma_i \\ i\sigma_i & 0 \end{pmatrix}, \quad i = 1, 2, 3, \qquad \gamma_4 = \begin{pmatrix} 0 & \mathbb{1}_2 \\ \mathbb{1}_2 & 0 \end{pmatrix}, \qquad \gamma_5 = \begin{pmatrix} \mathbb{1}_2 & 0 \\ 0 & -\mathbb{1}_2 \end{pmatrix}, \quad (2.9)$$

while σ_i are the three Pauli matrices. The Dirac matrices satisfy the SO(5) Clifford algebra,

$$\{\gamma_i, \gamma_j\} = 2\delta_{ij}\mathbb{1}_4. \tag{2.10}$$

The ten commutators of the five G-matrices,

$$G_{ij} \equiv \frac{1}{2} \left[G_i, G_j \right], \tag{2.11}$$

furnish a d_G -dimensional (anti-hermitian) irreducible representation of $SO(5) \simeq Sp(4)$:

$$[G_{ij}, G_{kl}] = 2\left(\delta_{jk}G_{il} + \delta_{il}G_{jk} - \delta_{ik}G_{jl} - \delta_{jl}G_{ik}\right).$$
(2.12)

This way, the domain wall solution (2.5)-(2.6) shares the global bosonic symmetry of the SO(5) symmetric D3-D7 system. More properties of the fuzzy S⁴ matrices (2.8) can be found in references [33, 34].

To compute defect CFT correlation functions at weak coupling we assign vacuum expectation values (vevs) to the gauge fields. The vevs originate from the breaking of gauge symmetry across the defect; the values of the vevs are given by the fuzzy funnel solutions. By using the fuzzy funnel solutions (2.1)-(2.2) and (2.5)-(2.6), tree-level one-point functions of chiral primary operators were computed in [29] for both dCFTs which are dual to the D3-D7 probe-brane system. In the double scaling limit λ/k^2 , $\lambda/n^2 \rightarrow 0$, the weak-coupling results so computed agreed with the corresponding supergravity calculation at strong coupling with Witten diagrams.

The computation of tree-level one-point functions of non-protected scalar operators, which led eventually to the discovery of a closed-form determinant formula for all treelevel one-point functions of scalar operators in the SO(5) symmetric D3-D7 dCFT, was carried out in [17, 35]. These works followed [36] which introduced the use of the Bethe ansatz for the computation of defect CFT correlation functions. The spectra of quantum fluctuations and one-loop corrections to the one-point functions of the vacuum state were worked out in [37, 38]. Agreement with the corresponding calculations at strong coupling was once more reported in the double scaling limit λ/k^2 , $\lambda/n^2 \rightarrow 0$. Non-integrability of the $SU(2) \times SU(2)$ symmetric dCFT was shown in [15], while leading-order integrability of the SO(5) symmetric dCFT was proven in [16]. The latter result was based on a (quench) integrability criterion which was formulated by Piroli, Pozsgay and Vernier in [39] and followed from on the seminal work of Ghoshal and Zamolodchikov [40]. In a follow-up paper [18], Gombor and Bajnok claimed that integrability of the SO(5) symmetric setup breaks down beyond leading order. More information can be found in the reviews [41-43].

3 Energy-momentum tensor

Having set up the domain wall descriptions of the two defect CFTs which are dual to the D3-D7 probe-brane system, we are now in position to compute correlation functions of the energy-momentum tensor at weak coupling. From these we can directly extract the correlation functions of the displacement operator and the corresponding boundary anomaly coefficients, as we will see in the next section. First off, let us go through the computation of ambient two and three-point functions. **Generalities** The form of two and three-point functions of the energy-momentum tensor in any *d*-dimensional CFT is completely fixed by symmetry, while one-point functions vanish [4, 5]. Two-point functions are in particular given by

$$\left\langle \Theta_{\mu\nu}\left(\mathbf{x}_{1}\right)\Theta_{\rho\sigma}\left(\mathbf{x}_{2}\right)\right\rangle =\frac{C_{T}}{\mathbf{x}_{12}^{2d}}\cdot\mathcal{I}_{\mu\nu\rho\sigma}\left(\mathbf{x}_{1}-\mathbf{x}_{2}\right),\qquad\mathbf{x}_{12}\equiv\left|\mathbf{x}_{1}-\mathbf{x}_{2}\right|,\tag{3.1}$$

where the energy-momentum tensor has been improved (i.e. it is symmetric, conserved and traceless, cf. (A.8)) and the inversion tensors $\mathcal{I}_{\mu\nu}$, $\mathcal{I}_{\mu\nu\rho\sigma}$ are defined as

$$\mathcal{I}_{\mu\nu}\left(\mathbf{x}\right) \equiv g_{\mu\nu} - \frac{2\,\mathbf{x}_{\mu}\mathbf{x}_{\nu}}{\mathbf{x}^{2}}, \quad \mathcal{I}_{\mu\nu\rho\sigma}\left(\mathbf{x}\right) \equiv \frac{1}{2}\left(\mathcal{I}_{\mu\rho}\left(\mathbf{x}\right)\mathcal{I}_{\nu\sigma}\left(\mathbf{x}\right) + \mathcal{I}_{\mu\sigma}\left(\mathbf{x}\right)\mathcal{I}_{\nu\rho}\left(\mathbf{x}\right)\right) - \frac{1}{d}g_{\mu\nu}g_{\rho\sigma}.$$
 (3.2)

A similar (albeit more complicated) expression holds for three-point functions. In $\mathcal{N} = 4$ super Yang-Mills (SYM) theory, two and three-point functions of the energy-momentum tensor are protected [44–46]. In other words, the leading-order values of the (improved) energy-momentum tensor two and three-point functions do not receive quantum corrections. As such they are both given by their free-field values (although it is quite straightforward to obtain the result by performing all the Wick contractions with the Feynman rules of $\mathcal{N} = 4$ SYM). For two-point functions [4],

$$C_T = \left(N_0 + 3N_{1/2} + 12N_1\right) \cdot \frac{d}{(d-1)} \cdot S_{d-1}^{-2}, \qquad S_d \equiv \frac{2\pi^{\frac{d}{2}}}{\Gamma\left(\frac{d}{2}\right)}, \tag{3.3}$$

where N_0 , $N_{1/2}$ and N_1 are respectively the numbers of free real scalar, Majorana spin one-half and spin-one fields, while S_{d-1} is the surface of a unit d-1 sphere. For $\mathcal{N} = 4$ SYM, $N_0 = 6N_c^2$, $N_{1/2} = 4N_c^2$, $N_1 = N_c^2$ in d = 4, so that

$$C_T = \frac{10N_c^2}{\pi^4}.$$
 (3.4)

In the presence of a codimension-1 boundary at z = 0 (in *d* spacetime dimensions), one-point functions of the (improved) energy-momentum tensor vanish,

$$\left\langle \Theta_{\mu\nu}\left(\mathbf{x}_{1}\right)\right\rangle = 0,\tag{3.5}$$

while the form of two-point functions is once more fully specified by symmetry [26, 27]:

$$\langle \Theta_{\mu\nu} \left(\mathbf{x}_{1} \right) \Theta_{\rho\sigma} \left(\mathbf{x}_{2} \right) \rangle = \frac{1}{\mathbf{x}_{12}^{2d}} \cdot \left\{ \left(X_{\mu} X_{\nu} - \frac{g_{\mu\nu}}{d} \right) \left(X_{\rho}^{\prime} X_{\sigma}^{\prime} - \frac{g_{\rho\sigma}}{d} \right) A \left(v \right) + \left(X_{\mu} X_{\rho}^{\prime} I_{\nu\sigma} + X_{\mu} X_{\sigma}^{\prime} I_{\nu\rho} + X_{\nu} X_{\sigma}^{\prime} I_{\mu\rho} + X_{\nu} X_{\rho}^{\prime} I_{\mu\sigma} - \frac{4}{d} g_{\mu\nu} X_{\rho}^{\prime} X_{\sigma}^{\prime} - \frac{4}{d} g_{\rho\sigma} X_{\mu} X_{\nu} + \frac{4}{d^{2}} g_{\mu\nu} g_{\rho\sigma} \right) B \left(v \right) + \mathcal{I}_{\mu\nu\rho\sigma} C \left(v \right) \right\},$$
(3.6)

however this time there is an explicit dependence on the dCFT invariant ratio v which can be formed out of only 2 ambient points. The dCFT invariant ratio v, which enters the defect two-point function (3.6) through the functions A(v), B(v), C(v), is defined as:

$$\xi \equiv \frac{\mathbf{x}_{12}^2}{4|\mathbf{z}_1||\mathbf{z}_2|}, \qquad v^2 \equiv \frac{\xi}{\xi+1} = \frac{\mathbf{x}_{12}^2}{\mathbf{x}_{12}^2 + 4|\mathbf{z}_1||\mathbf{z}_2|}.$$
(3.7)

The definitions of the inversion tensors $\mathcal{I}_{\mu\nu}$ and $\mathcal{I}_{\mu\nu\rho\sigma}$ can be found in (3.2) above. For simplicity, we have omitted their arguments $(x_1 - x_2)$ in (3.6). We have also set,

$$X_{\mu} \equiv z_{1} \cdot \frac{v}{\xi} \frac{\partial \xi}{\partial x_{1}^{\mu}} = v \left(\frac{2z_{1}}{x_{12}^{2}} \left(x_{1\mu} - x_{2\mu} \right) - \eta_{\mu} \right)$$
(3.8)

$$X'_{\rho} \equiv z_2 \cdot \frac{v}{\xi} \frac{\partial \xi}{\partial x_2^{\rho}} = -v \left(\frac{2z_2}{x_{12}^2} \left(x_{1\rho} - x_{2\rho} \right) + \eta_{\rho} \right),$$
(3.9)

where $\eta \equiv (0, 1)$ is the unit normal to the defect at z = 0. We also note that X, X' obey,

$$X_{\mu}X_{\mu} = X'_{\rho}X'_{\rho} = 1, \qquad X'_{\rho} = \mathcal{I}_{\rho\mu}X_{\mu}.$$
 (3.10)

The functions A(v), B(v) and C(v) of the codimension-1 defect two-point function (3.6) are not completely independent, but instead they satisfy the following condition [27]:

$$\left(v\frac{d}{dv}-d\right)\alpha\left(v\right) = 2(d-1)\gamma\left(v\right),\tag{3.11}$$

where the functions $\alpha(v)$ and $\gamma(v)$ are defined in terms of A(v), B(v) and C(v) as follows:

$$\alpha(v) \equiv \frac{d-1}{d^2} \cdot \left[(d-1) \left(A(v) + 4B(v) \right) + dC(v) \right], \qquad \gamma(v) \equiv -B(v) - \frac{C(v)}{2}. \quad (3.12)$$

Far away from the boundary (where $z_{1,2} \to \infty$ and $\xi, v \to 0$), conformal symmetry is restored and the dCFT two-point function (3.6) reduces to the CFT one in (3.1). Therefore, A(0) = B(0) = 0 and $C(0) = C_T$. Moreover, (3.12) tells us that $\alpha(0) = (d-1)C_T/d$.

Perturbative expansion Let us now employ the domain wall description of the D3-D7 dCFT to calculate the two-point function of the energy-momentum tensor. Before being able to do that, we need to expand the fields of $\mathcal{N} = 4$ SYM theory (A.1) around either of the classical solutions (2.1)–(2.2) and (2.5)–(2.6):

$$A_{\mu} = \tilde{A}^{a}_{\mu}T^{a}, \quad \psi_{\alpha,m} = \tilde{\psi}^{a}_{\alpha,m}T^{a}, \quad \varphi_{i}\left(x\right) = \varphi_{i}^{cl}\left(x_{3}\right) + \tilde{\varphi}_{i}\left(x\right), \quad \tilde{\varphi}_{i}\left(x\right) = \tilde{\varphi}^{a}_{i}\left(x\right)T^{a}, \quad (3.13)$$

where T^a are the $(N_c \times N_c)$ generators of $U(N_c)$ in the fundamental representation. By inserting the perturbation (3.13) into the expression of the improved energy-momentum tensor of $\mathcal{N} = 4$ SYM in (A.7), we are led to the following perturbative expansion:

$$\Theta_{\mu\nu}(x) = \sum_{n=0}^{4} \Theta_{\mu\nu}^{(n)}(x) = \Theta_{\mu\nu}^{(0)} + \Theta_{\mu\nu}^{(1)} + \Theta_{\mu\nu}^{(2)} + \Theta_{\mu\nu}^{(3)}(x) + \Theta_{\mu\nu}^{(4)}, \qquad (3.14)$$

where the superscripts in the parentheses denote the number of perturbed fields.

The first term $\Theta_{\mu\nu}^{(0)}$ in the perturbative expansion (3.14) can be obtained by simply plugging the fuzzy-funnel solutions (2.1)–(2.2), (2.5)–(2.6) into the expression (A.7) for the improved energy-momentum tensor of the ambient theory, that is $\mathcal{N} = 4$ SYM. Only the scalar part (A.9) of the energy-momentum tensor (A.7) is relevant, since the fermion and the vector boson fields have no vevs. We find that the classical value of the energy-momentum tensor vanishes,

$$\Theta_{\mu\nu}^{\rm cl} \equiv \Theta_{\mu\nu}^{(0)} = 0, \qquad (3.15)$$

for both the $SU(2) \times SU(2)$ symmetric domain wall (2.1)–(2.2), and the SO(5) symmetric one (2.5)–(2.6). Obviously, so do the corresponding one-point functions, in full accordance with the analysis of McAvity-Osborn in [26, 27] and Liendo-Rastelli-van Rees in [28].

To compute the second term $\Theta_{\mu\nu}^{(1)}$ in the perturbative expansion (3.14), we adopt the following unifying notation for the D3-D7 fuzzy funnel solutions (2.1)–(2.2) and (2.5)–(2.6):

$$\varphi_{i} = \varphi_{i}^{\text{cl}}(x_{3}) = \frac{1}{x_{3}} \cdot \begin{bmatrix} (\tau_{i})_{k \times k} & 0_{k \times (N_{c} - k)} \\ 0_{(N_{c} - k) \times k} & 0_{(N_{c} - k) \times (N_{c} - k)} \end{bmatrix}_{N_{c} \times N_{c}}, \quad i = 1, \dots, 6.$$
(3.16)

We dub the upper-left, $\mathbf{k} \times \mathbf{k}$ dimensional diagonal block as the "massive" one, and the lowerright, $(N_c - \mathbf{k}) \times (N_c - \mathbf{k})$ diagonal block as "massless". The two remaining, $\mathbf{k} \times (N_c - \mathbf{k})$ and $(N_c - \mathbf{k}) \times \mathbf{k}$ dimensional massless blocks of the $N_c \times N_c$ matrices (3.16) will be called "off-diagonal". The same nomenclature can also be applied to the perturbed fields:

so that the fields in the "massive" block are Wick-contracted with one set of (massive) defect propagators, the fields in the off-diagonal blocks are Wick-contracted with another set of (massive) defect propagators, while the fields in the massless block are Wick-contracted with the massless propagators of $\mathcal{N} = 4$ SYM.

The leading contribution to the perturbative expansion (3.14) takes the following form:

$$\Theta_{\mu\nu}^{(1)}(x) = \frac{1}{g_{\rm YM}^2} \frac{4}{3x_3^2} \cdot \operatorname{tr} \left\{ \left(\frac{1}{x_3} \eta_\mu \eta_\nu \tilde{\varphi}_i + \eta_\mu \partial_\nu \tilde{\varphi}_i + \eta_\nu \partial_\mu \tilde{\varphi}_i - \frac{g_{\mu\nu}}{2} \partial_3 \tilde{\varphi}_i + \frac{x_3}{2} \partial_\mu \partial_\nu \tilde{\varphi}_i \right) \cdot \tau_i + \frac{g_{\mu\nu}}{2x_3} \left(2\tau_j \tau_i \tau_j - \tau_i \tau_j \tau_j - \tau_j \tau_j \tau_i \right) \cdot \tilde{\varphi}_i \right\}.$$

$$(3.18)$$

For the $SU(2) \times SU(2)$ symmetric D3-D7 domain wall (2.1)–(2.2), $\tau_i = t_i^{(k_1)} \otimes \mathbb{1}_{k_2}$ and $\tau_{i+3} = \mathbb{1}_{k_1} \otimes t_i^{(k_2)}$ (for i = 1, 2, 3) in (3.16), so that the leading correction (3.18) becomes:

$$\Theta_{\mu\nu}^{(1)}(x) = \frac{1}{g_{\rm YM}^2} \frac{4}{3x_3^2} \cdot \operatorname{tr}\left\{ \left(\frac{1}{x_3} \left(\eta_{\mu} \eta_{\nu} - g_{\mu\nu} \right) \tilde{\varphi}_i + \eta_{\mu} \partial_{\nu} \tilde{\varphi}_i + \eta_{\nu} \partial_{\mu} \tilde{\varphi}_i - \frac{g_{\mu\nu}}{2} \partial_3 \tilde{\varphi}_i + \frac{x_3}{2} \partial_{\mu} \partial_{\nu} \tilde{\varphi}_i \right) \cdot \tau_i \right\}, \qquad (3.19)$$

where we have used

$$\tau_i \tau_i = (c_1 + c_2) \cdot \mathbb{1}_k, \qquad \tau_j \tau_i \tau_j = (c_1 + c_2 - 1) \tau_i, \qquad c_1 \equiv \frac{k_1^2 - 1}{4}, \quad c_2 \equiv \frac{k_2^2 - 1}{4}.$$
 (3.20)

For the SO(5) symmetric D3-D7 domain wall (2.5)–(2.6), $\tau_i = G_i/\sqrt{8}$ (i = 1, ..., 5) and the leading correction (3.18) in this case reads:

$$+\frac{x_3}{2}\partial_{\mu}\partial_{\nu}\tilde{\varphi}_i\Big)\cdot\frac{G_i}{\sqrt{8}}\bigg\},\qquad(3.21)$$

where we have used the following properties of G-matrices [33, 34] in (3.18):

$$G_i G_i = n(n+4) \cdot \mathbb{1}_{d_G}, \qquad G_j G_i G_j = (n(n+4) - 8) G_i.$$
 (3.22)

Because all the terms of the leading corrections $\Theta_{\mu\nu}^{(1)}$ in (3.19) and (3.21) are essentially traces of products of unperturbed fields like (3.16) and perturbed fields like (3.17), it is only the fields in the "massive" k × k blocks which contribute to $\Theta_{\mu\nu}^{(1)}$.

Two-point function Once we have at our disposal a perturbative framework (3.14) for the computation of the energy-momentum tensor in the dCFT that is dual to the D3-D7 probe-brane system, we can start computing correlation functions. To get the n-point connected correlation function, we multiply n copies of the (improved) energy-momentum tensor (3.14) (evaluated at different ambient points, x_1, \ldots, x_n) and Wick-contract all the perturbed fields with the defect propagators. We proceed order-by-order in the number of perturbed fields, starting with a single Wick contraction (since $\Theta_{\mu\nu}^{(0)} = 0$) and continuing until all four perturbed fields $\tilde{\varphi}_i$ (which can be present in the $\Theta_{\mu\nu}^{(4)}$ term of the energymomentum tensor (A.7)) are Wick-contracted.

Since each entry of the energy-momentum tensor contributes a factor of $1/\lambda$ and each Wick contraction is proportional to λ , the leading-order term of the **n**-point function will be proportional to λ^{1-n} , while each subsequent term will get an extra Wick contraction and an additional factor of λ . For the (connected) two-point function, the perturbative expansion takes the following form:²

$$\left\langle \Theta_{\mu\nu}\left(\mathbf{x}_{1}\right)\Theta_{\rho\sigma}\left(\mathbf{x}_{2}\right)\right\rangle = \mathbf{A}^{-1} + \mathbf{A}^{0} + \mathbf{A}^{0}$$

The leading contribution to the connected part of the energy-momentum tensor twopoint function $\langle \Theta_{\mu\nu}^{(1)} \Theta_{\rho\sigma}^{(1)} \rangle$, consists of a single Wick contraction and it is of order λ^{-1} . Based on what we discussed above, only the scalar fields in the massive k × k block contribute. The corresponding propagators for both versions of the D3-D7 domain wall can be found in appendix B (cf. (B.2) and (B.6)–(B.7) for the $SU(2) \times SU(2)$ symmetric case, (B.21) and (B.24) for SO(5)). Using these, we compute the following Wick-contracted quantities:

$$\operatorname{tr}[\tau_i \tilde{\varphi_i}] \cdot \operatorname{tr}[\tau_j \tilde{\varphi_j}] = k \left(c_1 + c_2\right) K^{5/2} \left(x, y\right) = k \left(c_1 + c_2\right) \frac{g_{\mathrm{YM}}^2}{320\pi^2} \frac{1}{x_3 y_3} \frac{{}_2F_1\left(2, 3, 6; -\xi^{-1}\right)}{\xi^3 \left(1 + \xi\right)} \left(3.24\right)$$

$$\operatorname{tr}[G_{i}\tilde{\varphi}_{i}] \cdot \operatorname{tr}[G_{j}\tilde{\varphi}_{j}] = c_{n} \cdot K^{5/2}(x,y) = c_{n} \cdot \frac{g_{\mathrm{YM}}^{2}}{320\pi^{2}} \frac{1}{x_{3}y_{3}} \frac{{}_{2}F_{1}\left(2,3,6;-\xi^{-1}\right)}{\xi^{3}\left(1+\xi\right)}, \quad (3.25)$$

where we have defined,

$$c_n \equiv \operatorname{tr} [G_i G_i] = \frac{1}{6} \cdot n(n+1)(n+2)(n+3)(n+4), \quad n = 1, 2, \dots$$
 (3.26)

 $^{^{2}}$ We ignore lollipop diagram contributions which have been shown to vanish in the dCFT which is dual to the D3-D5 probe-brane system [47].

To compute the leading-order contribution to the energy-momentum tensor two-point function, we only need the leading-order terms $\Theta_{\mu\nu}^{(1)}$ of the $\Theta_{\mu\nu}$ perturbative expansion (3.14). These are given by (3.19) for the $SU(2) \times SU(2)$ symmetric domain wall and by (3.21) for the SO(5) symmetric domain wall. Multiplying out two from each of these terms and performing all the Wick contractions we arrive at (for both domain wall systems):

$$\left\langle \Theta_{\mu\nu}^{(1)}\left(\mathbf{x}_{1}\right)\Theta_{\rho\sigma}^{(1)}\left(\mathbf{x}_{2}\right)\right\rangle = \frac{1}{\mathbf{x}_{12}^{8}} \left\{ \left(X_{\mu}X_{\nu} - \frac{g_{\mu\nu}}{4}\right)\left(Y_{\rho}Y_{\sigma} - \frac{g_{\rho\sigma}}{4}\right)A\left(v\right) + \left(X_{\mu}Y_{\rho}I_{\nu\sigma} + X_{\mu}Y_{\sigma}I_{\nu\rho} + X_{\nu}Y_{\sigma}I_{\mu\rho} + X_{\nu}Y_{\sigma}I_{\mu\sigma} - g_{\mu\nu}Y_{\rho}Y_{\sigma} - g_{\rho\sigma}X_{\mu}X_{\nu} + \frac{1}{4}g_{\mu\nu}g_{\rho\sigma}\right)B\left(v\right) + \mathcal{I}_{\mu\nu\rho\sigma}C\left(v\right) \right\}.$$

$$(3.27)$$

The result (3.27) we find, maintains the generic form of the energy-momentum tensor twopoint functions (3.6) which was specified in [26, 27], with

$$\mathcal{I}_{\mu\nu\rho\sigma}\left(\mathbf{x}\right) \equiv \frac{1}{2} \left(\mathcal{I}_{\mu\rho}\left(\mathbf{x}\right) \mathcal{I}_{\nu\sigma}\left(\mathbf{x}\right) + \mathcal{I}_{\mu\sigma}\left(\mathbf{x}\right) \mathcal{I}_{\nu\rho}\left(\mathbf{x}\right) \right) - \frac{1}{4} g_{\mu\nu} g_{\rho\sigma}.$$
 (3.28)

In both versions of the D3-D7 domain wall system, the functions of the dCFT invariant ratio A(v), B(v) and C(v) are given by

$$A(v) = 4\gamma_{\rm k} \left(6v^6 + 3v^4 + v^2\right) \tag{3.29}$$

$$B(v) = -\gamma_{k} \left(3v^{6} - v^{4} - 2v^{2} \right)$$
(3.30)

$$C(v) = \gamma_{k}v^{2} \left(v^{2} - 1\right)^{2}, \qquad (3.31)$$

where

$$\gamma_{\rm k} \equiv \frac{32c_{\rm k}}{9\pi^2 g_{\rm YM}^2}, \qquad c_{\rm k} = \begin{cases} k \left(c_1 + c_2 \right) & SU(2) \times SU(2) \\ c_n/8 & SO(5), \end{cases}$$
(3.32)

while $k \equiv k_1 k_2$, and the $SU(2) \times SU(2)$ constants c_1 and c_2 were defined in (3.20). See (3.26) for the definition of the SO(5) constant c_n . As in the case of the D3-D5 domain wall which was recently studied in [10], the functions A(v), B(v) and C(v) in (3.29)–(3.31) satisfy the condition (3.11).

4 Anomaly coefficients

In the present section we will compute the B-type boundary anomaly coefficient b_2 for both versions of the dCFT which is dual to the D3-D7 probe-brane system. As we have already seen in (1.3)-(1.4), these anomaly coefficients are directly related to the two-point function structure constants of the displacement operator. In turn, the displacement operator is related to the energy-momentum tensor in a simple way as we will see below, so that we can use the results we have found above for the two-point function of the energy-momentum tensor in order to extract the corresponding two-point function of the displacement operator and its structure constants.

Let us first briefly go through ambient anomalies. It has long been known that the

ambient Weyl anomaly in four dimensions (given by the ambient terms of (1.2)) is one-loop exact for theories with $\mathcal{N} = 4$ superconformal symmetry [46]. As a result, the corresponding (ambient) anomaly coefficients a and c are given by their free-field values, just like the structure constants C_T of the energy-momentum two-point functions in (3.3). The exact expressions for the ambient anomaly coefficients are [48]:

$$c = \frac{N_0 + 3N_{1/2} + 12N_1}{120} = \frac{\pi^4 C_T}{40}, \qquad a = \frac{2N_0 + 11N_{1/2} + 124N_1}{720}.$$
 (4.1)

In the case of $\mathcal{N} = 4$ SYM $(N_0 = 6N_c^2, N_{1/2} = 4N_c^2, N_1 = N_c^2)$, the ambient anomaly coefficients are given by,

$$a = c = \frac{N_c^2}{4} = \frac{\pi^4 C_T}{40}.$$
(4.2)

We will now specify the boundary, B-type anomaly coefficient b_2 , to leading order in the 't Hooft coupling constant λ , for both variants of the D3-D7 domain wall. Following [2, 7], we will read off the coefficient from the two-point function of the displacement operator, cf. (1.3)–(1.4). But first let us define the displacement operator. The displacement operator \mathcal{D} is a scalar operator which is localized on the boundary/defect at $x_3 = 0$ and quantifies the breaking of translation invariance across it, as we have already mentioned. It is defined by the divergence of the improved energy-momentum tensor as follows:

$$\partial^{\mu}\Theta_{\mu\nu}\left(x\right) = \delta(x_{3})\,\eta_{\nu}\,\mathcal{D}\left(\mathbf{x}\right),\tag{4.3}$$

where η_{μ} is the unit normal to the boundary at $x_3 = 0$, and $x = (\mathbf{x}, x_3)$. Integrating the normal coordinate x_3 from 0^- to 0^+ , by also taking into account the conformal invariance of the defect, we find

$$\mathcal{D}(\mathbf{x}) = \lim_{x_3 \to 0+} \Theta_{33}(\mathbf{x}, x_3) - \lim_{x_3 \to 0-} \Theta_{33}(\mathbf{x}, x_3), \qquad (4.4)$$

where $\mathbf{x} \equiv (x_0, x_1, x_2)$. The leading-order contribution to the displacement operator twopoint function follows directly from (3.27), (4.4):

$$\left\langle \mathcal{D}^{(1)}\left(\mathbf{x}_{1}\right)\mathcal{D}^{(1)}\left(\mathbf{x}_{2}\right)\right\rangle = \lim_{\mathbf{x}_{1},\mathbf{x}_{2}\to0^{+}}\left\langle \Theta_{33}^{(1)}\left(\mathbf{x}_{1}\right)\Theta_{33}^{(1)}\left(\mathbf{x}_{2}\right)\right\rangle = \frac{c_{\eta\eta}}{\mathbf{x}_{12}^{8}},$$
(4.5)

where the structure constant $c_{\eta\eta}$ is given by

$$c_{\eta\eta} \equiv \frac{15 \, b_2}{2\pi^4} = \frac{80 \, c_{\rm k} \, N_c}{\pi^2 \lambda}.\tag{4.6}$$

Therefore the boundary B-type anomaly coefficient b_2 takes the following values for each of the two versions of the D3-D7 domain wall, to leading order in perturbation theory:

$$b_{2} = \frac{32\pi^{2}c_{k}N_{c}}{3\lambda} + \mathcal{O}\left(\lambda^{0}\right), \qquad c_{k} = \begin{cases} k\left(c_{1}+c_{2}\right) & SU(2) \times SU(2) \\ c_{n}/8 & SO(5), \end{cases}$$
(4.7)

where again $k \equiv k_1 k_2$, the D3-D7 domain wall constants c_1 , c_2 were defined in (3.20), and c_n in (3.26).

5 Conclusions

We have computed the B-type boundary anomaly coefficient b_2 , to leading order in weak 't Hooft coupling λ , for both versions of the (non-supersymmetric) defect CFT that is dual to the D3-D7 probe-brane system. The dCFTs are described by domain walls; one domain wall has a global $SU(2) \times SU(2)$ symmetry and the other one has an SO(5) global symmetry.

Our result for the anomaly coefficient is (4.7). The 4-dimensional B-type anomaly coefficients c and b_2 satisfy,

$$c = \frac{\pi^4}{30} \cdot \alpha(0), \qquad b_2 = \frac{2\pi^4}{15} \cdot \alpha(1),$$
 (5.1)

where $\alpha(v)$ was defined in (3.12) above. The relation for the 4-dimensional B-type ambient anomaly coefficient c is a direct consequence of $c = \pi^4 C_T/40$ in (4.1) and the identity $\alpha(0) = (d-1)C_T/d$ which we showed in section 3. The relation for the 4-dimensional boundary anomaly coefficient b_2 follows from the identification (1.4), property (4.4) of the displacement operator and the generic form (3.6) of energy-momentum tensor two-point functions in codimension-1 dCFTs (see e.g. [27]).

By using the values (3.29)-(3.31) for the functions A(v), B(v) and C(v) that we found for the D3-D7 domain wall we can verify that b_2 in (4.7) does indeed satisfy (5.1). For the ambient anomaly coefficient we find c = 0 which is also to be expected, given that our leading-order result (3.27) for the energy-momentum tensor two-point function consists of a single Wick contraction. On the other hand, to obtain the full result (4.2) for the ambient anomaly coefficient c, two Wick contractions between the fields of $\mathcal{N} = 4$ SYM are needed. As we have mentioned in the introduction, in free dCFTs the ambient and boundary B-type anomaly coefficients c and b_2 are not independent as they satisfy $b_2 = 8c$. This identity is clearly not valid in our fully interacting setup, since $b_2 \sim \lambda^{-1} > 8c = 0$ (at least as long as $k_{1,2} > 1$), to leading order in the coupling constant $\lambda \to 0$.

Working out higher-order perturbative corrections to the b_2 boundary anomaly coefficient would allow us to check whether $b_2 = 8c$ remains true when k = 0 or $k_1 = k_2 = 1$ (besides being an interesting computation in itself). In other words, we would be able to find out whether the boundary B-type anomaly coefficient b_2 is also one-loop exact (for k = 0 and $k_1 = k_2 = 1$) like the corresponding ambient anomaly coefficients c and a. In the same vein, it would be interesting to calculate the B-type boundary anomaly coefficient b_1 in (1.4). This calculation is significantly harder of course, since now the three-point function of the energy-momentum tensor and the displacement operator are required.

The AdS/dCFT correspondence [30, 31] opens the possibility for a wide range of computations of boundary anomaly coefficients in holographic defect CFTs. Holographic defects of varying codimensions can be studied, both at weak [10] and strong coupling (following e.g. the recent works [19, 49]). For supersymmetric defects, localization methods [20–23] are also available. The program of computing new boundary anomaly coefficients in the context of the AdS/dCFT correspondence will be described in more detail elsewhere [50].

Acknowledgments

The author is thankful to M. de Leeuw, C. Kristjansen, P. Panopoulos, I. Papadimitriou, M. Roberts, M. Volk, and K. Zarembo for discussions. This work was supported by the National Development Research and Innovation Office (NKFIH) research grant K134946. The work of G.L. was supported in part by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT) (No. 2023R1A2C1006975), as well as by an appointment to the JRG Program at the APCTP through the Science and Technology Promotion Fund and Lottery Fund of the Korean Government.

A $\mathcal{N} = 4$ super Yang-Mills

Lagrangian The Lagrangian density of $\mathcal{N} = 4$ super Yang-Mills (SYM) theory reads:

$$\mathcal{L}_{\mathcal{N}=4} = \frac{2}{g_{\rm YM}^2} \cdot \operatorname{tr} \left\{ -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} - \frac{1}{2} \left(D_\mu \varphi_i \right)^2 + i \, \bar{\psi}_\alpha \not{D} \psi_\alpha + \frac{1}{4} \left[\varphi_i, \varphi_j \right]^2 + \sum_{i=1}^3 \mathcal{G}^i_{\alpha\beta} \bar{\psi}_\alpha \left[\varphi_i, \psi_\beta \right] + \sum_{i=4}^6 \mathcal{G}^i_{\alpha\beta} \bar{\psi}_\alpha \gamma_5 \left[\varphi_i, \psi_\beta \right] \right\}, \quad (A.1)$$

where $\bar{\psi}_{\alpha} \equiv \psi_{\alpha}^{\dagger} \gamma^{0}$, $\not{D} \equiv \gamma^{\mu} D_{\mu}$. In the present work, we adopt a mostly-plus (-+++) signature convention for the Minkowski metric. The definitions of the four 4×4 Minkowski Dirac matrices γ^{μ} in four dimensions (Weyl/chiral basis) and the six 4×4 matrices \mathcal{G}^{i} , which enter the $\mathcal{N} = 4$ SYM Lagrangian density (A.1), can be found in many places, see e.g. the appendixes of [19]. The fields of the Lagrangian (A.1) carry adjoint $U(N_{c})$ color indices for which,

$$A_{\mu} = A^a_{\mu} T^a, \qquad \varphi_i = \varphi^a_i T^a, \qquad \psi_{\alpha,m} = \psi^a_{\alpha,m} T^a, \qquad a = 1, \dots, N^2_c, \qquad (A.2)$$

where $\mu = 0, ..., 3$, i = 1, ..., 6, $\alpha = 1, ..., 4$ and m = 1, ..., 4. The equations of motion that follow from the action (A.1) are:

$$D^{\mu}F_{\mu\nu} = i \begin{bmatrix} D_{\nu}\varphi_i, \varphi_i \end{bmatrix}, \qquad D^{\mu}D_{\mu}\varphi_i = \begin{bmatrix} \varphi_j, \begin{bmatrix} \varphi_j, \varphi_i \end{bmatrix} \end{bmatrix},$$
(A.3)

Energy-momentum tensor To compute the energy-momentum tensor of $\mathcal{N} = 4$ SYM, we may employ either the canonical prescription,

$$T_{\mu\nu} = \frac{\partial \mathcal{L}}{\partial \partial^{\mu} A_{\rho}} \partial_{\nu} A_{\rho} + \frac{\partial \mathcal{L}}{\partial \partial^{\mu} \varphi_{i}} \partial_{\nu} \varphi_{i} + \frac{\partial \mathcal{L}}{\partial \partial^{\mu} \bar{\psi}_{\alpha}} \partial_{\nu} \bar{\psi}_{\alpha} + \frac{\partial \mathcal{L}}{\partial \partial^{\mu} \psi_{\alpha}} \partial_{\nu} \psi_{\alpha} - g_{\mu\nu} \mathcal{L}, \quad (A.5)$$

or the covariant prescription (see e.g. [48]),

$$T_{\mu\nu} \equiv \frac{2}{\sqrt{-g}} \cdot \frac{\delta S}{\delta g^{\mu\nu}}, \qquad S = \int dx^4 \sqrt{-g} \mathcal{L}, \qquad (A.6)$$

which leads to a manifestly symmetric energy-momentum tensor. On the other hand, the canonical energy-momentum tensor (A.5) is neither symmetric, traceless, or conserved and

it needs to be improved [51]. By applying a series of transformations to the canonical recipe (A.5) we are led to

$$\Theta_{\mu\nu} = \frac{2}{g_{\rm YM}^2} \cdot \operatorname{tr} \left\{ -F_{\mu}{}^{\varrho}F_{\nu\varrho} - \frac{2}{3} \left(D_{\mu}\varphi_i \right) \left(D_{\nu}\varphi_i \right) + \frac{1}{3} \varphi_i D_{(\mu}D_{\nu)}\varphi_i + \frac{i}{2} \bar{\psi}_{\alpha}\gamma_{(\mu}\overleftrightarrow{D}_{\nu)}\psi_{\alpha} \right\} - \frac{2}{g_{\rm YM}^2} \cdot \operatorname{tr} \left\{ -\frac{1}{4} F_{\mu\nu}F^{\mu\nu} - \frac{1}{6} \left(D_{\mu}\varphi_i \right)^2 - \frac{1}{12} \left[\varphi_i, \varphi_j \right]^2 \right\} \cdot g_{\mu\nu},$$
(A.7)

where $a_{(\mu\nu)} \equiv (a_{\mu\nu} + a_{\nu\mu})/2$, and $f \overleftrightarrow{\partial}_{\mu} g \equiv f(\partial_{\mu} g) - (\partial_{\mu} f) g$. Details of the improvement procedure can be found in the appendixes of [10] and in many other places, such as e.g. the set of lectures [52]. The improved energy-momentum tensor so constructed (A.7) is manifestly symmetric, on-shell traceless and conserved:

$$\Theta_{\mu\nu} = \Theta_{\nu\mu}, \qquad g^{\mu\nu}\Theta_{\mu\nu} = 0, \qquad \partial^{\mu}\Theta_{\mu\nu} = 0.$$
 (A.8)

In the present work we will be mainly interested in the scalar part of improved energymomentum tensor which is obtained from (A.7) by setting the fermions and the vector bosons to zero:

$$\Theta_{\mu\nu}^{\text{scalar}} = \frac{2}{g_{\text{YM}}^2} \text{tr} \left\{ -\frac{2}{3} \left(\partial_\mu \varphi_i \right) \left(\partial_\nu \varphi_i \right) + \frac{1}{3} \varphi_i \left(\partial_\mu \partial_\nu \varphi_i \right) + \frac{1}{6} g_{\mu\nu} \left[\left(\partial_\varrho \varphi_i \right)^2 + \frac{1}{2} \left[\varphi_i, \varphi_j \right]^2 \right] \right\}. (A.9)$$

The scalar part of the improved energy-momentum tensor (A.9) is obviously symmetric, on-shell traceless and conserved.

B Scalar propagators in the D3-D7 dCFT

The present appendix includes the expressions of all the scalar propagators that we use in this paper, for both versions of the defect CFT which is holographically dual to the D3-D7 probe brane system. As we have explained in section 3, we only need the propagators in the "massive" blocks. More details, as well as derivations of the results included here can be found in the original papers [37, 38].

 $SU(2)_{k_1} \times SU(2)_{k_2}$ symmetric dCFT We start off with the $SU(2)_{k_1} \times SU(2)_{k_2}$ symmetric domain wall [37]. The fluctuations of the scalar fields (in the "massive" $k \times k$ block) are decomposed in fuzzy SU(2) spherical harmonics as follows:

$$[\tilde{\varphi}_i]_{n_1,n_2} = \sum_{\ell_1=0}^{k_1-1} \sum_{\ell_2=0}^{k_2-1} \sum_{m_1=-\ell_1}^{\ell_1} \sum_{m_2=-\ell_2}^{\ell_2} (\tilde{\varphi}_i)_{\ell_1,m_1;\ell_2,m_2} \cdot \left[\hat{Y}_{\ell_1}^{m_1} \otimes \hat{Y}_{\ell_2}^{m_2} \right]_{n_1,n_2}, \quad (B.1)$$

where i = 1, ..., 6 and $n_1, n_2 = 1, ..., k \equiv k_1 k_2$. All in all, there are three different kinds of scalar propagators, (namely $\langle \tilde{\varphi}_i \tilde{\varphi}_{j+3} \rangle$, $\langle \tilde{\varphi}_i \tilde{\varphi}_j \rangle$, $\langle \tilde{\varphi}_{i+3} \tilde{\varphi}_{j+3} \rangle$, for i, j = 1, 2, 3), depending on which branch of the $SU(2) \times SU(2)$ symmetric domain wall (i.e. (2.1) or (2.2)) we are. In the mixed sector we find,

$$\langle (\tilde{\varphi}_i)_{\ell_1 m_1; \ell_2 m_2} (\tilde{\varphi}_{j+3})_{\ell'_1 m'_1; \ell'_2 m'_2} \rangle = (-1)^{m'_1 + m'_2} \delta_{\ell_1 \ell'_1} \delta_{\ell_2 \ell'_2} [t_i^{(\ell_1)}]_{m_1, -m'_1} [t_{j+3}^{(\ell_2)}]_{m_2, -m'_2} K^{\varphi}_{\text{opp}},$$
(B.2)

where the function K_{opp}^{φ} is defined as:

$$K_{\rm opp}^{\varphi} \equiv \frac{K_{\rm opp}^{m_{-}^{2}}}{N_{-}} - \frac{K_{\rm op}^{m_{0}^{2}}}{N_{0}} + \frac{K_{\rm op}^{m_{+}^{2}}}{N_{+}}, \qquad N_{\pm} \equiv \lambda_{\mp} \left(\lambda_{\mp} - \lambda_{\pm}\right), \quad N_{0} \equiv -\lambda_{+} \lambda_{-} \tag{B.3}$$

$$\lambda_{\pm} \equiv -\frac{1}{2} \pm \sqrt{\ell_1(\ell_1 + 1) + \ell_2(\ell_2 + 1) + \frac{1}{4}}, \quad (B.4)$$

and the masses are given by,

$$m_0^2 \equiv \ell_1(\ell_1 + 1) + \ell_2(\ell_2 + 1) + 2, \qquad m_{\pm}^2 \equiv \ell_1(\ell_1 + 1) + \ell_2(\ell_2 + 1) - 2\lambda_{\pm}.$$
 (B.5)

See (B.38) below for the functional form of the propagators $K^{m_{0,\pm}^2}$. In the two pure sectors the following formulas hold:

$$\langle (\tilde{\varphi}_i)_{\ell_1 m_1; \ell_2 m_2} (\tilde{\varphi}_j)_{\ell'_1 m'_1; \ell'_2 m'_2} \rangle = (-1)^{m'_1 + m'_2} \delta_{\ell_1 \ell'_1} \delta_{\ell_2 \ell'_2} \delta_{m_2 + m'_2} \left[\delta_{ij} \delta_{m_1 + m'_1} K^{\varphi, \ell_1}_{\text{sing}} - i\epsilon_{ijk} [t_k^{(\ell_1)}]_{m_1, -m'_1} K^{\varphi, \ell_1}_{\text{anti}} - [t_i^{(\ell_1)} t_j^{(\ell_1)}]_{m_1, -m'_1} K^{\varphi, \ell_1}_{\text{sym}} \right]$$
(B.6)

$$\langle (\tilde{\varphi}_{i+3})_{\ell_1 m_1; \ell_2 m_2} (\tilde{\varphi}_{j+3})_{\ell'_1 m'_1; \ell'_2 m'_2} \rangle = (-1)^{m'_1 + m'_2} \delta_{\ell_1 \ell'_1} \delta_{\ell_2 \ell'_2} \delta_{m_1 + m'_1} \left[\delta_{ij} \delta_{m_2 + m'_2} K^{\varphi, \ell_2}_{\text{sing}} - i\epsilon_{ijk} [t_k^{(\ell_2)}]_{m_2, -m'_2} K^{\varphi, \ell_2}_{\text{anti}} - [t_i^{(\ell_2)} t_j^{(\ell_2)}]_{m_2, -m'_2} K^{\varphi, \ell_2}_{\text{sym}} \right], (B.7)$$

where the functions $K_{\text{sing}}^{\varphi,\ell_i}$, $K_{\text{anti}}^{\varphi,\ell_i}$ and $K_{\text{sym}}^{\varphi,\ell_i}$ are defined as follows, for i = 1, 2:

$$K_{\text{sing}}^{\varphi,\ell_i} \equiv \frac{\ell_i + 1}{2\ell_i + 1} \cdot K^{m_{i,+}^2} + \frac{\ell_i}{2\ell_i + 1} \cdot K^{m_{i,-}^2}, \qquad K_{\text{anti}}^{\varphi,\ell_i} \equiv \frac{K^{m_{i,+}^2}}{2\ell_i + 1} - \frac{K^{m_{i,-}^2}}{2\ell_i + 1}$$
(B.8)

$$K_{\rm sym}^{\varphi,\ell_1} \equiv \frac{K^{m_{1,+}^2}}{(2\ell_1+1)(\ell_1+1)} + \frac{K^{m_{1,-}^2}}{(2\ell_1+1)\ell_1} - \frac{\ell_2(\ell_2+1)}{\ell_1(\ell_1+1)} \cdot \frac{K^{m_0^2}}{N_0} - \frac{K^{m_-^2}}{N_-} - \frac{K^{m_+^2}}{N_+} \quad (B.9)$$

$$K_{\rm sym}^{\varphi,\ell_2} \equiv \frac{K^{m_{2,+}^2}}{(2\ell_2+1)(\ell_2+1)} + \frac{K^{m_{2,-}^2}}{(2\ell_2+1)\ell_2} - \frac{\ell_1(\ell_1+1)}{\ell_2(\ell_2+1)} \cdot \frac{K^{m_0^2}}{N_0} - \frac{K^{m_-^2}}{N_-} - \frac{K^{m_+^2}}{N_+}, (B.10)$$

and the masses are given by:

$$m_{1,+}^2 = \ell_1(\ell_1 - 1) + \ell_2(\ell_2 + 1), \qquad m_{1,-}^2 = (\ell_1 + 1)(\ell_1 + 2) + \ell_2(\ell_2 + 1)$$
(B.11)

$$m_{2,+}^2 = \ell_1(\ell_1+1) + \ell_2(\ell_2-1), \qquad m_{2,-}^2 = \ell_1(\ell_1+1) + (\ell_2+1)(\ell_2+2).$$
 (B.12)

The propagator functions $K^{m_{0,\pm}^2}$ and $K^{m_{i,\pm}^2}$ can again be found in (B.38) below.

For the matrix elements of the $(2\ell + 1) \times (2\ell + 1)$ dimensional representation of the SU(2) generators $t_i^{(2\ell+1)}$, we are using the following shorthand notation:

$$[t_i^{(\ell)}]_{m,-m'} \equiv [t_i^{(2\ell+1)}]_{\ell-m+1,\ell+m'+1}, \qquad i = 1, 2, 3, \tag{B.13}$$

where the $k \times k$ generators are given in terms of the fuzzy SU(2) spherical harmonics \hat{Y}_{ℓ}^m by the following formulae:

$$t_1^{(k)} = \frac{(-1)^{k+1}}{2} \sqrt{\frac{k(k^2 - 1)}{6}} \cdot \left(\hat{Y}_1^{-1} - \hat{Y}_1^1\right)$$
(B.14)

$$t_2^{(k)} = \frac{i(-1)^{k+1}}{2} \sqrt{\frac{k(k^2 - 1)}{6}} \cdot \left(\hat{Y}_1^{-1} + \hat{Y}_1^1\right) \tag{B.15}$$

$$t_3^{(k)} = \frac{(-1)^{k+1}}{2} \sqrt{\frac{k(k^2 - 1)}{3}} \cdot \hat{Y}_1^0.$$
(B.16)

Moreover, the fuzzy SU(2) spherical harmonics satisfy the following identities:

$$(\hat{Y}_{\ell}^{m})^{\dagger} = (-1)^{m} \hat{Y}_{\ell}^{-m} \qquad \& \qquad \operatorname{tr} [\hat{Y}_{\ell}^{m} \hat{Y}_{\ell'}^{m'}] = (-1)^{m'} \delta_{\ell\ell'} \delta_{m+m'}.$$
 (B.17)

 $SO(5)_n$ symmetric dCFT In the case of the $SO(5)_n$ symmetric domain wall, we decompose the scalar fields of the "massive" $d_G \times d_G$ block in fuzzy SO(5) spherical harmonics as follows [38]:

$$\left[\tilde{\varphi}_{i}\right]_{n_{1},n_{2}} = \sum_{\mathbf{L}} \left(\tilde{\varphi}_{i}\right)_{\mathbf{L}} \cdot \left[\hat{Y}_{\mathbf{L}}\right]_{n_{1},n_{2}}, \qquad \operatorname{tr}\left(\hat{Y}_{\mathbf{L}'}^{\dagger}\hat{Y}_{\mathbf{L}}\right) = \delta_{\mathbf{L}',\mathbf{L}}, \qquad (B.18)$$

where i = 1, ..., 5 and $n_1, n_2 = 1, ..., d_G \equiv (n+1)(n+2)(n+3)/6$. We use the bold symbol **L** to denote all the quantum numbers which specify states in representations of SO(5). In particular, $\mathbf{L} = (L_1, L_2)\ell_1\ell_2m_1m_2$, where the quantum numbers L_1 and L_2 label the representation and the orbital/magnetic quantum numbers ℓ_i and m_i obey,

$$-L_1 + L_2 \le \ell_1 - \ell_2 \le L_1 - L_2 \le \ell_1 + \ell_2 \le L_1 + L_2$$
(B.19)

$$\ell_1 + \ell_2 \in \mathbb{Z}, \qquad m_i = -\ell_i, \dots, \ell_i. \tag{B.20}$$

Based on the description (2.5)–(2.6) of the SO(5) symmetric D3-D7 domain wall, we obtain two kinds of scalar propagators in the "massive" block, namely $\langle \tilde{\varphi}_i \tilde{\varphi}_j \rangle$ ("complicated" fields, $i = 1, \ldots, 5$), and $\langle \tilde{\varphi}_6 \tilde{\varphi}_6 \rangle$ ("easy" fields). For the complicated scalars we have,

$$\langle (\tilde{\varphi}_i)_{\mathbf{L}} (\tilde{\varphi}_j)_{\mathbf{L}'}^{\dagger} \rangle = \delta_{ij} \delta_{\mathbf{L},\mathbf{L}'} \hat{f}^{\operatorname{sing}} + \langle \mathbf{L} | L_{ij} | \mathbf{L}' \rangle \hat{f}^{\operatorname{lin}} + \langle \mathbf{L} | \{L_{ik}, L_{jl} \} L_{kl} | \mathbf{L}' \rangle \hat{f}^{\operatorname{cubic}} + \\ + \langle \mathbf{L} | \{L_{ik}, L_{kj} \} | \mathbf{L}' \rangle \hat{f}_5^{\operatorname{sym}} + \langle \mathbf{L} | \{L_{i6}, L_{6j} \} | \mathbf{L}' \rangle \cdot \left[\delta_{L_1, L_1'} \delta_{L_2, L_2'} \hat{f}_6^{\operatorname{sym}} + \\ + \delta_{L_1', L_1 \pm 1} \delta_{L_2', L_2 \mp 1} \hat{f}^{\operatorname{opp}} \right],$$

$$(B.21)$$

where the expressions of the propagator functions f^{sing} , f^{lin} , f^{cubic} , f_5^{sym} , f_6^{sym} , and f^{opp} can be found in appendix C of [38]. In (B.21) we have defined,

$$\langle \mathbf{L} | L_{ij} | \mathbf{L}' \rangle = \operatorname{tr} \left(\hat{Y}_{\mathbf{L}}^{\dagger} L_{ij} \hat{Y}_{\mathbf{L}'} \right) = \frac{i}{2} \cdot \operatorname{tr} \left(\hat{Y}_{\mathbf{L}}^{\dagger} [\hat{Y}_{\mathbf{L}'}, G_{ij}] \right), \qquad G_{i6} = -G_{6i} \equiv G_i, \quad (B.22)$$

for i, j, k = 1, ..., 5 and the commutators G_{ij} of the *G*-matrices were defined in (2.11) above. Also, $L_{ij} = -L_{ji}$ are the usual generators of SO(5) which satisfy,

$$[L_{ij}, L_{kl}] = i \left(\delta_{ik} L_{jl} + \delta_{jl} L_{ik} - \delta_{jk} L_{il} - \delta_{il} L_{jk} \right), \quad i, j, k, l = 1, \dots, 5.$$
(B.23)

For the propagators of the easy scalars we have:

$$\langle \langle (\tilde{\varphi}_6)_{\mathbf{L}} (\tilde{\varphi}_6)_{\mathbf{L}'}^{\dagger} \rangle = \delta_{\mathbf{L},\mathbf{L}'} \cdot K^{\hat{m}_{\text{easy}}^2}, \qquad \hat{m}_{\text{easy}}^2 \equiv 2L_1L_2 + L_1 + 2L_2.$$
(B.24)

As in the case of the SU(2) generators which were expressed in (B.14)–(B.16) by means of the SU(2) fuzzy spherical harmonics, the SO(5) G-matrices (2.8) can be expressed in terms of the SO(5) spherical harmonics as follows:

$$G_1 = \sqrt{\frac{c_n}{10}} \cdot \left(\hat{Y}_{++} + \hat{Y}_{--} \right), \qquad G_2 = -i\sqrt{\frac{c_n}{10}} \cdot \left(\hat{Y}_{++} - \hat{Y}_{--} \right), \qquad (B.25)$$

$$G_3 = -\sqrt{\frac{c_n}{10}} \cdot \left(\hat{Y}_{-+} - \hat{Y}_{+-}\right), \qquad G_4 = -i\sqrt{\frac{c_n}{10}} \cdot \left(\hat{Y}_{-+} + \hat{Y}_{+-}\right), \qquad (B.26)$$

$$G_5 = -\sqrt{\frac{c_n}{5}} \cdot \hat{Y}_{00}, \tag{B.27}$$

where the constant c_n has been defined in (3.26) above, and we have set

$$\hat{Y}_{\alpha\beta} \equiv \hat{Y}_{(\frac{1}{2},\frac{1}{2})\frac{1}{2}\frac{1}{2}\alpha\beta}, \qquad \hat{Y}_{00} \equiv \hat{Y}_{(\frac{1}{2},\frac{1}{2})0000}.$$
(B.28)

By using the relations (B.25)-(B.27) between the *G*-matrices and the fuzzy SO(5) spherical harmonics, we compute the contraction,

$$\overline{\tilde{\varphi}_i \cdot \operatorname{tr}[G_j \tilde{\varphi}_j]} = K^{5/2}(x, y) \cdot G_i.$$
(B.29)

Scalar propagators in AdS The bulk-to-bulk propagator of a massive scalar field in Euclidean AdS_{d+1} is given by the following formula (see e.g. the set of lectures [53]):

$$G_{\Delta}(x,z;y,w) = \frac{\Gamma(\Delta)\tilde{\eta}^{\Delta}}{2^{\Delta+1}\pi^{d/2}\Gamma\left(\Delta-\frac{d}{2}+1\right)} \cdot {}_{2}F_{1}\left(\frac{\Delta}{2},\frac{\Delta+1}{2},\Delta-\frac{d}{2}+1,\tilde{\eta}^{2}\right), \quad (B.30)$$

where Δ is the scaling dimension of the scalar field and \tilde{m} its mass, while we have defined,

$$\tilde{\eta} \equiv \frac{2zw}{z^2 + w^2 + (x - y)^2}, \qquad \tilde{m}^2 \equiv \Delta \left(\Delta - d\right) \equiv m^2 - \frac{d^2 - 1}{4}.$$
 (B.31)

As before, the sets of coordinates $(x, z) = (x_0, \ldots, x_{d-2}, z)$ and $(y, w) = (y_0, \ldots, y_{d-2}, w)$ parametrize (Euclidean) AdS_{d+1} in the Poincaré frame. We may obtain an alternative expression for the scalar propagator (B.30) by using the so-called quadratic transformation formulas for the hypergeometric function (see [54], eq. 15.3.16). We find [55],

$${}_{2}F_{1}\left(\frac{\Delta}{2},\frac{\Delta+1}{2},\Delta-\frac{d}{2}+1,\tilde{\eta}^{2}\right) = \left(1+\frac{1}{2s}\right)^{\nu+\frac{d}{2}} {}_{2}F_{1}\left(\nu+\frac{d}{2},\nu+\frac{1}{2},2\nu+1,-s^{-1}\right), (B.32)$$

where we have defined,

$$s \equiv \frac{1 - \tilde{\eta}}{2\tilde{\eta}} = \frac{(z - w)^2 + (x - y)^2}{4zw}, \qquad \nu \equiv \Delta - \frac{d}{2} = \sqrt{m^2 + \frac{1}{4}}.$$
 (B.33)

The scalar propagator (B.32) can be further transformed by means of the linear transformation formulas of hypergeometric functions (see [54], eq. 15.3.3). We get,

$$_{2}F_{1}\left(\frac{\Delta}{2},\frac{\Delta+1}{2},\Delta-\frac{d}{2}+1,\tilde{\eta}^{2}\right) =$$

$$= \left(1 + \frac{1}{s}\right)^{\frac{1-d}{2}} \left(1 + \frac{1}{2s}\right)^{\nu + \frac{d}{2}} {}_{2}F_{1}\left(\nu - \frac{d}{2} + 1, \nu + \frac{1}{2}, 2\nu + 1, -s^{-1}\right),$$
(B.34)

so that by plugging (B.34) into the expression for the scalar propagator (B.30), we get:

$$G_{\Delta}(x,z;y,w) = \frac{1}{2^{d}\pi^{\frac{d+1}{2}}} \frac{\Gamma\left(\nu + \frac{d}{2}\right)\Gamma\left(\nu + \frac{3}{2}\right)}{\Gamma\left(2\nu + 2\right)} \cdot \frac{{}_{2}F_{1}\left(\nu - \frac{d}{2} + 1,\nu + \frac{1}{2},2\nu + 1,-s^{-1}\right)}{(1+s)^{\frac{d-1}{2}}s^{\nu + \frac{1}{2}}},$$
(B.35)

by also using the Legendre duplication formula for the gamma function,

$$\Gamma\left(\nu+1\right) = \frac{\sqrt{\pi}}{2^{2\nu+1}} \cdot \frac{\Gamma\left(2\nu+2\right)}{\Gamma\left(\nu+\frac{3}{2}\right)}.$$
(B.36)

As it turns out [30, 47], the propagators of the various fields in the (codimension-1) defect CFTs we are examining are related to the corresponding propagators in AdS, where the holographic directions z and w are replaced by the coordinates which are normal to the codimension-1 boundary. In our case the scalar fields propagate inside AdS₄, so that we set d = 3 in the above formulas for the propagators, and $(z, w) \rightarrow (x_3, y_3)$ for a boundary that is located at $x_3 = y_3 = 0$. By defining,

$$K^{m^{2}}(x,y) \equiv \frac{g_{\rm YM}^{2}}{2} \cdot (x_{3}y_{3})^{-\frac{d-1}{2}} \cdot G_{\Delta}\left(\left\{x_{0}, x_{1}, x_{2}\right\}, x_{3}; \left\{y_{0}, y_{1}, y_{2}\right\}, y_{3}\right), \qquad (B.37)$$

we may use the expression (B.35) for the scalar propagator in AdS₄ to obtain,

$$K^{\nu}(x,y) = \frac{g_{\rm YM}^2}{16\pi^2} \frac{1}{\binom{2\nu+1}{\nu+\frac{1}{2}}} \frac{{}_2F_1\left(\nu - \frac{1}{2},\nu + \frac{1}{2},2\nu + 1;-\xi^{-1}\right)}{(1+\xi)\xi^{\nu+\frac{1}{2}}} \cdot \frac{1}{x_3y_3},\tag{B.38}$$

where m is the mass of the scalar field and the dCFT invariant ratio ξ was defined in (3.7):

$$\xi \equiv \frac{|x-y|^2}{4x_3y_3}, \qquad \nu \equiv \sqrt{m^2 + \frac{1}{4}},$$
 (B.39)

for $x = (x_0, \dots, x_3)$ and $y = (y_0, \dots, y_3)$.

References

- C.P. Herzog, K.-W. Huang and K. Jensen, Universal entanglement and boundary geometry in conformal field theory, JHEP 01 (2016) 162 [1510.00021].
- [2] C. Herzog, K.-W. Huang and K. Jensen, Displacement operators and constraints on boundary central charges, Phys. Rev. Lett. 120 (2018) 021601 [1709.07431].
- [3] A. Chalabi, C.P. Herzog, A. O'Bannon, B. Robinson and J. Sisti, Weyl anomalies of four dimensional conformal boundaries and defects, JHEP 02 (2022) 166 [2111.14713].
- [4] H. Osborn and A.C. Petkou, Implications of conformal invariance in field theories for general dimensions, Annals Phys. 231 (1994) 311 [hep-th/9307010].
- [5] J. Erdmenger and H. Osborn, Conserved currents and the energy momentum tensor in conformally invariant theories for general dimensions, Nucl. Phys. B483 (1997) 431 [hep-th/9605009].

- [6] N. Drukker, D. Martelli and I. Shamir, The energy-momentum multiplet of supersymmetric defect field theories, JHEP 08 (2017) 010 [1701.04323].
- [7] C.P. Herzog and K.-W. Huang, Boundary conformal field theory and a boundary central charge, JHEP 10 (2017) 189 [1707.06224].
- [8] D. Fursaev, Conformal anomalies of CFT's with boundaries, JHEP 12 (2015) 112 [1510.01427].
- [9] S.N. Solodukhin, Boundary terms of conformal anomaly, Phys. Lett. B752 (2016) 131 [1510.04566].
- [10] M. de Leeuw, C. Kristjansen, G. Linardopoulos and M. Volk, B-type anomaly coefficients for the D3-D5 domain wall, Phys. Lett. B846 (2023) 138235 [2307.10946].
- [11] J.L. Davis, P. Kraus and A. Shah, Gravity dual of a quantum Hall plateau transition, JHEP 11 (2008) 020 [0809.1876].
- [12] S.-J. Rey, String theory on thin semiconductors: Holographic realization of Fermi points and surfaces, Prog. Theor. Phys. Suppl. 177 (2009) 128 [0911.5295].
- [13] R.C. Myers and M.C. Wapler, Transport properties of holographic defects, JHEP 12 (2008) 115 [0811.0480].
- [14] O. Bergman, N. Jokela, G. Lifschytz and M. Lippert, Quantum Hall effect in a holographic model, JHEP 10 (2010) 063 [1003.4965].
- [15] M. de Leeuw, C. Kristjansen and K.E. Vardinghus, A non-integrable quench from AdS/dCFT, Phys. Lett. B798 (2019) 134940 [1906.10714].
- [16] M. de Leeuw, C. Kristjansen and G. Linardopoulos, Scalar one-point functions and matrix product states of AdS/dCFT, Phys. Lett. B781 (2018) 238 [1802.01598].
- [17] M. de Leeuw, T. Gombor, C. Kristjansen, G. Linardopoulos and B. Pozsgay, Spin chain overlaps and the twisted Yangian, JHEP 01 (2020) 176 [1912.09338].
- [18] T. Gombor and Z. Bajnok, Boundary states, overlaps, nesting and bootstrapping AdS/dCFT, JHEP 10 (2020) 123 [2004.11329].
- [19] G. Linardopoulos, String theory methods for defect CFTs, 2501.11985.
- [20] B. Robinson and C.F. Uhlemann, Supersymmetric D3/D5 for massive defects on curved space, JHEP 12 (2017) 143 [1709.08650].
- B. Robinson, Supersymmetric localization and probe branes in the AdS/CFT correspondence, Ph.D. thesis, Washington U., Seattle, 2017.
- [22] Y. Wang, Taming defects in $\mathcal{N} = 4$ super-Yang-Mills, JHEP 08 (2020) 021 [2003.11016].
- [23] S. Komatsu and Y. Wang, Non-perturbative defect one-point functions in planar $\mathcal{N} = 4$ super-Yang-Mills, Nucl. Phys. B958 (2020) 115120 [2004.09514].
- [24] M. Beccaria and A. Cabo-Bizet, 1/N expansion of the D3-D5 defect CFT at strong coupling, JHEP 02 (2023) 208 [2212.12415].
- [25] D. He and C.F. Uhlemann, One-point functions for doubly-holographic BCFTs and backreacting defects, 2501.07630.
- [26] D.M. McAvity and H. Osborn, Energy momentum tensor in conformal field theories near a boundary, Nucl. Phys. B406 (1993) 655 [hep-th/9302068].

- [27] D.M. McAvity and H. Osborn, Conformal field theories near a boundary in general dimensions, Nucl. Phys. B455 (1995) 522 [cond-mat/9505127].
- [28] P. Liendo, L. Rastelli and B.C. van Rees, The bootstrap program for boundary CFT_d , JHEP 07 (2013) 113 [1210.4258].
- [29] C. Kristjansen, G.W. Semenoff and D. Young, Chiral primary one-point functions in the D3-D7 defect conformal field theory, JHEP 01 (2013) 117 [1210.7015].
- [30] K. Nagasaki, H. Tanida and S. Yamaguchi, *Holographic interface-particle potential*, JHEP 01 (2012) 139 [1109.1927].
- [31] K. Nagasaki and S. Yamaguchi, Expectation values of chiral primary operators in holographic interface CFT, Phys. Rev. D86 (2012) 086004 [1205.1674].
- [32] N.R. Constable, R.C. Myers and O. Tafjord, The noncommutative bion core, Phys. Rev. D61 (2000) 106009 [hep-th/9911136].
- [33] N.R. Constable, R.C. Myers and O. Tafjord, Non-abelian brane intersections, JHEP 06 (2001) 023 [hep-th/0102080].
- [34] J. Castelino, S. Lee and W. Taylor, Longitudinal 5-branes as 4-spheres in matrix theory, Nucl. Phys. B526 (1998) 334 [hep-th/9712105].
- [35] M. de Leeuw, C. Kristjansen and G. Linardopoulos, One-point functions of non-protected operators in the SO(5) symmetric D3-D7 dCFT, J. Phys. A50 (2017) 254001 [1612.06236].
- [36] M. de Leeuw, C. Kristjansen and K. Zarembo, One-point functions in defect CFT and integrability, JHEP 08 (2015) 098 [1506.06958].
- [37] A. Gimenez-Grau, C. Kristjansen, M. Volk and M. Wilhelm, A quantum check of non-supersymmetric AdS/dCFT, JHEP 01 (2019) 007 [1810.11463].
- [38] A. Gimenez-Grau, C. Kristjansen, M. Volk and M. Wilhelm, A quantum framework for AdS/dCFT through fuzzy spherical harmonics on S⁴, JHEP 04 (2020) 132 [1912.02468].
- [39] L. Piroli, B. Pozsgay and E. Vernier, What is an integrable quench?, Nucl. Phys. B925 (2017) 362 [1709.04796].
- [40] S. Ghoshal and A.B. Zamolodchikov, Boundary S-matrix and boundary state in two-dimensional integrable quantum field theory, Int. J. Mod. Phys. A9 (1994) 3841 [hep-th/9306002].
- [41] M. de Leeuw, One-point functions in AdS/dCFT, J. Phys. A53 (2020) 283001 [1908.03444].
- [42] G. Linardopoulos, Solving holographic defects, PoS Corfu2019 (2020) 141 [2005.02117].
- [43] C. Kristjansen and K. Zarembo, Integrable holographic defect CFTs, 2401.17144.
- [44] D. Anselmi, D.Z. Freedman, M.T. Grisaru and A.A. Johansen, Nonperturbative formulas for central functions of supersymmetric gauge theories, Nucl. Phys. B526 (1998) 543
 [hep-th/9708042].
- [45] D.Z. Freedman, S.D. Mathur, A. Matusis and L. Rastelli, Correlation functions in the CFT_d/AdS_{d+1} correspondence, Nucl. Phys. B546 (1999) 96 [hep-th/9804058].
- [46] P.S. Howe, E. Sokatchev and P.C. West, Three point functions in $\mathcal{N} = 4$ Yang-Mills, *Phys. Lett.* B444 (1998) 341 [hep-th/9808162].
- [47] I. Buhl-Mortensen, M. de Leeuw, A.C. Ipsen, C. Kristjansen and M. Wilhelm, A quantum check of AdS/dCFT, JHEP 01 (2017) 098 [1611.04603].

- [48] N.D. Birrell and P.C.W. Davies, *Quantum fields in curved space*, Cambridge University Press, Cambridge (1999).
- [49] G. Georgiou, G. Linardopoulos and D. Zoakos, *Holographic correlators of semiclassical states in defect CFTs*, *Phys. Rev.* D108 (2023) 046016 [2304.10434].
- [50] G. Linardopoulos, B-type anomaly coefficients from holographic defects (to appear), PoS Corfu2024 (2025).
- [51] C.G. Callan Jr., S.R. Coleman and R. Jackiw, A new improved energy-momentum tensor, Annals Phys. 59 (1970) 42.
- [52] H. Osborn, Lectures on conformal field theories in more than two dimensions, 2019.
- [53] E. D'Hoker and D.Z. Freedman, Supersymmetric gauge theories and the AdS/CFT correspondence, hep-th/0201253.
- [54] M. Abramowitz and I. Stegun, eds., Handbook of mathematical functions, Dover, New York (1972).
- [55] E. D'Hoker and D.Z. Freedman, General scalar exchange in AdS(d+1), Nucl. Phys. B550 (1999) 261 [hep-th/9811257].