Exact Results for SYM on $Y^{p,q}$ and $S^2 \times S^2$ with Conical Singularities

Lorenzo Ruggeri¹

¹Yau Mathematical Sciences Center, Tsinghua University, Beijing, 100084, China

E-mail: ruggeri@mail.tsinghua.edu.cn

ABSTRACT: We compute the full partition function, including flux and instanton contributions, for an $\mathcal{N} = 1$ theory of vector multiplets and hypermultiplets on five-dimensional toric Sasakian manifolds $Y^{p,q}$. Dimensionally reducing, we obtain the partition function on a class of manifolds whose topology is $S^2 \times S^2$, for equivariant Donaldson-Witten and Pestun-like theories. Generalizing the procedure to branched covers of $Y^{p,q}$, and exploiting an equivalence with spaces containing orbifold singularities, we compute, for instance, the partition function of an $\mathcal{N} = 2$ theory on the product of two spindles.

Contents

| 2 Toric Sasakian Geometry Toric Sasakian Manifolds Branched Covers and Orbifolds 3 5d Partition Function Localization Perturbative Partition Function 4 Fluxes Via Dimensional Reduction Dimensional Reduction d to 5d d to 5d d to 4d Examples 5 Factorized Expressions and Instantons 6 Discussion | 1 |
|---|----|
| 2.1 Toric Sasakian Manifolds 2.2 Branched Covers and Orbifolds 3 5d Partition Function 3.1 Localization 3.2 Perturbative Partition Function 4 Fluxes Via Dimensional Reduction 4.1 Dimensional Reduction 4.2 6d to 5d 4.3 5d to 4d 4.4 Examples 5 Factorized Expressions and Instantons 6 Discussion | 3 |
| 2.2 Branched Covers and Orbifolds 3 5d Partition Function 3.1 Localization 3.2 Perturbative Partition Function 4 Fluxes Via Dimensional Reduction 4.1 Dimensional Reduction 4.2 6d to 5d 4.3 5d to 4d 4.4 Examples 5 Factorized Expressions and Instantons 6 Discussion | 3 |
| 3 5d Partition Function 3.1 Localization 3.2 Perturbative Partition Function 4 Fluxes Via Dimensional Reduction 4.1 Dimensional Reduction 4.2 6d to 5d 4.3 5d to 4d 4.4 Examples 5 Factorized Expressions and Instantons 6 Discussion | 6 |
| 3.1 Localization 3.2 Perturbative Partition Function 4 Fluxes Via Dimensional Reduction 4.1 Dimensional Reduction 4.2 6d to 5d 4.3 5d to 4d 4.4 Examples 5 Factorized Expressions and Instantons 6 Discussion | 8 |
| 3.2 Perturbative Partition Function Fluxes Via Dimensional Reduction 4.1 Dimensional Reduction 4.2 6d to 5d 4.3 5d to 4d 4.4 Examples 5 Factorized Expressions and Instantons 6 Discussion | 9 |
| Fluxes Via Dimensional Reduction 4.1 Dimensional Reduction 4.2 6d to 5d 4.3 5d to 4d 4.4 Examples 5 Factorized Expressions and Instantons 6 Discussion | 10 |
| 4.1 Dimensional Reduction 4.2 6d to 5d 4.3 5d to 4d 4.4 Examples 5 Factorized Expressions and Instantons 6 Discussion | 14 |
| 4.2 6d to 5d 4.3 5d to 4d 4.4 Examples 5 Factorized Expressions and Instantons 6 Discussion | 14 |
| 4.3 5d to 4d 4.4 Examples 5 Factorized Expressions and Instantons 6 Discussion | 16 |
| 4.4 Examples 5 Factorized Expressions and Instantons 6 Discussion | 18 |
| 5 Factorized Expressions and Instantons6 Discussion | 20 |
| 6 Discussion | 21 |
| | 24 |

1 Introduction

The study of supersymmetric quantum field theories (SQFTs) on non-trivial geometric backgrounds, pioneered in [1, 2], has provided profound insights into both physical phenomena and mathematical structures. Among these, five-dimensional toric Sasakian geometries [3–5] have emerged as a particularly rich setting due to their applications in the AdS/CFT correspondence [6–8], their connection to refined topological strings [9] and as a tool to compute partition function of 4d theories via dimensional reduction [10–12]. In [13–15], the partition function of an $\mathcal{N} = 1$ SQFT has been computed on the simplest instance of toric Sasakian manifold, the five-sphere S^5 . Later, these results have been extended in [16, 17] to manifolds which are topologically $S^2 \times S^3$, namely $Y^{p,q}$ [8] and $L^{a,b,c}$ [18]. Eventually, the partition function on a generic toric Sasakian manifold has been obtained in [19, 20].

In these results, the gauge configurations considered are either trivial connections or contact instantons. However, whenever a manifold has non-trivial two cycles, it is expected that gauge configurations with flux are allowed and the partition function is a sum over topological sector labelled by their first Chern class. For instance, such gauge configurations are expected on $Y^{p,q}$ and $L^{a,b,c}$, as $H_2(S^2 \times S^3, \mathbb{Z}) \simeq \mathbb{Z}$. One of the main results of this work is to compute their contributions to the partition function. Hence, we are able to provide the full (integrand of the) partition function for an $\mathcal{N} = 1$ vector multiplet coupled to a hypermultiplet in a representation R of the gauge group on both $Y^{p,q}$ and $L^{a,b,c}$.

Focusing first on the trivial contact instanton sector, we compute the one-loop determinant around flux configurations. The idea is to uplift the cohomological complex obtained via localization on $Y^{p,q}$ (or $L^{a,b,c}$) [16] to $S^3 \times S^3$ and compute the equivariant index of the resulting complex in 6d. The index can be decomposed in contributions from different representations under a freely acting S_Y^1 . Keeping only the trivial representation, the authors of [16] computed the perturbative partition function on $(S^3 \times S^3)/S_Y^1 \simeq Y^{p,q}$. It has been recently shown in¹ [11, 12] that, when dimensionally reducing along a non-trivially fibered S^1 , modes with non-zero charge for rotations under the S^1 contribute to different flux sectors on the base manifold. Hence, to compute the one-loop determinant at a given flux sector on $Y^{p,q}$ we simply have to restrict the 6d equivariant index to contributions with a given representation under S_Y^1 . Note that, rather than shrinking the radius of S_Y^1 , we perform the dimensional reduction by introducing a finite quotient $(S^3 \times S^3)/\mathbb{Z}_h$ and then taking the large h limit.

The manifold $Y^{p,q}$ is a non-trivial principal bundle over a four-manifold B whose topology is that of the product of two spheres and $H_2(B,\mathbb{Z}) \simeq \mathbb{Z}^2$. The non-trivial fibration has been exploited in [12] to compute, for an $\mathcal{N} = 2$ vector multiplet, the one-loop determinant on B around a single flux starting from the perturbative partition function on $Y^{p,q}$ [16]. Since we have computed the one-loop determinant around fluxes in 5d, the dimensional reduction produces the one-loop determinant around both fluxes on B. Moreover, with respect to a choice of the Killing vector in 5d, there exist two inequivalent S^1 -actions on $Y^{p,q}$. Therefore, dimensionally reducing we obtain two theories: one which is an equivariant version of Donaldson-Witten theory [22] and another which is a generalization of Pestun's theory on S^4 [1] to more generic four-manifolds [10, 23].

To compute the dependence on fluxes of the instanton part, we first show how in 5d (4d) the one-loop determinant around fluxes can be factorized in contributions from the fixed fibers (fixed points) under the torus action of the manifold [24, 25], with fluxes entering as a shift of the Coulomb branch parameter. This behaviour has been shown to hold in [11, 12] for a single flux on a large class of quasi-toric four-manifolds. It is conjectured that the same shift encodes also the dependence of the instanton part [10, 23, 26]. Under this assumption, we can write down the full partition function², including contact instantons on $Y^{p,q}$ and $L^{a,b,c}$, and instantons on B.

Another class of spaces that has drawn a lot of attention recently, following the finding of black hole solutions with near horizon geometries containing orbifold singularities [30– 32], is that of weighted projective spaces \mathbb{CP}_p^r , also known as spindles in 2d. Recently [33], the partition function of an $\mathcal{N} = 2$ theory on $\mathbb{CP}_p^1 \times S^1$ [34, 35] has been shown to reproduce the entropy of an AdS₄ black hole with spindle horizon, paving the way for studying the

¹See [21] for an earlier result, where the large h limit of the $\mathcal{N} = 2$ partition function on lens spaces L(h, -1) has been shown to reproduce the $\mathcal{N} = (2, 2)$ one-loop determinant around fluxes on S^2 .

²Note that full partition functions on complex toric surfaces have been studied also in [27-29]. However, in that case fluxes are equivariant fluxes further constrained by stability conditions. In our work all expressions are in terms of physical fluxes.

AdS/CFT correspondence in a setup with singularities.

These recent developments demand for a more detailed understanding of SQFTs on spaces with orbifold singularities. In [36]³, the full partition function on \mathbb{CP}_p^2 has been computed exploiting a conjectured equivalence with the partition function on a branched cover of \mathbb{CP}^2 . This conjecture has been further motivated in [38] for a vector multiplet and a hypermultiplet in the adjoint representation, and it relies on the interpretation of SQFTs near orbifold singularities in terms of multi-valued fields. Hence, by starting from the equivariant index on the branched cover of $S^3 \times S^3$, and repeating the previous steps, we compute the full partition function also on orbifolds and branched covers whose underlying spaces are $Y^{p,q}$, $L^{a,b,c}$ and B. In particular, in 4d we study $\mathcal{N} = 2$ theories on a space whose topology is that of the product of two spindles.

The outlook is as follows. In section 2 we introduce some basic facts about toric Sasakian geometry, explaining in some details the dimensional reductions from $S^3 \times S^3$ down to $Y^{p,q}$ and B. We further discuss the corresponding spaces with either surplus or deficit angles. Then, in section 3, we present the perturbative partition function on the five-dimensional spaces, also in the setup with singularities. In section 4 we perform the dimensional reduction, first from 6d to 5d and then from 5d to 4d, which enables us to compute the one-loop determinant around fluxes on these spaces. The factorisation properties of the partition functions are studied in section 5 where, employing the shifts derived from the one-loop part around fluxes, we can write the full partition function, including instanton contributions. We summarise our results in section 6 and comment on possible future directions.

2 Toric Sasakian Geometry

In this section, we briefly recall the geometric properties of certain toric Sasakian manifolds, $Y^{p,q}$ [8] and $L^{a,b,c}$ [18], whose topology is $S^2 \times S^3$. We refer to [3–5, 39] for a detailed account of Sasakian and toric geometry. Here, we focus on showing how both $Y^{p,q}$ and $L^{a,b,c}$ arise via free S^1 -quotients of $S^3 \times S^3$. We will exploit this fact to compute the one-loop determinant at all flux sectors on these manifolds in section 4. We also introduce branched covers and orbifolds whose underlying spaces are $Y^{p,q}$ and $L^{a,b,c}$, but with conical singularities associated to either surplus or deficit angles.

Finally, we will introduce $B = Y^{p,q}/S^1$, a class of quasi-toric manifolds whose topology is that of a product of two spheres, and the corresponding branched covers and orbifolds. In the latter case, the resulting spaces have the topology of the product of two spindles.

2.1 Toric Sasakian Manifolds

Sasakian manifolds (M, g) are defined as Riemannian manifolds whose metric cone $(C(M), \bar{g})$ is Kähler

$$C(M) = M \times \mathbb{R}_{>0}, \qquad \bar{g} = dr^2 + r^2 g. \tag{2.1}$$

³See also [37] for similar results concerning the equivariant index on \mathbb{CP}^2_{p} .

The existence of an effective, holomorphic and Hamiltonian torus action on the Kähler cone leads to the definition of toric Sasakian manifolds. Moreover, C(M) can be obtained as a Kähler quotient

$$C(M) = \mathbb{C}^m / / U(1)^{m-3}, \tag{2.2}$$

where $U(1)^{m-3}$ acts with charges

$$\vec{Q}_a^M = [Q_a^1, \dots, Q_a^m], \qquad a = 1, \dots, m - 3,$$
(2.3)

on \mathbb{C}^m . If the charges satisfy the m-3 conditions $Q_a^1 + \cdots + Q_a^m = 0$, the cone is Calabi-Yau and M is said to be Sasaki-Einstein.

Restricting to five-dimensional simply connected toric Sasakian manifolds, the simplest example, with m = 3, is S^5 whose metric cone is $\mathbb{C}^3/\{0\}$. Moving to m = 4, there are two classes of manifolds, $Y^{p,q}$ and $L^{a,b,c}$, whose metric cones are defined by the charges of the U(1) in (2.2)

$$\mathbb{C}^{m}//S_{Y}^{1} = C(Y^{p,q}): \qquad \vec{Q}^{Y} = [p+q, p-q, -p, -p], \quad p > q > 0, \\ \mathbb{C}^{m}//S_{L}^{1} = C(L^{a,b,c}): \qquad \vec{Q}^{L} = [a, b, -c, -a - b + c], \quad a, b, a + b - c > 0,$$
(2.4)

where gcd(p,q) = gcd(a,b) = gcd(a,c) = gcd(b,c) = 1. Each component of these fourvectors rotates the phase of each \mathbb{C} factor. As the sum of the charges in (2.4) vanishes in both cases, these two families of manifolds are Sasaki-Einstein.

Let us explicitly construct the class of $Y^{p,q}$ manifolds starting from \mathbb{C}^4 , following the presentation in [16, 20]. The Kähler quotient consists in

$$C(Y^{p,q}) = \mu_{S_Y^1}^{-1}(0)/S_Y^1, \tag{2.5}$$

where the moment map for S_Y^1 is given by

$$\mu_{S_Y^1} = (p+q)|z_1|^2 + (p-q)|z_2|^2 - p|z_3|^2 - p|z_4|^2.$$
(2.6)

To further obtain $Y^{p,q}$ from (2.5) we need to fix the scaling freedom in (2.1), which we achieve by restricting to a hypersurface

$$H_{\omega} = \{ z_i \in \mathbb{C} \mid \sum_{i=1}^{4} \omega_i |z_i|^2 = 1 \},$$
(2.7)

where $\omega_i \in \mathbb{R}$. We choose all $\omega_i > 0$, so that the intersection $\mu_{S_Y^1}^{-1}(0) \cap H_{\omega}$ is topologically $S^3 \times S^3$. Moreover, as both the loci $z_1 = z_2 = 0$ and $z_3 = z_4 = 0$ are not part of the intersection, the S_Y^1 -action acts freely on $\mu_{S_Y^1}^{-1}(0) \cap H_{\omega}$. Hence, we find

$$Y^{p,q} = (\mu_{S_Y^1}^{-1}(0) \cap H_\omega) / S_Y^1.$$
(2.8)

A possible choice for the hypersurface is the following

$$(p+q)|z_1|^2 + (p-q)|z_2|^2 + p|z_3|^2 + p|z_4|^2 = 1,$$
(2.9)



Figure 1. Toric diagrams for $Y^{p,q}$ and $L^{a,b,c}$. Left side: $Y^{4,3}$. Right side: $L^{1,5,2}$. The facets C_i label the codimension two submanifolds kept fixed by an S^1 in the T^3 Cartan isometry. The four fixed fibers of each manifold are at the intersection of two facets.

so that $S^3 \times S^3 \simeq \mu_{S^1_Y}^{-1}(0) \cap H_\omega$ is given by

$$S^{3} \times S^{3} \simeq \{(z_{1}, z_{2}, z_{3}, z_{4}) \in \mathbb{C}^{4} \mid (p+q)|z_{1}|^{2} + (p-q)|z_{2}|^{2} = \frac{1}{2}, \ p|z_{3}|^{2} + p|z_{4}|^{2} = \frac{1}{2}\}.$$
(2.10)

The same construction can be applied to $L^{a,b,c}$ manifolds, simply by using the charges \vec{Q}^L in (2.4).

Exploiting the toric structure, both $Y^{p,q}$ and $L^{a,b,c}$ can be seen as a Lagrangian T^3 fibration over their corresponding moment polytopes [40], shown in Figure 1. In both cases, each of the four facets C_i represents a codimension two submanifold where a circle subgroup of the T^3 Cartan isometry degenerates. In particular, for $Y^{p,q}$, these are given by lens spaces

$$C_1 \cong C_2 \cong S^3 / \mathbb{Z}_p, \qquad C_3 \cong S^3 / \mathbb{Z}_{p+q}, \qquad C_4 \cong S^3 / \mathbb{Z}_{p-q}.$$
 (2.11)

There are also four codimension four submanifolds which are fixed under a $T^2 \subset T^3$ action. These are given by the intersection of two facets $C_i \cap C_{i+1}$ where, locally, the manifold looks like $\mathbb{C}^2 \times S^1$.

Whenever a toric Sasakian manifolds M admits a globally free S^1 -action, the quotient leads to a quasi-toric four-manifold M/S^1 [10–12]. As $L^{a,b,c}$ does not admit any free S^1 action, we focus on $Y^{p,q}$. This manifold is a principal S^1 -bundle over the product of an axially squashed and a round sphere $B = \mathbb{CP}^1 \times \mathbb{CP}^1$, such that the first Chern number on the standard two-cycles are p and q.

To explicitly write the fibers, we perform a change of basis which better suits this setup with a reduced torus action. So far, we wrote four-vectors in terms of a basis $[e_1, e_2, e_3, e_4]$, where each e_i is a vector field generating rotations of the phase of each \mathbb{C} factor. We introduce a new basis $[\tilde{e}_1, \tilde{e}_2, \tilde{e}_3, e_u]$ where the first three components generate the T^3 -action in $Y^{p,q}$ and e_t generates the free S_Y^1 -action (2.4) in $S^3 \times S^3$. Thus, when studying the five-dimensional setup, we can simply forget about the last component and write vectors in terms of $\tilde{e}_1, \tilde{e}_2, \tilde{e}_3$].

On $Y^{p,q}$ there are two inequivalent⁵ free S^1 -actions generated by⁶

$$\vec{\mathbf{x}}^{\text{top}} = [1, 1, 0], \qquad \vec{\mathbf{x}}^{\text{ex}} = [1, -1, 0].$$
 (2.12)

Upon dimensional reduction to 4d, they lead to different theories ("top", "ex" for topological, exotic; see discussion in [10, 23]). The quotient $B = Y^{p,q}/S^1$ can be seen as a T^2 -fibration over the polytope on the left-hand side in Figure 1. In this case, the facets label different \mathbb{CP}^1 's and their four intersections label the fixed points of the manifold.

2.2 Branched Covers and Orbifolds

We now introduce spaces homeomorphic and locally isometric to the classes of toric Sasakian manifolds we introduced above. These spaces are characterized by conical singularities, associated to either surplus angles (branched covers) or deficit angles (orbifolds), at the facets C_i of the toric diagrams in Figure 1. SQFTs on spaces with conical singularities and underlying manifold the five-sphere have been discussed in [36, 41–43].

Branched Covers. Let us start from spaces with surplus angles, which we denote $\widehat{Y}_{p}^{p,q}$ and $\widehat{L}_{p}^{a,b,c}$, with $p = (p_1, p_2, p_3, p_4)$, $|p| = p_1 p_2 p_3 p_4$, and all p_i are pairwise coprimes. These branched covers can be seen as a T^3 -fibration over the same polytope of the corresponding toric Sasakian manifold in Figure 1. In the interior of the polytope these spaces are a |p|-fold cover of either $Y^{p,q}$ or $L^{a,b,c}$. However, on \mathring{C}_i , the interior of a facet, they are a $|p|/p_i$ -fold cover of the underlying manifold, with branching index p_i . At the intersection of two facets $C_i \cap C_{i+1}$, they are a $|p|/p_i p_{i+1}$ -fold cover with branching index $p_i p_{i+1}$. The branching indices for $\widehat{Y}_{p}^{p,q}$ and $\widehat{L}_{p}^{a,b,c}$ are shown in Figure 2. For $\widehat{Y}_{p}^{p,q}$, the facets are thus branched covers of the lens spaces in (2.11)

$$C_{1} \cong \widehat{S}_{(p_{2}p_{3}, p_{4}p_{3})}^{3} / \mathbb{Z}_{p}, \qquad C_{2} \cong \widehat{S}_{(p_{3}p_{4}, p_{1}p_{4})}^{3} / \mathbb{Z}_{p}, C_{3} \cong \widehat{S}_{(p_{4}p_{1}, p_{2}p_{1})}^{3} / \mathbb{Z}_{p+q}, \qquad C_{4} \cong \widehat{S}_{(p_{1}p_{2}, p_{3}p_{2})}^{3} / \mathbb{Z}_{p-q}.$$

$$(2.13)$$

Notice that the branching indices of these lens spaces are not coprimes.

We now repeat the steps above that led us to view $Y^{p,q}$ and $L^{a,b,c}$ as S^1 -quotients of $S^3 \times S^3$ starting, this time, from a **p**-branched cover of \mathbb{C}^4 . This space is obtained from \mathbb{C}^4 by enlarging the periodicities of the angles $\tilde{\theta}_i \in [0, 2\pi p_i)$ and it has conical singularities

⁴This choice of basis for the T^3 -action on $Y^{p,q}$ is related to that in [12, 20] via an $SL(3, \mathbb{Z})$ transformation. The explicit matrix is shown in section 3.2 of [19].

⁵They are inequivalent with respect to a choice of Killing vector employed in the localization computation [12].

⁶To compute these vectors one has to first find the inward-pointing normals \vec{v}_i of the cone given by $\mu(C(Y^{p,q})) \cup \{0\}$. In our conventions these are: $\vec{v}_1 = [1,0,0], \vec{v}_2 = [(p+q)/p, (p-q)/p, -1], \vec{v}_3 = [0,1,0]$ and $\vec{v}_4 = [0,0,1]$. Then, the vector field generating the free S^1 -action is determined solving $\vec{x} \cdot (\vec{v}_i \times \vec{v}_{i+1}) = \pm 1$, $\forall i = 1, \ldots, m$. The \pm sign determines whether upon dimensional reduction the field strength localizes to instantons or anti-instantons at the *i*-th fixed point [10].



Figure 2. Left side: branching structure of $\widehat{Y}_{p}^{4,3}$. Right side: branching structure of $\widehat{L}_{p}^{1,5,2}$.

whenever one of the $\tilde{z}_i = \rho_i e^{i\tilde{\theta}_i}$ vanishes. The five-dimensional branched covers are found as follows

$$\widehat{Y}^{p,q}_{p} = (\mu^{-1}_{S^{1}_{\widehat{Y}}}(0) \cap H_{\omega})/S^{1}_{\widehat{Y}},
\widehat{L}^{a,b,c}_{p} = (\mu^{-1}_{S^{1}_{\widehat{L}}}(0) \cap H_{\omega})/S^{1}_{\widehat{L}},$$
(2.14)

where the $S_{\widehat{Y}}^1$ and $S_{\widehat{L}}^1$ -action have charges

$$\vec{Q}^{\hat{Y}} = \left[\frac{p+q}{p_1}, \frac{p-q}{p_2}, -\frac{p}{p_3}, -\frac{p}{p_4}\right],$$

$$\vec{Q}^{\hat{L}} = \left[\frac{a}{p_1}, \frac{b}{p_2}, -\frac{c}{p_3}, -\frac{a+b-c}{p_4}\right].$$
(2.15)

The action is scaled with respect to (2.4) to take into account the larger periodicities of the angles $\tilde{\theta}_i$.

We focus on $\widehat{Y}_{p}^{p,q}$, whose corresponding moment map is given by

$$\mu_{S_{\hat{Y}}^1} = \frac{(p+q)}{p_1} |\tilde{z}_1|^2 + \frac{(p-q)}{p_2} |\tilde{z}_2|^2 - \frac{p}{p_3} |\tilde{z}_3|^2 - \frac{p}{p_4} |\tilde{z}_4|^2.$$
(2.16)

We modify accordingly also the intersection of $\widehat{\mathbb{C}}_{p}^{4}$ with the hypersurface H_{ω} , which is now a branched cover \widehat{S}_{p}^{7} of a seven-sphere

$$\frac{(p+q)}{p_1}|\tilde{z}_1|^2 + \frac{(p-q)}{p_2}|\tilde{z}_2|^2 + \frac{p}{p_3}|\tilde{z}_3|^2 + \frac{p}{p_4}|\tilde{z}_4|^2 = 1.$$
(2.17)

Combining, as before, (2.16)-(2.17), we find

$$\mu_{S^1_{\widehat{Y}}}^{-1}(0) \cap \widehat{S}^7_{\boldsymbol{p}} \simeq \widehat{S}^3_{\boldsymbol{p}_A} \times \widehat{S}^3_{\boldsymbol{p}_B}, \tag{2.18}$$

where each factor is a branched cover of S^3 and we defined $\boldsymbol{p}_A = (p_1, p_2)$ and $\boldsymbol{p}_B = (p_3, p_4)$. For the same argument as in the smooth case, the $S^1_{\hat{Y}}$ -action is globally free and $(\widehat{S}^3_{\boldsymbol{p}_A} \times \widehat{S}^3_{\boldsymbol{p}_B})/S^1_{\hat{Y}} = \widehat{Y}^{p,q}_{\boldsymbol{p}}$. The generalization to $\widehat{L}^{a,b,c}_{\boldsymbol{p}}$ is straightforward.

Finally, considering the quotient $\widehat{Y}_{p}^{p,q}/S^{1}$ we find a branched cover \widehat{B}_{p} whose topology is that of $\widehat{\mathbb{CP}}_{p_{A}}^{1} \times \widehat{\mathbb{CP}}_{p_{B}}^{1}$, that is the product of two branched covers of a sphere. To write the components of the vector field generating this S^{1} -action, we perform the same change of basis discussed in the smooth case. Moreover, as in (2.15), we need to adjust (2.12) to take into account the larger periodicities of the angles. Thus, we find

$$\vec{\mathbf{x}}_{p}^{\text{top}} = \left[\frac{1}{p_{1}}, \frac{1}{p_{2}}, 0\right], \qquad \vec{\mathbf{x}}_{p}^{\text{ex}} = \left[\frac{1}{p_{1}}, -\frac{1}{p_{2}}, 0\right].$$
 (2.19)

These four-dimensional branched covers can be viewed as T^2 -fibrations over the polytope in Figure 2. At a generic point in the interior, this space is a $|\mathbf{p}|$ -fold cover of $\mathbb{CP}^1 \times \mathbb{CP}^1$. Each interior of a facet C_i , which is a branched cover of \mathbb{CP}^1 , has an associated branching index p_i . At the fixed points $C_i \cap C_{i+1}$ the branching indices are $p_i p_{i+1}$ and the space locally looks like $\widehat{\mathbb{C}}_{p_i} \times \widehat{\mathbb{C}}_{p_{i+1}}$.

Orbifolds. According to the results in [38], partition functions of vector multiplets and hypermultiplets transforming in the adjoint of the gauge group on branched covers match those computed on orbifolds with the same underlying manifold and the same singularity structure. Thus, we will also consider spaces $Y_p^{p,q}$ and $L_p^{a,b,c}$ which are equivalent to the underlying manifolds at a generic point in the interior of the polytopes in Figure 1. However, in the interior of each facet C_i there is an orbifold singularity and the space locally looks like $(\mathbb{C}/\mathbb{Z}_{p_i}) \times \mathbb{C} \times S^1$. At the intersection of two facets, the space is locally $(\mathbb{C}/\mathbb{Z}_{p_i}) \times (\mathbb{C}/\mathbb{Z}_{p_{i+1}}) \times S^1$. We consider spaces whose distribution of orbifold singularities is that appearing in Figure 2 although, in this case, the p_i 's label deficit angles.

Also in this case, $Y_{\boldsymbol{p}}^{p,q}$ is a principal S^1 -orbibundle over a four-dimensional orbifold $B_{\boldsymbol{p}}$ whose topology is that of the product of two spindles $\mathbb{CP}^1_{\boldsymbol{p}_A} \times \mathbb{CP}^1_{\boldsymbol{p}_B}$.

3 5d Partition Function

In this section we consider an $\mathcal{N} = 1$ vector multiplet and a hypermultiplet in a representation R of the gauge group G on two classes of toric Sasakian manifolds, $Y^{p,q}$ and $L^{a,b,c}$, and on the corresponding branched covers $\widehat{Y}_{p}^{p,q}$ and $\widehat{L}_{p}^{a,b,c}$. In the smooth case, the perturbative partition function has been computed in [16]. Here, we briefly summarize their derivation which we then generalize, first to compute the perturbative partition function on $\widehat{Y}_{p}^{p,q}$ and $\widehat{L}_{p}^{a,b,c}$ and then, in section 4, to derive the one-loop determinant at all flux sectors. Contact instantons also contribute to the partition function but their analysis is left for section 5.

We will work with cohomologically twisted theory throughout this work, see [13] for a definition of the cohomological variables and the corresponding complex. Defining cohomological variables for hypermultiplets [44] requires the existence of a spin, or a spin^c, structure. The toric Sasakian manifolds we consider, and the 4d quotients, are spin. Hence, one can switch between the physical and cohomologically twisted theories bijectively. Moreover, we will assume that a spin^c structure can be defined on the branched covers and orbifolds where we will place our theories.

Finally, as shown in [38], all the results we compute on branched covers, for a vector multiplet or a hypermultiplet in the adjoint representation⁷, hold also for the corresponding spaces with orbifold singularities introduced in section 2.

3.1 Localization

The full partition function on a branched cover of a generic toric Sasakian manifold⁸ \widehat{M}_{p} is given by

$$\mathcal{Z}_{\widehat{M}_{p}} = \sum_{\mathfrak{m}_{1},\dots,\mathfrak{m}_{b_{2}}} \int_{\mathfrak{h}} \mathrm{d}a \, e^{-S_{\mathrm{cl}}} \cdot Z_{\widehat{M}_{p}} \cdot Z_{\widehat{M}_{p}}^{\mathrm{inst}}.$$
(3.1)

The sum is over gauge fluxes $\mathfrak{m}_i = \operatorname{diag}(m_{i\,1}, \ldots, m_{i\,\mathrm{rk}\,G}) \in \mathbb{Z}^{\operatorname{rk}\,G \times \operatorname{rk}\,G}$

$$\frac{1}{2\pi} \int_{\Sigma_i} F = \mathfrak{m}_i, \qquad i = 1, \dots, b_2,$$
(3.2)

where b_2 is the second Betti number of \widehat{M}_p and Σ_i is a basis of $H_2(\widehat{M}_p, \mathbb{Z})$. The spaces we consider have non trivial second homology group, $H_2(Y^{p,q}, \mathbb{Z}) \simeq H_2(L^{a,b,c}, \mathbb{Z}) \simeq \mathbb{Z}$, and thus we expect a sum over a single flux in the partition function⁹. Moreover, the integral with respect to the Coulomb branch parameter a is over the Cartan subalgebra \mathfrak{h} of the gauge group G.

The classical contribution is the non Q-exact action evaluated on the BPS locus. At the trivial flux sector, this is given by

$$S_{\rm cl}|_{(\mathfrak{m}_1,\dots,\mathfrak{m}_{b_2})=(0,\dots,0)} = -\frac{8\pi^3}{g_{\rm YM}^2} \rho \operatorname{tr}(a^2), \qquad \rho = \frac{\operatorname{Vol}_{\widehat{M}_{\boldsymbol{p}}}}{\operatorname{Vol}_{S^5}}.$$
(3.3)

The other terms in the integrand are the 1-loop determinant and the contact instanton contributions at each flux sector.

In the absence of conical singularities, and restricting to the trivial flux sector, the perturbative partition function is given by [16, 20]

$$Z_M|_{(\mathfrak{m}_1,\dots,\mathfrak{m}_{b_2})=(0,\dots,0)} = \frac{\det_{\mathrm{adj}}' S_3^{\Lambda_M}(\mathrm{i}a|\vec{\omega})}{\det_R S_3^{\Lambda_M}(\mathrm{i}a+\mathrm{i}\hat{m}+\frac{1}{2}\bar{\omega}|\vec{\omega})},\tag{3.4}$$

where the numerator and denominator are, respectively, the vector multiplet and hypermultiplet contributions. Moreover, we denoted $\sum_{i=1}^{m} \omega_i \equiv \bar{\omega}$, and \hat{m} , the hypermultiplet mass, enters as a shift or a. The generalized triple sine function is defined as follows

$$S_3^{\Lambda_M}(x|\vec{\omega}) = \prod_{\vec{n}\in\Lambda_M} (\vec{n}\cdot\vec{\omega}+x) \prod_{\vec{n}\in\Lambda_M^\circ} (\vec{n}\cdot\vec{\omega}-x).$$
(3.5)

⁷We expect the equivalence to hold in general, e.g. for a hypermultiplet in the fundamental representation of G.

⁸Clearly, for $\boldsymbol{p} = (1, \dots, 1)$ this space is a smooth toric Sasakian manifold M.

⁹Earlier results for the partition function on $Y^{p,q}$ and $L^{a,b,c}$ [16, 17, 20] only considered the trivial flux sector.

Here, $\vec{n} = (n_1, \ldots, n_m)$ is the vector of charges of the modes under the T^m -action of \mathbb{C}^m in (2.2) and $\vec{\omega} = (\omega_1, \ldots, \omega_m)$ is the squashing of M. Finally, the product over \vec{n} takes values in a lattice

$$\Lambda_M = \{ \vec{n} \in \mathbb{Z}_{\geq 0}^m \, | \, \vec{Q}_a^M \cdot \vec{n} = 0, \ a = 1, \dots, m - 3 \}$$
(3.6)

and in its interior Λ_M° . The m-3 conditions in the definition of Λ_M make sure that the partition function only depends on charges of the modes under the T^3 -action on M. In the literature [19, 20], the lattice often appears as a product over these three integers only.

Instead of using the det'_{adj}-notation, we write the perturbative part of the partition function explicitly as a product over the root set Δ of the gauge algebra \mathfrak{g}

$$Z_M|_{(\mathfrak{m}_1,\dots,\mathfrak{m}_{b_2})=(0,\dots,0)} \frac{\prod_{\alpha\in\Delta} S_3^{\Lambda_M}(\mathrm{i}\alpha(a)|\vec{\omega})}{\prod_{\rho\in R} S_3^{\Lambda_M}(\mathrm{i}a+\mathrm{i}\hat{m}+\frac{1}{2}\bar{\omega}|\vec{\omega})}.$$
(3.7)

Thus, we omit possible factors arising from fermionic zero-modes that would cancel a Vandermonde determinant in the integral over a. Momentarily, we will show how to derive (3.7) for $Y^{p,q}$ and $L^{a,b,c}$.

Instanton contributions are obtained by gluing Nekrasov partition functions [45, 46] at each of the m fixed fibers of \widehat{M}_p . The changes to the Nekrasov partition function in presence of conical singularities are discussed for branched covers in [43] and for orbifolds in [38]. Their dependence on the flux sector will be discussed in section 5.

3.2 Perturbative Partition Function

We now briefly recall the method employed in [16] to compute the perturbative partition function (3.7) on $Y^{p,q}$ and $L^{a,b,c}$. This method, which relies on viewing these manifolds as quotients of $S^3 \times S^3$, can be immediately generalized to compute the perturbative partition function also on the corresponding branched covers.

Manifolds. Following standard arguments [1], the perturbative partition function of certain SQFTs can be equivalently computed in terms of the equivariant index of a cohomological complex obtained via localization. On $Y^{p,q}$ such complex can be uplifted to $S^3 \times S^3$ as the extra S_Y^1 acts freely [16]. Then, once the index in 6d is computed, the index on $Y^{p,q}$ is obtained keeping only modes with trivial representation under S_Y^1 [47]. The last step requires to translate the index into a one-loop determinant. We focus on the complex associated to a vector multiplet; the hypermultiplet case can be treated analogously.

The complex on $S^3 \times S^3$ splits into individual complexes of the two S^3 factors [16]. The Cartan of the isometry group on $S^3 \times S^3$ is $T^4 = U(1)_A^2 \times U(1)_B$ where the two factors act on each S^3 separately. We denote s_1, s_2 and t_1, t_2 the coordinates on the two copies of $U(1)^2$. The relevant index of each individual complex is given by

$$\operatorname{index}_{U(1)_{A}^{2}}[\overline{\partial}] = \sum_{n_{1}, n_{2} \ge 0} s_{1}^{-n_{1}} s_{2}^{-n_{2}} - \sum_{n_{1}, n_{2} < 0} s_{1}^{-n_{1}} s_{2}^{-n_{2}},$$

$$\operatorname{index}_{U(1)_{B}^{2}}[\overline{\partial}] = \sum_{n_{3}, n_{4} \ge 0} t_{1}^{-n_{3}} t_{2}^{-n_{4}} - \sum_{n_{3}, n_{4} < 0} t_{1}^{-n_{3}} t_{2}^{-n_{4}}.$$
(3.8)

The index on $S^3 \times S^3$ then follows

$$\begin{aligned} \operatorname{index}_{T^{4}}[\partial] &= \operatorname{index}_{U(1)_{A}^{2}}[\partial] \cdot \operatorname{index}_{U(1)_{B}^{2}}[\partial] \\ &= \sum_{\substack{n_{1}, n_{2} \geq 0 \\ n_{3}, n_{4} \geq 0}} s_{1}^{-n_{1}} s_{2}^{-n_{2}} t_{1}^{-n_{3}} t_{2}^{-n_{4}} - \sum_{\substack{n_{1}, n_{2} \geq 0 \\ n_{3}, n_{4} < 0}} s_{1}^{-n_{1}} s_{2}^{-n_{2}} t_{1}^{-n_{3}} t_{2}^{-n_{4}} \\ &- \sum_{\substack{n_{1}, n_{2} < 0 \\ n_{3}, n_{4} \geq 0}} s_{1}^{-n_{1}} s_{2}^{-n_{2}} t_{1}^{-n_{3}} t_{2}^{-n_{4}} + \sum_{\substack{n_{1}, n_{2} < 0 \\ n_{3}, n_{4} < 0}} s_{1}^{-n_{1}} s_{2}^{-n_{2}} t_{1}^{-n_{3}} t_{2}^{-n_{4}}. \end{aligned}$$
(3.9)

The charge for rotation along S_Y^1 is (2.4)

$$u = (p+q)n_1 + (p-q)n_2 - pn_3 - pn_4.$$
(3.10)

Thus, keeping only the trivial representation and neglecting non-trivial KK modes, is equivalent to impose

$$u = 0 \Rightarrow n_4 = \frac{p+q}{p}n_1 + \frac{p-q}{p}n_2 - n_3.$$
 (3.11)

As p > q > 0, the only terms in (3.9) which satisfy this condition are those with either all $n_i \ge 0$ or with all $n_i < 0$.

Hence, the index on $Y^{p,q}$ is given by the quadruples \vec{n} in (3.9) satisfying u = 0. This defines a lattice

$$\Lambda_Y = \{ \vec{n} \in \mathbb{Z}_{\geq 0}^4 \, | \, u = 0 \}, \tag{3.12}$$

and equivalently for negative n_i . Adding the character for the adjoint representation for the vector multiplet, and translating into a one-loop determinant, we find¹⁰

$$Z_{Y^{p,q}}^{\text{vm}}|_{\mathfrak{m}_{1}=0} = \prod_{\alpha \in \Delta} \prod_{\vec{n} \in \Lambda_{Y}} \left(\vec{n} \cdot \vec{\omega} + i\alpha(a) \right) \prod_{\alpha \in \Delta} \prod_{\vec{n} \in \Lambda_{Y}^{\circ}} \left(\vec{n} \cdot \vec{\omega} - i\alpha(a) \right).$$
(3.13)

This expression reproduces the perturbative partition function in (3.7) with the lattice in (3.6) given by the condition

$$u = \vec{Q}^Y \cdot \vec{n} = 0 \tag{3.14}$$

in (3.12). Along the same lines, one can also derive the perturbative partition function for the hypermultiplet [16]. The main difference is an overall minus sign in (3.9) which leads to a term at the denominator in (3.7).

We can use the condition u = 0 (3.11) to substitute for n_4 in (3.13). We find

$$Z_{Y^{p,q}}^{\mathrm{vm}}|_{\mathfrak{m}_{1}=0} = \prod_{\alpha \in \Delta} \prod_{\vec{n}_{3\mathrm{d}} \in \mathcal{C}_{Y}} \left(\vec{n}_{3\mathrm{d}} \cdot \vec{\omega}_{3\mathrm{d}} + \mathrm{i}\alpha(a) \right) \prod_{\alpha \in \Delta} \prod_{\vec{n}_{3\mathrm{d}} \in \mathcal{C}_{Y}^{\circ}} \left(\vec{n}_{3\mathrm{d}} \cdot \vec{\omega}_{3\mathrm{d}} - \mathrm{i}\alpha(a) \right).$$
(3.15)

This rewriting depends on charges $\vec{n}_{3d} = (n_1, n_2, n_3)$ under the T^3 -action on $Y^{p,q}$. Moreover, we defined $\vec{\omega}_{3d} = (\tilde{\omega}_1, \tilde{\omega}_2, \tilde{\omega}_3)$ with

$$\tilde{\omega}_1 = \omega_1 + \frac{p+q}{p}\omega_4, \quad \tilde{\omega}_2 = \omega_2 + \frac{p-q}{p}\omega_4, \quad \tilde{\omega}_3 = \omega_3 - \omega_4.$$
(3.16)

¹⁰In the second factor, we flipped the sign of the contributions in (3.9) with $n_i < 0$ while also flipping the sign of $i\alpha(a)$.

The lattice in (3.15) is now given by

$$\mathcal{C}_Y = \{ \vec{n}_{3d} \in \mathbb{Z}_{\geq 0}^3 \mid \exists n_4 \in \mathbb{Z}_{\geq 0} : \frac{p+q}{p} n_1 + \frac{p-q}{p} n_2 - n_3 = n_4 \},$$
(3.17)

which is the dual moment map cone of $Y^{p,q}$. As mentioned earlier (cf. footnote 4), C_Y is related by an $SL(3,\mathbb{Z})$ transformation to the lattice appearing in [20].

As observed in [16], we can generalize to $L^{a,b,c} = (S^3 \times S^3)/S_L^1$ this approach to compute the perturbative partition function. The corresponding charge for rotations along the free S_L^1 (2.4) is given by

$$\tilde{u} = an_1 + bn_2 - cn_3 - (a+b-c)n_4, \tag{3.18}$$

and the index of the relevant complex from localization is obtained from (3.9) imposing $\tilde{u} = 0$. As a, b, c > 0 and a + b > c, also in this case only the first and and last terms in (3.9) contribute. The one-loop determinant then is a product over a lattice

$$\Lambda_{L^{a,b,c}} = \{ \vec{n} \in \mathbb{Z}_{\geq 0}^4 \, | \, \tilde{u} = 0 \}.$$
(3.19)

Branched Covers. On five-dimensional branched covers of toric Sasakian manifolds, the perturbative partition function has been computed on \hat{S}_p^5 [41–43]. The result is found to be equivalent to that of a smooth S^5 , with squashing parameters $\omega_i = p_i^{-1}$. We will find that on branched covers of $Y^{p,q}$ and $L^{a,b,c}$ the dependence is similar.

According to the same argument as in the smooth case [16], on $\widehat{Y}_{p}^{p,q}$ and $\widehat{L}_{p}^{a,b,c}$ the cohomological complex obtained via localization can be uplifted to $\widehat{S}_{p_{A}}^{3} \times \widehat{S}_{p_{B}}^{3}$, where recall that $p_{A} = (p_{1}, p_{2})$ and $p_{B} = (p_{3}, p_{4})$. These branched covers of $S^{3} \times S^{3}$ have been introduced in section 2. Hence, as in (3.8)-(3.9), the equivariant index can be computed taking the product of the individual indices on $\widehat{S}_{p_{A}}^{3}$ and $\widehat{S}_{p_{B}}^{3}$. On the two factors the index is given by that computed on a smooth S^{3} with squashing parameters given by [48]

$$\omega_A = (p_1^{-1}, p_2^{-1}), \qquad \omega_B = (p_3^{-1}, p_4^{-1}).$$
 (3.20)

Thus, we find

$$\operatorname{index}_{U(1)_{A}^{2}}[\overline{\partial}] = \sum_{n_{1},n_{2}\geq0} s_{1}^{-\frac{n_{1}}{p_{1}}} s_{2}^{-\frac{n_{2}}{p_{2}}} - \sum_{n_{1},n_{2}<0} s_{1}^{-\frac{n_{1}}{p_{1}}} s_{2}^{-\frac{n_{2}}{p_{2}}},$$

$$\operatorname{index}_{U(1)_{B}^{2}}[\overline{\partial}] = \sum_{n_{3},n_{4}\geq0} t_{1}^{-\frac{n_{3}}{p_{3}}} t_{2}^{-\frac{n_{4}}{p_{4}}} - \sum_{n_{3},n_{4}<0} t_{1}^{-\frac{n_{3}}{p_{3}}} t_{2}^{-\frac{n_{4}}{p_{4}}},$$

(3.21)

and their product gives the index on $\widehat{S}^3_{p_A} \times \widehat{S}^3_{p_B}$

$$\begin{aligned} \operatorname{index}_{T^4}[\overline{\partial}] &= \operatorname{index}_{U(1)_A^2}[\overline{\partial}] \cdot \operatorname{index}_{U(1)_B^2}[\overline{\partial}] \\ &= \sum_{\substack{n_1, n_2 \ge 0 \\ n_3, n_4 \ge 0}} s_1^{-\frac{n_1}{p_1}} s_2^{-\frac{n_2}{p_2}} t_1^{-\frac{n_3}{p_3}} t_2^{-\frac{n_4}{p_4}} - \sum_{\substack{n_1, n_2 \ge 0 \\ n_3, n_4 < 0}} s_1^{-\frac{n_1}{p_1}} s_2^{-\frac{n_2}{p_2}} t_1^{-\frac{n_3}{p_3}} t_2^{-\frac{n_4}{p_4}} \\ &- \sum_{\substack{n_1, n_2 < 0 \\ n_3, n_4 \ge 0}} s_1^{-\frac{n_1}{p_1}} s_2^{-\frac{n_2}{p_2}} t_1^{-\frac{n_3}{p_3}} t_2^{-\frac{n_4}{p_4}} + \sum_{\substack{n_1, n_2 < 0 \\ n_3, n_4 < 0}} s_1^{-\frac{n_1}{p_1}} s_2^{-\frac{n_2}{p_2}} t_1^{-\frac{n_3}{p_3}} t_2^{-\frac{n_4}{p_4}}. \end{aligned}$$
(3.22)

Focusing on $\hat{Y}_{p}^{p,q}$, we consider the globally free $S_{\hat{Y}}^{1}$ -action on $\hat{S}_{p_{A}}^{3} \times \hat{S}_{p_{B}}^{3}$ (2.15) whose corresponding charges are

$$\frac{u_{\boldsymbol{p}}}{|\boldsymbol{p}|} = n_1 \frac{p+q}{p_1} + n_2 \frac{p-q}{p_2} - n_3 \frac{p}{p_3} - n_4 \frac{p}{p_4}.$$
(3.23)

The index on $\widehat{Y}_{p}^{p,q} = S_{p_A}^3 \times S_{p_B}^3 / S_{\widehat{Y}}^1$ is obtained from (3.22) restricting to contributions with $u_p = 0$.

Translating into a one-loop determinant, we find that the perturbative partition function is given by

$$Z_{\widehat{Y}_{p}^{p,q}}^{\text{vm}}|_{\mathfrak{m}_{1}=0} = \prod_{\alpha \in \Delta} \prod_{\vec{n} \in \Lambda_{\widehat{Y}}} \left(\vec{n} \cdot \vec{\omega}_{p} + i\alpha(a) \right) \prod_{\vec{n} \in \Lambda_{\widehat{Y}}^{\circ}} \left(\vec{n} \cdot \vec{\omega}_{p} - i\alpha(a) \right), \tag{3.24}$$

where

$$\vec{\omega}_{\boldsymbol{p}} = \left(\frac{\omega_1}{p_1}, \dots, \frac{\omega_4}{p_4}\right),\tag{3.25}$$

and

$$\Lambda_{\widehat{Y}} = \{ \vec{n} \in \mathbb{Z}_{\geq 0}^4 \, | \, u_p = 0 \}.$$
(3.26)

As for the branched cover of S^5 [41–43], the branching indices p_i enter as squashing parameters. However, on $\hat{Y}_p^{p,q}$, also the lattice $\Lambda_{\hat{Y}}$ depends on them.

The perturbative partition function for the hypermultiplet is obtained keeping track of the overall minus sign in (3.22), the dependence on the mass \hat{m} in (3.7) and on the constant shift by $\sum_{i=1}^{4} \frac{\omega_i}{p_i} \equiv \bar{\omega}_p$. We find

$$Z_{\widehat{Y}_{p}^{p,q}}^{\operatorname{hm}}|_{\mathfrak{m}=0} = \prod_{\rho \in R} \prod_{\vec{n} \in \Lambda_{\widehat{Y}}} \left(\vec{n} \cdot \vec{\omega}_{p} + \mathrm{i}\hat{m} + \frac{1}{2}\bar{\omega}_{p} + \mathrm{i}\rho(a) \right)^{-1} \prod_{\vec{n} \in \Lambda_{\widehat{Y}}^{\circ}} \left(\vec{n} \cdot \vec{\omega}_{p} + \mathrm{i}\hat{m} + \frac{1}{2}\bar{\omega}_{p} - \mathrm{i}\rho(a) \right)^{-1},$$

$$(3.27)$$

where the infinite products, as expected, appear at the denominator.

As in (3.15), we substitute for n_4 and find

$$Z_{\widehat{Y}_{p}^{p,q}}^{\mathrm{vm}}|_{\mathfrak{m}_{1}=0} = \prod_{\alpha \in \Delta} \prod_{\vec{n}_{3\mathrm{d}} \in \mathcal{C}_{\widehat{Y}}} \left(\vec{n}_{3\mathrm{d}} \cdot \vec{\omega}_{p,3\mathrm{d}} + \mathrm{i}\alpha(a) \right) \prod_{\alpha \in \Delta} \prod_{\vec{n}_{3\mathrm{d}} \in \mathcal{C}_{\widehat{Y}}^{\circ}} \left(\vec{n}_{3\mathrm{d}} \cdot \vec{\omega}_{p,3\mathrm{d}} - \mathrm{i}\alpha(a) \right), \quad (3.28)$$

where

$$\vec{\omega}_{\boldsymbol{p},3\mathrm{d}} = \left(\frac{\tilde{\omega}_1}{p_1}, \frac{\tilde{\omega}_2}{p_2}, \frac{\tilde{\omega}_3}{p_3}\right),\tag{3.29}$$

and

$$\mathcal{C}_{\widehat{Y}} = \{ \vec{n}_{3d} \in \mathbb{Z}_{\geq 0}^3 \mid \exists n_4 \in \mathbb{Z}_{\geq 0} : \frac{p+q}{p} \frac{p_4}{p_1} n_1 + \frac{p-q}{p} \frac{p_4}{p_2} n_2 - \frac{p_4}{p_3} n_3 = n_4 \}.$$
(3.30)

The same procedure applies to $\hat{L}_{p}^{a,b,c}$, simply quotienting $S_{p_{A}}^{3} \times S_{p_{B}}^{3}$ by the $S_{\hat{L}}^{1}$ -action in (2.15) with charges

$$\frac{\tilde{u}_{\boldsymbol{p}}}{|\boldsymbol{p}|} = n_1 \frac{a}{p_1} + n_2 \frac{b}{p_2} - n_3 \frac{c}{p_3} - n_4 \frac{a+b-c}{p_4}.$$
(3.31)

The perturbative partition function is a product over the following lattice

$$\Lambda_{\widehat{L}} = \{ \vec{n} \in \mathbb{Z}_{\geq 0}^4 \, | \, \tilde{u}_p = 0 \}. \tag{3.32}$$

4 Fluxes Via Dimensional Reduction

In this section we present the main results of our work. First, focusing on the trivial instanton sector in (3.1), we compute the one-loop determinant at all flux sectors on $\hat{Y}_{p}^{p,q}$ and $\hat{L}_{p}^{a,b,c}$. Along the lines of [11, 12, 36], we obtain these contributions keeping non-trivial representations under the free $S_{\hat{Y}}^1$ - and $S_{\hat{L}}^1$ -action when dimensionally reducing from $\hat{S}_{p_A}^3 \times \hat{S}_{p_B}^3$. In the smooth case, our results improve those for the trivial flux sector derived in [16] and presented in the previous section.

In the second part, we further dimensionally reduce the $\mathcal{N} = 1$ theory to an $\mathcal{N} = 2$ theory on $\hat{B}_{p} = \hat{Y}_{p}^{p,q}/S^{1}$. This will enable us to compute the dependence, of the one-loop determinant, on fluxes through both independent two-cycles of \hat{B}_{p} . For manifolds, this improves the result for the one-loop determinant computed in [12] for fluxes through only one two-cycle.

Following the results in [38], the partition functions we compute in this section on branched covers hold, for vector multiplets and hypermultiplets in the adjoint representation, one the corresponding spaces with orbifold singularities discussed in section 2.

4.1 Dimensional Reduction

The method we use to compute one-loop determinants at each flux sector relies on a dimensional reduction along a globally free non-trivial S^1 -fiber

$$S^1 \hookrightarrow X^{d+1}_{\boldsymbol{p}} \to X^d_{\boldsymbol{p}},\tag{4.1}$$

where X_{p}^{d+1} and X_{p}^{d} can contain conical singularities associated to either surplus or deficit angles. The first example of such dimensional reduction appeared in [21], where the oneloop determinant at all flux sectors of an $\mathcal{N} = (2, 2)$ theory on \mathbb{CP}^{1} has been reproduced from the partition function of an $\mathcal{N} = 2$ theory on lens spaces L(h, -1) [49], taking the large *h* limit. More recently, this method has been thoroughly analysed in [11, 12], where it has been applied to the dimensional reduction of gauge theories on toric Sasakian manifolds down to quasi-toric four-manifolds. Moreover, it has been applied to compute flux contributions on $\widehat{\mathbb{CP}}_{p}^{2}$ dimensionally reducing from \widehat{S}_{p}^{5} [36].

Let us briefly summarize the mechanism through which fluxes arise via dimensional reduction; a detailed exposition can be found in [12]. The starting point is the perturbative partition function of an SQFT on X_p^{d+1} , which depends on the charges \vec{n} of the modes under the torus action on X_p^{d+1} (see e.g. (3.15)). We can rewrite such object in terms of the charge for rotations along the free S^1 -fiber, $t = f(\vec{n})$, where $f(\cdot)$ is a linear function with integer coefficients.

Rather than shirking the radius of the S^1 -fiber, we introduce a quotient space X_p^{d+1}/\mathbb{Z}_h with $\pi_1(X_p^{d+1}/\mathbb{Z}_h) = \mathbb{Z}_h$. To each element in π_1 there exists a non-trivial line bundle and an associated flat gauge connection over which one needs to sum in the partition function. At each topological sector, the SQFT contains only modes satisfying the projection condition

$$t = \alpha(\mathfrak{m}) \mod h, \qquad \mathfrak{m} = \operatorname{diag}(n_1, \dots, m_{\operatorname{rk} G}), \quad 0 \le n_i < h - 1, \tag{4.2}$$

where \mathfrak{m} are the winding numbers of the flat gauge connection. Therefore, the one-loop determinant at each topological sector is found by imposing (4.2) on the perturbative partition function on X_p^{d+1} .

At large h, we obtain the space $X_{p}^{d} = X_{p}^{d+1}/S^{1}$, where $H_{2}(X_{p}^{d}, \mathbb{Z}) \simeq H_{2}(X_{p}^{d+1}, \mathbb{Z}) \times \mathbb{Z}$. The sum over flat connections in the partition function gives rise to a sum over fluxes \mathfrak{m} labelled by the first Chern class of the line bundle. The one-loop determinant in each flux sector is obtained restricting to

$$t = \alpha(\mathfrak{m}) \tag{4.3}$$

in the perturbative partition function on X_p^{d+1} . Thus, unlike standard Kaluza-Klein dimensional reduction, when reducing along a non-trivial S^1 -fiber all KK modes are present in the *d*-dimensional theory, and distribute into the new topological sectors on the base space.

The dimensional reduction in section 3, which we employed to compute the perturbative partition function on $\widehat{Y}_{p}^{p,q}$, is along a non-trivial S^1 -fiber. There, we restricted the equivariant index on $\widehat{S}_{p_A}^3 \times \widehat{S}_{p_B}^3$ to the trivial charge under the $S_{\widehat{Y}}^1$ -action (3.23). Therefore, keeping also non-trivial charges, we will find the one-loop determinant at all flux sectors on the base space. We simply have to restrict the equivariant index in (3.22) to

$$\begin{aligned}
\widehat{Y}^{p,q}_{\boldsymbol{p}} : & u_{\boldsymbol{p}} = \alpha(\mathfrak{m}_{1}), \\
\widehat{L}^{a,b,c}_{\boldsymbol{p}} : & \widetilde{u}_{\boldsymbol{p}} = \alpha(\mathfrak{m}_{1}).
\end{aligned}$$
(4.4)

For what concerns the classical contribution at a given flux sector, we will assume that the expression at the trivial flux sector (3.3) continues to hold. The same is shown to be true dimensionally reducing from 5d to 4d [11, 12]. In those cases, a flux dependence of the classical contributions can be achieved via a shift of the Coulomb branch parameter.

Furthermore, starting now from the one-loop determinant at all flux sectors on $\widehat{Y}_{p}^{p,q}$, we can compute the one-loop determinant of an $\mathcal{N} = 2$ theory on \widehat{B}_{p} . This result will contain contributions from two fluxes as $H_2(\widehat{B}_p, \mathbb{Z}) \simeq \mathbb{Z}^2$, and it is obtained simply by restricting the equivariant index (3.22) to contributions with

$$u_{\boldsymbol{p}} = \alpha(\mathfrak{m}_1), \qquad t_{\boldsymbol{p}} = \alpha(\mathfrak{m}_2).$$
 (4.5)

On these 4d spaces, the partition function is given by

$$\mathcal{Z}_{\widehat{B}_{p}} = \sum_{\mathfrak{m}_{1},\mathfrak{m}_{2}} \int_{\mathfrak{h}} \mathrm{d}a \, e^{-S_{\mathrm{cl}}} \cdot Z_{\widehat{B}_{p}} \cdot Z_{\widehat{B}_{p}}^{\mathrm{inst}}.$$
(4.6)

As mentioned above, the classical action is not affect by the flux sector. Thus, we find¹¹

$$S_{\rm cl} = -\frac{4\pi^2}{g_{\rm YM,4d}^2(x)} \varrho_{\rm 4d} \,{\rm tr}(a^2), \tag{4.7}$$

¹¹See [12] for how to implement the limit of large h at the level of the classical contribution.

where $\rho|_{4d}$ is a function of $\vec{\omega}$ and the coupling constant depends on the position¹² $x \in \hat{B}_p$ [50].

Let us briefly remark that the two choices of S^1 -action in (2.12) lead to two inequivalent theories in 4d. The former is an equivariant deformation of Witten's $\mathcal{N} = 2$ topological twisting [22]. The latter, which we denote exotic [10], is the generalization of Pestun's theory [1] to four-dimensional spaces, possibly with conical singularities. One of the main differences between the two theories is that the field strength localizes to instantons at all fixed points for a topologically twisted theory. Instead, for exotic theories, the field strength localizes to anti-instantons at one or more fixed points. An extensive presentation of these two theories, and how they arise as two instances of a unique framework, can be found in [10, 23].

4.2 6d to 5d

We start our journey through dimensional reductions in 6d. Here, we will show how to use the equivariant index on $\widehat{S}^3_{p_A} \times \widehat{S}^3_{p_B}$ to compute the one-loop determinant, at all flux sectors, on $\widehat{Y}^{p,q}_p$ and $\widehat{L}^{a,b,c}_p$. We will first deal with the smooth case and later consider singular spaces.

Manifolds. The quotients of $S^3 \times S^3$ by the free S^1 -actions (2.4), lead to $Y^{p,q}$ and $L^{a,b,c}$. Focusing, first, on $Y^{p,q}$, we need to restrict the equivariant index (3.9) to charges \vec{n} satisfying

$$u = (p+q)n_1 + (p-q)n_2 - pn_3 - pn_4 = \alpha(\mathfrak{m}_1).$$
(4.8)

For $\alpha(\mathfrak{m}_1) = 0$ we argued, in the previous section, that only contributions with all $n_i \ge 0$ or all $n_i < 0$ contribute. For generic $\alpha(\mathfrak{m}) \ne 0$, two new phenomena occur.

- Terms in (3.9) with $n_1, n_2 \ge 0$ and $n_3, n_4 < 0$ (or $n_1, n_2 < 0$ and $n_3, n_4 \ge 0$) contribute. These come with a minus sign in the index and will appear at the denominator for a vector multiplet. Instead, they will be at the numerator for a hypermultiplet.
- There is no symmetry $\vec{n} \to -\vec{n}$ among the terms in (3.9) contributing. To reinstate such symmetry, we would have to restrict the product over positive roots.

Following this discussion, we define the following lattices

$$\Lambda_{Y}^{++}(\mathfrak{m}_{1}) = \{ \vec{n} \in \mathbb{Z}_{\geq 0}^{4} \mid u = \alpha(\mathfrak{m}_{1}) \},
\Lambda_{Y}^{+-}(\mathfrak{m}_{1}) = \{ (n_{1}, n_{2}) \in \mathbb{Z}_{\geq 0}^{2}, (n_{3}, n_{4}) \in \mathbb{Z}_{<0}^{2} \mid u = \alpha(\mathfrak{m}_{1}) \},
\Lambda_{Y}^{-+}(\mathfrak{m}_{1}) = \{ (n_{1}, n_{2}) \in \mathbb{Z}_{<0}^{2}, (n_{3}, n_{4}) \in \mathbb{Z}_{\geq 0}^{2} \mid u = \alpha(\mathfrak{m}_{1}) \},
\Lambda_{Y}^{--}(\mathfrak{m}_{1}) = \{ \vec{n} \in \mathbb{Z}_{<0}^{4} \mid u = \alpha(\mathfrak{m}_{1}) \}.$$
(4.9)

Correspondingly, for a vector multiplet the one-loop determinant at a flux sector is

$$Z_{Y^{p,q}}^{\text{vm}} = \prod_{\alpha \in \Delta} \frac{\prod_{\vec{n} \in \Lambda_Y^{++}(\mathfrak{m}_1)} (\vec{n} \cdot \vec{\omega} + i\alpha(a)) \prod_{\vec{n} \in \Lambda_Y^{--}(\mathfrak{m}_1)} (\vec{n} \cdot \vec{\omega} + i\alpha(a))}{\prod_{\vec{n} \in \Lambda_Y^{+-}(\mathfrak{m}_1)} (\vec{n} \cdot \vec{\omega} + i\alpha(a)) \prod_{\vec{n} \in \Lambda_Y^{-+}(\mathfrak{m}_1)} (\vec{n} \cdot \vec{\omega} + i\alpha(a))}.$$
(4.10)

¹²This is due to the fiber employed in dimensional reduction not being the Reeb vector whenever the toric Sasakian space is not regular.

For a hypermultiplet we find

$$Z_{Y^{p,q}}^{\operatorname{hm}} = \prod_{\rho \in R} \frac{\prod_{\vec{n} \in \Lambda_Y^{+-}(\mathfrak{m}_1)} \left(\vec{n} \cdot \vec{\omega} + \mathrm{i}\hat{m} + \frac{1}{2}\bar{\omega} + \mathrm{i}\rho(a) \right) \prod_{\vec{n} \in \Lambda_Y^{-+}(\mathfrak{m}_1)} \left(\vec{n} \cdot \vec{\omega} + \mathrm{i}\hat{m} + \frac{1}{2}\bar{\omega} + \mathrm{i}\rho(a) \right)}{\prod_{\vec{n} \in \Lambda_Y^{-+}(\mathfrak{m}_1)} \left(\vec{n} \cdot \vec{\omega} + \mathrm{i}\hat{m} + \frac{1}{2}\bar{\omega} + \mathrm{i}\rho(a) \right) \prod_{\vec{n} \in \Lambda_Y^{--}(\mathfrak{m}_1)} \left(\vec{n} \cdot \vec{\omega} + \mathrm{i}\hat{m} + \frac{1}{2}\bar{\omega} + \mathrm{i}\rho(a) \right)}.$$

$$(4.11)$$

These expressions generalize the results of [16, 17] computed at trivial flux sector.

As for the trivial flux (3.15), we can substitute for

$$n_4 = \frac{p+q}{p}n_1 + \frac{p-q}{p}n_2 - n_3 - \frac{\alpha(\mathfrak{m}_1)}{p}, \qquad (4.12)$$

which leads to

$$Z_{Y^{p,q}}^{\text{vm}} = \prod_{\alpha \in \Delta} \frac{\prod_{\vec{n}_{3d} \in \mathcal{C}_{Y}^{++}(\mathfrak{m}_{1})} \left(\vec{n}_{3d} \cdot \vec{\omega}_{3d} + i\alpha(a) - \frac{\omega_{4}}{p} \alpha(\mathfrak{m}_{1}) \right)}{\prod_{\vec{n}_{3d} \in \mathcal{C}_{Y}^{+-}(\mathfrak{m}_{1})} \left(\vec{n}_{3d} \cdot \vec{\omega}_{3d} + i\alpha(a) - \frac{\omega_{4}}{p} \alpha(\mathfrak{m}_{1}) \right)}$$

$$\frac{\prod_{\vec{n}_{3d} \in \mathcal{C}_{Y}^{--}(\mathfrak{m}_{1})} \left(\vec{n}_{3d} \cdot \vec{\omega}_{3d} + i\alpha(a) - \frac{\omega_{4}}{p} \alpha(\mathfrak{m}_{1}) \right)}{\prod_{\vec{n}_{3d} \in \mathcal{C}_{Y}^{-+}(\mathfrak{m}_{1})} \left(\vec{n}_{3d} \cdot \vec{\omega}_{3d} + i\alpha(a) - \frac{\omega_{4}}{p} \alpha(\mathfrak{m}_{1}) \right)}.$$

$$(4.13)$$

The product is over the lattices

$$\mathcal{C}_{Y}^{++}(\mathfrak{m}_{1}) = \{ \vec{n}_{3d} \in \mathbb{Z}_{\geq 0}^{3} \mid \exists n_{4} \in \mathbb{Z}_{\geq 0} : u = \alpha(\mathfrak{m}_{1}) \}, \\
\mathcal{C}_{Y}^{+-}(\mathfrak{m}_{1}) = \{ (n_{1}, n_{2}) \in \mathbb{Z}_{\geq 0}^{2}, n_{3} \in \mathbb{Z}_{< 0} \mid \exists n_{4} \in \mathbb{Z}_{< 0} : u = \alpha(\mathfrak{m}_{1}) \}, \\
\mathcal{C}_{Y}^{-+}(\mathfrak{m}_{1}) = \{ (n_{1}, n_{2}) \in \mathbb{Z}_{< 0}^{2}, n_{3} \in \mathbb{Z}_{\geq 0} \mid \exists n_{4} \in \mathbb{Z}_{\geq 0} : u = \alpha(\mathfrak{m}_{1}) \}, \\
\mathcal{C}_{Y}^{--}(\mathfrak{m}_{1}) = \{ \vec{n}_{3d} \in \mathbb{Z}_{< 0}^{3} \mid \exists n_{4} \in \mathbb{Z}_{< 0} : u = \alpha(\mathfrak{m}_{1}) \}.$$
(4.14)

As usual, one can derive the same expression also for hypermultiplets.

Similar results on $L^{a,b,c}$ are obtained straightforwardly simply employing the corresponding S_L^1 -action in (2.4).

Branched Covers. To compute the one-loop determinant at all flux sectors on $\widehat{Y}_{p}^{p,q}$ we simply have to employ the charges for rotations along the free $S_{\widehat{Y}}^{1}$ (2.15)

$$\frac{u_{\boldsymbol{p}}}{|\boldsymbol{p}|} = n_1 \frac{p+q}{p_1} + n_2 \frac{p-q}{p_2} - n_3 \frac{p}{p_3} - n_4 \frac{p}{p_4}.$$
(4.15)

Then, in analogy with (4.9), we define the following lattices

$$\Lambda_{\widehat{Y}}^{++}(\mathfrak{m}_{1}) = \{ \vec{n} \in \mathbb{Z}_{\geq 0}^{4} \mid \frac{u_{p}}{|p|} = \alpha(\mathfrak{m}_{1}) \},
\Lambda_{\widehat{Y}}^{+-}(\mathfrak{m}_{1}) = \{ (n_{1}, n_{2}) \in \mathbb{Z}_{\geq 0}^{2}, (n_{3}, n_{4}) \in \mathbb{Z}_{<0}^{2} \mid \frac{u_{p}}{|p|} = \alpha(\mathfrak{m}_{1}) \},
\Lambda_{\widehat{Y}}^{-+}(\mathfrak{m}_{1}) = \{ (n_{1}, n_{2}) \in \mathbb{Z}_{<0}^{2}, (n_{3}, n_{4}) \in \mathbb{Z}_{\geq 0}^{2} \mid \frac{u_{p}}{|p|} = \alpha(\mathfrak{m}_{1}) \},
\Lambda_{\widehat{Y}}^{--}(\mathfrak{m}_{1}) = \{ \vec{n} \in \mathbb{Z}_{<0}^{4} \mid \frac{u_{p}}{|p|} = \alpha(\mathfrak{m}_{1}) \},$$
(4.16)

which contribute to the one-loop determinant

$$Z_{\widehat{Y}_{p}^{p,q}}^{\mathrm{vm}} = \prod_{\alpha \in \Delta} \frac{\prod_{\vec{n} \in \Lambda_{\widehat{Y}}^{++}(\mathfrak{m}_{1})} (\vec{n} \cdot \vec{\omega} + \mathrm{i}\alpha(a)) \prod_{\vec{n} \in \Lambda_{\widehat{Y}}^{--}(\mathfrak{m}_{1})} (\vec{n} \cdot \vec{\omega} + \mathrm{i}\alpha(a))}{\prod_{\vec{n} \in \Lambda_{\widehat{Y}}^{+-}(\mathfrak{m}_{1})} (\vec{n} \cdot \vec{\omega} + \mathrm{i}\alpha(a)) \prod_{\vec{n} \in \Lambda_{\widehat{Y}}^{-+}(\mathfrak{m}_{1})} (\vec{n} \cdot \vec{\omega} + \mathrm{i}\alpha(a))}.$$
(4.17)

As in (3.28) we can substitute for n_4 and find

$$Z_{Y^{p,q}}^{\text{vm}} = \prod_{\alpha \in \Delta} \frac{\prod_{\vec{n}_{3d} \in \mathcal{C}_{\hat{Y}}^{++}(\mathfrak{m}_{1})} \left(\vec{n}_{3d} \cdot \vec{\omega}_{\boldsymbol{p},3d} + i\alpha(a) - \frac{\omega_{4}}{p} \frac{\alpha(\mathfrak{m}_{1})}{|\boldsymbol{p}|} \right)}{\prod_{\vec{n}_{3d} \in \mathcal{C}_{\hat{Y}}^{+-}(\mathfrak{m}_{1})} \left(\vec{n}_{3d} \cdot \vec{\omega}_{\boldsymbol{p},3d} + i\alpha(a) - \frac{\omega_{4}}{p} \frac{\alpha(\mathfrak{m}_{1})}{|\boldsymbol{p}|} \right)} \frac{\prod_{\vec{n}_{3d} \in \mathcal{C}_{\hat{Y}}^{--}(\mathfrak{m}_{1})} \left(\vec{n}_{3d} \cdot \vec{\omega}_{\boldsymbol{p},3d} + i\alpha(a) - \frac{\omega_{4}}{p} \frac{\alpha(\mathfrak{m}_{1})}{|\boldsymbol{p}|} \right)}{\prod_{\vec{n}_{3d} \in \mathcal{C}_{\hat{Y}}^{-+}(\mathfrak{m}_{1})} \left(\vec{n}_{3d} \cdot \vec{\omega}_{\boldsymbol{p},3d} + i\alpha(a) - \frac{\omega_{4}}{p} \frac{\alpha(\mathfrak{m}_{1})}{|\boldsymbol{p}|} \right)} \right)}$$

$$(4.18)$$

The lattices $C_{\widehat{Y}}^{\pm\pm}$ are given by (4.14) replacing u with u_p . Finally, expressions on $\widehat{L}_p^{a,b,c}$ and for hypermultiplets follow straightforwardly.

4.3 5d to 4d

As discussed in section 2, $L^{a,b,c}$ and the corresponding branched cover do not admit a free S^1 -action; hence we focus on $Y^{p,q}$. With respect to a choice of Killing vector in 5d there are two inequivalent choices of S^1 -action on $Y^{p,q}$ (2.12), whose corresponding charges are

$$t^{\text{top}} = n_1 + n_2,$$

 $t^{\text{ex}} = n_1 - n_2.$
(4.19)

Similarly, for $\widehat{Y}_{p}^{p,q}$ there are two choices (2.19) with charges

$$\frac{t_{\boldsymbol{p}}^{\text{top}}}{|\boldsymbol{p}_{3d}|} = \frac{n_1}{p_1} + \frac{n_2}{p_2},$$

$$\frac{t_{\boldsymbol{p}}^{\text{ex}}}{|\boldsymbol{p}_{3d}|} = \frac{n_1}{p_1} - \frac{n_2}{p_2},$$
(4.20)

where $|\mathbf{p}_{3d}| = p_1 p_2$. The two choices of t lead to, respectively, a topologically twisted and an exotic (or Pestun-like) theory.

We will only discuss the case with conical singularities. To recover the smooth setup it is enough to set all p_i to one. We take, as starting point, the expressions on $\widehat{Y}_{p}^{p,q}$ where we have substituted for n_4 : the perturbative partition function (3.28) and the oneloop determinant at a flux sector \mathfrak{m}_1 (4.18). Contributions at a given flux sector \mathfrak{m}_2 are obtained restricting to charges satisfying

$$t_{\boldsymbol{p}} = \alpha(\mathfrak{m}_2). \tag{4.21}$$

Let us start analysing the perturbative partition function in 5d (3.28). In this expression we substitute

top:
$$\frac{n_2}{p_2} = + \frac{\alpha(\mathbf{m}_2)}{|\mathbf{p}_{3d}|} - \frac{n_1}{p_1},$$

ex: $\frac{n_2}{p_2} = - \frac{\alpha(\mathbf{m}_2)}{|\mathbf{p}_{3d}|} + \frac{n_1}{p_1}.$ (4.22)

Further defining $\vec{\epsilon} = (\epsilon_1, \epsilon_2)$ with

top:
$$\epsilon_1^{\text{top}} = \frac{1}{p_1} (\tilde{\omega}_1 - \tilde{\omega}_2), \quad \epsilon_2^{\text{top}} = \frac{\tilde{\omega}_3}{p_3},$$

ex: $\epsilon_1^{\text{ex}} = \frac{1}{p_1} (\tilde{\omega}_1 + \tilde{\omega}_2), \quad \epsilon_2^{\text{ex}} = \frac{\tilde{\omega}_3}{p_3},$ (4.23)

we find for the topological theory and exotic theories

$$Z_{\widehat{B}_{p}}^{\mathrm{vm,top}}|_{\mathfrak{m}_{1}=0} = \prod_{\alpha \in \Delta} \prod_{\vec{n}_{2\mathrm{d}} \in \mathcal{B}_{\widehat{B}}(\mathfrak{m}_{2})} \left(\vec{n}_{2\mathrm{d}} \cdot \vec{\epsilon} + \mathrm{i}\alpha(a) + \tilde{\omega}_{2} \frac{\alpha(\mathfrak{m}_{2})}{|\boldsymbol{p}_{3\mathrm{d}}|} \right)$$
$$\prod_{\vec{n}_{2\mathrm{d}} \in \mathcal{B}_{\widehat{B}}^{\circ}(\mathfrak{m}_{2})} \left(\vec{n}_{2\mathrm{d}} \cdot \vec{\epsilon} - \mathrm{i}\alpha(a) + \tilde{\omega}_{2} \frac{\alpha(\mathfrak{m}_{2})}{|\boldsymbol{p}_{3\mathrm{d}}|} \right),$$
(4.24)

$$Z_{\widehat{B}_{p}}^{\mathrm{vm,ex}}|_{\mathfrak{m}_{1}=0} = \prod_{\alpha \in \Delta} \prod_{\vec{n}_{2\mathrm{d}} \in \mathcal{B}_{\widehat{B}}(\mathfrak{m}_{2})} \left(\vec{n}_{2\mathrm{d}} \cdot \vec{\epsilon} + \mathrm{i}\alpha(a) - \tilde{\omega}_{2} \frac{\alpha(\mathfrak{m}_{2})}{|\boldsymbol{p}_{3\mathrm{d}}|} \right)$$
$$\prod_{\vec{n}_{2\mathrm{d}} \in \mathcal{B}_{\widehat{B}}^{\circ}(\mathfrak{m}_{2})} \left(\vec{n}_{2\mathrm{d}} \cdot \vec{\epsilon} - \mathrm{i}\alpha(a) - \tilde{\omega}_{2} \frac{\alpha(\mathfrak{m}_{2})}{|\boldsymbol{p}_{3\mathrm{d}}|} \right).$$
(4.25)

Here, $\vec{n}_{2d} = (n_1, n_3)$ and we have defined the lattices

top:
$$\mathcal{B}_{\widehat{B}}(\mathfrak{m}_2) = \left\{ \vec{n}_{2d} \in \mathbb{Z}^2_{\geq 0} \, | \, \exists \, (n_2, n_4) \in \mathbb{Z}^2_{\geq 0} : \, u_p = 0 \text{ and } t_p^{\text{top}} = \alpha(\mathfrak{m}_2) \right\},\$$

ex: $\mathcal{B}_{\widehat{B}}(\mathfrak{m}_2) = \left\{ \vec{n}_{2d} \in \mathbb{Z}^2_{\geq 0} \, | \, \exists \, (n_2, n_4) \in \mathbb{Z}^2_{\geq 0} : \, u_p = 0 \text{ and } t_p^{\text{ex}} = \alpha(\mathfrak{m}_2) \right\}.$ (4.26)

It is possible to write (4.24)-(4.25) in terms of generalized Υ -functions [12, 23, 36, 51]. For vanishing branching indices $p_i = 1$, these expressions reproduce those found in [12], up to an $SL(2,\mathbb{Z})$ transformation.

Moving to the one-loop around fluxes on $\widehat{Y}_{p}^{p,q}$ (4.18), upon dimensional reduction we find

$$Z_{\hat{B}_{p}}^{\text{vm,top}} = \prod_{\alpha \in \Delta} \frac{\prod_{\vec{n}_{2d} \in \mathcal{B}_{\hat{B}}^{++}(\mathfrak{m}_{1},\mathfrak{m}_{2})} \left(\vec{n}_{2d} \cdot \vec{\epsilon} + i\alpha(a) + \tilde{\omega}_{2} \frac{\alpha(\mathfrak{m}_{2})}{|p_{3d}|} - \frac{\omega_{4}}{p} \frac{\alpha(\mathfrak{m}_{1})}{|p|} \right)}{\prod_{\vec{n}_{2d} \in \mathcal{B}_{\hat{B}}^{+-}(\mathfrak{m}_{1},\mathfrak{m}_{2})} \left(\vec{n}_{2d} \cdot \vec{\epsilon} + i\alpha(a) + \tilde{\omega}_{2} \frac{\alpha(\mathfrak{m}_{2})}{|p_{3d}|} - \frac{\omega_{4}}{p} \frac{\alpha(\mathfrak{m}_{1})}{|p|} \right)}{\prod_{\vec{n}_{2d} \in \mathcal{B}_{\hat{B}}^{--}(\mathfrak{m}_{1},\mathfrak{m}_{2})} \left(\vec{n}_{2d} \cdot \vec{\epsilon} + i\alpha(a) + \tilde{\omega}_{2} \frac{\alpha(\mathfrak{m}_{2})}{|p_{3d}|} - \frac{\omega_{4}}{p} \frac{\alpha(\mathfrak{m}_{1})}{|p|} \right)}{\prod_{\vec{n}_{2d} \in \mathcal{B}_{\hat{B}}^{-+}(\mathfrak{m}_{1},\mathfrak{m}_{2})} \left(\vec{n}_{2d} \cdot \vec{\epsilon} + i\alpha(a) + \tilde{\omega}_{2} \frac{\alpha(\mathfrak{m}_{2})}{|p_{3d}|} - \frac{\omega_{4}}{p} \frac{\alpha(\mathfrak{m}_{1})}{|p|} \right)},$$

$$(4.27)$$

$$Z_{\hat{B}_{p}}^{\text{vm,ex}} = \prod_{\alpha \in \Delta} \frac{\prod_{\vec{n}_{2d} \in \mathcal{B}_{\hat{B}}^{++}(\mathfrak{m}_{1},\mathfrak{m}_{2})} \left(\vec{n}_{2d} \cdot \vec{\epsilon} + i\alpha(a) - \tilde{\omega}_{2} \frac{\alpha(\mathfrak{m}_{2})}{|\mathbf{p}_{3d}|} - \frac{\omega_{4}}{p} \frac{\alpha(\mathfrak{m}_{1})}{|\mathbf{p}|} \right)}{\prod_{\vec{n}_{2d} \in \mathcal{B}_{\hat{B}}^{+-}(\mathfrak{m}_{1},\mathfrak{m}_{2})} \left(\vec{n}_{2d} \cdot \vec{\epsilon} + i\alpha(a) - \tilde{\omega}_{2} \frac{\alpha(\mathfrak{m}_{2})}{|\mathbf{p}_{3d}|} - \frac{\omega_{4}}{p} \frac{\alpha(\mathfrak{m}_{1})}{|\mathbf{p}|} \right)}{\prod_{\vec{n}_{2d} \in \mathcal{B}_{\hat{B}}^{--}(\mathfrak{m}_{1},\mathfrak{m}_{2})} \left(\vec{n}_{2d} \cdot \vec{\epsilon} + i\alpha(a) - \tilde{\omega}_{2} \frac{\alpha(\mathfrak{m}_{2})}{|\mathbf{p}_{3d}|} - \frac{\omega_{4}}{p} \frac{\alpha(\mathfrak{m}_{1})}{|\mathbf{p}|} \right)}{\prod_{\vec{n}_{2d} \in \mathcal{B}_{\hat{B}}^{-+}(\mathfrak{m}_{1},\mathfrak{m}_{2})} \left(\vec{n}_{2d} \cdot \vec{\epsilon} + i\alpha(a) - \tilde{\omega}_{2} \frac{\alpha(\mathfrak{m}_{2})}{|\mathbf{p}_{3d}|} - \frac{\omega_{4}}{p} \frac{\alpha(\mathfrak{m}_{1})}{|\mathbf{p}|} \right)},$$

$$(4.28)$$

where

$$\mathcal{B}_{\hat{B}}^{++} = \left\{ \vec{n}_{2d} \in \mathbb{Z}_{\geq 0}^{2} \mid \exists (n_{2}, n_{4}) \in \mathbb{Z}_{\geq 0}^{2} : u_{p} = \alpha(\mathfrak{m}_{1}) \text{ and } t_{p}^{\mathrm{top}} = \alpha(\mathfrak{m}_{2}) \right\},\$$

$$\mathcal{B}_{\hat{B}}^{+-} = \left\{ n_{1} \in \mathbb{Z}_{\geq 0}, n_{3} \in \mathbb{Z}_{<0} \mid \exists n_{2} \in \mathbb{Z}_{\geq 0} \text{ and } n_{4} \in \mathbb{Z}_{<0} : u_{p} = \alpha(\mathfrak{m}_{1}) \text{ and } t_{p}^{\mathrm{top}} = \alpha(\mathfrak{m}_{2}) \right\},\$$

$$\mathcal{B}_{\hat{B}}^{-+} = \left\{ n_{1} \in \mathbb{Z}_{<0}, n_{3} \in \mathbb{Z}_{\geq 0} \mid \exists n_{2} \in \mathbb{Z}_{<0} \text{ and } n_{4} \in \mathbb{Z}_{\geq 0} : u_{p} = \alpha(\mathfrak{m}_{1}) \text{ and } t_{p}^{\mathrm{top}} = \alpha(\mathfrak{m}_{2}) \right\},\$$

$$\mathcal{B}_{\hat{B}}^{--} = \left\{ \vec{n}_{2d} \in \mathbb{Z}_{<0}^{2} \mid \exists (n_{2}, n_{4}) \in \mathbb{Z}_{<0}^{2} : u_{p} = \alpha(\mathfrak{m}_{1}) \text{ and } t_{p}^{\mathrm{top}} = \alpha(\mathfrak{m}_{2}) \right\},\$$

$$(4.29)$$

and similarly for lattices associated to the exotic theory.

Expressions for the one-loop determinant of hypermultiplets at a generic flux sector $(\mathfrak{m}_1, \mathfrak{m}_2)$ follow from those presented above.

4.4 Examples

Focusing on vector multiplets, we now present two explicit examples, first a manifold $Y^{3,1}$ and then a branched cover $\hat{Y}^{2,1}_{(3,2,1,4)}$. This will enable us to draw the 2d lattices contributing to the one-loop determinant on \hat{B}_p for both topologically twisted and exotic theories. While the lattices for hypermultiplets are the same, contributions at the numerator appear at the denominator and viceversa.

Manifold. We start considering the case with $\alpha(\mathfrak{m}_1) = 0$: the lattices in (4.26) contributing to the partition functions (4.24)-(4.25) are shown, for both topological and exotic theories, in Figure 3 for various values of \mathfrak{m}_2 . These plots reproduce those appearing in [12] up to an $SL(2,\mathbb{Z})$ transformation. As expected, the lattices are compact for the topologically twisted theory and non-compact for the exotic theory. This property is due to the complexes of the two theories being, respectively, elliptic and transversally elliptic.

Moving to generic values of $(\mathfrak{m}_1, \mathfrak{m}_2)$, we plot in Figure 4 the lattices \mathcal{B}_Y^{++} and \mathcal{B}_Y^{+-} (4.29) contributing to (4.27)-(4.28). Plots for lattices \mathcal{B}_Y^{--} and \mathcal{B}_Y^{-+} are specular to those shown. Recall that contributions from regions \mathcal{B}_Y^{+-} and \mathcal{B}_Y^{-+} , where n_1 and n_3 have opposite sign, appear at the denominator in the one-loop determinant.

For the topological theory, increasing the value of $\alpha(\mathfrak{m}_1)$, keeping fixed $\alpha(\mathfrak{m}_2)$, the lattice moves towards the bottom, as it can be seen from the blue and red regions. At a certain value of $\alpha(\mathfrak{m}_1)$, contributions start appearing from the quadrant with $n_1 \geq 0$, $n_3 < 0$, as shown in the green region. Further increasing the value of $\alpha(\mathfrak{m}_1)$, one finds



Figure 3. Lattices $\mathcal{B}_Y(\mathfrak{m}_2)$ for $Y^{3,1}$. Left side: topologically twisted theory for $\alpha(\mathfrak{m}_2) = 3$ (blue) and $\alpha(\mathfrak{m}_2) = 6$ (red). Right side: exotic theory for $\alpha(\mathfrak{m}_2) = 0$ (blue), $\alpha(\mathfrak{m}_2) = -3$ (red) and $\alpha(\mathfrak{m}_2) = 3$ (green).



Figure 4. Lattices $\mathcal{B}_Y^{++}(\mathfrak{m}_1,\mathfrak{m}_2)$, contributing at the numerator, and $\mathcal{B}_Y^{+-}(\mathfrak{m}_1,\mathfrak{m}_2)$, contributing at the denominator, for $Y^{3,1}$. Left side: topologically twisted theory for $(\alpha(\mathfrak{m}_1), \alpha(\mathfrak{m}_2)) = (-3, 3)$ (red), $(\alpha(\mathfrak{m}_1), \alpha(\mathfrak{m}_2)) = (3, 3)$ (blue) and $(\alpha(\mathfrak{m}_1), \alpha(\mathfrak{m}_2)) = (9, 3)$ (green). Right side: exotic theory for $(\alpha(\mathfrak{m}_1), \alpha(\mathfrak{m}_2)) = (-6, 0)$ (red), $(\alpha(\mathfrak{m}_1), \alpha(\mathfrak{m}_2)) = (0, 0)$ (blue), $(\alpha(\mathfrak{m}_1), \alpha(\mathfrak{m}_2)) = (6, 0)$ (green).

that there are no contributions left from the quadrant $(n_1, n_3) \in \mathbb{Z}_{\geq 0}$. A similar behaviour occurs for the exotic theory.

Branched Cover. The behaviour for the branched cover is very similar. The main difference is that the parameters p affect the shape of the regions. As before we plot first the case where $\alpha(\mathfrak{m}_1) = 0$ in Figure 5 and later the case $\alpha(\mathfrak{m}_1) \neq 0$ in Figure 6.

5 Factorized Expressions and Instantons

We show how the one-loop determinant at different flux sectors, both in 5d and 4d, can be computed gluing contributions from different fixed fibers in 5d and different fixed points in 4d. This factorization property was shown to hold for the perturbative partition function in 5d [19] and in 4d [10, 52], and for the one-loop determinant around \mathfrak{m}_2 on quasi-toric four-manifolds [12] and branched covers of \mathbb{CP}^2 [36]. In this section we only treat the vector multiplet on manifolds without singularities. However, the same steps continue to hold also



Figure 5. Lattices $\mathcal{B}_{\widehat{Y}}(\mathfrak{m}_1)$ for $\widehat{Y}_{(3,2,1,4)}^{2,1}$. Left side: topologically twisted theory for $\alpha(\mathfrak{m}_2) = 6$ (blue) and $\alpha(\mathfrak{m}_2) = 12$ (red). Right side: exotic theory for $\alpha(\mathfrak{m}_2) = 0$ (blue), $\alpha(\mathfrak{m}_2) = -12$ (red) and $\alpha(\mathfrak{m}_2) = 4$ (green).



Figure 6. Lattices $\mathcal{B}_{\widehat{Y}}^{++}(\mathfrak{m}_1,\mathfrak{m}_2)$, contributing at the numerator, and $\mathcal{B}_{\widehat{Y}}^{+-}(\mathfrak{m}_1,\mathfrak{m}_2)$, contributing at the denominator, for $\widehat{Y}_{(3,2,1,4)}^{2,1}$. Left side: topologically twisted theory for $(\alpha(\mathfrak{m}_1),\alpha(\mathfrak{m}_2)) = (-240, 12)$ (blue). Right side: exotic theory $(\alpha(\mathfrak{m}_1), \alpha(\mathfrak{m}_2)) = (96, 0)$ (red).

for hypermultiplets, and on branched covers and orbifolds of $Y^{p,q}$ and B. The required modifications are along the lines as those appearing in [36] for the branched cover of \mathbb{CP}^2 .

Five Dimensions. The factorized expression for the one-loop determinant on $Y^{p,q}$ (4.13) depends on a choice of imaginary part of $\tilde{\omega}_i$. The following rewriting holds

$$Z_{Y^{p,q}}^{\text{vm}} = \prod_{\alpha \in \Delta} \prod_{i=1}^{4} \prod_{t} \Upsilon_{i}(i\alpha(a) + \beta_{1i}^{-1}\alpha(\mathfrak{m}_{1}) + \beta_{2i}^{-1}t|\epsilon_{1}^{i}, \epsilon_{2}^{i})^{s_{i}},$$
(5.1)

where we defined the Υ -function [12, 23, 51]

$$\Upsilon_i(z|\epsilon_1,\epsilon_2) = \prod_{(j,k)\in\mathcal{D}_i} (\epsilon_1 j + \epsilon_2 k + z) \prod_{(j,k)\in\mathcal{D}'_i} (\epsilon_1 j + \epsilon_2 k + \bar{z}).$$
(5.2)

The product over four contributions in (5.1) is from neighbourhoods $\mathbb{C}^2_{\epsilon_1^i,\epsilon_2^i} \times S^1$ around each fixed fiber of $Y^{p,q}$. The local equivariant parameters for the T^3 -action are¹⁴ $\epsilon_1^i, \epsilon_2^1, \beta_{2i}^{-1}$. These, together with the shift in front of $\alpha(\mathfrak{m}_2)$ and the parameters s_i can be read in Table 1. The two choices appearing in the table are completely equivalent up to this point,

¹³Each individual contribution is affected by this choice. However, after gluing all contributions, the partition function is independent.

¹⁴The relation between the $\tilde{\omega}_i$ and the ϵ_i can be found in (4.23).

| i | 1 | 2 | 3 | 4 |
|-------------------------------|--|--|--|--|
| $\epsilon_1^{{\rm top},i}$ | ϵ_1 | ϵ_2 | $-p\epsilon_1 - 2q\epsilon_2$ | $-\epsilon_2$ |
| $\epsilon_2^{\mathrm{top},i}$ | ϵ_2 | $-\epsilon_1$ | $-\epsilon_2$ | $p\epsilon_1 + 2q\epsilon_2$ |
| β_{1i}^{-1} | $\frac{\omega_4}{p \mathbf{p} }$ | $\frac{\omega_4}{p \boldsymbol{p} }$ | $\left \frac{\omega_4}{p oldsymbol{p} } - rac{\epsilon_2}{p+q} ight $ | $rac{\omega_4}{p oldsymbol{p} }-rac{\epsilon_2}{p+q}$ |
| β_{2i}^{-1} | $\left \frac{\tilde{\omega}_2}{ \boldsymbol{p}_{ m 3d} } \right $ | $\epsilon_1 + \frac{\tilde{\omega}_2}{ \boldsymbol{p}_{ m 3d} }$ | $\epsilon_1 + \frac{1}{p}(p+q)\epsilon_2 + \frac{\tilde{\omega}_2}{ p_{3d} }$ | $\frac{1}{p}(p-q)\epsilon_2 + \frac{\tilde{\omega}_2}{ \boldsymbol{p}_{ m 3d} }$ |
| s_i | 1 | -1 | 1 | -1 |
| $\epsilon_1^{\mathrm{ex},i}$ | ϵ_1 | ϵ_2 | $-\epsilon_1 - 2\epsilon_2$ | ϵ_2 |
| $\epsilon_2^{\mathrm{ex},i}$ | $-\epsilon_2$ | $-\epsilon_1$ | $-\epsilon_2$ | $\epsilon_1 + 2\epsilon_2$ |
| β_{1i}^{-1} | $\frac{\omega_4}{p \boldsymbol{p} }$ | $rac{\omega_4}{p oldsymbol{p} }$ | $rac{\omega_4}{p m{p} }-rac{\epsilon_2}{p+q}$ | $rac{\omega_4}{p oldsymbol{p} }-rac{\epsilon_2}{p+q}$ |
| β_{2i}^{-1} | $ rac{	ilde{\omega}_2}{ m{p}_{ m 3d} } $ | $\epsilon_1 + rac{	ilde{\omega}_2}{ m{p}_{ m 3d} }$ | $\epsilon_1 + \frac{1}{p}(p+q)\epsilon_2 + \frac{\tilde{\omega}_2}{ \boldsymbol{p}_{3d} }$ | $\frac{1}{p}(p-q)\epsilon_2 + \frac{\tilde{\omega}_2}{ \boldsymbol{p}_{3\mathrm{d}} }$ |
| s_i | -1 | -1 | 1 | 1 |

Table 1. Local equivariance parameters in 5d, and in 4d for the topological (top half) and exotic theory (bottom half). In 5d the two choices lead to equivalent factorizations.

the difference will arise upon dimensional reduction to 4d. The lattices $\mathcal{D}_i, \mathcal{D}'_i$, defined in [12], and the parameters s_i , are determined by different choices of regularization at the fixed fibers. Once the local equivariant parameters are known also on $L^{a,b,c}$, the factorization takes the same form as in (5.1).

As on quasi-toric four-manifolds [12, 23, 26], also in 5d the dependence on \mathfrak{m}_1 is via a shift of the Coulomb branch parameter. This allows us to conjecture the contact instanton contributions to the full partition function on $Y^{p,q}$ (3.1) to be the following

$$Z_{Y^{p,q}}^{\text{inst}} = \prod_{i=1}^{4} Z_{\mathbb{C}^2 \times S^1}^{Nek} (\mathrm{i}a + \beta_{1i}^{-1} \mathfrak{m}_1 | \beta_{2i}^{-1}, \epsilon_1^i, \epsilon_2^i, q),$$
(5.3)

where $q = \exp(2\pi i \tau)$ is the instanton counting parameter.

Four Dimensions. Once the factorization of the one-loop determinant on $Y^{p,q}$ is determined, the one-loop contributions on B follow simply by restricting t

$$Z_B^{\rm vm} = \prod_{\alpha \in \Delta} \prod_{i=1}^4 \Upsilon_i(\mathrm{i}\alpha(a) + \beta_{1i}^{-1}\alpha(\mathfrak{m}_1) + \beta_{2i}^{-1}\alpha(\mathfrak{m}_2)|\epsilon_1^i, \epsilon_2^i)^{s_i}.$$
 (5.4)

The local equivariant parameters for the topologically twisted and exotic theories appear in Table 1. The flux dependence enters as a shift of the Coulomb branch parameter for both \mathfrak{m}_1 and \mathfrak{m}_2 . Therefore, we can write down the instanton contributions to the full partition function (4.6) on B

$$Z_{B}^{\text{inst,top}} = \prod_{i=1}^{4} Z_{\mathbb{C}^{2}}^{Nek} (ia + \beta_{1i}^{-1} \mathfrak{m}_{1} + \beta_{2i}^{-1} \mathfrak{m}_{2} | \epsilon_{1}^{i}, \epsilon_{2}^{i}, q),$$

$$Z_{B}^{\text{inst,ex}} = \prod_{i=1,4} Z_{\mathbb{C}^{2}}^{Nek} (ia + \beta_{1i}^{-1} \mathfrak{m}_{1} + \beta_{2i}^{-1} \mathfrak{m}_{2} | \epsilon_{1}^{i}, \epsilon_{2}^{i}, q) \prod_{i=2,3} Z_{\mathbb{C}^{2}}^{Nek} (ia + \beta_{1i}^{-1} \mathfrak{m}_{1} + \beta_{2i}^{-1} \mathfrak{m}_{2} | \epsilon_{1}^{i}, \epsilon_{2}^{i}, q)$$
(5.5)

Note that the distribution of (anti-)instantons is different for the two theories as the exotic theory localizes to anti-instantons at two fixed points.

6 Discussion

In this work we computed partition functions of $\mathcal{N} = 1$ theories on five-dimensional toric Sasakian manifolds $Y^{p,q}$ and $L^{a,b,c}$, including previously neglected contributions from gauge configurations with flux. We extended these results to branched covers of $Y^{p,q}$ and $L^{a,b,c}$. Following the results in [38], for vector multiplets and hypermultiplets in the adjoint representation, the partition function on these spaces reproduces that on orbifolds under the matching of deficit and surplus angles. We expect the identification to hold also for matter in the fundamental representation.

We further employed these results to compute the full partition function for an $\mathcal{N} = 2$ SQFT on a manifold *B* whose topology is that of the product of two spheres, and on the corresponding branched cover and orbifold. In particular, our results provide the partition function on a class of orbifolds whose topology is that of the product of two spindles.

We computed all these results exploiting a sequence of non-trivial S^1 -fibrations, that is $Y^{p,q} = (S^3 \times S^3)/S_Y^1$ (similar for $L^{a,b,c}$) and $B = Y^{p,q}/S^1$. Thus, we have found that the equivariant index of a certain complex on $S^3 \times S^3$ completely determines the one-loop determinant around both fluxes on B. The same holds on branched covers, starting from the equivariant index on $\hat{S}^3_{p_A} \times \hat{S}^3_{p_B}$.

Future Directions.

- It would be interesting to compute all flux contributions on a generic (branched covers of) toric Sasakian manifolds. Once such a result is know, it would be possible to dimensionally reduce to (branched covers of) quasi-toric manifolds, along the lines of [12, 36] for a single flux sector. In general, the number of flux sectors is m 3 in 5d and m 2 in 4d, where m appears in (2.2).
- Recently, the large N limit of the spindle index [34, 35] has been shown to reproduce the entropy of certain accelerating black holes [33]. A technical difficulty in their computations is that fluxes for the gauge field contain a fractional part, which needs to be carefully treated. The entropy has also been computed for AdS_6 supergravity solution with near horizon geometry given by a space whose topology is $AdS_2 \times \mathbb{CP}_{p_A}^1 \times \mathbb{CP}_{p_B}^1$ [53]. This entropy is expected to be reproduced by the partition function of an $\mathcal{N} = 1$ theory on $\mathbb{CP}_{p_A}^1 \times \mathbb{CP}_{p_B}^1 \times S^1$, at large N and taking care of the fractional parts for both fluxes $\mathfrak{m}_1, \mathfrak{m}_2$. This partition function is obtained from (4.27)-(4.28), adding the contributions of modes along the extra S^1 .
- It is known that the partition function of $\mathcal{N} = 1$ SQCD with SU(2) gauge group and four flavours on S^5 , tuning the masses to specific values, reproduces the partition function of SQED with four flavours on S^3 [54, 55]. In particular, two of the three contributions from the fixed fibers of S^5 reduce to vortex partition functions, while the third trivializes. The two remaining contributions pair up to give the partition

function on S^3 [24, 25], corresponding to a facet of the toric diagram of S^5 . It would be interesting to generalize this result to $Y^{p,q}$ where one expects that the partition function reduces, at specific values of the masses, to the partition function on one of the facets of the corresponding toric diagram, that is a lens space.

Acknowledgments We are grateful to Roman Mauch for stimulating discussions and enjoyable collaboration on related topics. LR acknowledges support from the Shuimu Tsinghua Scholar Program.

References

- V. Pestun, "Localization of gauge theory on a four-sphere and supersymmetric Wilson loops," Commun. Math. Phys. 313 (2012) 71-129, arXiv:0712.2824 [hep-th].
- [2] G. Festuccia and N. Seiberg, "Rigid Supersymmetric Theories in Curved Superspace," JHEP 06 (2011) 114, arXiv:1105.0689 [hep-th].
- [3] W. Fulton, Introduction to Toric Varieties. Princeton University Press, Oxford, 1993.
- [4] C. P. Boyer and K. Galicki, "A note on toric contact geometry," Journal of Geometry and Physics 35 no. 4, (2000) 288-298, arXiv:math/9907043.
- [5] C. P. Boyer and K. Galicki, Sasakian Geometry. Oxford University Press, Oxford, 2008.
- [6] I. R. Klebanov and E. Witten, "Superconformal field theory on three-branes at a Calabi-Yau singularity," Nucl. Phys. B 536 (1998) 199-218, arXiv:hep-th/9807080.
- [7] D. R. Morrison and M. R. Plesser, "Nonspherical horizons. 1.," Adv. Theor. Math. Phys. 3 (1999) 1-81, arXiv:hep-th/9810201.
- [8] D. Martelli and J. Sparks, "Toric geometry, Sasaki-Einstein manifolds and a new infinite class of AdS/CFT duals," *Commun. Math. Phys.* 262 (2006) 51–89, arXiv:hep-th/0411238.
- [9] G. Lockhart and C. Vafa, "Superconformal Partition Functions and Non-perturbative Topological Strings," JHEP 10 (2018) 051, arXiv:1210.5909 [hep-th].
- [10] G. Festuccia, J. Qiu, J. Winding, and M. Zabzine, "Transversally Elliptic Complex and Cohomological Field Theory," J. Geom. Phys. 156 (2020) 103786, arXiv:1904.12782 [hep-th].
- [11] J. Lundin and L. Ruggeri, "SYM on quotients of spheres and complex projective spaces," *JHEP* 03 (2022) 204, arXiv:2110.13065 [hep-th].
- [12] J. Lundin, R. Mauch, and L. Ruggeri, "From 5d Flat Connections to 4d Fluxes (the Art of Slicing the Cone)," arXiv:2305.02313 [hep-th].
- [13] J. Källén and M. Zabzine, "Twisted supersymmetric 5D Yang-Mills theory and contact geometry," JHEP 05 (2012) 125, arXiv:1202.1956 [hep-th].
- [14] Y. Imamura, "Perturbative partition function for squashed S⁵," *PTEP* 2013 no. 7, (2013) 073B01, arXiv:1210.6308 [hep-th].
- [15] H.-C. Kim, J. Kim, and S. Kim, "Instantons on the 5-sphere and M5-branes," arXiv:1211.0144 [hep-th].
- [16] J. Qiu and M. Zabzine, "5D Super Yang-Mills on Y^{p,q} Sasaki-Einstein manifolds," Commun. Math. Phys. 333 no. 2, (2015) 861-904, arXiv:1307.3149 [hep-th].

- [17] J. Qiu and M. Zabzine, "Factorization of 5D super Yang-Mills theory on Y^{p,q} spaces," Phys. Rev. D 89 no. 6, (2014) 065040, arXiv:1312.3475 [hep-th].
- [18] M. Cvetic, H. Lu, D. N. Page, and C. N. Pope, "New Einstein-Sasaki spaces in five and higher dimensions," *Phys. Rev. Lett.* **95** (2005) 071101, arXiv:hep-th/0504225.
- [19] J. Qiu, L. Tizzano, J. Winding, and M. Zabzine, "Gluing Nekrasov partition functions," Commun. Math. Phys. 337 no. 2, (2015) 785-816, arXiv:1403.2945 [hep-th].
- [20] J. Qiu and M. Zabzine, "Review of localization for 5d supersymmetric gauge theories," J. Phys. A 50 no. 44, (2017) 443014, arXiv:1608.02966 [hep-th].
- [21] F. Benini and S. Cremonesi, "Partition Functions of $\mathcal{N} = (2, 2)$ Gauge Theories on S² and Vortices," *Commun. Math. Phys.* **334** no. 3, (2015) 1483–1527, arXiv:1206.2356 [hep-th].
- [22] E. Witten, "Topological Quantum Field Theory," Commun. Math. Phys. 117 (1988) 353.
- [23] G. Festuccia, J. Qiu, J. Winding, and M. Zabzine, "Twisting with a Flip (the Art of Pestunization)," Commun. Math. Phys. 377 no. 1, (2020) 341-385, arXiv:1812.06473 [hep-th].
- [24] S. Pasquetti, "Factorisation of N = 2 Theories on the Squashed 3-Sphere," JHEP 04 (2012) 120, arXiv:1111.6905 [hep-th].
- [25] C. Beem, T. Dimofte, and S. Pasquetti, "Holomorphic Blocks in Three Dimensions," JHEP 12 (2014) 177, arXiv:1211.1986 [hep-th].
- [26] N. A. Nekrasov, "Localizing gauge theories," in 14th International Congress on Mathematical Physics, pp. 645–654. 7, 2003.
- [27] M. Bershtein, G. Bonelli, M. Ronzani, and A. Tanzini, "Exact results for $\mathcal{N} = 2$ supersymmetric gauge theories on compact toric manifolds and equivariant Donaldson invariants," *JHEP* **07** (2016) 023, arXiv:1509.00267 [hep-th].
- [28] M. Bershtein, G. Bonelli, M. Ronzani, and A. Tanzini, "Gauge theories on compact toric surfaces, conformal field theories and equivariant Donaldson invariants," J. Geom. Phys. 118 (2017) 40-50, arXiv:1606.07148 [hep-th].
- [29] G. Bonelli, F. Fucito, J. F. Morales, M. Ronzani, E. Sysoeva, and A. Tanzini, "Gauge theories on compact toric manifolds," *Lett. Math. Phys.* 111 no. 3, (2021) 77, arXiv:2007.15468 [hep-th].
- [30] P. Ferrero, J. P. Gauntlett, J. M. Pérez Ipiña, D. Martelli, and J. Sparks, "D3-Branes Wrapped on a Spindle," *Phys. Rev. Lett.* **126** no. 11, (2021) 111601, arXiv:2011.10579 [hep-th].
- [31] P. Ferrero, J. P. Gauntlett, J. M. P. Ipiña, D. Martelli, and J. Sparks, "Accelerating black holes and spinning spindles," *Phys. Rev. D* 104 no. 4, (2021) 046007, arXiv:2012.08530 [hep-th].
- [32] P. Ferrero, J. P. Gauntlett, and J. Sparks, "Supersymmetric spindles," JHEP 01 (2022) 102, arXiv:2112.01543 [hep-th].
- [33] E. Colombo, S. M. Hosseini, D. Martelli, A. Pittelli, and A. Zaffaroni, "Microstates of accelerating and supersymmetric AdS₄ black holes from the spindle index," arXiv:2404.07173 [hep-th].
- [34] M. Inglese, D. Martelli, and A. Pittelli, "The Spindle Index from Localization," arXiv:2303.14199 [hep-th].

- [35] M. Inglese, D. Martelli, and A. Pittelli, "Supersymmetry and Localization on Three-Dimensional Orbifolds," arXiv:2312.17086 [hep-th].
- [36] R. Mauch and L. Ruggeri, "Super Yang-Mills on Branched Covers and Weighted Projective Spaces," arXiv:2404.11600 [hep-th].
- [37] D. Martelli and A. Zaffaroni, "Equivariant localization and holography," arXiv:2306.03891 [hep-th].
- [38] R. Mauch and L. Ruggeri, "Codimension Two Defects and SYM on Orbifolds," to appear.
- [39] J. Sparks, "Sasaki-Einstein Manifolds," Surveys Diff. Geom. 16 (2011) 265-324, arXiv:1004.2461 [math.DG].
- [40] E. Lerman, "Contact toric manifolds," J. Symplectic Geom. 1 no. 4, (2001) 785, arXiv:math/0107201.
- [41] L. F. Alday, P. Richmond, and J. Sparks, "The holographic supersymmetric Renyi entropy in five dimensions," *JHEP* 02 (2015) 102, arXiv:1410.0899 [hep-th].
- [42] N. Hama, T. Nishioka, and T. Ugajin, "Supersymmetric Rényi entropy in five dimensions," JHEP 12 (2014) 048, arXiv:1410.2206 [hep-th].
- [43] T. Nishioka and I. Yaakov, "Supersymmetric Rényi entropy and defect operators," JHEP 11 (2017) 071, arXiv:1612.02894 [hep-th].
- [44] G. Festuccia, A. Gorantis, A. Pittelli, K. Polydorou, and L. Ruggeri, "Cohomological localization of $\mathcal{N} = 2$ gauge theories with matter," *JHEP* **09** (2020) 133, arXiv:2005.12944 [hep-th].
- [45] N. A. Nekrasov, "Seiberg-Witten prepotential from instanton counting," Adv. Theor. Math. Phys. 7 no. 5, (2003) 831-864, arXiv:hep-th/0206161.
- [46] N. Nekrasov and A. Okounkov, "Seiberg-Witten theory and random partitions," *Prog. Math.* 244 (2006) 525–596, arXiv:hep-th/0306238.
- [47] M. F. Atiyah, *Elliptic operators and compact groups*, vol. 401. Springer, 1974.
- [48] T. Nishioka and I. Yaakov, "Supersymmetric Renyi Entropy," JHEP 10 (2013) 155, arXiv:1306.2958 [hep-th].
- [49] L. F. Alday, M. Fluder, and J. Sparks, "The Large N limit of M2-branes on Lens spaces," JHEP 10 (2012) 057, arXiv:1204.1280 [hep-th].
- [50] G. Festuccia, J. Qiu, J. Winding, and M. Zabzine, " $\mathcal{N} = 2$ supersymmetric gauge theory on connected sums of $S^2 \times S^2$," *JHEP* **03** (2017) 026, arXiv:1611.04868 [hep-th].
- [51] N. Hama and K. Hosomichi, "Seiberg-Witten Theories on Ellipsoids," JHEP 09 (2012) 033, arXiv:1206.6359 [hep-th]. [Addendum: JHEP 10, 051 (2012)].
- [52] R. Mauch and L. Ruggeri, "Index of the transversally elliptic complex in Pestunization," J. Phys. A 55 no. 29, (2022) 295401, arXiv:2112.10658 [hep-th].
- [53] F. Faedo, A. Fontanarossa, and D. Martelli, "Branes wrapped on quadrilaterals," arXiv:2402.08724 [hep-th].
- [54] D. Gaiotto, L. Rastelli, and S. S. Razamat, "Bootstrapping the superconformal index with surface defects," JHEP 01 (2013) 022, arXiv:1207.3577 [hep-th].
- [55] F. Nieri, S. Pasquetti, F. Passerini, and A. Torrielli, "5D partition functions, q-Virasoro systems and integrable spin-chains," *JHEP* **12** (2014) 040, arXiv:1312.1294 [hep-th].