

---

# Robust Counterfactual Inference in Markov Decision Processes

---

Jessica Lally<sup>1</sup>Milad Kazemi<sup>1</sup>Nicola Paoletti<sup>1</sup><sup>1</sup>Department of Informatics, King’s College London, London, UK

## Abstract

This paper addresses a key limitation in existing counterfactual inference methods for Markov Decision Processes (MDPs). Current approaches assume a specific causal model to make counterfactuals identifiable. However, there are usually many causal models that align with the observational and interventional distributions of an MDP, each yielding different counterfactual distributions, so fixing a particular causal model limits the validity (and usefulness) of counterfactual inference. We propose a novel *non-parametric* approach that computes tight bounds on counterfactual transition probabilities across all compatible causal models. Unlike previous methods that require solving prohibitively large optimisation problems (with variables that grow exponentially in the size of the MDP), our approach provides closed-form expressions for these bounds, making computation highly efficient and scalable for non-trivial MDPs. Once such an *interval* counterfactual MDP is constructed, our method identifies robust counterfactual policies that optimise the worst-case reward w.r.t. the uncertain interval MDP probabilities. We evaluate our method on various case studies, demonstrating improved robustness over existing methods.

## 1 INTRODUCTION

Markov Decision Processes (MDPs) are a fundamental model of computation for representing sequential decision-making processes under uncertainty, including reinforcement learning (RL) problems. However, evaluating RL-learned policies can be challenging, especially in safety-critical domains like healthcare, where testing these policies directly on patients would be risky and unethical.

Counterfactual inference of MDPs enables offline policy evaluation, i.e., without “deploying” the alternative policy into the environment. Given an observed sequence of actions and outcomes (e.g., from an existing clinician’s policy), counterfactual inference estimates what the outcome would have been if different actions had been taken. Counterfactual outcomes yielding higher rewards than the observation can serve as explanations for how the observed policy could be improved.

There is a rapidly growing body of literature on counterfactual inference for MDPs [Oberst and Sontag, 2019, Lorberbom et al., 2021, Tsirtsis et al., 2021, Benz and Rodriguez, 2022, Noorbakhsh and Rodriguez, 2022, Killian et al., 2022, Zhu et al., 2020, Tsirtsis and Rodriguez, 2024]. Most of these papers assume a specific causal model of the system (i.e., the Gumbel-max structural causal model (SCM) [Oberst and Sontag, 2019]) to identify counterfactuals, but, in general, counterfactuals in MDPs are *non-identifiable*: many causal models would fit the given observational and interventional data, each yielding different counterfactual probabilities [Zhang et al., 2022]. In the MDP context, the observational distribution will be a single observed path of the MDP, and the interventional distribution is the MDP’s transition probabilities for a given state and action. As a result, any counterfactual analysis based on a presumed causal model may not be valid, which is particularly concerning in safety-critical domains.

*Partial counterfactual inference* methods derive bounds (as opposed to sharp values) on counterfactual probabilities, accounting for *all* causal models compatible with given observational and interventional data, avoiding reliance on a fixed (and possibly erroneous) model. Our work is inspired by the canonical SCM approach developed by Zhang et al. [2022], which formulates partial counterfactual inference as an optimisation problem. However, this optimisation procedure is very inefficient even for small causal graphs (with variables and constraints that are exponential in the size of the MDP), which has encouraged the development of approximate algorithms [Zhang et al., 2022, Duarte et al.,

**Contributions** In this paper, we derive and prove exact analytical bounds for MDP counterfactual probabilities, thereby solving this inefficiency problem and enabling precise counterfactual analysis of non-trivial MDP models in a fully non-parametric way. We demonstrate how Zhang et al. [2022]’s partial counterfactual inference approach can be applied to MDPs, prove this optimisation reduces to exact analytical bounds, and show how these bounds can be extended to incorporate additional assumptions to generate more realistic counterfactual probabilities. Next, we use these bounds to construct interval counterfactual MDPs, which we solve using pessimistic value iteration [Mathiesen et al., 2024] to derive robust counterfactual policies that optimise the worst-case counterfactual rewards induced by all possible causal models compatible with the given data. Finally, we evaluate our approach on three MDP benchmarks (two Grid World environments and a model of Sepsis treatment), demonstrating that our policies are more robust to uncertainty about the causal model than those from the Gumbel-max SCM.

The most similar work to this paper is by Li and Pearl [2024], who identified counterfactual probability bounds for categorical SCMs given observational and interventional distributions. We show that their bounds can be reformulated for MDPs and correspond to a special case of our analytical bounds. However, our approach offers greater flexibility, allowing for the inclusion of additional properties to ensure more realistic counterfactual transition probabilities.

## 2 BACKGROUND

In this section, we provide background on counterfactual inference and an overview of related work on counterfactual inference in MDPs and partial counterfactual inference.

**Markov Decision Processes** MDPs are a class of stochastic models for representing sequential decision-making processes. In an MDP  $\mathcal{M}$ , at each step  $t$ , an agent in state  $s_t$  performs some action  $a_t$  determined by a policy  $\pi$ , ending up in state  $s_{t+1} \sim P(\cdot | s_t, a_t)$  and receiving reward  $R(s_t, a_t)$ . Formally, an MDP is a tuple  $\mathcal{M} = (\mathcal{S}, \mathcal{A}, P, P_I, R)$  where  $\mathcal{S}$  is the discrete *state space*,  $\mathcal{A}$  is the set of *actions*,  $P : (\mathcal{S} \times \mathcal{A} \times \mathcal{S}) \rightarrow [0, 1]$  is the *transition probability* function,  $P_I : \mathcal{S} \rightarrow [0, 1]$  is the *initial state distribution*, and  $R : (\mathcal{S} \times \mathcal{A}) \rightarrow \mathbb{R}$  is the *reward function*. A (deterministic) *policy*  $\pi$  for  $\mathcal{M}$  is a function  $\pi : \mathcal{S} \rightarrow \mathcal{A}$ . A path  $\tau$  of  $\mathcal{M}$  under policy  $\pi$  is a sequence  $\tau = (s_0, a_0), \dots, (s_{T-1}, a_{T-1})$  where  $T = |\tau|$  is the path length,  $P_I(s_0) > 0$ ,  $a_t = \pi(s_t)$  for all  $t = 0, \dots, T-1$ , and  $P(s_{t+1} | s_t, a_t) > 0$  for all  $t = 0, \dots, T-2$ .

**Counterfactual Inference** Structural causal models (SCMs) [Pearl, 2009, Halpern and Pearl, 2005] provide a mathematical framework for causal inference. Formally, a SCM is a tuple  $\mathcal{C} = (\mathbf{U}, \mathbf{V}, \mathcal{F}, P(\mathbf{U}))$ , where  $\mathbf{V}$  is the set of endogenous (observed) variables,  $\mathbf{U}$  is a set of exogenous (unobserved) variables,  $P(\mathbf{U})$  is a joint distribution over the possible values of each  $U \in \mathbf{U}$ , and  $\mathcal{F}$  is a set of structural equations where each  $f_V \in \mathcal{F}$  determines the value of endogenous variable  $V$ , given a fixed realisation of  $U_V \in \mathbf{U}$  and set of direct causes (parents)  $\text{PA}_V \subseteq \mathbf{V}^1$ .

Using SCMs, we can apply an *intervention*  $X \leftarrow x$  on an endogenous variable  $X$ , and measure the effect on the value of some other endogenous variable  $Y$ . We can also perform *counterfactual inference* to estimate what causal effect applying an intervention in the past would have had on an observation. This involves inferring the value of the exogenous variables present in the observation  $\mathbf{v}$ , i.e.,  $P(\mathbf{U} | \mathbf{v})$ , then evaluating the structural equations, replacing  $P(\mathbf{U})$  with the inferred  $P(\mathbf{U} | \mathbf{v})$ , and applying the intervention. Counterfactual inference can be challenging because many SCMs may fit the observed and interventional distributions but yield different counterfactual probabilities [Zhang et al., 2022]. As a result, many approaches require additional assumptions about the causal model for real-world application.

**Counterfactual Inference in MDPs** The Gumbel-max SCM was developed by Oberst and Sontag [2019] to perform counterfactual inference in MDPs and has been widely used in existing work [Lorberbom et al., 2021, Benz and Rodriguez, 2022, Noorbakhsh and Rodriguez, 2022, Killian et al., 2022, Zhu et al., 2020, Kazemi et al., 2025, 2024] due to its desirable property of counterfactual stability (see Definition 3.3). The Gumbel-max SCM for an MDP is expressed as:

$$\begin{aligned} S_{t+1} &= f(S_t, A_t, U_t = (G_{s,t})_{s \in \mathcal{S}}) \\ &= \arg \max_{s \in \mathcal{S}} \{\log(P_{\mathcal{M}}(s | S_t, A_t)) + G_{s,t}\} \end{aligned} \quad (1)$$

where  $P_{\mathcal{M}}$  is the MDP’s transition probabilities,  $S_t, A_t$  and  $S_{t+1}$  are endogenous variables representing the state, action and next state, and the values of the exogenous variable  $U_t = (G_{s,t})_{s \in \mathcal{S}}$  are sampled from the standard Gumbel distribution<sup>2</sup>. In this model, the Gumbel values  $G_{s,t}$  represent the exogenous noise. We can perform (approximate) posterior inference of  $P((G_{s,t})_{s \in \mathcal{S}} | s_t, a_t, s_{t+1})$  through

<sup>1</sup>In general, we denote variables with capital letters and specific values of the variables in lowercase.

<sup>2</sup>This formulation relies on the Gumbel-max trick [Madison et al., 2014] by which sampling from a categorical distribution with  $n$  categories is equivalent to sampling  $n$  values  $g_0, \dots, g_n$  from the standard Gumbel distribution and evaluating  $\arg \max_j \{\log(P(Y = j)) + g_j\}$  over all the categories, where  $Y$  is the output.

rejection sampling [Oberst and Sontag, 2019] or top-down Gumbel sampling [Maddison et al., 2014].

We can define a so-called *counterfactual MDP*  $\mathcal{M}_\tau$  by solving the SCM (1) for each transition along an observed path  $\tau$  in an MDP  $\mathcal{M}$ . The counterfactual probability for each transition is defined, for  $t = 0, \dots, T - 1$ , as:

$$\begin{aligned} P_{\mathcal{M},t,\tau}(s' | s, a) &= P(s' = \arg \max_{q \in \mathcal{S}} \{\log(P_{\mathcal{M}}(q | s, a)) + G'_{q,t}\}) \\ &\approx \frac{1}{N} \sum_{j=0}^N \mathbb{1} \left( s' = \arg \max_{q \in \mathcal{S}} \left\{ \log(P_{\mathcal{M}}(q | s, a)) + G'_{q,t}^{(j)} \right\} \right) \end{aligned} \quad (2)$$

where we sample  $N$  values of  $G'_{q,t}^{(j)}$  from the true posterior distribution  $G'_{q,t}$  through rejection sampling or top-down Gumbel sampling. The indicator function  $\mathbb{1}(\mathbb{X})$  takes the value 1 if the condition  $\mathbb{X}$  is satisfied and 0 otherwise.

Tsirtsis et al. [2021] used this counterfactual MDP to derive alternative sequences of actions (i.e., counterfactual paths) that would have produced a higher reward than an observed path. Counterfactual paths can be seen as *counterfactual explanations* for how the observed policy could have been improved. However, since the Gumbel-max SCM is just one of many possible SCMs compatible with any given MDP, this counterfactual MDP may be inaccurate, leading to unreliable counterfactual explanations. To address this limitation, we apply partial counterfactual inference methods to solve this problem of counterfactual unidentifiability.

**Partial Counterfactual Inference** Partial counterfactual inference methods bound counterfactual outcomes by considering all SCMs compatible with observational and interventional data. Key works are summarised in Table 1. Notably, Zhang et al. [2022] developed the first approach capable of handling all causal graph structures. Their *canonical SCM* approach derives a linear or polynomial optimisation problem (depending on the causal graph structure) producing exact bounds. However, this optimisation is very inefficient, as the number of linear constraints grows exponentially with the number and cardinality of the endogenous variables. Research has thus shifted from exact methods to developing approximation methods. One notable exception is recent work by Li and Pearl [2024], who identified analytical/closed-form counterfactual probability bounds for systems with categorical variables. We discuss this work further in Section 4.2, as their bounds are equivalent to ours in specific cases.

### 3 PARTIAL COUNTERFACTUAL INFERENCE VIA CANONICAL SCMS

<sup>3</sup>Zhang et al. [2022] provide both an exact and approximation method in their paper.

Table 1: Related works on partial counterfactual inference

Outcomes Bounds	Binary	Categorical
	<b>Exact</b>	Balke and Pearl [1994], Kang and Tian [2006], Cai et al. [2008]
<b>Approximate</b>	Robins [1989], Manski [1990]	Zaffalon et al. [2020, 2021, 2022, 2023, 2024], Duarte et al. [2023], Zhang et al. [2022] <sup>3</sup>

Zhang et al. [2022] introduced a family of canonical SCMs that can model all possible counterfactual distributions for any causal graph. Their partial counterfactual inference approach optimises over these canonical SCMs to identify counterfactual probability bounds. In this section, we explain how their approach can be applied to MDPs, and how additional assumptions can be incorporated to ensure counterfactuals are realistic, addressing issues in existing work.

#### 3.1 OPTIMISATION PROCEDURE

MDPs can be represented as SCMs with the following structural equations:

$$S_{t+1} = f(S_t, A_t, U_t); \quad A_t = \pi(S_t); \quad S_0 = f_0(U_0) \quad (3)$$

We can convert this SCM to its equivalent canonical SCM to perform partial counterfactual inference. To this end, we need to find the c-component of the exogenous variable  $U_t$  in the MDP causal graph (see Figure 1), which is defined as follows:




Figure 1: MDP causal graph

**Definition 3.1 (c-component [Tian and Pearl, 2002])**

Given a causal graph  $\mathcal{G}$ , a subset of its endogenous variables  $\mathbf{C} \subseteq \mathbf{V}$  is a **c-component** if any two variables  $V_i, V_j \in \mathbf{C}$  are connected by a sequence of bi-directed edges  $V_i \leftarrow U_k$  and  $V_j \leftarrow U_k$ , where each  $U_k$  is an exogenous parent shared by  $V_i$  and  $V_j$ .

**Definition 3.2 (Canonical SCM [Zhang et al., 2022])** A *canonical SCM* is a tuple  $\mathcal{C} = \langle \mathbf{V}, \mathbf{U}, \mathcal{F}, P \rangle$ , where:

1. For every endogenous variable  $V \in \mathbf{V}$ , its values  $v$  are determined by a structural equation  $v \leftarrow f_V(pa_V, u_V)$  where, for any  $pa_V$  and  $u_V$ ,  $f_V(pa_V, u_V)$  is contained within a finite domain  $\Omega_V$ .

2. For every exogenous  $U \in \mathcal{U}$ , its values  $u$  are drawn from a finite domain  $\Omega_U$ , where the cardinality of  $\Omega_U$  is equal to the total number of functions that map all possible inputs  $pa_V \in \Omega_{PA_V}$  to values  $v \in V$  for every endogenous  $V$  in the  $c$ -component covering  $U^4$ , i.e.,  $|\Omega_U| = \prod_{V \in \mathcal{C}(U)} |\Omega_{PA_V} \mapsto \Omega_V|$

The  $c$ -component covering the single exogenous variable  $U_t$  in an MDP is  $\mathcal{C}(U_t) = \{S_{t+1}\}$ . Therefore, for any MDP,  $|\Omega_U|$  is equal to the total number of functions that map all possible values of  $S_t$  and  $A_t$  (i.e., all possible combinations of states  $s \in \mathcal{S}$  and actions  $a \in \mathcal{A}$ ) to all possible values of  $S_{t+1}$  ( $\mathcal{S}$ ). Thus, the cardinality of  $U_t$  in a canonical MDP SCM is  $|\Omega_{U_t}| = |\mathcal{S}|^{|\mathcal{S}| \times |\mathcal{A}|}$ . Each value  $u_t \in U_t$  indexes a unique structural equation, which deterministically maps all possible state-action pairs  $(s, a)$  to a next state  $s'$ .

Given the canonical SCM representation of an MDP and an observed transition  $s_t, a_t \rightarrow s_{t+1}$ , we can define an optimisation procedure to find the minimum and maximum counterfactual probabilities for every transition [Zhang et al., 2022]. The first step is to define a mapping between exogenous values  $u_t \in U_t$  and the structural equation they index. We define an indicator variable  $\mu \in \{0, 1\}^{\mathcal{S} \times \mathcal{A} \times |\Omega_{U_t}| \times \mathcal{S}}$  such that, for any  $s_t, a_t, u_t$  and  $s_{t+1}$ :

$$\mu_{s_t, a_t, u_t, s_{t+1}} = \begin{cases} 1 & \text{if } f(s_t, a_t, u_t) = s_{t+1} \\ 0 & \text{otherwise} \end{cases}$$

Next, we define a vector  $\theta \in \mathbb{R}^{|\Omega_{U_t}|}$ , where each  $\theta_{u_t}$  represents the probability  $P(U_t = u_t)$ . Each instantiation of  $\theta$  defines a unique SCM. Given an observed transition  $s_t, a_t \rightarrow s_{t+1}$ , we can write the counterfactual probability  $\tilde{P}_t(\tilde{s}' | \tilde{s}, \tilde{a})$  of any transition  $\tilde{s}, \tilde{a} \rightarrow \tilde{s}'$  in terms of  $\mu$  and  $\theta$ :

$$\tilde{P}_t(\tilde{s}' | \tilde{s}, \tilde{a}) = \frac{\sum_{u_t=1}^{|\Omega_{U_t}|} \mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} \cdot \mu_{s_t, a_t, u_t, s_{t+1}} \cdot \theta_{u_t}}{P(s_{t+1} | s_t, a_t)}$$

To find the counterfactual probability bounds, we search over all values of  $\theta$  consistent with the interventional data (i.e., the MDP's transition probabilities):

$$\begin{aligned} \min / \max_{\theta} \quad & \sum_{u_t=1}^{|\Omega_{U_t}|} \mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} \cdot \mu_{s_t, a_t, u_t, s_{t+1}} \cdot \theta_{u_t} \\ \text{s.t.} \quad & \sum_{u_t=1}^{|\Omega_{U_t}|} \mu_{s, a, u_t, s'} \cdot \theta_{u_t} = P(s' | s, a), \forall s, a, s' \quad (4) \\ & 0 \leq \theta_{u_t} \leq 1, \forall u_t, \quad \sum_{u_t=1}^{|\Omega_{U_t}|} \theta_{u_t} = 1 \end{aligned}$$

<sup>4</sup> $\Omega_{PA_V} \mapsto \Omega_V$  denotes the set of all possible mappings from values in  $\Omega_{PA_V}$  to values in  $\Omega_V$




Figure 2: Example MDP where Gumbel-Max produces un-intuitive CF probabilities. The observed path is  $s_0 \rightarrow s_1$ .

### 3.2 INCORPORATING EXTRA ASSUMPTIONS

While this optimisation correctly considers all SCMs consistent with the interventional and observational data, this can result in wide or trivial  $[0, 1]$  bounds. We now explain how assumptions can be added to the optimisation problem to tighten the counterfactual probability bounds, and produce more realistic counterfactual probabilities. The first assumption is *counterfactual stability* [Oberst and Sontag, 2019], which, in the context of an MDP, is defined as:

**Definition 3.3 (Counterfactual stability)** *An MDP SCM (3) satisfies counterfactual stability if, given we observed the next state  $S_{t+1} = s_{t+1}$  after observing the state-action pair  $(s_t, a_t)$ , the counterfactual outcome under a different state-action pair  $(\tilde{s}, \tilde{a})$  will not change to some  $S_{t+1} = \tilde{s}' \neq s_{t+1}$  unless  $\frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} < \frac{P(\tilde{s}' | \tilde{s}, \tilde{a})}{P(\tilde{s}' | s_t, a_t)}$ .*

However, even with counterfactual stability, the Gumbel-max SCM yields unrealistic counterfactual probabilities. In the example in Figure 2, the counterfactual probability of transition  $s = 1, a = 0 \rightarrow s' = 2$  (highlighted in Table 2) is greater than its nominal probability, even though state  $s' = 2$  was reachable from the observed state  $s_t = 0$ , but not observed (see Appendix A for further discussion). Arguably, a state that was possible but did not unfold in the factual world should not be more likely in the counterfactual world. To formalise this, we introduce *monotonicity*<sup>5</sup>:

Table 2: Counterfactual transition probabilities produced by the optimisation in (4) vs. the Gumbel-max SCM (2) vs. the optimisation in (4) with the extra assumptions (5).

$s$	$a$	$s'$	$P(s'   s, a)$	Optimisation (4)		Gumbel-Max (2)	Optimisation (4) + (5)	
				LB	UB		LB	UB
0	0	0	0.3	0.0	0.0	0.0	0.0	0.0
0	0	1	0.4	1.0	1.0	1.0	1.0	1.0
0	0	2	0.3	0.0	0.0	0.0	0.0	0.0
1	0	0	0.4	0.0	1.0	0.35	0.4	0.4
1	0	1	0.0	0.0	0.0	0.0	0.0	0.0
<b>1</b>	<b>0</b>	<b>2</b>	<b>0.6</b>	<b>0.0</b>	<b>1.0</b>	<b>0.65</b>	<b>0.6</b>	<b>0.6</b>
2	0	0	0.0	0.0	0.0	0.0	0.0	0.0
2	0	1	0.0	0.0	0.0	0.0	0.0	0.0
2	0	2	1.0	1.0	1.0	1.0	1.0	1.0

<sup>5</sup>This is different from other definitions of monotonicity, which assume an ordering on interventions and outcomes, e.g., [Vlontzos et al., 2023].

**Definition 3.4 (Monotonicity)** An MDP SCM (3) satisfies monotonicity if, upon observing the transition  $(s_t, a_t, s_{t+1})$ , then,  $\forall s \in \mathcal{S}, a \in \mathcal{A}$ : 1) the counterfactual probability  $\tilde{P}_t(s_{t+1} | s, a)$  cannot be smaller than the nominal probability  $P(s_{t+1} | s, a)$  (i.e., observing an outcome cannot make it less likely in the counterfactual world), and 2)  $\forall \tilde{s}' \neq s_{t+1}$  with  $P(\tilde{s}' | s_t, a_t) > 0$ , the counterfactual probability  $\tilde{P}_t(\tilde{s}' | s, a)$  cannot be bigger than the nominal probability  $P(\tilde{s}' | s, a)$  (i.e., not observing a possible outcome cannot make it more likely in the counterfactual world).

The counterfactual stability (CS) and monotonicity assumptions (Mon1 + Mon2) can be added as linear constraints to the optimisation problem in (4) as follows:

$$\begin{aligned}
& \text{(CS)} \quad \tilde{P}_t(s_{t+1} | \tilde{s}, \tilde{a}) = 0 \\
& \text{if } \frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} > \frac{P(\tilde{s}' | \tilde{s}, \tilde{a})}{P(\tilde{s}' | s_t, a_t)} \text{ and } P(\tilde{s}' | s_t, a_t) > 0 \\
& \text{(Mon1)} \quad \tilde{P}_t(s_{t+1} | \tilde{s}, \tilde{a}) \geq P(s_{t+1} | \tilde{s}, \tilde{a}) \\
& \hspace{15em} \text{if } P(s_{t+1} | \tilde{s}, \tilde{a}) > 0 \\
& \text{(Mon2)} \quad \tilde{P}_t(\tilde{s}' | \tilde{s}, \tilde{a}) \leq P(\tilde{s}' | \tilde{s}, \tilde{a}) \\
& \text{if } P(\tilde{s}' | \tilde{s}, \tilde{a}) > 0 \text{ and } P(\tilde{s}' | s_t, a_t) > 0, \forall \tilde{s}' \neq s_{t+1} \quad (5)
\end{aligned}$$

## 4 DERIVING ANALYTICAL BOUNDS

While the optimisation in (4) and (5) yields exact bounds, it is very inefficient as the number of constraints grows exponentially with the sizes of the state and action spaces. However, for MDPs, we have proven this linear optimisation problem always reduces to exact closed-form solutions.

### 4.1 ANALYTICAL BOUNDS

Given the observed transition  $s_t, a_t \rightarrow s_{t+1}$ , the counterfactual probability of any transition  $\tilde{s}, \tilde{a} \rightarrow \tilde{s}'$  depends only on whether the state-action pair  $(\tilde{s}, \tilde{a})$  is equal to the observed state-action pair  $(s_t, a_t)$ , or if the support of its nominal transition probabilities,  $P(\cdot | \tilde{s}, \tilde{a})$ , is disjoint from or overlaps with the support of the observed state-action pair,  $P(\cdot | s_t, a_t)$ . The lower and upper bounds, denoted by  $\tilde{P}_t^{LB}$  and  $\tilde{P}_t^{UB}$ , for these cases are given in Theorems 4.1-4.3 (with proofs provided in Appendix B.1).

**Theorem 4.1** For the observed state-action pair  $(s_t, a_t)$ , the linear program will produce the following bounds:

$$\tilde{P}_t^{LB}(s_{t+1} | s_t, a_t) = \tilde{P}_t^{UB}(s_{t+1} | s_t, a_t) = 1$$

$$\forall \tilde{s}' \in \mathcal{S} \setminus \{s_{t+1}\}, \tilde{P}_t^{LB}(\tilde{s}' | s_t, a_t) = \tilde{P}_t^{UB}(\tilde{s}' | s_t, a_t) = 0$$

**Theorem 4.2** For state-action pairs  $(\tilde{s}, \tilde{a})$  which have disjoint support from the observed  $(s_t, a_t)$ , the linear program will produce,  $\forall \tilde{s}' \in \mathcal{S}$ , the following bounds:

$$\begin{aligned}
& \tilde{P}_t^{UB}(\tilde{s}' | \tilde{s}, \tilde{a}) = \\
& \begin{cases} \frac{P(\tilde{s}' | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)}, & \text{if } P(\tilde{s}' | \tilde{s}, \tilde{a}) < P(s_{t+1} | s_t, a_t) \\ 1, & \text{otherwise} \end{cases}
\end{aligned}$$

$$\begin{aligned}
& \tilde{P}_t^{LB}(\tilde{s}' | \tilde{s}, \tilde{a}) = \\
& \begin{cases} \frac{P(\tilde{s}' | \tilde{s}, \tilde{a}) - (1 - P(s_{t+1} | s_t, a_t))}{P(s_{t+1} | s_t, a_t)} & \text{if } P(\tilde{s}' | \tilde{s}, \tilde{a}) > (1 - P(s_{t+1} | s_t, a_t)) \\ 0, & \text{otherwise} \end{cases}
\end{aligned}$$

**Theorem 4.3** For state-action pairs  $(\tilde{s}, \tilde{a}) \neq (s_t, a_t)$  which have overlapping support with the observed  $(s_t, a_t)$ , the linear program will produce,  $\forall \tilde{s}' \in \mathcal{S}$ , the bounds

$$\begin{aligned}
& \tilde{P}_t^{UB}(\tilde{s}' | \tilde{s}, \tilde{a}) = \\
& \begin{cases} \frac{\min(P(s_{t+1} | s_t, a_t), P(s_{t+1} | \tilde{s}, \tilde{a}))}{P(s_{t+1} | s_t, a_t)}, & \text{if } \tilde{s}' = s_{t+1} \\ 0, & \text{if CS conditions hold (*)} \\ \min(P(\tilde{s}' | \tilde{s}, \tilde{a}), 1 - P(s_{t+1} | \tilde{s}, \tilde{a})), & \text{if } P(\tilde{s}' | s_t, a_t) > 0 \\ \min\left(1 - P(s_{t+1} | \tilde{s}, \tilde{a}), \frac{P(\tilde{s}' | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)}\right), & \text{otherwise} \end{cases}
\end{aligned}$$

$$\begin{aligned}
& \tilde{P}_t^{LB}(\tilde{s}' | \tilde{s}, \tilde{a}) = \\
& \begin{cases} \max\left(P(\tilde{s}' | \tilde{s}, \tilde{a}), 1 - \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} \tilde{P}_t^{UB}(s' | \tilde{s}, \tilde{a})\right), & \text{if } \tilde{s}' = s_{t+1} \\ 0, & \text{if CS conditions hold (*)} \\ \max\left(0, 1 - \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} \tilde{P}_t^{UB}(s' | \tilde{s}, \tilde{a})\right), & \text{otherwise} \end{cases}
\end{aligned}$$

where the counterfactual stability (CS) conditions (\*) are  $P(\tilde{s}' | s_t, a_t) > 0$  and  $\frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} > \frac{P(\tilde{s}' | \tilde{s}, \tilde{a})}{P(\tilde{s}' | s_t, a_t)}$

### 4.2 EQUIVALENCE WITH EXISTING WORK

The most similar work to ours is by Li and Pearl [2024], who derived tight counterfactual probability bounds for categorical models, given observational and interventional distributions. In contrast, our bounds are defined for one specific observation. In Appendix C, we prove our bounds are equivalent, but only when the counterfactual stability and monotonicity assumptions are removed, or equivalently, when the counterfactual state-action pair has *disjoint support* with the observed state-action pair<sup>6</sup>. This equivalence

<sup>6</sup>When the state-action pair has disjoint support, counterfactual stability and monotonicity hold vacuously.

reaffirms the correctness of our bounds and suggests we could reformulate them for an observational distribution, which would be an interesting future direction.

## 5 METHODOLOGY

With our analytical bounds, we can compute a non-stationary interval counterfactual MDP for any MDP  $\mathcal{M}$  and observed path  $\tau$ . Formally, an interval Markov decision process (IMDP) [Givan et al., 2000] is a tuple  $(\mathcal{S}, \mathcal{A}, P_{\downarrow}, \mathcal{R})$ , extending MDPs with an uncertain transition probability function  $P_{\downarrow}$  that maps each transition to a probability within its bounds  $P_{\downarrow}(s' | s, a) = [P^{LB}(s' | s, a), P^{UB}(s' | s, a)]$ . To construct an interval counterfactual MDP, we compute the counterfactual probability bounds for every transition in the MDP, given every observed transition in  $\tau$ .

### Definition 5.1 (Interval Counterfactual MDP (ICFMDP))

Given an observed path  $\tau$  and MDP  $\mathcal{M}$ , the counterfactual probability for each transition in the interval counterfactual MDP (ICFMDP)  $\tilde{\mathcal{M}}_{\tau}^{\downarrow}$  is defined, for  $t = 0, \dots, T - 1$ , as  $\tilde{P}_t^{\downarrow}(s' | \tilde{s}, \tilde{a}) = [\tilde{P}_t^{LB}(s' | \tilde{s}, \tilde{a}), \tilde{P}_t^{UB}(s' | \tilde{s}, \tilde{a})]$ .

To derive optimal counterfactual policies for this ICFMDP, we apply robust value iteration [Mathiesen et al., 2024], which pessimistically optimises the expected reward

$$V^*(s) = \max_{\pi} \min_{\tilde{P}_t \in \tilde{\mathcal{P}}_t^{\downarrow}} \mathbb{E}_{s' \sim \tilde{P}_t(\cdot | s, \pi(s))} [R(s, \pi(s)) + V^*(s')].$$

In Appendix B.2, we prove that any CFMDP entailed by the ICFMDP will be valid (i.e., satisfying (4) + (5)), so this policy will be robust across all possible causal models (i.e., the ICFMDP does not introduce spurious models).

## 6 EVALUATION

We apply our methodology to derive counterfactual policies for various MDPs, addressing three main research questions: (1) how does our policy’s performance compare to the Gumbel-max SCM approach; (2) how do the counterfactual stability and monotonicity assumptions impact the probability bounds; and (3) how fast is our approach compared with the Gumbel-max SCM method?

### 6.1 EXPERIMENTAL SETUP

To compare policy performance, we measure the average rewards of counterfactual paths induced by our policy and the Gumbel-max policy by uniformly sampling 200 counterfactual MDPs from the ICFMDP and generating 10,000 counterfactual paths over each sampled CFMDP. Since the interval CFMDP depends on the observed path, we select

4 paths of varying optimality to evaluate how the observed path impacts the performance of both policies: an optimal path, a slightly suboptimal path that could reach the optimal reward with a few changes, a catastrophic path that enters a catastrophic, terminal state with low reward, and an almost catastrophic path that was close to entering a catastrophic state. When measuring the average probability bound widths and execution time needed to generate the ICFMDPs, we averaged over 20 randomly generated observed paths <sup>7</sup>.

### 6.2 GRIDWORLD

The GridWorld MDP is a  $4 \times 4$  grid where an agent must navigate from the top-left corner to the goal state in the bottom-right corner, avoiding a dangerous terminal state in the centre. At each time step, the agent can move up, down, left, or right, but there is a small probability (controlled by hyper-parameter  $p$ ) of moving in an unintended direction. As the agent nears the goal, the reward for each state increases, culminating in a reward of +100 for reaching the goal. Entering the dangerous state results in a penalty of -100. We use two versions of GridWorld: a less stochastic version with  $p = 0.9$  (i.e., 90% chance of moving in the chosen direction) and a more stochastic version with  $p = 0.4$ .

**GridWorld ( $p = 0.9$ )** When  $p = 0.9$ , the counterfactual probability bounds are typically narrow (see Table 3 for average measurements). Consequently, as shown in Figure 3, both policies are nearly identical and perform similarly well across the optimal, slightly suboptimal, and catastrophic paths. However, for the almost catastrophic path, the interval CFMDP path is more conservative and follows the observed path more closely (as this is where the probability bounds are narrowest), which typically requires one additional step to reach the goal state than the Gumbel-max SCM policy.

**GridWorld ( $p = 0.4$ )** When  $p = 0.4$ , the GridWorld environment becomes more uncertain, increasing the risk of entering the dangerous state even if correct actions are chosen. Thus, as shown in Figure 4, the interval CFMDP policy adopts a more conservative approach, avoiding deviation from the observed policy if it cannot guarantee higher counterfactual rewards (see the slightly suboptimal and almost catastrophic paths), whereas the Gumbel-max SCM is inconsistent: it can yield higher rewards, but also much lower rewards, reflected in the wide error bars. For the catastrophic path, both policies must deviate from the observed path to achieve a higher reward and, in this case, perform similarly.

<sup>7</sup>Further training details are provided in Appendix D, and the code is provided at <https://github.com/ddv-lab/robust-cf-inference-in-MDPs>.




Figure 3: Average instant reward of CF paths induced by policies on GridWorld  $p = 0.9$ . Error bars denote the standard deviation in reward at each time step, and below each graph, we note the lowest cumulative reward across all counterfactual paths induced by each policy.

### 6.3 SEPSIS

The Sepsis MDP [Oberst and Sontag, 2019] simulates trajectories of Sepsis patients. Each state consists of four vital signs (heart rate, blood pressure, oxygen concentration, and glucose levels), categorised as low, normal, or high, and three treatments that can be toggled on/off at each time step (8 actions in total). Unlike Oberst and Sontag [2019], we scale rewards based on the number of out-of-range vital signs, between  $-1000$  (patient dies) and  $1000$  (patient discharged). Like the GridWorld  $p = 0.4$  experiment, the Sepsis MDP is highly uncertain, as many states are equally likely to lead to optimal and poor outcomes. Thus, as shown in Figure 5, both policies follow the observed optimal and almost catastrophic paths to guarantee rewards are no worse than the observation. However, improving the catastrophic path requires deviating from the observation. Here, the Gumbel-max SCM policy, on average, performs better than the interval CFMDP policy. But, since both policies have lower bounds clipped at  $-1000$ , neither policy reliably improves over the observation. In contrast, for the slightly suboptimal path, the interval CFMDP policy performs significantly better, shown by its higher lower bounds. Moreover, in these two cases, the worst-case counterfactual path generated by the interval CFMDP policy is better than that of the Gumbel-max SCM policy, indicating its greater robustness.

Table 3: Mean width of counterfactual probability bounds

Environment	Assumptions		
	CS + M	CS	None <sup>8</sup>
GridWorld ( $p = 0.9$ )	0.0817	0.0977	0.100
GridWorld ( $p = 0.4$ )	0.552	0.638	0.646
Sepsis	0.138	0.140	0.140

### 6.4 INTERVAL CFMDP BOUNDS

Table 3 presents the mean counterfactual probability bound widths (excluding transitions where the upper bound is 0) for each MDP, averaged over 20 observed paths. We compare the bounds under counterfactual stability (CS) and monotonicity (M) assumptions, CS alone, and no assumptions. This shows that the assumptions marginally reduce the bound widths, indicating the assumptions tighten the bounds without excluding too many causal models, as intended.

### 6.5 EXECUTION TIMES

Table 4 compares the average time needed to generate the interval CFMDP vs. the Gumbel-max SCM CFMDP for 20 observations. The GridWorld algorithms were run single-threaded, while the Sepsis experiments were run in parallel. Generating the interval CFMDP is significantly faster as it uses exact analytical bounds, whereas the Gumbel-max

<sup>8</sup>Equivalent to Li and Pearl [2024]’s bounds (see Section 4.2).




Figure 4: Average instant reward of CF paths induced by policies on GridWorld  $p = 0.4$ .




Figure 5: Average instant reward of CF paths induced by policies on Sepsis.



Table 4: Mean execution time to generate CFMDPs

Environment	Mean Execution Time (s)	
	Interval CFMDP	Gumbel-max CFMDP
GridWorld ( $p = 0.9$ )	0.261	56.1
GridWorld ( $p = 0.4$ )	0.336	54.5
Sepsis	688	2940

CFMDP requires sampling from the Gumbel distribution to estimate counterfactual transition probabilities. Since constructing the counterfactual MDP models is the main bottleneck in both approaches, ours is more efficient overall and suitable for larger MDPs.

## 7 CONCLUSION

This paper addresses a key limitation in existing counterfactual inference methods for MDPs, introducing the first partial counterfactual inference approach that can efficiently handle large state and action spaces in MDPs, eliminating the need for an assumed causal model. A major contribution of this work is deriving exact analytical counterfactual probability bounds, which makes our method significantly faster than the (widely-used) Gumbel-max SCM method. Moreover, counterfactual policies derived from the interval CFMDP offer greater robustness to uncertainty about the true counterfactual MDP, especially in environments with high stochasticity. As a result, our approach provides a more reliable explanation of how an observed policy could be improved, which is crucial for safety-critical applications.

**Future Work** Like most existing methods, we assume access to the MDP’s true transition probabilities and full state observability. Future work will focus on relaxing these assumptions by learning transition probabilities from data (with uncertainty bounds) and developing approaches for partially observable and continuous-state MDPs.

## Acknowledgements

The authors would like to thank Frederik Mathiesen and Luca Laurenti for their support with using the IntervalMDP.jl package. This work was supported by UK Research and Innovation [grant number EP/S023356/1] in the UKRI Centre for Doctoral Training in Safe and Trusted Artificial Intelligence ([www.safeandtrustedai.org](http://www.safeandtrustedai.org)); and UK Research and Innovation [grant number EP/W014785/2].

## References

Alexander Balke and Judea Pearl. Counterfactual probabilities: Computational methods, bounds and applications.

In *Uncertainty in artificial intelligence*, pages 46–54. Elsevier, 1994.

Nina L Corvelo Benz and Manuel Gomez Rodriguez. Counterfactual inference of second opinions. In *Uncertainty in Artificial Intelligence*, pages 453–463. PMLR, 2022.

Zhihong Cai, Manabu Kuroki, Judea Pearl, and Jin Tian. Bounds on direct effects in the presence of confounded intermediate variables. *Biometrics*, 64(3):695–701, 2008.

Guilherme Duarte, Noam Finkelstein, Dean Knox, Jonathan Mummolo, and Ilya Shpitser. An automated approach to causal inference in discrete settings. *Journal of the American Statistical Association*, pages 1–16, 2023.

Robert Givan, Sonia Leach, and Thomas Dean. Bounded-parameter Markov decision processes. *Artificial Intelligence*, 122(1):71–109, 2000. ISSN 0004-3702. doi: [https://doi.org/10.1016/S0004-3702\(00\)00047-3](https://doi.org/10.1016/S0004-3702(00)00047-3).

Joseph Y Halpern and Judea Pearl. Causes and explanations: A structural-model approach. part II: Explanations. *The British journal for the philosophy of science*, 2005.

Changsung Kang and Jin Tian. Inequality constraints in causal models with hidden variables. In *Proceedings of the Twenty-Second Conference on Uncertainty in Artificial Intelligence*, pages 233–240, 2006.

Milad Kazemi, Jessica Lally, Ekaterina Tishchenko, Hana Chockler, and Nicola Paoletti. Counterfactual influence in Markov decision processes. *arXiv preprint arXiv:2402.08514*, 2024.

Milad Kazemi, Jessica Lally, and Nicola Paoletti. Causal temporal reasoning for Markov decision processes. *Research Directions: Cyber-Physical Systems*, page 1–23, 2025. doi: 10.1017/cbp.2025.2.

Taylor W Killian, Marzyeh Ghassemi, and Shalmali Joshi. Counterfactually guided policy transfer in clinical settings. In *Conference on Health, Inference, and Learning*, pages 5–31. PMLR, 2022.

Ang Li and Judea Pearl. Probabilities of causation with nonbinary treatment and effect. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 38, pages 20465–20472, 2024.

Guy Lorberbom, Daniel D Johnson, Chris J Maddison, Daniel Tarlow, and Tamir Hazan. Learning generalized Gumbel-max causal mechanisms. *Advances in Neural Information Processing Systems*, 34:26792–26803, 2021.

- Chris J Maddison, Daniel Tarlow, and Tom Minka. A\* sampling. *Advances in Neural Information Processing Systems*, 27, 2014.
- Charles F Manski. Nonparametric bounds on treatment effects. *The American Economic Review*, 80(2):319–323, 1990.
- Frederik Baymler Mathiesen, Morteza Lahijanian, and Luca Laurenti. Intervalmdp.jl: Accelerated value iteration for interval Markov decision processes. volume 58, pages 1–6, 2024. doi: <https://doi.org/10.1016/j.ifacol.2024.07.416>. 8th IFAC Conference on Analysis and Design of Hybrid Systems ADHS 2024.
- Kimia Noorbakhsh and Manuel Rodriguez. Counterfactual temporal point processes. *Advances in Neural Information Processing Systems*, 35:24810–24823, 2022.
- Michael Oberst and David Sontag. Counterfactual off-policy evaluation with Gumbel-max structural causal models. In *ICML*, 2019.
- Judea Pearl. *Causality*. Cambridge University Press, 2<sup>nd</sup> edition, 2009. doi: [10.1017/CBO9780511803161](https://doi.org/10.1017/CBO9780511803161).
- James M Robins. The analysis of randomized and non-randomized aids treatment trials using a new approach to causal inference in longitudinal studies. *Health service research methodology: a focus on AIDS*, pages 113–159, 1989.
- Jin Tian and Judea Pearl. A general identification condition for causal effects. In *AAAI/IAAI*, pages 567–573, 2002.
- Stratis Tsirtsis and Manuel Rodriguez. Finding counterfactually optimal action sequences in continuous state spaces. *Advances in Neural Information Processing Systems*, 36, 2024.
- Stratis Tsirtsis, Abir De, and Manuel Rodriguez. Counterfactual explanations in sequential decision making under uncertainty. *Advances in Neural Information Processing Systems*, 34:30127–30139, 2021.
- Athanasios Vlontzos, Bernhard Kainz, and Ciarán M Gilligan-Lee. Estimating categorical counterfactuals via deep twin networks. *Nature Machine Intelligence*, 5(2): 159–168, 2023.
- Marco Zaffalon, Alessandro Antonucci, and Rafael Cabañas. Structural causal models are (solvable by) credal networks. In *International Conference on Probabilistic Graphical Models*, pages 581–592. PMLR, 2020.
- Marco Zaffalon, Alessandro Antonucci, and Rafael Cabañas. Causal expectation-maximisation. In *WHY-21 Workshop*, 2021.
- Marco Zaffalon, Alessandro Antonucci, Rafael Cabañas, David Huber, and Dario Azzimonti. Bounding counterfactuals under selection bias. In *International Conference on Probabilistic Graphical Models*, pages 289–300. PMLR, 2022.
- Marco Zaffalon, Alessandro Antonucci, Rafael Cabañas, and David Huber. Approximating counterfactual bounds while fusing observational, biased and randomised data sources. *International Journal of Approximate Reasoning*, 162:109023, 2023.
- Marco Zaffalon, Alessandro Antonucci, Rafael Cabañas, David Huber, and Dario Azzimonti. Efficient computation of counterfactual bounds. *International Journal of Approximate Reasoning*, page 109111, 2024.
- Junzhe Zhang, Jin Tian, and Elias Bareinboim. Partial counterfactual identification from observational and experimental data. In Kamalika Chaudhuri, Stefanie Jegelka, Le Song, Csaba Szepesvari, Gang Niu, and Sivan Sabato, editors, *Proceedings of the 39th International Conference on Machine Learning*, volume 162 of *Proceedings of Machine Learning Research*, pages 26548–26558. PMLR, 17–23 Jul 2022.
- Qingfu Zhu, Weinan Zhang, Ting Liu, and William Yang Wang. Counterfactual off-policy training for neural dialogue generation. In *Proceedings of the 2020 Conference on Empirical Methods in Natural Language Processing (EMNLP)*, pages 3438–3448, 2020.

# Robust Counterfactual Inference in Markov Decision Processes (Supplementary Material)

Jessica Lally<sup>1</sup>

Milad Kazemi<sup>1</sup>

Nicola Paoletti<sup>1</sup>

<sup>1</sup>Department of Informatics, King’s College London, London, UK

## A EXPLANATION OF UNINTUITIVE COUNTERFACTUAL PROBABILITIES PRODUCED BY GUMBEL-MAX SCMS

Take the example MDP from Figure 2, which contains three states and a single action. In this example, we observe the transition  $s_t = 0, a_t = 0 \rightarrow s_{t+1} = 1$ . Neither state 0 nor state 2 was observed, although both were possible. This means that the observed transition gives us no information as to whether state 0 or state 2 would have been more likely where, counterfactually,  $s_t = 1$ , so intuitively these counterfactual probabilities should be equal to the nominal probabilities. However, the Gumbel-max SCM scales the probabilities, making state 2 even more likely than in the nominal MDP, and vice versa for state 0. This happens because the Gumbel distributions that the Gumbel values are drawn from for each transition are truncated in proportion to the log probability of the transition’s nominal probability (instead of in proportion to their nominal probabilities).




Figure 6: Example of MDP where Gumbel-Max produces unintuitive CF probabilities. The observed path  $s_0 \rightarrow s_1$  is highlighted in grey.

Table 5: Counterfactual transition probabilities produced by the optimisation in (4) (without assumptions) vs. probabilities produced by Gumbel-max SCM (2) vs. the optimisation in (4) with the additional assumptions in (5).

$s$	$a$	$s'$	$P(s'   s, a)$	Optimisation (4)		Gumbel-Max (2)	Optimisation (4) + (5)	
				LB	UB		LB	UB
0	0	0	0.3	0.0	0.0	0.0	0.0	0.0
0	0	1	0.4	1.0	1.0	1.0	1.0	1.0
0	0	2	0.3	0.0	0.0	0.0	0.0	0.0
1	0	0	0.4	0.0	1.0	0.35	0.4	0.4
1	0	1	0.0	0.0	0.0	0.0	0.0	0.0
<b>1</b>	<b>0</b>	<b>2</b>	<b>0.6</b>	<b>0.0</b>	<b>1.0</b>	<b>0.65</b>	<b>0.6</b>	<b>0.6</b>
2	0	0	0.0	0.0	0.0	0.0	0.0	0.0
2	0	1	0.0	0.0	0.0	0.0	0.0	0.0
2	0	2	1.0	1.0	1.0	1.0	1.0	1.0

## B PROOFS

**Optimisation Problem** For a given counterfactual transition  $s, a \rightarrow s'$  and given observed transition  $s_t, a_t \rightarrow s_{t+1}$ , the optimisation problem is defined as follows:

$$\min / \max_{\theta} \sum_{u_t=1}^{|U_t|} \mu_{s,a,u_t,s'} \cdot \mu_{s_t,a_t,u_t,s_{t+1}} \cdot \theta_{u_t} \quad (6)$$

$$\text{s.t.} \sum_{u_t=1}^{|U_t|} \mu_{\tilde{s},\tilde{a},u_t,\tilde{s}'} \cdot \theta_{u_t} = P(\tilde{s}' | \tilde{s}, \tilde{a}), \forall \tilde{s}, \tilde{a}, \tilde{s}' \quad (7)$$

$$\tilde{P}_t(s_{t+1} | \tilde{s}, \tilde{a}) \geq P(s_{t+1} | \tilde{s}, \tilde{a}) \text{ if } P(s_{t+1} | \tilde{s}, \tilde{a}) > 0, \forall \tilde{s}, \tilde{a} \text{ (Mon1)} \quad (8)$$

$$\tilde{P}_t(\tilde{s}' | \tilde{s}, \tilde{a}) \leq P(\tilde{s}' | \tilde{s}, \tilde{a}) \text{ if } P(\tilde{s}' | \tilde{s}, \tilde{a}) > 0 \text{ and } P(\tilde{s}' | s_t, a_t) > 0, \forall \tilde{s}, \tilde{a}, \tilde{s}' \neq s_{t+1} \text{ (Mon2)} \quad (9)$$

$$\tilde{P}_t(\tilde{s}' | \tilde{s}, \tilde{a}) = 0 \text{ if } \frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} > \frac{P(\tilde{s}' | \tilde{s}, \tilde{a})}{P(\tilde{s}' | s_t, a_t)} \text{ and } P(\tilde{s}' | s_t, a_t) > 0, \forall \tilde{s}, \tilde{a}, \tilde{s}' \text{ (CS)} \quad (10)$$

$$0 \leq \theta_{u_t} \leq 1, \forall u_t \quad (11)$$

$$\sum_{u_t=1}^{|U_t|} \theta_{u_t} = 1 \quad (12)$$

where  $\theta$  and  $\mu$  are defined as follows:

$$\begin{aligned} \theta &\in \mathbb{R}^{|\Omega_{U_t}|} \\ \mu &\in \{0, 1\}^{\mathcal{S} \times \mathcal{A} \times |\Omega_{U_t}| \times \mathcal{S}} \\ \forall s \in \mathcal{S}, a \in \mathcal{A}, s' \in \mathcal{S}, u_t \in U_t, \mu_{s,a,u_t,s'} &= \begin{cases} 1 & \text{if } f(s, a, u_t) = s' \\ 0 & \text{otherwise} \end{cases} \end{aligned}$$

**Counterfactual Probabilities** For any transition  $s, a \rightarrow s'$  and observed transition  $s_t, a_t \rightarrow s_{t+1}$ , the counterfactual transition probability  $\tilde{P}_t(s' | s, a)$  can be calculated as follows:

$$\tilde{P}_t(s' | s, a) = \frac{\sum_{u_t=1}^{|U_t|} \mu_{s,a,u_t,s'} \cdot \mu_{s_t,a_t,u_t,s_{t+1}} \cdot \theta_{u_t}}{P(s_{t+1} | s_t, a_t)} \quad (13)$$

### B.1 COUNTERFACTUAL TRANSITION PROBABILITY BOUNDS

In this section, we prove that the above optimisation problem reduces to the exact analytical bounds for counterfactual transition probabilities given in Section 4, for any MDP  $\mathcal{M}$ .

**Theorem B.1** *For any MDP  $\mathcal{M}$  and path  $\tau$ , the optimisation problem defined in (6)-(12) reduces to the exact analytical bounds given in Theorems 4.1-4.3 for the counterfactual transition probabilities of every transition in  $\mathcal{M}$ .*

For any state-action pair  $(\tilde{s}, \tilde{a})$  in  $\mathcal{M}$ , the counterfactual probabilities for all transitions from  $(\tilde{s}, \tilde{a})$  depend only on the other transitions from  $(\tilde{s}, \tilde{a})$ , and the observed state-action pair  $(s_t, a_t)$  (as shown in the equation for calculating counterfactual probabilities, (13)). Since  $P(s_{t+1} | s_t, a_t)$  is fixed, the counterfactual probability of each transition  $\tilde{s}, \tilde{a} \rightarrow \tilde{s}'$  only depends on

$$\sum_{u_t=1}^{|U_t|} \mu_{s,a,u_t,s'} \cdot \mu_{s_t,a_t,u_t,s_{t+1}} \cdot \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t}$$

We need to prove that the constraints of the optimisation problem hold for *all* state-action pairs in the MDP. We will use induction to show that new values of the state-action pairs can be added to the causal model (while adjusting  $\theta$  to meet the constraints for the new pair) without affecting whether the constraints for previously included pairs are satisfied.

Therefore, if we can derive and prove closed-form solutions for the probability bounds for transitions from each state-action pair independently, we know that we can combine these solutions to produce a valid assignment of  $\theta$  across all pairs (e.g., by taking the upper bound of one transition with non-zero probability from all state-action pairs). Section B.1.1 contains the inductive proof, and Section B.1.2 provides the closed-form solutions.

### B.1.1 Inductive Proof

By induction, we will prove, given any MDP  $\mathcal{M}$ , that if we can assign  $\theta$  such that the constraints of the optimisation problem (7)-(12) are satisfied for all  $|\mathcal{S}| \times |\mathcal{A}|$  state-action pairs in  $\mathcal{M}$  separately, we know these can be combined to form a single valid  $\theta$  across all  $|\mathcal{S}| \times |\mathcal{A}|$  state-action pairs in  $\mathcal{M}$ .

Take an arbitrary MDP  $\mathcal{M}$  with state space  $|\mathcal{S}|$  and action space  $|\mathcal{A}|$ . In total, there are  $N = |\mathcal{S}| \times |\mathcal{A}|$  state-action pairs in the MDP, meaning there are  $|U_t| = |\mathcal{S}|^{|\mathcal{S}| \times |\mathcal{A}|}$  possible unique structural equation mechanisms. At each stage of the inductive proof, we will add a new state-action pair value from the MDP  $\mathcal{M}$  to the causal model, and adjust  $\theta$  to satisfy the constraints for this new state-action pair without affecting whether the constraints hold for all state-action pair values already considered in the causal model. When we have considered  $k$  state-action pairs, this results in  $\theta \in \mathbb{R}^{|\mathcal{S}|^k}$ . Therefore, once we have added all  $|\mathcal{S}| \times |\mathcal{A}|$  state-action pairs, we will have  $\theta \in \mathbb{R}^{|U_t|}$ , as required by the optimisation problem.

**Base Case 1** Assume  $(s_t, a_t)$  (the observed state-action pair) is the only state-action pair value we want to consider in the causal model (this always exists, as we always have an observed transition). We require a structural equation mechanism for each possible next state (of which there are  $|\mathcal{S}|$ ). This leads to an assignment of  $\theta^1 \in \mathbb{R}^{|\mathcal{S}|^1}$  as follows, which is the only possible assignment that satisfies all the constraints (7)-(12):

$$\frac{\theta_{u_1}^1}{P(s_1 | s_t, a_t)} \quad \Bigg| \quad \frac{\theta_{u_2}^1}{P(s_2 | s_t, a_t)} \quad \Bigg| \quad \dots \quad \Bigg| \quad \frac{\theta_{u_{|\mathcal{S}|}}^1}{P(s_{|\mathcal{S}|} | s_t, a_t)}$$

$\theta^1 = [P(s_1 | s_t, a_t), P(s_2 | s_t, a_t), \dots, P(s_{|\mathcal{S}|} | s_t, a_t)]$ . Given  $\theta^1$ , the counterfactual transition probability for each transition  $s_t, a_t \rightarrow s'$  can be calculated with:

$$\tilde{P}_t(s' | s_t, a_t) = \frac{\sum_{u_t=1}^{|U_t|} \mu_{s_t, a_t, u_t, s'} \cdot \mu_{s_t, a_t, u_t, s_{t+1}} \cdot \theta_{u_t}}{P(s_{t+1} | s_t, a_t)}$$

Therefore, given  $\theta^1$ ,  $\tilde{P}_t(s_{t+1} | s_t, a_t) = 1$ , and  $\forall s' \in \mathcal{S} \setminus \{s_{t+1}\}$ ,  $\tilde{P}_t(s' | s_t, a_t) = 0$ . These are the only possible counterfactual probabilities, so these probabilities are produced by the closed-form bounds in Section B.1.2. These counterfactual probabilities satisfy the monotonicity and counterfactual stability constraints of the optimisation problem, as follows:

- Because  $\frac{P(s_{t+1} | s_t, a_t)}{P(s_{t+1} | s_t, a_t)} = 1$  and  $\forall s' \in \mathcal{S}$ ,  $\frac{P(s' | s_t, a_t)}{P(s' | s_t, a_t)} = 1$ ,  

$$\forall s' \in \mathcal{S}, \frac{P(s_{t+1} | s_t, a_t)}{P(s_{t+1} | s_t, a_t)} = \frac{P(s' | s_t, a_t)}{P(s' | s_t, a_t)}$$

so CS (10) is always vacuously true for all transitions from  $(s_t, a_t)$ .


- Because  $\tilde{P}_t(s_{t+1} | s_t, a_t) = 1 \geq P(s_{t+1} | s_t, a_t)$ , this satisfies Mon1 (8).
- Because  $\forall s' \in \mathcal{S} \setminus \{s_{t+1}\}$ ,  $\tilde{P}_t(s' | s_t, a_t) = 0 \leq P(s' | s_t, a_t)$ , this satisfies Mon2 (9).

---

<sup>1</sup>The superscript of each  $\theta^k$  indicates the number of state-action pair values we have considered in the causal model. When we have all  $|\mathcal{S}| \times |\mathcal{A}|$  state-action pairs, we will have our final  $\theta = \theta^{|\mathcal{S}| \times |\mathcal{A}|}$


**Base Case 2** Let us assume that we only have the observed state-action pair in the causal model, and assume we have a valid  $\theta^1$  with  $|\mathcal{S}|$  structural equation mechanisms.

Now, take the state-action pair for which we wish to calculate the probability bounds,  $(s, a)$ , to add to the causal model. We now require  $|\mathcal{S}|^2$  structural equation mechanisms, one for each possible combination of transitions from the two state-action pairs. We can view this as separating each of the existing structural equation mechanisms from Base Case 1 into  $|\mathcal{S}|$  new structural equation mechanisms, one for every possible transition from the new state-action pair,  $(s, a)$ :



where, for  $1 \leq i \leq |\mathcal{S}|, 1 \leq n \leq |\mathcal{S}|$ ,  $u_{i,n}$  leads to the same next state for  $(s_t, a_t)$  as  $u_i$ , and produces the transition  $s, a \rightarrow s_n$ .

In the same way, we can split each  $\theta^1_{u_1}, \dots, \theta^1_{u_{|\mathcal{S}|}}$  across these new structural equation mechanisms to obtain  $\theta^2 \in \mathbb{R}^{|\mathcal{S}|^2}$ :



By splitting  $\theta^1$  in this way, we guarantee that the total probability across each set of mechanisms (where each set produces a different transition from  $(s_t, a_t)$ ) remains the same, i.e.,

$$\forall i \in \{1, \dots, |\mathcal{S}|\}, \sum_{n=1}^{|\mathcal{S}|} \theta^2_{u_{i,n}} = \theta^1_{u_i}$$

This means that the counterfactual probabilities of all transitions from  $(s_t, a_t)$  will be exactly the same when using  $\theta^2$  as with  $\theta^1$ . Because we know all the constraints were satisfied when using  $\theta^1$ , this guarantees that all the constraints of the optimisation problem will continue to hold for  $(s_t, a_t)$  with  $\theta^2$ .

Now, we only need to ensure we assign  $\theta^2$  such that all the constraints hold for the transitions from  $(s, a)$ . Firstly, we need  $\sum_{i=1}^{|\mathcal{S}|} \theta^2_{u_{i,n}} = P(s_n | s, a)$  for every possible next state  $s_n \in \mathcal{S}$ , to satisfy the interventional constraint (7). For each transition  $s, a \rightarrow s_n$ , there is a  $\theta^2_{u_{i,n}}$  for every  $u_i$ , and for every  $\theta^2_{u_{i,n}}, 0 \leq \theta^2_{u_{i,n}} \leq \theta^1_{u_i}$ . Therefore,  $0 \leq \sum_{i=1}^{|\mathcal{S}|} \theta^2_{u_{i,n}} \leq 1, \forall s_n \in \mathcal{S}$ . We also know that:

$$\begin{aligned} \sum_{n=1}^{|\mathcal{S}|} \sum_{i=1}^{|\mathcal{S}|} \theta^2_{u_{i,n}} &= \sum_{i=1}^{|\mathcal{S}|} \theta^1_{u_i} \text{ (because we split the probabilities of each } \theta^1_{u_i}\text{)} \\ &= \sum_{i=1}^{|\mathcal{S}|} P(s_i | s_t, a_t) \\ &= 1 \end{aligned}$$

and


$$\sum_{n=1}^{|\mathcal{S}|} \sum_{i=1}^{|\mathcal{S}|} \theta_{u_{i,n}}^2 = \sum_{n=1}^{|\mathcal{S}|} P(s_n | s, a) = 1$$

Therefore, no matter the assignment of  $\theta^1$ , we can satisfy the interventional probability constraints (7) for all transitions from  $(s, a)$ . In Section B.1.2, we prove valid assignments of  $\theta$  that would minimise and maximise the counterfactual probability for some transition from  $(s, a)$ , and satisfy all the constraints (7)-(12). Therefore, we can pick the assignment of  $\theta^2$  from these cases to minimise/maximise the counterfactual probability of the transition in question.

**Inductive Case** Let us assume that we now have  $k$  state-action pair values from the MDP  $\mathcal{M}$  in our causal model, and we have found a valid assignment  $\theta^k \in \mathbb{R}^{|\mathcal{S}|^k}$  that satisfies all the constraints for these  $k$  state-action pairs, including  $(s_t, a_t)$ . Because there are  $k$  state-action pairs, there are  $|\mathcal{S}|^k$  possible unique structural equation mechanisms for these state-action pairs<sup>2</sup>, with probabilities as follows:


$$\theta_{u_1}^k \quad \dots \quad \theta_{u_{|\mathcal{S}|}}^k \quad \dots \quad \theta_{u_{|\mathcal{S}|^2}}^k \quad \dots \quad \theta_{u_{|\mathcal{S}|^k}}^k$$

Now, we wish to add the  $k+1^{th}$  state-action pair from the MDP to the causal model, resulting in  $|\mathcal{S}|^{k+1}$  possible structural equation mechanisms. Let the  $k+1^{th}$  state-action pair be  $(s, a)$  arbitrarily. We can view this as separating the existing structural equation mechanisms each into  $|\mathcal{S}|$  new structural equation mechanisms, one for every possible transition from the  $k+1^{th}$  state-action pair:



where each  $u_{i,n}$  produces the same next states for all of the first  $k$  state-action pairs as  $u_i$ , and produces the transition  $s, a \rightarrow s_n$ .

In the same way, we can split each  $\theta_{u_1}^k, \dots, \theta_{u_{|\mathcal{S}|}}^k, \dots$  across these new structural equation mechanisms to find  $\theta^{k+1} \in \mathbb{R}^{|\mathcal{S}|^{k+1}}$ .



By splitting  $\theta^k$  in this way, we have  $\forall i \in \{1, \dots, |\mathcal{S}|^k\}, \sum_{n=1}^{|\mathcal{S}|} \theta_{u_{i,n}}^{k+1} = \theta_{u_i}^k$ . Each  $\theta_{u_{i,n}}^{k+1}$  is only different to  $\theta_{u_i}^k$  in the transition from  $(s, a)$ . Therefore, this guarantees that for every state-action pair  $(s', a') \neq (s, a)$  in the first  $k$  pairs added to the causal model, the total probability across each set of mechanisms (where each set produces a different transition from  $(s', a')$ ) remains the same in  $\theta^{k+1}$  as in  $\theta^k$ . As a result, the counterfactual probabilities of all transitions from each  $(s', a')$  will be exactly the same when using  $\theta^{k+1}$  as with  $\theta^k$ . Because we know all the constraints were satisfied when using  $\theta^k$ , this guarantees that all the constraints of the optimisation problem will continue to hold for  $(s', a')$  with  $\theta^{k+1}$ .

<sup>2</sup>Note we have changed the indexing of each  $\theta_{u_{i,n}}^k$  to  $\theta_j^k$ , where  $j = (i-1) \cdot |\mathcal{S}| + n$ .

Now, we only need to make sure that we can assign  $\theta^{k+1}$  such that the constraints also hold for all the transitions from  $(s, a)$ . We need  $\sum_{i=1}^{|\mathcal{S}|^k} \theta_{u_i, n}^{k+1} = P(s_n | s, a)$  for every possible next state  $s_n \in \mathcal{S}$ , to satisfy the interventional probability constraints (7). For each transition  $s, a \rightarrow s_n$ , there is a  $\theta_{u_i, n}^{k+1}$  for every  $u_i$ , and for every  $\theta_{u_i, n}^{k+1}$ ,  $0 \leq \theta_{u_i, n}^{k+1} \leq \theta_{u_n}^k$ , therefore  $0 \leq \sum_{i=1}^{|\mathcal{S}|^k} \theta_{u_i, n}^{k+1} \leq 1, \forall s_n \in \mathcal{S}$ . We also know that:

$$\begin{aligned} \sum_{n=1}^{|\mathcal{S}|} \sum_{i=1}^{|\mathcal{S}|^k} \theta_{u_i, n}^{k+1} &= \sum_{i=1}^{|\mathcal{S}|^k} \theta_{u_i}^k \text{ (because we split the probabilities of each } \theta_{u_i}^k \text{)} \\ &= 1 \end{aligned}$$

and

$$\sum_{n=1}^{|\mathcal{S}|} \sum_{i=1}^{|\mathcal{S}|^k} \theta_{u_i, n}^2 = \sum_{n=1}^{|\mathcal{S}|} P(s_n | s, a) = 1$$

Therefore, no matter the assignment of  $\theta^k$ , we can satisfy the interventional probability constraints (7) for all transitions from  $(s, a)$ . In Section B.1.2, we show that we can assign  $\theta^{k+1}$  such that it would minimise and maximise the counterfactual probability for some transition from  $(s, a)$ , and satisfy all the constraints (7)-(12). Therefore, we know we can assign  $\theta^{k+1}$  such that it satisfies the constraints for all  $k + 1$  state-action pairs in the causal model.

**Conclusion** By induction, we have proven that as long as we can satisfy the constraints for each state-action pair separately, we can combine these into a single valid  $\theta$  that satisfies the constraints for all state-action pairs, and minimises/maximises the counterfactual probability for a particular transition in question. Because the state-action pairs are added in an arbitrary order in this inductive proof (other than the observed state-action pair and the state-action pair for the transition in question, which both must always exist), this works for any MDP.



### B.1.2 Closed-form Probability Bounds

For compactness in the proofs, we adjust the notation of the optimisation problem in (6)-(12) as follows.  $\forall s, s' \in \mathcal{S}, \forall a \in \mathcal{A}, \forall u_t \in U_t$

$$\begin{aligned}
\sum_{u_t=1}^{|U_t|} \mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} \cdot \mu_{s_t, a_t, u_t, s_{t+1}} \cdot \theta_{u_t} &= \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1}}} \mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} \cdot \theta_{u_t} \text{ (by def, } \mu_{s_t, a_t, u_t, s_{t+1}} = 1 \iff f(s_t, a_t, u_t) = s_{t+1}) \\
&= \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} \text{ (by def, } \mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} = 1 \iff f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}')
\end{aligned} \tag{14}$$

Take an arbitrary transition  $\tilde{s}, \tilde{a} \rightarrow \tilde{s}'$  from the MDP. We need to consider the minimum/maximum possible value of  $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t}$  to find the lower and upper bounds, subject to the following constraints over the state-action pair,  $(\tilde{s}, \tilde{a})$ :

- $\forall \tilde{s}' \in \mathcal{S}, \sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = P(\tilde{s}' | \tilde{s}, \tilde{a})$  to satisfy (7)
- $\tilde{P}_t(s_{t+1} | \tilde{s}, \tilde{a}) \geq P(s_{t+1} | \tilde{s}, \tilde{a})$  to satisfy Mon1 (8)
- $\forall s' \in \mathcal{S} \setminus \{s_{t+1}\}, \tilde{P}_t(s' | \tilde{s}, \tilde{a}) \leq P(s' | \tilde{s}, \tilde{a})$  to satisfy Mon2 (9)
- $\forall \tilde{s}' \in \mathcal{S}, \tilde{P}_t(\tilde{s}' | \tilde{s}, \tilde{a})$  satisfies CS (10)
- $0 \leq \theta_{u_t} \leq 1, \forall u_t$  (11)
- $\sum_{u_t=1}^{|U_t|} \theta_{u_t} = 1$  (12)

From (7), we also derive the following constraints over the state-action pair  $(s_t, a_t)$ , which must hold when calculating the counterfactual probability bounds for every state-action pair.

- $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1}}} \theta_{u_t} = P(s_{t+1} | s_t, a_t)$  to satisfy (7)
- $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1}}} \theta_{u_t} = \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} P(s' | s_t, a_t) = 1 - P(s_{t+1} | s_t, a_t)$  to satisfy (7)

Consider the following distinct cases:

1.  $(\tilde{s}, \tilde{a}) = (s_t, a_t)$
2.  $(\tilde{s}, \tilde{a}) \neq (s_t, a_t)$  and  $(\tilde{s}, \tilde{a})$  has disjoint support with  $(s_t, a_t)$
3.  $(\tilde{s}, \tilde{a}) \neq (s_t, a_t), (\tilde{s}, \tilde{a})$  has overlapping support with  $(s_t, a_t)$

These cases are disjoint, and cover all possible state-action pairs.

### B.1.3 CF Bounds for Observed $(s_t, a_t)$

**Theorem B.2** For outgoing transitions from the observed state-action pair  $(s_t, a_t)$ , the linear program will produce bounds of:

$$\tilde{P}_t^{LB}(s_{t+1} \mid s_t, a_t) = \tilde{P}_t^{UB}(s_{t+1} \mid s_t, a_t) = 1$$

for the observed next state  $s_{t+1}$ , and

$$\tilde{P}_t^{LB}(\tilde{s}' \mid s_t, a_t) = \tilde{P}_t^{UB}(\tilde{s}' \mid s_t, a_t) = 0$$

for all other  $\tilde{s}' \in \mathcal{S} \setminus \{s_{t+1}\}$ .

**Proof B.1** If  $(\tilde{s}, \tilde{a}) = (s_t, a_t)$ , then it is impossible to have a  $u_t$  such that  $f(s_t, a_t, u_t) = s'$  and  $f(\tilde{s}, \tilde{a}, u_t) \neq s', \forall s' \in \mathcal{S}$ . Therefore, the only possible assignment of  $\theta$  is:

$$\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1}}} \theta_{u_t} = P(s_{t+1} \mid \tilde{s}, \tilde{a})$$

$$\forall s' \in \mathcal{S} \setminus \{s_{t+1}\}, \quad \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s'}} \theta_{u_t} = P(s' \mid \tilde{s}, \tilde{a})$$

This assignment satisfies the interventional probability constraints:

- $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1}}} \theta_{u_t} = P(s_{t+1} \mid s_t, a_t)$  satisfying (7).
- $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1}}} \theta_{u_t} = 1 - P(s_{t+1} \mid s_t, a_t)$  satisfying (7).
- $\forall s' \in \mathcal{S}, \sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = P(s' \mid \tilde{s}, \tilde{a})$  satisfying (7).
- Because of Lemma B.3,  $\tilde{P}_t(s_{t+1} \mid s_t, a_t) = 1 \geq P(s_{t+1} \mid s_t, a_t)$ , satisfying Mon1 (8), and  $\forall s' \in \mathcal{S} \setminus \{s_{t+1}\}, \tilde{P}_t(s' \mid s_t, a_t) = 0 \leq P(s' \mid s_t, a_t)$ , satisfying Mon2 (9).
- Because  $\frac{P(s_{t+1} \mid s_t, a_t)}{P(s_{t+1} \mid s_t, a_t)} = 1$  and  $\forall s' \in \mathcal{S}, \frac{P(s' \mid s_t, a_t)}{P(s' \mid s_t, a_t)} = 1$ ,

$$\forall s' \in \mathcal{S}, \frac{P(s_{t+1} \mid s_t, a_t)}{P(s_{t+1} \mid s_t, a_t)} = \frac{P(s' \mid s_t, a_t)}{P(s' \mid s_t, a_t)}$$

so CS (10) is always vacuously true for all transitions from  $(s_t, a_t)$ .

- $0 \leq \theta_{u_t} \leq 1, \forall u_t$ , satisfying (11).
- $\sum_{u_t=1}^{|U_t|} \theta_{u_t} = \sum_{s' \in \mathcal{S}} P(s' \mid s_t, a_t) = 1$ , satisfying (12).

Therefore,

$$\tilde{P}_t^{LB}(s_{t+1} \mid s_t, a_t) = \tilde{P}_t^{UB}(s_{t+1} \mid s_t, a_t) = 1$$

and  $\forall \tilde{s}' \in \mathcal{S} \setminus \{s_{t+1}\}$ :

$$\tilde{P}_t^{LB}(\tilde{s}' \mid s_t, a_t) = \tilde{P}_t^{UB}(\tilde{s}' \mid s_t, a_t) = 0$$

#### B.1.4 CF Bounds for Disjoint $(\tilde{s}, \tilde{a})$

To enhance readability, we split the proof for Theorem 4.2 into two separate theorems: Theorem B.3 proves the upper counterfactual probability bound, and Theorem B.4 proves the lower counterfactual probability bound.

**Theorem B.3** For outgoing transitions from state-action pairs  $(\tilde{s}, \tilde{a})$  which have completely disjoint support from the observed  $(s_t, a_t)$ , the linear program will produce an upper bound of:

$$\tilde{P}_t^{UB}(\tilde{s}' | \tilde{s}, \tilde{a}) = \begin{cases} \frac{P(\tilde{s}' | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} & \text{if } P(\tilde{s}' | \tilde{s}, \tilde{a}) < P(s_{t+1} | s_t, a_t) \\ 1 & \text{otherwise} \end{cases}$$

for all next states  $\tilde{s}' \in \mathcal{S}$ .

**Proof B.2** Take an arbitrary transition  $\tilde{s}, \tilde{a} \rightarrow \tilde{s}'$  where  $(\tilde{s}, \tilde{a})$  has disjoint support with the observed state-action pair  $(s_t, a_t)$ . By definition:

$$\tilde{P}_t^{UB}(\tilde{s}' | \tilde{s}, \tilde{a}) = \max_{\theta} \left( \frac{\sum_{u_t=1}^{|U_t|} \mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} \cdot \mu_{s_t, a_t, u_t, s_{t+1}} \cdot \theta_{u_t}}{P(s_{t+1} | s_t, a_t)} \right)$$

Because  $P(s_{t+1} | s_t, a_t)$  (the interventional probability of the observed transition) is fixed, to find the upper bound for  $\tilde{P}_t(\tilde{s}' | \tilde{s}, \tilde{a})$  we need to maximise  $\sum_{u_t=1}^{|U_t|} \mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} \cdot \mu_{s_t, a_t, u_t, s_{t+1}} \cdot \theta_{u_t}$ . From Lemma B.2, we have:

$$\sum_{u_t=1}^{|U_t|} \mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} \cdot \mu_{s_t, a_t, u_t, s_{t+1}} \cdot \theta_{u_t} \leq \min(P(s_{t+1} | s_t, a_t), P(\tilde{s}' | \tilde{s}, \tilde{a}))$$

Therefore, if we can show that there exists a  $\theta$  such that

$$\sum_{u_t=1}^{|U_t|} \mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} \cdot \mu_{s_t, a_t, u_t, s_{t+1}} \cdot \theta_{u_t} = \min((P(s_{t+1} | s_t, a_t), P(\tilde{s}' | \tilde{s}, \tilde{a})))$$

and  $\theta$  satisfies the constraints of the optimisation problem, we know that this results in the maximum possible counterfactual transition probability  $\tilde{P}_t(\tilde{s}' | \tilde{s}, \tilde{a})$ . Since  $\forall s, s' \in \mathcal{S}, \forall a \in \mathcal{A}, \forall u_t \in U_t, \mu_{s, a, u_t, s'} = 1 \iff f(s, a, u_t) = s'$  by definition of  $\mu$ , this is equivalent to showing that there exists a  $\theta$  such that:

$$\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = \min((P(s_{t+1} | s_t, a_t), P(\tilde{s}' | \tilde{s}, \tilde{a})))$$

As these notations are equivalent, we will use this second notation for brevity and clarity. Consider the following disjoint cases:

- $P(\tilde{s}' | \tilde{s}, \tilde{a}) < P(s_{t+1} | s_t, a_t)$
- $P(\tilde{s}' | \tilde{s}, \tilde{a}) \geq P(s_{t+1} | s_t, a_t)$

**Case 1:**  $P(\tilde{s}' | \tilde{s}, \tilde{a}) < P(s_{t+1} | s_t, a_t)$

**Proof B.3** In this case, assign  $\theta$  as follows:

$$\left\{ \begin{array}{l} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = P(\tilde{s}' \mid \tilde{s}, \tilde{a}) \\ \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq \tilde{s}'}} \theta_{u_t} = P(s_{t+1} \mid s_t, a_t) - P(\tilde{s}' \mid \tilde{s}, \tilde{a}) \\ \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = 0 \\ \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq \tilde{s}'}} \theta_{u_t} = 1 - P(s_{t+1} \mid s_t, a_t) \end{array} \right.$$

This assignment of  $\theta$  satisfies the constraints of the linear optimisation problem, as follows:

- $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1}}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq \tilde{s}'}} \theta_{u_t} = P(s_{t+1} \mid s_t, a_t)$   
satisfying (7).
- $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1}}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq \tilde{s}'}} \theta_{u_t} = 1 - P(s_{t+1} \mid s_t, a_t)$  satisfying (7).
- $\sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = P(\tilde{s}' \mid \tilde{s}, \tilde{a})$  satisfying (7).
- $\sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} \sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq \tilde{s}'}} \theta_{u_t}$   
 $= 1 - P(\tilde{s}' \mid \tilde{s}, \tilde{a}) = \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} P(s' \mid \tilde{s}, \tilde{a})$ , therefore it is possible to assign  $\theta$  such that  $\forall s' \in \mathcal{S} \setminus \{\tilde{s}'\}, \sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = P(s' \mid \tilde{s}, \tilde{a})$  satisfying (7).
- Mon1 (8), Mon2 (9) and CS (10) are all vacuously true for all transitions from  $(\tilde{s}, \tilde{a})$  due to Lemma B.1.
- $0 \leq \theta_{u_t} \leq 1, \forall u_t$ , satisfying (11).
- $\sum_{u_t=1}^{|U_t|} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t}$   
 $+ \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq \tilde{s}'}} \theta_{u_t} = 1$ , satisfying (12).

Therefore,

$$\tilde{P}_t^{UB}(\tilde{s}' \mid \tilde{s}, \tilde{a}) = \left( \max_{\theta} \frac{\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t}}{P(s_{t+1} \mid s_t, a_t)} \right) = \frac{P(\tilde{s}' \mid \tilde{s}, \tilde{a})}{P(s_{t+1} \mid s_t, a_t)}$$

**Case 2:**  $P(\tilde{s}' \mid \tilde{s}, \tilde{a}) \geq P(s_{t+1} \mid s_t, a_t)$

**Proof B.4** In this case, assign  $\theta$  as follows:

$$\left\{ \begin{array}{l} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = P(s_{t+1} \mid s_t, a_t) \\ \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq \tilde{s}'}} \theta_{u_t} = 0 \\ \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = P(\tilde{s}' \mid \tilde{s}, \tilde{a}) - P(s_{t+1} \mid s_t, a_t) \\ \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq \tilde{s}'}} \theta_{u_t} = 1 - P(\tilde{s}' \mid \tilde{s}, \tilde{a}) \end{array} \right.$$

This assignment of  $\theta$  satisfies the constraints of the linear optimisation problem, as follows:

- $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1}}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq \tilde{s}'}} \theta_{u_t} = P(s_{t+1} | s_t, a_t)$   
satisfying (7).
- $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1}}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq \tilde{s}'}} \theta_{u_t} = 1 - P(s_{t+1} | s_t, a_t)$  satisfying (7).
- $\sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = P(\tilde{s}' | \tilde{s}, \tilde{a})$  satisfying (7).
- $\sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} \sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq \tilde{s}'}} \theta_{u_t}$   
 $= 1 - P(\tilde{s}' | \tilde{s}, \tilde{a}) = \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} P(s' | \tilde{s}, \tilde{a})$ , therefore it is possible to assign  $\theta$  such that  $\forall s' \in \mathcal{S} \setminus \{\tilde{s}'\}, \sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = P(s' | \tilde{s}, \tilde{a})$  satisfying (7).
- Mon1 (8), Mon2 (9) and CS (10) are all vacuously true for all transitions from  $(\tilde{s}, \tilde{a})$  due to Lemma B.1.
- $0 \leq \theta_{u_t} \leq 1, \forall u_t$ , satisfying (11).
- $\sum_{u_t=1}^{|U_t|} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t}$   
 $+ \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq \tilde{s}'}} \theta_{u_t} = 1$ , satisfying (12).

All the constraints are satisfied, so this is a valid assignment of  $\theta$  for this state-action pair. Therefore,

$$\tilde{P}_t^{UB}(\tilde{s}' | \tilde{s}, \tilde{a}) = \max_{\theta} \left( \frac{\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t}}{P(s_{t+1} | s_t, a_t)} \right) = \frac{P(s_{t+1} | s_t, a_t)}{P(s_{t+1} | s_t, a_t)} = 1$$

These two cases can be combined as follows:

$$\tilde{P}_t^{UB}(\tilde{s}' | \tilde{s}, \tilde{a}) = \begin{cases} \frac{P(\tilde{s}' | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} & \text{if } P(\tilde{s}' | \tilde{s}, \tilde{a}) < P(s_{t+1} | s_t, a_t) \\ 1 & \text{otherwise} \end{cases}$$

for all next states  $\tilde{s}' \in \mathcal{S}$ .

**Theorem B.4** For outgoing transitions from state-action pairs  $(\tilde{s}, \tilde{a})$  which have completely disjoint support from the observed  $(s_t, a_t)$ , the linear program will produce a lower bound of:

$$\tilde{P}_t^{LB}(\tilde{s}' | \tilde{s}, \tilde{a}) = \begin{cases} \frac{P(\tilde{s}' | \tilde{s}, \tilde{a}) - (1 - P(s_{t+1} | s_t, a_t))}{P(s_{t+1} | s_t, a_t)} & \text{if } P(\tilde{s}' | \tilde{s}, \tilde{a}) > 1 - P(s_{t+1} | s_t, a_t) \\ 0 & \text{otherwise} \end{cases}$$

for all possible next states  $\tilde{s}'$ .

**Proof B.5** Take an arbitrary transition  $\tilde{s}, \tilde{a} \rightarrow \tilde{s}'$  where  $(\tilde{s}, \tilde{a})$  is disjoint from the observed state-action pair  $(s_t, a_t)$ . By definition:

$$\tilde{P}_t^{LB}(\tilde{s}' | \tilde{s}, \tilde{a}) = \min_{\theta} \left( \frac{\sum_{u_t=1}^{|U_t|} \mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} \cdot \mu_{s_t, a_t, u_t, s_{t+1}} \cdot \theta_{u_t}}{P(s_{t+1} | s_t, a_t)} \right)$$

Because  $P(s_{t+1} | s_t, a_t)$  (the interventional probability of the observed transition) is fixed, to find the lower bound for  $\tilde{P}_t(\tilde{s}' | \tilde{s}, \tilde{a})$  we need to minimise  $\sum_{u_t=1}^{|U_t|} \mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} \cdot \mu_{s_t, a_t, u_t, s_{t+1}} \cdot \theta_{u_t}$ . Because  $\sum_{u_t=1}^{|U_t|} \mu_{s_t, a_t, u_t, s_{t+1}} \cdot \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{u_t=1}^{|U_t|} \mu_{s_t, a_t, u_t, s'} \cdot \theta_{u_t} = 1$ , this is equivalent to maximising the following equation:

$$\max_{\theta} \left( \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{u_t=1}^{|U_t|} \mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} \cdot \mu_{s_t, a_t, u_t, s'} \cdot \theta_{u_t} \right)$$

From the constraints of the optimisation problem, we have:

$$\sum_{u_t=1}^{|U_t|} \mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} \cdot \theta_{u_t} = P(\tilde{s}' | \tilde{s}, \tilde{a}) \quad (15)$$

and

$$\sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{u_t=1}^{|U_t|} \mu_{s_t, a_t, u_t, s'} \cdot \theta_{u_t} = \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} P(s' | s_t, a_t) = 1 - P(s_{t+1} | s_t, a_t) \quad (16)$$

$\forall s, s' \in \mathcal{S}, \forall a \in \mathcal{A}, \forall u_t \in U_t, \mu_{s, a, u_t, s'} \in \{0, 1\}$  (from its definition). Therefore,

$\sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{u_t=1}^{|U_t|} \mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} \cdot \mu_{s_t, a_t, u_t, s'} \cdot \theta_{u_t}$  is maximal when the probability assigned to all  $u_t$  where  $\mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} = 1$  and  $\mu_{s_t, a_t, u_t, s_{t+1}} = 0$  is maximal:

$$\begin{aligned} \sum_{u_t=1}^{|U_t|} (\mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} \cdot \theta_{u_t}) \cdot \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \mu_{s_t, a_t, u_t, s'} &\leq \sum_{u_t=1}^{|U_t|} \mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} \cdot \theta_{u_t} = P(\tilde{s}' | \tilde{s}, \tilde{a}) \\ \sum_{u_t=1}^{|U_t|} \mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} \cdot \left( \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \mu_{s_t, a_t, u_t, s'} \cdot \theta_{u_t} \right) &\leq \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{u_t=1}^{|U_t|} \mu_{s_t, a_t, u_t, s'} \cdot \theta_{u_t} = 1 - P(s_{t+1} | s_t, a_t) \end{aligned}$$

Because of Eq. (15) and Eq. (16), the probability assigned to all  $u_t$  where  $\mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} = 1$  and  $\mu_{s_t, a_t, u_t, s_{t+1}} = 0$  cannot be greater than  $P(\tilde{s}' | \tilde{s}, \tilde{a})$  or  $1 - P(s_{t+1} | s_t, a_t)$ . So,

$$\sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{u_t=1}^{|U_t|} \mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} \cdot \mu_{s_t, a_t, u_t, s'} \cdot \theta_{u_t} \leq \min(P(\tilde{s}' | \tilde{s}, \tilde{a}), (1 - P(s_{t+1} | s_t, a_t)))$$

Therefore, if we can show that there exists a  $\theta$  such that

$$\sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{u_t=1}^{|U_t|} \mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} \cdot \mu_{s_t, a_t, u_t, s'} \cdot \theta_{u_t} = \min(P(\tilde{s}' | \tilde{s}, \tilde{a}), (1 - P(s_{t+1} | s_t, a_t)))$$

and  $\theta$  satisfies the constraints of the optimisation problem, we know that this results in the minimum possible counterfactual transition probability  $\tilde{P}_t(\tilde{s}' | \tilde{s}, \tilde{a})$ . Since  $\forall s, s' \in \mathcal{S}, \forall a \in \mathcal{A}, \forall u_t \in U_t, \mu_{s,a,u_t,s'} = 1 \iff f(s, a, u_t) = s'$  by definition of  $\mu$ , this is equivalent to showing that there exists a  $\theta$  such that:

$$\sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s' \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = \min(P(\tilde{s}' | \tilde{s}, \tilde{a}), (1 - P(s_{t+1} | s_t, a_t)))$$

As these notations are equivalent, we will use this second notation for brevity and clarity. Consider the following disjoint cases:

- $P(\tilde{s}' | \tilde{s}, \tilde{a}) > 1 - P(s_{t+1} | s_t, a_t)$
- $P(\tilde{s}' | \tilde{s}, \tilde{a}) \leq 1 - P(s_{t+1} | s_t, a_t)$

**Case 1:**  $P(\tilde{s}' | \tilde{s}, \tilde{a}) > 1 - P(s_{t+1} | s_t, a_t)$

**Proof B.6** In this case,

$$\sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s' \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} \leq 1 - P(s_{t+1} | s_t, a_t)$$

and so

$$\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} \geq \sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} - \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s' \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = P(\tilde{s}' | \tilde{s}, \tilde{a}) - (1 - P(s_{t+1} | s_t, a_t))$$

If we can show that there exists a  $\theta$  such that

$$\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = P(\tilde{s}' | \tilde{s}, \tilde{a}) - (1 - P(s_{t+1} | s_t, a_t))$$

and  $\theta$  satisfies the constraints in the optimisation problem, we know that this results in the minimum possible counterfactual transition probability for  $\tilde{P}_t(\tilde{s}' | \tilde{s}, \tilde{a})$ . We can assign  $\theta$  as follows:

$$\left\{ \begin{array}{l} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = P(\tilde{s}' | \tilde{s}, \tilde{a}) - (1 - P(s_{t+1} | s_t, a_t)) \\ \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq \tilde{s}'}} \theta_{u_t} = 1 - P(\tilde{s}' | \tilde{s}, \tilde{a}) \\ \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = 1 - P(s_{t+1} | s_t, a_t) \\ \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq \tilde{s}'}} \theta_{u_t} = 0 \end{array} \right.$$

This assignment of  $\theta$  satisfies the constraints of the linear optimisation problem, as follows:

- $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1}}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq \tilde{s}'}} \theta_{u_t} = P(s_{t+1} | s_t, a_t)$   
satisfying (7).

- $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1}}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq \tilde{s}'}} \theta_{u_t} = 1 - P(s_{t+1} | s_t, a_t)$  satisfying (7).
- $\sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = P(\tilde{s}' | \tilde{s}, \tilde{a})$  satisfying (7).
- $\sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} \sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq \tilde{s}'}} \theta_{u_t}$   
 $= 1 - P(\tilde{s}' | \tilde{s}, \tilde{a}) = \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} P(s' | \tilde{s}, \tilde{a})$ , therefore it is possible to assign  $\theta$  such that  $\forall s' \in \mathcal{S} \setminus \{\tilde{s}'\}$ ,  $\sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = P(s' | \tilde{s}, \tilde{a})$  satisfying (7).
- Mon1 (8), Mon2 (9) and CS (10) are all vacuously true for all transitions from  $(\tilde{s}, \tilde{a})$  due to Lemma B.1.
- $0 \leq \theta_{u_t} \leq 1, \forall u_t$ , satisfying (11).
- $\sum_{u_t=1}^{|U_t|} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t}$   
 $+ \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq \tilde{s}'}} \theta_{u_t} = 1$ , satisfying (12).

All the constraints are satisfied, so this is a valid assignment of  $\theta$  for this state-action pair. Therefore,

$$\tilde{P}_t^{LB}(\tilde{s}' | \tilde{s}, \tilde{a}) = \frac{\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t}}{P(s_{t+1} | s_t, a_t)} = \frac{P(\tilde{s}' | \tilde{s}, \tilde{a}) - (1 - P(s_{t+1} | s_t, a_t))}{P(s_{t+1} | s_t, a_t)}$$

**Case 2:**  $P(\tilde{s}' | \tilde{s}, \tilde{a}) \leq 1 - P(s_{t+1} | s_t, a_t)$

**Proof B.7** In this case,

$$\sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s' \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} \leq P(\tilde{s}' | \tilde{s}, \tilde{a})$$

and so

$$\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} \geq \sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} - \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s' \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = P(\tilde{s}' | \tilde{s}, \tilde{a}) - P(\tilde{s}' | \tilde{s}, \tilde{a}) = 0$$

If we can show that there exists a  $\theta$  such that

$$\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = 0$$

and  $\theta$  satisfies the constraints in the optimisation problem, we know that this results in the minimum possible counterfactual transition probability for  $\tilde{P}_t(\tilde{s}' | \tilde{s}, \tilde{a})$ . We can assign  $\theta$  as follows:



$$\left\{ \begin{array}{l} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = 0 \\ \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq \tilde{s}'}} \theta_{u_t} = P(s_{t+1} \mid s_t, a_t) \\ \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = P(\tilde{s}' \mid \tilde{s}, \tilde{a}) \\ \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq \tilde{s}'}} \theta_{u_t} = 1 - P(s_{t+1} \mid s_t, a_t) - P(\tilde{s}' \mid \tilde{s}, \tilde{a}) \end{array} \right.$$

This assignment of  $\theta$  satisfies the constraints of the linear optimisation problem, as follows:

- $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1}}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq \tilde{s}'}} \theta_{u_t} = P(s_{t+1} \mid s_t, a_t)$   
satisfying (7).
- $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1}}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq \tilde{s}'}} \theta_{u_t} = 1 - P(s_{t+1} \mid s_t, a_t)$  satisfying (7).
- $\sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = P(\tilde{s}' \mid \tilde{s}, \tilde{a})$  satisfying (7).
- $\sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} \sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq \tilde{s}'}} \theta_{u_t}$   
 $= 1 - P(\tilde{s}' \mid \tilde{s}, \tilde{a}) = \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} P(s' \mid \tilde{s}, \tilde{a})$ , therefore it is possible to assign  $\theta$  such that  $\forall s' \in \mathcal{S} \setminus \{\tilde{s}'\}, \sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = P(s' \mid \tilde{s}, \tilde{a})$  satisfying (7).
- Mon1 (8), Mon2 (9) and CS (10) are all vacuously true for all transitions from  $(\tilde{s}, \tilde{a})$  due to Lemma B.1.
- $0 \leq \theta_{u_t} \leq 1, \forall u_t$ , satisfying (11).
- $\sum_{u_t=1}^{|U_t|} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t}$   
 $+ \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq \tilde{s}'}} \theta_{u_t} = 1$ , satisfying (12).

All the constraints are satisfied, so this is a valid assignment of  $\theta$  for this state-action pair. Therefore,

$$\tilde{P}_t^{LB}(s' \mid \tilde{s}, \tilde{a}) = \frac{\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t}}{P(s_{t+1} \mid s_t, a_t)} = \frac{0}{P(s_{t+1} \mid s_t, a_t)} = 0$$

These two cases can be combined as follows:

$$\tilde{P}_t^{LB}(s' \mid \tilde{s}, \tilde{a}) = \begin{cases} \frac{P(\tilde{s}' \mid \tilde{s}, \tilde{a}) - (1 - P(s_{t+1} \mid s_t, a_t))}{P(s_{t+1} \mid s_t, a_t)} & \text{if } P(\tilde{s}' \mid \tilde{s}, \tilde{a}) > 1 - P(s_{t+1} \mid s_t, a_t) \\ 0 & \text{otherwise} \end{cases}$$

for all next states  $s' \in \mathcal{S}$ .

### B.1.5 CF Bounds for $(\tilde{s}, \tilde{a})$ with Overlapping Support with $(s_t, a_t)$

To enhance readability, we split the proof for Theorem 4.3 into two separate theorems: Theorem B.5 proves the upper counterfactual probability bounds, and Theorem B.6 proves the lower counterfactual probability bounds.

**Theorem B.5** For outgoing transitions from state-action pairs  $(\tilde{s}, \tilde{a})$  which have overlapping support with the observed  $(s_t, a_t)$ , but  $(\tilde{s}, \tilde{a}) \neq (s_t, a_t)$ , the linear program will produce an upper bound of:

$$\tilde{P}_t^{UB}(\tilde{s}' | \tilde{s}, \tilde{a}) = \begin{cases} \frac{\min(P(s_{t+1} | s_t, a_t), P(s_{t+1} | \tilde{s}, \tilde{a}))}{P(s_{t+1} | s_t, a_t)} & \text{if } \tilde{s}' = s_{t+1} \\ 0 & \text{if } P(\tilde{s}' | s_t, a_t) > 0 \\ & \text{and } \frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} > \frac{P(\tilde{s}' | \tilde{s}, \tilde{a})}{P(\tilde{s}' | s_t, a_t)} \\ \min(P(\tilde{s}' | \tilde{s}, \tilde{a}), 1 - P(s_{t+1} | \tilde{s}, \tilde{a})) & \text{if } P(\tilde{s}' | s_t, a_t) > 0 \\ \min(1 - P(s_{t+1} | \tilde{s}, \tilde{a}), \frac{P(\tilde{s}' | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)}) & \text{otherwise} \end{cases}$$

for all possible next states  $\tilde{s}'$ .

**Proof B.8** Take an arbitrary transition  $\tilde{s}, \tilde{a} \rightarrow \tilde{s}'$  where  $(\tilde{s}, \tilde{a})$  has overlapping support with the observed state-action pair  $(s_t, a_t)$  (and  $(\tilde{s}, \tilde{a}) \neq (s_t, a_t)$ ). Consider the following distinct cases:

- $\tilde{s}' = s_{t+1}$
- $\tilde{s}' \neq s_{t+1}$ ,  $\frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} > \frac{P(\tilde{s}' | \tilde{s}, \tilde{a})}{P(\tilde{s}' | s_t, a_t)}$  and  $P(\tilde{s}' | s_t, a_t) > 0$
- $\tilde{s}' \neq s_{t+1}$ ,  $\frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} \leq \frac{P(\tilde{s}' | \tilde{s}, \tilde{a})}{P(\tilde{s}' | s_t, a_t)}$  and  $P(\tilde{s}' | s_t, a_t) > 0$
- $\tilde{s}' \neq s_{t+1}$  and  $P(\tilde{s}' | s_t, a_t) = 0$

These cases are disjoint and cover all possible situations.

**Case 1:**  $\tilde{s}' = s_{t+1}$

**Proof B.9** Consider the following disjoint cases:

- $P(s_{t+1} | \tilde{s}, \tilde{a}) = 0$
- $P(s_{t+1} | \tilde{s}, \tilde{a}) > 0$  and  $P(s_{t+1} | \tilde{s}, \tilde{a}) \geq P(s_{t+1} | s_t, a_t)$
- $P(s_{t+1} | \tilde{s}, \tilde{a}) > 0$  and  $P(s_{t+1} | \tilde{s}, \tilde{a}) < P(s_{t+1} | s_t, a_t)$

**Case 1(a):**  $P(s_{t+1} | \tilde{s}, \tilde{a}) = 0$

**Proof B.10** If  $P(s_{t+1} | \tilde{s}, \tilde{a}) = 0$ , then

$\sum_{u_t=1}^{|U_t|} \mu_{\tilde{s}, \tilde{a}, u_t, s_{t+1}} \cdot \theta_{u_t} = 0$  to satisfy (7). Therefore, the maximum counterfactual probability  $\tilde{P}_t(s_{t+1} | \tilde{s}, \tilde{a})$  is:

$$\max_{\theta} \left( \sum_{u_t=1}^{|U_t|} \mu_{\tilde{s}, \tilde{a}, u_t, s_{t+1}} \cdot \mu_{s_t, a_t, u_t, s_{t+1}} \cdot \theta_{u_t} = 0 \right)$$

If we can show that there exists a  $\theta$  such that

$$\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} = 0$$

and  $\theta$  satisfies the constraints in the optimisation problem, we know that this results in the maximum possible counterfactual transition probability for  $\tilde{P}_t(s_{t+1} | \tilde{s}, \tilde{a})$ . We can assign  $\theta$  as follows:

$$\forall s' \in \mathcal{S}, \quad \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = P(s' | \tilde{s}, \tilde{a}) \cdot P(s_{t+1} | s_t, a_t)$$

$$\forall s' \in \mathcal{S}, \quad \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = P(s' | \tilde{s}, \tilde{a}) \cdot (1 - P(s_{t+1} | s_t, a_t))$$

which results in a counterfactual probability of  $\tilde{P}_t(s_{t+1} | \tilde{s}, \tilde{a}) = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1}}} \theta_{u_t}$   
 $= \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} = P(s_{t+1} | \tilde{s}, \tilde{a}) \cdot P(s_{t+1} | s_t, a_t) = 0.$

This assignment of  $\theta$  satisfies the constraints of the linear optimisation problem, as follows:

- $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1}}} \theta_{u_t} = \sum_{s' \in \mathcal{S}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = P(s_{t+1} | s_t, a_t)$ , satisfying (7).
- $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1}}} \theta_{u_t} = \sum_{s' \in \mathcal{S}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = 1 - P(s_{t+1} | s_t, a_t)$ , satisfying (7).
- $\forall s' \in \mathcal{S}, \sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = P(s' | \tilde{s}, \tilde{a})$ , satisfying (7).
- $\tilde{P}_t(s_{t+1} | \tilde{s}, \tilde{a}) = \frac{\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t}}{P(s_{t+1} | \tilde{s}, \tilde{a})} = \frac{P(s_{t+1} | \tilde{s}, \tilde{a}) \cdot P(s_{t+1} | s_t, a_t)}{P(s_{t+1} | \tilde{s}, \tilde{a})} = P(s_{t+1} | \tilde{s}, \tilde{a}) \geq P(s_{t+1} | \tilde{s}, \tilde{a})$  so Mon1 (8) holds.
- $\forall s' \in \mathcal{S} \setminus \{s_{t+1}\}$ :

$$\begin{aligned} \tilde{P}_t(s' | \tilde{s}, \tilde{a}) &= \frac{\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t}}{P(s_{t+1} | \tilde{s}, \tilde{a})} \\ &= \frac{P(s' | \tilde{s}, \tilde{a}) \cdot P(s_{t+1} | s_t, a_t)}{P(s_{t+1} | \tilde{s}, \tilde{a})} \\ &= P(s' | \tilde{s}, \tilde{a}) \\ &\leq P(s' | \tilde{s}, \tilde{a}) \end{aligned}$$

so Mon2 (9) holds.

- Because  $P(s_{t+1} | \tilde{s}, \tilde{a}) = 0$  and  $P(s_{t+1} | s_t, a_t) > 0$  (since this was observed):

$$\frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} = 0$$

Therefore,  $\forall s' \in \mathcal{S}$

$$\frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} \leq \frac{P(s' | \tilde{s}, \tilde{a})}{P(s' | s_t, a_t)}$$

so CS (10) holds  $\forall s' \in \mathcal{S}$ .

- $0 \leq \theta_{u_t} \leq 1, \forall u_t$ , satisfying (11).

- $\sum_{u_t=1}^{|U_t|} \theta_{u_t} = \sum_{s' \in \mathcal{S}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} + \sum_{s' \in \mathcal{S}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = 1$ , satisfying (12).

Therefore,

$$\tilde{P}_t^{UB}(\tilde{s}' \mid \tilde{s}, \tilde{a}) = \max_{\theta} \left( \frac{\sum_{u_t=1}^{|U_t|} \mu_{\tilde{s}, \tilde{a}, u_t, s_{t+1}} \cdot \mu_{s_t, a_t, u_t, s_{t+1}} \cdot \theta_{u_t}}{P(s_{t+1} \mid s_t, a_t)} \right) = \frac{P(s_{t+1} \mid \tilde{s}, \tilde{a})}{P(s_{t+1} \mid s_t, a_t)}$$

**Case 1(b):**  $P(s_{t+1} \mid \tilde{s}, \tilde{a}) > 0$  and  $P(s_{t+1} \mid \tilde{s}, \tilde{a}) \geq P(s_{t+1} \mid s_t, a_t)$

**Proof B.11** From Lemma B.2, we have:

$$\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} \leq \min(P(s_{t+1} \mid s_t, a_t), P(s_{t+1} \mid \tilde{s}, \tilde{a}))$$

Therefore,

$$\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} \leq P(s_{t+1} \mid s_t, a_t)$$

If we can show that there exists a  $\theta$  such that

$$\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} = P(s_{t+1} \mid s_t, a_t)$$

and  $\theta$  satisfies the constraints in the optimisation problem, we know that this results in the maximum possible counterfactual transition probability for  $\tilde{P}_t(s_{t+1} \mid \tilde{s}, \tilde{a})$ . We can assign  $\theta$  as follows:

$$\begin{aligned} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} &= P(s_{t+1} \mid s_t, a_t) \\ \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} &= P(s_{t+1} \mid \tilde{s}, \tilde{a}) - P(s_{t+1} \mid s_t, a_t) \\ \forall s' \in \mathcal{S} \setminus \{s_{t+1}\} \quad \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} &= 0 \\ \forall s' \in \mathcal{S} \setminus \{s_{t+1}\} \quad \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} &= P(s' \mid \tilde{s}, \tilde{a}) \end{aligned}$$

This assignment of  $\theta$  satisfies the constraints of the linear optimisation problem, as follows:

- $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1}}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq s_{t+1}}} \theta_{u_t} = P(s_{t+1} \mid s_t, a_t)$ , satisfying (7).

- $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1}}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq s_{t+1}}} \theta_{u_t} = 1 - P(s_{t+1} | s_t, a_t)$ , satisfying (7).
- $\forall s' \in \mathcal{S}$ ,  $\sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = P(s' | \tilde{s}, \tilde{a})$ , satisfying (7).
- $\tilde{P}_t(s_{t+1} | \tilde{s}, \tilde{a}) = \frac{\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t}}{P(s_{t+1} | s_t, a_t)} = \frac{P(s_{t+1} | s_t, a_t)}{P(s_{t+1} | s_t, a_t)} = 1 \geq P(s_{t+1} | \tilde{s}, \tilde{a})$  so Mon1 (8) holds.
- $\forall s' \in \mathcal{S} \setminus \{s_{t+1}\}$ :

$$\begin{aligned} \tilde{P}_t(s' | \tilde{s}, \tilde{a}) &= \frac{\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t}}{P(s_{t+1} | s_t, a_t)} \\ &= \frac{0}{P(s_{t+1} | s_t, a_t)} \\ &= 0 \\ &\leq P(s' | \tilde{s}, \tilde{a}) \end{aligned}$$

so Mon2 (9) holds.

- CS (10) vacuously holds for the transition  $\tilde{s}, \tilde{a} \rightarrow s_{t+1}$ , because  $\frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} = \frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)}$ . Also,  $\forall s' \in \mathcal{S} \setminus \{s_{t+1}\}$ ,  $\tilde{P}_t(s' | \tilde{s}, \tilde{a}) = 0$  which satisfies CS (10).
- $0 \leq \theta_{u_t} \leq 1, \forall u_t$ , satisfying (11).
- $\sum_{u_t=1}^{|U_t|} \theta_{u_t} = \sum_{s' \in \mathcal{S}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} + \sum_{s' \in \mathcal{S}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = 1$ , satisfying (12).

Therefore,

$$\max_{\theta} \left( \sum_{u_t=1}^{|U_t|} \mu_{\tilde{s}, \tilde{a}, u_t, s_{t+1}} \cdot \mu_{s_t, a_t, u_t, s_{t+1}} \cdot \theta_{u_t} \right) = P(s_{t+1} | s_t, a_t)$$

and

$$\tilde{P}_t^{UB}(\tilde{s}' | \tilde{s}, \tilde{a}) = \max_{\theta} \left( \frac{\sum_{u_t=1}^{|U_t|} \mu_{\tilde{s}, \tilde{a}, u_t, s_{t+1}} \cdot \mu_{s_t, a_t, u_t, s_{t+1}} \cdot \theta_{u_t}}{P(s_{t+1} | s_t, a_t)} \right) = \frac{P(s_{t+1} | s_t, a_t)}{P(s_{t+1} | s_t, a_t)} = 1$$

**Case 1(c):**  $P(s_{t+1} | \tilde{s}, \tilde{a}) > 0$  and  $P(s_{t+1} | \tilde{s}, \tilde{a}) < P(s_{t+1} | s_t, a_t)$

**Proof B.12** From Lemma B.2, we have:

$$\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} \leq \min(P(s_{t+1} | s_t, a_t), P(s_{t+1} | \tilde{s}, \tilde{a}))$$

Therefore,

$$\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} \leq P(s_{t+1} | \tilde{s}, \tilde{a})$$

If we can show that there exists a  $\theta$  such that

$$\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} = P(s_{t+1} | \tilde{s}, \tilde{a})$$

and  $\theta$  satisfies the constraints in the optimisation problem, we know that this results in the maximum possible counterfactual transition probability for  $\tilde{P}_t(s_{t+1} \mid \tilde{s}, \tilde{a})$ . We can assign  $\theta$  such that:

$$\begin{aligned}
\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} &= P(s_{t+1} \mid \tilde{s}, \tilde{a}) \\
\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq s_{t+1}}} \theta_{u_t} &= P(s_{t+1} \mid s_t, a_t) - P(s_{t+1} \mid \tilde{s}, \tilde{a}) \\
\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} &= 0 \\
\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq s_{t+1}}} \theta_{u_t} &= \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} P(s' \mid \tilde{s}, \tilde{a}) - \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq s_{t+1}}} \theta_{u_t}
\end{aligned}$$

However, it is not immediately clear whether this assignment of  $\theta$  is possible, because the Mon2 (9) and CS (10) constraints limit  $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq s'}} \theta_{u_t}$  for all  $s' \in \mathcal{S} \setminus \{s_{t+1}\}$ . In particular, there exists a set  $S_{CS} \subset \mathcal{S}$  of states which must have a

counterfactual transition probability  $\tilde{P}_t(s' \mid \tilde{s}, \tilde{a}) = 0$  to satisfy CS (10) (i.e.,  $\forall s' \in S_{CS}, \frac{P(s_{t+1} \mid \tilde{s}, \tilde{a})}{P(s_{t+1} \mid s_t, a_t)} > \frac{P(s' \mid \tilde{s}, \tilde{a})}{P(s' \mid s_t, a_t)}$  and  $P(s' \mid s_t, a_t) > 0$ ). This means that  $\forall s' \in S_{CS}$  we must assign  $\theta$  such that:

$$\begin{aligned}
\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} &= 0 \\
\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} &= P(s' \mid \tilde{s}, \tilde{a})
\end{aligned}$$

As this results in  $\tilde{P}_t(s' \mid \tilde{s}, \tilde{a}) = 0$ :

$$\begin{aligned}
\tilde{P}_t(s' \mid \tilde{s}, \tilde{a}) &= \frac{\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t}}{P(s_{t+1} \mid s_t, a_t)} \\
&= \frac{0}{P(s_{t+1} \mid s_t, a_t)} \\
&= 0
\end{aligned} \tag{17}$$

Let  $S_{other}$  be the set of all other states  $s \in \mathcal{S} \setminus \{s_{t+1}\} \cup S_{CS}$ .  $\forall s' \in S_{other}$ , if  $P(s' \mid \tilde{s}, \tilde{a}) > 0$  and  $P(s' \mid s_t, a_t) > 0$ , then  $\tilde{P}_t(s' \mid \tilde{s}, \tilde{a}) \leq P(s' \mid \tilde{s}, \tilde{a})$  to satisfy Mon2 (9). Because we do not know whether  $P(s' \mid \tilde{s}, \tilde{a}) > 0$  and  $P(s' \mid s_t, a_t) > 0$  for all  $s' \in S_{other}$ , we have to assume that this condition holds for all  $s' \in S_{other}$ , and assign  $\theta$  with this assumption. This means,  $\forall s' \in S_{other}$ :

$$\begin{aligned}
\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} &\leq P(s' | \tilde{s}, \tilde{a}) \cdot P(s_{t+1} | s_t, a_t) \\
\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} &\geq P(s' | \tilde{s}, \tilde{a}) - (P(s' | \tilde{s}, \tilde{a}) \cdot P(s_{t+1} | s_t, a_t))
\end{aligned} \tag{18}$$

as this ensures each counterfactual transition probability satisfies Mon2 (9), as follows:

$$\begin{aligned}
\tilde{P}_t(s' | \tilde{s}, \tilde{a}) &= \frac{\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t}}{P(s_{t+1} | s_t, a_t)} \\
&\leq \frac{P(s' | \tilde{s}, \tilde{a}) \cdot P(s_{t+1} | s_t, a_t)}{P(s_{t+1} | s_t, a_t)} \\
&= P(s' | \tilde{s}, \tilde{a})
\end{aligned}$$

Therefore, to show that there exists a valid assignment of  $\theta$ , we must prove that under the Mon2 (9) and CS (10) constraints:

$$\max_{\theta} \left( \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} \right) \geq P(s_{t+1} | s_t, a_t) - P(s_{t+1} | \tilde{s}, \tilde{a})$$

This is proven in Lemma B.4. Now, we can prove that this assignment of  $\theta$  satisfies the constraints of the linear optimisation problem, as follows:

- $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1}}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq s_{t+1}}} \theta_{u_t} = P(s_{t+1} | s_t, a_t)$ , satisfying (7).
- $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1}}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) \neq s_{t+1}}} \theta_{u_t} = 1 - P(s_{t+1} | s_t, a_t)$ , satisfying (7).
- $\sum_{s' \in \mathcal{S}} \sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = \sum_{s' \in \mathcal{S}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} + \sum_{s' \in \mathcal{S}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t}$   
 $= P(s' | \tilde{s}, \tilde{a})$ , therefore it is possible to assign  $\theta$  such that  $\forall s' \in \mathcal{S} \setminus \{s'\}, \sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = P(s' | \tilde{s}, \tilde{a})$ , satisfying (7).
- $\tilde{P}_t(s_{t+1} | \tilde{s}, \tilde{a}) = \frac{\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t}}{P(s_{t+1} | s_t, a_t)} = \frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} \geq P(s_{t+1} | \tilde{s}, \tilde{a})$  so Mon1 (8) holds.
- Lemma B.4 proves that we can assign  $\theta$  such that Mon2 (9) holds.
- CS (10) vacuously holds for the transition  $\tilde{s}, \tilde{a} \rightarrow s_{t+1}$ , because  $\frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} = \frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)}$ . Also, Lemma B.4 proves that we can assign  $\theta$  such that CS (10) holds for all other transitions.
- $0 \leq \theta_{u_t} \leq 1, \forall u_t$ , satisfying (11).
- $\sum_{u_t=1}^{|U_t|} \theta_{u_t} = \sum_{s' \in \mathcal{S}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} + \sum_{s' \in \mathcal{S}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = 1$ , satisfying (12).

Therefore,

$$\max_{\theta} \left( \sum_{u_t=1}^{|U_t|} \mu_{\tilde{s}, \tilde{a}, u_t, s_{t+1}} \cdot \mu_{s_t, a_t, u_t, s_{t+1}} \cdot \theta_{u_t} \right) = P(s_{t+1} | \tilde{s}, \tilde{a})$$

and

$$\tilde{P}_t^{UB}(\tilde{s}' | \tilde{s}, \tilde{a}) = \frac{\max_{\theta} \sum_{u_t=1}^{|U_t|} \mu_{\tilde{s}, \tilde{a}, u_t, s_{t+1}} \cdot \mu_{s_t, a_t, u_t, s_{t+1}} \cdot \theta_{u_t}}{P(s_{t+1} | s_t, a_t)} = \frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)}$$

These three sub-cases can be simplified to:

$$\tilde{P}_t^{UB}(\tilde{s}' | \tilde{s}, \tilde{a}) = \frac{\min(P(s_{t+1} | s_t, a_t), P(s_{t+1} | \tilde{s}, \tilde{a}))}{P(s_{t+1} | s_t, a_t)}$$

**Case 2:**  $\tilde{s}' \neq s_{t+1}$  and  $\frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} > \frac{P(\tilde{s}' | \tilde{s}, \tilde{a})}{P(\tilde{s}' | s_t, a_t)}$  and  $P(\tilde{s}' | s_t, a_t) > 0$

**Proof B.13** Because  $\frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} > \frac{P(\tilde{s}' | \tilde{s}, \tilde{a})}{P(\tilde{s}' | s_t, a_t)}$  and  $P(\tilde{s}' | s_t, a_t) > 0$ , to satisfy CS (10) we must always have  $\tilde{P}_t(\tilde{s}' | \tilde{s}, \tilde{a}) = 0$ , therefore  $\tilde{P}_t^{UB}(\tilde{s}' | \tilde{s}, \tilde{a}) = 0$ . From Case 1, we know that we can find an assignment of  $\theta$  that satisfies CS (10) for all transitions, when finding the upper bound of the transition  $\tilde{s}, \tilde{a} \rightarrow s_{t+1}$ . Therefore, we have already proven in Case 1 that there exists an assignment of  $\theta$  where  $\tilde{P}_t(\tilde{s}' | \tilde{s}, \tilde{a}) = 0$ , that satisfies the constraints (7)-(12). Therefore, there exists an assignment of  $\theta$  that satisfies  $\tilde{P}_t^{UB}(\tilde{s}' | \tilde{s}, \tilde{a}) = 0$ .



**Case 3:**  $\tilde{s}' \neq s_{t+1}$ ,  $\frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} \leq \frac{P(\tilde{s}' | \tilde{s}, \tilde{a})}{P(\tilde{s}' | s_t, a_t)}$ , and  $P(\tilde{s}' | s_t, a_t) > 0$

**Proof B.14** Because  $P(\tilde{s}' | s_t, a_t) > 0$ ,  $\tilde{P}_t^{UB}(\tilde{s}' | \tilde{s}, \tilde{a}) \leq P(\tilde{s}' | \tilde{s}, \tilde{a})$  to satisfy Mon2 (9). Therefore,

$$\sum_{u_t=1}^{|U_t|} \mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} \cdot \mu_{s_t, a_t, u_t, s_{t+1}} \cdot \theta_{u_t} \leq P(\tilde{s}' | \tilde{s}, \tilde{a}) \cdot P(s_{t+1} | s_t, a_t) \quad (19)$$

so that

$$\tilde{P}_t^{UB}(\tilde{s}' | \tilde{s}, \tilde{a}) = \frac{\sum_{u_t=1}^{|U_t|} \mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} \cdot \mu_{s_t, a_t, u_t, s_{t+1}} \cdot \theta_{u_t}}{P(s_{t+1} | s_t, a_t)} \leq \frac{P(\tilde{s}' | \tilde{s}, \tilde{a}) \cdot P(s_{t+1} | s_t, a_t)}{P(s_{t+1} | s_t, a_t)} = P(\tilde{s}' | \tilde{s}, \tilde{a})$$

Clearly, this is less than the maximum possible counterfactual probability proven in Lemma B.2:

$$\sum_{u_t=1}^{|U_t|} \mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} \cdot \mu_{s_t, a_t, u_t, s_{t+1}} \cdot \theta_{u_t} \leq P(\tilde{s}' | \tilde{s}, \tilde{a}) \cdot P(s_{t+1} | s_t, a_t) \leq \min(P(s_{t+1} | s_t, a_t), P(\tilde{s}' | \tilde{s}, \tilde{a}))$$

Also,  $\tilde{P}_t(s_{t+1} | \tilde{s}, \tilde{a}) \geq P(s_{t+1} | \tilde{s}, \tilde{a})$  to satisfy Mon1 (8). Therefore,

$$\sum_{u_t=1}^{|U_t|} \mu_{\tilde{s}, \tilde{a}, u_t, s_{t+1}} \cdot \mu_{s_t, a_t, u_t, s_{t+1}} \cdot \theta_{u_t} \geq P(s_{t+1} | \tilde{s}, \tilde{a}) \cdot P(s_{t+1} | s_t, a_t) \quad (20)$$

Since  $\forall s, s' \in \mathcal{S}, \forall a \in \mathcal{A}, \forall u_t \in U_t, \mu_{s, a, u_t, s'} = 1 \iff f(s, a, u_t) = s'$  by definition of  $\mu$ , Eq. (19) and Eq. (20) are equivalent to:

$$\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} \leq P(\tilde{s}' | \tilde{s}, \tilde{a}) \cdot P(s_{t+1} | s_t, a_t) \quad (21)$$

and

$$\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} \geq P(s_{t+1} | \tilde{s}, \tilde{a}) \cdot P(s_{t+1} | s_t, a_t) \quad (22)$$

respectively. As these notations are equivalent, we will use this second notation for brevity and clarity. Consider the following disjoint cases:

- $P(s_{t+1} | s_t, a_t) - (P(s_{t+1} | s_t, a_t) \cdot P(s_{t+1} | \tilde{s}, \tilde{a})) < P(\tilde{s}' | \tilde{s}, \tilde{a}) \cdot P(s_{t+1} | s_t, a_t)$
- $P(s_{t+1} | s_t, a_t) - (P(s_{t+1} | s_t, a_t) \cdot P(s_{t+1} | \tilde{s}, \tilde{a})) \geq P(\tilde{s}' | \tilde{s}, \tilde{a}) \cdot P(s_{t+1} | s_t, a_t)$

**Case 3(a):**  $P(s_{t+1} | s_t, a_t) - (P(s_{t+1} | s_t, a_t) \cdot P(s_{t+1} | \tilde{s}, \tilde{a})) < P(\tilde{s}' | \tilde{s}, \tilde{a}) \cdot P(s_{t+1} | s_t, a_t)$

**Proof B.15** We can assign  $\theta$  as follows:

$$\left\{ \begin{array}{l} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} = P(s_{t+1} | s_t, a_t) \cdot P(s_{t+1} | \tilde{s}, \tilde{a}) \\ \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = P(s_{t+1} | s_t, a_t) - (P(s_{t+1} | s_t, a_t) \cdot P(s_{t+1} | \tilde{s}, \tilde{a})) \\ \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = 0 \\ \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} = P(s_{t+1} | \tilde{s}, \tilde{a}) - \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} \\ \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = P(\tilde{s}' | \tilde{s}, \tilde{a}) - \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} \\ \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} P(\tilde{s}' | \tilde{s}, \tilde{a}) \end{array} \right.$$

This assignment of  $\theta$  satisfies the constraints of the linear optimisation problem, as follows:

- $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1}}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t}$   
 $= P(s_{t+1} | s_t, a_t)$ , satisfying (7).

- $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1}}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t}$   
 $= 1 - P(s_{t+1} | s_t, a_t)$ , satisfying (7).

- $\sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = P(\tilde{s}' | \tilde{s}, \tilde{a})$ ,  
satisfying (7).

- $\sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} = P(s_{t+1} | \tilde{s}, \tilde{a})$ ,  
satisfying (7).

- $\sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t}$   
 $+ \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = 1 - P(\tilde{s}' | \tilde{s}, \tilde{a}) = \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} P(s' | \tilde{s}, \tilde{a})$ ,

therefore it is possible to assign  $\theta$  such that

$$\forall s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}, \sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = P(s' | \tilde{s}, \tilde{a}), \text{ satisfying (7).}$$

- (22) holds, therefore Mon1 (8) is satisfied.

- (21) holds, therefore  $\tilde{P}_t(\tilde{s}' | \tilde{s}, \tilde{a}) \leq P(\tilde{s}' | \tilde{s}, \tilde{a})$ . Also, because

$$\sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = 0, \text{ the counterfactual probability of all other}$$

transitions  $\tilde{s}, \tilde{a} \rightarrow s'$  where  $s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}$  will be  $\tilde{P}_t(s' | \tilde{s}, \tilde{a}) = 0 \leq P(s' | \tilde{s}, \tilde{a})$ . Therefore, Mon2 (9) is satisfied.

- CS (10) is vacuously satisfied for the transitions  $\tilde{s}, \tilde{a} \rightarrow \tilde{s}'$  and  $\tilde{s}, \tilde{a} \rightarrow s_{t+1}$  because

$$\frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} \leq \frac{P(\tilde{s}' | \tilde{s}, \tilde{a})}{P(\tilde{s}' | s_t, a_t)} \text{ (from conditions of Case 3) and } \frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} = \frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)}. \text{ Also, because}$$

$$\sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = 0, \text{ the counterfactual probability of all other transitions } \tilde{s}, \tilde{a} \rightarrow s' \text{ where}$$

$s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}$  will be  $\tilde{P}_t(s' | \tilde{s}, \tilde{a}) = 0$ , which satisfies CS (10) for these transitions.

•  $0 \leq \theta_{u_t} \leq 1, \forall u_t$ , satisfying (11).

$$\begin{aligned} & \sum_{u_t=1}^{|U_t|} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} \\ & + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = 1, \text{ satisfying (12).} \end{aligned}$$

All the constraints are satisfied, so this is a valid assignment of  $\theta$  for this state-action pair.

**Case 3(b):**  $P(s_{t+1} | s_t, a_t) - (P(s_{t+1} | s_t, a_t) \cdot P(s_{t+1} | \tilde{s}, \tilde{a})) \geq P(\tilde{s}' | \tilde{s}, \tilde{a}) \cdot P(s_{t+1} | s_t, a_t)$

**Proof B.16** If  $P(s_{t+1} | s_t, a_t) - (P(s_{t+1} | s_t, a_t) \cdot P(s_{t+1} | \tilde{s}, \tilde{a})) \geq P(\tilde{s}' | \tilde{s}, \tilde{a}) \cdot P(s_{t+1} | s_t, a_t)$ , this implies  $P(s_{t+1} | s_t, a_t) \geq P(\tilde{s}' | \tilde{s}, \tilde{a}) \cdot P(s_{t+1} | s_t, a_t)$ . Therefore, we can assign  $\theta$  as follows:

$$\left\{ \begin{array}{l} P(s_{t+1} | s_t, a_t) \cdot P(s_{t+1} | \tilde{s}, \tilde{a}) \leq \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} \leq P(s_{t+1} | \tilde{s}, \tilde{a}) \\ \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = P(\tilde{s}' | \tilde{s}, \tilde{a}) \cdot P(s_{t+1} | s_t, a_t) \\ 0 \leq \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} \leq \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} P(s' | \tilde{s}, \tilde{a}) \cdot P(s_{t+1} | s_t, a_t) \\ \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} = P(s_{t+1} | \tilde{s}, \tilde{a}) - \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} \\ \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = P(\tilde{s}' | \tilde{s}, \tilde{a}) - \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} \\ \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} P(s' | \tilde{s}, \tilde{a}) - \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} \end{array} \right.$$

$$\text{s.t. } \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = P(s_{t+1} | s_t, a_t).$$

We do not know exactly how much probability will be assigned to  $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t}$  and

$\sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t}$ . However, if  $\theta$  satisfies the above assignment then  $\theta$  will always satisfy the

constraints of the optimisation problem, as follows:

$$\begin{aligned} & \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1}}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} \\ & = P(s_{t+1} | s_t, a_t), \text{ satisfying (7).} \\ & \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1}}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} \\ & = 1 - P(s_{t+1} | s_t, a_t), \text{ satisfying (7)} \\ & \sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = P(\tilde{s}' | \tilde{s}, \tilde{a}), \\ & \text{satisfying (7).} \\ & \sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} = P(s_{t+1} | \tilde{s}, \tilde{a}), \\ & \text{satisfying (7).} \end{aligned}$$

- $\sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t}$   
 $= 1 - P(\tilde{s}' \mid \tilde{s}, \tilde{a}) = \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} P(s' \mid \tilde{s}, \tilde{a})$ , therefore it is possible to assign  $\theta$  such that  $\forall s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}, \sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = P(s' \mid \tilde{s}, \tilde{a})$ , satisfying (7).

- (22) holds, therefore Mon1 (8) is satisfied.

- (21) holds, so Mon2 (9) holds for the transition  $\tilde{s}, \tilde{a} \rightarrow \tilde{s}'$ .

- CS (10) is vacuously satisfied for the transitions  $\tilde{s}, \tilde{a} \rightarrow \tilde{s}'$  and  $\tilde{s}, \tilde{a} \rightarrow s_{t+1}$  because  $\frac{P(s_{t+1} \mid \tilde{s}, \tilde{a})}{P(s_{t+1} \mid s_t, a_t)} \leq \frac{P(\tilde{s}' \mid \tilde{s}, \tilde{a})}{P(\tilde{s}' \mid s_t, a_t)}$

(from conditions of Case 3) and  $\frac{P(s_{t+1} \mid \tilde{s}, \tilde{a})}{P(s_{t+1} \mid s_t, a_t)} = \frac{P(s_{t+1} \mid \tilde{s}, \tilde{a})}{P(s_{t+1} \mid s_t, a_t)}$ .

- To ensure that Mon2 (9) and CS (10) hold for all other transitions, we must consider two possible cases:

- If  $P(s_{t+1} \mid \tilde{s}, \tilde{a}) \geq P(s_{t+1} \mid s_t, a_t)$ , then we can assign  $\theta$  such that:

$$\sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = 0$$

and

$$\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} = P(s_{t+1} \mid s_t, a_t) - \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t}$$

which must be possible in our given assignment of  $\theta$  because from the condition of Case 3(b) we have:

$$P(s_{t+1} \mid s_t, a_t) - (P(s_{t+1} \mid s_t, a_t) \cdot P(s_{t+1} \mid \tilde{s}, \tilde{a})) \geq P(\tilde{s}' \mid \tilde{s}, \tilde{a}) \cdot P(s_{t+1} \mid s_t, a_t)$$

$$\implies P(s_{t+1} \mid s_t, a_t) - (P(\tilde{s}' \mid \tilde{s}, \tilde{a}) \cdot P(s_{t+1} \mid s_t, a_t)) \geq P(s_{t+1} \mid s_t, a_t) \cdot P(s_{t+1} \mid \tilde{s}, \tilde{a})$$

Because  $\sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = 0, \forall s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}, \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t}$

= 0, and so  $\forall s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}, \tilde{P}_t(s' \mid \tilde{s}, \tilde{a}) = 0$ . This satisfies the Mon2 (9) (since  $0 \leq \tilde{P}_t(s' \mid \tilde{s}, \tilde{a})$ ) and CS (10) constraints for all other transitions  $\tilde{s}, \tilde{a} \rightarrow s'$ .

- Otherwise, we must have  $P(s_{t+1} \mid \tilde{s}, \tilde{a}) < P(s_{t+1} \mid s_t, a_t)$ . We need to prove that there exists an assignment of  $\theta$  where the Mon2 (9) and CS (10) constraints are satisfied for all transitions  $\tilde{s}, \tilde{a} \rightarrow s', \forall s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}$ , which equates to proving:

$$\max_{\theta} \left( \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} \right) \geq \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} P(s' \mid \tilde{s}, \tilde{a}) \cdot P(s_{t+1} \mid s_t, a_t)$$

since

$$\sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} \leq \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} P(s' \mid \tilde{s}, \tilde{a}) \cdot P(s_{t+1} \mid s_t, a_t)$$

in our given assignment of  $\theta$ . From Lemma B.4, we have:

$$\max_{\theta} \left( \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} \right) \geq P(s_{t+1} \mid s_t, a_t) - P(s_{t+1} \mid \tilde{s}, \tilde{a}) \quad (23)$$

under the Mon2 (9) and CS (10) constraints. Since we have assigned the maximum possible probability to  $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = P(\tilde{s}' | \tilde{s}, \tilde{a}) \cdot P(s_{t+1} | s_t, a_t)$  under the Mon2 (9) and CS (10) constraints in our given assignment of  $\theta$ , (23) implies:

$$\max_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} \geq P(s_{t+1} | s_t, a_t) - P(s_{t+1} | \tilde{s}, \tilde{a}) - P(\tilde{s}' | \tilde{s}, \tilde{a}) \cdot P(s_{t+1} | s_t, a_t)$$

Therefore, we know that we can assign  $\theta$  such that  $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = P(s_{t+1} | s_t, a_t)$ , and all transitions satisfy the Mon2 (9) and CS (10) constraints.

•  $0 \leq \theta_{u_t} \leq 1, \forall u_t$ , satisfying (11).

$$\begin{aligned} & \bullet \sum_{u_t=1}^{|U_t|} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} \\ & + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = 1, \text{ satisfying (12).} \end{aligned}$$

All the constraints are satisfied, so this is a valid assignment of  $\theta$  for this state-action pair.

These two cases can be simplified to:

$$\begin{aligned} \tilde{P}_t^{UB}(\tilde{s}' | \tilde{s}, \tilde{a}) &= \frac{\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t}}{P(s_{t+1} | s_t, a_t)} \\ &= \frac{\min(P(s_{t+1} | s_t, a_t) - (P(s_{t+1} | s_t, a_t) \cdot P(s_{t+1} | \tilde{s}, \tilde{a})), P(\tilde{s}' | \tilde{s}, \tilde{a}) \cdot P(s_{t+1} | s_t, a_t))}{P(s_{t+1} | s_t, a_t)} \\ &= \min(1 - P(s_{t+1} | \tilde{s}, \tilde{a}), P(\tilde{s}' | \tilde{s}, \tilde{a})) \end{aligned}$$

**Case 4:**  $\tilde{s}' \neq s_{t+1}$  and  $P(\tilde{s}' | s_t, a_t) = 0$

**Proof B.17** Because  $P(\tilde{s}' | s_t, a_t) = 0$ , then the monotonicity and counterfactual stability constraints are vacuously satisfied for  $\tilde{s}, \tilde{a} \rightarrow \tilde{s}'$ . Therefore, as proven in Lemma B.2:

$$\sum_{u_t=1}^{|U_t|} \mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} \cdot \mu_{s_t, a_t, u_t, s_{t+1}} \cdot \theta_{u_t} \leq \min(P(\tilde{s}' | \tilde{s}, \tilde{a}), P(s_{t+1} | s_t, a_t)) \quad (24)$$

Also,  $\tilde{P}_t(s_{t+1} | \tilde{s}, \tilde{a}) \geq P(s_{t+1} | \tilde{s}, \tilde{a})$  because of the monotonicity constraint, therefore:

$$\sum_{u_t=1}^{|U_t|} \mu_{\tilde{s}, \tilde{a}, u_t, s_{t+1}} \cdot \mu_{s_t, a_t, u_t, s_{t+1}} \cdot \theta_{u_t} \geq P(s_{t+1} | \tilde{s}, \tilde{a}) \cdot P(s_{t+1} | s_t, a_t) \quad (25)$$

Since  $\forall s, s' \in \mathcal{S}, \forall a \in \mathcal{A}, \forall u_t \in U_t, \mu_{s, a, u_t, s'} = 1 \iff f(s, a, u_t) = s'$  by definition of  $\mu$ , Eq. (24) and Eq. (25) are equivalent to:

$$\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} \leq \min(P(\tilde{s}' | \tilde{s}, \tilde{a}), P(s_{t+1} | s_t, a_t)) \quad (26)$$

and

$$\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} \geq P(s_{t+1} | \tilde{s}, \tilde{a}) \cdot P(s_{t+1} | s_t, a_t) \quad (27)$$

respectively. As these notations are equivalent, we will use this second notation for brevity and clarity. Consider the following disjoint cases:

- $P(s_{t+1} | s_t, a_t) - (P(s_{t+1} | s_t, a_t) \cdot P(s_{t+1} | \tilde{s}, \tilde{a})) < P(\tilde{s}' | \tilde{s}, \tilde{a})$
- $P(s_{t+1} | s_t, a_t) - (P(s_{t+1} | s_t, a_t) \cdot P(s_{t+1} | \tilde{s}, \tilde{a})) \geq P(\tilde{s}' | \tilde{s}, \tilde{a})$

**Case 4(a):**  $P(s_{t+1} | s_t, a_t) - (P(s_{t+1} | s_t, a_t) \cdot P(s_{t+1} | \tilde{s}, \tilde{a})) < P(\tilde{s}' | \tilde{s}, \tilde{a})$

**Proof B.18** We can assign  $\theta$  as follows:

$$\left\{ \begin{array}{l} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} = P(s_{t+1} | s_t, a_t) \cdot P(s_{t+1} | \tilde{s}, \tilde{a}) \\ \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = P(s_{t+1} | s_t, a_t) - (P(s_{t+1} | s_t, a_t) \cdot P(s_{t+1} | \tilde{s}, \tilde{a})) \\ \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = 0 \\ \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} = P(s_{t+1} | \tilde{s}, \tilde{a}) - \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} \\ \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = P(\tilde{s}' | \tilde{s}, \tilde{a}) - \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} \\ \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} P(\tilde{s}' | \tilde{s}, \tilde{a}) \end{array} \right.$$

This assignment of  $\theta$  satisfies the constraints of the linear optimisation problem, as follows:

- $\sum_{f(s_t, a_t, u_t) = s_{t+1}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t}$   
 $= P(s_{t+1} | s_t, a_t)$ , satisfying (7).
- $\sum_{f(s_t, a_t, u_t) \neq s_{t+1}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t}$   
 $= 1 - P(s_{t+1} | s_t, a_t)$ , satisfying (7).
- $\sum_{f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = P(\tilde{s}' | \tilde{s}, \tilde{a})$ ,  
satisfying (7).
- $\sum_{f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} = P(s_{t+1} | \tilde{s}, \tilde{a})$ ,  
satisfying (7).
- $\sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t}$   
 $= 1 - P(\tilde{s}' | \tilde{s}, \tilde{a}) = \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} P(s' | \tilde{s}, \tilde{a})$ , therefore it is possible to assign  $\theta$  such that  $\forall s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}, \sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = P(s' | \tilde{s}, \tilde{a})$ , satisfying (7).
- (27) holds, therefore Mon1 (8) is satisfied.
- Because  $P(\tilde{s}' | s_t, a_t) = 0$ , Mon2 (9) holds for the transition  $\tilde{s}, \tilde{a} \rightarrow \tilde{s}'$ . Also, because  $\sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = 0$ , the counterfactual probability of all other transitions  $\tilde{s}, \tilde{a} \rightarrow s'$  where  $s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}$  will be  $\tilde{P}_t(s' | \tilde{s}, \tilde{a}) = 0$ . Therefore, Mon2 (9) is satisfied.
- CS (10) is vacuously satisfied for the transitions  $\tilde{s}, \tilde{a} \rightarrow \tilde{s}'$  and  $\tilde{s}, \tilde{a} \rightarrow s_{t+1}$  because  $P(\tilde{s}' | s_t, a_t) = 0$  and  $\frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} = \frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)}$ . Also, because  $\sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = 0$ , the counterfactual probability of all other transitions  $\tilde{s}, \tilde{a} \rightarrow s'$  where  $s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}$  will be  $\tilde{P}_t(s' | \tilde{s}, \tilde{a}) = 0$ , which satisfies CS (10) for all other transitions.
- $0 \leq \theta_{u_t} \leq 1, \forall u_t$ , satisfying (11).

$$\begin{aligned}
& \bullet \sum_{u_t=1}^{|U_t|} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} \\
& + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = 1, \text{ satisfying (12).}
\end{aligned}$$

All the constraints are satisfied, so this is a valid assignment of  $\theta$  for this state-action pair.

**Case 4(b):**  $P(s_{t+1} | s_t, a_t) - (P(s_{t+1} | s_t, a_t) \cdot P(s_{t+1} | \tilde{s}, \tilde{a})) \geq P(\tilde{s}' | \tilde{s}, \tilde{a})$

**Proof B.19** We can assign  $\theta$  as follows:

$$\left\{ \begin{array}{l}
P(s_{t+1} | s_t, a_t) \cdot P(s_{t+1} | \tilde{s}, \tilde{a}) \leq \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} \leq P(s_{t+1} | \tilde{s}, \tilde{a}) \\
\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = P(\tilde{s}' | \tilde{s}, \tilde{a}) \\
0 \leq \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} \leq \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} P(s' | \tilde{s}, \tilde{a}) \cdot P(s_{t+1} | s_t, a_t) \\
\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} = P(s_{t+1} | \tilde{s}, \tilde{a}) - \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} \\
\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = 0 \\
\sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} P(\tilde{s}' | \tilde{s}, \tilde{a}) - \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t}
\end{array} \right.$$

$$\begin{aligned}
& \text{s.t. } \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} \\
& = P(s_{t+1} | s_t, a_t).
\end{aligned}$$

We do not know exactly how much probability will be assigned to  $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t}$  and  $\sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t}$ . However, if  $\theta$  satisfies the above assignment then  $\theta$  will always satisfy the constraints of the optimisation problem, as follows:

$$\begin{aligned}
& \bullet \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1}}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} \\
& = P(s_{t+1} | s_t, a_t), \text{ satisfying (7).} \\
& \bullet \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1}}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} \\
& = 1 - P(s_{t+1} | s_t, a_t), \text{ satisfying (7).} \\
& \bullet \sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = P(\tilde{s}' | \tilde{s}, \tilde{a}), \\
& \text{satisfying (7).} \\
& \bullet \sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} = P(s_{t+1} | \tilde{s}, \tilde{a}), \\
& \text{satisfying (7).}
\end{aligned}$$



- $\sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t}$   
 $= 1 - P(\tilde{s}' \mid \tilde{s}, \tilde{a}) = \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} P(s' \mid \tilde{s}, \tilde{a})$ , therefore it is possible to assign  $\theta$  such that  $\forall s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}, \sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = P(s' \mid \tilde{s}, \tilde{a})$ , satisfying (7).

- (27) holds, therefore Mon1 (8) is satisfied.

- Because  $P(\tilde{s}' \mid s_t, a_t) = 0$ , Mon2 (9) holds for  $\tilde{s}, \tilde{a} \rightarrow \tilde{s}'$ .

- CS (10) is vacuously satisfied for the transitions  $\tilde{s}, \tilde{a} \rightarrow \tilde{s}'$  and  $\tilde{s}, \tilde{a} \rightarrow s_{t+1}$  because  $P(\tilde{s}' \mid s_t, a_t) = 0$  and  $\frac{P(s_{t+1} \mid \tilde{s}, \tilde{a})}{P(s_{t+1} \mid s_t, a_t)} = \frac{P(s_{t+1} \mid \tilde{s}, \tilde{a})}{P(s_{t+1} \mid s_t, a_t)}$ .

- To ensure that Mon2 (9) and CS (10) hold for all other transitions, we must consider two possible cases:

- If  $P(s_{t+1} \mid \tilde{s}, \tilde{a}) \geq P(s_{t+1} \mid s_t, a_t)$ , then we can assign  $\theta$  such that:

$$\sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = 0$$

and

$$\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} = P(s_{t+1} \mid s_t, a_t) - \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t}$$

which must be possible in our given assignment of  $\theta$  because, from the condition of Case 4(b), we have:

$$P(s_{t+1} \mid s_t, a_t) - (P(s_{t+1} \mid s_t, a_t) \cdot P(s_{t+1} \mid \tilde{s}, \tilde{a})) \geq P(\tilde{s}' \mid \tilde{s}, \tilde{a})$$

$$\implies P(s_{t+1} \mid s_t, a_t) - P(\tilde{s}' \mid \tilde{s}, \tilde{a}) \geq P(s_{t+1} \mid s_t, a_t) \cdot P(s_{t+1} \mid \tilde{s}, \tilde{a})$$

Because  $\sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = 0, \forall s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}, \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = 0$ , and so

$\forall s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}, \tilde{P}_t(s' \mid \tilde{s}, \tilde{a}) = 0$ . This satisfies the Mon2 (9) (since  $0 \leq \tilde{P}_t(s' \mid \tilde{s}, \tilde{a})$ ) and CS (10) constraints for all other transitions  $\tilde{s}, \tilde{a} \rightarrow s'$ .

- Otherwise, we must have  $P(s_{t+1} \mid \tilde{s}, \tilde{a}) < P(s_{t+1} \mid s_t, a_t)$ . We need to prove that there exists an assignment of  $\theta$  where the Mon2 (9) and CS (10) constraints are satisfied for all transitions  $\tilde{s}, \tilde{a} \rightarrow s', \forall s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}$ , which equates to proving:

$$\max_{\theta} \left( \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} \right) \geq \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} P(s' \mid \tilde{s}, \tilde{a}) \cdot P(s_{t+1} \mid s_t, a_t)$$

since

$$\sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} \leq \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} P(s' \mid \tilde{s}, \tilde{a}) \cdot P(s_{t+1} \mid s_t, a_t)$$

in our given assignment of  $\theta$ . From Lemma B.4, we have:

$$\max_{\theta} \left( \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} \right) \geq P(s_{t+1} \mid s_t, a_t) - P(s_{t+1} \mid \tilde{s}, \tilde{a}) \quad (28)$$

under the Mon2 (9) and CS (10) constraints. Since we have assigned the maximum possible probability to  $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = P(\tilde{s}' | \tilde{s}, \tilde{a})$  under the Mon2 (9) and CS (10) constraints in our given assignment of  $\theta$ , (28) implies:

$$\max_{\theta} \left( \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} \right) \geq P(s_{t+1} | s_t, a_t) - P(s_{t+1} | \tilde{s}, \tilde{a}) - P(\tilde{s}' | \tilde{s}, \tilde{a})$$

Therefore, we know that we can assign  $\theta$  such that  $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = P(s_{t+1} | s_t, a_t)$ , and all transitions satisfy the Mon2 (9) and CS (10) constraints.

•  $0 \leq \theta_{u_t} \leq 1, \forall u_t$ , satisfying (11).

•  $\sum_{u_t=1}^{|U_t|} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = 1$ , satisfying (12).

All the constraints are satisfied, so this is a valid assignment of  $\theta$  for this state-action pair.

Therefore, these two cases can be simplified to:

$$\begin{aligned} \tilde{P}_t^{UB}(\tilde{s}' | \tilde{s}, \tilde{a}) &= \frac{\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t}}{P(s_{t+1} | s_t, a_t)} \\ &= \frac{\min(P(s_{t+1} | s_t, a_t) - (P(s_{t+1} | s_t, a_t) \cdot P(s_{t+1} | \tilde{s}, \tilde{a})), P(\tilde{s}' | \tilde{s}, \tilde{a}))}{P(s_{t+1} | s_t, a_t)} \\ &= \min \left( 1 - P(s_{t+1} | \tilde{s}, \tilde{a}), \frac{P(\tilde{s}' | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} \right) \end{aligned}$$

We have proven all the cases for Theorem B.5, therefore Theorem B.5 holds.

**Theorem B.6** For outgoing transitions from state-action pairs  $(\tilde{s}, \tilde{a})$  which have overlapping support with the observed  $(s_t, a_t)$ , but  $(\tilde{s}, \tilde{a}) \neq (s_t, a_t)$ , the linear program will produce a lower bound of:

$$\tilde{P}_t^{LB}(\tilde{s}' | \tilde{s}, \tilde{a}) = \begin{cases} \max(P(\tilde{s}' | \tilde{s}, \tilde{a}), 1 - \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \tilde{P}_t^{UB}(s' | \tilde{s}, \tilde{a})) & \text{if } \tilde{s}' = s_{t+1} \\ 0 & \text{if } P(\tilde{s}' | s_t, a_t) > 0 \\ & \text{and } \frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} > \frac{P(\tilde{s}' | \tilde{s}, \tilde{a})}{P(\tilde{s}' | s_t, a_t)} \\ \max(0, 1 - \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} \tilde{P}_t^{UB}(s' | \tilde{s}, \tilde{a})) & \text{otherwise} \end{cases}$$

**Proof B.20** Take an arbitrary transition  $\tilde{s}, \tilde{a} \rightarrow \tilde{s}'$  where  $(\tilde{s}, \tilde{a})$  has overlapping support with the observed state-action pair  $(s_t, a_t)$ . Consider the following disjoint cases:

- $\tilde{s}' = s_{t+1}$
- $\tilde{s}' \neq s_{t+1}$ ,  $\frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} > \frac{P(\tilde{s}' | \tilde{s}, \tilde{a})}{P(\tilde{s}' | s_t, a_t)}$  and  $P(\tilde{s}' | s_t, a_t) > 0$
- $\tilde{s}' \neq s_{t+1}$ , and  $\frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} \leq \frac{P(\tilde{s}' | \tilde{s}, \tilde{a})}{P(\tilde{s}' | s_t, a_t)}$  or  $P(\tilde{s}' | s_t, a_t) = 0$

These cases are disjoint and cover all possible situations.

**Case 1:**  $\tilde{s}' = s_{t+1}$

**Proof B.21** To satisfy Mon1 (8),  $\tilde{P}_t^{LB}(s_{t+1} | \tilde{s}, \tilde{a}) \geq P(s_{t+1} | \tilde{s}, \tilde{a})$ . To do this, we must assign  $\theta$  such that:

$$\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} \geq P(s_{t+1} | s_t, a_t) \cdot P(s_{t+1} | \tilde{s}, \tilde{a}) \quad (29)$$

Consider the following disjoint cases:

1.  $P(s_{t+1} | \tilde{s}, \tilde{a}) \geq 1 - \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \tilde{P}_t^{UB}(s' | \tilde{s}, \tilde{a})$
2.  $P(s_{t+1} | \tilde{s}, \tilde{a}) < 1 - \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \tilde{P}_t^{UB}(s' | \tilde{s}, \tilde{a})$

**Case 1(a):**  $P(s_{t+1} | \tilde{s}, \tilde{a}) \geq 1 - \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \tilde{P}_t^{UB}(s' | \tilde{s}, \tilde{a})$

**Proof B.22** In this case,  $P(s_{t+1} | \tilde{s}, \tilde{a}) \geq 1 - \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \tilde{P}_t^{UB}(s' | \tilde{s}, \tilde{a})$ , therefore:

$$P(s_{t+1} | s_t, a_t) \cdot P(s_{t+1} | \tilde{s}, \tilde{a}) \geq P(s_{t+1} | s_t, a_t) - P(s_{t+1} | s_t, a_t) \cdot \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \tilde{P}_t^{UB}(s' | \tilde{s}, \tilde{a}) \quad (30)$$

Therefore, we can assign  $\theta$  as follows:

$$\left\{ \begin{array}{l} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} = P(s_{t+1} | s_t, a_t) \cdot P(s_{t+1} | \tilde{s}, \tilde{a}) \\ \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = P(s_{t+1} | s_t, a_t) - \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} \\ \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} = P(s_{t+1} | \tilde{s}, \tilde{a}) - \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} \\ \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} P(\tilde{s}' | \tilde{s}, \tilde{a}) - \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} \end{array} \right.$$

This assignment of  $\theta$  satisfies the constraints of the linear optimisation problem, as follows:

- $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1}}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t}$   
 $= P(s_{t+1} \mid s_t, a_t)$ , satisfying (7).
- $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1}}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t}$   
 $= 1 - P(s_{t+1} \mid s_t, a_t)$ , satisfying (7).
- $\sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} = P(s_{t+1} \mid \tilde{s}, \tilde{a})$ ,  
satisfying (7).
- $\sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t}$   
 $= 1 - P(\tilde{s}' \mid \tilde{s}, \tilde{a}) = \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} P(s' \mid \tilde{s}, \tilde{a})$ , therefore it is possible to assign  $\theta$  such that  $\forall s' \in \mathcal{S} \setminus \{s_{t+1}\}$ ,  $\sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = P(s' \mid \tilde{s}, \tilde{a})$ , satisfying (7).
- Eq. (29) holds, so Mon1 (8) is satisfied.
- CS (10) always holds vacuously for the transition  $\tilde{s}, \tilde{a} \rightarrow s_{t+1}$ , because trivially  $\frac{P(s_{t+1} \mid \tilde{s}, \tilde{a})}{P(s_{t+1} \mid s_t, a_t)} = \frac{P(s_{t+1} \mid \tilde{s}, \tilde{a})}{P(s_{t+1} \mid s_t, a_t)}$ .
- From (30), we get:

$$P(s_{t+1} \mid s_t, a_t) - P(s_{t+1} \mid s_t, a_t) \cdot P(\tilde{s}' \mid \tilde{s}, \tilde{a}) < \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \tilde{P}_t^{UB}(s' \mid \tilde{s}, \tilde{a}) \cdot P(s_{t+1} \mid s_t, a_t)$$

This means that  $\theta$  can be assigned such that  $\forall s' \in \mathcal{S} \setminus \{s_{t+1}\}$ , the counterfactual probability of each transition  $\tilde{s}, \tilde{a} \rightarrow s'$  will be  $\tilde{P}_t(s' \mid \tilde{s}, \tilde{a}) \leq \tilde{P}_t^{UB}(s' \mid \tilde{s}, \tilde{a})$ . We have already proven that each  $\tilde{P}_t^{UB}(s' \mid \tilde{s}, \tilde{a})$  satisfies the (9) and CS (10) constraints, and because  $\tilde{P}_t(s' \mid \tilde{s}, \tilde{a}) \leq \tilde{P}_t^{UB}(s' \mid \tilde{s}, \tilde{a})$ ,  $\tilde{P}_t(s' \mid \tilde{s}, \tilde{a})$  will as well.

- $0 \leq \theta_{u_t} \leq 1, \forall u_t$ , satisfying (11).
- $\sum_{u_t=1}^{|U_t|} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t}$   
 $+ \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = 1$ , satisfying (12).

All the constraints are satisfied, so this is a valid assignment of  $\theta$  for this state-action pair.

**Case I(b):**  $P(s_{t+1} \mid \tilde{s}, \tilde{a}) < 1 - \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \tilde{P}_t^{UB}(s' \mid \tilde{s}, \tilde{a})$

**Proof B.23** In this case  $P(s_{t+1} \mid \tilde{s}, \tilde{a}) < 1 - \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \tilde{P}_t^{UB}(s' \mid \tilde{s}, \tilde{a})$ , therefore:

$$P(s_{t+1} \mid s_t, a_t) \cdot P(s_{t+1} \mid \tilde{s}, \tilde{a}) < P(s_{t+1} \mid s_t, a_t) \cdot \left(1 - \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \tilde{P}_t^{UB}(s' \mid \tilde{s}, \tilde{a})\right) \quad (31)$$

Therefore, we can assign  $\theta$  as follows:

$$\left\{ \begin{array}{l} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} = P(s_{t+1} \mid s_t, a_t) \cdot \left(1 - \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \tilde{P}_t^{UB}(s' \mid \tilde{s}, \tilde{a})\right) \\ \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = P(s_{t+1} \mid s_t, a_t) \cdot \left(\sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \tilde{P}_t^{UB}(s' \mid \tilde{s}, \tilde{a})\right) \\ \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} = P(s_{t+1} \mid \tilde{s}, \tilde{a}) - \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} \\ \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} P(s' \mid \tilde{s}, \tilde{a}) - \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} \end{array} \right.$$

This assignment of  $\theta$  satisfies the constraints of the linear optimisation problem, as follows:

- $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1}}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t}$   
 $= P(s_{t+1} | s_t, a_t)$ , satisfying (7).
- $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1}}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t}$   
 $= 1 - P(s_{t+1} | s_t, a_t)$ , satisfying (7).
- $\sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} = P(s_{t+1} | \tilde{s}, \tilde{a})$ ,  
satisfying (7).
- $\sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t}$   
 $= 1 - P(\tilde{s}' | \tilde{s}, \tilde{a}) = \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} P(s' | \tilde{s}, \tilde{a})$ , therefore it is possible to assign  $\theta$  such that  $\forall s' \in \mathcal{S} \setminus \{s_{t+1}\}, \sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = P(s' | \tilde{s}, \tilde{a})$ , satisfying (7).
- From (31), we get:

$$\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} = P(s_{t+1} | s_t, a_t) \cdot \left(1 - \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \tilde{P}_t^{UB}(s' | \tilde{s}, \tilde{a})\right) > P(s_{t+1} | s_t, a_t) \cdot P(s_{t+1} | \tilde{s}, \tilde{a})$$

Therefore, (29) holds, so Mon1 (8) is satisfied.

- CS (10) always holds vacuously for the transition  $\tilde{s}, \tilde{a} \rightarrow s_{t+1}$ , because trivially  $\frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} = \frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)}$ .
- Because  $\sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = P(s_{t+1} | s_t, a_t) \cdot (\sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \tilde{P}_t^{UB}(s' | \tilde{s}, \tilde{a}))$ , this means that  $\theta$  can be assigned such that  $\forall s' \in \mathcal{S} \setminus \{s_{t+1}\}$ , its counterfactual probability  $\tilde{P}_t(s' | \tilde{s}, \tilde{a}) = \tilde{P}_t^{UB}(s' | \tilde{s}, \tilde{a})$ , which we have already proven to satisfy Mon2 (9) and CS (10).
- $0 \leq \theta_{u_t} \leq 1, \forall u_t$ , satisfying (11).
- $\sum_{u_t=1}^{|U_t|} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t}$   
 $+ \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = 1$ , satisfying (12).

All the constraints are satisfied, so this is a valid assignment of  $\theta$  for this state-action pair.

Therefore, these two cases can be simplified as:

$$\begin{aligned} \tilde{P}_t^{LB}(\tilde{s}' | \tilde{s}, \tilde{a}) &= \frac{\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t}}{P(s_{t+1} | s_t, a_t)} \\ &= \frac{\max\left(P(s_{t+1} | s_t, a_t) \cdot P(s_{t+1} | \tilde{s}, \tilde{a}), P(s_{t+1} | s_t, a_t) \cdot \left(1 - \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \tilde{P}_t^{UB}(s' | \tilde{s}, \tilde{a})\right)\right)}{P(s_{t+1} | s_t, a_t)} \\ &= \max\left(P(\tilde{s}' | \tilde{s}, \tilde{a}), 1 - \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \tilde{P}_t^{UB}(s' | \tilde{s}, \tilde{a})\right) \end{aligned}$$

**Case 2:**  $\tilde{s}' \neq s_{t+1}$ ,  $\frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} > \frac{P(\tilde{s}' | \tilde{s}, \tilde{a})}{P(\tilde{s}' | s_t, a_t)}$  and  $P(\tilde{s}' | s_t, a_t) > 0$

**Proof B.24** Because  $\frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} > \frac{P(\tilde{s}' | \tilde{s}, \tilde{a})}{P(\tilde{s}' | s_t, a_t)}$  and  $P(\tilde{s}' | s_t, a_t) > 0$ , to satisfy CS (10)  $\tilde{P}_t^{LB}(\tilde{s}' | \tilde{s}, \tilde{a}) = 0$ . We have already proven in Case 2 of the proof for Theorem B.5 that there exists an assignment of  $\theta$  that satisfies all the constraints, where  $\tilde{P}_t(\tilde{s}' | \tilde{s}, \tilde{a}) = 0$ . Therefore, the same assignment of  $\theta$  will satisfy the constraints of the optimisation problem, and result in  $\tilde{P}_t^{LB}(\tilde{s}' | \tilde{s}, \tilde{a}) = 0$ .

**Case 3:**  $\tilde{s}' \neq s_{t+1}$ , and  $\frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} \leq \frac{P(\tilde{s}' | \tilde{s}, \tilde{a})}{P(\tilde{s}' | s_t, a_t)}$  or  $P(\tilde{s}' | s_t, a_t) = 0$

**Proof B.25** Consider the following disjoint cases:

- $1 - \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} \tilde{P}_t^{UB}(s' | \tilde{s}, \tilde{a}) \geq 0$
- $1 - \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} \tilde{P}_t^{UB}(s' | \tilde{s}, \tilde{a}) < 0$

**Case 3(a):**  $1 - \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} \tilde{P}_t^{UB}(s' | \tilde{s}, \tilde{a}) \geq 0$

**Proof B.26** We can assign  $\theta$  as follows:

$$\left\{ \begin{array}{l} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = P(s_{t+1} | s_t, a_t) \cdot (1 - \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} \tilde{P}_t^{UB}(s' | \tilde{s}, \tilde{a})) \\ \forall s' \in \mathcal{S} \setminus \{\tilde{s}'\}, \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = P(s_{t+1} | s_t, a_t) \cdot \tilde{P}_t^{UB}(s' | \tilde{s}, \tilde{a}) \\ \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = P(\tilde{s}' | \tilde{s}, \tilde{a}) - \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} \\ \forall s' \in \mathcal{S} \setminus \{\tilde{s}'\}, \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = P(s' | \tilde{s}, \tilde{a}) - \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} \end{array} \right.$$

This assignment of  $\theta$  satisfies the constraints of the linear optimisation problem, as follows:

- $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1}}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t}$   
 $= P(s_{t+1} | s_t, a_t)$ , satisfying (7).
- $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1}}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t}$   
 $= 1 - P(s_{t+1} | s_t, a_t)$ , satisfying (7).
- $\sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = P(\tilde{s}' | \tilde{s}, \tilde{a})$ ,  
satisfying (7).
- $\sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} \sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t}$   
 $= 1 - P(\tilde{s}' | \tilde{s}, \tilde{a}) = \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} P(s' | \tilde{s}, \tilde{a})$ , therefore it is possible to assign  $\theta$  such that  $\forall s' \in \mathcal{S} \setminus \{\tilde{s}'\}, \sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = P(s' | \tilde{s}, \tilde{a})$ , satisfying (7).
- $\forall s' \in \mathcal{S} \setminus \{\tilde{s}'\}, \tilde{P}_t(s' | \tilde{s}, \tilde{a}) = \frac{\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t}}{P(s_{t+1} | s_t, a_t)} = \frac{P(s_{t+1} | s_t, a_t) \cdot \tilde{P}_t^{UB}(s' | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} = \tilde{P}_t^{UB}(s' | \tilde{s}, \tilde{a})$ , which we have already shown in the proof for Theorem B.5 satisfies Mon1 (8), Mon2 (9) and CS (10).

- Mon2 (9) is also satisfied for the transition  $\tilde{s}, \tilde{a} \rightarrow \tilde{s}'$ , as follows:

– If  $P(\tilde{s}' | s_t, a_t) = 0$ , then this assignment of  $\theta$  vacuously satisfies Mon2 (9).

– Otherwise, we have already proven that there exists an assignment of  $\theta$  that satisfies all the constraints (as proven in Theorem B.5). If  $1 - \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} \tilde{P}_t^{UB}(\tilde{s}' | \tilde{s}, \tilde{a}) > 0$ , then the minimum possible counterfactual transition probability of  $\tilde{s}, \tilde{a} \rightarrow \tilde{s}'$  is  $\tilde{P}_t(\tilde{s}' | \tilde{s}, \tilde{a}) = 1 - \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} \tilde{P}_t^{UB}(\tilde{s}' | \tilde{s}, \tilde{a})$  (because the upper bounds must sum to 1). Therefore,  $1 - \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} \tilde{P}_t^{UB}(\tilde{s}' | \tilde{s}, \tilde{a}) \leq P(\tilde{s}' | \tilde{s}, \tilde{a})$ , or there would be no valid assignment of  $\theta$  that satisfies Mon2 (9).

- CS (10) will be vacuously satisfied for the transition  $\tilde{s}, \tilde{a} \rightarrow \tilde{s}'$ , because  $\frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} \leq \frac{P(\tilde{s}' | \tilde{s}, \tilde{a})}{P(\tilde{s}' | s_t, a_t)}$  or  $P(\tilde{s}' | s_t, a_t) = 0$ .

- $0 \leq \theta_{u_t} \leq 1, \forall u_t$ , satisfying (11).

- $$\sum_{u_t=1}^{|U_t|} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t}$$

$$+ \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = 1, \text{ satisfying (12).}$$

All the constraints are satisfied, so this is a valid assignment of  $\theta$  for this state-action pair.

**Case 3(b):**  $1 - \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} \tilde{P}_t^{UB}(s' | \tilde{s}, \tilde{a}) < 0$

**Proof B.27** Because  $1 - \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} \tilde{P}_t^{UB}(s' | \tilde{s}, \tilde{a}) < 0$ , we have:

$$P(s_{t+1} | s_t, a_t) \cdot \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} \tilde{P}_t^{UB}(s' | \tilde{s}, \tilde{a}) > P(s_{t+1} | s_t, a_t)$$

Therefore, we can assign  $\theta$  as follows:

$$\left\{ \begin{array}{l} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = 0 \\ \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = P(s_{t+1} | s_t, a_t) \\ \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = P(\tilde{s}' | \tilde{s}, \tilde{a}) \\ \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} P(s' | \tilde{s}, \tilde{a}) - \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} \end{array} \right.$$

This assignment of  $\theta$  satisfies the constraints of the linear optimisation problem, as follows:

- $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1}}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t}$   
 $= P(s_{t+1} | s_t, a_t)$ , satisfying (7).
- $\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1}}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t}$   
 $= 1 - P(s_{t+1} | s_t, a_t)$ , satisfying (7).
- $\sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t} = P(\tilde{s}' | \tilde{s}, \tilde{a})$ ,  
satisfying (7).
- $\sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} \sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t}$   
 $= 1 - P(\tilde{s}' | \tilde{s}, \tilde{a}) = \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} P(s' | \tilde{s}, \tilde{a})$ , therefore it is possible to assign  $\theta$  such that  $\forall s' \in \mathcal{S} \setminus \{\tilde{s}'\}, \sum_{\substack{u_t \in U_t \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = P(s' | \tilde{s}, \tilde{a})$ , satisfying (7).
- From this assignment of  $\theta$ ,  $\tilde{P}_t(\tilde{s}' | \tilde{s}, \tilde{a}) = \frac{\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t}}{P(s_{t+1} | s_t, a_t)} = \frac{0}{P(s_{t+1} | s_t, a_t)} = 0 \leq P(\tilde{s}' | \tilde{s}, \tilde{a})$ , which immediately satisfies the Mon2 (9) and CS (10) constraints for the transition  $\tilde{s}, \tilde{a} \rightarrow \tilde{s}'$ .
- Because  $1 - \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} \tilde{P}_t^{UB}(s' | \tilde{s}, \tilde{a}) < 0$ :

$$P(s_{t+1} | s_t, a_t) < \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} \tilde{P}_t^{UB}(s' | \tilde{s}, \tilde{a}) \cdot P(s_{t+1} | s_t, a_t)$$

This guarantees that we can assign  $\theta$  such that the Mon1 (8), Mon2 (9) and CS (10) constraints are satisfied for all transitions  $\tilde{s}', \tilde{a}' \rightarrow s', \forall s' \in \mathcal{S} \setminus \{\tilde{s}'\}$ . In particular, we should assign  $\theta$  such that  $\tilde{P}_t(s_{t+1} | \tilde{s}, \tilde{a}) = \tilde{P}_t^{UB}(s_{t+1} | \tilde{s}, \tilde{a})$  (Theorem B.5 proves this  $\theta$  exists), as we have proven that Mon1 (8) and CS (10) constraints are satisfied for  $\tilde{P}_t^{UB}(s_{t+1} | \tilde{s}, \tilde{a})$ . Then, we can assign  $\theta$  such that  $\forall s' \in \mathcal{S} \setminus \{\tilde{s}', s_{t+1}\}, \tilde{P}_t(s' | \tilde{s}, \tilde{a}) \leq \tilde{P}_t^{UB}(s' | \tilde{s}, \tilde{a})$ , and because each  $\tilde{P}_t^{UB}(s' | \tilde{s}, \tilde{a})$  has been proven to satisfy the Mon2 (9) and CS (10) constraints, each  $\tilde{P}_t(s' | \tilde{s}, \tilde{a})$  must as well.

- $0 \leq \theta_{u_t} \leq 1, \forall u_t$ , satisfying (11).



$$\begin{aligned}
\bullet \sum_{u_t=1}^{|U_t|} \theta_{u_t} &= \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} + \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} + \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s_{t+1}}} \theta_{u_t} \\
&+ \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) \neq s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} = 1, \text{ satisfying (12).}
\end{aligned}$$

All the constraints are satisfied, so this is a valid assignment of  $\theta$  for this state-action pair.

Therefore, these two cases can be simplified as:

$$\begin{aligned}
\tilde{P}_t^{LB}(\tilde{s}' \mid \tilde{s}, \tilde{a}) &= \frac{\sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = \tilde{s}'}} \theta_{u_t}}{P(s_{t+1} \mid s_t, a_t)} \\
&= \frac{\max\left(0, P(s_{t+1} \mid s_t, a_t) \cdot (1 - \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} \tilde{P}_t^{UB}(s' \mid \tilde{s}, \tilde{a}))\right)}{P(s_{t+1} \mid s_t, a_t)} \\
&= \max\left(0, 1 - \sum_{s' \in \mathcal{S} \setminus \{\tilde{s}'\}} \tilde{P}_t^{UB}(s' \mid \tilde{s}, \tilde{a})\right)
\end{aligned}$$

We have proven all the cases for Theorem B.6, therefore Theorem B.6 holds.

## B.2 ANY CFMDP ENTAILED BY AN INTERVAL CFMDP IS A VALID CFMDP

In our experiments, we evaluate the policies by sampling example CFMDPs from the interval CFMDPs. But, even if  $\theta$ s exist that would produce each sampled CF transition probability separately, this does not necessarily guarantee that there exists a single  $\theta$  that produces all of the CF transition probabilities in the CFMDP. We can prove that, given any CFMDP  $\tilde{\mathcal{M}}_t$ , there exists a  $\theta$  that produces all the counterfactual probabilities in  $\tilde{\mathcal{M}}_t$ , as follows.

**Theorem B.7** *Given any arbitrary interval counterfactual MDP, any counterfactual MDP  $\tilde{\mathcal{M}}_t$  entailed by this interval counterfactual MDP will be a valid counterfactual MDP.*

**Proof B.28** *Let us assume that we have an interval CFMDP, and an arbitrary CFMDP  $\tilde{\mathcal{M}}_t$  entailed by this interval CFMDP for each time-step  $t$  from this interval CFMDP. First, we need to prove that, for every transition  $s, a \rightarrow s'$  in  $\tilde{\mathcal{M}}_t$ , there exists a  $\theta$  that produces its counterfactual probability  $\tilde{P}_t(s' | s, a)$ . Next, we need to show that, for all state-action pairs  $(s, a)$ , there exists a  $\theta$  that satisfies the constraints of the optimisation problem (7)-(12) and produces the entailed CF probability distribution  $\tilde{P}_t(\cdot | s, a)$ . Finally, similarly to the proof in Section B.1, we can prove by induction that these assignments of  $\theta$  for each state-action pair can be combined to form a single  $\theta$  that produces  $\tilde{\mathcal{M}}_t$ .*

### B.2.1 Existence of $\theta$ for Each $\tilde{P}_t(s' | s, a)$

The counterfactual transition probability bounds are produced by a convex optimisation problem (or using the analytical bounds, which we have proven produce the same exact bounds as the optimisation problem). Because the optimisation problem is convex, the feasible set for each of the counterfactual transition probabilities is also convex (i.e., there are no gaps in the probability bounds). This means there is guaranteed to be at least one  $\theta$  that can produce each of the entailed counterfactual transition probabilities in  $\tilde{\mathcal{M}}_t$ .

### B.2.2 Existence of $\theta$ for Each $\tilde{P}_t(\cdot | s, a)$

Next, we need to show that, for all state-action pairs  $(s, a)$ , there exists a  $\theta$  that satisfies the constraints of the optimisation problem (7)-(12) and can produce the entailed CF probability distribution  $\tilde{P}_t(\cdot | s, a)$ . Take any state-action pair  $(s, a)$  in  $\tilde{\mathcal{M}}_t$  arbitrarily. Each CF transition probability  $\tilde{P}_t(\cdot | s, a)$  arises from the probability over the mechanisms that produce each transition  $s, a \rightarrow s'$  and observed transition  $s_t, a_t \rightarrow s_{t+1}$  vs. the total probability assigned to  $P(s_{t+1} | s_t, a_t)$ . Therefore, we need to show that the total probability assigned across all mechanisms containing each transition and the observed transition sums to  $P(s_{t+1} | s_t, a_t)$ . Because  $\tilde{P}_t(\cdot | s, a)$  is a valid probability distribution, we know

$$\sum_{s' \in \mathcal{S}} \tilde{P}_t(s' | s, a) = 1$$

This means

$$\sum_{s' \in \mathcal{S}} \frac{\sum_{u_t=1}^{|U_t|} \mu_{s,a,u_t,s'} \cdot \mu_{s_t,a_t,u_t,s_{t+1}} \cdot \theta_{u_t}}{P(s_{t+1} | s_t, a_t)} = 1$$

and

$$\sum_{s' \in \mathcal{S}} \sum_{u_t=1}^{|U_t|} \mu_{s,a,u_t,s'} \cdot \mu_{s_t,a_t,u_t,s_{t+1}} \cdot \theta_{u_t} = P(s_{t+1} | s_t, a_t)$$

Because  $\mu_{s,a,u_t,s'} = 1$  for exactly one next state,  $s'$ , and 0 for the remaining states, we must have

$$\sum_{u_t=1}^{|U_t|} \mu_{s_t,a_t,u_t,s_{t+1}} \cdot \theta_{u_t} = P(s_{t+1} | s_t, a_t)$$

so the total probability assigned across all mechanisms containing each transition and the observed transition sums to  $P(s_{t+1} | s_t, a_t)$ .

Next, we need to show that there exists an assignment of  $\theta$  which satisfies the interventional probability constraints (7) of the optimisation problem. We know that the interventional probabilities of all transitions from  $(s, a)$  must sum to 1:

$$\sum_{s' \in \mathcal{S}} P(s' | s, a) = 1$$

Because  $0 \leq \theta_{u_t} \leq 1, \forall u_t$  (11) and  $\sum_{u_t=1}^{|U_t|} \theta_{u_t} = 1$  (12) it must be possible to assign  $\theta$  such that

$$\sum_{u_t=1}^{|U_t|} \mu_{s,a,u_t,s'} \cdot \theta_{u_t} = P(s' | s, a)$$

and

$$\forall s' \in \mathcal{S}, \sum_{u_t=1}^{|U_t|} \mu_{s,a,u_t,s'} \cdot \mu_{s_t,a_t,u_t,s_{t+1}} \cdot \theta_{u_t} = \tilde{P}_t(s' | s, a) \cdot P(s_{t+1} | s_t, a_t)$$

and

$$\forall s' \in \mathcal{S}, \sum_{\tilde{s}' \neq s_{t+1}} \sum_{u_t=1}^{|U_t|} \mu_{s,a,u_t,s'} \cdot \mu_{s_t,a_t,u_t,\tilde{s}'} \cdot \theta_{u_t} = P(s' | s, a) - \tilde{P}_t(s' | s, a) \cdot P(s_{t+1} | s_t, a_t)$$

Therefore, this assignment satisfies

$$\sum_{u_t=1}^{|U_t|} \mu_{s,a,u_t,s'} \cdot \theta_{u_t} = P(s' | s, a)$$

$$\sum_{u_t=1}^{|U_t|} \mu_{s_t,a_t,u_t,s_{t+1}} \cdot \theta_{u_t} = P(s_{t+1} | s_t, a_t)$$

and

$$\sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{u_t=1}^{|U_t|} \mu_{s_t,a_t,u_t,s'} \cdot \theta_{u_t} = 1 - P(s_{t+1} | s_t, a_t)$$

which satisfies the interventional probability constraints (7). Finally, because each counterfactual transition probability was sampled from its bounds, we know that the entailed counterfactual transition probabilities satisfy the Mon1 (8), Mon (9) and CS (10) constraints as well.

### B.2.3 Existence of $\theta$ for $\tilde{\mathcal{M}}_t$

We have shown that there exists a  $\theta$  that meets the constraints of the optimisation problem for each state-action pair separately in the MDP. Now, we need to prove that these assignments can be combined to form a valid  $\theta$  for all state-action pairs. Let us prove inductively that adding a new state-action pair from the MDP to the causal model (and changing  $\theta$  to meet the constraints for this new state-action pair) does not affect whether the constraints of the other state-action pairs are satisfied. If this is the case, we can consider how  $\theta$  can be assigned to satisfy the constraints for each state-action pair separately, as we know these can be combined to form a valid assignment of  $\theta$  over all of the state-action pairs.

**Base Case 1** Assume  $(s_t, a_t)$  (the observed state-action pair) is the only state-action pair currently considered in the causal model (this always exists, as we always have an observed transition). We require a structural equation mechanism for each possible next state (of which there are  $|\mathcal{S}|$ ). This leads to an assignment of  $\theta^1$  as follows, which is the only possible assignment that satisfies the constraints (7)-(12):

$$\frac{\theta_{u_1}^1}{P(s_1 | s_t, a_t)} \quad \Big| \quad \frac{\theta_{u_2}^1}{P(s_2 | s_t, a_t)} \quad \Big| \quad \dots \quad \Big| \quad \frac{\theta_{u_{|\mathcal{S}|}}^1}{P(s_{|\mathcal{S}|} | s_t, a_t)}$$

$\theta^1 = [P(s_1 | s_t, a_t), P(s_2 | s_t, a_t), \dots, P(s_{|\mathcal{S}|} | s_t, a_t)]$ . Given  $\theta^1$ , the counterfactual transition probability for each transition  $s_t, a_t \rightarrow s'$  can be calculated with:

$$\tilde{P}_t(s' | s_t, a_t) = \frac{\sum_{u_t=1}^{|\mathcal{U}_t|} \mu_{s_t, a_t, u_t, s'} \cdot \mu_{s_t, a_t, u_t, s_{t+1}} \cdot \theta_{u_t}}{P(s_{t+1} | s_t, a_t)}, \forall s'$$

Therefore, given  $\theta^1$ ,  $\tilde{P}_t(s_{t+1} | s_t, a_t) = 1$ , and  $\forall s' \in \mathcal{S} \setminus \{s_{t+1}\}$ ,  $\tilde{P}_t(s' | s_t, a_t) = 0$ . This is the only possible counterfactual probability distribution for  $(s_t, a_t)$ , so this is guaranteed to be the distribution entailed from the interval CFMDP. These counterfactual probabilities also satisfy the monotonicity and counterfactual stability constraints (6-8) of the optimisation problem, as follows:


- Because  $\frac{P(s_{t+1} | s_t, a_t)}{P(s_{t+1} | s_t, a_t)} = 1$  and  $\forall s' \in \mathcal{S}$ ,  $\frac{P(s' | s_t, a_t)}{P(s' | s_t, a_t)} = 1$ ,  
 $\forall s' \in \mathcal{S}$ ,  $\frac{P(s_{t+1} | s_t, a_t)}{P(s_{t+1} | s_t, a_t)} = \frac{P(s' | s_t, a_t)}{P(s' | s_t, a_t)}$

so CS (10) is always vacuously true for all transitions from  $(s_t, a_t)$ .

- Because  $\tilde{P}_t(s_{t+1} | s_t, a_t) = 1 \geq P(s_{t+1} | s_t, a_t)$ , this satisfies Mon1 (8).
- Because  $\forall s' \in \mathcal{S} \setminus \{s_{t+1}\}$ ,  $\tilde{P}_t(s' | s_t, a_t) = 0 \leq P(s' | s_t, a_t)$ , this satisfies Mon2 (9).

**Base Case 2** Let us assume that we only have the observed state-action pair in the causal model, and assume we have a valid  $\theta^1$  with  $|\mathcal{S}|$  structural equation mechanisms.

Now, take another state-action pair  $(s, a)$  arbitrarily from the MDP to add to the causal model. We now require  $|\mathcal{S}|^2$  structural equation mechanisms, one for each possible combination of transitions from the two state-action pairs. We can view this as separating each of the existing structural equation mechanisms from Base Case 1 into  $|\mathcal{S}|$  new structural equation mechanisms, one for every possible transition from the new state-action pair,  $(s, a)$ :




where, for  $1 \leq i \leq |\mathcal{S}|, 1 \leq n \leq |\mathcal{S}|$ ,  $u_{i,n}$  leads to the same next state for  $(s_t, a_t)$  as  $u_i$ , and produces the transition  $s, a \rightarrow s_n$ .

In the same way, we can split each  $\theta_{u_1}^1, \dots, \theta_{u_{|\mathcal{S}|}}^1$  across these new structural equation mechanisms to obtain  $\theta^2$ :

By splitting  $\theta^1$  in this way, we guarantee that the total probability across each set of mechanisms (where each set produces a different transition from  $(s_t, a_t)$ ) remains the same, i.e.,

$$\forall i \in \{1, \dots, |\mathcal{S}|\}, \sum_{n=1}^{|\mathcal{S}|} \theta_{u_{i,n}}^2 = \theta_{u_i}^1$$



This means that the counterfactual probabilities of all transitions from  $(s_t, a_t)$  will be exactly the same when using  $\theta^2$  as with  $\theta^1$ . Because we assume all the constraints were satisfied when using  $\theta^1$ , this guarantees that all the constraints of the optimisation problem will continue to hold for  $(s_t, a_t)$  with  $\theta^2$ .

Now, we only need to ensure we assign  $\theta^2$  such that all the constraints hold for the transitions from  $(s, a)$ . Firstly, we need  $\sum_{i=1}^{|\mathcal{S}|} \theta_{u_{i,n}}^2 = P(s_n | s, a)$  for every possible next state  $s_n \in \mathcal{S}$ , to satisfy the interventional constraint (7). For each transition  $s, a \rightarrow s_n$ , there is a  $\theta_{u_{i,n}}^2$  for every  $u_i$ , and for every  $\theta_{u_{i,n}}^2$ ,  $0 \leq \theta_{u_{i,n}}^2 \leq \theta_{u_i}^1$ . Therefore,  $0 \leq \sum_{i=1}^{|\mathcal{S}|} \theta_{u_{i,n}}^2 \leq 1, \forall s_n \in \mathcal{S}$ . We also know that:

$$\begin{aligned} \sum_{n=1}^{|\mathcal{S}|} \sum_{i=1}^{|\mathcal{S}|} \theta_{u_{i,n}}^2 &= \sum_{i=1}^{|\mathcal{S}|} \theta_{u_i}^1 \text{ (because we split the probabilities of each } \theta_{u_i}^1 \text{)} \\ &= \sum_{i=1}^{|\mathcal{S}|} P(s_i | s_t, a_t) \\ &= 1 \end{aligned}$$

and

$$\sum_{n=1}^{|\mathcal{S}|} \sum_{i=1}^{|\mathcal{S}|} \theta_{u_{i,n}}^2 = \sum_{i=1}^{|\mathcal{S}|} P(s_i | s, a) = 1$$

Therefore, no matter the assignment of  $\theta^1$ , we can satisfy all the constraints (7)-(12) for all transitions from  $(s, a)$ . In Section B.2.2, we prove that, for any  $(s, a)$ , we can assign  $\theta^2$  such that it produces the entailed counterfactual probability distribution.

**Inductive Case** Let us assume that we now have  $k$  state-action pair values from the MDP  $\mathcal{M}$  in our causal model, and we have found a valid assignment  $\theta^k$  that satisfies all the constraints for  $k$  state-action pairs from the MDP  $\mathcal{M}$ , including  $(s_t, a_t)$ . Because there are  $k$  state-action pairs, there are  $|\mathcal{S}|^k$  possible unique structural equation mechanisms for these state-action pairs<sup>3</sup>, with probabilities as follows:


$$\theta_{u_1}^k \quad \dots \quad \theta_{u_{|S|}}^k \quad \dots \quad \theta_{u_{|S|^2}}^k \quad \dots \quad \theta_{u_{|S|^k}}^k$$

Now, we wish to add the  $k+1^{\text{th}}$  state-action pair from the MDP to the causal model, resulting in  $|\mathcal{S}|^{k+1}$  possible structural equation mechanisms. Let the  $k+1^{\text{th}}$  state-action pair be  $(s, a)$  arbitrarily. We can view this as separating the existing structural equation mechanisms each into  $|\mathcal{S}|$  new structural equation mechanisms, one for every possible transition from the  $k+1^{\text{th}}$  state-action pair:

where each  $u_{i,n}$  produces the same next states for all of the first  $k$  state-action pairs as  $u_i$ , and produces the transition  $s, a \rightarrow s_n$ .

In the same way, we can split each  $\theta_{u_1}^k, \dots, \theta_{u_{|S|}}^k, \dots$  across these new structural equation mechanisms to find  $\theta^{k+1}$ .

<sup>3</sup>Note we have changed the indexing of each  $\theta_{i,n}^k$  to  $\theta_j^k$ , where  $j = (i-1) \cdot |\mathcal{S}| + n$



By splitting  $\theta^k$  in this way, we have  $\forall i \in \{1, \dots, |\mathcal{S}|\}, \sum_{n=1}^{|\mathcal{S}|} \theta_{u_i,n}^{k+1} = \theta_{u_i}^k$ . Each  $\theta_{u_i,n}^{k+1}$  is only different to  $\theta_{u_i}^k$  in the transition from  $(s, a)$ . Therefore, this guarantees that for every state-action pair  $(s', a') \neq (s, a)$  in the first  $k$  pairs added to the causal model, the total probability across each set of mechanisms (where each set produces a different transition from  $(s', a')$ ) remains the same in  $\theta^{k+1}$  as in  $\theta^k$ . As a result, the counterfactual probabilities of all transitions from each  $(s', a')$  will be exactly the same when using  $\theta^{k+1}$  as with  $\theta^k$ . Because we assume all the constraints were satisfied when using  $\theta^k$ , this guarantees that all the constraints of the optimisation problem will continue to hold for  $(s', a')$  with  $\theta^{k+1}$ .

Now, we only need to make sure that we can assign  $\theta^{k+1}$  such that the constraints also hold for all the transitions from  $(s, a)$ . We need  $\sum_{i=1}^{|\mathcal{S}|^k} \theta_{u_i,n}^{k+1} = P(s_n | s, a)$  for every possible next state  $s_n \in \mathcal{S}$ , to satisfy the interventional probability constraints (7). For each transition  $s, a \rightarrow s_n$ , there is a  $\theta_{u_i,n}^{k+1}$  for every  $u_i$ , and for every  $\theta_{u_i,n}^{k+1}$ ,  $0 \leq \theta_{u_i,n}^{k+1} \leq \theta_{u_i}^k$ , therefore  $0 \leq \sum_{i=1}^{|\mathcal{S}|^k} \theta_{u_i,n}^{k+1} \leq 1, \forall s_n \in \mathcal{S}$ . We also know that:

$$\begin{aligned} \sum_{n=1}^{|\mathcal{S}|} \sum_{i=1}^{|\mathcal{S}|^k} \theta_{u_i,n}^{k+1} &= \sum_{i=1}^{|\mathcal{S}|^k} \theta_{u_i}^k \text{ (because we split the probabilities of each } \theta_{u_i}^k \text{)} \\ &= 1 \end{aligned}$$

and

$$\sum_{n=1}^{|\mathcal{S}|} \sum_{i=1}^{|\mathcal{S}|^k} \theta_{u_i,n}^2 = \sum_{n=1}^{|\mathcal{S}|} P(s_n | s, a) = 1$$

Therefore, no matter the assignment of  $\theta^k$ , we can satisfy the interventional probability constraints (7) for all transitions from  $(s, a)$ . In Section B.2.2, we prove that, for any  $(s, a)$ , we can assign  $\theta^{k+1}$  such that it produces the entailed counterfactual probability distribution (which we prove always satisfies the constraints of the optimisation problem). Therefore, we know we can assign  $\theta^{k+1}$  such that it produces the counterfactual probability distributions for all  $k + 1$  state-action pairs in the causal model.

**Conclusion** By induction, we have proven that as long as we can find a  $\theta$  that produces the entailed counterfactual probability distribution for each state-action pair separately, we can combine these into a single valid  $\theta$  that produces the entailed CFMDP  $\tilde{\mathcal{M}}_t$ .

### B.3 LEMMAS

**Lemma B.1** *If  $(s_t, a_t)$  and  $(\tilde{s}, \tilde{a})$  have disjoint support, the monotonicity constraints (8) and (9) and counterfactual stability (10) will be satisfied for all possible counterfactual transition probabilities for transitions from  $(\tilde{s}, \tilde{a})$ .*

**Proof B.29** *Because  $(s_t, a_t)$  and  $(\tilde{s}, \tilde{a})$  have disjoint support,  $P(s_{t+1} | \tilde{s}, \tilde{a}) = 0$  and, for all  $\tilde{s}'$  where  $P(\tilde{s}' | \tilde{s}, \tilde{a}) > 0$ ,  $P(\tilde{s}' | s_t, a_t) = 0$ . This is because for all  $\tilde{s}'$  where  $P(\tilde{s}' | s_t, a_t) > 0$ ,  $P(\tilde{s}' | \tilde{s}, \tilde{a}) = 0$ , and, for all  $\tilde{s}'$  where  $P(\tilde{s}' | \tilde{s}, \tilde{a}) > 0$ ,  $P(\tilde{s}' | s_t, a_t) = 0$ . Because for all  $\tilde{s}'$  where  $P(\tilde{s}' | \tilde{s}, \tilde{a}) > 0$ ,  $P(\tilde{s}' | s_t, a_t) = 0$ , CS (10) and Mon2 (9) will be vacuously true for all transitions from  $(\tilde{s}, \tilde{a})$ . Also, because  $P(s_{t+1} | \tilde{s}, \tilde{a}) = 0$ , Mon1 (8) is vacuously true.*

**Lemma B.2** *For any observed transition  $s_t, a_t \rightarrow s_{t+1}$  and counterfactual transition  $\tilde{s}, \tilde{a} \rightarrow \tilde{s}'$ :*

$$\sum_{u_t=1}^{|U_t|} \mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} \cdot \mu_{s_t, a_t, u_t, s_{t+1}} \cdot \theta_{u_t} \leq \min(P(s_{t+1} | s_t, a_t), P(\tilde{s}' | \tilde{s}, \tilde{a}))$$

**Proof B.30** *From our interventional probability constraints (7), we have:*

$$\sum_{u_t=1}^{|U_t|} \mu_{s_t, a_t, u_t, s_{t+1}} \cdot \theta_{u_t} = P(s_{t+1} | s_t, a_t) \quad (32)$$

and

$$\sum_{u_t=1}^{|U_t|} \mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} \cdot \theta_{u_t} = P(\tilde{s}' | \tilde{s}, \tilde{a}) \quad (33)$$

$\mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} \in \{0, 1\}$  (from its definition).  $\sum_{u_t=1}^{|U_t|} \mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} \cdot \mu_{s_t, a_t, u_t, s_{t+1}} \cdot \theta_{u_t}$  is maximal when the probability assigned to all  $u_t$  where  $\mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} = 1$  and  $\mu_{s_t, a_t, u_t, s_{t+1}} = 1$  is maximal.

$$\sum_{u_t=1}^{|U_t|} \mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} \cdot (\mu_{s_t, a_t, u_t, s_{t+1}} \cdot \theta_{u_t}) \leq \sum_{u_t=1}^{|U_t|} \mu_{s_t, a_t, u_t, s_{t+1}} \cdot \theta_{u_t} = P(s_{t+1} | s_t, a_t)$$

$$\sum_{u_t=1}^{|U_t|} (\mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} \cdot \theta_{u_t}) \cdot \mu_{s_t, a_t, u_t, s_{t+1}} \leq \sum_{u_t=1}^{|U_t|} \mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} \cdot \theta_{u_t} = P(\tilde{s}' | \tilde{s}, \tilde{a})$$

Because of Eq. (32) and Eq. (33), the probability assigned to all  $u_t$  where  $\mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} = 1$  and  $\mu_{s_t, a_t, u_t, s_{t+1}} = 1$  cannot be greater than  $P(s_{t+1} | s_t, a_t)$  or  $P(\tilde{s}' | \tilde{s}, \tilde{a})$ . Therefore,

$$\sum_{u_t=1}^{|U_t|} \mu_{\tilde{s}, \tilde{a}, u_t, \tilde{s}'} \cdot \mu_{s_t, a_t, u_t, s_{t+1}} \cdot \theta_{u_t} \leq \min(P(s_{t+1} | s_t, a_t), P(\tilde{s}' | \tilde{s}, \tilde{a}))$$

This also means that the upper bound of any counterfactual transition probability

$$\tilde{P}_t^{UB}(\tilde{s}' | \tilde{s}, \tilde{a}) \leq \frac{\min(P(\tilde{s}' | \tilde{s}, \tilde{a}), P(s_{t+1} | s_t, a_t))}{P(s_{t+1} | s_t, a_t)}.$$

**Lemma B.3** *For any observed transition  $s_t, a_t \rightarrow s_{t+1}$ ,  $\tilde{P}_t(s_{t+1} | s_t, a_t) = 1$  and,  $\forall \tilde{s}' \in \mathcal{S}$  where  $\tilde{s}' \in \mathcal{S} \setminus \{s_{t+1}\}$ ,  $\tilde{P}_t(\tilde{s}' | s_t, a_t) = 0$ .*

**Proof B.31** From (13), we have:

$$\begin{aligned}
\tilde{P}_t(s_{t+1} | s_t, a_t) &= \frac{\sum_{u_t=1}^{|U_t|} \mu_{s_t, a_t, u_t, s_{t+1}} \cdot \mu_{s_t, a_t, u_t, s_{t+1}} \cdot \theta_{u_t}}{P(s_{t+1} | s_t, a_t)} \\
&= \frac{\sum_{u_t=1}^{|U_t|} \mu_{s_t, a_t, u_t, s_{t+1}} \cdot \theta_{u_t}}{P(s_{t+1} | s_t, a_t)} \\
&= \frac{P(s_{t+1} | s_t, a_t)}{P(s_{t+1} | s_t, a_t)} && \text{(from the interventional probability constraint (7))} \\
&= 1 && \text{(since } s_t, a_t \rightarrow s_{t+1} \text{ was observed, } P(s_{t+1} | s_t, a_t) > 0)
\end{aligned}$$

$$\begin{aligned}
\tilde{P}_t(\tilde{s}' | s_t, a_t) &= \frac{\sum_{u_t=1}^{|U_t|} \mu_{s_t, a_t, u_t, \tilde{s}'} \cdot \mu_{s_t, a_t, u_t, s_{t+1}} \cdot \theta_{u_t}}{P(s_{t+1} | s_t, a_t)} \\
&= \frac{0}{P(s_{t+1} | s_t, a_t)} && \text{(since at most one of } \mu_{s_t, a_t, u_t, \tilde{s}'} \text{ and } \mu_{s_t, a_t, u_t, s_{t+1}} \text{ can equal 1)} \\
&= 0 && \text{(since } s_t, a_t \rightarrow s_{t+1} \text{ was observed, } P(s_{t+1} | s_t, a_t) > 0)
\end{aligned}$$

These results are as expected, because, given the same exogenous noise and input, we must get the same output in the counterfactual world.

**Lemma B.4** Take an arbitrary counterfactual state-action pair  $(\tilde{s}, \tilde{a})$  and observed state-action pair from an MDP, where  $(\tilde{s}, \tilde{a}) \neq (s_t, a_t)$  and  $(\tilde{s}, \tilde{a})$  has overlapping support with  $(s_t, a_t)$  and  $P(s_{t+1} | \tilde{s}, \tilde{a}) < P(s_{t+1} | s_t, a_t)$ . Let  $s_{t+1}$  be the observed next state. It can be shown that:

$$\max_{\theta} \left( \sum_{s' \in \mathcal{S} \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} \geq P(s_{t+1} | s_t, a_t) - P(s_{t+1} | \tilde{s}, \tilde{a}) \right)$$

**Proof B.32** First, let  $S_{CS} \subset \mathcal{S}$  be the set of states  $s' \in S_{CS}$  which must have a counterfactual transition probability  $\tilde{P}_t(s' | \tilde{s}, \tilde{a}) = 0$  because of the counterfactual stability constraint (i.e., where  $\forall s' \in S_{CS}, \frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} > \frac{P(s' | \tilde{s}, \tilde{a})}{P(s' | s_t, a_t)}$  and  $P(s' | s_t, a_t) > 0$ ). Consider the total interventional probability of across all states  $s' \in S_{CS}, \forall s' \in S_{CS}$ :

$$\begin{aligned}
\frac{P(s' | \tilde{s}, \tilde{a})}{P(s' | s_t, a_t)} &< \frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} \text{ (from CS (10))} \\
P(s' | \tilde{s}, \tilde{a}) &< \frac{P(s_{t+1} | \tilde{s}, \tilde{a}) \cdot P(s' | s_t, a_t)}{P(s_{t+1} | s_t, a_t)}
\end{aligned}$$

The sum of the interventional probabilities of these transitions is:

$$\sum_{s' \in S_{CS}} P(s' | \tilde{s}, \tilde{a}) \leq \sum_{s' \in S_{CS}} \frac{P(s' | s_t, a_t) \cdot P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)}$$

Let  $S_{other}$  be the set of all other states  $s' \in \mathcal{S} \setminus \{s_{t+1}\} \cup S_{CS}$ . We can calculate the sum of the interventional probabilities of the transitions going to states  $s' \in S_{other}$ :



$$\begin{aligned}
\sum_{s' \in S_{\text{Other}}} P(s' | \tilde{s}, \tilde{a}) &= 1 - P(s_{t+1} | \tilde{s}, \tilde{a}) - \sum_{s'' \in S_{CS}} P(s'' | \tilde{s}, \tilde{a}) \\
&\geq 1 - P(s_{t+1} | \tilde{s}, \tilde{a}) - \sum_{s'' \in S_{CS}} \frac{P(s'' | s_t, a_t) \cdot P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} \\
&= 1 - P(s_{t+1} | \tilde{s}, \tilde{a}) - \frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} \cdot \sum_{s'' \in S_{CS}} P(s'' | s_t, a_t) \\
&= 1 - P(s_{t+1} | \tilde{s}, \tilde{a}) \\
&\quad - \frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} \cdot (1 - P(s_{t+1} | s_t, a_t) - \sum_{s' \in S_{\text{Other}}} P(s' | s_t, a_t)) \\
&= 1 - P(s_{t+1} | \tilde{s}, \tilde{a}) - \frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} \\
&\quad + P(s_{t+1} | \tilde{s}, \tilde{a}) + \frac{P(s_{t+1} | \tilde{s}, \tilde{a}) \cdot \sum_{s' \in S_{\text{Other}}} P(s' | s_t, a_t)}{P(s_{t+1} | s_t, a_t)} \\
&= 1 - \frac{P(s_{t+1} | \tilde{s}, \tilde{a})}{P(s_{t+1} | s_t, a_t)} + \frac{P(s_{t+1} | \tilde{s}, \tilde{a}) \cdot \sum_{s' \in S_{\text{Other}}} P(s' | s_t, a_t)}{P(s_{t+1} | s_t, a_t)}
\end{aligned}$$

Because of the Mon2 (9) and CS (10) constraints, we know that  $\forall s'' \in S_{CS}, \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s''}} \theta_{u_t} = 0$ , and  $s' \in$

$S_{\text{Other}}, \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} \leq P(s' | \tilde{s}, \tilde{a}) \cdot P(s_{t+1} | s_t, a_t)$ . So,

$$\begin{aligned}
&\max_{\theta} \left( \sum_{s'' \in S_{CS}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s''}} \theta_{u_t} \right) = 0 \\
&\max_{\theta} \left( \sum_{s' \in S_{\text{Other}}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} \right) \leq \sum_{s' \in S_{\text{Other}}} P(s' | \tilde{s}, \tilde{a}) \cdot P(s_{t+1} | s_t, a_t)
\end{aligned}$$

Therefore, the maximum  $\sum_{s' \in S \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t}$  is as follows:

$$\begin{aligned}
&\max_{\theta} \left( \sum_{s' \in S \setminus \{s_{t+1}\}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} \right) = \max_{\theta} \left( \sum_{s'' \in S_{CS}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s''}} \theta_{u_t} \right) + \max_{\theta} \left( \sum_{s' \in S_{\text{Other}}} \sum_{\substack{u_t \in U_t \\ f(s_t, a_t, u_t) = s_{t+1} \\ f(\tilde{s}, \tilde{a}, u_t) = s'}} \theta_{u_t} \right) \\
&= \sum_{s'' \in S_{\text{Other}}} P(s'' | \tilde{s}, \tilde{a}) \cdot P(s_{t+1} | s_t, a_t) \\
&\geq P(s_{t+1} | s_t, a_t) - P(s_{t+1} | \tilde{s}, \tilde{a}) + P(s_{t+1} | \tilde{s}, \tilde{a}) \cdot \sum_{s' \in S_{\text{Other}}} P(s' | s_t, a_t) \\
&\geq P(s_{t+1} | s_t, a_t) - P(s_{t+1} | \tilde{s}, \tilde{a})
\end{aligned}$$

(34)

## C EQUIVALENCE WITH EXISTING WORK

Li and Pearl [2024]’s bounds for the probability of causation are provided below. We can show that these are equivalent to our analytical bounds where the support of the counterfactual state-action pair  $(\tilde{s}, \tilde{a})$  is disjoint from the support of the observed state-action pair  $(s_t, a_t)$ , given in Theorems B.3 and B.4.

**Theorem C.1 (Bounds for Probability of Causation)** *Suppose  $X$  has  $m$  values  $x_1, \dots, x_m$  and  $Y$  has  $n$  values  $y_1, \dots, y_n$ , the probability of causation  $P(y_{i_{x_j}}, y_k, x_p)$  where  $1 \leq i, k \leq n$ ,  $1 \leq j, p \leq m$ ,  $j \neq p$  is given by:*

$$P^{LB}(y_{i_{x_j}}, y_k, x_p) = \max(0, P(y_{i_{x_j}}) + P(x_p, y_k) - 1 + P(x_j) - P(x_j, y_i))$$

$$P^{UB}(y_{i_{x_j}}, y_k, x_p) = \min(P(y_{i_{x_j}}) - P(x_j, y_i), P(x_p, y_k))$$

**Proof C.1** *We show that Li and Pearl [2024]’s upper and lower bounds for the probability of causation are equivalent to our bounds in Theorems B.3 and B.4, as follows.*

$$\begin{aligned} P^{UB}(y_{i_{x_j}}, y_k, x_p) &= \min(P(x_p, y_k), P(y_{i_{x_j}}) - P(x_j, y_i)) \\ P^{UB}(y_{i_{x_j}} | y_k, x_p) &= \frac{\min(P(x_p, y_k), P(y_{i_{x_j}}) - P(x_j, y_i))}{P(x_p, y_k)} \\ &= \min(1, \frac{P(y_{i_{x_j}}) - P(x_j, y_i)}{P(x_p, y_k)}) \\ &= \min(1, \frac{P(y_{i_{x_j}}) - P(y_{i_{x_j}}, x_j)}{P(y_{k_{x_p}}, x_p)}) \\ &= \min(1, \frac{P(y_{i_{x_j}}) - P(y_{i_{x_j}}) \cdot P(x_j)}{P(y_{k_{x_p}}) \cdot P(x_p)}) \\ &= \min(1, \frac{P(y_{i_{x_j}})}{P(y_{k_{x_p}})}) \text{ because } P(x_p) = 1 \text{ and } P(x_j) = 0 \end{aligned} \tag{35}$$

which is equivalent to the upper bound in Theorem B.3 where  $x_p = (s_t, a_t)$ ,  $y_k = s_{t+1}$ ,  $x_j = (\tilde{s}, \tilde{a})$  and  $y_i = \tilde{s}'$ .

$$\begin{aligned} P^{LB}(y_{i_{x_j}}, y_k, x_p) &= \max(0, P(y_{i_{x_j}}) + P(x_p, y_k) - 1 + P(x_j) - P(x_j, y_i)) \\ P^{LB}(y_{i_{x_j}} | y_k, x_p) &= \frac{\max(0, P(y_{i_{x_j}}) + P(x_p, y_k) - 1 + P(x_j) - P(x_j, y_i))}{P(x_p, y_k)} \\ &= \max(0, \frac{P(y_{i_{x_j}}) + P(x_p, y_k) - 1 + P(x_j) - P(x_j, y_i)}{P(x_p, y_k)}) \\ &= \max(0, \frac{P(y_{i_{x_j}}) + P(y_{k_{x_p}}, x_p) - 1 + P(x_j) - P(y_{i_{x_j}}, x_j)}{P(y_{k_{x_p}}, x_p)}) \\ &= \max(0, \frac{P(y_{i_{x_j}}) + P(y_{k_{x_p}}) \cdot P(x_p) - 1 + P(x_j) - P(y_{i_{x_j}}) \cdot P(x_j)}{P(y_{k_{x_p}}) \cdot P(x_p)}) \\ &= \max(0, \frac{P(y_{i_{x_j}}) + P(y_{k_{x_p}}) - 1}{P(y_{k_{x_p}})}) \text{ because } P(x_p) = 1 \text{ and } P(x_j) = 0 \\ &= \max(0, \frac{P(y_{i_{x_j}}) - (1 - P(y_{k_{x_p}}))}{P(y_{k_{x_p}})}) \end{aligned} \tag{36}$$

which is equivalent to the lower bound in Theorem B.4 where  $x_p = (s_t, a_t)$ ,  $y_k = s_{t+1}$ ,  $x_j = (\tilde{s}, \tilde{a})$  and  $y_i = \tilde{s}'$ .

## D TRAINING DETAILS

Our algorithm was executed on a 128-core machine with an Intel Xeon CPU and 512 GB RAM. The interval CFMDP generation part of the algorithm is implemented in Python 3.10, and the value iteration algorithm over the interval CFMDP was implemented in Julia 1.9.4, using the Julia Interval MDP package (v.0.2.1) [Mathiesen et al., 2024]. Most of the algorithm was run single-threaded, except for generating the Sepsis interval CFMDP and Sepsis Gumbel-max SCM CFMDP, which were run in parallel. 10 threads were needed to generate the Sepsis interval CFMDP, and 32 threads were required to calculate the Gumbel-max SCM CFMDP.