# A class of parabolic reaction-diffusion systems governed by spectral fractional Laplacians : Analysis and numerical simulations

Maha DAOUD

POEMS, ENSTA Paris, Institut Polytechnique de Paris, Palaiseau, France mahaadaoud@gmail.com

February 20, 2025

#### Abstract

In this paper, we prove the global-in-time existence of strong solutions to a class of fractional parabolic reaction-diffusion systems set in a bounded open subset of  $\mathbb{R}^N$ . The diffusion operators are of the form  $u_i \mapsto d_i(-\Delta)_{Sp}^{s_i} u_i$ , where  $0 < s_i < 1$ . The operator  $(-\Delta)_{Sp}^s$  stands for the commonly called spectral fractional Laplacian. Moreover, the nonlinear reactive terms are assumed to fulfill natural structural conditions that ensure the nonnegativity of the solutions and provide uniform control of the total mass. We establish the global existence of strong solutions under the assumption that the nonlinearities exhibit at most polynomial growth. Our results extend previous results obtained when the diffusion operators are of the form  $u_i \mapsto d_i(-\Delta)^s u_i$ , where  $(-\Delta)^s$  denotes the widely known as regional fractional Laplacian. Furthermore, we use numerical simulations to investigate the global existence of solutions to the fractional version of the so-called "Brusselator" system, a theoretical question that remains open to date.

Mathematics Subject Classification (2020): 35R11, 35J62. 30G50, 47H10, 35B45.

**Keywords.** Reaction-diffusion system, fractional diffusion, strong solution, global existence, numerical simulation.

## 1 Introduction

In this work, we study the global existence-in-time of nonnegative strong solutions to a class of fractional parabolic reaction-diffusion systems. These systems are governed by spectral fractional Laplacians of different orders and take the following form:

$$(S) \begin{cases} \forall i = 1, \dots, m, \\ \partial_t u_i(t, \mathbf{x}) + d_i(-\Delta)_{Sp}^{s_i} u_i(t, \mathbf{x}) &= f_i(u_1(t, \mathbf{x}), \dots, u_m(t, \mathbf{x})), & (t, \mathbf{x}) \in (0, T) \times \Omega, \\ \mathcal{B}[u_i(t, \mathbf{x})] &= 0, & (t, \mathbf{x}) \in (0, T) \times \partial\Omega, \\ u_i(0, \mathbf{x}) &= u_{0i}(\mathbf{x}), & \mathbf{x} \in \Omega, \end{cases}$$

where  $\Omega$  is a bounded regular open subset of  $\mathbb{R}^N$  with  $N \in \mathbb{N}^*$ ,  $m \geq 2$  and for any  $i \in [\![1,m]\!]$ ,  $0 < s_i < 1, d_i > 0$  and  $f_i$  is locally Lipschitz continuous. Moreover,  $(-\Delta)_{Sp}^s$  represents a spectral fractional Laplacian associated with either Dirichlet or Neumann homogeneous boundary conditions. As such, the operator  $(-\Delta)_{Sp}^s$  is dependent to an homogeneous boundary condition:

$$\mathcal{B}[u] = u \quad ext{or} \quad \mathcal{B}[u] = rac{\partial u}{\partial 
u} \quad ext{on} \quad \partial \Omega.$$

Let  $\{\lambda_k, e_k\}_{k \in \mathbb{N}^*}$  be the eigenpairs of the "classical" Dirichlet or Neumann Laplacian. Then, the corresponding spectral fractional Laplacian is defined for  $s \in (0, 1)$  by

(1.1) 
$$(-\Delta)_{Sp}^{s} u(\mathbf{x}) := \sum_{k=1}^{+\infty} \lambda_{k}^{s} u_{k} e_{k}(\mathbf{x}),$$

where  $u_k = \int_{\Omega} u(\mathbf{x}) e_k(\mathbf{x}) d\mathbf{x}$  and  $||e_k||_{L^2(\Omega)} = 1$ .

Parabolic reaction-diffusion systems have long been employed to model a wide range of phenomena, from chemical reactions to biological processes (see, for instance [67, 72, 76, 81] and references therein. The "classical" Laplacian operator, commonly used to describe diffusion, has proven effective in many contexts. Further discussion can be found in [3, 25, 35–37, 45, 56– 58, 72, 75, 86, 87] and their bibliographies. However, in many cases, this operator fails to capture nonlocal interactions that are critical for accurately simulating complex phenomena. This has led to a surge of interest in fractional-type Laplacian operators, which offer a more suitable framework for modeling nonlocal diffusive processes. For further details, we refer the interested reader, for instance, to [2, 21, 30, 48, 63, 85, 90–92] and references cited therein. More broadly, fractional partial differential equations have gained prominence in recent years and have become fundamental in many scientific areas, see [52, 77] and their references.

In open bounded subsets of  $\mathbb{R}^N$ , there are various fractional Laplacians, each with its own characteristics and applications. Among theses different operators, two main classes are widely studied: the spectral fractional Laplacian (*SFL*) (typically associated with Dirichlet homogeneous boundary condition) and the regional fractional Laplacian (*RFL*) defined as

(1.2) 
$$(-\Delta)^{s} u(\mathbf{x}) := a_{N,s} \operatorname{P.V.} \int_{\mathbb{R}^{N}} \frac{u(\mathbf{x}) - u(\mathbf{y})}{\|\mathbf{x} - \mathbf{y}\|^{N+2s}} d\mathbf{y}, \quad \mathbf{x} \in \Omega,$$

where  $u(\mathbf{x}) = 0$  in  $\mathbb{R}^N \setminus \Omega$ ,  $a_{N,s} := \frac{s2^{2s}\Gamma(\frac{N+2s}{2})}{\pi^{\frac{N}{2}}\Gamma(1-s)}$ ,  $\|\cdot\|$  is the Euclidean norm of  $\mathbb{R}^N$  and P.V. stands for the Cauchy principal value. While both operators generalize the classical Laplacian,

stands for the Cauchy principal value. While both operators generalize the classical Laplacian, they differ in that one incorporates information from the boundary, while the other considers influences from the exterior of the domain, leading to different stochastic processes. For more details about the striking differences between the two operators, see [84] (see also [66, Section 5.3], [2, 44] and references therein). Furthermore, the author and her collaborator exhibit in [30, Section 4] several simple yet significant one-dimensional examples that effectively highlight the difference between the two operators. In this context, the same authors and their collaborator have previously studied in [31] System (S) with the (RFL). Overall, we have observed that the two operators lead to notable differences in the properties of the systems they govern: they require the adoption of tailored methodologies for each operator and careful consideration of specific technical aspects.

Recently, considerable studies have explored the existence of solutions to fractional reactiondiffusion problems, particularly for single parabolic equations (*i.e.* m = 1) in open bounded subsets of  $\mathbb{R}^N$ . For further details, we refer the reader to [18, 26, 33, 54, 68, 88] for the *(SFL)* and to [7, 14, 15, 39, 41, 42, 62] for the *(RFL)*. However, there are relatively few works addressing systems ( $m \ge 2$ ) governed by fractional Laplacians. For systems with m = 2 involving the *(SFL)*, we refer to [83, 94], while for the *(RFL)*, relevant studies can be found in [9–12, 64]. As far as we know, the only existing work for  $m \ge 2$  is [31], where the author and her collaborators analyzed System (S) with the *(RFL)* and  $s_1 = \cdots = s_m$ .

One should question the intriguing case of considering fractional Laplacians of different orders, *i.e.* at least for two different  $i, j \in [\![1, m]\!] s_i \neq s_j$ . This is motivated, for example, by stochastic considerations. Specifically, previous studies have investigated equations involving a sum of two fractional Laplacians with different orders. Such sums naturally arise from the superposition of two stochastic processes, each with distinct random walks and Lévy flights. This approach has numerous applications, including chemical reaction design, plasma physics, biophysics and population dynamics. It also serves to model heart anomalies caused by arterial issues. For more details, we refer to [27, 34] and references included therein.

Let us go back to System (S) with the (SFL). In this paper, our main goal is to extend the results obtained in [31] to a more general setting, specifically by incorporating other different fractional operators with varying orders.

Since the unknown  $\mathbf{u} = (u_1, \dots, u_m)$  represents quantities such as chemical concentrations or population densities, we focus on nonnegative solutions. Therefore, the initial data have to be chosen nonnegative, i.e.,  $u_{0i} \ge 0$  for any  $i \in [\![1,m]\!]$ . Moreover, it is well-known that solutions, as long as they exist, remain nonnegative if the reaction terms  $f_i$  satisfy the so-called "quasipositivity" property, that is:

(**P**) 
$$\forall i \in [\![1,m]\!], \quad f_i(r_1,\cdots,r_{i-1},0,r_{i+1},\cdots,r_m) \ge 0, \quad \forall \mathbf{r} = (r_1,\cdots,r_m) \in [0,+\infty)^m.$$

In addition, in many cases, the model inherently includes a control or even preservation of the total mass, specifically

(1.3) 
$$\sum_{i=1}^{m} \int_{\Omega} u_i(t, \mathbf{x}) \, d\mathbf{x} \le C \sum_{i=1}^{m} \int_{\Omega} u_{0i}(\mathbf{x}) \, d\mathbf{x} \quad \text{for some } C > 0.$$

The control (1.3) is satisfied if

(**M**) 
$$\exists (a_1, \cdots a_m) \in (0, +\infty)^m \text{ such that } \sum_{i=1}^m a_i f_i \le 0$$

or, more broadly, if this sum grows at most linearly with respect to its variables, namely

(**M'**) 
$$\forall \mathbf{r} = (r_1, \dots, r_m) \in [0, +\infty)^m, \ \sum_{i=1}^m a_i f_i(\mathbf{r}) \le C \Big[ 1 + \sum_{i=1}^m r_i \Big] \text{ for some } C \ge 0.$$

It is worth highlighting that properties (**P**) and (**M**) (or (**M'**)) are frequently observed in various applications, including models that describe evolutionary processes involving spatial diffusion and chemical reactions. Nevertheless, It turns out that these properties do not keep the solution from blowing up in  $L^{\infty}$ -norm, even in finite time (see [74]). Hence, in addition to the structure (**P**)+ (**M**), some growth restrictions and extra structure on the nonlinearities are needed if one expects global existence of strong solutions. This issue has been intensively studied in the case of the "classical" Laplacian, especially when the initial data are bounded. However, many open and challenging problems remain. Indeed, the construction of global solutions is primarily hindered by two major obstructions: (i) when the the  $f_i$ 's have quadratic, even faster, growth for large values of the  $u_i$ 's ; (ii) when the  $d_i$ 's are very different.

To situate our work within the existing body of research and underscore the novelty of this paper, let us provide a concise review of the literature concerning the global existence of strong solutions to system (S).

• Case where the diffusions are governed by classical Laplacian (i.e. all  $s_i$ 's = 1)

As mentioned above, for bounded initial data, some sufficient conditions on the fi's are required to guarantee the global existence of a strong solution (see Definition 2.2). More precisely,

- (i) polynomial growth and the so-called *triangular structure* (see (TS)). For further details, see [72] and its references;
- (ii) quadratic growth. For this setting, we refer, e.g., to [25, 36, 37, 45, 75, 86]);
- (iii) diffusion coefficients close to one another as illustrated, for instance, in [24, 35] (sometimes referred to as *quasi-uniform* in some works like [36, 37]);
- (iv) when the growth of the  $f_i$ 's barely exceed polynomial growth, limited research has been conducted, and solely for m = 2. The most thoroughly examined model is  $f_2(u_1, u_2) = -f_1(u_1, u_2) = u_1 e^{u_2^{\beta}}$ . In this case, the global existence of strong solutions has been established in [46] for  $\beta < 1$ , and in [13] for  $\beta = 1$  yet under a restrictive assumption on the size of  $u_{01}$ . However, the problem remains open for  $\beta > 1$ .

Let us emphasize that our primary interest lies in examining strong solutions. It goes without saying that there is a substantial body of research on weak solutions, including notable works such as [19, 28, 55, 57–60, 71–73]. On the other hand, we would like to underscore that the cited references do not encompass the extensive literature on the subject. For an exhaustive review of findings up to 2010, we refer interested readers to [72]. This survey not only provides a general overview of the problem but also offers an additional insightful commentary on the mathematical challenges posed by such systems. For some further recent results, see [24, 25, 35–37, 56, 75, 82, 86] and references included therein.

• Case where the diffusions are governed by fractional Laplacians (i.e.  $0 < s_i < 1$ )

#### ◊ With regional fractional Laplacians

— Systems with two equations (i.e. m = 2)

In the case where  $\Omega$  is bounded, the author and her coworkers have studied the following System

(1.4)

$$\begin{split} \dot{\partial}_{t} u_{1}(t, \mathbf{x}) &+ (-\Delta)^{s_{1}} u_{1}(t, \mathbf{x}) &= \|\nabla u_{2}(t, \mathbf{x})\|^{q} + h_{1}(t, \mathbf{x}), \quad (t, \mathbf{x}) \in (0, T) \times \Omega, \\ \partial_{t} u_{2}(t, \mathbf{x}) &+ (-\Delta)^{s_{2}} u_{2}(t, \mathbf{x}) &= \|\nabla u_{1}(t, \mathbf{x})\|^{p} + h_{2}(t, \mathbf{x}), \quad (t, \mathbf{x}) \in (0, T) \times \Omega, \\ u_{1}(t, \mathbf{x}) &= u_{2}(t, \mathbf{x}) &= 0, \quad (t, \mathbf{x}) \in (0, T) \times (\mathbb{R}^{N} \setminus \Omega), \\ u_{1}(0, \mathbf{x}) &= u_{01}(\mathbf{x}) \geq 0, \quad \mathbf{x} \in \Omega, \\ u_{2}(0, \mathbf{x}) &= u_{02}(\mathbf{x}) \geq 0, \quad \mathbf{x} \in \Omega, \end{split}$$

where  $h_1, h_2 \ge 0$  and p, q > 1 (see [9] for the case  $s_1 = s_2$  and [12] for the case  $s_1 \ne s_2$ ). Clearly, the right-hand sides in this case do not fall within the framework of (S), nor do they satisfy **(M)**. It is worth highlighting that the same authors have also explored two elliptic versions of System (1.4), namely: the case of identical fractional orders in [10], and a generalized setting with different orders in [11].

On the other hand, in the case  $\Omega = \mathbb{R}^N$ , the only works known to us are those of [4–6, 53].

— Systems with  $m \ge 2$  equations

As per our knowledge, the only existing works pertain to the case of equal fractional orders:

(i) In[31], the author and her collaborators investigated System (S) posed in a bounded open subset  $\Omega$ . First, they have treated the case of reversible chemical reactions of three components (see  $(S_{\alpha,\beta,\gamma})$  below). Second, they have examined the existence of strong global nonnegative solutions to System (S) along with properties (**P**)+(**M**), where the right-hand-sides are of polynomial growth and fulfill (TS). Furthermore, they have numerically investigated the global existence of solutions to the following system

 $(S_{\exp})$ 

$$\begin{cases} \partial_t u_1(t, \mathbf{x}) + d_1(-\Delta)^s u_1(t, \mathbf{x}) &= -u_1(t, \mathbf{x})e^{(u_2(t, \mathbf{x}))^{\beta}}, \quad (t, \mathbf{x}) \in (0, T) \times \Omega, \\ \partial_t u_2(t, \mathbf{x}) + d_2(-\Delta)^s u_2(t, \mathbf{x}) &= u_1(t, \mathbf{x})e^{(u_2(t, \mathbf{x}))^{\beta}}, \quad (t, \mathbf{x}) \in (0, T) \times \Omega, \\ u_1(t, \mathbf{x}) = u_2(t, \mathbf{x}) &= 0, \qquad (t, \mathbf{x}) \in (0, T) \times (\mathbb{R}^N \setminus \Omega), \\ u_1(0, \mathbf{x}) &= u_{01}(\mathbf{x}), \qquad \mathbf{x} \in \Omega, \\ u_2(0, \mathbf{x}) &= u_{02}(\mathbf{x}), \qquad \mathbf{x} \in \Omega, \end{cases}$$

where  $\beta > 1$ . Let us emphasize that this system arises in the modeling of exothermic combustion in gases, see, *e.g.*, [49].

(ii) In [69], the authors addressed System (S) in the case  $\Omega = \mathbb{R}^N$ .

#### $\diamond$ With spectral fractional Laplacians

In this case, the sole available studies concern System (S) with m = 2:

(i) In [85], the authors introduce a fractional model to describe coral growth in a tank, applied to modeling the pattern formation of coral reefs. More specifically, the model is characterized by the parameters  $s_1 = s_2$ ,  $f_1(u_1, u_2) = 1 - u_1 - \lambda u_1 u_2^2$  and  $f_2(u_1, u_2) = -\mu u_2 + \lambda u_1 u_2^2$ , with  $\lambda, \mu > 0$ .

(ii) In [94], the authors proposed another fractional model to study the long-range geographical spread of infectious diseases. To elaborate, the model is described by the parameters  $s_1 \neq s_2$ ,  $f_1(u_1, u_2) = \varphi(\mathbf{x})u_1 - \phi(\mathbf{x})u_1^2 - \frac{\psi(\mathbf{x})u_1u_2}{u_1 + u_2} + h(\mathbf{x})u_2$  and  $f_2(u_1, u_2) = \frac{\psi(\mathbf{x})u_1u_2}{u_1 + u_2} - h(\mathbf{x})u_2$ , where the functions  $\varphi$ ,  $\phi$ ,  $\psi$  and h are Hölder continuous in  $\overline{\Omega}$ .

To the best of our knowledge, System (S) with the reaction terms satisfying  $(\mathbf{P})+(\mathbf{M})$  and of polynomial growth has not yet been thoroughly investigated. In this work, we aim to address this by establishing the global existence of nonnegative strong solutions to System (S). This is the main purpose of this work. More precisely, we aim to generalize within the fractional setting  $(0 < s_i < 1)$  two main well-established results in the classical case  $(i.e. \text{ all } s_i) = 1$ :

1. The first one concerns the typical case of reversible chemical reactions for three species, that is

$$\begin{cases} (S_{\alpha,\beta,\gamma}) \\ \forall i = 1,2,3, \\ \partial_t u_i(t,\mathbf{x}) + d_i(-\Delta)_{Sp}^{s_i} u_i(t,\mathbf{x}) &= f_i(u_1(t,\mathbf{x}), u_2(t,\mathbf{x}), u_3(t,\mathbf{x})), \quad (t,\mathbf{x}) \in (0,T) \times \Omega, \\ \mathcal{B}[u_i(t,\mathbf{x})] &= 0, \quad (t,\mathbf{x}) \in (0,T) \times \partial\Omega, \\ u_i(0,\mathbf{x}) &= u_{0i}(\mathbf{x}), \quad \mathbf{x} \in \Omega, \end{cases}$$

where  $f_1 = \alpha g$ ,  $f_2 = \beta g$ ,  $f_3 = -\gamma g$  with  $g = u_3^{\gamma} - u_1^{\alpha} u_2^{\beta}$  and  $1 \leq \alpha, \beta, \gamma < +\infty$ . It is worth noting that this system naturally emerges in chemical kinetics when modeling the following reversible reaction

(1.5) 
$$\alpha_1 U_1 + \alpha_2 U_2 \rightleftharpoons \alpha_3 U_3$$

where  $u_1$ ,  $u_2$ ,  $u_3$  stand for the density of  $U_1$ ,  $U_2$  and  $U_3$  respectively, and  $\alpha_1$ ,  $\alpha_2$ ,  $\alpha_3$  are the stoichiometric coefficients. Moreover, let us point out that in addition to satisfying (**P**), the functions  $f_1$ ,  $f_2$  and  $f_3$  also fulfill (**M**), *i.e.*  $\beta\gamma f_1 + \alpha\gamma f_2 + 2\alpha\beta f_3 = 0$ .

It should also be emphasized that global existence of strong nonnegative global solution to System  $(S_{\alpha,\beta,\gamma})$  in the classical case  $s_1 = s_2 = s_3$  has been established in [56].

2. The second one addresses the case where System (S) consists of  $m \ge 2$  equations and the  $f_i$ 's fulfill a triangular structure (see (TS)). Let us mention that the classical version (*i.e.* all  $s_i$ 's=1) of this problem has been studied in [72].

To establish our main theorems, we will adapt several well-known tools from the classical setting to the fractional framework. These include the maximal regularity theorem (see Theorem 2.2), a Lamberton-type estimation in  $L^p$  (see Theorem 2.5), the so-called Pierre's duality Lemma (see Lemma 3.2). As emphasized in [31], these results are noteworthy in themselves. That being said, it should be noted that not all results from the classical case can be directly extended to the fractional setting; and when it is, it often entails substantial challenges. To illustrate, in a different context, we refer to [9–12].

On the other hand, we will present some numerical simulations to examine the global existence of solutions to the following fractional version of the "Brusselator"  $(S_{\text{Bruss}})$ 

$$\begin{array}{rcl} \mathcal{B}_{2}^{(n,n)} & (t,\mathbf{x}) + d_{1}(-\Delta)_{Sp}^{s_{1}}u_{1}(t,\mathbf{x}) &= -u_{1}(t,\mathbf{x})(u_{2}(t,\mathbf{x}))^{2} + bu_{2}(t,\mathbf{x}), & (t,\mathbf{x}) \in (0,T) \times \Omega, \\ \partial_{t}u_{2}(t,\mathbf{x}) + d_{2}(-\Delta)_{Sp}^{s_{2}}u_{2}(t,\mathbf{x}) &= u_{1}(t,\mathbf{x})(u_{2}(t,\mathbf{x}))^{2} - (b+1)u_{2}(t,\mathbf{x}) + a, & (t,\mathbf{x}) \in (0,T) \times \Omega, \\ \mathcal{B}[u_{1}(t,\mathbf{x})] = \mathcal{B}[u_{2}(t,\mathbf{x})] &= 0, & (t,\mathbf{x}) \in (0,T) \times \partial\Omega, \\ u_{1}(0,\mathbf{x}) &= u_{01}(\mathbf{x}), & \mathbf{x} \in \Omega, \\ u_{2}(0,\mathbf{x}) &= u_{02}(\mathbf{x}), & \mathbf{x} \in \Omega, \end{array}$$

where  $d_1, d_2, a, b > 0$  and  $(u_{01}, u_{02}) \in (L^{\infty}(\Omega)^+)^2$ . We can easily check that the right-hand-sides in System (S<sub>Bruss</sub>) fulfill (**P**) + (**M**'). Let us mention that such systems appear in the modeling of chemical morphogenetic processes, see [78, 79, 89].

Let us emphasize that global existence of strong solutions to System  $(S_{\text{Bruss}})$  with the classical Laplacian, has been established in [50]; see also [65, 80]. Nevertheless, in the fractional case, it remains an open theoretical question.

Before concluding the introduction, it is worth noting that the global existence of strong solutions to System (S) with regional fractional Laplacians of different orders, will be addressed in a forthcoming work, in the case where  $\Omega$  is either bounded or equal to  $\mathbb{R}^N$ .

The structure of this article is as follows. In Section 2, we provide the necessary background for the subsequent sections. First, we give two equivalent definitions of the (SFL), the adequate framework and some fundamental properties of the corresponding semigroup. Then, we address a fractional evolution equation. More precisely, We investigate the criteria for ensuring the boundedness of the solution. Additionally, we focus on relevant concepts such as the dual problem, maximal regularity and the comparison principle. Section 3 is divided into four subsections. In subsection 3.1, we begin by investigating the local existence of strong solutions to System (S). Subsection 3.2 is dedicated to extending Pierre's duality lemma to the fractional case with different orders. To do so, we will establish a result that contributes to this extension. Subsections 3.3 and 3.4 are devoted respectively to the proofs of our two main global existence of solutions to System  $(S_{\text{Bruss}})$ .

Now let us fix some notations that will be used throughout this work.

#### Notations.

- $\Omega$  is a bounded regular open subset of  $\mathbb{R}^N$  with  $N \in \mathbb{N}^*$ .
- For any T > 0,  $Q_T := (0, T) \times \Omega$  and  $\Sigma_T := (0, T) \times \partial \Omega$ .

$$- \text{ For any } p \in [1, +\infty), \|\phi\|_{L^p(\Omega)} = \left(\int_{\Omega} |\phi(\mathbf{x})|^p d\mathbf{x}\right)^{\frac{1}{p}} \text{ and } \|\psi\|_{L^p(Q_T)} = \left(\int_{0}^{T} \int_{\Omega} |\psi(t, \mathbf{x})|^p dt d\mathbf{x}\right)^{\frac{1}{p}} .$$
$$- \|\phi\|_{L^{\infty}(\Omega)} = \operatorname{ess \, sup}_{\mathbf{x} \in \Omega} |\phi(\mathbf{x})| \text{ and } \|\psi\|_{L^{\infty}(Q_T)} = \operatorname{ess \, sup}_{(t, \mathbf{x}) \in Q_T} |\psi(t, \mathbf{x})|.$$

— The boundary operator  $\mathcal{B}$  is defined as:

$$\mathcal{B}[u] = u \quad \text{or} \quad \mathcal{B}[u] = \frac{\partial u}{\partial \nu} \quad \text{on} \quad \partial \Omega.$$

## 2 Preliminaries

In this section, we briefly review some results that will be relevant for later use. First, we recall the spectral fractional Laplacian and its semigroup on  $L^2(\Omega)$ . Subsequently, we summarize existence and regularity results for solutions to a fractional evolution problem, along with a Lamberton-type estimate related to the dual problem.

#### **2.1** Spectral fractional Laplacian (*SFL*)

First, we set up the functional framework. Then, we recall two equivalent definitions of the spectral fractional Laplacian and discuss properties of the corresponding semigroup on  $L^2(\Omega)$ .

#### 2.1.1 Functional framework

Let  $\{\lambda_k\}_{k\in\mathbb{N}^*}$  with  $0 < \lambda_1 \leq \lambda_2 \leq \lambda_3 \leq \dots \uparrow +\infty$  be the sequence of eigenvalues of  $-\Delta$  in  $\Omega$  with homogeneous Dirichlet or Neumann boundary conditions, and  $\{e_k\}_{k\in\mathbb{N}^*}$  the corresponding eigenfunctions. Namely, for any  $k\in\mathbb{N}^*$ ,  $e_k$  is the solution to

$$\begin{cases} -\Delta e_k(\mathbf{x}) &= \lambda_k e_k(\mathbf{x}), \quad \mathbf{x} \in \Omega, \\ \mathcal{B}[e_k(\mathbf{x})] &= 0, \qquad \mathbf{x} \in \partial \Omega. \end{cases}$$

Furthermore, the eigenfunctions are normalized in  $L^2(\Omega)$ , that is  $\int_{\Omega} e_i(\mathbf{x}) e_j(\mathbf{x}) d\mathbf{x} = \delta_{ij}$ , where  $\delta_{ij}$  is the Kronecker symbol. For more details, see, for instance, [47, Theorem 1.2.2 and Theorem 1.2.8].

In this setting, we can write, for any  $u \in L^2(\Omega)$ ,  $u(\mathbf{x}) = \sum_{k=1}^{+\infty} u_k e_k(\mathbf{x})$  such that  $u_k = \int_{\Omega} u(\mathbf{x}) e_k(\mathbf{x}) d\mathbf{x}$ . Moreover, let us introduce the following space

$$\mathcal{X}^{s}(\Omega) := \{ u \in L^{2}(\Omega) \text{ such that } \sum_{k=1}^{+\infty} \lambda_{k}^{s} u_{k}^{2} < +\infty \}.$$

We easily verify that  $(\mathcal{X}^s(\Omega), \langle \cdot, \cdot \rangle_{\mathcal{X}^s(\Omega)})$  is a Hilbert space, where

$$\langle u, v \rangle_{\mathcal{X}^s(\Omega)} := \sum_{k=1}^{+\infty} \lambda_k^s u_k v_k, \text{ for any } u, v \in \mathcal{X}^s(\Omega).$$

For further details on this space, we refer the interested reader to, e.g., [51, Section 1] or [17, Subsection 3.1].

#### **2.1.2** First definition of the (SFL)

Let  $\mathbf{x} \in \Omega$  and  $s \in (0, 1)$ . The spectral fractional Laplacian (SFL) is defined as

(2.1) 
$$(-\Delta)_{Sp}^{s} u(\mathbf{x}) := \sum_{k=1}^{+\infty} \lambda_{k}^{s} u_{k} e_{k}(\mathbf{x}),$$

and its domain is given by

$$D((-\Delta)_{Sp}^{s}) := \{ u \in L^{2}(\Omega) \text{ such that } \|(-\Delta)_{Sp}^{s}u\|_{L^{2}(\Omega)}^{2} = \sum_{k=1}^{+\infty} \lambda_{k}^{2s}u_{k}^{2} < +\infty \}$$
$$= \mathcal{X}^{2s}(\Omega).$$

Obviously, when s = 0 we recover the identity, and when s = 1 we obtain the classical Laplacian.

#### 2.1.3 Semigroup properties

Let us consider the operator  $A^s$  on  $L^2(\Omega)$ , defined by

$$\begin{cases} D(A^s) &= \mathcal{X}^{2s}(\Omega), \\ A^s u &= -d(-\Delta)^s_{Sp} u, \ d > 0, \end{cases}$$

for any  $u \in D(A^s)$ . It is well-known that  $A^s$  generates a strongly continuous semigroup  $\{T_s(t)\}_{t\geq 0} := \{e^{tA^s}\}_{t\geq 0}$  on  $L^2(\Omega)$ . For readers who may be unfamiliar with the notion of semigroups, we suggest consulting [93, Chapter 7] or [70, Chapter 1]. Furthermore, the semigroup  $\{T_s(t)\}_{t\geq 0}$  is submarkovian (positivity-preserving and  $L^{\infty}$ -contractive), namely

$$T_s(t)u \ge 0$$
 for any  $t \ge 0$ , as  $0 \le u \in L^2(\Omega)$ 

and

$$|T_s(t)u||_{L^p(\Omega)} \le ||u||_{L^p(\Omega)}$$
 for any  $u \in L^p(\Omega) \cap L^2(\Omega), \ p \in [1, +\infty].$ 

As a consequence,  $\{T_s(t)\}_{t\geq 0}$  can be extended to a contraction semigroup on  $L^p(\Omega)$  for any  $p \in [1, +\infty]$ . Each semigroup is strongly continuous if  $p \in [1, +\infty)$  and bounded analytic if  $p \in (1, +\infty)$  (see, for instance, [32, Theorem 1.4.1] for more details). In addition,  $\{T_s(t)\}_{t\geq 0}$  is ultracontractive, in the sense that for any  $1 \leq p \leq q \leq +\infty$ , the operator  $T_s(t)$  maps  $L^p(\Omega)$  into  $L^q(\Omega)$ . In other words, for any  $u \in L^p(\Omega)$  there exists C > 0 such that

(2.2) 
$$\|T_s(t)u\|_{L^q(\Omega)} \le Ct^{-\frac{N}{2s}\left(\frac{1}{p} - \frac{1}{q}\right)} \|u\|_{L^p(\Omega)}, \quad \forall t > 0.$$

For further details and proofs regarding the properties of  $\{T_s(t)\}_{t\geq 0}$ , see [43, Chapter 2].

#### **2.1.4** Second definition of the (SFL)

The operator (SFL) can also be expressed as an integral with respect to an appropriate kernel, along with an additional linear term, see [1, Definition 1 and Lemma 38] and [2, Subsection 2.3]. More precisely, for any  $u \in \mathcal{X}^{2s}(\Omega)$  and for *a.e.*  $\mathbf{x} \in \Omega$ , we have

(2.3) 
$$(-\Delta)_{Sp}^{s} u(\mathbf{x}) = \text{P.V.} \int_{\Omega} [u(\mathbf{x}) - u(\mathbf{y})] J(\mathbf{x}, \mathbf{y}) d\mathbf{y} + \kappa(\mathbf{x}) u(\mathbf{x}),$$

where

$$J(\mathbf{x}, \mathbf{y}) := \frac{s}{\Gamma(1-s)} \int_0^{+\infty} \frac{p_{\Omega}(t, \mathbf{x}, \mathbf{y})}{t^{1+s}} dt \ge 0,$$

and  $p_{\Omega}(t, \mathbf{x}, \mathbf{y})$  is the heat kernel of the Dirichlet or Neumann Laplacian. Furthermore,

$$\kappa(\mathbf{x}) := \frac{s}{\Gamma(1-s)} \int_0^{+\infty} \left( 1 - \int_{\Omega} p_{\Omega}(t, \mathbf{x}, \mathbf{y}) d\mathbf{y} \right) \frac{dt}{t^{1+s}} \ge 0,$$

in the case of the Dirichlet Laplacian, and  $\kappa(\mathbf{x}) = 0$  in the case of the Neumann one. For more details, see [33, Subsection 1.1 and Subsection 1.2].

#### **2.2** Evolution Problem governed by the (SFL) and its dual

In this subsection, we review some existence results and key properties of solutions to the fractional evolution problem (2.4). Furthermore, we examine the dual of this problem to derive a useful Lamberton-type estimate.

#### 2.2.1 Fractional evolution problem

First, let us consider the following problem:

(2.4) 
$$\begin{cases} \partial_t w(t, \mathbf{x}) + d(-\Delta)^s_{Sp} w(t, \mathbf{x}) &= h(t, \mathbf{x}), \quad (t, \mathbf{x}) \in Q_T, \\ \mathcal{B}[w(t, \mathbf{x})] &= 0, \quad (t, \mathbf{x}) \in \Sigma_T, \\ w(0, \mathbf{x}) &= w_0(\mathbf{x}) \ge 0, \quad \mathbf{x} \in \Omega, \end{cases}$$

where d > 0,  $w_0 \in L^p(\Omega)$  and  $h \in L^p(Q_T)$  with p > 1.

**Definition 2.1** (Weak solution). Let  $h \in L^1(0,T; L^p(\Omega))$ . A function  $w \in C([0,T]; L^p(\Omega))$  is called a weak solution to Problem (2.4) if for any  $(t, \mathbf{x}) \in [0, T) \times \Omega$ ,

(2.5) 
$$w(t,\mathbf{x}) = T_s(t)w_0(\mathbf{x}) + \int_0^t T_s(t-\tau)h(\tau,\mathbf{x})d\tau.$$

**Definition 2.2** (Strong solution). A function w is called a strong solution to Problem (2.4) if : (i)  $w \in \mathcal{C}([0,T); L^p(\Omega)) \cap \mathcal{C}^1(0,T; L^p(\Omega))$ ; (ii) for any  $t \in (0,T)$ ,  $w(t, \cdot) \in \mathcal{X}^{2s}(\Omega)$ ;

(iii) Problem (2.4) is satisfied a.e. in  $Q_T$ .

The following theorem guarantees the existence of a unique weak solution.

**Theorem 2.1** ([70, Chapter 4]). Let  $h \in L^1(0,T; L^p(\Omega))$  and  $w_0 \in L^p(\Omega)$ . Then, Problem (2.4) admits a unique weak solution.

**Remark 2.1.** Let us recall that any "weak solution" taking values in  $L^{\infty}(Q_T)$  is, in fact, regular enough to be considered a "strong solution".

The regularity of the weak solution to Problem (2.4) relies on the three subsequent theorems.

1) A maximal regularity result in  $L^{\infty}$ :

**Theorem 2.2.** Let  $w_0 \in L^{\infty}(\Omega)^+$  and  $h \in L^p(Q_T)$  with p > 1. Moreover, let w be the weak solution to Problem (2.4). Then, for any  $p > \frac{N+2s}{2s}$  we have

$$||w||_{L^{\infty}(Q_T)} \le ||w_0||_{L^{\infty}(\Omega)} + C||h||_{L^p(Q_T)}, \text{ fore some } C > 0.$$

*Proof.* By applying the ultracontractivity estimate (2.2), the proof follows as in [31, Theorem 3.1].  $\Box$ 

2) A first comparison principle result :

**Theorem 2.3.** Let  $w_0 \in L^{\infty}(\Omega)^+$  and  $h \in L^p(Q_T)$  with p > 1. Further, let w be the weak solution to Problem (2.4). If moreover  $h \ge 0$  a.e. in  $Q_T$ , then  $w \ge 0$  a.e. in  $Q_T$ .

*Proof.* Using the definition (2.3), the proof proceeds similarly to [38, Proposition 3.18].

3) A second comparison principle result :

**Theorem 2.4.** Let  $w_0 \in L^{\infty}(\Omega)^+$ ,  $h \in L^p(Q_T)$  with p > 1, and w be the weak solution to Problem (2.4). In addition, let z be the weak solution to

(2.6) 
$$\begin{cases} \partial_t z(t, \mathbf{x}) + d(-\Delta)^s_{Sp} z(t, \mathbf{x}) &= g(t, \mathbf{x}), \quad (t, \mathbf{x}) \in Q_T, \\ \mathcal{B}[z(t, \mathbf{x})] &= 0, \quad (t, \mathbf{x}) \in \Sigma_T, \\ z(0, \mathbf{x}) &= z_0(\mathbf{x}) \ge 0, \quad \mathbf{x} \in \Omega, \end{cases}$$

where  $z_0 \in L^{\infty}(\Omega)$  and  $g \in L^q(Q_T)$  for some q > 1. Suppose that  $h \leq g$  and  $w_0 \leq z_0$  a.e. in  $Q_T$ and  $\Omega$  respectively. Then  $w \leq z$  a.e. in  $\Omega$  for  $t \in [0, T)$ .

*Proof.* Let us consider the following problem

(2.7) 
$$\begin{cases} \partial_t (z-w)(t,\mathbf{x}) + d(-\Delta)^s_{Sp}(z-w)(t,\mathbf{x}) &= (g-h)(t,\mathbf{x}), \quad (t,\mathbf{x}) \in Q_T, \\ \mathcal{B}[(z-w)(t,\mathbf{x})] &= 0, \quad (t,\mathbf{x}) \in \Sigma_T, \\ (z-w)(0,\mathbf{x}) &= (z_0-w_0)(\mathbf{x}), \quad \mathbf{x} \in \Omega. \end{cases}$$

To achieve the proof, we apply Theorem 2.3 to z - w.

#### **2.2.2 Dual of Problem** (2.4)

The dual of Problem (2.4) is given by

$$(P_{\phi,T}) \begin{cases} -\partial_t \mathcal{V}(t,\mathbf{x}) + d(-\Delta)^s_{Sp} \mathcal{V}(t,\mathbf{x}) &= \phi(t,\mathbf{x}), \quad (t,\mathbf{x}) \in Q_T, \\ \mathcal{B}[\mathcal{V}(t,\mathbf{x})] &= 0, \qquad (t,\mathbf{x}) \in \Sigma_T, \\ \mathcal{V}(T,\mathbf{x}) &= 0, \qquad \mathbf{x} \in \Omega, \end{cases}$$

where  $\phi$  is a regular function.

The following theorem outlines the conditions for the existence of a solution to Problem  $(P_{\phi,T})$ , and ensures it meets a Lamberton-type estimation in  $L^p$ .

**Theorem 2.5.** Assume that  $\phi \in L^p(Q_T)$  with p > 1. Then, Problem  $(P_{\phi,T})$  admits a unique weak solution. Moreover, there exists a constant C := C(p,T) > 0 such that

(2.8) 
$$\|\partial_t \mathcal{V}\|_{L^p(Q_T)} + \|\mathcal{V}\|_{L^p(Q_T)} + d \|(-\Delta)^s \mathcal{V}\|_{L^p(Q_T)} + \|\mathcal{V}_0\|_{L^p(\Omega)} \le C \|\phi\|_{L^p(Q_T)},$$

where  $\mathcal{V}_0 := \mathcal{V}(0, \cdot)$ .

*Proof.* By leveraging the properties of the semigroup  $\{T_s(t)\}_{t\geq 0}$ , the proof can be constructed by following the steps of the proof of [31, Theorem 3.3], relying on [61, Theorem 1].

r	-	-	-	
L				
L				
L,	_	_	_	

## 3 Global existence for reaction-diffusion systems governed by spectral fractional Laplacians

Let us recall our main system:

(S) 
$$\begin{cases} \forall i = 1, \dots, m, \\ \partial_t u_i(t, \mathbf{x}) + d_i(-\Delta)_{Sp}^{s_i} u_i(t, \mathbf{x}) &= f_i(u_1(t, \mathbf{x}), \dots, u_m(t, \mathbf{x})), \quad (t, \mathbf{x}) \in Q_T, \\ \mathcal{B}[u_i(t, \mathbf{x})] &= 0, \qquad (t, \mathbf{x}) \in \Sigma_T, \\ u_i(0, \mathbf{x}) &= u_{0i}(\mathbf{x}), \qquad \mathbf{x} \in \Omega, \end{cases}$$

where  $m \ge 2$  and for any  $i \in [1, m]$ ,  $0 < s_i < 1$ ,  $d_i > 0$  and  $f_i$  is locally Lipschitz continuous.

As stated in the introduction, our main contribution is to extend two fundamental results from the classical case to the fractional setting. This is the aim of this section. For clarity, it is structured into four subsections. In the first one, we establish the local existence. In the second, we prove a result that enables the extension of Pierre's duality lemma to the fractional case with different orders, and we subsequently demonstrate this extension. Finally, Subsections 4.2 and 4.3 are respectively devoted to the proofs of our main global existence theorems.

#### 3.1 Local existence

The local existence lemma is well-established. Here, we will recall its statement and omit the proof, as it is straightforward.

**Lemma 3.1.** Let  $(u_{01}, \ldots, u_{0m}) \in (L^{\infty}(\Omega))^m$ . Assume that the  $f_i$ 's are locally Lipschitz continuous. Then, there exists  $T_{\max} > 0$  and  $(\varphi_1, \ldots, \varphi_m) \in \mathcal{C}([0, T_{\max}), [0, +\infty)^m)$  such that : (i) System (S) admits a unique nonnegative strong solution  $(u_1, \ldots, u_m)$  in  $Q_{T_{\max}}$ ; (ii) for any  $i \in [\![1, m]\!]$ 

(3.1) 
$$\|u_i(t,\cdot)\|_{L^{\infty}(\Omega)} \leq \varphi_i(t) \text{ for any } t \in (0,T_{\max});$$

(iii) if  $T_{\max} < +\infty$ , then  $\lim_{t \nearrow T_{\max}} \sum_{i=1}^{m} \|u_i(t, \cdot)\|_{L^{\infty}(\Omega)} = +\infty$ ; (iv) if, in addition, the  $f_i$ 's satisfy (**P**), then

$$\left(\forall i \in \llbracket 1, m \rrbracket, \ u_{0i}(.) \ge 0\right) \Longrightarrow \left(\forall i \in \llbracket 1, m \rrbracket, \ u_i(t, .) \ge 0 \ \forall t \in [0, T_{\max})\right).$$

**Remark 3.1.** In light of (iii), to prove global existence of strong solutions to System (S), it suffices to establish an a priori estimate of the form:

(3.2) 
$$\forall t \in [0, T_{\max}), \ \sum_{i=1}^{m} \|u_i(t, \cdot)\|_{L^{\infty}(\Omega)} < \psi(t),$$

where  $\psi : [0, +\infty) \to [0, +\infty)$  is a continuous function.

**Remark 3.2.** It turns out that establishing an estimate like (3.2) is far from straightforward, except in the trivial case  $d_1 = \cdots = d_m$ ,  $s_1 = \cdots = s_m$ , and the  $f_i$ 's fulfill (**P**). For this particular instance, System (S) admits a global strong solution regardless of the growth of the  $f_i$ 's. The proof can be directly constructed following [31, Subsection 4.1]. Nonetheless, the situation becomes considerably more complicated when the diffusion coefficients differ. Additional assumptions on the growth of the source terms are required to guarantee the global existence of strong solutions, even in the case  $s_1 = \cdots = s_m = 1$ . Moreover, the complexity increases further when different fractional orders are taken into account.

#### 3.2 Fractional version of Pierre's duality lemma

To prove our global existence theorems, we will rely heavily on the following lemma. Let us mention that when the fractional powers tend to 1, the following result is essentially a generalization of Pierre's duality lemma (see [24, 35, 50, 55, 56, 72] and references therein).

**Lemma 3.2.** Let T > 0 and  $s_1, s_2 \in (0, 1)$  with  $s_1 \leq s_2$ . Consider the following  $2 \times 2$  system :

$$(S_{2\times 2}) \qquad \begin{cases} \partial_t u_1(t, \mathbf{x}) + d_1(-\Delta)_{Sp}^{s_1} u_1(t, \mathbf{x}) &= f_1(u_1(t, \mathbf{x}), u_2(t, \mathbf{x})), \quad (t, \mathbf{x}) \in Q_T, \\ \partial_t u_2(t, \mathbf{x}) + d_2(-\Delta)_{Sp}^{s_2} u_2(t, \mathbf{x}) &= f_2(u_1(t, \mathbf{x}), u_2(t, \mathbf{x})), \quad (t, \mathbf{x}) \in Q_T, \\ \mathcal{B}[u_1(t, \mathbf{x})] = \mathcal{B}[u_2(t, \mathbf{x})] &= 0, \qquad (t, \mathbf{x}) \in \Sigma_T, \\ u_1(0, \mathbf{x}) &= u_{01}(\mathbf{x}), \qquad \mathbf{x} \in \Omega, \\ u_2(0, \mathbf{x}) &= u_{02}(\mathbf{x}), \qquad \mathbf{x} \in \Omega, \end{cases}$$

where  $f_1$  and  $f_2$  satisfy  $(\mathbf{P}) + (\mathbf{M})$ . Let  $(u_1, u_2)$  be the strong solution to System  $(S_{2\times 2})$  in  $Q_T$ . Then, for any  $p \in (1, +\infty)$ , there exists a constant  $\lambda > 0$  such that

(3.3) 
$$\|u_j\|_{L^p(Q_T)} \le \lambda \left(1 + \|u_i\|_{L^p(Q_T)}\right),$$

where  $i, j \in \{1, 2\}$  with  $i \neq j$  and  $s_i \leq s_j$ .

Before giving the proof of Lemma 3.2, let us first state and prove a useful estimate that will be employed in the subsequent.

**Proposition 3.1.** Let  $p \in (1, +\infty)$  and  $s_1, s_2 \in (0, 1)$  with  $s_1 \leq s_2$ . We have

(3.4) 
$$\|(-\Delta)_{Sp}^{s_1}u\|_{L^p(\Omega)} \le \|(-\Delta)_{Sp}^{s_2}u\|_{L^p(\Omega)}^{\frac{s_1}{s_2}} \|u\|_{L^p(\Omega)}^{\frac{s_2-s_1}{s_2}},$$

for any  $u \in \mathcal{X}^{2s_2}(\Omega) \cap L^p(\Omega)$  such that  $(-\Delta)_S^{s_2} u \in L^p(\Omega)$ .

*Proof.* To start, the case  $s_1 = s_2$  is trivial. Now, let us assume that  $s_1 < s_2$ . For any  $\mathbf{x} \in \Omega$ , we have

$$\left| (-\Delta)_{Sp}^{s_1} u(\mathbf{x}) \right| = \left| \sum_{k=1}^{+\infty} \lambda_k^{s_1} u_k e_k(\mathbf{x}) \right| \le \sum_{k=1}^{+\infty} \left| \lambda_k^{s_1} u_k^{\frac{s_1}{s_2}} u_k^{\frac{s_2-s_1}{s_2}} (e_k(\mathbf{x}))^{\frac{s_1}{s_2}} (e_k(\mathbf{x}))^{\frac{s_2-s_1}{s_2}} \right|.$$

Now we use Hölder's inequality for sums by taking the exponents  $a = \frac{s_2}{s_1}$  and  $b = \frac{a}{a-1} = \frac{s_2}{s_2-s_1}$ . We get

$$\begin{aligned} \left| (-\Delta)_{Sp}^{s_1} u(\mathbf{x}) \right| &\leq \left( \sum_{k=1}^{+\infty} \left| \lambda_k^{s_1} u_k^{\frac{s_1}{s_2}} (e_k(\mathbf{x}))^{\frac{s_1}{s_2}} \right|^{\frac{s_2}{s_1}} \right)^{\frac{s_1}{s_2}} \left( \sum_{k=1}^{+\infty} \left| u_k^{\frac{s_2-s_1}{s_2}} (e_k(\mathbf{x}))^{\frac{s_2-s_1}{s_2}} \right|^{\frac{s_2-s_1}{s_2}} \right)^{\frac{s_2-s_1}{s_2}} \\ &\leq \left( \sum_{k=1}^{+\infty} \left| \lambda_k^{s_2} u_k e_k(\mathbf{x}) \right| \right)^{\frac{s_1}{s_2}} \left( \sum_{k=1}^{+\infty} \left| u_k e_k(\mathbf{x}) \right| \right)^{\frac{s_2-s_1}{s_2}} \\ &\leq \left| \sum_{k=1}^{+\infty} \lambda_k^{s_2} u_k e_k(\mathbf{x}) \right|^{\frac{s_1}{s_2}} \left| \sum_{k=1}^{+\infty} u_k e_k(\mathbf{x}) \right|^{\frac{s_2-s_1}{s_2}} = \left| (-\Delta)_{Sp}^{s_2} u(\mathbf{x}) \right|^{\frac{s_1}{s_2}} \times \left| u(\mathbf{x}) \right|^{\frac{s_2-s_1}{s_2}} \end{aligned}$$

as  $\frac{s_1}{s_2} < 1$  and  $\frac{s_2-s_1}{s_2} < 1$ . Therefore, by taking the p power and integrating over  $\Omega$  we obtain

$$\begin{split} \int_{\Omega} \left| (-\Delta)_{Sp}^{s_1} u(\mathbf{x}) \right|^p d\mathbf{x} &\leq \int_{\Omega} \left| (-\Delta)_{Sp}^{s_2} u(\mathbf{x}) \right|^{\frac{ps_1}{s_2}} \times |u|^{\frac{p(s_2-s_1)}{s_2}} d\mathbf{x} \\ &\leq \left( \int_{\Omega} \left| (-\Delta)_{Sp}^{s_2} u(\mathbf{x}) \right|^p d\mathbf{x} \right)^{\frac{s_1}{s_2}} \times \left( \int_{\Omega} |u(\mathbf{x})|^p d\mathbf{x} \right)^{\frac{s_2-s_1}{s_2}}, \end{split}$$

where the last inequality is obtained by using Hölder's inequality for integrals with the same exponents a and b above. Finally,

$$\|(-\Delta)_{Sp}^{s_1}u\|_{L^p(\Omega)} \le \|(-\Delta)_{Sp}^{s_2}u\|_{L^p(\Omega)}^{\frac{s_1}{s_2}} \|u\|_{L^p(\Omega)}^{\frac{s_2-s_1}{s_2}}.$$

Proof of Lemma 3.2. Let  $T \in (0, T_{\max})$  and  $(t, \mathbf{x}) \in (0, T] \times \Omega$ . Furthermore, let  $i, j \in \{1, 2\}$  with  $i \neq j$  and assume that  $0 < s_i \leq s_j < 1$ . By hypothesis,  $f_i$  and  $f_j$  satisfy (**M**), then there exists  $a_i, a_j \in (0, +\infty)$  such that  $a_i f_i(t, \mathbf{x}) + a_j f_j(t, \mathbf{x}) \leq 0$ . By multiplying the *i*-th (resp. the *j*-th) equation of System  $(S_{2\times 2})$  by  $a_i$  (resp.  $a_j$ ) and summing the two equations, we get for any  $(t, \mathbf{x}) \in Q_T$ 

(3.5) 
$$\partial_t (a_i u_i(t, \mathbf{x}) + a_j u_j(t, \mathbf{x})) + a_i d_i (-\Delta)^{s_i}_{Sp} u_i(t, \mathbf{x}) + a_j d_j (-\Delta)^{s_j}_{Sp} u_j(t, \mathbf{x}) \le 0.$$

Now, let  $\phi$  be a nonnegative regular function and let  $\mathcal{V}$  be the solution to Problem  $(P_{\phi,T})$  with  $s = s_j$  and  $d = d_j$ . First, let us multiply (3.5) by  $\mathcal{V}$  and integrate over  $Q_T$ . Then, we obtain

$$(3.6) \begin{cases} -\iint_{Q_T} (a_i u_i(t, \mathbf{x}) + a_j u_j(t, \mathbf{x})) \partial_t \mathcal{V}(t, \mathbf{x}) d\mathbf{x} dt + a_j d_j \iint_{Q_T} u_j(t, \mathbf{x}) (-\Delta)_{Sp}^{s_j} \mathcal{V}(t, \mathbf{x}) d\mathbf{x} dt \\ \leq \int_{\Omega} (a_i u_{0i} + a_j u_{0j})(\mathbf{x}) \mathcal{V}_0(\mathbf{x}) d\mathbf{x} - a_i d_i \iint_{Q_T} u_i(t, \mathbf{x}) (-\Delta)_{Sp}^{s_i} \mathcal{V}(t, \mathbf{x}) d\mathbf{x} dt. \end{cases}$$

Then, by combining the terms, we get

$$(3.7) \begin{cases} a_j \iint_{Q_T} u_j(t, \mathbf{x}) \Big[ -\partial_t \mathcal{V}(t, \mathbf{x}) + d_j(-\Delta)_{Sp}^{s_j} \mathcal{V}(t, \mathbf{x}) \Big] d\mathbf{x} dt - a_i \iint_{Q_T} u_i(t, \mathbf{x}) \partial_t \mathcal{V}(t, \mathbf{x}) d\mathbf{x} dt \\ \leq \int_{\Omega} (a_i u_{0i} + a_j u_{0j})(\mathbf{x}) \mathcal{V}_0(\mathbf{x}) d\mathbf{x} - a_i d_i \iint_{Q_T} u_i(t, \mathbf{x}) (-\Delta)_{Sp}^{s_i} \mathcal{V}(t, \mathbf{x}) d\mathbf{x} dt. \end{cases}$$

Hence,

$$(3.8) \begin{cases} a_j \iint_{Q_T} u_j(t, \mathbf{x}) \phi(t, \mathbf{x}) d\mathbf{x} dt \\ \leq \int_{\Omega} (a_1 u_{01} + a_j u_{0j})(\mathbf{x}) \mathcal{V}_0(\mathbf{x}) d\mathbf{x} + a_i \iint_{Q_T} u_i(t, \mathbf{x}) \left[ \partial_t \mathcal{V}(t, \mathbf{x}) - d_i (-\Delta)_{Sp}^{s_i} \mathcal{V}(t, \mathbf{x}) \right] d\mathbf{x} dt. \end{cases}$$

Now, let p > 1. By applying Hölder's inequality and using (2.8), we have

(3.9) 
$$\int_{\Omega} (a_i u_{0i} + a_j u_{0j})(\mathbf{x}) \mathcal{V}_0(\mathbf{x}) d\mathbf{x} \le a_0 C \|u_{0i} + u_{0j}\|_{L^p(\Omega)} \|\phi\|_{L^{p'}(Q_T)},$$

where  $a_0 := \max\{a_1, a_2\}$  and  $p' := \frac{p}{p-1}$ . Moreover, (3.10)

$$a_{i} \iint_{Q_{T}} u_{i}(t,\mathbf{x}) \left[ \partial_{t} \mathcal{V}(t,\mathbf{x}) - d_{i}(-\Delta)^{s_{i}}_{Sp} \mathcal{V}(t,\mathbf{x}) \right] d\mathbf{x} dt \leq a_{i} \|u_{i}\|_{L^{p}(Q_{T})} \left( C \|\phi\|_{L^{p'}(Q_{T})} + d_{i} \|(-\Delta)^{s_{i}}_{Sp} \mathcal{V}\|_{L^{p'}(Q_{T})} \right).$$

As  $s_i \leq s_j$ , thanks to the estimate (3.4), we get

$$\|(-\Delta)_{Sp}^{s_i}\mathcal{V}\|_{L^{p'}(Q_T)} \le \|(-\Delta)_{Sp}^{s_j}\mathcal{V}\|_{L^{p'}(Q_T)}^{\frac{s_i}{s_j}} \|\mathcal{V}\|_{L^{p'}(Q_T)}^{\frac{s_j-s_i}{s_j}} \le C \|\phi\|_{L^{p'}(Q_T)}^{\frac{s_i}{s_j}} \|\phi\|_{L^{p'}(Q_T)}^{\frac{s_j-s_i}{s_j}} = C \|\phi\|_{L^{p'}(Q_T)},$$

$$a_{j} \iint_{Q_{T}} u_{j}(t,\mathbf{x})\phi(t,\mathbf{x})d\mathbf{x}dt \leq C \Big[a_{0} \|u_{0i} + u_{0j}\|_{L^{p}(\Omega)} \|\phi\|_{L^{p'}(Q_{T})} + a_{i}(1+d_{i})\|u_{i}\|_{L^{p}(Q_{T})} \|\phi\|_{L^{p'}(Q_{T})}\Big].$$

Lastly, we deduce by duality that

$$\|u_j\|_{L^p(Q_T)} \le \frac{C}{a_j} \left( a_0 \|u_{0i} + u_{0j}\|_{L^p(\Omega)} + a_i \left(1 + d_i\right) \|u_i\|_{L^p(Q_T)} \right),$$

which implies

$$\|u_j\|_{L^p(Q_T)} \le \frac{C}{a_j} \max\left\{a_0 \|u_{0i} + u_{0j}\|_{L^p(\Omega)}, a_i(1+d_i)\right\} \left(1 + \|u_i\|_{L^p(Q_T)}\right).$$

Accordingly, we get the desired estimation.

#### **3.3** Global existence : case of reversible chemical reaction with three species

In this section, we will state and prove our first global existence related to the following system:

$$(S_{\alpha,\beta,\gamma}) \begin{cases} \forall i = 1, 2, 3, \\ \partial_t u_i(t, \mathbf{x}) + d_i(-\Delta)_{Sp}^{s_i} u_i(t, \mathbf{x}) &= f_i(u_1(t, \mathbf{x}), u_2(t, \mathbf{x}), u_3(t, \mathbf{x})), & (t, \mathbf{x}) \in Q_T, \\ \mathcal{B}[u_i(t, \mathbf{x})] &= 0, & (t, \mathbf{x}) \in \Sigma_T, \\ u_i(0, \mathbf{x}) &= u_{0i}(\mathbf{x}), & \mathbf{x} \in \Omega, \end{cases}$$

where  $0 < s_i < 1$ ,  $f_1 = \alpha g$ ,  $f_2 = \beta g$ ,  $f_3 = -\gamma g$  with  $g = u_3^{\gamma} - u_1^{\alpha} u_2^{\beta}$  and  $1 \le \alpha, \beta, \gamma < +\infty$ . Let us mention that for  $(r_1, r_2, r_3) \in [0, +\infty)^3$ , we have

$$f_1(0, r_2, r_3) = \alpha r_3^{\gamma} \ge 0, \quad f_2(r_1, 0, r_3) = \beta r_3^{\gamma} \ge 0 \quad \text{et } f_3(r_1, r_2, 0) = \gamma r_1^{\alpha} r_2^{\beta} \ge 0$$

and

$$\beta\gamma f_1(r_1, r_2, r_3) + \alpha\gamma f_2(r_1, r_2, r_3) + 2\alpha\beta f_3(r_1, r_2, r_3) = 0.$$

Then, System  $(S_{\alpha,\beta,\gamma})$  fulfill (**P**) and (**M**).

Our first result of global existence is as follows:

**Theorem 3.1.** For any  $i \in [\![1,3]\!]$ , let  $s_i \in (0,1)$  and  $u_{0i} \in L^{\infty}(\Omega)^+$ . Moreover, let  $(\alpha, \beta, \gamma) \in [1, +\infty)^3$ . Then, System  $(S_{\alpha,\beta,\gamma})$  admits a unique nonnegative global strong solution if one the following assumptions holds:

(i)  $s_3 \leq \min\{s_1, s_2\}$  and  $\gamma > \alpha + \beta$ ;

(ii)  $s_3 \ge \max\{s_1, s_2\}$  and  $(\alpha, \beta, \gamma) \in [1, +\infty)^2 \times \{1\}.$ 

**Comment.** It is noteworthy that the global existence of strong solutions when  $2 < \gamma \leq \alpha + \beta$  has remained an open question since 2011, even in the case of the classical Laplacian, as discussed in [56]. This is particularly surprising from a chemical perspective, as the reaction is expected to be reversible, suggesting that a strong solution should exist globally, similar to the case  $\alpha + \beta < \gamma$ . More broadly, this question remains largely unresolved. For instance, the global existence of a strong solution to a system modeling a reversible reaction of the form  $\alpha U_1 + \beta U_2 \rightleftharpoons \gamma U_3 + \delta U_4$  has only recently been established, and solely in the specific case  $\alpha = \beta = \gamma = \delta = 1$ , see [25, 36, 37, 86].

*Proof.* By Lemma 3.1, System  $(S_{\alpha,\beta,\gamma})$  admits a unique strong solution  $(u_1, u_2, u_3)$  in  $Q_{T_{\text{max}}}$ . It remains to prove that this solution is global, *i.e.*  $T_{\text{max}} = +\infty$ . We closely follow the proofs of [56, Theorems 1 and 3] and [31, Theorem 4.1]. Now, for any  $T \in (0, T_{\text{max}})$  and  $t \in (0, T]$ , we proceed to prove (i) and (ii).

(i) Let  $s_3 \leq \min\{s_1, s_2\}$  and  $(\alpha, \beta, \gamma) \in [1, +\infty)^3$  such that  $\gamma > \alpha + \beta$ . Moreover, let us consider the two following problems

$$(P_{\gamma,1}) \qquad \begin{cases} \partial_t v(t,\mathbf{x}) + d_1(-\Delta)_{Sp}^{s_1} v(t,\mathbf{x}) &= (u_3(t,\mathbf{x}))^{\gamma}, \quad (t,\mathbf{x}) \in Q_T, \\ \mathcal{B}[v(t,\mathbf{x})] &= 0, \qquad (t,\mathbf{x}) \in \Sigma_T, \\ v(0,\mathbf{x}) &= u_{01}(\mathbf{x}), \qquad \mathbf{x} \in \Omega, \end{cases}$$

and

$$(P_{\gamma,2}) \qquad \begin{cases} \partial_t z(t,\mathbf{x}) + d_2(-\Delta)^{s_2}_{Sp} z(t,\mathbf{x}) &= (u_3(t,\mathbf{x}))^{\gamma}, \quad (t,\mathbf{x}) \in Q_T, \\ \mathcal{B}[z(t,\mathbf{x})] &= 0, \qquad (t,\mathbf{x}) \in \Sigma_T, \\ z(0,\mathbf{x}) &= u_{02}(\mathbf{x}), \qquad \mathbf{x} \in \Omega. \end{cases}$$

As  $u_1, u_2$  and  $u_3$  are nonnegative, then the right hand sides of the equations for i = 1, 2 of System  $(S_{\alpha,\beta,\gamma})$  fulfill

$$f_1(u_1(t, \mathbf{x}), u_2(t, \mathbf{x}), u_3(t, \mathbf{x})) \le (u_3(t, \mathbf{x}))^{\gamma}$$

and

$$f_2(u_1(t, \mathbf{x}), u_2(t, \mathbf{x}), u_3(t, \mathbf{x})) \le (u_3(t, \mathbf{x}))^{\gamma}.$$

Therefore, using Theorem 2.4, we have for  $a.e. \mathbf{x} \in \Omega$  and for  $t \in [0, T)$ ,  $u_1(t, \mathbf{x}) \leq v(t, \mathbf{x})$  and  $u_2(t, \mathbf{x}) \leq z(t, \mathbf{x})$ . Hence, to establish global existence, it is enough to prove that  $u_3 \in L^p(Q_T)$  for sufficiently large p.

Now, let q > 1. Then, let us multiply the equation for i = 3 of System  $(S_{\alpha,\beta,\gamma})$  by  $(u_3(t,\mathbf{x}))^q$ , and then integrate over  $Q_T$ . We obtain (3.12)

$$\begin{cases} \frac{1}{q+1} \int_{\Omega} (u_3(T, \mathbf{x}))^{q+1} d\mathbf{x} + d_3 \iint_{Q_T} \left[ (-\Delta)^{s_3}_{Sp} u_3(t, \mathbf{x}) \right] (u_3(t, \mathbf{x}))^q d\mathbf{x} dt + \gamma \iint_{Q_T} (u_3(t, \mathbf{x}))^{q+\gamma} d\mathbf{x} dt \\ = \gamma \iint_{Q_T} (u_1(t, \mathbf{x}))^{\alpha} (u_2(t, \mathbf{x}))^{\beta} (u_3(t, \mathbf{x}))^q d\mathbf{x} dt + \frac{1}{q+1} \int_{\Omega} (u_{03}(\mathbf{x}))^{q+1} d\mathbf{x}. \end{cases}$$

Thanks to Stroock-Varopoulos inequality (see [16, Lemma 6.2]), we have (3.13)

$$\iint_{Q_T} \left[ (-\Delta)_{Sp}^{s_3} u_3(t, \mathbf{x}) \right] (u_3(t, \mathbf{x}))^q d\mathbf{x} dt \ge \frac{4q}{(q+1)^2} \iint_{Q_T} \left| (-\Delta)_{Sp}^{\frac{s_3}{2}} (u_3(t, \mathbf{x}))^{\frac{q+1}{2}} \right|^2 d\mathbf{x} dt \ge 0.$$

In addition, using Hölder's inequality, we get

$$(3.14) \quad \iint_{Q_T} (u_1(t,\mathbf{x}))^{\alpha} (u_2(t,\mathbf{x}))^{\beta} (u_3(t,\mathbf{x}))^q d\mathbf{x} dt \le \|u_1\|_{L^{\alpha\sigma_1}(Q_T)}^{\alpha} \|u_2\|_{L^{\beta\sigma_2}(Q_T)}^{\beta} \|u_3\|_{L^{\gamma+q}(Q_T)}^q,$$

where

$$\frac{1}{\sigma_1} + \frac{1}{\sigma_2} + \frac{q}{q+\gamma} = 1.$$

Then, we may choose  $(\sigma_1, \sigma_2)$  such that  $\sigma_1 \alpha \leq q + \gamma$  and  $\sigma_2 \beta \leq q + \gamma$ . This choice is valid, as by hypothesis  $\gamma > \alpha + \beta$ . Therefore, there exists  $C_1 > 0$  such that (3.15)

$$\iint_{Q_T} (u_1(t,\mathbf{x}))^{\alpha} (u_2(t,\mathbf{x}))^{\beta} (u_3(t,\mathbf{x}))^q d\mathbf{x} dt \le C_1 \|u_1\|_{L^{\gamma+q}(Q_T)}^{\alpha} \|u_2\|_{L^{\gamma+q}(Q_T)}^{\beta} \|u_3\|_{L^{\gamma+q}(Q_T)}^q.$$

Since  $s_3 \leq \min\{s_1, s_2\}$ , in accordance with Lemma 3.2, we get

(3.16) 
$$\|u_1\|_{L^{\gamma+q}(Q_T)} \le C_2 \left(1 + \|u_3\|_{L^{\gamma+q}(Q_T)}\right)$$

and

(3.17) 
$$\|u_2\|_{L^{\gamma+q}(Q_T)} \le C_3 \left(1 + \|u_3\|_{L^{\gamma+q}(Q_T)}\right).$$

Now, using the last two estimates (3.16) and (3.17), we can rewrite (3.15) as

(3.18) 
$$\iint_{Q_T} (u_1(t,\mathbf{x}))^{\alpha} (u_2(t,\mathbf{x}))^{\beta} (u_3(t,\mathbf{x}))^q d\mathbf{x} dt \le C_4 \left(1 + \|u_3\|_{L^{\gamma+q}(Q_T)}\right)^{q+\alpha+\beta}.$$

If  $||u_3||_{L^{\gamma+q}(Q_T)} \leq 1$ , the proof is complete. If not, further calculations are required. Indeed, we have

(3.19) 
$$\iint_{Q_T} (u_1(t, \mathbf{x}))^{\alpha} (u_2(t, \mathbf{x}))^{\beta} (u_3(t, \mathbf{x}))^q d\mathbf{x} dt \le C_5 \|u_3\|_{L^{\gamma+q}(Q_T)}^{q+\alpha+\beta}.$$

Then, thanks to (3.12), we get

(3.20) 
$$\iint_{Q_T} (u_3(t, \mathbf{x}))^{q+\gamma} d\mathbf{x} dt \le C_6 \|u_3\|_{L^{\gamma+q}(Q_T)}^{q+\alpha+\beta} + \frac{1}{q+1} \int_{\Omega} (u_{03}(\mathbf{x}))^{q+1} d\mathbf{x}$$

In the subsequent, we will apply the Young's inequality  $ab \leq \varepsilon a^{\lambda} + C(\varepsilon)b^{\mu}$ , where  $\frac{1}{\lambda} + \frac{1}{\mu} = 1$ and  $\varepsilon$  is a small positive number. As  $q + \gamma > q + \alpha + \beta$ , then by choosing  $\lambda := \frac{q+\gamma}{q+\alpha+\beta}$ , Young's inequality yields

(3.21) 
$$(1-\varepsilon) \times \iint_{Q_T} (u_3(t,\mathbf{x}))^{q+\gamma} d\mathbf{x} dt \le \frac{1}{q+1} \int_{\Omega} (u_{03}(\mathbf{x}))^{q+1} d\mathbf{x} + C_7$$

Thus, for any  $\varepsilon \in (0, 1)$ , we have

(3.22) 
$$||u_3||_{L^{\gamma+q}(Q_T)} \le C_8.$$

Let us choose  $\frac{\gamma + q}{\gamma_3} > \max\left\{\frac{N + 2s_1}{2s_1}, \frac{N + 2s_2}{2s_2}\right\}$ . Returning to Problems  $(P_{\gamma,1})$  et  $(P_{\gamma,2})$ , by applying Theorem 2.2 we obtain

$$(3.23) ||u_1||_{L^{\infty}(Q_T)} \le C_9$$

and

$$(3.24) ||u_2||_{L^{\infty}(Q_T)} \le C_{10}$$

Now, let us go back to the equation for i = 3 of System  $(S_{\alpha,\beta,\gamma})$ . Thanks to the estimates (3.23) and (3.24), we conclude that

$$(3.25) ||u_3||_{L^{\infty}(Q_T)} \le C_{11}.$$

Hence, we get  $T_{\max} = +\infty$ .

(ii) Let  $s_3 \ge \max\{s_1, s_2\}$  and  $(\alpha, \beta, \gamma) \in [1, +\infty)^2 \times \{1\}$ . Going pack to Problem  $(P_{\gamma,1})$ , we have for any  $(t, \mathbf{x}) \in Q_T$ ,

(3.26) 
$$v(t, \mathbf{x}) = T_{s_1}(t)u_{01}(\mathbf{x}) + \int_0^t T_{s_1}(t-\tau)u_3(\tau, \mathbf{x})d\tau.$$

By taking the  $L^p$ -norm and using the fact that  $\{T_{s_1}(t)\}_{t\geq 0}$  is a contraction semigroup on  $L^p(\Omega)$ , we obtain

(3.27) 
$$\begin{cases} \|v(t,\cdot)\|_{L^{p}(\Omega)} \leq \|T_{s_{1}}(t)u_{01}\|_{L^{p}(\Omega)} + \int_{0}^{t} \|T_{s_{1}}(t-\tau)u_{3}(\tau,\cdot)\|_{L^{p}(\Omega)}d\tau. \\ \leq \|u_{01}\|_{L^{p}(\Omega)} + \int_{0}^{t} \|u_{3}(\tau,\cdot)\|_{L^{p}(\Omega)}d\tau. \end{cases}$$

As for a.e.  $\mathbf{x} \in \Omega$  and for any  $t \ge 0$ , we have  $u_1(t, \mathbf{x}) \le v(t, \mathbf{x})$ , we deduce that

(3.28) 
$$\|u_1(t,\cdot)\|_{L^p(\Omega)} \le \|u_{01}\|_{L^p(\Omega)} + \int_0^t \|u_3(\tau,\cdot)\|_{L^p(\Omega)} d\tau$$

By applying Hölder's inequality, we have

(3.29) 
$$\int_0^t \|u_3(\tau,\cdot)\|_{L^p(\Omega)} d\tau \le t^{\frac{1}{p'}} \left(\int_0^t \int_\Omega (u_3(\tau,\mathbf{x}))^p d\tau d\mathbf{x}\right)^{\frac{1}{p}},$$

where  $p' = \frac{p}{p-1}$ . Moreover, by using Lemma 3.2 and the fact that  $s_3 \ge s_1$ , we get

(3.30) 
$$\int_{0}^{t} \|u_{3}(\tau, \cdot)\|_{L^{p}(\Omega)} d\tau \leq C_{12} t^{\frac{1}{p'}} \left[ 1 + \left( \int_{0}^{t} \int_{\Omega} (u_{1}(\tau, \mathbf{x}))^{p} d\tau d\mathbf{x} \right)^{\frac{1}{p}} \right]$$

Now, for any  $t \in (0,T]$ , let us denote  $g(t) := \int_{\Omega} |u_1(t,\mathbf{x})|^p d\mathbf{x}$ . Thus, we can rewrite (3.28) as

(3.31) 
$$g(t)^{\frac{1}{p}} \le C_{13} \left[ 1 + \left( \int_0^t \psi(\tau) d\tau \right)^{\frac{1}{p}} \right]$$

This yields

(3.32) 
$$g(t) \le C_{14} \left[ 1 + \int_0^t \psi(\tau) d\tau \right].$$

As a result of using Gronwall's Lemma, we determine that

$$(3.33) ||u_1||_{L^p(Q_T)} \le C_{15}.$$

By applying the same reasoning to  $u_2$  in place of  $u_1$  and using the fact that  $s_3 \ge s_2$ , we derive that

$$(3.34) ||u_2||_{L^p(Q_T)} \le C_{16}.$$

Thanks to the estimates (3.33) and (3.34), we may choose  $q > \frac{N+2s_3}{2s_3}$  such that

$$\|u_1^{\alpha}u_2^{\beta}\|_{L^q(Q_T)} \le C_{17}.$$

Now, let us go back to the equation for i = 3 of System  $(S_{\alpha,\beta,\gamma})$ . By using Theorem 2.2, we obtain

$$(3.36) ||u_3||_{L^{\infty}(Q_T)} \le C_{18}.$$

Finally  $T_{\text{max}} = +\infty$ , which achieves the proof.

### 3.4 Global existence : case of triangular structure

Our second result on global existence pertains to System (S) with  $m \ge 2$ , under an additional assumption on the source terms; referred to as the *triangular structure*.

**Theorem 3.2.** For any  $i \in [\![1,m]\!]$ , with  $m \geq 2$ , let  $u_{0i} \in L^{\infty}(\Omega)^+$ . Moreover, assume that  $s_i \in (0,1)$ , where  $\forall \ell \in [\![1,m-1]\!]$ ,  $s_\ell \leq s_{\ell+1}$ . In addition to **(P)**, let us assume that the  $f_i$ 's are at most polynomial satisfying the following so-called "triangular structure": (TS)

There exist a vector  $\mathbf{b} \in \mathbb{R}^m_+$  and a lower triangular invertible matrix  $Q \in \mathcal{M}_m(\mathbb{R}_+)$   $(q_{ii} \neq 0)$ , such that  $\forall \mathbf{r} = (r_1, \dots, r_m) \in \mathbb{R}^m_+$ ,  $Q\mathbf{f}(\mathbf{r}) \leq \left[1 + \sum_{i=1}^m r_i\right]\mathbf{b}$  where  $\mathbf{f}(\mathbf{r}) = (f_1(\mathbf{r}), \dots, f_m(\mathbf{r}))^T$ .

Then, System (S) admits a unique nonnegative global strong solution.

*Proof.* Let us denote  $Q = (q_{ij})_{i,j=1,...,m}$  and  $\mathbf{b} = (b_1, \ldots, b_m)^T$ . Thanks to Lemma 3.1, System (S) has a unique strong solution  $(u_1, \cdots, u_m)$  in  $Q_{T_{\max}}$ . In the subsequent, we will prove that this solution is global, meaning that  $T_{\max} = +\infty$ . Let  $T \in (0, T_{\max})$  and  $t \in (0, T]$ . For any  $i \in [1, m]$ , let  $v_i$  be the weak solution to

$$(P_i) \begin{cases} \partial_t v_i(t, \mathbf{x}) + d_i(-\Delta)_{Sp}^{s_i} v_i(t, \mathbf{x}) &= \left[1 + \sum_{j=1}^m u_j(t, \mathbf{x})\right] \frac{b_i}{q_{ii}}, & (t, \mathbf{x}) \in Q_T, \\ \mathcal{B}[v_i(t, \mathbf{x})] &= 0, & (t, \mathbf{x}) \in \Sigma_T, \\ v_i(0, \mathbf{x}) &= 0, & \mathbf{x} \in \Omega. \end{cases}$$

Therefore, we have

(3.37) 
$$v_i(t, \mathbf{x}) = \int_0^t T_{s_i}(t-\tau) \Big[ 1 + \sum_{j=1}^m u_j(\tau, \mathbf{x}) \Big] \frac{b_i}{q_{ii}} d\tau, \quad (t, \mathbf{x}) \in [0, T) \times \Omega.$$

Given (TS), the following inequality holds:

(3.38) 
$$\sum_{j=1}^{i} q_{ij} f_j(u_1(t, \mathbf{x}), \dots, u_m(t, \mathbf{x})) \le \left[1 + \sum_{j=1}^{m} u_j(t, \mathbf{x})\right] b_i.$$

— Let us multiply both the first equation of  $(P_1)$  and the equation for i = 1 of System (S) by  $q_{11}$ , then subtract the former from the latter. We obtain

(3.39) 
$$q_{11} \Big[ \partial_{\tau} (u_1 - v_1)(\tau, \mathbf{x}) + d_1 (-\Delta)^{s_1}_{Sp} (u_1 - v_1)(\tau, \mathbf{x}) \Big] \le 0$$

after replacing t by  $\tau$ . In light of the duality concept outlined in Lemma 3.2, let  $\phi$  be a nonnegative regular function and let  $\mathcal{V}(\tau, \mathbf{x})$  be the solution to Problem  $(P_{\phi,t})$  with  $s = s_1$  and  $d = d_1$ . Subsequently, we multiply both sides of inequality (3.39) by  $\mathcal{V}$  and integrate over  $Q_t$ . As  $q_{11} > 0$ , we get

(3.40) 
$$\iint_{Q_t} \partial_t (u_1 - v_1)(\tau, \mathbf{x}) \mathcal{V}(\tau, \mathbf{x}) d\tau d\mathbf{x} + \iint_{Q_t} d_1 (-\Delta)_{Sp}^{s_1} (u_1 - v_1)(\tau, \mathbf{x}) \mathcal{V}(\tau, \mathbf{x}) d\tau d\mathbf{x} \le 0.$$

Then,

$$\iint_{Q_t} -(u_1 - v_1)(\tau, \mathbf{x}) \partial_\tau \mathcal{V}(\tau, \mathbf{x}) d\tau d\mathbf{x} + \iint_{Q_t} d_1 (u_1 - v_1)(\tau, \mathbf{x}) (-\Delta)^{s_1}_{Sp} \mathcal{V}(\tau, \mathbf{x}) d\tau d\mathbf{x} \le \int_{\Omega} u_{01}(\mathbf{x}) \mathcal{V}_0(\mathbf{x}) d\mathbf{x}.$$

Therefore, we have

(3.42) 
$$\iint_{Q_t} (u_1 - v_1)(\tau, \mathbf{x}) \phi(\tau, \mathbf{x}) d\tau d\mathbf{x} \le \int_{\Omega} u_{01}(\mathbf{x}) \mathcal{V}_0(\mathbf{x}) d\mathbf{x}$$

Consider a sufficiently large p > 1 and let  $p' = \frac{p}{p-1}$ . By applying Hölder's inequality and the estimate (2.8), we obtain

(3.43) 
$$\iint_{Q_t} (u_1 - v_1)(\tau, \mathbf{x}) \phi(\tau, \mathbf{x}) d\tau d\mathbf{x} \le \|u_{01}\|_{L^p(\Omega)} \|\mathcal{V}_0\|_{L^{p'}(\Omega)} \le C_{19} \|\phi\|_{L^{p'}(Q_t)}.$$

By duality, it follows that

$$(3.44) ||(u_1 - v_1)^+||_{L^p(Q_t)} \le C_{19}.$$

Consequently, we obtain

$$(3.45) ||u_1||_{L^p(Q_t)} \le ||v_1||_{L^p(Q_t)} + ||(u_1 - v_1)^+||_{L^p(Q_t)} \le C_{20}[1 + ||v_1||_{L^p(Q_t)}].$$

— Now, consider  $i \ge 2$ . By multiplying the first equation of  $(P_i)$  and the equation for i of System (S) by  $q_{ii}$ , and then subtracting the former from the latter, we get

$$(3.46) \ q_{ii} \Big[ \partial_{\tau} (u_i - v_i)(\tau, \mathbf{x}) + d_i (-\Delta)_{Sp}^{s_i} (u_i - v_i)(\tau, \mathbf{x}) \Big] \le -\sum_{j=1}^{i-1} q_{ij} \Big[ \partial_{\tau} u_j(\tau, \mathbf{x}) + d_j (-\Delta)_{Sp}^{s_j} u_j(\tau, \mathbf{x}) \Big],$$

after replacing t by  $\tau$ . As above, let  $\phi$  be a nonnegative regular function and  $\mathcal{V}(\tau, \mathbf{x})$  be the solution to Problem  $(P_{\phi,t})$  with  $s = s_i$  and  $d = d_i$ . Next, we multiply both sides of inequality (3.46) by  $\mathcal{V}$  and then integrate over  $Q_t$ , resulting in

$$(3.47) \begin{cases} q_{ii} \iint_{Q_t} \partial_{\tau} (u_i - v_i)(\tau, \mathbf{x}) \mathcal{V}(\tau, \mathbf{x}) d\tau d\mathbf{x} + q_{ii} \iint_{Q_t} d_i (-\Delta)^{s_i}_{Sp} (u_i - v_i)(\tau, \mathbf{x}) \mathcal{V}(\tau, \mathbf{x}) d\tau d\mathbf{x} \\ \leq -\sum_{j=1}^{i-1} q_{ij} \iint_{Q_t} \partial_{\tau} u_j(\tau, \mathbf{x}) \mathcal{V}(\tau, \mathbf{x}) d\tau d\mathbf{x} - \sum_{j=1}^{i-1} q_{ij} d_j \iint_{Q_t} (-\Delta)^{s_j}_{Sp} u_j(\tau, \mathbf{x}) \mathcal{V}(\tau, \mathbf{x}) d\tau d\mathbf{x}. \end{cases}$$

Thus, we get (3.48)

$$\begin{cases} q_{ii} \iint_{Q_t} (u_i - v_i)(\tau, \mathbf{x}) \phi(\tau, \mathbf{x}) d\tau d\mathbf{x} \\ \leq \sum_{j=1}^{i-1} q_{ij} \int_{\Omega} u_{0j}(\mathbf{x}) \mathcal{V}_0(\mathbf{x}) d\mathbf{x} + \sum_{j=1}^{i-1} q_{ij} \iint_{Q_t} u_j(\tau, \mathbf{x}) \left( \partial_t \mathcal{V}(\tau, \mathbf{x}) - d_j(-\Delta)_{Sp}^{s_j} \mathcal{V}(\tau, \mathbf{x}) \right) d\tau d\mathbf{x}. \end{cases}$$

Consistent with the above, let us choose p > 1 sufficiently large. Applying Hölder's inequality along with the estimate (2.8) yields (3.49)

$$\begin{cases} (J)_{Q_{t}}^{(J,\mathbf{r},\mathbf{r})} \int_{Q_{t}}^{(J,\mathbf{r},\mathbf{r})} (u_{i} - v_{i})(\tau,\mathbf{x}) \phi(\tau,\mathbf{x}) d\tau d\mathbf{x} \\ \leq C_{21} \sum_{j=1}^{i-1} \|u_{0j}\|_{L^{p}(\Omega)} \|\mathcal{V}_{0}\|_{L^{p'}(\Omega)} + C_{21} \sum_{j=1}^{i-1} \|u_{j}\|_{L^{p}(Q_{t})} \left( \|\partial_{\tau}\mathcal{V}\|_{L^{p'}(Q_{t})} + d_{j} \left\| (-\Delta)_{Sp}^{s_{j}}\mathcal{V} \right\|_{L^{p'}(Q_{t})} \right) \\ \leq C_{22} \sum_{j=1}^{i-1} \|u_{0j}\|_{L^{p}(\Omega)} \|\phi\|_{L^{p'}(Q_{t})} + C_{22} \sum_{j=1}^{i-1} \|u_{j}\|_{L^{p}(Q_{t})} \left( \|\phi\|_{L^{p'}(Q_{t})} + d_{j} \left\| (-\Delta)_{Sp}^{s_{j}}\mathcal{V} \right\|_{L^{p'}(Q_{t})} \right), \end{cases}$$

By assumption, we have  $0 < s_1 \leq \cdots \leq s_m < 1$ . Then, for any fixed  $i \in [\![2,m]\!]$  and any  $j \in [\![1, i-1]\!]$ , it follows that  $s_j \leq s_i$ . Therefore, thanks to Proposition 3.1, for any  $j \in [\![1, i-1]\!]$ , (3.50)

$$\left\| (-\Delta)_{Sp}^{s_j} \mathcal{V} \right\|_{L^{p'}(Q_t)} \le \left\| (-\Delta)_{Sp}^{s_i} \mathcal{V} \right\|_{L^{p'}(Q_t)}^{\frac{s_j}{s_i}} \| \mathcal{V} \|_{L^{p'}(Q_t)}^{\frac{s_i - s_j}{s_i}} \le C \| \phi \|_{L^{p'}(Q_t)}^{\frac{s_j}{s_i}} \| \phi \|_{L^{p'}(Q_t)}^{\frac{s_i - s_j}{s_i}} = C \| \phi \|_{L^{p'}(Q_t)},$$

where the second inequatity follows from the estimate (2.8). Consequently, we obtain (3.51)

$$\begin{cases} \iint_{Q_t} (u_i - v_i)(\tau, \mathbf{x}) \phi(\tau, \mathbf{x}) d\tau d\mathbf{x} &\leq C_{22} \sum_{j=1}^{i-1} \|u_{0j}\|_{L^p(\Omega)} \|\phi\|_{L^{p'}(Q_t)} + C_{23} \sum_{j=1}^{i-1} \|u_j\|_{L^p(Q_t)} \|\phi\|_{L^{p'}(Q_t)} \\ &\leq C_{24} \Big( 1 + \sum_{j=1}^{i-1} \|u_j\|_{L^p(Q_t)} \Big) \|\phi\|_{L^{p'}(Q_t)}. \end{cases}$$

Since the function  $\phi$  is nonnegative and regular, we deduce by duality that

(3.52) 
$$\|(u_i - v_i)^+\|_{L^p(Q_t)} \le C_{24} \Big(1 + \sum_{j=1}^{i-1} \|u_j\|_{L^p(Q_t)}\Big).$$

Therefore, given  $u_i \leq v_i + (u_i - v_i)^+$  and applying elementary induction on *i*, we get

(3.53) 
$$\|u_i\|_{L^p(Q_t)} \le C_{25} \left(1 + \sum_{j=1}^i \|v_j\|_{L^p(Q_t)}\right),$$

for any  $i \in [\![2, m]\!]$ . Thanks to (3.45), inequality (3.53) is valid for any  $i \in [\![1, m]\!]$ . Hence, summing over i and raising to the *p*th power yields

(3.54) 
$$\left\|\sum_{i=1}^{m} u_i\right\|_{L^p(Q_t)}^p \le C_{26} \left(1 + \sum_{i=1}^{m} \|v_i\|_{L^p(Q_t)}^p\right).$$

By taking the  $L^p(\Omega)$ -norm of (3.37) and using the estimate (2.2) with q = p together with Hölder's inequality, we obtain

(3.55) 
$$\|v_i(t,\cdot)\|_{L^p(\Omega)}^p \le C_{27} \Big[1 + \Big\|\sum_{i=1}^m u_i\Big\|_{L^p(Q_t)}^p\Big].$$

Then, by using (3.54) and applying Gronwall's Lemma, we derive that

(3.56) 
$$\sum_{i=1}^{m} \|v_i(t,\cdot)\|_{L^p(\Omega)}^p \le C_{28} \left[ 1 + \int_0^t \sum_{i=1}^m \|v_i(\tau,\cdot)\|_{L^p(\Omega)}^p d\tau \right] < \psi(t),$$

where  $\psi(t) : [0, +\infty) \to [0, +\infty)$  is a nondecreasing and continuous function. Hence, for any  $i \in [\![1,m]\!], v_i(t,\cdot)$  is bounded in  $L^p(\Omega)$  for any  $p < +\infty$ . Returning to (3.54),  $\sum_{i=1}^m u_i(t,\cdot)$  is also bounded in  $L^p(\Omega)$  for any  $p < +\infty$ , and consequently, so is  $u_i(t,\cdot)$ .

Since the  $f_i$ 's are at most polynomial, it follows that  $f_i \in L^q(Q_T)$  for sufficiently large q. Therefore, we can select  $q > \max_{i \in [\![1,m]\!]} \left\{ \frac{N+2s_i}{2s_i} \right\}$  to ensure, by Theorem 2.2, that for any  $i \in [\![1,m]\!]$ 

(3.57) 
$$||u_i||_{L^{\infty}(Q_T)} \le C_{29}$$

Finally,  $T_{\max} = +\infty$ .

## 4 Numerical simulations of a fractional "Brusselator" model

The aim of this section is to numerically investigate a fractional version of the classical so-called "Brusselator", *namely* 

$$(S_{\text{Bruss}}) \begin{cases} \partial_t u_1(t, \mathbf{x}) + d_1(-\Delta)_{S_p}^{s_1} u_1(t, \mathbf{x}) &= -u_1(t, \mathbf{x})(u_2(t, \mathbf{x}))^2 + bu_2(t, \mathbf{x}), & (t, \mathbf{x}) \in Q_T, \\ \partial_t u_2(t, \mathbf{x}) + d_2(-\Delta)_{S_p}^{s_2} u_2(t, \mathbf{x}) &= u_1(t, \mathbf{x})(u_2(t, \mathbf{x}))^2 - (b+1)u_2(t, \mathbf{x}) + a, & (t, \mathbf{x}) \in Q_T, \\ \mathcal{B}[u_1(t, \mathbf{x})] = \mathcal{B}[u_2(t, \mathbf{x})] &= 0, & (t, \mathbf{x}) \in \Sigma_T, \\ u_1(0, \mathbf{x}) &= u_{01}(\mathbf{x}), & \mathbf{x} \in \Omega, \\ u_2(0, \mathbf{x}) &= u_{02}(\mathbf{x}), & \mathbf{x} \in \Omega, \end{cases}$$

where  $0 < s_1, s_2 < 1, d_1, d_2, a, b > 0$  and  $(u_{01}, u_{02}) \in (L^{\infty}(\Omega)^+)^2$ . Let us denote

$$f_1(u_1, u_2) := -u_1 u_2^2 + b u_2, \quad f_2(u_1, u_2) := u_1 u_2^2 - (b+1)u_2 + a$$

and consider  $(r_1, r_2) \in [0, +\infty)^2$ . Then,

$$f_1(0, r_2) = br_2 \ge 0, \quad f_2(r_1, 0) = a > 0, \quad af_1(r_1, r_2) + r_2 f_2(r_1, r_2) \le a_2$$

so that  $(\mathbf{P}) + (\mathbf{M}')$  holds.

It is worth recalling that, in the classical case (*i.e.*  $s_1 = s_2 = 1$ ), the global existence of strong solutions to System ( $S_{\text{Bruss}}$ ) was established in [50]; see also [72] and references therein. In contrast, for the fractional case, there is a notable lack of mathematical tools to rigorously prove either global existence or finite-time blow-up. This section aims to address this matter numerically.

Nonetheless, it should be emphasized that the numerical approximation of such models is challenging and imposes significant computational constraints. To tackle these challenges, we adopt a Fourier spectral method introduced in [22], which provides an appealing and straightforward approach for solving fractional problems of type (S) in bounded rectangular domains of  $\mathbb{R}^N$ , for  $N \in [1,3]$ . Indeed, their proposed schemes yield a fully diagonal representation of the SFL, enhancing both accuracy and computational efficiency.

Now, let us give a brief overview of the chosen numerical method in the case of the homogeneous Dirichlet boundary condition. Let us underline that the method also applies when considering the homogeneous Neumann boundary condition instead of the Dirichlet one.

#### Description of the numerical method

• Discretisation in space. To clearly convey the fundamental idea of the space discretisation, we will restrict the description to 1D. More precisely, let us consider the following fractional heat equation in  $(0, T) \times (c_1, c_2)$ ,

(4.1) 
$$\begin{cases} \partial_t w(t,x) + d(-\Delta)^s_{Sp} w(t,x) = 0, & (t,x) \in (0,T) \times (c_1,c_2), \\ w(t,c_1) = w(t,c_2) = 0, & t \in (0,T), \\ w(0,x) = w_0(x), & x \in (c_1,c_2), \end{cases}$$

where d > 0 and  $w_0 \in L^{\infty}(c_1, c_2)^+$ . Therefore, the analytical solution to Problem (4.1) could be written as

(4.2) 
$$w(t,x) = \sum_{k=1}^{+\infty} w_k(t)e_k(x) = \sum_{k=1}^{+\infty} w_k(0)e^{-d\lambda_k^s t}e_k(x),$$

where  $\lambda_k = \left(\frac{k\pi}{c_2-c_1}\right)^2$  and  $e_k = \sqrt{\frac{2}{c_2-c_1}} \sin\left(\frac{k\pi(x-c_1)}{c_2-c_1}\right)$ . The Fourier spectral method under discussion involves approximating the series expansion of

The Fourier spectral method under discussion involves approximating the series expansion of (4.2) by using a finite set of orthonormal trigonometric eigenfunctions  $\{e_k\}_{k\in\mathbb{N}^*}$ , where the number of these functions is equal to the number of discretization points. Then, we have

(4.3) 
$$w(t,x) \approx \sum_{k=1}^{K} w_k(0) e^{-d\lambda_k^s t} e_k(x).$$

The coefficients  $w_k(0)$  in (4.3), along with the inverse reconstruction of w, can be efficiently calculated using established algorithms such as the direct and inverse Discrete Sine/Cosine Transforms, see, for instance, [20, 23]. In [22], the authors have well illustrated the simplicity of applying this method. Additionally, the mesh points are defined by  $x_j := c_1 + jh_x$ , where  $h_x := \frac{c_2 - c_1}{K+1}$  and  $j \in [\![1, K]\!]$ .

• **Discretisation in time.** Let us go back to System  $(S_{\text{Bruss}})$ . Let us consider a discretization of time as  $0 = t_0 < t_1 < \cdots < t_{\widetilde{N}} = T$  with  $h_t := t_{n+1} - t_n$  for  $n \in [\![0, \widetilde{N} - 1]\!]$ . Following the approach outlined in [22], we use a backward Euler discretisation of the time derivative presented. Let us consider  $u_i^n(\cdot) \approx u_i(t_n, \cdot)$  for i = 1, 2. In each time interval  $[t_n, t_{n+1}]$ , the nonlinear terms are handled using the following fixed point iteration: given  $(u_1^n, u_2^n)$ , initialize  $(u_1^{n+1,0}, u_2^{n+1,0}) := (u_1^n, u_2^n)$ , and for  $\ell \in [\![1, L]\!]$  find  $(u_1^{n+1,\ell}, u_2^{n+1,\ell})$  such that

(4.4) 
$$\frac{u_i^{n+1,\ell} - u_i^n}{h_t} = -d_i(-\Delta)_{Sp}^{s_i} u_i^{n+1,\ell} + f_i\left(u_1^{n+1,\ell-1}, u_2^{n+1,\ell-1}\right)$$

where i = 1, 2 and L is to be determined later. Notably, L = 1 corresponds to a fully explicit treatment of the nonlinear term, whereas a sufficiently large L makes the method fully implicit.

Now, applying the Fourier transform to both sides of (4.4) yields, for  $k \in [\![1, K]\!]$  and i = 1, 2,

$$\frac{u_{i_k}^{n+1,\ell} - u_{i_k}^n}{h_t} = -d_i \lambda_k^{s_i} u_{i_k}^{n+1,\ell} + f_{i_k} \left( u_1^{n+1,\ell-1}, u_2^{n+1,\ell-1} \right)$$

where  $f_{ik}$  is the kth Fourier coefficient of the  $f_i$ . Hence, after rearranging the terms, we obtain

$$u_{ik}^{n+1,\ell} = \frac{1}{1+d_i\lambda_k^{s_i}h_t} \left[ u_{ik}^n + h_t f_{ik} \left( u_1^{n+1,\ell-1}, u_2^{n+1,\ell-1} \right) \right].$$

Before presenting our numerical simulations, let us point out that in [22], the authors effectively tested the method by comparing the numerical solution with the exact one in cases where the latter is known. Furthermore, they assessed the convergence of the schemes in both time and space.

#### Numerical simulations in 2D

Now, let us showcase some numerical simulations that illustrate the performance of the chosen numerical scheme. To do so, let us go back to System  $(S_{\text{Bruss}})$  with  $\Omega = (-1,1) \times (-1,1)$ ,  $\mathcal{B}[u_i] = u_i, u_{0i}(\mathbf{x}) = u_{0i}(x,y) = (1-x)^{s_i}(1-y)^{s_i}$  for i = 1, 2, a = 2 and b = 1. Our main aim is to examine the existence of a nonnegative solution to System  $(S_{\text{Bruss}})$ , which may persist globally in time. To this end, we aim to numerically solve the system until a sufficiently large final time  $(t_{\text{final}} = 5 \times 10^{10} \text{ in our case}).$ 

The two following figures display the solution graphs obtained at  $t_{\text{final}} = 5 \times 10^{10}$  for different values of  $s_1$ ,  $s_2$ ,  $d_1$  and  $d_2$ , while keeping  $h_x = 10^{-3}$ ,  $h_t = 10^{-2}$  and L = 3 fixed. More precisely, **Figure 1** and **Figure 2** correspond to the first and second components of the solution, respectively.



**Figure 1.** First component of the solution for different values of  $(s_1, s_2, d_1, d_2)$  at  $t_{\text{final}} = 5 \times 10^{10}$ 



Figure 2. Second component of the solution for different values of  $(s_1, s_2, d_1, d_2)$  at  $t_{\text{final}} = 5 \times 10^{10}$ 

Besides visualizing the solution, we compute the  $L^{\infty}$ -norm of  $u_1$  and  $u_2$  at  $t_{\text{final}} = 5 \times 10^{10}$  for different values of  $h_x$ . Specifically, we fix the time step  $h_t = 10^{-2}$  while varying the spatial mesh size  $h_x$  and the parameter L. The computed norms are presented in **Table 1** and **Table 2**. These tables serve to verify that the solution remains stable and does not exhibit significant changes as the mesh is refined. As shown, the norms remain relatively constant across different values of  $h_x$  and L, suggesting that the numerical solution is consistent.

	$s_1 = 0.25, s_2 =$	$0.75,  d_1 = 3,  d_2 = 5$	$s_1 = 0.9, s_2 = 0.5, d_1 = 2, d_2 = 4$		
$h_x, L$	$  u_1  _{\infty}$	$  u_2  _{\infty}$	$  u_1  _{\infty}$	$  u_2  _{\infty}$	
$h_x = 10^{-2}, L = 1$	0.034593927	0.147334012	0.032958290	0.231046944	
$h_x = 5 \times 10^{-3}, L = 2$	0.034597215	0.147343745	0.032968899	0.231071583	
$h_x = 2 \times 10^{-3}, L = 2$	0.034597791	0.147345473	0.032970613	0.231076136	
$h_x = 10^{-3}, L = 3$	0.034597830	0.147345587	0.032970734	0.231076469	

**Table 1.** Table of  $L^{\infty}$ -errors of  $u_1$  and  $u_2$  at  $t_{\text{final}} = 5 \times 10^{10}$ 

	$s_1 = 0.35, s_2 = 0.8, d_1 = 1, d_2 = 3$		$s_1 = 0.8, s_2 = 0.6, d_1 = 5, d_2 = 1$		
$h_x, M$	$  u_1  _{\infty}$	$  u_2  _{\infty}$	$  u_1  _{\infty}$	$  u_2  _{\infty}$	
$h_x = 10^{-2}, L = 1$	0.128912072	0.215751920	0.035781626	0.539477321	
$h_x = 5 \times 10^{-3}, L = 2$	0.128930880	0.215770510	0.035797500	0.539571499	
$h_x = 2 \times 10^{-3}, L = 2$	0.128934168	0.215773656	0.035800230	0.539588620	
$h_x = 10^{-2}, L = 3$	0.128934401	0.215773860	0.035800423	0.539589842	

**Table 2.** Table of  $L^{\infty}$ -errors of  $u_1$  and  $u_2$  at  $t_{\text{final}} = 5 \times 10^{10}$ 

In summary, we have numerically investigated the global existence of nonnegative solutions to System  $(S_{\text{Bruss}})$ . Our numerical simulations indicate that the system has nonnegative solutions that can be accurately computed over a large final time. This suggests the potential existence of a global solution in time for System  $(S_{\text{Bruss}})$ . However, the question of whether this solution persists for all time remains theoretically open. We intend to further explore this aspect in future studies.

## References

- N. Abatangelo, L. Dupaigne, Nonhomogeneous boundary conditions for the spectral fractional Laplacian. Ann. Inst. H. Poincaré Anal. Non Linéaire 34(2) (2017), 439–467,
- [2] N. Abatangelo, E. Valdinoci, Getting Acquainted with the Fractional Laplacian. In: Dipierro, S. (eds) Contemporary Research in Elliptic PDEs and Related Topics. Springer INdAM Series, vol 33. Springer, 2019.
- [3] B. Abdellaoui, A. Attar, R. Bentifour, E.-H. Laamri, Existence results to a class of nonlinear parabolic systems involving potential and gradient terms. Mediterranean Journal of Mathematics 17(119) (2020), 1–30.
- [4] B. Ahmad, M.S. Alhothuali, H.H. Alsulami, M. Kirane, S. Timoshin, On nonlinear nonlocal systems of reaction diffusion equations. Abstr. Appl. Anal. 2 (2014), 1–6.
- [5] B. Ahmad, A. Alsaedi, D. Hnaien, M. Kirane, On a semi-linear system of nonlocal time and space reaction diffusion equations with exponential nonlinearities. J. Integral Equa. and Applications **30** (2018), 17–40.
- [6] A. Alsaedi, M. Al-Yami, M. Kirane, F. Momenkhan, A triangular nonlinear reactionfractional diffusion system with a balance law. Math. Methods Appl. Sci. 41 (2018), 1825– 1830.
- [7] F. Andreu-Vaillo, J.M. Mazón, J.D. Rossi, J.J. Toledo-Melero, Nonlocal Diffusion Problems. Mathematical Surveys and Monographs, 165, American Mathematical Society, Providence, RI; Real Sociedad Matemàtica Española, Madrid, 2010.
- [8] W. Arendt, *Heat Kernels*. ISEM course 2005/06.
- [9] S. Atmani, K. Biroud, M. Daoud, E.-H. Laamri, On some nonlocal parabolic reactiondiffusion systems with gradient source terms. Fract. Calc. Appl. Anal. 27, 2644–2687 (2024).
- [10] S. Atmani, K. Biroud, M. Daoud, E.-H. Laamri, On some nonlocal elliptic systems with gradient source terms. Acta Appl. Math. 181(9) (2022).
- [11] S. Atmani, K. Biroud, M. Daoud, E.-H. Laamri, Fractional elliptic reaction-diffusion systems with coupled gradient terms and different diffusion. Hal-04123233. Submitted.
- [12] S. Atmani, K. Biroud, M. Daoud, E.-H. Laamri, Fractional parabolic reaction-diffusion systems with gradient terms and different diffusion. In preparation.
- [13] A. Barabanova, On the global existence of solutions of a reaction-diffusion system with exponential nonlinearity. Proc. Amer. Math. Soc. 122 (1994), 827-831.
- [14] U. Biccari, M. Warma, E. Zuazua, Local Regularity for Fractional Heat Equations. Recent Advances in PDEs: Analysis, Numerics and Control, SEMA SIMAI Springer Ser. 17, Springer, Cham, 2018, 233–249.

- [15] U. Biccari, M. Warma, E. Zuazua, Control and numerical approximation of fractional diffusion equations. Handbook of Numerical Analysis XXIII. Numerical Control: Part A, 23 (2022), 1–58.
- [16] M. Bonforte, P. Ibarrondo, M. Ispizua, The Cauchy-Dirichlet problem for singular nonlocal diffusions on bounded domains. Discrete Contin. Dyn. Syst., 43(3&4) (2023), 1090–1142.
- [17] M. Bonforte, Y. Sire and J.L. Vázquez, Existence, uniqueness and asymptotic behavior for fractional porous medium equations on bounded domains. Discrete Contin. Dyn. Syst. 35 (2015), 5725–5767.
- [18] A. Bonito, W. Lei, J.E. Pasciak, The approximation of parabolic equations involving fractional powers of elliptic operators. J. Comput. Appl. Math. 315 (2017), 32–48.
- [19] N. Boudiba, Existence globale pour des systèmes de réaction-diffusion avec contrôle de masse. Ph.D. thesis, Université de Rennes 1, France, 1999.
- [20] W.L. Briggs, V.E. Henson, *The DFT: an owner's manual for the discrete Fourier transform*. SIAM, Philadelphia, 2000.
- [21] C. Bucur, E. Valdinoci, <u>Nonlocal diffusion and applications</u>. ISBN 978-3-319-28738-6. Lecture Notes of the Unione Matematica Italiana, 2016.
- [22] A. Bueno-Orovio, D. Kay, K. Burrage, Fourier spectral methods for fractional-in-space reaction-diffusion equations. Bit Numer Math, 54 (2014), 937–954.
- [23] A. Bueno-Orovio, V.M. Pérez-García, Spectral smoothed boundary methods: the role of external boundary conditions. Numer. Meth. Part. Differ. Equ., 22 (2006), 435–448.
- [24] J.A. Cañizo, L. Desvillettes, K. Fellner, Improved Duality Estimates and Applications to Reaction-Diffusion Equations. Commun. Partial Differ. Equ. 39(6) (2014), 1185–1204.
- [25] C. Caputo, T. Goudon, A. Vasseur, Solutions of the 4-species quadratic reaction-diffusion system are bounded and C<sup>∞</sup>-smooth, in any space dimension. Anal. PDE **12**(7) (2019), 1773– 1804.
- [26] H. Chan, D. Gómez-Castro, J.L. Vázquez, Singular solutions for fractional parabolic boundary value problems. Rev. Real Acad. Cienc. Exactas Fis. Nat. Ser. A-Mat 116(159) (2022).
- [27] H. Chen, M. Bhakta, H. Hajaiej, On the bounds of the sum of eigenvalues for a Dirichlet problem involving mixed fractional Laplacians. J. Diff. Eq. 317 (2022), 1–31.
- [28] G. Ciavolella, B. Perthame, Existence of a global weak solution for a reaction-diffusion problem with membrane conditions. J. Evol. Equ. 21 (2021), 1513–1540.
- [29] B. Claus, M. Warma, Realization of the fractional Laplacian with nonlocal exterior conditions via forms method. J. Evol. Equ. 20 (2020), 1597–1631.
- [30] M. Daoud, E.-H. Laamri, Fractional Laplacians : A short survey. Discrete Contin. Dyn. Syst.-S 15(1) (2022), 95–116.
- [31] M. Daoud, E.-H. Laamri, A. Baalal, A class of parabolic fractional reaction-diffusion systems with control of total mass: Theory and numerics. J. Pseudo-Differ. Oper. Appl. 15(18) (2024). DOI: 10.1007/s11868-023-00576-w.
- [32] E.B. Davies, <u>Heat kernels and spectral theory</u>. Volume 92 of Cambridge Tracts in Mathematics. Cambridge University Press, Cambridge, 1990.

- [33] S. Dipierro, G. Giacomin, E. Valdinoci, Diffusive processes modeled on the spectral fractional Laplacian with Dirichlet and Neumann boundary conditions. Posted on 2022, DOI: web.ma.utexas.edu/mp\_arc/c/22/22-12.
- [34] S. Dipierro, E.P. Lippi, C. Sportelli, E. Valdinoci, Some nonlinear problems for the superposition of fractional operators with Neumann boundary conditions. arXiv preprint (2024). arXiv:2404.11091.
- [35] K. Fellner, E.-H. Laamri, Exponential decay towards equilibrium and global classical solutions for nonlinear reaction-diffusion systems. J. Evol. Equ. 16(3) (2016), 681–704.
- [36] K. Fellner, J. Morgan, B.Q. Tang, Global classical solutions to quadratic systems with mass control in arbitrary dimensions. Ann. Inst. H. Poincaré Anal. Non Linéaire 37(2) (2020), 281–307.
- [37] K. Fellner, J. Morgan, B.Q. Tang, Uniform-in-time for quadratic reaction-diffusion systems with mass dissipation in higher dimensions. Discrete Contin. Dyn. Syst. -S 14(2) (2021), 635-651.
- [38] X. Fernandez-Real, *Boundary regularity for the fractional heat equation*. Bachelor's degree thesis, Polytechnic University of Catalonia, 2014.
- [39] X. Fernandez-Real, X. Ros-Oton, Boundary regularity for the fractional heat equation. Rev. Acad. Cienc. Ser. A Math. 110 (2016), 49–64.
- [40] M. Fukushima, Y. Oshima, M. Takeda, <u>Dirichlet forms and symmetric Markov processes</u>. Berlin, New York: De Gruyter, 1994.
- [41] C.G. Gal, M. Warma, Reaction-diffusion equations with fractional diffusion on non-smooth domains with various boundary conditions. Discrete Contin. Dyn. Syst. 36(3) (2016), 1279– 1319.
- [42] C.G. Gal, M. Warma, Nonlocal transmission problems with fractional diffusion and boundary conditions on nonsmooth interfaces. Commun. Partial Differ. Equ. 42(4) (2017), 579–625.
- [43] C.G. Gal, M. Warma, *Fractional-in-Time Semilinear Parabolic Equations and Applications*. Springer Nature Switzerland AG, 2020.
- [44] P. Garbaczewski, V. Stephanovich, Fractional Laplacians in bounded domains: Killed, reflected, censored and taboo Lévy flights. Phys. Rev. E99 (2019), 042126.
- [45] T. Goudon, A. Vasseur, Regularity Analysis for Systems of Reaction-Diffusion Equations. Annales Sci. ENS (4) 43(1) (2010), 117–142.
- [46] A. Haraux, A. Youkana, On a result of K. Masuda concerning reaction-diffusion equations. Tôhoku Math. J. 40 (1988), 159–163.
- [47] A. Henrot, <u>Extremum Problems for Eigenvalues of Elliptic Operators</u>. Frontiers in Mathematics, Birkhäuser Verlag, Basel, Switzerland, 2006.
- [48] B.I. Henry, T.A.M. Langlands, P. Straka, An Introduction to Fractional Diffusion. Complex Physical, Biophysical and Econophysical Systems (2010), 37–89.
- [49] M.A. Herrero, A.A. Lacey and J.L. Velàzquez, Global existence for reaction-diffusion systems modelling ignition. Arch. Rat. Mech. Anal. 142 (1998), 219–251.
- [50] S.L. Hollis, R.H. Martin, M. Pierre, Global existence and boundedness in reaction-diffusion systems. SIAM J. Math. Ana. 18 (1987), 744–761.

- [51] J. Hulshof, R. Van Der Vorst, Differential Systems with Strongly Indefinite Variational Structure. Journal of Functional Analysis 114(1) (1993), 32-58.
- [52] A.A. Kilbas, H.M. Srivastava, J.J. Trujillo, <u>Theory and Applications of Fractional Diff-</u> erential Equations. Elsevier, Amsterdam, 2006.
- [53] M. Kirane, A. Alsaedi, B. Ahmad, On systems of reaction-diffusion equations with a balance law: The sequel. Comput. Math. with Appl. 78 (2019), 1244–1260.
- [54] H. Koba, H. Matsuoka, Generalized quasi-reversibility method for a backward heat equation with a fractional Laplacian. Analysis 35(1) (2015), 47–57.
- [55] E.-H. Laamri, Existence globale pour des systèmes de réaction-diffusion dans L<sup>1</sup>, Ph.D thesis, Université de Nancy 1, France, 1988.
- [56] E.-H. Laamri, Global existence of classical solutions for a class of reaction-diffusion systems. Acta Appl. Math. 115(2) (2011), 153–165.
- [57] E.-H. Laamri, B. Perthame, Reaction-diffusion systems with initial data of low regularity. J. Differ. Equ. 269(11) (2020), 9310–9335.
- [58] E.-H. Laamri, M. Pierre, Global existence for reaction-diffusion systems with nonlinear diffusion and control of mass. Ann. Inst. H. Poincaré Anal. Non Linéaire 34(3) (2017), 571–591.
- [59] E.-H. Laamri, M. Pierre, Stationary reaction-diffusion systems in  $L^1$ . M3AS **28**(11) (2018), 2161–2190.
- [60] E.-H. Laamri, M. Pierre, Stationary reaction-diffusion systems in L<sup>1</sup> revisited. Discrete Contin. Dyn. Syst. -S 14(2) (2021), 455–464.
- [61] D. Lamberton, Equations d'évolution linéaires associées à des semi-groupes de contractions dans les espace L<sup>p</sup>. J. Funct. Anal. 72 (1987), 252–262.
- [62] A. Léculier, S. Mirrahimi, J.M. Roquejoffre, Propagation in a fractional reaction-diffusion equation in a periodically hostile environment. J. Dyn. Diff. Equat. 33 (2021), 863–890.
- [63] A. Lischke, G. Pang, M. Gulian, F. Song, C. Glusa, X. Zheng, Z. Mao, W. Cai, M.M. Meerschaert, M. Ainsworth, G.E. Karniadakis, *What is the Fractional Laplacian?* Journal of Computational Physics 404 (2020), 109009.
- [64] C. Louis-Rose, Null controllability from the exterior of fractional parabolic-elliptic coupled systems. EJDE 26 (2020), 1–18.
- [65] R.H. Martin, M. Pierre, Nonlinear reaction-diffusion systems. In Nonlinear Equations in the Applied Sciences, W.F. Ames and C. Rogers ed., Math. Sci. Eng. 185, Acad. Press, New York 1991.
- [66] G. Molica Bisci, V.D. Radulescu, R. Servadei, <u>Variational Methods for Nonlocal Fractional Problems</u>. Encyclopedia of Mathematics and its Applications, Vol. 162, Cambridge University Press, Cambridge, 2016, with a foreword by Jean Mawhin.
- [67] J.D. Murray, <u>Mathematical biology : I. An introduction</u>. Springer-Verlag, Berlin, Heidlberg, 2002.
- [68] E. Nane, N.H. Tuan, Approximate Solutions of Inverse Problems for Nonlinear Space Fractional Diffusion Equations with Randomly Perturbed Data. SIAM/ASA Journal on Uncertainty Quantification 6(1) (2018), 302–338.
- [69] P.-T. Nguyen, B.Q. Tang, Well-Posedness for Non-local Reaction-Diffusion Systems with Mass Dissipation in ℝ<sup>N</sup>. arXiv:2501.02603 (2025).

- [70] A. Pazy, Semigroups of Linear Operators and Applications to Partial Differential Equations. Springer, New York, 1983.
- [71] M. Pierre, An L<sup>1</sup> method to prove global existence in some reaction-diffusion systems. In "Contributions to Nonlinear Partial Differential Equations", Vol. II, Pitman Research notes, J.I. Diaz and P.L. Lions ed., 155 (1987), 220–231.
- [72] M. Pierre, Global Existence in Reaction-Diffusion Systems with Control of Mass : a Survey. Milan J. Math. 78 (2010), 417–455.
- [73] M. Pierre, Weak solutions and supersolutions in L<sup>1</sup> for reaction-diffusion systems. J. Evol. Equ. 3 (2003), 153–168.
- [74] M. Pierre, D. Schmitt, Blowup in reaction-diffusion systems with dissipation of mass. SIAM Rev. 42(1) (2000), 93–10.
- [75] M. Pierre, T. Suzuki, Y. Yamada, Dissipative reaction diffusion systems with quadratic growth. Indiana Univ. Math. J. 68(1) (2019), 291–322.
- [76] L.M. Pismen, <u>Patterns and interfaces in dissipative dynamics</u>. Springer Science & Business Media, 2006.
- [77] I. Podlubny, Fractional Differential Equations. Academic Press, San Diego, 1999.
- [78] I. Prigogine, R. Lefever, Symmetry breaking instabilities in dissipative systems. J. Chem. Phys., 48 (1968), 1665–1700.
- [79] I. Prigogine, G. Nicolis, Biological order, structure and instabilities. Quart. Rev. Biophys. 4 (1971), 107–148.
- [80] F. Rothe, <u>Global Solutions of Reaction-Diffusion Systems</u>. Lecture Notes in Mathematics 1072, Springer, Berlin 1984.
- [81] P. Scherer, S.F. Fischer, <u>Reaction-Diffusion Systems</u>. In: Theoretical Molecular Biophysics. Biological and Medical Physics, Biomedical Engineering. Springer, Berlin, Heidelberg, 2010.
- [82] P. Quittner, Ph. Souplet, <u>Superlinear Parabolic Problems: Blow-up</u>, <u>Global Existence and</u> Steady States. Advanced Texts, Birkhäuser, (2019).
- [83] P. Shi, W.-T. Li, F.-Y. Yang, Spatiotemporal dynamics in a fractional diffusive SIS epidemic model with mass action infection mechanism. J. Math. Biol. 89(54) (2024).
- [84] R. Servadei, E. Valdinoci, On the spectrum of two different fractional operators. Proc. R. Soc. Edinb., Sect. A 144 (2014), 831–855.
- [85] L.W. Somathilake, K. Burrage, A space-fractional-reaction-diffusion model for pattern formation in coral reefs. Cogent Math. Stat. 5 (2018), 1426524.
- [86] Ph. Souplet, Global existence for reaction-diffusion systems with dissipation of mass and quadratic growth. J. Evol. Equ. 18(4) (2018), 1713–1720.
- [87] T. Suzuki, Y. Yamada, Global-in-time behavior of Lotka-Volterra system with diffusion: skewsymmetric case. Indiana Univ. Math. J. **64**(1) (2015), 181–216.
- [88] N.H. Tuan, V.V. Tri, D. O'Regan, On a nonlinear parabolic equation with fractional Laplacian and integral conditions. J. Appl. Anal. 101(17) (2022), 5974–5988.
- [89] A.M. Turing, The chemical basis of morphogenesis. Philos. Trans. R. Soc. London Ser. B 237 (1952), 37–72.

- [90] E. Valdinoci, From the long jump random walk to the fractional Laplacian. Bol. Soc. Esp. Mat. Apl. SeMA 49 (2009), 33–44.
- [91] J.L. Vázquez, <u>Nonlinear Diffusion with Fractional Laplacian Operators</u>. In: Holden, H., Karlsen, K. (eds) Nonlinear Partial Differential Equations. Abel Symposia 7. Springer, Berlin, Heidelberg, 2012.
- [92] J.L. Vázquez, The mathematical theories of diffusion. Nonlinear and fractional diffusion. Springer Lecture Notes in Mathematics, CIME Subseries, 2017.
- [93] I.I. Vrabie, C<sub>0</sub>-Semigroups and Applications. Elsevier Science B.V., 2003.
- [94] G. Zhao, S. Ruan, Spatiotemporal dynamics in epidemic models with Lévy flights: A fractional diffusion approach. J. Math. Pures Appl. 173 (2023), 243–277.