Testing New Physics in Oscillations at a Neutrino Factory

Peter B. Denton,¹ Julia Gehrlein,² and Chui-Fan Kong^{3,4}

- ¹High Energy Theory Group, Physics Department, Brookhaven National Laboratory, Upton, NY 11973, USA
- ²Physics Department, Colorado State University, Fort Collins, CO 80523, USA
- ³Tsung-Dao Lee Institute & School of Physics and Astronomy, Shanghai Jiao Tong University, China
- ⁴Key Laboratory for Particle Astrophysics and Cosmology (MOE) & Shanghai Key Laboratory for Particle Physics and Cosmology, Shanghai Jiao Tong University, Shanghai 200240, China E-mail: pdenton@bnl.gov, julia.gehrlein@colostate.edu,

kongcf@sjtu.edu.cn

ABSTRACT: A neutrino factory is a potential successor to the upcoming generation of neutrino oscillation experiments and a possible precursor to next-generation muon colliders. Such a machine would provide a well characterized beam of ν_{μ} , $\bar{\nu}_{\mu}$, ν_{e} , and $\bar{\nu}_{e}$ neutrinos with comparable statistics. Here we show the sensitivity of a neutrino factory to new oscillation physics scenarios such as vector neutrino non-standard interactions and CPT violation. We study two different potential setups for a neutrino factory with different assumptions on charge identification in the far detector. We find that a neutrino factory can improve over most of the current constraints on these scenarios and over forecasted constraints by DUNE, even when doubling DUNE's statistics. Additionally, we find that a neutrino factory can break degeneracies between the standard oscillation parameters and neutrino non-standard interaction parameters present at DUNE.

³0009-0007-7010-5085

¹0000-0002-5209-872X

²0000-0002-1235-0505

Contents

1	Introduction	1
2	Experimental Setup	2
3	Neutrino non-standard Interactions	4
	3.1 Framework	4
	3.2 Results for non-standard vector interactions	5
4	CPT violation	10
	4.1 Framework	10
	4.2 Results for the CPT framework	11
5	Conclusion	15
\mathbf{A}	Additional NSI results	15

1 Introduction

Neutrino physics has entered the precision era after a non-zero θ_{13} mixing angle was measured [1–6]. Still, there are three main unknowns in the standard three-flavor neutrino oscillation framework: the neutrino mass ordering, the octant of θ_{23} , and the value of δ_{CP} [7, 8]. The next-generation neutrino oscillation experiments like JUNO [9], Hyper-Kamiokande [10], and DUNE [11], are expected to address these unknowns. However, the experimental sensitivities to the standard oscillation parameters can be reduced in the presence of physics beyond the Standard Model (BSM), such as non-standard interaction [12, 13] and CPT violation [14]. In addition, given the open theoretical and experimental questions related to neutrinos, it is important to search for additional new physics in the neutrino sector wherever possible.

However, there exist degeneracies between many of the new physics parameters and the standard oscillation parameters, which may make it challenging to search for new physics even with the upcoming planned experiments. There are several experimental proposals beyond the upcoming JUNO, Hyper-Kamiokande, and DUNE programs that aim to break these degeneracies and improve the sensitivities to both BSM parameters and the standard oscillation parameters. First, a multi-baseline setup such as the T2HKK experiment [15], long-baseline experiments targetting the second maximum such as ESSnuSB [16], and atmospheric neutrino experiments [17, 18] can resolve the degeneracies by measuring neutrino oscillations at different baselines. Secondly, similar to the DUNE experiment [11] the P2O experiment [19] can search for differences between BSM and standard oscillations in a broad energy spectrum. Thirdly, an additional low-energy neutrino source from muon

decay at rest [20] can reduce the new physics impact and provide a clean measurement of standard oscillation parameters, which in turn gives a better sensitivity to new physics parameters when combined with a high-energy neutrino beam. Finally, performing a joint analysis of all experiments is desirable to resolve degeneracies and benefit from complementarities to provide stringent constraints on BSM scenarios (as done in [21] in the case of new interactions).

In this study, we consider another possibility for a potential future neutrino oscillation experiment, namely a neutrino factory (NF) which uses LAr far detectors to measure highenergy $\mathcal{O}(\text{GeV}) \stackrel{(-)}{\nu}_{\mu}$ and $\stackrel{(-)}{\nu}_{e}$ beams coming from muon decays [22–25]. Recently, standard three-flavor oscillations at a modern neutrino factory setup have been studied [26] as well as non-oscillation new physics in [27]. In this paper we study new physics in oscillations at a neutrino factory. We focus on vector neutrino non-standard interaction (NSI) and CPT violation where long-baseline experiments can provide the strongest bounds. Other BSM scenarios, like scalar NSI [28, 29] are much better constrained by solar neutrino experiments due to the significant amplification of this effect inside the sun [30] and the sterile neutrino scenario is typically best constrained by short-baseline experiments [31-33] and atmospheric neutrinos [34, 35]. A neutrino factory mainly benefits from three perspectives to constrain NSI and CPT violation compared to existing and planned oscillation experiments, like DUNE or Hyper-Kamiokande, which use neutrinos produced in a fixed target setup: i) a larger statistics of the oscillation channel from the electron flavor to the electron/muon flavor makes it sensitive to more new physics parameters, ii) the higher neutrino energy can amplify the new physics effects such as NSI to make it easier to be observed, and iii) the simultaneous presence of equally large amounts of neutrinos and antineutrinos, albeit with a different energy spectrum, in the beam allows for studies of new physics in the neutrino and anti-neutrino sectors separately.

The paper is outlined as follows. In Sec. 2, we discuss the experimental setup of a neutrino factory, in sec. 3, 4 we present our results for the the scenarios of new neutrino interactions, and CPT violation before we conclude in sec. 5.

2 Experimental Setup

At a neutrino factory an intense neutrino beam with $\mathcal{O}(\text{GeV})$ energy can be produced via the decays of positively (negatively) charged muons inside a muon storage ring. Due to the 3-body muon decay process, the flavor composition and energy spectrum of the neutrinos are well understood. Moreover, different from traditional accelerator neutrino experiments, there exists a large fraction of ν_e ($\bar{\nu}_e$) from μ^+ (μ^-) decay which allows studies of $\nu_e \rightarrow \nu_e, \nu_\mu$ ($\bar{\nu}_e \rightarrow \bar{\nu}_e, \bar{\nu}_\mu$) oscillation channels with large statistics. We consider two possible setups for a neutrino factory, motivated by existing accelerator complexes in the US.¹ Either the neutrinos are produced at i) Fermilab (FNAL) or at ii) Brookhaven National Laboratory (BNL) [26]. We assume 10²¹ muon decays per year with equal running time for muons and anti-muons. For the FNAL-SURF (BNL-SURF) baseline we use as maximal neutrino

¹See [36] for a study of a neutrino factory based at J-PARC.

energy $E_{\mu} = 5(8)$ GeV. The choice of energy and running time is motivated by previous results from [26] as these values optimize the precision of the CP phase δ in the absence of new physics at a neutrino factory; for different goals such as new physics one may prefer different muon energies. We assume that the far detector is the DUNE far detector with a total of 40 kt LAr fiducial mass to detect neutrinos produced at FNAL or BNL, which have traveled a baseline of L = 1284.9 km or L = 2542.3 km, respectively. For both configurations we use a matter density of 2.848 g/cm^3 and we assume a 2% uncertainty on matter density which is consistent with DUNE TDR [37]. Realistically, the matter density for the BNL configuration is somewhat higher due to its longer baseline however, similarly to DUNE, a neutrino factory is expected to be sensitive to the matter density at the 30% level [38], thus a slightly different value due to the longer baseline will not significantly affect our results.

The beam at a neutrino factory has a large intrinsic backgrounds, for example in the ν_e disappearance channel coming from $\bar{\nu}_{\mu} \rightarrow \bar{\nu}_e$ appearance. It is possible to differentiate them at some level due to the different initial spectrum combined with the (presumably) different oscillation probability. The excellent energy reconstruction abilities of LArTPC can partially differentiate these components to the far detector flux [39] and additionally a detector with charge identification (CID) capabilities allows for improved background rejection. To study the role of CID we consider three scenarios of the detector performance: i) no charge identification, ii) with 100% charge identification of electrons (eCID), and iii) with 100% charge identification of muons (μ CID). We also include the oscillated tau neutrinos with the charged-current detection process at the far detector. Due to the large uncertainties on the tau neutrino signal efficiency at LAr detectors, however, we treat the tau neutrino events as background only, using the cross section from [40]. We consider only normalization uncertainties at the far detector for a neutrino factory where we use 2.5% for both the muon and electron neutrino normalization uncertainties. For a neutrino factory the statistics of the ν_e , ν_{μ} in the FD is comparable, therefore we treat the normalization uncertainty for a neutrino factory slightly different than for LBNF neutrinos [37], where the muon and electron neutrino signal normalization uncertainties are 5% and 2%, respectively, while a common 5% uncertainty is used for muon and electron neutrino background. Moreover, we take a 10% uncertainty on the neutral-current background, 20% uncertainty on the tau background. We use the GLoBES framework [41, 42] for our studies.

To compare the potential of a NF to planned future experiments we also take DUNE into account², where the experimental details can be found in [11, 37]. We consider the following experimental setups: i) 10 years of DUNE (DUNE-10), ii) 20 years of (DUNE-20)³, or iii) 10 years of DUNE plus 10 years of a neutrino factory at either FNAL (NF-FNAL) and BNL (NF-BNL) baseline. Each of the two NF scenarios has three CID scenarios as mentioned above. Therefore we have eight different experimental setups in total.

²We note that there has been proposed a high energy tune for DUNE designed to enhance its sensitivity for tau neutrinos, NSIs, and other scenarios [43–48], but would do so at the cost of shifting away from the CP optimized beam. The NF beam parameters we have chosen here were selected to maximize the sensitivity to CP violation [26]. Thus for a direct comparison, we focus on DUNE in the nominal configuration.

³We assume 1.1×10^{21} POT per year of DUNE [37].

Table 1. Overview of the neutrino oscillation parameters and matter density used in this study. The oscillation parameters come from the global fit [49]. We assume the current best fit values as the true values for these oscillation parameters.

	Best Fit	1σ Range
$\sin^2 \theta_{12}$	0.307	Fixed
$\sin^2 heta_{23}$	0.572	Free
$\sin^2 heta_{13}$	0.02203	$\pm 5.8 \times 10^{-4}$
$\delta_{ m CP}$	-90°	Free
$\Delta m_{21}^2 \ [10^{-5} {\rm eV}^2]$	7.41	Fixed
$\Delta m_{31}^2 \ [10^{-3} {\rm eV}^2]$	2.511	Free
Matter density $[g/cc]$	2.848	2%

We summarize the assumed true values for the standard oscillation parameters in tab. 1, taken from [49]. ⁴ We assume true normal ordering throughout this study. For δ_{CP} we assume its true value corresponds to the largest allowed CP violation given the existing measurements, $\delta_{CP} = -90^{\circ}$, unless otherwise stated. This value is also somewhat preferred by T2K [51]. We assume no prior on δ_{CP} . Note that the preferred value is in minor tension with the NO ν A result [52, 53], where the best fit is around 150° for normal ordering and -90° for inverted ordering. We fix the solar mixing angle θ_{12} and mass-squared difference Δm_{21}^2 to their current best fit values from [49] as long-baseline accelerator experiments have limited sensitivity to them [54]. Since the reactor mixing angle θ_{13} has been measured very precisely by reactor experiments, we fix its best fit value with prior knowledge. We take θ_{23} and Δm_{31} parameters as free parameters without any prior since they will be well measured at LBL experiments.

3 Neutrino non-standard Interactions

We first introduce the framework for (vector) NSI and then show the sensitivity of DUNE and a NF to the NSI parameters and the standard oscillation parameters in the presence of NSI. We focus on $\delta_{CP}^{\text{true}} = -90^{\circ}$, the results for $\delta_{CP}^{\text{true}} = 0^{\circ}$ can be found in the appendix.

3.1 Framework

Neutrinos could have interactions with other fermions due to the presence of new mediators. A general framework to incorporate this effect is neutrino non-standard interactions which, for example, originate from dimension-6 operators. First proposed by Wolfenstein [55], NSIs are often considered to have a vector mediator. In fact, such new neutrino interactions are natural features in many neutrino mass models with new mediators [56–58]. The strength of the NSI depends on the specific model and large NSI is predicted in some models, see e.g. [58–65]. In general, vector NSI can be of charged-current (CC) or neutral-current (NC) type [12, 13]. Both affect neutrinos during their propagation through matter while only CC

⁴Updated global fit results [50] do not significantly change our results as either DUNE or a NF will measure most parameters relevant for new physics better than the existing constraints.

NSI affects neutrinos at the point of production and detection.⁵ Long-baseline experiments like DUNE and a NF are strongly affected by matter effects, and thus are expected to have exceptional sensitivity to the NC NSI. To this end, we take the NC NSI into consideration below in the context of propagation.⁶

In the low-energy regime, neutrino NC NSI with matter fields can be formulated in terms of the effective four-fermion Lagrangian term [68],

$$\mathcal{L}_{\rm NSI} = -2\sqrt{2}G_F \sum_{f,\alpha,\beta} \epsilon^{f,V}_{\alpha\beta} (\bar{\nu}_{\alpha}\gamma^{\mu}P_L\nu_{\beta}) (\bar{f}\gamma_{\mu}Pf) , \qquad (3.1a)$$

where G_F is the Fermi constant. The dimensionless coefficient $\epsilon_{\alpha\beta}^{f,V}$ quantifies the size of the neutrino NSI with matter fermions $f \in \{e, u, d\}$ relative to the standard weak interaction. The above effective Lagrangian introduces a new neutrino matter potential which modifies the Hamiltonian,

$$H = \frac{1}{2E} \left[U \begin{pmatrix} 0 & 0 & 0 \\ 0 & \Delta m_{21}^2 & 0 \\ 0 & 0 & \Delta m_{31}^2 \end{pmatrix} U^{\dagger} + a \begin{pmatrix} 1 + \epsilon_{ee} & \epsilon_{e\mu} & \epsilon_{e\tau} \\ \epsilon_{e\mu}^* & \epsilon_{\mu\mu} & \epsilon_{\mu\tau} \\ \epsilon_{e\tau}^* & \epsilon_{\mu\tau}^* & \epsilon_{\tau\tau} \end{pmatrix} \right],$$
(3.2)

where U is the PMNS mixing matrix [69, 70] and $\Delta m_{ij}^2 \equiv m_i^2 - m_j^2$ are the mass squared differences. Here the first term is the vacuum component H_{vac} , while $a \equiv 2\sqrt{2}G_F N_e E$ with N_e being the electron number density in the second term describes the strength of the matter effect. Finally, we relate the Lagrangian level NSI parameters to the Hamiltonian level ones via $\epsilon_{\alpha\beta} \equiv \sum_f N_f \epsilon_{\alpha\beta}^{f,V} / N_e$ with the fermion matter density N_f .

The diagonal NSI parameters are real, while the off-diagonal NSI parameters may be complex and can be parametrized as,

$$\epsilon_{\alpha\beta} = |\epsilon_{\alpha\beta}| e^{\mathbf{i}\phi_{\alpha\beta}}.\tag{3.3}$$

Therefore, there are a total of nine new real NSI parameters. However, one of the diagonal parameters can be subtracted without affecting the oscillation behavior. For definiteness, we adopt the notation, $a \times \text{Diag}(1 + \epsilon_{ee} - \epsilon_{\mu\mu}, 0, \epsilon_{\tau\tau} - \epsilon_{\mu\mu})$, for the diagonal part of the matter potential in our study.

3.2 Results for non-standard vector interactions

We start with the simplest case, where all NSI parameters are assumed to be real and only one NSI parameter is considered at one time. While this will miss some of the more interesting physics cases presented below, it is valuable in highlighting the relative role of each experiment. In Fig. 1, we show the expected constraints on the diagonal and the real part of the off-diagonal NSI parameters at the 90% confidence level (C.L.) in the left and right panel, respectively. As discussed in Sec. 2, there are eight experimental scenarios under consideration, the allowed regions are shown by patches of different colors. For comparison, we also include the current global fit data [21] with an error bar. Note

⁵Note that NC NSI does affect the detection of neutrinos in NC channels, see e.g. [66, 67].

 $^{^6 \}mathrm{One}$ could also consider NC NSI in the detection of the NC channel at a NF.



Figure 1. Constraints on the diagonal (left) NSI parameters and off-diagonal NSI parameters (taken to be real) (right) at 90% confidence level assume 10 years of DUNE (red), 20 years of DUNE, and 10 years of DUNE combined with 10 years of a NF located at FNAL or BNL (hatched). The purple, yellow, and green color correspond to no CID at the NF far detector, 100% electron CID, and 100% μ CID. We compare the results to the global fit results from [21]. There are three allowed regions of $\epsilon_{ee} - \epsilon_{\mu\mu}$ from the global fit, where two of them are shown by error bars while the remaining one [-3.1, -2.8] is shown with numbers.

however that for the global fit results the NSI parameters not shown have been marginalized over, in contrast to our approach.

We find that for all five NSI parameters, DUNE combined with a neutrino factory can reach better sensitivities than DUNE alone, even with doubled running time. Especially for a neutrino factory at BNL with e/μ CID, the sensitivity can be improved by a factor of ~ 2 compared to DUNE with 20 years running time. We also generally find that it is possible to improve beyond global bounds even in cases where non-accelerator experiments are expected to dominate. The first of which is $\epsilon_{ee} - \epsilon_{\mu\mu}$ which is strongly constrained by the combination of solar and reactor data, and yet DUNE and a NF may be better than existing constraints. The second of which are the two parameters tightly constrained in atmospheric experiments: $\epsilon_{\tau\tau} - \epsilon_{\mu\mu}$ and $\epsilon_{\mu\tau}$. We find that DUNE and a NF can actually outperform the existing constraints on $\epsilon_{\mu\tau}$. Finally, we note that in terms of the NSI parameters expected to be best probed by long-baseline accelerator experiments, $\epsilon_{e\mu}$ and $\epsilon_{e\tau}$, we find that DUNE with a NF can outperform, in some cases easily, existing constraints.

There are two separate regions in the $\epsilon_{ee} - \epsilon_{\mu\mu}$ parameter space due to the LMA-Dark solution [71–75]. This is a degeneracy in the oscillation Hamiltonian in matter in the presence of NSI. The Hamiltonian in the case of standard matter effect is physically equivalent to the Hamiltonian H in the presence of NSI by taking $\Delta m_{21}^2 \rightarrow -\Delta m_{21}^2$, $\Delta m_{31}^2 \rightarrow -\Delta m_{31}^2$, $\delta_{CP} \rightarrow -\delta_{CP}$, $\epsilon_{ee} - \epsilon_{\mu\mu} = -2$, $\epsilon_{\tau\tau} - \epsilon_{\mu\mu} = 0$, and $\epsilon_{\alpha\beta} \in \mathbb{R}$.

To study the degeneracies between different NSI parameters, we extend our study to the two-parameter space which we anticipate should cover the majority of the degeneracy space. We first consider the case where the off-diagonal NSI parameters are assumed to be real and the experimental constraints at 90% C.L. are shown in Fig. 2. There exists a significant degeneracy between $\Re \epsilon_{e\mu}$ and $\Re \epsilon_{e\tau}$, see e.g. [76, 77], which is because the only



Figure 2. Expected constraints on different combinations of two non-zero NSI parameters at 90% confidence level. All NSI parameters are assumed to be real, that is, the complex phase is $\in \{0, \pi\}$. We compare the expected constraints from DUNE only (with 10 and 20 years of running time) and DUNE combined with a NF for different assumptions on CID. Note that the $\epsilon_{e\mu}$ - $\epsilon_{e\tau}$ panel has different axes.

relevant combinations of NSI parameters in the $\nu_e/\bar{\nu}_e$ appearance channel are $\epsilon_{e\mu} + \epsilon_{e\tau}$ and $\epsilon_{e\mu} - \epsilon_{e\tau}$, assuming real NSI parameters [78]. Especially when $\delta_{CP} = -90^{\circ}$, only the combination of $\epsilon_{e\mu} - \epsilon_{e\tau}$ survives [78], which leads to the positive correlation between $\epsilon_{e\mu}$ and $\epsilon_{e\tau}$ as shown in the figure. In all cases except the $\epsilon_{\tau\tau} - \epsilon_{\mu\mu}$ - $\epsilon_{ee} - \epsilon_{\mu\mu}$ panel, DUNE combined with a neutrino factory at BNL with μ CID configuration gives the most stringent constraints on the two-parameter space due to the longer baseline of the BNL setup. In the $\epsilon_{\tau\tau} - \epsilon_{\mu\mu}$ - $\epsilon_{ee} - \epsilon_{\mu\mu}$ panel, the eCID and μ CID cases are both very similar and both provide the strongest constraint.

For a more general case, we take the off-diagonal NSI parameters to be complex and marginalize over the corresponding phases in Fig. 3. Due to the free complex phase, the degeneracies can be enlarged and significantly reduce the experimental sensitivities to the NSI parameters. Especially, the degeneracy between $\epsilon_{ee} - \epsilon_{\mu\mu}$ and $|\epsilon_{e\tau}|$ leads to a wing-like structure for the original DUNE configuration which comes from the non-trivial dependence among the $\phi_{e\tau}$, δ_{CP} and θ_{23} parameters [79]. To better understand this degeneracy we show in the left panel of Fig. 4 the oscillation probabilities without NSI and with non-zero $\epsilon_{ee} - \epsilon_{\mu\mu}$



Figure 3. The experimental constraint on different combinations of two NSI parameters at 90% confidence level, where the corresponding complex phase of NSI off-diagonal parameter and δ_{CP} are marginalized over.

and $\epsilon_{e\tau}$. For the standard case without NSI, we use the values of oscillation parameters from Tab. 1, while we take the best fit from that minimizes the χ^2 function for the NSI case. More explicitly, we choose the point ($\epsilon_{ee} - \epsilon_{\mu\mu} = 1.2$ and $|\epsilon_{e\tau}| = 0.3$) in the wing-like structure panel of Fig. 3 as the input and a parameter set $\phi_{e\tau} = -152^{\circ}$, $\theta_{23} = 42^{\circ}$, and $\delta_{CP} = -53^{\circ}$ the values that lead to a χ^2 minimum, which indicates this parameter set has a degeneracy with the standard case without NSI. As DUNE only measures ν_e appearance and ν_{μ} disappearance channels, this degeneracy significantly reduces DUNE's sensitivity to these NSI parameters. It is exciting to see the addition of a neutrino factory can break this degeneracy and significantly enhance the sensitivity in the $\epsilon_{ee} - \epsilon_{\mu\mu} - |\epsilon_{e\tau}|$ parameter space. This is because a neutrino factory can measure the ν_e disappearance channel and the ν_{μ} appearance channel, which help break the existing degeneracy in the ν_e appearance and ν_{μ} disappearance channels.

Another important degeneracy is the island-like structure that occurs for DUNE in the $\epsilon_{\tau\tau} - \epsilon_{\mu\mu}$ and $|\epsilon_{e\tau}|$ parameter space, see e.g. [77]. This also arises from the non-trivial dependence on the $\phi_{e\tau}$, δ_{CP} and θ_{23} parameters as shown in the right panel of Fig. 4. Still, the ν_e disappearance channel and ν_{μ} appearance channel can break the degeneracy in the original DUNE configuration. As a result, a neutrino factory can constrain these two parameters to a narrow region. For other combinations, notable improvements are also made by adding a neutrino factory.

In addition to the expected constraints on NSI parameters, we also study the precision of the Dirac CP phase as well as the sensitivity to θ_{23} in the presence of NSI. To quantify the expected precision on to δ_{CP} , we define $\Delta \delta_{CP} \equiv \max \{ |\delta_{CP}^{\text{fit}} - \delta_{CP}^{\text{true}}| \}$ to be the maximum absolute value of the difference between δ_{CP}^{fit} and $\delta_{CP}^{\text{true}}$, where δ_{CP}^{fit} is the fitted Dirac CP



Figure 4. The oscillation probability of $\nu_e \rightarrow \nu_e$ (orange), and $\nu_e \rightarrow \nu_\mu$ (green) with and without NSI using the best-fit values as summarized in Tab. 1. In the NSI case, we take $\epsilon_{ee} - \epsilon_{\mu\mu} =$ $1.2, |\epsilon_{e\tau}| = 0.3, \phi_{e\tau} = -152^{\circ}, \theta_{23} = 42^{\circ}, \text{ and } \delta_{CP} = -53^{\circ}$ in the left panel, while we take $\epsilon_{\tau\tau} - \epsilon_{\mu\mu} =$ $-0.55, |\epsilon_{e\tau}| = 0.125, \phi_{e\tau} = -133^{\circ}, \theta_{23} = 45^{\circ}, \text{ and } \delta_{CP} = -74^{\circ}$ in the right panel. Note that in the NSI cases the values of oscillation parameters not specifically mentioned are the same as those listed in Tab. 1.

phase which gives $\Delta \chi^2 = 2.71$ (90% C.L.) and $\delta_{CP}^{\text{true}}$ is the fixed true Dirac CP phase. We take the benchmark value of $\delta_{CP}^{\text{true}} = 0^{\circ}$ or -90° , and three benchmark values of $\theta_{23}^{\text{true}} = 42^{\circ}$, 45° , or 49° for completeness. The results for $\delta_{CP}^{\text{true}} = 0$ are shown in the appendix which show qualitatively similar improvements to the precision of δ_{CP} with the addition of a neutrino factory. Similar to the standard oscillation framework without NSI, the precision on $\delta_{CP}^{\text{true}} = 0^{\circ}$ is better than for the $\delta_{CP}^{\text{true}} = -90^{\circ}$ case while the octant of θ_{23} has only a mild impact on the CP precision. In the $\delta_{CP}^{\text{true}} = 0^{\circ}$ case, the increase in precision by combing DUNE with a neutrino factory can be enhanced by a factor of 2 (1.5) compared with the original DUNE configuration in the presence of diagonal (off-diagonal) NSI parameter. A sizable improvement exists for $\delta_{CP}^{\text{true}} = -90^{\circ}$, as seen in Fig. 5.

In Fig. 6 we show the sensitivity to θ_{23} for different benchmark values of $\delta_{CP}^{\text{true}}$ and $\theta_{23}^{\text{true}}$ as discussed above. Here we take the $\epsilon_{e\mu}$ parameter for illustration since the neutrino factory has a significant improvement in this case, while the results for other NSI parameters are shown in the appendix. In the figure, $\epsilon_{e\mu}$ is marginalized over its modulus together with its phase while the true value is taken to be zero. For benchmark values of $\theta_{23}^{\text{true}} = 42^{\circ}$ and 49° , the θ_{23} sensitivity of DUNE can be significantly reduced if one also considers NSI. As a result, the θ_{23} octant cannot be determined with over $\sim 3\sigma$ C.L. at the benchmarks considered. With the addition of a neutrino factory, however, the sensitivity to θ_{23} octant even when minimizing over $\epsilon_{e\mu}$ with a phase can reach over 5σ . We also find that the true value of $\delta_{CP}^{\text{true}}$ has only a small impact on the sensitivity.



Figure 5. The expected uncertainty on δ_{CP} at 90% confidence level, where the corresponding NSI parameter is minimized. In this figure, the off-diagonal NSI parameters are taken as complex and the corresponding phase is minimized together with its modulus at a time. We assume $\delta_{CP}^{\text{true}} = -90^{\circ}$ and $\theta_{23}^{\text{true}} = 42^{\circ}$, 45° , 49° . See Fig. 11 in the appendix for $\delta_{CP} = 0$.



Figure 6. The experimental sensitivity to θ_{23} for different true values of θ_{23} and we assume no NSI effect in the true case. For the fitting parameter, we consider the complex off-diagonal NSI parameter $\epsilon_{e\mu}$, which is minimized over including its complex phase. See Figs. 12-15 in the appendix.

4 CPT violation

4.1 Framework

The CPT symmetry, which connects three discrete symmetries: charge conjugation (C), parity (P), and time reversal (T), is one of the most essential symmetries in particle physics and its invariance has been a guiding tool to build models. The CPT theorem is based on several basic assumptions on models, including Lorentz invariance, hermiticity of the Hamiltonian, and local commutativity. CPT violation can appear in several classes of more complete models and may affect neutrino oscillations [80–88]. Certainly, the discovery of

the CPT violation would indicate that at least one basic assumption should be rejected and all our model building strategy need to be revisited. Therefore, testing the CPT symmetry is one of the major tasks in particle physics.

In the neutrino sector, neutrino oscillation provides a practical way to test CPT violation since a typical long-baseline neutrino experiment detects both neutrinos and antineutrinos. As a consequence of CPT violation, different mass terms for neutrinos and anti-neutrinos can be present, which allows for different mixing parameters. For clarity, we consider the flavor oscillations of neutrinos are described by $x \in \left\{\theta_{ij}, \delta_{CP}, \Delta m_{ij}^2\right\}$, while anti-neutrinos are associated with $\bar{x} \in \left\{\bar{\theta}_{ij}, \bar{\delta}_{CP}, \Delta \overline{m}_{ij}^2\right\}$. If CPT is conserved, the parameters should coincide except for a minus sign difference of the Dirac CP phase term. The constraint on CPT violation has been analyzed using data taken from the accelerator (T2K and NO ν A) and reactor (Daya Bay and RENO) neutrino experiments [89]. A synergy between two on-going experiments (T2K and NO ν A) and one upcoming reactor experiment (JUNO) has been studied in [90]. A specific bound on the CPT violation in the solar neutrino sector has been studied using the Kamland and solar neutrino data [91]. Moreover, projected constraints are discussed for next-generation experiments which include JUNO, DUNE, Hyper-Kamiokande, and ESSnuSB [91–94].

Since there are two sets of the neutrino oscillation parameters in the presence of CPT violation, we take the following χ^2 function to calculate experimental sensitivities [91],

$$\chi^{2} \equiv \chi^{2}_{\nu}(x) + \chi^{2}_{\bar{\nu}}(\bar{x}).$$
(4.1)

Here x (\bar{x}) corresponds to the oscillation parameter set in the neutrino (antineutrino) oscillation probabilities.

4.2 Results for the CPT framework

We now present our numerical results. In Fig. 7 we show the results for $x - \bar{x}$ at 90% C.L. for $x = \delta_{CP}$, θ_{13} , θ_{23} and both mass splittings assuming $\delta_{CP}^{\text{true}} = 0$, -90° and no CPT violation. Since a LBL experiment is not very sensitive to the solar mixing angle θ_{12} , we do not show the CPT results for it for simplicity. In general, the addition of a NF to DUNE improves the precision on the oscillation parameters for neutrinos and antineutrinos simultaneously, in particular when CID is implemented. An exception to this is the results for Δm_{31}^2 which are similar for 20 years of DUNE and DUNE+NF, nearly independent of the choice of CID. We find that the neutrino oscillation parameters can be more precisely constrained than the anti-neutrino parameters due to the larger statistics.

Accelerator experiments do not have a large sensitivity to the solar parameters, they are most precisely determined by solar and reactor experiments. Concerning CPT violation we see that DUNE alone cannot exclude non-zero $\bar{\Delta}m_{21}^2$ at 90% C.L. while this parameter in neutrino mode can be constrained to be non-zero. Adding a neutrino factory, in particular with the FNAL baseline allows to exclude zero $\bar{\Delta}m_{21}^2$.

For θ_{13} DUNE only cannot distinguish $\theta_{13} = 8.5^{\circ}$ from $\theta_{13} = 9.5^{\circ}$ at 90% C.L. under the assumption of CPT violation even when doubling the running time. This is because the free θ_{23} angle and CP phase can degenerate with θ_{13} . As shown in the left panel



Figure 7. The experimental sensitivity to the difference between the oscillation parameters for neutrino and anti neutrinos $x-\bar{x}$ at 90% confidence level for different experimental setups assuming CPT conservation. We mark the best-fit value with a star. The dashed line shows the CPT conserving line.



Figure 8. The expected experimental constraint on θ_{23} - $\bar{\theta}_{23}$ at 90% confidence level. Left panel: both θ_{13} and $\bar{\theta}_{13}$ are free parameters without any prior. Right panel: both θ_{13} and $\bar{\theta}_{13}$ are free parameters, but only the prior of $\bar{\theta}_{13}$ is included.



Figure 9. Left: The oscillation probability of $\nu_{\mu} \rightarrow \nu_{\mu}$ (blue), $\nu_{\mu} \rightarrow \nu_{e}$ (red), $\nu_{e} \rightarrow \nu_{e}$ (orange), and $\nu_{e} \rightarrow \nu_{\mu}$ (green) for two oscillation parameter sets: case 1) the values of oscillation parameters are same as Tab. 1 (solid lines), and case 2) $\theta_{13} = 9.5^{\circ}, \theta_{23} = 43^{\circ}, \delta_{CP} = -83^{\circ}$ (dashed lines). **Right:** Same as the left panel but for the anti-neutrino mode and the parameter values in case 2: $\bar{\theta}_{23} = 42^{\circ}, \bar{\theta}_{13} = 9.9^{\circ}, \bar{\delta}_{CP} = -94^{\circ}$. Note that the values of oscillation parameters not specifically mentioned are the same as those listed in Tab. 1.



Figure 10. CPT violation benchmark plot. Upper left panel: $\sin^2 \theta_{13}^{\text{true}} = 0.032$, $\sin^2 \bar{\theta}_{13}^{\text{true}} = 0.022$. Upper right panel: $\theta_{23}^{\text{true}} = 49^\circ$, $\bar{\theta}_{23}^{\text{true}} = 42^\circ$ with $\theta_{13}/\bar{\theta}_{13}$ priors. Lower left panel: same as the upper right one but without $\theta_{13}/\bar{\theta}_{13}$ priors. Lower right panel: same as the upper right one but only $\bar{\theta}_{13}$ prior is included.

of Fig. 9, for $\nu_{\mu} \rightarrow \nu_{e}$ and $\nu_{\mu} \rightarrow \nu_{\mu}$ channels which are main signals at DUNE, the oscillation probabilities with ($\theta_{13} = 9.5^{\circ}$, $\theta_{23} = 43^{\circ}$, $\delta_{CP} = -83^{\circ}$) are very close to that of ($\theta_{13} = 8.5^{\circ}, \theta_{23} = 49^{\circ}, \delta_{CP} = -90^{\circ}$), leading to the island-like structure in Fig. 7. However, the $\nu_{e} \rightarrow \nu_{e}$ channel is almost independent of the CP phase and θ_{23} angle, which means a measurement of ν_{e} disappearance channel can significantly improve the θ_{13} sensitivity, in a similar fashion to a typical medium baseline reactor neutrino experiment, except with the added channel of $\nu_{e} \rightarrow \nu_{e}$ instead of just $\bar{\nu}_{e} \rightarrow \bar{\nu}_{e}$. The precision on $\bar{\theta}_{13}$ is $\pm 2.5^{\circ}$. The addition of a NF excludes the $\theta_{13} = 9.5^{\circ}$ and the precision on $\bar{\theta}_{13}$ also improves.

For θ_{23} DUNE alone or combined with a NF cannot resolve the octant at 90% C.L. in the CPT violating scenario unless the NF has good μ CID. As shown in the right panel of Fig. 9, the ν_{μ} disappearance channel is sensitive to $\sin^2 2\theta_{23}$ and hence not sensitive to the θ_{23} octant [10]. However, ν_{μ} appearance is sensitive to $\sin^2 \theta_{23}$, so the octant can be resolved. In the case of neutrino factory, μ CID is required to provide a clean measurement of the ν_{μ} appearance channel. Consequently, the sensitivity of θ_{23} octant can be largely improved. The octant can be resolved for the neutrino parameter due to the larger statistics in the neutrino channel.

Of interest is also the impact of θ_{13} in the determination of θ_{23} octant because the main contribution to the oscillation probability $P_{\mu e}$ is $4\sin^2\theta_{23}\sin^2\theta_{13}\cos^2\theta_{13}\sin^2\Delta_{31}$ with $\Delta_{31} \equiv \Delta m_{31}^2 L/4E_{\nu}$ [10], which means θ_{13} can affect the θ_{23} octant. In Fig. 8 we study the expected constraints without using a prior on neither θ_{13} nor $\bar{\theta}_{13}$. DUNE alone cannot resolve the octant for neither θ_{23} nor $\bar{\theta}_{23}$. When a NF is added only two degenerate solutions for $\bar{\theta}_{23}$ survive due to the less statistics in the antineutrino mode. The prior on $\bar{\theta}_{13}$ is the driving force behind removing the wrong octant solutions for θ_{23} for a NF with μ CID as discussed above.

So far there is no experimental evidence that CPT is violated in nature. However, there is a mild discrepancy in the measurement of θ_{13} and $\bar{\theta}_{13}$ coming from the measurement of solar [95] and accelerator [51] neutrinos compared to reactor antineutrinos [6, 96, 97]. We now consider benchmark values of $\sin^2 \theta_{13} = 0.032$, $\sin^2 \bar{\theta}_{13} = 0.022$, which are consistent with the solar neutrino measurement and reactor neutrino measurement, respectively. To study the impact of CPT for θ_{13} we show in Fig. 10 the expected sensitivities to θ_{13} and $\bar{\theta}_{13}$ assuming the true values to be CPT violating. We again see that the addition of a NF excludes the higher solution for θ_{13} and the precision on both parameters is drastically improved due to the measurement of ν_e disappearance channel as discussed above; this would allow for the discovery of CPT violation if this hint was realized in nature. We now consider a similar scenario but with the octant of θ_{23} . If we assume that $\theta_{23}^{true} = 49^\circ$, $\bar{\theta}_{23}^{true} = 42^\circ$ we again find that having a prior on $\bar{\theta}_{13}$ is crucial to exclude the wrong-octant solutions with a NF. In some cases a CPT violating signal in the octant could be discovered with a NF with given the inclusion of other oscillation data.

5 Conclusion

We have studied the expected sensitivity of a neutrino factory to two new physics scenarios, vector new neutrino interactions and CPT violation. A neutrino factory with a long baseline and LArTPC far detectors is particularly well suited to study these new physics scenarios due to its long baseline, excellent energy reconstruction (including potential charge identification abilities), and access to six oscillation channels plus their CP conjugate channels.

We find that combining 10 years of DUNE with 10 years of a neutrino factory will improve over the constraints on these new physics scenarios from DUNE alone even when considering 20 years of DUNE. Furthermore, the addition of a neutrino factory can remove degeneracies in the DUNE-only constraints due to the presence of new oscillation channels, especially the ν_e disappearance and ν_{μ} appearance channels.

Acknowledgments

PBD acknowledges support by the United States Department of Energy under Grant Contract No. DE-SC0012704. JG acknowledges support by the U.S. Department of Energy Office of Science under award number DE-SC0025448. CFK acknowledges the usage of INPAC cluster at Shanghai Jiao Tong University.

A Additional NSI results

In Fig. 11 we show the precision on δ_{CP} assuming $\delta_{CP}^{\text{true}} = 0$ in the presence of NSI. Figs. 12, 13, 14, 15 we show the sensitivity to θ_{23} in the presence of non-zero NSI parameters, similar to Fig. 6 in the main text.

References

- DAYA BAY collaboration, Observation of electron-antineutrino disappearance at Daya Bay, Phys. Rev. Lett. 108 (2012) 171803 [1203.1669].
- [2] RENO collaboration, Observation of Reactor Electron Antineutrino Disappearance in the RENO Experiment, Phys. Rev. Lett. 108 (2012) 191802 [1204.0626].
- [3] DOUBLE CHOOZ collaboration, Improved measurements of the neutrino mixing angle θ_{13} with the Double Chooz detector, JHEP 10 (2014) 086 [1406.7763].
- [4] RENO collaboration, Measurement of Reactor Antineutrino Oscillation Amplitude and Frequency at RENO, Phys. Rev. Lett. 121 (2018) 201801 [1806.00248].
- [5] DOUBLE CHOOZ collaboration, Double Chooz θ_{13} measurement via total neutron capture detection, Nature Phys. 16 (2020) 558 [1901.09445].
- [6] DAYA BAY collaboration, Precision Measurement of Reactor Antineutrino Oscillation at Kilometer-Scale Baselines by Daya Bay, Phys. Rev. Lett. 130 (2023) 161802 [2211.14988].
- [7] P.B. Denton, M. Friend, M.D. Messier, H.A. Tanaka, S. Böser, J.a.A.B. Coelho et al., Snowmass Neutrino Frontier: NF01 Topical Group Report on Three-Flavor Neutrino Oscillations, 2212.00809.



Figure 11. The expected uncertainty on δ_{CP} assuming $\delta_{CP}^{\text{true}} = 0$ at 90% confidence level, where the corresponding NSI parameter is minimized at a time. In this figure, the off-diagonal NSI parameters are taken as complex and the corresponding phase is minimized together with its modulus at a time. See Fig. 5 in the main text for $\delta_{CP} = -90^{\circ}$.



Figure 12. The experimental sensitivity to θ_{23} , where the $\epsilon_{ee} - \epsilon_{\mu\mu}$ is minimized at a time.

- [8] P.B. Denton, Neutrino Oscillations in the Three Flavor Paradigm, 2501.08374.
- [9] JUNO collaboration, Neutrino Physics with JUNO, J. Phys. G 43 (2016) 030401 [1507.05613].
- [10] HYPER-KAMIOKANDE collaboration, Hyper-Kamiokande Design Report, 1805.04163.
- [11] DUNE collaboration, Deep Underground Neutrino Experiment (DUNE), Far Detector Technical Design Report, Volume II: DUNE Physics, 2002.03005.
- [12] Y. Farzan and M. Tortola, Neutrino oscillations and Non-Standard Interactions, Front. in Phys. 6 (2018) 10 [1710.09360].
- [13] Neutrino Non-Standard Interactions: A Status Report, vol. 2, 2019. 10.21468/SciPostPhysProc.2.001.



Figure 13. The experimental sensitivity to θ_{23} , where the $\epsilon_{\tau\tau} - \epsilon_{\mu\mu}$ is minimized at a time.



Figure 14. The experimental sensitivity to θ_{23} , where the $\epsilon_{\mu\tau}$ is taken as complex and the corresponding phase is minimized together with its modulus at a time.

- [14] C.A. Argüelles et al., Snowmass white paper: beyond the standard model effects on neutrino flavor: Submitted to the proceedings of the US community study on the future of particle physics (Snowmass N2021), Eur. Phys. J. C 83 (2023) 15 [2203.10811].
- [15] HYPER-KAMIOKANDE collaboration, Physics potentials with the second Hyper-Kamiokande detector in Korea, PTEP 2018 (2018) 063C01 [1611.06118].
- [16] ESSNUSB collaboration, Updated physics performance of the ESSnuSB experiment: ESSnuSB collaboration, Eur. Phys. J. C 81 (2021) 1130 [2107.07585].
- [17] S. Razzaque and A.Y. Smirnov, Super-PINGU for measurement of the leptonic CP-phase with atmospheric neutrinos, JHEP 05 (2015) 139 [1406.1407].
- [18] K.J. Kelly, P.A. Machado, I. Martinez Soler, S.J. Parke and Y.F. Perez Gonzalez, Sub-GeV Atmospheric Neutrinos and CP-Violation in DUNE, Phys. Rev. Lett. 123 (2019) 081801 [1904.02751].



Figure 15. The experimental sensitivity to θ_{23} , where the $\epsilon_{e\tau}$ is taken as complex and the corresponding phase is minimized together with its modulus at a time.

- [19] A.V. Akindinov et al., Letter of Interest for a Neutrino Beam from Protvino to KM3NeT/ORCA, Eur. Phys. J. C 79 (2019) 758 [1902.06083].
- [20] S.-F. Ge and A.Y. Smirnov, Non-standard interactions and the CP phase measurements in neutrino oscillations at low energies, JHEP 10 (2016) 138 [1607.08513].
- [21] P. Coloma, M.C. Gonzalez-Garcia, M. Maltoni, J.a.P. Pinheiro and S. Urrea, Global constraints on non-standard neutrino interactions with quarks and electrons, JHEP 08 (2023) 032 [2305.07698].
- [22] S. Geer, Neutrino beams from muon storage rings: Characteristics and physics potential, Phys. Rev. D 57 (1998) 6989 [hep-ph/9712290].
- [23] A. Blondel et al., The neutrino factory: Beam and experiments, Nucl. Instrum. Meth. A 451 (2000) 102.
- [24] A. Blondel, A. Cervera-Villanueva, A. Donini, P. Huber, M. Mezzetto and P.E. Strolin, Future neutrino oscillation facilities, Acta Phys. Polon. B 37 (2006) 2077 [hep-ph/0606111].
- [25] E. Fernandez Martinez, T. Li, S. Pascoli and O. Mena, Improvement of the low energy neutrino factory, Phys. Rev. D 81 (2010) 073010 [0911.3776].
- [26] P.B. Denton and J. Gehrlein, A modern look at the oscillation physics case for a neutrino factory, Nucl. Phys. B 1012 (2025) 116818 [2407.02572].
- [27] A. Bogacz et al., The Physics Case for a Neutrino Factory, in Snowmass 2021, 3, 2022 [2203.08094].
- [28] S.-F. Ge and S.J. Parke, Scalar Nonstandard Interactions in Neutrino Oscillation, Phys. Rev. Lett. 122 (2019) 211801 [1812.08376].
- [29] K.S. Babu, G. Chauhan and P.S. Bhupal Dev, Neutrino nonstandard interactions via light scalars in the Earth, Sun, supernovae, and the early Universe, Phys. Rev. D 101 (2020) 095029 [1912.13488].
- [30] P.B. Denton, A. Giarnetti and D. Meloni, Solar neutrinos and the strongest oscillation constraints on scalar NSI, JHEP 01 (2025) 097 [2409.15411].

- [31] P.A. Machado, O. Palamara and D.W. Schmitz, The Short-Baseline Neutrino Program at Fermilab, Ann. Rev. Nucl. Part. Sci. 69 (2019) 363 [1903.04608].
- [32] M. Dentler, A. Hernández-Cabezudo, J. Kopp, P.A.N. Machado, M. Maltoni,
 I. Martinez-Soler et al., Updated Global Analysis of Neutrino Oscillations in the Presence of eV-Scale Sterile Neutrinos, JHEP 08 (2018) 010 [1803.10661].
- [33] S. Böser, C. Buck, C. Giunti, J. Lesgourgues, L. Ludhova, S. Mertens et al., Status of Light Sterile Neutrino Searches, Prog. Part. Nucl. Phys. 111 (2020) 103736 [1906.01739].
- [34] H. Nunokawa, O.L.G. Peres and R. Zukanovich Funchal, Probing the LSND mass scale and four neutrino scenarios with a neutrino telescope, Phys. Lett. B 562 (2003) 279 [hep-ph/0302039].
- [35] (ICECUBE COLLABORATION)||, ICECUBE collaboration, Search for an eV-Scale Sterile Neutrino Using Improved High-Energy νμ Event Reconstruction in IceCube, Phys. Rev. Lett. 133 (2024) 201804 [2405.08070].
- [36] R. Kitano, J. Sato and S. Sugama, *T violation at a future neutrino factory*, *JHEP* **12** (2024) 014 [2407.05807].
- [37] DUNE collaboration, Experiment Simulation Configurations Approximating DUNE TDR, 2103.04797.
- [38] K.J. Kelly and S.J. Parke, Matter Density Profile Shape Effects at DUNE, Phys. Rev. D 98 (2018) 015025 [1802.06784].
- [39] P. Huber and T. Schwetz, A Low energy neutrino factory with non-magnetic detectors, Phys. Lett. B 669 (2008) 294 [0805.2019].
- [40] B. Yaeggy, Structure Functions and Tau Neutrino Cross Section at DUNE Far Detector, Phys. Sci. Forum 8 (2023) 64.
- [41] P. Huber, M. Lindner and W. Winter, Simulation of long-baseline neutrino oscillation experiments with GLoBES (General Long Baseline Experiment Simulator), Comput. Phys. Commun. 167 (2005) 195 [hep-ph/0407333].
- [42] P. Huber, J. Kopp, M. Lindner, M. Rolinec and W. Winter, New features in the simulation of neutrino oscillation experiments with GLoBES 3.0: General Long Baseline Experiment Simulator, Comput. Phys. Commun. 177 (2007) 432 [hep-ph/0701187].
- [43] M. Masud, M. Bishai and P. Mehta, Extricating New Physics Scenarios at DUNE with Higher Energy Beams, Sci. Rep. 9 (2019) 352 [1704.08650].
- [44] M. Masud, S. Roy and P. Mehta, Correlations and degeneracies among the NSI parameters with tunable beams at DUNE, Phys. Rev. D 99 (2019) 115032 [1812.10290].
- [45] A. De Gouvêa, K.J. Kelly, G.V. Stenico and P. Pasquini, *Physics with Beam Tau-Neutrino Appearance at DUNE*, *Phys. Rev. D* 100 (2019) 016004 [1904.07265].
- [46] J. Rout, S. Roy, M. Masud, M. Bishai and P. Mehta, Impact of high energy beam tunes on the sensitivities to the standard unknowns at DUNE, Phys. Rev. D 102 (2020) 116018 [2009.05061].
- [47] A. Ghoshal, A. Giarnetti and D. Meloni, On the role of the ν_{τ} appearance in DUNE in constraining standard neutrino physics and beyond, JHEP 12 (2019) 126 [1906.06212].
- [48] K. Siyeon, S. Kim, M. Masud and J. Park, Probing large extra dimension at DUNE using beam tunes, JHEP 11 (2024) 141 [2409.08620].

- [49] I. Esteban, M.C. Gonzalez-Garcia, M. Maltoni, T. Schwetz and A. Zhou, The fate of hints: updated global analysis of three-flavor neutrino oscillations, JHEP 09 (2020) 178 [2007.14792].
- [50] I. Esteban, M.C. Gonzalez-Garcia, M. Maltoni, I. Martinez-Soler, J.a.P. Pinheiro and T. Schwetz, NuFit-6.0: updated global analysis of three-flavor neutrino oscillations, JHEP 12 (2024) 216 [2410.05380].
- [51] T2K collaboration, Measurements of neutrino oscillation parameters from the T2K experiment using 3.6 × 10²¹ protons on target, Eur. Phys. J. C 83 (2023) 782 [2303.03222].
- [52] NOVA collaboration, Improved measurement of neutrino oscillation parameters by the NOvA experiment, Phys. Rev. D 106 (2022) 032004 [2108.08219].
- [53] NOVA collaboration, Latest Three-Flavor Neutrino Oscillation Results from NOvA, .
- [54] P.B. Denton and J. Gehrlein, Solar parameters in long-baseline accelerator neutrino oscillations, JHEP 06 (2023) 090 [2302.08513].
- [55] L. Wolfenstein, Neutrino Oscillations in Matter, Phys. Rev. D 17 (1978) 2369.
- [56] T. Ohlsson, Status of non-standard neutrino interactions, Rept. Prog. Phys. 76 (2013) 044201 [1209.2710].
- [57] O.G. Miranda and H. Nunokawa, Non standard neutrino interactions: current status and future prospects, New J. Phys. 17 (2015) 095002 [1505.06254].
- [58] K.S. Babu, P.S.B. Dev, S. Jana and A. Thapa, Non-Standard Interactions in Radiative Neutrino Mass Models, JHEP 03 (2020) 006 [1907.09498].
- [59] D.V. Forero and W.-C. Huang, Sizable NSI from the SU(2)_L scalar doublet-singlet mixing and the implications in DUNE, JHEP 03 (2017) 018 [1608.04719].
- [60] P.B. Denton, Y. Farzan and I.M. Shoemaker, Activating the fourth neutrino of the 3+1 scheme, Phys. Rev. D 99 (2019) 035003 [1811.01310].
- [61] U.K. Dey, N. Nath and S. Sadhukhan, Non-Standard Neutrino Interactions in a Modified ν2HDM, Phys. Rev. D 98 (2018) 055004 [1804.05808].
- [62] K.S. Babu, A. Friedland, P.A.N. Machado and I. Mocioiu, *Flavor Gauge Models Below the Fermi Scale*, JHEP 12 (2017) 096 [1705.01822].
- [63] Y. Farzan and J. Heeck, Neutrinophilic nonstandard interactions, Phys. Rev. D 94 (2016) 053010 [1607.07616].
- [64] Y. Farzan and I.M. Shoemaker, Lepton Flavor Violating Non-Standard Interactions via Light Mediators, JHEP 07 (2016) 033 [1512.09147].
- [65] Y. Farzan, A model for large non-standard interactions of neutrinos leading to the LMA-Dark solution, Phys. Lett. B 748 (2015) 311 [1505.06906].
- [66] P. Coloma, P.B. Denton, M.C. Gonzalez-Garcia, M. Maltoni and T. Schwetz, Curtailing the Dark Side in Non-Standard Neutrino Interactions, JHEP 04 (2017) 116 [1701.04828].
- [67] J. Gehrlein, P.A.N. Machado and J.a.P. Pinheiro, Constraining non-standard neutrino interactions with neutral current events at long-baseline oscillation experiments, 2412.08712.
- [68] P.B. Denton and J. Gehrlein, New constraints on the dark side of non-standard interactions from reactor neutrino scattering data, Phys. Rev. D 106 (2022) 015022 [2204.09060].

- [69] B. Pontecorvo, Inverse beta processes and nonconservation of lepton charge, Zh. Eksp. Teor. Fiz. 34 (1957) 247.
- [70] Z. Maki, M. Nakagawa and S. Sakata, Remarks on the unified model of elementary particles, Prog. Theor. Phys. 28 (1962) 870.
- [71] O.G. Miranda, M.A. Tortola and J.W.F. Valle, Are solar neutrino oscillations robust?, JHEP 10 (2006) 008 [hep-ph/0406280].
- [72] F.J. Escrihuela, O.G. Miranda, M.A. Tortola and J.W.F. Valle, Constraining nonstandard neutrino-quark interactions with solar, reactor and accelerator data, Phys. Rev. D 80 (2009) 105009 [0907.2630].
- [73] P. Bakhti and Y. Farzan, Shedding light on LMA-Dark solar neutrino solution by medium baseline reactor experiments: JUNO and RENO-50, JHEP 07 (2014) 064 [1403.0744].
- [74] P. Coloma and T. Schwetz, Generalized mass ordering degeneracy in neutrino oscillation experiments, Phys. Rev. D 94 (2016) 055005 [1604.05772].
- [75] P.B. Denton and S.J. Parke, Parameter symmetries of neutrino oscillations in vacuum, matter, and approximation schemes, Phys. Rev. D 105 (2022) 013002 [2106.12436].
- [76] J. Liao, D. Marfatia and K. Whisnant, Degeneracies in long-baseline neutrino experiments from nonstandard interactions, Phys. Rev. D 93 (2016) 093016 [1601.00927].
- [77] S.S. Chatterjee, P.S.B. Dev and P.A.N. Machado, Impact of improved energy resolution on DUNE sensitivity to neutrino non-standard interactions, JHEP 08 (2021) 163 [2106.04597].
- [78] P.B. Denton, A. Giarnetti and D. Meloni, How to identify different new neutrino oscillation physics scenarios at DUNE, JHEP 02 (2023) 210 [2210.00109].
- [79] P. Coloma, Non-Standard Interactions in propagation at the Deep Underground Neutrino Experiment, JHEP 03 (2016) 016 [1511.06357].
- [80] V.A. Kostelecky and S. Samuel, Spontaneous Breaking of Lorentz Symmetry in String Theory, Phys. Rev. D 39 (1989) 683.
- [81] V.A. Kostelecky and S. Samuel, Phenomenological Gravitational Constraints on Strings and Higher Dimensional Theories, Phys. Rev. Lett. 63 (1989) 224.
- [82] V.A. Kostelecky and S. Samuel, Gravitational Phenomenology in Higher Dimensional Theories and Strings, Phys. Rev. D 40 (1989) 1886.
- [83] M.M. Sheikh-Jabbari, C, P, and T invariance of noncommutative gauge theories, Phys. Rev. Lett. 84 (2000) 5265 [hep-th/0001167].
- [84] S.M. Carroll, J.A. Harvey, V.A. Kostelecky, C.D. Lane and T. Okamoto, Noncommutative field theory and Lorentz violation, Phys. Rev. Lett. 87 (2001) 141601 [hep-th/0105082].
- [85] O.W. Greenberg, CPT violation implies violation of Lorentz invariance, Phys. Rev. Lett. 89 (2002) 231602 [hep-ph/0201258].
- [86] G. Barenboim and J.D. Lykken, A Model of CPT Violation for Neutrinos, Phys. Lett. B 554 (2003) 73 [hep-ph/0210411].
- [87] A. De Gouvea, Can a CPT Violating Ether Solve ALL Electron (Anti)Neutrino Puzzles?, Phys. Rev. D 66 (2002) 076005 [hep-ph/0204077].
- [88] S.-F. Ge and H. Murayama, Apparent CPT Violation in Neutrino Oscillation from Dark Non-Standard Interactions, 1904.02518.

- [89] M.A. Tórtola, G. Barenboim and C.A. Ternes, CPT and CP, an entangled couple, JHEP 07 (2020) 155 [2005.05975].
- [90] T.V. Ngoc, S. Cao, N.T.H. Van and P.T. Quyen, Stringent constraint on CPT violation with the synergy of T2K-II, NOvA extension, and JUNO, Phys. Rev. D 107 (2023) 016013 [2210.13044].
- [91] G. Barenboim, P. Martínez-Miravé, C.A. Ternes and M. Tórtola, Neutrino CPT violation in the solar sector, Phys. Rev. D 108 (2023) 035039 [2305.06384].
- [92] G. Barenboim, C.A. Ternes and M. Tórtola, Neutrinos, DUNE and the world best bound on CPT invariance, Phys. Lett. B 780 (2018) 631 [1712.01714].
- [93] A. de Gouvêa and K.J. Kelly, Neutrino vs. Antineutrino Oscillation Parameters at DUNE and Hyper-Kamiokande, Phys. Rev. D 96 (2017) 095018 [1709.06090].
- [94] R. Majhi, D.K. Singha, K.N. Deepthi and R. Mohanta, Constraining CPT violation with Hyper-Kamiokande and ESSnuSB, Phys. Rev. D 104 (2021) 055002 [2101.08202].
- [95] SUPER-KAMIOKANDE collaboration, Solar neutrino measurements using the full data period of Super-Kamiokande-IV, Phys. Rev. D 109 (2024) 092001 [2312.12907].
- [96] RENO collaboration, Recent Results from RENO, PoS ICHEP2020 (2021) 177.
- [97] DOUBLE CHOOZ collaboration, Precision Neutrino Mixing Angle Measurement with the Double Chooz Experiment and Latest Results, PoS **TAUP2023** (2024) 228.