# Generalization Certificates for Adversarially Robust Bayesian Linear Regression

Mahalakshmi Sabanayagam<sup>1</sup>

**Russell Tsuchida**\*3

Cheng Soon Ong<sup>4,5</sup>

Debarghya Ghoshdastidar<sup>1,2</sup>

<sup>1</sup>School of Computation, Information and Technology, Technical University of Munich, Germany
 <sup>2</sup>Munich Data Science Institute, Technical University of Munich, Germany
 <sup>3</sup>Monash University, <sup>4</sup>Data61, CSIRO, <sup>5</sup>College of Systems and Society, Australian National University, Australia

sabanaya@in.tum.de, russell.tsuchida@monash.edu, chengsoon.ong@anu.edu.au, ghoshdas@in.tum.de

#### Abstract

Adversarial robustness of machine learning models is critical to ensuring reliable performance under data perturbations. Recent progress has been on point estimators, and this paper considers distributional predictors. First, using the link between exponential families and Bregman divergences, we formulate an adversarial Bregman divergence loss as an adversarial negative log-likelihood. Using the geometric properties of Bregman divergences, we compute the adversarial perturbation for such models in closed-form. Second, under such losses. we introduce *adversarially robust posteriors*, by exploiting the optimization-centric view of generalized Bayesian inference. Third, we derive the first rigorous generalization certificates in the context of an adversarial extension of Bayesian linear regression by leveraging the PAC-Bayesian framework. Finally, experiments on real and synthetic datasets demonstrate the superior robustness of the derived adversarially robust posterior over Bayes posterior, and also validate our theoretical guarantees.

# **1** INTRODUCTION

Machine learning models are vulnerable to adversarial inputs, where small, carefully crafted perturbations to the input data can significantly degrade model performance. These perturbations, though imperceptible to humans (e.g. in computer vision contexts), can cause models to make incorrect predictions with high confidence. Significant progress has been made in understanding and improving adversarial robustness of point predictors, with efforts in defense mechanisms, attack strategies, and the trade-offs between robustness and generalization [Szegedy et al., 2014, Shafahi et al., 2019, Li et al., 2023]. A key insight from this body of research is that models susceptible to adversarial attacks often exhibit near-perfect empirical generalization — achieving similar performance on training and test data [Goodfellow et al., 2015]. This observation suggests that deriving formal guarantees is critical to understanding the interplay between generalization and adversarial robustness.

Probabilistic models offer an alternative paradigm that quantifies uncertainty, a property that can detect adversarial inputs and reject uncertain predictions. However, despite these advantages, probabilistic models have received significantly less attention than their non-probabilistic counterparts in adversarial settings [Bradshaw et al., 2017, Grosse et al., 2018]. While they have been studied for robustness against outliers [Kim and Ghahramani, 2008], label noise [Hernández-Lobato et al., 2011], and domain shifts [Ovadia et al., 2019], their susceptibility to adversarial attacks remains largely unexplored. There exists no notion of adversarially robust probabilistic inference, and hence no formal generalization guarantees, thus raising a fundamental question:

How can we define and develop adversarially robust probabilistic inference and derive generalization certificates (formal guarantees) for such models?

In this work, we address this question by first introducing the notion of adversarially robust posteriors. We achieve this by formulating an adversarial variant of the negative log-likelihood (NLL) loss - drawing inspiration from adversarial training, one of the most effective defense strategies against adversarial attacks in standard machine learning [Madry et al., 2018] — and taking an optimizationcentric perspective of (generalized) Bayesian inference [Alquier et al., 2016]. In doing so, we obtain a posterior that is robust to adversarial perturbations. We leverage the PAC-Bayesian framework, a powerful tool for deriving datadependent generalization bounds for Bayesian predictors [McAllester, 1998, Catoni and Picard, 2004], and derive the first rigorous certificates for the robust posterior on linear regression. Furthermore, we also derive PAC-Bayesian based generalization certificates for Bayes posterior obtained using the standard negative log-likelihood loss.

<sup>\*</sup>work partially done while at Data61, CSIRO

	Standard generalization NLL $\ell$ $R(\theta) = \underset{\mathcal{D} \sim \mathcal{P}}{\mathbb{E}} [\ell(\theta, \mathcal{D})]$	Adversarial generalization Adversarial NLL $\ell_{\hat{\delta}}$ $R_{\hat{\delta}}(\theta) = \mathop{\mathbb{E}}_{\mathcal{D}\sim\mathcal{P}} \left[\ell_{\hat{\delta}}(\theta, \mathcal{D})\right]$
Bayes posterior q	$\mathbb{E}_{\theta \sim q} \left[ R(\theta) \right]$ Theorem 4.3	$\frac{\mathbb{E}}{\theta \sim q} \left[ R_{\hat{\delta}}(\theta) \right) \right]$ Theorem 4.4
Robust posterior $q_{\delta}$	$\frac{\mathbb{E}}{\substack{\theta \sim q_{\delta}}} [R(\theta)]$ Theorem 4.5	$\frac{\mathbb{E}}{_{\theta \sim q_{\delta}}} \left[ R_{\hat{\delta}}(\theta) \right]$ Theorems 4.6 and 4.7

Table 1: Overview of our derived generalization certificates. The guarantees are derived for the standard *Bayes posterior* q and the novel *robust posterior*  $q_{\delta}$ , where  $\delta$  denotes the training adversarial allowance. We consider generalization to standard NLL loss (*standard generalization*) as well as adversarial NLL (*adversarial generalization*), with a potentially different adversarial allowance  $\hat{\delta}$ .

In summary, our main contributions are as follows.

- (i) In exponential families, we review a one-to-one correspondence between the adversarial negative log-likelihood loss and the class of Bregman divergences. Based on this correspondence, we introduce a novel adversarial negative log-likelihood loss in Section 3. This probabilistic-geometric connection allows us to solve the adversarial perturbation problem in closed-form, for all exponential families, allowing for an adversarially robust formulation of generalized linear models.
- (ii) We define the adversarially *robust posterior* as minimizing a variational objective with this loss, thereby extending Bayesian inference to adversarially robust generalized linear models settings, in Section 3.
- (iii) In Section 4, focusing on the case of a Gaussian family, we derive the PAC-Bayesian generalization certificates for the Bayes posterior and the robust posterior under two settings: a) standard generalization: guarantees for standard negative log-likelihood loss (Theorems 4.3 and 4.4), and b) adversarial generalization: guarantees for adversarial negative log-likelihood loss (Theorems 4.5 to 4.7). Table 1 gives an overview of our bounds. We experimentally validate the derived certificates in Section 5 showing non-trivial guarantees.

We discuss the practical significance of the bounds, several technicalities, and touch on related works in Section 6; and conclude in Section 7.

# **2 PRELIMINARIES**

Our work combines elements from adversarial robustness and probabilistic inference. We briefly outline these topics here, and establish some preliminaries.

**Notation** We represent the entry-wise absolute value of matrix M as |M|, and vector Euclidean norm as  $\|\cdot\|$ . We use  $I_n$  for identity matrix of size  $n \times n$  and  $1_n$  for a vector of size n with all ones. We denote by  $\ell(\theta, (x, y))$  a loss

evaluated on parameter  $\theta$  and single (x, y) data pair, and use  $\mathcal{L}(\theta, \mathcal{D}) = \sum_{i=1}^{n} \ell(\theta, (x_i, y_i))$  for the sum of the losses over a dataset. The expected and empirical average errors are  $R(\theta) = \underset{(x,y)\sim\mathcal{P}}{\mathbb{E}} \left[\ell(\theta, (x, y))\right]$  and  $r(\theta) = \frac{1}{n}\mathcal{L}(\theta, \mathcal{D})$ .

#### 2.1 ADVERSARIAL ROBUSTNESS

We are given *n* labeled data samples drawn i.i.d. from an unknown probability measure  $\mathcal{P}$ , denoted by  $\mathcal{D} = \{(x_i, y_i)\}_{i=1}^n$  with  $x_i \in \mathbb{X} \subseteq \mathbb{R}^d$  representing the feature vector with the corresponding label  $y_i \in \mathbb{R}$ . We later use  $X \in \mathbb{R}^{n \times d}$  to refer to the matrix of all feature vectors, and  $Y \in \mathbb{R}^n$  for the vector of all *n* labels. We follow the standard setting of supervised learning [Bishop, 2007, Deisenroth et al., 2020], where we minimize the empirical loss  $\ell$ on the training set  $\mathcal{D} = (X, Y) = \{x_i, y_i\}_{i=1}^n$  with respect to the parameters  $\theta$  of our model,

$$\theta^* = \underset{\theta}{\operatorname{argmin}} \sum_{i=1}^n \ell(\theta, (x_i, y_i)) = \underset{\theta}{\operatorname{argmin}} \mathcal{L}(\theta, \mathcal{D}).$$

For the adversarially robust setting, we consider perturbed data  $\tilde{x}_i$  but not perturbed labels. Following typical adversarial constructs [Szegedy et al., 2014, for example] we consider perturbations whose  $\ell_2$  distance is bounded by a user-defined constant  $\delta$ .

$$\theta^* = \underset{\theta}{\operatorname{argmin}} \sum_{i=1}^n \max_{\|\widetilde{x}_i - x_i\| \le \delta} \ell \Big( \theta, (\widetilde{x}_i, y_i) \Big).$$

Here we focus on the parametric supervised setting, where a parameter  $\theta$  is mapped to a prediction  $f_{\theta}(x_i)$  on a feature  $x_i$  with some parameterized function  $f_{\theta} : \mathbb{X} \to \mathbb{F}$ .

#### 2.2 PROBABILISTIC INFERENCE

**Bayesian inference** Of central interest in Bayesian inference is the posterior  $q(\theta \mid D)$ . The posterior reflects a belief of an unknown quantity of interest  $\theta$  updated from a prior belief  $p(\theta)$  in light of the likelihood  $p(D \mid \theta) = \prod_{i=1}^{n} p(y_i \mid x_i, \theta)$  of observations under the model. One way to compute this update is via Bayes' rule,  $q(\theta \mid D) \propto p(D \mid \theta)p(\theta)$ . An alternate optimization-centric perspective of Bayesian inference, introduced by Csiszár [1975], Donsker and Varadhan [1983], reformulates the objective of deriving the Bayesian posterior as solving an optimization problem. Specifically, the Bayesian posterior distribution  $q(\theta \mid D)$  is obtained by minimizing a variational objective,

$$q(\theta \mid \mathcal{D}) = \arg\min_{\rho \in \Pi} \mathbb{E}_{\theta \sim \rho} \left[ -\log p(\mathcal{D} \mid \theta) \right] + KL(\rho \| \pi),$$
(1)

where  $\Pi$  is the space of all probability measures,  $\pi$  is the prior on  $\theta$ ,  $-\log p(\mathcal{D} \mid \theta)$  is the negative log-likelihood loss on the data, and  $KL(\rho \parallel \pi) = \underset{\theta \sim \rho}{\mathbb{E}} \left[ \log \frac{\rho(\theta)}{\pi(\theta)} \right]$  is the Kullback-Leibler (KL) divergence.

Generalized Bayesian inference Unfortunately, even under the optimization-centric view, Bayesian inference suffers from some limitations. First, the normalizing constant and/or optimization problem can be intractable. Second, the prior is often chosen for convenience and, particularly in large models, may not be truly calibrated to the statistician's prior beliefs. Third, the likelihood is often also chosen for convenience, very often corresponding with losses which are not robust. The Rule of Three (ROT) [Knoblauch et al., 2022] generalizes standard Bayesian inference via the optimization view, addressing the limitations above. Generalizing (1), the ROT replaces the negative log likelihood (NLL) with an arbitrary loss function  $\mathcal{L}$ , the KL divergence KL with an arbitrary divergence D, and the space  $\Pi$  of all probability measures with a subset of all probability measures  $\Lambda$ ,

$$q(\theta \mid \mathcal{D}) = \arg\min_{\rho \in \Lambda} \mathop{\mathbb{E}}_{\theta \sim \rho} \left[ \mathcal{L}(\theta, \mathcal{D}) \right] + D(\rho \| \pi).$$
(2)

The ROT has axiomatic foundations and also comes with guarantees on estimation procedures. The variational objective balances two competing terms: (i) the expected loss term, which encourages the posterior to assign a higher probability to parameters that fit the data well, and (ii) the divergence term, which regularizes the posterior by penalizing deviations from the prior.

**Gibbs Bayesian inference** As a special case of (2), the Gibbs posterior addresses the problem of mis-specified and non-robust likelihoods, and also partially addresses the problem of intractability. The Gibbs posterior is obtained by retaining the KL divergence and space of all probability measures II from (1) in (2), but using a general loss  $\mathcal{L}(\theta, \mathcal{D})$  in place of the NLL - log  $p(\mathcal{D} | \cdot)$ . In this case, the minimizer (2) is called the *Gibbs posterior*, and admits a closed-form (up to the normalizing constant) [Alquier et al., 2016, Knoblauch et al., 2022, for example],

$$q(\theta \mid \mathcal{D}) = \arg\min_{\rho \in \Pi} \mathop{\mathbb{E}}_{\theta \sim \rho} \left[ \mathcal{L}(\theta, \mathcal{D}) \right] + KL(\rho \| \pi)$$
$$= \frac{\exp\left( - \mathcal{L}(\theta, \mathcal{D}) \right) \pi(\theta)}{\int \exp\left( - \mathcal{L}(\theta', \mathcal{D}) \right) \pi(\theta') d\theta'}.$$
(3)

**Gaussian linear regression** A notable special case arises when the loss is chosen as the negative log-likelihood  $\mathcal{L}(\theta, \mathcal{D}) = -\log p(\mathcal{D} \mid \theta) = \sum_{i=1}^{n} -\log p(y_i \mid x_i, \theta)$ with isotropic Gaussian prior  $\pi \sim \mathcal{N}(0, \sigma_p^2 I_d)$ , as it recovers the standard Bayes posterior  $q(\theta) = \mathcal{N}(\hat{\theta}, \Sigma)$ , where  $\Sigma = \frac{1}{\sigma^2} X^T X + \frac{1}{\sigma_p^2} I_d$  and  $\hat{\theta} = \frac{1}{\sigma^2} \Sigma^{-1} X^T Y$  [Bishop, 2007]. In this work, we consider an isotropic Gaussian prior of mean zero and variance  $\sigma_p^2: \theta \sim \mathcal{N}(0, \sigma_p^2 I)$  and denote the negative log-likelihood loss on  $\mathcal{D}$  as  $\ell(\theta, \mathcal{D})$  which is

$$\ell(\theta, \mathcal{D}) = \frac{n}{2} \log \left( 2\pi\sigma^2 \right) + \frac{1}{2\sigma^2} \|Y - X\theta\|^2.$$

## 2.3 LINKING LOSS FUNCTIONS AND PROBABILITY DISTRIBUTIONS

We consider probabilistic models  $p(y | f_{\theta}(x))$  that belong to an exponential family and associate the NLL with the notion of empirical loss  $\ell$  via Bregman divergences, and vice versa. When such losses are later subject to adversarial perturbation, this allows us to make use of the geometrical properties of the Bregman divergence (more specifically, the law of cosines) to study adversarial extensions of probabilistic models in Lemmas 3.1 and 3.2.

**Exponential families** Generalizing Gausssian families, exponential families provide a flexible and theoretically tractable class of probability distributions [Deisenroth et al., 2020, § 6.6.3]. For our purposes, it suffices to consider 1 dimensional (and therefore minimal) exponential families. Let  $t : \mathbb{F} \to \mathbb{R}$  be a measurable function called a *sufficient statistic*. Let  $\mu$  be a nonnegative measure, called the *base measure*, defined on some appropriate sigma algebra generated by  $\mathbb{F}$ . An exponential family is the set of all probability distributions (with respect to base measure  $\mu$ ) parameterized by natural parameter  $\eta$  of the form

$$p(y \mid \eta) = \exp\left(\eta t(y) - \phi(\eta)\right)$$

where  $\phi(\eta) = \log \int_{\mathbb{F}} \exp(\eta t(y)) \mu(dy)$  is called the log normalizing constant, such that  $\phi(\eta) \in \mathbb{R}$ . We assume an extremely mild condition on exponential families, that they are *regular*. Regular means that the set of all  $\eta$  such that  $\phi(\eta) \in \mathbb{R}$  is an open set. Proposition 2 of Wainwright et al. [2008] then states that  $\phi$  is a strictly convex function.

**Bregman divergence** Generalizing the squared Euclidean distance, Bregman divergences allow for a natural class of loss functions for use in a wide variety of supervised and unsupervised applications. Assume  $\mathbb{F}$  is a convex set and let  $\phi : \mathbb{F} \to \mathbb{R}$  be a continuously differentiable and strictly convex function (so called generator). The *Bregman divergence*  $d_{\phi} : \mathbb{F} \times \mathbb{F} \to \mathbb{R}$  generated by  $\phi$  is defined by

$$d_{\phi}(y_1, y_2) = \phi(y_1) - \phi(y_2) - \nabla \phi(y_2)^{\top} (y_1 - y_2),$$

and is strictly convex in its first argument.

A link between geometric loss functions in Bregman divergences and probabilistic loss functions in NLLs is provided through the fact that (informally speaking) every NLL of an exponential family is a Bregman divergence. More precisely, in our current context, if  $p(y | f_{\theta}(x))$  belongs to a regular exponential family with log normalizing function  $\phi$  and natural parameter  $\eta = f_{\theta}(x)$ , then by Banerjee et al. [2005, Theorem 4],

$$-\log p(y \mid f_{\theta}(x)) = d_{\phi}(f_{\theta}(x), y^{*}) + C(y)$$
(4)

where C(y) is an additive constant independent of  $\theta$  and x available in closed-form, and  $y^* = (\nabla \phi)^{-1}(y)$  is the dual

coordinate of y. Note that some technical care is required in ensuring that the dual coordinate  $y^*$  lies in the effective domain of the divergence  $d_{\phi}$ , and (4) is a slight abuse of notation since  $y^*$  may be  $\pm \infty$ , but nevertheless the divergence  $d_{\phi}$  itself remains well defined on an appropriate extension of its domain. See Banerjee et al. [2005, Example 8] for an example. The special and uniquely symmetric case of squared Euclidean distance is obtained when  $\phi(y) = ||y||_2^2$ .

# 3 ADVERSARIALLY ROBUST GENERALIZED LINEAR MODELS

In this section, we derive the adversarially robust posterior  $q_{\delta}(\theta)$  when the likelihood is respectively a Gaussian likelihood, and more generally an exponential family likelihood. The result in Lemma 3.2 may be of independent interest for studying adversarially robust models even in the setting of point-estimation. It allows, for example, an adversarially robust extension of logistic regression (binary-valued data), Poisson regression (count-valued data), and exponential or gamma regression (positive-valued data). More generally, any generalized linear model [McCullagh and Nelder, 1989] with canonical link function may be adversarialized.

Adversarial negative log likelihood We consider adversarial losses  $\ell_{\delta}(\theta, D)$  of the form

$$\ell_{\delta}(\theta, (x, y)) = \max_{\|\widetilde{x} - x\|_{2} \le \delta} -\log p(y \mid f_{\theta}(\widetilde{x}))$$
(5)

$$= \max_{\|\widetilde{x} - x\|_2 \le \delta} d_{\psi} \left( f_{\theta}(\widetilde{x}), y^* \right) + C(y), \quad (6)$$

where  $\delta$  controls the allowable perturbation in the features. Considering a linear predictor and Gaussian likelihood (squared error Bregman divergence) allows us to derive the robust loss in closed-form.

**Lemma 3.1** (Robust loss in closed-form for Gaussian likelihood). Under a linear predictor  $f_{\theta}(x) = \theta^{\top} x$  and in the case where the exponential family is a Gaussian family,

$$\ell_{\delta}(\theta, \mathcal{D}) = \frac{n}{2} \log \left(2\pi\sigma^2\right) + \frac{1}{2\sigma^2} \left\| |Y - X\theta| + \delta \|\theta\|_{1_n} \right\|^2$$

and the adversarial perturbation of the sample x is  $\widetilde{x} = \delta \operatorname{sign}(\theta^{\top}x - y) \frac{\theta}{\|\theta\|_2} + x = \delta \operatorname{sign}(\theta^{\top}\widetilde{x} - y) \frac{\theta}{\|\theta\|_2} + x.$ 

More generally, a linear predictor with any exponential family likelihood (Bregman divergence) allows us to derive the robust loss in closed-form.

**Lemma 3.2** (Robust loss in closed-form for exponential family likelihood). Under a linear predictor  $f_{\theta}(x) = \theta^{\top} x$  and an exponential family likelihood, the robust loss is

$$\ell_{\delta} \left( \theta, (x, y) \right)$$
  
= 
$$\max_{s \in \{-1, 1\}} \psi(s\delta \|\theta\|_{2} + \theta^{\top} x) - \psi(\theta^{\top} x) - ys\delta \|\theta\|_{2}$$
  
+ 
$$d_{\psi}(\theta^{\top} x, y^{*}) + C(y),$$

and the adversarial perturbation of the sample x is  $\tilde{x} = \delta \operatorname{sign} \left( \nabla (\psi(\theta^{\top} \tilde{x}) - y) \frac{\theta}{\|\theta\|_2} + x \right)$ .

Note that in the case of a general exponential family likelihood, a trivial maximization problem over  $s \in \{-1, 1\}$ must be solved. All other terms in Lemma 3.2 are available in closed-form. In practice, this optimization problem can be solved by simply evaluating the objective for s = 1and s = -1, and picking the result with the highest value. Lemmas 3.1 and 3.2 are proved in Appendix A.

We are now ready to define our robust posterior for Bayesian generalized linear models.

**Corollary 3.3** (Robust posterior). *The Gibbs posterior* (3) *obtained by setting the loss*  $\mathcal{L}$  *to be an adversarially perturbed exponential family NLL* (5) (*or equivalently, an adversarially perturbed Bregman divergence* (6)) *under a linear model*  $f_{\theta}(x) = \theta^{\top} x$  *is given by* 

$$q_{\delta}(\theta) = \frac{\exp\left(-\sum_{i=1}^{N} \ell_{\delta}(\theta, (x_i, y_i))\right)\pi(\theta)}{\int \exp\left(-\sum_{i=1}^{N} \ell_{\delta}(\theta', (x_i, y_i))\right)\pi(\theta')d\theta'}$$

where  $\ell_{\delta}(\theta, (x_i, y_i))$  is as in Lemma 3.2, or in the special case of a Gaussian (or squared loss), Lemma 3.1.

We note that this notion of a robust posterior is not the only choice, however this choice does lead to tractable losses derived from adversarial likelihoods, and also allows us to derive generalization guarantees in Section 4. See Appendix A for a discussion of other choices.

# 4 STANDARD AND ADVERSARIAL GENERALIZATION CERTIFICATES

In this section, we focus on Bayesian linear regression (i.e. a robustified squared error loss or Gaussian NLL) for the robust posterior in Corollary 3.3. We consider labels generated using a true parameter  $\theta^*$ ,  $y_i = x_i^{\top} \theta^* + \epsilon_i$  where  $\mathbb{E}[x_i] = 0$ ,  $\mathbb{E}[||x_i||^2] = \sigma_x^2$  and  $\epsilon_i \sim \mathcal{N}(0, \sigma^2)$ .

**PAC-Bayesian generalization certificates** Unlike traditional generalization bounds based on uniform convergence such as VC-dimension [Vapnik and Chervonenkis, 1971], Rademacher complexity [Shalev-Shwartz and Ben-David, 2014], and information-theory [Zhang, 2006], PAC-Bayes [McAllester, 1998] focuses on Bayesian predictors rather than a single deterministic hypothesis class. This perspective allows PAC-Bayes to provide *data-dependent* generalization guarantees, that are computed on training samples without relying on the test data. As such, all certificates computed in this section depend on the data X, Y in a non-obvious way. While other approaches based on uniform convergence or information theory result in a worst-case guarantee, PAC-Bayesian offers fine-grained analysis by taking advantage of informed prior choice leading to a tighter certificate. **Data and constants in bounds** In addition to the data X, Y, each of the bounds in this section also rely on constants such as the training and testing perturbation allowances  $\delta$  and  $\hat{\delta}$ . We intuitively describe the role of these data and constant terms following the presentation of each of the bounds, even though the main purpose of the bounds is as computable data-dependent certificates.

Standard and adversarial generalization We derive certificates for both standard and adversarial generalization (columns in Table 1). In order to formalize this, let us define the expected and empirical errors for standard loss  $\ell$ as  $R(\theta) = \mathbb{E}_{(x,y)\sim \mathcal{P}} \left[ \ell(\theta, (x,y)) \right]$  and  $r(\theta) = \frac{1}{n} \mathcal{L}(\theta, \mathcal{D}),$ respectively. Similarly, the expected and empirical adversarial errors under perturbation  $\delta$  are defined as  $R_{\delta}(\theta) =$  $\mathbb{E}_{(x,y)\sim\mathcal{P}}\left[\ell_{\delta}\left(\theta,(x,y)\right)\right] \text{ and } r_{\delta}(\theta) = \frac{1}{n}\mathcal{L}_{\delta}(\theta,\mathcal{D}), \text{ respec-}$ tively. In the context of Bayesian inference, the standard generalization risk certificate quantifies the expected loss of the posterior  $\rho(\theta)$  on unperturbed test data x:  $\underset{\theta \sim \rho}{\mathbb{E}} [R(\theta)]$ . Similarly, the adversarial generalization certificate quantifies the performance of the test data under adversarial perturbation  $\hat{\delta}: \mathop{\mathbb{E}}_{\theta \sim \rho} \left[ R_{\hat{\delta}}(\theta) \right]$ . We derive the certificates by bounding the respective quantity for both the standard Bayes posterior  $q(\theta)$  and adversarially robust posterior  $q_{\delta}(\theta)$ . Note that we allow the perturbation  $\delta$  used for inference (i.e. calculation of  $q_{\delta}(\theta)$ ) to be different to the perturbation  $\delta$  at test-time.

#### 4.1 CUMULANT GENERATING FUNCTION

To derive the standard and adversarial generalization certificates, we leverage the PAC-Bayesian theorem for any loss with bounded cumulant generating function (CGF) in Banerjee and Montúfar [2021]. We state the result in Theorem 4.1, which requires a bounded CGF. We then show that the CGFs of the standard and adversarial losses corresponding with Gaussian NLLs are bounded in Lemma 4.2.

**Theorem 4.1** (Theorem 6 in Banerjee and Montúfar [2021]). Consider data  $\mathcal{D}$  and any loss  $\ell(\theta, (x, y))$  with its corresponding expected and empirical generalization errors  $R(\theta) = \underset{(x,y)\sim\mathcal{P}}{\mathbb{E}} \left[ \ell(\theta, (x, y)) \right]$  and  $r(\theta) = \frac{1}{n}\mathcal{L}(\theta, \mathcal{D})$ , respectively. Let the CGF of the loss  $\psi(t) = \log \mathbb{E} \left[ \exp \left( t \left( \mathbb{E} [\ell] - \ell \right) \right) \right]$  be bounded, where for some constant c > 0,  $t \in (0, 1/c)$ . Then, we have, with probability at least  $1 - \beta$ , for all densities  $\rho(\theta)$ ,

$$\mathop{\mathbb{E}}_{\theta \sim \rho} \left[ R(\theta) \right] \leq \mathop{\mathbb{E}}_{\theta \sim \rho} \left[ r(\theta) \right] + \frac{1}{t} \left[ \frac{\operatorname{KL}(\rho \| \pi) + \log \frac{1}{\beta}}{n} + \psi(t) \right].$$

Before stating the CGF bound for the losses, we first define a *sub-gamma* random variable. A random variable with variance  $s^2$  and scale parameter c is said to be sub-gamma if its CGF  $\psi$  satisfies the following upper bound:

$$\psi(t) \le \frac{s^2 t^2}{2(1-ct)} \text{ for all } 0 < t < 1/c.$$

We state the CGF bounds for both standard and adversarial losses in Lemma 4.2 and provide the proof in Appendix B.

**Lemma 4.2** (CGF bounds for standard and adversarial losses). The standard and adversarial losses are both subgamma with the following variance  $s^2$  and scale factor c. In the case of standard loss,

$$c = \frac{\sigma_p^2 \sigma_x^2}{\sigma^2}, \quad s^2 = \frac{1}{t} \left( cd - ct + 1 + \frac{\sigma_x^2 ||\theta^*||^2}{\sigma^2} \right).$$
(7)

For adversarial loss with  $\hat{\delta}$  perturbation,

$$c = \frac{2\sigma_p^2 \left(\sigma_x^2 + \hat{\delta}^2\right)}{\sigma^2},$$
  

$$s^2 = \frac{2}{t} \left(cd - ct + 1 + \frac{\sigma_x^2 \|\theta^*\|^2}{\sigma^2}\right).$$
(8)

Using the sub-gamma property of the losses and applying their CGF bounds in Theorem 4.1, we derive the standard and adversarial generalizations of Bayes posterior  $q(\theta)$  in Section 4.2, and robust posterior  $q_{\delta}(\theta)$  in Section 4.3, and present the proofs in Appendix C. Each of the bounds depends on parameters c and  $s^2$ , and intuitively larger values of either lead to worse bounds, as the losses are subject to higher variability.

#### 4.2 GENERALIZATION CERTIFICATES FOR BAYES POSTERIOR

Using the sub-gamma property of the standard loss, the certificate for the standard generalization of the Bayes posterior  $q(\theta)$  is derived in Germain et al. [2016] by setting the free variable t in CGF to 1. We restate this result in Theorem 4.3, expressing it explicitly in terms of the data. This contrasts with the formulation in Germain et al. [2016, Corollary 5], where the bound is expressed in terms of the posterior (which in turn depends on the data).

**Theorem 4.3** (Standard generalization of Bayes posterior, adapted from Germain et al. [2016]). Consider c and s as defined in (7) with t = 1,  $\sigma_p^2 < \frac{\sigma^2}{\sigma_x^2}$ ,  $W_d = I_d + \frac{\sigma_p^2}{\sigma^2} X^\top X$  and  $W_n = I_n + \frac{\sigma_p^2}{\sigma^2} X X^\top$ . Then, with probability at least  $1 - \beta$ , we have the following certificate for standard generalization of the Bayes posterior  $q(\theta)$ :

$$\mathbb{E}_{\theta \sim q} \left[ R(\theta) \right] \leq \frac{1}{n} \log \sqrt{\det\left(W_d\right)} + \frac{1}{2n\sigma^2} Y^\top W_n^{-1} Y + \frac{1}{n} \log \frac{1}{\beta} + \frac{s^2}{2(1-c)}.$$
(9)

Increasing sub-gamma variability parameters  $s^2$  and c increase the bound (9), as expected. Informally, the term depending on log det  $W_d$  is a sum of d log eigenvalues of  $W_d$ , and if  $X^{\top}X$  is low-rank, most of these log eigenvalues are close to 0. Hence the first term decreases like 1/n. The other data dependent term is essentially the product of  $Y^{\top}$  and the average training error of ridge regression, which should be small if the dataset is large and the model is well-specified.

Next, we derive the adversarial generalization certificate for Bayes posterior similar to standard generalization.

**Theorem 4.4** (Adversarial generalization of Bayes posterior). Consider c and s as defined in (8) with t = 1,  $\sigma_p^2 < \frac{\sigma^2}{2(\sigma_x^2 + \hat{\delta}^2)}$ ,  $W_d = I_d + \frac{\sigma_p^2}{\sigma^2} X^\top X$  and  $W_n = I_n + \frac{\sigma_p^2}{\sigma^2} X X^\top$ . Then, with probability at least  $1 - \beta$ , we have the following certificate for adversarial generalization of the Bayes posterior  $q(\theta)$ :

$$\mathbb{E}_{\theta \sim q} \left[ R_{\hat{\delta}}(\theta) \right] \leq \frac{2}{n} \log \sqrt{\det \left( W_d \right)} + \frac{1}{n\sigma^2} Y^\top W_n^{-1} Y \\
+ \frac{1}{n} \log \frac{1}{\beta} + \frac{s^2}{2(1-c)} + \frac{d\hat{\delta}^2 \sigma_p^2}{\sigma^2 - 2n\hat{\delta}^2 \sigma_p^2}.$$
(10)

While Theorems 4.3 and 4.4 are upper bounds and are incomparable, we note that the main difference in Theorem 4.4 is the additional constant term dependent on the perturbation radius  $\hat{\delta}$  and the data-dependent terms are scaled by 2. The additional constant term captures the effect of testing the model adversarially, increasing the bound. This term behaves linearly in  $\hat{\delta}$  for small  $\hat{\delta}$ .

#### 4.3 GENERALIZATION CERTIFICATES FOR ROBUST POSTERIOR

First we derive the standard generalization of robust posterior. While this setting may not be of practical interest, we provide the result for completeness.

**Theorem 4.5** (Standard generalization of robust posterior). Consider c and s as defined in (7) with t = 1,  $\sigma_p^2 < \frac{\sigma^2}{\sigma_x^2}$ ,  $k_{\delta} = \frac{2n\delta^2 \sigma_p^2}{\sigma^2} + 1$ ,  $U_d = k_{\delta}I_d + \frac{2\sigma_p^2}{\sigma^2}X^{\top}X$ ,  $U_n = k_{\delta}I_n + \frac{2\sigma_p^2}{\sigma^2}XX^{\top}$ ,  $V_d = k_{\delta}I_d + \frac{\sigma_p^2}{\sigma^2}X^{\top}X$ , and  $V_n = k_{\delta}I_n + \frac{\sigma_p^2}{\sigma^2}XX^{\top}$ . Then, with probability at least  $1 - \beta$ , we have the following certificate for standard generalization of the robust posterior  $q_{\delta}(\theta)$ :

$$\underset{\theta \sim q_{\delta}}{\mathbb{E}} \left[ R(\theta) \right] \leq \frac{2}{n} \log \sqrt{\det(U_d)} + \frac{2}{nk_{\delta}\sigma^2} Y^{\top} U_n^{-1} Y 
- \frac{1}{n} \log \sqrt{\det(V_d)} - \frac{k_{\delta}}{n\sigma^2} Y^{\top} V_n^{-1} Y 
+ \frac{1}{n} \log \frac{1}{\beta} + \frac{s^2}{2(1-c)}.$$
(11)

The terms involving det  $U_d$  and det  $V_d$  are sums of  $d \log$  eigenvalues divided by n, so they scale like 1/n. As in the previous bounds, the remaining data dependent bounds resemble the product of  $Y^{\top}$  and the average error of ridge regression with an effective regularization parameter.

For the robust posterior, we consider the cases  $\hat{\delta} = \delta$  and  $\hat{\delta} \neq \delta$  separately. We derive a tighter bound for the special case when the allowed adversarial perturbation at train and test-time are the same, i.e.,  $\hat{\delta} = \delta$ .

**Theorem 4.6** (Adversarial generalization of robust posterior with  $\hat{\delta} = \delta$ ). Consider c and s as defined in (8) with t = 1,  $\sigma_p^2 < \frac{\sigma^2}{2(\sigma_x^2 + \hat{\delta}^2)}$ ,  $k_{\delta} = \frac{2n\delta^2 \sigma_p^2}{\sigma^2} + 1$ ,  $U_d = k_{\delta}I_d + \frac{2\sigma_p^2}{\sigma^2}X^{\top}X$ , and  $U_n = k_{\delta}I_n + \frac{2\sigma_p^2}{\sigma^2}XX^{\top}$ . Then, with probability at least  $1 - \beta$ , we have the following certificate for adversarial generalization of the robust posterior  $q_{\delta}(\theta)$ :

$$\mathbb{E}_{\theta \sim q_{\delta}} \left[ R_{\delta}(\theta) \right] \leq \frac{1}{n} \log \sqrt{\det\left(U_{d}\right)} + \frac{1}{nk_{\delta}\sigma^{2}}Y^{\top}U_{n}^{-1}Y + \frac{1}{n}\log\frac{1}{\beta} + \frac{s^{2}}{2(1-c)}.$$
(12)

Compared with Theorem 4.5, Theorem 4.6 only includes 1 times the  $U_d$  and  $U_n$  dependent terms, instead of 2 times the  $U_d$  and  $U_n$  dependent terms minus the  $V_d$  and  $V_d$  dependent terms. Empirically, in Section 5, we find that this leads to a favorable bound.

Finally, using a different analysis we derive the adversarial generalization focusing on a general setting where the adversarial perturbation radius at test-time  $\hat{\delta}$  is not the same as the radius used to learn the posterior  $\delta$ .

**Theorem 4.7** (Adversarial generalization of robust posterior). Consider c and s as defined in (8) with t = 1,  $\sigma_p^2 < \frac{1}{4(\sigma_x^2 + \delta^2)}$ ,  $k_\delta = \frac{2n\delta^2 \sigma_p^2}{\sigma^2} + 1$ ,  $U_d = k_\delta I_d + \frac{2\sigma_p^2}{\sigma^2} X^\top X$ , and  $U_n = k_\delta I_n + \frac{2\sigma_p^2}{\sigma^2} X X^\top$ . Then, with probability at least  $1 - \beta$ , we have the certificate for adversarial generalization of the robust posterior  $q_\delta(\theta)$ :

$$\underset{\theta \sim q_{\delta}}{\mathbb{E}} \left[ R_{\hat{\delta}}(\theta) \right] \leq \frac{2}{n} \log \sqrt{\det\left(U_d\right)} + \frac{2}{nk_{\delta}\sigma^2} Y^{\top} U_n^{-1} Y + \frac{1}{n} \log \frac{1}{\beta} + \frac{s^2}{2(1-c)} + \frac{\left(\hat{\delta}^2 - \delta^2\right)\sigma_p^2 d}{\sigma^2 - 2n\left(\hat{\delta}^2 - \delta^2\right)\sigma_p^2}. \tag{13}$$

# **5 EXPERIMENTAL RESULTS**

In this section, we present (i) adversarial robustness of Bayes and robust posteriors on real datasets; (ii) validation of the derived generalization certificates for the posteriors, and compare it to the prior work of [Germain et al., 2016].

Dataset	Standard generalization (NLL) $\ell$		Adversarial generalization (adv-NLL) $\ell_{\hat{\delta}}$	
	Bayes posterior q	Robust posterior $q_{\delta}$	Bayes posterior q	Robust posterior $q_{\delta}$
Abalone	$1.1586 \pm 0.013$	$1.1729{\scriptstyle~\pm 0.015}$	$1.2539 \pm \textbf{0.012}$	$\boldsymbol{1.2178} \pm 0.015$
Air Foil	$1.1656 \pm 0.008$	$1.1690 \pm 0.008$	$1.2194 \pm 0.008$	$1.2175 \pm 0.008$
Air Quality	$0.9665 \pm 0.002$	$0.9670{\scriptstyle~\pm 0.002}$	$0.9826 \pm \textbf{0.003}$	$\textbf{0.9792} \pm 0.002$
Auto MPG	$1.0231  \pm 0.006$	$\boldsymbol{1.0228} \pm 0.007$	$1.0552 \pm 0.006$	$1.0469 \pm 0.008$
California Housing	$1.1193 \pm 0.003$	$1.1267{\scriptstyle~\pm 0.005}$	$1.1910 \pm 0.003$	$1.1769 \pm 0.005$
Energy Efficiency	$\textbf{0.9709} \pm 0.006$	$0.9731  \pm 0.007$	$0.9996 \pm 0.007$	$\textbf{0.9945} \pm 0.008$
Wine Quality	$1.2339{\scriptstyle~\pm~0.007}$	$1.2322 \pm 0.006$	$1.2696 \pm 0.008$	$1.2627 \pm 0.006$

Table 2: Test NLL and adversarial NLL of Bayes and robust posteriors on real datasets. The prior variance is set to  $\sigma_p^2 = \frac{1}{d}$ . The robust posterior is trained with  $\delta = 0.1$ , and adversarial generalization is evaluated using the same training-time perturbation ( $\hat{\delta} = 0.1$ ). The adversarial generalization results demonstrate that the robust posterior  $q_{\delta}$  is consistently more robust than the Bayes posterior q. For both standard and adversarial generalization, the best-performing model is bold.

Code to reproduce all experiments is provided <sup>1</sup>, and implementation and hardware details are given in Appendix D.

ject to the severe robustness-accuracy trade-off commonly observed in non-Bayesian models [Tsipras et al., 2019].

**Datasets and hyperparameters** We consider the following regression datasets with 70-30 train-test split: Abalone [Nash et al., 1994], Air Foil [Brooks et al., 1989], Air Quality [Vito, 2008], Auto MPG [Quinlan, 1993], California Housing [Pedregosa et al., 2011], Energy Efficiency [Tsanas and Xifara, 2012], Wine Ouality [Cortez et al., 2009]. The datasets are standardized to zero mean and unit variance. We provide results for prior variance  $\sigma_p^2 = \{\frac{1}{100}, \frac{1}{9}, \frac{1}{d}\}$ where d is the data feature dimension. For certificate validation, we use a synthetic dataset where the data features  $x \sim \mathcal{N}(0, \sigma_x^2 I_d)$  and  $y = \theta^* x^\top + \epsilon$  with d = 5 and  $\epsilon \sim \mathcal{N}(0, \sigma^2)$  with  $\sigma^2 = \frac{1}{9}$  and  $\|\theta^*\|^2 = 0.5$ . We fix prior variance  $\sigma_p^2 = 0.01$  and consider a range of training samples from 10 to  $10^4$  and  $10^4$  test samples. All results are averaged over 5 seeds and reported with standard deviation. We employ Hamiltonian Monte Carlo (HMC) to efficiently sample from the posterior distribution. Specifically, we utilize the No-U-Turn Sampler (NUTS) [Hoffman et al., 2014], an adaptive variant of HMC that automatically tunes step sizes and trajectory lengths for improved sampling efficiency.

#### **RESULTS ON REAL DATA** 5.1

We present the standard and adversarial generalization results for Bayes and robust posteriors evaluated on real data in Table 2, using a prior variance of  $\sigma_p^2 = \frac{1}{d}$ . Additional results for  $\sigma_p^2 = \{\frac{1}{100}, \frac{1}{9}\}$  are in Appendix E. Our findings clearly demonstrate that the robust posterior  $q_{\delta}$  consistently outperforms the standard Bayes posterior q in terms of adversarial robustness. Moreover, for certain choices of prior, the robust posterior also achieves superior standard generalization (see Tables 2, 4 and 5). This suggests that adversarial robustness in Bayesian models can potentially be enhanced through probabilistic inference and may not always be sub-

dc9034bb2e323a87a7a4

5.2 VALIDATION OF CERTIFICATES

We validate our derived generalization certificates in Figure 1 by plotting the PAC-Bayesian bounds as a function of the number of training samples. Since these bounds provide rigorous upper estimates on the generalization error, they are not directly comparable to each other but rather serve as theoretical guarantees. While the bounds may appear conservative, it is important to note that these are the first rigorous PAC-Bayesian bounds for adversarial robustness. Additionally, we compare our standard generalization bound for the Bayes posterior with the prior work [Germain et al., 2016] in Figure 1 (left). Although both approaches leverage PAC-Bayesian principles, the bound from [Germain et al., 2016] is numerically lower than ours because they approximate the expected training loss in Theorem 4.1 using the empirical loss, while we compute the actual expected training loss.

#### **DISCUSSION AND RELATED WORK** 6

Generalization bounds for probabilistic models Estimating the average generalization performance as the availability of training data increases, commonly referred to as learning curves, has been extensively studied for probabilistic models such as Gaussian processes (GPs) and Bayesian linear models. In GP regression, significant strides have been made in understanding generalization performance over the past two decades. For instance, Sollich and Halees [2002], Sollich [1998] estimated learning curves by bounding the prediction variance, while Opper and Vivarelli [1998] analyzed learning curves through bounds on prediction error. Further developments include Sollich [2001], who investigated learning curves under mismatched models, and Jin et al. [2022], who provided a more realistic analysis by assuming the eigenspectrum of the prior and the eigenexpansion coefficients of the target function follow a power-

<sup>&</sup>lt;sup>1</sup>https://figshare.com/s/



Figure 1: Validation of the derived generalization certificates Theorems 4.3 to 4.6. (left to right) Standard generalization of Bayes posterior with comparison to prior work Germain et al. [2016], standard generalization of the robust posterior, adversarial generalization of the robust posterior.

law distribution. Notably, Williams and Vivarelli [2000] derived non-trivial upper and lower bounds for GPs, offering key insights into their generalization capabilities. More recently, Savvides et al. [2024] bounded both the variance of the predictions and the bias. While learning curves provide valuable average-case insights, they do not offer certificates.

**Generalization certificates and their practical significance** In the context of GPs, PAC-Bayesian bounds are derived for GP classification Seeger [2002] and GP regression [Suzuki, 2012]. Beyond GPs, PAC-Bayesian theory has been instrumental in explaining the generalization capabilities of neural networks where the uniform convergence type of analyses fail [Dziugaite and Roy, 2017, Lotfi et al., 2022]. Practically, these bounds are useful in hyperparameter optimization [Cherian et al., 2020] and improving model training [Reeb et al., 2018, Wang et al., 2023] by directly minimizing the bound.

**Robustness certificates for GPs** While guaranteeing robustness against adversarial perturbations remain a relatively underexplored area, it is important to note that even heuristic methods aimed at improving the robustness of probabilistic models are less developed [Hernández-Lobato et al., 2011, Bradshaw et al., 2017, Grosse et al., 2018]. Moreover, there exists no notion of adversarially robust posteriors or GPs. In the context of GP classification, Blaas et al. [2020] provided robustness guarantees for the standard Bayes posterior by computing upper and lower bounds for the maximum and minimum of GP classification probabilities under adversarial perturbations. Similar analysis of robustness certification for standard GP regression has been investigated in works such as Patane et al. [2022] and Cardelli et al. [2019].

**Other robust posteriors** In addition to alternative robust posteriors under the optimization-centric view of Bayes rule (as discussed in Appendix A) existing and distinct notions of robust posterior are also available. Focusing on the categorical distribution (a special exponential families), Wicker et al. [2021] define a robust likelihood by marginalizing out perturbed softmax probability distributions with respect to

a distribution on the perturbation allowance. This is then used as a standard likelihood inside Bayesian inference to building Bayesian neural networks.

Adversarially robust optimization Although adversarially robust optimization might appear conceptually similar to our proposed adversarially robust posterior formalism, it addresses a fundamentally different problem. In Bayesian optimization, the goal is to select  $x_t$  such that it yields a high value even under adversarial perturbations, i.e., maximizing  $f(\tilde{x}_t)$ , where f is an unknown function [Bogunovic et al., 2018, Kirschner et al., 2020]. The fundamental distinction lies in the fact that f is not explicitly known in adversarially robust optimization. Moreover, the objective of sequentially choosing x to learn the unknown f is different from learning a posterior from a given data that is adversarially robust.

# 7 CONCLUSION

We consider the problem of adversarially robust probabilistic inference. Using the generalized Bayesian inference framework, we propose adversarially robust posteriors. We show that for exponential family models, closed-form adversarial NLLs result in posteriors that are robust to adversarial perturbations. We derive PAC-Bayes generalization bounds for the four cases as summarized in Table 1. The  $2 \times 2$  table corresponds to combinations of the following settings: (*i*) standard NLL  $\ell$ , and adversarial NLL  $\ell_{\hat{\delta}}$ ; (*ii*) the classical Bayes posterior q, and the robust posterior  $q_{\delta}$ . Our experiments validate that the derived PAC-Bayes bounds capture the empirical behavior, and demonstrate that the robust posterior  $q_{\delta}$  consistently improves adversarial robustness.

Our work primarily focuses on Bayesian linear regression, with the exception of Lemma 3.2 which may be of independent interest. This result opens the possibility of extending our analysis to other generalized linear models. We hope that our notion of adversarially robust posterior will lead to further results in other machine learning problems, and provide useful analysis for practical adversarial learning tasks in the real world.

#### Acknowledgements

This work was partially done when MS visited Data61, CSIRO, and the authors would like to thank Peter Caley for facilitating the visit to Australia. Additionally, this research has been supported by the TUM Georg Nemetschek Institute Artificial Intelligence for the Built World.

#### References

- Pierre Alquier, James Ridgway, and Nicolas Chopin. On the properties of variational approximations of gibbs posteriors. *Journal of Machine Learning Research*, 17(236): 1–41, 2016.
- Arindam Banerjee, Srujana Merugu, Inderjit S Dhillon, and Joydeep Ghosh. Clustering with bregman divergences. *Journal of machine learning research*, 6(10), 2005.
- Pradeep Kr. Banerjee and Guido Montúfar. Information complexity and generalization bounds. In *IEEE International Symposium on Information Theory, ISIT*, 2021.
- Eli Bingham, Jonathan P. Chen, Martin Jankowiak, Fritz Obermeyer, Neeraj Pradhan, Theofanis Karaletsos, Rohit Singh, Paul A. Szerlip, Paul Horsfall, and Noah D. Goodman. Pyro: Deep universal probabilistic programming. *Journal of Machine Learning Research*, 2019.
- Christopher M. Bishop. Pattern recognition and machine learning, 5th Edition. Information science and statistics. Springer, 2007. ISBN 9780387310732.
- Arno Blaas, Andrea Patane, Luca Laurenti, Luca Cardelli, Marta Kwiatkowska, and Stephen Roberts. Adversarial robustness guarantees for classification with gaussian processes. In *International Conference on Artificial Intelligence and Statistics*. PMLR, 2020.
- Ilija Bogunovic, Jonathan Scarlett, Stefanie Jegelka, and Volkan Cevher. Adversarially robust optimization with gaussian processes. *Advances in neural information processing systems*, 2018.
- John Bradshaw, Alexander G de G Matthews, and Zoubin Ghahramani. Adversarial examples, uncertainty, and transfer testing robustness in gaussian process hybrid deep networks. *arXiv preprint arXiv:1707.02476*, 2017.
- Thomas Brooks, D. Pope, and Michael Marcolini. Airfoil Self-Noise. UCI Machine Learning Repository, 1989.
- Luca Cardelli, Marta Kwiatkowska, Luca Laurenti, and Andrea Patane. Robustness guarantees for bayesian inference with gaussian processes. In *Proceedings of the AAAI conference on artificial intelligence*, 2019.

- Olivier Catoni and Jean Picard. Statistical learning theory and stochastic optimization ecole d'eté de probabilités de saint-flour xxxi-2001. 2004.
- John J. Cherian, Andrew G. Taube, Robert T. McGibbon, Panagiotis Angelikopoulos, Guy Blanc, Michael Snarski, Daniel D. Richman, John L. Klepeis, and David E. Shaw. Efficient hyperparameter optimization by way of PAC-Bayes bound minimization. *CoRR*, abs/2008.06431, 2020. URL https://arxiv.org/abs/2008.06431.
- Paulo Cortez, A. Cerdeira, F. Almeida, T. Matos, and J. Reis. Wine Quality. UCI Machine Learning Repository, 2009.
- Imre Csiszár. I-divergence geometry of probability distributions and minimization problems. *The annals of probability*, 1975.
- Marc Peter Deisenroth, A Aldo Faisal, and Cheng Soon Ong. *Mathematics for Machine Learning*. Cambridge University Press, 2020.
- Monroe D Donsker and SR Srinivasa Varadhan. Asymptotic evaluation of certain markov process expectations for large time. iv. *Communications on pure and applied mathematics*, 1983.
- Gintare Karolina Dziugaite and Daniel M. Roy. Computing nonvacuous generalization bounds for deep (stochastic) neural networks with many more parameters than training data. In Gal Elidan, Kristian Kersting, and Alexander Ihler, editors, *Proceedings of the Thirty-Third Conference on Uncertainty in Artificial Intelligence, UAI 2017, Sydney, Australia, August 11-15, 2017, 2017.*
- Pascal Germain, Francis Bach, Alexandre Lacoste, and Simon Lacoste-Julien. PAC-Bayesian theory meets Bayesian inference. *Advances in Neural Information Processing Systems*, 2016.
- Ian J. Goodfellow, Jonathon Shlens, and Christian Szegedy. Explaining and harnessing adversarial examples. In Yoshua Bengio and Yann LeCun, editors, *International Conference on Learning Representations*, *ICLR*, 2015.
- Kathrin Grosse, David Pfaff, Michael Thomas Smith, and Michael Backes. The limitations of model uncertainty in adversarial settings. *arXiv preprint arXiv:1812.02606*, 2018.
- Daniel Hernández-Lobato, Jose Hernández-Lobato, and Pierre Dupont. Robust multi-class gaussian process classification. *Advances in neural information processing systems*, 2011.
- Matthew D Hoffman, Andrew Gelman, et al. The no-u-turn sampler: adaptively setting path lengths in hamiltonian monte carlo. *Journal of Machine Learning Research*, 2014.

- Hui Jin, Pradeep Kr. Banerjee, and Guido Montúfar. Learning curves for gaussian process regression with power-law priors and targets. In *The Tenth International Conference on Learning Representations, ICLR*, 2022.
- Hyun-Chul Kim and Zoubin Ghahramani. Outlier robust gaussian process classification. In *Structural, Syntactic,* and Statistical Pattern Recognition: Joint IAPR International Workshop, SSPR & SPR 2008, Orlando, USA, December 4-6, 2008. Proceedings. Springer, 2008.
- Johannes Kirschner, Ilija Bogunovic, Stefanie Jegelka, and Andreas Krause. Distributionally robust bayesian optimization. In *International Conference on Artificial Intelligence and Statistics*. PMLR, 2020.
- Jeremias Knoblauch, Jack Jewson, and Theodoros Damoulas. An optimization-centric view on bayes' rule: Reviewing and generalizing variational inference. *Journal of Machine Learning Research*, 2022.
- Ang Li, Yifei Wang, Yiwen Guo, and Yisen Wang. Adversarial examples are not real features. In Alice Oh, Tristan Naumann, Amir Globerson, Kate Saenko, Moritz Hardt, and Sergey Levine, editors, *Neural Information Processing Systems (NeurIPS)*, 2023.
- Sanae Lotfi, Marc Finzi, Sanyam Kapoor, Andres Potapczynski, Micah Goldblum, and Andrew G Wilson. PAC-Bayes compression bounds so tight that they can explain generalization. *Advances in Neural Information Processing Systems*, 2022.
- Aleksander Madry, Aleksandar Makelov, Ludwig Schmidt, Dimitris Tsipras, and Adrian Vladu. Towards deep learning models resistant to adversarial attacks. In *International Conference on Learning Representations, ICLR*, 2018.
- Kelly Markelle, Longjohn Rachel, and Nottingham Kolby. The uci machine learning repository. URL https:// archive.ics.uci.edu.
- David A McAllester. Some PAC-Bayesian theorems. In *Proceedings of the eleventh annual conference on Computational learning theory*, 1998.
- P. McCullagh and J.A. Nelder. *Generalized Linear Models, Second Edition*. Monographs on Statistics and Applied Probability Series. Chapman & Hall, 1989.
- Warwick Nash, Tracy Sellers, Simon Talbot, Andrew Cawthorn, and Wes Ford. Abalone. UCI Machine Learning Repository, 1994.
- Frank Nielsen. On geodesic triangles with right angles in a dually flat space. In *Progress in Information Geometry: Theory and Applications*, pages 153–190. Springer, 2021.

- Manfred Opper and Francesco Vivarelli. General bounds on bayes errors for regression with gaussian processes. Advances in Neural Information Processing Systems, 1998.
- Yaniv Ovadia, Emily Fertig, Jie Ren, Zachary Nado, David Sculley, Sebastian Nowozin, Joshua Dillon, Balaji Lakshminarayanan, and Jasper Snoek. Can you trust your model's uncertainty? evaluating predictive uncertainty under dataset shift. *Advances in neural information processing systems*, 2019.
- Adam Paszke, Sam Gross, Soumith Chintala, Gregory Chanan, Edward Yang, Zachary DeVito, Zeming Lin, Alban Desmaison, Luca Antiga, and Adam Lerer. Automatic differentiation in PyTorch. In *NIPS-W*, 2017.
- Andrea Patane, Arno Blaas, Luca Laurenti, Luca Cardelli, Stephen Roberts, and Marta Kwiatkowska. Adversarial robustness guarantees for gaussian processes. *Journal of Machine Learning Research*, 2022.
- Fabian Pedregosa, Gaël Varoquaux, Alexandre Gramfort, Vincent Michel, Bertrand Thirion, Olivier Grisel, Mathieu Blondel, Peter Prettenhofer, Ron Weiss, Vincent Dubourg, Jake VanderPlas, Alexandre Passos, David Cournapeau, Matthieu Brucher, Matthieu Perrot, and Edouard Duchesnay. Scikit-learn: Machine learning in python. Journal of Machine Learning Research, 2011.
- R. Quinlan. Auto MPG. UCI Machine Learning Repository, 1993.
- David Reeb, Andreas Doerr, Sebastian Gerwinn, and Barbara Rakitsch. Learning gaussian processes by minimizing PAC-Bayesian generalization bounds. *Advances in Neural Information Processing Systems*, 2018.
- Rafael Savvides, Hoang Phuc Hau Luu, and Kai Puolamäki. Error bounds for any regression model using gaussian processes with gradient information. In *International Conference on Artificial Intelligence and Statistics*. PMLR, 2024.
- Matthias Seeger. PAC-Bayesian generalisation error bounds for gaussian process classification. *Journal of machine learning research*, 2002.
- Ali Shafahi, W. Ronny Huang, Christoph Studer, Soheil Feizi, and Tom Goldstein. Are adversarial examples inevitable? In *International Conference on Learning Representations, ICLR*, 2019.
- Shai Shalev-Shwartz and Shai Ben-David. Understanding Machine Learning: From Theory to Algorithms. Cambridge University Press, 2014.
- Peter Sollich. Learning curves for gaussian processes. Advances in neural information processing systems, 1998.

- Peter Sollich. Gaussian process regression with mismatched models. *Advances in Neural Information Processing Systems*, 2001.
- Peter Sollich and Anason S. Halees. Learning curves for gaussian process regression: Approximations and bounds. *Neural Computation*, 2002.
- Taiji Suzuki. PAC-Bayesian bound for gaussian process regression and multiple kernel additive model. In *Conference on Learning Theory*. JMLR Workshop and Conference Proceedings, 2012.
- Christian Szegedy, Wojciech Zaremba, Ilya Sutskever, Joan Bruna, Dumitru Erhan, Ian J. Goodfellow, and Rob Fergus. Intriguing properties of neural networks. In *International Conference on Learning Representations, ICLR*, 2014.
- Athanasios Tsanas and Angeliki Xifara. Energy Efficiency. UCI Machine Learning Repository, 2012.
- Dimitris Tsipras, Shibani Santurkar, Logan Engstrom, Alexander Turner, and Aleksander Madry. Robustness may be at odds with accuracy. In *International Conference on Learning Representations*, 2019.
- Joaquin Vanschoren, Jan N. van Rijn, Bernd Bischl, and Luis Torgo. Openml: Networked science in machine learning. *SIGKDD Explorations*, 2013.
- VN Vapnik and A Ya Chervonenkis. On the uniform convergence of relative frequencies of events to their probabilities. *Theory of Probability and its Applications*, 1971.
- Saverio Vito. Air Quality. UCI Machine Learning Repository, 2008.
- Martin J Wainwright, Michael I Jordan, et al. Graphical models, exponential families, and variational inference. *Foundations and Trends*® *in Machine Learning*, 1(1–2): 1–305, 2008.
- Zifan Wang, Nan Ding, Tomer Levinboim, Xi Chen, and Radu Soricut. Improving robust generalization by direct PAC-Bayesian bound minimization. In *Proceedings of the IEEE/CVF Conference on Computer Vision and Pattern Recognition*, 2023.
- Matthew Wicker, Luca Laurenti, Andrea Patane, Zhuotong Chen, Zheng Zhang, and Marta Kwiatkowska. Bayesian inference with certifiable adversarial robustness. In *International Conference on Artificial Intelligence and Statistics*, pages 2431–2439. PMLR, 2021.
- Christopher K. I. Williams and Francesco Vivarelli. Upper and lower bounds on the learning curve for gaussian processes. *Machine Learning*, 2000.
- Tong Zhang. Information-theoretic upper and lower bounds for statistical estimation. *IEEE Transactions on Information Theory*, 2006.

# GENERALIZATION CERTIFICATES FOR ADVERSARIALLY ROBUST BAYESIAN LINEAR REGRESSION (SUPPLEMENTARY MATERIAL)

## A ADVERSARIAL ROBUST LOSS

#### A.1 THE ROBUST POSTERIOR

The variational form of exact Bayes inference can be obtained by minimising the KL divergence of the notional posterior q' from the true posterior q over all probability density functions, as follows.

$$q(\theta \mid \mathcal{D}) = \underset{q' \in \Pi}{\operatorname{argmin}} KL(q' \mid q)$$

$$= \underset{q' \in \Pi}{\operatorname{argmin}} \underset{q'(\theta \mid \mathcal{D})}{\mathbb{E}} \left[ \log \frac{q'(\theta \mid \mathcal{D}) \int p(Y \mid X, \theta') \pi(\theta') d\theta'}{p(Y \mid X, \theta) \pi(\theta)} \right]$$

$$= \underset{q' \in \Pi}{\operatorname{argmin}} \underset{q'(\theta \mid \mathcal{D})}{\mathbb{E}} \left[ \sum_{i=1}^{n} -\log \frac{p(y_i \mid x_i, \theta)}{\int p(y_i \mid x_i, \theta) dy_i} \right] + \underbrace{\log \int \prod_{i=1}^{n} p(y_i \mid x_i, \theta') \pi(\theta') d\theta'}_{\operatorname{const. w.r.t. q'}} + KL(q' \mid \pi)$$

$$= \underset{q' \in \Pi}{\operatorname{argmin}} \underset{q'(\theta \mid \mathcal{D})}{\mathbb{E}} \left[ \sum_{i=1}^{n} -\log p(y_i \mid x_i, \theta) \right] + KL(q' \mid \pi).$$
(14)

#### A.2 OTHER NOTIONS OF ROBUST POSTERIOR

Note that in (14), we explicitly convey that they likelihood is a proper likelihood (i.e. integrates to 1) and the posterior is a proper posterior, possessing a normalizing constant which is independent of the functional variable q'. Thus, from an optimisation-centric view, both terms can be ignored. When we generalize to Gibbs Bayes posteriors by changing the negative log likelihood to an adversarial loss, we therefore obtain

$$q(\theta \mid \mathcal{D}) = \underset{q' \in \Pi}{\operatorname{argmin}} \mathbb{E}_{q'(\theta \mid \mathcal{D})} \left[ \sum_{i=1}^{n} \max_{\|\widetilde{x}_{i} - x_{i}\|_{2} \leq \delta} -\log p(y_{i} \mid x_{i}, \theta) \right] + KL(q' \mid \pi) = \frac{\exp\left(-\mathcal{L}(\theta, \mathcal{D})\right) \pi(\theta)}{\int \exp\left(-\mathcal{L}(\theta', \mathcal{D})\right) \pi(\theta') d\theta'}, \quad (15)$$

with  $\mathcal{L}(\theta, \mathcal{D}) = \sum_{i=1}^{n} \max_{\|\tilde{x}_i - x_i\|_2 \leq \delta} -\log p(y_i \mid x_i, \theta).$ 

We note however that this is not the only natural choice of "robustifying" posterior inference. In particular, starting from (14), there are four choices depending on whether we ignore or do not ignore the normalising constants of the likelihood and the posterior when inserting the operator  $\max_{\|\tilde{x}_i - x_i\|_2 \leq \delta}$ . These four choices all lead to the same standard Bayesian posterior (i.e.  $\delta = 0$ ), but lead to different notions of robust posterior. In addition to these four choices, one may also consider robustifying the likelihood before the normalizing step, leading to a true likelihood (and thus, standard Bayesian inference) Our choice (15) allows for a tractable loss term, leads to a satisfying theory, and good empirical performance. Other choices do not immediately lead to a tractable loss term, and we leave this and investigation of their theory and empirical performance for future work.

#### A.3 PROOF OF Lemmas 3.1 and 3.2

*Proof.* We begin with the Gaussian case, then consider the more general exponential family case, and then return to the Gaussian case. Choosing a Gaussian likelihood and a linear predictor, up to some constant,

$$\begin{aligned} \max_{\widetilde{x}_i: \|x_i - \widetilde{x}_i\| \le \delta} -\log p(y_i \mid \widetilde{x}_i, \theta) &= \max_{\widetilde{x}_i: \|x_i - \widetilde{x}_i\| \le \delta} (y_i - \widetilde{x}_i^\top \theta)^2 \\ &= \max_{\widetilde{x}_i: \|x_i - \widetilde{x}_i\| \le \delta} \left( y_i - x_i^\top \theta - \theta^\top (\widetilde{x}_i - x_i) \right)^2 \\ &= \max_{\widetilde{x}_i: \|x_i - \widetilde{x}_i\| \le \delta} \left( y_i - x_i^\top \theta - \|\theta\|_2 \delta \cos \gamma \right)^2, \end{aligned}$$

where  $\gamma$  is the angle between  $\theta$  and  $\tilde{x}_i - x_i$ . The argument of  $(\cdot)^2$  is maximally positive or negative when  $\cos \gamma$  is  $\pm 1$  and shares the same sign as  $y_i - x_i^{\top} \theta$ . Thus

$$\max_{\widetilde{x}_i: \|x_i - \widetilde{x}_i\| \le \delta} -\log p(y_i \mid \widetilde{x}_i, \theta) = \left( |y_i - x_i^\top \theta| + \|\theta\| \delta \right)^2$$
$$= (y_i - x_i^\top \theta)^2 + 2\delta \|\theta\| |y_i - x_i^\top \theta| + \|\theta\|^2 \delta^2,$$

That is,  $\tilde{x}_i = \delta \operatorname{sign}(x_i^{\top} \theta - y_i) \frac{\theta}{\|\theta\|} + x_i.$ 

Consider the Bregman divergence and apply the law of cosines (e.g. Property 1 of Nielsen [2021]),

$$\begin{aligned} \max_{\widetilde{x}: \|x - \widetilde{x}\| \le \delta} d_{\phi} (\theta^{\top} \widetilde{x}, y^{*}) &= \max_{\widetilde{x}: \|x - \widetilde{x}\| \le \delta} d_{\phi} (\theta^{\top} \widetilde{x}, \theta^{\top} x) + d_{\phi} (\theta^{\top} x, y^{*}) - \theta^{\top} (\widetilde{x} - x) (\nabla \phi(y^{*}) - \nabla \phi(\theta^{\top} x)) \\ &= \max_{\widetilde{x}: \|x - \widetilde{x}\| \le \delta} \phi(\theta^{\top} \widetilde{x}) - \phi(\theta^{\top} x) - \nabla \phi(y^{*}) \theta^{\top} (\widetilde{x} - x) + d_{\phi} (\theta^{\top} x, y^{*}). \end{aligned}$$

This is a convex objective on a convex constraint set  $||x - \tilde{x}||_2^2 \le \delta^2$ , so there exists a unique maximum on an extremal point on the constraint set. The KKT conditions give that at the optimal,

$$\left(\nabla(\phi(\theta^{\top}\widetilde{x}) - \nabla\phi(y^{*}))\theta - 2\lambda(\widetilde{x} - x) = 0,\right)$$

for Lagrange multiplier  $\lambda \leq 0$ . Therefore,  $\tilde{x}$  satisfies the implicit equation

$$\widetilde{x} = \frac{\left(\nabla(\phi(\theta^{\top}\widetilde{x}) - \nabla\phi(y^*)\right)\theta}{2\lambda} + x.$$
(16)

We must have the solution on the extremal, so

$$\delta = \Big| \frac{\left( \nabla(\phi(\theta^{\top} \widetilde{x}) - \nabla \phi(y^*) \right)}{2\lambda} \Big| \|\theta\|_2 \quad \text{and so} \quad \lambda = \Big| \frac{\nabla(\phi(\theta^{\top} \widetilde{x}) - \nabla \phi(y^*)}{2\delta} \Big| \|\theta\|_2.$$

Plugging  $\lambda$  back into (16), we find that the optimal  $\tilde{x}$  is a linear combination of  $\theta$  and x,

$$\widetilde{x} = \delta \underbrace{\operatorname{sign}\left(\nabla(\phi(\theta^{\top}\widetilde{x}) - \nabla\phi(y^*)\right)}_{:=s \in \{-1,1\}} \|\theta\|_2^{-1} \theta + x.$$

The maximum value is then

$$\phi(s\delta\|\theta\|_2 + \theta^\top x) - \phi(\theta^\top x) - \nabla\phi(y^*)s\delta\|\theta\|_2 + d_\phi(\theta^\top x, y^*).$$

Finally, note that  $\nabla \phi(y^*) = \nabla \phi((\nabla \phi)^{-1}(y)) = y$ . We may then compute the maximum by testing the two  $s \in \{-1, 1\}$ , and choosing the value of s which gives the maximum result. In the case of Gaussian loss (i.e. squared error), the parameters are self-dual and we have

$$\begin{aligned} \phi(s\delta \|\theta\|_{2} + \theta^{\top}x) &- \phi(\theta^{\top}x) - \nabla\phi(y)s\delta \|\theta\|_{2} + d_{\phi}(\theta^{\top}x,y) \\ &= \delta^{2} \|\theta\|_{2}^{2} + 2s\delta \|\theta\|_{2}(\theta^{\top}x) - 2ys\delta \|\theta\|_{2} + \|y - \theta^{\top}x\|_{2}^{2} \\ &= \delta^{2} \|\theta\|_{2}^{2} + 2s\delta \|\theta\|_{2} ((\theta^{\top}x) - y) + \|y - \theta^{\top}x\|_{2}^{2}, \end{aligned}$$

the maxima being  $\delta^2 \|\theta\|_2^2 + 2\delta \|\theta\|_2 |(\theta^\top x) - y| + \|y - \theta^\top x\|_2^2$  with  $s = \operatorname{sign}(\theta^\top x - y)$ .

# B PROOF OF CGF BOUNDS FOR STANDARD AND ADVERSARIAL NLL LOSSES IN Lemma 4.2

We first derive the following helpful Lemmas B.1 and B.2 to derive the CGF bounds and the generalization certificates.

**Lemma B.1** (Upper and lower bounds for adversarial NLL loss). Using the closed-form of the adversarial NLL loss, the upper and lower bounds are

$$\ell_{\delta}(\theta, \mathcal{D}) = \sum_{i=1}^{n} \left( \frac{1}{2\sigma^{2}} \left( |y_{i} - x_{i}^{\top}\theta| + ||\theta||\delta \right)^{2} + \frac{1}{2} \log \left(2\pi\sigma^{2}\right) \right)$$
  
$$\leq \sum_{i=1}^{n} \left( \frac{1}{2\sigma^{2}} \left( 2(y_{i} - x_{i}^{\top}\theta)^{2} + 2||\theta||^{2}\delta^{2} \right) + \frac{1}{2} \log \left(2\pi\sigma^{2}\right) \right) \quad ; (a-b)^{2} \geq 0 \implies a^{2} + b^{2} \geq 2ab$$
  
$$\ell_{\delta}(\theta, \mathcal{D}) \geq \sum_{i=1}^{n} \left( \frac{1}{2\sigma^{2}} \left( (y_{i} - x_{i}^{\top}\theta)^{2} + ||\theta||^{2}\delta^{2} \right) + \frac{1}{2} \log \left(2\pi\sigma^{2}\right) \right)$$

**Lemma B.2** (Standard Gaussian integral). The integral of the form  $\int \exp(-\theta^{\top} M\theta + 2b^{\top}\theta) d\theta$  evaluates to  $\sqrt{\frac{\pi^d}{\det M}} \exp(b^{\top} M^{-1}b)$ .

Proof.

$$-\theta^{\top}M\theta + 2b^{\top}\theta = -(\theta - M^{-1}b)^{\top}M(\theta - M^{-1}b) + b^{\top}M^{-1}b \quad ; \text{Completing the square}$$
  
Therefore, integral becomes  $\int \exp\left(-(\theta - M^{-1}b)^{\top}M(\theta - M^{-1}b) + b^{\top}M^{-1}b\right)d\theta$ 
$$= \exp\left(b^{\top}M^{-1}b\right)\int \exp\left(-(\theta - M^{-1}b)^{\top}M(\theta - M^{-1}b)\right)d\theta$$
$$= \exp\left(b^{\top}M^{-1}b\right)\sqrt{\frac{\pi^{d}}{\det M}} \quad ; \int \exp\left(-\phi^{\top}M\phi\right)d\phi = \sqrt{\frac{\pi^{d}}{\det M}}$$

**Proof of CGF bounds for standard NLL loss** Now we derive the CGF bound for standard NLL loss in the following. Note that similar derivation is done in Germain et al. [2016].

Proof.

$$\begin{split} &\log \mathbb{E}_{\theta} \mathbb{E}_{x_{i}} \underset{y_{i}|x_{i}}{\mathbb{E}} \left[ \exp \left( t \left( \mathbb{E}_{x_{i}} \underset{y_{i}|x_{i}}{\mathbb{E}} \left[ \frac{1}{2\sigma^{2}} (y_{i} - x_{i}^{\top} \theta)^{2} \right] - \frac{1}{2\sigma^{2}} (y_{i} - x_{i}^{\top} \theta)^{2} \right) \right) \right] \\ &= \log \mathbb{E}_{\theta} \mathbb{E}_{x_{i}} \underset{y_{i}|x_{i}}{\mathbb{E}} \left[ \exp \left( \frac{t}{2\sigma^{2}} \left( \mathbb{E}_{x_{i}} \underset{y_{i}|x_{i}}{\mathbb{E}} \left[ (y_{i} - x_{i}^{\top} \theta)^{2} \right] \right) \right) \right] \\ &\leq \log \mathbb{E}_{\theta} \mathbb{E}_{x_{i}} \underset{y_{i}|x_{i}}{\mathbb{E}} \left[ \exp \left( \frac{t}{2\sigma^{2}} \left( \mathbb{E}_{x_{i}} \underset{y_{i}|x_{i}}{\mathbb{E}} \left[ (y_{i} - x_{i}^{\top} \theta)^{2} \right] \right) \right) \right] \\ &= \log \mathbb{E}_{\theta} \mathbb{E}_{x_{i}} \underset{\epsilon_{i}}{\mathbb{E}} \left[ \exp \left( \frac{t}{2\sigma^{2}} \left( \mathbb{E}_{x_{i}} \underset{x_{i}}{\mathbb{E}} \left[ (x_{i}^{\top} (\theta^{*} - \theta) + \epsilon_{i})^{2} \right] \right) \right) \right] \\ &= \log \mathbb{E}_{\theta} \mathbb{E}_{x_{i}} \underset{\epsilon_{i}}{\mathbb{E}} \left[ \exp \left( \frac{t}{2\sigma^{2}} \left( \sigma_{x}^{2} \| \theta^{*} - \theta \|^{2} + \sigma^{2} \right) \right) \right] \quad ; \epsilon_{i} \sim \mathcal{N}(0, \sigma^{2}), \mathbb{E}[x_{i}] = 0, \mathbb{E}[\|x_{i}\|^{2}] = \sigma_{x}^{2} \\ &= \log \int \exp \left( \frac{t}{2\sigma^{2}} \left( \sigma_{x}^{2} \theta^{*} - 2\sigma_{x}^{2} \theta^{*} \theta^{*} \theta + \sigma_{x}^{2} \| \theta^{*} \|^{2} + \sigma^{2} \right) - \frac{1}{2\sigma_{p}^{2}} \theta^{\top} \theta \right) \frac{1}{\sqrt{2\pi\sigma_{p}^{2}}} d\theta \quad ; \theta \sim \mathcal{N}(0, \sigma_{p}^{2}I) \\ &= \log \int \exp \left( - \left( \frac{1 - t\sigma_{p}^{2}\sigma_{x}^{2}/\sigma^{2}}{2\sigma_{p}^{2}} \right) \theta^{\top} \theta - \frac{t\sigma_{x}^{2}}{\sigma^{2}} \theta^{*} \theta^{*} \theta + \frac{t\sigma_{x}^{2}}{2\sigma^{2}} \| \theta^{*} \|^{2} + \frac{t}{2} \right) \frac{1}{\sqrt{2\pi\sigma_{p}^{2}}} d\theta \end{split}$$

$$= \log \sqrt{\frac{\pi 2\sigma_p^2}{1 - t\sigma_p^2 \sigma_x^2 / \sigma^2}} \exp \left( \frac{t^2 \sigma_x^4 \|\theta^*\|^2 \sigma_p^2 / 2\sigma^2}{1 - t\sigma_p^2 \sigma_x^2 / \sigma^2} + \frac{t\sigma_x^2}{2\sigma^2} \|\theta^*\|^2 + \frac{t}{2} \right) \frac{1}{\sqrt{2\pi\sigma_p^2}} \quad ; t < \frac{\sigma^2}{\sigma_p^2 \sigma_x^2}, \text{Lemma B.2}$$

$$= \frac{d}{2} \log \frac{1}{1 - t\sigma_p^2 \sigma_x^2 / \sigma^2} + \frac{t\sigma_x^2 \|\theta^*\|^2 / 2\sigma^2}{1 - t\sigma_p^2 \sigma_x^2 / \sigma^2} + \frac{t}{2}$$

$$\leq \frac{t\sigma_p^2 \sigma_x^2 d / \sigma^2}{2\left(1 - t\sigma_p^2 \sigma_x^2 / \sigma^2\right)} + \frac{t\sigma_x^2 \|\theta^*\|^2 / \sigma^2}{2\left(1 - t\sigma_p^2 \sigma_x^2 / \sigma^2\right)} + \frac{t}{2} \quad ; -\log(1 - x) \leq \frac{x}{1 - x}$$

$$= \frac{t^2 s^2}{2(1 - tc)}$$

$$\text{above we get } s^2 = \frac{1}{t} \left( \frac{\sigma_p^2 \sigma_x^2 d}{\sigma^2} + \frac{\sigma_x^2 \|\theta^*\|^2}{\sigma^2} + \left(1 - \frac{t\sigma_p^2 \sigma_x^2}{\sigma^2}\right) \right), c = \frac{\sigma_p^2 \sigma_x^2}{\sigma^2} \text{ and } t \in (0, \frac{\sigma^2}{\sigma^2\sigma^2}).$$

From above we get  $s^2 = \frac{1}{t} \left( \frac{\sigma_p^2 \sigma_x^2 d}{\sigma^2} + \frac{\sigma_x^2 \|\theta^*\|^2}{\sigma^2} + \left( 1 - \frac{t\sigma_p^2 \sigma_x}{\sigma^2} \right) \right), c = \frac{\sigma_p^2 \sigma_x^2}{\sigma^2} \text{ and } t \in (0, \frac{\sigma^2}{\sigma_p^2 \sigma_x^2}).$ 

**Proof of CGF bounds for adversarial NLL loss** The cumulant generating function of the adversarial loss can be bounded similar to standard loss as follows.

Proof.

$$\begin{split} &\log \mathbb{E}_{\theta} \left[ \exp \left( \frac{t}{2\sigma^{2}} \mathbb{E}_{x_{i}} \mathbb{E}_{y_{i}|x_{i}} \left[ (y_{i} - x_{i}^{\top}\theta)^{2} + 2\delta \|\theta\| |y_{i} - x_{i}^{\top}\theta| + \|\theta\|^{2}\delta^{2} \right] \right) \right] \\ &\leq \log \mathbb{E}_{\theta} \left[ \exp \left( \frac{t}{2\sigma^{2}} \mathbb{E}_{x_{i}} \mathbb{E}_{y_{i}|x_{i}} \left[ 2(y_{i} - x_{i}^{\top}\theta)^{2} + 2\|\theta\|^{2}\delta^{2} \right] \right) \right] \\ &= \log \mathbb{E}_{\theta} \left[ \exp \left( \frac{t}{2\sigma^{2}} \mathbb{E}_{x_{i}} \mathbb{E}_{i} \left[ 2(x_{i}^{\top} (\theta^{*} - \theta) + \epsilon_{i})^{2} + 2\|\theta\|^{2}\delta^{2} \right) \right] \\ &= \log \mathbb{E}_{\theta} \left[ \exp \left( \frac{t}{\sigma^{2}} \sigma_{x}^{2} \|\theta^{*} - \theta\|^{2} + t + \frac{t}{\sigma^{2}} \|\theta\|^{2}\delta^{2} \right) \right] \\ &= \log \int \exp \left( \left( \frac{t\sigma_{x}^{2}}{\sigma^{2}} + \frac{t\delta^{2}}{\sigma^{2}} - \frac{1}{2\sigma_{p}^{2}} \right) \|\theta\|^{2} - \frac{2t\sigma_{x}^{2}}{\sigma^{2}} \theta^{*\top}\theta + \frac{t\sigma_{x}^{2}}{\sigma^{2}} \|\theta^{*}\|^{2} + t \right) \frac{1}{\sqrt{2\pi\sigma_{p}^{2}}} d\theta \\ &= \log \sqrt{\frac{\pi 2\sigma_{p}^{2}}{1 - 2\sigma_{p}^{2}t (\sigma_{x}^{2} + \delta^{2}) / \sigma^{2}}} \exp \left( \frac{t^{2}\sigma_{x}^{4} \|\theta^{*}\|^{2} 2\sigma_{p}^{2} / \sigma^{2}}{\sigma^{2}} + \frac{t\sigma_{x}^{2}}{\sigma^{2}} \|\theta^{*}\|^{2} + t \left( 1 - \frac{2\sigma_{p}^{2}t}{\sigma^{2}} (\sigma_{x}^{2} + \delta^{2}) \right) \right) \\ &\leq \frac{d}{2} \log \frac{1}{1 - 2\sigma_{p}^{2}t (\sigma_{x}^{2} + \delta^{2}) / \sigma^{2}}}{1 - 2\sigma_{p}^{2}t (\sigma_{x}^{2} + \delta^{2}) / \sigma^{2}} + \frac{\frac{t\sigma_{x}^{2}}{\sigma^{2}} \|\theta^{*}\|^{2} + t \left( 1 - \frac{2\sigma_{p}^{2}t}{\sigma^{2}} (\sigma_{x}^{2} + \delta^{2}) \right) }{1 - 2\sigma_{p}^{2}t (\sigma_{x}^{2} + \delta^{2}) / \sigma^{2}} \\ &\leq \frac{\sigma_{p}^{2} t d (\sigma_{x}^{2} + \delta^{2}) / \sigma^{2}}{1 - 2\sigma_{p}^{2}t (\sigma_{x}^{2} + \delta^{2}) / \sigma^{2}} + \frac{\frac{t\sigma_{x}^{2}}{\sigma^{2}} \|\theta^{*}\|^{2} + t \left( 1 - \frac{2\sigma_{p}^{2}t}{\sigma^{2}} (\sigma_{x}^{2} + \delta^{2}) \right) }{1 - 2\sigma_{p}^{2}t (\sigma_{x}^{2} + \delta^{2}) / \sigma^{2}} \\ &= \frac{t^{2}s^{2}}{2(1 - tc)} \end{aligned}$$

For adversarial loss,  $s^2 = \frac{2}{t} \left( cd + \frac{\sigma_x^2 \|\theta^*\|^2}{\sigma^2} + (1 - ct) \right), c = \frac{2\sigma_p^2 \left(\sigma_x^2 + \delta^2\right)}{\sigma^2} \text{ and } t \in \left(0, \frac{\sigma^2}{2\sigma_p^2 (\sigma_x^2 + \delta^2)}\right).$ 

# C PROOF OF THEOREMS IN Section 4.3

In this section, we derive Theorems 4.3 to 4.7 using Theorem 4.1. From the PAC-Bayesian bounds for bounded CGF loss theorem, it is clear that we need to bound the expected training risk plus the KL divergence between the posterior and prior. In the bound derivation, we require to bound the negative log normalizing constants of the posteriors which will be presented first in Lemmas C.1 and C.2.

Lemma C.1 (Negative log normalizing constant of the Bayes posterior). The normalizing constant z of Bayes posterior q is

$$\begin{aligned} z &= \int \exp\left(-\mathcal{L}(\theta, \mathcal{D})\right) \pi(\theta) d\theta \\ &= \int \exp\left(-\frac{1}{2\sigma^2} \sum_{i=1}^n (y_i - x_i^\top \theta)^2\right) \pi(\theta) d\theta \\ &= \frac{\exp(-Y^\top Y/2\sigma^2)}{\sqrt{2\pi\sigma_p^2}^d} \int \exp\left(-\theta^\top \left(\frac{1}{2\sigma^2} X^\top X + \frac{1}{2\sigma_p^2} I\right) \theta + \frac{1}{2\sigma^2} 2Y^\top X \theta\right) \right) d\theta \\ &= \frac{\exp\left(-\frac{1}{2\sigma^2} Y^\top Y + \frac{1}{2\sigma^2} Y^\top X \left(\frac{1}{\sigma^2} X^\top X + \frac{1}{\sigma_p^2} I\right)^{-1} X^\top Y \frac{1}{\sigma^2}\right)}{\sqrt{\det\left(\frac{\sigma_p^2}{\sigma^2} X^\top X + I\right)}} \\ \log \frac{1}{z} &= \frac{1}{2} \log \det\left(\frac{\sigma_p^2}{\sigma^2} X^\top X + I\right) + \frac{1}{2\sigma^2} \left(Y^\top Y - Y^\top X \left(\frac{1}{\sigma^2} X^\top X + I\right)^{-1} X^\top Y \frac{1}{\sigma^2}\right) \\ &= \frac{1}{2} \log \det\left(\frac{\sigma_p^2}{\sigma^2} X^\top X + I\right) + \frac{1}{2\sigma^2} Y^\top \left(I + \frac{\sigma_p^2}{\sigma^2} X X^\top\right)^{-1} Y \quad ; Woodbury Matrix Identity \end{aligned}$$

Similar to above Lemma C.1, we derive the negative log normalizing constant of the robust posterior in the following. Lemma C.2 (Negative log normalizing constant of the robust posterior). The normalizing constant  $z_{\delta}$  of robust posterior  $q_{\delta}$  is

$$\begin{split} z_{\delta} &= \int \exp\left(-\mathcal{L}_{\delta}(\theta,\mathcal{D})\right) \pi(\theta) d\theta \\ &\geq \int \exp\left(-\frac{1}{\sigma^{2}} \sum_{i=1}^{n} \left((y_{i} - x_{i}^{\top}\theta)^{2} + \|\theta\|^{2}\delta^{2}\right)\right) \pi(\theta) d\theta \\ &= \frac{1}{\sqrt{2\pi\sigma_{p}^{2}}} \int \exp\left(-\left(\frac{n\delta^{2}}{\sigma^{2}}\|\theta\|^{2} + \frac{1}{2\sigma_{p}^{2}}\|\theta\|^{2} + \frac{1}{\sigma^{2}}\|Y - X\theta\|^{2}\right)\right) d\theta \quad ; \pi(\theta) \sim \mathcal{N}(0, \sigma_{p}^{2}I) \\ &= \frac{1}{\sqrt{2\pi\sigma_{p}^{2}}} \int \exp\left(-\left(\frac{n\delta^{2}}{\sigma^{2}}\theta^{\top}\theta + \frac{1}{2\sigma_{p}^{2}}\theta^{\top}\theta + \frac{1}{\sigma^{2}}\left(\|Y\|^{2} - 2Y^{\top}X\theta + \theta^{\top}X^{\top}X\theta\right)\right)\right) d\theta \\ &= \frac{1}{\sqrt{2\pi\sigma_{p}^{2}}} \exp\left(-\frac{1}{\sigma^{2}}\|Y\|^{2}\right) \int \exp\left(-\theta^{\top}\left((\frac{n\delta^{2}}{\sigma^{2}} + \frac{1}{2\sigma_{p}^{2}})I + \frac{X^{\top}X}{\sigma^{2}}\right) \theta + \frac{2Y^{\top}X\theta}{\sigma^{2}}\right) d\theta \quad ; Lemma B.2 \\ &= \sqrt{\frac{1}{\sqrt{2\pi\sigma_{p}^{2}}}} \exp\left(-\frac{1}{\sigma^{2}}(1-\frac{\sigma^{2}}{\sigma^{2}})I + \frac{1}{\sigma^{2}}(1-\frac{\sigma^{2}}{\sigma^{2}})I + \frac{X^{\top}Y}{\sigma^{2}}\right) d\theta \quad ; Lemma B.2 \\ &= \sqrt{\frac{1}{\sqrt{2\pi\sigma_{p}^{2}}}} \exp\left(-\frac{1}{\sigma^{2}}(1-\frac{\sigma^{2}}{\sigma^{2}})I + \frac{1}{\sigma^{2}}I + \frac{1}{\sigma^{$$

With the above bounds derived, we are now ready to derive the generalization certificate using Theorem 4.1. Following holds for all the cases:

(*i*) The CGF bound of respective loss-NLL in case of standard generalization and adversarial NLL in case of adversarial generalization)-appears in the theorem. This directly implies that the constants c and s should follow Lemma 4.2 according to the considered setting.

(*ii*) The free parameter in the CGF bound, t is choosen as t = 1, which is also done in the prior work [Germain et al., 2016]. This means  $1 \in (0, 1/c)$ , that is, c < 1.

Consequently, the effective Theorem 4.1 is stated below for clarity.

**Theorem C.3** (Theorem 4.1). Consider data  $\mathcal{D}$  and any loss  $\ell(\theta, (x, y))$  with its corresponding expected and empirical generalization errors  $R(\theta) = \underset{(x,y)\sim\mathcal{P}}{\mathbb{E}} \left[\ell(\theta, (x, y))\right]$  and  $r(\theta) = \frac{1}{n}\mathcal{L}(\theta, \mathcal{D})$ , respectively. Let the CGF of the loss  $\psi(t) \leq \frac{1}{n}\mathcal{L}(\theta, \mathcal{D})$ .

 $\frac{t^2s^2}{2(1-tc)}$  be bounded, where for some constant  $c \in (0, .1)$  and  $t \in (0, 1/c)$ . Then, we have, with probability at least  $1 - \beta$ , for all densities  $\rho(\theta)$ ,

$$\mathbb{E}_{\theta \sim \rho}[R(\theta)] \le \mathbb{E}_{\theta \sim \rho}[r(\theta)] + \frac{1}{n} \mathrm{KL}(\rho \| \pi) + \frac{1}{n} \log \frac{1}{\beta} + \frac{s^2}{2(1-c)}$$

Therefore, we only need to bound the following for  $\rho$  being standard Bayes q and robust posterior  $q_{\delta}$ , and  $\ell$  being NLL and adversarial NLL.

$$\mathop{\mathbb{E}}_{\theta \sim \rho} \left[ r(\theta) \right] + \frac{1}{n} \mathrm{KL}(\rho \| \pi). \tag{17}$$

#### C.1 STANDARD GENERALIZATION OF BAYES POSTERIOR (Theorem 4.3)

In this case,  $\rho = q$  and  $\ell$  is NLL in (17) which reduces the expression to the following.

$$\mathbb{E}_{\theta \sim q} [r(\theta)] + \frac{1}{n} \mathrm{KL}(q \| \pi) = \frac{1}{n} \int \frac{1}{z} \ell(\theta, \mathcal{D}) \exp(-\ell(\theta, \mathcal{D})) \pi(\theta) d\theta + \frac{1}{n} \int \frac{1}{z} \exp(-\ell(\theta, \mathcal{D})) \pi(\theta) \log\left(\frac{\exp(-\ell(\theta, \mathcal{D}))\pi(\theta)}{z\pi(\theta)}\right) d\theta$$

$$= \frac{1}{n} \int \frac{1}{z} \exp(-\ell(\theta, \mathcal{D})) \pi(\theta) \log\left(\frac{1}{z}\right) d\theta$$

$$= \frac{1}{n} \log \frac{1}{z}$$
(18)

Therefore, we obtain Theorem 4.3 by substituting Lemma C.1 in (18) and combining it with the Theorem C.3.  $\Box$ 

## C.2 ADVERSARIAL GENERALIZATION OF ROBUST POSTERIOR $\delta = \hat{\delta}$ (Theorem 4.6)

Notice that the above derivation in Appendix C.1 holds for  $\rho = q_{\delta}$  and  $\ell$  is adversarial loss with  $\delta$  perturbation with  $z_{\delta}$  as the normalization constant. That is, the perturbation at train and test-time are the same  $\delta = \hat{\delta}$ .

$$\mathop{\mathbb{E}}_{\theta \sim q_{\delta}}\left[r(\theta)\right] + \frac{1}{n}\mathrm{KL}(q_{\delta}\|\pi) = \frac{1}{n}\log\frac{1}{z_{\delta}}$$
(19)

Therefore, we obtain Theorem 4.6 by substituting Lemma C.2 in (19) and combining it with the Theorem C.3.  $\Box$ 

#### C.3 ADVERSARIAL GENERALIZATION OF BAYES POSTERIOR (Theorem 4.4)

In this case,  $\rho = q$  and  $\ell$  is adversarial NLL with  $\hat{\delta}$  perturbation in (17). We upper bound it as follows.

For ease of notation, only in the following, we use short hand notations for  $\ell$  to denote  $\ell(\theta, D)$  and  $\ell_{\hat{\delta}}$  to denote  $\ell_{\hat{\delta}}(\theta, D)$ .

$$\begin{split} \mathbb{E}_{\theta \sim q} \left[ \frac{1}{n} \ell_{\delta}(\theta, \mathcal{D}) \right] + \frac{1}{n} \mathrm{KL}(q \| \pi) &= \frac{1}{n} \int \frac{1}{z} \ell_{\delta} \exp(-\ell) \pi(\theta) d\theta + \frac{1}{n} \int \frac{1}{z} \exp(-\ell) \pi(\theta) \log\left(\frac{\exp(-\ell)\pi(\theta)}{z\pi(\theta)}\right) d\theta \\ &= \frac{1}{n} \int \frac{1}{z} \left( \ell_{\delta} - \ell \right) \exp(-\ell) \pi(\theta) d\theta + \frac{1}{n} \int \frac{1}{z} \exp(-\ell) \pi(\theta) \log\left(\frac{1}{z}\right) d\theta \\ &\leq \frac{1}{n} \int \frac{1}{z} \left( 2\ell + \frac{1}{2\sigma^2} 2n \| \theta \|^2 \delta^2 - \ell \right) \exp(-\ell) \pi(\theta) d\theta + \frac{1}{n} \log \frac{1}{z} \quad ; \text{Lemma B.1} \\ &\leq \frac{1}{n} \log \int \frac{1}{z} \exp\left(\ell + \frac{1}{2\sigma^2} 2n \| \theta \|^2 \delta^2\right) \exp(-\ell) \pi(\theta) d\theta + \frac{1}{n} \log \frac{1}{z} \quad ; \text{Jensen's ineq.} \\ &= \frac{1}{n} \log \frac{1}{z} \int \exp\left(\left(\frac{2n\delta^2}{2\sigma^2} - \frac{1}{2\sigma_p^2}\right) \| \theta \|^2\right) \frac{1}{\sqrt{2\pi\sigma_p^2}} d\theta + \frac{1}{n} \log \frac{1}{z} \\ &= \frac{1}{n} \log \frac{1}{z} \sqrt{\frac{2\pi\sigma_p^2\sigma^2}{\sigma^2 - 2n\delta^2\sigma_p^2}} \frac{1}{\sqrt{2\pi\sigma_p^2}} \frac{1}{\pi} \log \frac{1}{z} \\ &= \frac{2}{n} \log \frac{1}{z} + \frac{d}{2n} \log \frac{1}{1 - \frac{2}{\sigma^2} n\delta^2\sigma_p^2} \quad ; -\log(1 - x) \leq \frac{x}{1 - x} \\ &= \frac{2}{n} \log \frac{1}{z} + \frac{d\delta^2\sigma_p^2/\sigma^2}{1 - 2n\delta^2\sigma_p^2/\sigma^2} \end{aligned}$$

We obtain Theorem 4.4 by substituting Lemma C.1 in (20) and combining it with the Theorem C.3.  $\Box$ 

# C.4 STANDARD GENERALIZATION OF ROBUST POSTERIOR (Theorem 4.5)

In this case,  $\rho = q_{\delta}$  and  $\ell$  is NLL in (17). We upper bound it as follows.

For ease of notation, only in the following, we use short hand notations for  $\ell$  to denote  $\ell(\theta, D)$  and  $\ell_{\delta}$  to denote  $\ell_{\delta}(\theta, D)$ .

$$\begin{aligned}
& \left[\frac{1}{n}\ell(\theta,\mathcal{D})\right] + \frac{1}{n}\mathrm{KL}(q_{\delta}\|\pi) = \frac{1}{n}\int\frac{1}{z_{\delta}}\ell\exp(-\ell_{\delta})\pi(\theta)d\theta + \frac{1}{n}\int\frac{1}{z_{\delta}}\exp(-\ell_{\delta})\pi(\theta)\log\left(\frac{\exp(-\ell_{\delta})\pi(\theta)}{z_{\delta}\pi(\theta)}\right)d\theta \\
& = \frac{1}{n}\int\frac{1}{z_{\delta}}\left(-\ell_{\delta}+\ell\right)\exp(-\ell_{\delta})\pi(\theta)d\theta + \frac{1}{n}\int\frac{1}{z_{\delta}}\exp(-\ell_{\delta})\pi(\theta)\log\left(\frac{1}{z_{\delta}}\right)d\theta \\
& \leq \frac{1}{n}\int\frac{1}{z_{\delta}}\left(-\ell-\frac{1}{2\sigma^{2}}n\|\theta\|^{2}\delta^{2}+\ell\right)\exp(-\ell_{\delta})\pi(\theta)d\theta + \frac{1}{n}\log\frac{1}{z_{\delta}} \quad ;\text{Lemma B.1} \\
& \leq \frac{1}{n}\log\int\frac{1}{z_{\delta}}\exp\left(-\frac{1}{2\sigma^{2}}n\|\theta\|^{2}\delta^{2}\right)\exp(-\ell_{\delta})\pi(\theta)d\theta + \frac{1}{n}\log\frac{1}{z_{\delta}} \quad ;\text{Jensen's ineq.} \\
& = \frac{1}{n}\log\int\exp\left(\left(-\frac{1}{2\sigma^{2}}2n\delta^{2}-\frac{1}{2\sigma_{p}^{2}}\right)\|\theta\|^{2}-\ell\right)\frac{1}{\sqrt{2\pi\sigma_{p}^{2}}}d\theta + \frac{2}{n}\log\frac{1}{z_{\delta}} \quad (21)
\end{aligned}$$

The evaluation of the above integral  $\int \exp\left(\left(-\frac{1}{2\sigma^2}2n\delta^2 - \frac{1}{2\sigma_p^2}\right)\|\theta\|^2 - \ell\right) \frac{1}{\sqrt{2\pi\sigma_p^2}^d}d\theta + \frac{2}{n}\log\frac{1}{z_\delta}$  is as follows:

$$\begin{split} &= \frac{1}{\sqrt{2\pi\sigma_{p}^{2}}} \int \exp\left(-\frac{1}{2} \left(\frac{1}{2\sigma^{2}} 4n \|\theta\|^{2} \delta^{2} + \frac{1}{\sigma_{p}^{2}} \|\theta\|^{2} + \frac{2}{2\sigma^{2}} \|Y - X\theta\|^{2}\right)\right) d\theta \quad ;\pi(\theta) \sim \mathcal{N}(0, \sigma_{p}^{2}I) \\ &= \frac{1}{\sqrt{2\pi\sigma_{p}^{2}}} \int \exp\left(-\frac{1}{2} \left(\frac{1}{2\sigma^{2}} 4n \delta^{2} \theta^{\top} \theta + \frac{1}{\sigma_{p}^{2}} \theta^{\top} \theta + \frac{1}{2\sigma^{2}} \left(2\|Y\|^{2} - 4Y^{\top} X\theta + 2\theta^{\top} X^{\top} X\theta\right)\right) \right) d\theta \\ &= \frac{1}{\sqrt{2\pi\sigma_{p}^{2}}} \exp\left(-\frac{1}{2\sigma^{2}} \|Y\|^{2}\right) \int \exp\left(-\theta^{\top} \left((\frac{1}{2\sigma^{2}} 2n \delta^{2} + \frac{1}{2\sigma_{p}^{2}}\right)I + \frac{1}{2\sigma^{2}} X^{\top} X\right) \theta + \frac{2}{2\sigma^{2}} Y^{\top} X\theta\right) d\theta \quad ;\text{Lemma B.2} \\ &= \sqrt{\frac{1}{\det\left(\frac{\sigma_{p}^{2}}{\sigma^{2}} X^{\top} X + \left(\frac{\sigma_{p}^{2}}{\sigma^{2}} 2n \delta^{2} + 1\right)I\right)}} \exp\left(-\frac{1}{2\sigma^{2}} \|Y\|^{2}\right) \exp\left(\frac{1}{2\sigma^{4}} Y^{T} X\left(\frac{1}{\sigma^{2}} X^{\top} X + \left(\frac{1}{\sigma^{2}} 2n \delta^{2} + \frac{1}{\sigma^{p}^{2}}\right)I\right)^{-1} X^{T} Y\right) \\ &= \sqrt{\frac{1}{\det\left(\frac{\sigma_{p}^{2}}{\sigma^{2}} X^{\top} X + \left(\frac{\sigma_{p}^{2}}{\sigma^{2}} 2n \delta^{2} + 1\right)I\right)}} \exp\left(-\frac{1}{2\sigma^{2}} Y^{T} \left(I - \frac{1}{\sigma^{2}} X\left(\frac{1}{\sigma^{2}} X^{\top} X + \left(\frac{1}{\sigma^{2}} 2n \delta^{2} + \frac{1}{\sigma^{p}^{2}}\right)I\right)^{-1} X^{T}\right) Y\right) \\ &= \sqrt{\frac{1}{\det\left(\frac{\sigma_{p}^{2}}{\sigma^{2}} X^{\top} X + \left(\frac{\sigma_{p}^{2}}{\sigma^{2}} 2n \delta^{2} + 1\right)I\right)}}} \exp\left(-\frac{1}{2\sigma^{2}} Y^{T} \left(I + \frac{1}{\sigma^{2}} X X^{T} \left(\frac{\sigma^{p}^{2} \sigma^{2}}{\sigma^{2} + 2n \delta^{2} \sigma^{p}^{2}}\right)\right)^{-1} Y\right)$$
(22)

Therefore, we obtain Theorem 4.5 by substituting (22) in (21) and combining it with the Theorem C.3.

# C.5 ADVERSARIAL GENERALIZATION OF ROBUST POSTERIOR WHEN $\delta \neq \hat{\delta}$ (Theorem 4.7)

In this case,  $\rho = q_{\delta}$  and  $\ell$  is adversarial NLL with  $\hat{\delta}$  perturbation in (17). We upper bound it as follows. For ease of notation, only in the following, we use short hand notations for  $\ell_{\delta}$  to denote  $\ell_{\delta}(\theta, D)$  and  $\ell_{\hat{\delta}}$  to denote  $\ell_{\hat{\delta}}(\theta, D)$ .

$$\begin{split} \mathbb{E}_{\sigma \sim q_{\delta}} \left[ \frac{1}{n} \ell_{\delta}(\theta, \mathcal{D}) \right] + \frac{1}{n} \mathrm{KL}(q_{\delta} \| \pi) &= \frac{1}{n} \int \frac{1}{z_{\delta}} \ell_{\delta} \exp(-\ell_{\delta}) \pi(\theta) d\theta + \frac{1}{n} \int \frac{1}{z_{\delta}} \exp(-\ell_{\delta}) \pi(\theta) \log\left(\frac{\exp(-\ell_{\delta})\pi(\theta)}{z_{\delta}\pi(\theta)}\right) d\theta \\ &= \frac{1}{n} \int \frac{1}{z_{\delta}} \left( \ell_{\delta} - \ell_{\delta} \right) \exp(-\ell_{\delta})\pi(\theta) d\theta + \frac{1}{n} \int \frac{1}{z_{\delta}} \exp(-\ell_{\delta})\pi(\theta) \log\left(\frac{1}{z_{\delta}}\right) d\theta \\ &\leq \frac{1}{n} \int \frac{1}{z_{\delta}} \left( 2\ell + \frac{2n \|\theta\|^{2} \delta^{2}}{2\sigma^{2}} - \ell - \frac{n \|\theta\|^{2} \delta^{2}}{2\sigma^{2}} \right) \exp(-\ell_{\delta})\pi(\theta) d\theta + \frac{1}{n} \log \frac{1}{z_{\delta}} \quad ; \text{Lemma B.1} \\ &\leq \frac{1}{n} \log \int \frac{1}{z_{\delta}} \exp\left(\ell + \frac{n \|\theta\|^{2} \left(2\delta^{2} - \delta^{2}\right)}{2\sigma^{2}}\right) \exp(-\ell_{\delta})\pi(\theta) d\theta + \frac{1}{n} \log \frac{1}{z_{\delta}} \quad ; \text{Jensen's ineq} \\ &= \frac{1}{n} \log \frac{1}{z_{\delta}} \int \exp\left(\left(\frac{n \left(2\delta^{2} - 2\delta^{2}\right)}{2\sigma^{2}} - \frac{1}{2\sigma_{p}^{2}}\right) \|\theta\|^{2}\right) \frac{1}{\sqrt{2\pi\sigma_{p}^{2}}} d\theta + \frac{1}{n} \log \frac{1}{z_{\delta}} \\ &= \frac{1}{n} \log \frac{1}{z_{\delta}} \int \frac{2\pi\sigma_{p}^{2}}{1 - 2n \left(\delta^{2} - \delta^{2}\right) \sigma_{p}^{2} / \sigma^{2}} \frac{1}{\sqrt{2\pi\sigma_{p}^{2}}} + \frac{1}{n} \log \frac{1}{z_{\delta}} \\ &= \frac{2}{n} \log \frac{1}{z_{\delta}} + \frac{d}{2n} \log \frac{1}{1 - 2n \left(\delta^{2} - \delta^{2}\right) \sigma_{p}^{2} / \sigma^{2}} \\ &\leq \frac{2}{n} \log \frac{1}{z_{\delta}} + \frac{d}{2n} \frac{2n \left(\delta^{2} - \delta^{2}\right) \sigma_{p}^{2} / \sigma^{2}}{1 - 2n \left(\delta^{2} - \delta^{2}\right) \sigma_{p}^{2} / \sigma^{2}}} \quad ; -\log(1 - x) \leq \frac{x}{1 - x} \\ &= \frac{2}{n} \log \frac{1}{z_{\delta}} + \frac{\left(\delta^{2} - \delta^{2}\right) \sigma_{p}^{2} / \sigma^{2}}{1 - 2n \left(\delta^{2} - \delta^{2}\right) \sigma_{p}^{2} / \sigma^{2}}} \tag{23}$$

Substituting Lemma C.2 in (23) and combining it with the Theorem C.3 proves Theorem 4.7.

# D DATASETS AND IMPLEMENTATION DETAILS

The real datasets used in the experiments are given in Table 3, and they are available in UCI repository [Markelle et al.] or OpenML [Vanschoren et al., 2013]. We use it 70-30 train-test split to learn and test the posteriors. The code is developed in Pytorch [Paszke et al., 2017] and use Pyro package [Bingham et al., 2019] for NUTS distribution sampler. All the experiments are run on CPU of Apple M1 chip with 16GB memory. The run time is between seconds upto a few minutes.

Dataset	Number of samples	Data dimension
Abalone	4177	10
Air Foil	1503	5
Air Quality	7355	11
Auto MPG	393	9
California Housing	20640	8
Energy Efficiency	768	8
Wine Quality	1599	11

Table 3: Real datasets

# **E** ADDITIONAL EXPERIMENTAL RESULTS ON REAL DATA

We provide the additional results on the real datasets for prior variance  $\sigma_p^2 = \frac{1}{100}$  in Table 4 and  $\frac{1}{9}$  in Table 5. The experimental results are consistent with the results in Table 2. We observe that informed choice of prior favors robust posterior in terms of adversarial generalization.

Dataset	Standard generalization (NLL) $\ell$		Adversarial generalization (adv-NLL) $\ell_{\hat{\delta}}$	
	Bayes posterior $q$	Robust posterior $q_{\delta}$	Bayes posterior $q$	Robust posterior $q_{\delta}$
Abalone	$1.1664 \pm 0.014$	$1.1797  \pm  0.014$	$1.2221\ \pm 0.014$	$1.2172 \pm 0.015$
Air Foil	$1.1714 \pm 0.008$	$1.1788 \pm 0.009$	$\boldsymbol{1.2183} \pm 0.009$	$1.2219 \pm 0.009$
Air Quality	$\textbf{0.9668} \pm 0.002$	$0.9674  \pm  0.002$	$0.9800 \pm 0.002$	$\textbf{0.9791} \pm 0.002$
Auto MPG	$1.0295 \pm 0.010$	$1.0305  \pm  0.010$	$1.0471 \pm 0.011$	$1.0474 \pm 0.011$
California Housing	$1.1195 \pm 0.003$	$1.1280 \pm 0.005$	$1.1862 \pm 0.003$	$\boldsymbol{1.1768} \pm 0.005$
Energy Efficiency	$0.9834 \pm 0.009$	$0.9838 \pm 0.009$	$1.0007  \pm  0.010$	$1.0006 \pm 0.010$
Wine Quality	$1.2323 \pm 0.006$	$1.2329_{0.006}$	$1.2642 \pm 0.006$	$1.2614 \pm 0.006$

Table 4: Test NLL and adversarial NLL of Bayes and robust posteriors on real datasets. The prior variance is set to  $\sigma_p^2 = \frac{1}{100}$ . The robust posterior is trained with  $\delta = 0.1$  in the adversarial NLL loss, and adversarial generalization is evaluated using the same training-time perturbation ( $\hat{\delta} = 0.1$ ). The adversarial generalization results demonstrate that the robust posterior  $q_{\delta}$  is consistently more robust than the Bayes posterior q. For both standard and adversarial generalization, the best-performing model for each dataset is highlighted in bold.

Dataset	Standard generalization (NLL) $\ell$		Adversarial generalization (adv-NLL) $\ell_{\hat{\delta}}$	
	Bayes posterior $q$	Robust posterior $q_{\delta}$	Bayes posterior $q$	Robust posterior $q_{\delta}$
Abalone	$1.1585 \pm 0.013$	$1.1726 \pm \textbf{0.014}$	$1.2554 \pm 0.011$	$1.2176 \pm 0.015$
Air Foil	$1.1657 \pm 0.009$	$1.1694{\scriptstyle~\pm 0.009}$	$1.2191 \pm 0.009$	$1.2176 \pm 0.009$
Air Quality	$0.9665 \pm 0.002$	$0.9670{\scriptstyle~\pm 0.002}$	$0.9826 \pm 0.003$	$0.9791 \pm 0.002$
Auto MPG	$1.0231  \pm 0.006$	$\boldsymbol{1.0228} \pm 0.007$	$1.0552 \pm 0.006$	$1.0469 \pm 0.008$
California Housing	$1.1193 \pm 0.003$	$1.1267{\scriptstyle~\pm 0.005}$	$1.1909  \pm  0.003$	$\boldsymbol{1.1768} \pm 0.005$
Energy Efficiency	$0.9712 \pm 0.006$	$0.9733{\scriptstyle~\pm 0.007}$	$0.9990 \pm 0.007$	$\textbf{0.9947} \pm 0.008$
Wine Quality	$1.2338  \pm  0.006$	$1.2321 \pm 0.006$	$1.2697 \pm 0.008$	$1.2625 \pm 0.006$

Table 5: Test NLL and adversarial NLL of Bayes and robust posteriors on real datasets. The prior variance is set to  $\sigma_p^2 = \frac{1}{9}$ . The robust posterior is trained with  $\delta = 0.1$  in the adversarial NLL loss, and adversarial generalization is evaluated using the same training-time perturbation ( $\hat{\delta} = 0.1$ ). The adversarial generalization results demonstrate that the robust posterior  $q_{\delta}$  is consistently more robust than the Bayes posterior q. For both standard and adversarial generalization, the best-performing model for each dataset is highlighted in bold.