

# Analysis of $q_{\text{rec}}^2$ -distribution for $B \rightarrow KM_X$ and $B \rightarrow K^*M_X$ decays in a scalar-mediator dark-matter scenario

Alexander Berezhnoy<sup>a</sup>, Wolfgang Lucha<sup>b</sup> and Dmitri Melikhov<sup>a,c,d</sup>

<sup>a</sup>*D. V. Skobeltsyn Institute of Nuclear Physics, M. V. Lomonosov Moscow State University, 119991, Moscow, Russia*

<sup>b</sup>*Institute of High Energy Physics, Austrian Academy of Sciences, Nikolsdorfergasse 18, A-1050 Vienna, Austria*

<sup>c</sup>*Joint Institute for Nuclear Research, 141980 Dubna, Russia*

<sup>d</sup>*Faculty of Physics, University of Vienna, Boltzmanngasse 5, A-1090 Vienna, Austria*

(Dated: February 21, 2025)

We demonstrate that the scalar-mediator dark-matter scenario is consistent with the experimental data on the decay  $B \rightarrow KM_X$  and provides a good description of the shape of the observed excess. Within this scenario, the interaction with dark-matter particles leads to approximately the same excess in  $\Gamma(B \rightarrow K^*M_X)$  and  $\Gamma(B \rightarrow KM_X)$  compared to the Standard Model; also the differential distributions of the excess events are similar in shape in the variable  $q_{\text{rec}}^2$  measured by experiment.

## I. INTRODUCTION

A recent Belle II observation [1] of  $B \rightarrow KM_X$  — for which also the notation  $B \rightarrow K\cancel{E}$ , with  $\cancel{E}$  the missing energy, is used — at a level much exceeding the Standard-Model (SM) prediction for  $B \rightarrow K\nu\bar{\nu}$  [2],

$$\mathcal{B}(B^+ \rightarrow K^+M_X) = (2.3 \pm 0.7) \times 10^{-5} \simeq (5.4 \pm 1.5) \mathcal{B}(B^+ \rightarrow K^+\nu\bar{\nu})_{\text{SM}}, \quad (1.1)$$

opened the window for immediate discussions of possible new-physics effects capable to explain this result (see, e.g., the recent publications [2–28]). One of the most popular discussed options is the decay into Dark-Matter (DM) particles [6] with multiple scenarios for the content of these particles and the possible mediators.

In our previous paper [28], we noted that combining the current Belle II result for the decay  $B \rightarrow KM_X$  with the hypothesis of a DM origin of the enhancement of the decay width  $\Gamma(B \rightarrow KM_X)$ , where the DM particles couple to SM particles via a scalar-mediator field, leads to rigorous constraints on  $\Gamma(B \rightarrow K^*M_X)$  that are independent of further details of the DM model.

Extending the results of [28], this analysis provides a study of the differential distributions in the decays  $B \rightarrow KM_X$  and  $B \rightarrow K^*M_X$  in the experimental variable  $q_{\text{rec}}^2$  [1]. By confronting the calculated distributions with the data, we (i) obtain constraints on the numerical parameters of the DM model — such as the mass and width of the scalar mediator, the mass of the light DM fermions, as well as the corresponding couplings — and (ii) report that within the scalar mediator scenario, the  $q_{\text{rec}}^2$  differential distributions in  $B \rightarrow KM_X$  and  $B \rightarrow K^*M_X$  are equal to each other within a few-percent accuracy, independently of the numerical parameters of the DM model.

## II. $B \rightarrow (K, K^*)\bar{\chi}\chi$ DECAYS VIA A SCALAR MEDIATOR $\phi$

Following [28], we focus on a rather simple representative of the really vast class of DM models. To this end, we consider a rather popular model involving an interaction of DM fermions  $\chi$  with the top quark  $t$  by exchange of a scalar-mediator field  $\phi$ , governed by the interaction Lagrangian [29, 30]

$$\mathcal{L}_{\text{int}} = -\frac{ym_t}{v}\phi\bar{t}t - \kappa\phi\bar{\chi}\chi. \quad (2.1)$$

The emerging effective Lagrangian encoding the flavor-changing neutral-current (FCNC) vertex  $b \rightarrow s\phi$  then reads [29, 30]

$$\mathcal{L}_{b \rightarrow s\phi} = g_{b \rightarrow s\phi}\phi\bar{s}_L b_R + \text{h.c.}, \quad g_{b \rightarrow s\phi} = \frac{ym_b}{v} \frac{3\sqrt{2}G_F m_t^2 V_{ts}^* V_{tb}}{16\pi^2}. \quad (2.2)$$

This Lagrangian enters in the amplitudes controlling the  $\phi$ -mediated decay  $B \rightarrow (K, K^*)\bar{\chi}\chi$ :

$$\begin{aligned} A(B(p) \rightarrow K^{(*)}(p-q)\bar{\chi}(k)\chi(q-k)) &= -i\langle K^{(*)}(p-q)\bar{\chi}(k)\chi(q-k) | \mathcal{L}_{b \rightarrow s\phi} | B(p) \rangle \\ &= \langle \bar{\chi}\chi | \bar{\chi}\chi | 0 \rangle \kappa \frac{1}{M_\phi^2 - q^2 + iM_\phi\Gamma_\phi(q^2)} g_{b \rightarrow s\phi} \langle K^{(*)}(p-q) | \bar{s}_L b_R | B(p) \rangle, \end{aligned} \quad (2.3)$$

with the abbreviation

$$\langle \bar{\chi}\chi|\bar{\chi}\chi|0\rangle \equiv \langle \bar{\chi}(k)\chi(q-k)|\bar{\chi}(0)\chi(0)|0\rangle. \quad (2.4)$$

For the present analysis, we take the liberty to assume  $M_\phi > 2m_\chi$  and, as a consequence, to introduce a  $q^2$ -dependent width of the mediator  $\phi$  according to

$$\Gamma_\phi(q^2) = \left( \frac{q^2 - 4m_\chi^2}{M_\phi^2 - 4m_\chi^2} \right)^{\frac{3}{2}} \Theta(q^2 - 4m_\chi^2) \Gamma_\phi^0. \quad (2.5)$$

Here,  $q$  is the momentum of the outgoing  $\chi\bar{\chi}$  pair of unobserved DM particles and  $M_\chi^2 \equiv q^2$  denotes the missing mass squared. Equation (2.5) provides a simplified parametrization of the  $q^2$ -dependent width that reproduces the correct threshold behavior of the decay rate of a scalar particle  $\phi$  into two spin-1/2 fermions  $\chi\bar{\chi}$ . The on-shell width  $\Gamma_\phi^0$  is proportional to  $\kappa^2$ , see Eq. (5.1) below.

Using the QCD equations of motion, the required amplitudes can be straightforwardly calculated, yielding

$$\begin{aligned} \langle K|\bar{s}_L b_R|B\rangle &= \frac{1}{2}\langle K|\bar{s}(1-\gamma_5)b|B\rangle = \frac{1}{2}\langle K|\bar{s}b|B\rangle = \frac{1}{2}\frac{M_B^2 - M_K^2}{m_b - m_s} f_0^{B\rightarrow K}(q^2), \\ \langle K^*|\bar{s}_L b_R|B\rangle &= \frac{1}{2}\langle K^*|\bar{s}(1-\gamma_5)b|B\rangle = -\frac{1}{2}\langle K^*|\bar{s}\gamma_5 b|B\rangle = -i(\epsilon q)\frac{M_{K^*}}{m_b + m_s} A_0^{B\rightarrow K^*}(q^2), \end{aligned} \quad (2.6)$$

with well-known dimensionless form factors  $f_0$  and  $A_0$  parametrizing the amplitudes  $\langle K|\bar{s}\gamma_\mu b|B\rangle$  and  $\langle K^*|\bar{s}\gamma_\mu\gamma_5 b|B\rangle$  [31].

Using these amplitudes and the recursive formula for the phase space

$$d\Phi_3(M_B^2, p_K, p_\chi, p_{\bar{\chi}}) = \Phi_2(M_B^2, p_K, p_\chi + p_{\bar{\chi}}) \frac{dq^2}{2\pi} d\Phi_2(q^2, p_\chi, p_{\bar{\chi}}) = \frac{\lambda^{1/2}(M_B^2, M_K^2, q^2)}{8\pi M_B^2} \frac{dq^2}{16\pi^2} \sqrt{1 - \frac{4m_\chi^2}{q^2}}, \quad (2.7)$$

where  $\lambda$  is defined by  $\lambda(a, b, c) \equiv (a - b - c)^2 - 4bc$ , we can factorize the process into the decay of the  $B$  meson to the  $K$  meson and a virtual  $\phi^*$ , followed by the decay of  $\phi^*$  to a  $\chi\bar{\chi}$  pair. The  $d\Gamma(B \rightarrow (K, K^*)\bar{\chi}\chi)/dq^2$  distributions can then be expressed as

$$\begin{aligned} \frac{d\Gamma}{dq^2}(B \rightarrow K\bar{\chi}\chi) &= \frac{\sum_{\chi \text{ polar}} |A|^2}{2M_B} \frac{\lambda^{1/2}(M_B^2, M_K^2, q^2)}{8\pi M_B^2} \frac{1}{16\pi^2} \sqrt{1 - \frac{4m_\chi^2}{q^2}} \\ &= \frac{\lambda^{1/2}(M_B^2, M_K^2, q^2)}{16\pi M_B^3} |\langle K|\bar{s}_L b_R|B\rangle|^2 \frac{g_{b\rightarrow s\phi}^2 \kappa^2}{(M_\phi^2 - q^2)^2 + M_\phi^2 \Gamma_\phi^2(q^2)} \frac{1}{16\pi^2} \sqrt{1 - \frac{4m_\chi^2}{q^2}} \\ &\times \sum_{\chi \text{ polar}} |\langle \bar{\chi}(k)\chi(q-k)|\bar{\chi}(0)\chi(0)|0\rangle|^2. \end{aligned} \quad (2.8)$$

The expression for  $B \rightarrow K^*\bar{\chi}\chi$  is obtained by the obvious replacements  $K \rightarrow K^*$  and  $M_K \rightarrow M_{K^*}$  in Eq. (2.8).

Taking into account that

$$\sum_{\chi \text{ polar}} |\langle \bar{\chi}(k)\chi(q-k)|\bar{\chi}(0)\chi(0)|0\rangle|^2 = q^2 - 4m_\chi^2 \quad (2.9)$$

and

$$\sum_{K^* \text{ polar}} |\langle K^*|\bar{s}_L b_R|B\rangle|^2 = \frac{\lambda(M_B^2, M_{K^*}^2, q^2)}{4(m_b + m_s)^2} |A_0^{B\rightarrow K^*}(q^2)|^2, \quad (2.10)$$

we obtain the following expressions for  $d\Gamma(B \rightarrow K\bar{\chi}\chi)/dq^2$  and  $d\Gamma(B \rightarrow K^*\bar{\chi}\chi)/dq^2$ :

$$\begin{aligned} \frac{d\Gamma}{dq^2}(B \rightarrow K\bar{\chi}\chi) &= \frac{\lambda^{1/2}(M_B^2, M_K^2, q^2)}{16\pi M_B^3} \frac{(M_B^2 - M_K^2)^2 |f_0^{B\rightarrow K}(q^2)|^2}{4(m_b - m_s)^2} \frac{g_{b\rightarrow s\phi}^2 \kappa^2}{(M_\phi^2 - q^2)^2 + M_\phi^2 \Gamma_\phi^2(q^2)} \frac{q^2}{16\pi^2} \left(1 - \frac{4m_\chi^2}{q^2}\right)^{3/2}, \\ \frac{d\Gamma}{dq^2}(B \rightarrow K^*\bar{\chi}\chi) &= \frac{\lambda^{3/2}(M_B^2, M_{K^*}^2, q^2)}{16\pi M_B^3} \frac{|A_0^{B\rightarrow K^*}(q^2)|^2}{4(m_b + m_s)^2} \frac{g_{b\rightarrow s\phi}^2 \kappa^2}{(M_\phi^2 - q^2)^2 + M_\phi^2 \Gamma_\phi^2(q^2)} \frac{q^2}{16\pi^2} \left(1 - \frac{4m_\chi^2}{q^2}\right)^{3/2}. \end{aligned} \quad (2.11)$$

A powerful probe of the DM scenario considered is provided by the ratio [28]

$$R_{K^*/K}^{(\phi)}(q^2) = \frac{d\Gamma(B \rightarrow K^* \chi \bar{\chi})/dq^2}{d\Gamma(B \rightarrow K \chi \bar{\chi})/dq^2} = \frac{\lambda^{3/2}(M_B^2, M_{K^*}^2, q^2) |A_0^{B \rightarrow K^*}(q^2)|^2}{\lambda^{1/2}(M_B^2, M_K^2, q^2) |f_0^{B \rightarrow K}(q^2)|^2} \frac{(m_b - m_s)^2}{(M_B^2 - M_K^2)^2 (m_b + m_s)^2}. \quad (2.12)$$

Clearly, the dependence on the specific parameters of the DM model cancels out in this ratio. We shall see that this ratio remains largely unaffected by the averaging procedure adopted in the experiment.

The form factors  $f_0$  and  $A_0$  required by Eq. (2.6) may be specified by implementing the results of [32, 33] in the form of rather convenient parametrizations [34]:

$$f_0(q^2) = \frac{0.33}{1 - 0.7 r_V + 0.27 r_V^2}, \quad r_V \equiv q^2/M_{B_s}^2; \quad (2.13)$$

$$A_0(q^2) = \frac{0.37}{(1 - 0.46 r_P)(1 - r_P)}, \quad r_P \equiv q^2/M_{B_s}^2. \quad (2.14)$$

Numerically, these parametrizations yield, for the ratio of differential distributions in the reactions  $B \rightarrow K \phi \rightarrow K \chi \bar{\chi}$  and  $B \rightarrow K^* \phi \rightarrow K^* \chi \bar{\chi}$ , the prediction depicted in Fig. 1 of [28]. More recent parametrizations of these form factors (for instance, those from [35, 36]) entail, for  $R_{K^*/K}^{(\phi)}$ , an uncertainty of about 2% [28].

### III. KINEMATICS

The Belle II experiment, as described in [1], analyzes partially reconstructed events to increase the statistical sample. For such events, the direction of the  $B$  meson cannot be determined. Therefore, instead of  $q^2 = (p_B - p_K)^2$  the variable  $q_{\text{rec}}^2$  is used:

$$q_{\text{rec}}^2 = E_B^2 + M_K^2 - 2E_B E_K, \quad (3.1)$$

where  $E_B$  and  $E_K$  are the energies of the  $B$  and  $K$  mesons in the center-of-mass frame of the  $B\bar{B}$ -meson pair produced in  $\Upsilon(4S)$  decays.

The variable  $q_{\text{rec}}^2$  can be expressed through the three-momenta  $\mathbf{p}_B$  and  $\mathbf{p}_K$  of the  $B$  and  $K$  mesons, and then through the velocity  $\mathbf{v}$  of the  $B$  meson in the  $B\bar{B}$  center-of-mass frame:

$$q_{\text{rec}}^2 = E_B^2 + M_K^2 - 2E_B E_K = q^2 + (E_B^2 - M_B^2) - 2(\mathbf{p}_K \cdot \mathbf{p}_B) = q^2 + \frac{M_B^2 v^2}{1 - v^2} - 2p_K^z \frac{M_B v}{\sqrt{1 - v^2}}, \quad (3.2)$$

where  $\mathbf{p}_K^z$  is the projection of  $\mathbf{p}_K$  onto the direction of the  $B$ -meson motion.

It is convenient to express  $\mathbf{p}_K^z$  as follows:

$$\mathbf{p}_K^z = \frac{\cos \Theta \mathbf{p}_K^0 - v E_K^0}{\sqrt{1 - v^2}}. \quad (3.3)$$

Here,  $\Theta$  is the angle between the direction of the  $K$ -meson motion in the  $B$ -meson rest frame and the direction of the  $B$ -meson motion in the  $B\bar{B}$  center-of-mass frame, while  $\mathbf{p}_K^0$  and  $E_K^0$  are, respectively, the projection of the  $K$ -meson three-momentum onto the  $B$ -meson direction, and the  $K$ -meson energy in the  $B$ -meson rest frame:

$$\mathbf{p}_K^0 = \frac{\lambda^{1/2}(M_B^2, M_K^2, q^2)}{2M_B}, \quad E_K^0 = \frac{M_B^2 + M_K^2 - q^2}{2M_B}. \quad (3.4)$$

By combining Eqs. (3.2), (3.3) and (3.4),  $q_{\text{rec}}^2$  becomes, in terms of  $q^2$ ,  $\Theta$  and  $v$ ,

$$q_{\text{rec}}^2 = \frac{q^2 - M_K^2 v^2 - \cos \Theta v \lambda^{1/2}(M_B^2, M_K^2, q^2)}{1 - v^2}. \quad (3.5)$$

This equation has an exact solution for  $q^2$ :

$$q^2 = \frac{1}{1 - \cos^2 \Theta v^2} \left\{ (1 - v^2) q_{\text{rec}}^2 + (1 - \cos^2 \Theta) v^2 M_K^2 - \cos^2 \Theta v^2 M_B^2 + \right. \\ \left. + \cos \Theta v \left[ \lambda \left( M_B^2, (1 - v^2) M_K^2, (1 - v^2) q_{\text{rec}}^2 \right) - 4(1 - \cos^2 \Theta) v^2 M_B^2 M_K^2 \right]^{1/2} \right\}. \quad (3.6)$$

Neglecting terms proportional to  $v^2$ , we obtain

$$q^2 \approx q_{\text{rec}}^2 + \cos \Theta v \lambda^{1/2} (M_B^2, M_K^2, q_{\text{rec}}^2). \quad (3.7)$$

Further, upon neglecting  $M_K$  we arrive at an even simpler relation between  $q^2$  and  $q_{\text{rec}}^2$ , which matches the previous expression within a few percent:

$$q^2 \approx q_{\text{rec}}^2 + (M_B^2 - q_{\text{rec}}^2) v \cos \Theta. \quad (3.8)$$

Since the  $B$  meson decays uniformly over  $\Theta$ , we have

$$\Gamma = \iint \frac{d^2 \Gamma(q^2, \cos \Theta)}{dq^2 d \cos \Theta} dq^2 d \cos \Theta = \frac{1}{2} \iint \frac{d\Gamma(q^2(q_{\text{rec}}^2, \cos \Theta))}{dq^2} \frac{dq^2(q_{\text{rec}}^2, \cos \Theta)}{dq_{\text{rec}}^2} dq_{\text{rec}}^2 d \cos \Theta. \quad (3.9)$$

Thus,

$$\frac{d\Gamma}{dq_{\text{rec}}^2} = \frac{1}{2} \int \frac{d\Gamma(q^2(q_{\text{rec}}^2, \cos \Theta))}{dq^2} \frac{dq^2(q_{\text{rec}}^2, \cos \Theta)}{dq_{\text{rec}}^2} d \cos \Theta. \quad (3.10)$$

Therefore, any decay characteristic  $\mathcal{A}(q_{\text{rec}}^2)$  at a given  $q_{\text{rec}}^2$  is the average of  $\mathcal{A}(q^2)$  over the range

$$q^2 \in [q_{\text{rec}}^2 - v \lambda^{1/2} (M_B^2, M_K^2, q_{\text{rec}}^2); q^2 + v \lambda^{1/2} (M_B^2, M_K^2, q_{\text{rec}}^2)]. \quad (3.11)$$

In particular, the averaged differential decay rate has the form

$$\overline{\frac{d\Gamma(q_{\text{rec}}^2)}{dq_{\text{rec}}^2}} \approx \left\langle \frac{d\Gamma(q^2)}{dq^2} \right\rangle_{q_{\text{rec}}^2 \pm v \lambda^{1/2} (M_B^2, M_K^2, q_{\text{rec}}^2)}. \quad (3.12)$$

In order to give an idea of the size of the averaging interval, we note that  $v \lambda^{1/2} (M_B^2, M_K^2, 0 \text{ GeV}^2) \approx 1.7 \text{ GeV}^2$  and  $v \lambda^{1/2} (M_B^2, M_K^2, 10 \text{ GeV}^2) \approx 1 \text{ GeV}^2$ .

It is instructive to consider the real  $\phi$ -mediator with a negligible decay width, in which case one finds

$$\frac{d\Gamma}{dq^2} (B \rightarrow K^{(*)} \phi) = \Gamma(B \rightarrow K^{(*)} \phi) \cdot \delta(q^2 - M_\phi^2). \quad (3.13)$$

So, in this case, the distributions over  $q_{\text{rec}}^2$  become rectangular functions with a width of  $2v \lambda^{1/2} (M_B^2, M_{K^{(*)}}^2, q_{\text{rec}}^2)$ :

$$\frac{d\Gamma}{dq_{\text{rec}}^2} (B \rightarrow K^{(*)} \phi) = \Gamma(B \rightarrow K^{(*)} \phi) \frac{\Theta(v \lambda^{1/2} (M_B^2, M_{K^{(*)}}^2, q_{\text{rec}}^2) - |q_{\text{rec}}^2 - M_\phi^2|)}{2v \lambda^{1/2} (M_B^2, M_{K^{(*)}}^2, q_{\text{rec}}^2)}. \quad (3.14)$$

#### IV. DATA FITTING

The Belle II experiment does not apply efficiency corrections to its data [1]. Consequently, in order to fit the data, we adopt the  $q^2$ -dependent efficiency  $\varepsilon(q^2)$  estimated in [23] (and provided by Fig. 2 of [23]),

$$\overline{\frac{d\Gamma_{\text{eff}}(q_{\text{rec}}^2)}{dq_{\text{rec}}^2}} = \left\langle \varepsilon(q^2) \frac{d\Gamma(q^2)}{dq^2} \right\rangle_{q_{\text{rec}}^2 \pm \lambda^{1/2} (M_B^2, M_K^2, q_{\text{rec}}^2)}. \quad (4.1)$$

To determine the optimal fit parameters, we minimize the  $\chi^2$  value, defined as

$$\chi^2 = \sum_i \frac{(n_{\text{exp}}^i - n_{\text{theor}}^i)^2}{(\Delta_{\text{exp}}^i)^2} + \frac{(N_{\text{exp}} - N_{\text{theor}})^2}{\Delta_{\text{exp}}^2}, \quad (4.2)$$

where the sum runs over the experimental bins,  $n_{\text{exp}}^i$  and  $n_{\text{theor}}^i$  are the measured and the predicted numbers of excess events over the SM expectation in the  $i$ -th bin, and  $\Delta_{\text{exp}}^i$  is the experimental error in the  $i$ -th bin;  $N_{\text{exp}}$  and  $N_{\text{theor}}$  are the full numbers of excess events,  $N_{\text{exp}} = \sum_i n_{\text{exp}}^i$  and  $N_{\text{theor}} = \sum_i n_{\text{theor}}^i$ , and  $\Delta_{\text{exp}}^2 = \sum_i (\Delta_{\text{exp}}^i)^2$ .

We fit the data under the following three assumptions:

1. The mediator  $\phi$  predominantly decays into a  $\chi\bar{\chi}$  pair.
2. The mass  $M_\phi$  of the mediator  $\phi$  is greater than the mass of the  $\chi\bar{\chi}$  pair:  $M_\phi > 2m_\chi$ .
3. The decay width  $\Gamma_\phi^0$  of the mediator  $\phi$  is smaller than its mass:  $\Gamma_\phi^0 < M_\phi$ .




Figure 1: The  $\chi^2$  distributions over the following variables: (a) the mass  $M_\phi$  of the mediator  $\phi$ , (b) the mass  $m_\chi$  of the light dark-matter fermion  $\chi$ , (c) the width  $\Gamma_\phi^0$  of the mediator  $\phi$ , and (d) the two-dimensional  $\chi^2$  distribution over  $M_\phi$  and  $m_\chi$ . The  $\chi^2$  minimum value,  $\chi_{\min}^2 = 4.65$ , has been subtracted. For each plot, the variables not being varied are fixed at their values corresponding to the  $\chi^2$  minimum:  $M_\phi = 2.8$  GeV,  $\Gamma_\phi^0 = 2.8$  GeV, and  $m_\chi = 0$  GeV.

## V. RESULTS

The fitting of the data indicates that the scenario under investigation is consistent with the experimental data and provides a good description of the shape of the observed excess. However, the limited statistics prevents the extraction of parameter values with high precision. Indeed, the regions with  $3\sigma$  significance are too broad to allow for a definitive choice. This is clearly illustrated in Fig. 1, which presents  $\chi^2$  distributions over  $\phi$  mass,  $\chi$  mass, and  $\phi$  width, as well as the two-dimensional  $\chi^2$  distribution over  $\phi$  mass and  $\chi$  mass.

Nevertheless, two preferred parameter regions with  $1\sigma$  significance can be identified:

1. The first region [located in the bottom-left corner of Fig. 1(d)] corresponds to the case where the observed excess is described by a broad resonance structure:  $2.2 \text{ GeV} < M_\phi < 3.1 \text{ GeV}$ ,  $1.8 \text{ GeV} < \Gamma_\phi^0 < 5 \text{ GeV}$ ,  $m_\chi < 0.8 \text{ GeV}$ .
2. The second region [located in the bottom-right corner of Fig. 1(d)] corresponds to the opposite case, where the observed excess is described by a non-resonant structure with  $M_\phi > 20 \text{ GeV}$  and  $m_\chi < 0.4 \text{ GeV}$ . In this scenario, the bell-shaped data distribution is reproduced by the phase space modulated by the efficiency. The propagator in (2.11) becomes independent of  $q^2$  for such parameter values and effectively acts as a constant.

Both parameter regions provide a very satisfactory description of the data, as demonstrated in Fig. 2.

Considering the first case, we assume that the mediator  $\phi$  decays predominantly into the  $\chi\bar{\chi}$  pair. This implies

$$\Gamma_\phi^0 = \frac{\kappa^2}{8\pi} M_\phi \left(1 - \frac{4m_\chi^2}{M_\phi^2}\right)^{3/2} \approx \frac{\kappa^2}{8\pi} M_\phi \quad (5.1)$$




Figure 2: The Belle II data on the decay  $B \rightarrow KM_X$  fitted by the DM model considered for different parameter values:  $M_\phi = 2.8$  GeV,  $\Gamma_\phi^0 = 2.8$  GeV and  $m_\chi = 0$  GeV (red solid curve);  $M_\phi = 20$  GeV,  $\Gamma_\phi^0 = 0$  GeV and  $m_\chi = 0$  GeV (green solid curve). The corresponding predictions for  $B \rightarrow K^*M_X$  are also shown, assuming the same detection efficiency (red dashed and green dashed curves). Only excess events over the SM expectations are shown.




Figure 3: Both predicted theoretical ratio  $R_{K^*/K}^{(\phi)}$  [which is independent of  $M_\phi$ ,  $\Gamma_\phi^0$  and  $m_\chi$  (blue solid curve)] and experimentally measurable ratio  $\overline{R_{K^*/K}^{(\phi)}}$  for different parameter values [ $M_\phi = 2.8$  GeV,  $\Gamma_\phi^0 = 2.8$  GeV and  $m_\chi = 0$  GeV (red dashed curve), and  $M_\phi = 20$  GeV,  $\Gamma_\phi^0 = 0$  GeV and  $m_\chi = 0$  GeV (black dotted-dashed curve)] vs.  $q_{\text{rec}}^2$ .

and

$$\kappa \approx \left( 8\pi \frac{\Gamma_\phi^0}{M_\phi} \right)^{1/2} \sim 5. \quad (5.2)$$

Taking into account the extracted value of  $\kappa$ , the total number of  $B$  mesons produced at Belle II ( $N_{\text{tot}} = 3.99 \cdot 10^8$ ), and the observed excess of approximately 170 events, we can estimate the coupling  $g_{b \rightarrow s\phi}$  in (2.11) as  $g_{b \rightarrow s\phi} \sim 7 \cdot 10^{-8}$ .

For the second, non-resonant case, where  $M_\phi$  is large and  $\Gamma_\phi^0$  is negligible, we obtain

$$\frac{\kappa g_{b \rightarrow s\phi}}{M_\phi^2} \sim 6 \cdot 10^{-8} \text{ GeV}^{-2}. \quad (5.3)$$

Our analysis shows that, if assuming the same detection efficiency for  $B \rightarrow K^*M_X$  and  $B \rightarrow KM_X$  decays, the ratio  $\overline{R_{K^*/K}^{(\phi)}}(q_{\text{rec}}^2)$  of excess events in the experimentally measured differential distributions in  $B \rightarrow K^*M_X$  and  $B \rightarrow KM_X$  decays,

$$\overline{R_{K^*/K}^{(\phi)}}(q_{\text{rec}}^2) = \frac{\overline{d\Gamma^{\text{eff}}(B \rightarrow K^*\chi\bar{\chi})/dq_{\text{rec}}^2}}{\overline{d\Gamma^{\text{eff}}(B \rightarrow K\chi\bar{\chi})/dq_{\text{rec}}^2}}, \quad (5.4)$$

is practically independent of the model parameters, such as  $\phi$  mass,  $\chi$  mass and  $\phi$  width, since the dependence on these parameters approximately cancels out in the ratio. [Recall that the dependence on these parameters cancels exactly in the theoretical ratio  $R_{K^*/K}^{(\phi)}$  of Eq. (2.12).]

Consequently, we have an approximate relation between the measured  $\overline{R_{K^*/K}^{(\phi)}}(q_{\text{rec}}^2)$  and the theoretical  $R_{K^*/K}^{(\phi)}(q_{\text{rec}}^2)$ , which is fulfilled with very high accuracy in a broad range of momentum transfers (see Fig. 3):

$$\overline{R_{K^*/K}^{(\phi)}}(q_{\text{rec}}^2) \simeq R_{K^*/K}^{(\phi)}(q_{\text{rec}}^2). \quad (5.5)$$

This useful result is not surprising, given the relation in (3.12). Notice, however, that if the reconstruction efficiencies for the decays  $B \rightarrow K^*M_X$  and  $B \rightarrow KM_X$  differ significantly, then (5.5) must be adjusted accordingly.

It is important to note that the experimental  $q_{\text{rec}}^2$  distribution reaches its maximum at  $q_{\text{rec}}^2 \approx 5 \text{ GeV}^2$ , where  $R_{K^*/K}^{(\phi)}(q_{\text{rec}}^2) \approx 1$ . This observation implies that the number of excess events in the decays  $B \rightarrow K^*M_X$  and  $B \rightarrow KM_X$  — assuming the same detection efficiency for both processes — should be approximately equal in magnitude and the corresponding differential distributions in  $q_{\text{rec}}^2$  should have similar shapes. This behavior is illustrated in Fig. 2, where predictions for the expected excess events for the  $B \rightarrow K^*M_X$  decay are represented by dashed curves.

## VI. CONCLUSIONS

In the current study, our aim was limited to the analysis of the impact of the enhancement in the  $B \rightarrow KM_X$  decay, observed by Belle II [1], on some DM scenario involving a scalar mediator. For this model, the following conclusions, based solely on Belle II data, hold:

- The model discussed is consistent with the experimental data and provides a good description of the shape of the excess observed. However, the limited statistics prevents the extraction of parameter values with high precision.
- An acceptable description of the data can be achieved for a scalar  $\phi$ -mediator with intermediate mass  $\sim 2.8 \text{ GeV}$ , as well as for a heavy  $\phi$ -mediator with a mass several times greater than the  $B$ -meson mass. In the first case, the data are described by a resonance structure; in the second case, the bell-shaped data distribution is reproduced by the phase space modulated by the detection efficiency. In both scenarios, a small mass value has to be chosen for the DM fermion  $\chi$ .
- Both the shape and the normalization of the ratio of the excess events which may be measured experimentally,  $\overline{R_{K^*/K}^{(\phi)}}(q_{\text{rec}}^2)$ , may be well approximated by the theoretical ratio (2.12) evaluated at the same  $q_{\text{rec}}^2$ ,  $R_{K^*/K}^{(\phi)}(q_{\text{rec}}^2)$ :

$$\overline{R_{K^*/K}^{(\phi)}}(q_{\text{rec}}^2) \simeq R_{K^*/K}^{(\phi)}(q_{\text{rec}}^2).$$

Consequently, the shape of  $\overline{R_{K^*/K}^{(\phi)}}(q_{\text{rec}}^2)$  is largely independent on the DM model parameters — such as  $M_\phi$ ,  $\Gamma_\phi$  and  $m_\chi$  — and provides a clear signature of the DM scenario based on the scalar mediator field.

- Within the framework of the model considered, the yields of the excess events over the SM background for the decays  $B \rightarrow K^*M_X$  and  $B \rightarrow KM_X$  should be approximately equal in magnitude and exhibit a qualitatively similar shape in the  $q_{\text{rec}}^2$  distribution, assuming the same detection efficiency for both reactions.

These constraints on DM particles may be combined with constraints coming from other phenomena (see, e.g., [6]) but such kind of analysis is beyond the scope of our interest in this analysis.

*Acknowledgments.* The research was carried out within the framework of the program *Particle Physics and Cosmology* of the National Center for Physics and Mathematics.

- 
- [1] I. Adachi et al. (Belle-II), *Evidence for  $B^+ \rightarrow K^+ \nu \nu$  decays*, Phys. Rev. D **109**, 112006 (2024), arXiv:2311.14647 [hep-ex].  
[2] L. Allwicher, D. Becirevic, G. Piazza, S. Rosauero-Alcaraz, and O. Sumensari, *Understanding the first measurement of  $B(B \rightarrow K \nu \nu)$* , Phys. Lett. B **848**, 138411 (2024), arXiv:2309.02246 [hep-ph].  
[3] P. Athron, R. Martinez, and C. Sierra, *B meson anomalies and large  $B \rightarrow K \nu \nu$  in non-universal  $U(1)'$  models*, JHEP **02**, 121 (2024), arXiv:2308.13426 [hep-ph].

- [4] R. Bause, H. Gisbert, and G. Hiller, *Implications of an enhanced  $B \rightarrow K\nu\nu$  branching ratio*, Phys. Rev. D **109**, 015006 (2024), arXiv:2309.00075 [hep-ph].
- [5] T. Felkl, A. Giri, R. Mohanta, and M. A. Schmidt, *When energy goes missing: new physics in  $b \rightarrow s\nu\nu$  with sterile neutrinos*, Eur. Phys. J. C **83**, 1135 (2023), arXiv:2309.02940 [hep-ph].
- [6] M. Abdughani and Y. Reyimuaji, *Constraining light dark matter and mediator with  $B \rightarrow K\nu\nu$  data*, Phys. Rev. D **110**, 055013 (2024), arXiv:2309.03706 [hep-ph].
- [7] H. K. Dreiner, J. Y. Guenther, and Z. S. Wang, *The Decay  $B \rightarrow K\nu\nu$  at Belle II and a Massless Bino in R-parity-violating Supersymmetry*, Eur. Phys. J. C **85**, 66 (2025), arXiv:2309.03727 [hep-ph].
- [8] X.-G. He, X.-D. Ma, and G. Valencia, *Revisiting models that enhance  $B \rightarrow K\nu\nu$  in light of the new Belle II measurement*, Phys. Rev. D **109**, 075019 (2024), arXiv:2309.12741 [hep-ph].
- [9] L. Calibbi, T. Li, L. Mukherjee, and M. A. Schmidt, *Is Dark Matter the origin of the  $B \rightarrow K\nu\nu$  excess at Belle II?*, arXiv:2502.04900 [hep-ph].
- [10] W. Altmannshofer, S. A. Gadam, and K. Toner, *New Strategies for New Physics Search with  $\Lambda_b \rightarrow \Lambda\nu\nu$  Decays*, arXiv:2501.10652 [hep-ph].
- [11] C.-Q. Zhang, J. Sun, Z.-P. Xing, and R.-L. Zhu, *Probing  $B \rightarrow K^{(*)}$  semi-leptonic FCNC decay with new physics effects under PQCD approach*, arXiv:2501.00512 [hep-ph].
- [12] B. Bhattacharya, A. Datta, G. Faisel, S. Khalil, and S. Roy, *Flavor Violations in B-Mesons within Non-Minimal SU(5)*, arXiv:2412.16115 [hep-ph].
- [13] W. Altmannshofer and S. Roy, *A joint explanation of the  $B \rightarrow \pi K$  puzzle and the  $B \rightarrow K\nu\nu$  excess*, arXiv:2411.06592 [hep-ph].
- [14] H. Davoudiasl and M. Schnubel, *Bringing the Peccei-Quinn mechanism down to Earth*, Phys. Rev. D **110**, 075014 (2024), arXiv:2406.19455 [hep-ph].
- [15] C. S. Kim, D. Sahoo, and K. N. Vishnudath, *Searching for signatures of new physics in  $B \rightarrow K\nu\nu$  to distinguish between Dirac and Majorana neutrinos*, Eur. Phys. J. C **84**, 882 (2024), arXiv:2405.17341 [hep-ph].
- [16] D. Marzocca, M. Nardecchia, A. Stanzione, and C. Toni, *Implications of  $B \rightarrow K\nu\nu$  under rank-one flavor violation hypothesis*, Eur. Phys. J. C **84**, 1217 (2024), arXiv:2404.06533 [hep-ph].
- [17] P. D. Bolton, S. Fajfer, J. F. Kamenik, and M. Novoa-Brunet, *Signatures of light new particles in  $B \rightarrow K^{(*)}E_{\text{miss}}$* , Phys. Rev. D **110**, 055001 (2024), arXiv:2403.13887 [hep-ph].
- [18] X.-G. He, X.-D. Ma, M. A. Schmidt, G. Valencia, and R. R. Volkas, *Scalar dark matter explanation of the excess in the Belle II  $B \rightarrow K+$  invisible measurement*, JHEP **07**, 168 (2024), arXiv:2403.12485 [hep-ph].
- [19] B.-F. Hou, X.-Q. Li, M. Shen, Y.-D. Yang, and X.-B. Yuan, *Deciphering the Belle II data on  $B \rightarrow K\nu\nu$  decay in the (dark) SMEFT with minimal flavour violation*, JHEP **06**, 172 (2024), arXiv:2402.19208 [hep-ph].
- [20] E. Gabrielli, L. Marzola, K. Muursepp, and M. Raidal, *Explaining the  $B \rightarrow K\nu\nu$  excess via a massless dark photon*, Eur. Phys. J. C **84**, 460 (2024), arXiv:2402.05901 [hep-ph].
- [21] F. Loporco, *A New Look at  $b \rightarrow s$  Observables in 331 Models*, Particles **7**, 161 (2024), arXiv:2401.11999
- [22] S.-Y. Ho, J. Kim, and P. Ko, *Recent  $B \rightarrow K\nu\nu$  Excess and Muon  $g - 2$  Illuminating Light Dark Sector with Higgs Portal*, arXiv:2401.10112 [hep-ph].
- [23] K. Fridell, M. Ghosh, T. Okui, and K. Tobioka, *Decoding the  $B \rightarrow K\nu\nu$  excess at Belle II: Kinematics, operators, and masses*, Phys. Rev. D **109**, 115006 (2024), arXiv:2312.12507 [hep-ph].
- [24] D. McKeen, J. N. Ng, and D. Tuckler, *Higgs portal interpretation of the Belle II  $B \rightarrow K\nu\nu$  measurement*, Phys. Rev. D **109**, 075006 (2024), arXiv:2312.00982 [hep-ph].
- [25] W. Altmannshofer, A. Crivellin, H. Haigh, G. Inguglia, and J. Martin Camalich, *Light new physics in  $B \rightarrow K^{(*)}\nu\nu$ ?*, Phys. Rev. D **109**, 075008 (2024), arXiv:2311.14629 [hep-ph].
- [26] Z. S. Wang, H. K. Dreiner, and J. Y. Guenther, *The decay  $B \rightarrow K\nu\nu$  at Belle II and a massless bino in R-parity-violating supersymmetry*, Eur. Phys. J. C **85**, 66 (2025), arXiv:2309.03727 [hep-ph].
- [27] X. G. He, X. D. Ma, J. Tandean and G. Valencia,  *$B \rightarrow K+$ invisible, dark matter, and CP violation in hyperon decays*, arXiv:2502.09603 [hep-ph].
- [28] A. Berezhnoy and D. Melikhov,  *$B \rightarrow K^*M_X$  vs  $B \rightarrow KM_X$  as a probe of a scalar-mediator dark matter scenario*, EPL **145**, 14001 (2024), arXiv:2309.17191 [hep-ph].
- [29] B. Batell, M. Pospelov, and A. Ritz, *Multi-lepton Signatures of a Hidden Sector in Rare B Decays*, Phys. Rev. D **83**, 054005 (2011), arXiv:0911.4938 [hep-ph].
- [30] K. Schmidt-Hoberg, F. Staub, and M. W. Winkler, *Constraints on light mediators: confronting dark matter searches with B physics*, Phys. Lett. B **727**, 506 (2013), arXiv:1310.6752 [hep-ph].
- [31] M. Wirbel, B. Stech, and M. Bauer, *Exclusive Semileptonic Decays of Heavy Mesons*, Z. Phys. C **29**, 637 (1985).
- [32] P. Ball and R. Zwicky, *New results on  $B \rightarrow \pi, K, \eta$  decay formfactors from LCSR*, Phys. Rev. D **71**, 014015 (2005).
- [33] P. Ball and R. Zwicky,  *$B_{d,s} \rightarrow \rho, \omega, K^*, \phi$  decay form-factors from LCSR*s revisited, Phys. Rev. D **71**, 014029 (2005).
- [34] D. Melikhov and B. Stech, *Weak form-factors for heavy meson decays: an update*, Phys. Rev. D **62**, 014006 (2000).
- [35] D. Becirevic, G. Piazza, and O. Sumensari, *Revisiting  $B \rightarrow K^{(*)}\nu\bar{\nu}$  decays in the Standard Model and beyond*, Eur. Phys. J. C **83**, 252 (2023).
- [36] W. G. Parrott, C. Bouchard, and C. T. H. Davies [HPQCD collaboration], *Standard Model predictions for  $B \rightarrow Kl^+l^-$ ,  $B \rightarrow Kl_1^-l_2^+$ , and  $B \rightarrow K\nu\bar{\nu}$  using form factors from  $N_f = 2 + 1 + 1$  lattice QCD*, Phys. Rev. D **107**, 014511 (2023); **107**, 119903(E) (2023).