

Computing the Polytope Diameter is Even Harder than NP-hard (Already for Perfect Matchings)

Lasse Wulf

Section of Algorithms, Logic and Graphs, Technical University of Denmark, Kongens Lyngby, Denmark, lawu@dtu.dk

Abstract

The diameter of a polytope is a fundamental geometric parameter that plays a crucial role in understanding the efficiency of the simplex method. Despite its central nature, the computational complexity of computing the diameter of a given polytope is poorly understood. Already in 1994, Frieze and Teng [Comp. Compl.] recognized the possibility that this task could potentially be harder than NP-hard, and asked whether the corresponding decision problem is complete for the second stage of the polynomial hierarchy, i.e. Π_2^P -complete. In the following years, partial results could be obtained. In a cornerstone result, Frieze and Teng themselves proved weak NP-hardness for a family of custom defined polytopes. Sanità [FOCS18] in a break-through result proved that already for the much simpler fractional matching polytope the problem is strongly NP-hard. Very recently, Steiner and Nöbel [SODA25] generalized this result to the even simpler bipartite perfect matching polytope and the circuit diameter. In this paper, we finally show that computing the diameter of the bipartite perfect matching polytope is Π_2^P -hard. Since the corresponding decision problem is also trivially contained in Π_2^P , this decidedly answers Frieze and Teng’s 30 year old question. As a consequence, computation of the diameter can be compared to the computation of other notoriously hard parameters like the generalized Ramsey number.

Our results in particular hold even when the constraint matrix of the given polytope is totally unimodular. They also hold when the diameter is replaced by the circuit diameter.

As our second main result, we prove that for some $\varepsilon > 0$ the (circuit) diameter of the bipartite perfect matching polytope cannot be approximated by a factor better than $(1 + \varepsilon)$. This answers a recent question by Nöbel and Steiner. It is the first known inapproximability result for the circuit diameter, and extends Sanità’s inapproximability result of the diameter to the totally unimodular case.

Keywords: Diameter, circuit diameter, computational complexity, polynomial hierarchy, second level, bipartite perfect matching polytope, combinatorial reconfiguration, Π_2^P , Σ_2^P , Hirsch conjecture, simplex method, 1-skeleton

Funding: This work was supported by Eva Rotenberg’s Carlsberg Foundation Young Researcher Fellowship CF21-0302 “Graph Algorithms with Geometric Applications”.

Acknowledgements: I am thankful to Eva Rotenberg for great encouragement and very helpful discussions.

1 Introduction

The *diameter* of a polytope is the maximum length of a shortest path (in terms of edges) between any two vertices of the polytope. This property plays a crucial role in understanding the complexity of algorithms that traverse polytopes, such as the simplex algorithm for linear programming.

The famous simplex algorithm, as invented in its basic form in the 1950s by George Dantzig, aims to solve *linear programming*, i.e. to maximize a linear functional $c^t x$ over a polytope P . Roughly speaking, the simplex algorithm works such that in each step, the algorithm visits one of the vertices of P , and a *pivot rule* decides which of the incident edges (in the 1-skeleton of P) to traverse in order to visit the next vertex. The performance of the algorithm is hence closely tied to the polytope's diameter. If the diameter is small relative to the polytope's size, fewer steps might be needed to find an optimal solution, making the algorithm more efficient. However, if the diameter is large relative to the polytope's size, then under *any* pivot rule the simplex algorithm needs to perform many steps in the worst case. More precisely, if $\text{diam}(P)$ denotes the diameter of P , there exists an initialization vertex of the simplex algorithm and a direction $c \in \mathbb{R}^n$, such that the algorithm needs to perform at least $\text{diam}(P)$ steps.

One of the most famous open questions of mathematical optimization, as re-iterated by Smale in his list of unsolved problems for the 21st century [Sma98], is whether there exists a strongly polynomial time algorithm for linear programming. Maybe surprisingly, the simplex algorithm is still a potential candidate for a positive resolution of this conjecture – while for most known pivot rules counterexamples have been found which cause the simplex algorithm to have superpolynomial running time (see e.g. [DFH23, KM72]), the existence of a pivot rule such that the simplex algorithm has strongly polynomial runtime still remains a famous open question.

Since the runtime of the simplex algorithm is lower-bounded by $\text{diam}(P)$, a necessary condition for such a strongly polynomial pivot rule would be that for every polytope $P \subseteq \mathbb{R}^d$ with n facets, its diameter is bounded by a polynomial function in n and d . This condition has become known under the name *polynomial Hirsch conjecture*.

$$\text{diam}(P) \leq \text{poly}(n, d) \text{ for all polytopes } P \subseteq \mathbb{R}^d \text{ with } n \text{ facets.}$$

Much like the existence of a strongly polynomial pivot rule, the polynomial Hirsch conjecture is a famous conjecture in the theory of mathematical optimization and convex geometry, and still remains wide open. It is a relaxation of the *classic Hirsch conjecture*, which was established by Hirsch in 1957 and stated that $\text{diam}(P) \leq n - d$ for all polytopes. The classic Hirsch conjecture has since been disproven, first for unbounded polytopes by Klee and Walkup [KW67] and finally for bounded polytopes in 2012 by Santos [San12]. However, even today the best known counterexamples are quite close to Hirsch' original conjecture, in the sense that Matschke, Santos and Weibel found polytopes P with $\text{diam}(P) \geq \frac{21}{20}(n - d)$ [MSW15]. In the important special case of 0/1 polytopes, i.e. polytopes with the property that all vertices lie on 0/1-integer coordinates, the Hirsch conjecture is known to be true, as shown by Naddef [Nad89].

The complexity of computing the diameter. We consider the problem DIAMETER of computing the diameter of a polytope. More specifically, since we are interested in questions regarding computational complexity, it is more convenient to talk about its corresponding decision version DIAMETER-DECISION. These problems are defined as follows.

Problem: DIAMETER

Input: $A \in \mathbb{Q}^{m \times d}$ and $b \in \mathbb{Q}^m$ describing the polytope $P = \{x \in \mathbb{R}^d : Ax \leq b\}$.

Task: Compute the diameter $\text{diam}(P)$.

Problem: DIAMETER-DECISION

Input: $A \in \mathbb{Q}^{m \times d}$ and $b \in \mathbb{Q}^m$ describing $P = \{x \in \mathbb{R}^d : Ax \leq b\}$, and some $t \in \mathbb{N}$.

Question: Is $\text{diam}(P) \leq t$?

Since the diameter of a polytope is the central object of study of the Hirsch conjecture, which has been established already in 1957, one would expect that the computational complexity of DIAMETER and DIAMETER-DECISION is well understood. However, that is not the case. As a cornerstone result, Frieze and Teng [FT94] proved that DIAMETER is weakly NP-hard for a family of custom defined polytopes. Sanità [San18] in a break-through result proved that already for the much simpler fractional matching polytope the problem DIAMETER is strongly NP-hard. Very recently, Nöbel and Steiner [NS25] generalized this result to the even simpler bipartite perfect matching polytope. However, none of these papers could establish whether the problem DIAMETER-DECISION is *actually contained* in the class NP. Indeed, this is not a coincidence. Already in 1994, Frieze and Teng recognized the possibility that the problem DIAMETER-DECISION could be even harder than NP-hard and asked

Question 1 (Frieze and Teng [FT94]). *Is DIAMETER-DECISION Π_2^P -complete?*

Here, the class $\Pi_2^P = \text{co-}\Sigma_2^P$ denotes the class in the second level of the *polynomial hierarchy* introduced by Stockmeyer [Sto76]. Roughly speaking, the class Π_2^P contains all the problems of the form: FOR ALL objects x , does there EXIST some object y such that some efficiently testable property $\mathcal{P}(x, y)$ holds? This is a natural fit to DIAMETER-DECISION, since $\text{diam}(P)$ is small if and only if FOR ALL vertex pairs, there EXISTS a short path in P connecting the pair. Similar to the $P \neq NP$ conjecture, complexity theorists believe that $\text{co-NP} \neq \Pi_2^P$, or equivalently, that the polynomial hierarchy does not collapse to the first level [Woe21]. If this belief is true, it means that every Π_2^P -complete is not contained in NP (or co-NP), since it is complete for a much larger complexity class. Indeed, we show in this paper that Question 1 has a positive answer, already for the bipartite perfect matching polytope. This provides an explanation why previous papers could not establish membership of DIAMETER-DECISION to NP. The precise result together with its implications will be explained below.

Bipartite perfect matchings. All our results hold already for the very well known and widely studied *bipartite perfect matching polytope*, which has countless important applications in combinatorial optimization. Given a bipartite graph $G = (V, E)$, let us define the characteristic vector $\chi^M \in \mathbb{R}^E$ of some perfect matching $M \subseteq E$ as the edge indexed vector $(x_e)_{e \in E}$ that has $x_e = 1$ if $e \in M$ and $x_e = 0$ otherwise. The bipartite perfect matching polytope of G is then defined as the convex hull of all characteristic vectors of perfect matchings.

$$P_G := \text{conv}\{\chi^M : M \subseteq E \text{ is a perfect matching of } G\}$$

In particular, the vertices of P_G correspond bijectively to perfect matchings of G . It is well-known that P_G has the following compact halfspace-encoding (see e.g. [KV11]). Here the term $\delta(v)$ denotes the set of edges incident to vertex v .

$$\begin{aligned} \sum_{e \in \delta(v)} x_e &= 1 & \forall v \in V \\ x_e &\geq 0 & \forall e \in E \end{aligned}$$

1.1 Our contribution

We present two main results. As our first main, result, we decidedly answer Frieze and Tengs' 30 year old question. Let BPM-DIAMETER-DECISION denote the restriction of DIAMETER-DECISION to only the bipartite perfect matching polytope, then

Theorem 1. BPM-DIAMETER-DECISION is Π_2^P -complete.

In particular, computing the diameter of a polytope, i.e. problem DIAMETER, is Π_2^P -hard, even when restricted to totally unimodular constraint matrices. Since $\Pi_2^P \supseteq \text{NP}$ is a larger complexity class than the class NP (and this inclusion is strict under the complexity-theoretic assumption that the polynomial hierarchy does not collapse to the first level), this result is stronger than the hardness results of [FT94, San18, NS25].

Is it also true that DIAMETER-DECISION $\in \Pi_2^P$ for all families of polytopes beside the bipartite perfect matching polytope? A standard argument shows that for all families \mathcal{F} of polytopes where it is known that the polynomial Hirsch conjecture holds, we indeed have DIAMETER-DECISION $\in \Pi_2^P$ restricted to that family \mathcal{F} . Hence if the polynomial Hirsch conjecture is true, we have that DIAMETER-DECISION is Π_2^P -complete and our hardness result cannot be improved further (than the class Π_2^P). Even without the assumption that the polynomial Hirsch conjecture is true, our hardness result cannot be improved further for the bipartite perfect matching polytope, and the family of all 0/1-polytopes, since here the Hirsch conjecture is already known to hold.

What are the consequences of a problem being Π_2^P -complete? Such problems are in general much harder to solve than even NP-complete problems. (An illustrative explanation of that fact is given e.g. by Woeginger [Woe21].) Since integer programming is contained in NP, unless the polynomial hierarchy collapses, our result implies the diameter of a polytope cannot be computed by a mixed integer program with only polynomially many variables and constraints. As a consequence, computation of the diameter can be compared to other notoriously hard-to-compute parameters, like the generalized Ramsey number (which Schaefer showed to be Π_2^P -complete [Sch01]).

Since exact computation of the diameter turns out to be very hard, we turn to approximation of the diameter and prove as our second main result

Theorem 2. *There exists $\varepsilon > 0$ such that the diameter of the bipartite perfect matching polytope cannot be approximated better than $(1 + \varepsilon)$ in polynomial time, unless $P \neq \text{NP}$.*

Our proof of Theorem 2 is an adaptation of a recent proof of Nöbel and Steiner [NS25]. Interestingly, Nöbel and Steiner suspected that their methods could probably not be adapted to show an inapproximability result. Hence we show that their suspicion is not the case, if only in the weak sense of a $(1 + \varepsilon)$ -approximation. Sanità showed that finding the diameter of the fractional matching polytope is an APX-hard problem [San18]. Hence our Theorem 2 can be understood as an extension of Sanità’s result to the bipartite perfect matching polytope (and thus the totally unimodular case).

1.2 Applications to the circuit diameter

In 2015, Borgwardt, Finhold and Hemmecke [BFH15] proposed a generalization of the diameter of a polytope, called the *circuit diameter*. This concept is motivated by so-called *circuit augmentation schemes*, which are generalizations of the simplex algorithm and have attracted a lot of attention over the last decade. The basic idea behind circuit augmentation schemes is that we would like to move more freely through some polytope P than only traversing its edges. These algorithms consider a set $\mathcal{C}(P)$ of potential directions, called the *circuits* of the polytope. For the sake of completeness, we include here the definition of the set $\mathcal{C}(P)$ (although it is not needed to understand the rest of the paper).

Definition (from [BGK⁺25]). *Let $P = \{x \in \mathbb{R}^d : Ax = a, Bx \leq b\}$ be a polyhedron. The set of circuits of P , denoted by $\mathcal{C}(P)$ with respect to its linear description A, B consists of all vectors $g \in \ker(A) \setminus \{0\}$ for which Bg is support-minimal in the set $\{Bx : x \in \ker(A) \setminus \{0\}\}$ and g has coprime integral components.*

It can be shown that $\mathcal{C}(P)$ always includes the original edge directions of P . A single *circuit move* starting from some $x \in P$ now selects some circuit $g \in \mathcal{C}(P)$ and moves maximally along direction g , i.e. to the point $x + \lambda g$ with the property that $\lambda \geq 0$ is maximal with respect to $x + \lambda g \in P$. Hence circuit moves are allowed to cross the interior of the polytope and end up at non-vertices of P . This fact makes circuit augmentation schemes more powerful than the simplex algorithm and a compelling subject of study. Indeed, many different pivot rules and algorithms for circuit augmentation schemes have been proposed by various authors over the last 10 years [DLHL15, BV20, BBFK21, LKS22, DKNV22]. Analogous to the regular distance and diameter, we can define the *circuit distance* between $x, y \in P$ as the minimum number of circuit moves necessary to reach y from x , and the *circuit diameter* $\text{cdiam}(P)$ as the maximum circuit distance over all vertex pairs of P . Analogously to the simplex algorithm, $\text{cdiam}(P)$ gives a lower bound for the worst-case runtime of circuit augmentation schemes under *any* pivot rule. Remarkably, De Loera, Kafer and Sanità came very close to proving the polynomial “circuit” Hirsch conjecture, in the sense that they proved that for every polytope P , its circuit diameter is bounded by a polynomial in n, d and the maximum encoding-length among the coefficients in its input encoding [LKS22]. The classic Hirsch conjecture adapted to the circuit diameter, as proposed by Borgwardt, Finhold and Hemmecke [BFH15] still remains open. In order to shed light on this question, multiple authors have studied the circuit diameter of different families of polytopes, e.g. in [SY15, BSY18, KPS19, DKNV22, BBB23].

The question about the computational complexity of computing the circuit diameter was raised by Sanità [San20] and reiterated by Kafer [Kaf22] and Borgwardt et al. [BGK⁺25]. This question was recently (partially) settled by Nöbel and Steiner [NS25], who showed strong NP-hardness. Analogously to the circuit diameter, they could not show containment in NP.

The main results in this paper have interesting consequences for the circuit diameter. This is because all our results are obtained for the perfect matching polytope, and for this polytope it is actually known that $\text{diam}(P) = \text{cdiam}(P)$ (see e.g. Corollary 4 in [BV22] or Lemma 2 in [CS23a]). Hence we immediately obtain the following corollaries of our main theorems. Let CDIAM-DECISION denote the decision problem corresponding to computing the circuit diameter, then

Corollary 3. *CDIAM-DECISION is Π_2^P -complete, even when restricted to the bipartite perfect matching polytope.*

This strengthens the recent NP-hardness result of Nöbel and Steiner [NS25] and shows that just like the diameter, the circuit diameter is even harder to compute than NP-hard. We remark that in contrast to Theorem 1, we automatically have the containment $\text{CDIAM-DECISION} \in \Pi_2^P$ due to the almost polynomial “circuit” Hirsch conjecture of [LKS22]. Hence this hardness result cannot be strengthened further than the class Π_2^P . As a corollary of our second main theorem, we have

Corollary 4. *There exists $\varepsilon > 0$, such that the circuit diameter of a polytope cannot be approximated by a factor better than $(1 + \varepsilon)$, unless $P = \text{NP}$.*

Corollary 4 provides the first known inapproximability result for the circuit diameter of a polytope. As a consequence, unless $P = \text{NP}$, no circuit augmentation scheme can find the optimal solution in worst-case runtime of $(1 + \varepsilon) \text{cdiam}(P)$ steps or less. This can be compared to the result of Cardinal and Steiner [CS23a], who show that no circuit augmentation scheme can have worst-case runtime $O(\log n / \log \log n)$ to find a circuit walk between two vertices of distance two (assuming ETH). It is a very interesting question to ask whether our inapproximability result can be improved to any constant factor.

1.3 Related work

There are several other works that deal not with the diameter, but rather with the problem of computing the *distance* of two given vertices of some polytope P . In the case of the distance this problem was shown to be NP-hard for the bipartite perfect matching polytope P_G independently by Aichholzer et al. [ACH⁺21] and Ito et al. [IKK⁺22]. Additionally, Borgward et al. [BGK⁺25] show NP-hardness of computing circuit distances in 0/1-network flow polytopes, and Cardinal and Steiner show the same for polymatroids [CS23b]. These results were strengthened by Cardinal and Steiner [CS23a] to a $O(\log n / \log \log n)$ -hardness of approximation result for computing a short path between two vertices of distance two on the bipartite perfect matching polytope, assuming ETH. As explained in [NS25], the techniques for handling polytope diameters and distances are very different, since distances can be very short (which is often helpful for proofs), but diameters are usually large.

Finally, a related series of work deals with *combinatorial reconfiguration* of perfect matchings [IDH⁺11, KMM12, GKM19, BHIM19, BBH⁺19, BMPT25]. In these papers, the notion of adjacency between two perfect matchings differs from the adjacency on the bipartite perfect matching polytope considered in the present paper.

2 Technical Overview

We give a quick overview of the arguments used to obtain the main result of the paper. Recall the following. A perfect matching (abbreviated PM) of a graph $G = (V, E)$ is a subset $M \subseteq E$ such that every vertex is touched exactly once. A path or cycle is *alternating* with respect to some PM M if it alternates between edges in M and not in M . An alternating path is explicitly allowed to start and/or end with an edge not in M . It is well-known that given two PMs M_1, M_2 , their *symmetric difference* $M_1 \triangle M_2 = (M_1 \setminus M_2) \cup (M_2 \setminus M_1)$ is a vertex-disjoint union of alternating cycles. Since we are concerned with shortest paths in the bipartite perfect matching polytope, we first need to understand, when two of its vertices are adjacent. This is a well-known fact first shown by Chvátal.

Lemma ([Chv75], see also [NS25]). *Let $G = (V, E)$ be a bipartite graph, and let $x = \chi^M$ and $y = \chi^N$ be two vertices of the polytope P_G , corresponding to two perfect matchings $M, N \subseteq E$. The vertices x and y are adjacent in the 1-skeleton of P if and only if the edges in the symmetric difference $M \triangle N$ form a single alternating cycle.*

Based on this insight, we define the following: *Flipping* an alternating cycle C means transforming the PM M into the PM $M \triangle C$. A *flip sequence* of length d from a PM M_1 to a PM M_2 is a sequence (C_1, \dots, C_d) such that for all $i \in [d]$, the cycle C_i is alternating with respect to the perfect matching $M \triangle C_1 \triangle \dots \triangle C_{i-1}$, and we have $M_2 = M_1 \triangle C_1 \triangle \dots \triangle C_d$. For two PMs M_1, M_2 , we define $\text{dist}(M_1, M_2)$ as the length of the shortest flip sequence from M_1 to M_2 . Finally, we define problem BPM-DIAMETER-DECISION as the restriction of problem DIAMETER-DECISION to the special case of the bipartite perfect matching polytope. This means that the input is a bipartite graph G , and some threshold $t \in \mathbb{N}$. The problem is to decide whether $\text{diam}(P_G) \leq t$, or alternatively whether

$$\max\{\text{dist}(M_1, M_2) : M_1, M_2 \text{ PM of } G\} \leq t.$$

Overview of the first main result. We wish to prove that BPM-DIAMETER-DECISION is Π_2^P -complete. As a starting point, one can look at the classic Π_2^P -complete problem $\forall\exists$ -SAT [Sto76]. Here, we are given a SAT-formula φ in conjunctive normal form, such that its variables are partitioned into two disjoint parts X, Y (and such that w.l.o.g. every clause contains exactly three literals). The question is whether for all assignments of X there exists an assignment of

Y that makes φ true, i.e.

$$\forall \alpha \in \{0, 1\}^X : \exists \beta \in \{0, 1\}^Y : \varphi(\alpha, \beta) = 1.$$

One can compare this to the definition of the diameter. The diameter of a polytope P is at most d if and only if for all vertex pairs of P there is a path of length at most d in the 1-skeleton connecting those two vertices. In the special case of the bipartite perfect matching polytope P_G of some graph G this means that

$$\forall M_1, M_2 \text{ a PM of } G : \exists \text{ a flip sequence of length at most } d \text{ from } M_1 \text{ to } M_2. \quad (1)$$

Our main idea is to combine recent advances in understanding of both the diameter of the bipartite perfect matching polytope, as well as recent understandings of the typical structure of Π_2^p -complete problems. In particular, the clever construction of Nöbel and Steiner [NS25] manages to transform a given graph H into another graph G such that short flip sequences in G relate closely to Hamiltonian cycles in H . They use this to obtain an elegant proof for the NP-hardness of DIAMETER. On the other hand, Grüne and Wulf [GW24] recently introduced a wide-ranging meta-theorem that characterizes a “typical” problem complete for the class $\Sigma_2^p = \text{co-}\Pi_2^p$. (A similar result is also presented in [GW25].) The study of [GW24] motivates the definition of the following Π_2^p -complete variant of the Hamiltonian cycle problem. Let $k \in \mathbb{N}$. Given a directed graph H together with a set $E' = \{e_1, \bar{e}_1, \dots, e_k, \bar{e}_k\}$ of $2k$ distinct arcs, consider a subset $P \subseteq E'$. We say that P is a *pattern*, if $|P \cap \{e_i, \bar{e}_i\}| = 1$ for all $i \in [k]$. Some Hamiltonian cycle C in H *respects* the pattern P , if $C \cap E' = P$. In other words, the pattern determines for each pair $\{e_i, \bar{e}_i\}$, which one of these two arcs the Hamiltonian cycle is allowed to use (and which one it is not allowed to use). We prove in Section 4.2 using techniques from [GW24] that the following problem is Π_2^p -complete:

Problem: $\forall\exists$ -HAMCYCLE

Input: Directed graph H , distinct vertices $v^{(1)}, \dots, v^{(k)}$ for some $k \in \mathbb{N}$, such that for all $i \in [k]$, the vertex $v^{(i)}$ has exactly two outgoing arcs, denoted by e_i, \bar{e}_i . The set $E' := \{e_1, \bar{e}_1, \dots, e_k, \bar{e}_k\}$.

Question: Is the following true:

\forall patterns $P \subseteq E' : \exists$ a Hamiltonian cycle C respecting P ?

We reduce $\forall\exists$ -HAMCYCLE to BPM-DIAMETER-DECISION. In Section 4.5, we show: Given an instance of $\forall\exists$ -HAMCYCLE, consisting out of a directed graph H on $n = |V(H)|$ vertices together with arcs $E' = \{e_1, \bar{e}_1, \dots, e_k, \bar{e}_k\}$, one can in polynomial time construct an undirected bipartite graph G_H on $\Theta(n^9)$ vertices such that the following two lemmas hold:

Lemma 5. *If for all patterns $P \subseteq E'$ the directed graph H contains a Hamiltonian cycle respecting P , then $\text{diam}(P_{G_H}) \leq 4n^4 + 46n$.*

Lemma 6. *If $\text{diam}(P_{G_H}) \leq 4n^4 + 46n$, then for all patterns $P \subseteq E'$ the directed graph H contains a Hamiltonian cycle respecting P .*

Together, these two lemmas prove our main theorem. In order to understand how our construction works, it is helpful to first understand the elegant idea of [NS25]. Nöbel and Steiner introduce so-called *tower-gadgets* (see Figure 1), and furthermore combine multiple towers into a gadget which for our purpose shall be called *city gadget* (see Figure 2). They show that there exist perfect matchings M_1 and M_2 such that any short flip sequence from M_1 to M_2 must contain at least one cycle that visits every city gadget. This is sufficient for their argument, but we require a stronger statement. Let us call a cycle C in the graph G_H *regular*, if C visits every city gadget.

We first show, that in a short enough flip sequence (C_1, \dots, C_d) , if n tends to infinity, then an arbitrarily large percentage of the cycles (C_1, \dots, C_d) must be regular. We remark that it is inherent to the arguments used in [NS25] that in general there can always be several cycles that are irregular, i.e. they do not visit every city gadget. Such irregular cycles are technically challenging for us to deal with, but their existence can not be excluded.

We significantly expand upon the ideas of [NS25] by introducing so-called *XOR-gadgets* (Figure 4), *ladder-gadgets* (Figure 7) and sophisticated \forall -gadgets (Figure 10). These gadgets have the property that if they interact with some regular cycle C , then their behavior is well understood. On the other hand, if they interact with some irregular cycle, then the “damage” done by this irregular cycle is in some sense restricted.

It is described in Section 4.5 how to define the desired bipartite undirected graph G_H given the directed graph H . The graph G_H contains many copies of the gadgets described above. Given two perfect matchings M_1, M_2 of G_H , one can ask what is their distance $\text{dist}(M_1, M_2)$ in the polytope P_{G_H} . We are able to show, that in a certain sense, this distance crucially depends on the structure of the matchings M_1, M_2 restricted to the \forall -gadgets. In particular, consider a fixed pattern $P \subseteq E'$ in H . We show that one can find PMs M_1, M_2 in G_H by carefully choosing their restriction to the \forall -gadgets, and completing this choice to the whole graph G_H in the right manner, such that $\text{dist}(M_1, M_2)$ is small if and only if there is a Hamiltonian cycle respecting the pattern P in the directed graph H . (This argument depends crucially on the asymmetric behavior of the set of perfect matchings of the ladder gadget, explained more closely in Section 4.4.) Note that this approach highlights the similarity between Problem (1) and $\forall\exists$ -HAMCYCLE, i.e. perfect matchings of G_H correspond to patterns $P \subseteq E'$ and vice versa. However, note that there is a major difference: Patterns are binary, that is for all $i \in [k]$ the set P either contains e_i or \bar{e}_i . In contrast, the set of perfect matchings of a \forall -gadget is not binary at all, rather it is exponentially large. Nonetheless, we are able to overcome this final technical difficulty. On a high level, we show that we can interpret a PM of a \forall -gadget as a “mixture” of so-called top-states and bottom-states, and make use of the property that *for all* patterns $P \subseteq E'$ the directed graph H has a Hamiltonian cycle respecting P .

We explain one more idea, which is also used in [NS25]. We will define a subset $V_s \subseteq V(G_H)$ of size $|V_s| = \Theta(n)$ of the $\Theta(n^9)$ vertices of G_H , called the *semi-default vertices*. We call a PM M of the graph G_H a *semi-default matching*, if M matches the semi-default vertices in a pre-described way. We show that the semi-default matchings lie “dense” in all perfect matchings. In particular, given an arbitrary PM M , one can find a semi-default PM M' such that $\text{dist}(M, M') \leq 23n \ll n^4$. The intuition we will use is that in order to transform any PM M_1 into any PM M_2 , we first look for semi-default representatives M'_1, M'_2 of M_1, M_2 . Then, we use at most $23n$ irregular cycles to transform M_1 to M'_1 , at most $4n^4$ regular cycles to transform M'_1 into M'_2 , and at most $23n$ irregular cycles to transform M'_2 into M_2 . This completes the proof sketch of the first main result. More details are given in Section 4.

Overview of the second main result. We wish to prove that the diameter is hard to approximate to a factor $(1 + \varepsilon)$. For this, we consider a similar reduction as for the first main result. We again make central use of the city gadgets introduced by [NS25] to “enforce” that most cycles in a short flip sequence (C_1, \dots, C_d) visit all cities. This is reminiscent of the Hamiltonian cycle problem. It is therefore natural to show a certain approximation version of the Hamiltonian cycle problem. It roughly states that we can w.l.o.g. consider problem instances such that in yes-instances it is possible to visit all cities, but in no-instances it is only possible to visit a $(1 - \varepsilon)$ fraction of all cities. We prove a result of this kind suited to our needs in Theorem 22. We achieve this as a consequence of the PCP theorem using established techniques. If we have a yes-instance, then the diameter of P_G is small, since we can efficiently visit all city gadgets. However, if it is only possible to visit a $(1 - \varepsilon)$ fraction of all cities, then the diameter of P_G must be large. This is because city gadgets (in the right configuration) have the property that each city must be visited many times by the cycles in (C_1, \dots, C_d) . This

completes the proof sketch of the second main result. More details are given in Section 5.

3 Preliminaries

In this paper, we use the notation $\mathbb{N} = \{1, 2, \dots\}$ and $[n] := \{1, \dots, n\}$. We use uv as a shorthand notation for the undirected edge $\{u, v\}$. All paths and cycles are simple. All graphs are assumed to be undirected, unless stated otherwise. Paths and cycles in a directed graph always adhere to the orientation of the edges. For $G = (V, E)$, we denote $V(G) := V$ and $E(G) := E$.

A *language* is a set $L \subseteq \{0, 1\}^*$. A language $L \subseteq \{0, 1\}^*$ is contained in Π_2^P if there exists some polynomial-time computable function V (verifier), such that for all $w \in \{0, 1\}^*$ for suitable $m_1, m_2 \leq \text{poly}(|w|)$

$$w \in L \Leftrightarrow \forall y_1 \in \{0, 1\}^{m_1} \exists y_2 \in \{0, 1\}^{m_2} : V(w, y_1, y_2) = 1.$$

A language L is Π_2^P -hard, if every $L' \in \Pi_2^P$ can be reduced to L with a polynomial-time many-one reduction. If L is both Π_2^P -hard and contained in Π_2^P , it is Π_2^P -complete.

An introduction to the polynomial hierarchy and the classes Π_2^P and $\Sigma_2^P = \text{co-}\Pi_2^P$ can be found in the book by Papadimitriou [Pap94], or the article by Woeginger [Woe21].

Let $X = \{x_1, \dots, x_n\}$ be a set of boolean variables. The corresponding set of *literals* is $\{x_1, \dots, x_n\} \cup \{\bar{x}_1, \dots, \bar{x}_n\}$. A *clause* is a disjunction of literals. A boolean formula φ is in conjunctive normal form (CNF) if it is a conjunction of clauses. An *assignment* is a map $\alpha : X \rightarrow \{0, 1\}$. We write $\varphi(\alpha) \in \{0, 1\}$ to denote the evaluation of φ under assignment α .

4 The Reduction

In this section, we present our proof of Π_2^P -completeness. We first show containment in the class Π_2^P in Section 4.1. Then we introduce the Π_2^P -complete problem $\forall\exists\text{-HAMCYCLE}$ in Section 4.2. We introduce a number of different gadgets in Sections 4.3 and 4.4, with the \forall -gadget being the most complicated one. In Section 4.5 we define the graph G_H , and in Section 4.6, we show that the diameter of the polytope P_{G_H} has the desired properties.

4.1 Containment

It is relatively easy to show the containment of DIAMETER-DECISION in the class Π_2^P , as long as the following assumption holds.

Lemma 7. *Let \mathcal{F} be a family of polytopes. If it holds that for each $P \in \mathcal{F}$ we have $\text{diam}(P) \leq \text{poly}(n, d, |P|)$, where n is its number of facets, d its dimension, and $|P|$ its encoding length, then DIAMETER-DECISION restricted to the family \mathcal{F} is contained in Π_2^P . An analogous statement holds for the circuit diameter $\text{cdiam}(P)$ and CDIAM-DECISION.*

Proof. Let $P = \{x \in \mathbb{R}^d : Ax \leq b\}$, and consider some $t \in \mathbb{N}$. We have $\text{diam}(P) \leq t$ if and only if

$\forall x, y$ vertices of $P : \exists (v_1, \dots, v_k) : x = v_1, y = v_k, k \leq t$, all v_i and v_{i+1} are neighboring vertices.

Furthermore, given a sequence (v_1, \dots, v_k) of vectors in \mathbb{R}^d , it is possible to test in polynomial time, whether v_1, \dots, v_k are actually vertices of the polytope and whether v_i is adjacent to v_{i+1} for all $i \in [k-1]$ (see e.g. [FT94]). In particular, we can w.l.o.g. restrict such sequences to length $k \leq \text{diam}(P) \leq \text{poly}(n, d, |P|)$. Therefore, the encoding space of the sequence (v_1, \dots, v_k) is at most polynomial in the input size. Thus by the definition of Π_2^P , we have DIAMETER-DECISION $\in \Pi_2^P$. For the circuit diameter, the same argument applies. We use the fact that given two points $v_i, v_{i+1} \in P$, one can test in polynomial time whether v_i, v_{i+1} are one circuit move apart [BGK⁺25]. \square

In particular, we have BPM-DIAMETER-DECISION $\in \Pi_2^p$, and CDIAM-DECISION $\in \Pi_2^p$.

4.2 The Base Problem

Consider problem $\forall\exists$ -HAMCYCLE, as defined in Section 2.

Lemma 8. $\forall\exists$ -HAMCYCLE is Π_2^p -complete (even when restricted to planar directed graphs of maximum in- and outdegree 2).

Proof. The problem is contained in Π_2^p , since it can be stated as:

$$\forall \text{ patterns } P \subseteq E' : \exists \text{ a Hamiltonian cycle } C \text{ respecting } P,$$

and given a directed graph H together with sets $P \subseteq E'$, $C \subseteq E(H)$, it can be checked in polynomial time whether C is a Hamiltonian cycle and respects the pattern P . So it remains to show Π_2^p -hardness. Our proof strategy is to first consider a classic reduction by Plesník [Ple79], which reduces 3SAT to the directed Hamiltonian cycle problem in planar digraphs of maximum in- and outdegree 2. We show that we can modify Plesník's reduction to be instead a reduction from $\forall\exists$ -SAT to $\forall\exists$ -HAMCYCLE. It is not necessary to re-iterate all the details of Plesník's reduction. Instead, it suffices to notice the following key properties: Given a 3SAT formula φ , the reduction constructs in polynomial time a (planar and 2-in/outdegree-bounded) directed graph H_φ . If φ has N variables and M clauses, the graph H_φ contains N so-called variable gadgets and M so-called clause gadgets. Plesník proceeds to prove that every Hamiltonian cycle must traverse the variable gadgets in order, and must make a “decision” for each variable gadget. In particular, the i -th variable gadget starts with some vertex $v^{(i)}$ of outdegree 2. Immediately after entering $v^{(i)}$, every Hamiltonian cycle needs to decide whether it leaves $v^{(i)}$ via the first outgoing arc and traverses the variable gadget in the “ $x_i = \text{true}$ ” configuration, or if it leaves $v^{(i)}$ via the second outgoing arc and traverses the variable gadget in the “ $x_i = \text{false}$ ” configuration. Let us define e_i as the first, and \bar{e}_i as the second outgoing arc of vertex $v^{(i)}$, and let $E'' = \{e_1, \bar{e}_1, \dots, e_N, \bar{e}_N\}$. By studying [Ple79] it can be checked that the following holds:

- For every $i \in \{1, \dots, N\}$, every Hamiltonian cycle C in H_φ uses exactly one of the two arcs e_i and \bar{e}_i .
- There is a direct correspondence between subsets of the arc set E'' that can be part of a Hamiltonian cycle and satisfying assignments for the formula φ . Formally: Let $Z = \{z_1, \dots, z_N\}$ be the variable set of φ . For an assignment $\alpha : Z \rightarrow \{0, 1\}$, let P_α denote the corresponding set of arcs defined by $P_\alpha = \{e_i : \alpha(z_i) = 1\} \cup \{\bar{e}_i : \alpha(z_i) = 0\}$. Given an assignment α , there exists a Hamiltonian cycle C in H_φ with $C \cap E'' = P_\alpha$ if and only if $\varphi(\alpha) = 1$.

Finally, we describe how to modify this reduction to a reduction from $\forall\exists$ -SAT to $\forall\exists$ -HAMCYCLE. Given a 3SAT formula $\varphi(X, Y)$, we assume w.l.o.g. that φ is a formula over $N = 2n$ variables partitioned into two parts X, Y of equal size $|X| = |Y| = n$. We consider the graph H_φ as above. Then, we consider the edges $E'' = \{e_1, \bar{e}_1, \dots, e_{2n}, \bar{e}_{2n}\}$, and let $E' = \{e_1, \bar{e}_1, \dots, e_n, \bar{e}_n\}$ be those arcs of E'' that appear in the variable gadgets corresponding to X (but not to Y). By the above observation, for a fixed pattern $P \subseteq E'$, there is a Hamiltonian cycle respecting P , if and only if it is possible to complete P with another pattern $P_2 \subseteq E'' \setminus E'$ and complete $P \cup P_2$ to a Hamiltonian cycle. Again, by the above correspondence, this implies that φ is a yes-instance of $\forall\exists$ -SAT if and only if (H_φ, E') is a yes-instance of $\forall\exists$ -HAMCYCLE. \square

We remark that the key properties described above are not specific to Plesník's proof. Similar properties appear in other folklore proofs for the NP-hardness of Hamiltonian cycle, and in plenty of NP-hardness proofs of other combinatorial problems. In [GW24] a systematic study of this property is performed, and its implications on completeness in the second level of the polynomial hierarchy are discussed.




Figure 1: A tower gadget of height h , and different states a perfect matching of the tower gadget may have.

4.3 Towers and Cities

Nöbel and Steiner [NS25] introduce so-called *tower gadgets*. For some $h \in \mathbb{N}$, a tower gadget of height h is an induced subgraph on the vertex set

$$\{v, w\} \cup \bigcup_{i=0}^h \{a_i, b_i\}$$

and edge set

$$\{va_0, b_0w\} \cup \bigcup_{i=0}^h \{a_i b_i\} \cup \bigcup_{i=1}^h \{a_i a_{i-1}, b_i b_{i-1}\}.$$

An example for $h = 6$ is depicted in Figure 1a. Recall that our plan is to use these tower gadgets as part of some larger graph G_H . We will maintain the invariant that every tower gadget is connected to the rest of G_H only via the two vertices v and w . If some tower gadget T is contained in G_H , and M is a PM of G_H , the PM M restricted to T can behave in many different ways. We give names to some of these behaviors. Consider Figures 1b to 1d. With respect to a fixed PM M , the tower gadget T is called in *default* state, if $\{va_0, b_0w\} \cup \bigcup_{i=1}^h \{a_i b_i\} \subseteq M$. The tower gadget T is called *locked*, if it is in default state, except for the 4 top vertices, where instead $a_h a_{h-1}, b_h b_{h-1} \in M$. The tower gadget T is called in *semi-default* state, if $va_0, b_0w \in M$ (and the rest of the tower can be matched in any arbitrary way).

Consider a tower gadget T , subgraph of G_H , which starts in the locked state, and which we want to transform into the default state using a flip sequence of alternating cycles (C_1, \dots, C_d) , where $C_i \subseteq G_H$ for $i \in [d]$. Of course, we can achieve this task by using a single alternating cycle C on the vertices $\{a_h, a_{h-1}, b_{h-1}, b_h\}$. However, if we choose to do this, it comes with a disadvantage: Even though we have achieved our goal using a single cycle C , the influence of the alternating cycle $C \subseteq G_H$ is limited to only T (but C could be useful also somewhere else). This motivates the following definition: Let T be a tower gadget in the graph G_H and M be a fixed PM of G_H . A cycle $C \subseteq G_H$ is called *well-behaved*¹ for T , if restricted to T it is a v - w -path. A sequence of paths (P_1, \dots, P_d) is a well-behaved flip sequence for T , if for all $i = 1, \dots, d$ the path P_i is a path from v to w inside T such that P_i is alternating with respect to the matching $M \triangle P_1 \triangle \dots \triangle P_{i-1}$.

Lemma 9 (adapted from [NS25]). *Let T be a tower gadget of height h for some $h \in \mathbb{N}$. If (P_1, \dots, P_d) is a well-behaved flip sequence for T which transforms T from the locked state into the default state, then $d \geq 2h - 2$.*

¹In [NS25] this behavior is called “touching the tower”. However, we would like to use the same name also for our more advanced gadgets later on, where the name “touching” could be misinterpreted.




Figure 2: A city gadget consists out of many tower gadgets.




Figure 3: A city gadget is called *matched*, or in *semi-default state*, if all of its towers are in semi-default state.

For the sake of completeness, we provide the proof sketch of Lemma 9. Let M_i be the PM after applying the first i paths P_1, \dots, P_i for $i = 0, \dots, d$. Let $\mathcal{H}(M_i) := \{j \in [h] : a_j b_j \in M_i\}$ be the set of indices corresponding to the horizontal edges of M_i . Because the flip sequence is well-behaved, every path can switch between the left and the right side of the tower only a single time. This means that $\mathcal{H}(M_{i+1})$ results from $\mathcal{H}(M_i)$ by adding or deleting only a single item. Furthermore, because every path C_i is also alternating, the first time P_i encounters some horizontal edge $a_j b_j$ of M_{i-1} , the path P_i must include it. Hence, in order to go from the locked to the default position, one must first remove the $h - 2$ horizontal edges of $\mathcal{H}(M_0)$, and then add all h horizontal edges. Since each requires a single path, we have $d \geq 2h - 2$.

Lemma 10 (adapted from [NS25]). *Let T be a tower gadget of height h , and let M_1, M_2 be two PMs of T in semi-default position. Then there exists a well-behaved flip sequence (P_1, \dots, P_d) which transforms M_1 into M_2 such that $d = 2h$.*

Here the intuition is that we can first remove all the at most h horizontal edges of M_1 , and then insert all the at most h horizontal edges of M_2 . It can be checked that this is possible with a well-behaved flip sequence $(P_1, \dots, P_{d'})$ for some $d' \leq 2h$, where in particular all paths are alternating with respect to the current matching. Since both M_1 and M_2 are in semi-default state, d' is an even number. Finally, we can assume w.l.o.g. that $d' = 2h$. Indeed, if $d' < 2h$, we can just choose any alternating path P from v to w (which always exists) and consider the flip sequence $(P_1, \dots, P_{d'}, P, P)$. Since flipping the same path twice does have no effect on the matching, this new flip sequence has the same end result.

We now combine multiple tower gadgets into so-called *city gadgets*. These were also used in [NS25], but not given an explicit name. A city gadget of width t and height h between vertices x and y is defined as follows: It consists out of t towers T_1, \dots, T_t , such that vertex x is identified with vertex v of the first tower T_1 , vertex y is identified with vertex w of the last tower T_t , and for $i = 2, \dots, t$, the edge $b_0 w$ of tower T_{i-1} is identified with edge va_0 of the next tower T_i . An example is depicted in Figure 2. Note that x and y are connected by a path of odd length. A city gadget is called in *semi-default state* or *matched*, if all of its towers are in semi-default state. Since every tower has an even number of vertices, a city gadget is matched if and only if the edge $xa_0^{(T_1)}$ of the first tower is in the matching. Similarly to before, we will maintain the

invariant, that a city gadget is connected to the rest of G_H only via the vertices x and y . City gadgets are useful since they “enforce” that certain locations must be visited. Let us say that some cycle C in the graph G_H *visits a city gadget g* , if the cycle C contains the first edge $xa_0^{(T_1)}$ of the city gadget g . Note that a cycle C visits a city if and only if C is well-behaved for every tower of the city.

Definition 11. Let G_H be a graph containing several city gadgets. A (not necessarily alternating) cycle C in G_H is called *regular*, if it visits every single city gadget of G_H .

Lemma 12. Let $n \in \mathbb{N}$. Let G_H be a graph which contains at most $17n$ city gadgets, with every city gadget having the same height $h = 2n^4$ and width $t = 4n^4 + 100n$. Let M_1 be a PM of G_H such that w.r.t. M_1 every tower of every city is in locked state, and M_2 be a PM of G_H such that every tower of every city is in default state. If (C_1, \dots, C_d) is a flip sequence from M_1 to M_2 with $d \leq 4n^4 + 46n$, then we also have $d \geq 4n^4 - 2$ and furthermore at most $1000n^2$ of the d cycles are not regular.

Proof. Consider a fixed city gadget g . We claim that the gadget g contains a tower T^* such that each of C_1, \dots, C_d either does not visit g at all or is well-behaved for T^* . Indeed, there can be at most d towers in g with the property that one of the cycles (C_1, \dots, C_d) is entirely contained in them. Since $t > d$, at least one tower T_j has the property that none of the cycles (C_1, \dots, C_d) is contained entirely in T_j . We let $T^* := T_j$. Now, observe that T^* starts in locked state, ends up in default state, and only interacts with well-behaved cycles. Due to Lemma 9, at least $2h - 2$ well-behaved cycles are required for this task. Hence at least $2h - 2$ cycles visit the city g . In particular $d \geq 4n^4 - 2$. If we define

$$\mathcal{C}_g := \{i \in [d] : C_i \text{ does not visit } g\},$$

then $|\mathcal{C}_g| \leq d - (2h - 2) \leq 46n + 2$. Furthermore if a cycle is not regular, then it is contained in \mathcal{C}_g for at least one city g , and so

$$|\{i \in [d] : C_i \text{ not regular}\}| \leq \sum_g |\mathcal{C}_g| \leq 17n(48n) \leq 1000n^2.$$

Here the index g runs over all cities of the graph. □

The XOR gadget. In the final construction of the graph G_H , we will choose the parameters such that Lemma 12 is applicable. The lemma then tells us that of the $\Theta(n^4)$ cycles in any short flip sequence, at most $O(n^2)$ are not regular, i.e. the majority of cycles visit all cities. Based on this insight, we introduce our next gadget, the so-called *XOR-gadget*. Given a graph G , and two edges $ab, uv \in E(G)$, a XOR gadget between ab and uv is created by subdividing the edge ab four times, creating four new vertices x_1, \dots, x_4 , and subdividing the edge uv four times, creating four new vertices y_1, \dots, y_4 . After that, we connect x_i and y_i with a city gadget for $i = 1, \dots, 4$. An example is shown in Figure 4. Note that if C is a regular cycle, then C traverses the XOR gadget either from u to v , but does not traverse the gadget from a to b , or traverses the gadget from a to b , but does not traverse the gadget from u to v . (This can be formally proven by noticing that every regular cycle uses either the edge x_1x_2 or the edge x_2x_3 .) In order to simplify future notation, let us say that in the first case the cycle C uses the edge uv , but does not use the edge ab , while in the second case C uses the edge uv , but does not use the edge ab (these formulations are slightly imprecise since technically the edges ab and uv do not exist in G_H). In conclusion, every regular cycle has to visit the XOR gadget and uses exactly one of the two edges ab and uv .

A XOR-gadget is called in *semi-default state* if every of its four city gadgets is in semi-regular state (see Figure 5). Observe that two XOR-gadgets can be applied to the same edge (see Figure 6).

Finally, we require the following technical lemma:




Figure 4: Definition of a XOR gadget. Note that if a cycle visits every city, then in every XOR gadget it “traverses” exactly one of the two edges ab and uv .




Figure 5: A XOR gadget is called in semi-default state, if every of its city gadgets is in semi-default state.




Figure 6: Multiple XOR gadgets can be applied to the same edge.




Figure 7: A ladder gadget. Ladder gadgets always have a fixed height of 5.

Lemma 13. *If G is a bipartite graph and e, e' are distinct edges, then one can add a XOR-gadget between e and e' such that the graph remains bipartite.*

Proof. Since every city gadget connects its two endpoints via an odd-length path, the XOR gadget itself is bipartite, such that vertex set $\{a, v\}$ and vertex set $\{u, b\}$ lie in different parts of the bipartition. Let vertices w_1, w_2 be the endpoints of edge e and vertices w'_1, w'_2 be the endpoints of edge e' in the graph G . Since G is bipartite, w_1 and w_2 lie in different parts, and w'_1 and w'_2 lie in different parts (w.r.t. the bipartition of G). We do a case distinction: If the vertex sets $\{w_1, w'_2\}$ and $\{w_2, w'_1\}$ lie in different parts of the bipartition of G , then we add the XOR-gadget in such a way that (w_1, w_2) is identified with (a, b) and (w'_1, w'_2) is identified with (u, v) . The resulting graph is bipartite, since a bipartite graph is inserted into another bipartite graph respecting the bipartition. In the other case, if the vertex sets $\{w_1, w'_1\}$ and $\{w_2, w'_2\}$ lie in different parts of the bipartition of G , then we add the XOR-gadget in such a way that (w_1, w_2) is still identified with (a, b) , but (w'_1, w'_2) is identified with (v, u) . This means that we effectively mirror the bottom edge of the XOR-gadget. The resulting graph is bipartite. \square

4.4 Ladders and the \forall -gadget

As explained in Section 2, the \forall -gadget needs to exhibit asymmetric behavior. In order to achieve this asymmetric behavior, we introduce yet another gadget, the *ladder gadget*. We show that the ladder gadget exhibits a very slight asymmetric behavior, and amplify this effect by combining many ladder gadgets together. A ladder gadget is an induced subgraph on the 14 vertices

$$\{a_0, \dots, a_6\} \cup \{b_0, \dots, b_6\}$$

and edge set

$$\bigcup_{i=1}^5 \{a_i b_i\} \cup \bigcup_{i=0}^5 \{a_i a_{i+1}, b_i b_{i+1}\}.$$

An example is depicted in Figure 7a. Ladder gadgets are similar to tower gadgets, but have a connection to both their top and bottom side. While tower gadgets have a variable height, a ladder gadget in this paper always has a fixed height of 5. We will maintain the invariant that a ladder is connected to the rest of the graph G_H only via the vertices a_0, b_0, a_6, b_6 . Consider a fixed PM M of G_H . With respect to M a ladder gadget is called in *semi-default state*, if the vertices a_0, b_0, a_6, b_6 are matched to vertices outside the gadget by M . It is in *default state*, if $\{a_1 b_1, \dots, a_5 b_5\} \subseteq M$. It is called the *bottom-open ladder*, if $\{a_5 b_5, a_4 a_3, b_4 b_3, a_2 a_1, b_2 b_1\} \subseteq M$. It is called the *top-open ladder*, if $\{a_5 a_4, b_5 b_4, a_3 a_2, b_3 b_2, a_1 b_1\} \subseteq M$ (see Figures 7b to 7e).

Similar to the case of tower gadgets, we again want to consider well-behaved cycles. Let L be some ladder gadget. Assume we have some (not necessarily alternating) cycle C in the graph




Figure 8: Ladder L_1 is transformed into ladder L_3 with two well-behaved alternating cycles from the top. Ladders L_1 and L_3 are in semi-default state, ladder L_2 is not.

G_H , such that C includes at least one vertex of L . The cycle C is called *well-behaved* for the ladder gadget L , if either

- both $a_0a_1, b_0b_1 \in E(C)$ and both $a_5a_6, b_5b_6 \notin E(C)$, or
- both $a_5a_6, b_5b_6 \in E(C)$ and both $a_0a_1, b_0b_1 \notin E(C)$.

In other words, the cycle C enters the ladder either from the top (i.e. via the edge a_5a_6 or b_5b_6) or from the bottom (i.e. via the edge a_0a_1 or b_0b_1), leaves from the the same side it entered, and after that does not visit the inner ladder (vertices a_i, b_i for $i \in \{1, \dots, 5\}$) again. In the first case, let us say that the well-behaved cycle C visits L *from the top*, in the second case, let us say that C visits L *from the bottom*. The next lemma establishes the already claimed asymmetry, by showing that in order to transform the bottom-open ladder (the top-open ladder, respectively) into the default ladder with well-behaved cycles, we need to primarily come from the bottom (the top, respectively). Similarly to before, a sequence (C_1, \dots, C_d) of cycles is called a *well-behaved flip sequence* for the ladder L that transforms M_1 into M_2 , if each of C_1, \dots, C_d is well-behaved, and restricted to L , we have that C_i is an alternating path with respect to $M \triangle C_1 \triangle \dots \triangle C_{i-1}$ for all $i \in [d]$, and $M_2 = M_1 \triangle C_1 \triangle \dots \triangle C_d$. Figure 8 showcases a well-behaved flip sequence of length 2.

Lemma 14. *Let L be a ladder gadget. If (C_1, \dots, C_d) is a well-behaved flip sequence for L that transforms the bottom-open ladder into the default state, then $d \geq 4$. Additionally if $d = 4$, then each of C_1, \dots, C_4 must come from the bottom. The analogous statement is true if L starts as the top-open ladder, except that now each of C_1, \dots, C_4 must come from the top.*

Proof. Observe that every well-behaved cycle C includes exactly one edge of the form a_ib_i , where $i \in [5]$. (This follows in particular since a well-behaved cycle visits a ladder only once, either from the top or from the bottom.) Let $\mathcal{H}(M) := \{i \in [5] : a_ib_i \in M\}$ be the index set of horizontal edges that some PM M contains. The previous fact implies that after flipping a well-behaved cycle, the set $\mathcal{H}(M)$ changes by only addition or deletion of a single element. Since for the default ladder $\mathcal{H}(M) = \{1, \dots, 5\}$, and for the bottom-open ladder $\mathcal{H}(M') = \{5\}$, we see that if (C_1, \dots, C_d) is a flip sequence transforming the former into the latter, then $d \geq 4$. Further, if $d = 4$, assume one of the cycles C_j for $j \in [4]$ comes from the top. Then either the cycle C_j or some cycle $C_{j'}$ for $j' < j$ contains the edge a_5b_5 as the only horizontal edge. This is a contradiction to the fact that we go from $\mathcal{H}(M) = \{1, \dots, 5\}$ to $\mathcal{H}(M') = \{5\}$ in only four steps. Finally, the statement about the top-open ladder holds analogously by vertical symmetry. \square

We remark that in the above lemma it is crucial that the cycles are well-behaved. In particular, if a cycle is allowed to enter a ladder from the top, but leave from the bottom, then one can show that it is possible to transform the bottom-open ladder into the default ladder with only two alternating cycles. Our construction of G_H will ensure that in a short enough flip sequence, the majority of all ladders only ever interact with well-behaved cycles.




Figure 9: All 8 semi-default ladders. Arrows indicate that one can be turned into the other by using two well-behaved cycles, either both from the top ($2t$) or both from the bottom ($2b$). In order to transform the top-open ladder into the default ladder, one requires at least 4 well-behaved cycles, all from the top.

The necessary condition of Lemma 14 for transforming a ladder into another is complemented by the following sufficient condition.

Lemma 15. *Let L be a ladder gadget. Let M_1 and M_2 be two semi-default states of L . There is a well-behaved flip sequence (C_1, \dots, C_4) of length 4 that transforms M_1 into M_2 . Furthermore there is one such sequence so that C_1, C_2 both come from the top, or both come from the bottom, and C_3, C_4 both come from the top, or both come from the bottom.*

Proof. Observe that there are 8 different semi-default ladders, since a semi-default ladder has either 1, 3 or 5 horizontal edges, and all possibilities to arrange these are displayed in Figure 9. Furthermore, Figure 9 shows a graph, where two ladders are connected with an edge, if one can be transformed into the other using two well-behaved cycles, either both from the top (denoted $2t$), or both from the bottom (denoted $2b$). Note that this relation is symmetric, i.e. if L_1 can be transformed into L_2 with two well-behaved cycles, then L_2 can also be transformed into L_1 . To prove the lemma it now suffices to observe that the graph of Figure 9 has diameter 2, i.e. every pair of ladders is connected by a path with at most two edges. Finally, we remark that if (C_1, \dots, C_d) with $d = 2$ or $d = 0$ is an even shorter flip sequence from M_1 to M_2 , we can also find a flip sequence of length exactly four: We can simply choose any well-behaved cycle C from the top or bottom (at least one always exists), and consider the sequence (C_1, \dots, C_d, C, C) . \square

\forall -gadgets. After we showed that a single ladder gadget can exhibit a slight asymmetry between top and bottom, we now show how to amplify this asymmetry by combining a lot of ladders into a single gadget. Assume we are given an undirected graph G_H , and three distinct vertices called $v_{\text{out}}, u_{\text{in}}, w_{\text{in}}$. Let $t \in \mathbb{N}$. A \forall -gadget of width t is the subgraph that is depicted in Figure 10 such that it is connected to the rest of G_H via only these three vertices. Precisely, we introduce 10 new vertices $\{x_1, \dots, x_{10}\}$. After that, we connect the four pairs $(x_1, x_2), (x_3, x_4), (x_5, x_6), (x_7, x_8)$ with a city gadget each. We add the edge set

$$\{v_{\text{out}}x_1, v_{\text{out}}x_5, x_1x_8, x_5x_4, x_2x_3, x_6x_7, x_4x_9, x_8x_9, x_9x_{10}, x_{10}u_{\text{in}}, x_{10}w_{\text{in}}\}.$$




Figure 10: A ∇ -gadget between the three vertices v_{out} , u_{in} , and w_{in} .





Figure 11: A ∇ -gadget is in semi-default state, if all its city gadgets and all its XOR-gadgets are in semi-default state and the edge x_9x_{10} is matched.

Three XOR-gadgets are applied to the edge pairs (x_2x_3, x_6x_7) , $(x_2x_3, x_{10}u_{\text{in}})$, and $(x_6x_7, x_{10}w_{\text{in}})$ respectively. Finally, a total amount of t ladder gadgets are added. For each of these t ladders, vertex a_6 in the ladder is identified with vertex x_2 in the ∇ -gadget. Likewise, b_6 is identified with x_3 , a_0 is identified with x_6 , and b_0 is identified with x_7 . This completes the description of the ∇ -gadget.


A ∇ -gadget is called in *semi-default state* with respect to a fixed PM M , if all its city-gadgets, and all its XOR-gadgets are in semi-default state, and $x_9x_{10} \in M$ (see Figure 11). We now describe the main functionality of a ∇ -gadget. Recall that a regular cycle is a (not necessarily alternating) cycle in G_H that visits all cities. Let A be a ∇ -gadget and C be a regular cycle. We say that the cycle C is in *top state* with respect to the gadget A , if C uses the edge $x_{10}u_{\text{in}}$. Analogously, we say that the cycle C is in *bottom state* with respect to the gadget A , if C uses the edge $x_{10}w_{\text{in}}$. Let us for the remainder of this section assume w.l.o.g. that the graph G_H contains at least one city gadget outside of A , such that every regular cycle needs to both enter and leave A . Let us say that C *visits* some ladder, if $V(C) \cap \left(\bigcup_{i=1}^5 \{a_i, b_i\}\right) \neq \emptyset$ for the vertices $a_0, \dots, a_6, b_0, \dots, b_6$ of that ladder.

Lemma 16. *If A is a ∇ -gadget, and C is a regular cycle in G_H , then C is either in top state or bottom state (but not both). Furthermore, the cycle C visits exactly one ladder L of the gadget, and C is well-behaved for L . If the cycle C is in top state (bottom state, respectively) then it visits L from the top (from the bottom, respectively).*

Proof. Since C is regular, the city-gadgets and XOR-gadgets behave as intended. In particular, the cycle C uses all four edges $x_1x_2, x_3x_4, x_5x_6, x_7x_8$, and exactly one of the two edges $x_{10}u_{\text{in}}$ and $x_{10}w_{\text{in}}$ (because they are connected by a chain of XOR-gadgets). Since we assumed that there is at least one city gadget outside of the ∇ -gadget A , and A is only connected to the rest



(a) Top state



(b) Bottom state

Figure 12: If C is a cycle visiting every city gadget, then a \forall -gadget is in one of the following two states: Either C contains the edge $x_{10}u_{\text{in}}$ and visits exactly one ladder coming from the top (*top state*), or C contains the edge $x_{10}w_{\text{in}}$ and visits exactly one ladder coming from the bottom (*bottom state*).

of the graph via the three vertices $v_{\text{out}}, u_{\text{in}}, w_{\text{in}}$, the cycle C enters the gadget in vertex v_{out} and leaves via either vertex u_{in} or w_{in} . We now distinguish between the following cases.

Case 1: C uses $x_{10}u_{\text{in}}$. In this case C also uses x_6x_7 , but does not use the two edges $x_2x_3, x_{10}u_{\text{in}}$. Since C enters the gadget at v_{out} , it uses exactly one of the two edges $v_{\text{out}}x_1$ and $v_{\text{out}}x_5$.

Case 1a: C uses $v_{\text{out}}x_1$. In this case, consider vertex x_5 . Since it has degree 3, and we already know that C uses x_5x_6 , but not $v_{\text{out}}x_5$, we deduce that C uses x_4x_5 . Then, combining all the information we have so far, we see that the cycle C looks like in Figure 12a. Precisely, C enters at v_{out} , goes to x_1 , traverses from x_1 to x_4 by entering a ladder gadget, goes to x_5 , and then visits vertices $x_6 - x_{10}$ in that order before leaving via u_{in} . In particular, the cycle C visits exactly one ladder. This is because in order to enter or exit a ladder, the cycle needs to visit one of the four vertices x_2, x_3, x_6 or x_7 . However, x_6 and x_7 are blocked. Therefore x_2 needs to be used to enter a ladder, and x_3 needs to be used to exit the same ladder, and no different ladder can be entered after that. Hence C visits exactly one ladder, coming from the top.

Case 1b: C uses $v_{\text{out}}x_5$. In that case, the argument is analogous to case 1a, with the exception that C uses the edge x_1x_8 . Still, it remains that x_6, x_7 are blocked, so C visits exactly one ladder coming from the top. To summarize, in both cases 1a and 1b, the cycle visits a ladder from the top and leaves the gadget via the edge $x_{10}u_{\text{in}}$.

Case 2: C uses $x_{10}w_{\text{in}}$. Then, due to vertical symmetry, the cycle visits exactly one ladder from the bottom, and leaves via the edge $x_{10}w_{\text{in}}$. \square

The above lemma shows that every regular cycle can only interact with one ladder in a well-behaved manner. But what about the irregular cycles? Since irregular cycles do not need to visit every city, they can potentially interact with our gadgets in a much less controlled way. Still, the “damage” done by an irregular cycle is restricted, as the following lemma shows.

Lemma 17. *If A is a \forall -gadget, and C is some cycle in G_H , then C visits at most 4 ladders inside A .*

Proof. Observe that in the graph $G_H - \{x_2, x_3, x_6, x_7\}$, each of the t ladders of A is its own connected component. Whenever the cycle C visits a ladder, it needs to enter and exit using one of these four vertices. Hence every ladder that C visits is associated to two vertices of x_2, x_3, x_6, x_7 . (We assume that C enters and exists the ladder, because if C is entirely contained in it, we are done). On the other hand, each of these four vertices is associated to at most two ladders. Hence C visits at most 4 ladders inside A . \square

4.5 Definition of G_H

We are now ready to describe the main construction. Assume we are given an instance of $\forall\exists$ -HAMCYCLE. This instance consists out of a directed graph H together with some $k \in \mathbb{N}$ and vertices $v^{(1)}, \dots, v^{(k)}$ in H , each having outdegree 2. Let $n := |V(H)|$. For $i \in [k]$, let $u^{(i)}$ and $w^{(i)}$ denote the two outneighbors of vertex $v^{(i)}$. Define the two arcs $e_i := (v^{(i)}, u^{(i)})$ and $\bar{e}_i := (v^{(i)}, w^{(i)})$, as well as the set $E' := \{e_1, \bar{e}_1, \dots, e_k, \bar{e}_k\}$. The question is whether for all patterns $P \subseteq E'$ the directed graph H has a Hamiltonian cycle respecting P . Given this instance, the undirected graph G_H is defined in two steps. In a first step, we consider the set of $2n$ vertices

$$\bigcup_{x \in V(H)} \{x_{\text{in}}, x_{\text{out}}\}.$$

In a second step, we add the following edges and gadgets:

- For $x \in V(H)$, we connect x_{in} to x_{out} with a city gadget.
- For $i \in [k]$, we connect the three vertices $v_{\text{out}}^{(i)}, u_{\text{in}}^{(i)}$ and $w_{\text{in}}^{(i)}$ with a \forall -gadget.




Figure 13: Construction of the graph G_H from H .

- For each arc $(x, y) \in E(H) \setminus E'$, we add the edge $x_{\text{out}}y_{\text{in}}$ to G_H .

Furthermore, we let all city gadgets, including those inside the XOR- and \forall -gadgets have a height of $2n^4$ and width of $2n^4 + 100n$. We let all \forall -gadgets have a width of $t = n^4$ (i.e. they contain exactly n^4 ladders each). This completes the description of G_H . A schematic of the construction is depicted in Figure 13. Observe that G_H contains a total of k \forall -gadgets, $3k$ XOR-gadgets, and $n + 16k \leq 17n$ city gadgets. Since each city gadget has $\Theta(n^8)$ vertices, and these are the majority of all vertices, we have $|V(G_H)| = \Theta(n^9)$. In particular G_H can be constructed in polynomial time given H .

Lemma 18. *The graph G_H is bipartite.*

Proof. First consider the graph G'_H obtained from G_H by deleting all the \forall -gadgets with exception of the vertices $v_{\text{out}}^{(i)}, u_{\text{in}}^{(i)}, w_{\text{in}}^{(i)}$. The graph G'_H is clearly bipartite, since every edge and every city gadget connects some vertex x_{out} to some vertex y_{in} , and the city gadgets connect their endpoints with an odd-length path. Consider Figure 10. By assigning for $i \in [10]$ the vertex x_i to color class $i \bmod 2$, and using Lemma 13, we see that a \forall -gadget is bipartite, such that the two sets $\{v_{\text{out}}\}$ and $\{u_{\text{in}}, w_{\text{in}}\}$ are contained in different parts of the bipartition. Since in G'_H the three vertices $v_{\text{out}}^{(i)}, u_{\text{in}}^{(i)}, w_{\text{in}}^{(i)}$ have the analogous behavior with respect to the bipartition of G'_H , we see that if we add the \forall -gadgets back into G'_H , the graph remains bipartite. Hence G_H is bipartite. \square

For the proof of our main theorem, we define a particular PM of G_H , called the *default PM* M_{def} . The PM $M_{\text{def}} \subseteq E(G_H)$ is defined by having

- all towers of all city gadgets in G_H in default state
- all \forall -gadgets in G_H in semi-default state
- all t ladders in every \forall -gadget in default state.

(By the term “all city gadgets in G_H ”, we mean all $n + 16k$ gadgets, including those inside other gadgets.) Observe that this is a well-defined perfect matching of G_H , since every vertex $x_{\text{in}}, x_{\text{out}}$ for $x \in V(H)$, as well as every vertex inside a city gadget or \forall -gadget is touched exactly once. Furthermore, let us say that some PM M' of G_H is in *semi-default* state, if with respect to M' , all the city gadgets in G_H and all \forall -gadgets in G_H are in semi-default state. The following shows that the semi-default PMs lie in a certain sense “dense” in the 1-skeleton of P_{G_H} .

Lemma 19. *For every PM M of G_H , there exists some PM M' of G_H in semi-default state, with flip distance $\text{dist}(M, M') \leq 23n$.*

Proof. Consider the vertex set $V_s \subseteq V(G_H)$ defined by

$$V_s := \bigcup_{x \in V(H)} \{x_{\text{in}}\} \cup \bigcup_X \{x_1^{(X)}, \dots, x_4^{(X)}\} \cup \bigcup_A \{x_1^{(A)}, \dots, x_{10}^{(A)}\}.$$

Here, the index X runs over all XOR gadgets in G_H , the index A runs over all \forall -gadgets in G_H , and $x_i^{(X)}$ denotes the corresponding vertex in the XOR-gadget (see Figure 4) and $x_i^{(A)}$ denotes the corresponding vertex in the \forall -gadget (see Figure 10). We make the following claim: A PM M' is in semi-default state if and only if M' assigns the same partner as the PM M_{def} to every vertex $v \in V_s$. Indeed, if we recall the definitions of a semi-default state of a city gadget, a XOR-gadget, and a \forall -gadget, we see that whenever for some PM M' the vertices of V_s are matched in the same way as in M_{def} , the PM M' must be in semi-default state (we use that every tower has an even number of vertices). We have $|V_s| = n + 12k + 10k \leq 23n$. Now, given an arbitrary PM M , consider the symmetric difference $M \triangle M_{\text{def}}$. Since $|V(G_H)| = \Theta(n^9)$, the set $M \triangle M_{\text{def}}$ could in general contain up to $O(n^9)$ vertex-disjoint cycles. Let C_1, \dots, C_t be those cycles of $M \triangle M_{\text{def}}$ that contain some vertex of V_s . Then $t \leq 23n$. If we flip the vertex-disjoint cycles C_1, \dots, C_t one after another, we transform M into a semi-default PM M' , because M' assigns the same partners as M_{def} to vertices in V_s . By construction $\text{dist}(M, M') \leq 23n$. \square

4.6 Proof of the main theorem

In this subsection we prove Lemmas 5 and 6, which together prove our main theorem. Consider throughout this subsection a fixed instance of $\forall\exists$ -HAMCYCLE consisting out of a directed graph H with vertices $v^{(1)}, \dots, v^{(k)}$ and edges $E' = \{e_1, \bar{e}_1, \dots, e_k, \bar{e}_k\}$. Let G_H be defined as in Section 4.5. As is standard in hardness reductions, we can assume a lower bound on the size of the input instance. Specifically, we assume w.l.o.g. that $n = |V(H)| \geq 83\,000$.

Lemma 5. *If for all patterns $P \subseteq E'$ the directed graph H contains a Hamiltonian cycle respecting P , then $\text{diam}(P_{G_H}) \leq 4n^4 + 46n$.*

Proof. For all patterns $P \subseteq E'$, let C_P be a Hamiltonian cycle respecting P in the directed graph H . Let M_1, M_2 be two arbitrary PMs of G_H . We show that $\text{dist}(M_1, M_2) \leq 4n^4 + 46n$. Since M_1, M_2 are arbitrary, this proves the lemma. First, due to Lemma 19, there exist PMs M'_1, M'_2 of G_H in semi-default state and $\text{dist}(M_1, M'_1) \leq 23n$ and $\text{dist}(M_2, M'_2) \leq 23n$. We claim $\text{dist}(M'_1, M'_2) \leq 4n^4$. If we can show this claim, we are done, since then

$$\text{dist}(M_1, M_2) \leq \text{dist}(M_1, M'_1) + \text{dist}(M'_1, M'_2) + \text{dist}(M'_2, M_2) \leq 4n^4 + 46n.$$

Let T be a fixed tower gadget in G_H . Since T has height $2n^4$ and T is in semi-default state with respect to both M'_1 and M'_2 , due to Lemma 10 there exists a well-behaved flip sequence $(P_1^{(T)}, \dots, P_d^{(T)})$ for T of length $d = 4n^4$ that transforms the PM M'_1 restricted to T into the PM M'_2 restricted to T .

Let A_j be a fixed \forall -gadget of G_H for some $j \in [k]$. Note that A_j contains n^4 ladders L_1, \dots, L_{n^4} . Let $i \in [n^4]$ and consider the ladder L_i . Since both M'_1 and M'_2 are in semi-default state, the ladder L_i is semi-default with respect to both M'_1 and M'_2 . Due to Lemma 15 there exists a sequence $(P_1^{(L_i)}, \dots, P_4^{(L_i)})$ of length 4 of well-behaved paths for L_i transforming the PM M'_1 restricted to L_i into the PM M'_2 restricted to L_i , and such that both $P_1^{(L_i)}, P_2^{(L_i)}$ come from the same direction (top/bottom) and both $P_3^{(L_i)}, P_4^{(L_i)}$ come from the same direction (top/bottom). We can represent this fact as a string $s(L_i) \in \{tt, bb\}^2$, i.e. one of the strings $tttt, ttbb, bbtt, bbbb$. More formally, the string $s(L_i)$ has four characters, indicating that if one can visit the ladder L_i with four cycles coming from either the top (t) or bottom (b) in the same sequence as in the string $s(L_i)$, then one can transform M'_1 into M'_2 restricted to the ladder L_i . We can concatenate the strings $s(L_i)$ for $i = 1$ up to $i = n^4$ and obtain a string

$$s(A) \in \{tt, bb\}^{2n^4}.$$

We can interpret the string $s(A_j)$ as the “total demand” of the \forall -gadget A_j . If we can manage to visit the ladders of A_j with $4n^4$ well-behaved cycles from the top or bottom in the same sequence as the string $s(A_j)$ describes, we can transform M'_1 into M'_2 (restricted to A_j). Let A_1, \dots, A_k be the k distinct \forall -gadgets contained in G_H . Observe that the demand strings $s(A_j), s(A_{j'})$ can in general be very different from each other for distinct $j, j' \in [k]$. However, we now crucially use the property that H has a Hamiltonian cycle for *all* patterns $P \subseteq E'$. Concretely, for all $j \in [k]$ consider the first two characters of the string $s(A_j)$, denoted by $p_j \in \{tt, bb\}$. We can define a pattern $P \subseteq E' \subseteq E(H)$ based on these characters in a natural way by letting

$$P := \{e_j : j \in [k], p_j = tt\} \cup \{\bar{e}_j : j \in [k], p_j = bb\}.$$

We define a cycle C in G_H depending on those characters as follows:

- For all $j \in [k]$, if $p_j = tt$, then C traverses the \forall -gadget A_j in top state and visits ladder L_1 in that gadget from the top.
- Otherwise, if $p_j = bb$, then C traverses the \forall -gadget A_j in bottom state and visits ladder L_1 in that gadget from the bottom.
- C visits all city gadgets in G_H
- In between the cities and \forall -gadgets, the cycle C follows globally the same path as the Hamiltonian cycle C_P . Formally, for all arcs $(a, b) \in E(C_P) \setminus E'$, we have $a_{\text{out}}b_{\text{in}} \in E(C)$.
- Restricted to each ladder gadget L , the cycle C is equal to $P_1^{(L)}$. Restricted to each tower gadget T , the cycle C is equal to $P_1^{(T)}$.

Claim 1: C is a well-defined cycle in G_H and is alternating with respect to M'_1 .

Proof of the claim. We split C into those segments inside and those segments outside some \forall -gadget. Restricted to some \forall -gadget A , the cycle C is a well-defined path, since both top-state and bottom-state have this property (see Figure 12). Furthermore, inside of A the cycle C is alternating with respect to M'_1 , since M'_1 is in semi-default state (see Figure 11). In particular, note that since the XOR-gadgets are in semi-default state, both ways in which C could interact with the XOR-gadget are alternating (see Figure 5). Restricted to the ladder L_1 inside of A , the cycle C is equal to $P_1^{(L_1)}$ and hence M'_1 -alternating as well. Observe that for all $j \in [k]$ if $p_j = tt$, then C traverses gadget A_j in top state, i.e. it goes from $v_{\text{out}}^{(j)}$ to $u_{\text{in}}^{(j)}$. We can interpret this as corresponding to the arc $e_j = (v^{(j)}, u^{(j)}) \in E(H)$. Likewise, if $p_j = bb$, we can interpret this as the arc $\bar{e}_j \in E(H)$.

Outside the \forall -gadgets, the cycle C is a well-defined cycle, because whenever C encounters some city gadget of some vertex $x \in V(H)$, it uses the gadget to go from x_{in} to x_{out} . In order to change from one city gadget to another, C uses either a \forall -gadget or some edges corresponding to arcs of $C_P \subseteq H$. If C uses a \forall -gadget it behaves just like the pattern P . Since C_P is a Hamiltonian cycle of H respecting the pattern P , we conclude that C is a single cycle in G_H that visits every city gadget. Since every city gadget is in semi-default state in M'_1 , and since for every tower T the cycle C is equal to $P_1^{(T)}$, the cycle C is M'_1 -alternating. This proves the claim.

We define a second cycle C' in G_H , by specifying that C' has exactly the same edges as C , with the exception of all tower gadgets and all ladders that C traverses. For all towers T , the cycle C' is now equal to $P_2^{(T)}$ instead of $P_1^{(T)}$. Analogously, for all ladders L that C visits, the new cycle C' is equal to $P_2^{(L)}$ instead of $P_1^{(L)}$.

Claim 2: C' is a well-defined cycle in G_H and is alternating with respect to $M'_1 \triangle C$, and $M'_1 \triangle C \triangle C'$ is a PM in semi-default state.

Proof of the claim. Since C is a well-defined cycle and C' differs from it only on the towers and ladders, C' is also a well-defined cycle (note that if C and C' differ on a ladder, both come from

the same direction). Consider all edges of C' not part of some ladder or tower. All these edges are also part of C , and C is M'_1 -alternating. Hence C' is alternating with respect to $M'_1 \triangle C$ on those edges. Inside some tower T , the cycle C' is also alternating with respect to $M'_1 \triangle C$, because $(P_1^{(T)}, P_2^{(T)}, \dots)$ is a well-behaved flip sequence. The same holds for ladders. Finally, $M'_1 \triangle C \triangle C'$ is in semi-default state, because M'_1 is in semi-default state, and since both the cycles C, C' traverse all city gadgets of the graph (as well as the edge x_9x_{10} of every \forall -gadget). Hence we can check that in every city gadget, every XOR-gadget, and every \forall -gadget the PM $M'_1 \triangle C \triangle C'$ is again in semi-default state (because, intuitively speaking, every city gadget and the edges x_9x_{10} were flipped twice). This proves the claim.

To summarize, we have found two cycles C, C' such that by flipping C, C' we have eliminated the first two characters of all demand strings $s(A_j)$, for $j \in [k]$ (which have length $4n^4$) and for each tower T somewhere in G_H we have successfully flipped the first two paths $P_1^{(T)}, P_2^{(T)}$ of its flip sequence (which has length $4n^4$). Furthermore the PM $M'_1 \triangle C \triangle C'$ is again in semi-default state. Iterating this argument shows that one can reach M'_2 from M'_1 by flipping $4n^4$ cycles, hence $\text{dist}(M'_1, M'_2) \leq 4n^4$. We remark that in the next iteration, the pattern P is in general different from the pattern in this iteration, but that is not a problem, since H contains a Hamiltonian cycle for all patterns. \square

Lemma 6. *If $\text{diam}(P_{G_H}) \leq 4n^4 + 46n$, then for all patterns $P \subseteq E'$ the directed graph H contains a Hamiltonian cycle respecting P .*

Proof. We give a short overview of the proof: Let $P \subseteq E'$ be an arbitrary pattern. We define depending on P a perfect matching M_P of G_H and consider it together with the default matching M_{def} . By assumption $\text{diam}(P_{G_H}) \leq 4n^4 + 46n$, so in particular $\text{dist}(M_P, M_{\text{def}}) \leq 4n^4 + 46n$. We then proceed to show that in any sufficiently short flip sequence (C_1, \dots, C_d) from M_P to M_{def} with $d \leq 4n^4 + 46n$, for some $i \in [d]$ a cycle C_i can be transformed into a Hamiltonian cycle respecting P in H . Roughly speaking, the fact that C_i mimics a Hamiltonian path will follow from the fact that most of the cycles C_1, \dots, C_d must be regular (i.e. visit every city). Similarly, the fact that C_i mimics a cycle respecting the pattern P will follow from the fact that a regular cycle must (usually) visit a top-open ladder from the top and a bottom-open ladder from the bottom. Hence the fact $\text{diam}(P_{G_H}) \leq 4n^4 + 46n$ implies that H contains a Hamiltonian cycle respecting P . Since P was arbitrary, this suffices to show.

Let now $P \subseteq E'$ be a pattern. Let A_1, \dots, A_k be the \forall -gadgets of G_H . We define a PM M_P of G_H as follows.

- M_P is in semi-default position.
- All tower gadgets are in locked position.
- For all $j \in [k]$, we distinguish whether $e_j \in P$ or $\bar{e}_j \in P$. If $e_j \in P$, then all the n^4 ladders of A_j are top-open ladders. Otherwise, if $\bar{e}_j \in P$, then all the n^4 ladders of A_j are bottom-open ladders.

We verify that M_P is a well-defined perfect matching of G_H . Since M_P is in semi-default position, all vertices outside tower and ladder gadgets are touched exactly once. All vertices inside some tower or ladder gadget are also touched exactly once.

Consider the PM M_{def} as defined in Section 4.5. By assumption there exists a flip sequence $\mathcal{C} = (C_1, \dots, C_d)$ with $d \leq 4n^4 + 46n$ from M_P to M_{def} . By Lemma 12 at most $1000n^2$ of these cycles are not regular, and \mathcal{C} contains at least $4n^4 - 1002n^2$ regular cycles. For some $j \in [k]$ consider some fixed \forall -gadget A_j . The gadget A_j contains n^4 ladders L_1, \dots, L_{n^4} . Let us call a ladder *corrupted*, if it is visited by at least one irregular cycle in \mathcal{C} . Consider the set

$$\mathcal{L}_j = \{L_i : i \in [n^4], L_i \text{ is not corrupted} \}$$

of all uncorrupted ladders in A_j . Due to Lemma 17 we have $|\mathcal{L}_j| \geq n^4 - 4000n^2$. Every ladder in \mathcal{L}_j gets only visited by regular cycles. On the other hand, every regular cycle visits only one ladder of the gadget A_j due to Lemma 16. Since we have $d \leq 4n^4 + 46n$, an average ladder from \mathcal{L}_j gets visited at most

$$\frac{4n^4 + 46n}{n^4 - 4000n^2}$$

times, i.e. roughly 4 times. More precisely, let

$$\mathcal{L}'_j := \{L \in \mathcal{L}_j : L \text{ gets visited exactly 4 times}\}.$$

Then we have

$$4|\mathcal{L}'_j| + 5|\mathcal{L}_j \setminus \mathcal{L}'_j| \leq d \leq 4n^4 + 46n.$$

Since of course $|\mathcal{L}'_j| + |\mathcal{L}_j \setminus \mathcal{L}'_j| = |\mathcal{L}_j|$, this implies

$$\begin{aligned} |\mathcal{L}_j \setminus \mathcal{L}'_j| &\leq 4n^4 + 46n - 4|\mathcal{L}_j| \leq 16\,000n^2 + 46n \\ \text{and} \quad |\mathcal{L}'_j| &= |\mathcal{L}_j| - |\mathcal{L}_j \setminus \mathcal{L}'_j| \geq n^4 - 20\,000n^2 - 46n. \end{aligned}$$

Now consider again the gadget A_j and how it interacts with some regular cycle C . Due to Lemma 16, the cycle C is either in top state or bottom state with respect to gadget A_j and visits only one ladder from the same direction. Let us say C respects the pattern P at A_j , if either both C is in top state and $e_j \in P$, or both C is in bottom state and $\bar{e}_j \in P$. Let

$$\mathcal{C}_j := \{C_i : i \in [d], C_i \text{ is regular and } C_i \text{ respects } P \text{ at } A_j\}.$$

Recall that we have defined M_P to contain only top-open ladders inside A_j if $e_j \in P$, and only bottom-open ladders inside A_j if $\bar{e}_j \in P$. Note that every ladder in \mathcal{L}'_j is visited only by regular (and therefore well-behaved) cycles, and exactly four of them. Hence the assumptions of Lemma 14 are met. Therefore these 4 cycles all need to come from the top if L is top-open, and they all need to come from the bottom if L is bottom-open. This shows

$$|\mathcal{C}_j| \geq 4|\mathcal{L}'_j| \geq 4n^4 - 81\,000n^2.$$

This implies that for all $j \in [k]$

$$|\mathcal{C} \setminus \mathcal{C}_j| \leq d - 4n^4 + 81\,000n^2 \leq 81\,000n^2 + 46n.$$

Finally, using $|\mathcal{C}| \geq 4n^4 - 1002n^2$ we arrive at

$$\bigcap_{j=1}^k \mathcal{C}_j \geq |\mathcal{C}| - \sum_{j=1}^k |\mathcal{C} \setminus \mathcal{C}_j| \geq 4n^4 - 83\,000n^3.$$

In particular, for $n = |V(H)|$ large enough, there exists some cycle $C \in \bigcap_{j=1}^k \mathcal{C}_j$. This cycle $C \subseteq G_H$ is regular, i.e. it traverses every city gadget corresponding to some $x \in V(H)$ from x_{in} to x_{out} . Furthermore, it traverses every \forall -gadget in the same manner as the pattern P . Therefore the directed graph H has a Hamiltonian cycle respecting the pattern P . \square

5 Inapproximability

In this section, we prove that for some $\varepsilon > 0$ the diameter (and the circuit diameter) of the bipartite perfect matching polytope cannot be approximated better than $(1 + \varepsilon)$ unless $P = NP$. We remark that our techniques inherently can show inapproximability only for a small $\varepsilon > 0$ (concretely we obtain $\varepsilon = 1/16\,226 \approx 6.1 \cdot 10^{-5}$), and we do not optimize our reductions to

obtain the best possible ε . To state our theorem completely formally, let us say an algorithm \mathcal{A} is an approximation algorithm for the diameter with approximation guarantee $(1 + \varepsilon)$, if \mathcal{A} never underestimates the diameter, and overestimates it by at most a factor $(1 + \varepsilon)$. This means

$$\text{diam}(P) \leq \mathcal{A}(P) \leq (1 + \varepsilon) \text{diam}(P).$$

(The same kind of approximation is considered in [San18]). The main result of this section is then stated as follows.

Theorem 2. *There exists $\varepsilon > 0$ such that the diameter of the bipartite perfect matching polytope cannot be approximated better than $(1 + \varepsilon)$ in polynomial time, unless $P \neq NP$.*

We remark that alternatively one could also consider approximation algorithms that are allowed to both over- and underestimate the true diameter, i.e. algorithms with

$$\max\left\{\frac{\mathcal{A}(P)}{\text{diam}(P)}, \frac{\text{diam}(P)}{\mathcal{A}(P)}\right\} \leq (1 + \varepsilon').$$

Since the proof of Theorem 2 is in terms of a gap-reduction, it follows that such two-sided approximation algorithms cannot have an approximation guarantee better than $(1 + \varepsilon')$, where $(1 + \varepsilon')^2 = (1 + \varepsilon)$, i.e. $\varepsilon' \approx \varepsilon/2$.

In the remainder of this section, the term 3SAT denotes the satisfiability problem, where the input is a boolean formula φ in conjunctive normal form (CNF) such that every clause has at most 3 literals. Max 3SAT is the maximization problem corresponding to 3SAT, where the goal is to find an assignment that satisfies as many of the clauses of φ as possible. The problem Max 3SAT- d is equal to problem Max 3SAT with the additional restriction that for each variable x_i , the number of clauses that contain the literal x_i or the literal \bar{x}_i is at most d .

The following is a direct consequence of the PCP theorem [ALM⁺98] in combination with constructions of certain expander graphs and is proven e.g. in Arora [Aro94, p.84], see also the survey of Trevisan [Tre04].

Theorem 20 (Theorem 7 in [Tre04]). *There are constants d and ε_1 and a polynomial time computable reduction from 3SAT to Max 3SAT- d such that if φ is satisfiable then $f(\varphi)$ is satisfiable, and if φ is not satisfiable then the optimum of $f(\varphi)$ is less than $1 - \varepsilon_1$ times the number of clauses.*

Using a standard strategy for inapproximability proofs, we can adapt Theorem 20 to our needs. Specifically, consider a graph $G = (V, E)$. A *closed walk* in G is a sequence $W = (v_1, \dots, v_k)$ of vertices, such that $v_i v_{i+1} \in E$ for all $i \in [k - 1]$ and also $v_k v_1 \in E$. In contrast to a cycle, a closed walk may visit some vertices more than once. Given a closed walk W in G and some $i \in \mathbb{N}_0$, we let

$$W_i := \{v \in V : W \text{ visits } v \text{ exactly } i \text{ times}\}.$$

Definition 21. *Let $\varepsilon > 0$. A walk W in a graph $G = (V, E)$ on $n = |V|$ vertices is called ε -good, if $|W_1| \geq (1 - \varepsilon)n$ (or equivalently $|V \setminus W_1| \leq \varepsilon n$).*

It seems likely that the following result is already known, however we were unable to locate a reference. It proves hardness of approximation for walks that are “almost Hamiltonian”, i.e. ε -good.

Theorem 22. *There is a constant $\varepsilon_2 > 0$ and a polynomial time computable reduction that takes a 3SAT formula φ and outputs an undirected graph H such that*




Figure 14: Folklore reduction of 3SAT to the Hamiltonian cycle problem. Arrows marked with a cross denote an XOR-gadget, as illustrated in (a). Vertices marked with a square are all connected in one clique.

- If φ is satisfiable, then H contains a Hamiltonian cycle
- If φ is not satisfiable, then H does not contain an ε_2 -good walk.

In order to prove Theorem 22, we rely on a folklore reduction from 3SAT to the Hamiltonian cycle problem. We show that this reduction without any modification satisfies also the stronger requirements of Theorem 22. The reduction is defined as follows: Assume we are given a 3SAT instance φ' in CNF with clauses C_1, \dots, C_m and variables x_1, \dots, x_k .

The reduction is depicted in Figure 14. Let ab and $a'b'$ be two edges. We first define XOR-gadgets between ab and $a'b'$ by introducing additional vertices y_1, \dots, y_{12} and connecting them like in Figure 14. They are similar to the XOR-gadgets in Section 4.3, with the exception that we replace the city gadgets with single vertices. Analogously to before, we can conclude that every Hamiltonian cycle uses exactly one of the two edges ab and $a'b'$ of a XOR-gadget. Given the 3SAT formula φ , we can then define an undirected graph H as follows: H contains k variable gadgets, m clause gadgets, and three additional vertices v_1, v_2, v_3 as depicted in Figure 14. Here, variable gadgets are defined as follows: The graph H includes vertices z_1, \dots, z_{2k} . For all $i \in [k]$ there are two parallel edges from z_{2i-1} to z_{2i} , which we denote by e_i and \bar{e}_i . We interpret them as the edges corresponding to literal x_i and literal \bar{x}_i . We connect e_i and \bar{e}_i with a XOR-gadget. Note that after the XOR gadget is applied, there is no multi-edge anymore, i.e. we still have that H is a simple graph. A clause gadget is defined as follows: For some clause C_j with $C_j = \ell_1 \vee \dots \vee \ell_{t_j}$ where $t_j \in \{1, 2, 3\}$, we introduce t_j vertices $u_1^{(j)}, \dots, u_{t_j}^{(j)}$ in H , and we connect them in a cycle of length t_j (i.e. a loop if $t_j = 1$, two parallel edges if $t_j = 2$, and a triangle if $t_j = 3$). For $i = 1, \dots, t_j$, a XOR-gadget connects the i -th edge of this cycle to the edge inside the variable gadget corresponding to the literal ℓ_i . Note that again this makes the graph simple. We add the edges $v_1v_2, v_2z_1, z_{2k}v_3$ and $z_{2i}z_{2i+1}$ for all $i = 1, \dots, k-1$. Finally, in the vertex set

$$V' := \bigcup_{j=1}^m \{u_1^{(j)}, \dots, u_{t_j}^{(j)}\} \cup \{v_1, v_3\},$$

every vertex is connected to each other (i.e. H is a clique induced on V'). The set V' is highlighted with square marks in Figure 14.

We quickly verify that this folklore reduction is indeed a correct reduction from 3SAT to Hamiltonian cycle. Indeed, every Hamiltonian cycle F visits v_2 (note that $v_2 \notin V'$). Hence every Hamiltonian cycle traverses the variable gadgets in order and uses exactly one of the two edges e_i or \bar{e}_i for $i \in [k]$. (For simplicity of notation, we say that the cycle F uses e_i even though due to the XOR-gadgets the edge e_i is not an actual edge of H .) If for some clause gadget corresponding to clause C_j the cycle F makes the wrong variable choices, i.e. includes none of the literals of C_j , then due to the XOR gadgets, the cycle F restricted to the clause gadget of C_j is already a cycle. But that is a contradiction, since a Hamiltonian cycle cannot contain a smaller cycle. Hence we conclude that if F is a Hamiltonian cycle in H , then φ' is satisfiable. On the other hand, due to V' inducing a clique, if φ is satisfiable, then H has a Hamiltonian cycle.

We now show that the same reduction also satisfies the requirements of Theorem 22. The main idea behind the proof is the following: Since every ε -good walk W has $|W_1| \geq (1 - \varepsilon)|V(H)|$, we have that the majority of all gadgets in H are entirely contained in W_1 , i.e. all their vertices are visited exactly once by W . We show that this implies that most of the gadgets work as intended, hence for ε_2 small enough, a ε_2 -good walk W implies that a fraction of at least $(1 - \varepsilon_1)$ clauses of φ can be made true.

Proof of Theorem 22. Assume we are given some 3SAT instance φ . Let ε_1, d be constants as in Theorem 20. In a first step we transform φ into an instance $\varphi' = f(\varphi)$ of Max 3SAT- d such that φ' has the properties described in Theorem 20. In a second step, we transform φ' into a graph $H = (V, E)$ as described above. We let

$$\varepsilon_2 := \frac{\varepsilon_1}{61(d+1)}.$$

If φ' is satisfiable, then H contains a Hamiltonian cycle, as proven above. So it remains to prove that if φ' is not satisfiable, then H does not contain a ε_2 -good walk. For the sake of contradiction, assume H contains a ε_2 -good walk W , we show that φ' is satisfiable. Let $n := |V(H)|$. Let m be the number of clauses of φ' . Note that the graph H contains k variable gadgets, m clause gadgets, at most $3m + k$ XOR-gadgets, and the three additional vertices v_1, v_2, v_3 . Since these in total are all the vertices of H , we have (using $k \leq 3m$ and w.l.o.g. $m \geq 4$)

$$n \leq 2k + 3m + k + 16(3m) + 3 \leq 60m + 3 < 61m.$$

Consider some XOR-gadget g in the graph H . We say that the XOR-gadget g consists of precisely the 16 vertices $V_g := \{a, b, a', b'\} \cup \{y_1, \dots, y_{12}\}$ as shown in Figure 14. Let $i \in [k]$ and consider the 3SAT variable x_i . Observe that the variable gadget of x_i contains the XOR-gadget g_0 between e_i, \bar{e}_i and in addition to that it is connected to at most d XOR-gadgets g_1, \dots, g_t with $t \leq d$. Let us say that with respect to the walk W , the variable x_i is *uncorrupted*, if

$$\left(\{z_{2i-1}, z_{2i}\} \cup \bigcup_{j=0}^t V_{g_j} \right) \subseteq W_1,$$

and it is corrupted otherwise. Since the above vertex sets are pairwise disjoint for all $i \in [k]$, we have that every vertex in $V \setminus W_1$ can corrupt at most one variable. Hence

$$|\{i \in [k] : x_i \text{ is corrupted}\}| \leq |V \setminus W_1| \leq \varepsilon_2 n.$$

We claim that if a variable x_i is uncorrupted, then the walk W traverses the variable gadget of x_i either only using edge e_i , or only using edge \bar{e}_i . Indeed, consider the walk W and its interaction with some XOR-gadget g . Since the XOR-gadget is contained entirely in W_1 (i.e. $V_g \subseteq W_1$), we can see that the walk W enters and leaves every of the vertices y_5, \dots, y_8 exactly

once. Furthermore, for each $y \in \{y_5, \dots, y_8\}$, the walk enters and leaves y using two different edges, since otherwise the walk would use a neighbor of y more than once (but all neighbors of y belong to W_1). Hence the walk W contains the subwalks (y_i, y_{i+4}, y_{i+8}) for all $i = 1, \dots, 4$. By extending this argument to y_1, \dots, y_4 and y_9, \dots, y_{12} , we see that W uses exactly one of the two edges ab and $a'b'$. Now, consider vertex z_{2i-1} . Since an uncorrupted XOR-gadget connects e_i and \bar{e}_i , the walk has to use one of the two edges e_i, \bar{e}_i . On the other hand, if the walk would use both edges e_i, \bar{e}_i , then because all XOR-gadgets (attached to either e_i or \bar{e}_i) work properly, the walk meets itself at z_{2i} . Hence $z_{2i} \notin W_1$, a contradiction. This proves the claim.

For $j \in [m]$ consider the clause C_j . It contains at most three literals, i.e. $C_j = \ell_1 \vee \dots \vee \ell_{t_j}$ with $t_j \in \{1, 2, 3\}$. Let us call the clause C_j corrupted, if one of the variables corresponding to its literals $\ell_1, \dots, \ell_{t_j}$ is corrupted, or if one of the t_j vertices $u_1^{(j)}, \dots, u_{t_j}^{(j)}$ of the clause gadget is not contained in W_1 . For the first reason at most $d\varepsilon_2 n$ clauses are corrupted, because a corrupted variable corrupts at most d clauses. For the second reason, at most $|V \setminus W_1| \leq \varepsilon_2 n$ clauses are corrupted, since the corresponding vertex sets are disjoint. Therefore

$$|\{j \in [m] : C_j \text{ is corrupted}\}| \leq d\varepsilon_2 n + \varepsilon_2 n < 61(d+1)\varepsilon_2 m = \varepsilon_1 m.$$

For $i \in [k]$, we can now consider the natural variable assignment

$$\alpha(x_i) := \begin{cases} 1 & \text{if } x_i \text{ uncorrupted, and } W \text{ uses } e_i \\ 0 & \text{if } x_i \text{ uncorrupted, and } W \text{ uses } \bar{e}_i \\ \text{arbitrary} & \text{if } x_i \text{ corrupted.} \end{cases}$$

For every uncorrupted clause C_j for some $j \in [m]$, we claim that the assignment α satisfies C_j . Indeed, note that all of its variables are uncorrupted, and so all of the XOR-gadgets attached to C_j work as intended. Therefore if α does not satisfy C_j , then the walk W describes a cycle restricted to the clause gadget, and so one of the vertices $u_1^{(j)}, \dots, u_{t_j}^{(j)}$ is visited at least twice by W , a contradiction. Since there are less than $\varepsilon_1 m$ corrupted clauses, due to Theorem 20, formula φ' has even an assignment that satisfies every clause. This was to show. \square

Theorem 2. *There exists $\varepsilon > 0$ such that the diameter of the bipartite perfect matching polytope cannot be approximated better than $(1 + \varepsilon)$ in polynomial time, unless $P \neq NP$.*

Proof. We show that there exists a polynomial-time computable reduction, that given a 3SAT formula φ returns a bipartite graph G , such that for some constant $n_0 \in \mathbb{N}$

- If φ is satisfiable, then $\text{diam}(P_G) \leq 2n^2 + 2n$
- If φ is not satisfiable, then $\text{diam}(P_G) > (1 + \varepsilon)(2n^2 + 2n)$ for all $n \geq n_0$.

This suffices to prove the theorem. Given a SAT formula φ , due to Theorem 22 there exists a constant $\varepsilon_2 > 0$ and a polynomial-time computable graph H such that if φ is satisfiable, H has a Hamiltonian cycle, and if φ is not satisfiable, H does not have an ε_2 -good walk. We choose some constant $0 < \delta < \varepsilon_2$ very close to zero and let

$$\varepsilon := \frac{1}{(1 + \delta)(1 - \varepsilon_2)} - 1.$$

Note that $\varepsilon > 0$, since $0 \leq (1 + \delta)(1 - \varepsilon_2) = 1 + \delta - \varepsilon_2 - \delta\varepsilon_2 < 1$. We construct a graph G from H as follows. We let $n := |V(H)|$. First we consider the vertex set

$$\bigcup_{v \in V(H)} \{v_{\text{in}}, v_{\text{out}}\}.$$

Then we connect for all $v \in V(H)$ the two vertices $v_{\text{in}}, v_{\text{out}}$ with a city gadget of height n^2 and width $4n^2 + 4n$. Finally, we add the edge set

$$\bigcup_{ab \in E(H)} \{a_{\text{out}}b_{\text{in}}, b_{\text{out}}a_{\text{in}}\}$$

to H . We remark that this construction of G resembles the original construction of [NS25]. It is similar to our construction of Section 4.5, with the difference that H is undirected (in Section 4.5 for technical reasons it is more convenient to work with a directed graph, while in the current section it is more convenient to work with an undirected graph). The graph G is bipartite, since city gadgets connect their endpoints via an odd-length path, and all other edges connect some vertex a_{out} to some vertex b_{in} . The proof of Theorem 2 is now completed with the following two Lemmas 23 and 24. \square

Lemma 23. *If φ is satisfiable, then $\text{diam}(P_G) \leq 2n^2 + 2n$.*

Proof. Let M_1, M_2 be arbitrary PMs of G , we will show $\text{dist}(M_1, M_2) \leq 2n^2 + 2n$. Since M_1, M_2 are arbitrary, this suffices to show. Call some PM M in *semi-default state*, if every city gadget is in semi-default state with respect to M . Observe that G contains certainly at least one PM M_{def} which is in semi-default state. Consider the vertex set $V_s := \{v_{\text{in}} : v \in V(H)\}$. We have $|V_s| = n$. Furthermore, a PM M is in semi-default state if and only if the PM M assigns the same partners to vertices of V_s as the matching M_{def} . Analogously to Lemma 19 we conclude that there exist semi-default PMs M'_1, M'_2 such that $\text{dist}(M_1, M'_1) \leq n$ and $\text{dist}(M_2, M'_2) \leq n$. Let T be fixed tower gadget in G . Since T has height n^2 , due to Lemma 10 there is a well-behaved flip sequence $(P_1^{(T)}, \dots, P_d^{(T)})$ of length $d = 2n^2$ that transforms M'_1 into M'_2 restricted to T . Since φ is satisfiable, G contains a Hamiltonian cycle F . We can consider for $i = 1, \dots, 2n^2$ the cycle C_i defined as follows: C_i visits every city gadget. Inside some tower T , the cycle C_i is equal to $P_i^{(T)}$. Outside the towers, it follows globally the same route as the Hamiltonian cycle F , i.e. it uses all the edges $\{a_{\text{out}}b_{\text{in}} : ab \in E(F)\}$. Analogously to Lemma 5, we see that for each $i \in [2n^2]$, the object C_i is a well-defined cycle. Since M'_1 is in semi-default state, one can show by induction that the cycle C_i is alternating with respect to $M'_1 \triangle C_1 \triangle \dots \triangle C_{i-1}$ for all $i \in [d]$. Furthermore, the flip sequence (C_1, \dots, C_d) transforms M'_1 into M'_2 . We conclude in total

$$\text{dist}(M_1, M_2) \leq \text{dist}(M_1, M'_1) + \text{dist}(M'_1, M'_2) + \text{dist}(M'_2, M_2) \leq n + 2n^2 + n.$$

\square

Lemma 24. *If φ is not satisfiable, then for all n larger than some constant n_0 , we have $\text{diam}(P_G) > (1 + \varepsilon)(2n^2 + 2n)$.*

Proof. We consider the contrapositive, that is we that assume $\text{diam}(P_G) \leq (1 + \varepsilon)(2n^2 + 2n)$, and we want show that φ is satisfiable. Consider PMs M_1, M_2 of G defined as follows. Both M_1, M_2 have all city gadgets in semi-default position. Furthermore, M_1 has all tower gadgets in locked position (see Figure 1c). In contrast, M_2 has all tower gadgets in default position (see Figure 1b). Note that M_1 and M_2 are well-defined PMs of G . Let $d := \text{dist}(M_1, M_2)$. By assumption on the diameter, we have $d \leq (1 + \varepsilon)(2n^2 + 2n)$ and there exists a flip sequence (C_1, \dots, C_d) from M_1 to M_2 . Consider for some vertex $v \in V(H)$ the corresponding city gadget g_v . By definition its width is $t = 4n^2 + 4n$. Due to $\varepsilon < 1$, we have $d < t$. This implies that among the t towers in city g_v , there exists one tower T_v^* with the property that none of the cycles C_1, \dots, C_d is entirely contained in T_v^* . Hence every cycle of (C_1, \dots, C_d) either is well-behaved for T_v^* or does not visit the city g_v at all. For each $i \in [d]$, let p_i be the number of different towers T_v^* that the cycle C_i visits, formally

$$p_i := |\{v \in V(H) : C_i \text{ visits } T_v^*\}|.$$

We apply Lemma 9 to the tower T_v^* , and see that at least $2n^2 - 2$ cycles need to visit T_v^* in order to transform M_1 into M_2 (restricted to T_v^*). On the other hand, a fixed cycle C_i can visit each tower T_v^* at most once. By double counting we obtain

$$\sum_{i=1}^d p_i = |\{(i, v) \in [d] \times V(H) : C_i \text{ visits } T_v^*\}| \geq (2n^2 - 2)n.$$

Since δ is a constant, for $n \geq n_0(\delta)$ large enough we have $(1 + \delta)(2n^3 - 2n) \geq (2n^3 + 2n^2)$. Then by the pigeonhole principle, there exists a $j \in [d]$ such that

$$p_j \geq \frac{2n^3 - 2n}{(1 + \varepsilon)(2n^2 + 2n)} \geq \frac{2n^3 + 2n^2}{(1 + \varepsilon)(1 + \delta)(2n^2 + 2n)} = \frac{n}{(1 + \varepsilon)(1 + \delta)} = (1 - \varepsilon_2)n.$$

Therefore the cycle C_j visits at least $(1 - \varepsilon_2)n$ city gadgets. We claim that we can modify the cycle $C_j \subseteq G$ to obtain an ε_2 -good walk W in H . We start with $W = \emptyset$ and modify W according to the following rule:

- Whenever C_j visits some vertex v_{in} in G (where $v \in V(H)$), the next step of C_j is either to traverse the city gadget and visit v_{out} , or to traverse some edge of the form $v_{\text{in}}w_{\text{out}}$. In the first case, we do not modify W , in the second case we add (v, w) to the walk W . Note that this is legal since $wv = vw$ is an edge in H .
- Analogously, whenever C_j visits some vertex v_{out} in G (where $v \in V(H)$), the next step of C_j is either to traverse the city gadget and visit v_{in} , or to traverse some edge of the form $v_{\text{out}}w_{\text{in}}$. In the first case, we do not modify W , in the second case we add (v, w) to the walk W . Again this is legal since vw is an edge in H .

Note that the above procedure indeed constructs a well-defined closed walk W in H . (If C_j is currently at one of the two vertices v_{in} or v_{out} , then W is currently at v). Furthermore, note that if C_j visits a city gadget g_v corresponding to some $v \in V(H)$, then $v \in W_1$. This is because C_j is a simple cycle and hence cannot visit a city twice. Since $p_j \geq (1 - \varepsilon_2)n$, we have $|W_1| \geq (1 - \varepsilon_2)n$, i.e. W is ε_2 -good. Since H was constructed according to Theorem 22, we have that the formula φ is satisfiable. \square

Concrete value of ε . Arora [Aro94] proves Theorem 20 for constants $d = 13$ and $\varepsilon_1 = 1/19$. Hence we obtain $\varepsilon_2 = 1/16\,226 \approx 6.1 \cdot 10^{-5}$ and finally $\varepsilon = \varepsilon_2$ (note that for δ arbitrarily small, ε can be made arbitrarily close to $1/(16\,226 - 1)$, where as a trade-off n_0 gets bigger). Since we did not optimize our reductions for the best possible ε , this can likely be improved.

6 The class Δ_3

For the sake of completeness, we discuss the following variant of Question 1: Frieze and Teng [FT94] consider the following variant DIAMETER', where the input are the constraints $A \in \mathbb{Q}^{m \times d}$, $b \in \mathbb{Q}^m$ of a polytope $P = \{x \in \mathbb{R}^d : Ax \leq b\}$ together with some $t \in \mathbb{N}$, and the question is whether $\text{diam}(P) = t$ (instead of $\text{diam}(P) \leq t$). Frieze and Teng ask whether DIAMETER' is complete for the complexity class $\Delta_3 = P^{\Sigma_2^p}$.

Unfortunately we are not able to completely answer this question. However, we can offer some insights. We derive these insights from an adaptation of Theorem 4.5 from [FT94]. Consider the problem HAMILTON' defined as follows: An instance of HAMILTON' is a tuple $I = (I_1, I_2)$ of two instances I_1, I_2 of $\forall\exists$ -HAMCYCLE. The instance I is a yes-instance if and only if I_1 is a yes-instance and I_2 is a no-instance of $\forall\exists$ -HAMCYCLE. By a standard argument, HAMILTON' is complete for Δ_3 . We give a reduction from HAMILTON' to DIAMETER'. By our main result,

Theorem 1, given I_1, I_2 one can construct in polynomial time graphs G_1, G_2 and polynomially bounded integers $n_1, n_2 \in \mathbb{N}$ such that for $i = 1, 2$

$$I_i \text{ is a yes-instance} \Leftrightarrow \text{diam}(P_{G_i}) \leq 4n_i^4 + 46n_i.$$

Let $N = |V(G_2)| + 1$. We let the graph G be the disjoint union out of N copies of G_1 and a single copy of G_2 . Since every alternating cycle of a flip sequence is contained in exactly one connected component of G , we have the equality

$$\text{diam}(G) = N \text{diam}(P_{G_1}) + \text{diam}(P_{G_2}).$$

Since $\text{diam}(P_{G_2}) < N$ holds trivially, we have $\text{diam}(P_{G_1}) = \lfloor \text{diam}(P_G)/N \rfloor$ and $\text{diam}(P_{G_2}) = \text{diam}(P_G) \bmod N$. Therefore one can decide HAMILTON' when given the precise value $\text{diam}(P_G)$ as an input. However, note that this reduction is not a standard many-one reduction from the decision problem HAMILTON' to the decision problem DIAMETER'. Instead it shows that HAMILTON' can be solved with a single oracle call to the optimization problem Π_O of computing $\text{diam}(P)$. Hence Π_O is Δ_3 -hard, but we cannot say anything about DIAMETER'. Assume that our main theorem could be modified, such that there are thresholds $t_i, i = 1, 2$ with the property that I_i is a yes-instance if and only if $\text{diam}(P_i) = t_i$ and $\text{diam}(P_i) = t_i - 1$ otherwise. Then the above immediately yields a many-one reduction from HAMILTON' to DIAMETER'. It is currently an open question whether such a modification is possible.

7 Conclusion

In this work, we proved that computing the diameter (the circuit diameter, respectively) of the bipartite perfect matching polytope is Π_2^P -complete. As a result, computation of the diameter can be compared to computation of other notoriously hard parameters like the generalized Ramsey number. Furthermore, we showed that the diameter (the circuit diameter, respectively) cannot be approximated in polynomial time with a factor better than $1 + \varepsilon$, unless $P = NP$. We remark that these two main results are incomparable. On the one hand the second result is stronger, because it is an inapproximability result. On the other hand, the second result is weaker, since it only shows NP-hardness instead of Π_2^P -hardness.

There are many open questions that remain. It seems likely that our value of $\varepsilon \approx 6.1 \cdot 10^{-5}$ can be improved, especially when comparing against the related result of Cardinal and Steiner for polytope distances [CS23a]. We hence ask the question, whether the (circuit) diameter of a polytope does or does not admit a constant-factor approximation. One could also ask, whether it is possible to combine our two main results, i.e. whether approximation of the (circuit) diameter is Π_2^P -hard. We remark however that even though theoretically valid, results of this kind seem to be very sparse in the literature. One reason for this might be that approximation algorithms are always required to be polynomial time, and so it could be argued that a completeness result for a higher class than NP offers less insights. The authors of this paper would be interested in such a result anyway.

Our Π_2^P -completeness result does not hold for the monotone diameter as defined in [NS25], since it follows from the arguments presented there that the decision problem for the monotone diameter of the bipartite perfect matching polytope is contained in NP. We therefore ask if for some other polytope it can be proven that computation of the monotone diameter is Π_2^P -hard. Furthermore, Kaibel and Pfetsch [KP03] ask for the complexity of computing the diameter of a *simple* polytope, i.e. a polytope $P \subseteq \mathbb{R}^d$ where every vertex has degree d . Since the bipartite perfect matching polytope is not simple, this question remains unanswered.

As a final question, we ask whether our Π_2^P -completeness and inapproximability results can also be extended to the case of bipartite, planar graphs of maximum degree 3, similar as in [IDH⁺11]. Our current Lemma 17 provides a barrier for this result, since it states that the

four vertices x_2, x_3, x_6, x_7 inside a \forall -gadget are essential to the proof. However, they have an arbitrarily high degree and together with their surroundings create a non-planar graph. Hence in order to prove hardness in this restricted case, a different approach may be needed.

References

- [ACH⁺21] Oswin Aichholzer, Jean Cardinal, Tony Huynh, Kolja Knauer, Torsten Mütze, Raphael Steiner, and Birgit Vogtenhuber. Flip distances between graph orientations. *Algorithmica*, 83(1):116–143, 2021.
- [ALM⁺98] Sanjeev Arora, Carsten Lund, Rajeev Motwani, Madhu Sudan, and Mario Szegedy. Proof verification and the hardness of approximation problems. *Journal of the ACM (JACM)*, 45(3):501–555, 1998.
- [Aro94] Sanjeev Arora. *Probabilistic checking of proofs and hardness of approximation problems*. University of California, Berkeley, 1994.
- [BBB23] Alexander E Black, Steffen Borgwardt, and Matthias Brugger. On the circuit diameter conjecture for counterexamples to the Hirsch conjecture. *arXiv preprint arXiv:2302.03977*, 2023.
- [BBFK21] Steffen Borgwardt, Cornelius Brand, Andreas Emil Feldmann, and Martin Koutecký. A note on the approximability of deepest-descent circuit steps. *Operations Research Letters*, 49(3):310–315, 2021.
- [BBH⁺19] Marthe Bonamy, Nicolas Bousquet, Marc Heinrich, Takehiro Ito, Yusuke Kobayashi, Arnaud Mary, Moritz Mühlenhaller, and Kunihiro Wasa. The perfect matching reconfiguration problem. In *44th International Symposium on Mathematical Foundations of Computer Science (MFCS 2019)*. Schloss Dagstuhl–Leibniz-Zentrum für Informatik, 2019.
- [BFH15] Steffen Borgwardt, Elisabeth Finhold, and Raymond Hemmecke. On the circuit diameter of dual transportation polyhedra. *SIAM J. Discret. Math.*, 29(1):113–121, 2015.
- [BGK⁺25] Steffen Borgwardt, Weston Grewe, Sean Kafer, Jon Lee, and Laura Sanità. On the hardness of short and sign-compatible circuit walks. *Discrete Applied Mathematics*, 367:129–149, 2025.
- [BHIM19] Nicolas Bousquet, Tatsuhiko Hatanaka, Takehiro Ito, and Moritz Mühlenhaller. Shortest reconfiguration of matchings. In Ignasi Sau and Dimitrios M. Thilikos, editors, *Graph-Theoretic Concepts in Computer Science - 45th International Workshop, WG 2019, Vall de Núria, Spain, June 19-21, 2019, Revised Papers*, volume 11789 of *Lecture Notes in Computer Science*, pages 162–174. Springer, 2019.
- [BMPT25] Carla Binucci, Fabrizio Montecchiani, Daniel Perz, and Alessandra Tappini. Flipping matchings is hard. *arXiv preprint arXiv:2503.02842*, 2025.
- [BSY18] Steffen Borgwardt, Tamon Stephen, and Timothy Yusun. On the circuit diameter conjecture. *Discrete & Computational Geometry*, 60:558–587, 2018.
- [BV20] Steffen Borgwardt and Charles Viss. An implementation of steepest-descent augmentation for linear programs. *Operations Research Letters*, 48(3):323–328, 2020.
- [BV22] Steffen Borgwardt and Charles Viss. Circuit walks in integral polyhedra. *Discret. Optim.*, 44(Part):100566, 2022.

- [Chv75] Vašek Chvátal. On certain polytopes associated with graphs. *Journal of Combinatorial Theory, Series B*, 18(2):138–154, 1975.
- [CS23a] Jean Cardinal and Raphael Steiner. Inapproximability of shortest paths on perfect matching polytopes. In Alberto Del Pia and Volker Kaibel, editors, *Integer Programming and Combinatorial Optimization - 24th International Conference, IPCO 2023, Madison, WI, USA, June 21-23, 2023, Proceedings*, volume 13904 of *Lecture Notes in Computer Science*, pages 72–86. Springer, 2023.
- [CS23b] Jean Cardinal and Raphael Steiner. Shortest paths on polymatroids and hypergraphic polytopes. *arXiv preprint arXiv:2311.00779*, 2023.
- [DFH23] Yann Disser, Oliver Friedmann, and Alexander V Hopp. An exponential lower bound for zadeh’s pivot rule. *Mathematical Programming*, 199(1):865–936, 2023.
- [DKNV22] Daniel Dadush, Zhuan Khye Koh, Bento Natura, and László A Végh. On circuit diameter bounds via circuit imbalances. In *International Conference on Integer Programming and Combinatorial Optimization*, pages 140–153. Springer, 2022.
- [DLHL15] Jesús A De Loera, Raymond Hemmecke, and Jon Lee. On augmentation algorithms for linear and integer-linear programming: From edmonds–karp to bland and beyond. *SIAM Journal on Optimization*, 25(4):2494–2511, 2015.
- [FT94] Alan M. Frieze and Shang-Hua Teng. On the complexity of computing the diameter of a polytope. *Comput. Complex.*, 4:207–219, 1994.
- [GKM19] Manoj Gupta, Hitesh Kumar, and Neeldhara Misra. On the complexity of optimal matching reconfiguration. In Barbara Catania, Rastislav Královic, Jerzy R. Nawrocki, and Giovanni Pighizzini, editors, *SOFSEM 2019: Theory and Practice of Computer Science - 45th International Conference on Current Trends in Theory and Practice of Computer Science, Nový Smokovec, Slovakia, January 27-30, 2019, Proceedings*, volume 11376 of *Lecture Notes in Computer Science*, pages 221–233. Springer, 2019.
- [GW24] Christoph Grüne and Lasse Wulf. Completeness in the polynomial hierarchy for many natural problems in bilevel and robust optimization. *arXiv preprint arXiv:2311.10540*, 2024.
- [GW25] Christoph Grüne and Lasse Wulf. The complexity of blocking all solutions. *arXiv preprint arXiv:2502.05348*, 2025.
- [IDH⁺11] Takehiro Ito, Erik D. Demaine, Nicholas J. A. Harvey, Christos H. Papadimitriou, Martha Sideri, Ryuhei Uehara, and Yushi Uno. On the complexity of reconfiguration problems. *Theor. Comput. Sci.*, 412(12-14):1054–1065, 2011.
- [IKK⁺22] Takehiro Ito, Naonori Kakimura, Naoyuki Kamiyama, Yusuke Kobayashi, and Yoshio Okamoto. Shortest reconfiguration of perfect matchings via alternating cycles. *SIAM J. Discret. Math.*, 36(2):1102–1123, 2022.
- [Kaf22] Sean Kafer. *Polyhedral diameters and applications to optimization*. PhD thesis, University of Waterloo, 2022.
- [KM72] Victor Klee and George J Minty. How good is the simplex algorithm. *Inequalities*, 3(3):159–175, 1972.
- [KMM12] Marcin Kamiński, Paul Medvedev, and Martin Milanič. Complexity of independent set reconfigurability problems. *Theoretical computer science*, 439:9–15, 2012.

- [KP03] Volker Kaibel and Marc E. Pfetsch. Some algorithmic problems in polytope theory. In Michael Joswig and Nobuki Takayama, editors, *Algebra, Geometry, and Software Systems [outcome of a Dagstuhl seminar]*, pages 23–47. Springer, 2003.
- [KPS19] Sean Kafer, Kanstantsin Pashkovich, and Laura Sanità. On the circuit diameter of some combinatorial polytopes. *SIAM Journal on Discrete Mathematics*, 33(1):1–25, 2019.
- [KV11] Bernhard H Korte and Jens Vygen. *Combinatorial optimization*, volume 1. Springer, 2011.
- [KW67] Victor Klee and David W Walkup. The d-step conjecture for polyhedra of dimension $d \leq 6$. *Acta Math*, 1967.
- [LKS22] Jesús A. De Loera, Sean Kafer, and Laura Sanità. Pivot rules for circuit-augmentation algorithms in linear optimization. *SIAM J. Optim.*, 32(3):2156–2179, 2022.
- [MSW15] Benjamin Matschke, Francisco Santos, and Christophe Weibel. The width of five-dimensional prisms. *Proceedings of the London Mathematical Society*, 110(3):647–672, 2015.
- [Nad89] Denis Naddef. The Hirsch conjecture is true for $(0, 1)$ -polytopes. *Mathematical Programming: Series A and B*, 45(1):109–110, 1989.
- [NS25] Christian Nöbel and Raphael Steiner. Complexity of polytope diameters via perfect matchings. In *Proceedings of the 2025 Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 2234–2251. SIAM, 2025.
- [Pap94] Christos H. Papadimitriou. *Computational complexity*. Addison-Wesley, 1994.
- [Ple79] Ján Plesník. The NP-completeness of the Hamiltonian cycle problem in planar digraphs with degree bound two. *Inf. Process. Lett.*, 8(4):199–201, 1979.
- [San12] Francisco Santos. A counterexample to the Hirsch conjecture. *Annals of mathematics*, pages 383–412, 2012.
- [San18] Laura Sanità. The diameter of the fractional matching polytope and its hardness implications. In Mikkel Thorup, editor, *59th IEEE Annual Symposium on Foundations of Computer Science, FOCS 2018, Paris, France, October 7-9, 2018*, pages 910–921. IEEE Computer Society, 2018.
- [San20] Laura Sanità. Diameter of polytopes: Algorithmic and combinatorial aspects. <https://www.lse.ac.uk/ipco-2020/summer-school/Sanita-complete.pdf>, 2020. Online; accessed 20 February 2025.
- [Sch01] Marcus Schaefer. Graph ramsey theory and the polynomial hierarchy. *J. Comput. Syst. Sci.*, 62(2):290–322, 2001.
- [Sma98] Steve Smale. Mathematical problems for the next century. *The mathematical intelligencer*, 20:7–15, 1998.
- [Sto76] Larry J. Stockmeyer. The polynomial-time hierarchy. *Theor. Comput. Sci.*, 3(1):1–22, 1976.
- [SY15] Tamon Stephen and Timothy Yusun. The circuit diameter of the Klee-Walkup polyhedron. *Electronic Notes in Discrete Mathematics*, 49:505–512, 2015.

- [Tre04] Luca Trevisan. Inapproximability of combinatorial optimization problems. *Electron. Colloquium Comput. Complex.*, TR04-065, 2004.
- [Woe21] Gerhard J. Woeginger. The trouble with the second quantifier. *JOR*, 19(2):157–181, 2021.