

# ON THE FUNCTIONAL MINKOWSKI PROBLEM

TOMER FALAH AND LIRAN ROTEM

ABSTRACT. To every log-concave function  $f$  one may associate a pair of measures  $(\mu_f, \nu_f)$  which are the surface area measures of  $f$ . These are a functional extension of the classical surface area measure of a convex body, and measure how the integral  $\int f$  changes under perturbations. The functional Minkowski problem then asks which pairs of measures can be obtained as the surface area measures of a log-concave function. In this work we fully solve this problem.

Furthermore, we prove that the surface area measures are continuous in correct topology: If  $f_k \rightarrow f$ , then  $(\mu_{f_k}, \nu_{f_k}) \rightarrow (\mu_f, \nu_f)$  in the appropriate sense. Finding the appropriate mode of convergence of the pairs  $(\mu_{f_k}, \nu_{f_k})$  sheds a new light on the construction of functional surface area measures. To prove this continuity theorem we associate to every convex function a new type of radial function, which seems to be an interesting construction on its own right.

Finally, we prove that the solution to functional Minkowski problem is continuous in the data, in the sense that if  $(\mu_{f_k}, \nu_{f_k}) \rightarrow (\mu_f, \nu_f)$  then  $f_k \rightarrow f$  up to translations.

## 1. INTRODUCTION

**1.1. Surface area measures and the Minkowski problem.** The Minkowski problem is one of the most famous problems in convex geometry and differential geometry. We begin with recalling some classical results about it, and about surface area measures in general. For more general background on convex geometry the reader may consult e.g. [31].

The Minkowski sum of two closed convex sets  $K, L \subseteq \mathbb{R}^n$  is defined as

$$K + L = \{x + y : x \in K, y \in L\}.$$

The set  $K + L$  is convex, and is closed if  $K$  or  $L$  are bounded. If in addition  $\lambda > 0$ , we define the dilation  $\lambda K = \{\lambda x : x \in K\}$ . The support function  $h_K : \mathbb{S}^{n-1} \rightarrow (-\infty, \infty]$  of a closed convex set  $K$  is given by  $h_K(y) = \sup_{x \in K} \langle x, y \rangle$ . These constructions interact well with each other, in the sense that  $\lambda K + \mu K = (\lambda + \mu)K$  and  $h_{\lambda K + \mu L} = \lambda h_K + \mu h_L$ . The (Lebesgue) volume of  $K$  will be denoted by  $|K|$ .

By a convex body we mean a compact convex set with non-empty interior, or equivalently a closed convex set  $K \subseteq \mathbb{R}^n$  such that  $0 < |K| < \infty$ . For every convex body  $K$  there exists a finite Borel measure  $S_K$  on the unit sphere  $\mathbb{S}^{n-1} = \{x \in \mathbb{R}^n : |x| = 1\}$  with the following property: For every closed convex set  $L \subseteq \mathbb{R}^n$  we have

$$\lim_{t \rightarrow 0^+} \frac{|K + tL| - |K|}{t} = \int_{\mathbb{S}^{n-1}} h_L dS_K.$$

The measure  $S_K$  is called the *surface area measure* of  $K$ .

The surface area measure  $S_K$  can also be defined explicitly. Let  $n_K : \partial K \rightarrow \mathbb{S}^{n-1}$  be the Gauss map, associating to every boundary point  $x \in \partial K$  the outer unit normal to  $K$  at  $x$ . It is known that this outer unit normal is unique  $\mathcal{H}^{n-1}$ -almost everywhere, where  $\mathcal{H}^{n-1}$  denotes the  $(n-1)$ -dimensional Hausdorff measure. We then have  $S_K = (n_K)_\# (\mathcal{H}^{n-1}|_{\partial K})$ , where  $\#$  denotes the push-forward of a measure. Explicitly, this means that for every (say continuous) function  $\rho : \mathbb{S}^{n-1} \rightarrow \mathbb{R}$  we have

$$\int_{\mathbb{S}^{n-1}} \rho dS_K = \int_{\partial K} (\rho \circ n_K) d\mathcal{H}^{n-1}.$$

For example, when  $K$  is a polytope the measure  $S_K$  is discrete, with atoms corresponding to the outer normals to the facets of  $K$ . When  $\partial K$  is smooth with non-vanishing Gauss curvature the measure  $S_K$  is continuous with respect to the  $\mathcal{H}^{n-1}|_{\mathbb{S}^{n-1}}$ , and  $\frac{dS_K}{d\mathcal{H}^{n-1}}(\theta) = \det D^2 h_K(\theta)$ , which is also the inverse of the Gauss curvature of  $\partial K$  at the point  $n_K^{-1}(\theta)$ . Here  $D^2 h_K = \nabla_{\mathbb{S}^{n-1}}^2 h_K + h_K \cdot \text{Id}$ , where  $\nabla_{\mathbb{S}^{n-1}}^2$  is the Riemannian Hessian on  $\mathbb{S}^{n-1}$ .

The Minkowski problem asks which measures on  $\mathbb{S}^{n-1}$  are the surface area measures of a convex body. Its solution, known as Minkowski's existence theorem, reads as follows:

**Theorem 1.1.** *Let  $\mu$  be a finite Borel measure on  $\mathbb{S}^{n-1}$ . Then  $\mu = S_K$  for some convex body  $K \subseteq \mathbb{R}^n$  if and only if  $\mu$  satisfies the following two conditions:*

- (1)  $\mu$  is centered, i.e.  $\int_{\mathbb{S}^{n-1}} \langle x, \theta \rangle d\mu(x) = 0$  for all  $\theta \in \mathbb{R}^n$ .
- (2)  $\mu$  is not supported on any hyperplane.

Moreover, the body  $K$  is unique up to translations: If  $S_K = S_L$  then  $K = L + v$  for some  $v \in \mathbb{R}^n$ .

We refer the reader to [31] for two proofs of Minkowski's theorem. One is Minkowski's original argument for polytopes, given in Theorem 8.2.1, followed by an approximation argument of Fenchel and Jessen in Theorem 8.2.2. An alternative proof due to Alexandrov appears (for a generalized problem) in Theorem 9.2.1. This latter proof is the one that serves as a motivation for our proof of Theorem 1.5. The reader may also be interested in the notes following Section 8.2 for a more comprehensive history of the problem, including many references.

When the body  $K$  is smooth and  $\mu$  has smooth density  $f$ , the equation  $S_K = \mu$  is equivalent to the Monge–Ampère type partial differential equation

$$\det(D^2 h_K) = f.$$

Theorem (1.1) is then a statement of existence and uniqueness of solutions to this equation, at least in a weak sense.

On the space of convex bodies we have a natural topology, given by the Hausdorff distance

$$\begin{aligned} d_H(K, L) &= \inf \{ r > 0 : K \subseteq L + \overline{B}_r(0) \text{ and } L \subseteq K + \overline{B}_r(0) \} \\ &= \max_{\theta \in \mathbb{S}^{n-1}} |h_K(\theta) - h_L(\theta)|, \end{aligned}$$

where  $\overline{B}_r(0)$  denotes the closed ball of radius  $r$  around the origin. The surface area measure is weakly continuous in this topology: If  $K_i \rightarrow K$  then  $S_{K_i} \rightarrow S_K$  weakly (see e.g. [31, Theorem

4.2.1] for a much more general result). It turns out that the converse to this statement is also true, i.e. the solution  $K$  to the Minkowski problem is continuous in the data  $\mu$ :

**Theorem 1.2.** *Let  $\{K_i\}_{i=1}^\infty, K \subseteq \mathbb{R}^n$  be convex bodies such that  $S_{K_i} \rightarrow S_K$  weakly. Then there exists translations  $\{v_i\}_{i=1}^\infty$  such that  $K_i + v_i \rightarrow K$  in the Hausdorff topology.*

Theorem 1.2 appears to be folklore. Several papers that prove extensions of this result refer to it simply as “well-known” (see e.g. [39], [35], [36]), and we were unable trace its origins. In any case, we are only stating Theorem 1.2 as a motivation and will not need to apply it at any point. A proof can be easily constructed by adapting our proof of Theorem 1.10.

**1.2. Functional surface area measures.** In this work we extend the results above from convex bodies to log-concave functions. Recall that a function  $f : \mathbb{R}^n \rightarrow [0, \infty)$  is called log-concave if for all  $x, y \in \mathbb{R}^n$  and  $0 \leq \lambda \leq 1$  we have

$$f((1 - \lambda)x + \lambda y) \geq f(x)^{1-\lambda} f(y)^\lambda.$$

In other words,  $f$  is log-concave if  $f = e^{-\phi}$  for a convex function  $\phi : \mathbb{R}^n \rightarrow (-\infty, \infty]$ . We will always assume our convex functions are lower semicontinuous and not identically  $+\infty$ , or equivalently that our log-concave functions are upper semicontinuous and not identically 0. For every closed convex set  $K \subseteq \mathbb{R}^n$  its indicator  $\mathbf{1}_K$  is a log-concave function, and we would like to consider log-concave functions as “generalized convex bodies”.

This approach, called “geometrization of probability” by V. Milman (see [24]), proved to be very useful in convexity over the last two decades. Several open problems in convex geometry, which seemed completely intractable using purely geometric tools, were solved by extending the problem to the functional setting and using analytic and probabilistic tools that only make sense when dealing with general log-concave functions. A survey of such results will take us too far from our goal, so let us only mention the recent solution of Bourgain’s slicing problem by Klartag and Lehec ([18]), that would have been completely impossible using purely geometric tools. Surface area measures are related to some of the most important open problems in convex geometry such as the log-Brunn–Minkowski problem (see [3]), and it seems natural that in order to attack such problem we need to develop the theory of functional surface area measures.

We define

$$\text{Cvx}_n = \left\{ \phi : \mathbb{R}^n \rightarrow (-\infty, \infty] : \begin{array}{l} \phi \text{ is convex, lower semicontinuous,} \\ \text{and } 0 < \int e^{-\phi} < \infty \end{array} \right\},$$

and  $\text{LC}_n = \{e^{-\phi} : \phi \in \text{Cvx}_n\}$ . This will be our extension of the class of convex bodies, with the integral  $\int f$  replacing the volume  $|K|$ . Note that if  $0 < \int e^{-\phi} < \infty$  then  $\phi$  is *coercive*, i.e.  $\phi(x) \geq a|x| + b$  for some  $a > 0$  and  $b \in \mathbb{R}$  (see e.g. [5, Lemma 2.5]). The sup-convolution of two log-concave functions  $f, g : \mathbb{R}^n \rightarrow [0, \infty)$  is defined by

$$(f \star g)(x) = \sup_{y \in \mathbb{R}^n} (f(y)g(x - y)).$$

This function is again log-concave, and is upper semi-continuous assuming  $f$  and  $g$  are and at least one of them belongs to  $\text{LC}_n$ . For  $\lambda > 0$  we define the dilation  $\lambda \cdot f$  by  $(\lambda \cdot f)(x) = f\left(\frac{x}{\lambda}\right)^\lambda$ . Finally, the support function of  $f$  is given by  $h_f = (-\log f)^*$ , where

$$\psi^*(y) = \sup_{x \in \mathbb{R}^n} (\langle x, y \rangle - \psi(x))$$

denotes the Legendre transform. There are several good characterization theorems that explain why these are the natural operations on the class of log-concave functions – see e.g. [1] and [27]. For now we just mention that  $(\lambda \cdot f) \star (\mu \cdot f) = (\lambda + \mu) \cdot f$  and that  $h_{(\lambda \cdot f) \star (\mu \cdot f)} = \lambda h_f + \mu h_g$ .

Given these basic operations we can now consider the same first variation as before:

**Definition 1.3.** For  $f \in \text{LC}_n$  and  $g : \mathbb{R}^n \rightarrow \mathbb{R}$  an upper semicontinuous log-concave function (not necessarily with  $0 < \int g < \infty$ ) we set

$$\delta(f, g) = \lim_{t \rightarrow 0^+} \frac{\int (f \star (t \cdot g)) - \int f}{t}.$$

**Theorem 1.4.** For all  $f = e^{-\phi} \in \text{LC}_n$  and every upper semicontinuous log-concave function  $g$  we have

$$(1.1) \quad \delta(f, g) = \int_{\mathbb{R}^n} h_g d\mu_f + \int_{\mathbb{S}^{n-1}} h_{\text{supp}(g)} d\nu_f.$$

Here  $\mu_f$  is the Borel measure on  $\mathbb{R}^n$  defined by  $\mu_f = (\nabla \phi)_\#(f dx)$  and  $\nu_f$  is the Borel measure on  $\mathbb{S}^{n-1}$  defined by  $\nu_f = (n_{\text{supp}(f)})_\#(f d\mathcal{H}^{n-1}|_{\partial \text{supp}(f)})$ . We refer to the pair  $(\mu_f, \nu_f)$  as the surface area measures of  $f$ .

A few comments are in order. First, the measures  $\mu_f$  and  $\nu_f$  are always well defined for all  $f \in \text{LC}_n$ . For  $\mu_f$ , this is because  $\phi$  is a convex function and therefore differentiable almost everywhere on the set  $\{x : \phi(x) < \infty\} = \{x : f(x) > 0\}$ . For  $\nu_f$ , note that the support

$$\text{supp}(f) = \overline{\{x \in \mathbb{R}^n : f(x) > 0\}}$$

is a closed convex set with non-empty interior (that may be unbounded). Therefore the Gauss map  $n_{\text{supp}(f)} : \partial \text{supp}(f) \rightarrow \mathbb{S}^{n-1}$  is well-defined  $\mathcal{H}^{n-1}$ -almost everywhere. Of course we may have  $\text{supp}(f) = \mathbb{R}^n$ , and then  $\partial \text{supp}(f) = \emptyset$ ; In this case we set  $\nu_f \equiv 0$ .

Regarding the history of Theorem 1.4, the first variation  $\delta(f, g)$  was first considered by Klartag and Milman in [19] in the special case  $f(x) = e^{-|x|^2/2}$ . This notion was further studied in [26], where (1.1) was proved for this special choice of a function  $f$ . At the same time and independently, Colesanti and Fragalà studied the first variation  $\delta(f, g)$  for general log-concave functions ([5]). Among other results, they proved formula (1.1) under some strong smoothness and regularity assumptions on the functions  $f$  and  $g$ . Cordero-Erausquin and Klartag ([11]) studied the measure  $\mu_f$  under the name “the moment measure of  $\phi = -\log f$ ”. They concentrated on the case  $\nu_f \equiv 0$ , and while they did not prove (1.1) in this case they proved a result similar in spirit. Note that  $\nu_f \equiv 0$  if and only if  $f \equiv 0$   $\mathcal{H}^{n-1}$ -almost everywhere on  $\partial \text{supp}(f)$ , which is equivalent to saying that  $f$  is continuous

$\mathcal{H}^{n-1}$ -almost everywhere. Therefore [11] named this condition *essential continuity*. Finally, formula (1.1) was proven in full generality by the second named author ([28], [29]).

Finally, we caution the reader that the support function appears in formula (1.1) in two different senses. The function  $h_g : \mathbb{R}^n \rightarrow (-\infty, \infty]$  is the support function in the sense of log-concave functions, i.e.  $h_g = (-\log g)^*$ . However,  $\text{supp}(g)$  is a closed convex set, and its support function  $h_{\text{supp}(g)} : \mathbb{S}^{n-1} \rightarrow \mathbb{R}$  is in the classical meaning. These two support functions are related though: the *horizon function*  $\bar{\psi} : \mathbb{S}^{n-1} \rightarrow (-\infty, \infty]$  of a convex function  $\psi : \mathbb{R}^n \rightarrow (-\infty, \infty]$  is defined as

$$(1.2) \quad \bar{\psi}(\theta) = \lim_{\lambda \rightarrow \infty} \frac{\psi(p + \lambda\theta)}{\lambda},$$

where  $p$  is an arbitrary point such that  $\psi(p) < \infty$ . Then  $\bar{\psi}$  is well-defined and independent of  $p$  (see [25, Theorem 3.21]). We then have  $h_{\text{supp}(g)} = \bar{h}_g$  ([25, Theorem 11.5]).

**1.3. Our main theorems.** We can now state our first main theorem, a complete solution of the Minkowski problem for functional surface area measures:

**Theorem 1.5.** *Let  $\mu$  be a finite Borel measure on  $\mathbb{R}^n$  and  $\nu$  be a finite Borel measure on  $\mathbb{S}^{n-1}$ . Then there exists a log-concave function  $f \in \text{LC}_n$  such that  $(\mu_f, \nu_f) = (\mu, \nu)$  if and only if the pair  $(\mu, \nu)$  satisfies the following conditions:*

- (1)  $\mu$  is not identically 0.
- (2)  $\mu$  has finite first moment, and the measure  $\mu + \nu$  is centered in the sense that for every  $\theta \in \mathbb{S}^{n-1}$  we have

$$\int_{\mathbb{R}^n} \langle x, \theta \rangle d\mu(x) + \int_{\mathbb{S}^{n-1}} \langle x, \theta \rangle d\nu(x) = 0.$$

- (3)  $\mu$  and  $\nu$  are not supported on a common hyperplane  $H \subseteq \mathbb{R}^n$ .

*In this case, the function  $f$  is unique up to translations: If  $g \in \text{LC}_n$  also satisfies  $(\mu_g, \nu_g) = (\mu, \nu)$  then there exists  $v \in \mathbb{R}^n$  such that  $g(x) = f(x + v)$ .*

The uniqueness part of Theorem 1.5 was already known – it was proved in [5] under regularity assumptions on  $f$  and  $g$ , in [11] in the essentially continuous case, and in [29] for general  $f, g \in \text{LC}_n$ . As for the existence part, when  $\nu \equiv 0$  Theorem 1.5 was proved by Cordero-Erausquin and Klartag ([11]), with another proof given by Santambrogio ([30]). Variants of the functional Minkowski problem, changing either the volume functional or the addition operation, were studied by Huang, Liu, Xi and Zhao ([15]), Fang, Ye, Zhang, Zhao ([14]), Fang, Ye, Zhang ([13]) and by the second named author ([28]). However, to the best of our knowledge all known results were partial results, in the sense that they deal with the essentially continuous case  $\nu \equiv 0$ . Dealing with general log-concave functions does require new ideas, as we'll see below.

The functional Minkowski problem may be written as a pair of partial differential equations. Indeed, assume  $\mu$  has density  $\rho_1 : \mathbb{R}^n \rightarrow \mathbb{R}$  and  $\nu$  has density  $\rho_2 : \mathbb{S}^{n-1} \rightarrow \mathbb{R}$ . If  $f = e^{-\phi} \in \text{LC}_n$  is a smooth

function on its smooth support  $K = \text{supp}(f)$ , then the change of variables formula shows that  $\phi$  and  $K$  solve the system

$$(1.3) \quad \begin{cases} \rho_1 (\nabla \phi(x)) \det ((\nabla^2 \phi)(x)) = e^{-\phi(x)} & \text{for all } x \in \text{int}(K) \\ \frac{\rho_2(n_K(x))}{\det((D^2 h_K)(n_K(x)))} = e^{-\phi(x)} & \text{for all } x \in \partial K. \end{cases}$$

Note that the domain  $K$  is not given, but both  $K$  and  $\phi$  are the unknowns in this system of equations. Also note that  $\det ((D^2 h_K)(n_K(x)))^{-1}$  is simply the Gauss curvature of  $K$  at the point  $x \in \partial K$ . Theorem 1.5 then states that under our assumptions on  $\rho_1$  and  $\rho_2$  the system (1.3) has a unique solution, at least in a weak sense. Studying the regularity of our solution and whether it is also a classical solution to (1.3) is an interesting problem beyond the scope of this paper. We refer the reader to [17] for some remarks and references regarding regularity in the case  $\nu \equiv 0$  and to [33] for a related result regarding the regularity of the weighted Minkowski problem.

Next, we discuss continuity of the surface area measures. On the class  $\text{Cvx}_n$  we have a natural notion of convergence called epi-convergence – see Definition 3.2 below and the discussion following it. This gives us a topology on  $\text{LC}_n$  in the obvious way. In [17], Klartag essentially proved that if  $f_k \rightarrow f$  then  $\mu_{f_k} \rightarrow \mu_f$  weakly (He claimed a weaker result, which does not use the notion of epi-convergence, but the same ideas can be used to prove the more general statement). However, the same cannot be true for the boundary measure  $\nu_f$ , as a simple example shows:

**Exercise 1.6.** Define  $\{\phi_k\}_{k=1}^\infty : \mathbb{R} \rightarrow (-\infty, \infty]$  by  $\phi_k(x) = \max(k|x| - k, 0)$ . Then  $\phi_k \rightarrow \phi$ , where

$$\phi(x) = \mathbf{1}_{[-1,1]}^\infty(x) = \begin{cases} 0 & x \in [-1, 1] \\ +\infty & \text{otherwise.} \end{cases}$$

However  $\nu_{e^{-\phi_k}} = 0$  and  $\nu_{e^{-\phi}} = \delta_1 + \delta_{-1} \neq 0$ , so clearly we cannot have  $\nu_{e^{-\phi_k}} \rightarrow \nu_{e^{-\phi}}$  in any reasonable sense.

Indeed, it makes no sense to treat the measures  $\mu_f$  and  $\nu_f$  separately. Instead, the correct claim should be that if  $f_k \rightarrow f$  then  $(\mu_{f_k}, \nu_{f_k}) \rightarrow (\mu, \nu)$  as a pair. We therefore make the following definitions:

**Definition 1.7.** We say a function  $\xi : \mathbb{R}^n \rightarrow \mathbb{R}$  is *cosmically continuous* if:

- (1)  $\xi$  is continuous in the usual sense on  $\mathbb{R}^n$ .
- (2) The limit  $\bar{\xi}(\theta) = \lim_{\lambda \rightarrow \infty} \frac{\xi(\lambda\theta)}{\lambda}$  exists (in the finite sense) *uniformly* in  $\theta \in \mathbb{S}^{n-1}$ .

**Definition 1.8.** Let  $\{\mu_k\}_{k=1}^\infty$  and  $\mu$  be finite Borel measures on  $\mathbb{R}^n$ , and let  $\{\nu_k\}_{k=1}^\infty$  and  $\nu$  be finite Borel measures on  $\mathbb{S}^{n-1}$ . We say that  $(\mu_k, \nu_k) \rightarrow (\mu, \nu)$  *cosmically* if for every cosmically continuous function  $\xi : \mathbb{R}^n \rightarrow \mathbb{R}$  we have

$$\int_{\mathbb{R}^n} \xi d\mu_k + \int_{\mathbb{S}^{n-1}} \bar{\xi} d\nu_k \xrightarrow{k \rightarrow \infty} \int_{\mathbb{R}^n} \xi d\mu + \int_{\mathbb{S}^{n-1}} \bar{\xi} d\nu.$$

A similar class of functions appeared in the work of Ulivelli ([34]) under the notation  $C_{rec}(\mathbb{R}^n)$ , for similar reasons. However, functions in  $C_{rec}(\mathbb{R}^n)$  satisfy the stronger assumption that  $|\xi - \bar{\xi}|$  is bounded, when  $\bar{\xi}$  is considered as a 1-homogeneous function on  $\mathbb{R}^n$ . We explain in Section 4 why our definitions appear to be the natural ones, as well as the origin of the name “cosmic”: The pair of measures  $(\mu, \nu)$  can be naturally identified with a single measure  $\lambda$  on a compactification of  $\mathbb{R}^n$  known as its cosmic closure ([25, Chapter 3A]). Cosmic convergence  $(\mu_k, \nu_k) \rightarrow (\mu, \nu)$  is then nothing more than standard weak convergence  $\lambda_k \rightarrow \lambda$ .

With the correct definition in place, we can state our second main theorem:

**Theorem 1.9.** *Fix  $\{f_k\}_{k=1}^\infty, f \in \text{LC}_n$  such that  $f_k \rightarrow f$ . Then  $(\mu_{f_k}, \nu_{f_k}) \rightarrow (\mu_f, \nu_f)$  cosmically.*

Finally, once we have Theorems 1.5 and 1.9, it is very natural to ask if the converse is also true, and if the solution  $f$  to the functional Minkowski problem is continuous in the data  $(\mu, \nu)$ . This turns out to be true, and is our final main result:

**Theorem 1.10.** *Fix  $\{f_k\}_{k=1}^\infty, f \in \text{LC}_n$  and assume that  $(\mu_{f_k}, \nu_{f_k}) \rightarrow (\mu_f, \nu_f)$  cosmically. Then there exists a sequence of translations  $\tilde{f}_k(x) = f_k(x + v_k)$  such that  $\tilde{f}_k \rightarrow f$ .*

In the very special case where  $\nu_{f_k} \equiv 0$  for all  $k$  and all the measures  $\mu_{f_k}$  are supported on a compact set (the same compact set for all  $k$ ), Theorem 1.10 was previously proved by Klartag ([17]).

**1.4. Proof ideas and the structure of this paper.** Very often, Minkowski type theorems are proven using a variational argument: One defines a functional  $F$ , and proves that  $F$  attains a minimum and that this minimum is the required solution to the Minkowski problem.

In our case, we define  $F : \text{Cvx}_n \rightarrow (-\infty, \infty]$  by

$$F(\phi) = \int_{\mathbb{R}^n} \phi^* d\mu + \int_{\mathbb{S}^{n-1}} \overline{\phi^*} d\nu - \mu(\mathbb{R}^n) \log \int_{\mathbb{R}^n} e^{-\phi}.$$

This is inspired by both Alexandrov’s proof of the classical Minkowski Theorem 1.1, and by the result of [11] that used the same functional in the case  $\nu = 0$  to prove the functional Minkowski theorem in this case.

In order to prove that  $F$  attains a minimum we equip  $\text{Cvx}_n$  with the topology of epi-convergence, and prove that every minimizing sequence for  $F$  must be uniformly coercive. From there we use known compactness results to prove that our minimizing sequence has a convergent sub-sequence, and therefore that  $F$  attains a minimum. We believe that thanks to the use of epi-convergence our argument here is simpler and more transparent than the corresponding argument of [11], even though we are dealing with general measures  $\mu$  and  $\nu$ .

Next, one needs to prove that the minimizer of  $F$  solves the Minkowski problem. This essentially requires computing the derivative  $\frac{d}{dt} \Big|_{t=0} F((\phi^* + t\xi)^*)$  for a large enough class of functions  $\xi : \mathbb{R}^n \rightarrow \mathbb{R}$ . In the case of convex bodies the computation of the corresponding derivative relies on the so-called Alexandrov lemma – see e.g. [31, Lemma 7.5.3]. We therefore need a functional version of this lemma, which is far from trivial. In [34] Ulivelli proved a functional Alexandrov lemma, using

a weighted version of Alexandrov’s lemma from [20]. Unfortunately, Ulivelli’s theorem has some assumptions that make it unsuitable for our goal, most crucially that the convex functions involved have compact domain. Luckily, we do not need to compute the derivative for all  $\xi$ , just for a large enough family of functions. We do so in Section 2, surprisingly using a theorem of Matheron about the Minkowski difference of convex bodies. Then in Section 3 we carry out the proof that  $F$  attains a minimizer and use the results of Section 2 to conclude that this minimizer is the sought after solution to the Minkowski theorem.

Next, in Section 4, we study cosmically continuous functions and cosmic convergence. We explain where these terms come from and how to view the pair  $(\mu_f, \nu_f)$  as a single measure on the cosmic closure of  $\mathbb{R}^n$ . This is crucial for the proof of Theorem 1.9, but we believe it is also of independent interest. For example, it allows us to rewrite the first variation formula (1.1) as

$$(1.4) \quad \delta(e^{-\phi}, g) = \int_{\partial \text{epi}(\phi)} \widehat{h}_g(n_{\text{epi}(\phi)}(x, t)) e^{-t} d\mathcal{H}^n(x, t),$$

where  $\widehat{h}_g$  is essentially the function  $h_g$  extended to the cosmic closure of  $\mathbb{R}^n$ . Note that in this formula we no longer have any explicit “boundary term”. Similar ideas appeared in [34], but as far as we can tell our definitions and the formula above are new.

In Section 5 we prove Theorem 1.9. To explain the main idea behind the proof it is useful to sketch a proof of the weak continuity of the surface area measure for convex bodies. This specific result can also be obtained using mixed volumes, but slightly modifying the problem makes this approach impossible, so various variants of the following argument appeared in the literature (See e.g. [16, Theorem 3.4], [34, Lemma 3.12], and [32, Lemma 6]). Assume  $K_i \rightarrow K$ . We want to prove that  $S_{K_i} \rightarrow S_K$ , or that for all continuous functions  $\rho : \mathbb{S}^{n-1} \rightarrow \mathbb{R}$  we have

$$\int_{\partial K_i} \rho(n_{K_i}(x)) d\mathcal{H}^{n-1}(x) = \int_{\mathbb{S}^{n-1}} \rho dS_{K_i} \rightarrow \int_{\mathbb{S}^{n-1}} \rho dS_K = \int_{\partial K} \rho(n_K(x)) d\mathcal{H}^{n-1}(x).$$

To do so we transform the domain of integration from  $\partial K$  (or  $\partial K_i$ ) to  $\mathbb{S}^{n-1}$  using the change of variables  $\theta \mapsto r_K(\theta)\theta$ , where  $r_K(\theta) = \max\{\lambda > 0 : \lambda\theta \in K\}$  is the radial function of  $K$ . Once all the integrals involved are on the same domain  $\mathbb{S}^{n-1}$ , it is enough to prove that the integrands converge almost everywhere and use the dominated convergence theorem.

We want to argue in a similar way using integrals of the form (1.4), so we need a cleverly chosen change of variables which transforms the domain of integration from  $\partial \text{epi}(\phi)$  to  $\mathbb{R}^n$ . Known constructions do not seem to be appropriate here, so we define a new kind of a radial function which we call the *curvilinear radial function* of  $\phi$ , and show that it can be used to parametrize  $\partial \text{epi}(\phi)$ . Constructing this new radial function and proving its basic properties take up the majority of Section 5, and once this is done Theorem 1.9 follows using the scheme mentioned above. We believe our new radial function can have more applications, for example in the proof of a more general Alexandrov-type lemma.

Finally, in Section 6 we prove Theorem 1.10. The argument uses again a compactness result for  $LC_n$ , Theorem 1.9, and the uniqueness part of Theorem 1.5. This is not very different than known arguments for theorems like Theorem 1.2, but working with log-concave functions does add some



difficulties. In particular, we note as a curious fact that the proof relies on an isoperimetric inequality for log-concave functions, proved in [23] – see Proposition 6.2. Another necessary ingredient is the computation of  $\delta(f, f)$  for  $f \in \text{LC}_n$  – for convex bodies this is trivial since volume is  $n$ -homogeneous, but it is less obvious for log-concave functions, and indeed  $\delta(f, f)$  is not proportional to  $\int f$ . Luckily the computation of  $\delta(f, f)$  was already carried out in [5] – see Proposition 6.1.

**1.5. Acknowledgements.** The authors were supported by ISF grant number 2574/24 and NSF-BSF grant number 2022707.

## 2. A PARTIAL ALEXANDROV LEMMA

In our solution of the Minkowski problem we will need to compute derivatives of the form

$$(2.1) \quad \left. \frac{d}{dt} \right|_{t=0} \int_{\mathbb{R}^n} e^{-(\phi^* + t\xi)^*},$$

where  $\phi \in \text{Cvx}_n$  and  $\xi : \mathbb{R}^n \rightarrow \mathbb{R}$  is a “nice enough” function. The function  $\phi^* + t\xi$  is not necessarily convex, but we still compute its Legendre transform using the standard formula, i.e.

$$(\phi^* + t\xi)^*(y) = \sup_{x \in \mathbb{R}^n} (\langle x, y \rangle - (\phi^*(x) + t\xi(x))).$$

Now that if  $\xi \in \text{Cvx}_n$  and if we define  $f = e^{-\phi}$  and  $g = e^{-\xi^*}$ , then  $e^{-(\phi^* + t\xi)^*} = f \star (t \cdot g)$  for all  $t > 0$ . Therefore in this case the derivative we are trying to compute is exactly the one from the first variation formula (1.1). However, formula (1.1) concerns only positive values of  $t$  (i.e. the right derivative), and for us it would be crucial to have a two-sided derivative.

In the case that  $\phi^*$  is the support function of a convex body and  $\xi$  is continuous and 1-homogeneous, the computation of (2.1) is exactly the well-known Alexandrov lemma, which is a standard ingredient in the proof of the classical Minkowski problem. Our goal is therefore to prove a functional extension of this lemma. As was already mentioned in the introduction such a functional Alexandrov lemma was recently proved in [34], but its assumptions are not suitable for our goals.

Computing the derivative (2.1) for *all* continuous functions  $\xi : \mathbb{R}^n \rightarrow \mathbb{R}$  is a non-trivial task, which we do not need for our goal so we will not carry out here. Instead we will only discuss two partial cases. The first case is relatively straightforward and was already computed in [29, Proposition 5.4]:

**Proposition 2.1.** *Fix  $\phi \in \text{Cvx}_n$  and define  $f = e^{-\phi}$ . Assume that  $\xi : \mathbb{R}^n \rightarrow \mathbb{R}$  is **bounded** and **continuous**. Then*

$$\left. \frac{d}{dt} \right|_{t=0} \int_{\mathbb{R}^n} e^{-(\phi^* + t\xi)^*} = \int_{\mathbb{R}^n} \xi d\mu_f.$$

The second case we will need is the case where  $\xi = h_L$  for a convex body  $L \subseteq \mathbb{R}^n$ . For this case we need to recall the definition of the *Minkowski difference*: For convex bodies  $A, B \subseteq \mathbb{R}^n$  their Minkowski difference is defined by

$$A \ominus B = \{x \in \mathbb{R}^n : x + B \subseteq A\} = \bigcap_{x \in B} (A - x).$$

In [22], Matheron proved the following facts about the Minkowski difference:

**Lemma 2.2.** *Fix convex bodies  $A, B \subseteq \mathbb{R}^n$  and consider the function*

$$\beta(t) = \begin{cases} |A + tB| & t \geq 0 \\ |A \ominus (|t|B)| & t < 0, \end{cases}$$

(defined on a ray  $(-t_0, \infty)$  such that  $\beta(t) > 0$  on  $(-t_0, \infty)$ ). Then:

- (1)  $\beta$  is (two-sided) differentiable at  $t = 0$ .
- (2)  $\beta$  is convex, so in particular  $|A + tB| - |A| \geq |A| - |A \ominus tB|$  for all small enough  $t > 0$ .

Using this lemma we prove:

**Theorem 2.3.** *Fix  $\phi \in \text{Cvx}_n$  and define  $f = e^{-\phi}$ . Then for all convex bodies  $L \subseteq \mathbb{R}^n$  we have*

$$(2.2) \quad \left. \frac{d}{dt} \right|_{t=0} \int_{\mathbb{R}^n} e^{-(\phi^* + th_L)^*} = \int_{\mathbb{R}^n} h_L d\mu_f + \int_{\mathbb{S}^{n-1}} h_L d\nu_f.$$

*Proof.* Recall that  $e^{-(\phi^* + th_L)^*} = f \star (t \cdot \mathbf{1}_L)$ , so the first variation formula (1.1) implies that

$$\left. \frac{d}{dt} \right|_{t=0+} \int_{\mathbb{R}^n} e^{-(\phi^* + th_L)^*} = \int_{\mathbb{R}^n} h_L d\mu_f + \int_{\mathbb{S}^{n-1}} h_L d\nu_f.$$

It is therefore enough to prove that the two-sided derivative exists, and we can save ourselves a bit of work by not re-computing this derivative.

Write  $f_t = e^{-(\phi^* + th_L)^*} = f \star \mathbf{1}_{tL}$ . For  $t > 0$  we have

$$f_t(x) = \sup_{y \in \mathbb{R}^n} (f(x - y) \mathbf{1}_{tL}(y)) = \max_{y \in tL} f(x - y).$$

If on the other hand  $t = -s < 0$  then we have

$$\begin{aligned} (\phi^* + th_L)^*(x) &= (\phi^* - h_{sL})^*(x) = \sup_{y \in \mathbb{R}^n} (\langle x, y \rangle - \phi^*(y) + h_{sL}(y)) \\ &= \sup_{y \in \mathbb{R}^n} \sup_{z \in sL} (\langle x, y \rangle - \phi^*(y) + \langle z, y \rangle) \\ &= \sup_{z \in sL} \sup_{y \in \mathbb{R}^n} (\langle x + z, y \rangle - \phi^*(y)) \\ &= \sup_{z \in sL} \phi^{**}(x + z) = \sup_{z \in sL} \phi(x + z), \end{aligned}$$

and so  $f_t(x) = \inf_{z \in sL} f(x + z)$ .

Let us rewrite these identities in terms of level sets. For  $u > 0$  we define

$$K_u = \{x \in \mathbb{R}^n : f(x) \geq u\},$$

a set we also denote by  $[f \geq u]$ . Then for  $t > 0$  we have

$$\begin{aligned} [f_t \geq u] &= \{x \in \mathbb{R}^n : \exists y \in tL, f(x - y) \geq u\} \\ &= \{x \in \mathbb{R}^n : \exists y \in tL, x - y \in K_u\} = K_u + tL, \end{aligned}$$

while for  $t = -s < 0$  we have

$$\begin{aligned} [f_t \geq u] &= \{x \in \mathbb{R}^n : \forall y \in sL, f(x+y) \geq u\} \\ &= \{x \in \mathbb{R}^n : \forall y \in sL, x+y \in K_u\} = K_u \ominus sL. \end{aligned}$$

We therefore consider the function

$$\beta_u(t) = \begin{cases} |K_u + tL| & t \geq 0 \\ |K_u \ominus sL| & t = -s < 0. \end{cases}$$

Write  $M = \max f$ , and note that by layer cake decomposition we have

$$\lim_{t \rightarrow 0} \frac{\int f_t - \int f}{t} = \lim_{t \rightarrow 0} \frac{\int_0^M |[f_t \geq u]| du - \int_0^M |[f \geq u]| du}{t} = \lim_{t \rightarrow 0} \int_0^M \frac{\beta_u(t) - \beta_u(0)}{t} du,$$

By Lemma 2.2(1) we know that

$$\lim_{t \rightarrow 0} \frac{\beta_u(t) - \beta_u(0)}{t}$$

exists. By the second part of the lemma we know that for all  $|t| < t_0$  we have

$$\left| \frac{\beta_u(t) - \beta_u(0)}{t} \right| \leq \frac{\beta_u(|t|) - \beta_u(0)}{|t|} \leq \frac{\beta_u(t_0) - \beta_u(0)}{t_0} < \infty,$$

so we may apply the bounded convergence theorem and exchange the limit and the integral (For the second inequality we used the known fact that for  $t > 0$  the function  $t \mapsto \frac{|K_u + tL| - |K_u|}{t}$  is increasing, which can be proven by writing  $|K_u + tL|$  as its Steiner polynomial – See e.g. Section 4.1 of [31]).

We conclude that

$$\lim_{t \rightarrow 0} \frac{\int f_t - \int f}{t} = \int_0^M \beta'_u(0) du,$$

so in particular the two-sided derivative exists and the proof is complete.  $\square$

### 3. THE MINKOWSKI PROBLEM

We now begin our proof of Theorem 1.5. We begin by showing that the conditions on the measures  $\mu$  and  $\nu$  are indeed necessary:

**Proposition 3.1.** *For every  $f \in \text{LC}_n$  we have:*

- (1)  $\mu_f$  is not identically 0.
- (2)  $\mu_f$  has finite first moment and for all  $\theta \in \mathbb{S}^{n-1}$  we have

$$\int_{\mathbb{R}^n} \langle x, \theta \rangle d\mu_f(x) + \int_{\mathbb{S}^{n-1}} \langle x, \theta \rangle d\nu_f(x) = 0.$$

- (3)  $\mu_f$  and  $\nu_f$  are not supported on a common hyperplane  $H \subseteq \mathbb{R}^n$ .

*Proof.* Property (1) is obvious since  $\mu_f(\mathbb{R}^n) = \int f > 0$ . The fact that  $\mu_f$  has a finite first moment (and that  $\mu_f$  and  $\nu_f$  are finite measures) was shown in [29, Proposition 1.6].

We will show the remaining part of property (2), as well as property (3), by choosing an appropriate function  $g$  in the variation formula (1.1). It will be convenient to use [29, Proposition 2.4]: If  $g = \mathbf{1}_L$  for a compact convex set  $L \subseteq \mathbb{R}^n$  then

$$\delta(f, g) = n \int_0^\infty V_1(F_s, L) ds,$$

where  $F_s = \{x \in \mathbb{R}^n : f(x) \geq s\}$  and  $V_1$  denotes the first mixed volume, i.e.

$$n \cdot V_1(K, L) = \lim_{t \rightarrow 0^+} \frac{|K + tL| - |K|}{t}.$$

Choosing  $L = \{\theta\}$  and  $g = \mathbf{1}_L$  we clearly have  $h_g(x) = \langle x, \theta \rangle$  and  $V_1(K, L) = 0$  for all  $K$ . Therefore

$$0 = \delta(f, g) = \int_{\mathbb{R}^n} \langle x, \theta \rangle d\mu_f(x) + \int_{\mathbb{S}^{n-1}} \langle x, \theta \rangle d\nu_f(x),$$

which proves (2).

To show (3), assume by contradiction that  $\mu_f$  and  $\nu_f$  are supported on  $H = \{x : \langle x, \theta \rangle = 0\}$ . Choose  $L = [-\theta, \theta]$  and  $g = h_L$ . Since  $h_g(x) = h_L(x) = |\langle x, \theta \rangle|$  it follows from our assumption and the variation formula (1.1) that

$$n \int_0^\infty V_1(F_s, [-\theta, \theta]) ds = \delta(f, g) = \int_{\mathbb{R}^n} |\langle x, \theta \rangle| d\mu_f(x) + \int_{\mathbb{S}^{n-1}} |\langle x, \theta \rangle| d\nu_f(x) = 0.$$

Therefore we must have  $V_1(F_s, [-\theta, \theta]) = 0$  for all  $s > 0$ . However it is known that  $V_1(K, [-\theta, \theta]) = \frac{2}{n} |\text{Proj}_{\theta^\perp} K|$ , where  $\text{Proj}$  denotes the orthogonal projection – this is for example a special case of [31, Theorem 5.3.1]. Therefore  $|\text{Proj}_{\theta^\perp} F_s| = 0$  for all  $s > 0$ , which implies that  $F_s$  has an empty interior so  $|F_s| = 0$  as well. But then  $\int f = \int_0^\infty |F_s| ds = 0$ , which is the required contradiction.  $\square$

We now begin our proof that these conditions are also sufficient for the existence of a solution to the Minkowski problem. Towards this goal we fix  $\mu$  and  $\nu$  that satisfy the conditions of Theorem 1.5 and define a functional  $F : \text{Cvx}_n \rightarrow (-\infty, \infty]$  by

$$(3.1) \quad F(\phi) = \int_{\mathbb{R}^n} \phi^* d\mu + \int_{\mathbb{S}^{n-1}} \overline{\phi^*} d\nu - \mu(\mathbb{R}^n) \log \int_{\mathbb{R}^n} e^{-\phi}.$$

We will show that  $F$  attains a minimum at some function  $\phi_0 \in \text{Cvx}_n$  and that our sought after solution is  $f = ce^{-\phi_0}$  for some  $c > 0$ . In order to prove that  $F$  attains a minimum it will be useful to equip the space  $\text{Cvx}_n$  with a topology:

**Definition 3.2.** Fix  $\{\phi_k\}_{k=1}^\infty, \phi \in \text{Cvx}_n$ . We say that  $\phi_k$  epi-converges to  $\phi$  as  $k \rightarrow \infty$  if:

- (1) For all  $x \in \mathbb{R}^n$  and all sequences  $x_k \rightarrow x$  we have  $\liminf_{k \rightarrow \infty} \phi_k(x_k) \geq \phi(x)$ .
- (2) For all  $x \in \mathbb{R}^n$  there exists a sequence  $x_k \rightarrow x$  such that  $\limsup_{k \rightarrow \infty} \phi_k(x_k) \leq \phi(x)$ .

In this case we simply write  $\phi_k \xrightarrow{k \rightarrow \infty} \phi$  or even  $\phi_k \rightarrow \phi$ .

The notion of epi-convergence is well-known in fields like convex analysis and optimization, and goes back to the work of Wijsman ([37, 38]) – see e.g. [25] for many results on epi-convergence and

historical remarks. In particular, to explain the name epi-convergence, recall that the epigraph of  $\phi \in \text{Cvx}_n$  is given by

$$(3.2) \quad \text{epi}(\phi) = \{(x, t) \in \mathbb{R}^n \times \mathbb{R} : \phi(x) \leq t\} \subseteq \mathbb{R}^{n+1}.$$

Then  $\phi_k \rightarrow \phi$  if and only if  $\text{epi}(\phi_k) \rightarrow \text{epi}(\phi)$  in the sense of Painlevé–Kuratowski ([25, Proposition 7.2]). For functions in  $\text{Cvx}_n$  this just means that  $\text{epi}(\phi_k) \cap \overline{B}_R(0) \rightarrow \text{epi}(\phi) \cap \overline{B}_R(0)$  in the Hausdorff metric for all large enough  $R > 0$  ([25, Exercise 4.16]).

In the field of convex geometry the notion of epi-convergence was used by Colesanti, Ludwig and Mussnig in [6] and subsequent papers ([10], [8], [7], [9]). These works present compelling evidence that epi-convergence should be used as the functional extension of the Hausdorff topology. Our work can be viewed as further evidence in this direction. Indeed, as mentioned above in the case  $\nu \equiv 0$  our functional  $F$  coincides with the one used by Cordero-Erausquin and Klartag ([11]). However, they only worked with pointwise convergence, and we believe that our use of epi-convergence in our proof leads to significant simplifications of the proof even in this special case.

One advantage of using epi-convergence is that there are known compactness theorems for this topology, serving as functional analogues of the Blaschke selection theorem. We will use the following formulation, which is due to Li and Mussnig ([21]):

**Theorem 3.3** ([21, Theorem 2.15]). *Fix a sequence  $\{\phi_k\}_{k=1}^\infty \subseteq \text{Cvx}_n$  such that:*

- (1)  $\sup_k (\min \phi_k) < \infty$ .
- (2) *The sequence is uniformly coercive: There exists  $a > 0$  and  $b \in \mathbb{R}^n$  such that  $\phi_k(x) \geq a|x| + b$  for all  $k$  and all  $x \in \mathbb{R}^n$ .*

*Then there exists a sub-sequence  $\{\phi_{k_\ell}\}_{\ell=1}^\infty$  such that  $\phi_{k_\ell} \rightarrow \phi$  for a coercive, lower semicontinuous convex function  $\phi : \mathbb{R}^n \rightarrow (-\infty, \infty]$ .*

As a technical point, it may happen that  $\int e^{-\phi} = 0$ , so  $\phi$  does not have to belong to  $\text{Cvx}_n$ . In our applications of Theorem 3.3 it will be easy to rule this option out.

We can now state the main technical lemma required to prove that  $F$  admits a minimizer:

**Lemma 3.4.** *Assume  $\mu$  and  $\nu$  satisfy the assumptions of Theorem 1.5. Then:*

- (1) *There exists a constant  $c > 0$  depending on  $\mu$  and  $\nu$  such that*

$$(3.3) \quad \int_{\mathbb{R}^n} |\langle x, y \rangle| d\mu(x) + \int_{\mathbb{S}^{n-1}} |\langle x, y \rangle| d\nu(x) \geq c|y|$$

*for all  $y \in \mathbb{R}^n$ .*

- (2) *Assume  $c > 0$  satisfies (3.3). Then for every  $\phi \in \text{Cvx}_n$  with  $\min \phi = \phi(0)$  and for all  $x \in \mathbb{R}^n$  we have*

$$(3.4) \quad \phi(x) \geq \frac{1}{\mu(\mathbb{R}^n)} \left( \frac{c}{2}|x| - \int_{\mathbb{R}^n} \phi^* d\mu - \int_{\mathbb{S}^{n-1}} \overline{\phi^*} d\nu \right).$$

*Proof.* For the first part, it is enough to observe that the function

$$p(y) = \int_{\mathbb{R}^n} |\langle x, y \rangle| d\mu(x) + \int_{\mathbb{S}^{n-1}} |\langle x, y \rangle| d\nu(x)$$

is a norm on  $\mathbb{R}^n$ . Indeed, it is clearly a semi-norm as the sum of semi-norms, and if  $p(y) = 0$  for  $y \neq 0$  then  $\mu$  and  $\nu$  are supported on  $y^\perp$  which contradicts our assumption. Since all norms on  $\mathbb{R}^n$  are equivalent it follows that  $p(y) \geq c|y|$  for some  $c > 0$ .

For the second assertion, set

$$G(\phi) = \int_{\mathbb{R}^n} \phi^* d\mu + \int_{\mathbb{S}^{n-1}} \overline{\phi^*} d\nu.$$

For all  $\lambda \in \mathbb{R}$  we have  $(\phi + \lambda)^* = \phi^* - \lambda$ , and therefore  $G(\phi + \lambda) = G(\phi) - \lambda\mu(\mathbb{R}^n)$ . It follows that the validity of (3.4) doesn't change when  $\phi$  is replaced by  $\phi + \lambda$ , so we may assume without loss of generality that  $\min \phi = \phi(0) = 0$ .

Fix  $x_0 \in \mathbb{R}^n$  with  $\phi(x_0) < \infty$ , and consider the function

$$\rho(x) = \begin{cases} t \cdot \phi(x_0) & x = tx_0, 0 \leq t \leq 1 \\ \infty & \text{otherwise.} \end{cases}$$

Since  $\phi$  is convex and  $\phi(0) = 0$  we clearly have  $\phi \leq \rho$ , and therefore  $\phi^* \geq \rho^*$ . A direct computation gives

$$\begin{aligned} \rho^*(x) &= \sup_{0 \leq t \leq 1} (\langle x, tx_0 \rangle - \rho(tx_0)) \\ &= \sup_{0 \leq t \leq 1} [t(\langle x, x_0 \rangle - \phi(x_0))] = [\langle x, x_0 \rangle - \phi(x_0)]_+, \end{aligned}$$

where  $a_+ = \max(a, 0)$ . Therefore  $\overline{\rho^*}(x) = \langle x, x_0 \rangle_+$ . We can now bound  $G(\phi)$  from below as

$$\begin{aligned} G(\phi) &\geq G(\rho) = \int_{\mathbb{R}^n} [\langle x, x_0 \rangle - \phi(x_0)]_+ d\mu(x) + \int_{\mathbb{S}^{n-1}} \langle x, x_0 \rangle_+ d\nu(x) \\ &\geq \int_{\mathbb{R}^n} \langle x, x_0 \rangle_+ d\mu(x) + \int_{\mathbb{S}^{n-1}} \langle x, x_0 \rangle_+ d\nu(x) - \phi(x_0) \cdot \mu(\mathbb{R}^n), \end{aligned}$$

where in the last inequality we used the fact that  $(a - b)_+ \geq a_+ - b$  whenever  $b \geq 0$ . Using the fact that  $a_+ = \frac{a + |a|}{2}$  and that  $\mu + \nu$  is centered we obtain

$$\begin{aligned} G(\phi) &\geq \int_{\mathbb{R}^n} \frac{\langle x, x_0 \rangle + |\langle x, x_0 \rangle|}{2} d\mu(x) + \int_{\mathbb{S}^{n-1}} \frac{\langle x, x_0 \rangle + |\langle x, x_0 \rangle|}{2} d\nu(x) - \phi(x_0) \cdot \mu(\mathbb{R}^n) \\ &= \frac{1}{2} \int_{\mathbb{R}^n} |\langle x, x_0 \rangle| d\mu(x) + \frac{1}{2} \int_{\mathbb{S}^{n-1}} |\langle x, x_0 \rangle| d\nu(x) - \phi(x_0) \mu(\mathbb{R}^n) \geq \frac{c}{2} |x_0| - \phi(x_0) \mu(\mathbb{R}^n), \end{aligned}$$

and rearranging we obtain  $\phi(x_0) \geq \frac{1}{\mu(\mathbb{R}^n)} \left( \frac{c}{2} |x_0| - G(\phi) \right)$  as claimed.  $\square$

We will also need the following simple invariance property:

**Lemma 3.5.** *Assume  $\mu + \nu$  is centered, and let  $F$  be the functional defined by (3.1). Fix  $\phi \in \text{Cvx}_n$ ,  $v \in \mathbb{R}^n$  and  $b \in \mathbb{R}$  and define  $\psi(x) = \phi(x + v) + b$ . Then  $F(\psi) = F(\phi)$ .*

*Proof.* We have

$$\begin{aligned}\psi^*(x) &= \sup_{y \in \mathbb{R}^n} [\langle x, y \rangle - \phi(y + v) - b] = \sup_{z \in \mathbb{R}^n} [\langle x, z - v \rangle - \phi(z) - b] \\ &= \phi^*(x) - \langle x, v \rangle - b.\end{aligned}$$

Therefore  $\overline{\psi^*}(x) = \overline{\phi^*}(x) - \langle x, v \rangle$ , and we may compute

$$\begin{aligned}F(\psi) &= \int_{\mathbb{R}^n} (\phi^* - \langle x, v \rangle - b) d\mu + \int_{\mathbb{S}^{n-1}} (\overline{\phi^*} - \langle x, v \rangle) d\nu - \mu(\mathbb{R}^n) \log \int_{\mathbb{R}^n} e^{-\phi-b} \\ &\stackrel{(*)}{=} \int_{\mathbb{R}^n} \phi^* d\mu - b\mu(\mathbb{R}^n) + \int_{\mathbb{S}^{n-1}} \overline{\phi^*} d\nu - \mu(\mathbb{R}^n) \left( \log \left( \int_{\mathbb{R}^n} e^{-\phi} \right) - b \right) = F(\phi)\end{aligned}$$

as claimed. Note that in the equality (\*) we used the fact that  $\mu + \nu$  is centered.  $\square$

We are ready to prove:

**Proposition 3.6.** *Assume  $\mu$  and  $\nu$  satisfy the assumptions of Theorem 1.5. Then the functional  $F : \text{Cvx}_n \rightarrow (-\infty, \infty]$  defined by (3.1) attains a minimum on  $\text{Cvx}_n$ .*

*Proof.* Choose a minimizing sequence  $\{\phi_k\}_{k=1}^\infty$ , i.e. a sequence such that  $F(\phi_k) \rightarrow \inf F$ . Using the invariance property of Lemma 3.5 we may assume that  $\int e^{-\phi_k} = 1$  and that  $\min \phi_k = \phi_k(0)$ . We may then apply Lemma 3.4 and deduce that

$$\phi_k(x) \geq \frac{1}{\mu(\mathbb{R}^n)} (c|x| - F(\phi_k))$$

for a constant  $c > 0$  independent of  $k$ . Since  $\{\phi_k\}_{k=1}^\infty$  is a minimizing sequence clearly  $\{F(\phi_k)\}_{k=1}^\infty$  is bounded from above, so it follows that  $\{\phi_k\}_{k=1}^\infty$  is uniformly coercive:  $\phi_k(x) \geq a|x| + b$  for all  $k$  and all  $x \in \mathbb{R}^n$ , where the constants  $a$  and  $b$  do not depend on  $k$ .

Next we claim that  $\sup_k (\min \phi_k) = \sup_k (\phi_k(0)) < \infty$ . Indeed, if this is not the case then there exists a subsequence  $\{\phi_{k_j}\}_{j=1}^\infty$  such that  $\min \phi_{k_j} \rightarrow \infty$ . But then

$$1 = \lim_{j \rightarrow \infty} \int e^{-\phi_{k_j}} \stackrel{(*)}{=} \int e^{-\lim_{j \rightarrow \infty} \phi_{k_j}} = \int 0 = 0,$$

which is a contradiction. The exchange of limit and integral in (\*) is justified by the dominated convergence theorem, since all the functions  $e^{-\phi_k}$  are bounded by the integrable function  $e^{-(a|x|+b)}$ .

It now follows from Theorem 3.3 that we may pass to a subsequence and assume without loss of generality that  $\phi_k \rightarrow \phi$  for a coercive, lower semicontinuous convex function  $\phi : \mathbb{R}^n \rightarrow (-\infty, \infty]$ . By Lemma [6, Lemma 15] we have  $\int e^{-\phi} = \lim_{k \rightarrow \infty} \int e^{-\phi_k} = 1$ , so  $\phi \in \text{Cvx}_n$ . It was shown already by Wijsman ([38, Theorem 6.2], see also [25, Theorem 11.34]) that we also have  $\phi_k^* \rightarrow \phi^*$ , and then by e.g. [25, Theorem 7.53] we also have  $\overline{\phi_k^*} \rightarrow \overline{\phi^*}$  (to make sense of this last convergence either extend the definition of epi-convergence to functions defined on  $\mathbb{S}^{n-1}$  in the obvious way, or consider the functions  $\overline{\phi_k^*}$  as 1-homogeneous functions on  $\mathbb{R}^n$ ). In particular, we have the pointwise estimate  $\phi^*(x) \leq \liminf_{k \rightarrow \infty} \phi_k^*(x)$  and similarly  $\overline{\phi^*}(x) \leq \liminf_{k \rightarrow \infty} \overline{\phi_k^*}(x)$ .

We note that the functions  $\{\phi_k^*\}$  are uniformly bounded from below, since

$$\inf_k (\inf \phi_k^*) = \inf_k (-\phi_k^{**}(0)) = -\sup_k \phi_k(0) > -\infty.$$

It follows that also  $\overline{\phi_k^*} \geq 0$ , and we may apply Fatou's lemma and deduce that

$$\begin{aligned} F(\phi) &\leq \int_{\mathbb{R}^n} \left( \liminf_{k \rightarrow \infty} \phi_k^* \right) d\mu + \int_{\mathbb{S}^{n-1}} \left( \liminf_{k \rightarrow \infty} \overline{\phi_k^*} \right) d\nu \\ &\leq \liminf_{k \rightarrow \infty} \left( \int_{\mathbb{R}^n} \phi_k^* d\mu + \int_{\mathbb{S}^{n-1}} \overline{\phi_k^*} d\nu \right) = \lim_{k \rightarrow \infty} F(\phi_k) = \inf F. \end{aligned}$$

Therefore  $\phi$  is the required minimizer.  $\square$

Using the existence of minimizer and the results of the previous section we can now prove Theorem 1.5:

*Proof of Theorem 1.5.* The fact that the conditions of the theorem are necessary is exactly Proposition 3.1, and the fact that the solution to the Minkowski problem is unique up to translations was previously proved in [29, Corollary 3.3]. Therefore we only need to prove the existence of a solution under our assumptions on  $\mu$  and  $\nu$ .

Let  $\phi$  be the minimizer of  $F$  over  $\text{Cvx}_n$ , whose existence is guaranteed by Proposition 3.6. By Lemma 3.5 we may add a constant to  $\phi$  and assume that  $\int_{\mathbb{R}^n} e^{-\phi} = \mu(\mathbb{R}^n)$ . Define  $f = e^{-\phi}$ . We will prove that  $f$  is the required solution, that is  $\mu_f = \mu$  and  $\nu_f = \nu$ .

Fix a continuous function  $\xi : \mathbb{R}^n \rightarrow \mathbb{R}$ , which is either bounded or the support function of a convex body. For  $t \in \mathbb{R}$  we define  $\phi_t = (\phi^* + t\xi)^* \in \text{Cvx}_n$ . Since  $\phi$  is a minimizer of  $F$  we have  $F(\phi_t) \geq F(\phi)$  for all  $t$ , so the function

$$\alpha(t) = F(\phi_t) = \int_{\mathbb{R}^n} \phi_t^* d\mu + \int_{\mathbb{S}^{n-1}} \overline{\phi_t^*} d\nu - \mu(\mathbb{R}^n) \log \int_{\mathbb{R}^n} e^{-\phi_t}$$

attains a minimum at  $t = 0$ . Since  $\phi_t^* = (\phi^* + t\xi)^{**} \leq \phi^* + t\xi$  it follows that the function

$$\beta(t) = \int_{\mathbb{R}^n} (\phi^* + t\xi) d\mu + \int_{\mathbb{S}^{n-1}} (\overline{\phi^*} + t\overline{\xi}) d\nu - \mu(\mathbb{R}^n) \log \int_{\mathbb{R}^n} e^{-\phi_t}$$

satisfies  $\alpha(t) \leq \beta(t)$  for all  $t$ , and  $\alpha(0) = \beta(0)$ . Therefore the function  $\beta$  also attains a minimum at  $t = 0$ , and  $\beta'(0) = 0$  whenever  $\beta$  is differentiable at  $t = 0$ .

Assume  $\xi$  is continuous and bounded, so  $\overline{\xi} = 0$ . It follows from Proposition 2.1 that  $\beta$  is differentiable at the origin, and that

$$0 = \beta'(0) = \int_{\mathbb{R}^n} \xi d\mu - \mu(\mathbb{R}^n) \cdot \frac{1}{\int_{\mathbb{R}^n} e^{-\phi}} \int_{\mathbb{R}^n} \xi d\mu_f = \int_{\mathbb{R}^n} \xi d\mu - \int_{\mathbb{R}^n} \xi d\mu_f.$$

Since this holds for all bounded and continuous  $\xi$ , it follows from the Riesz representation theorem that  $\mu_f = \mu$ .



Now assume  $\xi$  is a support function of a convex body, so  $\bar{\xi} = \xi$ . It follows from Theorem 2.3 that  $\beta$  is again differentiable at  $t = 0$  and

$$0 = \beta'(0) = \int_{\mathbb{R}^n} \xi d\mu + \int_{\mathbb{S}^{n-1}} \xi d\nu - \frac{\mu(\mathbb{R}^n)}{\int_{\mathbb{R}^n} e^{-\phi}} \left( \int_{\mathbb{R}^n} \xi d\mu_f + \int_{\mathbb{S}^{n-1}} \xi d\nu_f \right).$$

Using the fact that  $\mu(\mathbb{R}^n) = \int_{\mathbb{R}^n} e^{-\phi}$  and that  $\mu = \mu_f$  we deduce that for all support functions  $\xi$  we have

$$(3.5) \quad \int_{\mathbb{S}^{n-1}} \xi d\nu = \int_{\mathbb{S}^{n-1}} \xi d\nu_f.$$

By linearity (3.5) also holds for all differences of support functions, a class that includes all  $C^2$ -smooth functions on  $\mathbb{S}^{n-1}$ . As  $C^2$  functions are dense in the space of continuous functions (with the supremum norm) it follows that (3.5) holds for continuous functions  $\xi : \mathbb{S}^{n-1} \rightarrow \mathbb{R}$ , and so again by Riesz we have  $\nu = \nu_f$ . This completes the proof.  $\square$

#### 4. COSMIC CONVERGENCE

We now turn our attention to the continuity of functional surface area measures. To discuss continuity we need appropriate notions of convergence both on the class  $\text{LC}_n$  and on pairs of measures of the form  $(\mu_f, \nu_f)$ . The natural identification between  $\text{Cvx}_n$  and  $\text{LC}_n$  gives us a notion of convergence on  $\text{LC}_n$ :

**Definition 4.1.** Fix  $\{f_k\}_{k=1}^\infty, f \in \text{LC}_n$ . We say that  $f_k \rightarrow f$  if  $(-\log f_k) \rightarrow (-\log f)$  in the sense of epi-convergence.

Since the map  $x \mapsto -\log x$  is decreasing, our convergence on  $\text{LC}_n$  is not by itself epi-convergence, but the symmetric notion of hypo-convergence. Nonetheless we use the same simple notation  $f_k \rightarrow f$ , which should not cause any confusion.

For pairs of measures we use the (new) notion of cosmic convergence as was defined in Definition 1.8 of the introduction. Our first goal is to explain this convergence, and in particular to explain the name ‘‘cosmic convergence’’. Consider the embedding of  $\mathbb{R}^n$  into  $\mathbb{R}^{n+1}$  as  $\mathbb{R}^n \cong \mathbb{R}^n \times \{-1\} \subseteq \mathbb{R}^{n+1}$ . Also consider the lower half-sphere

$$\mathbb{S}_-^n = \{y = (y_1, y_2, \dots, y_{n+1}) \in \mathbb{S}^n : y_{n+1} < 0\} \subseteq \mathbb{R}^n.$$

We write a general point in  $\mathbb{R}^{n+1}$  as  $(x, t)$  for  $x \in \mathbb{R}^n$  and  $t \in \mathbb{R}$ . The spaces  $\mathbb{R}^n \cong \mathbb{R}^n \times \{-1\}$  and  $\mathbb{S}_-^n$  are then homeomorphic using the gnomonic projection  $\Phi : \mathbb{S}_-^n \rightarrow \mathbb{R}^n$  defined by  $\Phi(x, t) = -\frac{x}{t} = \frac{x}{|t|}$ . The inverse map  $\Phi^{-1} : \mathbb{R}^n \rightarrow \mathbb{S}_-^n$  is defined by

$$\Phi^{-1}(x) = \left( \frac{x}{\sqrt{1 + |x|^2}}, -\frac{1}{\sqrt{1 + |x|^2}} \right).$$

Now the space  $\mathbb{S}_-^n$  has a natural compactification, which is its usual closure  $\overline{\mathbb{S}_-^n}$  in  $\mathbb{R}^{n+1}$ , i.e. the closed lower half-sphere. Intuitively this is a compactification of  $\mathbb{R}^n$  with a point at infinity ‘‘in

every direction". This compactification is known as the cosmic closure of  $\mathbb{R}^n$ , and its identification with  $\overline{\mathbb{S}^n_-}$  is sometimes referred to as the hemispherical model – see [25, Chapter 3A].

To every function  $\xi : \mathbb{R}^n \rightarrow \mathbb{R}$  we associate a function  $\widehat{\xi} : \mathbb{S}^n_- \rightarrow \mathbb{R}$  in the following way: We first extend  $\xi$  from  $\mathbb{R}^n \cong \mathbb{R}^n \times \{-1\}$  to a 1-homogeneous function on the lower half-space  $\{(x, t) \in \mathbb{R}^{n+1} : t < 0\}$ , and then restrict this extension to  $\mathbb{S}^n_-$ . Explicitly, we have

$$(4.1) \quad \widehat{\xi}(x, t) = \frac{\xi(\Phi(x, t))}{|(\Phi(x, t), -1)|} = |t| \xi\left(\frac{x}{|t|}\right).$$

This operation is clearly a bijection between functions on  $\mathbb{R}^n$  and functions on  $\mathbb{S}^n_-$ , with the inverse operation given by

$$(4.2) \quad \xi(x) = |(x, -1)| \cdot \widehat{\xi}(\Phi^{-1}(x)) = \sqrt{|x|^2 + 1} \cdot \widehat{\xi}\left(\frac{x}{\sqrt{1 + |x|^2}}, -\frac{1}{\sqrt{1 + |x|^2}}\right).$$

We can now explain Definitions 1.7 and 1.8. We have:

**Proposition 4.2.** *A function  $\xi : \mathbb{R}^n \rightarrow \mathbb{R}$  is cosmically continuous in the sense of Definition 1.7 if and only if  $\widehat{\xi} : \mathbb{S}^n_- \rightarrow \mathbb{R}$  can be extended to a continuous function on the cosmic closure  $\overline{\mathbb{S}^n_-}$ . Moreover, in this case the extension  $\widehat{\xi} : \overline{\mathbb{S}^n_-} \rightarrow \mathbb{R}$  satisfies  $\widehat{\xi}(\theta, 0) = \bar{\xi}(\theta)$  for all  $\theta \in \mathbb{S}^{n-1}$ .*

In other words, cosmically continuous functions are simply continuous functions on  $\mathbb{R}^n$  that can be extended continuously to its cosmic closure (under our identification of  $\xi$  and  $\widehat{\xi}$ ). The proof of Proposition 4.2 is a straightforward exercise in topology, which we nonetheless include here for completeness:

*Proof.* Assume first that  $\widehat{\xi}$  can be extended to a continuous function  $\overline{\mathbb{S}^n_-}$ , which we also denote by  $\widehat{\xi}$ . Formula (4.2) immediately shows that  $\xi$  is continuous on  $\mathbb{R}^n$ . Moreover for all  $\theta \in \mathbb{S}^{n-1}$  we have

$$\begin{aligned} \lim_{t \rightarrow \infty} \frac{\xi(t\theta)}{t} &= \lim_{t \rightarrow \infty} \frac{\sqrt{t^2 + 1} \widehat{\xi}\left(\frac{t\theta}{\sqrt{t^2 + 1}}, -\frac{1}{\sqrt{t^2 + 1}}\right)}{t} \\ &= \lim_{t \rightarrow \infty} \frac{\sqrt{t^2 + 1}}{t} \cdot \lim_{t \rightarrow \infty} \widehat{\xi}\left(\frac{t}{\sqrt{t^2 + 1}}\theta, -\frac{1}{\sqrt{t^2 + 1}}\right) = \widehat{\xi}(\theta, 0) \end{aligned}$$

where we used the continuity of  $\widehat{\xi}$ . Moreover, since  $\widehat{\xi}$  is continuous on the compact set  $\overline{\mathbb{S}^n_-}$  it is uniformly continuous. Using this and the fact that

$$\lim_{t \rightarrow \infty} \sup_{\theta \in \mathbb{S}^{n-1}} \left| \left( \frac{t}{\sqrt{t^2 + 1}}\theta, -\frac{1}{\sqrt{t^2 + 1}} \right) - (\theta, 0) \right| = 0$$

we see that the  $\lim_{t \rightarrow \infty} \frac{\xi(t\theta)}{t}$  exists uniformly in  $\theta \in \mathbb{S}^{n-1}$ , and hence  $\xi$  is cosmically continuous in the sense of Definition 1.7.

For the converse, assume  $\xi : \mathbb{R}^n \rightarrow \mathbb{R}$  is cosmically continuous. Our goal is to prove that the function  $h : \overline{\mathbb{S}^n} \rightarrow \mathbb{R}$  defined by

$$h(x, t) = \begin{cases} \widehat{\xi}(x, t) & t < 0 \\ \bar{\xi}(x) & t = 0. \end{cases}$$

is the required continuous extension of  $\widehat{\xi}$ . It is clearly continuous on  $\mathbb{S}^n$  by (4.1), so we only need to check its continuity at every point of the form  $(\theta, 0)$ .

We note that  $\bar{\xi}$  is continuous on  $\mathbb{S}^{n-1}$  as a uniform limit of the continuous functions  $\theta \mapsto \frac{\xi(\lambda\theta)}{\lambda}$ . It is therefore enough to fix a sequence  $\{(x_k, t_k)\} \subseteq \mathbb{S}^n$  such that  $(x_k, t_k) \rightarrow (\theta, 0)$  and prove that  $h(x_k, t_k) \rightarrow h(\theta, 0)$ . Define  $\theta_k = \frac{x_k}{|x_k|} \rightarrow \theta$ , and write

$$|h(x_k, t_k) - h(\theta, 0)| = \left| \widehat{\xi}(x_k, t_k) - \bar{\xi}(\theta) \right| \leq \left| \widehat{\xi}(x_k, t_k) - \bar{\xi}(\theta_k) \right| + \left| \bar{\xi}(\theta_k) - \bar{\xi}(\theta) \right|.$$

By the continuity of  $\bar{\xi}$  we have  $|\bar{\xi}(\theta_k) - \bar{\xi}(\theta)| \rightarrow 0$ . For the first term we write

$$\begin{aligned} \left| \widehat{\xi}(x_k, t_k) - \bar{\xi}(\theta_k) \right| &= \left| |t_k| \xi \left( \frac{x_k}{|t_k|} \right) - \bar{\xi}(\theta_k) \right| \\ &= \left| |x_k| \frac{|t_k|}{|x_k|} \xi \left( \frac{|x_k| \theta_k}{|t_k|} \right) - |x_k| \bar{\xi}(\theta_k) + \bar{\xi}(\theta_k) (|x_k| - 1) \right| \\ &\leq |x_k| \cdot \left| \frac{\xi(s_k \theta_k)}{s_k} - \bar{\xi}(\theta_k) \right| + (\max \bar{\xi}) \cdot (|x_k| - 1) \\ &\xrightarrow{k \rightarrow \infty} 1 \cdot 0 + (\max \bar{\xi}) (1 - 1) = 0 \end{aligned}$$

where  $s_k = \frac{|x_k|}{|t_k|} \rightarrow \infty$ , and where we used the uniform convergence of  $\frac{\xi(\lambda\theta)}{\lambda}$  to  $\bar{\xi}$ . This shows that  $h(x_k, t_k) \rightarrow h(\theta, 0)$ , finishing the proof.  $\square$

We can now also better understand the notion of cosmic convergence. To every Borel measure  $\mu$  on  $\mathbb{R}^n$  we can associate a measure  $\widehat{\mu}$  on  $\mathbb{S}^n$  via the relation

$$(4.3) \quad \int_{\mathbb{R}^n} \xi d\mu = \int_{\mathbb{S}^{n-1}} \widehat{\xi} d\widehat{\mu}$$

for all cosmically continuous functions  $\xi : \mathbb{R}^n \rightarrow \mathbb{R}$ . Every Borel measure  $\nu$  on  $\mathbb{S}^{n-1}$  can also be considered as a measure  $\widehat{\nu}$  on the equator  $\overline{\mathbb{S}^n} \setminus \mathbb{S}^n = \{(x, 0) : x \in \mathbb{S}^{n-1}\}$  by identifying this equator with  $\mathbb{S}^{n-1}$  in the obvious way. Together  $\widehat{\mu} + \widehat{\nu}$  is a single measure on the cosmic closure  $\overline{\mathbb{S}^n}$  which is in a natural one-to-one correspondence with the pair  $(\mu, \nu)$ . We then have:

**Proposition 4.3.**  $(\mu_k, \nu_k) \rightarrow (\mu, \nu)$  cosmically if and only if  $\widehat{\mu}_k + \widehat{\nu}_k \rightarrow \widehat{\mu} + \widehat{\nu}$  weakly on  $\overline{\mathbb{S}^n}$ .

In other words, the cosmic convergence  $(\mu_k, \nu_k) \rightarrow (\mu, \nu)$  is the same as weak convergence on the cosmic closure  $\overline{\mathbb{S}^n}$  of  $\mathbb{R}^n$ , under our identification of the pair  $(\mu, \nu)$  with the measure  $\widehat{\mu} + \widehat{\nu}$ .

*Proof.* This is an exercise in expanding the definitions. By definition,  $(\mu_k, \nu_k) \rightarrow (\mu, \nu)$  cosmically if for every cosmically continuous function  $\xi : \mathbb{R}^n \rightarrow \mathbb{R}$  we have

$$(4.4) \quad \int_{\mathbb{R}^n} \xi d\mu_k + \int_{\mathbb{S}^{n-1}} \bar{\xi} d\nu_k \xrightarrow{k \rightarrow \infty} \int_{\mathbb{R}^n} \xi d\mu + \int_{\mathbb{S}^{n-1}} \bar{\xi} d\nu.$$

By (4.3) and the “moreover” part of Proposition 4.2 we have

$$\begin{aligned} \int_{\mathbb{R}^n} \xi d\mu + \int_{\mathbb{S}^{n-1}} \bar{\xi} d\nu &= \int_{\mathbb{S}_-^n} \widehat{\xi} d\widehat{\mu} + \int_{\mathbb{S}^{n-1}} \widehat{\xi}(\theta, 0) d\nu(\theta) \\ &= \int_{\mathbb{S}_-^n} \widehat{\xi} d\widehat{\mu} + \int_{\mathbb{S}_-^n \setminus \mathbb{S}_-^n} \widehat{\xi} d\widehat{\nu} = \int_{\mathbb{S}_-^n} \widehat{\xi} d(\widehat{\mu} + \widehat{\nu}). \end{aligned}$$

The same of course is true for  $(\mu_k, \nu_k)$ , so the cosmic convergence (4.4) is equivalent to

$$\int_{\mathbb{S}_-^n} \widehat{\xi} d(\widehat{\mu}_k + \widehat{\nu}_k) \rightarrow \int_{\mathbb{S}_-^n} \widehat{\xi} d(\widehat{\mu} + \widehat{\nu})$$

for all cosmically continuous function  $\xi : \mathbb{R}^n \rightarrow \mathbb{R}$ . By Proposition 4.2 this equivalent to saying that

$$\int_{\mathbb{S}_-^n} \rho d(\widehat{\mu}_k + \widehat{\nu}_k) \rightarrow \int_{\mathbb{S}_-^n} \rho d(\widehat{\mu} + \widehat{\nu})$$

for all continuous functions  $\rho : \overline{\mathbb{S}_-^n} \rightarrow \mathbb{R}$ , which precisely means that  $\widehat{\mu}_k + \widehat{\nu}_k \rightarrow \widehat{\mu} + \widehat{\nu}$  weakly on  $\overline{\mathbb{S}_-^n}$ .  $\square$

Given Proposition 4.3, it is natural to ask for an explicit description of the measure  $\widehat{\mu}_f + \widehat{\nu}_f$  for  $f = e^{-\phi} \in \text{Cvx}_n$ . Recall that the domain of a function  $\phi \in \text{Cvx}_n$  is

$$\text{dom}(\phi) = \{x \in \mathbb{R}^n : \phi(x) < \infty\} \subseteq \mathbb{R}^n,$$

and the epigraph of  $\phi$  is

$$\text{epi}(\phi) = \{(x, t) \in \mathbb{R}^{n+1} : x \in \text{dom}(\phi), t \geq \phi(x)\} \subseteq \mathbb{R}^{n+1}.$$

As usual, we denote by  $\mathcal{H}^n$  the  $n$ -dimensional Hausdorff measure. The epigraph  $\text{epi}(\phi)$  is an (unbounded) closed convex set with non-empty interior, so the Gauss map  $n_{\text{epi}(\phi)}$  is defined  $\mathcal{H}^n$ -almost everywhere. Moreover, for every  $(x, t) \in \partial \text{epi}(\phi)$  it is clear that  $(x, t') \in \text{epi}(\phi)$  for all  $t' \geq t$ , which implies that  $n_{\text{epi}(\phi)}(x, t) \in \overline{\mathbb{S}_-^n}$ . We can now state the result:

**Proposition 4.4.** *For every  $f = e^{-\phi} \in \text{LC}_n$  we have*

$$\widehat{\mu}_f + \widehat{\nu}_f = (n_{\text{epi}(\phi)})_{\#} \left( e^{-t} d\mathcal{H}^n(x, t) \Big|_{\partial \text{epi}(\phi)} \right).$$

*Explicitly, this means that for all continuous functions  $\rho : \overline{\mathbb{S}_-^n} \rightarrow \mathbb{R}$  one has*

$$\int_{\overline{\mathbb{S}_-^n}} \rho d(\widehat{\mu}_f + \widehat{\nu}_f) = \int_{\partial \text{epi}(\phi)} \rho(n_{\text{epi}(\phi)}(x, t)) e^{-t} d\mathcal{H}^n(x, t).$$

*Proof.* Write  $\rho = \widehat{\xi}$  for a cosmically continuous function  $\xi : \mathbb{R}^n \rightarrow \mathbb{R}$ . We partition  $\partial \text{epi}(\phi)$  as  $\partial \text{epi}(\phi) = A \cup B$  for

$$A = \{(x, \phi(x)) : x \in \text{dom}(\phi)\}$$

and

$$B = \{(x, t) : x \in \partial \text{dom}(\phi), t \geq \phi(x)\}.$$

We first prove that

$$(4.5) \quad \int_{\mathbb{S}_-^n} \widehat{\xi} d\widehat{\mu}_f = \int_A \widehat{\xi} (n_{\text{epi}(\phi)}(x, t)) e^{-t} d\mathcal{H}^n(x, t).$$

Indeed, by definition we have

$$\int_{\mathbb{S}_-^n} \widehat{\xi} d\widehat{\mu}_f = \int_{\mathbb{R}^n} \xi d\mu_f = \int_{\text{dom}(\phi)} \xi(\nabla\phi(u)) e^{-\phi(u)} du.$$

Perform the (locally Lipschitz) change of variables  $T : \text{dom}(\phi) \rightarrow A$  defined by  $Tu = (u, \phi(u))$ . The Jacobian of  $T$  is

$$JT(u) = \det(DT(u)^t \cdot DT(u)) = \sqrt{\det(Id + \nabla\phi \otimes \nabla\phi)} = \sqrt{1 + |\nabla\phi(u)|^2},$$

so by the change of variables formula for Lipschitz maps (see [12, Theorem 3.9])

$$(4.6) \quad \int_{\mathbb{S}_-^n} \widehat{\xi} d\widehat{\mu}_f = \int_A \xi(\nabla\phi(x)) e^{-t} \cdot \frac{1}{\sqrt{1 + |\nabla\phi(x)|^2}} d\mathcal{H}^n(x, t).$$

But for almost all  $(x, t) \in A$  we have  $n_{\text{epi}(\phi)}(x, t) = \frac{(\nabla\phi(x), -1)}{|(\nabla\phi(x), -1)|}$ . Therefore

$$(4.7) \quad \widehat{\xi}(n_{\text{epi}(\phi)}(x, t)) = \widehat{\xi}\left(\frac{(\nabla\phi(x), -1)}{|(\nabla\phi(x), -1)|}\right) = \frac{\xi(\nabla\phi(x))}{\sqrt{1 + |\nabla\phi(x)|^2}},$$

where we used formula (4.2). Equations (4.6) and (4.7) together clearly imply (4.5).

We now prove that

$$(4.8) \quad \int_{\mathbb{S}_-^n \setminus \mathbb{S}_-^n} \widehat{\xi} d\widehat{\nu}_f = \int_B \widehat{\xi}(n_{\text{epi}(\phi)}(x, t)) e^{-t} d\mathcal{H}^n(x, t).$$

Indeed, for almost every point  $(x, t) \in B$  we have  $n_{\text{epi}(\phi)}(x, t) = (n_{\text{dom}(\phi)}(x), 0)$ . Using this fact, Fubini's theorem, and the connection between  $\widehat{\xi}$  and  $\xi$  from Proposition 4.2 we have

$$\begin{aligned} \int_B \widehat{\xi}(n_{\text{epi}(\phi)}(x, t)) e^{-t} d\mathcal{H}^n(x, t) &= \int_{x \in \partial \text{dom}(\phi)} \int_{t=\phi(x)}^{\infty} \widehat{\xi}(n_{\text{epi}(\phi)}(x, t)) e^{-t} dt d\mathcal{H}^{n-1}(x) \\ &= \int_{x \in \partial \text{dom}(\phi)} \int_{t=\phi(x)}^{\infty} \widehat{\xi}(n_{\text{dom}(\phi)}(x), 0) e^{-t} dt d\mathcal{H}^{n-1}(x) \\ &= \int_{x \in \partial \text{dom}(\phi)} \bar{\xi}(n_{\text{dom}(\phi)}(x)) e^{-\phi(x)} d\mathcal{H}^{n-1}(x) \\ &= \int_{\mathbb{S}^{n-1}} \bar{\xi} d\nu_f = \int_{\mathbb{S}_-^n \setminus \mathbb{S}_-^n} \widehat{\xi} d\widehat{\nu}_f \end{aligned}$$

as claimed.

Since  $\mathcal{H}^n(A \cap B) = 0$ , the identities (4.5) and (4.8) together imply the result.  $\square$

We record for later use that what we've actually shown is that for every cosmically continuous function  $\xi : \mathbb{R}^n \rightarrow \mathbb{R}$  one has

$$(4.9) \quad \int_{\mathbb{R}^n} \xi d\mu_f + \int_{\mathbb{S}^{n-1}} \bar{\xi} d\nu_f = \int_{\partial \text{epi}(\phi)} \widehat{\xi}(n_{\text{epi}(\phi)}(x, t)) e^{-t} d\mathcal{H}^n(x, t).$$

Using equation (4.9) we can rewrite the first variation formula (1.1) in the following nice way:

**Corollary 4.5.** *For every  $f = e^{-\phi} \in \text{LC}_n$  and an upper semicontinuous log-concave function  $g : \mathbb{R}^n \rightarrow \mathbb{R}$  we have*

$$\delta(f, g) = \int_{\partial \text{epi}(\phi)} \widehat{h}_g(n_{\text{epi}(\phi)}(x, t)) e^{-t} d\mathcal{H}^n(x, t).$$

Here we define  $\widehat{h}_g : \overline{\mathbb{S}^n} \rightarrow (-\infty, \infty]$  in the natural way:  $h_g : \mathbb{R}^n \rightarrow (-\infty, \infty]$  is usually not cosmically continuous, but we may still define  $\widehat{h}_g$  on  $\mathbb{S}^n$  by formula (4.1), and then on the equator  $\overline{\mathbb{S}^n} \setminus \mathbb{S}^n$  define  $\widehat{h}_g(\theta, 0) = \overline{h}_g(\theta)$ . Even though  $h_g$  is not cosmically continuous, the proof that

$$\int h_g d\mu_f + \int \overline{h}_g d\nu_f = \int_{\partial \text{epi}(\phi)} \widehat{h}_g(n_{\text{epi}(\phi)}(x, t)) e^{-t} d\mathcal{H}^n(x, t)$$

still works in exactly the same way as the proof of (4.9). The corollary follows immediately from this identity and (1.1).

## 5. CONTINUITY OF FUNCTIONAL SURFACE AREA MEASURES

In this section we prove Theorem 1.9. Fix  $\{f_k\}_{k=1}^\infty, f \in \text{LC}_n$  such that  $f_k \rightarrow f$ . Write as usual  $f_k = e^{-\phi_k}$  and  $f = e^{-\phi}$ . Our goal is to prove that  $(\mu_{f_k}, \nu_{f_k}) \rightarrow (\mu_f, \nu_f)$  cosmically. By (4.9), we need to prove that for every cosmically continuous function  $\xi : \mathbb{R}^n \rightarrow \mathbb{R}$  one has

$$\int_{\partial \text{epi}(\phi_k)} \widehat{\xi}(n_{\text{epi}(\phi_k)}(x, t)) e^{-t} d\mathcal{H}^n(x, t) \rightarrow \int_{\partial \text{epi}(\phi)} \widehat{\xi}(n_{\text{epi}(\phi)}(x, t)) e^{-t} d\mathcal{H}^n(x, t).$$

The main idea of the proof is to perform a change of variables and transform the domain of integration to  $\mathbb{R}^n$ , using a new notion of a *curvilinear radial function* for convex functions. For technical reasons we will need to work with the following classes of convex functions:

**Definition 5.1.** For every  $\epsilon > 0$  we set

$$\text{Cvx}_n^{(\epsilon)} = \left\{ \phi \in \text{Cvx}_n : \phi(x) < \min \phi + \frac{1}{2} \text{ for all } x \in \mathbb{R}^n \text{ with } |x| \leq \epsilon \right\}.$$

The choice of the constant  $\frac{1}{2}$  in the definition of  $\text{Cvx}_n^{(\epsilon)}$  is immaterial, but choosing a constant smaller than 1 helps to eliminate some other constants later in the proof. The following rather technical properties of the classes  $\text{Cvx}_n^{(\epsilon)}$  will be crucial for the proof:

**Lemma 5.2.** (1) *For every  $\phi \in \text{Cvx}_n$  there exists  $v \in \mathbb{R}^n$  and  $\epsilon > 0$  such that  $\tau_v \phi \in \text{Cvx}_n^{(\epsilon)}$ , where  $(\tau_v \phi)(x) = \phi(x + v)$ .*

(2) *Assume that  $\{\phi_k\}_{k=1}^\infty \subseteq \text{Cvx}_n$  and  $\phi_k \rightarrow \phi \in \text{Cvx}_n^{(\epsilon)}$  for some  $\epsilon > 0$ . Then  $\phi_k \in \text{Cvx}_n^{(\epsilon/2)}$  for all large enough  $k$ .*

(3) *Assume  $\phi \in \text{Cvx}_n^{(\epsilon)}$ . Fix  $(x_0, t_0) \in \partial \text{epi}(\phi)$  and let  $v = (v_x, v_t) \in \mathbb{R}^{n+1}$  be an outer normal to  $\text{epi}(\phi)$  at  $(x_0, t_0)$ . Then*

$$\langle (x_0, -1), v \rangle = \langle x_0, v_x \rangle - v_t \geq \min \left( \epsilon, \frac{1}{2} \right) |v|^2.$$

*Proof.* (1) Fix  $w \in \mathbb{R}^n$  such that  $\phi(w) = \min \phi$ . If  $w \in \text{int}(\text{dom}(\phi))$  we just choose  $v = w$ . If  $w \in \partial \text{dom}(\phi)$ , then every  $v \in \text{int}(\text{dom}(\phi))$  close enough to  $w$  will satisfy

$$\phi(v) \leq \phi(w) + \frac{1}{4} = \min \phi + \frac{1}{4}.$$

Since  $\phi$  is continuous on  $\text{int}(\text{dom}(\phi))$  there exists  $\epsilon > 0$  such that for all  $x$  with  $|x - v| \leq \epsilon$  we have

$$\phi(x) < \phi(v) + \frac{1}{4} \leq \min \phi + \frac{1}{2}.$$

This shows that  $\tau_v \phi \in \text{Cvx}_n^{(\epsilon)}$ .

(2) Assume that  $\{\phi_k\}_{k=1}^\infty \subseteq \text{Cvx}_n$  and  $\phi_k \rightarrow \phi \in \text{Cvx}_n^{(\epsilon)}$ . Since  $\overline{B}_\epsilon(0) \subseteq \text{dom}(\phi)$  we have  $\overline{B}_{\epsilon/2}(0) \subseteq \text{int}(\text{dom}(\phi))$ , and it follows that  $\phi_k \rightarrow \phi$  uniformly on  $\overline{B}_{\epsilon/2}(0)$  – See [25, Theorem 7.17(c)]. Also by [6, Lemma 12] we have  $\min \phi_k \rightarrow \min \phi$ . Therefore

$$\lim_{k \rightarrow \infty} \left( \max_{\overline{B}_{\epsilon/2}(0)} \phi_k \right) = \max_{\overline{B}_{\epsilon/2}(0)} \phi < \min \phi + \frac{1}{2} = \lim_{k \rightarrow \infty} \phi_k + \frac{1}{2}.$$

It follows that indeed  $\max_{\overline{B}_{\epsilon/2}(0)} \phi_k < \phi_k + \frac{1}{2}$  for all large enough  $k$ .

(3) Recall that we always have  $v_t \leq 0$ . If  $v_x = 0$  then

$$\langle x_0, v_x \rangle - v_t = |v_t| = |v|$$

and there is nothing to prove. Otherwise, since  $\epsilon \frac{v_x}{|v_x|} \in \overline{B}_\epsilon(0)$  we know that  $\phi\left(\epsilon \frac{v_x}{|v_x|}\right) \leq \min \phi + \frac{1}{2}$ , and so  $(\epsilon \frac{v_x}{|v_x|}, \min \phi + \frac{1}{2}) \in \text{epi}(\phi)$ . It follows that

$$\left\langle \left( \epsilon \frac{v_x}{|v_x|}, \min \phi + \frac{1}{2} \right) - (x_0, t_0), (v_x, v_t) \right\rangle \leq 0.$$

But  $t_0 \geq \phi(x_0) \geq \min \phi$ , and so

$$\begin{aligned} \left\langle \left( \epsilon \frac{v_x}{|v_x|}, \min \phi + \frac{1}{2} \right) - (x_0, t_0), (v_x, v_t) \right\rangle &= \left\langle \epsilon \frac{v_x}{|v_x|} - x_0, v_x \right\rangle + \left( \min \phi + \frac{1}{2} - t_0 \right) v_t \\ &\geq \epsilon |v_x| - \langle x_0, v_x \rangle + \frac{1}{2} v_t. \end{aligned}$$

Therefore  $\epsilon |v_x| - \langle x_0, v_x \rangle + \frac{1}{2} v_t \leq 0$ , or

$$\langle x_0, v_x \rangle - v_t \geq \epsilon |v_x| + \frac{1}{2} |v_t| \geq \min \left( \epsilon, \frac{1}{2} \right) \cdot \sqrt{|v_x|^2 + v_t^2} = \min \left( \epsilon, \frac{1}{2} \right) |v|^2$$

as claimed. □

Following part (1) of the lemma, we remark that if  $\phi$  attains its minimum on  $\text{int}(\text{dom}(\phi))$  then after translation one can assume not only that  $\phi \in \text{Cvx}_n^{(\epsilon)}$  but also that  $\phi(0) = \min \phi$ . Using this property can greatly simplify some of the arguments below such as the proof of Proposition 5.3. Unfortunately, this simplification is not possible if  $\phi$  attains its minimum (only) on  $\partial \text{dom}(\phi)$ .

The following proposition establishes the main definition for this section:

**Proposition 5.3.** Fix  $\phi \in \text{Cvx}_n^{(\epsilon)}$  and  $u \in \mathbb{R}^n$ . Consider the curve  $\gamma_u : (0, \infty) \rightarrow \mathbb{R}^n \times \mathbb{R}$  defined by  $\gamma_u(s) = (su, \log(\frac{1}{s}))$ . Then there exists a number  $s_\phi(u) \in (0, \infty)$  with the following properties:

- (1)  $\gamma_u(s) \in \text{int}(\text{epi}(\phi))$  for all  $s < s_\phi(u)$ .
- (2)  $\gamma_u(s_\phi(u)) \in \partial \text{epi}(\phi)$ .
- (3)  $\gamma_u(s) \notin \text{epi}(\phi)$  for all  $s > s_\phi(u)$ .

The value  $s_\phi(u)$  is clearly unique. The function  $s_\phi : \mathbb{R}^n \rightarrow (0, \infty)$  mapping  $u$  to  $s_\phi(u)$  is called the curvilinear radial function of  $\phi$ .

*Proof.* Since  $\phi \in \text{Cvx}_n^{(\epsilon)}$  we know in particular that  $\phi$  is bounded on  $\overline{B}_\epsilon(0)$ . Since  $\gamma_u(s) \xrightarrow{s \rightarrow 0^+} (0, +\infty)$ , we clearly have  $\gamma_u(s) \in \text{int}(\text{epi}(\phi))$  for small enough  $s$ . On the other hand, since  $\phi$  is bounded from below and  $\log \frac{1}{s} \xrightarrow{s \rightarrow \infty} -\infty$ , we have  $\gamma_u(s) \notin \text{epi}(\phi)$  for large enough  $s$ . By continuity, there exists  $s_0 \in (0, \infty)$  such that  $\gamma_u(s_0) \in \partial \text{epi}(\phi)$ . We write  $\gamma_u(s_0) = (x_0, t_0)$ .

Our next claim is that there exists  $\delta > 0$  such that  $\gamma_u(s_0 - s) \in \text{int}(\text{epi}(\phi))$  and  $\gamma_u(s_0 + s) \notin \text{epi}(\phi)$  for all  $0 < s \leq \delta$ . To prove the claim, it is enough to show that for every outer normal  $v = (v_x, v_t) \in \mathbb{R}^{n+1}$  to  $\text{epi}(\phi)$  at  $\gamma_u(s_0)$  we have  $\langle \gamma'_u(s_0), v \rangle > 0$ . But  $\gamma'_u(s_0) = (u, -\frac{1}{s_0})$ , so

$$\langle \gamma'_u(s_0), v \rangle = \frac{1}{s_0} \langle (s_0 u, -1), v \rangle = \frac{1}{s_0} \langle (x_0, -1), v \rangle \geq \frac{1}{s_0} \min\left(\epsilon, \frac{1}{2}\right) |v|^2 > 0$$

by Lemma 5.2(3) and the claim is proved.

The rest of the proof is a straightforward topological argument. Assume by contradiction that there exists a point  $s_1 \neq s_0$  such that  $\gamma_u(s_1) \in \partial \text{epi}(\phi)$ . If  $s_1 < s_0$  then the set

$$\{s \in [s_1, s_0 - \delta] : \gamma_u(s) \in \partial \text{epi}(\phi)\}$$

is a non-empty compact set, so it has some maximum  $s_2$ . But then the same argument as above shows that  $\gamma_u(s_2 + \delta') \notin \text{epi}(\phi)$  for small enough  $\delta' > 0$ , and since  $\gamma_u(s_0 - \delta) \in \text{int}(\text{epi}(\phi))$  there must exist a point  $s_3 \in (s_2 + \delta', s_0 - \delta)$  such that  $\gamma_u(s_3) \in \partial \text{epi}(\phi)$ , contradicting the maximality of  $s_2$ . The case  $s_1 > s_0$  is the same. It follows that  $s_0$  is the unique point which satisfies  $\gamma_u(s_0) \in \text{epi}(\phi)$ .

If now  $\gamma_u(s) \notin \text{int}(\text{epi}(\phi))$  for  $s < s_0$ , then since  $\gamma_u(s_0 - \delta) \in \text{int}(\text{epi}(\phi))$  again by continuity we could find  $s \leq \tilde{s} < s_0 - \delta$  such that  $\gamma_u(\tilde{s}) \in \partial \text{epi}(\phi)$ , which is a contradiction. The same argument shows that  $\gamma_u(s) \notin \text{epi}(\phi)$  for all  $s > s_0$ , completing the proof.  $\square$

From the function  $s_\phi$  we obtain our desired parametrization of  $\partial \text{epi}(\phi)$ :

**Corollary 5.4.** For  $\phi \in \text{Cvx}_n^{(\epsilon)}$  define  $F_\phi : \mathbb{R}^n \rightarrow \partial \text{epi}(\phi)$  by

$$F_\phi(u) = \gamma_u(s_\phi(u)) = \left( s_\phi(u)u, \log\left(\frac{1}{s_\phi(u)}\right) \right).$$

Then  $F_\phi$  is a bijection between  $\mathbb{R}^n$  and  $\partial \text{epi}(\phi)$ , with the inverse map  $G_\phi : \partial \text{epi}(\phi) \rightarrow \mathbb{R}^n$  given by  $G_\phi(x, t) = e^t x$ .



*Proof.* The fact that  $F_\phi(u) \in \partial \text{epi}(\phi)$  follows from Proposition 5.3. It is immediate from the formulas that  $G_\phi(F_\phi(u)) = u$  for all  $u \in \mathbb{R}^n$ . Conversely, for every  $(x, t) \in \partial \text{epi}(\phi)$  we have

$$\gamma_{G_\phi(x,t)}(e^{-t}) = (x, t) \in \partial \text{epi}(\phi),$$

Which precisely means by definition that  $s_\phi(G_\phi(x, t)) = e^{-t}$  and  $F_\phi(G_\phi(x, t)) = (x, t)$ . This proves the claim.  $\square$

*Remark 5.5.* Assume  $\phi \in \text{Cvx}_n$  satisfies  $\min \phi = \phi(0) = 0$ . In [2], Artstein-Avidan and Milman defined the  $\mathcal{J}$ -transform of such functions, which can be considered as their Minkowski functional. Therefore the reciprocal  $\frac{1}{\mathcal{J}\phi}$  can be thought of as a radial function for  $\phi$ . Explicitly we have

$$\frac{1}{\mathcal{J}\phi}(u) = \sup \{s \geq 0 : (su, s) \in \text{epi}(\phi)\},$$

i.e.  $\frac{1}{\mathcal{J}\phi}(u)$  is the radial function of  $\text{epi}(\phi)$  in the direction  $(u, 1)$ . This is similar to our definition, with the non-linear curve  $\gamma_u(s)$  replaced by the linear curve  $\tilde{\gamma}_u(s) = (su, s)$ . In many ways the function  $\frac{1}{\mathcal{J}\phi}$  is simpler and more canonical than our function  $s_\phi$ . For example  $\mathcal{J}\phi$  is a convex function, while  $s_\phi$  does not seem to have any convexity properties. However, unlike Corollary 5.4, the corresponding map

$$\tilde{F}_\phi(s) = \tilde{\gamma}_u \left( \frac{1}{\mathcal{J}\phi}(u) \right) = \frac{(u, 1)}{(\mathcal{J}\phi)(u)}$$

does not always parametrize the boundary  $\partial \text{epi}(\phi)$ . As an example, for  $\phi(x) = |x|$  we have

$$(\mathcal{J}\phi)(u) = \mathbf{1}_{B_2^\infty}(u) = \begin{cases} 0 & |u| \leq 1 \\ \infty & |u| > 1, \end{cases}$$

so  $\tilde{F}_\phi$  is not even well-defined and in any case does not parametrize any part of  $\partial \text{epi}(\phi)$ . Therefore the curvilinear radial function  $s_\phi$  is more suitable for our purpose.

Our next goal is prove that  $s_\phi$  is a locally Lipschitz function and compute its gradient, which we will do using the implicit function theorem. Since  $\phi$  is not necessarily smooth we need a version of the implicit function theorem for Lipschitz functions, which is due to Clarke ([4]). We give the necessary definitions:

Consider a (locally) Lipschitz function  $\Phi : \mathbb{R}^N \rightarrow \mathbb{R}$ . The generalized gradient of  $\Phi$  at a point  $x_0$  can be defined by ([4, Theorem 2.5.1])

$$\partial\Phi(x_0) = \text{conv} \left\{ \lim_{i \rightarrow \infty} \nabla\Phi(x_i) : \begin{array}{l} x_i \rightarrow x, \Phi \text{ is differentiable at every } x_i \\ \text{and } \lim_{i \rightarrow \infty} \nabla\Phi(x_i) \text{ exists.} \end{array} \right\}.$$

If  $\Phi$  is convex, the generalized gradient agrees with the usual subgradient ([4, Proposition 2.2.7]). If we have a map  $\Phi : \mathbb{R}^N \times \mathbb{R}^M \rightarrow \mathbb{R}$ , which we write as  $\Phi(x, y)$ , we denote by  $\partial_y\Phi(x_0, y_0)$  the generalized gradient only in the  $y$  variable. Formally these are all vectors  $v \in \mathbb{R}^M$  such that  $(w | v) \in \partial\Phi(x_0, y_0)$  for some  $w \in \mathbb{R}^N$ . Generalized gradients satisfy the chain rule ([4, Theorem 2.3.10]): If  $\Psi : \mathbb{R}^N \rightarrow \mathbb{R}^N$  is smooth and  $\Phi : \mathbb{R}^N \rightarrow \mathbb{R}$  is Lipschitz then

$$\partial(\Phi \circ \Psi)(x_0) \subseteq \{v \cdot D\Psi(x_0) : v \in \partial\Phi(\Psi(x_0))\}$$

(in many cases including ours there is actually an equality, but we will not need this fact). The implicit function theorem then reads:

**Theorem 5.6** ([4, page 256]). *Assume  $\Phi : \mathbb{R}^N \times \mathbb{R} \rightarrow \mathbb{R}$  is a Lipschitz function and  $\Phi(x_0, y_0) = 0$ . Assume further that  $0 \notin \partial_y \Phi(x_0, y_0)$ . Then there exists a neighborhood  $U$  of  $x_0$  and a **Lipschitz** function  $\Psi : U \rightarrow \mathbb{R}$  such  $\Psi(x_0) = y_0$  and  $\Phi(x, \Psi(x)) = 0$  for all  $x \in U$ .*

Of course a similar theorem holds for functions  $\Phi : \mathbb{R}^N \times \mathbb{R}^M \rightarrow \mathbb{R}^M$ , but this version will suffice for our goals. Using these tools we prove:

**Proposition 5.7.** *For all  $\phi \in \text{Cvx}_n^{(\epsilon)}$  the function  $s_\phi$  is locally Lipschitz. Moreover, for almost every  $u_0 \in \mathbb{R}^n$  we have*

$$(5.1) \quad \nabla s_\phi(u_0) = -s_\phi(u_0)^2 \cdot \frac{n_{\text{epi}(\phi)}^{(x)}(x_0, t_0)}{\langle n_{\text{epi}(\phi)}(x_0, t_0), (x_0, -1) \rangle}.$$

Here  $(x_0, t_0) = F_\phi(u_0) \in \partial \text{epi}(\phi)$  and  $n_{\text{epi}(\phi)}^{(x)}$  denotes the first  $n$  coordinates of the outer unit normal  $n_{\text{epi}(\phi)}$ .

*Proof.* Fix an arbitrary point  $(\tilde{x}, \tilde{t}) \in \text{int}(\text{epi}(\phi))$ , and let  $\rho : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$  be the Minkowski functional of  $\text{epi}(\phi)$  with respect to  $(\tilde{x}, \tilde{t})$ :

$$\rho(x, t) = \inf \left\{ \lambda > 0 : \frac{(x, t) - (\tilde{x}, \tilde{t})}{\lambda} \in \text{epi}(\phi) \right\}.$$

Then  $\rho$  is convex, and  $\rho(x, t) = 1$  if and only if  $(x, t) \in \partial \text{epi}(\phi)$ . Moreover, at every  $(x, t) \in \partial \text{epi}(\phi)$  the elements of the subgradient  $\partial \rho(x, t)$  are outer normals to  $\text{epi}(\phi)$  at  $(x, t)$ .

Consider now the function  $\Phi : \mathbb{R}^n \times \mathbb{R}_+ \rightarrow \mathbb{R}$  defined by  $\Phi(u, s) = \rho(su, \log(\frac{1}{s}))$ . Fix a point  $u_0 \in \mathbb{R}^n$  and set  $s_0 = s_\phi(u_0)$ . Then by definition we have  $\Phi(u_0, s_0) = \rho(F_\phi(u_0)) = 1$ . Moreover, by the chain rule we have

$$\begin{aligned} \partial_s \Phi(u_0, s_0) &\subseteq \left\{ \langle v_x, u_0 \rangle - \frac{v_t}{s_0} : (v_x, v_t) \in \partial \rho \left( s_0 u_0, \log \left( \frac{1}{s_0} \right) \right) \right\} \\ &= \left\{ \langle v_x, u_0 \rangle - \frac{v_t}{s_0} : (v_x, v_t) \in \partial \rho(F_\phi(u_0)) \right\} \end{aligned}$$

Define  $(x_0, t_0) = F_\phi(u_0) \in \text{epi}(\phi)$ . Since every vector  $(v_x, v_t) \in \partial \rho(x_0, t_0)$  is an outer normal to  $\text{epi}(\phi)$  at  $(x_0, t_0)$ , it follows from Lemma 5.2(3) that

$$\langle v_x, u_0 \rangle - \frac{v_t}{s_0} = \frac{1}{s_0} (\langle v_x, x_0 \rangle - v_t) \geq \frac{1}{s_0} \cdot \min \left( \epsilon, \frac{1}{2} \right) |v|^2 > 0.$$

Therefore by the implicit function theorem the equation  $\Phi(u, s) = 1$  defines  $s$  as a Lipschitz function of  $u$  locally around  $u_0$ . But this function is exactly  $s_\phi$ , so  $s_\phi$  is locally Lipschitz.

To compute  $\nabla s_\phi$ , fix a point  $u_0 \in \mathbb{R}^n$  such that  $s_\phi$  is differentiable at  $u_0$  and  $\text{epi}(\phi)$  has a unique outer unit normal at  $(x_0, t_0) = F_\phi(u_0)$  – these conditions hold almost everywhere since  $s_\phi$  is locally Lipschitz and  $\text{epi}(\phi)$  is convex. Then  $\rho$  is differentiable at  $(x_0, t_0)$  and  $\nabla \rho(x_0, t_0) = (v_x, v_t) =$

$c \cdot n_{\text{epi}(\phi)}(x_0, t_0)$  for some  $c > 0$ . Differentiating the identity  $\Phi(u, s_\phi(u)) = 1$  at  $u_0$  using the (standard) chain rule we obtain

$$\begin{aligned} \nabla s_\phi(u_0) &= -\frac{(D_u \Phi)(u_0, s_\phi(u_0))}{(D_s \Phi)(u_0, s_\phi(u_0))} = -\frac{s_\phi(u_0) \cdot v_x}{\langle v_x, u_0 \rangle - \frac{v_t}{s_\phi(u_0)}} \\ &= -s_\phi(u_0)^2 \frac{v_x}{\langle (v_x, v_t), (s_\phi(u_0)u_0, -1) \rangle} = -s_\phi(u_0)^2 \cdot \frac{n_{\text{epi}(\phi)}^{(x)}(x_0, t_0)}{\langle n_{\text{epi}(\phi)}(x_0, t_0), (x_0, -1) \rangle} \end{aligned}$$

as claimed.  $\square$

Our next goal is to understand the behavior of the curvilinear radial function under epi-convergence:

**Proposition 5.8.** *Assume that  $\{\phi_k\}_{k=1}^\infty, \phi \in \text{Cvx}_n^{(\epsilon)}$  for some  $\epsilon > 0$  and that  $\phi_k \rightarrow \phi$ . Then:*

- (1) *For all  $u \in \mathbb{R}^n$  we have  $s_{\phi_k}(u) \rightarrow s_\phi(u)$  and  $F_{\phi_k}(u) \rightarrow F_\phi(u)$ .*
- (2) *For almost every  $u \in \mathbb{R}^n$  we have  $\nabla s_{\phi_k}(u) \rightarrow \nabla s_\phi(u)$  and  $DF_{\phi_k}(u) \rightarrow DF_\phi(u)$ .*

*Proof.* (1) Since  $\{\phi_k\}_{k=1}^\infty$  is convergent it is uniformly coercive, i.e.  $\phi_k \geq \psi_L$  for some function of the form  $\psi_L(x) = a|x| + b$ . We also know that for all  $x \in \overline{B}_\epsilon(0)$  we have

$$\phi_k(x) \leq \min \phi_k + \frac{1}{2} \rightarrow \min \phi + \frac{1}{2},$$

so the functions  $\{\phi_k\}_{k=1}^\infty$  are uniformly bounded on  $\overline{B}_\epsilon(0)$ . Therefore  $\phi_k \leq \psi_U$  for some function  $\psi_U$  of the form  $\psi_U = \mathbf{1}_{\overline{B}_\epsilon(0)}^\infty + M$ . Therefore  $0 < s_{\psi_U}(u) \leq s_{\phi_k}(u) \leq s_{\psi_L}(u) < \infty$  for all  $u$ , i.e. the sequence  $\{s_{\phi_k}(u)\}_{k=1}^\infty$  is bounded. It is therefore enough to prove that for every converging subsequence  $s_{\phi_{k_\ell}}(u) \rightarrow \tilde{s}$  we must have  $\tilde{s} = s_\phi(u)$ .

We have  $\gamma_u(s_{\phi_{k_\ell}}(u)) \in \text{epi}(\phi_{k_\ell})$  and  $\gamma_u(s_{\phi_{k_\ell}}(u)) \rightarrow \gamma_u(\tilde{s})$ . Since  $\text{epi}(\phi_{k_\ell}) \rightarrow \text{epi}(\phi)$ , it follows that  $\gamma_u(\tilde{s}) \in \text{epi}(\phi)$ . Therefore  $\tilde{s} \leq s_\phi(u)$ .

Assume by contradiction that  $\tilde{s} < s_\phi(u)$ , so  $\gamma_u(\tilde{s}) \in \text{int}(\text{epi}(\phi))$ . Choose a closed ball  $\overline{B}$  around  $\gamma_u(\tilde{s})$  such that  $\overline{B} \subseteq \text{int}(\text{epi}(\phi))$ . Since  $\gamma_u(s_{\phi_{k_\ell}}(u)) \rightarrow \gamma_u(\tilde{s})$  we clearly have  $\gamma_u(s_{\phi_{k_\ell}}(u)) \in \overline{B}$  for all large enough  $\ell$ . But  $\text{epi}(\phi_{k_\ell}) \rightarrow \text{epi}(\phi)$ , so by [25, Proposition 4.15], we know that  $\overline{B} \subseteq \text{int}(\text{epi}(\phi_{k_\ell}))$  for all large enough  $\ell$ . This means that for large enough  $\ell$  we have  $\gamma_u(s_{\phi_{k_\ell}}(u)) \in \text{int}(\text{epi}(\phi_{k_\ell}))$ , which is impossible. Therefore  $\tilde{s} = s_\phi(u)$  and the proof is complete.

The fact that we also have  $F_{\phi_k}(u) \rightarrow F_\phi(u)$  is now immediate from its definition.

- (2) Fix  $u$  such that all the functions  $s_{\phi_k}$  differentiable at  $u$ , every  $\text{epi}(\phi_k)$  has a unique outer unit normal at  $F_{\phi_k}(u)$ , and the same holds for  $s_\phi$  and  $\text{epi}(\phi)$ . We first claim that since  $\text{epi}(\phi_k) \rightarrow \text{epi}(\phi)$  and  $F_{\phi_k}(u) \rightarrow F_\phi(u)$  we must have

$$(5.2) \quad n_{\text{epi}(\phi_k)}(F_{\phi_k}(u)) \rightarrow n_{\text{epi}(\phi)}(F_\phi(u)).$$

Indeed, choose any converging subsequence  $n_{\text{epi}(\phi_{k_\ell})} \left( F_{\phi_{k_\ell}}(u) \right) \rightarrow \tilde{n}$ . Every  $(x, t) \in \text{int}(\text{epi}(\phi))$  also belongs to  $\text{epi}(\phi_k)$  for all large enough  $k$ , and therefore

$$\left\langle (x, t) - F_{\phi_{k_\ell}}(u), n_{\text{epi}(\phi_{k_\ell})} \left( F_{\phi_{k_\ell}}(u) \right) \right\rangle \leq 0.$$

Letting  $\ell \rightarrow \infty$  we have  $\langle (x, t) - F_\phi(u), \tilde{n} \rangle \leq 0$ , so  $\tilde{n} = n_{\text{epi}(\phi)}(F_\phi(u))$  since we assumed the outer unit normal is unique. This proves the claim.

Now the convergence  $\nabla s_{\phi_k}(u) \rightarrow \nabla s_\phi(u)$  follows immediately from the explicit formula (5.1). A direct differentiation shows that

$$(5.3) \quad DF_\phi(u) = \left( \begin{array}{c} \nabla s_\phi(u) \otimes u + s_\phi(u) \cdot I_n \\ \hline -\nabla s_\phi(u)/s_\phi(u) \end{array} \right)$$

from which it is also clear that  $DF_{\phi_k}(u) \rightarrow DF_\phi(u)$ .

The last ingredient we need for our proof is a bound on the curvilinear radial function of a simple convex function:  $\square$

**Lemma 5.9.** *Define  $\psi \in \text{Cvx}_n$  by  $\psi(x) = a|x| + b$  for  $a > 0$  and  $b \in \mathbb{R}$ . Then  $s_\psi(u) \leq C \frac{\log(1+|u|)}{|u|}$  for all  $u \in \mathbb{R}^n$ , where  $C > 0$  depends only on  $a$  and  $b$ .*

*Proof.* We choose  $C = \max \left\{ \frac{1}{a}, e^{-b} \right\}$ . Plugging  $a = -\frac{|u|}{1+|u|}$  into the standard inequality  $\log(1+a) \leq a$  we see that  $\log(1+|u|) \geq \frac{|u|}{1+|u|}$ . Therefore for  $\sigma = C \frac{\log(1+|u|)}{|u|}$  we have

$$\begin{aligned} \log\left(\frac{1}{\sigma}\right) &\leq \log\left(\frac{1+|u|}{C}\right) = \log(1+|u|) - \log C \\ &\leq aC \log(1+|u|) + b = \psi(\sigma u). \end{aligned}$$

Therefore  $s_\psi(u) \leq \sigma$  as claimed.  $\square$

With all the ingredient in place we are finally ready to prove Theorem 1.9:

*Proof of Theorem 1.9.* Fix  $\{f_k\}_{k=1}^\infty, f \in \text{LC}_n$  such that  $f_k \rightarrow f$ . Write as usual  $f_k = e^{-\phi_k}$  and  $f = e^{-\phi}$ . Our goal is to prove that for all cosmically continuous functions  $\xi : \mathbb{R}^n \rightarrow \mathbb{R}$  we have

$$\int_{\mathbb{R}^n} \xi d\mu_{f_k} + \int_{\mathbb{R}^n} \bar{\xi} d\mu_{f_k} \rightarrow \int_{\mathbb{R}^n} \xi d\mu_f + \int_{\mathbb{R}^n} \bar{\xi} d\mu_f,$$

which by (4.9) is equivalent to

$$\int_{\partial \text{epi}(\phi_k)} \widehat{\xi}(n_{\text{epi}(\phi_k)}(x, t)) e^{-t} d\mathcal{H}^n(x, t) \rightarrow \int_{\partial \text{epi}(\phi)} \widehat{\xi}(n_{\text{epi}(\phi)}(x, t)) e^{-t} d\mathcal{H}^n(x, t).$$

Using Lemma 5.2(1), we may translate all functions by the same vector  $v$  and assume that  $\phi \in \text{Cvx}_n^{(\epsilon)}$  for some  $0 < \epsilon < \frac{1}{2}$ . This does not change the measures  $(\mu_{f_k}, \nu_{f_k})$  or  $(\mu_f, \nu_f)$ . By part (2) of the same lemma we know that  $\phi_k \in \text{Cvx}_n^{(\epsilon/2)}$  for all large enough  $k$ . We may therefore perform the

change of variables  $(x, t) = F_{\phi_k}(u)$  on the left hand side and  $(x, t) = F_\phi(u)$  on the right hand side, and conclude that our goal is equivalent to

$$\int_{\mathbb{R}^n} \widehat{\xi}(n_{\text{epi}(\phi_k)}(F_{\phi_k}(u))) s_{\phi_k}(u) |JF_{\phi_k}(u)| du \rightarrow \int_{\mathbb{R}^n} \widehat{\xi}(n_{\text{epi}(\phi)}(F_\phi(u))) s_\phi(u) |JF_\phi(u)| du.$$

Here of course  $JF_\phi$  denotes the Jacobian of the map  $F_\phi$ .

By (5.2) we know that  $n_{\text{epi}(\phi_k)}(F_{\phi_k}(u)) \rightarrow n_{\text{epi}(\phi)}(F_\phi(u))$  almost everywhere, and since  $\widehat{\xi}$  is continuous  $\widehat{\xi}(n_{\text{epi}(\phi_k)}(F_{\phi_k}(u))) \rightarrow \widehat{\xi}(n_{\text{epi}(\phi)}(F_\phi(u)))$  almost everywhere. By Proposition 5.8 we also know that  $s_{\phi_k}(u) \rightarrow s_\phi(u)$  and

$$JF_{\phi_k}(u) = \sqrt{\det((DF_{\phi_k}(u))^t DF_{\phi_k}(u))} \rightarrow \sqrt{\det((DF_\phi(u))^t DF_\phi(u))} = JF_\phi(u)$$

almost everywhere. Therefore in order to conclude the proof it is enough to justify the use of the dominated convergence theorem.

Since  $\phi_k \rightarrow \phi$ , there exists  $\psi(x) = a|x| + b$  such that  $\phi_k \geq \psi$  for all  $k$ , and then by Lemma 5.9

$$s_{\phi_k}(u) \leq s_\psi(u) \leq C \frac{\log(1 + |u|)}{|u|}$$

for all  $u \in \mathbb{R}^n$ . Here and after we use  $C > 0$  to denote some constant that does not depend on  $k$  or  $u$ , whose exact value may change from line to line.

We saw in (5.3) that  $DF_{\phi_k}$  is an  $(n+1) \times n$  matrix of the form  $DF_\phi(u) = \begin{pmatrix} A \\ w \end{pmatrix}$  where  $A = \nabla s_\phi(u) \otimes u + s_\phi(u) \cdot I_n$  and  $w = -\nabla s_\phi(u)/s_\phi(u)$ . Therefore

$$(5.4) \quad \begin{aligned} JF_{\phi_k}(u) &= \sqrt{\det((DF_{\phi_k}(u))^t DF_{\phi_k}(u))} = \sqrt{\det(A^t A + w \otimes w)} \leq \|A^t A + w \otimes w\|^{\frac{n}{2}} \\ &\leq \left(\|A\|^2 + |w|^2\right)^{\frac{n}{2}} \leq \left((|\nabla s_\phi(u)| |u| + s_\phi(u))^2 + |w|^2\right)^{\frac{n}{2}}. \end{aligned}$$

Here  $\|\cdot\|$  denotes the operator norm, or the largest singular value.

Using the explicit formula (5.1) and Lemma 5.2(3) we see that

$$|w| = \frac{|\nabla s_{\phi_k}(u)|}{s_{\phi_k}(u)} = s_{\phi_k}(u) \cdot \frac{|n_{\text{epi}(\phi_k)}^{(x)}(x, t)|}{\langle n_{\text{epi}(\phi_k)}(x, t), (x, -1) \rangle} \leq C \frac{\log(1 + |u|)}{|u|} \cdot \frac{1}{\epsilon/2} = C \frac{\log(1 + |u|)}{|u|}.$$

Similarly

$$|\nabla s_{\phi_k}(u)| |u| \leq C \left( \frac{\log(1 + |u|)}{|u|} \right)^2 |u| \leq C \frac{\log^2(1 + |u|)}{|u|}.$$

Plugging these estimates into (5.4) and using the fact that  $\widehat{\xi}$  is bounded on  $\overline{\mathbb{S}^n}$  we obtain

$$\begin{aligned} \left| \widehat{\xi}(n_{\text{epi}(\phi_k)}(F_{\phi_k}(u))) s_{\phi_k}(u) |JF_{\phi_k}(u)| \right| &\leq C \frac{\log(1 + |u|)}{|u|} \cdot \left( \frac{\log^2(1 + |u|) + \log(1 + |u|)}{|u|} \right)^n \\ &\leq C \frac{\max\{\log^{2n+1}(1 + |u|), \log^{n+1}(1 + |u|)\}}{|u|^{n+1}}. \end{aligned}$$

As this function is integrable on  $\mathbb{R}^n$  the use of dominated convergence is justified, and the proof is complete.  $\square$

## 6. CONTINUITY OF SOLUTION TO THE MINKOWSKI PROBLEM

In this section we prove Theorem 1.10. We begin by collecting some preliminary facts. First, we will need the following result of Colesanti and Fragalà:

**Proposition 6.1** ([5, Proposition 3.11]). *For all  $f \in \text{LC}_n$  we have*

$$\delta(f, f) = n \int f + \int f \log f.$$

Next, we need an isoperimetric inequality for log-concave functions, first shown in [23]:

**Proposition 6.2** ([23, Proposition 27]). *For all  $f \in \text{LC}_n$  we have*

$$\delta(f, \mathbf{1}_{B_2^n}) \geq c_n (\max f)^{\frac{1}{n}} \left( \int f \right)^{1 - \frac{1}{n}},$$

for some constant  $c_n > 0$  that depends only on the dimension  $n$ .

In fact, it was shown in [23] that optimal value of the constant  $c_n$  is attained when  $f(x) = e^{-|x|}$ . For us however the exact value of  $c_n$  and the nature of the minimizers will not play any role.

Finally, we need the following strengthening of Lemma 3.4(1):

**Lemma 6.3.** *Fix  $\{f_k\}_{k=1}^\infty, f \in \text{LC}_n$  and assume that  $(\mu_{f_k}, \nu_{f_k}) \rightarrow (\mu_f, \nu_f)$  cosmically. Then there exists a constant  $c > 0$ , such that for all  $y \in \mathbb{R}^n$  and all  $k$  we have*

$$\int_{\mathbb{R}^n} |\langle x, y \rangle| d\mu_{f_k}(x) + \int_{\mathbb{S}^{n-1}} |\langle x, y \rangle| d\nu_{f_k}(x) \geq c |y|.$$

Of course, the main point of this strengthening is that the constant  $c$  is not allowed to depend on  $k$ .

*Proof.* Define a sequence of functions  $\Phi_k : \mathbb{R}^n \rightarrow \mathbb{R}$  by

$$\Phi_k(y) = \int_{\mathbb{R}^n} |\langle x, y \rangle| d\mu_{f_k}(x) + \int_{\mathbb{S}^{n-1}} |\langle x, y \rangle| d\nu_{f_k}(x),$$

and similarly define

$$\Phi(y) = \int_{\mathbb{R}^n} |\langle x, y \rangle| d\mu_f(x) + \int_{\mathbb{S}^{n-1}} |\langle x, y \rangle| d\nu_f(x).$$

By Lemma 3.4, there exist positive constants  $\{c_k\}_{k=0}^\infty$  such that for all  $y \in \mathbb{R}^n$  we have  $\Phi(y) \geq c_0 |y|$  and  $\Phi_k(y) \geq c_k |y|$ . Since the function  $\ell_y(x) = |\langle x, y \rangle|$  is cosmically continuous and  $\overline{\ell_y} = \ell_y$ , it follows from the definition of cosmic convergence that  $\Phi_k(y) \rightarrow \Phi(y)$  for all  $y \in \mathbb{R}^n$ . Since the functions  $\{\Phi_k\}_{k=1}^\infty$  are convex and finite, the pointwise convergence  $\Phi_k \rightarrow \Phi$  implies uniform convergence on compact subsets of  $\mathbb{R}^n$  (see e.g. [25, Theorem 7.17(c)]). In particular

$$\min_{y \in \mathbb{S}^{n-1}} \Phi_k(y) \rightarrow \min_{y \in \mathbb{S}^{n-1}} \Phi(y) \geq c_0.$$

Therefore there exists  $K > 0$  such that for all  $k > K$  we have and all  $y \in \mathbb{S}^{n-1}$  we have  $\Phi_k(y) \geq \frac{c_0}{2}$ . It is therefore enough to choose  $c = \min \left\{ \frac{c_0}{2}, c_1, c_2, \dots, c_K \right\}$ .  $\square$

We are now ready to prove Theorem 1.10:

*Proof of Theorem 1.10.* Assume by contradiction that we are given  $\{f_k\}_{k=1}^\infty, f \in \text{LC}_n$  such that  $(\mu_{f_k}, \nu_{f_k}) \rightarrow (\mu_f, \nu_f)$  cosmically but  $f_k \not\rightarrow f$ , even up to translations. Fix any metric  $d$  on  $\text{LC}_n$  which generates our notion of convergence (examples to such metrics were given in [21]), and define

$$\tilde{d}(f, g) = \inf_{v \in \mathbb{R}^n} d(f, \tau_v g)$$

where  $\tau_v g(x) = g(x + v)$ . Our assumption precisely means that  $\tilde{d}(f_k, f) \not\rightarrow 0$ . By passing to a subsequence we may assume without loss of generality that there exists an  $\epsilon > 0$  such that  $\tilde{d}(f_k, f) \geq \epsilon$  for all  $k$ .

Write as usual  $f_k = e^{-\phi_k}$  and  $f = e^{-\phi}$ . Since we allow translations, we may assume without loss of generality that  $\min \phi_k = \phi_k(0)$  for all  $k$ . Combining Lemma 6.3 with the second part of Lemma 3.4 we see that there exists a constant  $c > 0$  such for every  $k$ , every  $\psi \in \text{Cvx}_n$  with  $\min \psi = \psi(0)$  and every  $x \in \mathbb{R}^n$  we have

$$\psi(x) \geq \frac{1}{\mu_{f_k}(\mathbb{R}^n)} \left( c|x| - \int_{\mathbb{R}^n} \psi^* d\mu_{f_k} - \int_{\mathbb{S}^{n-1}} \overline{\psi^*} d\nu_{f_k} \right).$$

We now choose  $\psi = \phi_k$  and use the first variation formula 1.1 and Proposition 6.1 to conclude that

$$\begin{aligned} \phi_k(x) &\geq \frac{1}{\mu_{f_k}(\mathbb{R}^n)} \left( c|x| - \int_{\mathbb{R}^n} \phi_k^* d\mu_{f_k} - \int_{\mathbb{S}^{n-1}} \overline{\phi_k^*} d\nu_{f_k} \right) = \frac{1}{\mu_{f_k}(\mathbb{R}^n)} (c|x| - \delta(f_k, f_k)) \\ (6.1) \quad &= \frac{1}{\mu_{f_k}(\mathbb{R}^n)} \left( c|x| - n \int f_k - \int f_k \log f_k \right) = \frac{c}{\int f_k} |x| - n - \frac{\int f_k \log f_k}{\int f_k}. \end{aligned}$$

Note that we also used the fact that  $\mu_{f_k}(\mathbb{R}^n) = \int f_k$  which is obvious from the definition of  $\mu_{f_k}$ .

Using the fact that the constant function  $\mathbf{1}$  is cosmically continuous we obtain

$$\int f_k = \mu_{f_k}(\mathbb{R}^n) = \int \mathbf{1} d\mu_{f_k} + \int \overline{\mathbf{1}} d\nu_{f_k} \rightarrow \int \mathbf{1} d\mu_f + \int \overline{\mathbf{1}} d\nu_f = \mu_f(\mathbb{R}^n) = \int f,$$

so in particular  $\{\int f_k\}_{k=1}^\infty$  is bounded from above and from below by some positive constants. Similarly using the fact that  $x \mapsto |x|$  is cosmically continuous and (1.1) we have

$$\delta(f_k, \mathbf{1}_{B_2^n}) = \int |x| d\mu_{f_k} + \int |x| d\nu_{f_k} \rightarrow \int |x| d\mu_f + \int |x| d\nu_f = \delta(f, \mathbf{1}_{B_2^n}),$$

so the sequence  $\{\delta(f_k, \mathbf{1}_{B_2^n})\}_{k=1}^\infty$  is also bounded. Using Proposition 6.2 we have

$$\max f_k \leq C_n \frac{\delta(f_k, \mathbf{1}_{B_2^n})^n}{(\int f_k)^{n-1}},$$

so the sequence  $\{\max f_k\}_{k=1}^\infty$  is also bounded from above. Therefore

$$\frac{\int f_k \log f_k}{\int f_k} \leq \frac{\max(\log f_k) \int f_k}{\int f_k} = \log(\max f_k)$$

is also bounded from above. Plugging all these estimates into (6.1) we conclude that the sequence  $\{\phi_k\}_{k=1}^\infty$  is uniformly coercive. Since  $\{\int e^{-\phi_k}\}_{k=1}^\infty = \{\int f_k\}_{k=1}^\infty$  is bounded from below by a positive constant, the same argument as in Proposition 3.6 shows that  $\sup_k(\min \phi_k) < \infty$ .

Theorem 3.3 now allows us to pass to another sub-sequence and assume without loss of generality that  $f_k \rightarrow g$  for some  $g \in \text{LC}_n$  (the fact that  $\int g > 0$  was not part of the theorem, but it follows from the fact that  $\int g = \lim_{k \rightarrow \infty} \int f_k$ ). By Theorem 1.9 it follows that  $(\mu_{f_k}, \nu_{f_k}) \rightarrow (\mu_g, \nu_g)$  cosmically. But we also have  $(\mu_{f_k}, \nu_{f_k}) \rightarrow (\mu_f, \nu_f)$ , and since the cosmic limit is clearly unique we have  $(\mu_g, \nu_g) = (\mu_f, \nu_f)$ . But then by the uniqueness part of Theorem 1.5 we must have  $g = \tau_v f$  for some  $v \in \mathbb{R}^n$ . This implies that

$$\tilde{d}(f_k, f) = \tilde{d}(f_k, g) \leq d(f_k, g) \rightarrow 0,$$

contradicting our assumption that  $\tilde{d}(f_k, f) \geq \epsilon$  for all  $k$ . □

## REFERENCES

- [1] Shiri Artstein-Avidan and Vitali Milman. The concept of duality in convex analysis, and the characterization of the Legendre transform. *Annals of Mathematics*, 169(2):661–674, March 2009.
- [2] Shiri Artstein-Avidan and Vitali Milman. Hidden structures in the class of convex functions and a new duality transform. *Journal of the European Mathematical Society*, 13(4):975–1004, May 2011.
- [3] Károly J. Böröczky, Erwin Lutwak, Deane Yang, and Gaoyong Zhang. The log-Brunn–Minkowski inequality. *Advances in Mathematics*, 231(3-4):1974–1997, October 2012.
- [4] Frank H. Clarke. *Optimization and Nonsmooth Analysis*. Society for Industrial and Applied Mathematics, January 1990.
- [5] Andrea Colesanti and Ilaria Fragalà. The first variation of the total mass of log-concave functions and related inequalities. *Advances in Mathematics*, 244:708–749, September 2013.
- [6] Andrea Colesanti, Monika Ludwig, and Fabian Mussnig. Valuations on Convex Functions. *International Mathematics Research Notices*, 2019(8):2384–2410, April 2019.
- [7] Andrea Colesanti, Monika Ludwig, and Fabian Mussnig. The Hadwiger theorem on convex functions, III: Steiner formulas and mixed Monge–Ampère measures. *Calculus of Variations and Partial Differential Equations*, 61(5):181, October 2022.
- [8] Andrea Colesanti, Monika Ludwig, and Fabian Mussnig. The Hadwiger theorem on convex functions, II: Cauchy–Kubota formulas. (arXiv:2109.09434), May 2023.
- [9] Andrea Colesanti, Monika Ludwig, and Fabian Mussnig. The Hadwiger theorem on convex functions, IV: The Klain approach. *Advances in Mathematics*, 413:108832, January 2023.
- [10] Andrea Colesanti, Monika Ludwig, and Fabian Mussnig. The Hadwiger Theorem on Convex Functions, I. *Geometric and Functional Analysis*, 34(6):1839–1898, December 2024.
- [11] Dario Cordero-Erausquin and Bo’az Klartag. Moment measures. *Journal of Functional Analysis*, 268(12):3834–3866, June 2015.
- [12] Lawrence C. Evans and Ronald F. Gariepy. *Measure Theory and Fine Properties of Functions*. Textbooks in Mathematics. CRC press Taylor & Francis group, Boca Raton, revised edition edition, 2015.
- [13] Niufa Fang, Deping Ye, and Zengle Zhang. The Riesz  $\alpha$ -energy of log-concave functions and related Minkowski problem. (arXiv:2408.16141), August 2024.
- [14] Niufa Fang, Deping Ye, Zengle Zhang, and Yiming Zhao. Dual Orlicz curvature measures for log-concave functions and their Minkowski problems. *Calculus of Variations and Partial Differential Equations*, 64(2):44, March 2025.
- [15] Yong Huang, Jiaqian Liu, Dongmeng Xi, and Yiming Zhao. Dual curvature measures for log-concave functions. *Journal of Differential Geometry*, 128(2), October 2024.
- [16] Yong Huang, Dongmeng Xi, and Yiming Zhao. The Minkowski problem in Gaussian probability space. *Advances in Mathematics*, 385:107769, July 2021.
- [17] Bo’az Klartag. Logarithmically-Concave Moment Measures I. In Bo’az Klartag and Emanuel Milman, editors, *Geometric Aspects of Functional Analysis*, volume 2116, pages 231–260. Springer International Publishing, Cham, 2014.



- [18] Bo'az Klartag and Joseph Lehec. Affirmative Resolution of Bourgain's Slicing Problem using Guan's Bound. (arXiv:2412.15044), December 2024.
- [19] Bo'az Klartag and Vitali D. Milman. Geometry of Log-concave Functions and Measures. *Geometriae Dedicata*, 112(1):169–182, April 2005.
- [20] Liudmyla Kryvonos and Dylan Langharst. Weighted Minkowski's existence theorem and projection bodies. June 2023.
- [21] Ben Li and Fabian Mussnig. Metrics and Isometries for Convex Functions. *International Mathematics Research Notices*, 2022(18):14496–14563, September 2022.
- [22] Georges Matheron. La formule de Steiner pour les érosions. *Journal of Applied Probability*, 15(1):126–135, March 1978.
- [23] Vitali Milman and Liran Rotem. Mixed integrals and related inequalities. *Journal of Functional Analysis*, 264(2):570–604, January 2013.
- [24] Vitali D. Milman. Geometrization of Probability. In Mikhail Kapranov, Yuri Ivanovich Manin, Pieter Moree, Sergiy Kolyada, and Leonid Potyagailo, editors, *Geometry and Dynamics of Groups and Spaces*, volume 265, pages 647–667. Birkhäuser Basel, Basel, 2008.
- [25] R. Tyrrell Rockafellar and Roger J. B. Wets. *Variational Analysis*, volume 317 of *Grundlehren Der Mathematischen Wissenschaften*. Springer Berlin Heidelberg, Berlin, Heidelberg, 1998.
- [26] Liran Rotem. On the Mean Width of Log-Concave Functions. In Bo'az Klartag, Shahar Mendelson, and Vitali D. Milman, editors, *Geometric Aspects of Functional Analysis*, volume 2050, pages 355–372. Springer Berlin Heidelberg, Berlin, Heidelberg, 2012.
- [27] Liran Rotem. Support functions and mean width for  $\alpha$ -concave functions. *Advances in Mathematics*, 243:168–186, August 2013.
- [28] Liran Rotem. Surface area measures of log-concave functions. *Journal d'Analyse Mathématique*, 147(1):373–400, October 2022.
- [29] Liran Rotem. The Anisotropic Total Variation and Surface Area Measures. In Ronen Eldan, Bo'az Klartag, Alexander Litvak, and Emanuel Milman, editors, *Geometric Aspects of Functional Analysis*, volume 2327 of *Lecture Notes in Mathematics*, pages 297–312. Springer International Publishing, Cham, 2023.
- [30] Filippo Santambrogio. Dealing with moment measures via entropy and optimal transport. *Journal of Functional Analysis*, 271(2):418–436, July 2016.
- [31] Rolf Schneider. *Convex Bodies: The Brunn-Minkowski Theory*. Cambridge University Press, 2 edition, October 2013.
- [32] Rolf Schneider. A weighted Minkowski theorem for pseudo-cones. *Advances in Mathematics*, 450:109760, July 2024.
- [33] Jacopo Ulivelli. Entire Monge-Ampère equations and weighted Minkowski problems. (arXiv:2310.02787), October 2023.
- [34] Jacopo Ulivelli. First variation of functional Wulff shapes. (arXiv:2312.11172), May 2024.
- [35] Hejun Wang, Niufa Fang, and Jiazhu Zhou. Continuity of the solution to the dual Minkowski problem for negative indices. *Proceedings of the American Mathematical Society*, 147(3):1299–1312, December 2018.
- [36] Hejun Wang, Niufa Fang, and Jiazhu Zhou. Continuity of the solution to the even logarithmic Minkowski problem in the plane. *Science China Mathematics*, 62(7):1419–1428, July 2019.
- [37] Robert A. Wijsman. Convergence of sequences of convex sets, cones and functions. *Bulletin of the American Mathematical Society*, 70:186–188, 1964.
- [38] Robert A. Wijsman. Convergence of sequences of convex sets, cones and functions. II. *Transactions of the American Mathematical Society*, 123:32–45, 1966.
- [39] Guangxian Zhu. Continuity of the solution to the  $L_p$  Minkowski problem. *Proceedings of the American Mathematical Society*, 145(1):379–386, July 2016.

TECHNION ISRAEL INSTITUTE OF TECHNOLOGY, TECHNION CITY, HAIFA 3200003, ISRAEL

*Email address:* `fatomer@campus.technion.ac.il`

TECHNION ISRAEL INSTITUTE OF TECHNOLOGY, TECHNION CITY, HAIFA 3200003, ISRAEL

*Email address:* `lrotem@technion.ac.il`