

# CLASSICAL YANG-BAXTER EQUATIONS AND NIJENHUIS OPERATORS FOR LIE ALGEBRAS

HAIYING LI AND TIANSHUI MA \*

**ABSTRACT.** In this paper the conditions that when a Lie algebra is Nijenhuis are investigated. Furthermore all the Nijenhuis operators on  $\mathfrak{sl}_2$  under the standard Cartan-Weyl basis are given. On the other hand, the notion of classical  $P$ -Nijenhuis Yang-Baxter equations is introduced by means of the bialgebraic theory for Nijenhuis Lie algebras.

## 1. INTRODUCTION

One of the aims of this paper is to prove when a Lie algebra is Nijenhuis, which means that Lie algebras and Nijenhuis operators are closely related naturally. The other aim is to find the compatible conditions between classical Yang-Baxter equations and Nijenhuis operators for Lie algebras by establishing the bialgebraic theory for Nijenhuis Lie algebras.

As is well known that a pair  $(\mathfrak{Q}, [, ])$  is called a **Lie algebra** if the linear map  $[,] : \mathfrak{Q} \otimes \mathfrak{Q} \rightarrow \mathfrak{Q}$  satisfies

$$[x, y] = -[y, x], \quad (1)$$

$$[[x, y], z] + [[y, z], x] + [[z, x], y] = 0, \quad (2)$$

where  $x, y, z \in \mathfrak{Q}$ . The most best-known quantum group is  $U_q(\mathfrak{sl}_2)$ , which is deformation of the universal enveloping algebra of the 3-dimensional complex Lie algebra  $\mathfrak{sl}_2$ .  $\mathfrak{sl}_2$  is the smallest simple Lie algebra. Concretely, the structure  $[,]$  of  $\mathfrak{sl}_2$  with respect to the standard Cartan-Weyl basis  $\{e, f, g\}$  can be given as follows:

$$\begin{array}{c|ccc} [, ] & e & f & g \\ \hline e & 0 & g & -2e \\ f & -g & 0 & 2f \\ g & 2e & -2f & 0 \end{array}.$$

Recently Bai, Guo, Liu and the second named author established the bialgebraic theory for Rota-Baxter Lie algebras [2].

A **Nijenhuis Lie algebra** [7, 12] is a triple  $((\mathfrak{Q}, [, ]), N)$  including a Lie algebra  $(\mathfrak{Q}, [, ])$  and a Nijenhuis operator  $N$  on  $\mathfrak{Q}$ , i.e., a linear map  $N : \mathfrak{Q} \rightarrow \mathfrak{Q}$  satisfies

$$[N(x), N(y)] + N^2[x, y] = N[N(x), y] + N[x, N(y)], \quad \forall x, y \in \mathfrak{Q}. \quad (3)$$

*Date:* February 27, 2025.

*2020 Mathematics Subject Classification.* 17B38, 17B62, 16T25, 17A30.

*Key words and phrases.* classical Yang-Baxter equations; symplectic structure; Nijenhuis operators; Lie bialgebras.

\*Corresponding author.

Nijenhuis Lie algebras have applied to finite or infinite integrable systems [7, 10], cohomology and deformations theory [5, 12], almost complex structures [11], generalized Nijenhuis algebras [8], Poisson-Nijenhuis groupoid [4], etc.

A Lie bialgebra is composed of a Lie algebra and a Lie coalgebra together with a cocycle condition, which is introduced in the study of Hamiltonian mechanics and Poisson-Lie groups [6]. Lie bialgebras can be characterized by Manin triples of Lie algebras. The notion of classical  $r$ -matrix was introduced in the classical inverse scattering method as the natural foundation of the Hamiltonian approach. Semenov-Tian-Shansky first gave the relation between classical  $r$ -matrix and Rota-Baxter operator of weight 0 [1, 14]. Nijenhuis operators and Rota-Baxter operators are similar in form. Following the discussion above, there are two interesting questions: When a Lie (co)algebra is Nijenhuis? Are there suitable compatible conditions between classical  $r$ -matrix and Nijenhuis operator which can induce a bialgebraic structure on a Nijenhuis Lie algebra?

In this paper, we will discuss the questions above.

In Section 2, we give all the Nijenhuis operators on  $\mathfrak{sl}_2$  with respect to the standard Cartan-Weyl basis (see Theorem 2.1). Section 3 is devoted to finding that Lie algebras are very closely related to Nijenhuis operators. In subsection 3.1, all the quasitriangular structures on  $\mathfrak{sl}_2$  are provided (see Example 3.3). In subsection 3.2, we present all the symplectic structures on  $\mathfrak{sl}_2$  (see Example 3.9) and prove that a dual quasitriangular pre-Lie bialgebra can give rise to a symplectic Lie algebra (see Proposition 3.10). On the basis of the previous preparations, in subsection 3.3, we obtain that a symplectic Lie algebra together with both a quasitriangular and a dual quasitriangular Lie bialgebra will possess Nijenhuis operators (see Theorem 3.11). And meanwhile we apply the result in Theorem 3.11 to  $\mathfrak{sl}_2$  (see Example 3.12).

In order to establish the bialgebraic structure on a Nijenhuis Lie algebra, in Section 4, we present some useful explorations. In subsection 4.1, we introduce the notion of the representation of a Nijenhuis Lie algebra (see Definition 4.1) and study the dual representation (see Lemma 4.4). In subsection 4.2, we present the concepts of a Nijenhuis Lie coalgebra and its corepresentation through compatible Lie coalgebras (see Theorem 4.10, Definitions 4.11 and 4.13). In subsection 4.3, we consider the dual case of Theorem 3.11 (see Theorem 4.18), which provides a method to obtain Nijenhuis Lie coalgebras.

In Section 5, we focus on the structure of Nijenhuis Lie bialgebras. In subsection 5.1, we introduce the notion of Nijenhuis Lie bialgebras (see Definition 5.1) and also give the equivalent characterizations by using matched pairs of Nijenhuis Lie algebras (see Theorem 5.7) and Manin triples of Nijenhuis Frobenius Lie algebra (see Theorem 5.12), which are all compatible with the case of Lie algebras. Some examples of Nijenhuis Lie bialgebras on  $\mathfrak{sl}_2$  are presented (see Example 5.3). In subsection 5.2, we consider the quasitriangular case. Quasitriangular Nijenhuis Lie bialgebras and classical  $P$ -Nijenhuis Yang-Baxter equations are defined (see Theorem 5.15 and Definition 5.17). We introduce the notion of relative Rota-Baxter operators and prove that it can be used to construct Nijenhuis Lie bialgebras

(see Corollary 5.20), especially on semi-direct product Nijenhuis Lie algebras (see Theorem 5.22).

**Notations:** Throughout this paper, all vector spaces, tensor products, and linear homomorphisms are over the complex number field  $\mathbb{C}$ . We denote by  $\text{id}_M$  the identity map from  $M$  to  $M$ .

## 2. NIJENHUIS OPERATORS ON $\mathfrak{sl}_2$

In this section, we give all the Nijenhuis operators on  $\mathfrak{sl}_2$  with respect to the standard Cartan-Weyl basis.

**Theorem 2.1.** *Let  $\{e, f, g\}$  be the standard Cartan-Weyl basis. Then all the Nijenhuis operators on  $(\mathfrak{sl}_2, [, ])$  are given as follows.*

$$\begin{aligned}
 (1) & \begin{pmatrix} k_1 & 0 & 0 \\ 0 & k_5 & k_6 \\ 0 & 0 & k_5 \end{pmatrix}, & (2) & \begin{pmatrix} k_1 & 0 & k_3 \\ 0 & k_5 & -\frac{(k_1-k_9)(k_5-k_9)}{4k_3} \\ 0 & 0 & k_9 \end{pmatrix}, \\
 (3) & \begin{pmatrix} k_1 & k_2 & \frac{k_2(k_9-k_5)}{k_8} - \frac{k_8}{4} \\ 0 & k_5 & \frac{k_8(k_1-k_9)}{4k_2} \\ 0 & k_8 & k_9 \end{pmatrix}, & (4) & \begin{pmatrix} k_1 & k_2 & k_3 \\ 0 & k_5 & 0 \\ 0 & 0 & k_5 \end{pmatrix} \quad (k_2 \neq 0), \\
 (5) & \begin{pmatrix} k_1 & k_2 & \frac{k_2k_4}{k_7} \\ k_4 & k_5 & \frac{k_4(k_5-k_1)}{k_7} - \frac{k_7}{4} \\ k_7 & -\frac{4k_2k_4}{k_7} & k_5 \end{pmatrix}, & (6) & \begin{pmatrix} k_1 & k_2 & k_3 \\ 0 & k_5 & k_6 \\ -4k_6 & 0 & k_5 \end{pmatrix} \quad (k_6 \neq 0), \\
 (7) & \begin{pmatrix} k_1 & 0 & k_3 \\ k_4 & k_5 & k_6 \\ 0 & -4k_3 & k_1 \end{pmatrix}, \\
 (8) & \begin{pmatrix} k_1 & k_2 & k_3 \\ k_4 & k_9 + \frac{4k_3k_4}{k_7} - \frac{4k_2k_4^2}{k_7^2} & \frac{k_4(k_9-k_1)}{k_7} - \frac{k_7}{4} \\ k_7 & -\frac{4k_2k_4}{k_7} & k_9 \end{pmatrix} \quad (k_4 \neq 0, k_2 \neq \frac{k_3k_7}{k_4}),
 \end{aligned}$$

where  $k_i, i = 1, 2, \dots, 9$  are parameters.

*Proof.* By a direct computation, see appendix A. □

**Corollary 2.2.** *The diagonalisable Nijenhuis operator on  $(\mathfrak{sl}_2, [, ])$  under the standard Cartan-Weyl basis  $\{e, f, g\}$  must be*

$$\begin{pmatrix} k_1 & 0 & 0 \\ 0 & k_5 & 0 \\ 0 & 0 & k_5 \end{pmatrix} \text{ or } \begin{pmatrix} k_1 & 0 & 0 \\ 0 & k_5 & 0 \\ 0 & 0 & k_1 \end{pmatrix},$$

where  $k_1, k_5$  are parameters.

## 3. NIJENHUIS OPERATORS FROM SYMPLECTIC LIE ALGEBRAS

In order to obtain the main result in subsection 3.3, we need the following preparations in subsections 3.1 and 3.2.

**3.1. Quasitriangular Lie bialgebras.** Let us recall from [6] the definition of Lie bialgebra. A **Lie coalgebra** is a pair  $(\mathfrak{Q}, \delta)$ , where  $\mathfrak{Q}$  is a vector space and  $\delta : \mathfrak{Q} \longrightarrow \mathfrak{Q} \otimes \mathfrak{Q}$  (we use the Sweedler notation [13]: write  $\delta(x) = x_{(1)} \otimes x_{(2)}$  ( $= \sum x_{(1)} \otimes x_{(2)}$ )) is a linear map such that for all  $x \in \mathfrak{Q}$ ,

$$x_{(1)} \otimes x_{(2)} = -x_{(2)} \otimes x_{(1)}, \quad (4)$$

$$x_{(1)} \otimes x_{(2)(1)} \otimes x_{(2)(2)} + x_{(2)(1)} \otimes x_{(2)(2)} \otimes x_{(1)} + x_{(2)(2)} \otimes x_{(1)} \otimes x_{(2)(1)} = 0. \quad (5)$$

A **Lie bialgebra** is a triple  $(\mathfrak{Q}, [, ], \delta)$ , where the pair  $(\mathfrak{Q}, [, ])$  is a Lie algebra, and the pair  $(\mathfrak{Q}, \delta)$  is a Lie coalgebra, such that, for all  $x, y \in \mathfrak{Q}$ ,

$$\delta([x, y]) = [x, y_{(1)}] \otimes y_{(2)} + y_{(1)} \otimes [x, y_{(2)}] - [y, x_{(1)}] \otimes x_{(2)} - x_{(1)} \otimes [y, x_{(2)}]. \quad (6)$$

For a given element  $r = r^1 \otimes r^2 \in \mathfrak{Q} \otimes \mathfrak{Q}$ , define

$$\delta(x) := \delta_r(x) := r^1 \otimes [x, r^2] + [x, r^1] \otimes r^2, \quad \forall x \in \mathfrak{Q}. \quad (7)$$

**Remark 3.1.** [9] Let  $(\mathfrak{Q}, [, ])$  be a Lie algebra and  $r \in \mathfrak{Q} \otimes \mathfrak{Q}$ . If  $\delta : \mathfrak{Q} \longrightarrow \mathfrak{Q} \otimes \mathfrak{Q}$  is given by Eq.(7), then  $\delta$  satisfies Eq.(6).  $(\mathfrak{Q}, \delta)$  is a Lie coalgebra if and only if for all  $x \in \mathfrak{Q}$ ,

$$(ad_x \otimes \text{id} + \text{id} \otimes ad_x)(r + r^\sigma) = 0,$$

$$(ad_x \otimes \text{id} \otimes \text{id} - \text{id} \otimes ad_x \otimes \text{id} - \text{id} \otimes \text{id} \otimes ad_x)([r_{12}, r_{13}] + [r_{13}, r_{23}] + [r_{12}, r_{23}]) = 0,$$

where  $[r_{12}, r_{13}] = [r^1, \bar{r}^1] \otimes r^2 \otimes \bar{r}^2$ ,  $[r_{13}, r_{23}] = r^1 \otimes \bar{r}^1 \otimes [r^2, \bar{r}^2]$ ,  $[r_{23}, r_{12}] = r^1 \otimes [r^2, \bar{r}^1] \otimes \bar{r}^2$ ,  $\bar{r} = r$  and  $r^\sigma = r^2 \otimes r^1$ .

**Definition 3.2.** [9] Let  $(\mathfrak{Q}, [, ])$  be a Lie algebra. If  $r \in \mathfrak{Q} \otimes \mathfrak{Q}$  is an antisymmetric solution (in the sense of  $r = -r^\sigma$ ) of the **classical Yang-Baxter equation** (cYBe for short) in  $(\mathfrak{Q}, [, ])$ :

$$[r_{12}, r_{13}] + [r_{13}, r_{23}] + [r_{12}, r_{23}] = 0, \quad (8)$$

then  $(\mathfrak{Q}, [, ], \delta)$  is a Lie bialgebra, where the linear map  $\delta = \delta_r$  is defined by Eq. (7). In this case, we call this Lie bialgebra **quasitriangular** and denoted by  $(\mathfrak{Q}, [, ], r, \delta_r)$ .

By a direct computation, one can get

**Example 3.3.** Let  $\lambda$  be a parameter. All the quasitriangular Lie bialgebraic structures on  $(\mathfrak{sl}_2, [, ])$  are  $(\mathfrak{sl}_2, [, ], r, \delta_r)$ , where

$$r = \lambda(f \otimes g - g \otimes f)$$

and

$$\begin{cases} \delta_r(e) = 2\lambda(e \otimes f - f \otimes e), \\ \delta_r(f) = 0, \\ \delta_r(g) = 2\lambda(g \otimes f - f \otimes g) \end{cases}.$$

**3.2. Symplectic Lie algebras.** In this subsection, we prove that a dual quasitriangular Lie bialgebra can give rise to a symplectic Lie algebra and also derive all the symplectic structures on  $\mathfrak{sl}_2$ .

**Definition 3.4.** [9] Let  $(\mathfrak{Q}, \delta)$  be a Lie coalgebra and  $\omega \in (\mathfrak{Q} \otimes \mathfrak{Q})^*$ . We call the equation below a **classical co-Yang-Baxter equation in  $(\mathfrak{Q}, \delta)$** :

$$\omega(x_{(1)}, y)\omega(x_{(2)}, z) + \omega(x, z_{(1)})\omega(y, z_{(2)}) + \omega(x, y_{(1)})\omega(y_{(2)}, z) = 0, \quad (9)$$

where  $x, y, z \in \mathfrak{Q}$ .

**Example 3.5.** All the skew-symmetric solutions of the classical co-Yang-Baxter equation in the Lie coalgebra in Example 3.3 are given below:

$$\begin{array}{c|ccc} \omega & e & f & g \\ \hline e & 0 & \nu & \kappa \\ f & -\nu & 0 & \chi \\ g & -\kappa & -\chi & 0 \end{array},$$

where  $\nu, \kappa, \chi$  are parameters.

**Definition 3.6.** [9] Let  $(\mathfrak{Q}, \delta)$  be a Lie coalgebra and  $\omega \in (\mathfrak{Q} \otimes \mathfrak{Q})^*$ . If  $\omega$  is a skew-symmetric solution (in the sense of  $\omega(x, y) = -\omega(y, x)$ ) of the classical co-Yang-Baxter equation, then  $(\mathfrak{Q}, [, ]_\omega, \delta)$  is a Lie bialgebra, where  $[, ]_\omega$  is defined by

$$[x, y]_\omega = x_{(1)}\omega(x_{(2)}, y) - y_{(1)}\omega(x, y_{(2)}). \quad (10)$$

In this case, we call this Lie bialgebra **dual quasitriangular** and denoted by  $(\mathfrak{Q}, \delta, \omega, [, ]_\omega)$ .

**Proposition 3.7.** [9] Let  $(\mathfrak{Q}, \delta)$  be a coalgebra and  $\omega \in (\mathfrak{Q} \otimes \mathfrak{Q})^*$  skew-symmetric. Then the quadruple  $(\mathfrak{Q}, \delta, \omega, [, ]_\omega)$ , where  $[, ]_\omega$  is defined by Eq.(10), is a dual quasitriangular Lie bialgebra if and only if, for all  $x, y, z \in \mathfrak{Q}$ ,

$$\omega([x, y]_\omega, z) = \omega(x, z_1)\omega(y, z_2) \quad (11)$$

or

$$\omega(x, [y, z]_\omega) = -\omega(x_1, y)\omega(x_2, z) \quad (12)$$

holds.

Next we recall from [3] the notion of a symplectic Lie algebra.

**Definition 3.8.** Let  $(\mathfrak{Q}, [, ])$  be a Lie algebra and  $\omega \in (\mathfrak{Q} \otimes \mathfrak{Q})^*$  be skew-symmetric. If for all  $x, y, z \in \mathfrak{Q}$ , the equation below holds:

$$\omega([x, y], z) + \omega([y, z], x) + \omega([z, x], y) = 0, \quad (13)$$

then  $(\mathfrak{Q}, [, ], \omega)$  is a **symplectic Lie algebra** and denoted by  $(\mathfrak{Q}, [, ], \omega)$ .

**Example 3.9.** All the symplectic structures on  $(\mathfrak{sl}_2, [, ])$  are given below:

$$\begin{array}{c|ccc} \omega & e & f & g \\ \hline e & 0 & 0 & \kappa \\ f & 0 & 0 & -2\kappa \\ g & -\kappa & 2\kappa & 0 \end{array},$$

where  $\kappa$  is a parameter.

A symplectic Lie algebra can be obtained by a dual quasitriangular Lie bialgebra.

**Proposition 3.10.** *If  $(\mathfrak{Q}, \delta, \omega, [, ]_\omega)$ , where  $[, ]_\omega$  is defined by Eq.(10), is a dual quasitriangular Lie bialgebra, then  $(\mathfrak{Q}, [, ]_\omega, \omega)$  is a symplectic Lie algebra.*

*Proof.* By Proposition 3.7, we know that  $(\mathfrak{Q}, [, ]_\omega)$  is a Lie algebra. For all  $x, y, z \in \mathfrak{Q}$ , we have

$$\begin{aligned} & \omega([x, y]_\omega, z) + \omega([y, z]_\omega, x) + \omega([z, x]_\omega, y) \\ & \stackrel{(11)}{=} \omega(x, z_{(1)})\omega(y, z_{(2)}) + \omega(y, x_{(1)})\omega(z, x_{(2)}) + \omega(z, y_{(1)})\omega(x, y_{(2)}) \\ & \stackrel{(4)}{=} \omega(x_{(1)}, y)\omega(x_{(2)}, z) + \omega(x, z_{(1)})\omega(y, z_{(2)}) + \omega(x, y_{(1)})\omega(y_{(2)}, z) \\ & \quad \text{(by the skew-symmetry of } \omega) \\ & \stackrel{(9)}{=} 0, \end{aligned}$$

as we needed. □

**3.3. When Lie algebras are Nijenhuis.** A Nijenhuis Lie algebra can be investigated by the following way.

**Theorem 3.11.** *Let  $(\mathfrak{Q}, [, ], \omega)$  be a symplectic Lie algebra,  $r$  be an element in  $\mathfrak{Q} \otimes \mathfrak{Q}$  such that  $(\mathfrak{Q}, [, ], r, \delta_r)$  is a quasitriangular Lie bialgebra. If further,  $(\mathfrak{Q}, \delta_r, \omega, [, ]_\omega)$  is a dual quasitriangular Lie bialgebra, then  $((\mathfrak{Q}, [, ], N)$  is a Nijenhuis Lie algebra, where the Nijenhuis operator  $N : \mathfrak{Q} \rightarrow \mathfrak{Q}$  is defined by*

$$N(x) = \omega(x, r^1)r^2, \quad \forall x \in \mathfrak{Q}. \quad (14)$$

*Proof.* Because  $(\mathfrak{Q}, \delta_r, \omega, [, ]_\omega)$  is a dual quasitriangular Lie bialgebra, where  $\delta_r$  is defined by Eq.(7), and  $(\mathfrak{Q}, [, ], \omega)$  is a symplectic Lie algebra, one has

$$\begin{aligned} 0 &= \omega(x_{(1)}, y)\omega(x_{(2)}, z) + \omega(x, z_{(1)})\omega(y, z_{(2)}) + \omega(x, y_{(1)})\omega(y_{(2)}, z) \\ &= \omega(r^1, y)\omega([x, r^2], z) + \omega([x, r^1], y)\omega(r^2, z) + \omega(x, r^1)\omega(y, [z, r^2]) \\ & \quad + \omega(x, [z, r^1])\omega(y, r^2) + \omega(x, r^1)\omega([y, r^2], z) + \omega(x, [y, r^1])\omega(r^2, z) \\ &= \omega(r^1, y)\omega([x, r^2], z) + \omega(r^1, y)\omega([r^2, z], x) + \omega(x, r^1)\omega([r^2, z], y) \\ & \quad + \omega(x, r^1)\omega([y, r^2], z) + \omega([x, r^1], y)\omega(r^2, z) + \omega([r^1, y], x)\omega(r^2, z) \\ & \quad \text{(by the skew-symmetry of } \omega, \text{ the antisymmetry of } r \text{ and Eq.(1))} \\ &= \omega(y, r^1)\omega([z, x], r^2) + \omega(x, r^1)\omega(r^2, [y, z]) + \omega([x, y], r^1)\omega(r^2, z) \\ & \quad \text{(by the skew-symmetry of } \omega, \text{ and Eq.(1)).} \end{aligned} \quad (15)$$

In the rest, we check that the linear map  $N$  defined in Eq.(14) is a Nijenhuis operator on  $(\mathfrak{Q}, [, ])$ .

$$\begin{aligned}
 & [N(x), N(y)] - N([N(x), y] + [x, N(y)] - N([x, y])) \\
 &= \omega(x, r^1)\omega(y, \bar{r}^1)[r^2, \bar{r}^2] - \omega(x, r^1)\omega([r^2, y], \bar{r}^1)\bar{r}^2 - \omega(y, r^1)\omega([x, r^2], \bar{r}^1)\bar{r}^2 \\
 &\quad + \omega([x, y], r^1)\omega(r^2, \bar{r}^1)\bar{r}^2 \\
 &\stackrel{(15)}{=} \omega(x, r^1)\omega(y, \bar{r}^1)[r^2, \bar{r}^2] - \omega(x, r^1)\omega([r^2, y], \bar{r}^1)\bar{r}^2 - \omega(y, r^1)\omega([x, r^2], \bar{r}^1)\bar{r}^2 \\
 &\quad + \omega([x, y], r^1)\omega(r^2, \bar{r}^1)\bar{r}^2 - \omega(x, r^1)\omega([y, \bar{r}^1], r^2)\bar{r}^2 - \omega(y, r^1)\omega([\bar{r}^1, x], r^2)\bar{r}^2 \\
 &\quad - \omega(r^2, \bar{r}^1)\omega([x, y], r^1)\bar{r}^2 \\
 &\stackrel{(13)}{=} \omega(x, r^1)\omega(y, \bar{r}^1)[r^2, \bar{r}^2] + \omega(x, r^1)\omega([\bar{r}^1, r^2], y)\bar{r}^2 + \omega(y, r^1)\omega([r^2, \bar{r}^1], x)\bar{r}^2 \\
 &= \omega(x, r^1)\omega(y, \bar{r}^1)[r^2, \bar{r}^2] + \omega(x, r^1)\omega(y, [r^2, \bar{r}^1])\bar{r}^2 + \omega(x, [r^1, \bar{r}^1])\omega(y, r^2)\bar{r}^2 \\
 &\quad \text{(by the anti-symmetry of } r) \\
 &\stackrel{(8)}{=} 0.
 \end{aligned}$$

Thus,  $((\mathfrak{Q}, [, ]), N)$  is a Nijenhuis Lie algebra. These finish the proof.  $\square$

Next we apply Theorem 3.11 to  $(\mathfrak{sl}_2, [, ])$ .

**Example 3.12.** Based on Examples 3.3, 3.5, 3.9, set  $r = \lambda(f \otimes g - g \otimes f)$ ,

$$\begin{cases} \delta_r(e) = 2\lambda(e \otimes f - f \otimes e), \\ \delta_r(f) = 0, \\ \delta_r(g) = 2\lambda(g \otimes f - f \otimes g) \end{cases},$$

$$\begin{array}{c|ccc} \omega & e & f & g \\ \hline e & 0 & 0 & \kappa \\ f & 0 & 0 & -2\kappa \\ g & -\kappa & 2\kappa & 0 \end{array},$$

then  $(\mathfrak{sl}_2, [, ], r, \delta_r)$  is a quasitriangular Lie bialgebra,  $(\mathfrak{sl}_2, \delta_r, \omega, [, ]_\omega)$  is a dual quasitriangular Lie bialgebra and  $(\mathfrak{sl}_2, [, ], \omega)$  is a symplectic Lie algebra. Therefore all the possible Nijenhuis operators  $N : \mathfrak{sl}_2 \longrightarrow \mathfrak{sl}_2$  induced by Theorem 3.11 on  $(\mathfrak{sl}_2, [, ])$  are given below:

$$\begin{cases} N(e) = -\lambda\kappa f, \\ N(f) = 2\lambda\kappa f, \\ N(g) = 2\lambda\kappa g, \end{cases}$$

where  $\lambda, \kappa$  are parameters.

In Theorem 2.1, we give all the Nijenhuis operators on  $(\mathfrak{sl}_2, [, ])$ . Now we discuss the relation between Nijenhuis operators in Example 3.12 and in Theorem 2.1.

**Remark 3.13.** (1) Comparison between Nijenhuis operators in Example 3.12 and Nijenhuis operators in Theorem 2.1.

| Nij operators in Example 3.12   | Nij operators in Theorem 2.1   |
|---|--|
| $\begin{cases} N(e) = -\lambda\kappa f, \\ N(f) = 2\lambda\kappa f, \\ N(g) = 2\lambda\kappa g, \end{cases} \quad (\lambda\kappa \neq 0)$ | $k_1 = k_3 = 0, k_2 = -\lambda\kappa \neq 0, k_5 = 2\lambda\kappa$ in Case (4) |
| $\begin{cases} N(e) = 0 \\ N(f) = 0 \\ N(g) = 0 \end{cases} \quad (\lambda\kappa = 0)$  | $k_1 = k_5 = k_6 = 0$ in Case (1)  |

- (2) Although all the Nijenhuis operators on  $\mathfrak{sl}_2$  obtained by the methods in Theorem 3.11 are given in Example 3.12, they are only diagonalisable. Therefore we can get that the conditions in Theorem 3.11 are sufficient and not necessary.

#### 4. (CO)REPRESENTATIONS OF NIJENHUIS LIE (CO)ALGEBRAS

In this section, we discuss the representation of Nijenhuis Lie algebras and provide a construction of Nijenhuis operators on a Lie coalgebra.

**4.1. Dual representations.** Let  $(\mathfrak{L}, [, ])$  be a Lie algebra. A **representation of**  $(\mathfrak{L}, [, ])$  is a pair  $(V, \rho)$  where  $V$  is a vector space,  $\rho : \mathfrak{L} \rightarrow \text{End}(V)$  is a linear map such that, for all  $x, y \in \mathfrak{L}$  and  $v \in V$ ,

$$\rho([x, y])(v) = \rho(x)(\rho(y)v) - \rho(y)(\rho(x)v). \quad (16)$$

**Definition 4.1.** If  $((\mathfrak{L}, [, ]), N)$  is a Nijenhuis Lie algebra, then a **representation of**  $((\mathfrak{L}, [, ]), N)$  is a triple  $((V, \rho), \alpha)$  consisting of a representation  $(V, \rho)$  of  $(\mathfrak{L}, [, ])$  and a linear map  $\alpha : V \rightarrow V$  such that, for all  $x \in \mathfrak{L}$  and  $v \in V$ ,

$$\rho(N(x))\alpha(v) + \alpha^2(\rho(x)v) = \alpha(\rho(N(x))v) + \alpha(\rho(x)\alpha(v)). \quad (17)$$

**Example 4.2.** Let  $((\mathfrak{L}, [, ]), N)$  be a Nijenhuis Lie algebra and  $x \in \mathfrak{L}$ . Define a linear map  $ad : \mathfrak{L} \rightarrow \text{End}(\mathfrak{L})$  by  $ad_x y = [x, y]$ , then  $((\mathfrak{L}, ad), N)$  is a representation of  $((\mathfrak{L}, [, ]), N)$ , called **adjoint representation of**  $((\mathfrak{L}, [, ]), N)$ .

A representation of a Nijenhuis Lie algebra can be interpreted by a semi-direct product Nijenhuis Lie algebra.

**Proposition 4.3.** Let  $((\mathfrak{L}, [, ]), N)$  be a Nijenhuis Lie algebra,  $(V, \rho)$  be a representation of  $(\mathfrak{L}, [, ])$  and  $\alpha : V \rightarrow V$  a linear map. For all  $x, y \in \mathfrak{L}$  and  $u, v \in V$ , define a bracket on  $\mathfrak{L} \oplus V$  by

$$[x + u, y + v]_{\heartsuit} := [x, y] + \rho(x)v - \rho(y)u \quad (18)$$

and a linear map  $N_{\mathfrak{L} \oplus V} : \mathfrak{L} \oplus V \rightarrow \mathfrak{L} \oplus V$  by

$$N_{\mathfrak{L} \oplus V}(x + v) := (N + \alpha)(x + v) = N(x) + \alpha(v). \quad (19)$$

Then  $\mathfrak{Q} \oplus V$  equipped with the bracket product (18) and the linear map (19) turns to be a Nijenhuis Lie algebra, which is denoted by  $(\mathfrak{Q} \ltimes^\rho V, N + \alpha)$ , if and only if  $((V, \rho), \alpha)$  is a representation of  $((\mathfrak{Q}, [, ]), N)$ .

*Proof.* The proof can be seen as a special case of Proposition 5.5, so we skip it.  $\square$

**Lemma 4.4.** Let  $((\mathfrak{Q}, [, ]), N)$  be a Nijenhuis Lie algebra,  $(V, \rho)$  be a representation of  $(\mathfrak{Q}, [, ])$  and  $\beta : V \rightarrow V$  be a linear map. Define a linear map  $\rho^* : \mathfrak{Q} \rightarrow \text{End}(V^*)$  by

$$\langle \rho^*(x)v^*, v \rangle = -\langle v^*, \rho(x)v \rangle,$$

where  $x \in \mathfrak{Q}, v^* \in V^*$  and  $v \in V$ . Then  $((V^*, \rho^*), \beta^*)$  is a representation of  $((\mathfrak{Q}, [, ]), N)$  if and only if the following condition holds:

$$\beta(\rho(N(x))v) + \rho(x)\beta^2(v) = \rho(N(x))\beta(v) + \beta(\rho(x)\beta(v)), \quad (20)$$

where  $x \in \mathfrak{Q}$  and  $v \in V$ .

Especially, for a linear map  $P : \mathfrak{Q} \rightarrow \mathfrak{Q}$ ,  $((\mathfrak{Q}^*, \text{ad}^*), P^*)$  is a representation of  $((\mathfrak{Q}, [, ]), N)$  if and only if the following condition holds:

$$P([N(x), y]) + [x, P^2(y)] = [N(x), P(y)] + P([x, P(y)]), \quad (21)$$

where  $x, y \in \mathfrak{Q}$ .

*Proof.* First we note that  $(V^*, \rho^*)$  is a representation of  $(\mathfrak{Q}, [, ])$  since  $(V, \rho)$  is a representation of  $(\mathfrak{Q}, [, ])$ . Eq.(17) holds for  $((V^*, \rho^*), \beta^*)$  if and only if, for all  $x \in \mathfrak{Q}, v \in V, v^* \in V^*$ ,

$$\begin{aligned} & \langle \rho^*(N(x))\beta^*(v^*) + \beta^{*2}(\rho^*(x)v^*) - \beta^*(\rho^*(N(x))v^*) - \beta^*(\rho^*(x)\beta^*(v^*)), v \rangle \\ & = \langle m^*, -\beta(\rho(N(x))v) - \rho(x)\beta^2(v) + \rho(N(x))\beta(v) + \beta(\rho(x)\beta(v)) \rangle. \end{aligned}$$

Thus we finish the proof.  $\square$

**Definition 4.5.** Under the assumption in Lemma 4.4.  $((\mathfrak{Q}, [, ]), N)$  is  $\beta$ -admissible with respect to  $(V, \rho)$  if Eq.(20) holds. When Eq.(21) holds, we say  $((\mathfrak{Q}, [, ]), N)$  is  $P$ -admissible.

**Remark 4.6.** (a)  $((\mathfrak{Q}, [, ]), N)$  is not  $\alpha$ -admissible with respect to  $(V, \rho)$  even if  $((V, \rho), \alpha)$  is a representation of  $((\mathfrak{Q}, [, ]), N)$ .

(b)  $((\mathfrak{Q}, [, ]), N)$  is  $N^*$ -admissible.

(c) If  $((V, \rho), \alpha)$  is a representation of  $((\mathfrak{Q}, [, ]), N)$ , then when  $\beta = k\alpha + \ell \text{id}_V$ ,  $k, \ell \in K$ , Eq.(20) turns to be

$$k^2(\rho(x)\alpha^2(v) - \alpha(\rho(x)\alpha(v))) + k(\alpha^2(\rho(x)v) - \alpha(\rho(x)\alpha(v))) + k\ell(\rho(x)\alpha(v) - \alpha(\rho(x)v)) = 0,$$

where  $x \in \mathfrak{Q}$  and  $v \in V$ . Therefore  $((\mathfrak{Q}, [, ]), N)$  is  $\ell \text{id}_V$ -admissible with respect to  $(V, \rho)$ . If further  $\alpha(\rho(x)v) = \rho(x)\alpha(v)$ , then  $((\mathfrak{Q}, [, ]), N)$  is  $\beta = k\alpha + \ell \text{id}_V$ -admissible with respect to  $(V, \rho)$ .

(d) Let  $((\mathfrak{Q}, [, ]), N)$  be a Nijenhuis Lie algebra,  $(V, \rho)$  be a representation of  $(\mathfrak{Q}, [, ])$ ,  $\widetilde{N} = -\text{id}_\mathfrak{Q} - N$ ,  $\widetilde{\beta} = -\text{id}_V - \beta$ . Then  $((\mathfrak{Q}, [, ]), \widetilde{N})$  is a Nijenhuis Lie algebra and  $((\mathfrak{Q}, [, ]), N)$  is  $\beta$ -admissible with respect to  $(V, \rho)$  if and only if  $((\mathfrak{Q}, [, ]), \widetilde{N})$  is  $\widetilde{\beta}$ -admissible with respect to  $(V, \rho)$ .

**4.2. Corepresentations of Nijenhuis Lie coalgebras.** Let us consider compatible corepresentations over compatible Lie coalgebras and then introduce the notions of a Nijenhuis Lie coalgebra and its corepresentation.

**Definition 4.7.** [15] Let  $(\mathfrak{Q}, \delta)$  be a Lie coalgebra,  $V$  be a vector space,  $\gamma : V \rightarrow \mathfrak{Q} \otimes V$  (write  $\gamma(v) = v_{(-1)} \otimes v_{(0)}$ ) be a linear map. Then  $(V, \gamma)$  is a corepresentation of  $(\mathfrak{Q}, \delta)$  if

$$v_{(-1)(1)} \otimes v_{(-1)(2)} \otimes v_{(0)} = v_{(-1)} \otimes v_{(0)(-1)} \otimes v_{(0)(0)} - v_{(0)(-1)} \otimes v_{(-1)} \otimes v_{(0)(0)}, \quad (22)$$

where  $v \in V$ .

Let  $(V, \gamma)$  and  $(V', \gamma')$  be corepresentations of  $(\mathfrak{Q}, \delta)$  and  $(\mathfrak{Q}', \delta')$ , respectively. A **homomorphism from  $(V, \gamma)$  to  $(V', \gamma')$**  is a pair  $(f, g)$ , where  $f : \mathfrak{Q} \rightarrow \mathfrak{Q}'$  and  $g : V \rightarrow V'$  are linear maps such that for all  $x \in \mathfrak{Q}$  and  $v \in V$ ,

$$\delta'(f(x)) = (f \otimes f)(\delta(x)), \quad \gamma'(g(v)) = (f \otimes g)(\gamma(v)). \quad (23)$$

**Lemma 4.8.** Let  $\mathfrak{Q}$  be a vector spaces,  $\delta, \Delta : \mathfrak{Q} \rightarrow \mathfrak{Q} \otimes \mathfrak{Q}$  (write  $\delta(x) = x_{(1)} \otimes x_{(2)}$  and  $\Delta(x) = x_{[1]} \otimes x_{[2]}$ ) be two linear maps,  $s, t$  be two parameters. Set  $\delta_{s,t} = s\delta + t\Delta$ . Then  $(\mathfrak{Q}, \delta_{s,t})$  is a Lie coalgebra if and only if  $(\mathfrak{Q}, \delta), (\mathfrak{Q}, \Delta)$  are two Lie coalgebras and, for all  $x \in \mathfrak{Q}$ , the equation below holds:

$$\begin{aligned} & x_{(1)} \otimes x_{(2)[1]} \otimes x_{(2)[2]} + x_{[1]} \otimes x_{[2](1)} \otimes x_{[2](2)} + x_{(2)[1]} \otimes x_{(2)[2]} \otimes x_{(1)} \\ & + x_{[2](1)} \otimes x_{[2](2)} \otimes x_{[1]} + x_{(2)[2]} \otimes x_{(1)} \otimes x_{(2)[1]} + x_{[2](2)} \otimes x_{[1]} \otimes x_{[2](1)} = 0. \end{aligned}$$

*Proof.* It is direct by checking Eqs.(4) and (5) for  $\delta_{s,t}$ .  $\square$

**Lemma 4.9.** Let  $(\mathfrak{Q}, \delta_{s,t})$  be a Lie coalgebra, and  $\gamma, \Gamma : V \rightarrow \mathfrak{Q} \otimes V$  (write  $\gamma(v) = v_{(-1)} \otimes v_{(0)}$  and  $\Gamma(v) = v_{[-1]} \otimes v_{[0]}$ ) be two linear maps. Set  $\gamma_{s,t} = s\gamma + t\Gamma$ . Then  $(V, \gamma_{s,t})$  is a corepresentation of  $(\mathfrak{Q}, \delta_{s,t})$  if and only if  $(V, \gamma), (V, \Gamma)$  are corepresentations of  $(\mathfrak{Q}, \delta)$  and  $(\mathfrak{Q}, \Delta)$ , respectively, and further, for all  $v \in V$ , the equation below holds:

$$\begin{aligned} & v_{(-1)[1]} \otimes v_{(-1)[2]} \otimes v_{(0)} + v_{[-1](1)} \otimes v_{[-1](2)} \otimes v_{[0]} \\ & = v_{(-1)} \otimes v_{(0)[-1]} \otimes v_{(0)[0]} + v_{[-1]} \otimes v_{[0](-1)} \otimes v_{[0](0)} \\ & - v_{(0)[-1]} \otimes v_{(-1)} \otimes v_{(0)[0]} - v_{[0](-1)} \otimes v_{[-1]} \otimes v_{[0](0)}. \end{aligned}$$

*Proof.* It is direct by checking Eq.(22) for  $\delta_{s,t}$  and  $\gamma_{s,t}$ .  $\square$

**Theorem 4.10.** Let  $(V, \gamma)$  be a corepresentation of  $(\mathfrak{Q}, \delta)$ ,  $(V, \gamma_{s,t})$  be a corepresentation of  $(\mathfrak{Q}, \delta_{s,t})$ ,  $P : \mathfrak{Q} \rightarrow \mathfrak{Q}$  and  $\beta : V \rightarrow V$  be two linear maps. Then  $(\text{sid}_{\mathfrak{Q}} + tP, \text{sid}_V + t\beta)$  is a homomorphism from  $(V, \gamma)$  of  $(\mathfrak{Q}, \delta)$  to  $(V, \gamma_{s,t})$  of  $(\mathfrak{Q}, \delta_{s,t})$  if and only if for all  $x \in \mathfrak{Q}$  and  $v \in V$ , the equations below hold:

$$P(x)_{[1]} \otimes P(x)_{[2]} = P(x_{(1)}) \otimes P(x_{(2)}), \quad (24)$$

$$P(x)_{(1)} \otimes P(x)_{(2)} + x_{[1]} \otimes x_{[2]} = x_{(1)} \otimes P(x_{(2)}) + P(x_{(1)}) \otimes x_{(2)}, \quad (25)$$

$$\beta(v)_{[-1]} \otimes \beta(v)_{[0]} = P(v_{(-1)}) \otimes \beta(v_{(0)}), \quad (26)$$

$$\beta(v)_{(-1)} \otimes \beta(v)_{(0)} + v_{[-1]} \otimes v_{[0]} = v_{(-1)} \otimes \beta(v_{(0)}) + P(v_{(-1)}) \otimes v_{(0)}. \quad (27)$$

*Proof.* It is straightforward by checking Eq.(23) for  $(\text{sid}_{\mathfrak{L}} + tP, \text{sid}_V + t\beta)$ .  $\square$

Based on Eqs.(24) and (25), we have

**Definition 4.11.** Let  $(\mathfrak{L}, \delta)$  be a Lie coalgebra and  $P : \mathfrak{L} \rightarrow \mathfrak{L}$  a linear map. A **Nijenhuis Lie coalgebra** is a triple  $((\mathfrak{L}, \delta), P)$  such that for all  $x \in \mathfrak{L}$ ,

$$P(x_{(1)}) \otimes P(x_{(2)}) + P^2(x)_{(1)} \otimes P^2(x)_{(2)} = P(P(x)_{(1)}) \otimes P(x)_{(2)} + P(x)_{(1)} \otimes P(P(x)_{(2)}). \quad (28)$$

For a linear map  $\psi : M \rightarrow V$ , we denote the transpose map by  $\psi^* : V^* \rightarrow M^*$  given by

$$\langle \psi^*(v^*), m \rangle = \langle v^*, \psi(m) \rangle, \quad \forall m \in M, v^* \in V^*.$$

**Proposition 4.12.** (a) *If  $((\mathfrak{L}, \delta), P)$  is a Nijenhuis Lie coalgebra, then  $((\mathfrak{L}^*, \delta^*), P^*)$  is a Nijenhuis Lie algebra, where  $\delta^* : \mathfrak{L}^* \otimes \mathfrak{L}^* \rightarrow \mathfrak{L}^*$  is defined by  $\langle \delta^*(x^* \otimes y^*), z \rangle = \langle x^*, z_{(1)} \rangle \langle y^*, z_{(2)} \rangle$ .*

(b) *If  $((\mathfrak{L}, [, ]), N)$  is a finite dimensional Nijenhuis Lie algebra, then  $((\mathfrak{L}^*, [, ]^*), N^*)$  is a Nijenhuis Lie coalgebra, where  $[, ]^* : \mathfrak{L}^* \rightarrow \mathfrak{L}^* \otimes \mathfrak{L}^*$  is defined by  $\langle [, ]^*(x^*), y \otimes z \rangle = \langle x^*, [y, z] \rangle$ .*

*Proof.* We only verify Item (a). Item (b) is left to the readers. Since  $(\mathfrak{L}, \delta)$  is a Lie coalgebra, then  $(\mathfrak{L}^*, \delta^*)$  is a Lie algebra ([9]). Next we only need to check that Eq.(3) holds for  $\delta^*, P^*$ . In fact, for all  $x^*, y^* \in \mathfrak{L}^*$  and  $z \in \mathfrak{L}$ , one gets

$$\begin{aligned} & \langle \delta^*(P^*(x^*) \otimes P^*(y^*)) + P^{*2}(\delta^*(x^* \otimes y^*)) - P^*(\delta^*(P^*(x^*) \otimes y^*)) - P^*(\delta^*(x^* \otimes P^*(y^*))), z \rangle \\ &= \langle x^* \otimes y^*, P(z_{(1)}) \otimes P(z_{(2)}) + P^2(z)_{(1)} \otimes P^2(z)_{(2)} - P(P(z)_{(1)}) \otimes P(z)_{(2)} - P(z)_{(1)} \otimes P(P(z)_{(2)}) \rangle. \end{aligned}$$

We finish the proof.  $\square$

According to Eqs.(26) and (27), one can obtain

**Definition 4.13.** Let  $((\mathfrak{L}, \delta), P)$  be a Nijenhuis Lie coalgebra,  $(V, \gamma)$  be a corepresentation of  $(\mathfrak{L}, \delta)$  and  $\beta : V \rightarrow V$  a linear map. A **corepresentation of  $((\mathfrak{L}, \delta), P)$**  is a triple  $((V, \gamma), \beta)$  such that for all  $v \in V$ , the equation below holds:

$$P(v_{(-1)}) \otimes \beta(v_{(0)}) + \beta^2(v)_{(-1)} \otimes \beta^2(v)_{(0)} = P(\beta(v)_{(-1)}) \otimes \beta(v)_{(0)} + \beta(v)_{(-1)} \otimes \beta(\beta(v)_{(0)}).$$

**Proposition 4.14.** *Let  $((\mathfrak{L}, \delta), P)$  be a Nijenhuis Lie coalgebra,  $V$  a vector space,  $\gamma : V \rightarrow \mathfrak{L} \otimes V$ ,  $\beta : V \rightarrow V$  two linear maps. Define a coproduct*

$$\delta_{\mathfrak{L} \oplus V}(x + v) := x_{(1)} \otimes x_{(2)} + v_{(-1)} \otimes v_{(0)} - v_{(0)} \otimes v_{(-1)}, \quad \forall x \in \mathfrak{L}, v \in V$$

*on  $\mathfrak{L} \oplus V$  and a linear map  $P_{\mathfrak{L} \oplus V}$  on  $\mathfrak{L} \oplus V$  by  $P_{\mathfrak{L} \oplus V}(x + v) := P(x) + \beta(v)$ . Then  $((\mathfrak{L} \oplus V, \delta_{\mathfrak{L} \oplus V}), P_{\mathfrak{L} \oplus V})$  is a Nijenhuis Lie coalgebra if and only if  $((V, \gamma), \beta)$  is a corepresentation of  $((\mathfrak{L}, \delta), P)$ . In this case, we call this Nijenhuis Lie coalgebra **semi-direct coproduct of  $((\mathfrak{L}, \delta), P)$  by its corepresentation  $((V, \gamma), \beta)$**  and denote it by  $(\mathfrak{L} \times_{\gamma} V, P + \beta)$ .*

*Proof.* First we note that  $(\mathcal{Q} \oplus V, \delta_{\mathcal{Q} \oplus V})$  is a Lie coalgebra if and only if  $(V, \gamma)$  is a corepresentation of  $(\mathcal{Q}, \delta)$  ([9]). For all  $x \in \mathcal{Q}$  and  $v \in V$ , Eq.(28) holds for  $\delta_{\mathcal{Q} \oplus V}$  and  $P_{\mathcal{Q} \oplus V}$  if and only if

$$\begin{aligned} & P(x_{(1)}) \otimes P(x_{(2)}) + P(v_{(-1)}) \otimes \beta(v_{(0)}) - \beta(v_{(0)}) \otimes P(v_{(-1)}) \\ & + P^2(x)_{(1)} \otimes P^2(x)_{(2)} + \beta^2(v)_{(-1)} \otimes \beta^2(v)_{(0)} - \beta^2(v)_{(0)} \otimes \beta^2(v)_{(-1)} \\ = & P(P(x)_{(1)}) \otimes P(x)_{(2)} + P(\beta(v)_{(-1)}) \otimes \beta(v_{(0)}) - \beta(\beta(v_{(0)})) \otimes \beta(v)_{(-1)} \\ & + P(x)_{(1)} \otimes P(P(x)_{(2)}) + \beta(v)_{(-1)} \otimes \beta(\beta(v_{(0)})) - \beta(v_{(0)}) \otimes P(\beta(v)_{(-1)}). \end{aligned}$$

Thus we can finish the proof.  $\square$

Proposition 4.12 can be extended to

- Proposition 4.15.** (a) *If  $((V, \gamma), \beta)$  is a corepresentation of  $((\mathcal{Q}, \delta), P)$ , then  $((V^*, \gamma^*), \beta^*)$  is a representation of  $((\mathcal{Q}^*, \delta^*), P^*)$ , where  $\gamma^* : \mathcal{Q}^* \otimes V^* \rightarrow V^*$  is defined by  $\langle \gamma^*(x^* \otimes v^*), u \rangle = \langle x^*, u_{(-1)} \rangle \langle v^*, u_{(0)} \rangle$ .*
- (b) *If  $((V, \rho), \alpha)$  is a finite dimensional representation of a finite dimensional Nijenhuis Lie algebra  $((\mathcal{Q}, [, ], N)$ , then  $((V^*, \rho^*), \alpha^*)$  is a corepresentation of  $((\mathcal{Q}^*, [, ]^*), N^*)$ , where  $\rho^* : V^* \rightarrow \mathcal{Q}^* \otimes V^*$  is defined by  $\langle \rho^*(v^*), x \otimes u \rangle = \langle v^*, \rho(x)u \rangle$ .*

**4.3. A sufficient condition for Nijenhuis Lie coalgebra.** In this subsection, we consider the construction of Nijenhuis operators on a Lie coalgebra. First we give the dual concept of a symplectic Lie algebra.

**Definition 4.16.** Let  $(\mathcal{Q}, \delta)$  be a Lie coalgebra and  $r \in \mathcal{Q} \otimes \mathcal{Q}$  be an antisymmetric element. Assume that

$$r^1_1 \otimes r^1_2 \otimes r^2 + r^2 \otimes r^1_1 \otimes r^1_2 + r^1_2 \otimes r^2 \otimes r^1_1 = 0.$$

Then  $(\mathcal{Q}, \delta)$  is called a cosymplectic Lie coalgebra and denoted by  $(\mathcal{Q}, \delta, r)$ .

**Proposition 4.17.** *Let  $(\mathcal{Q}, \delta)$  be a Lie coalgebra and  $r \in \mathcal{Q} \otimes \mathcal{Q}$  be an antisymmetric element. If  $(\mathcal{Q}, [, ], r, \delta_r)$  is a quasitriangular Lie bialgebra, where  $\delta_r$  is defined by Eq.(7), then  $(\mathcal{Q}, \delta_r, r)$  is a cosymplectic Lie coalgebra.*

*Proof.* It is direct by Eqs.(7) and (8).  $\square$

**Theorem 4.18.** *Let  $(\mathcal{Q}, \delta, r)$  be a cosymplectic Lie coalgebra and  $\omega \in (\mathcal{Q} \otimes \mathcal{Q})^*$ . If  $(\mathcal{Q}, \delta, \omega, [, ]_\omega)$  is a dual quasitriangular Lie bialgebra with the multiplication  $[, ]_\omega$  defined in Eq.(10) and further,  $(\mathcal{Q}, [, ]_\omega, r, \delta_r)$  is a quasitriangular Lie bialgebra together with the comultiplication  $\delta_r$  defined in Eq.(7), then  $((\mathcal{Q}, \delta), P)$  is a Nijenhuis Lie coalgebra, where  $P$  is defined by*

$$P(x) = r^1 \omega(r^2, x), \quad \forall x \in \mathcal{Q}.$$

*Proof.* Similar to Theorem 3.11.  $\square$

## 5. NIJENHUIS LIE BIALGEBRAS

In this section, we establish the bialgebraic theory of Nijenhuis Lie algebras and introduce the notion of classical  $P$ -Nijenhuis Yang-Baxter equations. That is to say, we consider the following commutative diagram.

$$\begin{array}{ccc}
 \text{Lie bialgebras} & \xrightarrow{+\text{Nijenhuis operator}} & \text{Nijenhuis Lie bialgebras} \\
 \uparrow \text{Bialgebraization} & & \uparrow \text{Bialgebraization} \\
 \text{Lie algebras} & \xrightarrow{+\text{Nijenhuis operator}} & \text{Nijenhuis Lie algebras.}
 \end{array}$$

## 5.1. Nijenhuis Lie bialgebras and equivalent characterizations.

**Definition 5.1.** A Nijenhuis Lie bialgebra is a quintuple  $((\mathfrak{Q}, [, ]), \delta, N, P)$ , where

- (a)  $((\mathfrak{Q}, [, ]), N)$  is Nijenhuis Lie algebra;
- (b)  $((\mathfrak{Q}, \delta), P)$  is Nijenhuis Lie coalgebra;
- (c)  $(\mathfrak{Q}, [, ], \delta)$  is Lie bialgebra;
- (d) For all  $x, y \in \mathfrak{Q}$ , the equations below hold:

$$P([N(x), y]) + [x, P^2(y)] = [N(x), P(y)] + P([x, P(y)]), \quad (29)$$

$$P(N(x)_{(1)}) \otimes N(x)_{(2)} + x_{(1)} \otimes N^2(x)_{(2)} = P(x_{(1)}) \otimes N(x)_{(2)} + N(x)_{(1)} \otimes N(N(x)_{(2)}). \quad (30)$$

**Remark 5.2.** (a) Eq.(29) means that  $((\mathfrak{Q}, [, ]), N)$  is  $P$ -admissible;  
 (b) Eq.(30) means that  $((\mathfrak{Q}^*, \delta^*), P^*)$  is  $N^*$ -admissible.

**Example 5.3.** Let  $\mathfrak{Q} = \mathfrak{sl}_2$  be a Lie algebra given above. Set

$$\begin{cases} \delta(e) = 2\lambda(e \otimes f - f \otimes e), \\ \delta(f) = 0, \\ \delta(g) = 2\lambda(g \otimes f - f \otimes g) \end{cases}.$$

Then by Example 3.3,  $(\mathfrak{Q}, \delta)$  is a Lie coalgebra and further  $(\mathfrak{Q}, [, ], \delta)$  is a Lie bialgebra. Define the linear maps  $N, P : \mathfrak{Q} \rightarrow \mathfrak{Q}$  as follows ( $k_1, k_2, k_3$  and  $\ell$  are parameters).

$$\begin{aligned}
 \text{(a)} & \begin{cases} N(e) = k_1 e + k_2 f \\ N(f) = k_1 f \\ N(g) = k_1 g \end{cases}, \quad P = k_1 \text{id}; \\
 \text{(b)} & \begin{cases} N(e) = k_1 e + k_2 f \\ N(f) = k_1 f \\ N(g) = k_1 g \end{cases}, \quad \begin{cases} P(e) = k_1 e + \ell f \\ P(f) = k_1 f \\ P(g) = k_1 g \end{cases}; \\
 \text{(c)} & \begin{cases} N(e) = k_1 e + k_2 f + k_3 g \\ N(f) = k_1 f \\ N(g) = k_1 g \end{cases}, \quad \begin{cases} P(e) = k_1 e - k_2 f \\ P(f) = k_1 f \\ P(g) = k_1 g \end{cases}; \\
 \text{(d)} & \begin{cases} N(e) = k_1 e + k_2 f + k_3 g \\ N(f) = k_1 f \\ N(g) = k_1 g \end{cases}, \quad \begin{cases} P(e) = k_1 e - k_2 f - k_3 g \\ P(f) = k_1 f \\ P(g) = k_1 g \end{cases} \quad (k_2 \neq 0, k_3 \neq 0).
 \end{aligned}$$

Then by a direct computation we can find that  $((\mathfrak{Q}, [, ]), \delta, N, P)$  is a Nijenhuis Lie bialgebra, where the  $(N, P)$  is given in Item (a)-Item (d), respectively.

Nijenhuis Lie bialgebra can be characterized by Manin triple of Nijenhuis Lie algebras and matched pair of Lie algebras.

**Definition 5.4.** A **matched pair of Nijenhuis Lie algebras**  $((\mathfrak{Q}, [, ]_{\mathfrak{Q}}), N_{\mathfrak{Q}})$  and  $((\mathfrak{h}, [, ]_{\mathfrak{h}}), N_{\mathfrak{h}})$  is an octuple  $(((\mathfrak{Q}, [, ]_{\mathfrak{Q}}), N_{\mathfrak{Q}}), ((\mathfrak{h}, [, ]_{\mathfrak{h}}), N_{\mathfrak{h}}), \rho_{\mathfrak{Q}}, \rho_{\mathfrak{h}})$ , where  $((\mathfrak{h}, \rho_{\mathfrak{Q}}), N_{\mathfrak{h}})$  is a representation of  $((\mathfrak{Q}, [, ]_{\mathfrak{Q}}), N_{\mathfrak{Q}})$ ,  $((\mathfrak{Q}, \rho_{\mathfrak{h}}), N_{\mathfrak{Q}})$  is a representation of  $((\mathfrak{h}, [, ]_{\mathfrak{h}}), N_{\mathfrak{h}})$ , and  $((\mathfrak{Q}, [, ]_{\mathfrak{Q}}), (\mathfrak{h}, [, ]_{\mathfrak{h}}), \rho_{\mathfrak{Q}}, \rho_{\mathfrak{h}})$  is a matched pair of Lie algebras  $(\mathfrak{Q}, [, ]_{\mathfrak{Q}})$  and  $(\mathfrak{h}, [, ]_{\mathfrak{h}})$ , that is to say, the equations below hold:

$$\rho_{\mathfrak{h}}(a)[x, y]_{\mathfrak{Q}} - [\rho_{\mathfrak{h}}(a)x, y]_{\mathfrak{Q}} - [x, \rho_{\mathfrak{h}}(a)y]_{\mathfrak{Q}} = \rho_{\mathfrak{h}}(\rho_{\mathfrak{Q}}(y)a)x - \rho_{\mathfrak{h}}(\rho_{\mathfrak{Q}}(x)a)y, \quad (31)$$

$$\rho_{\mathfrak{Q}}(x)[a, b]_{\mathfrak{h}} - [\rho_{\mathfrak{Q}}(x)a, b]_{\mathfrak{h}} - [a, \rho_{\mathfrak{Q}}(x)b]_{\mathfrak{h}} = \rho_{\mathfrak{Q}}(\rho_{\mathfrak{h}}(b)x)a - \rho_{\mathfrak{Q}}(\rho_{\mathfrak{h}}(a)x)b, \quad (32)$$

where  $x, y \in \mathfrak{Q}$  and  $a, b \in \mathfrak{h}$ .

**Proposition 5.5.** Let  $((\mathfrak{Q}, [, ]_{\mathfrak{Q}}), N_{\mathfrak{Q}})$  and  $((\mathfrak{h}, [, ]_{\mathfrak{h}}), N_{\mathfrak{h}})$  be two Nijenhuis Lie algebras. Assume that  $((\mathfrak{h}, \rho_{\mathfrak{Q}}), N_{\mathfrak{h}})$  is a representation of  $((\mathfrak{Q}, [, ]_{\mathfrak{Q}}), N_{\mathfrak{Q}})$  and  $((\mathfrak{Q}, \rho_{\mathfrak{h}}), N_{\mathfrak{Q}})$  is a representation of  $((\mathfrak{h}, [, ]_{\mathfrak{h}}), N_{\mathfrak{h}})$ . Define a bracket  $[, ]_{\star}$  on  $\mathfrak{Q} \oplus \mathfrak{h}$  by

$$[x + a, y + b]_{\star} = [x, y]_{\mathfrak{Q}} + \rho_{\mathfrak{h}}(a)y - \rho_{\mathfrak{h}}(b)x + [a, b]_{\mathfrak{h}} + \rho_{\mathfrak{Q}}(x)b - \rho_{\mathfrak{Q}}(y)a \quad (33)$$

and a linear map  $N_{\star}$  on  $\mathfrak{Q} \oplus \mathfrak{h}$  by

$$N_{\star}(x + a) = N_{\mathfrak{Q}}(x) + N_{\mathfrak{h}}(a), \quad (34)$$

where  $x, y \in \mathfrak{Q}$  and  $a, b \in \mathfrak{h}$ . Then  $((\mathfrak{Q} \oplus \mathfrak{h}, [, ]_{\star}), N_{\star})$  is a Nijenhuis Lie algebra if and only if Eqs.(31) and (32) hold. In this case, we denote  $((\mathfrak{Q} \oplus \mathfrak{h}, [, ]_{\star}), N_{\star})$  by  $(\mathfrak{Q} \bowtie \mathfrak{h}, N_{\mathfrak{Q}} + N_{\mathfrak{h}})$ .

*Proof.* We note that  $(\mathfrak{Q} \oplus \mathfrak{h}, [, ]_{\star})$  is a Lie algebra if and only if Eqs.(31) and (32) hold ([9]). So we only need to check that  $N_{\star}$  satisfies Eq.(3). In fact, For all  $x, y \in \mathfrak{Q}$  and  $a, b \in \mathfrak{h}$ , we have

$$\begin{aligned} & [N_{\mathfrak{Q}}(x), N_{\mathfrak{Q}}(y)]_{\mathfrak{Q}} + \rho_{\mathfrak{h}}(N_{\mathfrak{h}}(a))N_{\mathfrak{Q}}(y) - \rho_{\mathfrak{h}}(N_{\mathfrak{h}}(b))N_{\mathfrak{Q}}(x) + [N_{\mathfrak{h}}(a), N_{\mathfrak{h}}(b)]_{\mathfrak{h}} + \rho_{\mathfrak{Q}}(N_{\mathfrak{Q}}(x))N_{\mathfrak{h}}(b) \\ & - \rho_{\mathfrak{Q}}(N_{\mathfrak{Q}}(y))N_{\mathfrak{h}}(a) + N_{\mathfrak{Q}}^2([x, y]_{\mathfrak{Q}}) + N_{\mathfrak{Q}}^2(\rho_{\mathfrak{h}}(a)y) - N_{\mathfrak{Q}}^2(\rho_{\mathfrak{h}}(b)x) + N_{\mathfrak{h}}^2([a, b]_{\mathfrak{h}}) + N_{\mathfrak{h}}^2(\rho_{\mathfrak{Q}}(x)b) \\ & - N_{\mathfrak{h}}^2(\rho_{\mathfrak{Q}}(y)a) \\ & \stackrel{(3)(17)}{=} N_{\mathfrak{Q}}([N_{\mathfrak{Q}}(x), y]_{\mathfrak{Q}}) + N_{\mathfrak{Q}}(\rho_{\mathfrak{h}}(N_{\mathfrak{h}}(a))y) - N_{\mathfrak{Q}}(\rho_{\mathfrak{h}}(b)N_{\mathfrak{Q}}(x)) + N_{\mathfrak{h}}([N_{\mathfrak{h}}(a), b]_{\mathfrak{h}}) + N_{\mathfrak{h}}(\rho_{\mathfrak{Q}}(N_{\mathfrak{Q}}(x))b) \\ & - N_{\mathfrak{h}}(\rho_{\mathfrak{Q}}(y)N_{\mathfrak{h}}(a)) + N_{\mathfrak{Q}}([x, N_{\mathfrak{Q}}(y)]_{\mathfrak{Q}}) + N_{\mathfrak{Q}}(\rho_{\mathfrak{h}}(a)N_{\mathfrak{Q}}(y)) - N_{\mathfrak{Q}}(\rho_{\mathfrak{h}}(N_{\mathfrak{h}}(b))x) + N_{\mathfrak{h}}([a, N_{\mathfrak{h}}(b)]_{\mathfrak{h}}) \\ & + N_{\mathfrak{h}}(\rho_{\mathfrak{Q}}(x)N_{\mathfrak{h}}(b)) - N_{\mathfrak{h}}(\rho_{\mathfrak{Q}}(N_{\mathfrak{Q}}(y))a). \end{aligned}$$

Then we finish the proof.  $\square$

**Remark 5.6.** If  $(((\mathfrak{Q}, [, ]_{\mathfrak{Q}}), N_{\mathfrak{Q}}), ((\mathfrak{h}, [, ]_{\mathfrak{h}}), N_{\mathfrak{h}}), \rho_{\mathfrak{Q}}, \rho_{\mathfrak{h}})$  is a matched pair of Nijenhuis Lie algebras  $((\mathfrak{Q}, [, ]_{\mathfrak{Q}}), N_{\mathfrak{Q}})$  and  $((\mathfrak{h}, [, ]_{\mathfrak{h}}), N_{\mathfrak{h}})$ , then one can obtain a Nijenhuis Lie algebra  $((\mathfrak{Q} \oplus \mathfrak{h}, [, ]_{\star}), N_{\star})$ .

**Theorem 5.7.** Let  $((\mathfrak{Q}, [, ]), N)$  and  $((\mathfrak{Q}^*, [, ]_{\circ} := \delta^*), P^*)$  be two Nijenhuis Lie algebras. Then the quintuple  $((\mathfrak{Q}, [, ]), \delta, N, P)$  is a Nijenhuis Lie bialgebra if and only if  $(((\mathfrak{Q}, [, ]), N), ((\mathfrak{Q}^*, [, ]_{\circ}), P^*), ad_{\mathfrak{Q}^*}^*, ad_{\mathfrak{Q}}^*)$  is a matched pair of Nijenhuis Lie algebras  $((\mathfrak{Q}, [, ]), N)$  and  $((\mathfrak{Q}^*, [, ]_{\circ}), P^*)$ .

*Proof.* For Lie algebras, we have ([2,9]):  $((\mathfrak{Q}, [, ]), \delta)$  is a Lie bialgebra if and only if  $((\mathfrak{Q}, [, ]), (\mathfrak{Q}^*, [, ]_{\circ}), ad_{\mathfrak{Q}^*}^*, ad_{\mathfrak{Q}}^*)$  is a matched pair of Lie algebras  $(\mathfrak{Q}, [, ])$  and  $(\mathfrak{Q}^*, [, ]_{\circ})$ , where Eq.(31)=Eq.(32)  $\Leftrightarrow$  Eq.(6). The rest can be obtained by Remark 5.2.  $\square$

Let  $(\mathfrak{Q}, [, ])$  be a Lie algebra,  $\mathfrak{B} : \mathfrak{Q} \otimes \mathfrak{Q} \rightarrow K$  be a nondegenerate bilinear form. If further  $\mathfrak{B}$  is invariant, i.e., for all  $x, y, z \in \mathfrak{Q}$ ,  $\mathfrak{B}([x, y], z) = (x, [y, z])$ . In this case, the triple  $((\mathfrak{Q}, [, ]), \mathfrak{B})$  is called a **Frobenius Lie algebra**. A Frobenius algebra  $((\mathfrak{Q}, [, ]), \mathfrak{B})$  is **symmetric** if  $\mathfrak{B}$  is symmetric.

**Definition 5.8.** A **Nijenhuis Frobenius Lie algebra** is a quadruple  $((\mathcal{Q}, [\cdot, \cdot]), N, \mathfrak{B})$  consisting of a Nijenhuis Lie algebra  $((\mathcal{Q}, [\cdot, \cdot]), N)$  and a Frobenius Lie algebra  $((\mathcal{Q}, [\cdot, \cdot]), \mathfrak{B})$ .

Let  $\hat{N} : \mathcal{Q} \rightarrow \mathcal{Q}$  denote the adjoint operator of  $N$  by the following way:

$$\mathfrak{B}(N(x), y) = \mathfrak{B}(x, \hat{N}(y)), \quad \forall x, y \in \mathcal{Q}.$$

**Proposition 5.9.** Let  $((\mathcal{Q}, [\cdot, \cdot]), N, \mathfrak{B})$  be a symmetric Nijenhuis Frobenius Lie algebra. Then  $((\mathcal{Q}, [\cdot, \cdot]), N)$  is  $\hat{N}$ -admissible.

*Proof.* It is natural that  $(\mathcal{Q}^*, ad^*)$  is a representation of  $(\mathcal{Q}, [\cdot, \cdot])$ . In what follows, we only need to prove that Eq.(21) holds for  $ad$  and  $\hat{N}$ . In fact, for all  $x, y, z \in \mathcal{Q}$ ,

$$\begin{aligned} 0 &\stackrel{(3)}{=} \mathfrak{B}([N(x), N(y)], z) + \mathfrak{B}(N^2([x, y]), z) - \mathfrak{B}(N([x, N(y)]), z) - \mathfrak{B}(N([N(x), y]), z) \\ &= \mathfrak{B}(N(x), [N(y), z]) + \mathfrak{B}([x, y], \hat{N}^2(z)) - \mathfrak{B}([x, N(y)], \hat{N}(z)) - \mathfrak{B}([N(x), y], \hat{N}(z)) \\ &= \mathfrak{B}(x, \hat{N}([N(y), z])) + \mathfrak{B}(x, [y, \hat{N}^2(z)]) - \mathfrak{B}(x, [N(y), \hat{N}(z)]) - \mathfrak{B}(x, \hat{N}([y, \hat{N}(z)])), \end{aligned}$$

so the proof is finished.  $\square$

**Definition 5.10.** Let  $((\mathcal{Q}, [\cdot, \cdot]), N)$  and  $((\mathcal{Q}^*, [\cdot, \cdot]_\circ), P^*)$  be two Nijenhuis Lie algebras. Assume that there is a Lie algebra structure  $[\cdot, \cdot]_\star$  on  $\mathcal{Q} \oplus \mathcal{Q}^*$  which contains both  $(\mathcal{Q}, [\cdot, \cdot])$  and  $(\mathcal{Q}^*, [\cdot, \cdot]_\circ)$  as Lie subalgebras. Define a bilinear form on  $\mathcal{Q} \oplus \mathcal{Q}^*$  by

$$\mathfrak{B}_d(x + a^*, y + b^*) = \langle a^*, y \rangle + \langle b^*, x \rangle, \quad \forall x, y \in \mathcal{Q}, a^*, b^* \in \mathcal{Q}^*. \quad (35)$$

If  $\mathfrak{B}_d$  is invariant, then  $((\mathcal{Q} \oplus \mathcal{Q}^*, [\cdot, \cdot]_\star), \mathfrak{B}_d)$  is a symmetric Frobenius Lie algebra. We call this Frobenius Lie algebra a **Manin triple of a Frobenius Lie algebra associated to  $(\mathcal{Q}, [\cdot, \cdot])$  and  $(\mathcal{Q}^*, [\cdot, \cdot]_\circ)$**  and denote it by  $(\mathcal{Q} \bowtie \mathcal{Q}^*, \mathfrak{B}_d)$ . If further  $((\mathcal{Q} \bowtie \mathcal{Q}^*, N + P^*), \mathfrak{B}_d)$  is a Nijenhuis Frobenius Lie algebra, then we call  $((\mathcal{Q} \bowtie \mathcal{Q}^*, N + P^*), \mathfrak{B}_d)$  a **Manin triple of a Nijenhuis Frobenius Lie algebra associated to  $(\mathcal{Q}, [\cdot, \cdot]), N$  and  $(\mathcal{Q}^*, [\cdot, \cdot]_\circ), P^*$** .

**Lemma 5.11.** Let  $((\mathcal{Q} \bowtie \mathcal{Q}^*, N + P^*), \mathfrak{B}_d)$  be a Manin triple of a Nijenhuis Frobenius Lie algebra associated to  $((\mathcal{Q}, [\cdot, \cdot]), N)$  and  $((\mathcal{Q}^*, [\cdot, \cdot]_\circ), P^*)$ . Then

- (a) The adjoint  $\widehat{N + P^*}$  of  $N + P^*$  with respect to  $\mathfrak{B}_d$  is  $P + N^*$ . Further  $(\mathcal{Q} \bowtie \mathcal{Q}^*, N + P^*)$  is  $(P + N^*)$ -admissible.
- (b)  $((\mathcal{Q}, [\cdot, \cdot]), N)$  is  $P$ -admissible.
- (c)  $((\mathcal{Q}^*, [\cdot, \cdot]_\circ), P^*)$  is  $N^*$ -admissible.

*Proof.* (a) For all  $x, y \in \mathcal{Q}$  and  $a^*, b^* \in \mathcal{Q}^*$ , we have

$$\begin{aligned} \mathfrak{B}((N + P^*)(x + a^*), y + b^*) &= \langle b^*, N(x) \rangle + \langle P^*(a^*), y \rangle \\ &= \langle N^*(b^*), x \rangle + \langle a^*, P(y) \rangle = \mathfrak{B}(x + a^*, (P + N^*)(y + b^*)). \end{aligned}$$

Then by Proposition 5.9, the proof of Item (a) is finished.

(b) Since  $(\mathcal{Q} \bowtie \mathcal{Q}^*, N + P^*)$  is  $(P + N^*)$ -admissible, by Eq.(21), for  $x, y \in \mathcal{Q}, a^*, b^* \in \mathcal{Q}^*$ , one has

$$\begin{aligned} &(P + N^*)[(N(x) + P^*(a^*)), (y + b^*)] + [(x + a^*), (P^2(y) + N^{*2}(b^*))] \\ &= [(N(x) + P^*(a^*)), (P(y) + N^*(b^*))] + (P + N^*)[(x + a^*), (P(y) + N^*(b^*))]. \end{aligned} \quad (36)$$

Then Item (b) can be obtained by letting  $a^* = b^* = 0$  in Eq. (36).

(c) Similarly, Item (c) can be proved by setting  $x = y = 0$  in Eq. (36).  $\square$

**Theorem 5.12.** Let  $((\mathcal{Q}, [\cdot, \cdot]), N)$  and  $((\mathcal{Q}^*, [\cdot, \cdot]_\circ), P^*)$  be two Nijenhuis Lie algebras. Then there is a Manin triple  $((\mathcal{Q} \bowtie \mathcal{Q}^*, N + P^*), \mathfrak{B}_d)$  of a Nijenhuis Frobenius Lie algebra associated to  $((\mathcal{Q}, [\cdot, \cdot]), N)$  and  $((\mathcal{Q}^*, [\cdot, \cdot]_\circ), P^*)$  if and only if  $((\mathcal{Q}, [\cdot, \cdot]), N), ((\mathcal{Q}^*, [\cdot, \cdot]_\circ), P^*), ad_{\mathcal{Q}^*}^*, ad_{\mathcal{Q}}^*$  is a matched pair of Nijenhuis Lie algebras  $((\mathcal{Q}, [\cdot, \cdot]), N)$  and  $((\mathcal{Q}^*, [\cdot, \cdot]_\circ), P^*)$ .

*Proof.* For Lie algebras, we have ([2, 9]): there is a Manin triple  $(\mathcal{Q} \bowtie \mathcal{Q}^*, \mathfrak{B}_d)$  of a Frobenius Lie algebra associated to  $(\mathcal{Q}, [, ])_{\circ}$  and  $(\mathcal{Q}^*, [, ]_{\circ})$  if and only if  $((\mathcal{Q}, [, ]), (\mathcal{Q}^*, [, ]_{\circ}), ad_{\mathcal{Q}^*}, ad_{\mathcal{Q}^*})$  is a matched pair of Lie algebras  $(\mathcal{Q}, [, ])$  and  $(\mathcal{Q}^*, [, ]_{\circ})$ . The rest can be obtained by Proposition 5.5 and Lemma 5.11.  $\square$

**5.2. Enrichment of classical Yang-Baxter equations.** Let  $((\mathcal{Q}, [, ]), N)$  be a  $P$ -admissible Nijenhuis Lie algebra,  $r$  be an antisymmetric solution of the cYBe in  $(\mathcal{Q}, [, ])$ . Then by Remark 3.1, we can get a quasitriangular Lie bialgebra  $(\mathcal{Q}, [, ], r, \delta_r)$ . Thus in this case, by Remark 5.2,  $((\mathcal{Q}, [, ]), \delta_r, N, P)$  is a Nijenhuis Lie bialgebra if and only if Eqs.(28) and (30) hold. In what follows, we continue to study these two conditions by using the element  $r$ .

**Theorem 5.13.** *Let  $((\mathcal{Q}, [, ]), N)$  be a  $P$ -admissible Nijenhuis Lie algebra and  $\delta := \delta_r$  given by Eq.(7). Then*

(a) *Eq.(28) holds if and only if, for all  $x \in \mathcal{Q}$ ,*

$$\begin{aligned} & (\text{id} \otimes P \circ ad_x - \text{id} \otimes ad_{P(x)})(P \otimes \text{id} - \text{id} \otimes N)(r) \\ & + (ad_{P(x)} \otimes \text{id} - P \circ ad_x \otimes \text{id})(N \otimes \text{id} - \text{id} \otimes P)(r) = 0. \end{aligned} \quad (37)$$

(b) *Eq.(30) holds if and only if, for all  $x \in \mathcal{Q}$ ,*

$$\begin{aligned} & (\text{id} \otimes ad_{N(x)} - \text{id} \otimes N \circ ad_x + ad_{N(x)} \otimes \text{id} + P \circ ad_x \otimes \text{id})(P \otimes \text{id} - \text{id} \otimes N)(r) \\ & = (ad_x \circ P^2 \otimes \text{id} - ad_x \otimes N^2)(r). \end{aligned} \quad (38)$$

*Proof.* (a) For all  $x \in \mathcal{Q}$ , we calculate

$$\begin{aligned} P(x_{(1)}) \otimes P(x_{(2)}) & \stackrel{(7)}{=} P(r^1) \otimes P([x, r^2]) + P([x, r^1]) \otimes P(r^2), \\ P^2(x)_{(1)} \otimes P^2(x)_{(2)} & \stackrel{(7)}{=} r^1 \otimes [P^2(x), r^2] + [P^2(x), r^1] \otimes r^2, \\ P(P(x)_{(1)}) \otimes P(x)_{(2)} & \stackrel{(7)}{=} P(r^1) \otimes [P(x), r^2] + P([P(x), r^1]) \otimes r^2 \\ & = P(r^1) \otimes [P(x), r^2] - P([r^1, P(x)]) \otimes r^2 \\ & \stackrel{(21)}{=} P(r^1) \otimes [P(x), r^2] + [N(r^1), P(x)] \otimes r^2 - P([N(r^1), x]) \otimes r^2 - [r^1, P^2(x)] \otimes r^2, \\ P(x)_{(1)} \otimes P(P(x)_{(2)}) & \stackrel{(7)}{=} r^1 \otimes P([P(x), r^2]) + [P(x), r^1] \otimes P(r^2) \\ & = -r^1 \otimes P([r^2, P(x)]) + [P(x), r^1] \otimes P(r^2) \\ & \stackrel{(21)}{=} r^1 \otimes [N(r^2), P(x)] - r^1 \otimes P([N(r^2), x]) - r^1 \otimes [r^2, P^2(x)] + [P(x), r^1] \otimes P(r^2), \end{aligned}$$

then Eq.(28)  $\Leftrightarrow$  Eq.(37).

(b) For all  $x \in \mathcal{Q}$ , by Eq.(7), one can obtain

$$\begin{aligned} P(N(x)_{(1)}) \otimes N(x)_{(2)} & = P(r^1) \otimes [N(x), r^2] + P([N(x), r^1]) \otimes r^2 \\ & \stackrel{(21)}{=} P(r^1) \otimes [N(x), r^2] + [N(x), P(r^1)] \otimes r^2 + P([x, P(r^1)]) \otimes r^2 - [x, P^2(r^1)] \otimes r^2, \\ x_{(1)} \otimes N^2(x)_{(2)} & = r^1 \otimes N^2([x, r^2]) + [x, r^1] \otimes N^2(r^2), \\ P(x)_{(1)} \otimes N(x)_{(2)} & = P(r^1) \otimes N([x, r^2]) + P([x, r^1]) \otimes N(r^2), \\ N(x)_{(1)} \otimes N(N(x)_{(2)}) & = r^1 \otimes N([N(x), r^2]) + [N(x), r^1] \otimes N(r^2) \\ & \stackrel{(3)}{=} r^1 \otimes [N(x), N(r^2)] + r^1 \otimes N^2([x, r^2]) - r^1 \otimes N([x, N(r^2)]) + [N(x), r^1] \otimes N(r^2), \end{aligned}$$

then Eq.(30)  $\Leftrightarrow$  Eq.(38).  $\square$

**Lemma 5.14.** (a) *If  $(P \otimes \text{id} - \text{id} \otimes N)(r) = 0$ , then  $(ad_x \circ P^2 \otimes \text{id} - ad_x \otimes N^2)(r) = 0$ .*

(b) If  $r$  is antisymmetric and  $(P \otimes \text{id} - \text{id} \otimes N)(r) = 0$ , then  $(N \otimes \text{id} - \text{id} \otimes P)(r) = 0$ .

*Proof.* Straightforward.  $\square$

By Theorem 5.13 and Lemma 5.14, one gets

**Theorem 5.15.** Let  $((\mathfrak{Q}, [, ]), N)$  be a  $P$ -admissible Nijenhuis Lie algebra and  $\delta := \delta_r$  given by Eq.(7). If  $r$  is an antisymmetric solution of the cYBe in  $(\mathfrak{Q}, [, ])$  and the equation below holds

$$(P \otimes \text{id} - \text{id} \otimes N)(r) = 0, \quad (39)$$

then  $((\mathfrak{Q}, [, ]), \delta_r, N, P)$  is a Nijenhuis Lie bialgebra. In this case, we call  $((\mathfrak{Q}, [, ]), \delta_r, N, P)$  **quasitriangular**.

**Example 5.16.** In fact, the Nijenhuis Lie bialgebras given in Example 5.3 are all quasitriangular, with  $r = \lambda(f \otimes g - g \otimes f)$ .

**Definition 5.17.** Let  $((\mathfrak{Q}, [, ]), N)$  be a  $P$ -admissible Nijenhuis Lie algebra. Then the cYBe (i.e. Eq.(8)) together with Eq.(39) is call a **classical  $P$ -Nijenhuis Yang-Baxter equation** (cPNYBe for short).

Let  $V$  be a vector space and  $r \in V \otimes V$ , define  $\Phi_r : V^* \rightarrow V$  by

$$\Phi_r(v^*) = \langle v^*, r^1 \rangle r^2, \quad \forall v^* \in V^*. \quad (40)$$

The element  $r$  is **nondegenerate** if  $\Phi_r$  is bijective.

**Theorem 5.18.** Let  $((\mathfrak{Q}, [, ]), N)$  be a Nijenhuis algebra,  $r \in \mathfrak{Q} \otimes \mathfrak{Q}$  antisymmetric and  $P : \mathfrak{Q} \rightarrow \mathfrak{Q}$  a linear map. Then  $r$  is a solution of the cPNYBe if and only if for all  $a^*, b^* \in \mathfrak{Q}^*$ ,  $\Phi_r$  satisfies

$$[\Phi_r(a^*), \Phi_r(b^*)] = \Phi_r(ad^*(\Phi_r(a^*))(b^*) - ad^*(\Phi_r(b^*))(a^*)) \quad (41)$$

and

$$N(\Phi_r(a^*)) = \Phi_r(P^*(a^*)). \quad (42)$$

*Proof.* By [2, 9], we know that  $r$  is a solution of the cYBe if and only if Eq.(41) holds. And for all  $a^* \in \mathfrak{Q}^*$ , one gets

$$\begin{aligned} \Phi_r(P^*(a^*)) &= \langle P^*(a^*), r^1 \rangle r^2 = \langle a^*, P(r^1) \rangle r^2, \\ P(\Phi_r(a^*)) &= \langle a^*, r^1 \rangle P(r^2). \end{aligned}$$

So Eq.(42)  $\Leftrightarrow$  Eq.(39). We finish the proof.  $\square$

**Definition 5.19.** Let  $((\mathfrak{Q}, [, ]), N)$  be a Nijenhuis Lie algebra,  $(V, \rho)$  a representation of  $(\mathfrak{Q}, [, ])$  and  $\alpha : V \rightarrow V$  a linear map. A linear map  $T : V \rightarrow \mathfrak{Q}$  is called a **weak relative Rota-Baxter operator (rRBO for short) associated to  $(V, \rho)$  and  $\alpha$**  if for all  $u, v \in V$ ,  $T$  satisfies

$$[T(u), T(v)] = T(\rho(T(u))(v) - \rho(T(v))(u)), \quad (43)$$

and

$$N(T(u)) = T(\alpha(u)). \quad (44)$$

Furthermore, if  $((V, \rho), \alpha)$  is a representation of  $((\mathfrak{Q}, [, ]), N)$ , then  $T$  is called a **rRBO associated to  $((V, \rho), \alpha)$** .

By Theorem 5.18, one can obtain

**Corollary 5.20.** *Let  $((\mathcal{Q}, [\cdot, \cdot]), N)$  be a Nijenhuis algebra,  $r \in \mathcal{Q} \otimes \mathcal{Q}$  antisymmetric and  $P : \mathcal{Q} \rightarrow \mathcal{Q}$  a linear map. Then  $r$  is a solution of the cPNYBe if and only if  $\Phi_r$  is a weak rRBO associated to  $(\mathcal{Q}^*, ad^*)$  and  $P^*$ . If furthermore,  $((\mathcal{Q}, [\cdot, \cdot]), N)$  is  $P$ -admissible, then  $r$  is a solution of the cPNYBe if and only if  $\Phi_r$  is a rRBO associated to  $(\mathcal{Q}^*, ad^*), P^*$ .*

Quasitriangular Nijenhuis Lie bialgebras can be derived from semi-product Lie algebra by using rRBOs.

**Lemma 5.21.** *Let  $((\mathcal{Q}, [\cdot, \cdot]), N)$  be a Nijenhuis Lie algebra,  $(V, \rho)$  a representation of  $(\mathcal{Q}, [\cdot, \cdot]), P : \mathcal{Q} \rightarrow \mathcal{Q}$  and  $\alpha, \beta : V \rightarrow V$  linear maps. Then TFAE:*

- (a)  $(\mathcal{Q} \ltimes_{\rho} V, N + \alpha)$  is a  $(P + \beta)$ -admissible Nijenhuis Lie algebra.
- (b)  $(\mathcal{Q} \ltimes_{\rho^*} V^*, N + \beta^*)$  is a  $(P + \alpha^*)$ -admissible Nijenhuis Lie algebra.
- (c) *The following are satisfied:*
  - (1)  $((V, \rho), \alpha)$  is a representation of  $((\mathcal{Q}, [\cdot, \cdot]), N)$ ;
  - (2)  $((\mathcal{Q}, [\cdot, \cdot]), N)$  is  $P$ -admissible;
  - (3)  $((\mathcal{Q}, [\cdot, \cdot]), N)$  is  $\beta$ -admissible associated to  $(V, \rho)$ ;
  - (4) For  $x \in \mathcal{Q}, v \in V$ , we have

$$\beta(\rho(x)(\alpha(v))) + \rho(P^2(x))(v) = \rho(P(x))(\alpha(v)) + \beta(\rho(P(x))(v)). \quad (45)$$

*Proof.* ((a)  $\Leftrightarrow$  (c)) By Proposition 4.3,  $(\mathcal{Q} \ltimes_{\rho} V, N + \alpha)$  is a Nijenhuis Lie algebra if and only if  $((V, \rho), \alpha)$  is a representation of  $((\mathcal{Q}, [\cdot, \cdot]), N)$ . Eq.(21) holds for  $[\cdot, \cdot]_{\circ}, N + \alpha, P + \beta$  if and only if, for  $x, y \in \mathcal{Q}, u, v \in V$ ,

$$\begin{aligned} & P([N(x), y]) + \beta(\rho(N(x))(v)) - \beta(\rho(y)(\alpha(u))) + [x, P^2(y)] + \rho(x)\beta^2(v) - \rho(P^2(y))u \\ &= [N(x), P(y)] + \rho(N(x))(\beta(v)) - \rho(P(y))(\alpha(u)) + P([x, P(y)]) + \beta(\rho(x)(\beta(v))) - \beta(\rho(P(y))(u)) \end{aligned}$$

if and only if  $((\mathcal{Q}, [\cdot, \cdot]), N)$  is  $P$ -admissible,  $((\mathcal{Q}, [\cdot, \cdot]), N)$  is  $\beta$ -admissible associated to  $(V, \rho)$  and Eq.(45) holds.

((b)  $\Leftrightarrow$  (c)) Based on (a)  $\Leftrightarrow$  (c), (b) is equivalent to

- (i)  $((V^*, \rho^*), \beta^*)$  is a representation of  $((\mathcal{Q}, [\cdot, \cdot]), N^*)$ , i.e.,  $((\mathcal{Q}, [\cdot, \cdot]), N)$  is  $\beta$ -admissible associated to  $(V, \rho)$ ;
- (ii)  $((\mathcal{Q}, [\cdot, \cdot]), N)$  is  $P$ -admissible;
- (iii)  $((\mathcal{Q}, [\cdot, \cdot]), N)$  is  $\alpha^*$ -admissible associated to  $(V^*, \rho^*)$ , i.e.,  $((V, \rho), \alpha)$  is a representation of  $((\mathcal{Q}, [\cdot, \cdot]), N)$ ;
- (iv) For  $x \in \mathcal{Q}, a^* \in V^*$ , we have

$$\alpha^*(\rho(x)(\beta^*(a^*))) + \rho(P^2(x))(a^*) = \rho(P(x))(\beta^*(a^*)) + \alpha^*(\rho(P(x))(a^*)),$$

i.e., Eq.(45) holds.

Thus the proof of Lemma 5.21 is finished.  $\square$

The following result provides a method to construct Nijenhuis Lie bialgebras on semi-direct product Nijenhuis Lie algebras.

**Theorem 5.22.** *Let  $((\mathcal{Q}, [\cdot, \cdot]), N)$  be a  $\beta$ -admissible Nijenhuis Lie algebra associated to  $(V, \rho)$ ,  $P : \mathcal{Q} \rightarrow \mathcal{Q}$ ,  $\alpha : V \rightarrow V$  be linear maps. Assume that  $T : V \rightarrow \mathcal{Q}$  is a linear map, which is identified as an element  $\sum_{i=1}^n T(e_i) \otimes e^i \in (\mathcal{Q} \ltimes_{\rho^*} V^*) \otimes (\mathcal{Q} \ltimes_{\rho^*} V^*)$ , where  $\{e_i\}_{i=1}^n$  is a basis of  $V$  and  $\{e^i\}_{i=1}^n$  is the dual basis.*

- (a) *The element  $r = \sum_{i=1}^n (T(e_i) \otimes e^i - e^i \otimes T(e_i))$  is an antisymmetric solution of the classical  $(P + \alpha^*)$ -Nijenhuis Yang-Baxter equation in the Nijenhuis Lie algebra  $(\mathcal{Q} \ltimes_{\rho^*} V^*, N + \beta^*)$  if and only if  $T$  is a weak rRBO associated to  $(V, \rho)$  and  $\alpha$ , and  $T \circ \beta = P \circ T$ .*

- (b) Assume that  $((V, \rho), \alpha)$  is a representation of  $((\mathfrak{L}, [\cdot, \cdot]), N)$ . If  $T$  is a  $r$ RBO associated to  $((V, \rho), \alpha)$  and  $T \circ \beta = P \circ T$ , then  $r = \sum_{i=1}^n (T(e_i) \otimes e^i - e^i \otimes T(e_i))$  is an antisymmetric solution of the classical  $(P + \alpha^*)$ -Nijenhuis Yang-Baxter equation in the Nijenhuis Lie algebra  $(\mathfrak{L} \ltimes_{\rho^*} V^*, N + \beta^*)$ . If further,  $((\mathfrak{L}, [\cdot, \cdot]), N)$  is  $P$ -admissible and Eq.(45) holds, then  $(\mathfrak{L} \ltimes_{\rho^*} V^*, N + \beta^*)$  is  $(P + \alpha^*)$ -admissible. In this case, there is a Nijenhuis Lie bialgebra  $((\mathfrak{L} \ltimes_{\rho^*} V^*, N + \beta^*), \delta_r, P + \alpha^*)$ , where  $\delta_r$  is given by Eq.(7) with  $r = \sum_{i=1}^n (T(e_i) \otimes e^i - e^i \otimes T(e_i))$ .

*Proof.* (a) By [1],  $r = \sum_{i=1}^n (T(e_i) \otimes e^i - e^i \otimes T(e_i))$  is an antisymmetric solution of the Eq.(8) in the Lie algebra  $\mathfrak{L} \ltimes_{\rho^*} V^*$  if and only if  $T$  satisfies Eq.(43) for  $(V, \rho)$ . It remains to prove that Eq.(39) for  $P + \alpha^*$  and  $N + \beta^* \Leftrightarrow T \circ \beta = P \circ T$  and Eq.(44) hold. In fact,

$$\begin{aligned} ((P + \alpha^*) \otimes \text{id} - \text{id} \otimes (N + \beta^*))(r) &= \sum_{i=1}^n (P(T(e_i)) \otimes e^i - \alpha^*(e^i) \otimes T(e_i) - T(e_i) \otimes \beta^*(e^i) + e^i \otimes N(T(e_i))) \\ &= \sum_{i=1}^n (P(T(e_i)) \otimes e^i - e^i \otimes T(\alpha(e_i)) - T(\beta(e_i)) \otimes e^i + e^i \otimes N(T(e_i))). \end{aligned}$$

Therefore we can finish the proof of (a).

(b) can be checked by Item (a) and Lemma 5.21.  $\square$

#### ACKNOWLEDGMENT

This work is supported by National Natural Science Foundation of China (No. 12471033) and Natural Science Foundation of Henan Province (No. 242300421389).

## Declarations

**Competing interests** The authors claim that there is no conflict of interests.

#### APPENDIX A. PROOF OF THEOREM 2.1

*Proof.* Let  $k_i, i = 1, 2, \dots, 9$  be parameters. Define the linear map  $N : \mathfrak{sl}_2 \rightarrow \mathfrak{sl}_2$  by

$$\begin{cases} N(e) = k_1 e + k_2 f + k_3 g \\ N(f) = k_4 e + k_5 f + k_6 g \\ N(g) = k_7 e + k_8 f + k_9 g \end{cases}.$$

Then  $N$  is Nijenhuis operator on  $(\mathfrak{sl}_2, [\cdot, \cdot])$  if and only if

$$\begin{cases} k_7 k_8 + 4k_2 k_4 = 0 \\ k_7(k_5 - k_9) - k_4(k_8 + 4k_3) = 0 \\ k_2 k_7 + 4k_2 k_6 - k_8(k_1 - k_9) = 0 \\ 4k_2(k_5 - k_9) + k_8(4k_3 + k_8) = 0 \\ k_7(k_7 + 4k_6) - 4k_4(k_1 - k_9) = 0 \\ (k_1 - k_9)(k_5 - k_9) - k_2 k_4 + k_3 k_7 + k_6(k_8 + 4k_3) = 0 \end{cases}.$$

(We note here that other three missing equations can be derived by these six equations.)

$$(I) \quad k_7 = 0, \text{ then } \begin{cases} k_2 k_4 = 0 \\ k_4(k_8 + 4k_3) = 0 \\ 4k_2 k_6 - k_8(k_1 - k_9) = 0 \\ k_4(k_1 - k_9) = 0 \\ 4k_2(k_5 - k_9) + k_8(4k_3 + k_8) = 0 \\ (k_1 - k_9)(k_5 - k_9) - k_2 k_4 + k_6(k_8 + 4k_3) = 0 \end{cases}.$$

$$(IA) \quad k_4 = 0, \text{ then } \begin{cases} 4k_2k_6 - k_8(k_1 - k_9) = 0 \\ 4k_2(k_5 - k_9) + k_8(4k_3 + k_8) = 0 \\ (k_1 - k_9)(k_5 - k_9) + k_6(4k_3 + k_8) = 0 \end{cases} .$$

$$(IA1) \quad k_2 = 0, \text{ then } \begin{cases} k_8(k_1 - k_9) = 0 \\ k_8(4k_3 + k_8) = 0 \\ (k_1 - k_9)(k_5 - k_9) + k_6(k_8 + 4k_3) = 0 \end{cases} .$$

$$(IA1a) \quad k_8 = 0, \text{ then } (k_1 - k_9)(k_5 - k_9) + 4k_3k_6 = 0.$$

$$(IA1ai) \quad k_3 = 0, \text{ we have } (k_1 - k_9)(k_5 - k_9) = 0.$$

$$(IA1ai\alpha) \quad k_1 = k_9. \text{ This case is contained in Item (IB2b).}$$

$$(IA1ai\beta) \quad k_5 = k_9, \text{ i.e.,}$$

$$\begin{cases} N(e) = k_1e \\ N(f) = k_5f + k_6g \\ N(g) = k_5g \end{cases} .$$

$$(IA1aii) \quad k_3 \neq 0, \text{ we have } k_6 = \frac{(k_1 - k_9)(k_5 - k_9)}{4k_3}. \text{ Then}$$

$$\begin{cases} N(e) = k_1e + k_3g \\ N(f) = k_5f - \frac{(k_1 - k_9)(k_5 - k_9)}{4k_3}g \\ N(g) = k_9g \end{cases} \quad (k_3 \neq 0).$$

$$(IA1b) \quad k_1 - k_9 = 0, \text{ we have } \begin{cases} k_8(4k_3 + k_8) = 0 \\ k_6(4k_3 + k_8) = 0 \end{cases} .$$

$$(IA1bi) \quad \begin{cases} k_8 = 0 \\ k_3k_6 = 0 \end{cases} .$$

$$(IA1bi\alpha) \quad k_3 = 0. \text{ This case is included in Item (IB2b).}$$

$$(IA1bi\beta) \quad k_6 = 0. \text{ This case is included in Item (IB2b).}$$

$$(IA1bii) \quad k_8 = -4k_3. \text{ This case is contained in Item (IB2b).}$$

$$(IA2) \quad k_2 \neq 0, \text{ we have } \begin{cases} k_6 = \frac{k_8(k_1 - k_9)}{4k_2} \\ 4k_2(k_5 - k_9) + k_8(4k_3 + k_8) = 0 \\ (k_1 - k_9)(k_5 - k_9) + \frac{k_8(k_1 - k_9)}{4k_2}(4k_3 + k_8) = 0 \end{cases} .$$

$$(IA2a) \quad k_8 = 0, \text{ then } k_6 = 0, k_5 = k_9 \text{ since } k_2 \neq 0. \text{ That is to say,}$$

$$\begin{cases} N(e) = k_1e + k_2f + k_3g \\ N(f) = k_5f \\ N(g) = k_5g \end{cases} \quad (k_2 \neq 0).$$

$$(IA2b) \quad k_8 \neq 0, \text{ we have } \begin{cases} k_6 = \frac{k_8(k_1 - k_9)}{4k_2} \\ k_3 = \frac{k_2(k_9 - k_5)}{k_8} - \frac{k_8}{4} \end{cases} . \text{ That is to say,}$$

$$\begin{cases} N(e) = k_1e + k_2f + \left(\frac{k_2(k_9 - k_5)}{k_8} - \frac{k_8}{4}\right)g \\ N(f) = k_5f + \frac{k_8(k_1 - k_9)}{4k_2}g \\ N(g) = k_8f + k_9g \end{cases} \quad (k_2, k_8 \neq 0).$$

$$(IB) \quad k_2 = 0, \text{ we have } \begin{cases} k_4(k_8 + 4k_3) = 0 \\ k_8(k_1 - k_9) = 0 \\ k_8(k_8 + 4k_3) = 0 \\ k_4(k_1 - k_9) = 0 \\ (k_1 - k_9)(k_5 - k_9) + k_6(k_8 + 4k_3) = 0 \end{cases} .$$

$$(IB1) \quad k_4 = 0, \text{ then } \begin{cases} k_8(k_1 - k_9) = 0 \\ k_8(k_8 + 4k_3) = 0 \\ (k_1 - k_9)(k_5 - k_9) + k_6(k_8 + 4k_3) = 0 \end{cases} .$$

(IB1a)  $k_8 = 0, (k_1 - k_9)(k_5 - k_9) + 4k_3k_6 = 0$ . This case is same to Item (IA1a).

$$(IB1b) \quad k_9 - k_1 = 0, \text{ then } \begin{cases} k_8(4k_3 + k_8) = 0 \\ k_6(4k_3 + k_8) = 0 \end{cases} . \text{ This case is same to Item (IA1b).}$$

$$(IB2) \quad k_8 = -4k_3, \text{ then } \begin{cases} k_3(k_1 - k_9) = 0 \\ k_4(k_1 - k_9) = 0 \\ (k_1 - k_9)(k_5 - k_9) = 0 \end{cases} .$$

$$(IB2a) \quad k_3 = 0, \text{ then } \begin{cases} k_4(k_1 - k_9) = 0 \\ (k_1 - k_9)(k_5 - k_9) = 0 \end{cases} .$$

(IB2ai)  $k_4 = 0, (k_1 - k_9)(k_5 - k_9) = 0$ . This case is same to Item (IA1ai).

(IB2aai)  $k_1 - k_9 = 0$ . This case is included in Item (IB2b).

(IB2b)  $k_1 - k_9 = 0$ , i.e.,

$$\begin{cases} N(e) = k_1e + k_3g \\ N(f) = k_4e + k_5f + k_6g \\ N(g) = -4k_3f + k_1g \end{cases} .$$

$$(II) \quad k_7 \neq 0, \text{ then } \begin{cases} k_8 = -\frac{4k_2k_4}{k_7} \\ k_7^2(k_5 - k_9) + 4k_4(k_2k_4 - k_3k_7) = 0 \\ k_7(k_7 + 4k_6) - 4k_4(k_1 - k_9) = 0 \\ (k_1k_5 - k_2k_4 + k_3k_7 - k_1k_9 - k_5k_9 + 4k_3k_6 + k_9^2)k_7 - 4k_2k_4k_6 = 0 \end{cases} .$$

$$(IIA) \quad k_5 - k_9 = 0, \text{ then } \begin{cases} k_8 = -\frac{4k_2k_4}{k_7} \\ k_4(k_2k_4 - k_3k_7) = 0 \\ k_7(k_7 + 4k_6) - 4k_4(k_1 - k_9) = 0 \\ (-k_2k_4 + k_3k_7 + 4k_3k_6)k_7 - 4k_2k_4k_6 = 0 \end{cases} .$$

$$(IIA1) \quad k_4 = 0, \text{ then } \begin{cases} k_8 = 0 \\ k_7 + 4k_6 = 0 \end{cases} . \text{ That is to say,}$$

$$\begin{cases} N(e) = k_1e + k_2f + k_3g \\ N(f) = k_5f + k_6g \\ N(g) = -4k_6e + k_5g \end{cases} \quad (k_6 \neq 0).$$

$$(IIA2) \quad k_2k_4 - k_3k_7 = 0, \text{ then } \begin{cases} k_8 = -\frac{4k_2k_4}{k_7} \\ k_3 = \frac{k_2k_4}{k_7} \\ k_6 = \frac{k_4(k_9 - k_1)}{k_7} - \frac{k_7}{4} \end{cases} . \text{ That is to say,}$$

$$\begin{cases} N(e) = k_1e + k_2f + \frac{k_2k_4}{k_7}g \\ N(f) = k_4e + k_5f + \left(\frac{k_4(k_5 - k_1)}{k_7} - \frac{k_7}{4}\right)g \\ N(g) = k_7e - \frac{4k_2k_4}{k_7}f + k_5g \end{cases} \quad (k_7 \neq 0).$$

$$(IIB) \quad k_5 - k_9 \neq 0, \text{ then } \begin{cases} k_8 = -\frac{4k_2k_4}{k_7} \\ k_5 = k_9 + \frac{4k_3k_4}{k_7} - \frac{4k_2k_4^2}{k_7^2} \\ k_6 = \frac{k_4(k_9 - k_1)}{k_7} - \frac{k_7}{4} \end{cases} . \text{ That is to say,}$$

$$\begin{cases} N(e) = k_1e + k_2f + k_3g \\ N(f) = k_4e + (k_9 + \frac{4k_3k_4}{k_7} - \frac{4k_2k_4^2}{k_7^2})f + (\frac{k_4(k_9-k_1)}{k_7} - \frac{k_7}{4})g \\ N(g) = k_7e - \frac{4k_2k_4}{k_7}f + k_9g \end{cases} \quad (k_4, k_7 \neq 0, k_2k_4 \neq k_3k_7).$$

□

## REFERENCES

- [1] C. M. Bai, A unified algebraic approach to the classical Yang-Baxter equation, *J. Phys. A: Math. Gen.* **40** (2007), 11073-11082. [2](#), [19](#)
- [2] C. M. Bai, L. Guo, G. L. Liu, T. S. Ma, Rota-Baxter Lie bialgebras, classical Yang-Baxter equations and special L-dendriform bialgebras. *Algebr. Represent. Theory* **27**(2024), 1347-1372. [1](#), [14](#), [16](#), [17](#)
- [3] I. Bajo, S. Benayadi, A. Medina, Symplectic structures on quadratic Lie algebras. *J. Algebra* **316**(2007), 174-188. [5](#)
- [4] A. Das, Poisson-Nijenhuis groupoids. *Rep. Math. Phys.* **84** (2019), 303-331. [2](#)
- [5] A. Das, Twisted Rota-Baxter families and NS-family algebras. *J. Algebra* **612** (2022), 577-615. [2](#)
- [6] V. G. Drinfeld, Hamiltonian structures on Lie groups, Lie bialgebras, and geometric meaning of the classical Yang-Baxter equations. *Dokl. Akad. Nauk SSSR* **268**(1983), 285-287; translation in *Soviet Math. Dokl.* **27** (1983), 222-225. [2](#), [4](#)
- [7] Y. Kosmann-Schwarzbach, F. Magri, Poisson-Nijenhuis structures. *Ann. Inst. Henri Poincaré* **53** (1990), 35-81. [1](#), [2](#)
- [8] L. Liu, A. Makhlof, C. Menini, F. Panaite, BiHom-NS-algebras, twisted Rota-Baxter operators and generalized Nijenhuis operators. *Results Math.* **78**(2023):251. [2](#)
- [9] S. Majid, Foundations of quantum group theory. *Cambridge University Press*, Cambridge, 1995. [4](#), [5](#), [11](#), [12](#), [14](#), [16](#), [17](#)
- [10] O. Mokhov, Compatible flat metrics. *J. Appl. Math.* **2** (2002), 337-370. [2](#)
- [11] A. Newlander, L. Nirenberg, Complex analytic coordinates in almost complex manifolds. *Ann. Math.*(2) **65**(1957), 391-404. [2](#)
- [12] A. Nijenhuis, R. W. Richardson, Cohomology and deformations in graded Lie algebras. *Bull. Amer. Math. Soc.* **72**(1)(1966), 1-29. [1](#), [2](#)
- [13] D. E. Radford, Hopf Algebras, *KE Series on Knots and Everything*, **Vol. 49**, World Scientific, New Jersey, 2012. [4](#)
- [14] M. Semonov-Tian-Shansky, What is the classical R-matrix? *Funct. Anal. Appl.* **17**(1983), 259-272. [2](#)
- [15] L. Y. Zhang, Lie comodules and the constructions of Lie bialgebras. *Sci. China Ser. A* **51**(2008), 1017-1026. [10](#)

SCHOOL OF MATHEMATICS AND STATISTICS, HENAN NORMAL UNIVERSITY, XINXIANG 453007, CHINA  
*Email address:* lihaiying@htu.edu.cn

SCHOOL OF MATHEMATICS AND STATISTICS, HENAN NORMAL UNIVERSITY, XINXIANG 453007, CHINA  
*Email address:* matianshui@htu.edu.cn