

Higher derivative holography and temperature dependence of QGP viscosities

Thomas Apostolidis^a, Umut Gürsoy^b, Edwan Préau^b

^a*Université Paris Cité, Astroparticule et Cosmologie, F-75006 Paris, France*

^b*Institute for Theoretical Physics and Center for Extreme Matter and Emergent Phenomena, Utrecht University, 3584 CC Utrecht, The Netherlands*

E-mail: apostolidis@apc.in2p3.fr, u.gursoy@uu.nl, e.c.m.preau@uu.nl

ABSTRACT: Recent Bayesian analyses of heavy ion collision data have established a non-trivial temperature dependence of the shear and bulk viscosity per entropy. Motivated by this, we consider higher derivative corrections to realistic, bottom-up holographic models of quark-gluon plasma based on five-dimensional Einstein-dilaton theories and determine the dilaton potentials in the higher derivative terms by matching the Bayesian analyses. A byproduct of our analysis is the bulk viscosity that follows from the holographic V-QCD theory. Higher derivative corrections when treated perturbatively lead to tension with existing data. We investigate possible resolutions.

Contents

1	Introduction	1
2	Bayesian analysis of Heavy Ion Collisions	4
3	Holographic shear and bulk viscosities	5
3.1	Generalities	5
3.2	Holographic setup	8
4	Fitting holographic models to Bayesian analyses	10
4.1	Shear viscosity	10
4.2	Bulk viscosity	13
4.2.1	Bulk viscosity of V-QCD theory	13
4.2.2	Bulk viscosity with curvature corrections	14
4.3	Global fits to both shear and bulk viscosity data	16
5	Discussion and Outlook	18
A	The vacuum solution	20
A.1	Equations of motion	20
A.2	UV Asymptotics	21
A.3	IR Asymptotics	22
A.3.1	Special case of $\gamma = -\frac{4}{3}$	23
B	Calculation of shear viscosity	24
C	Invariance of η/s under modification of curvature terms	27
D	Calculation of the bulk viscosity	29
E	Numerical computation of the bulk viscosity	32

1 Introduction

Heavy-ion collisions at RHIC and LHC provide a unique window into the transport properties of QCD; see, e.g. [1] for a recent review. The transport properties of the quark-gluon plasma produced in these experiments can be characterized by the transport coefficients that enter the hydrodynamic description of the fluid, most prominent of which are the shear and the bulk viscosities that are usually defined per entropy, i.e. η/s and ζ/s respectively¹. Knowledge of these coefficients requires computing stress tensor retarded Green's

¹Another coefficient at first-order in the hydrodynamic derivative expansion, conductivity, is relevant at finite quark density or in the presence of electromagnetic fields, which are ignored here.

functions at finite temperature. The only available first-principles method, lattice QCD, is not quite suitable for this task: it is based on the Euclidean formulation, and the analytic continuation to the Lorentzian signature contains large systematic errors, as this requires knowledge of the entire particle spectrum of QCD; see [2, 3].

Recently, a more phenomenological approach, based on Bayesian analysis of heavy-ion data, became available [4–6], see section 2. Here, one models the evolution of the quark-gluon plasma by hydrodynamics, e.g. 14-moment Müller-Israel-Stewart model [7], supplemented by initial conditions for the energy-momentum distribution of nucleons, e.g. provided by the Trento model [8] and a standard freezeout procedure [9] to obtain the hadron distributions that are subsequently matched onto experimental data. This provides a probability distribution for the transport coefficients that characterize the hydrodynamic model. This approach also contains significant systematic errors resulting mainly from the modeling of the initial conditions and hydrodynamic evolution and the choice of the data sets used for matching. This can be clearly seen by comparing Bayesian analyses of Jetscape [6] versus Trajectum [4, 5]; see Figures 1 and 2. A crucial aspect that resulted from these analyses is the non-trivial temperature dependence of η/s and ζ/s .

Yet another complementary tool is provided by the holographic correspondence [10–12]. A triumph of this approach was the calculation of η/s [13] at strong coupling that resulted in the universal value $1/(4\pi)$ that agrees well with the aforementioned Bayesian analyses [4–6]. Five-dimensional “bottom-up” holographic models of QCD-like large- N gauge theories have been developed to include the salient features of the theory such as confinement, gapped hadron spectrum, chiral symmetry breaking, QCD anomalies, etc. The type of models we will consider in this work are the large N improved holographic QCD (ihQCD) model [14, 15] and its extension to include fundamental matter [16] in the Veneziano limit [17], with large N and large number of flavors with their ratio fixed. These theories naturally incorporate the running of the ’t Hooft coupling through a non-trivial profile of the dilaton Φ in the holographic dimension. The latter is prompted by introduction of a dilaton potential $V(\Phi)$, whose large Φ asymptotics (corresponding to the IR of the QFT, and the deep interior of the bulk geometry) are judiciously chosen such that the theory is confining. The extension of ihQCD and V-QCD to finite temperature was studied in [18–20] and [21, 22] respectively, and incorporation of electromagnetic fields — relevant for off-central heavy-ion collisions and neutron star mergers — was considered in [23, 24].

All these holographic theories are built on two-derivative Einstein gravity, which universally fixes the value of $\eta/s = 1/(4\pi)$ [25]. In particular, the aforementioned non-trivial temperature dependence observed in experiment requires inclusion of $1/N$ or higher derivative corrections, realized, respectively, as loop and α' contributions in string theory. On the other hand, the temperature dependence of ζ/s is already present at the two-derivative level and was investigated in the holographic models of QCD-like theories in [26, 27]. See [26–37] for a non-exhaustive list of other literature on bulk viscosity in the context of holography. Hence, one needs to incorporate the higher derivative corrections to Einstein-dilaton type holographic models to capture the nontrivial temperature profile of both η/s and ζ/s , as first observed in [38].

Such corrections have been considered early on in the context of violations of the conjectured KSS bound $\eta/s \geq 1/(4\pi)$ [39, 40]; see e.g. [41–43] and [44] for a review. However, these investigations were concerned with holographic duals of conformal field theories which are not suitable for QCD. On the other hand, literature on stringy corrections to holographic theories of QCD-like non-conformal and confining theories is rather limited². Stringy corrections to ihQCD are discussed in [14, 46, 47] but not in the context of transport. Higher curvature corrections to shear transport in duals of Einstein-dilaton theories were studied in detail in [38], but no reliable temperature profile was available from heavy-ion data back then for neither η/s nor ζ/s , so these corrections were not sufficiently constrained. Recently, there has been a revival of interest in determining the transport coefficients in holographic theories, including higher curvature corrections. While [34, 35] introduced a novel method to compute all transport coefficients by varying the background solutions, which in principle can be extended directly to higher curvature theories [48], Buchel, Cremonini and Early [36] extended the standard techniques to beyond the supergravity approximation in generic theories, which is precisely the problem we want to address in this paper.

The question we start with is whether we can construct realistic (in the aforementioned sense) holographic theories for the quark-gluon plasma beyond the supergravity approximation, which postdict the non-trivial temperature profile of η/s and ζ/s observed in the Bayesian analyses [4–6]. This is a sensible approach to characterizing transport in strongly interacting quark-gluon matter, as it attempts to merge the two existing approaches, Bayesian analysis of heavy-ion data and holography, in the absence of a first-principle derivation. Specifically, we consider a 5D Einstein-dilaton theory with a dilaton potential $V(\Phi)$ and with fourth order derivative corrections. Among the three types of corrections, i.e. R^2 , $R_{\mu\nu}^2$ and $R_{\mu\nu\alpha\beta}^2$, one can show, by means of a field redefinition of the metric $g_{\mu\nu} \rightarrow g_{\mu\nu} + a g_{\mu\nu} R + b R_{\mu\nu}$ with a and b constants, that only the last one contributes to the shear-viscosity per entropy η/s when the coefficients of the curvature corrections are constant³. This is no longer true for the bulk-viscosity per entropy which receives contributions from all three terms. However, motivated by this, we will simplify our discussion and include only the $R_{\mu\nu\alpha\beta}^2$ correction, albeit with a dilaton-dependent coefficient $G(\Phi)$ to have sufficient freedom to match the temperature profiles in η/s and ζ/s .

Our main results are:

1. We find that a potential $G(\Phi)$ can be constructed to fit the Trajectum result for η/s [4, 5], using the analytic holographic expression for the shear viscosity per entropy derived in [38]. However, this implies that the corresponding corrections on the background would have to be large, thus invalidating our perturbative framework for the curvature corrections. This implies that the finite-coupling corrections are indispensable for transport in the quark-gluon plasma.
2. We investigated the influence of a (very mild) increase in temperature in η/s on the bulk viscosity. Our results indicate that ζ/s tends to decrease at low temperature,

²See [45] for the first such example.

³The argument, which was first formulated in [41], is reviewed in appendix C.

hence approaching the Trajectum result [4, 5]), whereas it becomes larger at high temperature, hence approaching JetScape [6]). This suggests that other types of correction are necessary to fit both η/s and ζ/s with reasonable accuracy.

3. Before even adding the curvature corrections we calculate the zeroth order result for ζ/s in the V-QCD model for the first time. Our findings hint towards a non-negligible break of conformality in the quark-gluon plasma.

We organized the rest of the paper as follows. In the next section, we summarize the relevant results from existing Bayesian analysis. In Section 3 we outline the holographic calculation of shear and bulk viscosity and the holographic models that we employ to fit the Bayesian data. The calculations in this section are detailed in Appendices A, B and C. Section 4 is devoted to fitting the holographic models to results of Bayesian analyses. We discuss our results and provide an outlook to future work in section 5. Several appendices A-E provide details of our calculations but also contain complementary information. In Appendix A, we analyze the background equations including the higher derivative corrections and work out the near boundary and deep interior limits of the background. In particular we obtain the asymptotic behavior of the G function such that it does not spoil the salient features, e.g. confinement, of our holographic QCD models. This is important also to justify our omission of curvature corrections to the background. For completeness, in Appendices B, D and E we recapitulate the calculations of shear and bulk viscosities including higher derivative corrections. In Appendix C we further justify our omission of different type of 4-derivative corrections by showing that they do not contribute to η/s .

2 Bayesian analysis of Heavy Ion Collisions

In complex systems such as the quark-gluon plasma, the correlation between microscopic parameters of the theory, e.g. the shear and bulk viscosity per entropy, and observables, e.g. the distribution of soft hadrons observed at the detector, is complicated. As a result, inferring the values of the microscopic parameters from observed data, i.e. the “inverse problem,” becomes extremely difficult. A systematic means to address this problem is provided by the Bayes’ theorem

$$\mathcal{P}(B|A) = \frac{\mathcal{P}(A|B)\mathcal{P}(B)}{\mathcal{P}(A)}, \quad (2.1)$$

where $\mathcal{P}(B|A)$ denotes the conditional probability that B occurs given condition A . For example, if A is a collection of observables in collisions and B is a set of microscopic variables, we can construct $\mathcal{P}(A|B)$ by running many simulations of the collisions based on the hydrodynamic model with parameters B . Bayes’ theorem then provides a probability distribution for the microscopic parameters given the actual observed data.

The Bayesian method is flawed by an inherent ambiguity, originating from the choice of parametrization for the microscopic parameters and the selection of observables. It would be surprising if quantitatively similar results are obtained for different choices of

parametrizing the models. Indeed, the two existing analyses [6] and [4, 5] differ both in the parametrization of temperature dependence for the viscosities and in selected observables. In particular, in [4], η/s is parametrized as

$$\frac{\eta}{s} = a + b(T - T_c) \left(\frac{T}{T_c} \right)^c, \quad (2.2)$$

where $T_c = 154 \times 10^{-3} \text{ GeV}$ is the deconfinement cross-over temperature and the coefficients a , b and c are to be determined by the Bayesian analysis. On the other hand, in [6] the parametrization was of the form

$$\frac{\eta}{s} = a_{\text{low}}(T - T_c)\Theta(T_c - T) + \eta_0 + a_{\text{high}}(T_c - T)\Theta(T - T_c), \quad (2.3)$$

with the parameters a_{low} , a_{high} and the kink η_0 to be determined. Bulk viscosity per entropy is parametrized as a skewed Cauchy distribution in both works

$$\frac{\zeta}{s} = d \left[\frac{e^2}{(T - T_\zeta)^2 + e^2} \right]. \quad (2.4)$$

While the constants d , e and T_ζ are left as parameters to be determined in [4, 5], [6] further parametrizes $e = w_\zeta(1 + \lambda_\zeta \text{sgn}(T - T_\zeta))$. As mentioned above, not only the parametrization of transport coefficients, but also the choice of observables differ in the two works. Whereas Jetscape uses p_T -integrated flow observables and includes only lead-lead collisions, Trajectum works with p_T -differential flow observables and includes both lead-lead and proton-lead collisions.

We present the most probable values of the parameters obtained in Trajectum analysis here for future reference:

$$a = 0.065_{-0.040}^{+0.038}, \quad b = 0.9_{-0.90}^{+0.81} \text{ GeV}^{-1}, \quad c = -0.04_{-0.85}^{+0.75}. \quad (2.5)$$

$$d = 0.0060_{-0.0060}^{+0.0058}, \quad e = 0.10_{-0.08}^{+0.15} \text{ GeV}, \quad T_\zeta = 0.202_{-0.039}^{+0.047} \text{ GeV}. \quad (2.6)$$

The resulting profiles for η/s along with a 90% confidence band are shown in Figure ??.

We also reproduce in Figure 2 the bulk viscosity per entropy obtained in the two studies. The Trajectum value of ζ/s is seen to be an order of magnitude smaller than that of Jetscape and consistent with zero bulk viscosity, which suggests a nearly conformal behavior. Both results have a large uncertainty, and they are in statistical tension only at low temperatures.

3 Holographic shear and bulk viscosities

3.1 Generalities

Kubo's linear response theory relates the shear and bulk viscosities to the stress tensor correlators. Shear deformation corresponds to a traceless deformation of the metric which




Figure 1: Posterior distribution for the shear viscosity to entropy ratio versus temperature for Trajectum (left) and Jetscape (right). For Trajectum the 90% confidence band is shown in blue and the dark blue curve is the mean. The gray area signifies the 90% confidence band for the prior distribution. For Jetscape see Figure 2 for definition of the confidence bands.




Figure 2: Posterior distribution for the bulk viscosity to entropy ratio versus temperature, with a 90% confidence band shown in blue, from Trajectum [4] (left) and Jetscape[6] (right).

we can take on the xy plane with no loss of generality, using rotational symmetry:

$$\eta = - \lim_{\omega \rightarrow 0} \frac{1}{\omega} \text{Im} G_{xy,xy}^R(\omega, k=0), \quad (3.1)$$

where the retarded Green's function is at vanishing momentum and in the thermal state. It is given by

$$G_{xy,xy}^R(\omega, \mathbf{k} = 0) = -i \int dt d\mathbf{x} e^{i\omega t} \theta(t) \langle [T_{xy}(x), T_{xy}(0)] \rangle. \quad (3.2)$$

Bulk deformations correspond to a scaling of the spatial volume, which is related to the trace of the metric fluctuation. The corresponding Kubo formula reads

$$\zeta = -\frac{4}{9} \lim_{\omega \rightarrow 0} \frac{1}{\omega} \text{Im} G_{ii}^R \omega, \quad (3.3)$$

where the Green's function is given by

$$G_{ii}^R(\omega) = -i \int dt d\mathbf{x} e^{i\omega t} \theta(t) \left\langle \left[\frac{1}{2} T_i^i(t, \mathbf{x}), \frac{1}{2} T_j^j(0, \mathbf{0}) \right] \right\rangle. \quad (3.4)$$

In holography, the $\omega \rightarrow 0$ limit of the Green's functions follows from the corresponding fluctuation of the black-brane background, with non-normalizable boundary condition at the boundary and infalling boundary condition at the horizon⁴ [49].

As mentioned in the introduction, we simplify our analysis by considering only the Riemann-square type corrections⁵. Hence the generic form of our action reads

$$S = \frac{1}{16\pi G_5} \int d^5x \sqrt{-g} \left[R - \frac{4}{3} (\nabla\Phi)^2 + V(\Phi) + \ell^2 \beta G(\Phi) R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} \right], \quad (3.5)$$

where G_5 is the 5D Newton constant, ℓ is the AdS length, β is a parameter which controls the magnitude of the higher curvature corrections and V and G are generic functions of the dilaton Φ . In particular, the background solution is pure AdS_5 for $V = 12/\ell^2$ and $G = 0$. For the action (3.5), the calculation of [38] implies that the shear viscosity per entropy⁶ can be expressed completely in terms of the potentials in (3.5) as

$$\frac{\eta}{s} = \frac{1}{4\pi} \left[1 + \frac{2}{3} \beta \ell^2 \left(-G(\Phi_h) V(\Phi_h) + \frac{9}{8} \partial_\Phi G(\Phi_h) \partial_\Phi V(\Phi_h) \right) \right], \quad (3.6)$$

where Φ_h is the value of the dilaton at the horizon. A similar derivation⁷ in the case of bulk viscosity per entropy (generalizing the result of [36]) leads to

$$\frac{\zeta}{s} = \frac{8z_0^2}{27\pi} \left[1 - \frac{1}{2} \beta \ell^2 \left(\partial_\Phi V(\Phi_h) \partial_\Phi G(\Phi_h) + \frac{(\partial_\Phi V(\Phi_h))^2 G(\Phi_h)}{V(\Phi_h)} - \frac{4}{3} V(\Phi_h) G(\Phi_h) \right) \right], \quad (3.7)$$

where z_0 is the horizon value of the gauge invariant dilaton fluctuation in the hydrodynamic limit (see appendix D). The expression (3.7) can readily be extended to the case where the

⁴An alternative derivation involves formulating the fluctuation equations as background variations which has the advantage of providing analytic formulae for both shear and bulk viscosity [34, 35].

⁵It can actually be shown that the shear viscosity per entropy is not modified by the other types of curvature squared corrections R^2 and $R_{\mu\nu}^2$. The argument is presented in appendix C.

⁶We note that the usual Bekenstein-Hawking formula for the entropy in terms of the area of the black brane horizon is modified by Wald's formula in the presence of higher curvature corrections [50].

⁷The details of the calculation can be found in appendix D.

action contains all three different types of 4-derivative corrections

$$S = \frac{1}{16\pi G_5} \int d^5x \sqrt{-g} \left[R - \frac{4}{3} (\nabla\Phi)^2 + V(\Phi) + \ell^2 \beta (G_1(\Phi) R^2 + G_2(\Phi) R_{\mu\nu} R^{\mu\nu} + G_3(\Phi) R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma}) \right]. \quad (3.8)$$

The same analysis in this case yields the following bulk viscosity to entropy ratio (see appendix D):

$$\frac{\zeta}{s} = \frac{8z_0^2}{27\pi} \left[1 - \frac{1}{2} \beta \ell^2 \left(\partial_\Phi V(\Phi_h) \partial_\Phi \tilde{G}(\Phi_h) + \frac{(\partial_\Phi V(\Phi_h))^2 \tilde{G}(\Phi_h)}{V(\Phi_h)} - \frac{4}{3} V(\Phi_h) \tilde{G}(\Phi_h) \right) \right], \quad (3.9)$$

where we defined $\tilde{G} \equiv 5G_1 + G_2 - G_3$.

The formulae (3.6) and (3.7) will form the basis for our comparison with the Bayesian analysis below. In particular, (3.6) can be seen as a differential equation for $G(\Phi_h)$, which implies that any given profile $\eta/s(T)$ can be reproduced⁸ with the appropriate potential G . The explicit solution takes the form

$$G(\Phi_h) = F^{-1}(\Phi_h) \int d\Phi_h F(\Phi_h) R(\Phi_h) + c_1 F^{-1}(\Phi_h), \quad (3.10)$$

where

$$F(\Phi_h) \equiv \exp \left\{ \int \left(-\frac{8}{9} \frac{V(\Phi_h)}{\partial_\Phi V(\Phi_h)} \right) d\Phi_h \right\}, \quad R(\Phi_h) \equiv \left[4\pi \frac{\eta}{s}(T(\Phi_h)) - 1 \right] \frac{4}{3\ell^2 \beta \partial_\Phi V(\Phi_h)}. \quad (3.11)$$

The integration constant c_1 which multiplies the homogeneous solution in (3.10) does not affect the viscosity since it cancels in (3.6).

3.2 Holographic setup

As our starting point, we will consider the V-QCD model [16], which is itself an extension of improved holographic QCD (ihQCD) [14, 15]. These models are based on the assumption that the operator product algebra of QCD is semi-closed in the IR, and consists of only a small subset of UV marginal and irrelevant operators [51–53]: the stress tensor $T_{\mu\nu}$, the scalar glueball operator $\text{tr}F^2$, chiral condensate $\bar{q}q$, left and right $\text{SU}(N_f)$ chiral currents L^μ and R^μ , and the $\text{U}(1)_B$ baryon number current V^μ ⁹. The corresponding holographic bulk fields are the metric, dilaton, open string tachyon and gauge fields. These fields propagate on a five-dimensional asymptotically AdS background, which is a solution of an Einstein-dilaton theory with space-filling flavor branes. See [54] for a review. A non-trivial dilaton potential $V(\Phi)$ yields a non-trivial profile for Φ in holographic coordinate r , which corresponds to the running of the QCD 't Hooft coupling λ_{QCD} in the IR. Accordingly, the

⁸Note that going from $\eta/s(T)$ to $\eta/s(\Phi_h)$ also requires to know the temperature as a function of the horizon location $T = T(\Phi_h)$. The latter can be extracted in a standard way from the background solution, by computing the surface gravity at the horizon.

⁹One can extend this to include the CP-odd operator $\text{tr}F \wedge F$ but this can be ignored in the large N_c limit at finite temperature.

“glue” part of the theory (pure Yang-Mills at large N_c), improved further by a Riemann-square correction, is given by the action

$$S_{glue} = M_p^3 N_c^2 \int d^5 x \sqrt{-g} \left[R - \frac{4}{3} (\partial\Phi)^2 + V_g(\Phi) + \ell^2 \beta G(\Phi) R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} \right], \quad (3.12)$$

where we wrote $16\pi G_5 = 1/(M_p^3 N_c^2)$, with N_c the number of colors and M_p the 5D Planck mass, which controls the overall normalization of the free energy. A judicious choice of V_g asymptotics for large and small λ ensures confinement and asymptotic freedom in the dual theory [14, 15]. Deconfinement is described by a Hawking-Page transition between the low temperature thermal graviton gas solution, and the high temperature black-brane solution [19]. Thermodynamics in the deconfined black-brane phase agrees remarkably well with large N_c extrapolations of lattice QCD simulations [18, 20, 55].

V-QCD is obtained from ihQCD by coupling the “flavor” part, which consists of a number N_f of space-filling $D4$ and $\overline{D4}$ branes with the DBI type action

$$S_{flavor} = -x_f M_p^3 N_c^2 \int d^5 x V_{f0}(\Phi) e^{-\tau^2} \sqrt{-\det \left(g_{\mu\nu} + \kappa(\Phi) \partial_\mu \tau \partial_\nu \tau + w(\Phi) \hat{F}_{\mu\nu} \right)}. \quad (3.13)$$

The endpoints of open strings on these branes model left and right-handed quarks and the condensation of the open-string tachyon field τ models chiral symmetry breaking holographically. We will be interested in the Veneziano limit [17]:

$$N_c \rightarrow \infty, \quad N_f \rightarrow \infty, \quad x_f \equiv \frac{N_f}{N_c} = \text{fixed}, \quad (3.14)$$

so that the back-reaction of the flavor fields on the color sector will be non-negligible. The background solutions will then be obtained by extremizing the total action $S_{glue} + S_{flavor}$.

The flavor action (3.13) depends on three additional potentials $\kappa(\Phi)$, $w(\Phi)$, $V_{f0}(\Phi)$, assumed to only depend on the dilaton Φ . Small and large Φ asymptotics of these potentials follow from the requirement to reproduce a number of salient features of QCD in the mesonic sector. Given these constraints, there remains some indeterminacy in the precise form of the potentials, which is however limited by fitting both heavy-ion collision experiments and neutron star mergers, see [22, 47, 54, 56–61]. Based on these fits we make the following choice for the potentials:

$$\begin{aligned} V_g(\Phi) &= 12 \left[1 + V_1 \lambda + \frac{V_2 \lambda^2}{1 + \lambda/\lambda_0} + V_{IR} e^{-\lambda_0/\lambda} (\lambda/\lambda_0)^{4/3} \sqrt{\log(1 + \lambda/\lambda_0)} \right], \\ V_{f0}(\Phi) &= W_0 + W_1 \lambda + \frac{W_2 \lambda^2}{1 + \lambda/\lambda_0} + 12 W_{IR} e^{-\lambda_0/\lambda} (\lambda/\lambda_0)^2, \end{aligned} \quad (3.15)$$

where we defined $\lambda = \exp \Phi$ to simplify notation. It is important to stress that all the parameters that appear above are fixed. On the UV side, they are matched to the RG flow of QCD perturbation theory, and on the IR side they are determined by comparing to lattice data. Below we present the parameter values that we use, corresponding to the potential set 7a in [54] which is a particular V-QCD model with intermediate stiffness of

the equation of state:

$$V_1 = \frac{11}{27\pi^2}, \quad V_2 = \frac{4619}{46656\pi^4}, \quad W_1 = \frac{8 + 3W_0}{9\pi^2}, \quad W_2 = \frac{6488 + 999W_0}{15552\pi^4} \quad (3.16)$$

$$\lambda_0 = 8\pi^2/3, \quad V_{IR} = 2.05, \quad W_0 = 2.5, \quad W_{IR} = 0.9, \quad \Lambda_{UV} = 211\text{MeV}.$$

The effective potential for V-QCD that enters the semi-analytic formulas for shear and bulk viscosity per entropy, given in section 3.1, is the combination

$$V_{eff}(\Phi) = V_g(\Phi) - x_f V_{f0}(\Phi). \quad (3.17)$$

In the following, the ratio $x_f = N_f/N_c$ is set to 1. Figure 6 provides a plot of the effective V-QCD potential (3.17) in this case, with potential parameters chosen as in (3.16).

Our setup is different from usual V-QCD, since it includes a higher derivative term in the glue action (3.12). In the following, it is assumed that this term only leads to small order $\mathcal{O}(\beta)$ corrections to the background, which do not alter dramatically the properties of the holographic-QCD theories. In particular, we require the IR asymptotics to preserve the confinement criteria established in [15]. The $\mathcal{O}(\beta)$ corrections to both the UV and IR asymptotics are worked out in appendix A, where it is shown in particular that confinement is preserved for a generic ansatz

$$G(\Phi) = G_{IR} e^{\gamma\Phi} \Phi^\delta, \quad \gamma \leq -\frac{4}{3}. \quad (3.18)$$

4 Fitting holographic models to Bayesian analyses

4.1 Shear viscosity

We explained at the end of section 3.1 how to fit any desired temperature profile for η/s by appropriately choosing the function $G(\Phi)$. However, the profiles that follow from the Bayesian analysis, see Figure 1, contain wide errors allowing for a wide range of choices for this function. We therefore start with the simplest choice, $G = 1$, and ask whether the data can be fitted within the 90% confidence bands.

For $G = 1$, (3.6) reduces to

$$\frac{\eta}{s} = \frac{1}{4\pi} \left[1 - \frac{2}{3} \beta V(\Phi_h) \right], \quad (4.1)$$

were we set $\ell = 1$. Since we work with a fixed V-QCD potential $V_{eff}(\Phi_h)$ (see (3.15)-(3.17)), the only free parameter is β . Assuming $\beta \ll 1$, the relation between the temperature and Φ_h can be determined from the black-brane surface gravity omitting the curvature corrections. The resulting profile for $T(\Phi_h)$ is shown in Figure 3, while Figure 4 shows the η/s temperature profile for the value of β that gives the best fit to data. This indicates that a constant $G(\Phi)$ does not provide a good fit. Even though the curve seems to be in agreement with the 90% confidence range of the Jetscape data, we do not obtain an η/s profile that monotonically increases with temperature as expected from the running of the gauge coupling.




Figure 3: Temperature in units of the confining scale Λ , as a function of the dilaton horizon value in V-QCD. The dotted line indicates the value of the critical temperature T_c/Λ , where the deconfining transition happens.




Figure 4: Shear viscosity to entropy ratio for V-QCD with $G(\Phi) = 1$. The blue area corresponds to the 90% confidence band for Trajectum while the green area corresponds to the 90% confidence band for Jetscape. The orange line shows the theoretical curve while the black line is $\eta/s = 1/(4\pi)$ for reference. The best fit is obtained for $\beta = -0.022$.

We now move on to the case of general $G(\Phi)$ and solve the differential equation (3.6) numerically to determine the optimal fit to Bayesian data. As an example we consider the data from Trajectum, see Figure 1. We set $\beta = 0.1$ without loss of generality and compute the function G which reproduces the mean of the Bayesian data, shown as the orange curve in Figure 5. Equation (3.10) gives the corresponding potential $G(\Phi)$, which is plotted in Figure 6 (right), along with the effective V-QCD potential $V_{eff}(\Phi)$ (left). A few remarks are in order:




Figure 5: Best fit for the shear viscosity to entropy ratio for V-QCD with $G(\Phi)$ shown in Figure 6 (β is set to 0.1). The orange line is the theoretical curve.




Figure 6: Dilaton potentials for the model with η/s as in Figure 5.

- As mentioned above, the integration constant c_1 in (3.10) is irrelevant to η/s . It will matter however for the fit to bulk viscosity. In Figure 6, c_1 was chosen¹⁰ such that $G(\Phi_h = 4.5) = 0.2$.
- Outside of the fitting region shown in Figure 6, we make sure that G drops off to zero according to the asymptotic behaviors derived in appendix A. These behaviors ensure that the order $\mathcal{O}(\beta^0)$ asymptotics of the background and the associated QCD-like properties of the model are not modified by the curvature corrections.
- More importantly, we find that the correction to the background solution for this choice of G violates our perturbative treatment. This means that, to fit Bayesian data with our holographic model, it is necessary to solve the metric equations including

¹⁰This choice was made such that G is flat for large Φ hence the derivatives remain small, so that we stay within the perturbative regime.

the curvature corrections. This indicates that finite coupling corrections $1/\lambda_{QCD}$ are significant. Below, we present an alternative fitting scheme for both viscosities which agrees less well with the Bayesian data but remains within the perturbative regime.

4.2 Bulk viscosity

In this section, we focus on comparison of the bulk viscosity per entropy with the Bayesian analysis. For simplicity, we again consider the V-QCD action with a single Riemann-square type correction with coupling β given by $S_{glue} + S_{flavor}$ (see (3.12) and (3.13)).

4.2.1 Bulk viscosity of V-QCD theory

As a first step we calculate ζ/s for the original V-QCD theory with no curvature corrections, i.e. with $\beta = 0$. The easiest way to compute the bulk viscosity in this case is to employ the Eling-Oz formula [30]:

$$\frac{\zeta}{s} = \frac{2}{3\pi} \left(s \frac{\partial \Phi_h}{\partial s} \right)^2. \quad (4.2)$$

Equivalence between this formula and the standard holographic computation through fluctuating the background has been established in [34].

The resulting V-QCD bulk viscosity is plotted in Figure 7 as a function of temperature, along with the 90% confidence bands of the two data sets. Some comments about this fit:




Figure 7: Bulk viscosity to entropy ratio as a function of temperature for V-QCD without curvature corrections. The orange line is the model prediction. The green region is the 90% confidence band from the Jetscape collaboration and is derived from combined RHIC and LHC data. The blue region comes from the analysis of the Trajectum collaboration.

- The fit has no additional free parameters since the V-QCD potential used is completely fixed by other requirements; see e.g. [54].




Figure 8: Bulk and shear viscosities for $\beta G = -0.002$ compared with Trajectum and explicit check of the magnitude of the corrections for the background and bulk perturbations. The bottom left plot is the relative change in the Ricci scalar near the horizon and boundary. The bottom right plot is the relative change in the gauge invariant dilaton fluctuations at the horizon. Both are largest in absolute value at the horizon. In all figures the temperature is in GeV.

- We observe a better agreement with the data from Jetscape for this zeroth order result. Thus, V-QCD without curvature corrections implies that bulk viscosity is not suppressed compared to shear viscosity, unlike what Trajectum data suggest. This hints at a soft but non-negligible breaking of conformality.

4.2.2 Bulk viscosity with curvature corrections

We now discuss the effect of curvature corrections on the V-QCD bulk viscosity per entropy. In particular, we would like to investigate whether these corrections can bring ζ/s closer to either of the bands from the Bayesian analyses. As for the shear viscosity, we can obtain some insight by starting from the simplest form of curvature corrections, namely a Riemann-square term with constant G . The Eling-Oz formula has not yet been generalized




Figure 9: Same as Figure 8, but for $\beta G = 0.001$ and comparing with Jetscape results.

to include higher curvature corrections, hence we need to follow the standard holographic method for this calculation, based on fluctuating the background [36]. Details can be found in appendix D and E.

Figure 8 shows the temperature profile of ζ/s for $\beta G = -0.002$. The latter is seen to shift closer to Trajectum at low enough temperatures while remaining within the regime of applicability of our perturbative calculation, as shown by the bottom figures. The profile can be brought into the Trajectum band for more negative values of βG , but this eventually induces large corrections to the background, outside of the perturbative regime.

The temperature profile of ζ/s can also be made compatible with the Jetscape band if we consider positive values of βG . This is shown in Figure 9, where we set $\beta G = 0.001$. The bottom plots indicate that this calculation also consistently remains within the perturbative regime, although it is not completely clear for the Z fluctuation.

As an indication, Figures 8 and 9 also show the corresponding profiles of $(\eta/s)(T)$, which are mostly temperature independent as we saw above.




Figure 10: Relative change in the bulk and shear viscosities with the $\mathcal{O}(\beta)$ result of V-QCD for a linearly increasing η/s (4.3) with a slope $\gamma = 0.001$. The bottom left plot is the relative change in the Ricci scalar at different points in the holographic coordinate. The bottom right plot is the relative change in the gauge invariant dilaton fluctuations at the horizon. Both are largest in absolute value at the horizon. In all the figures the temperature is in GeV.

4.3 Global fits to both shear and bulk viscosity data

The analysis of the previous subsection shows that the bulk viscosity can be brought close to either Trajectum or Jetscape data with a constant coefficient βG for the curvature corrections. In this case however, we saw in section 4.1 that the shear viscosity is essentially a constant, which is not compatible with the increase expected in QCD, nor with the Trajectum results. We also observed a tension between η/s increasing in T and our perturbative treatment of the curvature corrections.

In this subsection, we perform a joint analysis of the shear and bulk viscosity, where the potential $G(\Phi)$ is fixed from (3.6) by imposing that η/s increases with temperature while remaining within the perturbative regime. The goal is to observe the qualitative effect of curvature corrections on the bulk viscosity in this more realistic scenario. For this analysis, we consider a simple linear behavior for $(\eta/s)(T)$, of the form

$$\eta/s = 1/4\pi + \gamma(T - T^*), \quad (4.3)$$

with γ parametrizing the slope of the temperature profile. We find that a choice $\gamma \lesssim 0.001$ leads to corrections to the background that are small everywhere, so that our perturbative treatment is applicable.




Figure 11: Shear viscosity to entropy ratio for an AdS solution with constant higher derivative corrections. The value of $\beta = -0.064$ was picked such that it provides the best fit to the Trajectum central value of η/s , while at the same time requiring that the change in the background Ricci scalar induced by the higher curvature terms should obey $\delta R/R < 0.1$. The right plot is the relative difference in η/s with the central Trajectum value, $[(\eta/s)_{\text{Traj}} - \eta/s]/(\eta/s)_{\text{Traj}}$, as a function of temperature.

The resulting relative differences in the viscosities and the corrections to the background are shown in Figure 10. Unlike the constant G case, the qualitative change in ζ/s is observed to depend on temperature: it decreases at low temperatures but increases at higher temperatures. There is therefore no indication of ζ/s approaching either of the Bayesian data when η/s is imposed to increase with temperature, although a fully non-perturbative analysis may lead to different conclusions. This suggests that fitting both types of viscosities requires including other types of curvature corrections (R^2 and $R_{\mu\nu}^2$ at quadratic order).

As a final note, we would like to discuss the alternative of working with different holographic models. In particular, demanding that the effect of higher-curvature corrections on the background is negligible, the model used here seems to fit the Jetscape data better (which is compatible with constant η/s). Then, it is natural to ask whether one can fit the two separate data sets using different holographic models. In particular, it is tempting to use a constant potential V for Trajectum, as this data set is consistent with vanishing bulk viscosity, as arises for a conformal theory dual to pure AdS.

Taking G to be a constant preserves conformal invariance at the higher derivative level, so that we only need to fit η/s . However, constant V and G imply a constant η/s , so that a conformal model can only give a coarse approximation to the steep temperature profile of η/s in the Trajectum data. This is shown in more detail in Figure 11, which compares the Trajectum central value $(\eta/s)_{\text{Traj}}$ with the best constant fit compatible with

a perturbative treatment of the curvature corrections¹¹. In particular, the relative difference $[(\eta/s)_{\text{Traj}} - \eta/s]/(\eta/s)_{\text{Traj}}$ (right Figure 11) is seen to exceed 50% at the edges of the fitting range.

5 Discussion and Outlook

We explored various possibilities to fix the curvature corrections in the realistic holographic QCD model, V-QCD, in a way consistent with the temperature dependence of shear and bulk viscosity observed in the Bayesian analysis of heavy-ion collisions [4–6].

We treated curvature corrections in a perturbative manner. In particular, this allowed us to work with second order equations of motion, instead of the full fourth order problem. Solving the fourth order equations is in principle possible. However, it is well known that higher derivative terms in metric equations may lead to causality violation, see e.g. [62, 63]. It would be interesting to explore the causality of the 4-derivative corrections to holographic-QCD theories in general. Staying within the causal range, it would make it possible to go beyond the perturbative treatment. This could provide more flexibility in fitting both viscosity profiles.

However, one should then also ask whether such corrections are compatible with the previously established properties of ihQCD or V-QCD holographic models i.e., confinement, desired running of the 't Hooft coupling, asymptotics of glueball and meson spectra, chiral symmetry breaking, holographic realization of QCD anomalies, properties of the phase diagram at finite temperature and chemical potential, etc. In principle, we would need to make another global fit including both the potentials related to curvature corrections, G_1 , G_2 and G_3 and also the other potentials of V-QCD i.e. V_g , V_{f0} , κ and w , with the requirement of the aforementioned properties.

Even if we manage to reproduce the desired QCD features, there remains the problem of other higher curvature corrections in a non-perturbative treatment. That is, once we treat the four-derivative corrections as the same order as the original background, then why not include higher derivative corrections. One possible answer is to treat the theory with four-derivative corrections only, as a phenomenological model, which fits well some QCD observables (for example the viscosities per entropy), but is probably less good at reproducing other QCD properties away from the IR regime. Another possibility is to try to include a whole series of higher derivative corrections to the 5D non-critical string theory altogether by introducing an $O(d, d)$ invariant action as in the double field theory; see e.g. [64, 65]. In this case, one could also employ the constraints that arise in the “swampland” program [66, 67], see e.g. [68] for a review, which in general restricts the range of consistent UV-complete higher derivative theories. All these questions are clearly interesting but beyond the scope of this work.

Apart from these structural questions, an interesting extension of our work involves charge transport. Presence of dense magnetic fields in off-central collisions results in various charge-transport related effects in the quark-gluon plasma with observable consequences;

¹¹Our criterium is such that the change in the background Ricci scalar induced by the curvature squared terms should obey $\delta R/R < 10\%$.

see for example [69–77], and [78] for a review. Indeed, extensions of the Bayesian analyses to include charge transport coefficients and the presence of electromagnetic fields in off-central events would, most importantly, also provide a temperature profile for conductivity, σ . Now, there is also a universal formula for conductivity which leads to a model-independent constant for the ratio of σ and $s^{1/3}$ in two-derivative Einstein gravity [79]. To go beyond this and model a more generic temperature dependence in holography one also needs higher derivative corrections to extensions of holographic theories that already include magnetic fields e.g. [24, 80–84]; see [85] for a review. We wish to turn back to the points listed here in the future.

Acknowledgements

We are grateful to Alex Buchel, Sera Cremonini and Govert Nijs for useful correspondence. EP and UG are supported by the Netherlands Organization for Scientific Research (NWO) under the VICI grant VI.C.202.104. Work of UG is also supported by the ENW-XL program “Probing the QCD phase diagram” of NWO.

A The vacuum solution

We discuss in this appendix the zero-temperature solution corresponding to the vacuum of the theory. We start from presenting the equations of motion that it solves, before discussing the behavior in the UV and IR asymptotic limits. As mentioned in the main text, the asymptotics of the V-QCD potentials are chosen to reproduce several salient qualitative features of QCD. We describe how requiring that those features are not spoiled by the higher curvature corrections puts constraints on the additional potential $G(\Phi)$. In the following, we work with ihQCD but the extension to V-QCD is straightforward.

A.1 Equations of motion

The equations of motion are derived from the action (3.12). The dilaton equation takes the form

$$\frac{8}{3}\nabla_\mu\nabla^\mu\Phi + \ell^2\beta R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}\partial_\Phi G(\Phi) + \partial_\Phi V(\Phi) = 0, \quad (\text{A.1})$$

whereas the metric equation is given by

$$\begin{aligned} & R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R - \frac{1}{2}g_{\mu\nu}V(\Phi) - \frac{4}{3}(\nabla_\mu\Phi)(\nabla_\nu\Phi) + \frac{2}{3}g_{\mu\nu}(\nabla_\sigma\Phi)(\nabla^\sigma\Phi) + \\ & + \ell^2\beta\left[2G(\Phi)R_\mu^{\rho\sigma\alpha}R_{\nu\rho\sigma\alpha} - \frac{1}{2}g_{\mu\nu}G(\Phi)R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} + 2G(\Phi)\nabla^\rho\nabla^\sigma R_{\mu\rho\nu\sigma} + \right. \\ & + 2G(\Phi)\nabla^\sigma\nabla^\rho R_{\mu\rho\nu\sigma} + 4(\nabla^\rho\Phi)\nabla^\sigma R_{\mu\rho\nu\sigma}\partial_\Phi G(\Phi) + 4(\nabla^\rho\Phi)(\nabla^\sigma R_{\mu\sigma\nu\rho})\partial_\Phi G(\Phi) + \\ & + 2R_{\mu\rho\nu\sigma}(\nabla^\sigma\nabla^\rho\Phi)\partial_\Phi G(\Phi) + 2R_{\mu\rho\nu\sigma}(\nabla^\rho\nabla^\sigma\Phi)\partial_\Phi G(\Phi) + \\ & \left. + 4R_{\mu\rho\nu\sigma}(\nabla^\rho\Phi)(\nabla^\sigma\Phi)\partial_\Phi^2 G(\Phi)\right] = 0, \end{aligned} \quad (\text{A.2})$$

where $V(\Phi)$ is the ihQCD dilaton potential.

The background solutions that we are interested in are homogeneous, horizon-less configurations, for which an appropriate ansatz is given by

$$ds^2 = e^{2A(r)}(dr^2 - dt^2 + d\mathbf{x}^2), \quad \Phi = \Phi(r), \quad (\text{A.3})$$

where r is the holographic coordinate, which goes to zero at the boundary. Substituting this ansatz into the equations of motion (A.1)-(A.2) results in the following set of equations for the ansatz fields $A(r)$ and $\Phi(r)$

$$\begin{aligned} & 108\ell^2\beta G(\Phi)A'(r)^4 - 36\ell^2\beta A'(r)^3\partial_\Phi G(\Phi)\Phi'(r) - 9e^{2A(r)}A''(r) + \\ & + 48\ell^2\beta G(\Phi)A''(r)^2 + 3A'(r)^2\left(3e^{2A(r)} - 52\ell^2\beta G(\Phi)A''(r)\right) - \\ & - 4\Phi'(r)^2\left(e^{2A(r)} - 3\ell^2\beta A''(r)\partial_\Phi^2 G(\Phi)\right) \\ & + 12\ell^2\beta\partial_\Phi G(\Phi)A''(r)\Phi''(r) + 24\ell^2\beta\partial_\Phi G(\Phi)\Phi'(r)A^{(3)}(r) - \\ & - 36\ell^2\beta A'(r)(\partial_\Phi G(\Phi)\Phi'(r)A''(r) + G(\Phi)A^{(3)}(r)) + 12\ell^2\beta G(\Phi)A^{(4)}(r) = 0, \end{aligned} \quad (\text{A.4})$$

$$\begin{aligned}
& e^{4A(r)}V(\Phi) - 3e^{2A(r)}A''(r) - A'(r)^2 \left(9e^{2A(r)} + 20\ell^2\beta G(\Phi)A''(r) \right) - \\
& - 12\ell^2\beta A'(r)^3 \partial_\Phi G(\Phi) \Phi'(r) + 4\ell^2\beta \Phi'(r)^2 A'(r) \partial_\Phi^2 G(\Phi) + 4\ell^2\beta \partial_\Phi G(\Phi) A''(r) \Phi''(r) + \\
& + 8\ell^2\beta \partial_\Phi G(\Phi) \Phi'(r) A^{(3)}(r) + 20\ell^2\beta A'(r) \left(\partial_\Phi G(\Phi) \Phi'(r) A''(r) + G(\Phi) A^{(3)}(r) \right) + \\
& + 4\ell^2\beta G(\Phi) A^{(4)}(r) - 36\ell^2\beta G(\Phi) A'(r)^4 = 0,
\end{aligned} \tag{A.5}$$

where primes denote radial derivatives.

A.2 UV Asymptotics

We now analyze the UV limit ($r \rightarrow 0$) of the equations of motion (A.4)-(A.5). Our goal is to identify conditions on the potential $G(\Phi)$ such that the UV asymptotics of the vacuum are not modified by the curvature corrections. In particular, we require that the scale factor $A(r)$ follows the same UV expansion as in [15]

$$A(r) = -\log(r/\ell) + \frac{4}{9\log(r\Lambda)} + \mathcal{O}\left(\frac{-\log(\log(r\Lambda))}{\log(r\Lambda)^2}\right). \tag{A.6}$$

Substituting (A.6) into the equation of motion (A.4) gives the following relation at leading order in the $r \rightarrow 0$ limit

$$\begin{aligned}
& 3(\log(\Lambda r))^2 \left((\log(\Lambda r))^4 - (\log(\Lambda r))^6 r^2 \Phi'(r)^2 \right) - 4\beta G(\Phi) (\log(\Lambda r))^6 \\
& 3\beta \partial_\Phi G(\Phi) \left(6(\log(\Lambda r))^6 \Phi'(r) + 9\log(\Lambda r)^6 r \Phi''(r) \right) + 9\beta \partial_\Phi^2 G(\Phi) (\log(\Lambda r))^6 + \dots = 0,
\end{aligned} \tag{A.7}$$

where the dots indicate subleading terms in the $r \rightarrow 0$ limit. If $G(\Phi)$ is assumed to go to a constant G_0 in the UV limit ($\Phi \rightarrow -\infty$), then (A.7) determines the leading order behavior of the dilaton to be

$$\Phi(r) = -\log(-\log(r\Lambda)) \left(1 - \frac{4}{3}\beta G_0 \right) + \dots, \tag{A.8}$$

which is a non-perturbative result in β . In particular, requiring that the leading order UV behavior of the dilaton is not modified by the curvature corrections imposes that

$$G(\Phi) \xrightarrow[\Phi \rightarrow -\infty]{} 0. \tag{A.9}$$

Requiring that the subleading terms in the UV expansion of the dilaton (which are fixed in ihQCD by matching to the QCD β -function) are also unchanged, would put further constraints on the way G approaches 0 in the UV.

The other equation of motion (A.5) takes the following form at leading order in the UV limit

$$\frac{\ell^2 V(\Phi) - 12}{r^2} + \frac{8\beta G(\Phi)}{r^2} = \partial_\Phi G(\Phi) + \partial_\Phi^2 G(\Phi) + \dots \tag{A.10}$$

In the case where $G(\Phi)$ goes to a constant, (A.10) implies that the AdS curvature receives a correction proportional to β

$$V(\Phi) \rightarrow V_0 = \frac{12}{\ell^2} - \frac{8\beta G_0}{\ell^2}. \tag{A.11}$$

This correction to the UV geometry vanishes when (A.9) is imposed.

A.3 IR Asymptotics

We now discuss the IR asymptotics of the vacuum solution, corresponding to $r \rightarrow \infty$. The behavior of the background in this limit determines several key qualitative features of the model, including confinement and general properties of the hadronic spectrum. In particular, the best match with QCD properties is obtained when the ihQCD potential $V(\Phi)$ behaves in the IR as

$$V(\Phi) \underset{\Phi \rightarrow \infty}{\sim} V_{IR} e^{Q\Phi} \Phi^P. \quad (\text{A.12})$$

The purpose of our analysis is to understand what are the IR asymptotics of $G(\Phi)$ for which these properties hold, once the curvature corrections are introduced. For the remainder of this appendix, β will now be assumed to be small, such that the background fields follow the same asymptotics as in [14, 15], up to small corrections of order $\mathcal{O}(\beta)$

$$\begin{aligned} A(r) &= -Cr^\alpha + \dots + \beta\delta A(r) + \mathcal{O}(\beta^2), \quad \alpha > 0, C > 0 \\ \Phi(r) &= -\frac{3}{2}A(r) + \frac{3}{4}\log|A'(r)| + \dots + \beta\delta\Phi(r) + \mathcal{O}(\beta^2), \end{aligned} \quad (\text{A.13})$$

where

$$\alpha = \frac{1}{1-P}, \quad (\text{A.14})$$

and the dots refer to subleading terms in the $r \rightarrow \infty$ limit. We ask how $G(\Phi)$ should behave in the IR such that $\delta A(r)$ and $\delta\Phi(r)$ remain small up to $r \rightarrow \infty$.

Substituting (A.12)-(A.13) into the equations of motion (A.4)-(A.5) and keeping only the terms of order $\mathcal{O}(\beta)$, we find the following relation at leading order in $1/r$

$$\begin{aligned} \frac{8}{3}\alpha^2 C^2 \delta\Phi'(r) + (1-\alpha)r^{-2\alpha}((1-\alpha)\delta A'(r) + r\delta A''(r)) = \\ e^{2Cr^\alpha} r^{3(\alpha-1)} \times 6C^4 \ell^2 \alpha^4 (2\alpha C(2G(\Phi) - \partial_\Phi G(\Phi)) + (\alpha-1)r^{-\alpha} \partial_\Phi^2 G(\Phi)) + \dots + \mathcal{O}(\beta). \end{aligned} \quad (\text{A.15})$$

$$\begin{aligned} \frac{4}{3}\tilde{V}_{IR} r^{\alpha-1} \delta\Phi(r) + \delta A'(r) \left(18\alpha C - (\alpha C)^{-1} \tilde{V}_{IR} \right) - 3r^{1-\alpha} \delta A''(r) = \\ 18\ell^2 (\alpha C)^4 e^{2Cr^\alpha} r^{3(\alpha-1)} (-2G(\Phi) + \partial_\Phi G(\Phi)) + \dots + \mathcal{O}(\beta), \end{aligned} \quad (\text{A.16})$$

where we defined

$$\tilde{V}_{IR} \equiv V_{IR} (3C/2)^P (-\alpha C). \quad (\text{A.17})$$

Note that the right-hand side of (A.15) and (A.16) is multiplied by an exponential that diverges in the $r \rightarrow \infty$ limit, whereas the order $\mathcal{O}(\beta^0)$ background has a polynomial behavior (see (A.13)). So for the order $\mathcal{O}(\beta)$ corrections to remain sub-leading in the IR, the function $G(\Phi)$ should be such that the exponential divergence cancels in (A.15). Given the asymptotics of Φ (A.13), an appropriate ansatz for G is given by

$$G(\Phi) = G_{IR} e^{\gamma\Phi} \Phi^\delta. \quad (\text{A.18})$$

which implies the following behavior at leading order in β and $1/r$

$$G(\Phi(r)) = G_{IR} (\alpha C)^{\frac{3}{4}\gamma} \left(\frac{3}{2}C \right)^\delta e^{\frac{3}{2}\gamma Cr^\alpha} r^{\frac{3}{4}\gamma(\alpha-1) + \alpha\delta} + \dots + \mathcal{O}(\beta). \quad (\text{A.19})$$

Plugging this back into (A.15)-(A.16) gives

$$\begin{aligned} \frac{8}{3}\alpha^2 C^2 \delta\Phi'(r) + (1-\alpha)r^{-2\alpha} \left((1-\alpha)\delta A'(r) + r\delta A''(r) \right) = \\ \tilde{G}_{IR} e^{2Cr^\alpha(1+\frac{3}{4}\gamma)} r^{3(\alpha-1)(1+\frac{1}{4}\gamma)+\alpha\delta} + \dots + \mathcal{O}(\beta), \end{aligned} \quad (\text{A.20})$$

$$\begin{aligned} \frac{4}{3}\tilde{V}_{IR} r^{\alpha-1} \delta\Phi(r) + \delta A'(r) \left(18\alpha C - (\alpha C)^{-1} \tilde{V}_{IR} \right) - 3r^{1-\alpha} \delta A''(r) = \\ -\frac{3}{2}(\alpha C)^{-1} \tilde{G}_{IR} e^{2Cr^\alpha(1+\frac{3}{4}\gamma)} r^{3(\alpha-1)(1+\frac{1}{4}\gamma)+\alpha\delta} + \dots + \mathcal{O}(\beta), \end{aligned} \quad (\text{A.21})$$

where we defined

$$\tilde{G}_{IR} \equiv 12G_{IR}(2-\gamma)\ell^2 \left(\frac{3}{2}C \right)^\delta (\alpha C)^{5+\frac{3}{4}\gamma}. \quad (\text{A.22})$$

From (A.20)-(A.21), requiring that the corrections δA and $\delta\Phi$ behave sub-exponentially in the IR imposes that γ obeys

$$\gamma \leq -\frac{4}{3}. \quad (\text{A.23})$$

For the strictly smaller case, the corrections decay exponentially in the $r \rightarrow \infty$ limit. The case $\gamma = -\frac{4}{3}$ requires a refined analysis depending on the exponent δ , that is carried out below.

A.3.1 Special case of $\gamma = -\frac{4}{3}$

We now consider setting $\gamma = -\frac{4}{3}$, for which the right hand side of (A.20)-(A.21) behaves polynomially. In this case the potential $G(\Phi)$ is given at leading order by

$$G(\Phi(r)) = G_{IR} (\alpha C)^{-1} \left(\frac{3}{2}C \right)^\delta e^{-2Cr^\alpha} r^{-(\alpha-1)+\alpha\delta} + \dots + \mathcal{O}(\beta), \quad (\text{A.24})$$

and (A.20)-(A.21) become

$$\frac{8}{3}\alpha^2 C^2 \delta\Phi'(r) + (1-\alpha)r^{-2\alpha} \left((1-\alpha)\delta A'(r) + r\delta A''(r) \right) = \tilde{G}_{IR} r^{2(\alpha-1)+\alpha\delta} + \dots + \mathcal{O}(\beta), \quad (\text{A.25})$$

$$\begin{aligned} \frac{4}{3}\tilde{V}_{IR} r^{\alpha-1} \delta\Phi(r) + \delta A'(r) \left(18\alpha C - (\alpha C)^{-1} \tilde{V}_{IR} \right) - 3r^{1-\alpha} \delta A''(r) = \\ -\frac{3}{2}(\alpha C)^{-1} \tilde{G}_{IR} r^{2(\alpha-1)+\alpha\delta} + \dots + \mathcal{O}(\beta). \end{aligned} \quad (\text{A.26})$$

These equations imply that δA and $\delta\Phi$ have a polynomial behavior at $r \rightarrow \infty$

$$\begin{aligned} \delta A(r) &= C_A r^\zeta + \dots + \mathcal{O}(\beta), \\ \delta\Phi(r) &= C_\Phi r^\eta + \dots + \mathcal{O}(\beta), \end{aligned} \quad (\text{A.27})$$

where the exponents are determined in terms of δ and α as

$$\zeta = \alpha(3+\delta) - 1 \quad , \quad \eta = \zeta - \alpha. \quad (\text{A.28})$$

Requiring the corrections to remain small compared with the leading r^α behavior therefore imposes a constraint on δ

$$\delta < \frac{1}{\alpha} - 2. \quad (\text{A.29})$$

B Calculation of shear viscosity

In this appendix, we reproduce for completeness the holographic shear viscosity calculation of [38], which includes four-derivative corrections. We start with the action

$$S = \frac{1}{16\pi G_5} \int d^5x \sqrt{-g} \left[R - \frac{4}{3} (\nabla\Phi)^2 + V(\Phi) + \ell^2 \beta G(\Phi) R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} \right], \quad (\text{B.1})$$

and introduce a shear fluctuation h_{xy} as

$$g_{\mu\nu} = g_{\mu\nu}^{(0)} + h_{\mu\nu}, \quad (\text{B.2})$$

where $g_{\mu\nu}^{(0)}$ is the background metric, of the form

$$ds^2 = -a^2(u) dt^2 + c^2(u) du^2 + b^2(u) d\mathbf{x}^2. \quad (\text{B.3})$$

The background also contains a non-trivial profile for the dilaton, which depends solely on the holographic coordinate u

$$\Phi(x) = \Phi(u). \quad (\text{B.4})$$

Expanding the metric perturbation in Fourier modes

$$h_x^y = \int d^4k \phi_k(u) e^{-i\omega t + ikz}, \quad (\text{B.5})$$

and substituting in the action, up to second order in ϕ_k we obtain

$$S_{eff} = \int \frac{d^4k}{(2\pi)^4} du \left[A(u) \phi_k'' \phi_{-k} + B(u) \phi_k' \phi_{-k}' + C(u) \phi_k' \phi_{-k} \right. \\ \left. + D(u) \phi_k \phi_{-k} + E(u) \phi_k'' \phi_{-k}'' + F(u) \phi_k'' \phi_{-k}' \right], \quad (\text{B.6})$$

where primes denote u derivatives. From the Kubo relation, one arrives at the shear viscosity in terms of the horizon data [86, 87]

$$\eta = \frac{1}{8\pi G_5} \left[\sqrt{-\frac{g_{uu}}{g_{tt}}} \left(A - B + \frac{F'}{2} \right) + \left(E \left(\sqrt{-\frac{g_{uu}}{g_{tt}}} \right)' \right) \right] \Big|_{u=u_h}, \quad (\text{B.7})$$

where the whole expression is evaluated at the horizon.

The various coefficients that appear in the effective action (B.6) are given by

$$\begin{aligned}
A(u) &= \frac{2a(u)b^2(u)(b(u)c^3(u) + 4\ell^2\beta G(\Phi)(b'(u)c'(u) - c(u)b''(u)))}{c(u)^4}, \\
B(u) &= \frac{b(u)}{2a(u)c^5(u)} [4\ell^2\beta b^2(u)c^2(u)G(\Phi)(a'(u))^2 + a^2(u)(16\ell^2\beta c^2(u)G(\Phi)(b'(u))^2 \\
&\quad + b^2(u)(3c^4(u) + 4\ell^2\beta G(\Phi)(c'(u))^2) - 8\ell^2\beta b(u)c(u)G(\Phi)(b'(u)c'(u) + c(u)b''(u))], \\
C(u) &= -\frac{1}{c^5(u)a(u)} [-a(u)b^3(u)c^4(u)a'(u) + 4\ell\beta b^2(u)c^2(u)G(\Phi)(a'(u))^2 b'(u) + \\
&\quad + a^2(u)(8\ell^2\beta c^2(u)G(\phi)(b')^3 + b^3(u)c^3(u)c'(u) - \\
&\quad - 8\ell^2\beta b(u)c(u)G(\Phi)b'(u)(b'(u)c'(u) - c(u)b''(u)) - \\
&\quad - 4b^2(u)(c^4(u)b'(u) - \ell^2\beta G(\Phi)b'(u)(c'(u))^2 + \ell^2\beta c(u)G(\Phi)c'(u)b''(u))], \\
D(u) &= \frac{1}{2c^5(u)a(u)b(u)} [2a(u)b^3(u)c^3(u) \times \\
&\quad \times (-b(u)a'(u)c'(u) + c(u)(3a'(u)b'(u) + b(u)a''(u))) - \\
&\quad - 4\ell^2\beta b^2(u)G(\Phi)(b^2(u)(a'(u))^2(c'(u))^2 - 2b^2(u)c(u)a'(u)c'(u)a''(u) + \\
&\quad + c^2(u)(3(a'(u))^2(b'(u))^2 + b^2(u)(a''(u))^2)) + \\
&\quad + a^2(u)(b^4(u)c^6(u)(2\partial_u\Phi\partial^u\Phi - V(\Phi)) - 12\ell^2\beta c^2(u)G(\Phi)(b'(u))^4 + \\
&\quad + 6b^3(u)c^3(u)(-b'(u)c'(u) + c(u)b''(u)) + \\
&\quad + 6b^2(u)(c^4(u)(b'(u))^2 - 2\ell^2\beta G(\Phi)(b'(u))^2(c'(u))^2 + \\
&\quad + 4\ell^2\beta c(u)G(\Phi)b'(u)c'(u)b''(u) - 2\ell^2\beta c^2(u)G(\Phi)(b''(u))^2)], \\
E(u) &= \frac{2\ell^2\beta a(u)b^3(u)G(\Phi)}{c^3(u)}, \\
F(u) &= \frac{4\ell^2\beta a(u)b^2(u)G(\Phi)(2c(u)b'(u) - b(u)c'(u))}{c^4(u)},
\end{aligned} \tag{B.8}$$

where we have already applied the relevant limit in the Kubo relation, $\omega \rightarrow 0$ and $k = 0$.

For η/s we also need to compute the entropy density from horizon data. In the presence of curvature corrections, the black brane entropy S is computed from Wald's formula as

$$S = -2\pi \oint_{\Sigma} d^3x \sqrt{-h} \frac{\delta \mathcal{L}}{\delta R_{\mu\nu\rho\sigma}} \epsilon_{\mu\nu} \epsilon_{\rho\sigma}, \tag{B.9}$$

where Σ refers to the horizon, with induced metric h . The tensor $\epsilon_{\mu\nu}$ is binormal to Σ and is normalized as $\epsilon_{\mu\nu}\epsilon^{\mu\nu} = -2$ and $\epsilon_{\mu\nu} = -\epsilon_{\nu\mu}$. Using that

$$\frac{\delta\mathcal{L}}{\delta R_{\mu\nu\rho\sigma}} = \frac{1}{16\pi G_5} [g^{\nu\sigma}g^{\rho\mu} + 2\ell^2\beta G(\Phi)R^{\mu\nu\rho\sigma}] , \quad (\text{B.10})$$

and expanding the background (B.3)-(B.4) near the horizon as

$$\begin{aligned} a(u)^2 &= a_0(1-u) + a_1(1-u)^2 + a_2(1-u)^3 + \dots \\ b(u)^2 &= b_0(1+(1-u) + \dots) \\ c(u)^2 &= c_0(1-u)^{-1} + c_1 + c_2(1-u) + \dots \\ \Phi(u) &= \Phi_h + \varphi_1(1-u) + \varphi_2(1-u)^2 + \dots \end{aligned} \quad (\text{B.11})$$

we find

$$S = \frac{b_0^{3/2}}{4G_5} V_3 \left[1 - \frac{3a_1c_0 - a_0c_1}{a_0c_0^2} \ell^2 \beta G(\Phi_h) \right] , \quad (\text{B.12})$$

with V_3 the three-dimensional spatial volume. The entropy density s is thus given by

$$s = \frac{b_0^{3/2}}{4G_5} \left[1 - \frac{3a_1c_0 - a_0c_1}{a_0c_0^2} \ell^2 \beta G(\Phi_h) \right] . \quad (\text{B.13})$$

Employing (B.11) in (B.7) and dividing by (B.13) we finally arrive at [38, 88]

$$\frac{\eta}{s} = \frac{1}{4\pi} \left[1 - \frac{\beta\ell^2}{c_0} \left(G(\Phi_h) + 2\sqrt{\frac{3}{2}}\varphi_1\partial_\Phi G(\Phi_h) \right) \right] . \quad (\text{B.14})$$

This result can be fully expressed in terms of the potentials V and G at the horizon. To see this, we first change coordinates to

$$ds^2 = f^{-1}(r) dr^2 + e^{2A(r)} (d\mathbf{x}^2 - f(r) dt^2) , \quad \Phi = \Phi(r) , \quad (\text{B.15})$$

with near-horizon expansions

$$\begin{aligned} A(r) &= A_h + A_1(r - r_h) + \dots \\ f(r) &= f_1(r - r_h) + \dots \\ \Phi(r) &= \Phi_h + \Phi_1(r - r_h) + \dots \end{aligned} \quad (\text{B.16})$$

The relation to (B.11) is given by

$$\varphi_1 = \frac{1}{2A_1}\Phi_1, \quad c_0 = \frac{1}{2f_1A_1} . \quad (\text{B.17})$$

Substituting back into (B.14) we find

$$\frac{\eta}{s} = \frac{1}{4\pi} \left[1 - 2f_1A_1\beta\ell^2 \left(G(\Phi_h) + \sqrt{\frac{2}{3}}\frac{\Phi_1}{A_1}\partial_\Phi G(\Phi_h) \right) \right] . \quad (\text{B.18})$$

The metric equations can then be conveniently expressed as [19]

$$\begin{aligned}\frac{dA}{dr} &= A_1 = -\frac{C}{\ell} \frac{S^{\frac{1}{3}}}{T} V(\Phi_h) \\ \frac{d\Phi}{dr} &= \Phi_1 = \frac{3C}{4\ell} \frac{S^{\frac{1}{3}}}{T} \sqrt{\frac{3}{2}} \partial_\Phi V(\Phi_h) \\ f_1 &= -M_p (4\pi)^{\frac{4}{3}} \frac{T}{S^{\frac{1}{3}}},\end{aligned}\tag{B.19}$$

where S, T are the entropy and temperature and C is a constant that is unimportant for our purposes. Substituting A_1, Φ_1 and f_1 back into our expression for η/s (B.18) we obtain (3.6) which was first derived in [38].

C Invariance of η/s under modification of curvature terms

In this work, we considered a single type of curvature squared correction to the bulk Lagrangian, corresponding to the Riemann squared $R_{\mu\nu\rho\sigma}^2$. The most general action at this order in curvature however contains more terms and takes the form

$$S = \frac{1}{16\pi G_5} \int d^5x \sqrt{-g} \left[R - \frac{4}{3} (\nabla\Phi)^2 + V(\Phi) + \ell^2 \beta (\lambda_1 R^2 + \lambda_2 R_{\mu\nu} R^{\mu\nu} + \lambda_3 R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma}) \right],\tag{C.1}$$

where all the λ_i may be taken to depend on the dilaton Φ . In this appendix, we show that the shear viscosity per entropy η/s is actually independent of λ_1 and λ_2 , which implies that (3.6) is actually the most general expression for η/s at quadratic order in curvature.

As a warm-up, we start by reviewing the result of [41], which applies to constant coefficients λ_i . Following the same procedure as in appendix B, we find

$$\eta = \frac{b_0^{3/2}}{16\pi G} \left[1 - \ell^2 \beta \left(\frac{6a_0c_0 + 3a_1c_0 - a_0c_1}{a_0c_0^2} \lambda_1 + \frac{3(a_0 + a_1)c_0 - a_0c_1}{2a_0c_0^2} \lambda_2 + \frac{a_0c_0 + 3a_1c_0 - a_0c_1}{a_0c_0^2} \lambda_3 \right) \right],\tag{C.2}$$

and

$$s = \frac{b_0^{3/2}}{4G} \left[1 - \ell^2 \beta \left(\frac{6a_0c_0 + 3a_1c_0 - a_0c_1}{a_0c_0^2} \lambda_1 + \frac{3(a_0 + a_1)c_0 - a_0c_1}{2a_0c_0^2} \lambda_2 + \frac{3a_1c_0 - a_0c_1}{a_0c_0^2} \lambda_3 \right) \right].\tag{C.3}$$

Up to order $\mathcal{O}(\beta)$, the ratio η/s is then given by¹²

$$\begin{aligned}\frac{\eta}{s} &= \frac{1}{4\pi} \left[1 - \frac{\beta\ell^2}{c_0} \lambda_3 \right] + \mathcal{O}(\beta^2) \\ &= \frac{1}{4\pi} [1 - 2f_1 A_1 \beta \ell^2 \lambda_3] + \mathcal{O}(\beta^2) \\ &= \frac{1}{4\pi} \left[1 - \frac{2}{3} \beta \ell^2 \lambda_3 V(\Phi_h) \right] + \mathcal{O}(\beta^2).\end{aligned}\tag{C.4}$$

¹²We again refer to appendix B for the definitions of the various coefficients in the near-horizon expansion of the background.

This shows that η/s is indeed independent of λ_1 and λ_2 .

This property of the shear viscosity per entropy can be explained by considering redefinitions of the metric of the form

$$g_{\mu\nu} \rightarrow g_{\mu\nu} + a g_{\mu\nu} R + b R_{\mu\nu}. \quad (\text{C.5})$$

with a and b two constants of order $\mathcal{O}(\beta)$. Under such a redefinition, the action (C.1) is brought to the following form [41]

$$S = \frac{1}{16\pi G_5} \int d^5x \sqrt{-g} \left[\frac{R}{\kappa} - \frac{4}{3} (\nabla\Phi)^2 + V(\Phi) + \ell^2 \beta \left(\tilde{\lambda}_1 R^2 + \tilde{\lambda}_2 R_{\mu\nu} R^{\mu\nu} + \lambda_3 R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} \right) \right], \quad (\text{C.6})$$

where

$$\frac{1}{\kappa} = 1 + \frac{5a+b}{2} \left(V(\Phi) - \frac{4}{3} (\nabla\Phi)^2 \right), \quad \tilde{\lambda}_1 = \lambda_1 + \frac{3a+b}{2}, \quad \tilde{\lambda}_2 = \lambda_2 - b. \quad (\text{C.7})$$

By choosing $b = \lambda_2$ and $a = \frac{-2\lambda_1 - \lambda_2}{3}$, we are therefore left with an action containing only the Riemann squared term at the quadratic level, albeit with a rescaled Ricci scalar. This rescaling will not produce a deviation from the universal result of $1/4\pi$ since η/s does not depend on κ . This is because in formula (B.7) the change of κ only amounts to a rescaling of the A, B, E and F coefficients

$$\begin{aligned} A &\rightarrow A\kappa^{-1} + \mathcal{O}(\beta^2) \quad , \quad B \rightarrow B\kappa^{-1} + \mathcal{O}(\beta^2), \\ E &\rightarrow E + \mathcal{O}(\beta^2) \quad , \quad F \rightarrow F + \mathcal{O}(\beta^2). \end{aligned} \quad (\text{C.8})$$

This implies a shift in the shear viscosity

$$\eta \rightarrow \eta + \frac{b_0^{3/2}}{16\pi G} (\kappa^{-1} - 1) + \mathcal{O}(\beta^2). \quad (\text{C.9})$$

The entropy density however receives the same shift

$$s \rightarrow s + \frac{b_0^{3/2}}{16\pi G} (\kappa^{-1} - 1) + \mathcal{O}(\beta^2), \quad (\text{C.10})$$

so that the ratio η/s is invariant under the rescaling by κ up to order $\mathcal{O}(\beta)$.

We now turn to the case where the higher curvature couplings are functions of the dilaton $\lambda_i(\Phi)$. The general formula for the shear viscosity and entropy density with the action (C.1) can still be expressed in terms of horizon data through (B.7). The difference with constant coefficients can therefore only arise if η and s depend on derivatives of $\lambda_i(\Phi)$ at the horizon. It can be checked however that derivatives of $\lambda_1(\Phi)$ and $\lambda_2(\Phi)$ appear in the expressions of neither η nor s , so that η/s should remain independent of λ_1 and λ_2 even with dilaton dependence. This indicates that the metric redefinition is still valid even if the coefficients are not constant.

D Calculation of the bulk viscosity

We detail here the derivation of the expression for the bulk viscosity (3.7), which generalizes the calculation of [36] to the case where the coefficients of the curvature corrections depend on the dilaton. We start from a general ansatz for the background metric and dilaton

$$ds^2 = -c_1(r)^2 dt^2 + c_2^2(r) d\mathbf{x}^2 + c_3^2(r) dr^2, \quad \Phi = \Phi(r). \quad (\text{D.1})$$

Substituting this ansatz in the equations of motion presented in Appendix A gives the equations obeyed by the ansatz fields

$$\begin{aligned} \Phi'' + \frac{c_1' \Phi'}{c_1} + \frac{3c_2' \Phi'}{c_2} - \frac{c_3' \Phi'}{c_3} + \frac{3}{8} c_3^2 \partial_\Phi V + \mathcal{O}(\beta) &= 0, \\ (\Phi')^2 + \frac{3}{4} c_3^2 V - \frac{9c_1' c_2'}{2c_1 c_2} - \frac{9(c_2')^2}{2c_2^2} + \mathcal{O}(\beta) &= 0, \\ c_2'' + \frac{(c_2')^2}{c_2} - \frac{c_2' c_3'}{c_3} + \frac{2}{9} c_2 (\Phi')^2 - \frac{1}{6} c_2 c_3^2 V + \mathcal{O}(\beta) &= 0, \\ c_1'' + \frac{2c_1' c_2'}{c_2} - \frac{c_1' c_3'}{c_3} + \frac{c_1 (c_2')^2}{c_2^2} - \frac{1}{6} c_1 c_3^2 V + \frac{2}{9} c_1 (\Phi')^2 + \mathcal{O}(\beta) &= 0, \end{aligned} \quad (\text{D.2})$$

where the terms of order $\mathcal{O}(\beta)$ were not written explicitly to avoid clutter.

To compute the relevant retarded Green's function (see section 3) we expand the metric and dilaton, considering SO(3) invariant perturbations¹³ in the axial gauge $h_{rr} = h_{rt} = 0$

$$\begin{aligned} \delta c_1(r, t) &= \frac{h_{tt}(r, t)}{2c_1(r)}, \\ \delta c_2(r, t) &= \frac{h_{11}(r, t)}{2c_2(r)}, \\ \delta c_3(r, t) &= 0, \\ \delta \Phi(r, t) &= \Psi(r, t). \end{aligned} \quad (\text{D.3})$$

It is convenient to introduce new variables H_{11}, H_{tt} such that

$$h_{tt}(r, t) \equiv e^{-i\omega t} c_1^2 H_{00}(r), \quad h_{11}(r, t) \equiv e^{-i\omega t} c_2^2 H_{11}(r), \quad (\text{D.4})$$

and the gauge invariant scalar fluctuation Z defined by

$$\Psi(r, t) \equiv e^{-i\omega t} \left(Z(r) + \frac{\Phi'(r) c_2(r)}{2c_2'(r)} H_{11}(r) \right). \quad (\text{D.5})$$

The linearized equations of motion give the equations obeyed by the fluctuations

$$H' + \frac{8c_2 c_3 \Phi'}{9c_2'} Z + \mathcal{O}(\beta) = 0, \quad H \equiv \frac{c_2 c_3}{c_2'} H_{11}, \quad (\text{D.6})$$

$$\begin{aligned} H'_{00} - \frac{8c_2 \Phi'}{9c_2'} Z' + \frac{c_2}{27c_1 (c_2')^2} (24c_1' c_2' \Phi' - c_1 c_3^2 (9c_2' \partial_\Phi V + 8c_2 V \Phi')) Z \\ + \left(\frac{\omega^2 c_3}{c_1^2} - \frac{1}{3} c_3 V + \frac{(c_1')^2}{c_3 c_1^2} + \frac{3c_1' c_2'}{c_1 c_2 c_3} \right) H + \mathcal{O}(\beta) = 0, \end{aligned} \quad (\text{D.7})$$

¹³Our definition for h_{tt} differs from [36] by a minus sign.

where we made use of the background equations of motion to simplify the above expressions. As before, we only provide the zeroth order results as the order $\mathcal{O}(\beta)$ expressions are too cumbersome to present here. Remarkably, the equation for Z decouples from the other fluctuations once the constraints above are used in tandem with the background equations. The resulting equation for Z takes the form

$$Z'' + \left(\frac{c_1'}{c_1} + \frac{3c_2'}{c_2} - \frac{c_3'}{c_3} \right) Z' + c_3^2 \left(\frac{\omega^2}{c_1^2} + \frac{4}{3}V - \frac{2c_2^2 c_3^2 V^2}{9(c_2')^2} + \frac{4c_2 V c_1'}{3c_1 c_2'} + \frac{2c_2 \partial_\Phi V \Phi'}{3c_2'} + \frac{3}{8}V'' \right) Z + \mathcal{O}(\beta) = 0, \quad (\text{D.8})$$

where the $\mathcal{O}(\beta)$ terms are also only functions of Z .

The bulk viscosity can be computed from the effective action at quadratic order in perturbations¹⁴

$$S_c = \frac{1}{16\pi G_5} \int dr \mathcal{L}_c \{ h_{11,\omega}, h_{00,\omega}, p_\omega, h_{11,\omega}^*, h_{00,\omega}^*, p_\omega^* \}, \quad (\text{D.9})$$

where we defined

$$h_{tt}(t, r) = e^{-i\omega t} h_{00,\omega}(r), \quad h_{11}(t, r) = e^{-i\omega t} h_{11,\omega}(r), \quad \psi(t, r) = e^{-i\omega t} p_\omega(r). \quad (\text{D.10})$$

Since the expression for the effective action is very lengthy and does not add to the understanding of the derivation, we do not present it here and refer to [36] for more details. In the following we suppress the ω indices and instead use “*” to refer to conjugate fields. The effective action is invariant under a U(1) symmetry which acts on the mode functions in (D.10) by a simultaneous phase rotation

$$h_{00,\omega} \rightarrow e^{i\alpha} h_{00,\omega}, \quad h_{11,\omega} \rightarrow e^{i\alpha} h_{11,\omega}, \quad p_\omega \rightarrow e^{i\alpha} p_\omega. \quad (\text{D.11})$$

This symmetry is associated with the conserved current

$$J_N = -i(J_\omega - J_{-\omega}) = 2\text{Im}J_\omega. \quad (\text{D.12})$$

with

$$\begin{aligned} J_\omega = & \frac{\partial \mathcal{L}}{\partial h_{11}^{*'}} \delta h_{11}^* + \frac{\partial \mathcal{L}}{\partial h_{11}^{*''}} \delta h_{11}^{*'} - \left(\frac{\partial \mathcal{L}}{\partial h_{11}^{*''}} \right)' \delta h_{11}^* \\ & + \frac{\partial \mathcal{L}}{\partial h_{00}^{*'}} \delta h_{00}^* + \frac{\partial \mathcal{L}}{\partial h_{00}^{*''}} \delta h_{00}^{*'} - \left(\frac{\partial \mathcal{L}}{\partial h_{00}^{*''}} \right)' \delta h_{00}^* \\ & + \frac{\partial \mathcal{L}}{\partial p^{*'}} \delta p^* + \frac{\partial \mathcal{L}}{\partial p^{*''}} \delta p^{*'} - \left(\frac{\partial \mathcal{L}}{\partial p^{*''}} \right)' \delta p^*. \end{aligned} \quad (\text{D.13})$$

After some lengthy but straightforward calculations, we find:

$$\begin{aligned} J_\omega = & \frac{1}{12c_1^4 c_2^2 c_3} \left[9c_2^5 h_{00}^* h_{00} c_1' - 9c_1^2 c_2^3 h_{00}^* h_{11} c_1' - 3c_1 c_2^5 (h_{00} h_{00}^{*'} + h_{00}^* h_{00}') + \right. \\ & c_1^5 (9c_2 h_{11} h_{11}^{*'} + 9h_{11}^* (h_{11} c_2' - c_2 h_{11}')) + 16c_2^5 p^* p' + 24c_2^3 p^* h_{11} \Phi') + \\ & \left. c_1^3 c_2^2 (-9h_{11}^* h_{00} c_2' + c_2 (9h_{00} h_{11}^{*'} + 9h_{11} h_{00}^{*'} + 8c_2^2 p^* h_{00} \Phi')) \right] + \mathcal{O}(\beta). \end{aligned} \quad (\text{D.14})$$

¹⁴The action (D.9) is actually the *complexified* version of the quadratic on-shell action; see [36] for details.

For our purpose, we wish to compute the current (D.14) at leading order in the hydrodynamic approximation. Assuming $\mathfrak{w} = \frac{\omega}{2\pi T}$ to be much smaller than 1, the fluctuations can be expanded up to order $\mathcal{O}(\mathfrak{w})$ as

$$Z = \left(\frac{c_1}{c_2}\right)^{-i\mathfrak{w}} (z_0 + i\mathfrak{w}z_1) + \mathcal{O}(\mathfrak{w}^2) \quad , \quad H = H_0 + i\mathfrak{w}H_1 + \mathcal{O}(\mathfrak{w}^2), \quad (\text{D.15})$$

$$H_{00} = H_{00,0} + i\mathfrak{w}H_{00,1} + \mathcal{O}(\mathfrak{w}^2). \quad (\text{D.16})$$

To extract the bulk viscosity using the Kubo relation, we need the imaginary part of the current at order $\mathcal{O}(\mathfrak{w})$. After making use of the equations of motion for H, H_{00} we find that the expression simplifies significantly and the conserved current only depends on z_0, z_1 . Its imaginary part takes the form

$$\text{Im}J_\omega = -\mathfrak{w}J_1, \quad (\text{D.17})$$

with

$$\begin{aligned} J_1 = & \frac{4c_2^2 (c_1 z_0^2 c'_2 - c_2 (z_0^2 c'_1 + c_1 z_1 z'_0 - c_1 z_0 z'_1))}{3c_3} \\ & + \frac{16\ell^2 \beta}{27c_1^2 c_3^3 c_2^2} (c_1 z_0^2 c'_2 - c_2 (z_0^2 c'_1 + c_1 z_1 z'_0 - c_1 z_0 z'_1)) \left[108c_2^2 G(\Phi) c_1'^2 c_2'^2 - \right. \\ & 6c_1 c_2 c'_1 c'_2 (4c_2^2 c_3^2 G(\Phi) V(\Phi) - 21G(\Phi) c_2'^2 + 6c_2 c'_2 \partial_\Phi G(\Phi) \Phi') + \\ & c_1^2 (c_2^4 c_3^4 G(\Phi) V(\Phi)^2 - 9c_2^2 c_3^2 G(\Phi) V(\Phi) c_2'^2 + 18G(\Phi) c_2'^4 - \\ & \left. 18c_2 c_2'^3 \partial_\Phi G(\Phi) \Phi' + c_2^3 c_3^2 c'_2 (5V(\Phi) \partial_\Phi G(\Phi) + 2G(\Phi) \partial_\Phi V(\Phi)) \Phi' \right]. \end{aligned} \quad (\text{D.18})$$

A key observation in this calculation is that since the current is radially conserved one may evaluate it at any point in the holographic direction and in particular evaluate it at the horizon. This simplifies the expression and gives ζ only in terms of horizon data. Expanding around the horizon as in (B.11):

$$\begin{aligned} c_1^2(u) &= a_0(1-u) + a_1(1-u)^2 + a_2(1-u)^3 + \dots \\ c_2^2(u) &= b_0(1 + (1-u) + \dots) \\ c_3^2(u) &= c_0(1-u)^{-1} + c_1 + c_2(1-u) + \dots \\ \Phi(u) &= \Phi_h + \varphi_1(1-u) + \varphi_2(1-u)^2 + \dots \end{aligned} \quad (\text{D.19})$$

Regularity at the horizon yields the following constraints:

$$c_0 = \frac{3}{2V_h}, \quad \varphi_1 = -\frac{9\partial_\Phi V_h}{16V_h}, \quad c_1 = \frac{15}{4V_h} + \frac{81(\partial_\Phi V_h)^2}{128V_h^3}, \quad a_1 = a_0 \left(\frac{1}{2} + \frac{9(\partial_\Phi V_h)^2}{64V_h^2} \right), \quad (\text{D.20})$$

Expanding the current at the horizon and making use of the constraints, we find

$$J_{1h} = \frac{2}{3} \sqrt{\frac{a_0}{c_0}} b_0^{3/2} z_0^2 \left(1 + \frac{\ell^2 \beta}{2} \partial_\Phi V \left(\partial_\Phi G + \frac{G \partial_\Phi V}{V} \right) \right), \quad (\text{D.21})$$

where the entire expression is evaluated at the horizon. The Kubo relation

$$\zeta = -\frac{4}{9} \lim_{\omega \rightarrow 0} \frac{1}{\omega} \text{Im} G_R = -\frac{1}{18\pi G_5} \lim_{\omega \rightarrow 0} \frac{1}{\omega} \text{Im} J_\omega \quad (\text{D.22})$$

then yields

$$\zeta = \frac{\mathfrak{w}}{27\pi G_5} \sqrt{\frac{a_0}{c_0}} b_0^{3/2} z_0^2 \left(1 + \frac{\ell^2 \beta}{2} \partial_\Phi V \left(\partial_\Phi G + \frac{G \partial_\Phi V}{V} \right) \right), \quad (\text{D.23})$$

which is valid for a normalization of the metric fluctuations such that

$$H_{11}(r=0) = 1. \quad (\text{D.24})$$

Using (B.13) results in the following for the bulk viscosity per entropy

$$s = \frac{b_0^{3/2}}{4G_5} \left[1 - \frac{3a_1 c_0 - a_0 c_1}{a_0 c_0^2} \ell^2 \beta G \right] = \frac{b_0^{3/2}}{4G_5} \left[1 + \frac{2\ell^2 \beta}{3} GV \right]. \quad (\text{D.25})$$

Taking the ratio between the two expressions and keeping terms up to order $\mathcal{O}(\beta)$ we finally arrive at

$$\frac{\zeta}{s} = \frac{8z_0^2}{27\pi} \left[1 + \frac{1}{2} \ell^2 \beta \left(-\frac{4}{3} GV + \frac{G(\partial_\Phi V)^2}{V} + \partial_\Phi G \partial_\Phi V \right) \right]. \quad (\text{D.26})$$

One can easily generalize this to (3.9) in the presence of all 4-derivative corrections.

E Numerical computation of the bulk viscosity

We outline here our procedure for the numerical evaluation of ζ/s and to fit the Bayesian data. We find it more convenient for the numerical calculation to change coordinates as follows

$$c_1 \rightarrow \sqrt{f(r)} e^{A(r)}, \quad c_2 \rightarrow e^{A(r)}, \quad c_3 \rightarrow \frac{e^{A(r)}}{\sqrt{f(r)}}. \quad (\text{E.1})$$

In addition, we change coordinates to use the scale factor A instead of the radial coordinate r , for which it is useful to introduce the variable

$$q(A) \equiv \frac{e^A(r)}{A'(r)}. \quad (\text{E.2})$$

The first step to compute the fluctuation is to solve the background equations at zeroth order in β , where the ansatz functions are expanded as

$$f = f_0 + \beta f_1 + \mathcal{O}(\beta^2), \quad q = q_0 + \beta q_1 + \mathcal{O}(\beta^2), \quad \lambda = \lambda_0 + \beta \lambda_1 + \mathcal{O}(\beta^2), \quad (\text{E.3})$$

with λ the exponential of the dilaton

$$\lambda \equiv e^\Phi, \quad (\text{E.4})$$

which is a better variable for the numerical analysis. At leading order, the three equations that we solve are

$$\begin{aligned}
\frac{f_0''(A)}{\sqrt{f_0(A)}} + \frac{q_0(A)^2 V(\lambda_0) f_0'(A)}{3f_0(A)^{3/2}} - \frac{f_0'(A)^2}{f_0(A)^{3/2}} &= 0, \\
q_0(A) \left(-\frac{f_0'(A)}{f_0(A)} - 4 \right) + \frac{q_0(A)^3 V(\lambda_0)}{3f_0(A)} + q_0' &= 0, \\
\frac{1}{2} \sqrt{\frac{9f_0'(A) + 36f_0(A) - 3q_0(A)^2 V(\lambda_0)}{f_0(A)}} + \frac{\lambda_0'(A)}{\lambda_0(A)} &= 0,
\end{aligned} \tag{E.5}$$

subject to the following horizon boundary conditions

$$\begin{aligned}
f_0 &= f_{00}(A - A_h) + f_{01}(A - A_h)^2 + \dots \\
\lambda_0 &= \lambda_{00} + \lambda_{01}(A - A_h) + \dots \\
q_0 &= q_{00} + q_{01}(A - A_h) + \dots
\end{aligned} \tag{E.6}$$

Regularity at the horizon in these coordinates implies:

$$q_{00}^2 = \frac{3f_{00}}{V_h}. \tag{E.7}$$

Notice that in this form the equations (E.5) are invariant under a resealing of the type

$$f_0 \rightarrow \alpha f_0, \quad q_0 \rightarrow \sqrt{\alpha} q_0. \tag{E.8}$$

We use this in the numerical solution to scale $f_0 \rightarrow 1$ at the UV boundary. Thus, the procedure we follow is:

- Solve the system of equations numerically assuming the horizon behavior (E.6) with $f_{00} = 1$. $f(A)$ will then go to some value f_{UV} at the boundary.
- Rescale $f_0 \rightarrow f_0/f_{UV}$, $q_0 \rightarrow q_0/\sqrt{f_{UV}}$ using (E.8), so that the rescaled f_0 goes to 1 in the UV.

Then the solution is correctly normalized, with a fixed definition of the boundary time.

The next step in the bulk viscosity calculation is to solve numerically the equation of motion obeyed by z_0 , which is obtained from (D.8) at leading order in the hydrodynamic expansion. z_0 follows an expansion in powers of β similar to (E.3)

$$z_0 = z_{00} + \beta z_{01} + \mathcal{O}(\beta^2), \tag{E.9}$$

where the leading order term z_{00} obeys

$$\begin{aligned}
z_{00}'' + z_{00}' \left(\frac{f_0'}{f_0} - \frac{q_0'}{q_0} + 4 \right) + \\
z_{00} \frac{q_0^2 (3f_0 (64V(\lambda_0) + (16\lambda_0' + 9\lambda_0) V'(\lambda_0) + 9\lambda_0^2 V''(\lambda_0) - 16V(\lambda_0) (q_0^2 V(\lambda_0) - 3f_0'))}{72f_0^2} = 0.
\end{aligned} \tag{E.10}$$

z_{00} is assumed to be regular at the horizon, where it admits an expansion of the form:

$$z_{00} = c_0 + c_1 (A - A_h). \quad (\text{E.11})$$

Substituting this into the z_{00} equation along with the horizon expansion of the background (E.6), regularity at the horizon is found to impose a relation between the coefficients c_0 and c_1

$$c_1 = c_0 \frac{9}{8} \left(\frac{V'(\Phi_h)^2}{V(\Phi_h)^2} - \frac{V''(\Phi_h)}{8V(\Phi_h)} \right). \quad (\text{E.12})$$

In addition, we must require that the dilaton perturbation Ψ is not sourced, since only the metric should be sourced to compute the bulk viscosity. From (D.5) and (D.24), this fixes the leading behavior of z_{00} near the boundary to be

$$z_{00}|_{UV} \underset{A \rightarrow \infty}{\sim} 1/2A. \quad (\text{E.13})$$

With the two boundary conditions (E.12)-(E.13), the fluctuation equations admit a unique solution that can be found numerically. Since we use a shooting method from the horizon to solve the equations, implementing the UV condition (E.13) is not as straightforward as (E.12). It can be done with the following procedure:

- We solve the equation with the previously computed background solution, imposing (E.12) at the horizon, with $c_0 = 1$.
- We rescale the solution appropriately such that z_{00} satisfies (E.13). Notice that (E.10) is invariant under the rescaling of z_{00} .

Then z_{00} matches the required boundary conditions. The bulk viscosity then follows from

$$\frac{\zeta}{s} = \frac{8z_{00} (A_h)^2}{27\pi}. \quad (\text{E.14})$$

Once we have the order $\mathcal{O}(\beta^0)$ solution, the extension of the above procedure to include the order $\mathcal{O}(\beta)$ corrections is straightforward. The method for fixing the normalization will then be applied to the background functions $\{f, q, \lambda\}$ and the perturbation z_0 including the $\mathcal{O}(\beta)$ corrections.

References

- [1] W. Busza, K. Rajagopal, and W. van der Schee, *Heavy Ion Collisions: The Big Picture, and the Big Questions*, *Ann. Rev. Nucl. Part. Sci.* **68** (2018) 339–376, [[arXiv:1802.04801](#)].
- [2] H. B. Meyer, *A Calculation of the shear viscosity in $SU(3)$ gluodynamics*, *Phys. Rev. D* **76** (2007) 101701, [[arXiv:0704.1801](#)].
- [3] H. B. Meyer, *A Calculation of the bulk viscosity in $SU(3)$ gluodynamics*, *Phys. Rev. Lett.* **100** (2008) 162001, [[arXiv:0710.3717](#)].
- [4] G. Nijs, W. van der Schee, U. Gürsoy, and R. Snellings, *Transverse Momentum Differential Global Analysis of Heavy-Ion Collisions*, *Phys. Rev. Lett.* **126** (2021), no. 20 202301, [[arXiv:2010.15130](#)].

- [5] G. Nijs, W. van der Schee, U. Gürsoy, and R. Snellings, *Bayesian analysis of heavy ion collisions with the heavy ion computational framework Trajectum*, Phys. Rev. C **103** (2021), no. 5 054909, [[arXiv:2010.15134](#)].
- [6] **JETSCAPE** Collaboration, D. Everett et al., *Multisystem Bayesian constraints on the transport coefficients of QCD matter*, Phys. Rev. C **103** (2021), no. 5 054904, [[arXiv:2011.01430](#)].
- [7] G. S. Denicol, S. Jeon, and C. Gale, *Transport Coefficients of Bulk Viscous Pressure in the 14-moment approximation*, Phys. Rev. C **90** (2014), no. 2 024912, [[arXiv:1403.0962](#)].
- [8] J. S. Moreland, J. E. Bernhard, and S. A. Bass, *Alternative ansatz to wounded nucleon and binary collision scaling in high-energy nuclear collisions*, Phys. Rev. C **92** (2015), no. 1 011901, [[arXiv:1412.4708](#)].
- [9] F. Cooper and G. Frye, *Comment on the Single Particle Distribution in the Hydrodynamic and Statistical Thermodynamic Models of Multiparticle Production*, Phys. Rev. D **10** (1974) 186.
- [10] J. M. Maldacena, *The Large N limit of superconformal field theories and supergravity*, Adv. Theor. Math. Phys. **2** (1998) 231–252, [[hep-th/9711200](#)].
- [11] S. S. Gubser, I. R. Klebanov, and A. M. Polyakov, *Gauge theory correlators from noncritical string theory*, Phys. Lett. B **428** (1998) 105–114, [[hep-th/9802109](#)].
- [12] E. Witten, *Anti-de Sitter space and holography*, Adv. Theor. Math. Phys. **2** (1998) 253–291, [[hep-th/9802150](#)].
- [13] G. Policastro, D. T. Son, and A. O. Starinets, *The Shear viscosity of strongly coupled N=4 supersymmetric Yang-Mills plasma*, Phys. Rev. Lett. **87** (2001) 081601, [[hep-th/0104066](#)].
- [14] U. Gursoy and E. Kiritsis, *Exploring improved holographic theories for QCD: Part I*, JHEP **02** (2008) 032, [[arXiv:0707.1324](#)].
- [15] U. Gursoy, E. Kiritsis, and F. Nitti, *Exploring improved holographic theories for QCD: Part II*, JHEP **02** (2008) 019, [[arXiv:0707.1349](#)].
- [16] M. Jarvinen and E. Kiritsis, *Holographic Models for QCD in the Veneziano Limit*, JHEP **03** (2012) 002, [[arXiv:1112.1261](#)].
- [17] G. Veneziano, *Some Aspects of a Unified Approach to Gauge, Dual and Gribov Theories*, Nucl. Phys. B **117** (1976) 519–545.
- [18] U. Gursoy, E. Kiritsis, L. Mazzanti, and F. Nitti, *Deconfinement and Gluon Plasma Dynamics in Improved Holographic QCD*, Phys. Rev. Lett. **101** (2008) 181601, [[arXiv:0804.0899](#)].
- [19] U. Gursoy, E. Kiritsis, L. Mazzanti, and F. Nitti, *Holography and Thermodynamics of 5D Dilaton-gravity*, JHEP **05** (2009) 033, [[arXiv:0812.0792](#)].
- [20] U. Gursoy, E. Kiritsis, L. Mazzanti, and F. Nitti, *Improved Holographic Yang-Mills at Finite Temperature: Comparison with Data*, Nucl. Phys. B **820** (2009) 148–177, [[arXiv:0903.2859](#)].
- [21] T. Alho, M. Järvinen, K. Kajantie, E. Kiritsis, and K. Tuominen, *On finite-temperature holographic QCD in the Veneziano limit*, JHEP **01** (2013) 093, [[arXiv:1210.4516](#)].
- [22] T. Alho, M. Järvinen, K. Kajantie, E. Kiritsis, C. Rosen, and K. Tuominen, *A holographic model for QCD in the Veneziano limit at finite temperature and density*, JHEP **04** (2014) 124, [[arXiv:1312.5199](#)]. [Erratum: JHEP **02**, 033 (2015)].

- [23] U. Gürsoy, I. Iatrakis, M. Järvinen, and G. Nijs, *Inverse Magnetic Catalysis from improved Holographic QCD in the Veneziano limit*, JHEP **03** (2017) 053, [[arXiv:1611.06339](#)].
- [24] U. Gürsoy, M. Jarvinen, and G. Nijs, *Holographic QCD in the Veneziano Limit at a Finite Magnetic Field and Chemical Potential*, Phys. Rev. Lett. **120** (2018), no. 24 242002, [[arXiv:1707.00872](#)].
- [25] A. Buchel and J. T. Liu, *Universality of the shear viscosity in supergravity*, Phys. Rev. Lett. **93** (2004) 090602, [[hep-th/0311175](#)].
- [26] S. S. Gubser, S. S. Pufu, and F. D. Rocha, *Bulk viscosity of strongly coupled plasmas with holographic duals*, JHEP **08** (2008) 085, [[arXiv:0806.0407](#)].
- [27] U. Gürsoy, E. Kiritsis, G. Michalogiorgakis, and F. Nitti, *Thermal Transport and Drag Force in Improved Holographic QCD*, JHEP **12** (2009) 056, [[arXiv:0906.1890](#)].
- [28] A. Buchel, *Bulk viscosity of gauge theory plasma at strong coupling*, Phys. Lett. B **663** (2008) 286–289, [[arXiv:0708.3459](#)].
- [29] A. Buchel and C. Pagnutti, *Bulk viscosity of $N=2^*$ plasma*, Nucl. Phys. B **816** (2009) 62–72, [[arXiv:0812.3623](#)].
- [30] C. Eling and Y. Oz, *A Novel Formula for Bulk Viscosity from the Null Horizon Focusing Equation*, JHEP **06** (2011) 007, [[arXiv:1103.1657](#)].
- [31] A. Buchel, U. Gürsoy, and E. Kiritsis, *Holographic bulk viscosity: GPR versus EO*, JHEP **09** (2011) 095, [[arXiv:1104.2058](#)].
- [32] A. Buchel, *Violation of the holographic bulk viscosity bound*, Phys. Rev. D **85** (2012) 066004, [[arXiv:1110.0063](#)].
- [33] A. Ballon-Bayona, L. A. H. Mamani, A. S. Miranda, and V. T. Zanchin, *Effective holographic models for QCD: Thermodynamics and viscosity coefficients*, Phys. Rev. D **104** (2021), no. 4 046013, [[arXiv:2103.14188](#)].
- [34] T. Demircik, D. Gallegos, U. Gürsoy, M. Järvinen, and R. Lier, *A Novel Method for Holographic Transport*, [arXiv:2311.00042](#).
- [35] T. Demircik, D. Gallegos, U. Gürsoy, M. Järvinen, and R. Lier, *Holographic transport in anisotropic plasmas*, Phys. Rev. D **110** (2024), no. 6 066007, [[arXiv:2402.12224](#)].
- [36] A. Buchel, S. Cremonini, and L. Early, *Holographic transport beyond the supergravity approximation*, JHEP **04** (2024) 032, [[arXiv:2312.05377](#)].
- [37] J. Cruz Rojas, T. Gorda, C. Hoyos, N. Jokela, M. Järvinen, A. Kurkela, R. Paatelainen, S. Säppi, and A. Vuorinen, *Estimate for the Bulk Viscosity of Strongly Coupled Quark Matter Using Perturbative QCD and Holography*, Phys. Rev. Lett. **133** (2024), no. 7 071901, [[arXiv:2402.00621](#)].
- [38] S. Cremonini, U. Gürsoy, and P. Szepietowski, *On the Temperature Dependence of the Shear Viscosity and Holography*, JHEP **08** (2012) 167, [[arXiv:1206.3581](#)].
- [39] P. Kovtun, D. T. Son, and A. O. Starinets, *Holography and hydrodynamics: Diffusion on stretched horizons*, JHEP **10** (2003) 064, [[hep-th/0309213](#)].
- [40] P. Kovtun, D. T. Son, and A. O. Starinets, *Viscosity in strongly interacting quantum field theories from black hole physics*, Phys. Rev. Lett. **94** (2005) 111601, [[hep-th/0405231](#)].
- [41] Y. Kats and P. Petrov, *Effect of curvature squared corrections in AdS on the viscosity of the dual gauge theory*, JHEP **01** (2009) 044, [[arXiv:0712.0743](#)].

- [42] M. Brigante, H. Liu, R. C. Myers, S. Shenker, and S. Yaida, *Viscosity Bound Violation in Higher Derivative Gravity*, Phys. Rev. D **77** (2008) 126006, [[arXiv:0712.0805](#)].
- [43] R. C. Myers, M. F. Paulos, and A. Sinha, *Quantum corrections to η/s* , Phys. Rev. D **79** (2009) 041901, [[arXiv:0806.2156](#)].
- [44] S. Cremonini, *The Shear Viscosity to Entropy Ratio: A Status Report*, Mod. Phys. Lett. B **25** (2011) 1867–1888, [[arXiv:1108.0677](#)].
- [45] A. Buchel and S. Cremonini, *Viscosity Bound and Causality in Superfluid Plasma*, JHEP **10** (2010) 026, [[arXiv:1007.2963](#)].
- [46] E. Kiritsis, *Dissecting the string theory dual of QCD*, Fortsch. Phys. **57** (2009) 396–417, [[arXiv:0901.1772](#)].
- [47] T. Alho, M. Jarvinen, K. Kajantie, E. Kiritsis, and K. Tuominen, *Quantum and stringy corrections to the equation of state of holographic QCD matter and the nature of the chiral transition*, Phys. Rev. D **91** (2015), no. 5 055017, [[arXiv:1501.06379](#)].
- [48] D. Gallegos, “unpublished.”.
- [49] D. T. Son and A. O. Starinets, *Minkowski space correlators in AdS / CFT correspondence: Recipe and applications*, JHEP **09** (2002) 042, [[hep-th/0205051](#)].
- [50] R. M. Wald, *Black hole entropy is the Noether charge*, Phys. Rev. D **48** (1993), no. 8 R3427–R3431, [[gr-qc/9307038](#)].
- [51] M. A. Shifman, A. I. Vainshtein, and V. I. Zakharov, *QCD and Resonance Physics. Theoretical Foundations*, Nucl. Phys. B **147** (1979) 385–447.
- [52] V. A. Novikov, M. A. Shifman, A. I. Vainshtein, M. B. Voloshin, and V. I. Zakharov, *Use and Misuse of QCD Sum Rules, Factorization and Related Topics*, Nucl. Phys. B **237** (1984) 525–552.
- [53] V. A. Novikov, M. A. Shifman, A. I. Vainshtein, and V. I. Zakharov, *OPERATOR EXPANSION IN QUANTUM CHROMODYNAMICS BEYOND PERTURBATION THEORY*, Nucl. Phys. B **174** (1980) 378–396.
- [54] M. Järvinen, *Holographic modeling of nuclear matter and neutron stars*, Eur. Phys. J. C **82** (2022), no. 4 282, [[arXiv:2110.08281](#)].
- [55] M. Panero, *Thermodynamics of the QCD plasma and the large- N limit*, Phys. Rev. Lett. **103** (2009) 232001, [[arXiv:0907.3719](#)].
- [56] T. Demircik, C. Ecker, and M. Järvinen, *Dense and Hot QCD at Strong Coupling*, Phys. Rev. X **12** (2022), no. 4 041012, [[arXiv:2112.12157](#)].
- [57] N. Jokela, M. Järvinen, and J. Remes, *Holographic QCD in the NICER era*, Phys. Rev. D **105** (2022), no. 8 086005, [[arXiv:2111.12101](#)].
- [58] C. Hoyos, N. Jokela, M. Järvinen, J. G. Subils, J. Tarrío, and A. Vuorinen, *Holographic approach to transport in dense QCD matter*, Phys. Rev. D **105** (2022), no. 6 066014, [[arXiv:2109.12122](#)].
- [59] C. Ecker, M. Järvinen, G. Nijs, and W. van der Schee, *Gravitational waves from holographic neutron star mergers*, Phys. Rev. D **101** (2020), no. 10 103006, [[arXiv:1908.03213](#)].
- [60] N. Jokela, M. Järvinen, and J. Remes, *Holographic QCD in the Veneziano limit and neutron stars*, JHEP **03** (2019) 041, [[arXiv:1809.07770](#)].

- [61] D. Arean, I. Iatrakis, M. Järvinen, and E. Kiritsis, *V-QCD: Spectra, the dilaton and the S-parameter*, Phys. Lett. B **720** (2013) 219–223, [[arXiv:1211.6125](#)].
- [62] A. Buchel and R. C. Myers, *Causality of Holographic Hydrodynamics*, JHEP **08** (2009) 016, [[arXiv:0906.2922](#)].
- [63] M. Brigante, H. Liu, R. C. Myers, S. Shenker, and S. Yaida, *The Viscosity Bound and Causality Violation*, Phys. Rev. Lett. **100** (2008) 191601, [[arXiv:0802.3318](#)].
- [64] O. Hohm, C. Hull, and B. Zwiebach, *Background independent action for double field theory*, JHEP **07** (2010) 016, [[arXiv:1003.5027](#)].
- [65] O. Hohm, C. Hull, and B. Zwiebach, *Generalized metric formulation of double field theory*, JHEP **08** (2010) 008, [[arXiv:1006.4823](#)].
- [66] C. Vafa, *The String landscape and the swampland*, [hep-th/0509212](#).
- [67] H. Ooguri and C. Vafa, *On the Geometry of the String Landscape and the Swampland*, Nucl. Phys. B **766** (2007) 21–33, [[hep-th/0605264](#)].
- [68] E. Palti, *The Swampland: Introduction and Review*, Fortsch. Phys. **67** (2019), no. 6 1900037, [[arXiv:1903.06239](#)].
- [69] D. E. Kharzeev, L. D. McLerran, and H. J. Warringa, *The Effects of topological charge change in heavy ion collisions: 'Event by event P and CP violation'*, Nucl. Phys. A **803** (2008) 227–253, [[arXiv:0711.0950](#)].
- [70] V. Skokov, A. Y. Illarionov, and V. Toneev, *Estimate of the magnetic field strength in heavy-ion collisions*, Int. J. Mod. Phys. A **24** (2009) 5925–5932, [[arXiv:0907.1396](#)].
- [71] K. Tuchin, *Particle production in strong electromagnetic fields in relativistic heavy-ion collisions*, Adv. High Energy Phys. **2013** (2013) 490495, [[arXiv:1301.0099](#)].
- [72] V. Voronyuk, V. D. Toneev, W. Cassing, E. L. Bratkovskaya, V. P. Konchakovski, and S. A. Voloshin, *(Electro-)Magnetic field evolution in relativistic heavy-ion collisions*, Phys. Rev. C **83** (2011) 054911, [[arXiv:1103.4239](#)].
- [73] W.-T. Deng and X.-G. Huang, *Event-by-event generation of electromagnetic fields in heavy-ion collisions*, Phys. Rev. C **85** (2012) 044907, [[arXiv:1201.5108](#)].
- [74] K. Tuchin, *Time and space dependence of the electromagnetic field in relativistic heavy-ion collisions*, Phys. Rev. C **88** (2013), no. 2 024911, [[arXiv:1305.5806](#)].
- [75] L. McLerran and V. Skokov, *Comments About the Electromagnetic Field in Heavy-Ion Collisions*, Nucl. Phys. A **929** (2014) 184–190, [[arXiv:1305.0774](#)].
- [76] U. Gürsoy, D. Kharzeev, and K. Rajagopal, *Magnetohydrodynamics, charged currents and directed flow in heavy ion collisions*, Phys. Rev. C **89** (2014), no. 5 054905, [[arXiv:1401.3805](#)].
- [77] U. Gürsoy, D. Kharzeev, E. Marcus, K. Rajagopal, and C. Shen, *Charge-dependent Flow Induced by Magnetic and Electric Fields in Heavy Ion Collisions*, Phys. Rev. C **98** (2018), no. 5 055201, [[arXiv:1806.05288](#)].
- [78] D. Kharzeev, K. Landsteiner, A. Schmitt, and H.-U. Yee, eds., Strongly Interacting Matter in Magnetic Fields, vol. 871. 2013.
- [79] N. Iqbal and H. Liu, *Universality of the hydrodynamic limit in AdS/CFT and the membrane paradigm*, Phys. Rev. D **79** (2009) 025023, [[arXiv:0809.3808](#)].

- [80] E. D'Hoker and P. Kraus, *Magnetic Brane Solutions in AdS*, JHEP **10** (2009) 088, [[arXiv:0908.3875](#)].
- [81] A. Ballon-Bayona, *Holographic deconfinement transition in the presence of a magnetic field*, JHEP **11** (2013) 168, [[arXiv:1307.6498](#)].
- [82] R. Rougemont, R. Critelli, and J. Noronha, *Holographic calculation of the QCD crossover temperature in a magnetic field*, Phys. Rev. D **93** (2016), no. 4 045013, [[arXiv:1505.07894](#)].
- [83] S. I. Finazzo, R. Critelli, R. Rougemont, and J. Noronha, *Momentum transport in strongly coupled anisotropic plasmas in the presence of strong magnetic fields*, Phys. Rev. D **94** (2016), no. 5 054020, [[arXiv:1605.06061](#)]. [Erratum: Phys.Rev.D 96, 019903 (2017)].
- [84] U. Gürsoy, M. Järvinen, G. Nijs, and J. F. Pedraza, *On the interplay between magnetic field and anisotropy in holographic QCD*, JHEP **03** (2021) 180, [[arXiv:2011.09474](#)].
- [85] U. Gürsoy, *Holographic QCD and magnetic fields*, Eur. Phys. J. A **57** (2021), no. 7 247, [[arXiv:2104.02839](#)].
- [86] A. Buchel, J. T. Liu, and A. O. Starinets, *Coupling constant dependence of the shear viscosity in $N=4$ supersymmetric Yang-Mills theory*, Nucl. Phys. B **707** (2005) 56–68, [[hep-th/0406264](#)].
- [87] R. C. Myers, M. F. Paulos, and A. Sinha, *Holographic Hydrodynamics with a Chemical Potential*, JHEP **06** (2009) 006, [[arXiv:0903.2834](#)].
- [88] S. Cremonini and P. Szepietowski, *Generating Temperature Flow for eta/s with Higher Derivatives: From Lifshitz to AdS*, JHEP **02** (2012) 038, [[arXiv:1111.5623](#)].