

ON A GEOMETRIC EXTREMUM PROBLEM FOR CONVEX CONES

OLEG MUSHKAROV AND NIKOLAI NIKOLOV

ABSTRACT. We discuss the optimization problem for minimizing the $(n - 1)$ -volume of the intersection of a convex cone in \mathbb{R}^n with a hyperplane through a given point.

1. INTRODUCTION

Let K be a closed convex cone in \mathbb{R}^n ($n \geq 2$) with vertex at the origin O and non-empty interior K° . Assume that K is pointed, that is $K \cap (-K) = \{O\}$. In [4], the following optimization problem is considered:

(*) *Minimize the $(n - 1)$ -volume of the intersection of K with a hyperplane $\mathcal{H} \not\ni O$ through a given point A .*

Write \mathcal{H} as $\mathcal{H}(b) = \{x : (b, x) = 1\}$ ($b \neq 0$) and denote by $V(b)$ the $(n - 1)$ -volume of $K(b) = K \cap \mathcal{H}(b)$. Then (*) means to minimize $V(b)$ under the constraint $(b, a) = 1$ ($a = \overrightarrow{OA}$). Note that $V(b)$ is finite exactly when $K(b)$ is compact, i.e. b is an inner point of the dual cone K^* of K . Recall that $K^* = \{y : (y, x) \geq 0, \forall x \in K\}$. Since $K^* \cap (-K) = \{O\}$, we may assume $A \notin (-K)$ and we may consider only vectors $b \in K^+ := (K^*)^\circ = \{y : (y, x) > 0, \forall x \in K\}$. Call the respective hyperplanes admissible.

The main Result 3.1 in [4] states that b is stationary for V under the constraint $(b, a) = 1$, i.e. $D_b(V) = \lambda a$ (λ is the Lagrange multiplier), if and only if

$$(1) \quad \overrightarrow{AH} = n\overrightarrow{AG},$$

where H is the orthogonal projection of O on $\mathcal{H}(b)$ and G is the centroid of $K(b)$. We call $\mathcal{H}(b)$ a stationary hyperplane for A .

This result extends the classical case $n = 2$, where the respective stationary line is called a Philon line (for more details, see e.g. [2] and [4]).

Note that (1) implies that any admissible hyperplane \mathcal{H} is stationary exactly for one point from \mathcal{H} . On the other hand, denote by S the set of points $A \in \mathbb{R}^n$ for which there is a stationary hyperplane through A . Since (*) has a solution for any $A \in K^\circ$, we have that $K^\circ \subset S \subset \mathbb{R}^n \setminus (-K)$. In [4], a general question is posed, namely, to describe the set S and to characterize the respective stationary hyperplanes. In Proposition 3 we recall the answer to this question for $n = 2$ and note that the situation changes when $n \geq 3$.

The main purpose of this note is to show that if K is the non-negative orthant $(\mathbb{R}_0^+)^n$, then $S = K^\circ$ and any point from S lies in a unique stationary hyperplane (which solves (*)). Moreover, we describe this hyperplane in terms of the unique real root of an irrational equation (Proposition 4). Applying this result (for $n = 2$), we also find the length of the shortest segment cut from this orthant K by a line through a given point $A \in K^\circ$ (Proposition 6).

2. GEOMETRIC CHARACTERIZATION OF STATIONARY HYPERPLANES

In this section we give a geometric characterization of the stationary hyperplanes when K is a closed convex cone in \mathbb{R}^n spanned by n linearly independent vectors $\vec{e}_1, \vec{e}_2, \dots, \vec{e}_n$. Call such a cone hyperangle. Consider a hyperplane $\mathcal{H} = \mathcal{H}(b)$ such that $b \in K^+$. This

2020 *Mathematics Subject Classification.* 49K21, 51M16, 52A38.

Key words and phrases. Philon line, convex cone, simplex.

The second named author was partially supported by the Bulgarian National Science Fund, Ministry of Education and Science of Bulgaria under contract KP-06-N82/6.

means that \mathcal{H} intersects any ray $O\vec{e}_i$, say at A_i . Let C be the circumcenter of the $(n-1)$ -simplex $\mathcal{A} = A_1A_2 \dots A_n$. For $n \geq 3$, denote by M the Monge point of \mathcal{A} , i.e. the common point of the hyperplanes through the centroids of the $(n-3)$ -dimensional faces of \mathcal{A} which are perpendicular to the opposite edges of \mathcal{A} (see e.g. [3]). For $A \in \mathcal{H}$ define A' by

$$(2) \quad \overrightarrow{OA'} = \frac{n-1}{2} \overrightarrow{OA}.$$

Proposition 1. *For $n \geq 3$, any two of the following three conditions imply the third one:*

- (i) \mathcal{H} is a stationary hyperplane for A ;
- (ii) $A'A_1 = A'A_2 = \dots = A'A_n$;
- (iii) $H = M$.

Proof. Let A'' be the orthogonal projection of A' on \mathcal{H} . Then, by (2),

$$(3) \quad \overrightarrow{A''A} = \frac{n-3}{2} \overrightarrow{AH}.$$

On the other hand, it is a classical result (see e.g. [3]) that

$$(4) \quad \overrightarrow{CG} = \frac{n-2}{n} \overrightarrow{CM}.$$

The proposition follows by (1), (3), (4), and the fact that (ii) is equivalent to $A'' = C$. \square

Remarks. (i) If $n = 2$, (iii) is missing, i.e. (i) \Leftrightarrow (ii); in other words, l is a Philon line for A exactly when G is the midpoint of $[AH]$.

(ii) Recall that a k -simplex Δ ($k \geq 2$) is called orthocentric if the altitudes from the vertices to the opposite facets have a common point, called orthocenter. It turns out that this is equivalent to orthogonality of any edge to the respective $(k-2)$ -dimensional face. Note that the orthocenter of Δ coincides with its Monge point (see e.g. [3]).

Assume now that $OA_1A_2 \dots A_n$ is an orthocentric n -simplex ($n \geq 3$). Then $A_1A_2 \dots A_n$ is an orthocentric $(n-1)$ -simplex and $H = M$. Hence in this case (i) \Leftrightarrow (ii), too.

Proposition 2. *Let K be a hyperangle.*

- (i) *If $K^* \subset K$, then $S = K^\circ$.*
- (ii) *If an edge of K forms acute angles with the others, then there exists a point $A \in \partial K$ with a unique hyperplane solving problem (*); in particular, $K^\circ \subsetneq S$.*

Proof. Let \mathcal{H} be an admissible hyperplane and H be the orthogonal projection of O on \mathcal{H} . For a point $X \in H$, denote by (x_1, x_2, \dots, x_n) its barycentric coordinates w.r.t. to the $(n-1)$ -simplex \mathcal{A} defined at the beginning of this section.

(i) The condition $K^* \subset K$ is equivalent to $K^+ \subset K^\circ$ which means that $H \in \mathcal{A}^\circ$ for any admissible hyperplane \mathcal{H} , i.e. $h_i > 0$ for any $1 \leq i \leq n$.

Let now \mathcal{H} be a stationary hyperplane for A . Then (1) reads as

$$(5) \quad h_i + (n-1)a_i = 1, \quad 1 \leq i \leq n.$$

Hence $a_i < 1/(n-1)$ and then

$$a_k = 1 - \sum_{i \neq k}^n a_i > 1 - (n-1) \cdot \frac{1}{n-1} = 0.$$

This shows $A \in K^\circ$ and thus $S \subset K^\circ$.

(ii) We may assume $(e_1, e_i) > 0$ for any $2 \leq i \leq n$. Denote by $\mathcal{A}_1 = A_1A_2 \dots A_n$ the respective $(n-1)$ -simplex to the hyperplane $\mathcal{H}_1 = \mathcal{H}(e_1)$. Consider the point $A \in \partial K$ with barycentric coordinates $(0, \frac{1}{n-1}, \dots, \frac{1}{n-1})$ w.r.t. \mathcal{A}_1 .

Let \mathcal{H} be an admissible hyperplane for A and $\mathcal{A}' = A'_1A'_2 \dots A'_n$ be the orthogonal projection of the respective $(n-1)$ -simplex \mathcal{A} on H_1 . Then $A'_1 = A_1$, $A'_2 \in A_1A_2^\rightarrow, \dots, A'_n \in A_1A_n^\rightarrow$. Denote by K_1 the cone with edges these $n-1$ rays.

Recall now (see e.g. [4, Section (iv)]) that if the $(n-1)$ -volume of the cone segment of K_1 cut off by an $(n-2)$ -plane through A is minimal, then A is the centroid of the respective $(n-2)$ -simplex. Note that $A_2 \dots A_n$ is the unique $(n-2)$ -simplex with this property. So

$$V_{n-1}(\mathcal{A}) \geq V_{n-1}(\mathcal{A}') \geq V_{n-1}(\mathcal{A}_1)$$

and (ii) easily follows. \square

It is easy to see that $K \subset K^*$ (resp. $K \subset K^+$) if and only if the angles between the edges of K are non-obtuse (resp. acute). Hence the condition $K \subset K^+$ implies that in (ii) of Proposition 2. On the other hand, since $(K^*)^* = K$, we have that $K^* \subset K$ if and only if the angles between the inner normals to the facets of K are non-obtuse. We also claim that the non-negative orthant is self-dual, i.e. $K = K^*$.

Now we will describe the set S if $n = 2$. Set $\alpha = \angle(\vec{e}_1, \vec{e}_2)$. Then $K^* \subset K$ if $\alpha \geq 90^\circ$ and $K \subset K^+$ if $\alpha < 90^\circ$. So Proposition 1 implies that $S = K^\circ$ in the first case and $K^\circ \subsetneq S$ in the second one. In fact, we know much more (see e.g. [1] and [2]).

If $\alpha < 90^\circ$, set

$$\theta = \arctan \left(\frac{1 + \sin^2 \alpha / 2}{1 + \cos^2 \alpha / 2} \right)^{3/2}$$

and consider the closed cone T with vertex O , the same bisector as K , and aperture 2θ . Then $K \subsetneq T$.

Proposition 3. (i) If $\alpha \geq 90^\circ$, then $S = K^\circ$ and for any point from K° there exists one stationary line (global minima point).

(ii) If $\alpha < 90^\circ$, then $S = T \setminus \{O\}$. Moreover,

- for $A \in K$ there exists one stationary line (global minima point);
- for $A \in T^\circ \setminus K$ there exist two stationary lines (local minima and maxima points);
- for $A \in \partial T \setminus \{O\}$ there exists one stationary line (saddle point).

It is noted in [4, Section 4 (ii)] that the situation changes when $n \geq 3$. For example, if K is the trihedral angle spanned by the vectors $\vec{e}_1 = (1, -1, -0.2)$, $\vec{e}_2 = (1, 1, -0.2)$, and $\vec{e}_3 = (1, 0, 0.01)$, then three stationary planes exist for the point $A = (1, 0, 0) \in K^\circ$ (two local minima points and one saddle point).

3. THE NON-NEGATIVE ORTHANT

Let K be the non-negative orthant in \mathbb{R}^n , i.e. $K = (\mathbb{R}_0^+)^n$. We already know that $S = K^\circ$. For $A = (a_1, a_2, \dots, a_n) \in K^\circ$, set $b_i = \frac{n-1}{2}a_i$, $1 \leq i \leq n$, and

$$f_A(x) = \sum_{i=1}^n \frac{b_i}{b_i + \sqrt{b_i^2 + x}} - \frac{n-1}{2}.$$

This function is continuous and strictly decreasing on the interval $[-\min b_i^2, +\infty)$, $f(0) = \frac{1}{2}$ and $f(+\infty) = \frac{1-n}{2}$. Hence the equation $f_A(x) = 0$ has a unique real root λ_A , and $\lambda_A > 0$. Set $c_i = b_i + \sqrt{b_i^2 + \lambda_A}$, $1 \leq i \leq n$.

Proposition 4. For any $A \in K^\circ$ there exists a unique stationary hyperplane. For the respective $(n-1)$ -simplex \mathcal{A} we have that $\overrightarrow{OA_i} = c_i \vec{e}_i$, $1 \leq i \leq n$, and

$$V_{n-1}(\mathcal{A}) = \frac{\prod_{i=1}^n c_i}{(n-1)! \sqrt{\lambda_A}}.$$

Proof. Let \mathcal{H} be an admissible hyperplane. Then, by Remark (ii) after Proposition 1, \mathcal{H} is a stationary hyperplane for A if and only if $A'A_1 = A'A_2 = \dots = A'A_n$, where A' is given by (2). Let $\overrightarrow{OA_i} = x_i \vec{e}_i$, $1 \leq i \leq n$. Then we get that

$$(6) \quad x_i^2 - (n-1)x_i a_i = A'A_i^2 - A'O^2 =: \lambda, \quad 1 \leq i \leq n,$$

i.e.

$$1 - (n-1) \frac{a_i}{x_i} = \frac{\lambda}{x_i^2}.$$

Summing up these equalities and using that

$$(7) \quad \sum_{i=1}^n \frac{a_i}{x_i} = 1$$

(since $A, A_1, A_2, \dots, A_n \in \mathcal{H}$), we get that $\lambda \sum_{i=1}^n x_i^{-2} = 1$, i.e. $\lambda = d^2$, where d is the distance from O to \mathcal{H} . By (6), $x_i = b_i + \sqrt{b_i^2 + \lambda}$ and then (7) implies that $f_A(\lambda) = 0$. So $\lambda = \lambda_A$, $x_i = c_i$, and the formula for $V_{n-1}(S)$ follows. \square

Let us note that the circumcenter C of the simplex \mathcal{A} coincides with A if only if $n = 3$.

Applying Proposition 4 for $n = 2$ gives the following known fact.

Corollary 5. *The length of a shortest segment cut from the first quadrant in \mathbb{R}^2 by a line through its interior point $A(a_1, a_2)$ is equal to $(a_1^{2/3} + a_2^{2/3})^{3/2}$ and such a segment is unique.*

Proof. Let $\lambda = (a_1 a_2)^{2/3} (a_1^{2/3} + a_2^{2/3})$. Since

$$b_1^2 + \lambda = a_1^{2/3} (a_1^{2/3}/2 + a_2^{2/3})^2, \quad b_2^2 + \lambda = a_2^{2/3} (a_2^{2/3}/2 + a_1^{2/3})^2,$$

it easily follows that $\lambda = \lambda_A$. Then $c_1 = a_1 + a_1^{1/3} a_2^{2/3}$, $c_2 = a_2 + a_2^{1/3} a_1^{2/3}$, and hence $V_1 = (c_1^2 + c_2^2)^{1/2} = (a_1^{2/3} + a_2^{2/3})^{3/2}$. \square

Using the above corollary, we will generalize it in higher dimensions.

Proposition 6. *Let $A = A(a_1, a_2, \dots, a_n)$ with $0 < a_1 \leq a_2 \leq \dots \leq a_n$. Then the length of a shortest segment cut from the non-negative orthant K in \mathbb{R}^n by a line through A is equal to $(a_1^{2/3} + a_2^{2/3})^{3/2}$.*

Proof. We may assume that $n \geq 3$. We will first prove that the length of a shortest segment cut from the set $Q = (\mathbb{R}_0^+)^2 \times \mathbb{R}^{n-2} \supset K$ by a line through A is given by the above formula, such a segment is unique and it lies in K . To do this, set $L_1 = \mathbb{R}_0^+ \times \mathbb{R}^{n-1}$ and $L_2 = \mathbb{R} \times \mathbb{R}_0^+ \times \mathbb{R}^{n-2}$. Let a line through A cuts from Q a segment $[A_1 A_2]$, where $A_1 \in L_1$ and $A_2 \in L_2$. Denote by Π the plane through A which is orthogonal to $\{0\} \times \{0\} \times \mathbb{R}^{n-2}$. Let A'_1 and A'_2 be the orthogonal projections of A_1 and A_2 on Π . Then $A'_1 \in L_1$, $A'_2 \in L_2$, $O \in [A'_1 A'_2]$, and $|A'_1 A'_2| \leq |A_1 A_2|$ with equality if and only if $A'_1 = A_1$ and $A'_2 = A_2$. Hence the problem is reduced to that for the right angle $Ol_1 l_2 \subset K$, where $l_1 = L_1 \cap \Pi$ and $l_2 = L_2 \cap \Pi$. It remains to apply Corollary 5.

Let now $[BC]$ be a segment cut from K by a line through A . Setting $L_k = \mathbb{R}^{k-1} \times \mathbb{R}_0^+ \times \mathbb{R}_0^{n-k}$, $1 \leq k \leq n$, we may choose $i \neq j$ such that $B \in L_i$ and $C \in L_j$. Then the same reasoning as above shows that

$$|BC| \geq (a_i^{1/3} + a_j^{1/3})^{3/2} \geq (a_1^{1/3} + a_2^{1/3})^{3/2}$$

and the proposition is proved. \square

This proof shows that for $n \geq 3$ the number of lines through A which cut from K segments of minimal length is equal to $k-1$ if $a_1 < a_2 = a_k < a_{k+1}$, and to $\binom{k}{2}$ if $a_1 = a_2 = a_k < a_{k+1}$. In particular, such a line is unique if and only if $a_2 < a_3$.

On the other hand, it is clear that among all lines $l \not\ni O$ through $A \notin K^\circ$ there is no one which cuts from K a segment of minimal length.

REFERENCES

- [1] O. Bottema, *Verscheidenheden LXXIII. De uiterste waarden van een lijnstuk*, Euclides 44 (1968-1969), 20-26.
- [2] H. S. M. Coxeter, J. van de Craats, *Philon lines in non-Euclidean planes*, J. Geom. 48 (1993), 26-55.
- [3] A. Edmonds, M. Hajja, H. Martini, *Orthocentric simplices and their centers*, Results Math. 47 (2005), 266-295.
- [4] W. W. E. Wetterling, *Philon's line generalized: an optimization problem from geometry*. J. Optim. Theory Appl. 90 (1996), 517-521.

INSTITUTE OF MATHEMATICS AND INFORMATICS, BULGARIAN ACADEMY OF SCIENCES, ACAD. G. BONCHEV 8, 1113 SOFIA, BULGARIA
Email address: muskarov@math.bas.bg

INSTITUTE OF MATHEMATICS AND INFORMATICS, BULGARIAN ACADEMY OF SCIENCES, ACAD. G. BONCHEV 8, 1113 SOFIA, BULGARIA
FACULTY OF INFORMATION SCIENCES, STATE UNIVERSITY OF LIBRARY STUDIES AND INFORMATION TECHNOLOGIES, SHIPCHENSKI PROHOD 69A, 1574 SOFIA, BULGARIA
Email address: nik@math.bas.bg