

Color-magnetic correlations in SU(N) lattice QCD

Hideo Suganuma,* Atsuya Tokutake and Kei Tohme

*Department of Physics, Graduate School of Science, Kyoto University,
Kitashirakawa-oiwake, Sakyo, Kyoto 606-8502, Japan*

E-mail: suganuma@scphys.kyoto-u.ac.jp,

tokutake@gauge.scphys.kyoto-u.ac.jp, tohme@ruby.scphys.kyoto-u.ac.jp

Motivated by color-magnetic instabilities in QCD, we investigate field-strength correlations in both SU(2) and SU(3) lattice QCD. In the Euclidean Landau gauge, we numerically calculate the perpendicular-type color-magnetic correlation, $C_{\perp}(r) \equiv \langle H_z^a(s) H_z^a(s + r \hat{\perp}) \rangle$ with $\perp \equiv x, y$, and the parallel-type one, $C_{\parallel}(r) \equiv \langle H_z^a(s) H_z^a(s + r \hat{\parallel}) \rangle$ with $\parallel \equiv z, t$. In the Landau gauge, all two-point field-strength correlations $\langle G_{\mu\nu}^a(s) G_{\alpha\beta}^b(s') \rangle$ are described by these two quantities, due to the Lorentz and global SU(N_c) color symmetries. Curiously, the perpendicular-type color-magnetic correlation $C_{\perp}(r)$ is found to be always negative for arbitrary r , except for the same-point correlation. The parallel-type color-magnetic correlation $C_{\parallel}(r)$ is always positive. In the infrared region, $C_{\perp}(r)$ and $C_{\parallel}(r)$ strongly cancel each other, which leads to an approximate cancellation for the sum of the field-strength correlations as $\sum_{\mu,\nu} \langle G_{\mu\nu}^a(s) G_{\mu\nu}^a(s') \rangle \propto C_{\perp}(|s-s'|) + C_{\parallel}(|s-s'|) \sim 0$. Next, we decompose the perpendicular-type color-magnetic correlation $C_{\perp}(r)$ into quadratic, cubic and quartic terms of the gluon field A_{μ} . The quadratic term is always negative, which is explained by the Yukawa-type gluon propagator $\langle A_{\mu}(s) A_{\mu}(s') \rangle \propto e^{-mr}/r$ with $r \equiv |s - s'|$ in the Landau gauge. The quartic term gives a relatively small contribution. In the infrared region, the cubic term is positive and tends to cancel with the quadratic term, resulting in a small value of $C_{\perp}(r)$.

*The XVIth Quark Confinement and the Hadron Spectrum Conference (QCHSC24), 19-24 August, 2024,
Cairns Convention Centre, Cairns, Queensland, Australia*

*Speaker

1. Introduction

Since 1973, quantum chromodynamics (QCD) has been established as the fundamental theory of the strong interaction. Due to the asymptotic freedom in QCD, its coupling decreases with the renormalization scale, and perturbative QCD is applicable to the analysis of high-energy hadron reactions. At low energies, however, the coupling becomes strong, and QCD exhibits nonperturbative phenomena such as dynamical chiral symmetry breaking and color confinement [1].

In particular, due to the asymptotic freedom, QCD has a color-magnetic instability, which involves the spontaneous emergence of color-magnetic fields [2]. In other words, the system with zero color-magnetic field is energetically unstable. The non-zero color-magnetic QCD vacuum is called the Savvidy vacuum and/or the Copenhagen vacuum [3]. Actually, the gluon condensate $\frac{\alpha_s}{\pi} \langle G_{\mu\nu}^a G_a^{\mu\nu} \rangle$ is positive in the Minkowski space, implying significant excess of color-magnetic fields rather than color-electric fields. The color-magnetic system appears to be vortex-like, and the color-vortex systems form a fluctuating stochastic domain structure at a large scale.

Considering the fluctuating color fields in the QCD vacuum, Dosch and Simonov proposed the ‘‘stochastic vacuum model’’ for gauge-invariant field-strength correlators and showed that its infrared exponential damping leads to an asymptotic linear potential [4, 5]. Later, Di Giacomo et al. [6, 7] and Bali, Brambilla and Vairo [8] found that the gauge-invariant field-strength correlator exhibits infrared exponential damping in lattice QCD.

Motivated by these studies, we reconsider the field-strength correlation in QCD. We are interested in not only infrared behavior, but also in its overall behavior. In this study, using lattice QCD, we mainly investigate the color-magnetic correlation in the Landau gauge, which has many advantages in terms of symmetries and minimal gauge-field fluctuations.

2. Color-magnetic correlations in the Landau gauge

In Euclidean QCD, the Landau gauge has a global definition to minimize

$$R[A_\mu^a] \equiv \int d^4x \{A_\mu^a(x) A_\mu^a(x)\} \quad (1)$$

by the gauge transformation. In the global definition, the Landau gauge has a clear physical interpretation that it maximally suppresses artificial gauge-field fluctuations associated with the gauge degrees of freedom [9].

Taking into account the above-mentioned color-vortex structure in the QCD vacuum, we investigate the following type of color-magnetic correlation in lattice QCD in the Landau gauge:

1. Perpendicular-type color-magnetic correlation $C_\perp(r) \equiv \langle H_z^a(s) H_z^a(s + r \hat{\perp}) \rangle$
($\hat{\perp}$: lattice unit vector on the xy -plane),
2. Parallel-type color-magnetic correlation $C_\parallel(r) \equiv \langle H_z^a(s) H_z^a(s + r \hat{\parallel}) \rangle$
($\hat{\parallel}$: lattice unit vector on the zt -plane).

In the Euclidean metric, one finds $\langle H_z^a(s) H_z^a(s + r \hat{z}) \rangle = \langle H_z^a(s) H_z^a(s + r \hat{t}) \rangle$ due to the four-dimensional rotational invariance. In the Landau gauge, all two-point field-strength correlations $\langle G_{\mu\nu}^a(s) G_{\alpha\beta}^b(s') \rangle$ can be expressed with these two correlations $C_\perp(r)$ and $C_\parallel(r)$, due to the Lorentz and global $SU(N_c)$ color symmetries.

3. Lattice QCD setup

For the nonperturbative analysis of the color-magnetic correlation, we use $SU(2)$ and $SU(3)$ lattice QCD Monte Carlo calculations with the standard plaquette action at the quenched level. For the spatial correlation, we take both on-axis and off-axis lattice data. On the statistical error of the lattice data, the jackknife error estimate is adopted.

3.1 $SU(3)$ lattice QCD setup

For the $SU(3)$ lattice QCD calculations, we adopt the following three lattices:

- i) $\beta = 5.7$, $L^4 = 16^4$ (i.e., $a \simeq 0.186$ fm, $La \simeq 3.0$ fm)
- ii) $\beta = 5.8$, $L^4 = 16^4$ (i.e., $a \simeq 0.152$ fm, $La \simeq 2.4$ fm)
- iii) $\beta = 6.0$, $L^4 = 24^4$ (i.e., $a \simeq 0.104$ fm, $La \simeq 2.5$ fm)

Here, the lattice spacing a is determined so as to reproduce the string tension $\sigma = 0.89$ GeV [9, 10]. The gauge configurations are picked up with the interval of 1,000 sweeps, after the thermalization of 20,000 sweeps. In this study, 200 gauge configurations are used at $\beta=5.7$ and 5.8, and 800 gauge configurations at $\beta = 6.0$.

As for the Landau gauge fixing, we use the ordinary iterative maximization algorithm with an over-relaxation parameter of 1.6. In the Landau gauge, since the gluon field is globally minimized, we define $SU(3)$ gluon fields with the link-variable as

$$\mathcal{A}_\mu(s) \equiv \frac{1}{2iag} [U_\mu(s) - U_\mu^\dagger(s)] - \frac{1}{2iagN_c} \text{Tr}[U_\mu(s) - U_\mu^\dagger(s)] \in \text{su}(N_c) \quad (2)$$

in the fundamental representation. This definition is often used in the Landau gauge.

3.2 $SU(2)$ lattice QCD setup

For the $SU(2)$ lattice QCD calculations, we adopt the following three lattices:

- i) $\beta = 2.3$, $L^4 = 16^4$ (i.e., $a \simeq 0.18$ fm, $La \simeq 2.9$ fm)
- ii) $\beta = 2.4$, $L^4 = 24^4$ (i.e., $a \simeq 0.127$ fm, $La \simeq 3.0$ fm)
- iii) $\beta = 2.5$, $L^4 = 32^4$ (i.e., $a \simeq 0.09$ fm, $La \simeq 2.9$ fm)

The lattice spacing a is determined to reproduce the string tension $\sigma = 0.89$ GeV [11]. The gauge configurations are picked up with the interval of 200 sweeps, after the thermalization of 2,000 sweeps. We mainly use 400 gauge configurations at each β . The Landau gauge fixing is achieved by the ordinary iterative maximization algorithm with over-relaxation parameter of 1.7. In $SU(2)$ lattice QCD, the gluon field $A_\mu(s)$ is directly obtained from the link-variable $U_\mu(s) = e^{iagA_\mu(s)}$ using the general relation of $e^{i\tau^a\theta^a} = \cos\theta + i\tau^a\hat{\theta}^a \sin\theta$ with $\theta \equiv (\theta^a\theta^a)^{1/2}$ and $\hat{\theta}^a \equiv \theta^a/\theta$.

3.3 Landau-gauge gluon propagator

Before proceeding the color-magnetic correlation, we investigate the Landau-gauge gluon propagator in lattice QCD. In the Euclidean Landau gauge, all the gluon two-point functions $D_{\mu\nu}^{ab}(x-y) \equiv \langle A_\mu^a(x)A_\nu^b(y) \rangle$ are expressed with the scalar combination $\langle A_\mu^a(s)A_\mu^a(s') \rangle$, which is a single-valued function of the four-dimensional Euclidean space-time distance $r \equiv |s-s'|$. Figure 1 shows the gluon propagator in the Landau gauge in lattice QCD.




Figure 1: Landau-gauge gluon propagator $\langle A_\mu^a(s)A_\mu^a(s') \rangle$ plotted against $r \equiv |s - s'|$ in SU(2) (left) and SU(3) (right) lattice QCD. The curve is the best-fit Yukawa function. The right figure is taken from Ref. [9].

In both SU(2) and SU(3) QCD, the Landau-gauge gluon propagator is well described with a Yukawa-type function,

$$D(r) \equiv \frac{1}{3(N_c^2 - 1)} \langle A_\mu^a(s)A_\mu^a(s') \rangle \simeq D_{\text{Yukawa}}(r), \quad D_{\text{Yukawa}}(r) \equiv A \frac{m}{r} e^{-mr}, \quad r \equiv |s - s'|, \quad (3)$$

in the wide region of $r = 0.1 - 1.0$ fm. The gluonic mass parameter is estimated as $m \simeq 0.66$ GeV for SU(2) QCD and $m \simeq 0.624$ GeV for SU(3) QCD [9]. Since the Yukawa-type gluon propagation is natural in the three-dimensional space-time instead of the four-dimensional one, this might relate to some dimensional reduction hidden in nonperturbative QCD [12].

4. Lattice QCD result for color-magnetic correlations

In this section, we study the color-magnetic correlations, $C_\perp(r)$ and $C_\parallel(r)$, in the Landau gauge using SU(2) and SU(3) lattice QCD at the quenched level. Note again that all the two-point field-strength correlations $\langle G_{\mu\nu}^a(s)G_{\alpha\beta}^b(s') \rangle$ are expressed with these two correlations. In the adopted β region, $C_\perp(r)$ and $C_\parallel(r)$ are found to be approximate single-valued functions of r in lattice QCD.

4.1 Perpendicular-type color-magnetic correlation

To begin with, we investigate the perpendicular-type of the color-magnetic correlation

$$C_\perp(r) \equiv \langle H_z^a(s)H_z^a(s + r\hat{\perp}) \rangle \quad (\hat{\perp} : \text{lattice unit vector on the } xy\text{-plane}) \quad (4)$$

in the Landau gauge. Figure 2 shows the numerical result in SU(3) and SU(2) lattice QCD.

Curiously, the perpendicular-type color-magnetic correlation is *always negative*, $C_\perp(r) < 0$, except for the same-point correlation of $r = 0$, which seems to be contradict the simple vortex-like picture. Moreover, an “always negative” correlation would be rare in physics, while “always positive correlation” and “alternating correlation” have been observed in various areas of physics.

One might suspect that the gauge fixing has some unphysical effect. Then, we also examine gauge-invariant field-strength correlation extracted from the plaquette correlators, as shown in Fig. 3, and obtain a similar result. Indeed, the correlation corresponding to the perpendicular-type color-magnetic correlation is always negative.




Figure 2: The perpendicular-type color-magnetic correlation $C_{\perp}(r) \equiv \langle H_z^a(s) H_z^a(s + r \hat{\perp}) \rangle$ ($\hat{\perp} \equiv x, y$) in the Landau gauge in $SU(3)$ (left) and $SU(2)$ (right) lattice QCD.



Figure 3: An example of the plaquette correlator to extract the gauge-invariant field-strength correlation in lattice QCD.

4.2 Parallel-type color-magnetic correlation

Next, we show in Fig. 4 the parallel-type color-magnetic correlation in the Landau gauge,

$$C_{\parallel}(r) \equiv \langle H_z^a(s) H_z^a(s + r \hat{\parallel}) \rangle \quad (\hat{\parallel} : \text{lattice unit vector on the } zt\text{-plane}), \quad (5)$$

in $SU(3)$ and $SU(2)$ lattice QCD, together with the perpendicular-type one $C_{\perp}(r)$ and the sum of $C_{\perp}(r) + C_{\parallel}(r)$. The parallel-type color-magnetic correlation $C_{\parallel}(r)$ is always positive, $C_{\parallel}(r) > 0$.




Figure 4: The parallel-type color-magnetic correlation $C_{\parallel}(r) \equiv \langle H_z^a(s) H_z^a(s + r \hat{\parallel}) \rangle$ ($\hat{\parallel} \equiv z, t$) in the Landau gauge in $SU(3)$ (left) and $SU(2)$ (right) lattice QCD. The perpendicular-type color-magnetic correlation $C_{\perp}(r)$ (gray) and the sum of $C_{\perp}(r) + C_{\parallel}(r)$ (black) are also shown.

In the infrared region, the parallel-type color-magnetic correlation $C_{\parallel}(r)$ strongly cancels with the perpendicular-type one as

$$C_{\parallel}(r) \sim -C_{\perp}(r), \quad \text{e.g., } \langle H_z^a(s) H_z^a(s + r \hat{z}) \rangle \sim -\langle H_z^a(s) H_z^a(s + r \hat{x}) \rangle, \quad (6)$$

which leads to an approximate cancellation for the sum of Landau-gauge field-strength correlations,

$$\sum_{\mu,\nu} \langle G_{\mu\nu}^a(s) G_{\mu\nu}^a(s') \rangle = 6 [C_{\perp}(r) + C_{\parallel}(r)] \sim 0. \quad (7)$$

5. Analysis for color-magnetic correlations in QCD

In this section, we try to analyze the lattice QCD result of the color-magnetic correlation in the Landau gauge, particularly considering the origin of the negative correlation of $C_{\perp}(r) < 0$.

5.1 Decomposition of the field-strength correlation in terms of the gluon field

We decompose the field-strength correlation $\langle G_{\mu\nu}^a(s) G_{\alpha\beta}^a(s') \rangle$ into three parts, i.e., quadratic, cubic and quartic terms of the gluon field A_{μ} :

$$\langle G_{\mu\nu}^a(s) G_{\alpha\beta}^a(s') \rangle = \langle G_{\mu\nu}^a(s) G_{\alpha\beta}^a(s') \rangle_{\text{quad}} + \langle G_{\mu\nu}^a(s) G_{\alpha\beta}^a(s') \rangle_{\text{cubic}} + \langle G_{\mu\nu}^a(s) G_{\alpha\beta}^a(s') \rangle_{\text{quartic}}. \quad (8)$$

Here, the quadratic, cubic and quartic terms are defined by

$$\begin{aligned} \langle G_{\mu\nu}^a(s) G_{\alpha\beta}^a(s') \rangle_{\text{quad}} &\equiv \langle \text{Tr} (\partial_{\mu} A_{\nu} - \partial_{\nu} A_{\mu})(s) (\partial_{\alpha} A_{\beta} - \partial_{\beta} A_{\alpha})(s') \rangle, \\ \langle G_{\mu\nu}^a(s) G_{\alpha\beta}^a(s') \rangle_{\text{cubic}} &\equiv ig \langle \text{Tr} (\partial_{\mu} A_{\nu} - \partial_{\nu} A_{\mu})(s) [A_{\alpha}, A_{\beta}](s') \rangle \\ &\quad + ig \langle \text{Tr} (\partial_{\alpha} A_{\beta} - \partial_{\beta} A_{\alpha})(s) [A_{\mu}, A_{\nu}](s') \rangle, \\ \langle G_{\mu\nu}^a(s) G_{\alpha\beta}^a(s') \rangle_{\text{quartic}} &\equiv -g^2 \langle \text{Tr} [A_{\mu}, A_{\nu}](s) [A_{\alpha}, A_{\beta}](s') \rangle. \end{aligned} \quad (9)$$

Among the three terms, the quadratic term can be directly expressed with the gluon propagator $D_{\mu\nu}^{ab}(x-y) \equiv \langle A_{\mu}^a(x) A_{\nu}^b(y) \rangle$ as

$$\begin{aligned} \langle G_{\mu\nu}^a(x) G_{\alpha\beta}^b(y) \rangle_{\text{quad}} \\ = \partial_{\mu}^x \partial_{\alpha}^y D_{\nu\beta}^{ab}(x-y) - \partial_{\mu}^x \partial_{\beta}^y D_{\nu\alpha}^{ab}(x-y) - \partial_{\nu}^x \partial_{\alpha}^y D_{\mu\beta}^{ab}(x-y) + \partial_{\nu}^x \partial_{\beta}^y D_{\mu\alpha}^{ab}(x-y). \end{aligned} \quad (10)$$

In the Landau gauge, due to the Lorentz symmetry, we only have to consider scalar combination of the gluon propagator $\langle A_{\mu}^a(x) A_{\mu}^a(y) \rangle$ as a function of four-dimensional Euclidean distance $r = |x-y|$.

5.2 Decomposition of perpendicular-type color-magnetic correlation in the Landau gauge

For the color-magnetic correlation,

$$\begin{aligned} \langle H_z^a(s) H_z^a(s') \rangle &= \langle \text{Tr} (\partial_x A_y - \partial_y A_x)(s) (\partial_x A_y - \partial_y A_x)(s') \rangle \\ &\quad + 2ig \langle \text{Tr} (\partial_x A_y - \partial_y A_x)(s) [A_x, A_y](s') \rangle \\ &\quad - g^2 \langle \text{Tr} [A_x, A_y](s) [A_y, A_x](s') \rangle, \end{aligned} \quad (11)$$

the quadratic term can be described with the gluon propagator. Using the Yukawa-type gluon propagator $D_{\text{Yukawa}}(r)$ in Eq. (3) in the Landau gauge, we find that the quadratic term of the perpendicular-type color-magnetic correlation becomes always negative:

$$\begin{aligned} \langle H_z^a(r \hat{\perp}) H_z^a(0) \rangle_{\text{quad}} &= \langle \text{Tr} (\partial_x A_y - \partial_y A_x)(r \hat{\perp}) (\partial_x A_y - \partial_y A_x)(0) \rangle \\ &= -(N_c^2 - 1) Am^4 \frac{e^{-mr}}{mr} \left(1 + \frac{1}{mr} + \frac{1}{m^2 r^2} \right) < 0 \quad (\perp \equiv x, y). \end{aligned} \quad (12)$$

Then, if the quadratic term is dominant, the negative behavior of the perpendicular-type color-magnetic correlation, $C_{\perp}(r) < 0$, can be explained. However, the real situation is not so simple.

Figure 5 shows individual contributions of the quadratic, cubic and quartic terms of the perpendicular-type color-magnetic correlation $C_{\perp}(r)$ in lattice QCD. The quadratic term is always negative, as was deduced with the Yukawa-type gluon propagator in the Landau gauge. The cubic term is comparable to the quadratic term, while the quartic term gives a relatively small contribution. In the infrared region, the cubic term is positive and tends to cancel with the quadratic term, resulting in a small value of $C_{\perp}(r)$.




Figure 5: Each contribution of the quadratic (red), cubic (blue) and quartic (green) terms of the perpendicular-type color-magnetic correlation $C_{\perp}(r)$ (black) in the Landau gauge in $SU(3)$ (left) and $SU(2)$ (right) lattice QCD.

6. Summary and Conclusion

To examine color-magnetic instabilities in QCD, we have studied field-strength correlations in both $SU(2)$ and $SU(3)$ lattice QCD. In the Euclidean Landau gauge, we have numerically calculated the perpendicular-type color-magnetic correlation, $C_{\perp}(r) \equiv \langle H_z^a(s) H_z^a(s + r \hat{\perp}) \rangle$ ($\perp \equiv x, y$), and the parallel-type one, $C_{\parallel}(r) \equiv \langle H_z^a(s) H_z^a(s + r \hat{\parallel}) \rangle$ ($\parallel \equiv z, t$).

Curiously, we have found that the perpendicular-type color-magnetic correlation $C_{\perp}(r)$ is always negative for arbitrary r , except for the same-point correlation. We have found that the parallel-type color-magnetic correlation $C_{\parallel}(r)$ is always positive. In the infrared region, $C_{\perp}(r)$ and $C_{\parallel}(r)$ strongly cancel each other, which leads to an approximate cancellation for the sum of the field-strength correlations as $\sum_{\mu, \nu} \langle G_{\mu\nu}^a(s) G_{\mu\nu}^a(s') \rangle \propto C_{\perp}(|s - s'|) + C_{\parallel}(|s - s'|) \sim 0$.

Next, we have decomposed the perpendicular-type color-magnetic correlation $C_{\perp}(r)$ into the quadratic, cubic and quartic terms of the gluon field A_{μ} . The quadratic term is always negative, which can be explained with the Yukawa-type gluon propagator in the Landau gauge. The quartic term gives a relatively small contribution. In the infrared region, the cubic term is positive and tends to cancel with the quadratic term, resulting in small $C_{\perp}(r)$.

Finally, we consider the negativity of the perpendicular-type color-magnetic correlation. In abelian gauge theories, the negativity of the perpendicular-type magnetic correlation can be explained by the magnetic-flux conservation, although the ordinary QED vacuum is trivial. In non-abelian gauge theories, however, such an explanation cannot be applied, and the negative color-magnetic correlation of $C_{\perp}(r) < 0$ remains to be a puzzle in QCD.

Acknowledgment

We thank Professor Jeff Greensite for his useful comment and discussion on the case of abelian gauge theories. H.S. is supported in part by the Grants-in-Aid for Scientific Research [19K03869] from Japan Society for the Promotion of Science. K.T. is supported by SPRING Program at Kyoto University. The lattice QCD calculations have been performed by SQUID at Osaka University.

References

- [1] For a recent review article, H. SUGANUMA, *Quantum Chromodynamics, Quark Confinement and Chiral Symmetry Breaking*, "Handbook of Nuclear Physics", (Springer, 2023) 22-1.
- [2] G.K. SAVVIDY, *Infrared Instability of the Vacuum State of Gauge Theories and Asymptotic Freedom*, Phys. Lett. **B71**, 133-134 (1977).
- [3] J. AMBJORN and P. OLESEN, *On the Formation of a Random Color Magnetic Quantum Liquid in QCD*, Nucl. Phys. **B170**, 265-282 (1980).
- [4] H.G. DOSCH, *Gluon Condensate and Effective Linear Potential*, Phys. Lett. **B190**, 177-181 (1987); Yu. A. SIMONOV, *Vacuum Background Fields in QCD as a Source of Confinement*, Nucl. Phys. **B307**, 512-530 (1988).
- [5] H.G. DOSCH and Yu. A. SIMONOV, *The Area Law of the Wilson Loop and Vacuum Field Correlators*, Phys. Lett. **B205**, 339-344 (1988).
- [6] A. DI GIACOMO and H. PANAGOPoulos, *Field Strength Correlations in the QCD Vacuum*, Phys. Lett. **B285**, 133-136 (1992).
- [7] M. D'ELIA, A. DI GIACOMO and E. MEGGIOLARO, *Field Strength Correlators in Full QCD*, Phys. Lett. **B408**, 315-319, (1997); *Gauge Invariant Field Strength Correlators in Pure Yang-Mills and Full QCD at Finite Temperature*, Phys. Rev. **D67**, 114504 (2003).
- [8] G.S. BALI, N. BRAMBILLA and A. VAIRO, *A Lattice Determination of QCD Field Strength Correlators*, Phys. Lett. **B421**, 265-272 (1998).
- [9] T. IRITANI, H. SUGANUMA and H. IIDA, *Gluon-propagator Functional Form in the Landau Gauge in $SU(3)$ Lattice QCD: Yukawa-type Gluon Propagator and Anomalous Gluon Spectral Function*, Phys. Rev. **D80**, 114505 (2009).
- [10] T.T. TAKAHASHI, H. SUGANUMA, Y. NEMOTO and H. MATSUFURU, *Detailed Analysis of the Three-Quark Potential in $SU(3)$ lattice QCD*, Phys. Rev. **D65**, 114509 (2002).
- [11] K. AMEMIYA and H. SUGANUMA, *Off-diagonal Gluon Mass Generation and Infrared Abelian Dominance in the Maximally Abelian Gauge in Lattice QCD*, Phys. Rev. **D60**, 114509 (1999).
- [12] K. TOHME and H. SUGANUMA, *Dimensional Reduction Gauge and Effective Dimensional Reduction in Four-dimensional Yang-Mills Theory*, Phys. Rev. **D110**, 034505 (2024).